

## Supplement 1 of

Schultz et al., 2017,

“The Chemistry Climate Model ECHAM-HAMMOZ”

in Geosci. Model Dev. (2017), available at:

<http://www.geosci-model-dev.net>

Date: August 15, 2017

The supplement provides:

The Chemical Mechanism of HAMMOZ

List of Henry coefficients

# The Chemical Mechanism of ECHAM6.3-HAM2.3-MOZ1.0

JAM version: 002b

Number of reactions:

Tropospheric photolysis (S1):	97
Stratospheric photolysis (S2):	45
(Tropospheric) O <sub>x</sub> (S3):	7
(Tropospheric) HO <sub>x</sub> (S4):	17
NO <sub>x</sub> (S5):	25
C1 oxidation (S6):	20
C2 oxidation (S7):	48
C3 oxidation (S8):	31
C4 oxidation (S9):	63
C5 oxidation (S10):	137
C6 oxidation (S11):	31
C7 oxidation (S12):	20
C8 oxidation (S13):	13
C10/C15 oxidation (terpenes, sesqui-terpenes; S14):	36
Tropospheric halogen + organics (S15):	21
Sulfur (S16):	11
Stratospheric O(1D) (S17):	28
Stratospheric inorganic halogen (S18):	48
Stratospheric organic halogen (S19):	12
(Tropospheric) heterogeneous (S20):	7
Stratospheric Sulfate aerosol (S21):	6
Stratospheric Nitric acid dihydrate (S22):	5
Stratospheric Ice aerosol (S23):	6

---

All equations:	734
----------------	-----

Table S1: Tropospheric photolysis reactions

reaction	reference
$\text{O}_3 + h\nu \longrightarrow \text{O}_1\text{D} + \text{O}_2$	
$\text{O}_3 + h\nu \longrightarrow \text{O} + \text{O}_2$	
$\text{H}_2\text{O}_2 + h\nu \longrightarrow 2 \cdot \text{OH}$	
$\text{N}_2\text{O} + h\nu \longrightarrow \text{N}_2 + \text{O}_1\text{D}$	
$\text{NO} + h\nu \longrightarrow \text{N} + \text{O}$	
$\text{NO}_2 + h\nu \longrightarrow \text{NO} + \text{O}$	
$\text{NO}_3 + h\nu \longrightarrow \text{NO}_2 + \text{O}$	
$\text{NO}_3 + h\nu \longrightarrow \text{NO} + \text{O}_2$	
$\text{HNO}_3 + h\nu \longrightarrow \text{NO}_2 + \text{OH}$	
$\text{HONO} + h\nu \longrightarrow \text{NO} + \text{OH}$	
$\text{HO}_2\text{NO}_2 + h\nu \longrightarrow \text{NO}_3 + \text{OH}$	
$\text{HO}_2\text{NO}_2 + h\nu \longrightarrow \text{HO}_2 + \text{NO}_2$	
$\text{N}_2\text{O}_5 + h\nu \longrightarrow \text{NO}_2 + \text{NO}_3$	
$\text{N}_2\text{O}_5 + h\nu \longrightarrow \text{NO} + \text{O} + \text{NO}_3$	
$\text{CO}_2 + h\nu \longrightarrow \text{CO} + \text{O}$	
$\text{CH}_4 + h\nu \longrightarrow \text{CH}_3\text{O}_2 + \text{H}$	
$\text{CH}_4 + h\nu \longrightarrow 1.44 \cdot \text{H}_2 + 0.18 \cdot \text{CH}_2\text{O} + 0.18 \cdot \text{O} + 0.66 \cdot \text{OH} + 0.44 \cdot \text{CO}_2 + 0.38 \cdot \text{CO} + 0.05 \cdot \text{H}_2\text{O}$	
$\text{CH}_2\text{O} + h\nu \longrightarrow \text{CO} + 2 \cdot \text{H}$	
$\text{CH}_2\text{O} + h\nu \longrightarrow \text{CO} + \text{H}_2$	
$\text{CH}_3\text{OOH} + h\nu \longrightarrow \text{CH}_2\text{O} + \text{H} + \text{OH}$	
$\text{CH}_3\text{O}_2\text{NO}_2 + h\nu \longrightarrow \text{HO}_2 + \text{NO}_3 + \text{HCHO}$	
$\text{CH}_3\text{O}_2\text{NO}_2 + h\nu \longrightarrow \text{CH}_3\text{O}_2 + \text{NO}_2$	
$\text{CH}_3\text{CHO} + h\nu \longrightarrow \text{CH}_3\text{O}_2 + \text{CO} + \text{HO}_2$	
$\text{CH}_3\text{COOOH} + h\nu \longrightarrow \text{CH}_3\text{O}_2 + \text{OH} + \text{CO}_2$	
$\text{C}_2\text{H}_5\text{OOH} + h\nu \longrightarrow \text{CH}_3\text{CHO} + \text{HO}_2 + \text{OH}$	
$\text{PAN} + h\nu \longrightarrow 0.6 \cdot \text{CH}_3\text{CO}_3 + 0.6 \cdot \text{NO}_2 + 0.4 \cdot \text{CH}_3\text{O}_2 + 0.4 \cdot \text{NO}_3 + 0.4 \cdot \text{CO}_2$	
$\text{EOOH} + h\nu \longrightarrow \text{EO} + \text{OH}$	
$\text{GLYOXAL} + h\nu \longrightarrow 2 \cdot \text{CO} + 2 \cdot \text{HO}_2$	
$\text{GLYALD} + h\nu \longrightarrow 2 \cdot \text{HO}_2 + \text{CO} + \text{CH}_2\text{O}$	
$\text{HOCH}_2\text{CO}_3\text{H} + h\nu \longrightarrow \text{CH}_2\text{O} + \text{HO}_2 + \text{OH} + \text{CO}_2$	
$\text{HCOCO}_2\text{H} + h\nu \longrightarrow 2 \cdot \text{HO}_2 + \text{CO} + \text{CO}_2$	
$\text{HCOCO}_3\text{H} + h\nu \longrightarrow \text{CO} + \text{HO}_2 + \text{OH} + \text{CO}_2$	
$\text{CH}_3\text{COCH}_3 + h\nu \longrightarrow \text{CH}_3\text{CO}_3 + \text{CH}_3\text{O}_2$	

Table S1: Tropospheric photolysis reactions (... continued)

reaction	reference
$\text{C}_3\text{H}_7\text{OOH} + h\nu \longrightarrow 0.82 \cdot \text{CH}_3\text{COCH}_3 + \text{OH} + \text{HO}_2 + 0.27 \cdot \text{CH}_3\text{CHO}$	
$\text{POOH} + h\nu \longrightarrow \text{CH}_2\text{O} + \text{CH}_3\text{CHO} + \text{HO}_2 + \text{OH}$	
$\text{HYAC} + h\nu \longrightarrow \text{CH}_3\text{CO}_3 + \text{HO}_2 + \text{CH}_2\text{O}$	
$\text{CH}_3\text{COCHO} + h\nu \longrightarrow \text{CH}_3\text{CO}_3 + \text{CO} + \text{HO}_2$	
$\text{ROOH} + h\nu \longrightarrow \text{CH}_2\text{O} + \text{CH}_3\text{CO}_3 + \text{OH}$	
$\text{PR}_2\text{O}_2\text{HNO}_3 + h\nu \longrightarrow 0.83 \cdot \text{HO}_2 + 0.83 \cdot \text{NOA} + 0.17 \cdot \text{CH}_2\text{O} + 0.17 \cdot \text{CH}_3\text{CHO} + \text{OH}$	
$\text{NOA} + h\nu \longrightarrow \text{CH}_2\text{O} + \text{CH}_3\text{CO}_3 + \text{NO}_2$	Müller et al. (2014)
$\text{MEK} + h\nu \longrightarrow \text{C}_2\text{H}_5\text{O}_2 + \text{CH}_3\text{CO}_3$	
$\text{MEKOOH} + h\nu \longrightarrow \text{CH}_3\text{CO}_3 + \text{OH} + \text{CH}_3\text{CHO}$	
$\text{MEKNO}_3 + h\nu \longrightarrow \text{CH}_3\text{CHO} + \text{CH}_3\text{CO}_3 + \text{NO}_2$	Müller et al. (2014)
$\text{MACR} + h\nu \longrightarrow \text{HO}_2 + 0.5 \cdot \text{MCO}_3 + 0.5 \cdot \text{CH}_2\text{O} + 0.5 \cdot \text{CH}_3\text{CO}_3 + 0.5 \cdot \text{CO}$	
$\text{MACR} + h\nu \longrightarrow \text{HO}_2 + 0.5 \cdot \text{MCO}_3 + 0.5 \cdot \text{CH}_2\text{O} + 0.5 \cdot \text{CH}_3\text{CO}_3 + 0.5 \cdot \text{CO}$	
$\text{MACROOH} + h\nu \longrightarrow \text{HO}_2 + \text{HYAC} + \text{OH} + \text{CO}$	
$\text{MACROH} + h\nu \longrightarrow \text{CO} + \text{HYAC} + 2 \cdot \text{HO}_2 + \text{H}_2\text{O}$	
$\text{MPAN} + h\nu \longrightarrow \text{MCO}_3 + \text{NO}_2$	
$\text{MACO}_3\text{H} + h\nu \longrightarrow \text{CH}_2\text{O} + \text{CH}_3\text{CO}_3 + \text{OH} + \text{CO}_2$	
$\text{MVK} + h\nu \longrightarrow 0.5 \cdot \text{C}_3\text{H}_6 + 0.5 \cdot \text{CH}_3\text{CO}_3 + 0.5 \cdot \text{CH}_2\text{O} + \text{CO} + 0.5 \cdot \text{HO}_2$	
$\text{LHMKABOOH} + h\nu \longrightarrow 0.3 \cdot \text{CH}_3\text{COCHO} + \text{OH} + 0.3 \cdot \text{CH}_2\text{O} + 0.3 \cdot \text{HO}_2 + 0.7 \cdot \text{CH}_3\text{CO}_3 + 0.7 \cdot \text{GLYALD}$	
$\text{MVKN} + h\nu \longrightarrow \text{CH}_3\text{CO}_3 + \text{GLYALD} + \text{NO}_2$	Müller et al. (2014)
$\text{MACRN} + h\nu \longrightarrow \text{CO} + \text{HYAC} + \text{HO}_2 + \text{NO}_2$	Müller et al. (2014)
$\text{CO}_2\text{H}_3\text{CHO} + h\nu \longrightarrow \text{CH}_3\text{COCHO} + \text{CO} + 2 \cdot \text{HO}_2$	
$\text{CO}_2\text{H}_3\text{CO}_3\text{H} + h\nu \longrightarrow \text{CH}_3\text{COCHO} + \text{HO}_2 + \text{OH} + \text{CO}_2$	
$\text{BIACETOH} + h\nu \longrightarrow \text{CH}_3\text{CO}_3 + \text{HOCH}_2\text{CO}_3$	
$\text{ALKOOH} + h\nu \longrightarrow 0.4 \cdot \text{CH}_3\text{CHO} + 0.25 \cdot \text{CH}_2\text{O} + 0.25 \cdot \text{CH}_3\text{COCH}_3 + \text{HO}_2 + 0.8 \cdot \text{MEK} + \text{OH}$	
$\text{ALKNO}_3 + h\nu \longrightarrow 0.4 \cdot \text{CH}_3\text{CHO} + 0.25 \cdot \text{CH}_2\text{O} + 0.25 \cdot \text{CH}_3\text{COCH}_3 + \text{HO}_2 + 0.8 \cdot \text{MEK} + \text{NO}_2$	
$\text{LISOPACOOH} + h\nu \longrightarrow \text{HO}_2 + \text{LHC}_4\text{ACCHO} + \text{OH}$	
$\text{LISOPACNO}_3 + h\nu \longrightarrow \text{HO}_2 + \text{LHC}_4\text{ACCHO} + \text{NO}_2$	
$\text{HPALD} + h\nu \longrightarrow \text{LHC}_4\text{ACCO}_3 + \text{OH}$	D. Taraborrelli: $J(\text{MACR})/\phi(\text{MACR}) = 2 \cdot j_{\text{macr\_a}}/0.004 = 500 \cdot j_{\text{macr\_a}}$
$\text{PACALD} + h\nu \longrightarrow \text{OH} + 0.5 \cdot \text{HO}_2 + 0.5 \cdot \text{CO} + 0.5 \cdot \text{CH}_3\text{COCHO} + 0.5 \cdot \text{GLYOX} + 0.5 \cdot \text{CH}_3\text{CO}_3$	D. Taraborrelli: average of product yields of C5PACALD 1 and 2 in MCMv3.3.1, $2 \cdot J(\text{MACR})/\phi(\text{MACR}) = 2 \cdot j_{\text{hpald}} = 1000 \cdot j_{\text{macr\_a}}$

Table S1: Tropospheric photolysis reactions (... continued)

reaction	reference
$\text{LIECHO} + h\nu \longrightarrow \text{CO} + \text{HO}_2 + 0.6 \cdot \text{LHMKABO}_2 + 0.4 \cdot \text{MACRO}_2$	
$\text{LIECO}_3\text{H} + h\nu \longrightarrow 0.6 \cdot \text{LHMKABO}_2 + 0.4 \cdot \text{MACRO}_2 + \text{CO}_2 + \text{OH}$	
$\text{ISOPBOOH} + h\nu \longrightarrow \text{CH}_2\text{O} + \text{MVK} + \text{HO}_2 + \text{OH}$	
$\text{ISOPBNO}_3 + h\nu \longrightarrow \text{CH}_2\text{O} + \text{MVK} + \text{HO}_2 + \text{NO}_2$	
$\text{ISOPDOOH} + h\nu \longrightarrow \text{CH}_2\text{O} + \text{MACR} + \text{HO}_2 + \text{OH}$	
$\text{ISOPDNO}_3 + h\nu \longrightarrow \text{CH}_2\text{O} + \text{MACR} + \text{HO}_2 + \text{NO}_2$	
$\text{NISOPOOH} + h\nu \longrightarrow \text{HO}_2 + \text{NC}_4\text{CHO} + \text{OH}$	
$\text{NC}_4\text{CHO} + h\nu \longrightarrow \text{LHC}_4\text{ACCO}_3 + \text{NO}_2$	Müller et al. (2014)
$\text{LNISOOH} + h\nu \longrightarrow \text{NOA} + \text{OH} + 0.5 \cdot \text{GLYOXAL} + 0.5 \cdot \text{CO} + \text{HO}_2 + 0.5 \cdot \text{CO}_2$	
$\text{LHC}_4\text{ACCHO} + h\nu \longrightarrow 0.5 \cdot \text{LHC}_4\text{ACCO}_3 + 0.25 \cdot \text{HYAC} + 0.25 \cdot \text{GLYALD} + 0.25 \cdot \text{CH}_3\text{CO}_3 + 0.75 \cdot \text{CO} + 1.25 \cdot \text{HO}_2$	
$\text{LC}_{578}\text{OOH} + h\nu \longrightarrow 0.5 \cdot \text{HYAC} + 0.5 \cdot \text{CH}_3\text{COCHO} + 0.5 \cdot \text{GLYOXAL} + 0.5 \cdot \text{GLYALD} + \text{HO}_2 + \text{OH}$	
$\text{LHC}_4\text{ACCO}_3\text{H} + h\nu \longrightarrow 0.5 \cdot \text{HYAC} + 0.5 \cdot \text{GLYALD} + 0.5 \cdot \text{CH}_3\text{CO}_3 + 0.5 \cdot \text{CO} + 0.5 \cdot \text{HO}_2 + \text{OH} + \text{CO}_2$	
$\text{HCOC}_5 + h\nu \longrightarrow \text{CH}_2\text{O} + \text{CH}_3\text{CO}_3 + \text{HOCH}_2\text{CO}_3$	
$\text{C}_{59}\text{OOH} + h\nu \longrightarrow \text{HOCH}_2\text{CO}_3 + \text{HYAC} + \text{NO}_2 + \text{OH}$	
$\text{LISOPOOHOOH} + h\nu \longrightarrow 0.25 \cdot \text{CH}_3\text{COCHO} + 0.25 \cdot \text{GLYALD} + 0.25 \cdot \text{GLYOXAL} + 0.25 \cdot \text{HYAC} + 0.25 \cdot \text{CO} + 0.25 \cdot \text{MACROH} + 0.25 \cdot \text{CH}_2\text{O} + 0.25 \cdot \text{CO}_2\text{H}_3\text{CHO} + \text{HO}_2$	
$\text{LISOPNO}_3\text{OOH} + h\nu \longrightarrow \text{HOCH}_2\text{CO}_3 + \text{HYAC} + \text{NO}_2 + \text{OH}$	
$\text{LISOPNO}_3\text{NO}_3 + h\nu \longrightarrow \text{HOCH}_2\text{CO}_3 + \text{HYAC} + \text{NO}_2 + \text{NO}_2$	
$\text{MBOOH} + h\nu \longrightarrow \text{HO}_2 + \text{OH} + 0.67 \cdot \text{GLYALD} + 0.67 \cdot \text{CH}_3\text{COCH}_3 + 0.33 \cdot \text{IBUTALOH} + 0.33 \cdot \text{CH}_2\text{O}$	
$\text{IBUTALOH} + h\nu \longrightarrow 2 \cdot \text{HO}_2 + \text{CO} + \text{CH}_3\text{COCH}_3$	
$\text{IBUTALOOH} + h\nu \longrightarrow \text{CH}_3\text{COCH}_3 + \text{OH} + \text{CO}_2 + \text{HO}_2$	
$\text{BEPOMUC} + h\nu \longrightarrow \text{BIGALD}_1 + 1.5 \cdot \text{HO}_2 + 1.5 \cdot \text{CO}$	
$\text{BIGALD}_1 + h\nu \longrightarrow 0.6 \cdot \text{MALO}_2 + \text{HO}_2$	
$\text{TOLOOH} + h\nu \longrightarrow \text{OH} + 0.6 \cdot \text{GLYOXAL} + 0.4 \cdot \text{CH}_3\text{COCHO} + \text{HO}_2 + 0.2 \cdot \text{BIGALD}_1 + 0.2 \cdot \text{BIGALD}_2 + 0.2 \cdot \text{BIGALD}_3$	
$\text{TEPOMUC} + h\nu \longrightarrow 0.5 \cdot \text{CH}_3\text{CO}_3 + \text{HO}_2 + 1.5 \cdot \text{CO}$	
$\text{CATEC}_1\text{OOH} + h\nu \longrightarrow \text{CATEC}_1\text{O} + \text{OH}$	
$\text{BIGALD}_2 + h\nu \longrightarrow 0.6 \cdot \text{DICARBO}_2 + 0.6 \cdot \text{HO}_2$	
$\text{BIGALD}_3 + h\nu \longrightarrow 0.6 \cdot \text{CO} + 0.6 \cdot \text{HO}_2 + 0.6 \cdot \text{MDIALO}_2$	
$\text{BIGALD}_4 + h\nu \longrightarrow \text{CO} + \text{HO}_2 + \text{CH}_3\text{COCHO} + \text{CH}_3\text{CO}_3$	
$\text{TERPOOH} + h\nu \longrightarrow 0.4 \cdot \text{CH}_2\text{O} + 0.05 \cdot \text{CH}_3\text{COCH}_3 + 0.945 \cdot \text{TERPROD}_1 + \text{HO}_2 + \text{OH}$	

Table S1: Tropospheric photolysis reactions (... continued)

reaction	reference
$\text{TERPROD}_1 + h\nu \longrightarrow \text{CO} + \text{HO}_2 + \text{TERPROD}_2$	
$\text{TERP}_2\text{OOH} + h\nu \longrightarrow \text{OH} + 0.372 \cdot \text{CH}_2\text{O} + 0.3 \cdot \text{CH}_3\text{COCH}_3 + 0.25 \cdot \text{CO} + \text{CO}_2 +$	
$\text{TERPROD}_2 + \text{HO}_2 + 0.25 \cdot \text{GLYALD}$	
$\text{TERPROD}_2 + h\nu \longrightarrow 0.15 \cdot \text{CH}_3\text{COCH}_2\text{O}_2 + 0.68 \cdot \text{CH}_2\text{O} + 0.8 \cdot \text{CO}_2 + 0.5 \cdot \text{CH}_3\text{COCH}_3 +$	
$1.2 \cdot \text{HO}_2 + 1.7 \cdot \text{CO}$	
$\text{ISOPBNO}_3 + h\nu \longrightarrow \text{HO}_2 + \text{NO}_2 + \text{TERPROD}_1$	
$\text{NTERPNO}_3 + h\nu \longrightarrow \text{NO}_2 + \text{OH} + \text{TERPROD}_1$	
$\text{ELVOC} + h\nu \longrightarrow \text{HO}_2 + \text{OH} + \text{TERPROD}_2$	

Table S2: Stratospheric photolysis reactions

reaction	reference
$\text{O}_2 + h\nu \longrightarrow \text{O} + \text{O}_1\text{D}$	
$\text{O}_2 + h\nu \longrightarrow 2 \cdot \text{O}$	
$\text{H}_2\text{O} + h\nu \longrightarrow \text{H} + \text{OH}$	
$\text{H}_2\text{O} + h\nu \longrightarrow \text{H}_2 + \text{O}_1\text{D}$	
$\text{H}_2\text{O} + h\nu \longrightarrow 2 \cdot \text{H} + \text{O}$	
$\text{CL}_2 + h\nu \longrightarrow 2 \cdot \text{CL}$	
$\text{CL}_2\text{O}_2 + h\nu \longrightarrow 2 \cdot \text{CL}$	
$\text{CLO} + h\nu \longrightarrow \text{CL} + \text{O}$	
$\text{HCL} + h\nu \longrightarrow \text{CL} + \text{H}$	
$\text{HOCL} + h\nu \longrightarrow \text{CL} + \text{OH}$	
$\text{CLONO}_2 + h\nu \longrightarrow \text{CL} + \text{NO}_3$	
$\text{CLONO}_2 + h\nu \longrightarrow \text{CLO} + \text{NO}_2$	
$\text{OCLO} + h\nu \longrightarrow \text{CLO} + \text{O}$	
$\text{BRO} + h\nu \longrightarrow \text{BR} + \text{O}$	
$\text{HBR} + h\nu \longrightarrow \text{BR} + \text{H}$	
$\text{HOBR} + h\nu \longrightarrow \text{BR} + \text{OH}$	
$\text{BRONO} + h\nu \longrightarrow \text{BR} + \text{NO}_2$	50% branching ratio assigned to both possible channels. Cross-sections consistent with Burkholder and Orlando (2000)
$\text{BRONO} + h\nu \longrightarrow \text{BRO} + \text{NO}$	50% branching ratio assigned to both possible channels. Cross-sections consistent with Burkholder and Orlando (2000)
$\text{BRNO}_2 + h\nu \longrightarrow \text{BR} + \text{NO}_2$	
$\text{BRONO}_2 + h\nu \longrightarrow \text{BR} + \text{NO}_3$	

Table S2: Stratospheric photolysis reactions (... continued)

reaction	reference
$\text{BRONO}_2 + \text{h}\nu \longrightarrow \text{BRO} + \text{NO}_2$	
$\text{BR}_2 + \text{h}\nu \longrightarrow 2 \cdot \text{BR}$	
$\text{BRCL} + \text{h}\nu \longrightarrow \text{BR} + \text{CL}$	
$\text{HF} + \text{h}\nu \longrightarrow \text{F} + \text{H}$	
$\text{SF}_6 + \text{h}\nu \longrightarrow$	
$\text{CH}_3\text{BR} + \text{h}\nu \longrightarrow \text{BR} + \text{CH}_3\text{O}_2$	
$\text{CH}_2\text{BR}_2 + \text{h}\nu \longrightarrow 2 \cdot \text{BR}$	
$\text{CHBR}_3 + \text{h}\nu \longrightarrow 3 \cdot \text{BR}$	
$\text{CH}_3\text{CL} + \text{h}\nu \longrightarrow \text{CH}_3\text{O}_2 + \text{CL}$	
$\text{CH}_3\text{CCL}_3 + \text{h}\nu \longrightarrow 3 \cdot \text{CL}$	
$\text{CF}_3\text{BR} + \text{h}\nu \longrightarrow \text{BR} + \text{F} + \text{COF}_2$	
$\text{CF}_2\text{CLBR} + \text{h}\nu \longrightarrow \text{BR} + \text{CL} + \text{COF}_2$	
$\text{CCL}_4 + \text{h}\nu \longrightarrow 4 \cdot \text{CL} + \text{CO}_2$	
$\text{CFC}_{11} + \text{h}\nu \longrightarrow 2 \cdot \text{CL} + \text{COFCL}$	
$\text{CFC}_{12} + \text{h}\nu \longrightarrow 2 \cdot \text{CL} + \text{COF}_2$	
$\text{CFC}_{113} + \text{h}\nu \longrightarrow 2 \cdot \text{CL} + \text{COFCL} + \text{COF}_2$	
$\text{CFC}_{114} + \text{h}\nu \longrightarrow 2 \cdot \text{CL} + 2 \cdot \text{COF}_2$	
$\text{CFC}_{115} + \text{h}\nu \longrightarrow \text{CL} + \text{F} + 2 \cdot \text{COF}_2$	
$\text{HCFC}_{22} + \text{h}\nu \longrightarrow \text{CL} + \text{COF}_2$	
$\text{HCFC}_{141}\text{B} + \text{h}\nu \longrightarrow \text{CL} + \text{COFCL}$	
$\text{HCFC}_{142}\text{B} + \text{h}\nu \longrightarrow \text{CL} + \text{COF}_2$	
$\text{H}_{1202} + \text{h}\nu \longrightarrow 2 \cdot \text{BR} + \text{COF}_2$	
$\text{H}_{2402} + \text{h}\nu \longrightarrow 2 \cdot \text{BR} + 2 \cdot \text{COF}_2$	
$\text{COF}_2 + \text{h}\nu \longrightarrow 2 \cdot \text{F}$	
$\text{COFCL} + \text{h}\nu \longrightarrow \text{CL} + \text{F}$	

Table S3: (Tropospheric) O<sub>x</sub> reactions

reaction	rate coefficient	reference
$\text{O} + \text{O}_2 + \text{M} \longrightarrow \text{M} + \text{O}_3$	$\text{O}_{\text{O}_2}$	
$\text{O} + \text{O}_3 \longrightarrow 2 \cdot \text{O}_2$	$8.000 \cdot 10^{-12} \exp(-2060./T)$	JPL (2011)
$\text{O} + \text{O} + \text{M} \longrightarrow \text{M} + \text{O}_2$	$\text{O}_{\text{O}}$	not in JPL (2011)
$\text{O}_1\text{D} + \text{N}_2 \longrightarrow \text{N}_2 + \text{O}$	$2.150 \cdot 10^{-11} \exp(110./T)$	
$\text{O}_1\text{D} + \text{O}_2 \longrightarrow \text{O} + \text{O}_2$	$3.135 \cdot 10^{-11} \exp(55./T)$	

Table S3: (Tropospheric) O<sub>x</sub> reactions (... continued)

reaction	rate coefficient	reference
$\text{O}_1\text{D} + \text{O}_2 \longrightarrow \text{O} + \text{O}_2$	$1.650 \cdot 10^{-12} \exp(55./T)$	
$\text{O}_1\text{D} + \text{H}_2\text{O} \longrightarrow 2 \cdot \text{OH}$	$1.630 \cdot 10^{-10} \exp(60./T)$	JPL (2011)

Table S4: (Tropospheric) HO<sub>x</sub> reactions

reaction	rate coefficient	reference
$\text{H} + \text{O}_2 + \text{M} \longrightarrow \text{HO}_2 + \text{M}$	<i>ktroe</i> ( $4.400 \cdot 10^{-32}, 1.3, 7.500 \cdot 10^{-11}, -0.2, 0.6$ )	JPL (2011)
$\text{H} + \text{O}_3 \longrightarrow \text{O}_2 + \text{OH}$	$1.400 \cdot 10^{-10} \exp(-470./T)$	JPL (2011)
$\text{H} + \text{HO}_2 \longrightarrow 2 \cdot \text{OH}$	$7.200 \cdot 10^{-11}$	JPL (2011)
$\text{H} + \text{HO}_2 \longrightarrow \text{H}_2\text{O} + \text{O}$	$1.600 \cdot 10^{-12}$	JPL (2011)
$\text{H} + \text{HO}_2 \longrightarrow \text{H}_2 + \text{O}_2$	$6.900 \cdot 10^{-12}$	JPL (2011)
$\text{H}_2 + \text{O} \longrightarrow \text{H} + \text{OH}$	$1.600 \cdot 10^{-11} \exp(-4570./T)$	
$\text{H}_2 + \text{OH} \longrightarrow \text{H} + \text{H}_2\text{O}$	$2.800 \cdot 10^{-12} \exp(-1800./T)$	JPL (2011)
$\text{OH} + \text{O} \longrightarrow \text{H} + \text{O}_2$	$1.800 \cdot 10^{-11} \exp(180./T)$	JPL (2011)
$\text{OH} + \text{OH} \longrightarrow \text{H}_2\text{O} + \text{O}$	$1.800 \cdot 10^{-12}$	JPL (2011)
$\text{OH} + \text{OH} + \text{M} \longrightarrow \text{H}_2\text{O}_2 + \text{M}$	<i>ktroe</i> ( $6.900 \cdot 10^{-31}, 1., 2.600 \cdot 10^{-11}, 0., 0.6$ )	JPL (2011)
$\text{OH} + \text{O}_3 \longrightarrow \text{HO}_2 + \text{O}_2$	$1.700 \cdot 10^{-12} \exp(-940./T)$	JPL (2011)
$\text{HO}_2 + \text{O} \longrightarrow \text{O}_2 + \text{OH}$	$3.000 \cdot 10^{-11} \exp(200./T)$	JPL (2011)
$\text{HO}_2 + \text{OH} \longrightarrow \text{H}_2\text{O} + \text{O}_2$	$4.800 \cdot 10^{-11} \exp(250./T)$	JPL (2011)
$\text{HO}_2 + \text{O}_3 \longrightarrow \text{OH} + 2 \cdot \text{O}_2$		IUPAC (2004)
$\text{HO}_2 + \text{HO}_2 \longrightarrow \text{H}_2\text{O}_2 + \text{O}_2$	HO <sub>2</sub> –HO <sub>2</sub>	
$\text{H}_2\text{O}_2 + \text{O} \longrightarrow \text{HO}_2 + \text{OH}$	$1.400 \cdot 10^{-12} \exp(-2000./T)$	JPL (2011)
$\text{H}_2\text{O}_2 + \text{OH} \longrightarrow \text{H}_2\text{O} + \text{HO}_2$	$1.800 \cdot 10^{-12}$	JPL (2011)

Table S5: NO<sub>x</sub> reactions

reaction	rate coefficient	reference
$\text{N} + \text{OH} \longrightarrow \text{H} + \text{NO}$	$5.000 \cdot 10^{-11}$	
$\text{N} + \text{O}_2 \longrightarrow \text{NO} + \text{O}$	$1.500 \cdot 10^{-11} \exp(-3600./T)$	JPL (2011)
$\text{N} + \text{NO} \longrightarrow \text{N}_2 + \text{O}$	$2.100 \cdot 10^{-11} \exp(100./T)$	JPL (2011)
$\text{N} + \text{NO}_2 \longrightarrow 0.5 \cdot \text{N}_2\text{O} + 0.5 \cdot \text{O} + 0.5 \cdot \text{NO} + 0.25 \cdot \text{N}_2 + 0.25 \cdot \text{O}_2$	$5.800 \cdot 10^{-12} \exp(220./T)$	JPL (2011), products: D. Kinnison
$\text{NO} + \text{O} + \text{M} \longrightarrow \text{M} + \text{NO}_2$	<i>ktroe</i> ( $9.000 \cdot 10^{-32}, 1.5, 3.000 \cdot 10^{-11}, 0., 0.6$ )	JPL (2011)
$\text{NO} + \text{O}_3 \longrightarrow \text{NO}_2 + \text{O}_2$	$3.000 \cdot 10^{-12} \exp(-1500./T)$	JPL (2011)



Table S5: NOx reactions (... continued)

reaction	rate coefficient	reference
$\text{NO} + \text{HO}_2 \longrightarrow \text{NO}_2 + \text{OH}$	$3.300 \cdot 10^{-12} \exp(270./T)$	JPL (2011)
$\text{NO}_2 + \text{O} \longrightarrow \text{NO} + \text{O}_2$	$5.100 \cdot 10^{-12} \exp(210./T)$	JPL (2011)
$\text{NO}_2 + \text{O} + \text{M} \longrightarrow \text{M} + \text{NO}_3$	$ktroe(2.500 \cdot 10^{-31}, 1.8, 2.200 \cdot 10^{-11}, 0.7, 0.6)$	JPL (2011)
$\text{NO}_2 + \text{O}_3 \longrightarrow \text{NO}_3 + \text{O}_2$	$1.200 \cdot 10^{-13} \exp(-2450./T)$	JPL (2011)
$\text{NO}_2 + \text{H} \longrightarrow \text{NO} + \text{OH}$	$4.000 \cdot 10^{-10} \exp(-340./T)$	JPL (2011)
$\text{NO}_2 + \text{OH} + \text{M} \longrightarrow \text{HNO}_3 + \text{M}$	$ktroe(1.800 \cdot 10^{-30}, 3., 2.800 \cdot 10^{-11}, 0., 0.6)$	JPL (2011)
$\text{NO}_3 + \text{O} \longrightarrow \text{NO}_2 + \text{O}_2$	$1.000 \cdot 10^{-11}$	JPL (2011)
$\text{NO}_3 + \text{OH} \longrightarrow \text{HO}_2 + \text{NO}_2$	$2.200 \cdot 10^{-11}$	JPL (2011)
$\text{NO}_3 + \text{HO}_2 \longrightarrow \text{NO}_2 + \text{OH} + \text{O}_2$	$3.500 \cdot 10^{-12}$	JPL (2011)
$\text{NO}_3 + \text{NO} \longrightarrow 2 \cdot \text{NO}_2$	$1.500 \cdot 10^{-11} \exp(170./T)$	JPL (2011)
$\text{HNO}_3 + \text{OH} \longrightarrow \text{H}_2\text{O} + \text{NO}_3$		$X/(1 + X/(2.7 \cdot 10^{-17} \exp(2199/T))) + 2.4 \cdot 10^{-14} \exp(460/T); X = M6.5 \cdot 10^{-34} \exp(1335/T) - \text{JPL (2011)}$
$\text{NO} + \text{OH} \longrightarrow \text{HONO}$	$ktroe(7.000 \cdot 10^{-31}, 2.6, 3.600 \cdot 10^{-11}, 0.1, 0.6)$	JPL (2011)
$\text{HONO} + \text{OH} \longrightarrow \text{H}_2\text{O} + \text{NO}_2$	$1.800 \cdot 10^{-11} \exp(-390./T)$	JPL (2011)
$\text{NO}_2 + \text{HO}_2 + \text{M} \longrightarrow \text{HO}_2\text{NO}_2 + \text{M}$	$ktroe(2.000 \cdot 10^{-31}, 3.4, 2.900 \cdot 10^{-12}, 1.1, 0.6)$	JPL (2011)
$\text{HO}_2\text{NO}_2 + \text{OH} \longrightarrow \text{H}_2\text{O} + \text{NO}_2 + \text{O}_2$	$1.300 \cdot 10^{-12} \exp(380./T)$	JPL (2011)
$\text{HO}_2\text{NO}_2 + \text{M} \longrightarrow \text{HO}_2 + \text{NO}_2 + \text{M}$		$[\text{NO}_2 - \text{NO}_3] * \exp(-10900./T)/2.1 \cdot 10^{-27} \text{ JPL (2011)}$
$\text{NO}_2 + \text{NO}_3 + \text{M} \longrightarrow \text{M} + \text{N}_2\text{O}_5$	$ktroe(2.000 \cdot 10^{-30}, 4.4, 1.400 \cdot 10^{-12}, 0.7, 0.6)$	JPL (2011)
$\text{N}_2\text{O}_5 + \text{M} \longrightarrow \text{NO}_2 + \text{NO}_3 + \text{M}$	$kN2O5$	$[\text{NO}_2 - \text{NO}_3] * \exp(-11000/T)/2.7 \cdot 10^{-27} \text{ JPL (2011)}$
$\text{NH}_3 + \text{OH} \longrightarrow$	$1.700 \cdot 10^{-12} \exp(-710./T)$	JPL (2011)

Table S6: C1 oxidation

reaction	rate coefficient	reference
$\text{CO} + \text{OH} \longrightarrow \text{CO}_2 + \text{H}$	$kactiv(1.5 \cdot 10^{-13}, -0.6, 2.1 \cdot 10^9, -6.1)$	JPL (2011)
$\text{CO} + \text{OH} + \text{M} \longrightarrow \text{CO}_2 + \text{HO}_2 + \text{M}$	$ktroe(5.900 \cdot 10^{-33}, 1.4, 1.100 \cdot 10^{-12}, -1.3, 0.6)$	
$\text{CH}_4 + \text{OH} \longrightarrow \text{CH}_3\text{O}_2 + \text{H}_2\text{O}$	$2.450 \cdot 10^{-12} \exp(-1775./T)$	JPL (2011)
$\text{CH}_3\text{OH} + \text{OH} \longrightarrow \text{CH}_2\text{O} + \text{HO}_2 + \text{H}_2\text{O}$	$2.900 \cdot 10^{-12} \exp(-345./T)$	JPL (2011)
$\text{CH}_3\text{O}_2 + \text{NO} \longrightarrow \text{CH}_2\text{O} + \text{NO}_2 + \text{HO}_2$	$1.960 \cdot 10^{-12} \exp(403./T)$	Orlando and Tyndall (2012)
$\text{CH}_3\text{O}_2 + \text{HO}_2 \longrightarrow \text{CH}_3\text{OOH} + \text{O}_2$	$3.800 \cdot 10^{-13} \exp(730./T)$	Orlando and Tyndall (2012)
$\text{CH}_3\text{O}_2 + \text{CH}_3\text{O}_2 \longrightarrow 2 \cdot \text{CH}_2\text{O} + 2 \cdot \text{HO}_2$	$7.400 \cdot 10^{-13} \exp(-520./T)$	IUPAC (2006)

Table S6: C1 oxidation (... continued)

reaction	rate coefficient	reference
$\text{CH}_3\text{O}_2 + \text{CH}_3\text{O}_2 \longrightarrow \text{CH}_2\text{O} + \text{CH}_3\text{OH}$	$2.330 \cdot 10^{-14} \exp(678./T)$	own fit for $T = 240 - 300\text{K}$
$\text{CH}_2\text{O} + \text{O} \longrightarrow \text{HO}_2 + \text{OH} + \text{CO}$	$3.400 \cdot 10^{-11} \exp(-1600./T)$	JPL (2011)
$\text{CH}_2\text{O} + \text{OH} \longrightarrow \text{CO} + \text{H}_2\text{O} + \text{H}$	$5.500 \cdot 10^{-12} \exp(125./T)$	JPL (2011)
$\text{CH}_2\text{O} + \text{HO}_2 \longrightarrow \text{HOCH}_2\text{OO}$	$9.700 \cdot 10^{-15} \exp(625./T)$	IUPAC (2006)
$\text{CH}_2\text{O} + \text{NO}_3 \longrightarrow \text{CO} + \text{HO}_2 + \text{HNO}_3$	$6.000 \cdot 10^{-13} \exp(-2058./T)$	
$\text{CH}_3\text{OOH} + \text{OH} \longrightarrow 0.7 \cdot \text{CH}_3\text{O}_2 + 0.3 \cdot \text{OH} + 0.3 \cdot \text{CH}_2\text{O} + \text{H}_2\text{O}$	$3.800 \cdot 10^{-12} \exp(200./T)$	JPL (2011)
$\text{HCOOH} + \text{OH} \longrightarrow \text{CO}_2 + \text{HO}_2 + \text{H}_2\text{O}$	$4.000 \cdot 10^{-13}$	JPL (2011)
$\text{HOCH}_2\text{OO} \longrightarrow \text{CH}_2\text{O} + \text{HO}_2$	$2.400 \cdot 10^{12} \exp(-7000./T)$	Lamarque et al. (2012)
$\text{HOCH}_2\text{OO} + \text{NO} \longrightarrow \text{HCOOH} + \text{NO}_2 + \text{HO}_2$	$2.600 \cdot 10^{-12} \exp(265./T)$	Lamarque et al. (2012)
$\text{HOCH}_2\text{OO} + \text{HO}_2 \longrightarrow \text{H}_2\text{O} + \text{HCOOH}$	$7.500 \cdot 10^{-13} \exp(700./T)$	Lamarque et al. (2012)
$\text{HCN} + \text{OH} \longrightarrow \text{CO} + \text{NO} + \text{H}_2\text{O}$	$1.200 \cdot 10^{-13} \exp(-400./T)$	JPL (2011), products: Tyndall
$\text{CH}_3\text{O}_2 + \text{NO}_2 + \text{M} \longrightarrow \text{CH}_3\text{O}_2\text{NO}_2 + \text{M}$	$ktroe(1.000 \cdot 10^{-30}, 4.8, 7.200 \cdot 10^{-12}, 2.1, 0.6)$	JPL (2011)
$\text{CH}_3\text{O}_2\text{NO}_2 + \text{M} \longrightarrow \text{CH}_3\text{O}_2 + \text{NO}_2 + \text{M}$		

Table S7: C2 oxidation

reaction	rate coefficient	reference
$\text{C}_2\text{H}_2 + \text{OH} + \text{M} \longrightarrow 0.65 \cdot \text{GLYOXAL} + 0.65 \cdot \text{OH} + 0.35 \cdot \text{HCOOH} + 0.35 \cdot \text{HO}_2 + 0.35 \cdot \text{CO} + \text{M}$	$ktroe(5.500 \cdot 10^{-30}, 0., 8.300 \cdot 10^{-13}, -2., 0.6)$	JPL (2011)
$\text{C}_2\text{H}_4 + \text{OH} + \text{M} \longrightarrow \text{EO}_2 + \text{M}$	$ktroe(1.000 \cdot 10^{-28}, 4.5, 7.500 \cdot 10^{-12}, 0.85, 0.6)$	JPL (2011)
$\text{C}_2\text{H}_4 + \text{O}_3 \longrightarrow \text{CH}_2\text{O} + 0.65 \cdot \text{CO} + 0.15 \cdot \text{OH} + 0.15 \cdot \text{HO}_2 + 0.5 \cdot \text{H}_2\text{O} + 0.35 \cdot \text{HCOOH}$	$9.100 \cdot 10^{-15} \exp(-2580./T)$	IUPAC (2006)
$\text{C}_2\text{H}_6 + \text{OH} \longrightarrow \text{C}_2\text{H}_5\text{O}_2 + \text{H}_2\text{O}$	$7.660 \cdot 10^{-12} \exp(-1020./T)$	JPL (2011)
$\text{C}_2\text{H}_5\text{OH} + \text{OH} \longrightarrow \text{CH}_3\text{CHO} + \text{HO}_2 + \text{H}_2\text{O}$	$3.350 \cdot 10^{-12} \exp(0./T)$	JPL (2011)
$\text{CH}_3\text{CHO} + \text{OH} \longrightarrow \text{CH}_3\text{CO}_3 + \text{H}_2\text{O}$	$4.630 \cdot 10^{-12} \exp(350./T)$	JPL (2011)
$\text{CH}_3\text{CHO} + \text{NO}_3 \longrightarrow \text{CH}_3\text{CO}_3 + \text{HNO}_3$	$1.400 \cdot 10^{-12} \exp(-1900./T)$	JPL (2011)
$\text{CH}_3\text{COOOH} + \text{OH} \longrightarrow 0.5 \cdot \text{CH}_3\text{CO}_3 + \text{H}_2\text{O} + 0.5 \cdot \text{CH}_2\text{O} + 0.5 \cdot \text{CO}_2 + 0.5 \cdot \text{OH}$	$1.000 \cdot 10^{-12}$	Orlando (p.c.) added OH
$\text{C}_2\text{H}_5\text{O}_2 + \text{NO} \longrightarrow \text{CH}_3\text{CHO} + \text{HO}_2 + \text{NO}_2$	$2.620 \cdot 10^{-12} \exp(373./T)$	Orlando and Tyndall (2012)
$\text{C}_2\text{H}_5\text{O}_2 + \text{HO}_2 \longrightarrow \text{C}_2\text{H}_5\text{OOH} + \text{O}_2$	$7.400 \cdot 10^{-13} \exp(700./T)$	Orlando and Tyndall (2012)
$\text{C}_2\text{H}_5\text{O}_2 + \text{CH}_3\text{O}_2 \longrightarrow 0.7 \cdot \text{CH}_2\text{O} + 0.8 \cdot \text{CH}_3\text{CHO} + \text{HO}_2 + 0.3 \cdot \text{CH}_3\text{OH} + 0.2 \cdot \text{C}_2\text{H}_5\text{OH}$	$2.000 \cdot 10^{-13}$	Orlando (p.c.), products: Tyndall (p.c.)
$\text{C}_2\text{H}_5\text{O}_2 + \text{CH}_3\text{CO}_3 \longrightarrow \text{CH}_3\text{CHO} + \text{HO}_2 + \text{CH}_3\text{O}_2 + \text{CO}_2$	$1.800 \cdot 10^{-12} \exp(500./T)$	10% lower than $\text{CH}_3\text{O}_2 + \text{CH}_3\text{CO}_3$ ; Orlando and Tyndall (2012) only give k@298K

Table S7: C2 oxidation (... continued)

reaction	rate coefficient	reference
$\text{C}_2\text{H}_5\text{O}_2 + \text{C}_2\text{H}_5\text{O}_2 \longrightarrow 1.6 \cdot \text{CH}_3\text{CHO} + 1.2 \cdot \text{HO}_2 + 0.4 \cdot \text{C}_2\text{H}_5\text{OH}$	$7.600 \cdot 10^{-14}$	Orlando and Tyndall (2012)
$\text{C}_2\text{H}_5\text{OOH} + \text{OH} \longrightarrow 0.5 \cdot \text{C}_2\text{H}_5\text{O}_2 + 0.5 \cdot \text{CH}_3\text{CHO} + 0.5 \cdot \text{OH} + \text{H}_2\text{O}$	$3.800 \cdot 10^{-12} \exp(200./T)$	no data, analog to $\text{CH}_3\text{OOH} + \text{OH}$
$\text{CH}_3\text{CO}_3 + \text{NO} \longrightarrow \text{CH}_3\text{O}_2 + \text{CO}_2 + \text{NO}_2$	$7.500 \cdot 10^{-12} \exp(290./T)$	Orlando and Tyndall (2012)
$\text{CH}_3\text{CO}_3 + \text{NO}_2 + \text{M} \longrightarrow \text{M} + \text{PAN}$	$ktroe(2.700 \cdot 10^{-28}, 7.1, 1.200 \cdot 10^{-11}, 0.9, 0.6)$	
$\text{CH}_3\text{CO}_3 + \text{HO}_2 \longrightarrow 0.4 \cdot \text{CH}_3\text{COOOH} + 0.2 \cdot \text{CH}_3\text{COOH} + 0.2 \cdot \text{O}_3 + 0.4 \cdot \text{CH}_3\text{O}_2 + 0.4 \cdot \text{OH} + 0.4 \cdot \text{CO}_2$	$5.200 \cdot 10^{-13} \exp(980./T)$	Orlando and Tyndall (2012)
$\text{CH}_3\text{CO}_3 + \text{CH}_3\text{O}_2 \longrightarrow \text{CH}_2\text{O} + 0.9 \cdot \text{CH}_3\text{O}_2 + 0.9 \cdot \text{HO}_2 + 0.9 \cdot \text{CO}_2 + 0.1 \cdot \text{CH}_3\text{COOH}$	$2.000 \cdot 10^{-12} \exp(500./T)$	Orlando and Tyndall (2012)
$\text{CH}_3\text{CO}_3 + \text{CH}_3\text{CO}_3 \longrightarrow 2 \cdot \text{CH}_3\text{O}_2 + 2 \cdot \text{CO}_2$	$2.900 \cdot 10^{-12} \exp(500./T)$	Orlando and Tyndall (2012)
$\text{CH}_3\text{COOH} + \text{OH} \longrightarrow \text{CH}_3\text{O}_2 + \text{CO}_2 + \text{H}_2\text{O}$	$3.150 \cdot 10^{-14} \exp(920./T)$	JPL (2011)
$\text{PAN} + \text{M} \longrightarrow \text{CH}_3\text{CO}_3 + \text{NO}_2 + \text{M}$		
$\text{PAN} + \text{OH} \longrightarrow \text{CH}_2\text{O} + \text{CO}_2 + \text{NO}_3$	$4.000 \cdot 10^{-14}$	JPL (2011), includes implicit $\text{NO} \longrightarrow \text{NO}_2$ conversion
$\text{EO}_2 + \text{NO} \longrightarrow 0.75 \cdot \text{EO} + \text{NO}_2 + 0.5 \cdot \text{CH}_2\text{O} + 0.25 \cdot \text{HO}_2$	$4.200 \cdot 10^{-12} \exp(180./T)$	Lamarque et al. (2012)
$\text{EO}_2 + \text{HO}_2 \longrightarrow \text{EOOH}$	$7.500 \cdot 10^{-13} \exp(700./T)$	
$\text{EO}_2 + \text{CH}_3\text{O}_2 \longrightarrow 0.5 \cdot \text{EO} + 0.5 \cdot \text{O}_2 + 0.5 \cdot \text{CH}_2\text{O} + 0.5 \cdot \text{HO}_2 + 0.5 \cdot \text{CH}_3\text{OH} + 0.5 \cdot \text{GLYALD}$	$4.000 \cdot 10^{-12} \exp(1000./T)$	Tyndall (p.c.)
$\text{EO}_2 + \text{CH}_3\text{CO}_3 \longrightarrow \text{CH}_3\text{O}_2 + \text{EO} + \text{CO}_2$	$1.000 \cdot 10^{-11}$	
$\text{EO} + \text{O}_2 \longrightarrow \text{GLYALD} + \text{HO}_2$	$1.000 \cdot 10^{-14}$	Lamarque et al. (2012)
$\text{EO} \longrightarrow 2 \cdot \text{CH}_2\text{O} + \text{HO}_2$	$1.600 \cdot 10^{11} \exp(-4150./T)$	Lamarque et al. (2012)
$\text{GLYOXAL} + \text{OH} \longrightarrow 0.6 \cdot \text{HO}_2 + 1.2 \cdot \text{CO} + \text{H}_2\text{O} + 0.4 \cdot \text{HCOCO}_3$	$3.100 \cdot 10^{-12} \exp(340./T)$	MCM3.2
$\text{GLYOXAL} + \text{NO}_3 \longrightarrow 0.6 \cdot \text{HO}_2 + 1.2 \cdot \text{CO} + 0.4 \cdot \text{HCOCO}_3 + \text{HNO}_3$	$2.500 \cdot 10^{-12}$	Taraborrelli et al. (2009)
$\text{GLYALD} + \text{OH} \longrightarrow 0.2 \cdot \text{GLYOXAL} + 0.2 \cdot \text{HO}_2 + 0.8 \cdot \text{HOCH}_2\text{CO}_3 + \text{H}_2\text{O}$	$1.000 \cdot 10^{-11}$	Taraborrelli et al. (2009)
$\text{GLYALD} + \text{NO}_3 \longrightarrow \text{HNO}_3 + \text{HOCH}_2\text{CO}_3$	$1.440 \cdot 10^{-12} \exp(-1862./T)$	Taraborrelli et al. (2009)
$\text{HOCH}_2\text{CO}_3 + \text{NO}_2 + \text{M} \longrightarrow \text{CH}_2\text{O} + \text{CO}_2 + \text{HNO}_3 + \text{M}$	$ktroe(2.700 \cdot 10^{-28}, 7.1, 1.200 \cdot 10^{-11}, 0.9, 0.6)$	Orlando and Tyndall (2012)
$\text{HOCH}_2\text{CO}_3 + \text{HO}_2 \longrightarrow 0.4 \cdot \text{HOCH}_2\text{CO}_3\text{H} + 0.2 \cdot \text{HOCH}_2\text{CO}_2\text{H} + 0.2 \cdot \text{O}_3 + 0.4 \cdot \text{CO}_2 + 0.4 \cdot \text{OH} + 0.4 \cdot \text{HO}_2 + 0.4 \cdot \text{CH}_2\text{O}$	$4.300 \cdot 10^{-13} \exp(1040./T)$	Orlando (p.c.)
$\text{HOCH}_2\text{CO}_3 + \text{CH}_3\text{O}_2 \longrightarrow 2 \cdot \text{CH}_2\text{O} + \text{CO}_2 + 2 \cdot \text{HO}_2$	$1.000 \cdot 10^{-11}$	Taraborrelli et al. (2009)
$\text{HOCH}_2\text{CO}_3 + \text{CH}_3\text{CO}_3 \longrightarrow \text{CH}_3\text{O}_2 + 2 \cdot \text{CO}_2 + \text{HO}_2 + \text{CH}_2\text{O}$	$1.000 \cdot 10^{-11}$	Taraborrelli et al. (2009)

Table S7: C2 oxidation (... continued)

reaction	rate coefficient	reference
$\text{HOCH}_2\text{CO}_3 + \text{NO} \longrightarrow \text{HO}_2 + \text{NO}_2 + \text{CH}_2\text{O} + \text{CO}_2$	$8.100 \cdot 10^{-12} \exp(270./T)$	Taraborrelli et al. (2009)
$\text{HOCH}_2\text{CO}_3 + \text{NO}_3 \longrightarrow \text{HO}_2 + \text{NO}_2 + \text{CH}_2\text{O} + \text{CO}_2$	$4.000 \cdot 10^{-12}$	Taraborrelli et al. (2009)
$\text{HOCH}_2\text{CO}_2\text{H} + \text{OH} \longrightarrow \text{CH}_2\text{O} + \text{HO}_2 + \text{CO}_2 + \text{H}_2\text{O}$	$2.730 \cdot 10^{-12}$	Taraborrelli et al. (2009)
$\text{HOCH}_2\text{CO}_3\text{H} + \text{OH} \longrightarrow \text{H}_2\text{O} + \text{HOCH}_2\text{CO}_3$	$6.190 \cdot 10^{-12}$	Taraborrelli et al. (2009)
$\text{HCOCO}_3 + \text{CH}_3\text{O}_2 \longrightarrow \text{CO} + 2 \cdot \text{HO}_2 + \text{CO}_2 + \text{CH}_2\text{O}$	$1.000 \cdot 10^{-11}$	Taraborrelli et al. (2009)
$\text{HCOCO}_3 + \text{CH}_3\text{CO}_3 \longrightarrow \text{CO} + \text{HO}_2 + \text{CH}_3\text{O}_2 + 2 \cdot \text{CO}_2$	$1.000 \cdot 10^{-11}$	Taraborrelli et al. (2009)
$\text{HCOCO}_3 + \text{HO}_2 \longrightarrow 0.7 \cdot \text{HCOCO}_3\text{H} + 0.7 \cdot \text{O}_2 + 0.3 \cdot \text{HCOCO}_2\text{H} + 0.3 \cdot \text{O}_3$	$4.300 \cdot 10^{-13} \exp(1040./T)$	Taraborrelli et al. (2009)
$\text{HCOCO}_3 + \text{NO} \longrightarrow \text{CO} + \text{HO}_2 + \text{NO}_2 + \text{CO}_2$	$8.100 \cdot 10^{-12} \exp(270./T)$	Taraborrelli et al. (2009)
$\text{HCOCO}_3 + \text{NO}_3 \longrightarrow \text{CO} + \text{HO}_2 + \text{NO}_2 + \text{CO}_2$	$4.000 \cdot 10^{-12}$	Taraborrelli et al. (2009)
$\text{HCOCO}_2\text{H} + \text{OH} \longrightarrow \text{CO} + \text{HO}_2 + \text{CO}_2 + \text{H}_2\text{O}$	$1.230 \cdot 10^{-11}$	Taraborrelli et al. (2009)
$\text{HCOCO}_3\text{H} + \text{OH} \longrightarrow \text{H}_2\text{O} + \text{HCOCO}_3$	$1.580 \cdot 10^{-11}$	Taraborrelli et al. (2009)
$\text{CH}_3\text{CN} + \text{OH} \longrightarrow \text{CH}_2\text{O} + \text{H}_2\text{O} + \text{CO} + \text{NO}_2$	$7.800 \cdot 10^{-13} \exp(-1050./T)$	products: Tyndall

Table S8: C3 oxidation

reaction	rate coefficient	reference
$\text{C}_3\text{H}_6 + \text{OH} + \text{M} \longrightarrow \text{M} + \text{PO}_2$	<i>ktroe</i> ( $8.000 \cdot 10^{-27}, 3.5, 3.000 \cdot 10^{-11}, 0., 0.5$ )	IUPAC (2006)
$\text{C}_3\text{H}_6 + \text{O}_3 \longrightarrow 0.28 \cdot \text{CH}_3\text{O}_2 + 0.1 \cdot \text{CH}_4 + 0.075 \cdot \text{CH}_3\text{COOH} + 0.56 \cdot \text{CO} + 0.075 \cdot \text{HCOOH} + 0.09 \cdot \text{H}_2\text{O}_2 + 0.28 \cdot \text{HO}_2 + 0.2 \cdot \text{CO}_2 + 0.545 \cdot \text{CH}_3\text{CHO} + 0.545 \cdot \text{CH}_2\text{O} + 0.36 \cdot \text{OH}$	$5.500 \cdot 10^{-15} \exp(-1880./T)$	IUPAC (2006)
$\text{C}_3\text{H}_6 + \text{NO}_3 \longrightarrow \text{PRONO}_3\text{BO}_2$	$4.600 \cdot 10^{-13} \exp(-1156./T)$	IUPAC (2006)
$\text{C}_3\text{H}_8 + \text{OH} \longrightarrow \text{C}_3\text{H}_7\text{O}_2 + \text{H}_2\text{O}$	$7.600 \cdot 10^{-12} \exp(-585./T)$	IUPAC (2006)
$\text{CH}_3\text{COCH}_3 + \text{OH} \longrightarrow \text{CH}_3\text{COCH}_2\text{O}_2 + \text{H}_2\text{O}$	$\text{CH}_3\text{COCH}_3 - \text{OH}$	
$\text{C}_3\text{H}_7\text{O}_2 + \text{NO} \longrightarrow 0.82 \cdot \text{CH}_3\text{COCH}_3 + \text{NO}_2 + \text{HO}_2 + 0.27 \cdot \text{CH}_3\text{CHO}$	$2.900 \cdot 10^{-12} \exp(350./T)$	Orlando and Tyndall (2012)
$\text{C}_3\text{H}_7\text{O}_2 + \text{HO}_2 \longrightarrow \text{C}_3\text{H}_7\text{OOH} + \text{O}_2$	$7.500 \cdot 10^{-13} \exp(700./T)$	Lamarque et al. (2012)
$\text{C}_3\text{H}_7\text{O}_2 + \text{CH}_3\text{O}_2 \longrightarrow \text{CH}_2\text{O} + 2 \cdot \text{HO}_2 + 0.82 \cdot \text{CH}_3\text{COCH}_3 + 0.27 \cdot \text{CH}_3\text{CHO}$	$3.750 \cdot 10^{-13} \exp(-40./T)$	Lamarque et al. (2012)
$\text{C}_3\text{H}_7\text{O}_2 + \text{CH}_3\text{CO}_3 \longrightarrow 0.82 \cdot \text{CH}_3\text{COCH}_3 + \text{HO}_2 + 0.27 \cdot \text{CH}_3\text{CHO} + \text{CO}_2 + \text{CH}_3\text{O}_2$	$1.000 \cdot 10^{-11}$	Lamarque et al. (2012)
$\text{C}_3\text{H}_7\text{OOH} + \text{OH} \longrightarrow 0.41 \cdot \text{CH}_3\text{COCH}_3 + 0.5 \cdot \text{OH} + 0.5 \cdot \text{C}_3\text{H}_7\text{O}_2 + 0.5 \cdot \text{H}_2\text{O} + 0.135 \cdot \text{CH}_3\text{CHO}$	$3.800 \cdot 10^{-12} \exp(200./T)$	Lamarque et al. (2012); $1.5 \cdot \text{CH}_3\text{CHO}$ as surrogate for propanal
$\text{PO}_2 + \text{NO} \longrightarrow \text{CH}_2\text{O} + \text{CH}_3\text{CHO} + \text{HO}_2 + \text{NO}_2$	$4.200 \cdot 10^{-12} \exp(180./T)$	Lamarque et al. (2012)

Table S8: C3 oxidation (... continued)

reaction	rate coefficient	reference
$\text{PO}_2 + \text{NO}_3 \longrightarrow \text{CH}_2\text{O} + \text{CH}_3\text{CHO} + \text{HO}_2 + \text{NO}_2$	$2.500 \cdot 10^{-12}$	Taraborrelli et al. (2009)
$\text{PO}_2 + \text{HO}_2 \longrightarrow \text{O}_2 + \text{POOH}$	$7.500 \cdot 10^{-13} \exp(700./T)$	Lamarque et al. (2012)
$\text{PO}_2 + \text{CH}_3\text{O}_2 \longrightarrow 0.5 \cdot \text{CH}_3\text{CHO} + 1.25 \cdot \text{CH}_2\text{O} + \text{HO}_2 + 0.5 \cdot \text{HYAC} + 0.25 \cdot \text{CH}_3\text{OH}$	$8.300 \cdot 10^{-13}$	products: Tyndall (p.c.)
$\text{PO}_2 + \text{CH}_3\text{CO}_3 \longrightarrow \text{CH}_2\text{O} + \text{CH}_3\text{CHO} + \text{HO}_2 + \text{CH}_3\text{O}_2 + \text{CO}_2$	$1.000 \cdot 10^{-11}$	
$\text{POOH} + \text{OH} \longrightarrow 0.5 \cdot \text{PO}_2 + 0.5 \cdot \text{HYAC} + 0.5 \cdot \text{OH} + \text{H}_2\text{O}$	$3.800 \cdot 10^{-12} \exp(200./T)$	Lamarque et al. (2012)
$\text{HYAC} + \text{OH} \longrightarrow \text{CH}_3\text{COCHO} + \text{HO}_2 + \text{H}_2\text{O}$	$3.000 \cdot 10^{-12}$	IUPAC (2006)
$\text{CH}_3\text{COCHO} + \text{OH} \longrightarrow \text{CH}_3\text{CO}_3 + \text{CO} + \text{H}_2\text{O}$	$8.400 \cdot 10^{-13} \exp(830./T)$	Lamarque et al. (2012)
$\text{CH}_3\text{COCHO} + \text{NO}_3 \longrightarrow \text{CO} + \text{HNO}_3 + \text{CH}_3\text{CO}_3$	$1.400 \cdot 10^{-12} \exp(-1860./T)$	Lamarque et al. (2012)
$\text{CH}_3\text{COCH}_2\text{O}_2 + \text{NO} \longrightarrow \text{CH}_2\text{O} + \text{CH}_3\text{CO}_3 + \text{NO}_2$	$2.900 \cdot 10^{-12} \exp(300./T)$	Lamarque et al. (2012)
$\text{CH}_3\text{COCH}_2\text{O}_2 + \text{HO}_2 \longrightarrow 0.85 \cdot \text{O}_2 + 0.85 \cdot \text{ROOH} + 0.15 \cdot \text{CH}_2\text{O} + 0.15 \cdot \text{CH}_3\text{CO}_3 + 0.15 \cdot \text{OH} + 0.15 \cdot \text{H}_2\text{O}$	$8.600 \cdot 10^{-13} \exp(700./T)$	Lamarque et al. (2012) + MCM3.2
$\text{CH}_3\text{COCH}_2\text{O}_2 + \text{CH}_3\text{O}_2 \longrightarrow 0.3 \cdot \text{CH}_3\text{CO}_3 + 0.8 \cdot \text{CH}_2\text{O} + 0.3 \cdot \text{HO}_2 + 0.2 \cdot \text{HYAC} + 0.5 \cdot \text{CH}_3\text{COCHO} + 0.5 \cdot \text{CH}_3\text{OH}$	$7.100 \cdot 10^{-13} \exp(500./T)$	Lamarque et al. (2012)
$\text{CH}_3\text{COCH}_2\text{O}_2 + \text{CH}_3\text{CO}_3 \longrightarrow \text{CH}_2\text{O} + \text{CH}_3\text{CO}_3 + \text{CH}_3\text{O}_2 + \text{CO}_2$	$1.000 \cdot 10^{-11}$	
$\text{ROOH} + \text{OH} \longrightarrow \text{CH}_3\text{COCH}_2\text{O}_2 + \text{H}_2\text{O}$	$3.800 \cdot 10^{-12} \exp(200./T)$	Lamarque et al. (2012)
$\text{PRONO}_3\text{BO}_2 + \text{NO} \longrightarrow 0.83 \cdot \text{HO}_2 + 0.83 \cdot \text{NOA} + 0.17 \cdot \text{CH}_2\text{O} + 0.17 \cdot \text{CH}_3\text{CHO} + 1.17 \cdot \text{NO}_2$	$2.540 \cdot 10^{-12} \exp(360./T)$	Taraborrelli et al. (2009)
$\text{PRONO}_3\text{BO}_2 + \text{NO}_3 \longrightarrow 0.83 \cdot \text{HO}_2 + 0.83 \cdot \text{NOA} + 0.17 \cdot \text{CH}_2\text{O} + 0.17 \cdot \text{CH}_3\text{CHO} + 1.17 \cdot \text{NO}_2$	$2.500 \cdot 10^{-12}$	Taraborrelli et al. (2009)
$\text{PRONO}_3\text{BO}_2 + \text{HO}_2 \longrightarrow \text{PR}_2\text{O}_2\text{HNO}_3$	$1.320 \cdot 10^{-12} \exp(360./T)$	Taraborrelli et al. (2009)
$\text{PRONO}_3\text{BO}_2 + \text{CH}_3\text{O}_2 \longrightarrow 0.915 \cdot \text{HO}_2 + 0.915 \cdot \text{NOA} + 0.835 \cdot \text{CH}_2\text{O} + 0.085 \cdot \text{CH}_3\text{CHO} + 0.25 \cdot \text{CH}_3\text{OH}$	$1.000 \cdot 10^{-12}$	
$\text{PRONO}_3\text{BO}_2 + \text{CH}_3\text{CO}_3 \longrightarrow 0.83 \cdot \text{HO}_2 + 0.83 \cdot \text{NOA} + 0.17 \cdot \text{CH}_2\text{O} + 0.17 \cdot \text{CH}_3\text{CHO} + \text{CH}_3\text{O}_2 + \text{CO}_2$	$1.000 \cdot 10^{-11}$	
$\text{PR}_2\text{O}_2\text{HNO}_3 + \text{OH} \longrightarrow 0.5 \cdot \text{PRONO}_3\text{BO}_2 + 0.5 \cdot \text{NOA} + 0.5 \cdot \text{OH} + \text{H}_2\text{O}$	$7.000 \cdot 10^{-12}$	Taraborrelli et al. (2009)
$\text{NOA} + \text{OH} \longrightarrow \text{CH}_3\text{COCHO} + \text{NO}_2 + \text{H}_2\text{O}$	$1.300 \cdot 10^{-13}$	Taraborrelli et al. (2009)

Table S9: C4 oxidation

reaction	rate coefficient	reference
$\text{BIGENE} + \text{OH} \longrightarrow \text{ENEO}_2$	$5.400 \cdot 10^{-11}$	Lamarque et al. (2012)

Table S9: C4 oxidation (... continued)

reaction	rate coefficient	reference
$\text{MEK} + \text{OH} \longrightarrow \text{H}_2\text{O} + \text{MEKO}_2$	$2.300 \cdot 10^{-12} \exp(-170./T)$	Lamarque et al. (2012)
$\text{MEKO}_2 + \text{NO} \longrightarrow \text{CH}_3\text{CHO} + \text{CH}_3\text{CO}_3 + \text{NO}_2$	$4.032 \cdot 10^{-12} \exp(180./T)$	treated like $\text{MEKBO}_2$ from MCM3.2; $A = 4.200 \cdot 10^{-12} \cdot 0.96$ ; 4% nitrate yield as for MVKN
$\text{MEKO}_2 + \text{NO} \longrightarrow \text{MEKNO}_3$	$1.680 \cdot 10^{-13} \exp(180./T)$	$A = 4.200 \cdot 10^{-12} \cdot 0.04$ ; 4% nitrate yield as for MVKN
$\text{MEKO}_2 + \text{HO}_2 \longrightarrow \text{MEKOOH}$	$7.500 \cdot 10^{-13} \exp(700./T)$	Lamarque et al. (2012)
$\text{MEKO}_2 + \text{CH}_3\text{O}_2 \longrightarrow 0.3 \cdot \text{CH}_3\text{CHO} + 0.3 \cdot \text{CH}_3\text{CO}_3 + \text{CH}_2\text{O} + 0.3 \cdot \text{HO}_2 + 0.3 \cdot \text{O}_2 + 0.5 \cdot \text{BIACETOH} + 0.5 \cdot \text{CH}_3\text{OH} + 0.266 \cdot \text{HYAC}$	$1.000 \cdot 10^{-12}$	Tyndall (p.c.) added $\text{CH}_2\text{O}$ to first and third channel
$\text{MEKO}_2 + \text{CH}_3\text{CO}_3 \longrightarrow \text{CH}_3\text{CHO} + \text{CH}_3\text{CO}_3 + \text{CH}_3\text{O}_2 + \text{CO}_2$	$1.000 \cdot 10^{-11}$	
$\text{MEKOOH} + \text{OH} \longrightarrow \text{H}_2\text{O} + \text{MEKO}_2$	$3.800 \cdot 10^{-12} \exp(200./T)$	Lamarque et al. (2012)
$\text{ENEO}_2 + \text{NO} \longrightarrow \text{CH}_3\text{CHO} + 0.5 \cdot \text{CH}_2\text{O} + 0.5 \cdot \text{CH}_3\text{COCH}_3 + \text{HO}_2 + \text{NO}_2$	$4.200 \cdot 10^{-12} \exp(180./T)$	Lamarque et al. (2012)
$\text{ENEO}_2 + \text{HO}_2 \longrightarrow 1.333 \cdot \text{POOH} + \text{O}_2$	$7.500 \cdot 10^{-13} \exp(700./T)$	factor 4/3 to preserve carbon
$\text{ENEO}_2 + \text{CH}_3\text{O}_2 \longrightarrow 0.665 \cdot \text{HYAC} + 0.5 \cdot \text{CH}_3\text{OH} + 0.5 \cdot \text{CH}_3\text{CHO} + 0.25 \cdot \text{CH}_3\text{COCH}_3 + 0.75 \cdot \text{CH}_2\text{O} + \text{HO}_2$	$1.000 \cdot 10^{-12}$	products: Tyndall (p.c.)
$\text{ENEO}_2 + \text{CH}_3\text{CO}_3 \longrightarrow \text{CH}_3\text{CHO} + 0.5 \cdot \text{CH}_2\text{O} + 0.5 \cdot \text{CH}_3\text{COCH}_3 + \text{HO}_2 + \text{CH}_3\text{O}_2 + \text{CO}_2$	$1.000 \cdot 10^{-11}$	
$\text{MACR} + \text{OH} \longrightarrow 0.45 \cdot \text{MCO}_3 + 0.55 \cdot \text{MACRO}_2$	$1.860 \cdot 10^{-11} \exp(175./T)$	Tnydall (p.c.)
$\text{MACR} + \text{O}_3 \longrightarrow 0.59 \cdot \text{CH}_3\text{COCHO} + 0.41 \cdot \text{CH}_3\text{CO}_3 + 0.82 \cdot \text{CO} + 0.41 \cdot \text{HO}_2 + 0.82 \cdot \text{OH} + 0.033750 \cdot \text{HCOOH} + 0.556250 \cdot \text{CH}_2\text{O} + 0.123750 \cdot \text{H}_2\text{O}_2$	$1.360 \cdot 10^{-15} \exp(-2112./T)$	
$\text{MACR} + \text{NO}_3 \longrightarrow \text{HNO}_3 + \text{MCO}_3$	$2.880 \cdot 10^{-12} \exp(-1862./T)$	
$\text{MCO}_3 + \text{CH}_3\text{O}_2 \longrightarrow 0.315 \cdot \text{CH}_3\text{CO}_3 + 0.585 \cdot \text{CH}_3\text{O}_2 + 0.585 \cdot \text{CO} + 1.9 \cdot \text{CH}_2\text{O} + 0.9 \cdot \text{CO}_2 + 0.9 \cdot \text{HO}_2 + 0.1 \cdot \text{MACO}_2\text{H}$	$1.000 \cdot 10^{-11}$	$\text{MCO}_3$ is an acyl radical, therefore $\text{kRO}_2\text{CH}_3\text{CO}_3$
$\text{MCO}_3 + \text{CH}_3\text{CO}_3 \longrightarrow \text{CH}_2\text{O} + 0.35 \cdot \text{CH}_3\text{CO}_3 + 1.65 \cdot \text{CH}_3\text{O}_2 + 0.65 \cdot \text{CO} + 2 \cdot \text{CO}_2$	$1.000 \cdot 10^{-11}$	
$\text{MCO}_3 + \text{HO}_2 \longrightarrow 0.44 \cdot \text{OH} + 0.154 \cdot \text{CH}_3\text{CO}_3 + 0.286 \cdot \text{CH}_3\text{O}_2 + 0.286 \cdot \text{CO} + 0.44 \cdot \text{CH}_2\text{O} + 0.44 \cdot \text{CO}_2 + 0.15 \cdot \text{MACO}_2\text{H} + 0.15 \cdot \text{O}_3 + 0.41 \cdot \text{MACO}_3\text{H} + 0.41 \cdot \text{O}_2$	$4.300 \cdot 10^{-13} \exp(1040./T)$	
$\text{MCO}_3 + \text{NO} \longrightarrow \text{CH}_2\text{O} + 0.35 \cdot \text{CH}_3\text{CO}_3 + 0.65 \cdot \text{CH}_3\text{O}_2 + 0.65 \cdot \text{CO} + \text{NO}_2 + \text{CO}_2$	$8.700 \cdot 10^{-12} \exp(290./T)$	
$\text{MCO}_3 + \text{NO}_3 \longrightarrow \text{CH}_2\text{O} + 0.35 \cdot \text{CH}_3\text{CO}_3 + 0.65 \cdot \text{CH}_3\text{O}_2 + 0.65 \cdot \text{CO} + \text{NO}_2 + \text{CO}_2$	$4.000 \cdot 10^{-12}$	

Table S9: C4 oxidation (... continued)

reaction	rate coefficient	reference
$\text{MCO}_3 + \text{NO}_2 + \text{M} \longrightarrow \text{M} + \text{MPAN}$	$ktroe(2.700 \cdot 10^{-28}, 7.1, 1.200 \cdot 10^{-11}, 0.9, 0.3)$	
$\text{MACRO}_2 + \text{CH}_3\text{O}_2 \longrightarrow 0.75 \cdot \text{CO} + 0.75 \cdot \text{HYAC} + \text{CH}_2\text{O} + 1.5 \cdot \text{HO}_2 + 0.25 \cdot \text{MACROH} + 0.25 \cdot \text{O}_2$	$9.200 \cdot 10^{-14}$	products: Tyndall (p.c.)
$\text{MACRO}_2 + \text{CH}_3\text{CO}_3 \longrightarrow 0.85 \cdot \text{CO} + 0.85 \cdot \text{HYAC} + 0.15 \cdot \text{CH}_2\text{O} + 0.15 \cdot \text{CH}_3\text{COCHO} + \text{HO}_2 + \text{CH}_3\text{O}_2 + \text{CO}_2$	$1.000 \cdot 10^{-11}$	
$\text{MACRO}_2 + \text{NO} \longrightarrow \text{NO}_2 + 0.85 \cdot \text{CO} + 0.85 \cdot \text{HYAC} + \text{HO}_2 + 0.15 \cdot \text{CH}_2\text{O} + 0.15 \cdot \text{CH}_3\text{COCHO}$	$2.464 \cdot 10^{-12} \exp(360./T)$	$A = 2.54 \cdot 10^{-12} \cdot 0.97$ ; 3% MACRN-yield inferred from HYAC 42% yield at high-NO from Crounse et al. (2011) (Tyndall, p.c.)
$\text{MACRO}_2 + \text{NO} \longrightarrow \text{MACRN}$	$7.620 \cdot 10^{-14} \exp(360./T)$	$A = 2.54 \cdot 10^{-12} \cdot 0.03$ ; 3% MACRN-yield inferred from HYAC 42% yield at high-NO from Crounse et al. (2011) (Tyndall, p.c.)
$\text{MACRO}_2 + \text{NO}_3 \longrightarrow 0.85 \cdot \text{CO} + 0.85 \cdot \text{HYAC} + 0.15 \cdot \text{CH}_2\text{O} + 0.15 \cdot \text{CH}_3\text{COCHO} + \text{HO}_2 + \text{NO}_2$	$2.500 \cdot 10^{-12}$	
$\text{MACRO}_2 + \text{HO}_2 \longrightarrow 0.6 \cdot \text{MACROOH} + 0.4 \cdot \text{CO} + 0.4 \cdot \text{HYAC} + 0.4 \cdot \text{HO}_2 + 0.4 \cdot \text{OH}$	$1.820 \cdot 10^{-13} \exp(1300./T)$	
$\text{MACRO}_2 \longrightarrow \text{CO} + \text{HYAC} + \text{OH}$	$2.900 \cdot 10^7 \exp(-5297./T)$	Isomerisation according to Crounse et al. (2012)
$\text{MACROOH} + \text{OH} \longrightarrow \text{CO} + \text{HYAC} + \text{OH} + \text{H}_2\text{O}$	$1.800 \cdot 10^{-11}$	
$\text{MACROH} + \text{OH} \longrightarrow \text{CO}_2 + \text{HYAC} + \text{HO}_2 + \text{H}_2\text{O}$	$1.800 \cdot 10^{-11}$	
$\text{MPAN} + \text{M} \longrightarrow \text{MCO}_3 + \text{NO}_2 + \text{M}$		
$\text{MPAN} + \text{OH} \longrightarrow \text{CO} + \text{HYAC} + \text{NO}_2$	$3.200 \cdot 10^{-11}$	rate: Orlando et al. (2002)
$\text{MACO}_2\text{H} + \text{OH} \longrightarrow \text{CH}_2\text{O} + \text{CH}_3\text{CO}_3 + \text{CO}_2 + \text{H}_2\text{O}$	$1.510 \cdot 10^{-11}$	products should be pyruvic acid + $\text{CH}_2\text{O}$ + $\text{HO}_2$
$\text{MACO}_3\text{H} + \text{OH} \longrightarrow \text{H}_2\text{O} + \text{MCO}_3$	$1.870 \cdot 10^{-11}$	
$\text{MVK} + \text{OH} \longrightarrow \text{LHMKABO}_2$	$4.130 \cdot 10^{-12} \exp(452./T)$	
$\text{MVK} + \text{O}_3 \longrightarrow 0.85 \cdot \text{CH}_3\text{COCHO} + 0.85 \cdot \text{HCOOH} + 0.15 \cdot \text{CH}_3\text{CO}_3 + 0.15 \cdot \text{OH} + 0.15 \cdot \text{CO} + 0.15 \cdot \text{CH}_2\text{O}$	$7.510 \cdot 10^{-16} \exp(-1521./T)$	products according to IUPAC (2006)
$\text{LHMKABO}_2 + \text{CH}_3\text{O}_2 \longrightarrow 0.9 \cdot \text{CH}_2\text{O} + 0.35 \cdot \text{GLYALD} + 0.65 \cdot \text{HO}_2 + 0.35 \cdot \text{CH}_3\text{CO}_3 + 0.175 \cdot \text{BIACETOH} + 0.25 \cdot \text{CH}_3\text{OH} + 0.25 \cdot \text{MACROH} + 0.15 \cdot \text{CH}_3\text{COCHO} + 0.075 \cdot \text{CO}_2\text{H}_3\text{CHO}$	$1.000 \cdot 10^{-12}$	products: Tyndall (p.c.)
$\text{LHMKABO}_2 + \text{CH}_3\text{CO}_3 \longrightarrow 0.3 \cdot \text{CH}_3\text{COCHO} + 0.7 \cdot \text{GLYALD} + 0.7 \cdot \text{CH}_3\text{CO}_3 + 0.3 \cdot \text{CH}_2\text{O} + 0.3 \cdot \text{HO}_2 + \text{CH}_3\text{O}_2 + \text{CO}_2$	$1.000 \cdot 10^{-11}$	

Table S9: C4 oxidation (... continued)

reaction	rate coefficient	reference
$\text{LHMKABO}_2 + \text{HO}_2 \longrightarrow 0.34 \cdot \text{LHMKABOOH} + 0.66 \cdot \text{CH}_3\text{CO}_3 + 0.66 \cdot \text{OH} + 0.66 \cdot \text{GLYALD}$	$1.820 \cdot 10^{-13} \exp(1300./T)$	Praske et al. (2015)
$\text{LHMKABO}_2 + \text{NO} \longrightarrow \text{NO}_2 + 0.3 \cdot \text{CH}_2\text{O} + 0.3 \cdot \text{CH}_3\text{COCHO} + 0.3 \cdot \text{HO}_2 + 0.7 \cdot \text{CH}_3\text{CO}_3 + 0.7 \cdot \text{GLYALD}$	$2.438 \cdot 10^{-12} \exp(360./T)$	$A = 2.54 \cdot 10^{-12} \cdot 0.96$ ; 4% nitrate yield Praske et al. (2015)
$\text{LHMKABO}_2 + \text{NO} \longrightarrow \text{MVKN}$	$1.020 \cdot 10^{-13} \exp(360./T)$	$A = 2.54 \cdot 10^{-12} \cdot 0.04$ ; 4% nitrate yield Praske et al. (2015)
$\text{LHMKABO}_2 + \text{NO}_3 \longrightarrow 0.3 \cdot \text{CH}_3\text{COCHO} + 0.7 \cdot \text{GLYALD} + 0.7 \cdot \text{CH}_3\text{CO}_3 + 0.3 \cdot \text{CH}_2\text{O} + 0.3 \cdot \text{HO}_2 + \text{NO}_2$	$2.500 \cdot 10^{-12}$	
$\text{MVKN} + \text{OH} \longrightarrow \text{CH}_3\text{COCHO} + \text{HO}_2 + \text{CO}_2 + \text{NO}_2 + \text{H}_2\text{O}$	$5.600 \cdot 10^{-12}$	k from Paulot et al. (2009); simplified products
$\text{MACRN} + \text{OH} \longrightarrow \text{CO}_2 + \text{HYAC} + \text{NO}_2 + \text{H}_2\text{O}$	$5.000 \cdot 10^{-11}$	k from Paulot et al. (2009); simplified products
$\text{LHMKABOOH} + \text{OH} \longrightarrow 0.3 \cdot \text{CO}_2\text{H}_3\text{CHO} + \text{OH} + \text{H}_2\text{O} + 0.7 \cdot \text{BIACETOH}$	$4.500 \cdot 10^{-12}$	
$\text{CO}_2\text{H}_3\text{CHO} + \text{OH} \longrightarrow \text{CO}_2\text{H}_3\text{CO}_3 + \text{H}_2\text{O}$	$2.450 \cdot 10^{-11}$	
$\text{CO}_2\text{H}_3\text{CHO} + \text{NO}_3 \longrightarrow \text{CO}_2\text{H}_3\text{CO}_3 + \text{HNO}_3$	$5.760 \cdot 10^{-12} \exp(-1862./T)$	
$\text{CO}_2\text{H}_3\text{CO}_3 + \text{CH}_3\text{O}_2 \longrightarrow \text{CH}_3\text{COCHO} + 2 \cdot \text{HO}_2 + \text{CO}_2 + \text{CH}_2\text{O}$	$1.000 \cdot 10^{-11}$	products: Tyndall (p.c.)
$\text{CO}_2\text{H}_3\text{CO}_3 + \text{CH}_3\text{CO}_3 \longrightarrow \text{CH}_3\text{COCHO} + \text{HO}_2 + \text{CH}_3\text{O}_2 + 2 \cdot \text{CO}_2$	$1.000 \cdot 10^{-11}$	
$\text{CO}_2\text{H}_3\text{CO}_3 + \text{HO}_2 \longrightarrow 0.6 \cdot \text{CO}_2\text{H}_3\text{CO}_3\text{H} + 0.4 \cdot \text{CO}_2 + 0.4 \cdot \text{OH} + 0.4 \cdot \text{HO}_2 + 0.4 \cdot \text{CH}_3\text{COCHO}$	$4.300 \cdot 10^{-13} \exp(1040./T)$	
$\text{CO}_2\text{H}_3\text{CO}_3 + \text{NO} \longrightarrow \text{CH}_3\text{COCHO} + \text{HO}_2 + \text{NO}_2 + \text{CO}_2$	$8.100 \cdot 10^{-12} \exp(270./T)$	
$\text{CO}_2\text{H}_3\text{CO}_3 + \text{NO}_3 \longrightarrow \text{CH}_3\text{COCHO} + \text{HO}_2 + \text{NO}_2 + \text{CO}_2$	$4.000 \cdot 10^{-12}$	
$\text{CO}_2\text{H}_3\text{CO}_3\text{H} + \text{OH} \longrightarrow \text{CO}_2\text{H}_3\text{CO}_3 + \text{H}_2\text{O}$	$1.000 \cdot 10^{-12}$	Orlando (p.c.)
$\text{MALO}_2 + \text{NO}_2 + \text{M} \longrightarrow 0.8 \cdot \text{LC}_5\text{PAN}_{1719} + \text{M}$	$ktroe(2.700 \cdot 10^{-28}, 7.1, 1.200 \cdot 10^{-11}, 0.9, 0.6)$	Orlando and Tyndall (2012) - Same as $k(\text{CH}_3\text{CO}_3 + \text{NO}_2)$ ; $\text{LC}_5\text{PAN}$ as a surrogate
$\text{MALO}_2 + \text{NO} \longrightarrow 0.4 \cdot \text{GLYOXAL} + \text{HO}_2 + 0.4 \cdot \text{CO} + 0.4 \cdot \text{CO}_2 + \text{NO}_2 + 0.6 \cdot \text{CO}_2\text{H}_3\text{CHO}$	$7.500 \cdot 10^{-12} \exp(290./T)$	products: Tyndall (p.c.)
$\text{MALO}_2 + \text{HO}_2 \longrightarrow 0.16 \cdot \text{GLYOXAL} + \text{HO}_2 + 0.16 \cdot \text{CO} + 0.16 \cdot \text{CO}_2 + 0.16 \cdot \text{OH} + 0.84 \cdot \text{CO}_2\text{H}_3\text{CHO}$	$4.300 \cdot 10^{-13} \exp(1040./T)$	
$\text{MALO}_2 + \text{CH}_3\text{O}_2 \longrightarrow 0.4 \cdot \text{GLYOXAL} + 2 \cdot \text{HO}_2 + 0.4 \cdot \text{CO} + 0.4 \cdot \text{CO}_2 + 0.6 \cdot \text{CO}_2\text{H}_3\text{CHO} + \text{CH}_2\text{O}$	$1.000 \cdot 10^{-12}$	products: Tyndall (p.c.)
$\text{MALO}_2 + \text{CH}_3\text{CO}_3 \longrightarrow 0.4 \cdot \text{GLYOXAL} + \text{HO}_2 + 0.4 \cdot \text{CO} + 1.4 \cdot \text{CO}_2 + 0.6 \cdot \text{CO}_2\text{H}_3\text{CHO} + \text{CH}_3\text{O}_2$	$1.000 \cdot 10^{-11}$	



Table S9: C4 oxidation (... continued)

reaction	rate coefficient	reference
MDIALO <sub>2</sub> + HO <sub>2</sub> $\longrightarrow$ 0.4 · OH + 0.332 · HO <sub>2</sub> + 0.068 · CH <sub>3</sub> COCHO + 0.136 · CO + 0.068 · CH <sub>3</sub> O <sub>2</sub> + 0.068 · GLYOXAL	$4.300 \cdot 10^{-13} \exp(1040./T)$	
MDIALO <sub>2</sub> + NO $\longrightarrow$ NO <sub>2</sub> + 0.83 · HO <sub>2</sub> + 0.17 · CH <sub>3</sub> COCHO + 0.34 · CO + 0.17 · CH <sub>3</sub> O <sub>2</sub> + 0.17 · GLYOXAL	$7.500 \cdot 10^{-12} \exp(290./T)$	
MDIALO <sub>2</sub> + NO <sub>2</sub> + M $\longrightarrow$ M	$ktroe(2.700 \cdot 10^{-28}, 7.1, 1.200 \cdot 10^{-11}, 0.9, 0.6)$	Orlando and Tyndall (2012); Same as k(CH <sub>3</sub> CO <sub>3</sub> + NO <sub>2</sub> )
MDIALO <sub>2</sub> + CH <sub>3</sub> O <sub>2</sub> $\longrightarrow$ 1.83 · HO <sub>2</sub> + 0.17 · CH <sub>3</sub> COCHO + 0.34 · CO + 0.17 · CH <sub>3</sub> O <sub>2</sub> + 0.17 · GLYOXAL + CH <sub>2</sub> O	$1.000 \cdot 10^{-12}$	
MDIALO <sub>2</sub> + CH <sub>3</sub> CO <sub>3</sub> $\longrightarrow$ 0.83 · HO <sub>2</sub> + 0.17 · CH <sub>3</sub> COCHO + 0.34 · CO + 1.17 · CH <sub>3</sub> O <sub>2</sub> + 0.17 · GLYOXAL + CO <sub>2</sub>	$1.000 \cdot 10^{-11}$	

Table S10: C5 oxidation

reaction	rate coefficient	reference
BIGALKANE + OH $\longrightarrow$ ALKO <sub>2</sub> + H <sub>2</sub> O	$3.500 \cdot 10^{-12}$	
C <sub>5</sub> H <sub>8</sub> + OH $\longrightarrow$ 0.4 · LISOPACO <sub>2</sub> + 0.35 · ISOPBO <sub>2</sub> + 0.25 · ISOPDO <sub>2</sub>	$2.700 \cdot 10^{-11} \exp(390./T)$	Tyndall (p.c.); MCM3.2 has yields .25, .5, .25
C <sub>5</sub> H <sub>8</sub> + O <sub>3</sub> $\longrightarrow$ 0.051 · CH <sub>3</sub> O <sub>2</sub> + 0.1575 · CH <sub>3</sub> CO <sub>3</sub> + 0.054 · LHMVKABO <sub>2</sub> + 0.522 · CO + 0.068750 · HCOOH + 0.11 · H <sub>2</sub> O <sub>2</sub> + 0.324750 · MACR + 0.1275 · C <sub>3</sub> H <sub>6</sub> + 0.2625 · HO <sub>2</sub> + 0.255 · CO <sub>2</sub> + 0.749750 · CH <sub>2</sub> O + 0.041250 · MACO <sub>2</sub> H + 0.27 · OH + 0.244 · MVK	$7.860 \cdot 10^{-15} \exp(-1913./T)$	
C <sub>5</sub> H <sub>8</sub> + NO <sub>3</sub> $\longrightarrow$ NISOPO <sub>2</sub>	$3.030 \cdot 10^{-12} \exp(-446./T)$	
MBO + OH $\longrightarrow$ MBOO <sub>2</sub>	$8.100 \cdot 10^{-12} \exp(610./T)$	
MBO + O <sub>3</sub> $\longrightarrow$ 0.35 · CO + 0.5 · CH <sub>2</sub> O + 0.1 · CH <sub>3</sub> COCH <sub>3</sub> + 0.9 · IBUTALOH + 0.25 · HCOOH + 0.06 · HO <sub>2</sub> + 0.06 · OH	$1.000 \cdot 10^{-17}$	
MBO + NO <sub>3</sub> $\longrightarrow$ MBONO <sub>3</sub> O <sub>2</sub>	$4.600 \cdot 10^{-14} \exp(-400./T)$	
ALKO <sub>2</sub> + NO $\longrightarrow$ 0.4 · CH <sub>3</sub> CHO + 0.25 · CH <sub>2</sub> O + 0.25 · CH <sub>3</sub> COCH <sub>3</sub> + HO <sub>2</sub> + 0.8 · MEK + NO <sub>2</sub>	$3.780 \cdot 10^{-12} \exp(180./T)$	$A = 4.200 \cdot 10^{-12} \cdot 0.9$ ; 10% ALKNO <sub>3</sub> -yield
ALKO <sub>2</sub> + NO $\longrightarrow$ ALKNO <sub>3</sub>	$4.200 \cdot 10^{-13} \exp(180./T)$	10% ALKNO <sub>3</sub> -yield
ALKO <sub>2</sub> + HO <sub>2</sub> $\longrightarrow$ ALKOOH	$7.500 \cdot 10^{-13} \exp(700./T)$	
ALKO <sub>2</sub> + CH <sub>3</sub> O <sub>2</sub> $\longrightarrow$ 0.3 · CH <sub>3</sub> CHO + 1.1875 · CH <sub>2</sub> O + 0.1875 · CH <sub>3</sub> COCH <sub>3</sub> + 0.75 · HO <sub>2</sub> + 0.6 · MEK + 0.25 · ALKOH	$1.000 \cdot 10^{-12}$	

Table S10: C5 oxidation (... continued)

reaction	rate coefficient	reference
$\text{ALKO}_2 + \text{CH}_3\text{CO}_3 \longrightarrow 0.4 \cdot \text{CH}_3\text{CHO} + 0.25 \cdot \text{CH}_2\text{O} + 0.25 \cdot \text{CH}_3\text{COCH}_3 + \text{HO}_2 + 0.8 \cdot \text{MEK} + \text{CH}_3\text{O}_2 + \text{CO}_2$	$1.000 \cdot 10^{-11}$	
$\text{ALKOOH} + \text{OH} \longrightarrow \text{ALKO}_2 + \text{H}_2\text{O}$	$3.800 \cdot 10^{-12} \exp(200./T)$	
$\text{ALKOH} + \text{OH} \longrightarrow 1.25 \cdot \text{MEK} + \text{HO}_2 + \text{H}_2\text{O}$	$5.000 \cdot 10^{-12}$	Tyndall (p.c.), MEK yield to account for C
$\text{ALKNO}_3 + \text{OH} \longrightarrow 0.4 \cdot \text{CH}_3\text{CHO} + 0.25 \cdot \text{CH}_2\text{O} + 0.25 \cdot \text{CH}_3\text{COCH}_3 + \text{HO}_2 + 0.8 \cdot \text{MEK} + \text{NO}_2$	$2.000 \cdot 10^{-12}$	
$\text{LISOPACO}_2 + \text{HO}_2 \longrightarrow \text{LISOPACOOH}$	$2.050 \cdot 10^{-13} \exp(1300./T)$	add OH channel
$\text{LISOPACO}_2 + \text{NO} \longrightarrow \text{HO}_2 + 0.977 \cdot \text{LHC}_4\text{ACCHO} + \text{NO}_2 + 0.0277 \cdot \text{CH}_3\text{COCHO} + 0.0277 \cdot \text{GLYOXAL} + 0.0277 \cdot \text{HYAC} + 0.0277 \cdot \text{GLYALD}$	$2.235 \cdot 10^{-12} \exp(360./T)$	$A = 2.54 \cdot 10^{-12} \cdot 0.88$ ; average 12% nitrate yield from Paulot et al. (2009); Tyndall (p.c.); direct GLYOXAL channel from lab meas.
$\text{LISOPACO}_2 + \text{NO} \longrightarrow \text{LISOPACNO}_3$	$3.050 \cdot 10^{-13} \exp(360./T)$	$A = 2.54 \cdot 10^{-12} \cdot 0.12$ ; average 12% nitrate yield from Paulot et al. (2009)
$\text{LISOPACO}_2 + \text{NO}_3 \longrightarrow \text{HO}_2 + \text{LHC}_4\text{ACCHO} + \text{NO}_2$	$2.500 \cdot 10^{-12}$	
$\text{LISOPACO}_2 + \text{CH}_3\text{O}_2 \longrightarrow 0.75 \cdot \text{CH}_2\text{O} + 0.75 \cdot \text{LHC}_4\text{ACCHO} + 0.25 \cdot \text{CH}_3\text{OH} + 0.25 \cdot \text{ISOPAOH} + \text{HO}_2$	$2.400 \cdot 10^{-12}$	products: Tyndall (p.c.)
$\text{LISOPACO}_2 + \text{CH}_3\text{CO}_3 \longrightarrow \text{HO}_2 + \text{LHC}_4\text{ACCHO} + \text{CH}_3\text{O}_2 + \text{CO}_2$	$1.000 \cdot 10^{-11}$	
$\text{LISOPACO}_2 \longrightarrow \text{HO}_2 + \text{HPALD}$	$4.1 \cdot 10^8 \exp(-7700./T)$	Bulk isomerization rate constant for all ISOPO2 by Crounse et al. (2011)
$\text{ISOPBO}_2 \longrightarrow \text{HO}_2 + \text{HPALD}$	$4.1 \cdot 10^8 \exp(-7700./T)$	see note of [LISOPACO2]
$\text{ISOPDO}_2 \longrightarrow \text{HO}_2 + \text{HPALD}$	$4.1 \cdot 10^8 \exp(-7700./T)$	see note of [LISOPACO2]
$\text{LISOPACOOH} + \text{OH} \longrightarrow 0.415 \cdot \text{LIEPOX} + 0.415 \cdot \text{OH} + 0.415 \cdot \text{LISOPOOHO}_2 + 0.14 \cdot \text{LHC}_4\text{ACCHO} + 0.03 \cdot \text{H}_2\text{O} + 0.03 \cdot \text{LISOPACO}_2$	$1.540 \cdot 10^{-10}$	k from MCMv3.3.1 and OH-addition branching ratios estimated with site-specific SAR by Peeters et al. (2007) and H-abstraction channel assumed to be like the one for $\text{CH}_3\text{OOH} + \text{OH}$ reaction and abstraction from the alpha-hydroperoxyl allyl hydrogen estimated by SAR of MOM (Taraborrelli in prep.) being $2.12 \cdot 10^{-11}$
$\text{ISOPAOH} + \text{OH} \longrightarrow \text{LISOPOOHO}_2$	$9.300 \cdot 10^{-11}$	OH-addition to double bond and products approximated with the one from $\text{ISOPOOH} + \text{OH}$ reaction leading to similar SOA precursors.
$\text{LISOPACNO}_3 + \text{OH} \longrightarrow \text{LISOPNO}_3\text{O}_2$	$6.000 \cdot 10^{-11}$	
$\text{LIEPOX} + \text{OH} \longrightarrow 0.29 \cdot \text{IEC}_1\text{O}_2 + 0.71 \cdot \text{LIECHO} + 0.71 \cdot \text{HO}_2 + \text{H}_2\text{O}$	$1.500 \cdot 10^{-11}$	MCM3.2
$\text{LIECHO} + \text{OH} \longrightarrow \text{H}_2\text{O} + \text{LIECO}_3$	$1.760 \cdot 10^{-11}$	MCM3.2
$\text{LIECHO} + \text{NO}_3 \longrightarrow \text{HNO}_3 + \text{LIECO}_3$	$1.050 \cdot 10^{-11} \exp(-1860./T)$	MCM3.2

Table S10: C5 oxidation (... continued)

reaction	rate coefficient	reference
$\text{LIECO}_3 + \text{HO}_2 \longrightarrow 0.6 \cdot \text{LIECO}_3\text{H} + 0.4 \cdot \text{CO}_2 + 0.4 \cdot \text{OH} + 0.25 \cdot \text{LHMVKABO}_2 + 0.15 \cdot \text{MACRO}_2$	$5.200 \cdot 10^{-13} \exp(980./T)$	MCM3.2
$\text{LIECO}_3 + \text{NO} \longrightarrow 0.6 \cdot \text{LHMVKABO}_2 + 0.4 \cdot \text{MACRO}_2 + \text{NO}_2 + \text{CO}_2$	$7.500 \cdot 10^{-12} \exp(290./T)$	MCM3.2
$\text{LIECO}_3 + \text{NO}_3 \longrightarrow 0.6 \cdot \text{LHMVKABO}_2 + 0.4 \cdot \text{MACRO}_2 + \text{NO}_2 + \text{CO}_2$	$4.000 \cdot 10^{-12}$	MCM3.2
$\text{LIECO}_3\text{H} + \text{OH} \longrightarrow \text{H}_2\text{O} + \text{LIECO}_3$	$1.040 \cdot 10^{-11}$	MCM3.2
$\text{IEC}_1\text{O}_2 + \text{HO}_2 \longrightarrow \text{LIECO}_3\text{H}$	$2.050 \cdot 10^{-13} \exp(1300./T)$	MCM3.2 plus shortcut
$\text{IEC}_1\text{O}_2 + \text{NO} \longrightarrow \text{BIACETOH} + \text{NO}_2 + \text{CH}_2\text{O} + \text{HO}_2$	$2.700 \cdot 10^{-12} \exp(360./T)$	MCM3.2
$\text{IEC}_1\text{O}_2 + \text{NO}_3 \longrightarrow \text{BIACETOH} + \text{NO}_2 + \text{CH}_2\text{O} + \text{HO}_2$	$2.300 \cdot 10^{-12}$	MCM3.2
$\text{ISOPBO}_2 + \text{HO}_2 \longrightarrow \text{ISOPBOOH}$	$2.050 \cdot 10^{-13} \exp(1300./T)$	
$\text{ISOPBO}_2 + \text{NO} \longrightarrow \text{CH}_2\text{O} + \text{MVK} + \text{HO}_2 + \text{NO}_2$	$2.235 \cdot 10^{-12} \exp(360./T)$	$A = 2.54 \cdot 10^{-12} \cdot 0.88$ ; average 12% nitrate yield from Paulot et al. (2009);
$\text{ISOPBO}_2 + \text{NO} \longrightarrow \text{ISOPBNO}_3$	$3.050 \cdot 10^{-13} \exp(360./T)$	$A = 2.54 \cdot 10^{-12} \cdot 0.12$ ; average 12% nitrate yield from Paulot et al. (2009);
$\text{ISOPBO}_2 + \text{NO}_3 \longrightarrow \text{CH}_2\text{O} + \text{MVK} + \text{HO}_2 + \text{NO}_2$	$2.500 \cdot 10^{-12}$	
$\text{ISOPBO}_2 + \text{CH}_3\text{O}_2 \longrightarrow 0.75 \cdot \text{MVK} + 1.75 \cdot \text{CH}_2\text{O} + 1.5 \cdot \text{HO}_2 + 0.25 \cdot \text{ISOPBOH}$	$8.000 \cdot 10^{-13}$	products: Tyndall (p.c.)
$\text{ISOPBO}_2 + \text{CH}_3\text{CO}_3 \longrightarrow \text{CH}_2\text{O} + \text{MVK} + \text{HO}_2 + \text{CH}_3\text{O}_2 + \text{CO}_2$	$1.000 \cdot 10^{-11}$	
$\text{ISOPBOOH} + \text{OH} \longrightarrow 0.63 \cdot \text{LIEPOX} + 0.63 \cdot \text{OH} + 0.12 \cdot \text{LISOPOOHO}_2 + 0.15 \cdot \text{H}_2\text{O} + 0.15 \cdot \text{ISOPBO}_2$	$7.500 \cdot 10^{-11}$	St. Clair et al. (2015)
$\text{ISOPBOH} + \text{OH} \longrightarrow \text{LISOPOOHO}_2$	$3.850 \cdot 10^{-11}$	OH-addition to double bond and products approximated with the one from $\text{ISOPOOH} + \text{OH}$ reaction leading to similar SOA precursors.
$\text{ISOPBNO}_3 + \text{OH} \longrightarrow \text{LISOPNO}_3\text{O}_2$	$1.360 \cdot 10^{-11}$	
$\text{ISOPDO}_2 + \text{HO}_2 \longrightarrow \text{ISOPDOOH}$	$2.050 \cdot 10^{-13} \exp(1300./T)$	
$\text{ISOPDO}_2 + \text{NO} \longrightarrow \text{CH}_2\text{O} + \text{MACR} + \text{HO}_2 + \text{NO}_2$	$2.235 \cdot 10^{-12} \exp(360./T)$	$A = 2.54 \cdot 10^{-12} \cdot 0.88$ ; average 12% nitrate yield from Paulot et al. (2009);
$\text{ISOPDO}_2 + \text{NO} \longrightarrow \text{ISOPDNO}_3$	$3.050 \cdot 10^{-13} \exp(360./T)$	$A = 2.54 \cdot 10^{-12} \cdot 0.12$ ; average 12% nitrate yield from Paulot et al. (2009);
$\text{ISOPDO}_2 + \text{NO}_3 \longrightarrow \text{CH}_2\text{O} + \text{MACR} + \text{HO}_2 + \text{NO}_2$	$2.500 \cdot 10^{-12}$	
$\text{ISOPDO}_2 + \text{CH}_3\text{O}_2 \longrightarrow 0.5 \cdot \text{MACR} + 1.25 \cdot \text{CH}_2\text{O} + \text{HO}_2 + 0.25 \cdot \text{CH}_3\text{OH} + 0.25 \cdot \text{HCOC}_5 + 0.25 \cdot \text{ISOPDOH}$	$2.900 \cdot 10^{-12}$	products: Tyndall (p.c.)
$\text{ISOPDO}_2 + \text{CH}_3\text{CO}_3 \longrightarrow \text{CH}_2\text{O} + \text{MACR} + \text{HO}_2 + \text{CH}_3\text{O}_2 + \text{CO}_2$	$1.000 \cdot 10^{-11}$	

Table S10: C5 oxidation (... continued)

reaction	rate coefficient	reference
ISOPDOOH + OH $\longrightarrow$ 0.79 · LIEPOX + 0.79 · OH + 0.14 · LISOPOOHO <sub>2</sub> + 0.07 · H <sub>2</sub> O + 0.07 · ISOPDO <sub>2</sub>	$1.180 \cdot 10^{-10}$	St. Clair et al. (2015)
ISOPDOH + OH $\longrightarrow$ LISOPOOHO <sub>2</sub>	$7.380 \cdot 10^{-11}$	OH-addition to double bond and products approximated with the one from ISOPOOH + OH reaction leading to sim- ilar SOA precursors.
ISOPDNO <sub>3</sub> + OH $\longrightarrow$ LISOPNO <sub>3</sub> O <sub>2</sub>	$6.100 \cdot 10^{-11}$	OH-addition to double bond
NISOP <sub>2</sub> + HO <sub>2</sub> $\longrightarrow$ NISOPOOH	$2.050 \cdot 10^{-13} \exp(1300./T)$	
NISOP <sub>2</sub> + NO $\longrightarrow$ HO <sub>2</sub> + NC <sub>4</sub> CHO + NO <sub>2</sub>	$2.540 \cdot 10^{-12} \exp(360./T)$	
NISOP <sub>2</sub> + NO <sub>3</sub> $\longrightarrow$ HO <sub>2</sub> + NC <sub>4</sub> CHO + NO <sub>2</sub>	$2.500 \cdot 10^{-12}$	
NISOP <sub>2</sub> + CH <sub>3</sub> O <sub>2</sub> $\longrightarrow$ 0.75 · CH <sub>2</sub> O + 0.75 · NC <sub>4</sub> CHO + HO <sub>2</sub> + 0.25 · CH <sub>3</sub> OH + 0.25 · LISOPACNO <sub>3</sub>	$1.300 \cdot 10^{-12}$	products: Tyndall (p.c.)
NISOP <sub>2</sub> + CH <sub>3</sub> CO <sub>3</sub> $\longrightarrow$ HO <sub>2</sub> + NC <sub>4</sub> CHO + CH <sub>3</sub> O <sub>2</sub> + CO <sub>2</sub>	$1.000 \cdot 10^{-11}$	
NISOPOOH + OH $\longrightarrow$ NC <sub>4</sub> CHO + OH + H <sub>2</sub> O	$1.030 \cdot 10^{-10}$	
NC <sub>4</sub> CHO + OH $\longrightarrow$ H <sub>2</sub> O + LNISO <sub>3</sub>	$4.160 \cdot 10^{-11}$	
NC <sub>4</sub> CHO + O <sub>3</sub> $\longrightarrow$ 0.445 · NO <sub>2</sub> + 0.89 · CO + 0.075625 · H <sub>2</sub> O <sub>2</sub> + 0.034375 · HCOCO <sub>2</sub> H + 0.555 · NOA + 0.445 · HO <sub>2</sub> + 0.520625 · GLYOXAL + 0.89 · OH + 0.445 · CH <sub>3</sub> COCHO	$2.400 \cdot 10^{-17}$	
NC <sub>4</sub> CHO + NO <sub>3</sub> $\longrightarrow$ HNO <sub>3</sub> + LNISO <sub>3</sub>	$6.120 \cdot 10^{-12} \exp(-1862./T)$	
LNISO <sub>3</sub> + HO <sub>2</sub> $\longrightarrow$ 0.8 · LNISOOH + 0.2 · NOA + 0.2 · OH + 0.2 · CO <sub>2</sub> + 0.2 · CO + 0.2 · HO <sub>2</sub>	$1.930 \cdot 10^{-13} \exp(1300./T)$	products: Tyndall (p.c.)
LNISO <sub>3</sub> + NO $\longrightarrow$ NOA + 0.5 · GLYOXAL + 0.5 · CO + HO <sub>2</sub> + NO <sub>2</sub> + 0.5 · CO <sub>2</sub>	$4.270 \cdot 10^{-12} \exp(360./T)$	
LNISO <sub>3</sub> + NO <sub>3</sub> $\longrightarrow$ NOA + 0.5 · GLYOXAL + 0.5 · CO + HO <sub>2</sub> + NO <sub>2</sub> + 0.5 · CO <sub>2</sub>	$3.302 \cdot 10^{-12} \exp(360./T)$	
LNISO <sub>3</sub> + CH <sub>3</sub> O <sub>2</sub> $\longrightarrow$ 0.375 · GLYOXAL + 0.875 · NOA + CH <sub>2</sub> O + 1.75 · HO <sub>2</sub> + 0.625 · CO <sub>2</sub> + 0.0625 · MACRN + 0.0625 · MVKN + 0.5 · CO	$1.000 \cdot 10^{-12}$	products: Tyndall (p.c.)
LNISO <sub>3</sub> + CH <sub>3</sub> CO <sub>3</sub> $\longrightarrow$ NOA + 0.5 · GLYOXAL + 0.5 · CO + HO <sub>2</sub> + CH <sub>3</sub> O <sub>2</sub> + 1.5 · CO <sub>2</sub>	$1.000 \cdot 10^{-11}$	
LNISOOH + OH $\longrightarrow$ H <sub>2</sub> O + LNISO <sub>3</sub>	$2.650 \cdot 10^{-11}$	
LHC <sub>4</sub> ACCHO + OH $\longrightarrow$ 0.52 · LC <sub>578</sub> O <sub>2</sub> + 0.48 · LHC <sub>4</sub> ACCO <sub>3</sub> + H <sub>2</sub> O	$4.520 \cdot 10^{-11}$	

Table S10: C5 oxidation (... continued)

reaction	rate coefficient	reference
$\text{LHC}_4\text{ACCHO} + \text{O}_3 \longrightarrow 0.2225 \cdot \text{CH}_3\text{CO}_3 +$ $0.89 \cdot \text{CO} + 0.017188 \cdot \text{HOCH}_2\text{CO}_2\text{H} + 0.075625 \cdot \text{H}_2\text{O}_2 +$ $0.017188 \cdot \text{HCOCO}_2\text{H} + 0.2775 \cdot \text{HYAC} + 0.6675 \cdot \text{HO}_2 +$ $0.260313 \cdot \text{GLYOXAL} + 0.2225 \cdot \text{CH}_2\text{O} + 0.89 \cdot \text{OH} +$ $0.260313 \cdot \text{GLYALD} + 0.5 \cdot \text{CH}_3\text{COCHO}$	$2.400 \cdot 10^{-17}$	
$\text{LHC}_4\text{ACCHO} + \text{NO}_3 \longrightarrow \text{HNO}_3 + \text{LHC}_4\text{ACCO}_3$	$6.120 \cdot 10^{-12} \exp(-1862./T)$	
$\text{LC}_{578}\text{O}_2 + \text{NO} \longrightarrow 0.25 \cdot \text{CH}_3\text{COCHO} + 0.25 \cdot \text{GLYALD} +$ $0.25 \cdot \text{GLYOXAL} + 0.25 \cdot \text{HYAC} + 0.25 \cdot \text{CO} + 0.25 \cdot \text{MACROH} +$ $0.25 \cdot \text{CH}_2\text{O} + 0.25 \cdot \text{CO}_2\text{H}_3\text{CHO} + \text{HO}_2 + \text{NO}_2$	$2.540 \cdot 10^{-12} \exp(360./T)$	products: Tyndall (p.c.)
$\text{LC}_{578}\text{O}_2 + \text{NO}_3 \longrightarrow 0.25 \cdot \text{CH}_3\text{COCHO} + 0.25 \cdot \text{GLYALD} +$ $0.25 \cdot \text{GLYOXAL} + 0.25 \cdot \text{HYAC} + 0.25 \cdot \text{CO} + 0.25 \cdot \text{MACROH} +$ $0.25 \cdot \text{CH}_2\text{O} + 0.25 \cdot \text{CO}_2\text{H}_3\text{CHO} + \text{HO}_2 + \text{NO}_2$	$2.500 \cdot 10^{-12}$	products: Tyndall (p.c.)
$\text{LC}_{578}\text{O}_2 + \text{HO}_2 \longrightarrow \text{LC}_{578}\text{OOH}$	$2.050 \cdot 10^{-13} \exp(1300./T)$	add OH channel
$\text{LC}_{578}\text{O}_2 + \text{CH}_3\text{O}_2 \longrightarrow 0.156250 \cdot \text{CH}_3\text{COCHO} +$ $0.156250 \cdot \text{GLYALD} + 0.156250 \cdot \text{GLYOXAL} + 0.156250 \cdot \text{HYAC} +$ $0.156250 \cdot \text{CO} + 0.468750 \cdot \text{MACROH} + 1.031250 \cdot \text{CH}_2\text{O} +$ $0.3125 \cdot \text{CO}_2\text{H}_3\text{CHO} + 1.25 \cdot \text{HO}_2 + 0.125 \cdot \text{CH}_3\text{OH}$	$1.000 \cdot 10^{-12}$	products: Tyndall (p.c.)
$\text{LC}_{578}\text{O}_2 + \text{CH}_3\text{CO}_3 \longrightarrow 0.25 \cdot \text{CH}_3\text{COCHO} + 0.25 \cdot \text{GLYALD} +$ $0.25 \cdot \text{GLYOXAL} + 0.25 \cdot \text{HYAC} + 0.25 \cdot \text{CO} + 0.25 \cdot \text{MACROH} +$ $0.25 \cdot \text{CH}_2\text{O} + 0.25 \cdot \text{CO}_2\text{H}_3\text{CHO} + \text{CH}_3\text{O}_2 + \text{CO}_2$	$1.000 \cdot 10^{-11}$	
$\text{LC}_{578}\text{OOH} + \text{OH} \longrightarrow \text{H}_2\text{O} + \text{LC}_{578}\text{O}_2$	$3.160 \cdot 10^{-11}$	
$\text{LHC}_4\text{ACCO}_3 \longrightarrow \text{HO}_2 + \text{PACALD}$	$4.1 \cdot 10^8 \exp(-7700./T)$	see note of [LISOPACO2]
$\text{LHC}_4\text{ACCO}_3 + \text{CH}_3\text{O}_2 \longrightarrow \text{CH}_2\text{O} + 0.1 \cdot \text{LHC}_4\text{ACCO}_2\text{H} +$ $0.45 \cdot \text{GLYALD} + 0.45 \cdot \text{HYAC} + 0.45 \cdot \text{CH}_3\text{CO}_3 + 0.45 \cdot \text{CO} +$ $0.45 \cdot \text{HO}_2 + 0.9 \cdot \text{CO}_2$	$1.000 \cdot 10^{-11}$	products: Tyndall (p.c.)
$\text{LHC}_4\text{ACCO}_3 + \text{CH}_3\text{CO}_3 \longrightarrow 0.5 \cdot \text{HYAC} + 0.5 \cdot \text{GLYALD} +$ $0.5 \cdot \text{CH}_3\text{CO}_3 + 0.5 \cdot \text{CO} + 0.5 \cdot \text{HO}_2 + \text{CH}_3\text{O}_2 + 2 \cdot \text{CO}_2$	$1.000 \cdot 10^{-11}$	
$\text{LHC}_4\text{ACCO}_3 + \text{HO}_2 \longrightarrow 0.4 \cdot \text{LHC}_4\text{ACCO}_3\text{H} +$ $0.2 \cdot \text{LHC}_4\text{ACCO}_2\text{H} + 0.2 \cdot \text{O}_3 + 0.4 \cdot \text{CO}_2 + 0.4 \cdot \text{OH} +$ $0.2 \cdot \text{HYAC} + 0.2 \cdot \text{GLYALD} + 0.2 \cdot \text{CH}_3\text{CO}_3 + 0.2 \cdot \text{CO} +$ $0.2 \cdot \text{HO}_2$	$4.300 \cdot 10^{-13} \exp(1040./T)$	
$\text{LHC}_4\text{ACCO}_3 + \text{NO} \longrightarrow 0.5 \cdot \text{HYAC} + 0.5 \cdot \text{GLYALD} +$ $0.5 \cdot \text{CH}_3\text{CO}_3 + 0.5 \cdot \text{CO} + 0.5 \cdot \text{HO}_2 + \text{NO}_2 + \text{CO}_2$	$8.100 \cdot 10^{-12} \exp(270./T)$	
$\text{LHC}_4\text{ACCO}_3 + \text{NO}_3 \longrightarrow 0.5 \cdot \text{HYAC} + 0.5 \cdot \text{GLYALD} +$ $0.5 \cdot \text{CH}_3\text{CO}_3 + 0.5 \cdot \text{CO} + 0.5 \cdot \text{HO}_2 + \text{NO}_2 + \text{CO}_2$	$4.000 \cdot 10^{-12}$	

Table S10: C5 oxidation (... continued)

reaction	rate coefficient	reference
$\text{LHC}_4\text{ACCO}_3 + \text{NO}_2 + \text{M} \longrightarrow \text{LC}_5\text{PAN}_{1719} + \text{M}$	$ktroe(2.700 \cdot 10^{-28}, 7.1, 1.200 \cdot 10^{-11}, 0.9, 0.3)$	
$\text{LHC}_4\text{ACCO}_2\text{H} + \text{OH} \longrightarrow 0.5 \cdot \text{HYAC} + 0.5 \cdot \text{GLYALD} + 0.5 \cdot \text{CH}_3\text{CO}_3 + 0.5 \cdot \text{CO} + 0.5 \cdot \text{HO}_2 + \text{CO}_2 + \text{H}_2\text{O}$	$2.520 \cdot 10^{-11}$	
$\text{LHC}_4\text{ACCO}_3\text{H} + \text{OH} \longrightarrow \text{H}_2\text{O} + \text{LHC}_4\text{ACCO}_3$	$2.880 \cdot 10^{-11}$	
$\text{LC}_5\text{PAN}_{1719} + \text{M} \longrightarrow \text{LHC}_4\text{ACCO}_3 + \text{NO}_2 + \text{M}$		$k_0 = 4.9 \cdot 10^{-3} \exp(-12100./T) * M, k_{inf} = 5.4 \cdot 10^{16} \exp(-13830./T), f_c = 0.3$
$\text{LC}_5\text{PAN}_{1719} + \text{OH} \longrightarrow \text{CO} + \text{MACROH} + \text{NO}_2$	$2.520 \cdot 10^{-11}$	
$\text{HCOC}_5 + \text{OH} \longrightarrow \text{C}_{59}\text{O}_2$	$3.810 \cdot 10^{-11}$	
$\text{C}_{59}\text{O}_2 + \text{CH}_3\text{O}_2 \longrightarrow 0.75 \cdot \text{HOCH}_2\text{CO}_3 + 0.75 \cdot \text{HYAC} + \text{CH}_2\text{O} + 0.75 \cdot \text{HO}_2 + 0.3125 \cdot \text{MACROH}$	$1.000 \cdot 10^{-12}$	Tyndall (p.c.)
$\text{C}_{59}\text{O}_2 + \text{CH}_3\text{CO}_3 \longrightarrow \text{HOCH}_2\text{CO}_3 + \text{HYAC} + \text{CH}_3\text{O}_2 + \text{CO}_2$	$1.000 \cdot 10^{-11}$	
$\text{C}_{59}\text{O}_2 + \text{NO} \longrightarrow \text{HOCH}_2\text{CO}_3 + \text{HYAC} + \text{NO}_2$	$2.540 \cdot 10^{-12} \exp(360./T)$	
$\text{C}_{59}\text{O}_2 + \text{NO}_3 \longrightarrow \text{HOCH}_2\text{CO}_3 + \text{HYAC} + \text{NO}_2$	$2.500 \cdot 10^{-12}$	
$\text{C}_{59}\text{O}_2 + \text{HO}_2 \longrightarrow \text{C}_{59}\text{OOH}$	$2.050 \cdot 10^{-13} \exp(1300./T)$	
$\text{C}_{59}\text{OOH} + \text{OH} \longrightarrow \text{C}_{59}\text{O}_2 + \text{H}_2\text{O}$	$9.700 \cdot 10^{-12}$	
$\text{MBOO}_2 + \text{NO} \longrightarrow \text{HO}_2 + 0.67 \cdot \text{CH}_3\text{COCH}_3 + 0.67 \cdot \text{GLYALD} + 0.33 \cdot \text{CH}_2\text{O} + 0.33 \cdot \text{IBUTALOH} + \text{NO}_2$	$2.600 \cdot 10^{-12} \exp(365./T)$	
$\text{MBOO}_2 + \text{CH}_3\text{O}_2 \longrightarrow 0.9165 \cdot \text{CH}_2\text{O} + 0.625 \cdot \text{MACROH} + 0.25 \cdot \text{CH}_3\text{OH} + \text{HO}_2 + 0.3335 \cdot \text{CH}_3\text{COCH}_3 + 0.3335 \cdot \text{GLYALD} + 0.1665 \cdot \text{IBUTALOH}$	$3.750 \cdot 10^{-13} \exp(-40./T)$	
$\text{MBOO}_2 + \text{CH}_3\text{CO}_3 \longrightarrow 0.67 \cdot \text{CH}_3\text{COCH}_3 + 0.67 \cdot \text{GLYALD} + 0.33 \cdot \text{CH}_2\text{O} + 0.33 \cdot \text{IBUTALOH} + \text{HO}_2 + \text{CH}_3\text{O}_2 + \text{CO}_2$	$1.000 \cdot 10^{-11}$	
$\text{MBOO}_2 + \text{HO}_2 \longrightarrow \text{MBOOOH}$	$7.500 \cdot 10^{-13} \exp(700./T)$	
$\text{MBOOOH} + \text{OH} \longrightarrow 0.5 \cdot \text{MBOO}_2 + 0.625 \cdot \text{MACROH} + 0.5 \cdot \text{OH} + \text{H}_2\text{O}$	$3.800 \cdot 10^{-12} \exp(200./T)$	
$\text{IBUTALOH} + \text{OH} \longrightarrow \text{H}_2\text{O} + \text{IBUTALOHO}_2$	$1.400 \cdot 10^{-11}$	
$\text{IBUTALOHO}_2 + \text{NO} \longrightarrow \text{CO}_2 + \text{NO}_2 + \text{HO}_2 + \text{CH}_3\text{COCH}_3$	$2.600 \cdot 10^{-12} \exp(365./T)$	
$\text{IBUTALOHO}_2 + \text{HO}_2 \longrightarrow 0.6 \cdot \text{IBUTALOHOOH} + 0.4 \cdot \text{HO}_2 + 0.4 \cdot \text{OH} + 0.4 \cdot \text{CH}_3\text{COCH}_3 + 0.4 \cdot \text{CO}_2$	$4.300 \cdot 10^{-13} \exp(1040./T)$	
$\text{IBUTALOHO}_2 + \text{CH}_3\text{O}_2 \longrightarrow \text{CH}_3\text{COCH}_3 + 2 \cdot \text{HO}_2 + \text{CO}_2 + \text{CH}_2\text{O}$	$1.000 \cdot 10^{-12}$	products: Tyndall (p.c.)
$\text{IBUTALOHO}_2 + \text{CH}_3\text{CO}_3 \longrightarrow \text{CH}_3\text{COCH}_3 + \text{HO}_2 + \text{CH}_3\text{O}_2 + 2 \cdot \text{CO}_2$	$1.000 \cdot 10^{-11}$	
$\text{IBUTALOHOOH} + \text{OH} \longrightarrow \text{H}_2\text{O} + \text{IBUTALOHO}_2$	$1.000 \cdot 10^{-12}$	Tyndall (p.c.)

Table S10: C5 oxidation (... continued)

reaction	rate coefficient	reference
$\text{MBONO}_3\text{O}_2 + \text{HO}_2 \longrightarrow$	$4.300 \cdot 10^{-13} \exp(1040./T)$	
$\text{MBONO}_3\text{O}_2 + \text{NO} \longrightarrow 0.25 \cdot \text{CH}_2\text{O} + 0.25 \cdot \text{IBUTALOH} + 1.25 \cdot \text{NO}_2 + 0.500250 \cdot \text{NOA} + 0.75 \cdot \text{CH}_3\text{COCH}_3 + 0.75 \cdot \text{HO}_2$	$2.600 \cdot 10^{-12} \exp(365./T)$	
$\text{MBONO}_3\text{O}_2 + \text{NO}_3 \longrightarrow 0.25 \cdot \text{CH}_2\text{O} + 0.25 \cdot \text{IBUTALOH} + 1.25 \cdot \text{NO}_2 + 0.500250 \cdot \text{NOA} + 0.75 \cdot \text{CH}_3\text{COCH}_3 + 0.75 \cdot \text{HO}_2$	$2.400 \cdot 10^{-12}$	
$\text{MBONO}_3\text{O}_2 + \text{CH}_3\text{O}_2 \longrightarrow 0.875 \cdot \text{CH}_2\text{O} + 0.125 \cdot \text{IBUTALOH} + 0.125 \cdot \text{NO}_2 + 0.250125 \cdot \text{NOA} + 0.375 \cdot \text{CH}_3\text{COCH}_3 + 0.875 \cdot \text{HO}_2 + 0.25 \cdot \text{CH}_3\text{OH} + 0.625 \cdot \text{MACROH}$	$1.000 \cdot 10^{-12}$	Tyndall (p.c.)
$\text{MBONO}_3\text{O}_2 + \text{CH}_3\text{CO}_3 \longrightarrow 0.25 \cdot \text{CH}_2\text{O} + 0.25 \cdot \text{IBUTALOH} + 0.25 \cdot \text{NO}_2 + 0.500250 \cdot \text{NOA} + 0.75 \cdot \text{CH}_3\text{COCH}_3 + 0.75 \cdot \text{HO}_2 + \text{CH}_3\text{O}_2 + \text{CO}_2$	$1.000 \cdot 10^{-11}$	
$\text{DICARBO}_2 + \text{HO}_2 \longrightarrow 0.4 \cdot \text{OH} + 0.068 \cdot \text{CH}_3\text{COCHO} + 0.068 \cdot \text{HO}_2 + 0.068 \cdot \text{CO} + 0.4 \cdot \text{CO}_2 + 0.332 \cdot \text{CH}_3\text{CO}_3 + 0.332 \cdot \text{GLYOXAL}$	$4.300 \cdot 10^{-13} \exp(1040./T)$	
$\text{DICARBO}_2 + \text{NO} \longrightarrow 0.17 \cdot \text{CH}_3\text{COCHO} + 0.17 \cdot \text{HO}_2 + 0.17 \cdot \text{CO} + \text{CO}_2 + 0.83 \cdot \text{CH}_3\text{CO}_3 + 0.83 \cdot \text{GLYOXAL} + \text{NO}_2$	$7.500 \cdot 10^{-12} \exp(290./T)$	
$\text{DICARBO}_2 + \text{NO}_2 + \text{M} \longrightarrow \text{M}$	$ktroe(2.700 \cdot 10^{-28}, 7.1, 1.200 \cdot 10^{-11}, 0.9, 0.6)$	Orlando and Tyndall (2012), Same as $k(\text{CH}_3\text{CO}_3 + \text{NO}_2)$
$\text{DICARBO}_2 + \text{CH}_3\text{O}_2 \longrightarrow 0.17 \cdot \text{CH}_3\text{COCHO} + 1.17 \cdot \text{HO}_2 + 0.17 \cdot \text{CO} + \text{CO}_2 + 0.83 \cdot \text{CH}_3\text{CO}_3 + 0.83 \cdot \text{GLYOXAL} + \text{CH}_2\text{O}$	$1.000 \cdot 10^{-12}$	
$\text{DICARBO}_2 + \text{CH}_3\text{CO}_3 \longrightarrow 0.17 \cdot \text{CH}_3\text{COCHO} + 0.17 \cdot \text{HO}_2 + 0.17 \cdot \text{CO} + 2 \cdot \text{CO}_2 + 0.83 \cdot \text{CH}_3\text{CO}_3 + 0.83 \cdot \text{GLYOXAL} + \text{CH}_3\text{O}_2$	$1.000 \cdot 10^{-11}$	
$\text{HPALD} + \text{OH} \longrightarrow 0.641 \cdot \text{OH} + 0.385 \cdot \text{PACALD} + 0.256 \cdot \text{BIGALD}_3 + 0.359 \cdot \text{CH}_3\text{COCHO} + 0.359 \cdot \text{GLYOX} + 0.359 \cdot \text{HO}_2$	$5.200 \cdot 10^{-11}$	simplification of chemistry in MCMv3.3.1
$\text{PACALD} + \text{OH} \longrightarrow \text{CH}_3\text{COCHO} + \text{HCOCO}_3\text{H} + \text{HO}_2$	$4.720 \cdot 10^{-11}$	k and products for $\text{C}_5\text{PACALD}_2$ from MCMv3.3.1 assuming an implicit $\text{RO}_2 \longrightarrow \text{RO}$ conversion
$\text{LISOPOOHO}_2 + \text{HO}_2 \longrightarrow \text{LISOPOOHOOH}$	$2.050 \cdot 10^{-13} \exp(1300./T)$	
$\text{LISOPOOHO}_2 + \text{NO} \longrightarrow \text{CH}_2\text{O} + \text{HO}_2 + 0.5 \cdot \text{MACROOH} + 0.5 \cdot \text{LHMKABOOH} + \text{NO}_2$	$2.540 \cdot 10^{-12} \exp(360./T)$	$A = 2.54 \cdot 10^{-12} \cdot 0.12$ ; average 12% nitrate yield from Paulot et al. (2009); nitrate yield left for the moment equal to the one of the simple ISOPO2 (it should be higher)
$\text{LISOPOOHO}_2 + \text{NO} \longrightarrow \text{LISOPNO}_3\text{OOH}$	$3.050 \cdot 10^{-13} \exp(360./T)$	$A = 2.54 \cdot 10^{-12} \cdot 0.12$ ; average 12% nitrate yield from Paulot et al. (2009); nitrate yield left for the moment equal to the one of the simple ISOPO2 (it should be higher)

Table S10: C5 oxidation (... continued)

reaction	rate coefficient	reference
$\text{LISOPOOHO}_2 + \text{NO}_3 \longrightarrow \text{CH}_2\text{O} + \text{HO}_2 + 0.5 \cdot \text{MACROOH} + 0.5 \cdot \text{LHMKABOOH} + \text{NO}_2$	$2.500 \cdot 10^{-12}$	
$\text{LISOPOOHO}_2 + \text{CH}_3\text{O}_2 \longrightarrow 1.5 \cdot \text{CH}_2\text{O} + 0.75 \cdot \text{HO}_2 + 0.375 \cdot \text{MACROOH} + 0.375 \cdot \text{LHMKABOOH} + 0.25 \cdot \text{LISOPOOHOOH}$	$8.000 \cdot 10^{-13}$	
$\text{LISOPOOHO}_2 + \text{CH}_3\text{CO}_3 \longrightarrow \text{CH}_2\text{O} + \text{HO}_2 + 0.5 \cdot \text{MACROOH} + 0.5 \cdot \text{LHMKABOOH} + \text{CH}_3\text{O}_2 + \text{CO}_2$	$1.000 \cdot 10^{-11}$	
$\text{LISOPNO}_3\text{O}_2 + \text{HO}_2 \longrightarrow \text{LISOPNO}_3\text{OOH}$	$2.050 \cdot 10^{-13} \exp(1300./T)$	
$\text{LISOPNO}_3\text{O}_2 + \text{NO} \longrightarrow \text{CH}_2\text{O} + 0.5 \cdot \text{MACRN} + 0.5 \cdot \text{MVKN} + \text{HO}_2 + \text{NO}_2$	$2.235 \cdot 10^{-12} \exp(360./T)$	$A = 2.54 \cdot 10^{-12} \cdot 0.88$ ; average 12% nitrate yield from Paulot et al. (2009); nitrate yield left for the moment equal to the one of the simple ISOPO2 (it should be higher); previous $\text{RONO}_2$ split into 50% MACRN + 50% MVKN
$\text{LISOPNO}_3\text{O}_2 + \text{NO} \longrightarrow \text{LISOPNO}_3\text{NO}_3$	$3.050 \cdot 10^{-13} \exp(360./T)$	$A = 2.54 \cdot 10^{-12} \cdot 0.12$ ; average 12% nitrate yield from Paulot et al. (2009); nitrate yield left for the moment equal to the one of the simple ISOPO2 (it should be higher)
$\text{LISOPNO}_3\text{O}_2 + \text{NO}_3 \longrightarrow \text{CH}_2\text{O} + 0.5 \cdot \text{MACRN} + 0.5 \cdot \text{MVKN} + \text{HO}_2 + \text{NO}_2$	$2.500 \cdot 10^{-12}$	
$\text{LISOPNO}_3\text{O}_2 + \text{CH}_3\text{O}_2 \longrightarrow 0.3525 \cdot \text{MACRN} + 0.3525 \cdot \text{MVKN} + 1.75 \cdot \text{CH}_2\text{O} + 1.5 \cdot \text{HO}_2 + 0.25 \cdot \text{LISOPNO}_3\text{OOH}$	$8.000 \cdot 10^{-13}$	
$\text{LISOPNO}_3\text{O}_2 + \text{CH}_3\text{CO}_3 \longrightarrow \text{CH}_2\text{O} + 0.5 \cdot \text{MACRN} + 0.5 \cdot \text{MVKN} + \text{HO}_2 + \text{CH}_3\text{O}_2 + \text{CO}_2$	$1.000 \cdot 10^{-11}$	
$\text{LISOPOOHOOH} + \text{OH} \longrightarrow \text{H}_2\text{O} + \text{LISOPOOHO}_2$	$7.600 \cdot 10^{-12} \exp(200./T)$	twice the $k(\text{CH}_3\text{OOH} + \text{OH} \longrightarrow \text{CH}_3\text{O}_2)$
$\text{LISOPOOHOOH} + \text{OH} \longrightarrow \text{LC}_{578}\text{OOH} + \text{OH}$	$2.104 \cdot 10^{-11}$	$k$ for H-abstractions from SAR in MOM by a secondary carbon bearing a -OH group and a secondary and a tertiary carbon atoms bearing an -OOH group: $8.42 \cdot 10^{-13} \cdot 3.44 + (8.42 \cdot 10^{-13} + 1.75 \cdot 10^{-12}) \cdot 7$
$\text{LISOPNO}_3\text{OOH} + \text{OH} \longrightarrow \text{H}_2\text{O} + \text{LISOPNO}_3\text{O}_2$	$3.800 \cdot 10^{-12} \exp(200./T)$	
$\text{LISOPNO}_3\text{OOH} + \text{OH} \longrightarrow \text{C}_{59}\text{OOH} + \text{OH}$	$1.515 \cdot 10^{-11}$	$k$ for H-abstractions from SAR in MOM by a secondary carbon bearing a -OH group and a tertiary carbon atom bearing an -OOH group: $8.42 \cdot 10^{-13} \cdot 3.44 + 1.75 \cdot 10^{-12} \cdot 7$
$\text{LISOPNO}_3\text{NO}_3 + \text{OH} \longrightarrow 0.5 \cdot \text{MACRN} + 0.5 \cdot \text{MVKN} + \text{NO}_2 + \text{CH}_2\text{O} + \text{HO}_2$	$8.916 \cdot 10^{-12}$	$k$ for H-abstractions from SAR in MOM by a secondary and a tertiary carbon bearing a -OH group: $(8.42 \cdot 10^{-13} + 1.75 \cdot 10^{-12}) \cdot 3.44$ ; previous $\text{RONO}_2$ split into 50% MACRN + 50% MVKN



Table S11: C6 oxidation

reaction	rate coefficient	reference
$\text{BENZ} + \text{OH} \longrightarrow 0.53 \cdot \text{PHENOL} + 0.12 \cdot \text{BEPOMUC} + 0.65 \cdot \text{HO}_2 + 0.35 \cdot \text{BENZO}_2$	$2.300 \cdot 10^{-12} \exp(-193./T)$	
$\text{PHENOL} + \text{OH} \longrightarrow 0.14 \cdot \text{PHENO}_2 + 0.8 \cdot \text{HO}_2 + 0.8 \cdot \text{CATECHOL} + 0.06 \cdot \text{C}_6\text{H}_5\text{O}$	$4.700 \cdot 10^{-13} \exp(1220./T)$	
$\text{PHENOL} + \text{NO}_3 \longrightarrow 0.26 \cdot \text{PHENO}_2 + 0.74 \cdot \text{C}_6\text{H}_5\text{O} + 0.74 \cdot \text{HNO}_3$	$3.800 \cdot 10^{-12}$	NPHEO2 approximated with PHENO <sub>2</sub>
$\text{PHENO}_2 + \text{NO} \longrightarrow \text{HO}_2 + 0.7 \cdot \text{GLYOXAL} + \text{NO}_2$	$2.600 \cdot 10^{-12} \exp(365./T)$	
$\text{PHENO}_2 + \text{HO}_2 \longrightarrow \text{PHENOOH}$	$7.500 \cdot 10^{-13} \exp(700./T)$	
$\text{PHENO}_2 + \text{CH}_3\text{O}_2 \longrightarrow 2 \cdot \text{HO}_2 + 0.7 \cdot \text{GLYOXAL} + \text{CH}_2\text{O}$	$1.000 \cdot 10^{-12}$	
$\text{PHENO}_2 + \text{CH}_3\text{CO}_3 \longrightarrow \text{HO}_2 + 0.7 \cdot \text{GLYOXAL} + \text{CH}_3\text{O}_2 + \text{CO}_2$	$1.000 \cdot 10^{-11}$	
$\text{PHENOOH} + \text{OH} \longrightarrow \text{H}_2\text{O} + \text{PHENO}_2$	$3.800 \cdot 10^{-12} \exp(200./T)$	
$\text{C}_6\text{H}_5\text{O} + \text{NO}_2 \longrightarrow$	$2.100 \cdot 10^{-12}$	
$\text{C}_6\text{H}_5\text{O} + \text{O}_3 \longrightarrow \text{C}_6\text{H}_5\text{O}_2$	$2.800 \cdot 10^{-13}$	
$\text{C}_6\text{H}_5\text{O}_2 + \text{NO} \longrightarrow \text{C}_6\text{H}_5\text{O} + \text{NO}_2$	$2.600 \cdot 10^{-12} \exp(365./T)$	
$\text{C}_6\text{H}_5\text{O}_2 + \text{NO}_3 \longrightarrow \text{C}_6\text{H}_5\text{O} + \text{NO}_2$	$2.300 \cdot 10^{-12}$	MCM3.2
$\text{C}_6\text{H}_5\text{O}_2 + \text{HO}_2 \longrightarrow \text{C}_6\text{H}_5\text{OOH}$	$7.500 \cdot 10^{-13} \exp(700./T)$	
$\text{C}_6\text{H}_5\text{O}_2 + \text{CH}_3\text{O}_2 \longrightarrow \text{C}_6\text{H}_5\text{O} + \text{CH}_2\text{O} + \text{HO}_2$	$1.000 \cdot 10^{-12}$	
$\text{C}_6\text{H}_5\text{O}_2 + \text{CH}_3\text{CO}_3 \longrightarrow \text{C}_6\text{H}_5\text{O} + \text{CH}_3\text{O}_2 + \text{CO}_2$	$1.000 \cdot 10^{-11}$	
$\text{C}_6\text{H}_5\text{OOH} + \text{OH} \longrightarrow \text{C}_6\text{H}_5\text{O}_2$	$3.800 \cdot 10^{-12} \exp(200./T)$	
$\text{BENZO}_2 + \text{NO} \longrightarrow \text{GLYOXAL} + \text{NO}_2 + 0.5 \cdot \text{BIGALD}_1 + \text{HO}_2$	$2.600 \cdot 10^{-12} \exp(365./T)$	MCM3.2
$\text{BENZO}_2 + \text{HO}_2 \longrightarrow \text{BENZOOH}$	$7.500 \cdot 10^{-13} \exp(700./T)$	
$\text{BENZO}_2 + \text{CH}_3\text{O}_2 \longrightarrow \text{GLYOXAL} + 0.5 \cdot \text{BIGALD}_1 + 2 \cdot \text{HO}_2 + \text{CH}_2\text{O}$	$1.000 \cdot 10^{-12}$	
$\text{BENZO}_2 + \text{CH}_3\text{CO}_3 \longrightarrow \text{GLYOXAL} + 0.5 \cdot \text{BIGALD}_1 + \text{HO}_2 + \text{CH}_3\text{O}_2 + \text{CO}_2$	$1.000 \cdot 10^{-11}$	
$\text{BENZOOH} + \text{OH} \longrightarrow \text{BENZO}_2$	$3.800 \cdot 10^{-12} \exp(200./T)$	
$\text{CATECHOL} + \text{OH} \longrightarrow \text{CATEC}_1\text{O}$	$1.000 \cdot 10^{-10}$	
$\text{CATECHOL} + \text{NO}_3 \longrightarrow \text{CATEC}_1\text{O} + \text{HNO}_3$	$9.900 \cdot 10^{-11}$	
$\text{CATEC}_1\text{O} + \text{NO}_2 \longrightarrow$	$2.100 \cdot 10^{-12}$	
$\text{CATEC}_1\text{O} + \text{O}_3 \longrightarrow \text{CATEC}_1\text{O}_2$	$2.800 \cdot 10^{-13}$	
$\text{CATEC}_1\text{O}_2 + \text{HO}_2 \longrightarrow \text{CATEC}_1\text{OOH}$	$7.500 \cdot 10^{-13} \exp(700./T)$	
$\text{CATEC}_1\text{O}_2 + \text{NO} \longrightarrow \text{CATEC}_1\text{O} + \text{NO}_2$	$2.600 \cdot 10^{-12} \exp(365./T)$	
$\text{CATEC}_1\text{O}_2 + \text{NO}_3 \longrightarrow \text{CATEC}_1\text{O} + \text{NO}_2$	$2.300 \cdot 10^{-12}$	MCM3.2
$\text{CATEC}_1\text{O}_2 + \text{CH}_3\text{O}_2 \longrightarrow \text{CATEC}_1\text{O} + \text{CH}_2\text{O} + \text{HO}_2$	$1.000 \cdot 10^{-12}$	
$\text{CATEC}_1\text{O}_2 + \text{CH}_3\text{CO}_3 \longrightarrow \text{CATEC}_1\text{OCH}_3\text{O}_2 + \text{CO}_2$	$1.000 \cdot 10^{-11}$	
$\text{CATEC}_1\text{OOH} + \text{OH} \longrightarrow \text{CATEC}_1\text{O}_2$	$1.900 \cdot 10^{-12} \exp(190./T)$	

Table S12: C7 oxidation

reaction	rate coefficient	reference
$\text{TOL} + \text{OH} \longrightarrow 0.18 \cdot \text{CRESOL} + 0.1 \cdot \text{TEPOMUC} + 0.07 \cdot \text{BZOO} + 0.65 \cdot \text{TOLO}_2 + 0.28 \cdot \text{HO}_2$	$1.700 \cdot 10^{-12} \exp(352./T)$	
$\text{CRESOL} + \text{OH} \longrightarrow 0.2 \cdot \text{PHENO}_2 + 0.73 \cdot \text{HO}_2 + 0.73 \cdot \text{CATECHOL} + 0.07 \cdot \text{C}_6\text{H}_5\text{O}$	$4.700 \cdot 10^{-11}$	CATECHOL and PHENO <sub>2</sub> omits one CH <sub>3</sub> group of MCATECHOL and CRESO <sub>2</sub>
$\text{CRESOL} + \text{NO}_3 \longrightarrow 0.61 \cdot \text{PHENO}_2 + 0.39 \cdot \text{C}_6\text{H}_5\text{O} + 0.49 \cdot \text{HNO}_3$	$1.400 \cdot 10^{-11}$	CRESO <sub>2</sub> and NCRESO <sub>2</sub> approximated with PHENO <sub>2</sub> ; TOL <sub>1</sub> O with C <sub>6</sub> H <sub>5</sub> O (see MCM3.2 for details)
$\text{TOLO}_2 + \text{HO}_2 \longrightarrow \text{TOLOOH}$	$7.500 \cdot 10^{-13} \exp(700./T)$	
$\text{TOLO}_2 + \text{NO} \longrightarrow \text{NO}_2 + 0.6 \cdot \text{GLYOXAL} + 0.4 \cdot \text{CH}_3\text{COCHO} + \text{HO}_2 + 0.2 \cdot \text{BIGALD}_1 + 0.2 \cdot \text{BIGALD}_2 + 0.2 \cdot \text{BIGALD}_3$	$2.600 \cdot 10^{-12} \exp(365./T)$	
$\text{TOLO}_2 + \text{CH}_3\text{O}_2 \longrightarrow 0.6 \cdot \text{GLYOXAL} + 0.4 \cdot \text{CH}_3\text{COCHO} + 2 \cdot \text{HO}_2 + \text{CH}_2\text{O}$	$1.000 \cdot 10^{-12}$	
$\text{TOLO}_2 + \text{CH}_3\text{CO}_3 \longrightarrow 0.6 \cdot \text{GLYOXAL} + 0.4 \cdot \text{CH}_3\text{COCHO} + \text{HO}_2 + \text{CH}_3\text{O}_2 + \text{CO}_2 + 0.2 \cdot \text{BIGALD}_1 + 0.2 \cdot \text{BIGALD}_2 + 0.2 \cdot \text{BIGALD}_3$	$1.000 \cdot 10^{-11}$	
$\text{TOLOOH} + \text{OH} \longrightarrow \text{TOLO}_2$	$3.800 \cdot 10^{-12} \exp(200./T)$	
$\text{BZOO} + \text{HO}_2 \longrightarrow \text{BZOOH}$	$7.500 \cdot 10^{-13} \exp(700./T)$	
$\text{BZOO} + \text{NO} \longrightarrow \text{BZALD} + \text{NO}_2 + \text{HO}_2$	$2.600 \cdot 10^{-12} \exp(365./T)$	MCM3.2 forms 10% nitrate
$\text{BZOO} + \text{CH}_3\text{O}_2 \longrightarrow \text{BZALD} + 2 \cdot \text{HO}_2 + \text{CH}_2\text{O}$	$1.000 \cdot 10^{-12}$	
$\text{BZOO} + \text{CH}_3\text{CO}_3 \longrightarrow \text{BZALD} + \text{HO}_2 + \text{CH}_3\text{O}_2 + \text{CO}_2$	$1.000 \cdot 10^{-11}$	
$\text{BZOOH} + \text{OH} \longrightarrow \text{BZOO}$	$3.800 \cdot 10^{-12} \exp(200./T)$	
$\text{BZALD} + \text{OH} \longrightarrow \text{ACBZO}_2$	$5.900 \cdot 10^{-12} \exp(225./T)$	
$\text{ACBZO}_2 + \text{NO}_2 + \text{M} \longrightarrow \text{M} + \text{PBZNIT}$	$ktroe(2.700 \cdot 10^{-28}, 7.1, 1.200 \cdot 10^{-11}, 0.9, 0.6)$	Orlando and Tyndall (2012) - Same as k(CH <sub>3</sub> CO <sub>3</sub> + NO <sub>2</sub> )
$\text{PBZNIT} + \text{M} \longrightarrow \text{ACBZO}_2 + \text{NO}_2 + \text{M}$		
$\text{ACBZO}_2 + \text{NO} \longrightarrow \text{C}_6\text{H}_5\text{O}_2 + \text{NO}_2$	$7.500 \cdot 10^{-12} \exp(290./T)$	
$\text{ACBZO}_2 + \text{HO}_2 \longrightarrow 0.4 \cdot \text{C}_6\text{H}_5\text{O}_2 + 0.4 \cdot \text{OH}$	$4.300 \cdot 10^{-13} \exp(1040./T)$	
$\text{ACBZO}_2 + \text{CH}_3\text{O}_2 \longrightarrow \text{C}_6\text{H}_5\text{O}_2 + \text{CH}_2\text{O} + \text{HO}_2$	$1.000 \cdot 10^{-12}$	
$\text{ACBZO}_2 + \text{CH}_3\text{CO}_3 \longrightarrow \text{C}_6\text{H}_5\text{O}_2 + \text{CH}_3\text{O}_2 + \text{CO}_2$	$1.000 \cdot 10^{-11}$	

Table S13: C8 oxidation

reaction	rate coefficient	reference
$\text{XYL} + \text{OH} \longrightarrow 0.15 \cdot \text{XYLOL} + 0.23 \cdot \text{TEPOMUC} + 0.06 \cdot \text{BZOO} + 0.56 \cdot \text{XYLENO}_2 + 0.38 \cdot \text{HO}_2$	$1.700 \cdot 10^{-11}$	

Table S13: C8 oxidation (... continued)

reaction	rate coefficient	reference
$\text{XYLOL} + \text{OH} \longrightarrow 0.3 \cdot \text{XYLOLO}_2 + 0.63 \cdot \text{HO}_2 + 0.63 \cdot \text{CATECHOL} + 0.07 \cdot \text{C}_6\text{H}_5\text{O}$	$8.400 \cdot 10^{-11}$	CATECHOL omits two CH <sub>3</sub> groups of O-, M- and P-XYCATECH
$\text{XYLOL} + \text{NO}_3 \longrightarrow 0.61 \cdot \text{PHENO}_2 + 0.39 \cdot \text{C}_6\text{H}_5\text{O} + 0.49 \cdot \text{HNO}_3$	$3.200 \cdot 10^{-11}$	XYLOLO <sub>2</sub> and NXYLOLO <sub>2</sub> approximated with PHENO <sub>2</sub> ; XY1O with C <sub>6</sub> H <sub>5</sub> O (see MCM3.2 for details)
$\text{XYLOLO}_2 + \text{NO} \longrightarrow \text{HO}_2 + \text{NO}_2 + 0.17 \cdot \text{GLYOXAL} + 0.51 \cdot \text{CH}_3\text{COCHO}$	$2.600 \cdot 10^{-12} \exp(365./T)$	
$\text{XYLOLO}_2 + \text{HO}_2 \longrightarrow \text{XYLOLOOH}$	$7.500 \cdot 10^{-13} \exp(700./T)$	
$\text{XYLOLO}_2 + \text{CH}_3\text{O}_2 \longrightarrow 0.17 \cdot \text{GLYOXAL} + 0.51 \cdot \text{CH}_3\text{COCHO} + 2 \cdot \text{HO}_2 + \text{CH}_2\text{O}$	$1.000 \cdot 10^{-12}$	
$\text{XYLOLO}_2 + \text{CH}_3\text{CO}_3 \longrightarrow \text{HO}_2 + 0.17 \cdot \text{GLYOXAL} + 0.51 \cdot \text{CH}_3\text{COCHO} + \text{CH}_3\text{O}_2 + \text{CO}_2$	$1.000 \cdot 10^{-11}$	
$\text{XYLOLOOH} + \text{OH} \longrightarrow \text{XYLOLO}_2$	$3.800 \cdot 10^{-12} \exp(200./T)$	
$\text{XYLENO}_2 + \text{HO}_2 \longrightarrow \text{XYLENOOH}$	$7.500 \cdot 10^{-13} \exp(700./T)$	
$\text{XYLENO}_2 + \text{NO} \longrightarrow \text{HO}_2 + \text{NO}_2 + 0.34 \cdot \text{GLYOXAL} + 0.54 \cdot \text{CH}_3\text{COCHO} + 0.06 \cdot \text{BIGALD}_1 + 0.2 \cdot \text{BIGALD}_2 + 0.15 \cdot \text{BIGALD}_3 + 0.21 \cdot \text{BIGALD}_4$	$2.600 \cdot 10^{-12} \exp(365./T)$	
$\text{XYLENO}_2 + \text{CH}_3\text{O}_2 \longrightarrow 0.34 \cdot \text{GLYOXAL} + 0.54 \cdot \text{CH}_3\text{COCHO} + 2 \cdot \text{HO}_2 + \text{CH}_2\text{O} + 0.06 \cdot \text{BIGALD}_1 + 0.2 \cdot \text{BIGALD}_2 + 0.15 \cdot \text{BIGALD}_3 + 0.21 \cdot \text{BIGALD}_4$	$1.000 \cdot 10^{-12}$	
$\text{XYLENO}_2 + \text{CH}_3\text{CO}_3 \longrightarrow \text{HO}_2 + 0.34 \cdot \text{GLYOXAL} + 0.54 \cdot \text{CH}_3\text{COCHO} + 0.06 \cdot \text{BIGALD}_1 + 0.2 \cdot \text{BIGALD}_2 + 0.15 \cdot \text{BIGALD}_3 + 0.21 \cdot \text{BIGALD}_4 + \text{CH}_3\text{O}_2 + \text{CO}_2$	$1.000 \cdot 10^{-11}$	
$\text{XYLENOOH} + \text{OH} \longrightarrow \text{XYLENO}_2$	$3.800 \cdot 10^{-12} \exp(200./T)$	

Table S14: C10/C15 oxidation (terpenes, sesqui-terpenes)

reaction	rate coefficient	reference
$\text{APIN} + \text{OH} \longrightarrow \text{TERPO}_2$	$1.200 \cdot 10^{-11} \exp(440./T)$	
$\text{BPIN} + \text{OH} \longrightarrow \text{TERPO}_2$	$1.600 \cdot 10^{-11} \exp(470./T)$	
$\text{LIMON} + \text{OH} \longrightarrow \text{TERPO}_2$	$4.200 \cdot 10^{-11} \exp(400./T)$	
$\text{MYRC} + \text{OH} \longrightarrow \text{TERPO}_2$	$2.100 \cdot 10^{-10}$	
$\text{BCARY} + \text{OH} \longrightarrow \text{TERPO}_2$	$2.000 \cdot 10^{-10}$	
$\text{APIN} + \text{O}_3 \longrightarrow 0.07 \cdot \text{ELVOC} + 0.39 \cdot \text{TERPROD}_1 + 0.27 \cdot \text{TERPROD}_2 + 0.63 \cdot \text{OH} + 0.57 \cdot \text{HO}_2 + 0.23 \cdot \text{CO} + 0.27 \cdot \text{CO}_2 + 0.52 \cdot \text{CH}_3\text{COCH}_3 + 0.34 \cdot \text{CH}_2\text{O} + 0.05 \cdot \text{HCOOH} + 0.05 \cdot \text{BIGALKANE} + 0.06 \cdot \text{CH}_3\text{CO}_3 + 0.06 \cdot \text{CH}_3\text{COCH}_2\text{O}_2$	$6.300 \cdot 10^{-16} \exp(-580./T)$	7% ELVOC-yield according to Ehn et al. (2014) for endocyclic alkenes
$\text{BPIN} + \text{O}_3 \longrightarrow 0.43 \cdot \text{TERPROD}_1 + 0.3 \cdot \text{TERPROD}_2 + 0.63 \cdot \text{OH} + 0.57 \cdot \text{HO}_2 + 0.23 \cdot \text{CO} + 0.27 \cdot \text{CO}_2 + 0.52 \cdot \text{CH}_3\text{COCH}_3 + 0.34 \cdot \text{CH}_2\text{O} + 0.05 \cdot \text{HCOOH} + 0.05 \cdot \text{BIGALKANE} + 0.06 \cdot \text{CH}_3\text{CO}_3 + 0.06 \cdot \text{CH}_3\text{COCH}_2\text{O}_2$	$1.700$ $10^{-15} \exp(-1300./T)$	.

Table S14: C10/C15 oxidation (terpenes, sesqui-terpenes; ... continued)

reaction	rate coefficient	reference
$\text{LIMON} + \text{O}_3 \longrightarrow 0.07 \cdot \text{ELVOC} + 0.39 \cdot \text{TERPROD}_1 + 0.27 \cdot \text{TERPROD}_2 + 0.63 \cdot \text{OH} + 0.57 \cdot \text{HO}_2 + 0.23 \cdot \text{CO} + 0.27 \cdot \text{CO}_2 + 0.52 \cdot \text{CH}_3\text{COCH}_3 + 0.34 \cdot \text{CH}_2\text{O} + 0.05 \cdot \text{HCOOH} + 0.05 \cdot \text{BIGALKANE} + 0.06 \cdot \text{CH}_3\text{CO}_3 + 0.06 \cdot \text{CH}_3\text{COCH}_2\text{O}_2$	$3.000 \cdot 10^{-15} \exp(-780./T)$	7% ELVOC-yield according to Ehn et al. (2014) for endocyclic alkenes
$\text{MYRC} + \text{O}_3 \longrightarrow 0.43 \cdot \text{TERPROD}_1 + 0.3 \cdot \text{TERPROD}_2 + 0.63 \cdot \text{OH} + 0.57 \cdot \text{HO}_2 + 0.23 \cdot \text{CO} + 0.27 \cdot \text{CO}_2 + 0.52 \cdot \text{CH}_3\text{COCH}_3 + 0.34 \cdot \text{CH}_2\text{O} + 0.05 \cdot \text{HCOOH} + 0.05 \cdot \text{BIGALKANE} + 0.06 \cdot \text{CH}_3\text{CO}_3 + 0.06 \cdot \text{CH}_3\text{COCH}_2\text{O}_2$	$4.700 \cdot 10^{-16}$	
$\text{BCARY} + \text{O}_3 \longrightarrow 0.645 \cdot \text{TERPROD}_1 + 0.45 \cdot \text{TERPROD}_2 + 0.63 \cdot \text{OH} + 0.57 \cdot \text{HO}_2 + 0.23 \cdot \text{CO} + 0.27 \cdot \text{CO}_2 + 0.52 \cdot \text{CH}_3\text{COCH}_3 + 0.34 \cdot \text{CH}_2\text{O} + 0.05 \cdot \text{HCOOH} + 0.05 \cdot \text{BIGALKANE} + 0.06 \cdot \text{CH}_3\text{CO}_3 + 0.06 \cdot \text{CH}_3\text{COCH}_2\text{O}_2$	$1.200 \cdot 10^{-14}$	
$\text{APIN} + \text{NO}_3 \longrightarrow \text{NTERPO}_2$	$1.200 \cdot 10^{-12} \exp(490./T)$	
$\text{BPIN} + \text{NO}_3 \longrightarrow \text{NTERPO}_2$	$2.500 \cdot 10^{-12}$	
$\text{LIMON} + \text{NO}_3 \longrightarrow \text{NTERPO}_2$	$1.100 \cdot 10^{-11}$	
$\text{MYRC} + \text{NO}_3 \longrightarrow \text{NTERPO}_2$	$1.200 \cdot 10^{-11}$	
$\text{BCARY} + \text{NO}_3 \longrightarrow \text{NTERPO}_2 + 0.5 \cdot \text{TERPROD}_1$	$1.900 \cdot 10^{-11}$	
$\text{TERPO}_2 + \text{NO} \longrightarrow 0.26 \cdot \text{TERPNO}_3 + 0.74 \cdot \text{NO}_2 + 0.36 \cdot \text{CH}_2\text{O} + 0.045 \cdot \text{CH}_3\text{COCH}_3 + 0.695 \cdot \text{TERPROD}_1 + 0.74 \cdot \text{HO}_2$	$4.200 \cdot 10^{-12} \exp(180./T)$	alkyl nitrate yield according to Rindelaub et al. (2015) for alpha-pinene
$\text{TERPO}_2 + \text{HO}_2 \longrightarrow \text{TERPOOH}$	$7.500 \cdot 10^{-13} \exp(700./T)$	
$\text{TERPO}_2 + \text{CH}_3\text{O}_2 \longrightarrow 1.15 \cdot \text{CH}_2\text{O} + 0.05 \cdot \text{CH}_3\text{COCH}_3 + 0.945 \cdot \text{TERPROD}_1 + \text{HO}_2 + 0.25 \cdot \text{CH}_3\text{OH}$	$2.000 \cdot 10^{-12} \exp(500./T)$	
$\text{TERPO}_2 + \text{CH}_3\text{CO}_3 \longrightarrow 0.4 \cdot \text{CH}_2\text{O} + 0.05 \cdot \text{CH}_3\text{COCH}_3 + 0.945 \cdot \text{TERPROD}_1 + \text{HO}_2 + \text{CH}_3\text{O}_2 + \text{CO}_2$	$1.000 \cdot 10^{-11}$	
$\text{TERPOOH} + \text{OH} \longrightarrow \text{H}_2\text{O} + \text{TERPO}_2$	$3.300 \cdot 10^{-11}$	
$\text{TERPROD}_1 + \text{OH} \longrightarrow \text{TERP}_2\text{O}_2$	$5.700 \cdot 10^{-11}$	
$\text{TERPROD}_1 + \text{NO}_3 \longrightarrow 0.5 \cdot \text{TERP}_2\text{O}_2 + 0.5 \cdot \text{NTERPO}_2 + 0.5 \cdot \text{NO}_2$	$1.000 \cdot 10^{-12}$	
$\text{TERPNO}_3 + \text{OH} \longrightarrow \text{NO}_2 + \text{TERPROD}_1 + \text{H}_2\text{O}$	$3.500 \cdot 10^{-12}$	
$\text{TERP}_2\text{O}_2 + \text{NO} \longrightarrow 0.1 \cdot \text{TERPNO}_3 + 0.9 \cdot \text{NO}_2 + 0.34 \cdot \text{CH}_2\text{O} + 0.27 \cdot \text{CH}_3\text{COCH}_3 + 0.225 \cdot \text{CO} + 0.9 \cdot \text{CO}_2 + 0.9 \cdot \text{TERPROD}_2 + 0.9 \cdot \text{HO}_2 + 0.225 \cdot \text{GLYALD}$	$4.200 \cdot 10^{-12} \exp(180./T)$	
$\text{TERP}_2\text{O}_2 + \text{HO}_2 \longrightarrow \text{TERP}_2\text{OOH}$	$7.500 \cdot 10^{-13} \exp(700./T)$	
$\text{TERP}_2\text{O}_2 + \text{CH}_3\text{O}_2 \longrightarrow \text{TERPROD}_2 + 0.93 \cdot \text{CH}_2\text{O} + 0.25 \cdot \text{CH}_3\text{OH} + \text{HO}_2 + 0.5 \cdot \text{CO}_2 + 0.125 \cdot \text{CO} + 0.125 \cdot \text{GLYALD} + 0.15 \cdot \text{CH}_3\text{COCH}_3$	$2.000 \cdot 10^{-12} \exp(500./T)$	
$\text{TERP}_2\text{O}_2 + \text{CH}_3\text{CO}_3 \longrightarrow 0.34 \cdot \text{CH}_2\text{O} + 0.27 \cdot \text{CH}_3\text{COCH}_3 + 0.225 \cdot \text{CO} + 2 \cdot \text{CO}_2 + \text{TERPROD}_2 + \text{HO}_2 + 0.225 \cdot \text{GLYALD} + \text{CH}_3\text{O}_2$	$1.000 \cdot 10^{-11}$	
$\text{TERP}_2\text{OOH} + \text{OH} \longrightarrow \text{H}_2\text{O} + \text{TERP}_2\text{O}_2$	$2.300 \cdot 10^{-11}$	

Table S14: C10/C15 oxidation (terpenes, sesqui-terpenes; ... continued)

reaction	rate coefficient	reference
TERPROD <sub>2</sub> + OH $\longrightarrow$ 0.15 · CH <sub>3</sub> COCH <sub>2</sub> O <sub>2</sub> + 0.68 · CH <sub>2</sub> O + 1.8 · CO <sub>2</sub> + 0.5 · CH <sub>3</sub> COCH <sub>3</sub> + 0.65 · CH <sub>3</sub> CO <sub>3</sub> + 0.2 · HO <sub>2</sub> + 0.7 · CO	$3.400 \cdot 10^{-11}$	
NTERPO <sub>2</sub> + NO $\longrightarrow$ 0.26 · NTERPNO <sub>3</sub> + 1.48 · NO <sub>2</sub> + 0.74 · TERPROD <sub>1</sub>	$4.200 \cdot 10^{-12} \exp(180./T)$	alkyl nitrate yield according to Rindelaub et al. (2015) for alpha-pinene
NTERPO <sub>2</sub> + HO <sub>2</sub> $\longrightarrow$ NTERPNO <sub>3</sub>	$7.500 \cdot 10^{-13} \exp(700./T)$	
NTERPO <sub>2</sub> + CH <sub>3</sub> O <sub>2</sub> $\longrightarrow$ 0.5 · NTERPNO <sub>3</sub> + 0.75 · CH <sub>2</sub> O + 0.25 · CH <sub>3</sub> OH + 0.5 · HO <sub>2</sub> + 0.5 · TERPROD <sub>1</sub> + 0.5 · NO <sub>2</sub>	$2.000 \cdot 10^{-12} \exp(500./T)$	
NTERPO <sub>2</sub> + CH <sub>3</sub> CO <sub>3</sub> $\longrightarrow$ CH <sub>3</sub> O <sub>2</sub> + TERPROD <sub>1</sub> + CO <sub>2</sub> + NO <sub>2</sub>	$1.000 \cdot 10^{-11}$	
NTERPO <sub>2</sub> + NO <sub>3</sub> $\longrightarrow$ 2 · NO <sub>2</sub> + TERPROD <sub>1</sub>	$2.400 \cdot 10^{-12}$	
NTERPNO <sub>3</sub> + OH $\longrightarrow$ NO <sub>2</sub> + TERPROD <sub>1</sub> + H <sub>2</sub> O	$3.500 \cdot 10^{-12}$	
ELVOC + OH $\longrightarrow$ HO <sub>2</sub> + TERPROD <sub>1</sub>	$1.000 \cdot 10^{-11}$	a general rate coefficient for oxygenated VOC

Table S15: Tropospheric halogen + organics reactions

reaction	rate coefficient	reference
CL + CH <sub>2</sub> O $\longrightarrow$ HCL + HO <sub>2</sub> + CO	$8.100 \cdot 10^{-11} \exp(-30./T)$	
CL + CH <sub>4</sub> $\longrightarrow$ CH <sub>3</sub> O <sub>2</sub> + HCL	$7.300 \cdot 10^{-12} \exp(-1280./T)$	
CL + CH <sub>3</sub> CN $\longrightarrow$ CH <sub>2</sub> O + HCL + CO + NO	$1.600 \cdot 10^{-11} \exp(-2140./T)$	JPL (2011), products: Tyndall
CL + C <sub>2</sub> H <sub>2</sub> + M $\longrightarrow$ 0.1 · CL + 0.1 · GLYOXAL + 0.9 · HCL + 0.9 · HO <sub>2</sub> + 1.8 · CO + M	$ktroe(5.200 \cdot 10^{-30}, 2.4, 2.200 \cdot 10^{-10}, 0.7, 0.6)$	
CL + C <sub>2</sub> H <sub>4</sub> + M $\longrightarrow$ HO <sub>2</sub> + 2 · CO + HCL + M	$ktroe(1.600e-29, 3.3, 3.100 \cdot 10^{-10}, 1., 0.6)$	
CL + C <sub>2</sub> H <sub>6</sub> $\longrightarrow$ C <sub>2</sub> H <sub>5</sub> O <sub>2</sub> + HCL	$7.200 \cdot 10^{-11} \exp(-70./T)$	
CL + CH <sub>3</sub> O <sub>2</sub> $\longrightarrow$ 0.5 · CLO + 0.5 · CH <sub>2</sub> O + 0.5 · HCL + 0.5 · OH + 0.5 · CO + HO <sub>2</sub>	$1.600 \cdot 10^{-10}$	50% CH <sub>2</sub> OO is produced and assumed to be produced mostly in dry environments and thus decomposing
CL + CH <sub>3</sub> OH $\longrightarrow$ CH <sub>2</sub> O + HO <sub>2</sub> + HCL	$7.100 \cdot 10^{-11} \exp(-75./T)$	IUPAC (2008)
CL + CH <sub>3</sub> OOH $\longrightarrow$ CH <sub>2</sub> O + OH + HCL	$5.900 \cdot 10^{-11}$	IUPAC (2008)
CL + CH <sub>3</sub> CHO $\longrightarrow$ CH <sub>3</sub> CO <sub>3</sub> + HCL	$8.000 \cdot 10^{-11}$	IUPAC (2008)
CL + GLYALD $\longrightarrow$ 0.35 · GLYOXAL + 0.35 · HO <sub>2</sub> + 0.65 · HOCH <sub>2</sub> CO <sub>3</sub> + HCL	$7.600 \cdot 10^{-11}$	k by Bacher et al. (2001); products by Niki et al. (1987)
CL + GLYOXAL $\longrightarrow$ HCL + HCOCO <sub>3</sub>	$3.441 \cdot 10^{-11}$	k(298K) from Niki et al. (1985); k(CH <sub>2</sub> O + Cl) has been adjusted for the k(CH <sub>2</sub> O + Cl) by JPL (2011) used here
CL + C <sub>3</sub> H <sub>8</sub> $\longrightarrow$ C <sub>3</sub> H <sub>7</sub> O <sub>2</sub> + HCL	$1.400 \cdot 10^{-10}$	IUPAC (2008)

Table S15: Tropospheric halogen + organics reactions (... continued)

reaction	rate coefficient	reference
$\text{CL} + \text{CH}_3\text{COCH}_3 \longrightarrow \text{CH}_3\text{COCH}_2\text{O}_2 + \text{HCL}$	$1.500 \cdot 10^{-11} \exp(-590./T)$	IUPAC (2008)
$\text{CL} + \text{HYAC} \longrightarrow \text{CH}_3\text{COCHO} + \text{HO}_2 + \text{HCL}$	$5.400 \cdot 10^{-11}$	Calvert et al. (2008a)
$\text{CL} + \text{BIGALKANE} \longrightarrow \text{ALKO}_2 + \text{HCL}$	$1.935e - 10$	from the reaction with OH, BIGALKANE seems to be methyl-butane whose k is an average of k given by Qian et al. (2002) and Anderson et al. (2007)
$\text{CL} + \text{MEK} \longrightarrow \text{HCL} + \text{MEKO}_2$	$3.800 \cdot 10^{-11} \exp(16./T)$	Calvert et al. (2008b)
$\text{CLO} + \text{CH}_3\text{O}_2 \longrightarrow \text{CL} + \text{HO}_2 + \text{CH}_2\text{O}$	$3.300 \cdot 10^{-12} \exp(-115./T)$	
$\text{BR} + \text{CH}_2\text{O} \longrightarrow \text{HBR} + \text{HO}_2 + \text{CO}$	$1.700 \cdot 10^{-11} \exp(-800./T)$	
$\text{BR} + \text{CH}_3\text{CHO} \longrightarrow \text{CH}_3\text{CO}_3 + \text{HBR}$	$1.800 \cdot 10^{-11} \exp(-460./T)$	IUPAC (2008)
$\text{BRO} + \text{CH}_3\text{O}_2 \longrightarrow \text{HOBR} + \text{OH} + \text{HO}_2 + \text{CO}$	$2.420 \cdot 10^{-14} \exp(1617./T)$	Shallcross et al. (2015); $\text{CH}_2\text{OO}$ assumed to be produced mostly in dry environments and thus decomposing

Table S16: Sulfur reactions

reaction	rate coefficient	reference
$\text{SO}_2 + \text{OH} + \text{M} \longrightarrow \text{H}_2\text{SO}_4 + \text{HO}_2$	$ktroe(3.300 \cdot 10^{-31}, 4.3, 1.600 \cdot 10^{-12}, 0., 0.6)$	JPL (2011)
$\text{DMS} + \text{OH} \longrightarrow \text{CH}_3\text{SO}_2 + \text{HCHO}$	$1.130 \cdot 10^{-11} \exp(-253./T)$	
$\text{DMS} + \text{OH} \longrightarrow \text{DMSO} + \text{HO}_2$	$1. \cdot 10^{-9} \exp(5820./T) \cdot [\text{O}_2] / (1. \cdot 10^{30} + 5. \cdot \exp(6280./T) \cdot [\text{O}_2])$	
$\text{DMS} + \text{NO}_3 \longrightarrow \text{CH}_3\text{SO}_2 + \text{HNO}_3 + \text{HCHO}$	$1.900 \cdot 10^{-13} \exp(520./T)$	
$\text{DMS} + \text{CL} \longrightarrow \text{CH}_3\text{SO}_2 + \text{HCL} + \text{HCHO}$	$3.300 \cdot 10^{-10}$	
$\text{DMS} + \text{BR} \longrightarrow \text{CH}_3\text{SO}_2 + \text{HBR} + \text{HCHO}$	$9.000 \cdot 10^{-11} \exp(-2386./T)$	
$\text{DMS} + \text{BRO} \longrightarrow \text{BR} + \text{DMSO}$	$4.400 \cdot 10^{-13}$	
$\text{DMSO} + \text{OH} \longrightarrow 0.6 \cdot \text{SO}_2 + \text{HCHO} + 0.6 \cdot \text{CH}_3\text{O}_2 + 0.4 \cdot \text{HO}_2 + 0.4 \cdot \text{CH}_3\text{SO}_3\text{H}$	$1.000 \cdot 10^{-10}$	
$\text{CH}_3\text{SO}_2 \longrightarrow \text{CH}_3\text{O}_2 + \text{SO}_2$	$1.800 \cdot 10^{13} \exp(-8661./T)$	
$\text{CH}_3\text{SO}_2 + \text{O}_3 \longrightarrow \text{CH}_3\text{SO}_3$	$3.000 \cdot 10^{-13}$	
$\text{CH}_3\text{SO}_3 + \text{HO}_2 \longrightarrow \text{CH}_3\text{SO}_3\text{H}$	$5.000 \cdot 10^{-11}$	

Table S17: Stratospheric O(1D) reactions (... continued)

reaction	rate coefficient	reference
Table S17: Stratospheric O(1D) reactions		
reaction	rate coefficient	reference
$\text{O}_1\text{D} + \text{N}_2\text{O} \longrightarrow 2 \cdot \text{NO}$	$7.250 \cdot 10^{-11} \exp(20./T)$	
$\text{O}_1\text{D} + \text{N}_2\text{O} \longrightarrow \text{N}_2 + \text{O}_2$	$4.630 \cdot 10^{-11} \exp(20./T)$	
$\text{O}_1\text{D} + \text{O}_3 \longrightarrow 2 \cdot \text{O}_2$	$1.200 \cdot 10^{-10}$	
$\text{O}_1\text{D} + \text{CFC}_{11} \longrightarrow 2 \cdot \text{CL} + \text{COFCL}$	$2.020 \cdot 10^{-10}$	
$\text{O}_1\text{D} + \text{CFC}_{12} \longrightarrow 2 \cdot \text{CL} + \text{COF}_2$	$1.204 \cdot 10^{-10}$	
$\text{O}_1\text{D} + \text{CFC}_{113} \longrightarrow 2 \cdot \text{CL} + \text{COFCL} + \text{COF}_2$	$1.500 \cdot 10^{-10}$	
$\text{O}_1\text{D} + \text{CFC}_{114} \longrightarrow 2 \cdot \text{CL} + 2 \cdot \text{COF}_2$	$9.750 \cdot 10^{-11}$	
$\text{O}_1\text{D} + \text{CFC}_{115} \longrightarrow \text{CL} + \text{F} + 2 \cdot \text{COF}_2$	$1.500 \cdot 10^{-11}$	
$\text{O}_1\text{D} + \text{HCFC}_{22} \longrightarrow \text{CL} + \text{COF}_2$	$7.200 \cdot 10^{-11}$	
$\text{O}_1\text{D} + \text{HCFC}_{141}\text{B} \longrightarrow \text{CL} + \text{COFCL}$	$1.794 \cdot 10^{-10}$	
$\text{O}_1\text{D} + \text{HCFC}_{142}\text{B} \longrightarrow \text{CL} + \text{COF}_2$	$1.628 \cdot 10^{-10}$	
$\text{O}_1\text{D} + \text{CCL}_4 \longrightarrow 4 \cdot \text{CL}$	$2.840 \cdot 10^{-10}$	
$\text{O}_1\text{D} + \text{CH}_3\text{BR} \longrightarrow \text{BR}$	$1.674 \cdot 10^{-10}$	
$\text{O}_1\text{D} + \text{CF}_2\text{CLBR} \longrightarrow \text{BR} + \text{CL} + \text{COF}_2$	$9.600 \cdot 10^{-11}$	
$\text{O}_1\text{D} + \text{CF}_3\text{BR} \longrightarrow \text{BR} + \text{F} + \text{COF}_2$	$4.100 \cdot 10^{-11}$	
$\text{O}_1\text{D} + \text{H}_{1202} \longrightarrow 2 \cdot \text{BR} + \text{COF}_2$	$1.012 \cdot 10^{-10}$	
$\text{O}_1\text{D} + \text{H}_{2402} \longrightarrow 2 \cdot \text{BR} + 2 \cdot \text{COF}_2$	$1.200 \cdot 10^{-10}$	
$\text{O}_1\text{D} + \text{CHBR}_3 \longrightarrow 3 \cdot \text{BR}$	$4.490 \cdot 10^{-10}$	
$\text{O}_1\text{D} + \text{CH}_2\text{BR}_2 \longrightarrow 2 \cdot \text{BR}$	$2.570 \cdot 10^{-10}$	
$\text{O}_1\text{D} + \text{COF}_2 \longrightarrow 2 \cdot \text{F}$	$2.140 \cdot 10^{-11}$	
$\text{O}_1\text{D} + \text{COFCL} \longrightarrow \text{CL} + \text{F}$	$1.900 \cdot 10^{-10}$	
$\text{O}_1\text{D} + \text{CH}_4 \longrightarrow \text{CH}_3\text{O}_2 + \text{OH}$	$1.310 \cdot 10^{-10}$	
$\text{O}_1\text{D} + \text{CH}_4 \longrightarrow \text{CH}_2\text{O} + \text{H} + \text{HO}_2$	$3.500 \cdot 10^{-11}$	
$\text{O}_1\text{D} + \text{CH}_4 \longrightarrow \text{CH}_2\text{O} + \text{H}_2$	$9.000 \cdot 10^{-12}$	
$\text{O}_1\text{D} + \text{H}_2 \longrightarrow \text{H} + \text{OH}$	$1.200 \cdot 10^{-10}$	
$\text{O}_1\text{D} + \text{HCL} \longrightarrow \text{CL} + \text{OH}$	$1.500 \cdot 10^{-10}$	
$\text{O}_1\text{D} + \text{HBR} \longrightarrow \text{BR} + \text{OH}$	$1.200 \cdot 10^{-10}$	
$\text{O}_1\text{D} + \text{HCN} \longrightarrow \text{CO} + \text{OH} + \text{NO}$	$7.700 \cdot 10^{-11} \exp(100./T)$	Strekowski et al. (2001), products: Tyndall

Table S18: Stratospheric inorganic halogen reactions

reaction	rate coefficient	reference
$\text{CL} + \text{O}_3 \longrightarrow \text{CLO} + \text{O}_2$	$2.300 \cdot 10^{-11} \exp(-200./T)$	
$\text{CL} + \text{H}_2 \longrightarrow \text{H} + \text{HCL}$	$3.050 \cdot 10^{-11} \exp(-2270./T)$	
$\text{CL} + \text{H}_2\text{O}_2 \longrightarrow \text{HCL} + \text{HO}_2$	$1.100 \cdot 10^{-11} \exp(-980./T)$	
$\text{CL} + \text{HO}_2 \longrightarrow \text{HCL} + \text{O}_2$	$1.400 \cdot 10^{-11} \exp(270./T)$	
$\text{CL} + \text{HO}_2 \longrightarrow \text{CLO} + \text{OH}$	$3.600 \cdot 10^{-11} \exp(-375./T)$	
$\text{CL}_2\text{O}_2 + \text{M} \longrightarrow 2 \cdot \text{CLO} + \text{M}$	$k_{\text{CL2O2}}$	
$\text{CLO} + \text{O} \longrightarrow \text{CL} + \text{O}_2$	$2.800 \cdot 10^{-11} \exp(85./T)$	
$\text{CLO} + \text{OH} \longrightarrow \text{CL} + \text{HO}_2$	$7.400 \cdot 10^{-12} \exp(270./T)$	
$\text{CLO} + \text{OH} \longrightarrow \text{HCL} + \text{O}_2$	$6.000 \cdot 10^{-13} \exp(230./T)$	
$\text{CLO} + \text{HO}_2 \longrightarrow \text{HOCL} + \text{O}_2$	$2.600 \cdot 10^{-12} \exp(290./T)$	
$\text{CLO} + \text{NO} \longrightarrow \text{CL} + \text{NO}_2$	$6.400 \cdot 10^{-12} \exp(290./T)$	
$\text{CLO} + \text{NO}_2 + \text{M} \longrightarrow \text{CLONO}_2 + \text{M}$	$ktroe(1.800 \cdot 10^{-31}, 3.4, 1.500 \cdot 10^{-11}, 1.9, 0.6)$	
$\text{CLO} + \text{CLO} \longrightarrow 2 \cdot \text{CL} + \text{O}_2$	$3.000 \cdot 10^{-11} \exp(-2450./T)$	
$\text{CLO} + \text{CLO} \longrightarrow \text{CL}_2 + \text{O}_2$	$1.000 \cdot 10^{-12} \exp(-1590./T)$	
$\text{CLO} + \text{CLO} \longrightarrow \text{CL} + \text{OCLO}$	$3.500 \cdot 10^{-13} \exp(-1370./T)$	
$\text{CLO} + \text{CLO} + \text{M} \longrightarrow \text{CL}_2\text{O}_2 + \text{M}$	$ktroe(1.600 \cdot 10^{-32}, 4.5, 3.000 \cdot 10^{-12}, 2., 0.6)$	
$\text{HCL} + \text{OH} \longrightarrow \text{CL} + \text{H}_2\text{O}$	$1.800 \cdot 10^{-12} \exp(-250./T)$	
$\text{HCL} + \text{O} \longrightarrow \text{CL} + \text{OH}$	$1.000 \cdot 10^{-11} \exp(-3300./T)$	
$\text{HOCL} + \text{O} \longrightarrow \text{CLO} + \text{OH}$	$1.700 \cdot 10^{-13}$	
$\text{HOCL} + \text{CL} \longrightarrow \text{CLO} + \text{HCL}$	$3.400 \cdot 10^{-12} \exp(-130./T)$	
$\text{HOCL} + \text{OH} \longrightarrow \text{CLO} + \text{H}_2\text{O}$	$3.000 \cdot 10^{-12} \exp(-500./T)$	
$\text{CLONO}_2 + \text{O} \longrightarrow \text{CLO} + \text{NO}_3$	$3.600 \cdot 10^{-12} \exp(-840./T)$	
$\text{CLONO}_2 + \text{OH} \longrightarrow \text{HOCL} + \text{NO}_3$	$1.200 \cdot 10^{-12} \exp(-330./T)$	
$\text{CLONO}_2 + \text{CL} \longrightarrow \text{CL}_2 + \text{NO}_3$	$6.500 \cdot 10^{-12} \exp(135./T)$	
$\text{BR} + \text{O}_3 \longrightarrow \text{BRO} + \text{O}_2$	$1.600 \cdot 10^{-11} \exp(-780./T)$	
$\text{BR} + \text{HO}_2 \longrightarrow \text{HBR} + \text{O}_2$	$4.800 \cdot 10^{-12} \exp(-310./T)$	
$\text{BR} + \text{NO}_2 + \text{M} \longrightarrow 0.85 \cdot \text{BRONO} + 0.15 \cdot \text{BRNO}_2 + \text{M}$	$ktroe(4.200 \cdot 10^{-31}, 2.4, 2.700 \cdot 10^{-11}, 0., 0.6)$	
$\text{BRONO} + \text{M} \longrightarrow \text{BR} + \text{NO}_2 + \text{M}$	$1.648 \cdot 10^{11} \exp(-7399./T)$	fit to upper limit by Wine et al. (1993) and scaled 1.5 factor to match data by Orlando and Burkholder (2000)
$\text{BRO} + \text{O} \longrightarrow \text{BR} + \text{O}_2$	$1.900 \cdot 10^{-11} \exp(230./T)$	
$\text{BRO} + \text{OH} \longrightarrow \text{BR} + \text{HO}_2$	$1.700 \cdot 10^{-11} \exp(250./T)$	
$\text{BRO} + \text{HO}_2 \longrightarrow \text{HOBR} + \text{O}_2$	$4.500 \cdot 10^{-12} \exp(460./T)$	



Table S18: Stratospheric inorganic halogen reactions (... continued)

reaction	rate coefficient	reference
$\text{BRO} + \text{NO} \longrightarrow \text{BR} + \text{NO}_2$	$8.800 \cdot 10^{-12} \exp(260./T)$	
$\text{BRO} + \text{NO}_2 + \text{M} \longrightarrow \text{BRONO}_2 + \text{M}$	<i>ktroe</i> ( $5.200 \cdot 10^{-31}$ , 3.2, $6.900 \cdot 10^{-12}$ , 2.9, 0.6)	
$\text{BRO} + \text{CLO} \longrightarrow \text{BR} + \text{OCLO}$	$9.500 \cdot 10^{-13} \exp(550./T)$	
$\text{BRO} + \text{CLO} \longrightarrow \text{BR} + \text{CL} + \text{O}_2$	$2.300 \cdot 10^{-12} \exp(260./T)$	
$\text{BRO} + \text{CLO} \longrightarrow \text{BRCL} + \text{O}_2$	$4.100 \cdot 10^{-13} \exp(290./T)$	
$\text{BRO} + \text{BRO} \longrightarrow 2 \cdot \text{BR} + \text{O}_2$	$2.400 \cdot 10^{-12} \exp(40./T)$	
$\text{BRO} + \text{BRO} \longrightarrow \text{BR}_2 + \text{O}_2$	$2.800 \cdot 10^{-14} \exp(860./T)$	
$\text{HBR} + \text{OH} \longrightarrow \text{BR} + \text{H}_2\text{O}$	$5.500 \cdot 10^{-12} \exp(200./T)$	
$\text{HBR} + \text{O} \longrightarrow \text{BR} + \text{OH}$	$5.800 \cdot 10^{-12} \exp(-1500./T)$	
$\text{HOBR} + \text{O} \longrightarrow \text{BRO} + \text{OH}$	$1.200 \cdot 10^{-10} \exp(-430./T)$	
$\text{BRONO}_2 + \text{O} \longrightarrow \text{BRO} + \text{NO}_3$	$1.900 \cdot 10^{-11} \exp(215./T)$	
$\text{BRONO}_2 + \text{BR} \longrightarrow \text{BR}_2 + \text{NO}_3$	$5.805 \cdot 10^{-11}$	Average of k at 298K by Orlando and Tyndall (1996) and Harwood et al. (1998)
$\text{BR}_2 + \text{OH} \longrightarrow \text{BR} + \text{HOBR}$	$2.100 \cdot 10^{-11} \exp(240./T)$	
$\text{F} + \text{H}_2\text{O} \longrightarrow \text{HF} + \text{OH}$	$1.400 \cdot 10^{-11}$	
$\text{F} + \text{H}_2 \longrightarrow \text{H} + \text{HF}$	$1.400 \cdot 10^{-10} \exp(-500./T)$	
$\text{F} + \text{CH}_4 \longrightarrow \text{CH}_3\text{O}_2 + \text{HF}$	$1.600 \cdot 10^{-10} \exp(-260./T)$	
$\text{F} + \text{HNO}_3 \longrightarrow \text{HF} + \text{NO}_3$	$6.000 \cdot 10^{-12} \exp(400./T)$	

Table S19: Stratospheric organic halogen reactions

reaction	rate coefficient	reference
$\text{CH}_3\text{BR} + \text{OH} \longrightarrow \text{BR} + \text{H}_2\text{O} + \text{HO}_2$	$2.350 \cdot 10^{-12} \exp(-1300./T)$	
$\text{CH}_3\text{BR} + \text{CL} \longrightarrow \text{HCL} + \text{HO}_2 + \text{BR}$	$1.400 \cdot 10^{-11} \exp(-1030./T)$	
$\text{CH}_2\text{BR}_2 + \text{OH} \longrightarrow 2 \cdot \text{BR} + \text{H}_2\text{O}$	$2.000 \cdot 10^{-12} \exp(-840./T)$	
$\text{CHBR}_3 + \text{OH} \longrightarrow 3 \cdot \text{BR}$	$1.350 \cdot 10^{-12} \exp(-600./T)$	
$\text{CH}_2\text{BR}_2 + \text{CL} \longrightarrow 2 \cdot \text{BR} + \text{HCL}$	$6.300 \cdot 10^{-12} \exp(-800./T)$	
$\text{CHBR}_3 + \text{CL} \longrightarrow 3 \cdot \text{BR} + \text{HCL}$	$4.850 \cdot 10^{-12} \exp(-850./T)$	
$\text{CH}_3\text{CL} + \text{CL} \longrightarrow \text{CO} + \text{HO}_2 + 2 \cdot \text{HCL}$	$2.170 \cdot 10^{-11} \exp(-1130./T)$	
$\text{CH}_3\text{CL} + \text{OH} \longrightarrow \text{CO} + \text{HO}_2 + \text{HCL} + \text{H}_2\text{O}$	$2.400 \cdot 10^{-12} \exp(-1250./T)$	products: Tyndall(p.c.), implicitly includes $\text{NO} \longrightarrow \text{NO}_2$ conversion with $\text{CH}_2\text{ClO}_2$
$\text{CH}_3\text{CCL}_3 + \text{OH} \longrightarrow \text{H}_2\text{O} + 3 \cdot \text{CL}$	$1.640 \cdot 10^{-12} \exp(-1520./T)$	
$\text{HCFC}_{22} + \text{OH} \longrightarrow \text{CL} + \text{H}_2\text{O} + \text{COF}_2$	$1.050 \cdot 10^{-12} \exp(-1600./T)$	
$\text{HCFC}_{141}\text{B} + \text{OH} \longrightarrow \text{CL} + \text{COFCL}$	$1.250 \cdot 10^{-12} \exp(-1600./T)$	

Table S19: Stratospheric organic halogen reactions (... continued)

reaction	rate coefficient	reference
$\text{HCFC}_{142}\text{B} + \text{OH} \longrightarrow \text{CL} + \text{COF}_2$	$1.300 \cdot 10^{-12} \exp(-1770./T)$	

Table S20: (Tropospheric) heterogeneous reactions

reaction	reaction probability	reference
$\text{O}_3 \longrightarrow \text{HO}_2$	$\gamma = 10^{-6}$	Stadtler et al. (2017)
$\text{HO}_2 \longrightarrow 0.5 \cdot \text{H}_2\text{O}_2$	$\gamma = 0.2$	Stadtler et al. (2017)
$\text{NO}_3 \longrightarrow$	$\gamma = 0.001$	Stadtler et al. (2017)
$\text{NO}_2 \longrightarrow 0.5 \cdot \text{HONO}$	$\gamma = 10^{-4}$	Stadtler et al. (2017)
$\text{HNO}_3 \longrightarrow$	$\gamma_{SS} = 0.01, \gamma_{DU} = 0.1$	Stadtler et al. (2017)
$\text{N}_2\text{O}_5 \longrightarrow$	$\gamma = f(T, RH)$	Stadtler et al. (2017)

Table S21: Stratospheric Sulfate aerosol reactions

reaction	reaction probability	reference
$\text{N}_2\text{O}_5 \longrightarrow 2 \cdot \text{HNO}_3$	$\gamma = 0.04$	Lamarque et al. (2012)
$\text{CLONO}_2 \longrightarrow \text{HOCL} + \text{HNO}_3$	$f(\text{sulfuricacidwt}\%)$	Lamarque et al. (2012)
$\text{BRONO}_2 \longrightarrow \text{HOBR} + \text{HNO}_3$	$f(T, P, HCl, \text{H}_2\text{O}, r)$	Lamarque et al. (2012)
$\text{CLONO}_2 + \text{HCL} \longrightarrow \text{CL}_2 + \text{HNO}_3$	$f(T, P, \text{H}_2\text{O}, r)$	Lamarque et al. (2012)
$\text{HOCL} + \text{HCL} \longrightarrow \text{CL}_2 + \text{H}_2\text{O}$	$f(T, P, HCl, \text{H}_2\text{O}, r)$	Lamarque et al. (2012)
$\text{HOBR} + \text{HCL} \longrightarrow \text{BRCL} + \text{H}_2\text{O}$	$f(T, P, HCl, \text{HOBr}, \text{H}_2\text{O}, r)$	Lamarque et al. (2012)

Table S22: Stratospheric Nitric acid dihydrate reactions

reaction	reaction probability	reference
$\text{N}_2\text{O}_5 \longrightarrow 2 \cdot \text{HNO}_3$	$\gamma = 0.0004$	Lamarque et al. (2012)
$\text{CLONO}_2 \longrightarrow \text{HOCL} + \text{HNO}_3$	$\gamma = 0.004$	Lamarque et al. (2012)
$\text{CLONO}_2 + \text{HCL} \longrightarrow \text{CL}_2 + \text{HNO}_3$	$\gamma = 0.2$	Lamarque et al. (2012)
$\text{HOCL} + \text{HCL} \longrightarrow \text{CL}_2 + \text{H}_2\text{O}$	$\gamma = 0.1$	Lamarque et al. (2012)
$\text{BRONO}_2 \longrightarrow \text{HOBR} + \text{HNO}_3$	$\gamma = 0.3$	Lamarque et al. (2012)

Table S23: Stratospheric Ice aerosol reactions

reaction	reaction probability	reference
$\text{N}_2\text{O}_5 \longrightarrow 2 \cdot \text{HNO}_3$	$\gamma = 0.02$	Lamarque et al. (2012)
$\text{CLONO}_2 \longrightarrow \text{HOCL} + \text{HNO}_3$	$\gamma = 0.3$	Lamarque et al. (2012)
$\text{BRONO}_2 \longrightarrow \text{HOBR} + \text{HNO}_3$	$\gamma = 0.3$	Lamarque et al. (2012)
$\text{CLONO}_2 + \text{HCL} \longrightarrow \text{CL}_2 + \text{HNO}_3$	$\gamma = 0.3$	Lamarque et al. (2012)
$\text{HOCL} + \text{HCL} \longrightarrow \text{CL}_2 + \text{H}_2\text{O}$	$\gamma = 0.2$	Lamarque et al. (2012)
$\text{HOBR} + \text{HCL} \longrightarrow \text{BRCL} + \text{H}_2\text{O}$	$\gamma = 0.3$	Lamarque et al. (2012)

Table S24: Henry coefficients ( $H_0$  and temperature factor) and dry deposition reactivities for gas-phase species in ECHAM-HAMMOZ.

Species	H	Reactivity coefficient	Source
CO	$9.81 \cdot 10^{-4}$ , 1720.	0.	JPL (2011)
H <sub>2</sub>	$7.8 \cdot 10^{-4}$ , 500.	0.	Sander (1999)
H <sub>2</sub> O <sub>2</sub>	$8.44 \cdot 10^4$ , 7600.	1	JPL (2011)
HCN	12., 5000.	1	Sander (1999)
HNO <sub>3</sub>	$3.2 \cdot 10^{11}$ , 8700.	1	$H = 2.1 \cdot 10^5$ M/atm and $Ka = 15.4$ M (Schwartz and White, 1981). At an average cloud droplet pH=5 $\Rightarrow H^* = H \cdot (1 + Ka/[H+]) = 2.1 \cdot 10^5 \cdot (1 + 15.4/1 \cdot 10^{-5}) = 3.2 \cdot 10^{11}$
HO <sub>2</sub>	690., 0.	1	JPL (2006)
HO <sub>2</sub> NO <sub>2</sub>	$1.2 \cdot 10^4$ , 6900.	1	Sander (1999)
HONO	$5.05 \cdot 10^3$ , 4800.	0.1	Sander (1999): $H=49$ and $Ka = 5.1 \cdot 10^{-4} \Rightarrow H^* = 49(1 + 5.1 \cdot 10^{-4} * 1 \cdot 10^7) = 5.05 \cdot 10^3$
N <sub>2</sub> O <sub>5</sub>	2.1, 3400.	1	Sander (1999): 2 ref. give it as infinite and one as 2.1 M/atm
N <sub>2</sub> O <sub>5</sub>	$3.2 \cdot 10^{11}$ , 8700.	1	HNO <sub>3</sub> as proxy as done in GEOS-Chem
NH <sub>3</sub>	$1.02 \cdot 10^4$ , 4200.	0.	Sander (2015): $H(298) = 58$ M/atm and $H^*(298)$ calculated at pH=7 with $Kb = 1.75 \cdot 10^5$ M
NO	$1.92 \cdot 10^{-3}$ , 1790.	1	JPL (2006), comment
NO <sub>2</sub>	$12. \cdot 10^{-2}$ , 2360.	1	JPL (2011)
NO <sub>3</sub>	$3.8 \cdot 10^{-2}$ , 0.	1	JPL (2006)
O <sub>3</sub>	$1.03 \cdot 10^{-2}$ , 2830.	1	JPL (2006)
SO <sub>2</sub>	$2.45 \cdot 10^5$ , 3100.	0.	Sander (2015): $H(298) = 1.2$ M/atm and $H^*(298)$ calculated at pH=7 with $K1a = 1.23 \cdot 10^{-2}$ M and $K2a = 6.61 \cdot 10^{-8}$ M
H <sub>2</sub> SO <sub>4</sub>	$1.3 \cdot 10^{15}$ , 20000.	0.	Sander (2015)
CH <sub>4</sub>	$1.41 \cdot 10^{-3}$ , 2040.	0.	JPL (2011)
CH <sub>2</sub> O	$3.23 \cdot 10^3$ , 7100.	0.1	JPL (2011)
CH <sub>3</sub> OH	203., 5640.	0.	JPL (2011)
CH <sub>3</sub> OOH	300., 5280.	0.1	JPL (2011)
HCOOH	$8.9 \cdot 10^3$ , 6100.	0.	Sander (1999): measured value by Johnson et al. (1996)??, surface reactivity as in Nguyen et al. (2015)
CH <sub>3</sub> O <sub>2</sub> NO <sub>2</sub>	2.0, 4700.	0.1	Sander (1999): methyl nitrate as proxy species
C <sub>2</sub> H <sub>5</sub> OH	190., 6660.	0.	JPL (2011)
C <sub>2</sub> H <sub>5</sub> OOH	336., 5995.	1	JPL (2011)
CH <sub>3</sub> CHO	12.9, 5890.	0.1	JPL (2011)
CH <sub>3</sub> CN	52.8, 3970.	0.	JPL (2011)
PAN	2.8, 5730.	0.1	JPL (2011)
CH <sub>3</sub> COOH	$4.1 \cdot 10^3$ , 6200.	0.	JPL (2011)

Table S24: Henry coefficients (... continued)

Species	H	Reactivity coefficient	Source
CH <sub>3</sub> COOOH	837., 5310.	0.1	JPL (2011)
GLYALD	$4.1 \cdot 10^4$ , 4600.	0.1	Sander (1999)
GLYOXAL	$4.19 \cdot 10^5$ , 7480.	0.1	JPL (2011)
HCOCO <sub>2</sub> H	$1.1 \cdot 10^4$ , 4800.	0.1	Sander (2015)
HCOCO <sub>3</sub> H	$1.1 \cdot 10^4$ , 4800.	0.1	Sander (2015)
HOCH <sub>2</sub> CO <sub>2</sub> H	$2.8 \cdot 10^4$ , 4000.	0.1	Sander (2015)
HOCH <sub>2</sub> CO <sub>3</sub> H	$2.8 \cdot 10^4$ , 4000.	0.1	Sander (2015)
C <sub>3</sub> H <sub>7</sub> OOH	336., 5995.	1	
CH <sub>3</sub> COCH <sub>3</sub>	27.8, 5530.	0.	JPL (2011)
CH <sub>3</sub> COCHO	$3.7 \cdot 10^3$ , 7500.	0.1	Sander (1999)
HYAC	$7.7 \cdot 10^3$ , 0.	0.1	Sander (2015)
NOA	$1. \cdot 10^3$ , 0.	0.1	Sander (2015)
MACR	6.5, 5300.	0.1	Sander (1999)
MACROOH	$2.1 \cdot 10^5$ , 9900.	1	Sander (2015)
MACROH	$2.1 \cdot 10^5$ , 9900.	0.	Sander (2015)
MEK	37., 8200.	0.1	Sander (2015)
MEKOOH	$1.1 \cdot 10^3$ , 7300.	1	Sander (2015)
MPAN	1.7, 0.	0.1	Sander (1999)
MVK	$4.1 \cdot 10^1$ , 7800.	0.1	Sander (1999)
MACO <sub>2</sub> H	$1.5 \cdot 10^3$ , 6800.	0.1	Sander (2015)
MACO <sub>3</sub> H	$1.5 \cdot 10^3$ , 6800.	1	Sander (2015)
BIGALD <sub>1</sub>	$2.5 \cdot 10^5$ , 0.	0.1	Sander (2015)
CO <sub>2</sub> H <sub>3</sub> CHO	$4.1 \cdot 10^4$ , 4600.	0.1	Sander (1999)
CO <sub>2</sub> H <sub>3</sub> CO <sub>3</sub> H	$4.1 \cdot 10^4$ , 4600.	1	Sander (1999)
BIACETOH	$4.1 \cdot 10^4$ , 4600.	0.1	Sander (1999)
IBUTALOH	$4.1 \cdot 10^4$ , 4600.	0.1	Sander (1999)
IBUTALOHOOH	$4.1 \cdot 10^4$ , 4600.	1	Sander (1999)
LHMKABOOH	$2.1 \cdot 10^5$ , 9900.	1	Sander (2015)
MVKN	$5. \cdot 10^3$ , 0.	0.1	like for ISOPNO3 from Nguyen et al. (2015)
MACRN	$5. \cdot 10^3$ , 0.	0.1	like for ISOPNO3 from Nguyen et al. (2015)
MBO	64., 0.	0.1	Sander (2015)
HPALD	$4.1 \cdot 10^4$ , 4600.	1	Sander (1999)
PACALD	$4.1 \cdot 10^4$ , 4600.	1	Sander (1999)
ISOPBNO <sub>3</sub>	$5. \cdot 10^3$ , 0.	0.1	Nguyen et al. (2015)

Table S24: Henry coefficients (... continued)

Species	H	Reactivity coefficient	Source
ISOPDNO <sub>3</sub>	$5. \cdot 10^3$ , 0.	0.1	Nguyen et al. (2015)
LISOPACNO <sub>3</sub>	$5. \cdot 10^3$ , 0.	0.1	Nguyen et al. (2015)
LC <sub>5</sub> PAN <sub>1719</sub>	$5. \cdot 10^3$ , 0.	0.1	Nguyen et al. (2015)
LHC <sub>4</sub> ACCO <sub>2</sub> H	$2.1 \cdot 10^5$ , 9900.	0.1	Sander (2015)
LHC <sub>4</sub> ACCO <sub>3</sub> H	$2.1 \cdot 10^5$ , 9900.	1	Sander (2015)
LIECHO	$4.1 \cdot 10^4$ , 4600.	0.1	Sander (1999)
LIECO <sub>3</sub> H	$2.1 \cdot 10^5$ , 9900.	1	Sander (2015)
LIEPOX	$2.1 \cdot 10^5$ , 9900.	1	Sander (2015)
LNISOOH	$2.1 \cdot 10^5$ , 9900.	1	Sander (2015)
MBOOOH	$3. \cdot 10^{11}$ , 0.	1	Sander (2015)
NC <sub>4</sub> CHO	2.3, 0.	0.1	Sander (2015)
NISOPOOH	$3.6 \cdot 10^4$ , 0.	1	Sander (2015)
LHC <sub>4</sub> ACCHO	$4.1 \cdot 10^4$ , 4600.	0.1	Sander (1999)
HCOC <sub>5</sub>	$4.1 \cdot 10^4$ , 4600.	1	Sander (1999)
ISOPBOOH	$2.1 \cdot 10^5$ , 9900.	1	Sander (2015)
ISOPDOOH	$2.1 \cdot 10^5$ , 9900.	1	Sander (2015)
LISOPACOOH	$2.1 \cdot 10^5$ , 9900.	1	Sander (2015)
ISOPBOH	$2.1 \cdot 10^5$ , 9900.	1	Sander (2015)
ISOPDOH	$2.1 \cdot 10^5$ , 9900.	1	Sander (2015)
ISOPAOH	$2.1 \cdot 10^5$ , 9900.	1	Sander (2015)
LISOPOOHOOH	$2. \cdot 10^{16}$ , 0.	1	Sander (2015)
LISOPNO <sub>3</sub> OOH	$3. \cdot 10^{11}$ , 0.	1	Sander (2015)
LISOPNO <sub>3</sub> NO <sub>3</sub>	$2.1 \cdot 10^5$ , 9900.	1	Sander (2015)
LC <sub>578</sub> OOH	$3. \cdot 10^{11}$ , 0.	1	Sander (2015)
C <sub>59</sub> OOH	$3. \cdot 10^{11}$ , 0.	1	Sander (2015)
BIGALD <sub>2</sub>	$2.5 \cdot 10^5$ , 0.	0.1	Sander (2015)
BIGALD <sub>3</sub>	$2.5 \cdot 10^5$ , 0.	0.1	Sander (2015)
BZALD	38., 5500.	0.1	Sander (2015)
BZOOH	$2.9 \cdot 10^3$ , 0.	1	Sander (2015)
PHENOOH	$3. \cdot 10^{11}$ , 0.	1	Sander (2015)
C <sub>6</sub> H <sub>5</sub> OOH	$3. \cdot 10^3$ , 5900.	1	Sander (2015)
PHENOL	$3. \cdot 10^3$ , 5900.	0.1	Sander (2015)
CATECHOL	$4.6 \cdot 10^3$ , 0.	0.1	Sander (2015)
CATEC <sub>1</sub> OOH	$4.6 \cdot 10^3$ , 0.	1	Sander (2015)

Table S24: Henry coefficients (... continued)

Species	H	Reactivity coefficient	Source
BEPOMUC	$2.5 \cdot 10^5$ , 0.	0.1	Sander (2015)
BENZOOH	$2.1 \cdot 10^5$ , 9900.	1	Sander (2015)
PBZNIT	2.8, 5730.	0.1	PAN as proxy MCM3.2 name PBZN
BIGALD <sub>4</sub>	$2.5 \cdot 10^5$ , 0.	0.1	Sander (2015)
TOLOOH	$2.1 \cdot 10^5$ , 9900.	1	Sander (2015)
CRESOL	$1.1 \cdot 10^3$ , 6700.	0.1	Sander (2015)
TEPOMUC	$2.5 \cdot 10^5$ , 0.	0.1	Sander (2015)
TERPROD <sub>2</sub>	$2.5 \cdot 10^5$ , 0.	0.1	Sander (2015)
XYLENOOH	$2.1 \cdot 10^5$ , 9900.	1	Sander (2015)
XYLOL	$1.1 \cdot 10^3$ , 6700.	0.1	cresol as proxy MCM3.2 name OXYLOL
XYLOLOOH	$2.1 \cdot 10^5$ , 9900.	1	Sander (2015)
TERPOOH	$2.1 \cdot 10^5$ , 9900.	1	Sander (2015)
TERP <sub>2</sub> OOH	$2.1 \cdot 10^5$ , 9900.	1	Sander (2015)
TERPNO <sub>3</sub>	$5. \cdot 10^3$ , 0.	0.1	like an isoprene nitrate MCM3.2 name APINANO <sub>3</sub>
TERPROD <sub>1</sub>	$2.5 \cdot 10^5$ , 0.	1	Sander (2015)
NTERPNO <sub>3</sub>	$5. \cdot 10^3$ , 0.	0.1	like an isoprene nitrate MCM3.2 name NAPINAOOH
MTHOM	$2. \cdot 10^9$ , 0.	1	Sander (2015)
ALCOH	336., 5995.	0.1	
ALKOOH	336., 5995.	1	
POOH	300., 5280.	0.1	
ROOH	300., 5280.	1	
EOOH	300., 5280.	0.1	
ALKNO <sub>3</sub>	$5. \cdot 10^3$ , 0.	0.1	like for ISOPNO <sub>3</sub> from Nguyen et al. (2015) MCM3.2
HF	$1.3 \cdot 10^4$ , 0.	0.	Sander (2015)
HCL	$2.0 \cdot 10^{11}$ , 600.	0.	Fernandez et al. (2014)
HOCL	$6.6 \cdot 10^2$ , 5900.	0.	Sander (1999)
CL <sub>2</sub>	$9.2 \cdot 10^{-2}$ , 2000.	0.	Sander (1999)
CLONO <sub>2</sub>	$2. \cdot 10^{13}$ , 0.	1	Sander (1999) : listed as "infinite"
BR <sub>2</sub>	$7.2 \cdot 10^{-1}$ , 0.	0.	Sander (2015)
HBR	$7.2 \cdot 10^{13}$ , 6100.	0.	Fernandez et al. (2014)
HOBR	$6.1 \cdot 10^3$ , 0.	0.	Sander (1999)
BRONO <sub>2</sub>	$2. \cdot 10^{13}$ , 0.	1	Sander (1999) : listed as "infinite"
BRONO	$3. \cdot 10^{-1}$ , 0.	0.	Sander (2015)
BRNO <sub>2</sub>	$3. \cdot 10^{-1}$ , 0.	0.	Sander (2015)

Table S24: Henry coefficients (... continued)

Species	H	Reactivity coefficient	Source
BRCL	$9.7 \cdot 10^{-1}$ , 5600.	0.	Sander (2015)
DMS	0.54, 3460.	0.	JPL (2011)
DMSO	$9.8 \cdot 10^4$ , 0.	0.1	JPL (2011)
CH <sub>3</sub> SO <sub>3</sub> H	$1. \cdot 10^{30}$ , 0.	1	Jöckel et al. (2006)



## References

- Anderson, J. L., Dixon, J. K., and Brennecke, J. F.: Solubility of CO<sub>2</sub>, CH<sub>4</sub>, C<sub>2</sub>H<sub>6</sub>, C<sub>2</sub>H<sub>4</sub>, O<sub>2</sub>, and N<sub>2</sub> in 1-Hexyl-3-methylpyridinium Bis(trifluoromethylsulfonyl)imide: Comparison to Other Ionic Liquids, *Accounts of Chemical Research*, 40, 1208–1216, doi:10.1021/ar7001649, PMID: 17970599, 2007.
- Bacher, C., Tyndall, G. S., and Orlando, J. J.: The Atmospheric Chemistry of Glycolaldehyde, *Journal of Atmospheric Chemistry*, 39, 171–189, doi:10.1023/A:1010689706869, 2001.
- Burkholder, J. and Orlando, J.: UV absorption cross-sections of cis-BrONO, *CHEMICAL PHYSICS LETTERS*, 317, 603–608, doi:10.1016/S0009-2614(99)01412-8, 2000.
- Calvert, J. G., Derwent, R. G., Orlando, J. J., Tyndall, G. S., and Wallington, T. J.: *Mechanisms of Atmospheric Oxidation of the Alkanes*, Oxford University Press, Oxford, UK, 2008a.
- Calvert, J. G., Mellouki, A., Orlando, J. J., Pilling, M., and Wallington, T. J.: *The Mechanisms of Atmospheric Oxidation of the Oxygenates*, Oxford University Press, Oxford, UK, 2008b.
- Crounse, J. D., Paulot, F., Kjaergaard, H. G., and Wennberg, P. O.: Peroxy radical isomerization in the oxidation of isoprene, *PHYSICAL CHEMISTRY CHEMICAL PHYSICS*, 13, 13 607–13 613, doi:10.1039/c1cp21330j, 2011.
- Crounse, J. D., Knap, H. C., Ornsø, K. B., Jorgensen, S., Paulot, F., Kjaergaard, H. G., and Wennberg, P. O.: Atmospheric Fate of Methacrolein. 1. Peroxy Radical Isomerization Following Addition of OH and O<sub>2</sub>, *JOURNAL OF PHYSICAL CHEMISTRY A*, 116, 5756–5762, doi:10.1021/jp211560u, 2012.
- Ehn, M., Thornton, J. A., Kleist, E., Sipila, M., Junninen, H., Pullinen, I., Springer, M., Rubach, F., Tillmann, R., Lee, B., Lopez-Hilfiker, F., Andres, S., Acir, I.-H., Rissanen, M., Jokinen, T., Schobesberger, S., Kangasluoma, J., Kontkanen, J., Nieminen, T., Kurten, T., Nielsen, L. B., Jorgensen, S., Kjaergaard, H. G., Canagaratna, M., Dal Maso, M., Berndt, T., Petaja, T., Wahner, A., Kerminen, V.-M., Kulmala, M., Worsnop, D. R., Wildt, J., and Mentel, T. F.: A large source of low-volatility secondary organic aerosol, *NATURE*, 506, 476+, doi:10.1038/nature13032, 2014.
- Fernandez, R. P., Salawitch, R. J., Kinnison, D. E., Lamarque, J.-F., and Saiz-Lopez, A.: Bromine partitioning in the tropical tropopause layer: implications for stratospheric injection, *Atmospheric Chemistry and Physics*, 14, 13 391–13 410, doi:10.5194/acp-14-13391-2014, 2014.
- Harwood, M., Burkholder, J., and Ravishankara, A.: Photodissociation of BrONO<sub>2</sub> and N<sub>2</sub>O<sub>5</sub>: Quantum yields for NO<sub>3</sub> production at 248, 308, and 352.5 nm, *JOURNAL OF PHYSICAL CHEMISTRY A*, 102, 1309–1317, doi:10.1021/jp9729829, 1998.
- IUPAC, Atkinson, R. and Baulch, D. L. and Cox, R. A. and Crowley, J. N. and Hampson, R. F. and Hynes, R. G. and Jenkin, M. E. and Rossi, M. J. and Troe, J.: Evaluated kinetic and photochemical data for atmospheric chemistry: Volume I - gas phase reactions of O<sub>x</sub>, HO<sub>x</sub>, NO<sub>x</sub> and SO<sub>x</sub> species, *Atmospheric Chemistry and Physics*, 4, 1461–1738, doi:10.5194/acp-4-1461-2004, 2004.
- IUPAC, Atkinson, R. and Baulch, D. L. and Cox, R. A. and Crowley, J. N. and Hampson, R. F. and Hynes, R. G. and Jenkin, M. E. and Rossi, M. J. and Troe, J. and IUPAC Subcommittee.: Evaluated kinetic and photochemical data for atmospheric chemistry: Volume II - gas phase reactions of organic species, *Atmospheric Chemistry and Physics*, 6, 3625–4055, doi:10.5194/acp-6-3625-2006, 2006.

- IUPAC, Atkinson, R. and Baulch, D. L. and Cox, R. A. and Crowley, J. N. and Hampson, R. F. and Hynes, R. G. and Jenkin, M. E. and Rossi, M. J. and Troe, J. and Wallington, T. J.: Evaluated kinetic and photochemical data for atmospheric chemistry: Volume IV gas phase reactions of organic halogen species, *Atmospheric Chemistry and Physics*, 8, 4141–4496, doi:10.5194/acp-8-4141-2008, 2008.
- Jöckel, P., Tost, H., Pozzer, A., Brühl, C., Buchholz, J., Ganzeveld, L., Hoor, P., Kerkweg, A., Lawrence, M., Sander, R., et al.: The atmospheric chemistry general circulation model ECHAM5/MESSy1: consistent simulation of ozone from the surface to the mesosphere, *Atmospheric Chemistry and Physics Discussions*, 6, 6957–7050, doi:10.5194/acp-6-5067-2006, 2006.
- Johnson, M., Kuwata, K., Wong, C., and Okumura, M.: Vibrational spectrum of I-(H<sub>2</sub>O), *CHEMICAL PHYSICS LETTERS*, 260, 551–557, doi:10.1016/0009-2614(96)00911-6, 1996.
- JPL, S. P. Sander and J. Abbatt and J. R. Barker and J. B. Burkholder and R. R. Friedl and D. M. Golden and R. E. Huie and C. E. Kolb and M. J. Kurylo and G. K. Moortgat and V. L. Orkin and P. H. Wine.: Chemical Kinetics and Photochemical Data for Use in Atmospheric Studies, Evaluation No. 17, JPL Publication 10-6, Jet Propulsion Laboratory, Pasadena, URL <http://jpldataeval.jpl.nasa.gov>, 2011.
- JPL, S. P. Sander and R. R. Friedl and D. M. Golden and M. J. Kurylo and G. K. Moortgat and H. Keller-Rudek and P. H. Wine and A. R. Ravishankara and C. E. Kolb and M. J. Molina and B. J. Finlayson-Pitts and R. E. Huie and V. L. Orkin.: Chemical Kinetics and Photochemical Data for Use in Atmospheric Studies, Evaluation Number 15, JPL Publication 06-2, Jet Propulsion Laboratory, Pasadena, CA, URL <http://jpldataeval.jpl.nasa.gov>, 2006.
- Lamarque, J. F., Emmons, L. K., Hess, P. G., Kinnison, D. E., Tilmes, S., Vitt, F., Heald, C. L., Holland, E. A., Lauritzen, P. H., Neu, J., Orlando, J. J., Rasch, P. J., and Tyndall, G. K.: CAM-chem: description and evaluation of interactive atmospheric chemistry in the Community Earth System Model, *Geoscientific Model Development*, 5, 369–411, doi:10.5194/gmd-5-369-2012, 2012.
- Müller, J.-F., Peeters, J., and Stavrakou, T.: Fast photolysis of carbonyl nitrates from isoprene, *Atmospheric Chemistry and Physics*, 14, 2497–2508, doi:10.5194/acp-14-2497-2014, 2014.
- Nguyen, T. L., Lee, H., Matthews, D. A., McCarthy, M. C., and Stanton, J. F.: Stabilization of the Simplest Criegee Intermediate from the Reaction between Ozone and Ethylene: A High-Level Quantum Chemical and Kinetic Analysis of Ozonolysis, *The Journal of Physical Chemistry A*, 119, 5524–5533, doi:10.1021/acs.jpca.5b02088, PMID: 25945650, 2015.
- Niki, H., Maker, P., Savage, C., and Breitenbach, L.: AN FTIR STUDY OF THE CL-ATOM-INITIATED REACTION OF GLYOXAL, *INTERNATIONAL JOURNAL OF CHEMICAL KINETICS*, 17, 547–558, doi:10.1002/kin.550170507, 1985.
- Niki, H., Maker, P., Savage, C., and Hurley, M.: FOURIER-TRANSFORM INFRARED STUDY OF THE KINETICS AND MECHANISMS FOR THE CL-ATOM-INITIATED AND HO-RADICAL-INITIATED OXIDATION OF GLYCOLALDEHYDE, *JOURNAL OF PHYSICAL CHEMISTRY*, 91, 2174–2178, doi:10.1021/j100292a038, 1987.
- Orlando, J. and Burkholder, J.: Identification of BrONO as the major product in the gas-phase reaction of Br with NO<sub>2</sub>, *JOURNAL OF PHYSICAL CHEMISTRY A*, 104, 2048–2053, doi:10.1021/jp993713g, 2000.
- Orlando, J., Tyndall, G., Bertman, S., Chen, W., and Burkholder, J.: Rate coefficient for the reaction of OH with CH<sub>2</sub>=C(CH<sub>3</sub>)C(O)OONO<sub>2</sub> (MPAN), *ATMOSPHERIC ENVIRONMENT*, 36, 1895–1900, doi:10.1016/S1352-2310(02)00090-0, 2002.

- Orlando, J. J. and Tyndall, G. S.: Rate coefficients for the thermal decomposition of  $\text{BrONO}_2$  and the heat of formation of  $\text{BrONO}_2$ , *J. Phys. Chem.*, 100, 19 398–19 405, 1996.
- Orlando, J. J. and Tyndall, G. S.: Laboratory studies of organic peroxy radical chemistry: an overview with emphasis on recent issues of atmospheric significance, *CHEMICAL SOCIETY REVIEWS*, 41, 6294–6317, doi:10.1039/c2cs35166h, 2012.
- Paulot, F., Crounse, J. D., Kjaergaard, H. G., Kurten, A., St Clair, J. M., Seinfeld, J. H., and Wennberg, P. O.: Unexpected Epoxide Formation in the Gas-Phase Photooxidation of Isoprene, *Science*, 325, 730–733, doi:10.1126/science.1172910, 2009.
- Peeters, J., Boullart, W., Pultau, V., Vandenberg, S., and Vereecken, L.: Structure-activity relationship for the addition of OH to (poly)alkenes: Site-specific and total rate constants, *JOURNAL OF PHYSICAL CHEMISTRY A*, 111, 1618–1631, doi:10.1021/jp066973o, 2007.
- Praske, E., Crounse, J. D., Bates, K. H., Kurten, T., Kjaergaard, H. G., and Wennberg, P. O.: Atmospheric Fate of Methyl Vinyl Ketone: Peroxy Radical Reactions with NO and HO<sub>2</sub>, *JOURNAL OF PHYSICAL CHEMISTRY A*, 119, 4562–4572, doi:10.1021/jp5107058, 2015.
- Qian, H.-B., Turton, D., Seakins, P. W., and Pilling, M. J.: A laser flash photolysis/IR diode laser absorption study of the reaction of chlorine atoms with selected alkanes, *International Journal of Chemical Kinetics*, 34, 86–94, doi:10.1002/kin.10025, 2002.
- Rindelaub, J. D., McAvey, K. M., and Shepson, P. B.: The photochemical production of organic nitrates from alpha-pinene and loss via acid-dependent particle phase hydrolysis, *ATMOSPHERIC ENVIRONMENT*, 100, 193–201, doi:10.1016/j.atmosenv.2014.10.010, 2015.
- Sander, R.: Modeling atmospheric chemistry: Interactions between gas-phase species and liquid cloud/aerosol particles, *SURVEYS IN GEOPHYSICS*, 20, 1–31, doi:10.1023/A:1006501706704, 1999.
- Sander, R.: Compilation of Henry’s law constants (version 4.0) for water as solvent, *ATMOSPHERIC CHEMISTRY AND PHYSICS*, 15, 4399–4981, doi:10.5194/acp-15-4399-2015, 2015.
- Schwartz, S. E. and White, W. H.: Solubility equilibria of the nitrogen oxides and oxyacids in dilute aqueous solution, in: *Advances in Environmental Science and Engineering*, edited by Pfafflin, J. R. and Ziegler, E. N., vol. 4, pp. 1–45, Gordon and Breach Science Publishers, NY, 1981.
- Shallcross, D. E., Leather, K. E., Bacak, A., Xiao, P., Lee, E. P. F., Ng, M., Mok, D. K. W., Dyke, J. M., Hossaini, R., Chipperfield, M. P., Khan, M. A. H., and Percival, C. J.: Reaction between  $\text{CH}_3\text{O}_2$  and BrO radicals: a new source of upper troposphere lower stratosphere hydroxyl radicals, *J. Phys. Chem. A*, 119, 4618–4632, 2015.
- St. Clair, J., Moon, S., Holbrook, W. S., Perron, J. T., Riebe, C. S., Martel, S. J., Carr, B., Harman, C., Singha, K., and Richter, D. d.: Geophysical imaging reveals topographic stress control of bedrock weathering, *Science*, 350, 534–538, doi:10.1126/science.aab2210, 2015.
- Stadtler, S., Simpson, D., Schröder, S., Taraborrelli, D., Bott, A., and Schultz, M.: Ozone Impacts of Gas-Aerosol Uptake in Global Chemistry Transport Models, *Atmos. Chem. Phys. Discuss.*, 2017, 1–35, doi:10.5194/acp-2017-566, URL <https://www.atmos-chem-phys-discuss.net/acp-2017-566/>, 2017.
- Strekowski, R., Nicovich, I., and Wine, P.: Kinetic and mechanistic study of the reaction of O(D-1) with  $\text{CF}_2\text{HBr}$ , *INTERNATIONAL JOURNAL OF CHEMICAL KINETICS*, 33, 262–270, doi:10.1002/kin.1019, 2001.

- Taraborrelli, D., Lawrence, M. G., Butler, T. M., Sander, R., and Lelieveld, J.: Mainz Isoprene Mechanism 2 (MIM2): an isoprene oxidation mechanism for regional and global atmospheric modelling, *Atmospheric Chemistry and Physics*, 9, 2751–2777, 2009.
- Wine, P. H., Nicovich, J. M., Stickel, R. E., Zhao, Z., Shackelford, C. J., Kreutter, K. D., Daykin, E. P., and Wang, S.: Halogen and Sulphur Reactions Relevant to Polar Chemistry, *The Tropospheric Chemistry of Ozone in the Polar Regions*, 1993.