

Species	z_b (m)	$[\sigma_{zd}^{\uparrow}, \sigma_{zd}^{\downarrow}]$ (m)	No. of points	Obs. living range (m)	References
<i>G. ruber</i>	-10] $0, -30]$	130	0–120	Ganssen and Kroon (2000); Anand et al. (2003); Kuroyanagi and Kawahata (2004); Farmer et al. (2007)
<i>N. incompta</i>	-65	[$-35, -150]$	60	0–200	Pak and Kennett (2002); Pak et al. (2004); Kuroyanagi and Kawahata (2004); von Langen et al. (2005)
<i>T. sacculifer</i>	-100	[$-75, -200]$	46	0–200	Ganssen and Kroon (2000); Anand et al. (2003); Kuroyanagi and Kawahata (2004); Farmer et al. (2007)
<i>G. bulloides</i>	-400	[$-100, -\infty[$	123	0–300	Bauch et al. (2002); Ganssen and Kroon (2000); Pak and Kennett (2002); Pak et al. (2004); Kuroyanagi and Kawahata (2004)
<i>N. pachyderma</i>	-550 ¹	[$-275, -900]$	244	0–500	Kohfeld et al. (1996); Bauch et al. (1997, 2002); Simstich et al. (2003); Kuroyanagi and Kawahata (2004)