

Supplementary information to:

Partitioning soil organic carbon into its centennially active and stable fractions with machine-learning models based on Rock-Eval® thermal analysis (PARTY_{SOC}v2.0 and PARTY_{SOC}v2.0_{EU})

Lauric Cécillon^{1,2}, François Baudin³, Claire Chenu⁴, Bent T. Christensen⁵, Uwe Franko⁶, Sabine Houot⁴, Eva Kanari^{2,3}, Thomas Kätterer⁷, Ines Merbach⁸, Folkert van Oort⁴, Christopher Poeplau⁹, Juan Carlos Quezada^{10,11,12}, Florence Savignac³, Laure N. Soucémarianadin¹³, Pierre Barré²

¹Normandie Univ., UNIROUEN, INRAE, ECODIV, Rouen, France

²Laboratoire de Géologie, École normale supérieure, CNRS, PSL Univ., IPSL, Paris, France

³Institut des Sciences de la Terre de Paris, Sorbonne Université, CNRS, Paris, 75005, France

⁴UMR 1402 ECOSYS, INRAE, AgroParisTech, Univ. Paris Saclay, Thiverval-Grignon, 78850, France

⁵Department of Agroecology, Aarhus University, AU Foulum, 8830 Tjele, Denmark

⁶Department of soil system science, Helmholtz Centre for Environmental Research, UFZ, 06120 Halle Germany

⁷Department of Ecology, Swedish University of Agricultural Sciences, 75007 Uppsala, Sweden

⁸Department Community Ecology, Helmholtz Centre for Environmental Research, UFZ, 06246 Bad Lauchstädt, Germany

⁹Thünen Institute of Climate-Smart Agriculture, 38116 Braunschweig, Germany

¹⁰Laboratory of Ecological Systems ECOS and Laboratory of Plant Ecology Research PERL, School of Architecture, Civil and Environmental Engineering ENAC, École Polytechnique Fédérale de Lausanne EPFL, 1015 Lausanne, Switzerland

¹¹Swiss Federal Institute for Forest, Snow and Landscape Research WSL, 1015 Lausanne, Switzerland

¹²Ecosystem Management, Institute of Terrestrial Ecosystems, Department of Environmental Systems Science, ETHZ, 8092 Zürich, Switzerland

¹³ACTA – les instituts techniques agricoles, 75595 Paris, France

Correspondence to: Lauric Cécillon (lauric.cecillon@inrae.fr)




Figure S1: Sensitivity of the $\text{PARTY}_{\text{SOCv2.0}_{\text{EU}}}$ machine-learning model to the test set. Model sensitivity to the test set was assessed as its sensitivity to independent test samples (1) from a reference soil group not existing in the training set (*i.e.* excluding Chernozem soil samples from the test set); (2) that are unlikely to be encountered in agricultural soils (*i.e.* excluding from the test set soils sampled at late dates of bare fallow treatments, more than 25 years after the experiment onset, which cannot represent soils with regular carbon input). This figure is built with a test set composed exclusively of soils from independent Cambisols and Luvisols of Northwestern Europe. It should be compared to Fig. 2b (for $\text{PARTY}_{\text{SOCv2.0}_{\text{EU}}}$) that is built using the exact same training set and validation procedure (leave-one-site-out), but that uses a different test set including 32 additional independent test samples (15 Chernozem soils samples and 17 soil samples collected at late dates in bare fallow treatments).

Table S1: Main changes between the first version of PARTY_{SOC} and the second version of the model (PARTY_{SOC}v2.0 and PARTY_{SOC}v2.0_{EU}). Abbreviations: SOC, soil organic carbon; SOC_{EA}: soil organic carbon content determined by elemental analysis; TOC_{RE6}, soil organic carbon content determined by Rock-Eval® thermal analysis; MinC, soil inorganic carbon content determined by Rock-Eval® thermal analysis; LTBF, long-term bare fallow.

| | First version of PARTY_{SOC} (Cécillon et al., 2018) | PARTY_{SOC}v2.0 (This study) | PARTY_{SOC}v2.0_{EU} (This study) |
|---|---|---|---|
| Number of reference sites | 4 | 7 | 6 |
| Method to estimate the centennially stable SOC content at the reference sites | Exclusively inferred from plant-free LTBF treatments; and calculated exclusively by modelling the decline with time of SOC initially present at the onset of the experiment | Inferred from plant-free LTBF treatments (6 sites) and from one vegetated site (La Cabaña); and estimated by modelling the decline with time of SOC initially present at the onset of the experiment, or by using a measured value of SOC content or C ₄ -plant derived SOC that is lower than the parameter <i>c</i> of Eq. (1) | Exclusively inferred from plant-free LTBF treatments; and estimated by modelling the decline with time of SOC initially present at the onset of the experiment, or by using a measured value of SOC content or C ₄ -plant derived SOC that is lower than the parameter <i>c</i> of Eq. (1) |
| Method to estimate the centennially stable SOC proportion in reference topsoil samples | By dividing the site-specific content of the centennially stable SOC content by the SOC _{EA} content of topsoil samples | By dividing the site-specific content of the centennially stable SOC content by the TOC _{RE6} or the TOC _{RE6} +MinC content of topsoil samples | By dividing the site-specific content of the centennially stable SOC content by the TOC _{RE6} or the TOC _{RE6} +MinC content of topsoil samples |
| Number of reference topsoil samples (per site) | 118 (unequal number of samples per site) | 105 (15 samples per reference site) | 90 (15 samples per reference site) |
| Rock-Eval® predictor variables | 30, not directly related to SOC content, not necessarily correlated to the proportion of the centennially stable SOC fraction | 18, some of them are directly related to SOC content, all are highly correlated to the proportion of the centennially stable SOC fraction (Spearman's rho > 0.5) | 18, some of them are directly related to SOC content, all are highly correlated to the proportion of the centennially stable SOC fraction (Spearman's rho > 0.5) |
| Criteria regarding the inclusion of topsoils in the | None (all available reference topsoil samples are used) | Exclusion of reference topsoil samples (1) from treatments experiencing repeated | Exclusion of reference topsoil samples (1) from treatments experiencing repeated |

| | | | |
|--|---|--|---|
| reference sample set of the model | | applications of compost or manure; (2) having an organic carbon yield of Rock-Eval® analysis below 86% or above 116%. Then, selection of 15 reference topsoil samples per site (1) enhancing the range of the centennially stable SOC proportion at that site; (2) having the best organic carbon yield of Rock-Eval® analysis | applications of compost or manure; (2) having an organic carbon yield of Rock-Eval® analysis below 86% or above 116%. Then, selection of 15 reference topsoil samples per site (1) enhancing the range of the centennially stable SOC proportion; (2) having the best organic carbon yield of Rock-Eval® analysis |
| Validation procedure for the machine-learning model | Mostly “random splitting” procedure, “leave-one-site-out” procedure tested only for one site (Ultuna) | Fully independent “leave-one-site-out” procedure | Fully independent “leave-one-site-out” procedure |

Table S2: Basic characteristics of the seven reference sites used as the training set of PARTY_{soC}v2.0, and site-specific values of the parameter c in Eq. (1). Most data are taken from Barré et al. (2010), Cécillon et al. (2018), Franko and Merbach (2017) and Quezada et al. (2019). Abbreviations: SFE, static fertilization experiment (V120); FE, fallow experiment (V505a).

| Site, Country | Latitude, longitude [° min s] | Mean annual temperature [°C] | Mean annual precipitation [mm] | Soil type [WRB; FAO, 2014] | Land use before experiment | Sampling depth [cm] | pH (in H ₂ O) | Soil texture [%] (clay/silt/sand) | Parameter c in Eq. (1) (g C kg ⁻¹) |
|-----------------------------------|---|---------------------------------------|--------------------------------------|-------------------------------|----------------------------------|---------------------------|-----------------------------|---|--|
| Versailles, France | 48°48'12.76"N; 2°05'09.95"E | 10.7 | 628 | Haplic Luvisol | grassland | 0–25 | 5.6 to 6.4 | 17/57/26 | 6.22 |
| Rothamsted, England | 51°48'14.12"N; 0°21'41.39"E | 9.5 | 712 | Chromic Luvisol | grassland | 0–23 | 5.2 to 6.3 | 25/62/13 | 10.46 |
| Ultuna, Sweden | 59°48'46.00"N; 17°39'01.75"E | 5.5 | 533 | Eutric Cambisol | arable | 0–20 | 6.6 | 36/41/23 | 6.95 |
| Grignon, France | 48°51'01.35"N; 1°57'03.55"E | 10.7 | 649 | Calcaric Cambisol | grassland | 0–25 | 8.0 to 8.3 | 16/54/30 | 7.12 |
| Askov, Denmark | B3 55°28'12.89"N; 9°06'41.77"E B4 55°28'19.19"N; 9°07'00.02"E | 7.8 | 862 | Arenic Luvisol | arable | 0–20 | 5.5 to 6.5 | 10/11/79 | 5.10 |
| Bad Lauchstädt, Germany | SFE 51°23'24.34"N; 11°52'48.48"E FE 51°23'34.75"N; 11°52'39.61"E | 8.9 | 484 | Haplic Chernozem | arable | 0–20 (SFE) 0–30 (FE) | 6.9 to 7.4 | 21/68/11 | 16.22 |
| La Cabaña, Colombia | 4°16'N; 73°22'W | 27 | 3400 | Dystric Cambisol | grassland | 0–10 | 4.0 to 4.6 | 30/34/36 | 5.12 |

Table S3: List of the 105 reference topsoil samples retained as the training set of PARTY_{SOC}v2.0, including information on their reference site, land cover, agronomical treatment, sampling year and values for the 40 Rock-Eval® parameters considered in this study. See the Section 2.2 for a description of the 40 Rock-Eval® parameters and their units. Supplementary Table S3 is provided as a separate csv file.

Table S4: Basic statistics of the 40 Rock-Eval® parameters, calculated on the 105 reference topsoil samples of the training set of PARTY_{SOC}v2.0. See the Section 2.2 for a description of the 40 Rock-Eval® parameters. The 18 Rock-Eval® parameters retained as predictor variables for the second version of PARTY_{SOC} are shown in bold.

| Rock-Eval® parameter (unit) | Mean | Median | Minimum | Maximum | Standard deviation |
|-----------------------------------|------------|------------|------------|------------|--------------------|
| T10 _{HC_PYR} (°C) | 310 | 309 | 279 | 343 | 16 |
| T30 _{HC_PYR} (°C) | 372 | 372 | 330 | 402 | 15 |
| T50 _{HC_PYR} (°C) | 422 | 421 | 389 | 445 | 12 |
| T70_{HC_PYR} (°C) | 465 | 462 | 446 | 484 | 10 |
| T90_{HC_PYR} (°C) | 523 | 522 | 495 | 547 | 15 |
| T10 _{CO_PYR} (°C) | 307 | 305 | 295 | 318 | 5 |
| T30 _{CO_PYR} (°C) | 358 | 357 | 346 | 373 | 6 |
| T50 _{CO_PYR} (°C) | 405 | 403 | 391 | 423 | 7 |
| T70 _{CO_PYR} (°C) | 460 | 458 | 451 | 477 | 7 |
| T90 _{CO_PYR} (°C) | 525 | 524 | 519 | 539 | 4 |
| T10 _{CO2_PYR} (°C) | 286 | 284 | 276 | 300 | 7 |
| T30_{CO2_PYR} (°C) | 343 | 339 | 329 | 365 | 10 |
| T50_{CO2_PYR} (°C) | 392 | 387 | 375 | 419 | 13 |
| T70_{CO2_PYR} (°C) | 445 | 441 | 427 | 469 | 13 |
| T90_{CO2_PYR} (°C) | 511 | 510 | 497 | 526 | 8 |
| T10 _{CO_OX} (°C) | 323 | 317 | 299 | 383 | 16 |
| T30 _{CO_OX} (°C) | 374 | 364 | 331 | 495 | 32 |
| T50 _{CO_OX} (°C) | 421 | 400 | 366 | 578 | 46 |
| T70_{CO_OX} (°C) | 481 | 464 | 404 | 671 | 52 |
| T90 _{CO_OX} (°C) | 568 | 547 | 477 | 776 | 64 |
| T10 _{CO2_OX} (°C) | 327 | 326 | 318 | 358 | 6 |
| T30 _{CO2_OX} (°C) | 375 | 374 | 363 | 405 | 8 |
| T50_{CO2_OX} (°C) | 417 | 413 | 398 | 455 | 14 |
| T70_{CO2_OX} (°C) | 469 | 465 | 437 | 509 | 19 |
| T90_{CO2_OX} (°C) | 540 | 538 | 511 | 568 | 12 |
| I-index (unitless) | 0.18 | 0.17 | 0.02 | 0.37 | 0.08 |
| R-index (unitless) | 0.59 | 0.59 | 0.47 | 0.70 | 0.05 |
| TLHC-index (unitless) | 0.63 | 0.64 | 0.53 | 0.72 | 0.05 |

| | | | | | | |
|-----------------------------|--|-------------|-------------|-------------|-------------|-------------|
| HI | (mg HC g⁻¹ C) | 156 | 143 | 69 | 284 | 58 |
| OI _{RE6} | (mg O ₂ g ⁻¹ C) | 183 | 184 | 146 | 246 | 21 |
| TOC_{RE6} | (g C kg⁻¹) | 16.0 | 14.3 | 5.6 | 41.5 | 7.3 |
| MinC | (g C kg ⁻¹) | 2.7 | 1.7 | 0.6 | 12.5 | 2.8 |
| PC | (g C kg⁻¹) | 3.7 | 3 | 1.1 | 11 | 2.2 |
| S2 | (g C kg⁻¹) | 2.2 | 1.7 | 0.6 | 7.4 | 1.6 |
| PseudoS1 | (g C kg⁻¹) | 0.12 | 0.09 | 0.05 | 0.51 | 0.07 |
| PseudoS1/PC | (unitless) | 0.04 | 0.03 | 0.02 | 0.11 | 0.02 |
| PseudoS1/TOC _{RE6} | (unitless) | 0.008 | 0.007 | 0.003 | 0.023 | 0.004 |
| S2/PC | (unitless) | 0.56 | 0.54 | 0.42 | 0.76 | 0.09 |
| PC/TOC_{RE6} | (unitless) | 0.22 | 0.22 | 0.14 | 0.32 | 0.05 |
| HI/OI_{RE6} | (mg HC mg⁻¹ O₂) | 0.87 | 0.74 | 0.43 | 1.92 | 0.39 |

Table S5: Sensitivity of model performance to the reference sites included in the training set, using 15 topsoil samples from the sites of Grignon (a) or Versailles (b) as test sets. Site-specific performance statistics are calculated on a test set made of topsoil samples from Grignon or Versailles. Abbreviations are defined in Section 2.5.

| a | | | | | |
|---|--|---|--|------------------------------------|----------------------------|
| Grignon topsoil samples used as test set (n = 15) | | | | | |
| Composition of the training set | 6 reference sites | 5 reference sites | 4 reference sites | 3 reference sites | 2 reference sites |
| | (PARTY_{SOC}v2.0, Grignon out) | (PARTY_{SOC}v2.0_{EU}, Grignon out) | (Ultuna, Versailles, Rothamsted, Askov) | (Ultuna, Versailles, Askov) | (Versailles, Askov) |
| R²_{OOB} | 0.82 | 0.88 | 0.79 | 0.86 | 0.88 |
| RMSEP_{OOB} | 0.09 | 0.07 | 0.08 | 0.06 | 0.06 |
| R² | 0.56 | 0.44 | 0.86 | 0.82 | 0.87 |
| RMSEP | 0.11 | 0.18 | 0.07 | 0.06 | 0.07 |
| _RRMSEP | 0.18 | 0.28 | 0.11 | 0.10 | 0.10 |
| RPIQ | 0.74 | 0.47 | 1.22 | 1.36 | 1.27 |
| Bias | 0.088 | 0.157 | 0.003 | -0.035 | -0.042 |
| b | | | | | |
| Versailles topsoil samples used as test set (n = 15) | | | | | |
| Composition of the training set | 6 reference sites | 5 reference sites | 4 reference sites | 3 reference sites | 2 reference sites |
| | (PARTY_{SOC}v2.0, Versailles out) | (PARTY_{SOC}v2.0_{EU}, Versailles out) | (Ultuna, Grignon, Askov, Rothamsted) | (Ultuna, Grignon, Askov) | (Ultuna, Grignon) |
| R²_{OOB} | 0.85 | 0.89 | 0.83 | 0.89 | 0.85 |
| RMSEP_{OOB} | 0.08 | 0.07 | 0.06 | 0.04 | 0.05 |
| R² | 0.48 | 0.66 | 0.79 | 0.83 | 0.79 |
| RMSEP | 0.17 | 0.14 | 0.13 | 0.10 | 0.11 |
| _RRMSEP | 0.28 | 0.24 | 0.23 | 0.17 | 0.19 |
| RPIQ | 1.77 | 2.08 | 2.20 | 2.90 | 2.67 |
| Bias | 0.036 | 0.018 | 0.052 | 0.006 | 0.025 |

Supplementary references

Barré, P., Eglin, T., Christensen, B. T., Ciais, P., Houot, S., Kätterer, T., van Oort, F., Peylin, P., Poulton, P. R., Romanenkov, V. and Chenu, C.: Quantifying and isolating stable soil organic carbon using long-term bare fallow experiments, *Biogeosciences*, 7(11), 3839–3850, <https://doi.org/10.5194/bg-7-3839-2010>, 2010.

Cécillon, L., Baudin, F., Chenu, C., Houot, S., Jolivet, R., Kätterer, T., Lutfalla, S., Macdonald, A., van Oort, F., Plante, A. F., Savignac, F., Soucémarianadin, L. N. and Barré, P.: A model based on Rock-Eval thermal analysis to quantify the size of the centennially persistent organic carbon pool in temperate soils, *Biogeosciences*, 15(9), 2835–2849, <https://doi.org/10.5194/bg-15-2835-2018>, 2018.

FAO: World reference base for soil resources 2014: international soil classification system for naming soils and creating legends for soil maps., FAO, Rome., 2014.

Franke, U. and Merbach, I.: Modelling soil organic matter dynamics on a bare fallow Chernozem soil in Central Germany, *Geoderma*, 303, 93–98, <https://doi.org/10.1016/j.geoderma.2017.05.013>, 2017.

Quezada, J. C., Etter, A., Ghazoul, J., Buttler, A. and Guillaume, T.: Carbon neutral expansion of oil palm plantations in the Neotropics, *Sci. Adv.*, 5(11), eaaw4418, <https://doi.org/10.1126/sciadv.aaw4418>, 2019.