



Supplement of

Added value of EURO-CORDEX high-resolution downscaling over the Iberian Peninsula revisited – Part 2: Max and min temperature

João António Martins Careto et al.

Correspondence to: João António Martins Careto (jacareto@fc.ul.pt)

The copyright of individual parts of the supplement might differ from the article licence.

Table S1. Regional models forced by the ERA-Interim reanalysis from the European Centre for Medium-Range Weather Forecasts, for the 1989-2008 period.

RCM	Reference	acronym
CLMcom-CCLM4-8-17	Keuler et al. (2016)	CCLM
ETH-COSMO-crCLIM-v1-1	Pothapakula et al. (2020), Vautard et al. (2020)	ETH
CNRM-ALADIN53	Colin et al. (2010), Herrmann et al. (2011)	CNRM53
CNRM-ALADIN63	Daniel et al. (2019), Nabat et al. (2020)	CNRM63
DHMZ-RegCM4-2	Giorgi et al. (2012)	DHMZ
DMI-HIRHAM5	Christensen et al. (2007)	DMI
GERICS-REMO2015	Remedio et al. (2019)	GERICS
ICTP-RegCM4-6	Giorgi et al. (2012)	ICTP
IPSL-INNERIS-WRF381P	Vautard et al. (2013)	IPSL
KNMI-RACMO22E	van Meijgaard et al. (2008)	KNMI
MPI-CSC-REMO02009	Jacob et al. (2012)	MPI
SMHI-RCA4	Samuelsson et al. (2011)	SMHI
HadREM3-GA7-05	Tinker et al. (2015)	MOHC

Table S2. EURO-CORDEX Regional models driven by the CMIP5 GCMs. Also shown the approximate spatial resolution from each GCM taken from <https://portal.enes.org/data/enes-model-data/cmip5/resolution>. References: (A) Keuler et al. (2016) (B) Colin et al (2010), Herrmann et al (2011), (C) Daniel et al (2019), Nabat et al (2020), (D) Christensen et al. (2007), (E) Remedio et al. (2019), (F) Vautard et al. (2013), (G) van Meijgaard et al. (2008), (H) Samuelsson et al. (2011), (I) Tinker et al. (2015), (J) Giorgi et al. (2012), (K) Jacob et al. (2012).

CMIP5 GCM	Variant	Resolution	RCM	Reference	Acronym
CNRM-CERFACS-CNRM-CM5	r1i1p1	1.40° X 1.41°	CLMcom-CCLM4-8-17	A	CNRM-CCLM
			CNRM-ALADIN53	B	CNRM-CNRM53
			CNRM-ALADIN63	C	CNRM-CNRM63
			DMI-HIRHAM5	D	CNRM-DMI
			GERICS-REMO2015	E	CNRM-GERICS
			IPSL-WRF381P	F	CNRM-IPSL
			KNMI-RACMO22E	G	CNRM-KNMI
			SMHI-RCA4	H	CNRM-SMHI
ICHEC-EC-EARTH	r1i1p1	1.12° X 1.13°	DMI-HIRHAM5	D	ICHEC1-DMI
			KNMI-RACMO22E	G	ICHEC1-KNMI
			SMHI-RCA4	H	ICHEC1-SMHI
			CLMcom-CCLM4-8-17	A	ICHEC2-CCLM
IPSL-CM5A-LR	r1i1p1	1.89° X 3.75°	ETH-COSMO-crCLIM-v1-1	A	ICHEC2-ETH
			DMI-HIRHAM5	D	ICHEC2-DMI
			IPSL-WRF381P	F	ICHEC2-IPSL
			KNMI-RACMO22E	G	ICHEC2-KNMI
			MOHC-HadREM3-GA7-05	I	ICHEC2-MOHC
IPSL-CM5A-MR	r1i1p1	1.27° X 2.5°	SMHI-RCA4	H	ICHEC2-SMHI
			CLMcom-CCLM4-8-17	A	MPI1-CCLM
			ETH-COSMO-crCLIM-v1-1	A	MPI1-ETH
			CNRM-ALADIN63	C	MPI1-CNRM63
MPI-ESM-LR	r1i1p1	1.87° X 1.88°	DMI-HIRHAM5	D	MPI1-DMI
			ICTP-RegCM4-6	J	MPI1-ICTP
			KNMI-RACMO22E	G	MPI1-KNMI
			MPI-REMO2009	K	MPI1-MPI
			SMHI-RCA4	H	MPI1-SMHI
			ETH-COSMO-crCLIM-v1-1	A	MPI2-ETHZ
			MPI-REMO2009-MPI2-MPI	K	MPI2-MPI
			SMHI-RCA4	H	MPI2-SMHI
MOHC-HadGEM2-ES	r3i1p1	1.25° X 1.88°	ETH-COSMO-crCLIM-v1-1	A	MPI3-ETH
			GERICS-REMO2015	E	MPI3-GERICS
			SMHI-RCA4	H	MPI3-SMHI
			CLMcom-CCLM4-8-17	A	MOHC-CCLM
NCC-NorESM1-M	r1i1p1	1.89° X 2.5°	ETH-COSMO-crCLIM-v1-1	A	MOHC-ETH
			CNRM-ALADIN63	C	MOHC-CNRM
			DMI-HIRHAM5	D	MOHC-DMI
			ICTP-RegCM4-6	J	MOHC-ICTP
			IPSL-WRF381P	F	MOHC-IPSL
			KNMI-RACMO22E	G	MOHC-KNMI
			MOHC-HadREM3-GA7-05	I	MOHC-MOHC
			SMHI-RCA4	H	MOHC-SMHI
NOAA-GFDL-ESM2G	r1i1p1	2.02° X 2.00°	ETH-COSMO-crCLIM-v1-1	A	NCC-ETH
			DMI-HIRHAM5	D	NCC-DMI
			GERICS-REMO2015	E	NCC-GERICS
			IPSL-WRF381P	F	NCC-IPSL
			KNMI-RACMO22E	G	NCC-KNMI
			MOHC-HadREM3-GA7-05	I	NCC-MOHC
			SMHI-RCA4	H	NCC-SMHI
NOAA-GFDL-ESM2G			GERICS-REMO2015	E	NOAA-GERICS




Figure S1. Maximum (left) and minimum (right) daily temperature distributions taken from the hindcast EURO-CORDEX RCMs and ERA-Interim reanalysis (1989-2008) for the Iberian Peninsula. Also shown the Iberian Gridded Dataset distribution for the same domain and period. All RCM data was previously interpolated into the IGD 0.1° regular resolution. As for Era-Interim, two PDFs are shown, one for the original resolution of the low-resolution and other interpolated into the IGD resolution. The dash point and the value written refers to either the 90th percentile for max temperature or the 10th percentile for min temperature of the observations for NGD on the original resolution (blue) and interpolated into the ERA-Interim resolution (red). The time periods are (a) Year, (b) DJF, (c) MAM, (d) JJA and (e) SON.

(a)

Max Temperature Frequency 0-100

YEAR	-1.2	0.4	2.7	1.2	-5.6	-0.9	5.0	-2.5	6.0	1.7	5.0	-1.3	0.3
DJF	-5.0	2.6	8.8	8.1	-1.2	-8.4	8.0	-7.5	9.0	1.6	8.5	-3.6	-1.1
MAM	6.0	8.4	12.4	9.6	-5.8	2.0	13.3	-5.2	11.7	3.3	13.4	-3.4	3.7
JJA	8.1	5.9	4.5	4.8	3.7	6.3	9.9	1.3	13.4	11.0	9.4	14.7	10.9
SON	4.1	4.2	6.6	4.6	-0.9	2.3	5.0	-3.3	9.7	4.2	5.9	2.6	5.4

(b)

Min Temperature Frequency 0-100

YEAR	-4.9	-4.1	-10.2	-7.1	0.7	-6.7	-5.7	-1.3	-0.2	-2.4	-5.7	-4.4	-3.0
DJF	-8.1	-10.3	-20.1	-7.1	-3.3	-7.8	-12.6	-2.4	-5.6	-8.7	-12.9	-10.8	-1.3
MAM	-7.9	-10.6	-13.1	-2.2	-2.0	-12.7	-9.6	0.4	-3.0	-5.9	-10.0	-8.3	-0.7
JJA	-14.3	-6.1	-2.1	-14.2	6.2	-17.6	-12.3	-5.8	-0.9	6.1	-12.0	3.6	-7.3
SON	-7.6	-6.4	-6.1	-3.8	3.6	-10.4	-8.3	-0.1	0.5	0.9	-7.2	-1.4	0.8

(c)

Max Temperature Frequency 90-100

YEAR	2.2	-1.1	2.9	2.2	3.5	12.2	5.1	5.2	9.1	16.0	5.2	14.8	9.5
DJF	11.3	9.2	13.3	7.6	7.3	-5.4	14.3	-15.4	13.6	6.7	13.9	-4.0	0.2
MAM	4.8	-2.6	0.1	-1.9	2.0	2.5	7.6	3.0	2.7	6.3	8.0	4.9	2.9
JJA	8.5	5.2	15.0	11.2	16.8	22.1	12.3	8.0	17.3	24.2	12.3	24.3	21.6
SON	3.0	-3.2	2.6	-0.9	7.8	11.2	-3.2	1.2	10.9	7.8	-0.3	11.6	9.3

(d)

Min Temperature Frequency 0-10

YEAR	-9.0	-9.2	-4.1	-17.3	-8.5	-3.2	-26.1	-7.4	-9.1	-4.2	-26.3	-20.1	-10.2
DJF	0.1	-0.7	-0.1	-10.7	-3.4	-3.2	-17.0	-3.7	-4.1	-4.7	-20.4	-16.7	-12.4
MAM	-3.4	-1.4	-1.1	-3.8	2.3	0.1	-17.1	1.7	0.5	-5.0	-20.8	-13.4	1.3
JJA	1.4	1.4	-8.1	-1.4	-0.1	1.3	2.5	-2.9	0.8	1.1	2.9	-6.3	2.5
SON	-9.3	-10.1	-5.3	-11.1	-4.4	-6.8	-20.0	-2.3	-5.8	-1.6	-20.4	-14.4	-4.0




Figure S2. Yearly and seasonal distribution added values (DAV) of the Iberian Peninsula, between the RCMs and the ERA-Interim reanalysis for the 1989-2008 period, taken from the Hindcast EURO-CORDEX simulations, with the IGD regular dataset as a reference for (a) maximum daily temperature, considering the whole PDF, (b) minimum daily temperature considering the whole PDF. (c) maximum daily temperature extremes, only considering the values above the observational 90th percentile from maximum temperatures and (d) minimum temperature extremes only considering the values below the observational 10th percentile from minimum temperatures. All model data were previously interpolated to 0.1° regular resolution from the observations.




Figure S3. Maximum (left) and minimum (right) daily temperature distribution for the historical EURO-CORDEX RCMs driven by the CMIP5 GCMs, for the Iberian Peninsula, considering the 1971-2005 period, where all results were previously interpolated into the observational grid. The dash point and the value written refers to the 90th percentile of the observations for the maximum temperature and to the 10th percentile of the observations for the minimum temperature. (a) Year, (b) DJF, (c) MAM, (d) JJA and (e) SON.




Figure S4. Maximum (left) and minimum (right) daily temperature distributions for the historical driving CMIP5 GCMs and IGD observations interpolated into each GCM resolution for the Iberian Peninsula, considering the 1971-2005 period. Also shown the PDF from the IGD observations at the original resolution. (a) Year, (b) DJF, (c) MAM, (d) JJA and (e) SON.


(a)

Max Temperature Frequency 0-100



(b)

Min Temperature Frequency 0-100



(c)

Max Temperature Frequency 90-100



(d)

Min Temperature Frequency 0-10



Figure S5. Yearly and seasonal distribution added values (DAV) of the Iberian Peninsula, between the RCMs and the CMIP5 GCMs for the 1989-2008 period, taken from the Historical EURO-CORDEX simulations, with the IGD regular dataset as a reference for (a) maximum daily temperature, considering the whole PDF shown in the left panels of Figure S3, (b) minimum daily temperature considering the whole PDF shown in the right panels of Figure S3, (c) maximum daily temperature extremes, only considering the values above the observational 90th percentile from maximum temperatures and (d) minimum daily temperature extremes, only considering the values below the observational 10th percentile from minimum temperatures. All model data were previously interpolated to 0.1° regular resolution from the observations. The thick blue lines separate the RCMs driven by different GCMs.

References

- Christensen, O.B., Drews, M., Christensen, J.H., Dethloff, K., Ketelsen, K., Hebestadt, I. and Rinke, A.: The HIRHAM regional climate model. Version 5 (beta), 2007, <https://www.dmi.dk/fileadmin/Rapporter/TR/tr06-17.pdf>, last access: 29 April 2021.
- Colin, J., Déqué, M., Radu, R. and Somot, S.: Sensitivity study of heavy precipitation in Limited Area Model climate simulations: influence of the size of the domain and the use of the spectral nudging technique. *Tellus A: Dynamic Meteorology and Oceanography*, 62, 591-604, <https://doi.org/10.1111/j.1600-0870.2010.00467.x>, 2010.
- Daniel, M., Lemonsu, A., Déqué, M., Somot, S., Alias, A. and Masson, V.: Benefits of explicit urban parameterization in regional climate modeling to study climate and city interactions. *Climate Dynamics*, 52, 2745-2764, <https://doi.org/10.1007/s00382-018-4289-x>, 2019.
- Giorgi, F., Coppola, E., Solmon, F., Mariotti, L., Sylla, M.B., Bi, X., Elguindi, N., Diro, G.T., Nair, V., Giuliani, G. and Turuncoglu, U.U.: RegCM4: model description and preliminary tests over multiple CORDEX domains. *Climate Research*, 52, 7-29, <https://doi.org/10.3354/cr01018>, 2012.
- Herrmann, M., Somot, S., Calmanti, S., Dubois, C., and Sevault, F.: Representation of spatial and temporal variability of daily wind speed and of intense wind events over the Mediterranean Sea using dynamical downscaling: impact of the regional climate model configuration, *Nat. Hazards Earth Syst. Sci.*, 11, 1983–2001, <https://doi.org/10.5194/nhess-11-1983-2011>, 2011.
- Jacob, D., Elizalde, A., Haensler, A., Hagemann, S., Kumar, P., Podzun, R., Rechid, D., Remedio, A.R., Saeed, F., Sieck, K. and Teichmann, C.: Assessing the transferability of the regional climate model REMO to different coordinated regional climate downscaling experiment (CORDEX) regions. *Atmosphere*, 3, 181-199, <https://doi.org/10.3390/atmos3010181>, 2012.
- Keuler, K., Radtke, K., Kotlarski, S., and Lüthi, D.: Regional climate change over Europe in COSMO-CLM: Influence of emission scenario and driving global model, *Meteorologische Zeitschrift*, 25, 121–136, <https://doi.org/10.1127/metz/2016/0662>, 2016.
- Nabat, P., Somot, S., Cassou, C., Mallet, M., Michou, M., Bouniol, D., Decharme, B., Drugé, T., Roehrig, R., and Saint-Martin, D.: Modulation of radiative aerosols effects by atmospheric circulation over the Euro-Mediterranean region, *Atmos. Chem. Phys.*, 20, 8315–8349, <https://doi.org/10.5194/acp-20-8315-2020>, 2020.
- Pothapakula, P. K., Primo, C., Sørland, S., and Ahrens, B.: The synergistic impact of ENSO and IOD on the Indian Summer Monsoon Rainfall in observations and climate simulations - an information theory perspective, *Earth Syst. Dynam.*, 11, 903–923, <https://doi.org/10.5194/esd11-903-2020>, 2020.
- Remedio, A.R., Teichmann, C., Buntемeyer, L., Sieck, K., Weber, T., Rechid, D., Hoffmann, P., Nam, C., Kotova, L. and Jacob, D.: Evaluation of new CORDEX simulations using an updated Köppen–Trewartha climate classification. *Atmosphere*, 10, 726, <https://doi.org/10.3390/atmos10110726>, 2019.
- Samuelsson, P., Jones, C.G., Will’ En, U., Ullerstig, A., Gollvik, S., Hansson, U.L.F., Jansson, E., Kjellstro” M, C., Nikulin, G. and Wyser, K.: The Rossby Centre Regional Climate model RCA3: model description and performance. *Tellus A: Dynamic Meteorology and Oceanography*, 63, 4-23, <https://doi.org/10.1111/j.1600-0870.2010.00478.x>, 2011.

Tinker, J., Lowe, J., Holt, J., Pardaens, A. and Wiltshire, A.: Validation of an ensemble modelling system for climate projections for the northwest European shelf seas. *Progress in Oceanography*, 138, 211-237, <https://doi.org/10.1016/j.pocean.2015.07.002>, 2015.

Vautard, R., Gobiet, A., Jacob, D., Belda, M., Colette, A., Déqué, M., Fernández, J., García-Díez, M., Goergen, K., Güttler, I. and Halenka, T.: The simulation of European heat waves from an ensemble of regional climate models within the EURO-CORDEX project. *Climate Dynamics*, 4, 2555-2575, <https://doi.org/10.1007/s00382-013-1714-z>, 2013.

Vautard, R., Kadygrov, N., Iles, C., Boberg, F., Buonomo, E., Buelow, K., Coppola, E., Corre, L., van Meijgaard, E., Nogherotto, R., Sandstad, M., Schwingshakl, C., Somot, S., Aalbers, E. E., Christensen, O., Ciarlo, J., Demory, M.-E., Giorgi, F., Jacob, D., Jones, R. G., Keuler, K., Kjellström, E., Lenderink, G., Levavasseur, G., Nikulin, G., Sillmann, J., Solidoro, C., Sørland, S., Steger, C., Teichmann, C., Warrach-Sagi, K., and Wulfmeyer, V.: Evaluation of the large EURO-CORDEX regional climate model ensemble, *Journal of Geophysical Research: Atmospheres*, <https://doi.org/10.1029/2019JD032344>, 2020.

van Meijgaard, E., Van Ulft, L.H., Van de Berg, W.J., Bosveld, F.C., Van den Hurk, B.J.J.M., Lenderink, G. and Siebesma, A.P., 2008. The KNMI regional atmospheric climate model RACMO, version 2.1 (p. 43). De Bilt, Netherlands: KNMI, <http://citeserx.ist.psu.edu/viewdoc/download?doi=10.1.1.475.9494&rep=rep1&type=pdf>, last access: 29 April 2021.