



Supplement of

Modular Assessment of Rainfall–Runoff Models Toolbox (MARRMoT) v2.1: an object-oriented implementation of 47 established hydrological models for improved speed and readability

Luca Trotter et al.

Correspondence to: Luca Trotter (l.trotter@unimelb.edu.au)

The copyright of individual parts of the supplement might differ from the article licence.

Additional figures




Figure S1: Differences in daily storage between MARRMoT v1.4 and v2.0 of m_07_gr4j_4p_2s. Each grey line is one of 559 catchments, red line indicates their average.




Figure S2: Differences in daily storage between MARRMoT v1.4 and v2.0 of m_14_topmodel_7p_2s. Each grey line is one of 559 catchments, red line indicates their average.




Figure S3: Differences in daily storage between MARRMoT v1.4 and v2.0 of m_21_flexb_9p_3s. Each grey line is one of 559 catchments, red line indicates their average.




Figure S4: Differences in daily storage between MARRMoT v1.4 and v2.0 of m_26_flexi_10p_4s. Each grey line is one of 559 catchments, red line indicates their average.




Figure S5: Differences in daily storage between MARRMoT v1.4 and v2.0 of m_27_tank_12p_4s. Each grey line is one of 559 catchments, red line indicates their average.




Figure S6: Differences in daily storage between MARRMoT v1.4 and v2.0 of m_34_flexis_12p_5s. Each grey line is one of 559 catchments, red line indicates their average.