



INFERNO-peat v1.0.0: a representation of northern high-latitude peat fires in the JULES-INFERNNO global fire model

Katie R. Blackford^{1,2}, Matthew Kasoar^{1,2}, Chantelle Burton³, Eleanor Burke³, Iain Colin Prentice^{1,4}, and Apostolos Voulgarakis^{1,2,5}

¹The Leverhulme Centre for Wildfires, Environment, and Society, Imperial College London, London, SW7 2BX, UK

²Department of Physics, Imperial College London, SW7 2BX, UK

³Met Office Hadley Centre, Exeter, EX1 3PB, UK

⁴Department of Life Sciences, Imperial College London, London, SW7 2BX, UK

⁵School of Chemical and Environmental Engineering, Technical University of Crete, Kounoupidiana, 73100, Greece

Correspondence: Katie R. Blackford (k.blackford19@imperial.ac.uk)

Received: 17 October 2023 – Discussion started: 8 November 2023

Revised: 9 February 2024 – Accepted: 28 February 2024 – Published: 18 April 2024

Abstract. Peat fires in the northern high latitudes have the potential to burn vast amounts of carbon-rich organic soil, releasing large quantities of long-term stored carbon to the atmosphere. Due to anthropogenic activities and climate change, peat fires are increasing in frequency and intensity across the high latitudes. However, at present they are not explicitly included in most fire models. Here we detail the development of INFERNO-peat, the first parameterization of peat fires in the JULES-INFERNNO (Joint UK Land Environment Simulator INTERactive Fire and Emission algoRithm for Natural enviroNments) fire model. INFERNO-peat utilizes knowledge from lab and field-based studies on peat fire ignition and spread to be able to model peat burnt area, burn depth, and carbon emissions, based on data of the moisture content, inorganic content, bulk density, soil temperature, and water table depth of peat. INFERNO-peat improves the representation of burnt area in the high latitudes, with peat fires simulating on average an additional 0.305×10^6 km² of burn area each year, emitting 224.10 Tg of carbon. Compared to Global Fire Emissions Database version 5 (GFED5), INFERNO-peat captures $\sim 20\%$ more burnt area, whereas INFERNO underestimated burning by 50%. Additionally, INFERNO-peat substantially improves the representation of interannual variability in burnt area and subsequent carbon emissions across the high latitudes. The coefficient of variation in carbon emissions is increased from 0.071 in INFERNO to 0.127 in INFERNO-peat, an almost 80% increase. Therefore, explicitly modelling peat fires shows a

substantial improvement in the fire modelling capabilities of JULES-INFERNNO, highlighting the importance of representing peatland systems in fire models.

1 Introduction

Peatlands are a globally important store of carbon, housing approximately one-third of the world's soil carbon despite only covering 3% of the Earth's land surface (Xu et al., 2018; Yu et al., 2010). The high latitudes make up the vast majority of global peatland area, with $\sim 50\%$ occurring in Canada and Russia alone (UNEP, 2022). The northern high latitudes are therefore critical carbon stores containing 415 Pg C (Hugelius et al., 2020a), exerting a net cooling effect on the atmosphere (Frolking and Roulet, 2007). Peatlands are rich in carbon as peat forms under waterlogged anaerobic conditions, which reduces the decomposition rates of vegetation, allowing for the build-up of carbon-rich organic matter within the soil. However, peatlands are being increasingly threatened by both climate change and anthropogenic activity, with 54% of high-latitude peatlands drying over the last 200 years (Zhang et al., 2022). Peatlands are anticipated to continue to degrade with climate change, amplifying carbon loss with the potential to switch peatlands from being sinks to sources of carbon, creating a positive feedback loop (Loisel et al., 2021; Swindles et al., 2019; Hugelius et al., 2020; Tarnocai, 2009; Zhao and Zhuang, 2023). Further-

more, the degradation of peatlands resulting from both humans and climate change is increasing the frequency and extent of wildfires in peatlands (Turetsky et al., 2015; Dadap et al., 2019).

Peat fires are among the largest and most persistent wildfire phenomena on Earth (Rein, 2013). In the northern high latitudes, peat fires largely originate from lightning strikes (Wendler et al., 2011; McCarty et al., 2021; Xu et al., 2022). However, human activities such as timber and energy extraction, prescribed burning, and tourism can also lead to fires (McCarty et al., 2021). Peat fires are a fundamentally different phenomenon to vegetation fires as they burn predominantly by smouldering combustion, which is characterized by slow, low temperature and flameless burning with incomplete combustion (Rein, 2013; Huang and Rein, 2017; Rein, 2015). Smouldering is a volumetric phenomenon that also spreads downwards within the soil (Rein, 2013). Smouldering combustion is heavily influenced by soil properties (Rein, 2015; Archibald et al., 2018), the most important being soil moisture (Rein, 2013, 2015). In general, drier soils with deep water tables facilitate greater and deeper burning with high fuel consumption (Purnomo et al., 2020; Lin et al., 2019; Benscoter et al., 2011; Turetsky et al., 2011a; Che Azmi et al., 2021). However, fires can still be maintained at moisture contents as high as 160 % (Rein, 2013; Hu et al., 2019b; Rein, 2015; Purnomo et al., 2020), albeit reliant on other parameters such as bulk density and inorganic content, indicating the high combustion potential of peatlands. Inorganic content and bulk density also exert an important control on peat fire ignition and spread (Rein, 2013, 2015). Higher inorganic content levels in peat results in the slower spread rate of peat fires (Yang and Chen, 2018; Christensen et al., 2020), whilst increased peat bulk density is associated with increased fire spread (Huang and Rein, 2019). Soil temperature is also important, as a peat fire will continue to spread downward and laterally into the soil, existing and spreading underground for months, until it is too cold to maintain a fire (Lin et al., 2021). On average, peat fires burn 12 cm deep into the soil (Santoso et al., 2019) but have been shown to burn to 100 cm deep in laboratory experiments (Qin et al., 2023).

Due to the large quantities of carbon sequestered in peatlands, peat fires can release vast amounts of carbon, estimated to be roughly equivalent to 15 % of that of anthropogenic emissions (Lasslop et al., 2019; Loisel et al., 2021; Poulter et al., 2006; Rein, 2015). Carbon emissions from peat fires are heavily influenced by the depth of burn, as the deeper a peat fire burns, the larger the pool of carbon that is exposed to combustion (Lin et al., 2021; Huang and Rein, 2017; Che Azmi et al., 2021). Smouldering peat fires also emit a range of gas species, including CO₂, CO, CH₄, and NH₃, alongside a suite of aerosols and particulates (Hu et al., 2019a; Voulgarakis and Field, 2015), while they are also dominant in driving the interannual variability in global fire emissions and their consequent effects on global atmospheric composition (van der Werf et al., 2010; Voulgarakis et al., 2015). Aerosols and

particulates from peat fires result in the degradation of air quality and can lead to haze events (Turetsky et al., 2015; Hu et al., 2018), consequently disrupting transport, tourism, and agriculture (Hu et al., 2018; Heil and Goldammer, 2001). Haze also leads to respiratory and cardiovascular problems (WHO, 2006; Hu et al., 2018), with an estimated 25 000 to 50 000 premature deaths due to Arctic wildfire attributed PM_{2.5} (Silver et al., 2023). For example, in 2010, peat fires surrounding Moscow led to extreme air pollution and 11 000 additional deaths (Konovalov et al., 2011; Shaponshnikov et al., 2014). Therefore, peat fires are of major concern for the climate and air quality.

Peat fires also have widespread impacts on ecosystems through altered ecosystem composition and successional trajectories and changes to moisture and nutrient dynamics including increased evapotranspiration, which may alter the functioning of peatlands and further carbon losses (Kettridge et al., 2015, 2019; Mekonnen et al., 2021). In the high latitudes 50 % of peatlands are affected by permafrost (Hugelius et al., 2020a). Following a fire, permafrost can be exposed to warming, resulting in degradation, thermokarst development, and further carbon losses (Chen et al., 2021; Nitze et al., 2018). Therefore, peat fires have the potential to cause large shifts in ecosystem functioning and escalate carbon emissions from peatlands.

Peatlands are becoming increasingly vulnerable to fires (York et al., 2020). The Arctic is currently warming at twice the rate of the global average (Bruhwiler et al., 2021); this alongside decreased precipitation can lead to earlier snowmelt and increased water deficits, thus increasing peatland vulnerability to fires and burnt area (Talucci et al., 2022). Land use change, drainage, agriculture, and logging are also increasing peatland vulnerability (Rein, 2015; Langner and Siegert, 2009). Coincidentally, lightning frequency has increased substantially in the high latitudes (Veraverbeke et al., 2017), alongside an expansion of human populations and activities into the high latitudes (Bartsch et al., 2021). Increasing lightning and human ignitions, combined with amplified peatland vulnerability to wildfires, is resulting in an increase in fire activity across the high latitudes (McCarty et al., 2021) and risks switching peatlands from fire-resistant systems to fire-prone systems (Turetsky et al., 2015). For example, 2019, 2020, and 2021 saw the largest fire years on record in north-east Siberia, driven by increased summer temperatures, earlier snowmelt, and greater plant water stress (Descals et al., 2022; Scholten et al., 2022).

Climate change is expected to amplify the vulnerability of peatlands to wildfires through rising temperatures, increased frequency and intensity of droughts, and increases in fire weather (Thompson et al., 2019; Descals et al., 2022; Lund et al., 2023). Lightning strikes in the high latitudes are expected to increase by 113 % by 2100 (Chen et al., 2021). As a result, peat fires are expected to increase in frequency and severity (McCarty et al., 2021; Turetsky et al., 2015). Increased fire occurrence and severity leads to greater carbon

emissions from fires and may result in a positive feedback loop with the climate system and potentially a catastrophic loss of carbon from the northern high latitudes (Mack et al., 2011; Chen et al., 2021; Turetsky et al., 2015), potentially resulting in peatlands switching from a carbon sink to source by 2100 (Wilkinson et al., 2023).

Despite the importance of peat fires they are currently not explicitly incorporated into most fire models, meaning that the important climate and carbon feedbacks cannot be accurately assessed. At present, the Community Earth System Model (CESM) is the only model to represent peatland burning through its fire and land surface model CLM-Li (Li et al., 2013). The CESM approach was a major step forward, but it is limited by the fact that it does not consider the effects of soil properties on peat fires (Li et al., 2013). Fire models in general do not completely reproduce observed patterns of burnt area (Jones et al., 2022), in particular in the high latitudes. The absence of peat fires is often highlighted as a limiting factor in a model's ability to reproduce present-day burning (Mangeon et al., 2016; Teixeira et al., 2021). Therefore, at present the capacity of fire models to predict future fire activity is limited (Jones et al., 2022). Peat fire representation in models is also key to accurately representing the northern peatland carbon balance in Earth system models (Wilkinson et al., 2023).

The INteractive Fire and Emission algoRithm for Natural enviroNments (INFERNO) is a reduced-complexity fire model that is part of the Joint UK Land Environment Simulator (JULES) land surface model (Mangeon et al., 2016; Burton et al., 2019). INFERNO estimates plant functional type burnt area, utilizing lightning and population density to calculate ignitions and key variables such as relative humidity, precipitation, soil moisture, temperature, and fuel load to calculate flammability (Mangeon et al., 2016). INFERNO has been shown to accurately diagnose global burnt area compared to observational data from the Global Fire Emissions Database (GFED) (Mangeon et al., 2016) and compares well to other fire models on a global scale (Hantson et al., 2020). However, over the northern high latitudes INFERNO fails to capture a significant amount of burnt area, particularly in Canada, Alaska, and Siberian Russia. INFERNO also underestimates carbon emissions and fails to capture the interannual variability in these emissions (Mangeon et al., 2016). Mangeon et al. (2016) put these underestimates down to the lack of representation of peat fires in INFERNO. We address this gap here by developing a new peat fire parameterization in INFERNO.

2 Model description and developments

INFERNO-peat is a simplified peat fire model, which utilizes the existing JULES-INFERNO framework to add additional burnt area and carbon emissions from peatland burning. At present INFERNO-peat is an offline model run in

Python version 3.8, using outputs from JULES-INFERNO (detailed below). Figure 1 provides an overview of the model. In summary, INFERNO-peat utilizes ignition data from population density and lightning, along with plant functional type (PFT) flammability calculated by INFERNO and PFT fractions from JULES, to estimate the number of potential peat fire ignitions. The likelihood of those ignitions developing into a peat fire is represented by a parameterization of peat combustibility based on key relationships with soil moisture, inorganic content, and bulk density (Frandsen, 1997). We also parameterized the depth of burn in peatlands using critical soil temperature (Lin et al., 2021) and water table depth, with soil and hydrology simulated by JULES. A parameterization of peatland carbon emissions was also implemented using calculated peat burnt area, depth of burn, and the carbon content existing in the peat (Lin et al., 2021). In the subsections below, we present the individual steps of the peat fire simulation in more detail.

2.1 Peat fire ignitions (I_{peat})

The majority of peat fire ignitions results from a pre-existing flaming vegetation fire (Rein, 2013). To account for this, peat fire ignitions are based on the number of flaming vegetation fires in a grid box identified from INFERNO (Eq. 1). Here, the third ignition mode of INFERNO is used unchanged from Mangeon et al. (2016). Total ignitions (I_T) are comprised of human ignitions and suppressions, based on population density, and varying natural ignitions, based on cloud-to-ground lightning strikes. The flammability of each of the 13 PFTs represented in JULES is also calculated using the original equations from Mangeon et al. (2016) in INFERNO. Flammability of each PFT ($\text{Flamp}_{\text{PFT}}$) (0–1) is based on key climatic drivers of fires such as temperature, relative humidity, and precipitation, alongside fuel density and soil moisture (Mangeon et al., 2016). From Eq. (1), the number of peat fires (I_{peat}) in a particular PFT is given by the flammability ($\text{Flamp}_{\text{PFT}}$) multiplied by the gridbox ignition rate (I_T) and the fraction of the gridbox occupied by that PFT (Frac_{PFT}).

$$I_{\text{peat}} = \sum_{\text{PFT}} I_T \cdot \text{Flamp}_{\text{PFT}} \cdot \text{Frac}_{\text{PFT}} \quad (1)$$

2.2 Peat combustibility ($\text{Comb}_{\text{peat}}$)

For each grid box where peat is located, the combustibility of peat is calculated using Eq. (2) (Frandsen, 1997; Purnomo et al., 2020).

$$\text{Comb}_{\text{peat}} = \frac{1}{1 + \exp(-(B_0 + (B_1 \cdot \text{SM}) + (B_2 \cdot \text{IC}) + (B_3 \cdot \rho)))} \quad (2)$$

Combustibility describes the probability of a peat fire igniting and spreading. Peat combustibility depends on the peatland soil moisture (SM) (%), inorganic content (IC) (%), and




Figure 1. Schematic summarizing the new parameterizations introduced as part of INFERNO-peat, the input variables, and their sources.

bulk density (ρ) (kg m^{-3}). Soil moisture is the most important variable affecting the ignition and spread of peat fires (Rein, 2013, 2015). Here, soil moisture and peat combustibility exhibit a reverse sigmoid curve, where the likelihood of peat combusting is high at low SM and low at high SM (Frandsen, 1997; Fig. 2a). Fixed values are utilized for IC and ρ of 9.4 % and 222 kg m^{-3} , respectively (Frandsen, 1997), due to a lack of robust observational datasets of peatland-specific IC and BD on a global scale and to avoid adding additional sources of uncertainty into the model. B values represent constants identified by Frandsen (1997), where $B_0 = -19.8198$, $B_1 = -0.1169$, $B_2 = 1.0414$, and $B_3 = 0.0782$.

2.3 Peat burnt area (BA_{peat})

Calculating burnt area from peat fires uses a similar approach to how PFT burnt areas are calculated in JULES (Mangeon et al., 2016). In INFERNO, average burnt area values for each PFT were heuristically determined (Mangeon et al., 2016). However, here we obtain an average peat burnt area ($\overline{BA_{\text{peat}}}$) from Santoso et al. (2019), who estimated an average peat fire burnt area, depth, and emitted carbon for boreal peat fires based on reported field studies between 1983 and 2015. For use in INFERNO-peat, anomalously large values reported from Santoso et al. (2019) were omitted from the average. Therefore, an average peat burnt area of 381.7 km^2 was used. The peat burnt area is then calculated following Eq. (3), where BA_{peat} (peatland burnt area in km^2) results from combining I_{peat} , $\text{Comb}_{\text{peat}}$, $\overline{BA_{\text{peat}}}$, and grid box peatland fraction ($\text{Frac}_{\text{peat}}$).

$$BA_{\text{peat}} = I_{\text{peat}} \cdot \text{Comb}_{\text{peat}} \cdot \overline{BA_{\text{peat}}} \cdot \text{Frac}_{\text{peat}} \quad (3)$$

2.4 Depth of burn (BD_{peat})

We adapt the scheme used by Lin et al. (2021) to estimate the depth of burn (m) resulting from a peat fire. Here we estimate the critical soil temperature (T_{crit}) ($^{\circ}\text{C}$), which represents the minimum environmental temperature that can sustain a smouldering fire in peatlands and is driven by the moisture content (SM) (%) of the soil (Lin et al., 2021; Eq. 4). Lin et al. (2021) identified a linear relationship between T_{crit} and SM, with increasing SM increasing the T_{crit} , which allows dry peat to burn at extremely low temperatures (Fig. 2b). INFERNO-peat uses outputs of soil temperature from JULES to then locate at what depth within the soil column T_{crit} is reached. We then assume that a peat fire will burn to this depth or the depth of the water table (zw) if that is higher. A maximum depth of burn of 40 cm is used in INFERNO-peat, meaning that no fire can burn deeper than 40 cm into the soil. Field studies conducted across the high latitudes have shown that peat fires burn on average the top 5 to 20 cm of the soil (Turetsky et al., 2011b; Walker et al., 2020; Santoso et al., 2019; Lukenbach et al., 2015; Hokanson et al., 2016; Wilkinson et al., 2018; Davies et al., 2013). Although some lab-based studies have shown the possibility of peat burning up to 1 m into the soil (Qin et al., 2023), only 7.4 % of the 905 sites studied by Walker et al. (2020) showed burning deeper than 20 cm, with the maximum value recorded being 34.2 cm. Therefore, capping burn depth at

40 cm should still capture the variation seen in burning across the high latitudes whilst preventing unrealistically deep burning. A sensitivity analysis using different caps on burn depth showed that 40 cm produced the optimum model output compared to observations (Sect. S2 and Fig. S3 in the Supplement).

$$T_{\text{crit}} = (42 \times \text{SM}) - 28 \quad (4)$$

2.5 Carbon emissions (C_{peat})

Total emitted carbon from a peatland burning is calculated using Eq. (5) and reflects common carbon emission calculations (Lin et al., 2021; Che Azmi et al., 2021; Hu et al., 2018). Total emitted carbon from peat fires (C_{peat}) is obtained from peat burnt area (BA_{peat}), the depth of burn (BD_{peat}), a combustion completeness (CC) value, and the carbon content of the peat in that grid box (C). A combustion completeness of 0.8 was used in this study. On average, data from field studies collated by Walker et al. (2020) showed that on average the proportion of total C combusted that is attributed to below-ground carbon is 0.866. Similarly, smouldering combustion is often cited as having a combustion completeness of less than 0.9 (Wiggins et al., 2021; Urbanski, 2014). A sensitivity analysis was performed using multiple combustion completeness values which showed that using 0.8 for combustion completeness resulted in the best modelled carbon in comparison to GFED 500m belowground burning observations (Sect. S2, Fig. S3).

$$C_{\text{peat}} = \text{BA}_{\text{peat}} \times \text{BD}_{\text{peat}} \times C \times \text{CC} \quad (5)$$

3 Experimental set-up and evaluation

INFERNO-peat is run as an offline model using Python version 3.8, requiring output files from JULES and INFERNO. JULES-INFERNO outputs, for the majority of variables, were obtained from a TRENDY JULES simulation in JULES vn5.4 using the TRIFFID (Top-down Representation of Interactive Foliage and Flora Including Dynamics) dynamic vegetation model. For the peat soil variables (SM, water table depth (z_w), and soil temperature (t_{soil})), an experimental JULES run utilizing new peat module developments (Chadburn et al., 2022) was used which assumes from 50° N all soils are organic, therefore producing a better representation of the northern peatland soil physics and dynamics. Standard JULES operates in a similar manner but instead assumes that all soils are mineral. Within early INFERNO-peat testing this was identified to be causing a systematic bias in the model towards peatlands being drier than they should be and consequently resulting in inflated burnt area estimates. This was resolved when using organic soil moisture (Supplement Sect. S1). HYDE population density data

(Hurt et al., 2011) were used to calculate human ignitions and suppressions. For natural ignitions, we ran INFERNO-peat with two datasets. Firstly, as in the original INFERNO (Mangeon et al., 2016), we used a monthly lightning climatology from LIS-OTD (Lightning Imaging Sensor – Optical Transient Detector) (Christian et al., 2003). However, a lightning climatology does not represent year-to-year variation in lightning and therefore could contribute to inaccuracies in the model. Consequently, we also used the WGLC (Worldwide Lightning Location Network – WWLLN – Global Lightning Climatology and time series) lightning time series produced by WWLLN covering 2010–2020 for comparison (Kaplan and Lau, 2021). Gridded peatland fractions and their respective carbon contents were prescribed to the model from the Northern Peatland Dataset (Hugelius et al., 2020b).

INFERNO-peat was run at N96 resolution (1.25° latitude \times 1.875° longitude) at monthly time steps from 1997 to 2014 for the LIS-OTD lightning run and from 2010–2014 for the WGLC lightning run. Due to the availability of data from the organic soil moisture run, INFERNO-peat could only be run up until 2014.

To evaluate model performance, burnt area data from GFED4s (van der Werf et al., 2017), GFED5 (Chen et al., 2023a, b), and FireCCILT11 (Otón et al., 2021) were used. The ABoVE-FED (Potter et al., 2022) dataset was also used to evaluate burnt area in Alaska and Canada. Multiple observational datasets were used for evaluation due to known deficiencies in the ability of remote-sensing-based products, in particular MODIS, which is used in GFED4s, in being able to detect peat fires in the high latitudes (McCarty et al., 2021). GFED5 was therefore used as the most up-to-date product which, on top of MODIS burned area, utilizes high-resolution observations from Landsat and Sentinel-2, leading to a 61 % increase in burned area globally compared to GFED4s (Chen et al., 2023b). Additionally, FireCCILT11, which is based off Advanced Very-High-Resolution Radiometer (AVHRR) images, was used as a non-MODIS-based comparison (Otón et al., 2021). Spatial correlations using Pearson's R , alongside the RMSE metric, were used to diagnose spatial performance in burnt area. Temporal correlations were also conducted for the entire time series, alongside comparisons of variation metrics such as standard deviation and the coefficient of variation. Carbon emissions estimates were evaluated against total carbon emissions from GFED4s (van der Werf et al., 2017), GFED 500m (van Wees et al., 2022), and the Global Fire Assimilation System (GFAS; Kaiser et al., 2012). Similarly to burned area, multiple products were used in addition to GFED4s. GFED 500m enhances the accuracy of carbon emissions estimations by increasing the spatial resolution to 500 m, as well as differentiating between aboveground and belowground carbon emissions, allowing for enhanced analysis of INFERNO-peat's performance (Van Wees et al., 2022). GFAS, which is based on MODIS fire radiative power (FRP) observations, was used as an alternative to the GFED family of products (Kaiser et




Figure 2. Relationship between soil moisture and (a) peat combustibility and (b) critical temperature.

al., 2012). Temporal correlation and variation metrics were again examined for the carbon time series results. These analyses were also carried out on subregions of the high latitudes (Fig. S4). All datasets used for evaluation were resampled to a N96 grid. A null model (Standard JULES-INFERNO v1.5.4) without peat fires was also used for comparison.

4 Results

4.1 Burnt area


INFERNO-peat results in an overall improvement compared to the original INFERNO in total BA when evaluated against observations (Fig. 3). On average peat fires contribute an additional simulated $0.305 \times 10^6 \text{ km}^2$ of burnt area per year across the high latitudes, bringing the total INFERNO-peat burnt area to within $0.09 \times 10^6 \text{ km}^2$ of the GFED5 observations (Table 1). INFERNO-peat allows us to represent clusters of burning more accurately across the high latitudes compared to INFERNO, especially in western Canada and central and eastern Siberia. However, overestimations are clear in western Russia and eastern Canada, whilst in Alaska we are under-representing the burning occurring. When driving the lightning ignitions in the model with WGLC lightning time series, we capture significantly less burning than when using the LIS-OTD climatology (Fig. 3b, c). Whilst this brings the annual total closer to that seen in FireCCILT11, it results in a large underestimation compared to GFED5 (Table 1). There is a notable area of southern Russia where there is a high degree of burning in all observational datasets but which has minimal burning in INFERNO or either of the INFERNO-peat simulations. According to the land cover types modelled by JULES, this area is dominated by C₃ crops, and the Northern Peatland Dataset indicates minimal peatland coverage (Figs. S5 and S7). Therefore, the underestimations seen in southern Russia are likely a result of INFERNO underestimating cropland burning globally rather than representing region-specific agricultural fire management (Burton et al., 2021).

Over the entire model run (1997 to 2014), there are large improvements in the representation of interannual variability (IAV) in burnt area in INFERNO-peat compared to INFERNO (Fig. 4). We capture significantly higher IAV in INFERNO-peat, with the standard deviation increasing from 0.011 in INFERNO to 0.041 in INFERNO-peat, bringing the model much closer to the magnitude of IAV seen in the observations (Table 2). Furthermore, there is an improvement in the *R* value in INFERNO-peat across all observational datasets, meaning that we are also more accurately capturing the timing of the IAV in burnt area (Table 2). However, when compared to GFED5 there is still a noteworthy underestimation of burnt area IAV in INFERNO-peat, mainly between 2001 and 2011. In particular, the large spikes in burning occurring in 2003 and 2008 are not as pronounced in INFERNO-peat, even though they are captured in a qualitative sense.

Regional features and patterns are evident across the high latitudes. To evaluate this, seven subregions were examined (Fig. S6). Across all three North American subregions INFERNO-peat overestimates burning compared to the observations including the North America only dataset ABoVE-FED (Figs. 5 and S8). Whilst this is only minor in Alaska, the overestimation is particularly pronounced in western Canada, with burnt area in INFERNO-peat around almost 4 times greater than in GFED5. However, in all other subregions the opposite is true, especially when compared to GFED5. In central Russia INFERNO-peat burns on average $0.151 \times 10^6 \text{ km}^2 \text{ yr}^{-1}$, which is significantly lower than the 0.187×10^6 observed in GFED5. Similarly, we also see a large underestimation in burning in eastern Russia when compared to GFED5 and FireCCILT11. Despite this underestimation we do still see an improvement in RMSE in this region, which is not seen in others (Table S2). In accordance with the overall high-latitude results, INFERNO-peat captures more interannual variability in all regions compared to INFERNO (Tables S2 and S3). This increase is a lot more pronounced in western Canada, eastern Canada, central Rus-

Table 1. The 2010 to 2014 average annual burnt area (M km^2) fraction and statistics for the various models and observations.

Model	BA	<i>R</i>			RMSE		
		GFED4S	GFED5	FireCCILT11	GFED4S	GFED5	FireCCILT11
INFERNO	0.215	0.485	0.530	0.486	0.061	0.095	0.068
INFERNO-peat LIS-OTD	0.520	0.414	0.431	0.398	0.098	0.124	0.099
INFERNO-peat WGLC	0.282	0.475	0.510	0.467	0.071	0.099	0.076
GFED4s	0.172		0.820	0.868		0.068	0.028
GFED5	0.429	0.820		0.879	0.068		0.053
FireCCILT11	0.276	0.868	0.879		0.028	0.053	

**Figure 3.** The 2010 to 2014 average annual burnt area fraction for INFERNO (a), INFERNO-peat driven by LIS-OTD climatology (b), INFERNO-peat driven by WGLC time series (c), GFED4s (d), GFED5 (e), and FireCCILT11 (f).

sia, and eastern Russia, which are also the regions where we see the greatest change between the models.

4.2 Carbon emissions

On average, peat fires emit an additional 204.5 Tg of carbon per year in INFERNO-peat (Fig. 6), significantly more than the 103.28 Tg modelled by INFERNO. This brings emissions estimates closer to the 305.35 Tg C in GFED 500m and 248.57 Tg C in GFAS (Table 3). Not only are annual averages brought closer to the observations, but INFERNO-peat

also allows for more accurate representation in the interannual variability in carbon emissions. We also see an improved temporal correlation between INFERNO-peat and the GFAS and GFED 500m observations over this time period when compared to INFERNO.

The GFED 500m product is uniquely useful as it has a differentiation between aboveground and belowground carbon emissions. Belowground emissions come from the burning of organic matter within the soil, which occurs predominantly during peat burning. When compared to GFED 500m, we are representing fire emissions and their sources well.




Figure 4. The annual total burnt area across the high latitudes ($< 50^{\circ}$ N) from 1997 to 2014 compared between the models (indicated by solid lines) and observations (indicated by dashed lines).

Table 2. The 1997 to 2014 average annual burnt area, temporal standard deviation (SD), coefficient of variation (cv), and temporal correlation (R) for models and observations. Correlation coefficients for the WGLC period (2010–2014) are available in Table S1 in the Supplement.

Model	Average annual BA ($\times 10^6$ km 2)	SD	cv	R (GFED4s)	R (GFED5)	R (FireCCILT11)
INFERNO	0.181	0.011	0.058	0.463	0.206	0.370
INFERNO-peat LIS-OTD	0.434	0.041	0.094	0.676	0.323	0.394
INFERNO-peat WGLC	0.238	0.017	0.072			
GFED4s	0.146	0.050	0.340		0.759	0.709
GFED5	0.514	0.155	0.300	0.759		0.692
FireCCILT11	0.267	0.051	0.192	0.709	0.692	

Specifically, Fig. 7 shows the breakdown in above- and belowground emissions, as well as peat vs. non-peat emissions from INFERNO-peat. We can see that INFERNO, which only represents vegetation fires, does an adequate job at capturing the aboveground burning, albeit not capturing the interannual fluctuations in burning. Emissions from peat fires from INFERNO-peat fall relatively in-line with belowground burning in GFED 500m. This indicates that the changes we have implemented are successfully capturing the observed belowground emissions.


The carbon emissions modelled by INFERNO-peat vary greatly between subregions (Figs. 8 and S9, Tables S4, S5). For example, whilst burnt area total is very close to the observations in Alaska (Fig. 5), carbon emissions are underestimated. There is an even more pronounced underestimation in carbon emissions in eastern Russia, where emissions captured by the GFED 500m product are over 6 times greater than the emissions modelled by INFERNO-peat. However,

INFERNO-peat is overestimating carbon emissions by almost double in central Russia compared to GFED 500m and GFAS.

By examining the aboveground and belowground burning reported in the GFED 500m product, we can look at what is potentially driving these differences (Figs. 9 and S10). For example, in Alaska INFERNO is capturing the majority of GFED 500m aboveground emissions, but emissions from the peat model are negligible. This means that the deficit in carbon emissions estimate in Alaska is likely a result of not capturing the belowground burning happening in GFED 500m. Similarly in eastern Russia, peat fires modelled by INFERNO-peat emit on average only 5.127 Tg C yr $^{-1}$, as opposed to the 107.002 Tg of carbon observed from belowground burning in GFED 500m. In contrast in central Russia, carbon emissions are almost double that from peat fires as they are from belowground burning in GFED 500m, meaning that INFERNO-peat is likely over-representing peat burning

Table 3. Time series of annual carbon emissions, standard deviation (SD), coefficient of variation (cv), and temporal correlation R coefficients for models and observations over 2003 to 2014.

Model	Average annual C emissions (Tg)	SD	cv	R (GFED4s)	R (GFED 500m)	R (GFAS)
INFERNO	103.282	7.341	0.071	0.685	0.601	0.670
INFERNO-peat LIS-OTD	307.786	48.663	0.158	0.657	0.768	0.702
INFERNO-peat WGLC	149.731	16.533	0.110			
Peat-only LIS-OTD	204.503	45.587	0.223			
GFED4s	180.192	77.259	0.429		0.858	0.914
GFED 500m – total	305.350	111.106	0.364	0.858		0.912
GFED 500m – aboveground	102.718	29.674	0.289			
GFED 500m – belowground	202.632	86.844	0.429			
GFAS	248.574	121.469	0.489	0.914	0.912	

**Figure 5.** The average annual burnt area (2010–2014) from INFERNO and INFERNO-peat driven by the LIS-OTD and WGLC lightning data compared to the observations in each subregion.

in these regions. In western and eastern Canada, however, peat fire emissions modelled by INFERNO-peat LIS-OTD show only small overestimations compared to GFED 500m belowground burning.

Underestimations in burnt area and carbon emissions seen in eastern Russia and Alaska may be a result of inherent biases within JULES-INFERNO. In these regions, temperatures are low, which results in INFERNO underestimating the flammability of vegetation in these areas, which causes INFERNO-peat to underestimate the number of ignitions in these areas, resulting in underestimations in burnt area and emitted carbon. Furthermore, simulated vegetation bias in the TRIFFID dynamic vegetation model within JULES results in a high proportion of the land surface being covered by non-vegetative surface types, predominately bare soil ($\sim 67\%$ of the land surface north of 60° is classed as non-vegetative) (Fig. S7). Therefore, the number of possible

ignition events in INFERNO-peat is further reduced. However, the opposite may be true in central Russia and western Canada where we see large overestimations in burnt area and carbon emissions. In these regions the most common plant functional types are needleleaf deciduous trees, whereas, in reality, these regions contain a large number of herbaceous wetlands, a PFT which is not currently represented in JULES. This may be contributing to a possible inflation in the number of flaming vegetation fires in these regions, leading to overestimates in burnt area and consequently carbon emissions in INFERNO-peat.

5 Discussion

Through the explicit representation of peat fires in INFERNO-peat, we have improved INFERNO's ability to capture burning in the northern high latitudes and in particular improved simulated estimates of wildfire carbon emissions and their interannual variability. According to INFERNO-peat, peat fires accounted for, on average, 58% of burned area and 68% of carbon emissions north of 50° latitude; therefore peat fires have a large impact on simulated model performance. At present the only other fire model that represents peat burning is CLM-Li (Li et al., 2013). However, the inclusion of peat burning in CLM-Li did not show substantial improvements in the simulation of fire in the high latitudes of North America and eastern Siberia (Li et al., 2013). This was attributed to a wet simulation bias seen in CLM-Li, whereby the latent heat flux was underestimated leading to an inflation in the amount of water held by the land (Li et al., 2013). This is in contrast to our findings, where we see a substantial increase in burnt area, particularly in North America, presumably as a result of using organic soil moisture, allowing us to represent the hydrology of peatlands more accurately. However, in eastern Siberia, whilst INFERNO-peat does improve our estimations of burning, we still see much less burning compared to observations. Li et al. (2013) attributed a similar feature they found to low fuel loads in their




Figure 6. The annual total carbon emissions from fires across the high latitudes ($< 50^{\circ}$ N) from 1997 to 2014 compared between the models (indicated by solid lines) and observations (indicated by dashed lines).




Figure 7. The annual total carbon emissions from fires across the high latitudes ($< 50^{\circ}$ N) from 1997 to 2014 compared between the GFED 500m products above- and belowground burning and INFERNO-peat emissions split to peat fires only and INFERNO only.

model. Similarly, land cover fractions modelled by the TRIFID dynamic vegetation model in JULES show low levels of vegetation in eastern Siberia and in particular show a dominance of bare soil (Fig. S7). In reality, much of these tundra ecosystems in eastern Siberia are dominated by grass, moss, and lichen. At present there is no moss or lichen PFT in JULES, and therefore the amount of burnable area is significantly underestimated. Improvements in the simulation of

high-latitude ecosystems within dynamic vegetation models are vital to improving fire modelling in these regions.

There is significant regional variation in the performance of INFERNO-peat. One notable area of improvement is seen in the Northwest Territories region of Canada, where previously INFERNO has struggled to replicate observed patterns of burning. Simulating fires accurately in this region is of vital importance because they have been shown to have




Figure 8. The average annual carbon emissions (2010–2014) from INFERNO and INFERNO-peat driven by the LIS-OTD and WGLC lightning data compared to the observations in each subregion.




Figure 9. The average annual carbon emissions (1997–2014) from INFERNO and peat fires only from INFERNO-peat driven by the LIS-OTD and WGLC lightning data compared to the aboveground and belowground burning from the GFED 500m product in each subregion.

major climatic effects. For example, Canadian fires in 2013 led to high levels of black carbon deposition over Greenland, which resulted in a lowering of albedo and consequently a warming effect on the climate (Thomas et al., 2017). Similar improvements are also evident in Russia, where large peat fires are common. For example, fires in 2010 burned at least 40 000 ha of peatlands in the Moscow region (Sirin and Medvedeva, 2022). As shown here, peat fires can lead to large carbon emissions; for example during the summer of

1998, peat and boreal forest fires in Russia burnt 11×10^6 ha and emitted 176 Tg of carbon (Kajii et al., 2002), representing a substantial effect on the atmosphere, as well as peatland carbon stores.

However, INFERNO-peat also overestimates peat burning compared to observations in Canada and Fennoscandia. One potential cause of these overestimates may be a result of how humans are represented in INFERNO. At present most global fire models rely on simplistic relationships with population density, like INFERNO, or GDP, which fail to capture the highly complex relationship between humans and fire (Perkins et al., 2022; Teckentrup et al., 2019). Humans use fire as a land management tool around the world, altering fuel loads; fragmenting landscapes; and converting land to agriculture, pasture, or industry (Smith et al., 2022; Perkins et al., 2022; Archibald, 2016). Therefore, improvements are required in how we represent humans within INFERNO, potentially through recent approaches that have accounted for the Human Development Index (Teixeria et al., 2023) or on agent-based modelling (Perkins et al., 2022). Recent analysis has also shown land fragmentation metrics such as road density to exert a strong control on burnt area globally (Haas et al., 2022), and therefore this represents a potential future improvements that could be made to INFERNO.

A lack of interannual variability is a common deficiency across global fire models, with most models considered in the Fire Model Intercomparison Project (FireMIP) failing to simulate the interannual variability in fires (Li et al., 2019). However, through the inclusion of peat burning in INFERNO we see substantial increases in interannual variability in burnt area and carbon emissions in the northern high latitudes. Furthermore, we also see improvements in our ability to capture large fire years, for example in 1998, 2003, and 2012 (Fig. 6). Whilst INFERNO-peat cannot replicate real-life fire events and only relies on an estimate of the likelihood of burning, the simulated spikes in burning are also seen in observational data and coincide with record fire years. For example, until recently, 2003 was the largest fire season on record in Siberia burning 22×10^6 ha of land and emitting at least 72 Tg of CO (Talucci et al., 2022). Not only did the 2003 fire season have a major impact on Siberian populations and ecosystems, but haze and smoke plumes were reported to have reached Japan and the USA (Huang et al., 2009). Similarly in 2012, there were a reported 17 000 wildfires in July and August in Siberia, emitting 48 Tg of CO, with smoke reaching the Pacific Northwest of the USA (Teakles et al., 2017). Consequently, being able to accurately simulate these large-scale fire events and their subsequent emissions is highly important in the assessment of both local and global impacts of wildfires. Furthermore, climate change is anticipated to further increase peat fires, with significant impacts on climate, air quality, and the peatland carbon store (Mack et al., 2011; Chen et al., 2021; Turetsky et al., 2015). Therefore, it is important that fire models represent peat fires in order to better

anticipate future changes in burning across the high latitudes over the coming century.

A major challenge in modelling peat burning stems from an absence of robust observational datasets on peat fires, making model evaluation difficult. MODIS satellite products form the basis of the data of GFED4s, which is the most used observational dataset to evaluate fire model performance. However, MODIS and other satellite-based products likely omit a large number of peat fires, as such fires tend to burn below ground and at low temperatures, making them difficult to detect by remote sensing (McCarty et al., 2021). For example, MODIS was shown to be insufficient at detecting the peat fires that occurred in the Moscow region in 2010 (Sirin and Medvedeva, 2022). Further afield, burnt area estimated from Sentinel-2 over sub-Saharan Africa was estimated to be 80 % larger than MODIS through the improved detection of small fires (Roteta et al., 2019). If that pattern holds true for the high latitudes, then MODIS-based products could be failing to capture a substantial amount of burning and consequently carbon emissions. In this study, we have also utilized data from the ABoVE-FED dataset for analysis of North America, which uses the difference normalized burn ratio calculated from Landsat imagery to enhance MODIS estimates (Potter et al., 2023), which provides a slightly more accurate estimation of burning across Canada and Alaska. However, a similar more finely detailed satellite-based product does not currently exist for Russia, nor is there a ground-based dataset available (McCarty et al., 2021). Therefore, it is challenging to sufficiently evaluate the performance of INFERNO-peat due to an under-representation of peat fires in observational datasets.

Refinements and developments to INFERNO-peat could further improve the model's capabilities to capture peat burning and the associated emissions. For example, through the use of peat-specific emissions factors (Hu et al., 2018), the modelling scope could be extended to include emissions of other species such as CO₂, CH₄, and NO_x. This would allow for further investigation into the air quality impacts of peat fires and in particular the impacts on human health. There is also the possibility of the INFERNO-peat scheme being extended globally, with a particular focus on the tropics. Tropical peat fires release vast quantities of carbon to the atmosphere; for example, the 1997 peat fires in Indonesia released up to 2.57 Gt of carbon, equivalent to 40 % of the global annual carbon emissions from fossil fuel burning at that time (Page et al., 2002). Therefore, representing peatland burning in these regions would represent another substantial step forward in fire modelling capabilities. INFERNO-peat can also be used to model future changes in peatland burning over the coming century under different climate change scenarios, improving our ability to anticipate future fire regimes. It is also anticipated that in the near future the INFERNO-peat logic will be added into JULES-INFERNO as an optional switch. This will allow for broader usage by the JULES community, easier integration of the peat fire functionality with

future JULES developments, and full Earth system simulations accounting for peat fires using the UK Earth System Model (UKESM).

6 Conclusions

The explicit representation of peat fires in INFERNO-peat improves simulated burnt area estimates and has increased our ability to capture the interannual variability in carbon emissions across the northern high latitudes. Results presented here not only have addressed noted deficiencies in the INFERNO fire model (Mangeon et al., 2016), but also highlight the crucial need for representing peat burning in fire models in order to simulate the release of vast amounts of long-term stored carbon to the atmosphere. The high latitudes are warming at twice the rate of the global average (Bruhwiler et al., 2021), and continued climate change is expected to cause peatlands to dry out, lightning strikes to increase, and the frequency and severity of wildfires to escalate (Talucci et al., 2022; McCarty et al., 2021; Turetsky et al., 2015; Chen et al., 2021). Increases in peat fire frequency and severity may amplify carbon loss and create a positive feedback loop on the climate system (Rein, 2013; Hu et al., 2018; Turetsky et al., 2015), ultimately shifting peatlands from sinks to sources of carbon by the end of the century (Swindles et al., 2019; Turetsky et al., 2015; Loisel et al., 2021). Therefore, it is vitally important that fire models include a specific parameterization of peat fires in order to be able to replicate historical and present-day burning more accurately, allowing for more comprehensive assessments on the impacts of fire on the climate system, air quality, and the carbon cycle, both now and in the future.

Code and data availability. The JULES code, used for generating inputs into INFERNO-peat including all soil, vegetation, and fire variables, is freely available from JULES trunk version 5.4 onwards at <https://code.metoffice.gov.uk/trac/jules> (JULES collaboration, 2023). Outputs cannot be made available and can only be obtained from running the JULES code (registration required; last access: 12 October 2023).

Code for INFERNO-peat is available from <https://doi.org/10.5281/zenodo.10007362> (Blackford et al., 2023).

The peatland data used in INFERNO-peat are available from <https://doi.org/10.17043/hugelius-2020-peatland-2> (Hugelius et al., 2020b). Population density data are available from Hurtt et al. (2011). LIS-OTD lightning data are available from <https://doi.org/10.5067/LIS/LIS-OTD/DATA311> (Cecil, 2015). WGLC lightning data are available from <https://doi.org/10.5281/zenodo.6007052> (Kaplan and Lau, 2021).

Supplement. The supplement related to this article is available online at: <https://doi.org/10.5194/gmd-17-3063-2024-supplement>.

Author contributions. KRB developed the parameterization of peat fires in INFERNO and undertook model analysis and evaluation with help and advice from MK, CB, CP, and AV. CB provided the modelled outputs from JULES-INFERNO. EB provided modelled outputs from JULES organic soil moisture runs. KRB drafted the text and figures, with MK, CB, and AV reviewing drafts and contributing to the text.

Competing interests. The contact author has declared that none of the authors has any competing interests.

Disclaimer. Publisher's note: Copernicus Publications remains neutral with regard to jurisdictional claims made in the text, published maps, institutional affiliations, or any other geographical representation in this paper. While Copernicus Publications makes every effort to include appropriate place names, the final responsibility lies with the authors.

Acknowledgements. Katie R. Blackford, Matthew Kasoar, Colin Prentice, and Apostolos Voulgarakis were supported by the Leverhulme Trust through the Leverhulme Centre for Wildfires, Environment and Society, grant number RC-2018-023. Chantelle Burton was funded by the Met Office Climate Science for Service Partnership (CSSP) Brazil project, which is supported by the Department for Science, Innovation and Technology (DSIT). Colin Prentice acknowledges support from the LEMONTREE (Land Ecosystem Models based on New Theory, observations and Experiments) project, funded through the generosity of Eric and Wendy Schmidt by recommendation of the Schmidt Futures programme. Apostolos Voulgarakis has also been supported by the AXA Research Fund (project title "AXA Chair in Wildfires and Climate") and received funding from the Hellenic Foundation for Research and Innovation (HFRI) and the General Secretariat for Research and Technology (GSRT), under grant agreement no. 3453. The authors would like to thank Yang Chen and James Randerson for providing access to the GFED5 data.

Financial support. This research has been supported by the Leverhulme Trust (grant no. RC-2018-023).

Review statement. This paper was edited by Sam Rabin and reviewed by two anonymous referees.

References

- Archibald, S.: Managing the human component of fire regimes: lessons from Africa, *Philos. T. R. Soc. B.*, 371, 20150346, <https://doi.org/10.1098/rstb.2015.0346>, 2016.
- Archibald, S., Lehmann, C. E. R., Belcher, C. M., Bond, W. J., Bradstock, R. A., Daniau, A.-L., Dexter, K. G., Forrester, E. J., Greve, M., He, T., Higgins, S. I., Hoffmann, W. A., Lamon, B. B., McGlenn, D. J., Moncrieff, G. R., Osborne, C. P., Pausas, J. G.,

Price, O., Ripley, B. S., Rogers, B. M., Schwilk, D. W., Simon, M. F., Turetsky, M. R., Van der Werf, G. R., and Zanne, A. E.: Biological and geophysical feedbacks with fire in the Earth system, *Environ. Res. Lett.*, 13, 033003, <https://doi.org/10.1088/1748-9326/aa9ead>, 2018.

Bartsch, A., Pointner, G., Nitze, I., Efimova, A., Jakober, D., Ley, S., Hogstrom, E., Grosse, G., and Schweitzer, P.: Expanding infrastructure and growing anthropogenic impacts along Arctic coasts, *Environ. Res. Lett.*, 16, 115013, <https://doi.org/10.1088/1748-9326/ac3176>, 2021.

Benscoter, B. W., Thompson, D. K., Waddington, J. M., Flannigan, M., Wotton, M., Groot, W. J., and Turetsky, M. R.: Interactive effects of vegetation, soil moisture and bulk density on depth of burning of thick organic soils, *Int. J. Wildland Fire*, 20, 1–12, <https://doi.org/10.1071/WF08183>, 2011.

Blackford, K. R., Kasoar, M., Burton, C., Prentice, C., and Voulgarakis, A.: INFERNO-peat v1.0.0 (v1.0.0), Zenodo [code], <https://doi.org/10.5281/zenodo.10007362>, 2023.

Bruhwieler, L., Parmentier, F.-J. W., Crill, P., Leonard, M., and Palmer, P. I.: The Arctic Carbon Cycle and Its Response to Changing Climate, *Curr. Clim. Change Rep.*, 7, 14–34, <https://doi.org/10.1007/s40641-020-00169-5>, 2021.

Burton, C., Betts, R., Cardoso, M., Feldpausch, T. R., Harper, A., Jones, C. D., Kelley, D. I., Robertson, E., and Wiltshire, A.: Representation of fire, land-use change and vegetation dynamics in the Joint UK Land Environment Simulator vn4.9 (JULES), *Geosci. Model Dev.*, 12, 179–193, <https://doi.org/10.5194/gmd-12-179-2019>, 2019.

Burton, C., Kelley, D. I., Jones, C. D., Betts, R. A., Cardoso, M., and Anderson, L.: South American fires and their impacts on ecosystems increase with continued emissions, *Climate Res. Sustain.*, 1, e8, <https://doi.org/10.1002/cli2.8>, 2021.

Chadburn, S. E., Burke, E. J., Gallego-Sala, A. V., Smith, N. D., Bret-Harte, M. S., Charman, D. J., Drewer, J., Edgar, C. W., Euskirchen, E. S., Fortuniak, K., Gao, Y., Nakhavali, M., Pawlak, W., Schuur, E. A. G., and Westermann, S.: A new approach to simulate peat accumulation, degradation and stability in a global land surface scheme (JULES vn5.8_accumulate_soil) for northern and temperate peatlands, *Geosci. Model Dev.*, 15, 1633–1657, <https://doi.org/10.5194/gmd-15-1633-2022>, 2022.

Che Azmi, N. A., Apandi, N. M., and Rashid, A. S. A.: Carbon emissions from the peat fire problem – a review, *Environ. Sci. Pollut. R.*, 2, 16948–16961, <https://doi.org/10.1007/s11356-021-12886-x>, 2021.

Chen, Y., Romps, D. M., Seeley, J. T., Veraverbeke, S., Riley, W. J., Mekonnen, Z. A., and Randerson, J. T.: Future increases in Arctic lightning and fire risk for permafrost carbon, *Nat. Clim. Change*, 11, 404–410, <https://doi.org/10.1038/s41558-021-01011-y>, 2021.

Chen, Y., Hall, J., van Wees, D., Andela, N., Hantson, S., Giglio, L., van der Werf, G. R., Morton, D. C., and Randerson, J. T.: Global Fire Emissions Database (GFED5) Burned Area (0.1), Zenodo [data set], <https://doi.org/10.5281/zenodo.7668424>, 2023a.

Chen, Y., Hall, J., van Wees, D., Andela, N., Hantson, S., Giglio, L., van der Werf, G. R., Morton, D. C., and Randerson, J. T.: Multi-decadal trends and variability in burned area from the fifth version of the Global Fire Emissions Database (GFED5), *Earth Syst. Sci. Data*, 15, 5227–5259, <https://doi.org/10.5194/essd-15-5227-2023>, 2023b.

- Christensen, E. G., Fernandez-Anex, N., and Rein, G.: Influence of soil moisture conditions on the multidimensional spread of smouldering combustion in shallow layers, *Combust. Flame*, 214, 361–370, <https://doi.org/10.1016/j.combustflame.2019.11.001>, 2020.
- Christian, H. J., Blakeslee, R. J., Boccippio, D. J., Boeck, W. L., Buechler, D. E., Driscoll, K. T., Goodman, S. J., Hall, J. M., Koshak, W. J., Mach, D. M., and Stewart, M. F.: Global frequency and distribution of lightning as observed from space by the Optical Transient Detector, *J. Geophys. Res.-Atmos.*, 108, 4005, <https://doi.org/10.1029/2002jd002347>, 2003.
- Dadap, N. C., Cobb, A. R., Hoyt, A. M., Harvey, C. F., and Konings, A. G.: Satellite soil moisture observations predict burned area in Southeast Asian peatlands, *Environ. Res. Lett.*, 14, 094014, <https://doi.org/10.1088/1748-9326/ab3891>, 2019.
- Davies, M. G., Gray, A., Rein, G., and Legg, C. J.: Peat consumption and carbon loss due to smouldering wildfire in a temperature peatland, *Forest Ecol. Manage.*, 308, 169–177, <https://doi.org/10.1016/j.foreco.2013.07.051>, 2013.
- Descals, A., Gaveau, D. L. A., Verger, A., Sheil, D., Naito, D., and Peñuelas, J.: Unprecedented fire activity above the Arctic Circle linked to rising temperatures, *Science*, 378, 532–537, <https://doi.org/10.1126/science.abn9768>, 2022.
- Frandsen, W. H.: Ignition probability of organic soils, *Can. J. Forest Res.*, 27, 1471–1477, <https://doi.org/10.1139/x97-106>, 1997.
- Frolking, S. and Roulet, N. T.: Holocene radiative forcing impact of northern peatland carbon accumulation and methane emissions, *Glob. Change Biol.*, 13, 1079–1088, <https://doi.org/10.1111/j.1365-2486.2007.01339.x>, 2007.
- Haas, O., Prentice, I. C., and Harrison, S. P.: Global environmental controls on wildfire burnt area, size and intensity, *Environ. Res. Lett.*, 17, 065004, <https://doi.org/10.1088/1748-9326/ac6a69>, 2022.
- Hanson, S., Kelley, D. I., Armeth, A., Harrison, S. P., Archibald, S., Bachelet, D., Forrest, M., Hickler, T., Lasslop, G., Li, F., Manguon, S., Melton, J. R., Nieradzki, L., Rabin, S. S., Prentice, I. C., Sheehan, T., Sitch, S., Teckentrup, L., Voulgarakis, A., and Yue, C.: Quantitative assessment of fire and vegetation properties in simulations with fire-enabled vegetation models from the Fire Model Intercomparison Project, *Geosci. Model Dev.*, 13, 3299–3318, <https://doi.org/10.5194/gmd-13-3299-2020>, 2020.
- Heil, A. and Goldammer, J.: Smoke-haze pollution: a review of the 1997 episode in Southeast Asia, *Reg. Environ. Change*, 2, 24–37, <https://doi.org/10.1007/S101130100021>, 2001.
- Hokanson, K. J., Lukenbach, M. C., Devito, K. J., Kettridge, N., Petrone, R. M., and Waddington, J. M.: Groundwater connectivity controls peat burn severity in the boreal plains, *Ecohydrol.*, 9, 574–584, <https://doi.org/10.1002/eco.1657>, 2016.
- Hu, Y., Fernandez-Anex, N., Smith, T. E. L., and Rein, G.: Review of emissions from smouldering peat fires and their contribution to regional haze episodes, *Int. J. Wildland Fire*, 27, 293–312, <https://doi.org/10.1071/WF17084>, 2018.
- Hu, Y., Christensen, E., Restuccia, F., and Rein, G.: Transient gas and particle emissions from smouldering combustion of peat, *P. Combust. Inst.*, 37, 4035–4042, <https://doi.org/10.1016/j.proci.2018.06.008>, 2019a.
- Hu, Y., Christensen, E. G., Amin, H. M. F., Smith, T. E. L., and Rein, G.: Experimental study of moisture content effects on the transient gas and particle emissions from peat fires, *Combust. Flame*, 209, 408–417, <https://doi.org/10.1016/j.combustflame.2019.07.046>, 2019b.
- Huang, S., Siegert, F., Goldammer, J. G., and Sukhinin, A. I.: Satellite-derived 2003 wildfires in southern Siberia and their potential influence on carbon sequestration, *Int. J. Remote Sens.*, 30, 1479–1492, <https://doi.org/10.1080/01431160802541549>, 2009.
- Huang, X. and Rein, G.: Downward spread of smouldering peat fire: the role of moisture, density, and oxygen supply, *Int. J. Wildland Fire*, 26, 907–918, <https://doi.org/10.1071/WF16198>, 2017.
- Huang, X. and Rein, G.: Upward-and-downward spread of smouldering peat fire, *P. Combust. Inst.*, 37, 4025–4033, <https://doi.org/10.1016/j.proci.2018.05.125>, 2019.
- Hugelius, G., Loisel, J., Chadburn, S., Jackson, R. B., Jones, M., MacDonald, G., Marushchak, M., Olefeldt, D., Packalen, M., Siewert, M. B., Treat, C., Turetsky, M. R., Voigt, C., and Yu, Z.: Large stocks of peatland carbon and nitrogen are vulnerable to permafrost thaw, *P. Natl. Acad. Sci. USA*, 117, 20438–20446, <https://doi.org/10.1073/pnas.1916387117>, 2020a.
- Hugelius, G., Loisel, J., Chadburn, S., Jackson, R. B., Jones, M., MacDonald, G., Marushchak, M., Olefeldt, D., Packalen, M., Siewert, M. B., Treat, C., Turetsky, M. R., Voigt, C., and Yu, Z.: Maps of northern peatland extent, depth, carbon storage and nitrogen storage, Dataset version 2, Bolin Centre Database [data set], <https://doi.org/10.17043/hugelius-2020-peatland-2>, 2020b.
- Hurt, G., Chini, L. P., Frolking, S., Betts, R. A., Feddema, J., Fischer, G., Fisk, G., Hibbard, K., Houghton, R. A., Janetos, A., Jones, C. D., Kinderman, G., Kinoshita, T., Goldewijk, K. K., Riahi, K., Shevliakova, E., Smith, S., Stehfest, E., Thomson, A., Thornton, P., van Vuuren, D. P., and Wang, Y. P.: Harmonization of land-use scenarios for the period 1500–2100: 600 years of global gridded annual land-use transitions, wood harvest, and resulting secondary lands, *Climate Change*, 109, 117–161, <https://doi.org/10.1007/s10584-011-0153-2>, 2011.
- Jones, M. W., Abatzoglou, J. T., Veraverbeke, S., Andela, N., Lasslop, G., Forkel, M., Smith, A. J. P., Burton, C., Betts, R. A., van der Werf, G. R., Sitch, S., Canadell, J. G., Santín, C., Kolden, C., Doerr, S. H., and Le Quére, C.: Global and Regional Trends and Drivers of Fire Under Climate Change, *Rev. Geophys.*, 60, e2020RG000726, <https://doi.org/10.1029/2020RG000726>, 2022.
- JULES collaboration: JULES code, Met Office Science Repository Service [code], <https://code.metoffice.gov.uk/trac/jules>, last access: 12 October 2023.
- Kaiser, J. W., Heil, A., Andreae, M. O., Benedetti, A., Chubarova, N., Jones, L., Morcrette, J.-J., Razinger, M., Schultz, M. G., Suttie, M., and van der Werf, G. R.: Biomass burning emissions estimated with a global fire assimilation system based on observed fire radiative power, *Biogeosciences*, 9, 527–554, <https://doi.org/10.5194/bg-9-527-2012>, 2012.
- Kajii, Y., Kato, S., Streets, D. G., Tsai, N.-Y., Shvidenko, A., Nilsson, S., McCallum, I., Minko, N. P., Abushenko, N., Altyntsev, D., and Khodzer, T. V.: Boreal forest fires in Siberia in 1998: Estimation of area burned and emissions of pollutants by advanced very high resolution radiometer satellite data, *J. Geophys. Res.-Atmos.*, 174, 4745, <https://doi.org/10.1029/2001JD001078>, 2002.

- Kaplan, J. O. and Lau, K. H-K.: The WWLLN Global Lightning Climatology and timeseries (WGLC) v2022.0.0, Zenodo [data set], <https://doi.org/10.5281/zenodo.6007052>, 2021.
- Kettridge, N., Turetsky, M. R., Sherwood, J. H., Thompson, D. K., Miller, C. A., Benscoter, B. W., Flannigan, M. D., Wotton, B. M., and Waddington, J. M.: Moderate drop in water table increases peatland vulnerability to post-fire regime shift, *Sci. Rep.-UK*, 5, 8063, <https://doi.org/10.1038/srep08063>, 2015.
- Kettridge, N., Lukenbach, M. C., Hokanson, K. J., Devito, K. J., Petrone, R. M., Mendoza, C. A., and Waddington, J. M.: Severe wildfire exposes remnant peat carbon stocks to increased post-fire drying, *Sci. Rep.-UK*, 9, 3727, <https://doi.org/10.1038/s41598-019-40033-7>, 2019.
- Konovalov, I. B., Beekmann, M., Kuznetsova, I. N., Yurova, A., and Zvyagintsev, A. M.: Atmospheric impacts of the 2010 Russian wildfires: integrating modelling and measurements of an extreme air pollution episode in the Moscow region, *Atmos. Chem. Phys.*, 11, 10031–10056, <https://doi.org/10.5194/acp-11-10031-2011>, 2011.
- Langner, A. and Siegert, F.: Spatiotemporal fire occurrence in Borneo over a period of 10 years, *Glob. Change Biol.*, 15, 48–62, <https://doi.org/10.1111/j.1365-2486.2008.01828.x>, 2009.
- Lasslop, G., Coppola, A. I., Voulgarakis, A., Yue, C., and Vereverbeke, S.: Influence of Fire on the Carbon Cycle and Climate, *Curr. Clim. Change Rep.*, 5, 112–123, <https://doi.org/10.1007/s40641-019-00128-9>, 2019.
- Li, F., Levis, S., and Ward, D. S.: Quantifying the role of fire in the Earth system – Part 1: Improved global fire modeling in the Community Earth System Model (CESM1), *Biogeosciences*, 10, 2293–2314, <https://doi.org/10.5194/bg-10-2293-2013>, 2013.
- Lin, S., Sun, P., and Huang, X. Can peat fire support a flaming wildlife?, *Int. J. Wildland Fire*, 28, 601–613, <https://doi.org/10.1071/WF19018>, 2019.
- Lin, S., Liu, Y., and Huang, X. Climate-induced Arctic-boreal peatland fire and carbon loss in the 21st century, *Sci. Total Environ.*, 796, 148924, <https://doi.org/10.1016/j.scitotenv.2021.148924>, 2021.
- Loisel, J., Gallego-Sala, A. V., Amesbury, M. J., Magnan, G., Anshari, G., Beilman, D. W., Benavides, J. C., Blewett, J., Camill, P., Charman, D. J., Chawchai, S., Hedgpeth, A., Kleinen, T., Korhola, A., Large, D., Mansilla, C. A., Müller, J., van Bellen, S., West, J. B., Yu, Z., Bubier, J. L., Garneau, M., Moore, T., Sannel, A. B. K., Page, S., Välranta, M., Bechtold, M., Brovkin, V., Cole, L. E. S., Chanton, J. P., Christensen, T. R., Davies, M. A., De Vleeschouwer, F., Finkelstein, S. A., Frohling, S., Galka, M., Gandois, L., Girkin, N., Harris, L. I., Heinemeyer, A., Hoyt, A. M., Jones, M. C., Joos, F., Juutinen, S., Kaiser, K., Lacourse, T., Lamentowicz, M., Larmola, T., Leifeld, J., Lohila, A., Milner, A. M., Minkinen, K., Moss, P., Naafs, B. D. A., Nichols, J., O'Donnell, J., Payne, R., Philben, M., Piilo, S., Quillet, A., Ratnayake, A. S., Roland, T. P., Sjögersten, S., Sonntag, O., Swindles, G. T., Swinnen, W., Talbot, J., Treat, C., Valach, A. C., and Wu, J.: Expert assessment of future vulnerability of the global peatland carbon sink, *Nat. Clim. Change*, 11, 70–77, <https://doi.org/10.1038/s41558-020-00944-0>, 2021.
- Lukenbach, M. C., Hokanson, K. J., Moore, P. A., Devito, K. J., Kettridge, N., Thompson, D. K., Wotton, B. M., Petrone, R. M., and Waddington, J. M.: Hydrological controls on deep burning in a northern forested peatland, *Hydrol. Process.*, 29, 4114–4124, <https://doi.org/10.1002/hyp.10440>, 2015.
- Lund, M. T., Nordling, K., Gjelsvik, A. B., and Samset, B. H.: The influence of variability on fire weather conditions in high latitude regions under present and future global warming, *Environ. Res. Commun.*, 5, 065016, <https://doi.org/10.1088/2515-7620/acdfad>, 2023.
- Mack, M. C., Bret-Harte, M. S., Hollingsworth, T. N., Jandt, R. R., Schuur, E. A. G., Shaver, G. R., and Verbylas, D. L.: Carbon loss from an unprecedented Arctic tundra wildfire, *Nature*, 475, 489–492, <https://doi.org/10.1038/nature10283>, 2011.
- Mangeon, S., Voulgarakis, A., Gilham, R., Harper, A., Sitch, S., and Folberth, G.: INFERNO: a fire and emissions scheme for the UK Met Office's Unified Model, *Geosci. Model Dev.*, 9, 2685–2700, <https://doi.org/10.5194/gmd-9-2685-2016>, 2016.
- McCarty, J. L., Aalto, J., Paunu, V.-V., Arnold, S. R., Eckhardt, S., Klimont, Z., Fain, J. J., Evangeliou, N., Venäläinen, A., Tchepakova, N. M., Parfenova, E. I., Kupiainen, K., Soja, A. J., Huang, L., and Wilson, S.: Reviews and syntheses: Arctic fire regimes and emissions in the 21st century, *Biogeosciences*, 18, 5053–5083, <https://doi.org/10.5194/bg-18-5053-2021>, 2021.
- Mekonnen, Z. A., Riley, W. J., Berner, L. T., Bouskill, N. J., Torn, M. S., Iwahana, G., Breen, A. L., Myers-Smith, I. H., García Criado, M., and Liu, Y.: Arctic tundra shrubification: a review of mechanisms and impacts on ecosystem carbon balance, *Environ. Res. Lett.*, 16, 053001, <https://doi.org/10.1088/1748-9326/abf28b>, 2021.
- Nitze, I., Grosse, G., Jones, B. M., Romanovsky, V. E., and Boike, J.: Remote sensing quantifies widespread abundance of permafrost region disturbances across the Arctic and Subarctic, *Nat. Commun.*, 9, 5423, <https://doi.org/10.1038/s41467-018-07663-3>, 2018.
- Otón, G., Lizundia-Loiola, J., Pettinari, M. L., and Chuvieco, E.: Development of a consistent global long-term burned area product (1982–2018) based on AVHRR-LTDR data, *Int. J. Appl. Earth Obs.*, 103, 102473, <https://doi.org/10.1016/j.jag.2021.102473>, 2021.
- Page, S. E., Siegert, F., Rieley, J. O., Boehm, H.-D. V., Jaya, A., and Limin, S.: The amount of carbon released from peat and forest fires in Indonesia during 1997, *Nature*, 420, 61–65, <https://doi.org/10.1038/nature01131>, 2002.
- Perkins, O., Mateji, S., Erb, K., and Millington, J.: Towards a global behavioural model of anthropogenic fire: the spatiotemporal distribution of land-fire systems, *Socio-Environ. Syst. Model*, 4, 18130, <https://doi.org/10.18174/sesmo.18130>, 2022.
- Potter, S., Cooperdock, S., Veraverbeke, S., Walker, X., Mack, M. C., Goetz, S. J., Baltzer, J., Bourgeau-Chavez, L., Burrell, A., Dieleman, C., French, N., Hantson, S., Hoy, E. E., Jenkins, L., Johnstone, J. F., Kane, E. S., Natali, S. M., Randerson, J. T., Turetsky, M. R., Whitman, E., Wiggins, E., and Rogers, B. M.: Burned area and carbon emissions across northwestern boreal North America from 2001–2019, *Biogeosciences*, 20, 2785–2804, <https://doi.org/10.5194/bg-20-2785-2023>, 2023.
- Poulter, B., Christensen, N. L., and Halpin, P. N.: Carbon emissions from a temperate peat fire and its relevance to interannual variability of trace atmospheric greenhouse gases, *J. Geophys. Res.*, 111, D06301, <https://doi.org/10.1029/2005JD006455>, 2006.
- Purnomo, D. M. J., Bonner, M., Moafi, S., and Rein, G.: Using cellular automata to simulate field-scale flaming and smouldering

- wildfires in tropical peatlands, *Proc. Combust. Inst.*, 38, 5119–5127, <https://doi.org/10.1016/j.proci.2020.08.052>, 2020.
- Qin, Y., Musa, D. N. S., Lin, S., Huang, X.: Deep peat fire persistently smouldering for weeks: a laboratory demonstration, *Int. J. Wildland Fire*, 32, 86–98, <https://doi.org/10.1071/WF22143>, 2023.
- Rein, G.: Smouldering Fires and Natural Fuels, in: *Fire Phenomena and the Earth System: An Interdisciplinary Guide to Fire Science*, edited by: Belcher, C., Wiley and Sons, 15–33, <https://doi.org/10.1002/9781118529539.ch2>, 2013.
- Rein, G.: Smouldering peat megafires: the largest fires on earth, in: *Coal and peat fires: a global perspective, Volume 4: Peat – Geology, Combustion, and Case Studies*, edited by: Stracher, G. B., Prakash, A., and Rein, G., Elsevier Science, 1–11, <https://doi.org/10.1016/C2010-0-68845-6>, 2015.
- Roteta, E., Bastarrika, A., Padilla, M., Storm, T., and Chuvieco, E.: Development of a Sentinel-2 burned area algorithm: Generation of a small fire database for sub-Saharan Africa, *Remote Sens. Environ.*, 222, 1–17, <https://doi.org/10.1016/j.rse.2018.12.011>, 2019.
- Santoso, M. A., Huang, X., Prat-Guitart, N., Christensen, E., and Rein, G.: Smouldering fires and soil, in: *Fire Effects on Soil Properties*, edited by: Pereira, P., Mataix-Solera, J., Ubeda, X., Rein, G., and Ceradà, A., CSIRO Publishing, 203–216, <https://doi.org/10.1071/9781486308149>, 2019.
- Scholten, R., C., Coumou, D., Luo, F., and Veraverbeke, S.: Early snowmelt and polar jet dynamics co-influence recent extreme Siberian fire seasons, *Science*, 378, 1005–1009, <https://doi.org/10.1126/science.abn4419>, 2022.
- Shaposhnikov, D., Revich, B., Bellander, T., Bedada, G. B., Bottai, M., Kharkova, T., Kvasha, E., Lezina, E., Lind, T., Semutnikova, E., and Pershagen, G.: Mortality Related to Air Pollution with the Moscow Heat Wave and Wildfire of 2010, *Epidemiology*, 25, 359–364, <https://doi.org/10.1097/EDE.000000000000090>, 2014.
- Silver, B., Arnold, S., Emmons, L., Reddington, C., and Conibear, L.: Health impacts of wildfire smoke in the Arctic, *EGU General Assembly 2023*, Vienna, Austria, 24–28 Apr 2023, EGU23-9394, <https://doi.org/10.5194/egusphere-egu23-9394>, 2023.
- Sirin, A. and Medvedeva, M.: Remote sensing mapping of peat-fire-burnt areas: Identification among other wildfires, *Remote Sens.*, 14, 194, <https://doi.org/10.3390/rs14010194>, 2022.
- Smith, C., Perkins, O., and Mistry, J.: Global decline in subsistence-oriented and smallholder fire use, *Nat. Sustain.*, 5, 542–551, <https://doi.org/10.1038/s41893-022-00867-y>, 2022.
- Swindles, G. T., Morris, P. J., Mullan, D. J., Payne, R. J., Roland, T. P., Amesbury, M. J., Lamentowicz, M., Turner, T. E., Gallego-Sala, A., Sim, T., Barr, I. D., Blaauw, M., Blundell, A., Chambers, F. M., Charman, D. J., Feurdean, A., Galloway, J. M., Galka, M., Green, S. M., Kajukalo, K., Karofeld, E., Korhola, A., Lamentowicz, L., Langdon, P., Marcisz, K., Mauquoy, D., Mazei, Y. A., McKeown, M. M., Mitchell, E. A. D., Novenko, E., Plunkett, G., Roe, H. M., Schoning, K., Sillasoo, Ü., Tsyganov, A. N., van der Linden, M., Väliranta, M., and Warner, B.: Widespread drying of European peatlands in recent centuries, *Nat. Geosci.*, 12, 922–928, <https://doi.org/10.1038/s41561-019-0462-z>, 2019.
- Talucci, A. C., Loranty, M. M., and Alexander, H. D.: Siberian taiga and tundra fire regimes from 2001–2020, *Environ. Res. Lett.*, 17, 025001, <https://doi.org/10.1088/1748-9326/ac3f07>, 2022.
- Tarnocai, C.: The Impact of Climate Change on Canadian Peatlands. *Can. Water Resour. J.*, 34, 453–466, <https://doi.org/10.4296/cwrj3404453>, 2009.
- Teakles, A. D., So, R., Ainslie, B., Nissen, R., Schiller, C., Vinagarzan, R., McKendry, I., Macdonald, A. M., Jaffe, D. A., Bertram, A. K., Strawbridge, K. B., Leaitch, W. R., Hanna, S., Toom, D., Baik, J., and Huang, L.: Impacts of the July 2012 Siberian fire plume on air quality in the Pacific Northwest, *Atmos. Chem. Phys.*, 17, 2593–2611, <https://doi.org/10.5194/acp-17-2593-2017>, 2017.
- Teckentrup, L., Harrison, S. P., Hantson, S., Heil, A., Melton, J. R., Forrest, M., Li, F., Yue, C., Arneth, A., Hickler, T., Sitch, S., and Lasslop, G.: Response of simulated burned area to historical changes in environmental and anthropogenic factors: a comparison of seven fire models, *Biogeosciences*, 16, 3883–3910, <https://doi.org/10.5194/bg-16-3883-2019>, 2019.
- Teixeira, J. C., Folberth, G. A., O’Connor, F. M., Unger, N., and Voulgarakis, A.: Coupling interactive fire with atmospheric composition and climate in the UK Earth System Model, *Geosci. Model Dev.*, 14, 6515–6539, <https://doi.org/10.5194/gmd-14-6515-2021>, 2021.
- Teixeira, J. C. M., Burton, C., Kelly, D. I., Folberth, G. A., O’Connor, F. M., Betts, R. A., and Voulgarakis, A.: Representing socio-economic factors in the INFERNO global fire model using the Human Development Index, *Biogeosciences Discuss.* [preprint], <https://doi.org/10.5194/bg-2023-136>, in review, 2023.
- Thomas, J. L., Polashenski, C. M., Soja, A. J., Marelle, L., Casey, K. A., Choi, H. D., Raut, J.-C., Wiedinmyer, C., Emmons, L. K., Fast, J. D., Pelon, J., Law, K. S., Flanner, M. G., and Dibb, J. E.: Quantifying black carbon deposition over the Greenland ice sheet from forest fires in Canada, *Geophys. Res. Lett.*, 44, 7965–7974, <https://doi.org/10.1002/2017GL073701>, 2017.
- Thompson, D. K., Simpson, B. N., Whitman, E., Barber, Q. E., and Parisien, M.-A.: Peatland Hydrological Dynamics as A Driver of Landscape Connectivity and Fire Activity in the Boreal Plain of Canada, *Forests*, 10, 534, <https://doi.org/10.3390/f10070534>, 2019.
- Turetsky, M. R., Donahue, W. F., and Benscoter, B. W.: Experimental drying intensifies burning and carbon losses in a northern peatland, *Nat. Commun.*, 2, 514, <https://doi.org/10.1038/ncomms1523>, 2011a.
- Turetsky, M. R., Kane, E. S., Harden, J. W., Ottmar, R. D., Manies, K. L., Hoy, E., and Kasischke, E. S.: Recent acceleration of biomass burning and carbon losses in Alaskan forests and peatlands, *Nat. Geosci.*, 4, 27–31, <https://doi.org/10.1038/NGEO1027>, 2011b.
- Turetsky, M. R., Benscoter, B., Page, S., Rein, G., van der Werf, G. R., and Watts, A.: Global vulnerability of peatlands to fire and carbon loss, *Nat. Geosci.*, 8, 11–14, <https://doi.org/10.1038/ngeo2325>, 2015.
- UNEP: Global Peatlands Assessment – The State of the World’s Peatlands: Evidence for action toward the conservation, restoration, and sustainable management of peatlands. Main Report, Global Peatlands Initiative, United Nations Environment Programme, Nairobi, <https://doi.org/10.59117/20.500.11822/41222>, 2022.
- Urbanski, S.: Wildland fire emissions, carbon, and climate: Emission factors, *Forest Ecol. Manage.*, 217, 51–60, <https://doi.org/10.1016/j.foreco.2013.05.045>, 2014.

- van der Werf, G. R., Randerson, J. T., Giglio, L., Collatz, G. J., Mu, M., Kasibhatla, P. S., Morton, D. C., DeFries, R. S., Jin, Y., and van Leeuwen, T. T.: Global fire emissions and the contribution of deforestation, savanna, forest, agricultural, and peat fires (1997–2009), *Atmos. Chem. Phys.*, 10, 11707–11735, <https://doi.org/10.5194/acp-10-11707-2010>, 2010.
- van der Werf, G. R., Randerson, J. T., Giglio, L., van Leeuwen, T. T., Chen, Y., Rogers, B. M., Mu, M., van Marle, M. J. E., Morton, D. C., Collatz, G. J., Yokelson, R. J., and Kasibhatla, P. S.: Global fire emissions estimates during 1997–2016, *Earth Syst. Sci. Data*, 9, 697–720, <https://doi.org/10.5194/essd-9-697-2017>, 2017.
- van Wees, D., van der Werf, G. R., Randerson, J. T., Rogers, B. M., Chen, Y., Veraverbeke, S., Giglio, L., and Morton, D. C.: Global biomass burning fuel consumption and emissions at 500 m spatial resolution based on the Global Fire Emissions Database (GFED), *Geosci. Model Dev.*, 15, 8411–8437, <https://doi.org/10.5194/gmd-15-8411-2022>, 2022.
- Veraverbeke, S., Rogers, B. M., Goulden, M. L., Jandt, R. R., Miller, C. E., Wiggins, E. B., and Randerson, J. T.: Lightning as a major driver of recent large fire years in North American boreal forests, *Nat. Clim. Change*, 7, 529–534, <https://doi.org/10.1038/NCLIMATE3329>, 2017.
- Voulgarakis, A. and Field, R. D.: Fire Influences on Atmospheric Composition, Air Quality and Climate, *Curr. Pollut. Rep.*, 1, 70–81, <https://doi.org/10.1007/s40726-015-0007-z>, 2015.
- Voulgarakis, A., Marlier, M. E., Faluvegi, G., Shindell, D. T., Tsigaridis, K., and Mangeon, S.: Interannual variability of tropospheric trace gases and aerosols: The role of biomass burning emissions, *J. Geophys. Res.-Atmos.*, 12, 7157–7173, <https://doi.org/10.1002/2014JD022926>, 2015.
- Walker, X. J., Baltzer, J. L., Bourgeau-Chaves, L. L., Day, N. J., de Groot, W. J., Dieleman, C., Hoy, E. E., Johnstone, J. F., Kane, E. S., Parisien, M. A., Potter, S., Rogers, B. M., Turetsky, M. R., Veraverbeke, S., Whitman, E., and Mack, M. C.: ABoVE: Synthesis of Burned and Unburned Forest Site Data, AK and Canada, 1983–2016, Oak Ridge, Tennessee, USA, ORNL DAAC [data set], <https://doi.org/10.3334/ORNLDAAC/1744>, 2020.
- Wendler, G., Conner, J., Moore, B., Shulski, M., and Stuefer, M.: Climatology of Alaskan wildfires with special emphasis on the extreme year of 2004, *Theor. Appl. Climatol.*, 104, 459–472, <https://doi.org/10.1007/s00704-010-0357-9>, 2011.
- Wiggins, E. B., Andrews, A., Sweeney, C., Miller, J. B., Miller, C. E., Veraverbeke, S., Commane, R., Wofsy, S., Henderson, J. M., and Randerson, J. T.: Boreal forest fire CO and CH₄ emission factors derived from tower observations in Alaska during the extreme fire season of 2015, *Atmos. Chem. Phys.*, 21, 8557–8574, <https://doi.org/10.5194/acp-21-8557-2021>, 2021.
- Wilkinson, S. L., Moore, P. A., Flannigan, M. D., Wotton, B. M., Waddington, J. M.: Did enhanced afforestation cause high severity peat burn in the Fort McMurray Horse River wildfire, *Environ. Res. Lett.*, 13, 014018, <https://doi.org/10.1088/1748-9326/aaa136>, 2018.
- Wilkinson, S. L., Andersen, R., Moore, P. A., Davidson, S. J., Granath, G., and Waddington, J. M.: Wildfire and degradation accelerate northern peatland carbon release, *Nat. Clim. Change*, 13, 456–461, <https://doi.org/10.1038/s41558-023-01657-w>, 2023.
- World Health Organisation: Air quality guidelines: global update 2005: particulate matter, ozone, nitrogen dioxide and sulfur dioxide, World Health Organisation, Copenhagen, ISBN 92 890 2192 6, 2006.
- Xu, J., Morris, P. J., Liu, J., and Holden, J.: PEATMAP: Refining estimates of global peatland distribution based on a meta-analysis, *Catena*, 160, 134–140, <https://doi.org/10.1016/j.catena.2017.09.010>, 2018.
- Xu, W., Scholten, R. C., Hessilt, T. D., Liu, Y., and Veraverbeke, S.: Overwintering fires rising in eastern Siberia, *Environ. Res. Lett.*, 17, 045005, <https://doi.org/10.1088/1748-9326/ac59aa>, 2022.
- Yang, J. and Chen, H.: Natural Downward Smouldering of Peat: Effects of Inorganic Content and Piled Bed Height, *Fire Technol.*, 54, 1219–1247, <https://doi.org/10.1007/s10694-018-0737-8>, 2018.
- York, A., Bhatt, U. S., Gargulinski, E., Garbinski, Z., Jain, P., Soja, A., Thoman, R. L., and Ziel, R.: Wildland Fire in High Northern Latitudes, in: Arctic Report Card 2020, edited by: Thoman, R. L., Richter-Menge, J., and Druckenmiller, M. L., <https://doi.org/10.25923/2gef-3964>, 2020.
- Yu, Z., Loisel, J., Brosseau, D. P., Beilman, D. W., and Hunt, S. J.: Global peatland dynamics since the Last Glacial Maximum, *Geophys. Res. Lett.*, 37, L13402, <https://doi.org/10.1029/2010GL043584>, 2010.
- Zhang, H., Väiliranta, M., Swindles, G. T., Aquino-López, M. A., Mullan, D., Tan, N., Amesbury, M., Babeshko, K. V., Bao, K., Bobrov, A., Chernyshov, V., Davies, M. A., Diaconu, A. C., Feurdean, A., Finkelstein, S. A., Garneau, M., Guo, Z., Jones, M. C., Kay, M., Klein, E. S., Lamentowicz, M., Magnan, G., Marcisz, K., Mazei, N., Mazei, Y., Payne, R., Pelletier, N., Piilo, S. R., Pratte, S., Roland, T., Saldaev, D., Shoty, W., Sim, T. G., Sloan, T. J., Słowiński, M., Talbot, J., Taylor, L., Tsyganov, A. N., Wetterich, S., Xing, W., and Zhao, Y.: Recent climate change has driven divergent hydrological shifts in high-latitude peatlands, *Nat. Commun.*, 13, 4959, <https://doi.org/10.1038/s41467-022-32711-4>, 2022.
- Zhao, B. and Zhuang, Q.: Peatlands and their carbon dynamics in northern high latitudes from 1990 to 2300: a process-based biogeochemistry model analysis, *Biogeosciences*, 20, 251–270, <https://doi.org/10.5194/bg-20-251-2023>, 2023.