



Sources of uncertainty in the SPITFIRE global fire model: development of LPJmL-SPITFIRE1.9 and directions for future improvements

Luke Oberhagemann^{1,2}, Maik Billing², Werner von Bloh², Markus Drüke^{2,3}, Matthew Forrest⁴, Simon P. K. Bowring⁵, Jessica Hetzer⁴, Jaime Ribalaygua Batalla⁶, and Kirsten Thonicke²

¹Institute of Environmental Science and Geography, University of Potsdam, Karl-Liebknecht Str. 24/25, Potsdam, Germany

²Potsdam Institute for Climate Impact Research, Telegrafenberg A 31, 14473 Potsdam, Germany

³Deutscher Wetterdienst, Hydrometeorologie, Frankfurter Str. 135, 63067 Offenbach, Germany

⁴Senckenberg Biodiversity and Climate Research Centre, Senckenberganlage 25, 60325 Frankfurt, Germany

⁵Laboratoire des Sciences du Climat et de l'Environnement (LSCE), IPSL-CEA-CNRS-UVSQ, Université Paris-Saclay, Gif-sur-Yvette, France

⁶Meteogrid, Almansa 88, Madrid, Spain

Correspondence: Luke Oberhagemann (oberhagemann@uni-potsdam.de)

Received: 24 June 2024 – Discussion started: 13 August 2024

Revised: 9 December 2024 – Accepted: 2 January 2025 – Published: 27 March 2025

Abstract. Since its development in 2010, the SPITFIRE global fire model has had a substantial impact on the field of fire modelling using dynamic global vegetation models. It includes process-based representations of fire dynamics, including ignitions, fire spread, and fire effects, resulting in a holistic representation of fire on a global scale. Previously, work had been undertaken to understand the strengths and weaknesses of SPITFIRE and similar models by comparing their outputs against remotely sensed data. We seek to augment this work with new validation methods and extend it by completing a thorough review of the theory underlying the SPITFIRE model to better identify and understand sources of modelling uncertainty. We find several points of improvement in the model, the most impactful being an incorrect implementation of the Rothermel fire spread model that results in large positive biases in fire rate of spread and a live grass moisture parametrization that results in unrealistically dry grasses. The combination of these issues leads to excessively large and intense fires, particularly on the dry modelled grasslands. Because of the tall flames present in these intense fires, which can cause substantial damage to tree crowns, these issues bias SPITFIRE toward high tree mortality. We resolve these issues by correcting the implementation of the Rothermel model and implementing a new live grass mois-

ture parametrization, in addition to several other improvements, including a multi-day fire spread algorithm, and evaluate these changes in the European domain. Our model developments allow SPITFIRE to incorporate more realistic live grass moisture content and result in more accurate burnt area on grasslands and reduced tree mortality. This work provides a crucial improvement to the theoretical basis of the SPITFIRE model and a foundation upon which future model improvements may be built. In addition, this work further supports these model developments by highlighting areas in the model where high amounts of uncertainty remain, based on new analysis and existing knowledge about the SPITFIRE model, and by identifying potential means of mitigating them to a greater extent.

1 Introduction

Fire is an important component of the Earth system, influencing global carbon cycles and modifying vegetation (e.g. Archibald et al., 2018; Bowman et al., 2009). Because of these global-scale impacts, fire models based on dynamic global vegetation models (DGVMs) have been developed with the goal of simulating fire on a global scale and in-

corporating vegetation feedbacks in predictions of future fire regimes. One such model, the SPread and InTensity of FIRE (SPITFIRE) global fire model was first introduced by Thonicke et al. (2010). It is designed for use with DGVMs, and models fire processes on coarse temporal and spatial scales, most often at a 0.5° by 0.5° grid resolution, and daily time steps. It has been implemented in several DGVMs since its development for the Lund–Potsdam–Jena (LPJ) model, including the LPJ managed Land model (LPJmL), LPJ-GUESS, ORCHIDEE, and JS-BACH (Schaphoff et al., 2018a; von Bloh et al., 2018a; Lasslop et al., 2014; Lehsten et al., 2009, 2015; Yue et al., 2014, 2015). In addition to these direct implementations of the SPITFIRE model, the derived LPJ-LMfire model was developed in 2013 by Pfeiffer et al. (2013) and the derived model LPX was developed by Prentice et al. (2011). Ward et al. (2018) adapted the crown scorch component of SPITFIRE for a crown fire parametrization in the LM3-FINAL fire model. Detailed information on each model can be found in their respective basis papers, and a summary is available in Rabin et al. (2017). Some key differences between these applications and the original version include updated lightning ignitions (LPX, LPJ-LMfire, and ORCHIDEE), parametrizations for differences in human fire use (LPJ-LMfire), a population density effect on fire duration (JS-BACH), stochastic burning of vegetation patches (LPJ-GUESS), and empirically derived regional scaling of burnt area (ORCHIDEE).

The SPITFIRE model has been used in a number of model application studies, e.g. Wu et al. (2015), comparing LPJmL-SPITFIRE and LPJ-GUESS-SIMFIRE in the European domain; Drüke et al. (2023), examining the impact of fire on Amazon forest regrowth under future climate change using LPJmL-SPITFIRE; Felsberg et al. (2018), examining the impact of the temporal resolution of lightning data on JSBACH-SPITFIRE model results; and Hantson et al. (2015), using JSBACH-SPITFIRE to model anthropogenic influences on global fire size distributions. LPJ-LMfire has also been widely used in studies, e.g. Boulanger et al. (2022), analysing future performance of tree species in Québec, Canada; Chaste et al. (2018), comparing model results to observations in eastern boreal Canada; Emmett et al. (2021) developing a local-scale model based on LPJ-LMfire; and Kaplan et al. (2016), applying LPJ-LMfire to a historical study of Europe during the Last Glacial Maximum. Due to its widespread application in global fire modelling, SPITFIRE and its derived models form the basis of a substantial proportion of the models used for the Fire Model Intercomparison Project (FireMIP), with 4 of the 11 models used for the first phase of the project being SPITFIRE-based (Rabin et al., 2017). The FireMIP ensemble of models, in various configurations, has also been used in a number of studies, e.g. Hantson et al. (2020), analyzing the performance of FireMIP models; Li et al. (2019), using the model ensemble to simulate historical fire emissions; Lasslop et al. (2020), studying the effect of fire on tree cover in the FireMIP models; An-

dela et al. (2017), showing that FireMIP models generally do not represent the decline in burnt area observed in satellite data; Forkel et al. (2019a), comparing burnt area drivers in FireMIP models to a random forest model; and Teckentrup et al. (2019), performing sensitivity analyses on the FireMIP models.

The SPITFIRE model has therefore had a substantial impact on the field of predictive fire modelling on a global scale. Understanding the sources of uncertainty in the model is therefore important for contextualizing work done with the model itself and with ensembles in which the model plays a substantial role. Many of the studies discussed analyze these sources of uncertainty using an a posteriori approach, in which the results of model simulations are interpreted in the context of satellite observations (see in particular Forkel et al., 2019a; Hantson et al., 2020; Andela et al., 2017; Teckentrup et al., 2019). Here, we supplement and extend this work by examining the fundamental components of the SPITFIRE model in the context of the theory on which they are based. By returning to the underlying theory and examining its importance at DGVM scales, we help fill a critical research gap between those examining fire at local scales and those undertaking fire modelling at a global scale.

The sources of uncertainty in the SPITFIRE model can broadly be grouped into four classes: inaccuracies in the implementation of previous research into the model, unrealistic results from equations designed for the model, general modelling simplifications, and uncertainty associated with model inputs. In the first two cases, we analyze the nature and impacts of these sources of uncertainty and provide potential solutions or road maps toward solutions. In the third and fourth cases, we provide a general discussion of the nature of these simplifications to provide a better understanding of use cases for SPITFIRE. We focus in particular on two major issues that we identify in the model: an incorrect implementation of the Rothermel fire spread model and unrealistically low live grass moisture content. This paper is organized into a brief description of the model, the results of several validation tests, an identification of sources of uncertainty in each component of the model, an improved model version in the European domain, and an outlook discussing the current status of SPITFIRE with a road map for future developments. The results shown here also focus on SPITFIRE embedded in the LPJmL DGVM, in particular versions 4 and 5.7, and on the European domain. Because of this, some of the more detailed technical changes apply more specifically to the LPJmL implementation, while broader discussions of the equations underlying SPITFIRE are more general. We have highlighted differences in SPITFIRE implementations within some other DGVMs that alter the impacts of certain parameterizations.

The present work is organized into a brief methods section, giving a general description of the SPITFIRE model, the forcing data used, two new validation methods we have developed, and the model runs at the European scale that we

have performed to test these developments. The subsequent results and discussion section is organized into a discussion of the two major issues that were identified in SPITFIRE, followed by further discussions of individual model components, results of model tests in the European domain, and a section describing the current status of the model. The sections describing specific model developments and potential future developments are summarized in Fig. 1 for readers who are interested in individual developments or model components.

2 Methods

2.1 Model structure

SPITFIRE is a holistic fire model that models fire occurrence, spread, and effects (Thonicke et al., 2010; Schaphoff et al., 2018a; Drüke et al., 2019). The structure of the model is shown in Fig. 2. We provide a brief introduction to the model here; for a detailed description, see the referenced work.

The version of SPITFIRE that forms the basis of this work is that embedded in the LPJmL version 4.0 DGVM, i.e. the most recently published global version of SPITFIRE (Schaphoff et al., 2018a). We use this model version as the starting point of our work because it represents the published status of LPJmL-SPITFIRE before the changes made here. This DGVM divides global vegetation into 11 plant functional types (PFTs). These are divided into three tropical, four temperate, and four boreal types. The three tropical PFTs are TrBE – tropical broadleaved evergreen trees, TrBR – tropical broadleaved raingreen trees, and TrH – tropical herbaceous, i.e. tropical grasses. The four temperate PFTs are TNE – temperate needleleaved evergreen trees, TBE – temperate broadleaved evergreen trees, TBS – temperate broadleaved summergreen trees, and TH – temperate herbaceous, i.e. temperate grasses. The boreal PFTs are BBS – boreal broadleaved summergreen trees, BNS – boreal needleleaved summergreen trees, BNE – boreal needleleaved evergreen trees, and PH – polar herbaceous, i.e. polar grasses. In a given grid cell, each PFT has a dynamically calculated uniform size (in all dimensions including height and diameter), corresponding to an average individual of that PFT, and all PFTs contribute to the same, grid-cell-averaged fuel bed. Note that this description does not apply to other vegetation models in which SPITFIRE is implemented. For example, LPJ-GUESS uses a cohort approach with different age classes per PFT, and accounts for stochasticity by simulating multiple patches (e.g. Lehsten et al., 2009, 2015). The number and types of PFTs vary between DGVMs as well; for example the JSBACH version used in phase 1 of FireMIP has 12 PFTs, including shrubs (Rabin et al., 2017; Lasslop et al., 2014).

For new model developments, we use the most recent version of LPJmL – version 5.7 – to incorporate the most

recent updates (Wirth et al., 2024). The major differences between this version and version 4 are the implementation of the global nitrogen cycle, originally developed for LPJmL version 5.0, a description of which can be found in von Bloh et al. (2018a), and the implementation of a new litter parametrization that includes a new litter layer that decomposes over time and is renewed through vegetation mortality and leaf turnover (Lutz et al., 2019). This component is discussed in further detail in Sect. 3.2.5. The model code for versions 4 and 5 of LPJmL are available from Schaphoff et al. (2018b) and von Bloh et al. (2018b) respectively. Our simulations include human land use, which mainly affects fire in LPJmL through determining the total area of natural land within a grid cell.

As shown in Fig. 2, modelled ignitions are scaled by the fire danger, spread according to the fire spread component, and impact vegetation according to the fuel consumption and mortality components. Component 1 of the SPITFIRE model determines the number of ignitions that may potentially become spreading fires in a grid cell on a given day. These ignitions originate from two sources: (1) human-caused ignitions that are calculated using a function of population density and the tendency of the population of a given grid cell to start fires and (2) lightning-caused ignitions that are based on a stationary dataset. The lightning ignition component of the model assumes that 20 % of lightning strikes are cloud-to-ground strikes and that 4 % of these strikes cause ignitions.

Subsequently, in component 2, the number of ignitions is scaled using a fire danger index (FDI) with a value between 0 and 1. This index captures the proportion of ignitions that successfully become spreading fires. There are three versions of this FDI that have been implemented in different versions of SPITFIRE, shown in grey in Fig. 2. All versions aim to account for the role of moisture and fuel bed composition in determining the likelihood of a fire spreading following a potential ignition event.

The original version of the FDI, described in Thonicke et al. (2010), is a function of a previously existing fire danger index, the Nesterov index. The Nesterov index is calculated cumulatively over days in which there are less than 3 mm of precipitation and is a function of daily maximum temperature and dew point temperature. The weighted-average relative moisture content of the fuels being burnt is then calculated based on an exponential function of the Nesterov index and the geometry of the particles making up the fuel bed. The ratio of this moisture content to the moisture of extinction is used as the FDI.

This original parametrization was modified in Schaphoff et al. (2018a), with the parameters dependent on the surface-area-to-volume ratio replaced by tuning parameters that have separate values for each PFT. Finally, Drüke et al. (2019) introduced an FDI parametrization based on the vapour pressure deficit (VPD) that combines an earlier parametrization by Pechony and Shindell (2009) with PFT-specific tuning parameters. Both updates retain the original, fuel class-




Figure 1. Summary of model changes implemented in LPJmL5.7-SPITFIRE and general areas for potential future development in SPITFIRE that are identified in this work, organized according to the model's structure. The key changes discussed in this text are the repaired implementation of the Rothermel model and the realistic live grass moisture levels (the first two bullet points in the fire spread component). The individual model components are further described in Fig. 2.




Figure 2. Structure of the SPITFIRE model. Grey text indicates alternate versions of the fire danger index.

weighted Nesterov parametrization for the dead fuel moisture content from Thonicke et al. (2010), only applying the changes to the FDI.

The fire spread component, component 3, of SPITFIRE is based on the Rothermel fire spread model (Rothermel, 1972) with modifications by Albini (1976). The fuel loads required for the Rothermel model are provided by the DGVM in which SPITFIRE is embedded, fuel bulk densities are a function of PFT-specific values, and the surface-area-to-volume ratio of the fuels within a fuel class is assumed to be uniform across all PFTs. Live fuel moisture levels are calculated as a function of modelled soil moisture. The impact of slope on fire spread is neglected and the average daily wind speed across the grid cell is used to calculate the rate of spread. Each fire lasts for a duration that is calculated as a function of the FDI. All fires are assumed to be elliptical, with a size given by the rate of spread, the fire duration, and a wind-speed-dependent length-to-breadth ratio on a particular day in a grid cell. The burnt area for a given day is thus calculated as a product of the number of ignitions, the FDI, and the fire size for that grid cell and day (all fires on a given day in the same grid cell have the same input parameters and, therefore, size).

The effects of this burnt area on simulated vegetation, shown in component 4, is 2-fold: a portion of the modelled litter bed and live fuel is consumed due to the fire and modelled trees may undergo mortality due to either cambial damage or crown scorch. Modelled bark thickness, heights, and crown base heights are used in these calculations, generally calculated dynamically by the coupled vegetation model. The probability of mortality due to cambial damage is calculated based on the bark thickness and on modelled fire characteristics. The probability of mortality due to crown scorch is calculated based on the height at which calculated flames have an impact on tree crowns, the modelled crown base and top heights, an assumed cylindrical crown structure, and PFT-specific resistance parameters. The two probabilities of mortality are combined by assuming that they are independent.

The altered fuel beds and vegetation, in turn, impact subsequent simulated fires.

2.2 Model forcing data

The solar radiation, precipitation, temperature, wind, and humidity data required to run the model at the 0.5° scale were obtained from the WFDE5 bias-adjusted ERA5 dataset (Cucchi et al., 2020). For the 0.07° European domain runs we use ERA5 land data regridded to the FirEUrisk grid (Muñoz-Sabater et al., 2021; Chuvieco et al., 2023). Lightning data are derived from the LIS/OTD monthly historical lightning dataset (Christian, 2003), interpolated to a daily time step as described in Thonicke et al. (2010). Population density for these runs was obtained from the HYDE 3.1 dataset (Klein Goldewijk et al., 2010).

2.3 New validation methods

Validation of DGVM-based fire models often involves comparisons of maps and time series of model-computed burnt area to burnt area products from satellite-based datasets. We conduct this comparison for LPJmL-SPITFIRE using the GFED4s dataset (Randerson et al., 2015). In addition, we develop two new methods for validating SPITFIRE results. In the first, we split the global burnt area into grid cells that are dominated by tree PFTs and those that are dominated by grass PFTs, i.e. the foliar projective cover (FPC) of the tree or grass PFTs comprises over half of the cell. In the case of the modelled burnt area we use the modelled FPC for this division, and in the case of the satellite-based data we divide the burnt area using the PFT distributions determined by Forkel et al. (2019b). This allows us to gain additional insight into the modelled results by examining whether the global burnt area in the model arises from similar fire-vegetation dynamics as the observed data.

The total burnt area calculated by the model is the result of several model components, as shown in Fig. 2, and this poses a challenge for determining where errors in the burnt area arise. For example, halving the number of ignitions and

doubling the fire size results in the same final burnt area. To gain additional insight into the fire spread component, we introduce a second new validation method. For this, we add a feature to LPJmL-SPITFIRE that allows it to operate using prescribed fire starts (we use the term fire starts here to distinguish these successful ignitions that result in steady-state fire spread from the total modelled ignitions; i.e. these fire starts are not reduced by the fire danger index, whereas the ignitions are). We use the Global Fire Atlas dataset and start fires in the same grid cell and on the same day as the fires in the Global Fire Atlas (Andela et al., 2019). We then compare the burnt area calculated by aggregating all of the fires in a given grid cell from the Global Fire Atlas to the burnt area calculated by LPJmL4-SPITFIRE using these prescribed fire starts. By doing so we circumvent the considerable uncertainty due to the modelling of ignitions and fire danger to focus on the factors that affect the rate of spread and, therefore, fire size.

2.4 New model version at the European scale

To examine the impact of our model updates, we create a preliminary model version specifically for the European domain. We choose this area as a test case because we aim to restrict the amount of variability the model has to account for on a global scale and due to the involvement of the SPITFIRE model in the FirEURisk project (<https://fireurisk.eu>, last access: 19 March 2025). The new model version uses data available through the FirEURisk project at a 0.07° grid cell resolution, also allowing for less sub-grid variability than the usual 0.5° scale. A full new version of the SPITFIRE model is reserved for further work pending additional testing and operation on a global scale. We designate this updated model version as LPJmL-SPITFIRE1.9, reserving the label LPJmL-SPITFIRE2.0 for a version that has been tested at the global scale.

We compare the results of the new model version with the standard SPITFIRE model. For this comparison we implement both model versions in LPJmL5.7, with its included litter moisture (developed by Lutz et al., 2019, and described in detail in Sect. 3.2.5). To allow for a direct comparison of the fire spread and mortality processes, we apply the new tuning parameters and the multi-day fire spread that we develop in this work to the old model version as well. In all other respects, the new model version contains the new improvements and the old model version does not. We then analyze the differences in burnt area and rate of spread between the two model versions, also using the PFT-split burnt area validation method. Note that due to the lack of a prescribed fire starts input at 0.07° resolution, we do not perform the prescribed ignition validation for this smaller-scale version and reserve such tests for future larger-scale versions in which the preliminary fire duration function we develop in this work can be further updated as well.

3 Results and discussion

3.1 Examination of LPJmL4-SPITFIRE using new validation methods

LPJmL4-SPITFIRE shows a reasonable agreement in terms of annual global burnt area with the satellite-derived dataset GFED4s (Randerson et al., 2015), as shown in Fig. 3c. Spatially, as shown in Fig. 3a and b, there is some disagreement with the validation data, e.g. with India experiencing a substantially too-high burnt area while the burnt area in Australia and at high northern latitudes is underestimated (perceptually uniform colour maps in this paper were created using colorcet; see Kovési, 2015). Other spatial patterns, including the large burnt areas in parts of Africa, are reasonably well captured.

The validation using burnt area split by PFT is shown in Fig. 3d and e. Note that the sum of Fig. 3d and e is not the total burnt area shown in Fig. 3c because many grid cells where there is a non-negligible burnt area contain a substantial proportion of managed land in addition to the natural fraction, resulting in area fractions of trees and grasses that are both less than half of the grid cell. These time series reveal a substantial bias in the manner in which the global burnt area is distributed. Overall, the amount of burnt area in grid cells dominated by trees is substantially too low, while grid cells dominated by grasses show a large positive bias. Therefore, the burnt area, while showing some agreement on a global scale, does not appear to arise from the correct physical mechanisms. We show the equivalent figure for the LPJmL4 version of SPITFIRE incorporated in LPJmL57 in Fig. S1 in the Supplement. Because calibrating this version is outside the scope of the present work, the total burnt area is substantially too high. However, Fig. 3d and e show the same overall results, i.e. too much fire on grass-dominated grid cells and too little on grid cells dominated by trees.

Figures S2–S8 show additional details for this comparison. Figures S2 and S3 show a comparison between the observed and simulated grass PFT distribution and a mapped comparison of the burnt area in grid cells that are dominated by grasses respectively. They illustrate that while there is some disagreement between the reference PFT distribution data and those that are simulated, the grass-dominated grid cells that burn are, generally speaking, in broadly the same geographic regions. The burnt area in individual grass-dominated grid cells is also much higher in LPJmL4-SPITFIRE. Therefore, LPJmL4-SPITFIRE appears to have a propensity for predicting too much fire on grasslands.

Figures S4 and S5 show the same comparisons for tree-dominated grid cells. Figure S4, showing the reference and simulated tree PFT distributions, shows that there is a large reduction in simulated tree PFTs in fire-prone regions. This is further illustrated by the large burnt area in tree-dominated grid cells coming from a much larger area, shown in Fig. S5. Therefore, fire in LPJmL4-SPITFIRE appears to greatly re-




Figure 3. Comparison of LPJmL4-SPITFIRE burnt area with GFED4s validation data. Maps of burnt area in panels (a) and (b) show a general alignment in assigning large portions of the global burnt area, in hectares, to the African continent. However, there are regions of substantial geographic disagreement, particularly in India and Australia. The time series plot in panel (c) shows a reasonable agreement in global annual burnt area between modelled and validation datasets. Comparisons of burnt area in grid cells that are over 50% covered by tree PFTs in panel (d) show a substantially lower modelled burnt area. A comparison of burnt area in grid cells that are over 50% covered by grass PFTs in panel (e) shows a strong excess in modelled burnt area.

duce the amount of simulated tree cover relative to a reference dataset, suggesting excessive tree mortality. Figures S6–S8 show the information in Fig. 3d and e by individual PFTs.

The validation using prescribed fire starts is shown in Fig. 4. The model, in Fig. 4b, calculates a much lower burnt area given the prescribed fire starts than the cumulative burnt area of the respective fire sizes, shown in Fig. 4a. The only minor exception is a region in southern Africa, visible also as a small region of greater burnt area in the model in Fig. 4c. Andela et al. (2019) identify this region as having a high ignition density and small fire sizes (see their Fig. 8). That a high ignition density is required to produce appreciable burnt area in LPJmL4-SPITFIRE suggests that individual fires in the model are quite small and, therefore, a large number of them is required. An equivalent figure with LPJmL57, Fig. S9, shows a similarly strong reduction in burnt area when implementing prescribed fire starts. Therefore, the validation using prescribed fire starts offers additional insight into the model that was not available from the original burnt area maps. Motivated by the discrepancies found using the new validation methods, we thoroughly examine the model structure to identify sources of uncertainty that may contribute to these issues.

3.2 Improvements on errors and uncertainties in SPITFIRE

Two substantial errors in SPITFIRE are the incorrect weighting of parameters in the rate-of-spread calculation based on Rothermel (1972) that results in unrealistically severe fires that spread too rapidly and unrealistically low modelled live grass moisture levels. In combination with the incorrect weighting factors, this can cause a strong bias towards simulated fires on grasslands as opposed to forests. In some cases, e.g. due to model tuning, other parts of the model can compensate for these errors somewhat, but this balancing of errors is often insufficient to overcome the biases introduced.

3.2.1 Errors in the implementation of the Rothermel model

As described in Sect. 2.1, the rate-of-spread calculation, which implements the widely used Rothermel model, is a core component of SPITFIRE. We discovered several logical discrepancies between the Rothermel model, given in the original publication by Rothermel (1972) and updated by Albin (1976), and its implementation in SPITFIRE. These are described in detail in Appendix A. The main discrepancies arise from the manner in which different fuel components, separated by size and by dead or living status, are treated when they are combined in the model. Rothermel (1972) and




Figure 4. Comparison of burnt area per grid cell from fires in the Global Fire Atlas to the burnt area per grid cell when starting the same fires in the LPJmL4-SPITFIRE model (panels **a** and **b** respectively). Given the same fire starts LPJmL4-SPITFIRE produces a substantially lower burnt area across the vast majority of regions. The difference map in panel (**c**), showing the modelled burnt area minus the validation data, further illustrates this.

Albini (1976) introduced weighting schemes based on the contribution of each component to the overall surface area of the fuel bed, and dead and live fuels were treated separately. Conversely, SPITFIRE uses a weighting scheme based on contributions to the overall mass of the fuel bed or, in the case of fuel loads, neglects the weighting factors entirely, allowing for values up to 3 times the desired amount. The fuel classes are then combined into a version of the Rothermel model designed for uniform fuels, rather than the non-uniform fuels present; therefore, living and dead fuels are not treated separately, despite their substantially different characteristics.

To analyze the impact that these errors in the application of the Rothermel equation have on SPITFIRE model results, we compare the output of the Rothermel equation with and without errors. For this purpose, we implement the rate-of-spread calculations from SPITFIRE independently from LPJmL (“offline”) using MATLAB (The MathWorks Inc., 2021) and develop an additional code that correctly implements the Rothermel equation for non-uniform fuel beds, which we verify by reproducing several plots in Scott and Burgan (2005). We take the intermediate step of implementing this in MATLAB as it allows us to examine the behaviour of the SPITFIRE implementation of the Rothermel model in a much more efficient way than conducting large amounts of model runs with specified inputs, an application for which the SPITFIRE code was not set up. This MATLAB implementation also allows us to further verify our understanding of the Rothermel model and to produce results for specific fuel and wind speed configurations that we can use for model verification of the new SPITFIRE version, described in further detail later in this section.

To compare the two approaches, we focus on the rate of spread, as this is the main output of the Rothermel model; the fire size, as the sum of fire sizes makes up the burnt area, an important output of SPITFIRE; and the scorch height, the distance from the ground that flames have an effect on the vegetation, a major factor in determining vegetation mortality. In SPITFIRE, the probability of mortality due to cambial damage is set to the fraction of a tree crown for a given PFT that is below this scorch height. For further details, see Sect. 3.2.6. To compare the fire size between the Rothermel and SPITFIRE approaches, we use the elliptical fire spread approach from SPITFIRE, and to compare the scorch height we calculate the fireline intensity using the approach in Albini (1976). This approach is slightly different from the fuel-consumption-based approach used in SPITFIRE. However, we adopt it here for the sake of a more direct comparison. The equation for scorch height, in metres, is given by

$$SH = F \times I_B^{2/3}, \quad (1)$$

where I_B is the fireline intensity, in kW m^{-1} , and F is a PFT-specific scaling parameter.

We conduct this comparison for three Scott and Burgan fuel models that capture different balances between live and dead fuels (Scott and Burgan, 2005). The Scott and Burgan fuel models are a set of parameters that describe various types of fuel beds in a manner that can be input into the Rothermel equation. The three we use in this work are TL3, moderate load conifer litter, which includes only dead fuels; TU2, moderate load humid climate timber-shrub, which includes dead and living fuels; and GR6, moderate load humid climate grass, which is dominated by live fuels. The




Figure 5. Comparison between outputs of the SPITFIRE implementation of the Rothermel equation and the correct implementation created by Rothermel (1972) and Albini (1976). Comparisons are done for the TU2 and TL3 fuel models, moderate load humid climate timber-shrub, and moderate load conifer litter of Scott and Burgan (2005). Differently coloured lines indicate different dead fuel moisture content, following the moisture content scenarios of Scott and Burgan (2005). For all plots the herbaceous live fuel moisture content is set to 60 % and the woody live fuel moisture is set to 90 %.

choice of these fuel models was also motivated by the work of Aragonese et al. (2022), who found that these are the most widespread fuel models across Europe for the timber litter, timber understory, and grass fuel model types. Following the plots in Scott and Burgan (2005), we set the live fuel moisture levels to 60 % for the herbaceous fuels and 90 % for the woody fuels. For dead fuel moisture content, we reproduce the scenarios in Scott and Burgan (2005), where the very low fuel moisture scenario has moisture levels of 3 %, 4 %, and 5 % for the 1, 10, and 100 h dead fuel classes, and each subsequent scenario increases these moisture content by 3 percentage points.

The comparison between the SPITFIRE and Albini (1976) implementations of the Rothermel equation for fuel models TL3 and TU2 are shown in Fig. 5. Figure 5a and b show the impact of the errors on the difference between the SPITFIRE rate of spread and the correct rate of spread, with the SPITFIRE rate of spread being generally higher at low wind speeds and lower at higher wind speeds. The effect of this on fire size is shown in Fig. 5c and d. Because the rate-of-spread difference at low wind speeds is very large compared to the rate of spread, there is a very high peak, up to 2000 % for TL3

and 900 % for TU2, at lower wind speeds in terms of the percent difference ($\frac{\text{SPITFIRE-correct}}{\text{correct}} \times 100 \%$) between the SPITFIRE implementation and the correct one. This difference becomes slightly negative at high wind speeds. While the most extreme difference is only observed at low wind speeds, this may still pose a substantial problem for the model, as the daily and grid-cell-averaged wind speeds that are used as inputs are often in this lower range (the impact on actual model results can also be seen in Sect. 3.3).

This impact of the errors in the SPITFIRE Rothermel implementation on the rate of spread and sub-components of the Rothermel model also cause a substantial bias in the scorch height. As shown in Fig. 5e, in the TL3 model this reaches values up to 1900 % and, in contrast to the impact on fire size, remains high for all values of wind speed shown, with a minimum percent difference for the wind speeds we analyze of about 400 %. This same pattern, with somewhat lower biases, is also visible for the TU2 model in Fig. 5f. To illustrate this more tangibly we have calculated the difference between the scorch heights by setting the F factor in Eq. (1) to 0.1. This value is within 0.01 of the values for all PFTs aside from tropical broadleaved evergreen trees, trop-

ical broadleaved raingreen trees, and temperate broadleaved evergreen trees, which have values of 0.149, 0.061, and 0.371 respectively. As shown in Fig. 5g and h, the difference in scorch height rises rapidly, reaching several metres at wind speeds below 2 km h^{-1} for both of the fuel models shown. This strong bias towards high scorch heights is likely to lead to excessive tree mortality, as further shown in the comparative model runs in Sect. 3.3.

In addition to these upward biases, there is an additional dynamic that can be seen by comparing the fuel models in Fig. 5. Namely, the upward bias on fire size decreases with increasing presence of live fuel. TL3, which contains no live fuels, has the highest biases and TU2, which contains some live fuels has lower biases. This is to the extent that the mostly live GR6 fuel model, shown in Fig. S10, permits no fire spread at all. Therefore, realistic live grass moisture levels should often lead to complete fire extinction on grasslands in SPITFIRE. The reason why there is still fire on grasslands in LPJmL4-SPITFIRE despite this extreme damping is due to extremely low modelled live grass moisture levels, which we discuss in the subsequent section.

To rectify these issues, we have rewritten the implementation of the Rothermel equation in the SPITFIRE model to ensure that the SPITFIRE rate-of-spread calculations match those in Albini (1976). Due to the large amount of space that a detailed description of the Rothermel model would require, in Appendix A we have included only the equations from the Rothermel model relevant to the discussion above. For a full overview of the model we refer readers to Andrews (2018).

We have verified the corrected implementation by reproducing selected results in Scott and Burgan (2005). Specifically, we tested the model with several runs for each of the TL3, TU2, and GR6 models, conducted using both the MATLAB implementation of the Rothermel model that was, in turn, tested against Scott and Burgan (2005) and the new, corrected implementation of the Rothermel model in the SPITFIRE code. In all cases, we use the low dead fuel moisture scenario, with a live herbaceous fuel moisture of 60 %, a live woody fuel moisture of 90 %, and the wind speed limit from Andrews et al. (2013). The full results of this comparison can be found in Table S1 in the Supplement. The greatest difference in terms of rate of spread that we find is 2 % and the greatest difference in terms of Fireline intensity is 3 %. These differences occur in the TL3 fuel model at a wind speed of 300 m min^{-1} , which produces a relatively low intensity and rate of spread compared to the other fuel models tested. These differences amount to only 0.04 m min^{-1} in the rate of spread and 2.67 kW m^{-1} in the fireline intensity. These small differences can most likely be attributed to rounding errors in the model code. The updated implementation of the Rothermel model in SPITFIRE therefore performs accurately. For further technical details on the implementation of this parametrization in the model, please consult the model code provided with this article, and we also provide the MATLAB implementation of the Rothermel model.

3.2.2 Live fuel moisture parametrization

LPJmL-SPITFIRE only calculates fire spread in live herbaceous fuels; this section therefore focuses on the live grass moisture parametrization contained in the model.

Sources of error

As alluded to in the previous section and shown in Fig. S10, the manner in which the live and dead fuel moisture levels are combined in SPITFIRE leads to a combined M_f that is often quite high. This is to the extent that it can be above the maximum combined moisture of extinction used in SPITFIRE of 30 %, on an oven-dry mass basis, when calculated for realistic live fuel moisture levels and fuel beds that contain a large live fuel component (note that in NFDRS 2016, described in Andrews, 2018, and Scott and Burgan, 2005, 30 % is the minimum allowed live fuel moisture). However, as shown in Fig. 3, despite this tendency of the parametrization to produce high combined fuel moisture levels when given realistic live grass moisture levels, SPITFIRE simulates an excessive amount of fire in grasslands. To examine this apparent contradiction, we implement a live grass moisture output in LPJmL4-SPITFIRE and subsequent versions. We find that the live grass moisture is unrealistically low, often to an extent that it does not comport with the conditions required for grasses to survive.

Live grass moisture levels above 100 % on a dry mass basis are often observed in the literature (e.g. Mendiguren et al., 2015; Brown et al., 2022; Yebra et al., 2019). Because these values occur frequently and are a natural part of the dynamics of grass phenology, a successful live grass moisture parametrization must be able to reproduce this. By its construction, the SPITFIRE live grass moisture parametrization cannot exceed 100 %, as visible in Fig. 6a. Further, there are many regions, particularly on the Iberian Peninsula where the live grass moisture *on average* is below 10 %. Specifically, the live grass moisture in SPITFIRE is given by the following (Equation B2 in Thonicke et al., 2010):

$$M_{f,lg} = \max\left(0, \frac{10}{9}M_{s,1} - \frac{1}{9}\right), \quad (2)$$

where $M_{s,1}$ is the moisture content of the top soil layer relative to saturation (note that for consistency in our variables we adopt the variable naming from Andrews, 2018, rather than those of Thonicke et al., 2010). The use of a soil moisture level that is relative to its saturation point in Eq. (2), which has a maximum value of 100 % by definition, results in a maximum live grass moisture of 100 %. In addition, as shown in Fig. S11, the simulated live grass moisture is even lower when considering only months in which modelled fire occurs, illustrating the extremely low live grass moisture values that are required to permit fire spread in SPITFIRE.

Figure 6b and c show the mean live grass moisture during the three winter months of December to February and




Figure 6. Original and updated live grass moisture parametrizations in LPJmL5.7-SPITFIRE at 0.07° resolution in the European domain. Panels (a) through (c) show the original SPITFIRE parametrization from Thonicke et al. (2010), in percent of liquid mass to oven-dry mass. Panels (d) through (f) show a parametrization based on a growing season index for the live fuel moisture following NFDRS 2016. Panels (g) through (i) show a newly developed live grass moisture parametrization based on LPJmL phenology. The columns show, from left to right, the mean live grass moisture from 2003 to 2016, the mean moisture content during the winter months of December to February, and the mean moisture content during the summer months of June to August, during the same time period. The two new live grass moisture parametrizations show general agreement on average and during the months in which there are modelled fires. Note that the live grass moisture applies only to the component of the grasses that is not cured. Therefore, depending on the extent of curing and the dead fuel moisture, the overall moisture content of the grasses may be substantially lower.

the three summer months of June to August from 2003 to 2016. Because the live grass moisture is entirely dependent on the soil moisture in the current parametrization, dormancy of grasses during the winter has no impact on their moisture content, and the live grass moisture during the winter, (Fig. 6b) is, incorrectly, higher than in the summer (Fig. 6c) for most of the temperate and northern regions in Europe (see e.g. Bristiel et al., 2018; Keep et al., 2021; Sjöström and Granström, 2023). In the LPJ-GUESS implementation of SPITFIRE, the issue of these seasonality dynamics was improved upon by treating phenologically inactive grasses as dead fuel, assigning to them the dead fuel moisture.

Finally, the live grass moisture of extinction, M_x , of 20 % used in SPITFIRE is substantially lower than many realistic live grass fuel moisture levels. In Albin (1976), for example, the live fuel moisture of extinction is set to, at minimum, the dead fuel moisture of extinction (which in SPITFIRE is 30 %). Therefore, the value of 20 % is not supported by previous literature. In the subsequent section we introduce new parametrizations.

Improvements to the treatment of live grass moisture in SPITFIRE

The issue of high live grass moisture levels resulting in excessive fire extinction is solved through the corrected weighting scheme in the implementation of the Rothermel equation. Specifically, Eq. (A11) is replaced with a separate treatment of live and dead fuel moisture levels, allowing for fire at higher live grass moisture content. In addition, we have implemented the live fuel moisture of extinction parametrization from Albin (1976) to replace the previous, excessively low, live grass moisture of extinction. These corrections now allow for more accurate, higher, live grass moisture levels to be implemented into the model.

In addition to the more accurate live grass moisture levels described below, we introduce a frequently used curing function to account for transitions of grasses between dormant and active states. This curing function, originally developed for the 1978 version of the US National Fire Danger Rating system and published in Burgan (1979), and described in detail in Scott and Burgan (2005), transfers grass fuel loads from the living fuel category to an additional dead fuel category depending on its moisture content. Grasses that contain

over 120 % water relative to their oven-dry mass are considered fully green and are fully treated as live fuels, and grasses that contain less than 30 % water are considered fully cured and their load is transferred entirely to a dead fuel class. In between these endpoints, the proportion of fuel transferred is a linear function of the live grass moisture. The portion of the live grasses transferred to a dead fuel class is given the same moisture content as the finest (1 h) dead fuel class and the moisture of extinction of the dead fuels.

This treatment of live grass moisture can represent conditions in which grasses are dormant in winter, and therefore have moisture content that is passively determined by weather conditions, followed by a green-up period in spring. This green-up period is an important factor for fire behaviour in many grass-dominated regions because the onset of warmer temperatures does not immediately translate to green grasses, resulting in a well-aerated, dry, fine fuel-dominated litter bed that is prone to high rates of fire spread (see, e.g. Sjöström and Granström, 2023; Burgan, 1979). Note that in the subsequent discussions of live grass moisture parametrizations and in Fig. 6, the live grass moisture refers only to the moisture content of the portion of the grasses that is not cured. Therefore, in regions, such as temperate Europe in winter, with a live grass moisture content of 30 %, i.e. fully cured, the moisture content of the fuel is purely determined by the dead fuel moisture, and the live grass moisture is, therefore, not representative of the overall fuel moisture. Similar considerations apply when there are partially cured fuel beds in extremely dry regions. As an extreme example, a live grass moisture of 50 % on a day where the fine dead fuels are completely dry (this may occur due to the much faster response of dead fine fuels to local weather conditions compared to live grasses) corresponds to an overall moisture content of the cured and not cured grass components of 11 %. We include these examples to help clarify the subsequent results. However, it should be noted that because of the separate treatment of dead and live fuels, this overall moisture content is not directly used in the fire spread calculations, and it is possible for the dead fuel component to burn while the live fuels do not ignite.

To calculate these new, more accurate live grass moisture levels, we introduce two updated approaches. The first of these is a simple adoption of the herbaceous live fuel moisture parametrization from NFDRS 2016, the US National Fire Danger Rating System (described in Andrews, 2018). Because this parametrization is dependent only on weather data and not the output of the DGVM to which SPITFIRE is coupled, it can be easily ported to models other than LPJmL and is not affected by issues in modelled DGVM phenology. This parametrization is based on the growing season index (GSI), developed by Jolly et al. (2005), and it calculates a vegetation phenological status by combining three functions: one of temperature, one of photoperiod, and one of vapour pressure deficit. Each of these functions has a range of 0 to 1, and these functions are multiplied together so that each

individual equation can limit the phenological status of vegetation. This combination is shown in Eq. (3):

$$iGSI = iVPD \times iphoto \times itemp. \quad (3)$$

Here, *iGSI* refers to the daily growing season index. A 21 d running average of the *iGSI* is calculated to arrive at the GSI and this is converted into a live grass moisture parametrization by setting a threshold value of 0.5, below which the grass is dormant, and by using a linear relationship with endpoints of 30 % live grass moisture at a GSI of 0.5 and 250 % live grass moisture at a GSI of 1. As noted in Krueger et al. (2022), this parametrization, although in use in the US, has not yet been thoroughly tested. Therefore, we also introduce a new live grass moisture parametrization that can be used for European runs of the SPITFIRE model.

As stated in Krueger et al. (2022), and further supported by other work, including Brown et al. (2022), soil moisture plays an important role in shaping fire dynamics. One effect, of particular relevance here, is that the soil moisture is an important parameter in determining herbaceous live fuel moisture. The GSI, in essence, includes this effect by proxy, via the VPD function. The LPJmL DGVM has existing phenology functions that act in a similar manner to the GSI but use the soil moisture calculated in the model rather than the VPD (an approach that was also suggested as a potential adjustment for modelled phenology by Jolly et al., 2005). We develop a new live grass moisture parametrization by making use of the phenology functions of the grass PFTs in LPJmL, giving each function a green-up threshold, below which the grass is considered dormant, and fitting a linear function of the grass phenology to live grass moisture levels as reported by Forkel et al. (2023). These functions are shown in Fig. 7, and the equation for live grass moisture is

$$M_{f,lg} = k_{lg} \times phen, \quad (4)$$

subject to the condition that $30 \leq M_{f,lg} \leq 250$, which we adopt from the GSI approach. Here, k_{lg} is an empirically derived scaling factor and *phen* is the phenological status of the grass as determined in LPJmL (a description of how phenology is calculated in LPJmL can be found in Forkel et al., 2014). This phenological status is a function of mean daily air temperature, total daily shortwave radiation, and modelled daily soil water content. The scaling factor k_{lg} is determined by defining a point in the empirical data, i.e. a green-up threshold, where live grass moisture begins to increase with phenology.

To ensure that the observed live grass moisture levels used to arrive at this parametrization are not overly impacted by other live fuels, we select only grid cells that contain over 50 % cover by a specific grass PFT, and over 75 % natural vegetation, to fit these functions. This also has the benefit of focusing on grid cells in which the Forkel et al. (2023) dataset has been shown to be most accurate, i.e. non-forested areas. The shape of these derived functions, as seen in Fig. 7, agrees




Figure 7. New PFT-specific live grass moisture functions in LPJmL-SPITFIRE. Here, phenology refers to the numerical phenological status of the grasses, i.e. how active they are, as calculated in LPJmL using the parametrization from Forkel et al. (2014). Points indicate measured values.

well with data for temperate grasses, which is the herbaceous vegetation type present in the majority of Europe in the model (R^2 of 0.663). The function for polar grasses shows less strong agreement, with an R^2 of 0.29, perhaps due to the observed live fuel moisture being larger than 30 % for many points before the green-up threshold. Future parametrizations may experiment with different live grass moisture content at the beginning of green-up. In general, however, the observed data agree well with the conceptual framework for GSI-based live grass moisture from NFDRS 2016 that the live grass moisture increases linearly with increasing phenological status after a green-up threshold is reached.

The combined live grass moisture for a grid cell is then calculated using a mass-weighted average of the moisture content for each herbaceous PFT. For the current work, we limit this new parametrization to the European domain, reserving a

global parametrization for future work. The live grass moisture levels produced when these parametrizations are integrated into the vegetation model are shown in the second and third rows, Fig. 6d through i. Both parametrizations produce live grass moisture values that reflect the seasonal cycle of live grass moisture content for temperate and boreal regions in Europe in which the grass is dormant in winter and active in summer. The LPJmL phenology-based live grass moisture shows a lower minimum in this cycle, with generally drier grasses in winter than the GSI-based parametrization. The GSI-based parametrization shows a slight divide between eastern and western Europe in live grass moisture that is not present in the phenology-based LFMC. This may be due to differences in the VPD included in the GSI parametrization that do not carry over into the soil moisture and are therefore not present in the LPJmL phenology-based parametrization

(for this reason this divide is also not seen in the old live grass moisture parametrization).

While the new live grass moisture parametrizations are brought more into agreement with established methods and the newly introduced LPJmL phenology-based live grass moisture shows a reasonably strong fit to satellite observations, some spatial incongruencies with observed data remain. In particular, the live grass moisture in the summer in Mediterranean regions may be too high. As shown in the observations of Chuvieco et al. (2009) and Mendiguren et al. (2015), and examined experimentally for select species by Bristiel et al. (2018) and Keep et al. (2021), Mediterranean grasses often enter dormancy in the summer. This would correspond, in the case of dormancy of all grasses in a grid cell, to a live grass moisture of 30 %, i.e. the minimum value, and, through the curing function, all of the grass would be transferred to the dead fuel category. There are two possible reasons why this is not the case in the modelled values.

The first reason may be the parametrization of the grass PFTs in the LPJmL vegetation model version used for these results. In the model, the temperate herbaceous PFT covers much of Europe and, since there is only one phenology function for each PFT, albeit with different inputs given grid cell conditions, temperate grasses across Europe follow the same response to stresses. Because of this, differences in adaptation to drought stress may not be captured and modelled grasses in the Mediterranean may exhibit tendencies to remain active under hot and dry conditions, as were observed for individuals from further northern regions by Bristiel et al. (2018) and Keep et al. (2021). Therefore, the excessive activity during dry periods may be a result of the manner in which vegetation is divided into PFTs by LPJmL, not allowing for the differences at an intra-specific level observed by Bristiel et al. (2018) and Keep et al. (2021) to manifest in the modelled results. In addition to these intra-specific adaptations, the use of broad PFTs can also result in the loss of inter-specific differences (as discussed by Fischer et al., 2018). This topic of the division of PFTs and its impact on model results is discussed further in Appendix C3. The issue of differences in adaptation, of course, applies to the GSI-based grass moisture as well, which is currently based on a single phenology (although it may be possible to adjust the thresholds used in the GSI equations on a regional basis in the future).

The second reason for the high modelled grass moisture levels in Mediterranean summers may be due to overestimation of live grass fuel moisture content in remotely sensed data under very dry conditions. Our modelled values, while exceeding those observed by Mendiguren et al. (2015) on the ground, are in much better agreement with values that they calculated based on satellite observations of vegetation greenness and only depart substantially from the on-the-ground values during the summer months. This potentially suggests a general challenge in deriving grass moisture from remotely sensed data, and future versions of the live

grass moisture parametrization in SPITFIRE may be further improved by also including local observations (e.g. those in Yebra et al., 2019).

These uncertainties in the live grass moisture parametrization are mitigated somewhat by the fact that there are years, as shown in Fig. 2 of Chuvieco et al. (2009), in which Mediterranean grasses may remain active. These years are exceptions rather than the rule, but they indicate that the modelled error is an increase in the frequency of an infrequent occurrence rather than the creation of entirely unrealistic conditions. In addition, the separate treatment of live and dead fuels and the new live grass moisture of extinction, which is generally higher than the dead fuel moisture of extinction, result in live grasses that can still burn at higher moisture content and in fires that can continue to burn in dead fuels even if this higher moisture of extinction in live fuels is exceeded. Together with the curing function, which transfers a portion of grass to the dead category beginning at 120 %, meaning that Mediterranean grasses in the summertime are generally treated as partially cured, this substantially reduces the impact of the too-high grass moisture levels, as shown by the model output in Sect. 3.3, particularly when compared to the previous parametrization.

Due to the mitigating factors above, the generally good agreement with satellite observations, shown in Fig. 7, and the substantial improvement over the original live grass moisture parametrization in SPITFIRE, the updated live grass moisture levels shown here represent a strong development in the context of SPITFIRE and in including soil moisture levels in live grass moisture calculations. Because further refinements of these parametrizations likely require changes on the DGVM side as well, they are outside of the scope of the current work, in which our developments focus on the major issues in the existing SPITFIRE parametrizations.

We discuss the remaining (lesser) sources of uncertainty in the SPITFIRE model following the order of the model structure in Fig. 2.

3.2.3 Fire danger index

As mentioned previously, there are three versions of the FDI implemented in different versions of SPITFIRE. The first, original formulation given in Thonicke et al. (2010) is different from the formulation in Venevsky et al. (2002) upon which it is based. While Venevsky et al. (2002) use an exponential function of the Nesterov index together with a tuning parameter to arrive at the FDI directly, Thonicke et al. (2010) apply the Nesterov index to calculate the dead fuel moisture as an intermediate step in calculating the FDI (Eq. 6). To our knowledge, this equation remains untested against observed fuel moisture levels, and an examination of its ability to predict dead fuel moisture may be relevant to versions of SPITFIRE that depend on it. To improve upon these issues, we replace this untested parametrization with a new dead

fuel moisture parametrization that makes use of a dynamic LPJmL-based litter moisture calculation developed by Lutz et al. (2019) (described further in Sect. 3.2.5). This allows us to replace the Nesterov-based FDI with the VPD-based FDI that was more recently developed for SPITFIRE by Druke et al. (2019).

3.2.4 Fire duration function

The fire duration function in SPITFIRE is shown for reference in Fig. S12 and given by Eq. (14) in Thonicke et al. (2010):

$$t_{\text{fire}} = \frac{241}{1 + 240e^{-11.06 \times \text{FDI}}}. \quad (5)$$

This function has a maximum value of 241 min and the functional form has a step-like nature such that at low FDI values the fire durations are extremely low, followed by a sharp increase around FDI values of 0.5. This results in fires with durations that are often substantially below the 241 min maximum. In practice, fires in the LPJmL4-SPITFIRE model have a median duration of about 30 min. In addition, the SPITFIRE model does not allow for fires that continue to burn over several days, despite this being a common phenomenon (see e.g. Andela et al., 2019, where the median fire duration globally is 3 d). Dividing multi-day fires into separate single-day fires suffers from the issue that the elliptical fire spread in SPITFIRE results in a quadratic dependence of fire size on fire duration. Therefore, as an example, two single-day fires have a combined fire size only half as large as one 2 d fire, all else being equal. This substantial downward bias in fire size is a large compensating error that prevents large positive biases due to the incorrect weighting factors in the fire spread component of the model, from causing highly overestimated modelled burnt areas. Now that these upward biases have been corrected, however, it is possible to implement a more realistic fire duration function.

For this purpose, we have introduced a multi-day burning algorithm into SPITFIRE. This allows fires that begin burning on one day to continue into the next, with an adjustable maximum number of days. We have currently implemented an FDI-based condition for stopping the spread of these multi-day fires, where all fires in a given grid cell are extinguished if the FDI reaches a value below 0.005. This condition operates in addition to the previous condition in SPITFIRE that fires are extinguished if the calculated fire-line intensity is below a specified threshold. These conditions, in particular the FDI threshold, are set as initial placeholders, and further work is required to develop a fire extinction function that can properly capture the conditions under which fires no longer spread. One benefit of the multi-day fire approach we introduce is that it allows for the existence of fires that experience less fire spread on some days, e.g. due to lower wind speeds, but continue burning and experience more spread on subsequent days.

The principle mechanism underlying the multi-day fire algorithm is that the major axis of the elliptical fires in SPITFIRE grows according to the daily wind speed and fuel bed parameters. The length-to-breadth ratio is calculated using the mean wind speed over the full set of days that the fire spreads, assuming that the fire spread continues broadly in the same direction each day. These simplifications are necessary based on the current model inputs and the state of knowledge of fire spread at the SPITFIRE scales. The fundamental challenge is that including changes in spread direction would require the use of coarse grid-cell-averaged wind directions in addition to the current grid-cell-averaged wind speeds. These directions can be quite variable on the smaller scales at which a fire spreads and, therefore, may not be sufficiently represented by the grid cell average. Because of these simplifications, the algorithm we have introduced should be considered experimental and requires further development and research to apply fully. However, it represents a pathway for removing the bias caused by including only single-day fires.

In addition to the multi-day algorithm, we have implemented the possibility of setting a daily maximum fire duration more than 4 h in the model, allowing fires to burn longer on a given day. However, the step-like shape of the function and the fact that modelled FDI values are generally substantially lower than the location of the step in the function result in daily fire durations that are often less than 30 min. As a preliminary solution, we have also introduced the possibility of a user-specified minimum fire duration into the equation and replaced the factor of -11.06 , which was set so that an FDI of 0.5 corresponds to 1/2 of the maximum fire duration of 241 min, with an adjustable parameter. The new equation is

$$t_{\text{fire}} = \frac{t_{\text{fire,max}} + 1}{1 + (t_{\text{fire,max}}/t_{\text{fire,min}} - 1)e^{k_{\text{fireduration}} \times \text{FDI}}}, \quad (6)$$

where $t_{\text{fire,max}}$ and $t_{\text{fire,min}}$ are respectively the maximum and minimum fire durations, in minutes, and $k_{\text{fireduration}}$ is a tuning parameter that controls the slope of the function before it saturates towards the maximum duration. A plot of this function is shown in Fig. S13.

These updates improve upon a conceptual limitation in SPITFIRE and establish a technical framework in the model code for calculating multi-day fire spread. Therefore, they act as a strengthened platform upon which to build more detailed fire duration functions. These functions are written so that, if desired by the user, the previous SPITFIRE fire duration function can be simply recovered by adjusting options in the model. The updated multi-day fire spread algorithm also differs from the one implemented in LPJ-LMfire by Pfeiffer et al. (2013) in the duration of fires per day, since they use the original daily fire duration function, and in the extinction criteria, since we choose a more flexible criterion based on the FDI rather than one that is based only on changes in fuel moisture due to precipitation.

3.2.5 Fire spread

To resolve the uncertainty regarding the Nesterov-based fuel moisture content described in the FDI section above, we replace the fuel moisture parametrization with the new parametrization developed for LPJmL by Lutz et al. (2019). This parametrization contains a full water balance for the litter that includes interception of precipitation by the litter, infiltration and runoff from the litter, and a temperature-based evaporation function. In addition to providing a stronger theoretical basis for the dead fuel moisture content, this has the benefit of creating greater internal consistency in LPJmL-SPITFIRE. Further improvements to the dead fuel moisture content component of the model may be made in the future by distinguishing between the moisture content of different fuel size classes, e.g. by using the Nelson dead fuel moisture model (Nelson, 2000; Carlson et al., 2007). We have made provisions for this in the model code by treating the moisture content of the different fuel classes separately, giving each one the current uniform value. Values from a future parametrization can therefore simply be input into the model.

In addition, we discovered an LPJmL-SPITFIRE-specific bug in the fuel load of live grass in which the amount of live grass was scaled by the phenology to represent curing. However, the cured component of the grass was then not accounted for. Our introduction of the curing function has resolved this issue.

Another LPJmL-SPITFIRE-specific bug was fixed in which the total amount of burnable live grass is overwritten during a loop over each PFT. The former version resulted in the total amount of consumable live grass being equal to the amount given by the final PFT. This reduction may account in part for why the extremely severe fires caused by the biases discussed above did not result in non-physically high fire carbon emissions.

Finally, to avoid excessive burning on open grasslands, we adopt the wind limitation function previously applied to JSBACH-SPITFIRE by Lasslop et al. (2014) in LPJmL-SPITFIRE. This function is a version of the original wind limit from Rothermel (1972), updated by Andrews et al. (2013). We also include the original wind limit function as an alternative option. In our tests, due to the relatively low values of grid cell and daily averaged wind speeds, this wind limit is often not reached, but we introduce it to provide for future parametrizations that may include higher wind speeds.

3.2.6 Mortality

We identified several potential improvements in the cambial damage mortality function in SPITFIRE. The cambial damage function is largely based on the work of Peterson and Ryan (1986), with a few modifications. In contrast to Peterson and Ryan (1986), who use a weighted averaging scheme to arrive at the residence time, the residence time in SPITFIRE is calculated using a simple ratio of the amount of fuel

consumed to Rothermel's reaction velocity, Γ . In this ratio, the amount of fuel consumed is simply calculated using an average of the amount of 1, 10, and 100 h fuel consumed, i.e. a sum of the three values divided by three. In Peterson and Ryan (1986), the fuel consumption of different fuel classes is combined into a common value using a weighting based on the amount of fuel bed area that is covered by the individual components. To address this, we simplify the residence time equation to the simple function of fuel surface-area-to-volume ratio given by Albin (1976). The surface-area-to-volume ratio input into this parametrization results from the Rothermel weighting scheme described previously and therefore accounts for fuel bed heterogeneity in a more conceptually sound manner.

In the subsequent step of the cambial mortality function, the amount of time that the vegetation experiences lethal heat τ_L is calculated from the residence time. In Peterson and Ryan (1986) this value is 5 times the residence time, whereas in SPITFIRE this value is only 2 times the residence time. As there is no conceptual reason for this difference, we reinstate the value of 5. This substantially lower residence time may have also compensated somewhat for the excessive fire intensity caused by the biases in the application of the Rothermel equation.

The equation for the probability of mortality due to cambial damage that is based on this burning time is given in Eq. (19) of Thonicke et al. (2010). We discovered some disagreement between this equation and the cited literature, Peterson and Ryan (1986), on which it is based. Specifically, Peterson and Ryan (1986) combine cambial damage and crown scorch into a single equation, in their Eq. (11), whereas SPITFIRE treats the two separately and combines the probabilities of mortality due to cambial damage and crown scorch under the assumption that these are independent. This assumption does not necessarily agree with the physical processes of mortality wherein a more intense fire produces more heat as well as taller flames and is therefore more likely to both damage the cambium and scorch the crown of a tree. We therefore implement the Peterson and Ryan (1986) probability of mortality calculation as a simplified parametrization that removes the uncertainty associated with the theoretical basis of the SPITFIRE functions. This also allows for an easier interpretation of model results and forms a basis for future work improving the mortality parametrization. The equation is given by

$$P_m = c_k^{\tau_c/\tau_L - 0.5}, \quad (7)$$

where c_k is the fraction of the crown that is scorched, determined using the scorch height and tree geometry, τ_c is the critical time for cambial damage, in minutes, and τ_L is the amount of time, in minutes, that a tree experiences lethal heat.

Finally, the equation for the c_k parameter, the parameter describing the fraction of a tree crown that is killed by fire,

in Peterson and Ryan (1986) is based on a paraboloid crown structure. In SPITFIRE this was replaced with a cylindrical crown structure, reducing the amount of the crown that is near the ground (this change is also present in the adaptation of the c_k parameter for the crown fire parametrization in Ward et al., 2018). Because the new fire spread formulation avoids excessive crown scorch due to high-intensity fires, we return to the formulation of Peterson and Ryan (1986) for the sake of conceptual uniformity and the more realistic crown structure described therein.

3.3 Updated model version at the European scale

The results shown here are preliminary in nature and intended to ensure that the changes made to the model perform reasonably. In addition to implementing the changes described in previous sections, we have conducted a thorough code cleanup and have introduced a new change log in the model. For the new model version we have re-tuned the parameters for the European domain. We follow the standard tuning approach for the model and use the PFT-specific α parameters as the main tuning parameter but also tune various ignition and fire duration parameters listed below. The values of these parameters for each PFT are shown in Table 1. We set the fire duration to a minimum of 2 h and a maximum of 7 h. The maximum number of consecutive days that a fire can burn is set to 3. On managed grasslands, which forms a small portion of the modelled grid cells, these parameters are set to a minimum of 1 min, a maximum of 2 h, and a maximum of 1 d respectively. In the fire duration function, Eq. (6), the factor $k_{\text{fireduration}}$ multiplying the FDI is set to -8 from the previous -11.06 , allowing for a more gradual rise in fire duration with increasing FDI. Finally, in the human ignitions function, Eq. (B1), the coefficient at the start of the equation, which we now call $k_{\text{ignitions}}$, was changed from 30 to 165. We have applied a fairly liberal tuning to these parameters as there is no clearly established value for them currently, and our current aim is largely to compare the impact of the changes we have introduced.

The comparison between the new SPITFIRE version and the old SPITFIRE version implemented in LPJmL5.7 is shown in Fig. 8. The old model version shows the same biases in grasslands as the version implemented in LPJmL4 (shown in Fig. 3). Namely, there is an extreme over-burning in grassland-dominated grid cells, shown most clearly in Fig. 8f, despite forested grid cells, in this case, showing rough agreement for many years. The new model version shows a substantial improvement in this regard, showing better agreement in grass-dominated grid cells and reasonable agreement in tree-dominated grid cells. This improvement is visible in central areas of the Iberian Peninsula, for example, where modelled grasslands (see Fig. S14) result in a higher burnt area in the old model version than the new. The improvement in this division is particularly noteworthy since this division was not used as a target when tuning the

model, with only the broad spatial pattern and annual burnt area being targets. The new model version also results in a much lower average rate of spread, shown most clearly in Fig. 8i, and a reduction in the number of trees that undergo fire mortality, shown in Fig. S14. Therefore, the positive biases shown in Fig. 5 had a substantial impact on model results, which is now reduced. The higher burnt area in tree-dominated grid cells for the old model version in Europe, shown in Fig. 8e and in contrast to the global results in Fig. 3d, may suggest a less extreme tree mortality in Europe, also illustrated by the lack of any complete gaps in the tree FPC map in Fig. S14.

Some issues remain in the new model version, however. Most importantly, the model's lack of ability to reproduce regions of high burnt area, such as Portugal, illustrates that there remain factors determining burnt area that the model currently does not capture, including fire suppression, fragmentation effects, and greater flammability of some species of vegetation. In addition, there is a substantially too-low burnt area in eastern Europe in the new model version. This may be due to an underrepresentation of ignition factors in that region, as the rate of spread, shown in Fig. 8i, remains high in this area. A more detailed study of this region is outside the scope of this work, but future work may examine the modelling of these sources of ignition further.

The fixes to the implementation of the Rothermel equation result in substantial reductions in the rate of spread and the elimination of regions where there are extremely high rates of spread in the old model version. Due to the lack of reliable spread rates in satellite-based products, we reserve the testing of modelled rates of spread against local data for future work. Overall, the new model version shows a substantial improvement in the relative amount of fire in forested grid cells as opposed to those dominated by grassland and has a substantially corrected theoretical basis. It can therefore be adopted as a foundation upon which to build future model versions.

3.4 Model status

We have identified several sources of uncertainty in the SPITFIRE model. The most important of these are the inaccurate implementation of the Rothermel equation and the unrealistically low live grass moisture levels. We resolve these issues in part by correcting the implementation of the Rothermel equation in SPITFIRE and by introducing new live grass moisture parametrizations. These parametrizations produce much more realistic live grass moisture levels. The applied LPJmL phenology-based parametrization, in particular, represents an innovative approach to incorporating soil moisture in the calculation of herbaceous live fuel moisture, an important development, the need for which has been identified in previous studies (Brown et al., 2022; Krueger et al., 2022; Jolly et al., 2005). Our test of the new SPITFIRE version in the European domain shows that it is capable of broadly representing burnt area, similar to the previous model version,

Table 1. Tuning parameters for the European model version: α parameters are the FDI scaling parameters for each PFT, $t_{\text{fire,min}}$ and $t_{\text{fire,max}}$ are the minimum and maximum daily fire durations in minutes, $k_{\text{fire, duration}}$ is the slope factor applied to the daily fire duration, $k_{\text{ignitions}}$ is the scaling factor applied to human ignitions, and fire days is the maximum number of days for which fires are allowed to burn.

α_{TrH}	α_{TH}	α_{PH}	α_{TBE}	α_{TNE}	α_{TBS}	α_{BNE}	α_{BNS}	α_{BBS}	$t_{\text{fire,min}}$	$t_{\text{fire,max}}$	$k_{\text{fire, duration}}$	$k_{\text{ignitions}}$	fire days
4	4	7	10	10	10	15	15	15	120	480	−8	165	3




Figure 8. Impact of changes to the LPJmL5.7-SPITFIRE model in the European domain. Maps in panels (a) through (c) show mean burnt area per grid cell over the 2003–2014 simulation time period. Time series in panels (d) through (f) show total annual burnt area over the simulation domain. Tree burnt area and grass burnt area refer to the burnt areas in the grid cells made up of over 50 % tree PFTs and grass PFTs respectively. Panels (g) through (i) show maps and time series of the rate of spread calculated in the new and old model versions. The time series in this row contains the mean annual value rather than the annual sum as in the row above. The new model version shows a substantial improvement in the relative amounts of burnt area in forests compared to grasslands. It is also less volatile, with less extreme inter-annual variability.

while providing a substantially strengthened theoretical basis.

Because of the extensive changes we have made to the LPJmL-SPITFIRE model and the uncertainties we have discovered, it is relevant at this point to discuss the current status of the model. In addition to summarizing the changes that we have made, we include a discussion of sources of uncertainty that should be taken into account when using the model and that may be improved upon in the future. While many of these sources of uncertainty are familiar to those experienced with fire-enabled DGVMs, we include a detailed discussion here both for the sake of transparency and to support future work. It is our aim to clarify these sources of uncertainty for those unfamiliar with the SPITFIRE model but who may be users of model results, are new adopters of

the model, or may have expertise in other areas of fire research that could be used to help develop improved model parametrizations. In many cases the improved model foundations that we have presented here enable these improvements to be made to a greater extent. For example, the ignitions parametrizations can now be improved using more realistic data as they no longer need to compensate for the upward biases in the fire spread component of the model. It should be noted that the impact of these sources of uncertainty is dependent on the application to which the model is put, and the specific use case should be considered when interpreting their importance. The changes made to LPJmL-SPITFIRE and key areas that remain for future development, which are summarized in Fig. 1, are discussed in detail below.

3.4.1 Uncertainties and areas for future development in specific model components

Areas for future development in specific model components are discussed in Appendix B. For the ignitions component, we highlight the uncertainty in the human ignitions parametrization and the temporal averaging of lightning ignitions that lead to small fractions of one ignition each day that may not align with the timing of associated precipitation (Appendix B1). These components, in particular, have gained a greater scope for future developments through the model improvements in this work since they are no longer required in order to compensate for excessive fire spread. For the fire spread component, we highlight simplifications in the moisture of extinction and surface-area-to-volume ratios that may reduce model accuracy as well as aspects that could be important to fire spread that are currently not modelled (Appendix B2). A particular detail about the fire spread component that emerged in our literature analysis and where a new parametrization remains necessary is the calculation of fuel bulk density.

Fuel bulk density

A major outstanding issue in the fire spread component of the model is its treatment of the fuel bulk density, i.e. how tightly packed the fuel beds are, an important input in the Rothermel model. In its fire spread component, SPITFIRE uses literature-based values for the fuel bulk density for each plant functional type. These values are then modified by the following equation given in Table A1 in Thonicke et al. (2010):

$$\rho_b = \frac{1}{n} \sum_{\text{PFT}=1}^n \rho_{b,\text{PFT}}(w_{0,1\text{h}} + 0.2w_{0,10\text{h}} + w_{0,100\text{h}}). \quad (8)$$

Here, n is the number of plant functional types in a given grid cell, $\rho_{b,\text{PFT}}$ is the PFT-specific fuel bulk density value, in kg m^{-3} , and $w_{0,x}$ is the fuel load in fuel class x , in kg m^{-2} . The justification for this equation is based on a statement in Brown (1981) that when the 10 h fuel class is eliminated from calculations, the calculated bulk densities averaged 80 % of their previous values. Equation (8) does not have this effect. Rather, it leads to a scaling of the bulk density depending on how much total fuel load is present. Therefore, this question of parametrizing fuel bulk density in an aggregated grid-cell-scale manner remains open, and a new parametrization is required.

3.4.2 The impact of model resolution

In addition to the sources of uncertainty that we have highlighted above, there are several sources of uncertainty that arise due to the resolution of SPITFIRE and attached DGVMs. In several cases, which we highlight, these sources of uncertainty can be better mitigated due to the improved

theoretical basis for SPITFIRE that we have established in this work.

Model resolution is one area where global fire models necessarily differ from predictive models used in an operational context, such as FARSITE and FlamMap (Finney, 1998, 2006). There are three domains in which resolution plays an important role in the SPITFIRE model. The first is spatial resolution (i.e. the size of the grid cells), the second is the temporal resolution, and the third is the separation of global vegetation into PFTs. In all of these cases a certain coarseness is required to feasibly run the model on a global scale. We include a detailed discussion of these three domains as guidance for model users in Appendix C.

We particularly highlight the effects of the information given to SPITFIRE by associated DGVMs and cases where these differ from one DGVM to another. Questions of vegetation size, arrangement, and what times during the year fuel beds are replenished are particularly relevant, as well as the choice of PFTs and how they align with fire characteristics for the regions under study. Spatially, challenges may be posed by calculating fire spread in fuel beds that are homogenized across a grid cell and that therefore do not contain fragmentation effects or changes in fire behaviour between differently vegetated sub-grid regions (Appendix C1). Temporally, annual time steps of vegetation establishment and mortality may impact regions with bimodal fire seasons where vegetation is replenished in the middle of the year and regions where the fire season encompasses the turn of the year, resulting in unrealistic developments in modelled fuel beds (Appendix C2). A common issue in several DGVMs in which SPITFIRE is integrated is the lack of a shrub PFT for regions in which shrubs are important parts of the fire regime, e.g. chaparral-covered regions of California and parts of the Mediterranean where shrubs play a key role in post-fire dynamics (Weise et al., 2016; Baudena et al., 2020). To help address this, we have made provisions in our updated fire spread parametrizations to facilitate the inclusion of shrubs in the future (Appendix C3). As a representative example, one that is most relevant to the model improvements in this work, here we highlight the impact of model resolution on wind speeds.

Relevance for the incorporation of wind speeds

An area where the model improvements presented in this work can contribute substantially to future developments is the wind speeds input into the Rothermel model. These are particularly subject to the impacts of model resolution. Spatially, interactions between terrain and wind, in particular, could be examined more closely, as complex terrain can have a substantial effect on wind speeds local to the flame front in a manner that is not reflected by the grid-cell-averaged wind speed (e.g. see the strong wind speed variance across domains with complex terrain shown by Jung and Schindler, 2020). Temporally, wind speeds are often lower at night than

during the day (e.g. Ephrath et al., 1996; He et al., 2013). Therefore, the daily averaged wind speed currently used as an input to the model may not be reflective of the wind speed during the hours in which fires are most likely to spread. Future work could examine the feasibility of using averaged wind speeds over the hours during which the VPD meets some threshold for fire spread, as discussed in the context of fire duration in Sect. 3.2.4 or, more simply, whether midday wind speeds may be more appropriate as an input parameter than the daily average. With the removal, in the new SPITFIRE version, of the large positive biases in the rate of spread and scorch height caused by the previous implementation of the Rothermel model it may be possible to incorporate these higher wind speeds effectively.

3.4.3 Model tuning and validation

The fact that the issues we have highlighted, particularly the bias in the implementation of the Rothermel equation and the unreasonably low live grass moisture levels, were able to remain undetected in previous versions of SPITFIRE, suggests that more detailed validation procedures may be beneficial to future model versions. In this work, we developed two new validation methods, namely the division of burnt area by vegetation and the use of prescribed fire starts, which have shown an ability to identify underlying issues not revealed by standard validation procedures. This suggests that the proposed validation methods, in addition to the more detailed methods already in use in the FireMIP project (Rabin et al., 2017), may be highly useful in future developments. An additional element that was highlighted by the results in this work is the potential ability of model tuning to obfuscate biases by creating errors that compensate for one another (as illustrated, for example, by the reasonable agreement of mapped burnt area in Fig. 3 that did not arise from a reasonable balance of fire spread and ignitions as shown in Fig. 4).

Model tuning is a key component of the development process for global models in general and process-based global fire models in particular (e.g. Hourdin et al., 2017; Hantson et al., 2016). Because compounding errors can often arise, e.g. due to measurement uncertainties in observed input parameters or modelling uncertainties in modelled input parameters, tuning fire model equations based on observed data is often necessary if the goal is to align model output with these data. In the case of the SPITFIRE model, process parametrizations are generally tuned to achieve agreement between modelled and satellite-based mean burnt area maps, as well as annual time series of burnt area, over the modelled domain. Generally, the parameters used for model tuning are those which are the most uncertain due to a lack of data constraining their values. Most commonly, the number of human ignitions and the PFT-dependent scaling parameters applied to the fire danger index are the main levers by which model outputs are optimized. These parameters are adjusted by hand until the maps and time series in question

achieve acceptable agreement with their observed counterparts. It should also be noted that the DGVMs coupled to SPITFIRE contain their own tuning processes that are similar to this approach albeit with different targets (e.g. von Bloh et al., 2018a; Scheiter et al., 2013). Therefore, the manner and targets of the tuning procedure may be a point of consideration for future users of SPITFIRE, particularly if values from particular sub-components of the model are desired outputs. For example, applications for which the number of fires is an important output, in addition to the burnt area, would benefit from a tuning procedure that takes into account the prescribed fire starts validation at an intermediate step.

More generally, and importantly for future predictions, the fact that the validation dataset is used to inform the model tuning approach in an iterative manner results in a substantial transfer of information from the former to the latter that weakens the model validation, as the validation dataset is no longer an independent source of information. In objective tuning methods, as discussed by Hourdin et al. (2017) and applied to the SPITFIRE model for a South American domain by Drüke et al. (2019), it is common practice to divide data into separate datasets for training and for testing, resulting in a reduced transfer of information. The objective tuning approach is not a panacea, however, as it can result in tuning parameters that are not physically justified, and this must therefore be carefully monitored by modellers. However, if such unphysical parameters are arrived at, this can also be valuable information about the model, as it indicates model biases that may require re-parametrization (Hourdin et al., 2017).

The division of validation data into training and test datasets could also be implemented for cross-validation of the hand-tuning approach commonly used in SPITFIRE. The performance of SPITFIRE in grid cells that are not used in the tuning process would give a more robust estimate of model performance outside of its training data (e.g. see the general discussion of cross-validation in Morin and Davis, 2017). The issue of increasing uncertainty outside of the training data may also be visible in the strong agreement of global fire models, in general, within the time period for which satellite data are available, in contrast to their strong divergence outside of this time period (e.g. Teckentrup et al., 2019). While such upgrades to the model tuning approach in LPJmL-SPITFIRE are out of the scope of the current text, we have sought to discuss our model tuning approach transparently to allow for greater clarity in the interpretation of model results, and we recommend the cross-validation approach here for future full SPITFIRE versions.

3.4.4 Conditions for best performance of LPJmL-SPITFIRE1.9

Based on the qualitative and quantitative analysis in this work we can outline the conditions under which we anticipate the best model performance for model versions based on

LPJmL-SPITFIRE1.9. Note that not meeting all of these conditions does not preclude satisfactory model performance; they simply indicate ideal conditions. Also, because of the possibility for model tuning to balance errors in one spatial region with errors in another, these conditions do not necessarily correspond to the accuracy of final model results in specific geographic regions. However, a focus on these regions may be effective when conducting model developments. The conditions for best model performance given the current status of the model are ones where the following are true.

- The coupled vegetation model accurately represents fuel beds.
- The species that dominate the fire regime align with the plant functional types of the DGVM (e.g. the fire regime is not dominated by shrubs in a DGVM that does not model them).
- Potential human ignitions occur steadily throughout the year (although successful ignitions may still vary considerably due to meteorological conditions), and lightning ignitions do not vary substantially between years or are not a major ignition source.
- Fires are generally short in duration and are therefore less subject to the uncertainties in propagating longer duration fires.
- The fire regime in a given location is dominated by surface fires, and fire suppression does not have a strong impact on overall burnt area.
- The region has a subhumid or humid climate, as defined by Scott and Burgan (2005), to ensure the dead fuel moisture of extinction in SPITFIRE is accurate.
- Vegetation in a grid cell is homogeneous and evenly distributed.
- Terrain in a grid cell is flat.
- Fires occur toward the middle of the year in DGVMs that calculate annual establishment and mortality.
- Wind speeds are temporally and spatially homogeneous, and wind directions are steady.

4 Conclusions

We have undertaken a thorough review of the global fire model SPITFIRE to identify and better understand the sources of uncertainty in the model and the cause of several known issues in the model results. We have found that two major sources of error exist in the model. First, the

model contains an incorrect implementation of the Rothermel fire spread model that can result in substantially too-large and too-intense fires. Second, the model contains a live grass moisture parametrization that results in unrealistically low moisture content. We correct these issues with a corrected implementation of the Rothermel model and a novel live grass moisture parametrization for the European domain. Other sources of uncertainty are identified and partially corrected. This results in an updated version of the model that is more aligned with the physical basis of wildland fire spread and allows for future additions of more accurate parametrizations to other parts of the model. The updated model retains the ability of previous model versions to broadly represent spatial patterns of burnt area. Results in the European domain show a reduction in excessive burning on modelled grasslands and reduced tree mortality. Further work is required to test the SPITFIRE model in the global and European domains and to address challenges of the modelling scale and required re-parametrizations. By improving the theoretical basis of the SPITFIRE model and highlighting directions of future model development we have sought to create a foundation upon which such work can be built. In particular, the new live grass moisture parametrization we introduce here is an example of such a development, as the model no longer requires unrealistic live grass moisture levels for the fire spread component to function, allowing for a representation that better captures seasonal moisture dynamics.

Appendix A: Detailed errors in the SPITFIRE implementation of the Rothermel model

A1 Background on the Rothermel equation

The Rothermel equation, Eq. (A1), is a semi-empirical rate-of-spread equation first developed by Rothermel (1972). Fundamentally, it derives the rate of spread of a fire by dividing the rate at which energy is released towards a section of fuel by the amount of energy that is required for that section of fuel to ignite. Specifically,

$$R = \frac{I_R \xi (1 + \phi_w + \phi_s)}{\rho_b \epsilon Q_{ig}}, \quad (\text{A1})$$

where R is the rate of spread, in m s^{-1} ; I_R is the reaction intensity, i.e. the rate at which energy is released by the fire, in W m^{-2} ; ξ is the propagating flux ratio, the proportion of that energy that is transmitted to the subsequent part of the fuel bed; and ϕ_w and ϕ_s are factors that describe the effect of wind and slope on the propagating energy flux. In the denominator, ρ_b is the bulk density of the fuel, in kg m^{-3} ; ϵ is the effective heating number, i.e. the proportion of a fuel particle that must be heated to ignition for combustion to occur; and Q_{ig} is the amount of energy required to heat a kilogram of fuel to ignition, in J kg^{-1} (Rothermel, 1972; Andrews, 2018).

The individual parameters of Eq. (A1) are functions of further fuel bed parameters, including the fuel bed depth, surface-area-to-volume ratio of the fuel particles, fuel moisture content, fuel moisture of extinction, and heat content. In the case of a uniform fuel bed, the surface-area-to-volume ratio and moisture content implemented in the Rothermel equation are simply the moisture content and surface-area-to-volume ratio of the uniform fuel particle size. In the case of a non-uniform fuel bed, Rothermel (1972) introduced a weighting system based on the contribution of individual fuel size classes to the overall surface area of the fuel bed. The commonly used formulation of this weighting system is the updated version described in Albin (1976). In this system, the fuel bed is divided into living and dead fuel categories as well as different fuel size classes. Each category and size class has a weighting factor, f_{ij} , assigned to it, where i connotes the living and dead fuel categories and j connotes the different fuel size classes. These factors are calculated using

$$f_{ij} = \frac{\sigma_{ij} w_{0,ij} / \rho_{p,ij}}{\sum_j \sigma_{ij} w_{0,ij} / \rho_{p,ij}}, \quad (\text{A2})$$

where σ_{ij} is the surface-area-to-volume ratio of a fuel class, in m^{-1} ; $w_{0,ij}$ is the oven-dry fuel load of the fuel class, in kg m^{-2} ; and $\rho_{p,ij}$ is the particle density of the fuel in the fuel class, in kg m^{-3} . The summation in the denominator signifies a summation over all components within the dead or live fuel category, i . The numerator in Eq. (A2), therefore, is the total surface area of a given fuel class in a unit area of the fuel bed, and the denominator is the total surface area of all classes over that same unit area. This results in a weighting by fuel surface area. As an example, the combined representative surface-area-to-volume ratio of the dead fuel category ($i = 1$) would be

$$\sigma_1 = \sum_j f_{1,j} \sigma_{1,j}. \quad (\text{A3})$$

One subtlety, added to this approach by Albin (1976), is the inclusion of the g_{ij} weighting factors for combining fuel loads. These factors are similar to the f_{ij} factors, with the distinction that they gather the individual fuel classes into bins based on their surface-area-to-volume ratio and assign a single weighting factor to all fuel classes in a bin. This circumvents a conceptual issue in the original Rothermel equation where the f_{ij} factors result in differing combined fuel loads depending on how the fuel classes are partitioned (Albin, 1976; Andrews, 2018). The g factors for each size bin are calculated using

$$g_{ij} = \begin{cases} \sum_{\text{bin}} f_{ij}, & \text{if } \sigma \geq 0.54 \text{ cm}^{-1} \\ 0, & \text{otherwise} \end{cases}. \quad (\text{A4})$$

A2 Implementation of the Rothermel equation in SPITFIRE

Fundamentally, the main error in the implementation of the Rothermel equation in SPITFIRE results from an application of the Rothermel equation for a uniform fuel bed to the non-uniform fuel beds present in the vegetation models with which SPITFIRE operates. Rather than applying the form designed for a non-uniform fuel bed, the variance in fuel bed parameters is accounted for by combining them according to an incorrect weighting scheme. In this scheme, the surface-area-based weighting factors were neglected in the implementation of the Rothermel equation and a different weighting approach was taken, whereby most of the components are weighted by their contribution to the total fuel bed mass, e.g.

$$\sigma = \sum_i \sum_j \sigma_{ij} \frac{w_{0,ij}}{w_0}, \quad (\text{A5})$$

where w_0 is the total oven-dry fuel load of the fuel bed (Thonicke et al., 2010). The standard weighting system shown in Eq. (A2) is more heavily weighted toward fine fuels due to its weighting by the contribution of each fuel class to the total surface area of the fuel bed. The SPITFIRE weighting system in Eq. (A5), however, is more heavily weighted towards coarse fuels as they are heavier and therefore contribute to a greater proportion of the overall fuel bed's mass.

In addition to the lack of f_{ij} factors in SPITFIRE, the g_{ij} factors are omitted entirely, and the fuel load that is inserted into the Rothermel equation is a simple sum of the individual fuel loads (Thonicke et al., 2010):

$$w_0 = \sum_i \sum_j w_{0,ij}. \quad (\text{A6})$$

This contributes an even larger source of error since the replacement of a weighted average by a simple sum can increase the fuel load by a factor up to the number of individual fuel components there are (e.g. if there are three fuel classes the sum may be 3 times larger than the average).

Further, an element of the Rothermel equation that is lacking from SPITFIRE is the separate treatment of dead and live fuels. For example, the reaction intensity in the non-uniform fuel bed Rothermel equation is calculated using

$$I_R = \Gamma' \sum_i w_{0,i} (1 - S_{T,ij}) h_i \eta_{M,i} \eta_{s,i}, \quad (\text{A7})$$

where Γ' is the optimum reaction velocity of the fuel, in min^{-1} ; $S_{T,ij}$ is the mineral content of the fuel component; h_i is the weighted average heat content of the fuel bed component, in kJ kg^{-1} ; $\eta_{M,i}$ is a moisture dampening coefficient; and $\eta_{s,i}$ is a mineral damping coefficient (Andrews, 2018; Albin, 1976). The moisture damping coefficient is calculated using

$$\eta_{M,i} = 1 - 2.59 \frac{M_{f,i}}{M_{x,i}} + 5.11 \left(\frac{M_{f,i}}{M_{x,i}} \right)^2 - 3.52 \left(\frac{M_{f,i}}{M_{x,i}} \right)^3, \quad (\text{A8})$$

where $M_{f,i}$ is the moisture content of a fuel component, in kilogramme water per kilogramme oven-dry mass, and $M_{x,i}$ is the moisture of extinction of that fuel component. As shown in Eqs. (A7) and (A8), the moisture content of the dead fuel component and that of the live fuel component are treated separately before being combined in the Rothermel equation. In the implementation of the Rothermel equation in SPITFIRE, however, the moisture content of the dead fuel component and that of the live fuel component are combined using the following series of equations:

$$\alpha_2 = -\ln(M_{f,2})/\text{NI}, \quad (\text{A9})$$

$$M_{f,\text{combined}} = \exp\left(-1 \times \left(\alpha_1 \frac{w_{0,1}}{w_0} + \alpha_2 \frac{w_{0,2}}{w_0}\right) \times \text{NI}\right), \quad (\text{A10})$$

where NI is the Nesterov index (see Thonicke et al., 2010). Note these equations are present in the SPITFIRE code, documented in Schaphoff et al. (2018b) and alluded to in Pfeiffer et al. (2013). However, they are not explicitly given in Thonicke et al. (2010). Equations (8) and (9), together with Eq. (6) in Thonicke et al. (2010), can simply be combined into a single equation:

$$M_{f,\text{combined}} = M_{f,1}^{\frac{w_{0,1}}{w_0}} M_{f,2}^{\frac{w_{0,2}}{w_0}}. \quad (\text{A11})$$

Effectively, SPITFIRE uses a combined moisture content that is a weighted geometric mean of the dead and live fuel moisture content and places this in the Rothermel equation for uniform fuels, despite operating on non-uniform fuel beds.

Appendix B: Sources of uncertainty in specific model components

B1 Ignitions

B1.1 Human ignitions

Human ignitions in SPITFIRE are calculated using a simple function of population density and a parameter that describes the propensity of humans in a given grid cell to cause ignitions. The full equation is (combining Eqs. 3 and 4 in Thonicke et al., 2010)

$$n_{h,\text{ig}} = 30 P_D a(N_D) e^{-0.5 \times \sqrt{P_D}} / 100, \quad (\text{B1})$$

where P_D is the population density (in individuals per km²) and $a(N_D)$ is the propensity of people in a given grid cell to cause ignitions (in ignitions per individual per day). This function was based on theoretical considerations, including findings by Archibald et al. (2009) for southern Africa that suggest a relationship between population density and the number of ignitions that increases until reaching a maximum at intermediate population densities, followed by a decrease. The $a(N_D)$ parameter was derived for some regions using local fire databases and used as a tuning parameter for others.

This parametrization does not explicitly take into account different fire management practices and how these practices depend on local conditions. Connecting this parametrization to these measures, e.g. as has been explored by Perkins et al. (2022), is an avenue of future model development that may improve burnt area distribution in SPITFIRE.

B1.2 Lightning ignitions

The 5 years of the LIS/OTD monthly dataset used in SPITFIRE were averaged in the original parametrization to produce 1 year of monthly values that was then interpolated to a daily lightning dataset. This results in a consistent, low number of lightning ignitions, with a small fraction of one ignition on most days. Because of this, the lightning ignitions are also desynchronized from any precipitation that may accompany them. The impact of temporal resolution on lightning ignitions has also been explored by Felsberg et al. (2018), showing a small impact on a global scale but greater effects regionally. Alternative lightning ignition parametrizations for use in SPITFIRE-related models suggested by Pfeiffer et al. (2013) and Kelley et al. (2014) may help to remedy this. Because these models are SPITFIRE-related, testing their implementations in future versions of SPITFIRE may be straightforward. Further model development may also be based on newer lightning datasets, e.g. Kaplan and Lau (2021), that allow for a better understanding of inter-annual variability in lightning ignitions and their impact on burnt area. Given the newly available datasets and the improved model foundations here, revisiting the lightning parametrization in SPITFIRE may be a fruitful area of research, particularly for regions that are remote from human activity and in which lightning is, therefore, the dominant source of ignitions.

B2 Fire spread

B2.1 Surface-area-to-volume ratio

The SPITFIRE model undertakes some simplifications in its fuel bed parameters that may be built upon in future work. This includes uniform surface-area-to-volume ratios for fuels across all vegetation types. While this is common for the coarser size classes, the surface-area-to-volume ratios of the 1 h fuel classes generally show some distinctions, e.g. between leaf-dominated fuel beds and needle-dominated fuel beds. The Scott and Burgan (2005) fuel models, for example, have 1 h surface-area-to-volume ratios that range from 1500 to 2200 ft⁻¹ (49.2 to 72.2 cm⁻¹), not including shrub fuel models, whereas SPITFIRE uses a uniform value of 2021 ft⁻¹ (66 cm⁻¹). Since the surface-area-to-volume ratio is a key parameter for calculating many of the variables in the Rothermel model, potential model improvements may be made by varying the 1 h surface-area-to-volume ratio for different vegetation types.

B2.2 Dead fuel moisture of extinction

The dead fuel moisture of extinction in SPITFIRE is a uniform value of 30 %. This is distinct from the Scott and Burgan (2005) fuel models, for example, where $M_{x,dead}$ has a range of 15 %–40 %, with values depending on the type of vegetation and climate. Therefore, variable dead fuel moisture levels of extinction are another potential improvement for the SPITFIRE model. A challenge in this regard would be determining an appropriate manner for constructing a crosswalk between the Scott and Burgan (2005) fuel models and the LPJmL PFTs as the fuel models depend on both vegetation and climatic parameters.

B2.3 Crown and ground fires

Future model developments may expand the fire spread component of the SPITFIRE model to include other forms of fire behaviour. Currently, the model only models surface fire spread, but crown fires and ground fires are also major contributors to fire dynamics. Crown fires have been incorporated into DGVMs, e.g. by Ward et al. (2018), and are important due to the increased vegetation mortality they may cause and the high spread rates present in them. Ground fires, among other factors, play a substantial role in Arctic fire dynamics and may therefore be relevant to future modelling endeavours that focus on fires at high latitudes (McCarty et al., 2021).

B2.4 Convective and terrain-driven fires

Another dynamic that may be relevant for future inclusion in the SPITFIRE model is the effect of atmospheric stability on fire growth and the potential for larger and more severe fires during extreme pyroconvection events (e.g. Senande-Rivera et al., 2022). In addition, terrain can act to promote fire spread through increased rates of spread when fires are spreading up slopes (although, at a grid cell scale this effect may be counteracted somewhat through the barriers to fire spread that complex terrain creates).

B2.5 Fire suppression

Finally, fire suppression can impact the size and incidence of fires. It is currently not included in the model but may improve modelling accuracy in the future.

Appendix C: The impact of model resolution

C1 Spatial resolution

Wildland fires are an inherently multi-scale phenomenon, with conductive heat transfer that can occur on the scale of microns up to fire–atmosphere interactions that can occur on scales of tens of kilometres (e.g. Collin et al., 2011; Potter,

2012). Because of this it is often necessary to include empirically derived equations in process-based models to account for sub-grid-cell processes. In the case of SPITFIRE, a substantial factor that is not currently accounted for is the arrangement of fuel beds at a sub-grid level. Fuel bed heterogeneity may arise from terrain, the spatial arrangement of vegetation, and human influence, and it can have strong effects on determining the final shape and size of fires (e.g. Sharples, 2009; Brown, 1981; Narayananaraj and Wimberly, 2011). We therefore highlight this as an area of potential future model development.

The spatial arrangement of fuel beds can also be influenced by previous fires, and this leads to an issue in LPJmL-SPITFIRE specifically due to the manner in which fuel consumption is calculated. In the case of fuel consumption, the current approach in the model is to apply fuel consumption to the entire grid-cell-averaged fuel bed, scaled by the burnt area. Because of this if, e.g. 5 % of the grid cell is burnt with sufficient intensity to consume the entirety of the fuel, the model treats this as 5 % of the total grid cell fuel bed being “skimmed off.” This can substantially weaken fuel feedbacks in grid cells where a low portion of the grid cell is burnt due to the small reduction in fuel load. In LPJ-GUESS-SPITFIRE this issue is mitigated by including burnt patches as separate stands, allowing for strong local effects of fires. This approach may be an avenue for improvement of other SPITFIRE implementations, and it may be preferable for users of SPITFIRE to work with an implementation in LPJ-GUESS if such feedbacks are important to their use case.

C2 Temporal resolution

SPITFIRE operates on a daily time step and many processes in the DGVMs with which it is coupled generally operate on an annual time step. Processes on the daily time step generally use daily averaged values to calculate parameters such as the dead fuel moisture content. Because fire spread largely occurs during the day (e.g. Balch et al., 2022), there may be some bias in applying daily averaged values as inputs and model accuracy may be improved in the future by considering inputs on the basis of the times of day during which fire spread occurs. For example, the 1 h dead fuel moisture is highly subject to sub-daily variations and may be better captured if calculated using daytime values.

Regarding the time step of DGVMs, in LPJmL, for example, while mortality due to fire is calculated on a daily time step, the background mortality and establishment of vegetation occur on an annual time step. This is particularly relevant for fires in the Southern Hemisphere or areas in the tropics where fire spread occurs at the end of the year (see, e.g. Giglio et al., 2013), because it can result in fires on 31 December that experience substantially less standing vegetation and lower litter loads than fires on 1 January. In regions where depleted fuel beds are regenerated by a monsoon season, and which therefore show a bimodal fire season

(Archibald et al., 2009), the annual establishment time step may result in substantially under-loaded fuel beds in the second half of the year relative to their real-world counterparts. Therefore, this annual time step may impact the seasonality of fires in many parts of the world and should be considered if fire seasonality is a desired model output. Future work in this direction may improve this issue by spreading mortality and establishment around the year rather than confining them to the year's end.

C3 Division of global vegetation into PFTs

The final manner in which model resolution impacts results is the division of global vegetation into individual PFTs. This is necessary as it is infeasible to model the difference in every species and sub-species of vegetation globally. However, this can pose challenges, particularly since the PFTs in DGVMs are often not chosen to reflect the differences in their response to fire (e.g. Fischer et al., 2018). For example, the boreal needleleaved evergreen PFT represents a wide variety of tree species including spruces and pines. Because spruces generally have a fundamentally different structure, with branches that extend much closer to the ground, they are more likely to experience crown scorch (e.g. Lacand et al., 2023). Further research into fire-specific PFTs may include studies of specific vegetation adaptations to fire, e.g. as discussed by Harrison et al. (2021), or the inclusion of shrub PFTs in DGVMs that do not include them, since shrubs can be an important factor in fire and post-fire dynamics (e.g. Baudena et al., 2020). To allow for the inclusion of shrubs in future we have included a (currently empty) live woody fuel category in our implementation of the Rothermel model so that shrub parameters can simply be input into the fire spread calculations.

Another example key to LPJmL-SPITFIRE is the uniform tree height and diameter for a given PFT in a given grid cell that the LPJmL DGVM operates with (Schaphoff et al., 2018a). When combined with the mortality functions of SPITFIRE that are based on tree height and diameter, this poses a challenge to accurately representing the mortality of vegetation that can be quite heterogeneously sized in actual fact but which is uniform in the model. Future developments in DGVMs that allow for heterogeneous tree sizes, as has been incorporated into LPJ-GUESS, may reduce the uncertainty in this regard, and the importance of vegetation heterogeneity should be considered when applying SPITFIRE to a desired use case.

Code and data availability. The current version of LPJmL is archived on Zenodo at <https://doi.org/10.5281/zenodo.11105506> (Schaphoff et al., 2024) under the AGPLv3 license. The exact version of the model used to produce the results used in this paper is archived on Zenodo at <https://doi.org/10.5281/zenodo.1473450> (Oberhagemann et al., 2024), along with the MATLAB implemen-

tation of the Rothermel model and input data and scripts used to produce the plots for all of the simulations presented in this paper.

Supplement. The supplement related to this article is available online at <https://doi.org/10.5194/gmd-18-2021-2025-supplement>.

Author contributions. LO conceptualized the new validation methods and implemented the prescribed fire starts validation together with WvB and MD. Analysis of literature and its relevance to the SPITFIRE model was conducted by LO with support from MB, WvB, MD, MF, SPKB, JH, and KT. JH and JRB performed data curation on the input data for the European model runs. Model code for LPJmL-SPITFIRE1.9 was written by WvB, LO, MB, and MD. LO prepared the original draft of this work. All authors participated in discussions and manuscript revisions. Additionally, MB performed the model tuning for the European model version, led the development of the phenology-based live grass moisture parametrization, and prepared Fig. 7.

Competing interests. The contact author has declared that none of the authors has any competing interests.

Disclaimer. Publisher's note: Copernicus Publications remains neutral with regard to jurisdictional claims made in the text, published maps, institutional affiliations, or any other geographical representation in this paper. While Copernicus Publications makes every effort to include appropriate place names, the final responsibility lies with the authors.

Acknowledgements. This research was performed within the "Natural Hazards and Risks in a Changing World" (NatRiskChange) research training group funded by the Deutsche Forschungsgemeinschaft (DFG). This project has received funding from the FirEURisk research project, a European Union Horizon 2020 research and innovation programme. The authors gratefully acknowledge the European Regional Development Fund (ERDF), the German Federal Ministry of Education and Research, and the State of Brandenburg for supporting this project by providing resources on the high-performance computer system at the Potsdam Institute for Climate Impact Research. The authors thank Christoph Müller, Susanne Rolinski, and Sibyll Schaphoff for the development and continued maintenance of LPJmL version 5.

Financial support. This research has been supported by the Deutsche Forschungsgemeinschaft (grant no. GRK 2043/2) and the European Union Horizon 2020 programme (grant no. 101003890).

Review statement. This paper was edited by Makoto Saito and reviewed by two anonymous referees.

References

- Albini, F. A.: Computer-based models of wildland fire behavior: a user's manual, Tech. rep., USDA Forest Service, Intermountain Forest and Range Experiment Station, Ogden, UT, https://www.frames.gov/documents/behavplus/publications/Albini_1976_FIREMOD_ocr.pdf (last access: 21 March 2025), 1976.
- Andela, N., Morton, D. C., Giglio, L., Chen, Y., Van Der Werf, G. R., Kasibhatla, P. S., DeFries, R. S., Collatz, G. J., Hantson, S., Kloster, S., Bachelet, D., Forrest, M., Lasslop, G., Li, F., Mangeon, S., Melton, J. R., Yue, C., and Randerson, J. T.: A human-driven decline in global burned area, *Science*, 356, 1356–1362, <https://doi.org/10.1126/science.aal4108>, 2017.
- Andela, N., Morton, D. C., Giglio, L., Paugam, R., Chen, Y., Hantson, S., van der Werf, G. R., and Randerson, J. T.: The Global Fire Atlas of individual fire size, duration, speed and direction, *Earth Syst. Sci. Data*, 11, 529–552, <https://doi.org/10.5194/essd-11-529-2019>, 2019.
- Andrews, P. L.: The Rothermel surface fire spread model and associated developments: A comprehensive explanation, Gen. Tech. Rep. RMRS-GTR-371, Fort Collins, CO, US Department of Agriculture, Forest Service, Rocky Mountain Research Station, 121p., 371, <https://doi.org/10.2737/RMRS-GTR-371>, 2018.
- Andrews, P. L., Cruz, M. G., and Rothermel, R. C.: Examination of the wind speed limit function in the Rothermel surface fire spread model, *Int. J. Wildland Fire*, 22, 959–969, <https://doi.org/10.1071/WF12122>, 2013.
- Aragoneses, E., García, M., Salis, M., Ribeiro, L. M., and Chuvieco, E.: Classification and mapping of European fuels using a hierarchical, multipurpose fuel classification system, *Earth Syst. Sci. Data*, 15, 1287–1315, <https://doi.org/10.5194/essd-15-1287-2023>, 2023.
- Archibald, S., Roy, D. P., Van Wilgen, B. W., and Scholes, R. J.: What limits fire? An examination of drivers of burnt area in Southern Africa, *Glob. Change Biol.*, 15, 613–630, <https://doi.org/10.1111/j.1365-2486.2008.01754.x>, 2009.
- Archibald, S., Lehmann, C. E. R., Belcher, C. M., Bond, W. J., Bradstock, R. A., Daniau, A.-L., Dexter, K. G., Forrester, E. J., Greve, M., He, T., Higgins, S. I., Hoffmann, W. A., Lamont, B. B., McGlenn, D. J., Moncrieff, G. R., Osborne, C. P., Pausas, J. G., Price, O., Ripley, B. S., Rogers, B. M., Schwilk, D. W., Simon, M. F., Turetsky, M. R., van der Werf, G. R., and Zanne, A. E.: Biological and geophysical feedbacks with fire in the Earth system, *Environ. Res. Lett.*, 13, 033003, <https://doi.org/10.1088/1748-9326/aa9ead>, 2018.
- Balch, J. K., Abatzoglou, J. T., Joseph, M. B., Koontz, M. J., Mahood, A. L., McGlinchy, J., Cattau, M. E., and Williams, A. P.: Warming weakens the night-time barrier to global fire, *Nature*, 602, 442–448, <https://doi.org/10.1038/s41586-021-04325-1>, 2022.
- Baudena, M., Santana, V. M., Baeza, M. J., Bautista, S., Eppinga, M. B., Hemerik, L., Garcia Mayor, A., Rodriguez, F., Valdecantos, A., Vallejo, V. R., Vasques, A., and Rietkerk, M.: Increased aridity drives post fire recovery of Mediterranean forests towards open shrublands, *New Phytol.*, 225, 1500–1515, <https://doi.org/10.1111/nph.16252>, 2020.
- Boulanger, Y., Pascual, J., Bouchard, M., D'Orangeville, L., Périé, C., and Girardin, M. P.: Multi model projections of tree species performance in Quebec, Canada under future climate change, *Glob. Change Biol.*, 28, 1884–1902, <https://doi.org/10.1111/gcb.16014>, 2022.
- Bowman, D. M. J. S., Balch, J. K., Artaxo, P., Bond, W. J., Carlson, J. M., Cochrane, M. A., D'Antonio, C. M., DeFries, R. S., Doyle, J. C., Harrison, S. P., Johnston, F. H., Keeley, J. E., Krawchuk, M. A., Kull, C. A., Marston, J. B., Moritz, M. A., Prentice, I. C., Roos, C. I., Scott, A. C., Swetnam, T. W., Van Der Werf, G. R., and Pyne, S. J.: Fire in the Earth system, *Science*, 324, 481–484, <https://doi.org/10.1126/science.1163886>, 2009.
- Bristiel, P., Gillespie, L., Østrem, L., Balachowski, J., Violette, C., and Volaire, F.: Experimental evaluation of the robustness of the growth–stress tolerance trade off within the perennial grass *Dactylis glomerata*, *Funct. Ecol.*, 32, 1944–1958, <https://doi.org/10.1111/1365-2435.13112>, 2018.
- Brown, J. K.: Bulk densities of nonuniform surface fuels and their application to fire modeling, *For. Sci.*, 24, 667–683, 1981.
- Brown, T. P., Hoylman, Z. H., Conrad, E., Holden, Z., Jencso, K., and Jolly, W. M.: Decoupling between soil moisture and biomass drives seasonal variations in live fuel moisture across co-occurring plant functional types, *Fire Ecol.*, 18, 14, <https://doi.org/10.1186/s42408-022-00136-5>, 2022.
- Burgan, R. E.: Estimating live fuel moisture for the 1978 National Fire Danger Rating System, Gen. Tech. Rep. INT-226, US Department of Agriculture, Forest Service, Intermountain Forest and Range Experiment Station, Ogden, UT, <https://www.frames.gov/catalog/5573> (last access: 21 March 2025), 1979.
- Carlson, J. D., Bradshaw, L. S., Nelson Jr., R. M., Bensch, R. R., and Jabrzemski, R.: Application of the Nelson model to four timelag fuel classes using Oklahoma field observations: model evaluation and comparison with national fire danger rating system algorithms, *Int. J. Wildland Fire*, 16, 204–216, <https://doi.org/10.1071/wf06073>, 2007.
- Chaste, E., Girardin, M. P., Kaplan, J. O., Portier, J., Bergeron, Y., and Hély, C.: The pyrogeography of eastern boreal Canada from 1901 to 2012 simulated with the LPJ-LMfire model, *Biogeosciences*, 15, 1273–1292, <https://doi.org/10.5194/bg-15-1273-2018>, 2018.
- Christian, H. J.: Global frequency and distribution of lightning as observed from space by the optical transient detector, *J. Geophys. Res.*, 108, 4005, <https://doi.org/10.1029/2002JD002347>, 2003.
- Chuvieco, E., González, I., Verdú, F., Aguado, I., and Yebra, M.: Prediction of fire occurrence from live fuel moisture content measurements in a Mediterranean ecosystem, *Int. J. Wildland Fire*, 18, 430–441, <https://doi.org/10.1071/WF08020>, 2009.
- Chuvieco, E., Yebra, M., Martino, S., Thonicke, K., Gómez-Giménez, M., San-Miguel, J., Oom, D., Velea, R., Mouillot, F., Molina, J. R., Miranda, A. I., Lopes, D., Salis, M., Bugaric, M., Sofiev, M., Kadantsev, E., Gitas, I. Z., Stavrakoudis, D., Eftychidis, G., Bar-Massada, A., Neidermeier, A., Pampanoni, V., Pettinari, M. L., Arrogante-Funes, F., Ochoa, C., Moreira, B., and Viegas, D.: Towards an integrated approach to wildfire risk assessment: when, where, what and how may the landscapes burn, *Fire*, 6, 215, <https://doi.org/10.3390/fire6050215>, 2023.
- Collin, A., Bernardin, D., and Séro-Guillaume, O.: A physical-based cellular automaton model for forest-fire propagation, *Combust. Sci. Technol.*, 183, 347–369, <https://doi.org/10.1080/00102202.2010.508476>, 2011.

- Cocchi, M., Weedon, G. P., Amici, A., Bellouin, N., Lange, S., Müller Schmied, H., Hersbach, H., and Buontempo, C.: WFDE5: bias-adjusted ERA5 reanalysis data for impact studies, *Earth Syst. Sci. Data*, 12, 2097–2120, <https://doi.org/10.5194/essd-12-2097-2020>, 2020.
- Drüke, M., Forkel, M., von Bloh, W., Sakschewski, B., Cardoso, M., Bustamante, M., Kurths, J., and Thonicke, K.: Improving the LPJmL4-SPITFIRE vegetation–fire model for South America using satellite data, *Geosci. Model Dev.*, 12, 5029–5054, <https://doi.org/10.5194/gmd-12-5029-2019>, 2019.
- Drüke, M., Sakschewski, B., von Bloh, W., Billing, M., Lucht, W., and Thonicke, K.: Fire may prevent future Amazon forest recovery after large-scale deforestation, *Communications Earth and Environment*, 4, 1–10, <https://doi.org/10.1038/s43247-023-00911-5>, 2023.
- Emmett, K. D., Renwick, K. M., and Poulter, B.: Adapting a dynamic vegetation model for regional biomass, plant biogeography, and fire modeling in the greater Yellowstone ecosystem: evaluating LPJ-GUESS-LMfireCF, *Ecol. Model.*, 440, 109417, <https://doi.org/10.1016/j.ecolmodel.2020.109417>, 2021.
- Ephrath, J. E., Goudriaan, J., and Marani, A.: Modelling diurnal patterns of air temperature, radiation wind speed and relative humidity by equations from daily characteristics, *Agr. Syst.*, 51, 377–393, [https://doi.org/10.1016/0308-521X\(95\)00068-G](https://doi.org/10.1016/0308-521X(95)00068-G), 1996.
- Felsberg, A., Kloster, S., Wilkenskeld, S., Krause, A., and Lasslop, G.: Lightning forcing in global fire models: the importance of temporal resolution, *J. Geophys. Res.-Biogeo.*, 123, 168–177, <https://doi.org/10.1002/2017JG004080>, 2018.
- Finney, M. A.: FARSITE: Fire Area Simulator-model development and evaluation, Res. Pap. RMRS-RP-4, Revised 2004, Ogden, UT, US Department of Agriculture, Forest Service, Rocky Mountain Research Station. 47p., 4, <https://doi.org/10.2737/RMRS-RP-4>, 1998.
- Finney, M. A.: An overview of FlamMap fire modeling capabilities, in: Fuels Management-How to Measure Success: Conference Proceedings, USDA Forest Service, Rocky Mountain Research Station, Portland, OR, 213–220, <https://research.fs.usda.gov/treearch/25948> (last access: 21 March 2025), 2006.
- Fischer, R., Rödiger, E., and Huth, A.: Consequences of a reduced number of plant functional types for the simulation of forest productivity, *Forests*, 9, 460, <https://doi.org/10.3390/f9080460>, 2018.
- Forkel, M., Carvalhais, N., Schaphoff, S., v. Bloh, W., Migliavacca, M., Thurner, M., and Thonicke, K.: Identifying environmental controls on vegetation greenness phenology through model–data integration, *Biogeosciences*, 11, 7025–7050, <https://doi.org/10.5194/bg-11-7025-2014>, 2014.
- Forkel, M., Andela, N., Harrison, S. P., Lasslop, G., van Marle, M., Chuvieco, E., Dorigo, W., Forrest, M., Hantson, S., Heil, A., Li, F., Melton, J., Sitch, S., Yue, C., and Arneth, A.: Emergent relationships with respect to burned area in global satellite observations and fire-enabled vegetation models, *Biogeosciences*, 16, 57–76, <https://doi.org/10.5194/bg-16-57-2019>, 2019a.
- Forkel, M., Drüke, M., Thurner, M., Dorigo, W., Schaphoff, S., Thonicke, K., von Bloh, W., and Carvalhais, N.: Constraining modelled global vegetation dynamics and carbon turnover using multiple satellite observations, *Sci. Rep.-UK*, 9, 18757, <https://doi.org/10.1038/s41598-019-55187-7>, 2019b.
- Forkel, M., Schmidt, L., Zotta, R.-M., Dorigo, W., and Yebra, M.: Estimating leaf moisture content at global scale from passive microwave satellite observations of vegetation optical depth, *Hydro. Earth Syst. Sci.*, 27, 39–68, <https://doi.org/10.5194/hess-27-39-2023>, 2023.
- Giglio, L., Randerson, J. T., and Van Der Werf, G. R.: Analysis of daily, monthly, and annual burned area using the fourth generation global fire emissions database (GFED4), *J. Geophys. Res.-Biogeo.*, 118, 317–328, <https://doi.org/10.1002/jgrg.20042>, 2013.
- Hantson, S., Lasslop, G., Kloster, S., and Chuvieco, E.: Anthropogenic effects on global mean fire size, *Int. J. Wildland Fire*, 24, 589–596, <https://doi.org/10.1071/WF14208>, 2015.
- Hantson, S., Arneth, A., Harrison, S. P., Kelley, D. I., Prentice, I. C., Rabin, S. S., Archibald, S., Mouillot, F., Arnold, S. R., Artaxo, P., Bachelet, D., Ciais, P., Forrest, M., Friedlingstein, P., Hickler, T., Kaplan, J. O., Kloster, S., Knorr, W., Lasslop, G., Li, F., Manguon, S., Melton, J. R., Meyn, A., Sitch, S., Spessa, A., van der Werf, G. R., Voulgarakis, A., and Yue, C.: The status and challenge of global fire modelling, *Biogeosciences*, 13, 3359–3375, <https://doi.org/10.5194/bg-13-3359-2016>, 2016.
- Hantson, S., Kelley, D. I., Arneth, A., Harrison, S. P., Archibald, S., Bachelet, D., Forrest, M., Hickler, T., Lasslop, G., Li, F., Manguon, S., Melton, J. R., Nieradzick, L., Rabin, S. S., Prentice, I. C., Sheehan, T., Sitch, S., Teckentrup, L., Voulgarakis, A., and Yue, C.: Quantitative assessment of fire and vegetation properties in simulations with fire-enabled vegetation models from the Fire Model Intercomparison Project, *Geosci. Model Dev.*, 13, 3299–3318, <https://doi.org/10.5194/gmd-13-3299-2020>, 2020.
- Harrison, S. P., Prentice, I. C., Bloomfield, K. J., Dong, N., Forkel, M., Forrest, M., Ningthoujam, R. K., Pellegrini, A., Shen, Y., Baudena, M., Cardoso, A. W., Huss, J. C., Joshi, J., Oliveras, I., Pausas, J. G., and Simpson, K. J.: Understanding and modelling wildfire regimes: an ecological perspective, *Environ. Res. Lett.*, 16, 125008, <https://doi.org/10.1088/1748-9326/ac39be>, 2021.
- He, Y., Monahan, A. H., and McFarlane, N. A.: Diurnal variations of land surface wind speed probability distributions under clear sky and low cloud conditions, *Geophys. Res. Lett.*, 40, 3308–3314, <https://doi.org/10.1002/grl.50575>, 2013.
- Hourdin, F., Mauritsen, T., Gettelman, A., Golaz, J.-C., Balaji, V., Duan, Q., Folini, D., Ji, D., Klocke, D., Qian, Y., Rauser, F., Rio, C., Tomassini, L., Watanabe, M., and Williamson, D.: The art and science of climate model tuning, *B. Am. Meteorol. Soc.*, 98, 589–602, <https://doi.org/10.1175/BAMS-D-15-00135.1>, 2017.
- Jolly, W. M., Nemani, R., and Running, S. W.: A generalized, bioclimatic index to predict foliar phenology in response to climate, *Glob. Change Biol.*, 11, 619–632, <https://doi.org/10.1111/j.1365-2486.2005.00930.x>, 2005.
- Jung, C. and Schindler, D.: Integration of small-scale surface properties in a new high resolution global wind speed model, *Energ. Convers. Manage.*, 210, 112733, <https://doi.org/10.1016/j.enconman.2020.112733>, 2020.
- Kaplan, J. O. and Lau, K. H.-K.: The WGLC global gridded lightning climatology and time series, *Earth Syst. Sci. Data*, 13, 3219–3237, <https://doi.org/10.5194/essd-13-3219-2021>, 2021.
- Kaplan, J. O., Pfeiffer, M., Kolen, J. C. A., and Davis, B. A. S.: Large scale anthropogenic reduction of forest cover in

- last glacial maximum Europe, *PLoS One*, 11, e0166726, <https://doi.org/10.1371/journal.pone.0166726>, 2016.
- Keep, T., Sampoux, J., Barre, P., Blanco Pastor, J., Dehmer, K. J., Durand, J., Hegarty, M., Ledauphin, T., Muylle, H., Roldán Ruiz, I., Ruttink, T., Surault, F., Willner, E., and Volaire, F.: To grow or survive: Which are the strategies of a perennial grass to face severe seasonal stress?, *Funct. Ecol.*, 35, 1145–1158, <https://doi.org/10.1111/1365-2435.13770>, 2021.
- Kelley, D. I., Harrison, S. P., and Prentice, I. C.: Improved simulation of fire–vegetation interactions in the Land surface Processes and eXchanges dynamic global vegetation model (LPX-Mv1), *Geosci. Model Dev.*, 7, 2411–2433, <https://doi.org/10.5194/gmd-7-2411-2014>, 2014.
- Klein Goldewijk, K., Beusen, A., and Janssen, P.: Long-term dynamic modeling of global population and built-up area in a spatially explicit way: HYDE 3.1, *Holocene*, 20, 565–573, <https://doi.org/10.1177/0959683609356587>, 2010.
- Kovesi, P.: Good Colour Maps: How to Design Them, Tech. Rep., arXiv [preprint], <https://doi.org/10.48550/arXiv.1509.03700>, 2015.
- Krueger, E. S., Levi, M. R., Achieng, K. O., Bolten, J. D., Carlson, J. D., Coops, N. C., Holden, Z. A., Magi, B. I., Rigden, A. J., Ochsner, T. E., Krueger, E. S., Levi, M. R., Achieng, K. O., Bolten, J. D., Carlson, J. D., Coops, N. C., Holden, Z. A., Magi, B. I., Rigden, A. J., and Ochsner, T. E.: Using soil moisture information to better understand and predict wildfire danger: a review of recent developments and outstanding questions, *Int. J. Wildland Fire*, 32, 111–132, <https://doi.org/10.1071/WF22056>, 2022.
- Lacand, M., Asselin, H., Magne, G., Aakala, T., Remy, C. C., Seppä, H., and Ali, A. A.: Multimillennial fire history of northern Finland along a latitude/elevation gradient, *Quaternary Sci. Rev.*, 312, 108171, <https://doi.org/10.1016/j.quascirev.2023.108171>, 2023.
- Lasslop, G., Thonicke, K., and Kloster, S.: SPITFIRE within the MPI Earth system model: model development and evaluation, *J. Adv. Model. Earth Sy.*, 6, 740–755, <https://doi.org/10.1002/2013MS000284>, 2014.
- Lasslop, G., Hantson, S., Harrison, S. P., Bachelet, D., Burton, C., Forkel, M., Forrest, M., Li, F., Melton, J. R., Yue, C., Archibald, S., Scheiter, S., Arneth, A., Hickler, T., and Sitch, S.: Global ecosystems and fire: multi model assessment of fire induced tree cover and carbon storage reduction, *Glob. Change Biol.*, 26, 5027–5041, <https://doi.org/10.1111/gcb.15160>, 2020.
- Lehsten, V., Tansey, K., Balzter, H., Thonicke, K., Spessa, A., Weber, U., Smith, B., and Arneth, A.: Estimating carbon emissions from African wildfires, *Biogeosciences*, 6, 349–360, <https://doi.org/10.5194/bg-6-349-2009>, 2009.
- Lehsten, V., Arneth, A., Spessa, A., Thonicke, K., Moustakas, A., Lehsten, V., Arneth, A., Spessa, A., Thonicke, K., and Moustakas, A.: The effect of fire on tree–grass coexistence in savannas: a simulation study, *Int. J. Wildland Fire*, 25, 137–146, <https://doi.org/10.1071/WF14205>, 2015.
- Li, F., Val Martin, M., Andreae, M. O., Arneth, A., Hantson, S., Kaiser, J. W., Lasslop, G., Yue, C., Bachelet, D., Forrest, M., Kluzek, E., Liu, X., Mangeon, S., Melton, J. R., Ward, D. S., Darnenov, A., Hickler, T., Ichoku, C., Magi, B. I., Sitch, S., van der Werf, G. R., Wiedinmyer, C., and Rabin, S. S.: Historical (1700–2012) global multi-model estimates of the fire emissions from the Fire Modeling Intercomparison Project (FireMIP), *Atmos. Chem. Phys.*, 19, 12545–12567, <https://doi.org/10.5194/acp-19-12545-2019>, 2019.
- Lutz, F., Herzfeld, T., Heinke, J., Rolinski, S., Schaphoff, S., von Bloh, W., Stoorvogel, J. J., and Müller, C.: Simulating the effect of tillage practices with the global ecosystem model LPJmL (version 5.0-tillage), *Geosci. Model Dev.*, 12, 2419–2440, <https://doi.org/10.5194/gmd-12-2419-2019>, 2019.
- McCarty, J. L., Aalto, J., Paunu, V.-V., Arnold, S. R., Eckhardt, S., Klimont, Z., Fain, J. J., Evangeliou, N., Venäläinen, A., Tchebakova, N. M., Parfenova, E. I., Kupiainen, K., Soja, A. J., Huang, L., and Wilson, S.: Reviews and syntheses: Arctic fire regimes and emissions in the 21st century, *Biogeosciences*, 18, 5053–5083, <https://doi.org/10.5194/bg-18-5053-2021>, 2021.
- Mendiguren, G., Pilar Martín, M., Nieto, H., Pacheco-Labrador, J., and Jurdao, S.: Seasonal variation in grass water content estimated from proximal sensing and MODIS time series in a Mediterranean Fluxnet site, *Biogeosciences*, 12, 5523–5535, <https://doi.org/10.5194/bg-12-5523-2015>, 2015.
- Morin, K. and Davis, J. L.: Cross-validation: What is it and how is it used in regression?, *Commun. Stat. Theory*, 46, 5238–5251, <https://doi.org/10.1080/03610926.2015.1099672>, 2017.
- Muñoz-Sabater, J., Dutra, E., Agustí-Panareda, A., Albergel, C., Arduini, G., Balsamo, G., Boussetta, S., Choulga, M., Harrigan, S., Hersbach, H., Martens, B., Miralles, D. G., Piles, M., Rodríguez-Fernández, N. J., Zsoter, E., Buontempo, C., and Thépaut, J.-N.: ERA5-Land: a state-of-the-art global reanalysis dataset for land applications, *Earth Syst. Sci. Data*, 13, 4349–4383, <https://doi.org/10.5194/essd-13-4349-2021>, 2021.
- Narayananaraj, G. and Wimberly, M. C.: Influences of forest roads on the spatial pattern of wildfire boundaries, *Int. J. Wildland Fire*, 20, 792–803, <https://doi.org/10.1071/WF10032>, 2011.
- Nelson, R. M.: Prediction of diurnal change in 10-h fuel stick moisture content, *Can. J. Forest Res.*, 30, 1071–1087, <https://doi.org/10.1139/x00-032>, 2000.
- Oberhagemann, L., Billing, M., von Bloh, W., Drueke, M., Forrest, M., Bowering, S. P. K., Hetzer, J., Ribalaygua Batalla, J., and Thonicke, K.: Model Code and Data for “Sources of Uncertainty in the Global Fire Model SPITFIRE: Development of LPJmL-SPITFIRE1.9 and Directions for Future Improvements” (Version Version 1), Zenodo [data set], <https://doi.org/10.5281/zenodo.11473450>, 2024.
- Pechony, O. and Shindell, D. T.: Fire parameterization on a global scale, *J. Geophys. Res.-Atmos.*, 114, D16115, <https://doi.org/10.1029/2009JD011927>, 2009.
- Perkins, O., Matej, S., Erb, K., and Millington, J.: Towards a global behavioural model of anthropogenic fire: the spatiotemporal distribution of land-fire systems, *Socio-Environmental Systems Modelling*, 4, 18130–18130, <https://doi.org/10.18174/sesmo.18130>, 2022.
- Peterson, D. L. and Ryan, K. C.: Modeling postfire conifer mortality for long-range planning, *Environ. Manage.*, 10, 797–808, <https://doi.org/10.1007/BF01867732>, 1986.
- Pfeiffer, M., Spessa, A., and Kaplan, J. O.: A model for global biomass burning in preindustrial time: LPJ-LMfire (v1.0), *Geosci. Model Dev.*, 6, 643–685, <https://doi.org/10.5194/gmd-6-643-2013>, 2013.
- Potter, B. E.: Atmospheric interactions with wildland fire behaviour – I. Basic surface interactions, vertical profiles

- and synoptic structures, *Int. J. Wildland Fire*, 21, 779–801, <https://doi.org/10.1071/WF11128>, 2012.
- Prentice, I. C., Kelley, D. I., Foster, P. N., Friedlingstein, P., Harrison, S. P., and Bartlein, P. J.: Modeling fire and the terrestrial carbon balance, *Global Biogeochem. Cy.*, 25, GB3005, <https://doi.org/10.1029/2010GB003906>, 2011.
- Rabin, S. S., Melton, J. R., Lasslop, G., Bachelet, D., Forrest, M., Hantson, S., Kaplan, J. O., Li, F., Mangeon, S., Ward, D. S., Yue, C., Arora, V. K., Hickler, T., Kloster, S., Knorr, W., Nieradzik, L., Spessa, A., Folberth, G. A., Sheehan, T., Voulgarakis, A., Kelley, D. I., Prentice, I. C., Sitch, S., Harrison, S., and Arneth, A.: The Fire Modeling Intercomparison Project (FireMIP), phase 1: experimental and analytical protocols with detailed model descriptions, *Geosci. Model Dev.*, 10, 1175–1197, <https://doi.org/10.5194/gmd-10-1175-2017>, 2017.
- Randerson, J. T., Van Der Werf, G. R., Giglio, L., Collatz, G. J., and Kasibhatla, P. S.: Global Fire Emissions Database, Version 4.1 (GFEDv4), ORNL DAAC, <https://doi.org/10.3334/ORNLDAAC/1293>, 2015.
- Rothermel, R. C.: A Mathematical Model for Predicting Fire Spread in Wildland Fuels, Intermountain Forest and Range Experiment Station, Forest Service, US Department of Agriculture, https://www.fs.usda.gov/rm/pubs_int/int_rp115.pdf (last access: 21 March 2025), 1972.
- Schaphoff, S., von Bloh, W., Rammig, A., Thonicke, K., Biemans, H., Forkel, M., Gerten, D., Heinke, J., Jägermeyr, J., Knauer, J., Langerwisch, F., Lucht, W., Müller, C., Rolinski, S., and Waha, K.: LPJmL4 – a dynamic global vegetation model with managed land – Part 1: Model description, *Geosci. Model Dev.*, 11, 1343–1375, <https://doi.org/10.5194/gmd-11-1343-2018>, 2018a.
- Schaphoff, S., von Bloh, W., Thonicke, K., Biemans, H., Forkel, M., Gerten, D., Heinke, J., Jägermeyr, J., Müller, C., Rolinski, S., Waha, K., Stehfest, E., de Waal, L., Heyder, U., Gumpenberger, M., and Beringer, T.: LPJmL4 Model Code, Climate Institute for Climate Impact Research, <https://doi.org/10.5880/PIK.2018.002>, 2018b.
- Schaphoff, S., von Bloh, W., Rammig, A., Thonicke, K., Biemans, H., Forkel, M., Gerten, D., Heinke, J., Jägermeyr, J., Langerwisch, F., Lucht, W., Rolinski, S., Waha, K., Ostberg, S., Wirth, S. B., Fader, M., Drüke, M., Jans, Y., Lutz, F., Herzfeld, T., Minoli, S., Porwollik, V., Stehfest, E., de Waal, L., Beringer (Erbrecht), T., Rost (Jachner), S., Gumpenberger, M., Heyder, U., Werner, C., Braun, J., Breier, J., Stenzel, F., Mathesius, S., Hemmen, M., Billing, M., Oberhagemann, L., Sakschewski, B., and Müller, C.: LPJmL: central open-source github repository of LPJmL at PIK, Zenodo [code], <https://doi.org/10.5281/zenodo.11105506>, 2024.
- Scheiter, S., Langan, L., and Higgins, S. I.: Next generation dynamic global vegetation models: learning from community ecology, *New Phytol.*, 198, 957–969, <https://doi.org/10.1111/nph.12210>, 2013.
- Scott, J. H. and Burgan, R. E.: Standard fire behavior fuel models: a comprehensive set for use with Rothermel's surface fire spread model, Gen. Tech. Rep. RMRS-GTR-153, Fort Collins, CO, US Department of Agriculture, Forest Service, Rocky Mountain Research Station, 72p., 153, <https://doi.org/10.2737/RMRS-GTR-153>, 2005.
- Senande-Rivera, M., Insua-Costa, D., and Miguez-Macho, G.: Towards an atmosphere more favourable to firestorm development in Europe, *Environ. Res. Lett.*, 17, 094015, <https://doi.org/10.1088/1748-9326/ac85ce>, 2022.
- Sharples, J. J.: An overview of mountain meteorological effects relevant to fire behaviour and bushfire risk, *Int. J. Wildland Fire*, 18, 737–754, <https://doi.org/10.1071/WF08041>, 2009.
- Sjöström, J. and Granström, A.: A phenology-driven fire danger index for northern grasslands, *Int. J. Wildland Fire*, 32, 1332–1346, <https://doi.org/10.1071/WF23013>, 2023.
- Teckentrup, L., Harrison, S. P., Hantson, S., Heil, A., Melton, J. R., Forrest, M., Li, F., Yue, C., Arneth, A., Hickler, T., Sitch, S., and Lasslop, G.: Response of simulated burned area to historical changes in environmental and anthropogenic factors: a comparison of seven fire models, *Biogeosciences*, 16, 3883–3910, <https://doi.org/10.5194/bg-16-3883-2019>, 2019.
- The MathWorks Inc.: MATLAB version: 9.10.0 (R2021b), <https://www.mathworks.com> (last access: 21 March 2025), 2021.
- Thonicke, K., Spessa, A., Prentice, I. C., Harrison, S. P., Dong, L., and Carmona-Moreno, C.: The influence of vegetation, fire spread and fire behaviour on biomass burning and trace gas emissions: results from a process-based model, *Biogeosciences*, 7, 1991–2011, <https://doi.org/10.5194/bg-7-1991-2010>, 2010.
- Venevsky, S., Thonicke, K., Sitch, S., and Cramer, W.: Simulating fire regimes in human-dominated ecosystems: Iberian Peninsula case study, *Glob. Change Biol.*, 8, 984–998, <https://doi.org/10.1046/j.1365-2486.2002.00528.x>, 2002.
- von Bloh, W., Schaphoff, S., Müller, C., Rolinski, S., Waha, K., and Zaehle, S.: Implementing the nitrogen cycle into the dynamic global vegetation, hydrology, and crop growth model LPJmL (version 5.0), *Geosci. Model Dev.*, 11, 2789–2812, <https://doi.org/10.5194/gmd-11-2789-2018>, 2018a.
- von Bloh, W., Schaphoff, S., Müller, C., Rolinski, S., Waha, K., and Zaehle, S.: LPJmL5 Model Code, Climate Institute for Climate Impact Research, <https://doi.org/10.5880/PIK.2018.011>, 2018b.
- Ward, D. S., Shevliakova, E., Malyshev, S., and Rabin, S.: Trends and variability of global fire emissions due to historical anthropogenic activities, *Global Biogeochem. Cy.*, 32, 122–142, <https://doi.org/10.1002/2017GB005787>, 2018.
- Weise, D. R., Koo, E., Zhou, X., Mahalingam, S., Morandini, F., and Balbi, J.-H.: Fire spread in chaparral – a comparison of laboratory data and model predictions in burning live fuels, *Int. J. Wildland Fire*, 25, 980–994, <https://doi.org/10.1071/WF15177>, 2016.
- Wirth, S. B., Braun, J., Heinke, J., Ostberg, S., Rolinski, S., Schaphoff, S., Stenzel, F., von Bloh, W., and Müller, C.: Biological nitrogen fixation of natural and agricultural vegetation simulated with LPJmL 5.7.9, EGU sphere [preprint], <https://doi.org/10.5194/egusphere-2023-2946>, 2024.
- Wu, M., Knorr, W., Thonicke, K., Schurgers, G., Camia, A., and Arneth, A.: Sensitivity of burned area in Europe to climate change, atmospheric CO₂ levels, and demography: a comparison of two fire-vegetation models, *J. Geophys. Res.-Bioge.*, 120, 2256–2272, <https://doi.org/10.1002/2015JG003036>, 2015.
- Yeber, M., Scortechini, G., Badi, A., Beget, M. E., Boer, M. M., Bradstock, R., Chuvieco, E., Danson, F. M., Dennison, P., Resco de Dios, V., Di Bella, C. M., Forsyth, G., Frost, P., Garcia, M., Hamdi, A., He, B., Jolly, M., Kraaij, T., Martín, M. P., Mouillot, F., Newnham, G., Nolan, R. H., Pellizzaro, G., Qi, Y., Quan, X., Riaño, D., Roberts, D., Sow, M., and Ustin, S.: Globe-LFMC, a global plant water status database for vegetation eco-

- physiology and wildfire applications, *Scientific Data*, 6, 155, <https://doi.org/10.1038/s41597-019-0164-9>, 2019.
- Yue, C., Ciais, P., Cadule, P., Thonicke, K., Archibald, S., Poulter, B., Hao, W. M., Hantson, S., Mouillot, F., Friedlingstein, P., Maignan, F., and Viovy, N.: Modelling the role of fires in the terrestrial carbon balance by incorporating SPITFIRE into the global vegetation model ORCHIDEE – Part 1: simulating historical global burned area and fire regimes, *Geosci. Model Dev.*, 7, 2747–2767, <https://doi.org/10.5194/gmd-7-2747-2014>, 2014.
- Yue, C., Ciais, P., Cadule, P., Thonicke, K., and van Leeuwen, T. T.: Modelling the role of fires in the terrestrial carbon balance by incorporating SPITFIRE into the global vegetation model ORCHIDEE – Part 2: Carbon emissions and the role of fires in the global carbon balance, *Geosci. Model Dev.*, 8, 1321–1338, <https://doi.org/10.5194/gmd-8-1321-2015>, 2015.