



Supplement of

Soil nitrous oxide emissions from global land ecosystems and their drivers within the LPJ-GUESS model (v4.1)

Jianyong Ma et al.

Correspondence to: Jianyong Ma (jianyong.ma@kit.edu) and Stefan Olin (stefan.olin@nateko.lu.se)

The copyright of individual parts of the supplement might differ from the article licence.

Table S1. Literature-based natural vegetation sites used for model soil N₂O evaluation: Geographic location; Dominant species; Experimental period; Observed N₂O fluxes at the studied sites^a.

Geographic location	Lon	Lat	Vegetation types	Dominant species	Period of N ₂ O measurement	Observed N ₂ O fluxes (kg N ha ⁻¹ yr ⁻¹) ^b	Reference
Mooloolah (Australia)	152.93	-26.76	Subtropical rainforest	-	Mar.2007-Feb.2009	0.511±0.118 (2007/2008) 0.521±0.006 (2008/2009)	Rowlings et al. (2012)
Klausen-Leopoldsdorf (Austria)	16.05	48.12	Temperate deciduous broadleaf forest	Beech	May 2002-Apr.2004	0.538±0.031 (2002/2003) 0.671±0.021 (2003/2004)	Kitzler et al. (2006)
Achenkirch (Austria)	11.71	47.49	Temperate needleleaf forest	Spruce	Entire 2002-2003	0.402±0.036 (2002) 0.292±0.036 (2003)	Kesik et al. (2005)
Innsbruck (Austria)	11.45	-47.30	Temperate needleleaf forest	Spruce	Jul.1990-Jun.1991	0.081±0.008	Henrich and Haselwandter (1997)
Schottenwald (Austria)	16.25	48.23	Temperate deciduous broadleaf forest	Beech	May 2002-Apr.2004	0.76±0.003 (2002/2003) 0.82±0.005 (2003/2004)	Kitzler et al. (2006)
Manaus (Brazil)	-59.96	-2.89	Tropical rainforest	-	Apr.1987-Apr.1988	1.90	Luizão et al. (1989)
Tapajós National Forest (Brazil)	-54.95	-2.89	Tropical rainforest	-	Entire 2000-2004	2.10±0.70 (5-year average)	Davidson et al. (2008)
Paragominas (Brazil)	-47.52	-2.98	Tropical rainforest	-	Entire 1995-1996	2.43±0.22 (2-year average)	Verchot et al. (1999)
Ontario (Canada)	-80.67	49.05	Boreal needleleaf forest	Black spruce	Jun.1992-Aug.1992	0.044±0.067	Schiller and Hastie (1996)
Saskatchewan 1 (Canada)	-106.20	53.62	Boreal deciduous broadleaf forest	Trembling aspen	Apr.1994-Sep.1994	0.281±0.101	Simpson et al. (1997)
Saskatchewan 1 (Canada)	-106.20	53.62	Boreal deciduous broadleaf forest	Trembling aspen	May 2006-Oct.2006 May 2007-Oct.2007	0.12 (2006) 0.14 (2007)	Matson et al. (2009)
Saskatchewan 2 (Canada)	-105.12	53.99	Boreal needleleaf forest	Black spruce	May 2006-Oct.2006 May 2007-Oct.2007	0.13 (2006) -0.02 (2007)	Matson et al. (2009)
Saskatchewan 3 (Canada)	-104.69	53.92	Boreal needleleaf forest	Jack pine	May 2006-Oct.2006 May 2007-Oct.2007	0.05 (2006) 0.08 (2007)	Matson et al. (2009)

Mont St. Hilaire (Canada)	-73.13	45.52	Temperate deciduous broadleaf forest	American beech	May 2006- May 2008	0.307±0.081 (3-year average)	Ullah and Moore (2011)
Tibetan Plateau (China)	93.05	35.13	Alpine grassland	Stipa purpurea	Aug.2000-Jul.2001	0.044	Pei et al. (2004)
Changbai mountain (China)	126.92	41.38	Temperate mixed forest	Amur lime; Korean pine	Jun.1994-Oct.1995	0.557±0.483	Chen et al. (2000)
Inner Mongolia (China)	114.67	43.53	Temperate semi-arid grassland	Chinese rye grass	Entire 1995, 1998, 2001- 2003	0.58±0.16 (1995) 0.25±0.12 (1998) 1.62±0.53 (2001) 0.61±0.53 (2002) 0.58±0.36 (2003)	Du et al. (2006)
Xishuangbanna (China)	101.26	21.93	Tropical seasonal rainforest	Pometia tomentosa	Entire 2003	2.59±0.29	Yan et al. (2008)
Sorø (Denmark)	11.59	55.39	Temperate deciduous broadleaf forest	Beech	Entire 2003	0.548±0.036	Kesik et al. (2005)
Bombuscaro (Ecuador)	-78.97	-4.10	Tropical rainforest	Moraceae	May 2008-May 2009	0.279±0.119	Wolf et al. (2011)
Reserva San Francisco (Ecuador)	-79.07	-3.97	Tropical rainforest	Lauraceae	May 2008-May 2009	0.566±0.148	Wolf et al. (2011)
Padasjoki (Finland)	24.97	61.32	Boreal needleleaf forest	Picea abies	Jun.-Aug. in 2000, 2001, 2003	0.197±0.056 (3-year average)	Maljanen et al. (2006)
Kiel (Germany)	10.12	54.26	Temperate deciduous broadleaf forest	European beech	Entire 1993	0.40	Mogge et al. (1998)
Wildmooswald (Germany)	8.12	47.12	Temperate needleleaf forest	Norway spruce	Entire 2001-2002	0.675±0.359 (2001) 0.475±0.457 (2002)	Jungkunst et al. (2004)
Höglwald (Germany)	11.28	48.83	Temperate needleleaf forest	Norway spruce	Entire 1994-1995, 1997, 2000-2005, 2007-2009	0.4 (1994), 0.8 (1995) 0.7 (1997), 0.5 (2000) 0.4 (2001), 0.7 (2002) 0.4 (2003), 0.2 (2004) 1.0 (2005), 0.6 (2007) 0.8 (2008), 1.0 (2009)	Luo et al. (2012)
Jambi (Indonesia)	102.11	-1.12	Tropical rainforest	-	Sep.1997-Aug.1998	0.259±0.204	Ishizuka et al. (2002)
Puncak Dingin (Indonesia)	120.30	-1.27	Tropical rainforest	-	Nov.2002-Oct.2003	1.01±0.16	Purbopuspito et al. (2006)

Rorekatimbu (Indonesia)	120.30	-1.32	Tropical rainforest	-	Nov.2002-Oct.2003	0.29±0.02	Purbopuspito et al. (2006)
Wuasa (Indonesia)	120.30	-1.42	Tropical rainforest	-	Nov.2002-Oct.2003	1.11±0.17	Purbopuspito et al. (2006)
Palangka Raya (Indonesia)	114.03	-2.35	Tropical deciduous forest	Tetramerista glabra	Jun.2002-May 2004	2.51 (2-year average)	Takakai et al. (2006)
Parco Ticino (Italy)	8.74	45.36	Temperate deciduous broadleaf forest	Hardwoods	Entire 2002-2003	0.511±0.109 (2002) 0.183±0.001 (2003)	Kesik et al. (2005)
Sarawak (Malaysia)	111.85	2.82	Tropical rainforest	Gonystylus bancanus	Aug.2002-Jul.2003	0.70	Melling et al. (2007)
Christchurch (New Zealand)	171.68	-43.13	Temperate evergreen broadleaf forest	Black beech	Jan.1998-Apr.1999	0.01	Price et al. (2004)
Åmli (Norway)	8.5	58.82	Temperate evergreen needleleaf forest	Scots pine	Jul.1992-Aug.1992	0.124±0.031	Sitaula et al. (1995)
Gigante Peninsula (Panama)	-79.83	9.10	Tropical evergreen forests	Oenocarpus mapora	Entire 2006-2007	1.35±0.09 (2006) 0.98±0.10 (2007)	Koehler et al. (2009)
Gårdsjön (Sweden)	12.02	58.07	Temperate needleleaf forest	Norway spruce	Entire 1993-1994	0.095±0.024 (2-year average)	Klemedtsson et al. (1997)
Northumberland (UK)	-2.05	55.17	Grassland	-	Entire 2001-2002	0.191±0.064 (2001) 0.191±0.064 (2002)	Ball et al. (2007)
Harvard forest (USA)	-72.19	42.53	Temperate deciduous broadleaf forest	Black oak	May 1988-Apr.1989	0.017±0.017	Bowden et al. (1990)
Ithaca, New York (USA)	-76.45	42.44	Grassland	Timothy weed	May 1980-Apr.1981	0.9	Duxbury et al. (1982)
White Mountain National Forest (USA)	-71.75	43.91	Temperate deciduous broadleaf forest	American beech; Sugar maple	Entire 1998-2000	1.25±0.13 (1998) 0.55±0.04 (1999) 0.67±0.24 (2000)	Groffman et al. (2006)
Mercury, Nevada (USA)	-115.92	36.82	Arid grassland	Pleuraphis rigida	Entire 1999-2000	0.091±0.066 (2-year average)	Billings et al. (2002)
Saratoga (USA)	-106.85	41.44	Temperate semi-arid grassland	Festuca idahoensis	Entire 1986-1987	0.255 (1986) 0.280 (1987)	Matson et al. (1991)
Whiteface Mt., New York (USA)	-73.90	44.40	Temperate needleleaf forest	Spruce-fir	May 1990-Sep.1990	0.185	Castro et al. (1993)

Mt. Mansfield, Vermont (USA)	-72.80	44.50	Temperate needleleaf forest	Spruce-fir	May 1990-Sep.1990	0.171	Castro et al. (1993)
Acadia, Maine (USA)	-68.40	44.30	Temperate needleleaf forest	Spruce-fir	May 1990-Sep.1990	0.032	Castro et al. (1993)
Fort Collins (USA)	-104.75	40.80	Grassland	Blue gramma	Entire 1991-1994	0.130±0.017 (1991) 0.118±0.019 (1992) 0.177±0.049 (1993) 0.127±0.035 (1994)	Mosier et al. (1996)
Baraboo (USA)	-89.67	43.44	Grassland	Little bluestem	Sep.1981-Mar.1982	0.028	Cates and Keeney (1987)
Santa Barbara (Venezuela)	-63.45	9.75	Tropical herbaceous savanna	Trachypogon	Apr.2000-Mar.2001	0.913±0.073	Simona et al. (2004)
Harare (Zimbabwe)	30.91	-17.68	Subtropical woody savanna	Julbernardia globiflora	Oct.2000-Sep.2001	0.497±0.223	Rees et al. (2006)
Ankasa Conservation Area (Ghana)	-2.68	5.27	Tropical rainforest	-	Entire 2009-2010	2.33±0.20 (2-year average)	Castaldi et al. (2013)
Kakamega Forest National Park (Kenya)	34.86	0.25	Tropical rainforest	-	Apr.2004-Jun.2004	2.60±1.20	Werner et al. (2007)
Fukushima (Japan)	140.33	37.35	Temperate evergreen needleleaf forest	Japanese cedar	Entire 2003-2004	0.872 (2003) 1.077 (2004)	Morishita et al. (2007)
Oita (Japan)	131.28	33.18	Temperate evergreen needleleaf forest	Japanese cedar	Entire 2003-2004	0.979 (2003) 1.128 (2004)	Morishita et al. (2007)

a–Additional data collected from the original literature are available at <https://doi.org/10.5281/zenodo.14169306>; b–Observations of N₂O fluxes are represented as annual cumulative values ± standard errors, derived from field trial replicates reported in the literature.

Table S2. Literature-based cropland sites used for model soil N₂O evaluation: Geographic location; Cropping system and its growth calendar; Field management practices (such as irrigation, N fertilizer, and tillage) at the studied sites^a.

Geographic location	Lon	Lat	Cropping systems ^b	Sowing-harvest dates (DOY) ^c	Water and N fertilizer inputs ^d	Tillage, Residue Retention, Cover Crops	Period of N ₂ O measurement	Reference
Bretenierre (France)	5.11	47.23	SW	58-206	RF F: 131	NO, NO, YES	Mar.2012-Mar.2013	Vermue et al. (2016)
Reckenholz (Switzerland)	8.52	47.43	Maize	128-285	IRRI F: 160	YES, NO, NO	Mar.2014-Oct.2014	Hüppi et al. (2015)
DOK (Switzerland)	7.53	47.60	Maize	146-267	RF F: 0 170 183 335	YES, YES, YES	May 2013-Sep.2013	Skinner et al. (2019)
Aeschi (Switzerland)	7.66	47.17	Pea Maize	112-183, 2019 (Pea) 129-259, 2020 (Maize)	RF F:0 (Pea) 110 (Maize)	YES, NO, YES	May 2019-Jun.2019 May 2020-Sep.2020	Maier et al. (2022)
Jinzhou (China)	119.71	32.58	Rice	162-294	IRRI F: 0 150 250	YES, NO, NO	Jun.-Oct. in 2005, 2006, 2007	Yao et al. (2012)
Shuangqiao (China)	120.66	30.83	Rice	180-302	IRRI F: 0 90 180 270 360	YES, NO, NO	Jul.2007-Aug.2007	Iqbal (2009)
New Delhi (India)	77.20	28.67	Rice WW	180-302 (Rice) 315-89 (WW)	IRRI F: 0 120	YES, NO, NO	Jul.2000-Mar. 2001	Bhatia et al. (2005)
Cuttack (India)	85.93	20.43	Rice	193-293	IRRI F: 100	YES, NO, NO	Jul.2011-Oct.2011	Bhattacharyya et al. (2013)
Los Baños (Philippines)	121.26	14.16	Maize	55-164	IRRI F: 0 130 190	YES, NO, NO	Mar.2012-Jun.2012	Weller et al. (2015)
Suzhou (China)	120.92	31.53	Rice WW	174-292 (Rice) 323-139 (WW)	IRRI (Rice); RF (WW) F: 0 240	YES, NO, NO	Jul.2011-May 2014	Yang et al. (2015)
Changshu (China)	120.68	31.55	Rice	174-292	IRRI F: 0 225 270 300	YES, NO, NO	Jul.-Oct. in 2009, 2010	Ma et al. (2013)
Munich (Germany)	11.33	48.50	Maize WW	115-292 (Maize) 297-226 (WW)	RF F: 0 65 130 (Maize) F: 0 90 180 (WW)	YES, NO, NO	Mar.1995-Feb.1997	Ruser et al. (2001)
Jurong (China)	119.30	31.97	WW	346-163	RF F: 0 100 200 270	YES, NO, NO	Dec.2008-Jun.2009 Dec.2009-Jun.2010	Ji et al. (2012)

Yongji (China)	110.72	34.93	WW Maize	291-164 (WW) 166-288 (Maize)	IRRI F: 0 60 120 180 300 400 (WW) F: 0 75 150 250 350 450 (Maize)	YES, YES, NO	Nov.2009-Oct.2010	Liu et al. (2012)
Yanting (China)	105.28	31.16	WW Maize	299-130 (WW) 144-258 (Maize)	RF F: 0 150 250	YES, NO, NO	Oct.2004-Sep.2007	Zhou et al. (2013)
New Delhi (India)	77.20	28.67	WW	324-97	IRRI F: 0 120	YES, NO, NO	Dec.1998-Mar.1999	Majumdar et al. (2002)
Mundare (Canada)	-112.0	53.0	SW	144-268	RF F: 60	NO, NO, NO	Jun.1996-Sep.1996	Izaurrealde et al. (2004)
Swift Current (Canada)	-107.0	50.0	SW	144-272	RF F: 0 45 110	NO, NO, NO	Jun.1999-Sep.1999	Izaurrealde et al. (2004)
Huantai (China)	117.98	36.97	WW Maize	286-161 (WW) 173-270 (Maize)	IRRI F: 0 180 270 (WW) F: 0 220 330 (Maize)	YES, YES, NO	Oct.2008-Sep.2010	Yan et al. (2013)
Saint-Jean-sur- Richelieu (Canada)	-73.35	45.30	Maize Soybean	131-301 (Maize) 144-275 (Soybean)	RF F: 0 80 160 (Maize) F: 0 (Soybean)	YES, NO, NO NO, NO, NO	May-Oct. in 2004, 2005	Larouche (2006)
Honghe Farm (China)	131.52	47.58	Maize	153-264	RF F: 0 150 250	YES, NO, NO	Jun.2005-Sep.2005	Song and Zhang (2009)
Fengqiu (China)	114.40	35.00	Maize WW	158-262 (Maize) 282-156 (WW)	IRRI F: 0 150 250	YES, NO, NO	Jun.2004-May 2005	Ding et al. (2007)
Fort Collins (USA)	-104.98	40.65	Maize	124-287	IRRI F: 0 67 134 246	YES, NO, NO NO, YES, NO	May-Oct. in 2005, 2006	Halvorson et al. (2008)
Fort Collins (USA)	-104.98	40.65	Maize	116-290	IRRI F: 0 134 224	YES, NO, NO NO, YES, NO	May-Oct. in 2003, 2004	Liu et al. (2005)
Guelph (Canada)	-80.42	43.57	Maize	123-272 (2006) 129-268 (2007)	RF F: 0 60 90 150	YES, NO, NO	May and Jul. in 2006, 2007	Ma et al. (2010)
L'Île-Perrot (Canada)	-73.95	45.11	Maize Soybean	126-287 (Maize) 143-291 (Soybean)	RF F: 0 90 180 (Maize) F: 0 20 40 (Soybean)	YES, NO, NO	May 1994-Oct.1994	MacKenzie et al. (1997)
Wageningen (Netherlands)	5.67	51.96	Maize	126-281	RF F: 0 75 113 150 188	YES, NO, NO	May 2001-Oct.2001	van Groenigen et al. (2004)
Leeuwarden (Netherlands)	5.78	53.17	Maize	131-281	RF F: 0 75 113 150 188	YES, NO, NO	May 2001-Oct.2001	van Groenigen et al. (2004)

Eldorado do Sul (Brazil)	-51.07	-30.10	Maize	300-102	IRRI F: 0 180	NO, YES, YES	Oct.2004-Nov.2004	Bayer et al. (2015)
Beijing (China)	116.3	39.95	Maize	129-272	IRRI F: 0 150 300	YES, NO, NO	May 2002-Jun.2002	Liu et al. (2004)
KBS (USA)	-85.37	42.41	Maize	141-285 (2007) 135-297 (2008)	RF F: 0 45 90 135 180 225	YES, NO, NO	May-Oct. in 2007, 2008	Hoben et al. (2011)
Fairgrove (USA)	-83.64	43.52	Maize	111-281 (2007) 135-306 (2008)	RF F: 0 45 90 135 180 225	YES, NO, NO	Apr.-Oct. in 2007, 2008	Hoben et al. (2011)
Reese (USA)	-83.65	43.45	Maize	111-281 (2007) 135-306 (2008)	RF F: 0 45 90 135 180 225	YES, NO, NO	Apr.-Oct. in 2007, 2008	Hoben et al. (2011)
Mason (USA)	-84.51	42.47	Maize	127-295	RF F: 0 45 90 135 180 225	YES, NO, NO	May 2007-Oct.2007	Hoben et al. (2011)
Stockbridge (USA)	-84.27	42.48	Maize	143-287	RF F: 0 45 90 135 180 225	YES, NO, NO	May 2008-Sep.2008	Hoben et al. (2011)
Carlow (Ireland)	-6.92	52.85	SW	95-223 (2004) 92-217 (2005)	RF F: 0 70 140 (2004) F: 0 79 159 (2005)	YES, NO, NO NO, NO, NO	Apr.-Aug. in 2004, 2005	Abdalla et al. (2010)
Harlaka (Canada)	-71.32	46.77	Soybean	141-276 (2001) 141-258 (2002)	RF F: 0	YES, YES, NO	May -Oct. in 2001, 2002	Rochette et al. (2004)
Patancheru (India)	78.45	17.88	Sorghum	172-281	RF F: 0 90	YES, NO, NO	Jun.2010-Sep.2010	Ramu et al. (2012)
Dano (Burkina Faso)	-3.08	11.16	Sorghum	121-277	RF F: 0 52.5 140	YES, NO, NO	May 2006-Aug.2006	Brümmer et al. (2008)
Siribougou (Mali)	-6.43	13.27	Millet Bean	190-300 (Millet) 190-275 (Bean)	RF F: 0 24 (Millet) F: 0 (Bean)	YES, NO, NO	Entire 2004	Dick et al. (2008)
Göttingen (Germany)	9.93	51.49	WW	323-219	RF F: 0 180	YES, NO, NO	Mar.2016-Feb.2018	Wang et al. (2021)
Aranjuez (Spain)	-3.52	40.05	Maize	96-268	IRRI F: 140 170 190	YES, NO, NO YES, NO, YES	Oct.2013-Sep.2014	Guardia et al. (2016)
Willbriggie (Australia)	145.95	-34.48	Sorghum	356-119	IRRI F: 300	YES, NO, NO	Dec.2013-Apr.2014	Jamali et al. (2015)
Toowoomba (Australia)	151.78	-27.51	WW	161-298	IRRI F: 200	YES, NO, NO	Jun.2009-Oct.2009	Scheer et al. (2012)

St. Petersburg (Russia)	30.13	59.57	SW	132-242	RF F: 0 65 110	YES, NO, NO	May 2004-Aug.2004	Buchkina et al. (2010)
Ehime (Japan)	132.78	33.95	Rice	165-279	IRRI F: 80	YES, YES, NO	Jun.2012-Sep.2012	Toma et al. (2016)
Amir Temur Garden (Uzbekistan)	60.52	41.60	WW	282-171	IRRI F: 192	YES, NO, NO	Oct.2005-Jun.2006	Scheer et al. (2008)
Amir Temur Cum (Uzbekistan)	60.42	41.63	Rice	140-257 (2005) 143-252 (2006)	IRRI F: 270 (2005) 200 (2006)	YES, NO, NO	May-Oct. in 2005, 2006	Scheer et al. (2008)
Skierniewice (Poland)	20.17	51.95	SW	81-294	RF F: 0 90	YES, NO, NO	Apr.2012-Oct.2012	Sosulski et al. (2015)
Naples (Italy)	14.97	40.62	Maize	125-231	IRRI F: 130	YES, NO, NO NO, NO, NO	May 2008-Sep.2008	Forte et al. (2017)
Santa Maria (Brazil)	-53.71	-29.72	Maize WW	315-119 (Maize) 151-309 (WW)	IRRI F: 0 135 (Maize) F: 0 112 (WW)	YES, NO, YES YES, NO, NO	Nov.2010-Oct.2011	Aita et al. (2015)
Sinnamary (French Guiana)	-52.92	5.28	Maize	333-118	RF F: 169	YES, NO, YES NO, NO, YES	May 2010-Apr.2011	Petitjean et al. (2015)
Celaya (Mexico)	-101.32	20.73	Maize WW	104-258 (Maize) 287-78 (WW)	IRRI F: 240 (Maize) F: 300 (WW)	YES, YES, NO	Apr.2007-Mar.2008	Grageda-Cabrera et al. (2011)
Maseno (Kenya)	34.58	0.00	Maize	80-220	RF F: 0 50 75 100 200	YES, NO, NO	Mar.2010-Jun.2010	Hickman et al. (2014)
Harare (Zimbabwe)	31.01	-17.70	Maize	292-79	RF F: 0 60 120	YES, NO, NO	Oct.2008-Mar.2009	Nyamadzawo et al. (2017)
Kolero (Tanzania)	37.80	-7.25	Maize	76-189	RF F: 0 100	YES, NO, NO (F0) NO, NO, NO (F100)	Mar.2013-May 2014	Kimaro et al. (2016)
Kairanga (New Zealand)	175.65	-40.35	Maize	296-137	RF F: 163 186	YES, NO, NO	Nov.1998-Sep.1999	Choudhary et al. (2001)

a–The measured N₂O fluxes collected from the original literature are available at <https://doi.org/10.5281/zenodo.14169306>; b–Cropping systems: SW-spring wheat; WW-winter wheat; c–Sowing and harvest dates of crops are represented as DOY (day-of-year calendar); d–Water and N fertilizer management: RF-rain-fed; IRRI-irrigated; F-mineral N fertilizer application rate in kg N ha⁻¹.

Table S3. Modelled continent-level total soil N₂O emissions from natural vegetation, pasture, and cropland in the decades of the 1960s and 2010s (i.e., S1 run in Table 1 in the main text). The simulated results are represented as 10-year mean $\pm$ 1 standard deviation. The division of the 10 continents used in this study was given in Fig. S4.

	Natural vegetation (Tg N yr⁻¹)		Pasture (Tg N yr⁻¹)		Cropland (Tg N yr⁻¹)		All (Tg N yr⁻¹)	
	1960s	2010s	1960s	2010s	1960s	2010s	1960s	2010s
North America	0.58 $\pm$ 0.04	0.89 $\pm$ 0.09	0.04 $\pm$ 0.03	0.06 $\pm$ 0.04	0.09 $\pm$ 0.01	0.46 $\pm$ 0.06	0.71 $\pm$ 0.07	1.40 $\pm$ 0.12
South America	1.09 $\pm$ 0.06	0.94 $\pm$ 0.09	0.05 $\pm$ 0.01	0.05 $\pm$ 0.01	0.08 $\pm$ 0.01	0.18 $\pm$ 0.01	1.22 $\pm$ 0.06	1.17 $\pm$ 0.09
Europe	0.29 $\pm$ 0.02	0.71 $\pm$ 0.04	0.01 $\pm$ 0.00	0.02 $\pm$ 0.00	0.17 $\pm$ 0.06	0.49 $\pm$ 0.04	0.47 $\pm$ 0.08	1.21 $\pm$ 0.07
Africa	1.34 $\pm$ 0.09	1.30 $\pm$ 0.09	0.10 $\pm$ 0.02	0.12 $\pm$ 0.01	0.07 $\pm$ 0.01	0.19 $\pm$ 0.01	1.52 $\pm$ 0.09	1.61 $\pm$ 0.10
Oceania	0.17 $\pm$ 0.03	0.20 $\pm$ 0.06	0.07 $\pm$ 0.02	0.05 $\pm$ 0.02	0.05 $\pm$ 0.02	0.05 $\pm$ 0.01	0.28 $\pm$ 0.06	0.30 $\pm$ 0.08
West Asia	0.13 $\pm$ 0.02	0.19 $\pm$ 0.03	0.01 $\pm$ 0.00	0.01 $\pm$ 0.00	0.01 $\pm$ 0.00	0.09 $\pm$ 0.01	0.15 $\pm$ 0.02	0.29 $\pm$ 0.03
Central Asia	0.03 $\pm$ 0.00	0.07 $\pm$ 0.02	0.01 $\pm$ 0.00	0.02 $\pm$ 0.00	0.03 $\pm$ 0.02	0.11 $\pm$ 0.03	0.07 $\pm$ 0.02	0.19 $\pm$ 0.05
East Asia	0.20 $\pm$ 0.02	0.66 $\pm$ 0.03	0.02 $\pm$ 0.00	0.07 $\pm$ 0.01	0.04 $\pm$ 0.01	1.18 $\pm$ 0.12	0.25 $\pm$ 0.03	1.91 $\pm$ 0.13
South Asia	0.22 $\pm$ 0.02	0.40 $\pm$ 0.03	0.01 $\pm$ 0.00	0.01 $\pm$ 0.00	0.16 $\pm$ 0.01	0.79 $\pm$ 0.06	0.39 $\pm$ 0.03	1.20 $\pm$ 0.08
Southeast Asia	0.45 $\pm$ 0.02	0.50 $\pm$ 0.08	0.00 $\pm$ 0.00	0.01 $\pm$ 0.00	0.03 $\pm$ 0.00	0.14 $\pm$ 0.02	0.48 $\pm$ 0.02	0.65 $\pm$ 0.09
Global	4.51 $\pm$ 0.10	5.87 $\pm$ 0.13	0.31 $\pm$ 0.04	0.40 $\pm$ 0.05	0.73 $\pm$ 0.12	3.64 $\pm$ 0.18	5.55 $\pm$ 0.19	9.92 $\pm$ 0.28

Table S4. Modelled relative contributions ($\Delta N_2O\%$) of each factor to global soil N_2O emissions in the decades of the 1960s and 2010s. The modelled results are represented as a 10-year average (%). See S0–S6 runs in Table 1 in the main text for simulation setups and Eq. 24 for $\Delta N_2O\%$ calculation.

	Natural vegetation (%)		Pasture (%)		Cropland (%)		All (%)	
	1960s	2010s	1960s	2010s	1960s	2010s	1960s	2010s
Climate change	68.6	75.1	69.6	61.4	1.3	0.2	30.9	23.7
Rising CO_2	-73.9	-87.3	-34.1	-97.9	-1.8	-6.3	-30.3	-31.9
N fertilization	0.0	0.0	0.0	0.0	52.2	87.3	25.9	58.1
N deposition	174.7	125.9	63.7	116.7	8.9	9.0	76.5	46.4
Land use change	-89.2	-35.6	2.6	-2.3	46.0	15.6	-6.6	1.0
Interactive effects ^a	19.8	21.9	-1.8	22.1	-6.6	-5.8	3.6	2.8

a–The total effect of the five environmental factors did not sum to 100% due to their interactive effects in the factorial experiments. See Sect. 2.3.2 in the main text for more details.




Figure S1. Maps of (a) N deposition and (b) cropland N fertilization (chemical N fertilizer + manure) in the decades of the 1900s and the 2010s. Global total N inputs of deposition and cropland fertilization between 1901-2020 is shown in (c). N deposition data on global lands are taken from Tian et al. (2018). Mineral N fertilizer and manure application rate on croplands (1901-2015) are derived from Ag-GRID (Elliott et al., 2015) and Zhang et al. (2017), respectively.




Figure S2. Comparison of modelled and observed cumulative N₂O emissions on natural lands (top) and cropping systems (bottom) across all evaluated sites. Box plots denote the 5th and 95th percentiles by the whiskers, median and interquartile range are the box lines. ME (mean error) and MAE (mean absolute error) are shown as percentages.




Figure S3. Modelled daily fluxes of N_2 , nitrified N_2O , denitrified N_2O , and soil WFPS and temperature in the top layer (0–50 cm) for three different sites: (a) the Tapajós National Forest in Brazil ($54.9^{\circ}W$, $2.9^{\circ}S$; 2000–2004), (b) a temperate cedrus forest in Fukushima, Japan ($140.3^{\circ}E$, $37.4^{\circ}N$; 2003–2004), and (c) a semi-arid grassland in Inner Mongolia, China ($114.7^{\circ}E$, $43.5^{\circ}N$; 2001–2003). The thick lines represent the multi-year average of simulations, while open circles indicate the simulations for individual years.




Figure S4. A global division of geographical regions to analyze soil N₂O emissions at a continental level.




Figure S5. Modelled total N input and output in soils (Tg N yr⁻¹) across global terrestrial ecosystems in “All_Combined” and “Const_CO₂” runs in the 2010s. See Table 1 in the main text for simulation setups.

Reference

- Abdalla, M., Jones, M., Ambus, P. and Williams, M.: Emissions of nitrous oxide from Irish arable soils: Effects of tillage and reduced N input, *Nutr. Cycl. Agroecosystems*, 86(1), 53–65, doi:10.1007/s10705-009-9273-8, 2010.
- Aita, C., Schirrmann, J., Pujol, S. B., Giacomini, S. J., Rochette, P., Angers, D. A., Chantigny, M. H., Gonzatto, R., Giacomini, D. A. and Doneda, A.: Reducing nitrous oxide emissions from a maize-wheat sequence by decreasing soil nitrate concentration: Effects of split application of pig slurry and dicyandiamide, *Eur. J. Soil Sci.*, 66(2), 359–368, doi:10.1111/ejss.12181, 2015.
- Ball, T., Smith, K. A. and Moncrieff, J. B.: Effect of stand age on greenhouse gas fluxes from a Sitka spruce [*Picea sitchensis* (Bong.) Carr.] chronosequence on a peaty gley soil, *Glob. Chang. Biol.*, 13(10), 2128–2142, doi:10.1111/j.1365-2486.2007.01427.x, 2007.
- Bayer, C., Gomes, J., Zanatta, J. A., Vieira, F. C. B., Piccolo, M. de C., Dieckow, J. and Six, J.: Soil nitrous oxide emissions as affected by long-term tillage, cropping systems and nitrogen fertilization in Southern Brazil, *Soil Tillage Res.*, 146(PB), 213–222, doi:10.1016/j.still.2014.10.011, 2015.
- Bhatia, A., Pathak, H., Jain, N., Singh, P. K. and Singh, A. K.: Global warming potential of manure amended soils under rice-wheat system in the Indo-Gangetic plains, *Atmos. Environ.*, 39(37), 6976–6984, doi:10.1016/j.atmosenv.2005.07.052, 2005.
- Bhattacharyya, P., Roy, K. S., Neogi, S., Dash, P. K., Nayak, A. K., Mohanty, S., Baig, M. J., Sarkar, R. K. and Rao, K. S.: Impact of elevated CO₂ and temperature on soil C and N dynamics in relation to CH₄ and N₂O emissions from tropical flooded rice (*Oryza sativa* L.), *Sci. Total Environ.*, 461–462, 601–611, doi:10.1016/j.scitotenv.2013.05.035, 2013.
- Billings, S. A., Schaeffer, S. M. and Evans, R. D.: Trace N gas losses and N mineralization in Mojave desert soils exposed to elevated CO₂, *Soil Biol. Biochem.*, 34(11), 1777–1784, doi:10.1016/S0038-0717(02)00166-9, 2002.
- Bowden, R. D., Steudler, P. A., Melillo, J. M. and Aber, J. D.: Annual nitrous oxide fluxes from temperature forest soils in the northeastern United States, *J. Geophys. Res.*, 95, 13997–14005, doi:10.1029/JD095iD09p13997, 1990.
- Brümmer, C., Brüggemann, N., Butterbach-Bahl, K., Falk, U., Szarzynski, J., Vielhauer, K., Wassmann, R. and Papen, H.: Soil-atmosphere exchange of N₂O and NO in near-natural savanna and agricultural land in Burkina Faso (W. Africa), *Ecosystems*, 11(4), 582–600, doi:10.1007/s10021-008-9144-1, 2008.
- Buchkina, N. P., Balashov, E. V., Rizhiya, E. Y. and Smith, K. A.: Nitrous oxide emissions from a light-textured arable soil of North-Western Russia: Effects of crops, fertilizers, manures and climate parameters, *Nutr. Cycl. Agroecosystems*, 87(3), 429–442, doi:10.1007/s10705-010-9349-5, 2010.
- Castaldi, S., Bertolini, T., Valente, A., Chiti, T. and Valentini, R.: Nitrous oxide emissions from soil of an African rain forest in Ghana, *Biogeosciences*, 10(6), 4179–4187, doi:10.5194/bg-10-4179-2013, 2013.
- Castro, M. S., Steudler, P. A., Melillo, J. M., Aber, J. D. and Millham, S.: Exchange of N₂O and CH₄ between the atmosphere and soils in spruce-fir forests in the northeastern United States, *Biogeochemistry*, 18(3), 119–135, doi:10.1007/BF00003273, 1993.
- Cates, R. L. and Keeney, D. R.: Nitrous Oxide Emission from Native and Reestablished Prairies in Southern Wisconsin, *Am. Midl. Nat.*, 117(1), 35–42, doi:10.2307/2425705, 1987.
- Chen, G. X., Huang, B., Xu, H., Zhang, Y., Huang, G. H., Yu, K. W., Hou, A. X., Du, R., Han, S. J. and

- VanCleemput, O.: Nitrous oxide emissions from terrestrial ecosystems in China, *Chemosph. - Glob. Chang. Sci.*, 2(3–4), 373–378, doi:10.1016/S1465-9972(00)00036-2, 2000.
- Choudhary, M. A., Akramkhanov, A. and Saggar, S.: Nitrous oxide emissions in soils cropped with maize under long-term tillage and under permanent pasture in New Zealand, *Soil Tillage Res.*, 62(1–2), 61–71, doi:10.1016/S0167-1987(01)00208-2, 2001.
- Davidson, E. A., Nepstad, D. C., Ishida, F. Y. and Brando, P. M.: Effects of an experimental drought and recovery on soil emissions of carbon dioxide, methane, nitrous oxide, and nitric oxide in a moist tropical forest, *Glob. Chang. Biol.*, 14(11), 2582–2590, doi:10.1111/j.1365-2486.2008.01694.x, 2008.
- Dick, J., Kaya, B., Soutoura, M., Skiba, U., Smith, R., Niang, A. and Tabo, R.: The contribution of agricultural practices to nitrous oxide emissions in semi-arid Mali, *Soil Use Manag.*, 24(3), 292–301, doi:10.1111/j.1475-2743.2008.00163.x, 2008.
- Ding, W., Cai, Y., Cai, Z., Yagi, K. and Zheng, X.: Nitrous oxide emissions from an intensively cultivated maize-wheat rotation soil in the North China Plain, *Sci. Total Environ.*, 373(2–3), 501–511, doi:10.1016/j.scitotenv.2006.12.026, 2007.
- Du, R., Lu, D. and Wang, G.: Diurnal, seasonal, and inter-annual variations of N₂O fluxes from native semi-arid grassland soils of inner Mongolia, *Soil Biol. Biochem.*, 38(12), 3474–3482, doi:10.1016/j.soilbio.2006.06.012, 2006.
- Duxbury, J. M., Bouldin, D. R., Terry, R. E. and Tate, R. L.: Emissions of nitrous oxide from soils, *Nature*, 298, 462–464, doi:10.1038/298462a0, 1982.
- Elliott, J., Müller, C., Deryng, D., Chrysanthacopoulos, J., Boote, K. J., Büchner, M., Foster, I., Glotter, M., Heinke, J., Iizumi, T., Izaurralde, R. C., Mueller, N. D., Ray, D. K., Rosenzweig, C., Ruane, A. C. and Sheffield, J.: The Global Gridded Crop Model Intercomparison: Data and modeling protocols for Phase 1 (v1.0), *Geosci. Model Dev.*, 8(2), 261–277, doi:10.5194/gmd-8-261-2015, 2015.
- Forte, A., Fiorentino, N., Fagnano, M. and Fierro, A.: Mitigation impact of minimum tillage on CO₂ and N₂O emissions from a Mediterranean maize cropped soil under low-water input management, *Soil Tillage Res.*, 166, 167–178, doi:10.1016/j.still.2016.09.014, 2017.
- Grageda-Cabrera, O. A., Vera-Núñez, J. A., Aguilar-Acuña, J. L., Macías-Rodríguez, L., Aguado-Santacruz, G. A. and Peña-Cabriales, J. J.: Fertilizer dynamics in different tillage and crop rotation systems in a Vertisol in Central Mexico, *Nutr. Cycl. Agroecosystems*, 89(1), 125–134, doi:10.1007/s10705-010-9382-4, 2011.
- van Groenigen, J. W., G.J.Kasper, G.L.Velthof, van den Pol-van Dasselaar, A. and Kuikman, P. J.: Nitrous oxide emissions from silage maize fields under different mineral nitrogen fertilizer and slurry applications, *Plant Soil*, (263), 101–111, doi:10.1023/B, 2004.
- Groffman, P. M., Hardy, J. P., Discoll, C. T. and Fahey, T. J.: Snow depth, soil freezing, and fluxes of carbon dioxide, nitrous oxide and methane in a northern hardwood forest, *Glob. Chang. Biol.*, 12(9), 1748–1760, doi:10.1111/j.1365-2486.2006.01194.x, 2006.
- Guardia, G., Abalos, D., García-Marco, S., Quemada, M., Alonso-Ayuso, M., Cárdenas, L. M., Dixon, E. R. and Vallejo, A.: Effect of cover crops on greenhouse gas emissions in an irrigated field under integrated soil fertility management, *Biogeosciences*, 13(18), 5245–5257, doi:10.5194/bg-13-5245-2016, 2016.
- Halvorson, A. D., Del Grosso, S. J. and Reule, C. A.: Nitrogen, Tillage, and Crop Rotation Effects on Nitrous Oxide Emissions from Irrigated Cropping Systems, *J. Environ. Qual.*, 37(4), 1337–1344, doi:10.2134/jeq2007.0268, 2008.
- Henrich, M. and Haselwandter, K.: Denitrification and gaseous nitrogen losses from an acid spruce

forest soil, *Soil Biol. Biochem.*, 29(9–10), 1529–1537, doi:10.1016/S0038-0717(97)00010-2, 1997.

Hickman, J. E., Palm, C. A., Mutuo, P., Melillo, J. M. and Tang, J.: Nitrous oxide (N₂O) emissions in response to increasing fertilizer addition in maize (*Zea mays* L.) agriculture in western Kenya, *Nutr. Cycl. Agroecosystems*, 100(2), 177–187, doi:10.1007/s10705-014-9636-7, 2014.

Hoben, J. P., Gehl, R. J., Millar, N., Grace, P. R. and Robertson, G. P.: Nonlinear nitrous oxide (N₂O) response to nitrogen fertilizer in on-farm corn crops of the US Midwest, *Glob. Chang. Biol.*, 17(2), 1140–1152, doi:10.1111/j.1365-2486.2010.02349.x, 2011.

Hüppi, R., Felber, R., Neftel, A., Six, J. and Leifeld, J.: Effect of biochar and liming on soil nitrous oxide emissions from a temperate maize cropping system, *Soil*, 1(2), 707–717, doi:10.5194/soil-1-707-2015, 2015.

Iqbal, M. T.: Effects of nitrogen and phosphorous fertilisation on nitrous oxide emission and nitrogen loss in an irrigated rice field, *Malaysian J. Soil Sci.*, 13, 105–117, 2009.

Ishizuka, S., Tsuruta, H. and Murdiyarso, D.: An intensive field study on CO₂, CH₄, and N₂O emissions from soils at four land-use types in Sumatra, Indonesia, *Global Biogeochem. Cycles*, 16(3), doi:10.1029/2001gb001614, 2002.

Izaurrealde, R. C., Lemke, R. L., Goddard, T. W., McConkey, B. and Zhang, Z.: Nitrous Oxide Emissions from Agricultural Toposequences in Alberta and Saskatchewan, *Soil Sci. Soc. Am. J.*, (68), 1285–1294, doi:https://doi.org/10.2136/sssaj2004.1285, 2004.

Jamali, H., Quayle, W. C. and Baldock, J.: Reducing nitrous oxide emissions and nitrogen leaching losses from irrigated arable cropping in Australia through optimized irrigation scheduling, *Agric. For. Meteorol.*, 208, 32–39, doi:10.1016/j.agrformet.2015.04.010, 2015.

Ji, Y., Liu, G., Ma, J., Xu, H. and Yagi, K.: Effect of controlled-release fertilizer on nitrous oxide emission from a winter wheat field, *Nutr. Cycl. Agroecosystems*, 94(1), 111–122, doi:10.1007/s10705-012-9532-y, 2012.

Jungkunst, H. F., Fiedler, S. and Stahr, K.: N₂O emissions of a mature Norway spruce (*Picea abies*) stand in the Black Forest (southwest Germany) as differentiated by the soil pattern, *J. Geophys. Res. Atmos.*, 109(7), 1–11, doi:10.1029/2003jd004344, 2004.

Kesik, M., Ambus, P., Baritz, R., Brüggemann, N., Butterbach-Bahl, K., Damm, M., Duyzer, J., Horváth, L., Kiese, R., Kitzler, B., Leip, A., Li, C., Pihlatie, M., Pilegaard, K., Seufert, G., Simpson, D., Skiba, U., Smiatek, G., Vesala, T. and Zechmeister-Boltenstern, S.: Inventories of N₂O and NO emissions from European forest soils, *Biogeosciences*, 2(4), 353–375, doi:10.5194/bg-2-353-2005, 2005.

Kimaro, A. A., Mpanda, M., Rioux, J., Aynekulu, E., Shaba, S., Thiong'o, M., Mutuo, P., Abwanda, S., Shepherd, K., Neufeldt, H. and Rosenstock, T. S.: Is conservation agriculture 'climate-smart' for maize farmers in the highlands of Tanzania?, *Nutr. Cycl. Agroecosystems*, 105(3), 217–228, doi:10.1007/s10705-015-9711-8, 2016.

Kitzler, B., Zechmeister-Boltenstern, S., Holtermann, C., Skiba, U. and Butterbach-Bahl, K.: Nitrogen oxides emission from two beech forests subjected to different nitrogen loads, *Biogeosciences*, 3(3), 293–310, doi:10.5194/bg-3-293-2006, 2006.

Klemetsson, L., Kasimir Klemetsson, Å., Moldan, F. and Weslien, P.: Nitrous oxide emission from Swedish forest soils in relation to liming and simulated increased N-deposition, *Biol. Fertil. Soils*, 25(3), 290–295, doi:10.1007/s003740050317, 1997.

Koehler, B., Corre, M. D., Veldkamp, E., Wullaert, H. and Wright, S. J.: Immediate and long-term nitrogen oxide emissions from tropical forest soils exposed to elevated nitrogen input, *Glob. Chang. Biol.*, 15(8), 2049–2066, doi:10.1111/j.1365-2486.2008.01826.x, 2009.

Larouche, F.: Émissions De Protoxyde D ' Azote Dans Une Rotation Maïs/Soya Telles Qu '

- Influencées Par Le Travail Du Sol Et La Fertilisation Azotée, Laval University, Quebec, Canada., 2006.
- Liu, C., Wang, K. and Zheng, X.: Responses of N₂O and CH₄ fluxes to fertilizer nitrogen addition rates in an irrigated wheat-maize cropping system in northern China, *Biogeosciences*, 9(2), 839–850, doi:10.5194/bg-9-839-2012, 2012.
- Liu, X., Walsh, M., Ju, X., Zhang, F., Schimel, D. and Ojima, D.: NO and N₂O fluxes from agricultural soils in Beijing area, *Prog. Nat. Sci.*, 14(6), 489–494, doi:10.1080/10020070412331343841, 2004.
- Liu, X. J., Mosier, A. R., Halvorson, A. D. and Zhang, F. S.: Tillage and nitrogen application effects on nitrous and nitric oxide emissions from irrigated corn fields, *Plant Soil*, 276(1–2), 235–249, doi:10.1007/s11104-005-4894-4, 2005.
- Luizão, F., Matson, P., Livingston, G., Luizão, R. and Vitousek, P.: Nitrous oxide flux following tropical land clearing, *Global Biogeochem. Cycles*, (3), 281–285, doi:10.1029/GB003i003p00281, 1989.
- Luo, G. J., Brüggemann, N., Wolf, B., Gasche, R., Grote, R. and Butterbach-Bahl, K.: Decadal variability of soil CO₂, NO, N₂O, and CH₄ fluxes at the Höglwald Forest, Germany, *Biogeosciences*, 9(5), 1741–1763, doi:10.5194/bg-9-1741-2012, 2012.
- Ma, B. L., Wu, T. Y., Tremblay, N., Deen, W., Morrison, M. J., McLaughlin, N. B., Gregorich, E. G. and Stewart, G.: Nitrous oxide fluxes from corn fields: On-farm assessment of the amount and timing of nitrogen fertilizer, *Glob. Chang. Biol.*, 16(1), 156–170, doi:10.1111/j.1365-2486.2009.01932.x, 2010.
- Ma, Y. C., Kong, X. W., Yang, B., Zhang, X. L., Yan, X. Y., Yang, J. C. and Xiong, Z. Q.: Net global warming potential and greenhouse gas intensity of annual rice-wheat rotations with integrated soil-crop system management, *Agric. Ecosyst. Environ.*, 164, 209–219, doi:10.1016/j.agee.2012.11.003, 2013.
- MacKenzie, A. F., Fan, M. X. and Cadarin, F.: Nitrous oxide emission as affected by tillage, corn-soybean-alfalfa rotations and nitrogen fertilization, *Can. J. Soil Sci.*, 77(2), 145–152, doi:10.4141/S96-104, 1997.
- Maier, R., Hörtnagl, L. and Buchmann, N.: Greenhouse gas fluxes (CO₂, N₂O and CH₄) of pea and maize during two cropping seasons: Drivers, budgets, and emission factors for nitrous oxide, *Sci. Total Environ.*, 849, 157541, doi:10.1016/j.scitotenv.2022.157541, 2022.
- Majumdar, D., Pathak, H., Kumar, S. and Jain, M. C.: Nitrous oxide emission from a sandy loam Inceptisol under irrigated wheat in India as influenced by different nitrification inhibitors, *Agric. Ecosyst. Environ.*, 91(1–3), 283–293, doi:10.1016/S0167-8809(01)00223-7, 2002.
- Maljanen, M., Jokinen, H., Saari, A., Strömmer, R. and Martikainen, P. J.: Methane and nitrous oxide fluxes, and carbon dioxide production in boreal forest soil fertilized with wood ash and nitrogen, *Soil Use Manag.*, 22(2), 151–157, doi:10.1111/j.1475-2743.2006.00029.x, 2006.
- Matson, A., Pennock, D. and Bedard-Haughn, A.: Methane and nitrous oxide emissions from mature forest stands in the boreal forest, Saskatchewan, Canada, *For. Ecol. Manage.*, 258(7), 1073–1083, doi:10.1016/j.foreco.2009.05.034, 2009.
- Matson, P. A., Volkman, C., Coppinger, K. and Reiners, W. A.: Annual nitrous oxide flux and soil nitrogen characteristics in sagebrush steppe ecosystems, *Biogeochemistry*, 14(1), 1–12, doi:10.1007/BF00000883, 1991.
- Melling, L., Hatano, R. and Goh, K. J.: Nitrous oxide emissions from three ecosystems in tropical peatland of Sarawak, Malaysia, *Soil Sci. Plant Nutr.*, 53(6), 792–805, doi:10.1111/j.1747-0765.2007.00196.x, 2007.
- Mogge, B., Kaiser, E. A. and Munch, J. C.: Nitrous oxide emissions and denitrification N-losses from

forest soils in the bornhoved lake region (Northern Germany), *Soil Biol. Biochem.*, 30(6), 703–710, doi:10.1016/S0038-0717(97)00205-8, 1998.

Morishita, T., Sakata, T., Takahashi, M., Ishizuka, S., Mizoguchi, T., Inagaki, Y., Terazawa, K., Sawata, S., Igarashi, M., Yasuda, H., Koyama, Y., Suzuki, Y., Toyota, N., Muro, M., Kinjo, M., Yamamoto, H., Ashiya, D., Kanazawa, Y., Hashimoto, T. and Umata, H.: Methane uptake and nitrous oxide emission in Japanese forest soils and their relationship to soil and vegetation types, *Soil Sci. Plant Nutr.*, 53(5), 678–691, doi:10.1111/j.1747-0765.2007.00181.x, 2007.

Mosier, A. R., Parton, W. J., Valentine, D. W. and Schimel, D. S.: CH₄ and N₂O fluxes in the Colorado shortgrass steppe: 1. Impact of landscape and nitrogen addition, *Global Biogeochem. Cycles*, 10(3), 387–399, 1996.

Nyamadzawo, G., Shi, Y., Chirinda, N., Olesen, J. E., Mapanda, F., Wuta, M., Wu, W., Meng, F., Oelofse, M., de Neergaard, A. and Smith, J.: Combining organic and inorganic nitrogen fertilisation reduces N₂O emissions from cereal crops: a comparative analysis of China and Zimbabwe, *Mitig. Adapt. Strateg. Glob. Chang.*, 22(2), 233–245, doi:10.1007/s11027-014-9560-9, 2017.

Pei, Z., Hua, O., Zhou, C. and Xu, X.: N₂O Exchange Within a Soil and Atmosphere Profile in Alpine Grasslands on the Qinghai-Xizang Plateau, *Acta Bot. Sin.*, 46(1), 20–28, 2004.

Petitjean, C., Hénault, C., Perrin, A. S., Pontet, C., Metay, A., Bernoux, M., Jehanno, T., Viard, A. and Roggy, J. C.: Soil N₂O emissions in French Guiana after the conversion of tropical forest to agriculture with the chop-and-mulch method, *Agric. Ecosyst. Environ.*, 208, 64–74, doi:10.1016/j.agee.2015.04.004, 2015.

Price, S. J., Sherlock, R. R., Kelliher, F. M., McSeveny, T. M., Tate, K. R. and Condron, L. M.: Pristine New Zealand forest soil is a strong methane sink, *Glob. Chang. Biol.*, 10(1), 16–26, doi:10.1046/j.1529-8817.2003.00710x, 2004.

Purbopuspito, J., Veldkamp, E., Brumme, R. and Murdiyarso, D.: Trace gas fluxes and nitrogen cycling along an elevation sequence of tropical montane forests in Central Sulawesi, Indonesia, *Global Biogeochem. Cycles*, 20(3), 1–11, doi:10.1029/2005GB002516, 2006.

Ramu, K., Watanabe, T., Uchino, H., Sahrawat, K. L., Wani, S. P. and Ito, O.: Fertilizer induced nitrous oxide emissions from Vertisols and Alfisols during sweet sorghum cultivation in the Indian semi-arid tropics, *Sci. Total Environ.*, 438(2012), 9–14, doi:10.1016/j.scitotenv.2012.08.005, 2012.

Rees, R. M., Wuta, M., Furley, P. A. and Li, C.: Nitrous Oxide Fluxes from Savanna (Miombo) Woodlands in Zimbabwe, *J. Biogeogr.*, 33(3), 424–437, doi:10.1111/j.1365-2699.2005.01423.x, 2006.

Rochette, P., Angers, D. A., Bélanger, G., Chantigny, M. H., Prévost, D. and Lévesque, G.: Emissions of N₂O from Alfalfa and Soybean Crops in Eastern Canada, *Soil Sci. Soc. Am. J.*, 68(2), 493–506, doi:10.2136/sssaj2004.4930, 2004.

Rowlings, D. W., Grace, P. R., Kiese, R. and Weier, K. L.: Environmental factors controlling temporal and spatial variability in the soil-atmosphere exchange of CO₂, CH₄ and N₂O from an Australian subtropical rainforest, *Glob. Chang. Biol.*, 18(2), 726–738, doi:10.1111/j.1365-2486.2011.02563.x, 2012.

Ruser, R., Flessa, H., Schilling, R., Beese, F. and Munch, J. C.: Effect of crop-specific field management and N fertilization on N₂O emissions from a fine-loamy soil, *Nutr. Cycl. Agroecosystems*, 59(2), 177–191, doi:10.1023/A:1017512205888, 2001.

Scheer, C., Wassmann, R., Kienzler, K., Ibragimov, N., Lamers, J. P. A. and Martius, C.: Methane and nitrous oxide fluxes in annual and perennial land-use systems of the irrigated areas in the Aral Sea Basin, *Glob. Chang. Biol.*, 14(10), 2454–2468, doi:10.1111/j.1365-2486.2008.01631.x, 2008.

- Scheer, C., Grace, P. R., Rowlings, D. W. and Payero, J.: Nitrous oxide emissions from irrigated wheat in Australia: Impact of irrigation management, *Plant Soil*, 359(1–2), 351–362, doi:10.1007/s11104-012-1197-4, 2012.
- Schiller, C. L. and Hastie, D. R.: Nitrous oxide and methane fluxes from perturbed and unperturbed boreal forest sites in northern Ontario, *J. Geophys. Res.*, 101, 22767–22774, doi:10.1029/96JD01620, 1996.
- Simona, C., Ariangelo, D. P. R., John, G., Nina, N., Ruben, M. and José, S. J.: Nitrous oxide and methane fluxes from soils of the Orinoco savanna under different land uses, *Glob. Chang. Biol.*, 10(11), 1947–1960, doi:10.1111/j.1365-2486.2004.00871.x, 2004.
- Simpson, I. J., Edwards, G. C. and Thurtell, G. W.: Micrometeorological measurements of methane and nitrous oxide exchange above a boreal aspen forest, *J. Geophys. Res.*, 102, 29331–29341, doi:10.1029/97JD03181, 1997.
- Sitaula, B. K., Bakken, L. R. and Abrahamsen, G.: N-fertilization and soil acidification effects on N₂O and CO₂ emission from temperate pine forest soil, *Soil Biol. Biochem.*, 27(11), 1401–1408, doi:10.1016/0038-0717(95)00078-S, 1995.
- Skinner, C., Gattinger, A., Krauss, M., Krause, H. M., Mayer, J., van der Heijden, M. G. A. and Mäder, P.: The impact of long-term organic farming on soil-derived greenhouse gas emissions, *Sci. Rep.*, 9, 1702, doi:10.1038/s41598-018-38207-w, 2019.
- Song, C. and Zhang, J.: Effects of soil moisture, temperature, and nitrogen fertilization on soil respiration and nitrous oxide emission during maize growth period in northeast China, *Acta Agric. Scand. Sect. B Soil Plant Sci.*, 59(2), 97–106, doi:10.1080/09064710802022945, 2009.
- Sosulski, T., Szara, E., Stępień, W. and Rutkowska, B.: The influence of mineral fertilization and legumes cultivation on the N₂O soil emissions, *Plant, Soil Environ.*, 61(12), 529–536, doi:10.17221/229/2015-PSE, 2015.
- Takakai, F., Morishita, T., Hashidoko, Y., Darung, U., Kuramochi, K., Dohong, S., Limin, S. H. and Hatano, R.: Effects of agricultural land-use change and forest fire on N₂O emission from tropical peatlands, Central Kalimantan, Indonesia, *Soil Sci. Plant Nutr.*, 52(5), 662–674, doi:10.1111/j.1747-0765.2006.00084.x, 2006.
- Tian, H., Yang, J., Lu, C., Xu, R., Canadell, J. G., Jackson, R. B., Arneeth, A., Chang, J., Chen, G., Ciais, P., Gerber, S., Ito, A., Huang, Y., Joos, F., Lienert, S., Messina, P., Olin, S., Pan, S., Peng, C., Saikawa, E., Thompson, R. L., Vuichard, N., Winiwarter, W., Zaehle, S., Zhang, B., Zhang, K. and Zhu, Q.: The global N₂O model intercomparison project, *Bull. Am. Meteorol. Soc.*, 99(6), 1231–1251, doi:10.1175/BAMS-D-17-0212.1, 2018.
- Toma, Y., Oomori, S., Maruyama, A., Ueno, H. and Nagata, O.: Effect of the number of tillages in fallow season and fertilizer type on greenhouse gas emission from a rice (*Oryza sativa* L.) paddy field in Ehime, southwestern Japan, *Soil Sci. Plant Nutr.*, 62(1), 69–79, doi:10.1080/00380768.2015.1109999, 2016.
- Ullah, S. and Moore, T. R.: Biogeochemical controls on methane, nitrous oxide, and carbon dioxide fluxes from deciduous forest soils in eastern Canada, *J. Geophys. Res. Biogeosciences*, 116(3), 1–15, doi:10.1029/2010JG001525, 2011.
- Verchot, L. V., Davidson, E. a, Cattfinio, J. H., Ackerman, T. M. I. L., Erickson, H. E. and Keller, M.: Land use change and biogeochemical controls of nitrogen oxide emissions from soils in eastern Amazonia, *Global Biogeochem. Cycles*, 13, 31–46, doi:10.1029/1998GB900019, 1999.
- Vermue, A., Nicolardot, B. and Hénault, C.: High N₂O variations induced by agricultural practices in integrated weed management systems, *Agron. Sustain. Dev.*, 36(3), doi:10.1007/s13593-016-

0381-y, 2016.

Wang, H., Ma, S., Shao, G. and Dittert, K.: Use of urease and nitrification inhibitors to decrease yield-scaled N₂O emissions from winter wheat and oilseed rape fields: A two-year field experiment, *Agric. Ecosyst. Environ.*, 319(July), 107552, doi:10.1016/j.agee.2021.107552, 2021.

Weller, S., Kraus, D., Ayag, K. R. P., Wassmann, R., Alberto, M. C. R., Butterbach-Bahl, K. and Kiese, R.: Methane and nitrous oxide emissions from rice and maize production in diversified rice cropping systems, *Nutr. Cycl. Agroecosystems*, 101(1), 37–53, doi:10.1007/s10705-014-9658-1, 2015.

Werner, C., Kiese, R. and Butterbach-Bahl, K.: Soil-atmosphere exchange of N₂O, CH₄ and CO₂ and controlling environmental factors for tropical rain forest sites in western Kenya, *J. Geophys. Res.*, 112(3), 1–15, doi:10.1029/2006JD007388, 2007.

Wolf, K., Veldkamp, E., Homeier, J. and Martinson, G. O.: Nitrogen availability links forest productivity, soil nitrous oxide and nitric oxide fluxes of a tropical montane forest in southern Ecuador, *Global Biogeochem. Cycles*, 25(4), doi:10.1029/2010GB003876, 2011.

Yan, G., Zheng, X., Cui, F., Yao, Z., Zhou, Z., Deng, J. and Xu, Y.: Two-year simultaneous records of N₂O and NO fluxes from a farmed cropland in the northern China plain with a reduced nitrogen addition rate by one-third, *Agric. Ecosyst. Environ.*, 178, 39–50, doi:10.1016/j.agee.2013.06.016, 2013.

Yan, Y., Sha, L., Cao, M., Zheng, Z., Tang, J., Wang, Y., Zhang, Y., Wang, R., Liu, G., Wang, Y. and Sun, Y.: Fluxes of CH₄ and N₂O from soil under a tropical seasonal rain forest in Xishuangbanna, Southwest China, *J. Environ. Sci.*, 20(2), 207–215, doi:10.1016/S1001-0742(08)60033-9, 2008.

Yang, B., Xiong, Z., Wang, J., Xu, X., Huang, Q. and Shen, Q.: Mitigating net global warming potential and greenhouse gas intensities by substituting chemical nitrogen fertilizers with organic fertilization strategies in rice-wheat annual rotation systems in China: A 3-year field experiment, *Ecol. Eng.*, 81, 289–297, doi:10.1016/j.ecoleng.2015.04.071, 2015.

Yao, Z., Zheng, X., Dong, H., Wang, R., Mei, B. and Zhu, J.: A 3-year record of N₂O and CH₄ emissions from a sandy loam paddy during rice seasons as affected by different nitrogen application rates, *Agric. Ecosyst. Environ.*, 152, 1–9, doi:10.1016/j.agee.2012.02.004, 2012.

Zhang, B., Tian, H., Lu, C., Dangal, S., Yang, J. and Pan, S.: Manure nitrogen production and application in cropland and rangeland during 1860–2014: A 5-minute gridded global data set for Earth system modeling, *Earth Syst. Sci. Data*, 9, 667–678, doi:10.5194/essd-2017-11, 2017.

Zhou, M., Zhu, B., Butterbach-Bahl, K., Zheng, X., Wang, T. and Wang, Y.: Nitrous oxide emissions and nitrate leaching from a rain-fed wheat-maize rotation in the Sichuan Basin, China, *Plant Soil*, 362(1–2), 149–159, doi:10.1007/s11104-012-1269-5, 2013.