



# The Detection and Attribution Model Intercomparison Project (DAMIP v2.0) contribution to CMIP7

Nathan P. Gillett<sup>1</sup>, Isla R. Simpson<sup>2</sup>, Gabi Hegerl<sup>3</sup>, Reto Knutti<sup>4</sup>, Dann Mitchell<sup>5</sup>, Aurélien Ribes<sup>6</sup>, Hideo Shiogama<sup>7</sup>, Dáithí Stone<sup>8</sup>, Claudia Tebaldi<sup>9</sup>, Piotr Wolski<sup>10</sup>, Wenxia Zhang<sup>11</sup>, and Vivek K. Arora<sup>1</sup>

<sup>1</sup>Canadian Centre for Climate Modelling and Analysis, Environment and Climate Change Canada, Victoria, BC, Canada

<sup>2</sup>Climate and Global Dynamics Laboratory, NSF National Center for Atmospheric Research, Boulder, CO, USA

<sup>3</sup>School of GeoSciences, University of Edinburgh, Edinburgh, UK

<sup>4</sup>Institute for Atmospheric and Climate Science, ETH Zürich, Universitätstrasse 16, 8092 Zurich, Switzerland

<sup>5</sup>School of Geographical Sciences, University of Bristol, Bristol, UK

<sup>6</sup>CNRM, Université de Toulouse, Météo-France, CNRS, Toulouse, France

<sup>7</sup>National Institute for Environmental Studies, Tsukuba, 305-8506, Japan

<sup>8</sup>National Institute of Water and Atmospheric Research, Wellington, 6021, New Zealand

<sup>9</sup>Joint Global Change Research Institute, Pacific Northwest National Laboratory, College Park, MD, USA

<sup>10</sup>Climate System Analysis Group, University of Cape Town, Cape Town, South Africa

<sup>11</sup>State Key Laboratory of Numerical Modeling for Atmospheric Sciences and Geophysical Fluid Dynamics, Institute of Atmospheric Physics, Chinese Academy of Sciences, Beijing, China

**Correspondence:** Nathan P. Gillett (nathan.gillett@ec.gc.ca)

Received: 21 December 2024 – Discussion started: 7 February 2025

Revised: 24 April 2025 – Accepted: 26 April 2025 – Published: 22 July 2025

**Abstract.** The first version of the Detection and Attribution Model Intercomparison Project (DAMIP v1.0) coordinated key simulations exploring the role of individual forcings in past, current and future climate as part of the Coupled Model Intercomparison Project, Phase 6 (CMIP6). The simulations have been used extensively in the literature for detection and attribution of long-term changes, constraining projections of climate change, attributing extreme events and understanding drivers of past and future simulated climate changes. Attribution studies using DAMIP v1.0 simulations underpinned prominent assessments of human-induced warming in the Intergovernmental Panel on Climate Change (IPCC) Sixth Assessment Report. Here, we describe the set of DAMIP v2.0 simulations, proposed for the next phase of CMIP, CMIP7. Detection and attribution studies rely on pre-industrial control simulations and historical simulations, which will be part of the Diagnostic, Evaluation and Characterization of Klima (DECK) set of simulations for CMIP7. In addition, we identify the three highest-priority single-forcing experiments for CMIP7 to be run as “Assessment Fast Track” simulations in support of the Seventh Assess-

ment Report of the IPCC: simulations with natural forcings only, anthropogenic well-mixed greenhouse gases only and anthropogenic aerosols only. Beyond this, the DAMIP v2.0 experimental design includes full-column ozone-only simulations and land-use-only simulations, such that the set of individual forcing experiments, when these are considered together, represents the full set of historical forcings. While concentration-driven simulations are prioritised for attribution, emissions-driven versions of the DAMIP experiments are also proposed to support understanding of the influence of carbon-cycle feedbacks on the simulated responses to individual forcings.

## 1 Introduction

Research on the detection and attribution of climate change aims to identify and quantify the influence of particular forcings or subsets of forcings on the climate system, with a special focus on net human influence. Such research has underpinned progressively strengthening assessments of the role of

human influence in driving observed climate change in successive Intergovernmental Panel on Climate Change (IPCC) reports, including the assessment in the most recent report that “it is unequivocal that human influence has warmed the atmosphere, ocean and land” (Eyring et al., 2021). This research generally relies on climate model simulations of the response to individual forcings or subsets of external forcings, together with observations (Eyring et al., 2021; Stott et al., 2004). In particular, such analyses generally require simulations of historical climate change, including all major anthropogenic and natural influences (historical), long pre-industrial constant forcing simulations to characterise the influence of internal variability alone (piControl) and simulations with subsets of forcing agents. This paper describes the coordinated set of climate model simulations designed to support detection and attribution research that are proposed by the Detection and Attribution Model Intercomparison Project v2.0 (DAMIP v2.0), which is part of the Coupled Model Intercomparison Project, Phase 7 (CMIP7). In CMIP7, historical and pre-industrial control simulations will be coordinated as part of the Diagnostic, Evaluation and Characterization of Klima (DECK, Dunne et al., 2024) set of simulations, which all of the models must complete and which serve as a basis for model evaluation. Simulations with individual forcings or subsets of forcings are the focus of this paper.


## 2 Applications of DAMIP v1.0 CMIP6 simulations

DAMIP v2.0 follows on from DAMIP v1.0 (Gillett et al., 2016), the coordinated set of detection and attribution simulations conducted as part of the Coupled Model Intercomparison Project, Phase 6 (CMIP6) (Eyring et al., 2016). CMIP6 included historical simulations driven by both anthropogenic and natural forcings (historical) as well as constant pre-industrial forcing simulations (piControl). DAMIP v1.0 complemented these with historical simulations driven by subsets of the historical experiment forcings. The highest-priority Tier-1 simulations consisted of historical simulations driven by natural forcings only (hist-nat), historical simulations driven by well-mixed greenhouse gas changes only (hist-GHG) and historical simulations driven by aerosol and aerosol precursor emission changes only (hist-aer). These simulations were supplemented with lower-priority Tier-2 and Tier-3 simulations, including simulations driven by changes in stratospheric ozone only (hist-stratO3), volcanic aerosol only (hist-volc), solar irradiance only (hist-sol) and CO<sub>2</sub> only (hist-CO2). DAMIP v1.0 extended the individual historical forcing simulations up to 2100 using SSP2-4.5 (O’Neill et al., 2016) forcings to support analysis of the contributions of the different forcings to future changes (experiments ssp245-nat, ssp245-GHG, ssp245-aer and ssp245-stratO3). Since the publication of the original experimental design (Gillett et al., 2016), some additional experiments

were added to DAMIP v1.0: single-forcing experiments with CMIP5 forcings, to examine the effects of updates to the forcings from CMIP5 to CMIP6 (Fyfe et al., 2021), a simulation with ozone changes through the full atmospheric column (Shiogama et al., 2023) and a set of simulations to examine the response to COVID-induced changes in emissions (Jones et al., 2021).

Consistent with expectations, the Tier-1 simulations were carried out with the largest number of CMIP6 models, with hist-GHG simulations from 18 models, hist-aer simulations from 17 models and hist-nat simulations from 17 models published on the CMIP6 data portal as of 31 July 2024 (Fig. 1a). The simulations carried out with the fewest models were those added later in the CMIP6 cycle, namely the hist-GHG-cmip5, hist-aer-cmip5, hist-nat-cmip5, ssp245-cov-GHG and ssp245-cov-aer simulations, which were each carried out with only one model. Some modelling groups are now expanding their ensemble sizes through the Large Ensemble Single Forcing Model Intercomparison Project (LESFMIP) (Smith et al., 2022), which is particularly focused on the attribution of multi-annual to decadal changes in climate, including the effects of updates to forcing datasets. While all DAMIP v1.0 experiments were referred to in at least one publication, the Tier-1 simulations (hist-GHG, hist-aer and hist-nat) were by far the most cited, with hist-nat referred to in 245 publications (Fig. 1b). The DAMIP v1.0 simulations were also used extensively in the Sixth Assessment Report (AR6) of Working Group I of the IPCC, with data from these simulations used in the figures of five chapters of the report (Canadell et al., 2021; Doblas-Reyes et al., 2021; Eyring et al., 2021; Fox-Kemper et al., 2021; Szopa et al., 2021). Here also the Tier-1 simulations were also by far the most used. DAMIP v1.0 simulations were, for example, used in two attribution analyses of warming since pre-industrial times (Gillett et al., 2021; Ribes et al., 2021). These were two of the three main studies used to assess the anthropogenic contribution to observed warming (Eyring et al., 2021), which was a headline result in the Summary for Policymakers of the report (IPCC, 2021) and was also directly quoted in the Glasgow Climate Pact (UNFCCC, 2022). Since the publication of IPCC AR6, a selection of key climate indicators has been updated on a yearly basis (Forster et al., 2024). This includes warming attributable to human influence, calculated using DAMIP v1.0 simulations.

Beyond assessments of global temperature change, DAMIP simulations have been used to explore a wide variety of Earth system changes. They have been used to explore the role of individual forcings in modelled historical and projected future changes in the Atlantic Meridional Overturning Circulation (Menary et al., 2020). While the CMIP6 ensemble mean Atlantic Meridional Overturning Circulation response to forcings was rather linear in the forcing, this is not true of all models (Simpson et al., 2023). In addition, DAMIP simulations have been used to assess the relative roles of greenhouse gases and aerosols in historical changes



**Figure 1.** Bar graphs showing (a) the number of models with data published on the Earth System Grid Federation (ESGF) portal for each DAMIP v1.0 experiment and (b) the number of Google Scholar citations for each DAMIP v1.0 experiment (based on a search for “CMIP6” and each experiment name), as of 31 July 2024.

in drought frequency, duration and intensity (Chiang et al., 2021), the relative role of anthropogenic and natural forcings in contributing to increasing fire weather in the western United States (Zhuang et al., 2021) and the role of natural forcing, greenhouse gases and anthropogenic aerosols in historical changes in precipitation variability (Zhang et al., 2024b). They have been used to attribute the observed weakening of the summertime Eurasian jet stream in the historical record to anthropogenic aerosol forcing (Dong et al., 2022) and to isolate the relative role of greenhouse gases and aerosols in Northern Hemisphere summertime storm track trends (Kang et al., 2024). Despite this progress, we still do not adequately understand the relative role of forced changes and internal variability in observed circulation changes. Having now observed forced signals emerge for longer in observations and with continued improvements in process representation in Earth system models (ESMs), together with improved estimation of external forcings, further advances in this area may be achieved with the use of DAMIP v2.0 simulations.

As well as studies attributing long-term changes in climate, another application of DAMIP v1.0 simulations is in extreme event attribution, which aims to characterise how the probability of a single weather or climate event was altered by specific forcings, particularly anthropogenic forcing (Christidis et al., 2023; Herring et al., 2022; Lanet et al., 2024). This is an important use of DAMIP data and allows for wider global engagement in this field than those that have the capacity to run climate model experiments tailored to the specific event under analysis. However, it also presents an additional potential challenge. Approaches to extreme event attribution sometimes require ensembles of climate simulations, which are larger than the majority of DAMIP models will have (5–10 members) and up to thousands of ensemble members for very rare events (Schaller et al., 2014). Therefore, when DAMIP simulations are used for extreme event attribution analysis, methods for increasing the sample size might sometimes be needed. Such methods could include the use of model years beyond that of the specific event, combining different climate model ensembles (King, 2017) and increasing the sample size using extreme value theory (Sippel et al., 2015) or ensemble boosting (Fischer et al., 2023).

Increasingly, single-forcing climate simulations are being used in hazard and impacts research. DAMIP data have been used very effectively in going most of the way to explaining the observed impact changes for both trend-based impact attribution and event-based impact attribution. In many such studies, the authors are able to show a plausible explanation for the climate-influenced part of the observed impact. This has happened primarily in the fields of hydrology (e.g. Li et al., 2024a) and human health (e.g. Carlson, 2024; Chapman et al., 2022; Vicedo-Cabrera et al., 2021; Zhang et al., 2022). A challenge for impact detection and attribution research is to effectively integrate the relevant different socio-economic factors into the attribution framework, and this is particularly relevant in loss and damage, adaptation and legal settings (James et al., 2019).

### 3 Experimental design of DAMIP v2.0

In most respects, DAMIP v2.0 follows the experimental design of DAMIP v1.0 (Table 1, Fig. 2). As in DAMIP v1.0, we designate the highest-priority experiments as Tier-1 experiments, with Tier 2 and Tier 3 representing successively lower-priority experiments (see Fig. 2). There are two primary designs for detection and attribution experiments – namely individual forcing simulations and all-but-one simulations, in which all forcings except for the one of interest are included (Gillett et al., 2016; Smith et al., 2022). All-but-one simulations, together with historical simulations, may offer some advantages for detecting the presence of one particular forcing in the observations, in particular from a causality theory point of view (Hannart et al., 2016; Naveau et al., 2020), and they can be used together with individual

forcing simulations to test additivity (Marvel et al., 2015; Shiogama et al., 2013; Simpson et al., 2023). For example, aerosol-only simulations and all-but-aerosol simulations can be used together with simulations including all forcings to test whether the response to aerosols and the response to other forcings add up to give the response to all forcings combined (e.g. Simpson et al., 2023). LESFMIP proposed a comprehensive set of such simulations to investigate these questions (Smith et al., 2022). However, if the objective of an analysis is to characterise the response to one particular forcing, then individual forcing simulations will lead to reduced sampling uncertainties, because they do not require a difference between two sets of simulations to be taken, each of which has its own sampling uncertainties. For this reason, and for ease of comparison with previous CMIP generations, including the DAMIP v1.0 experiments, DAMIP v2.0 largely follows DAMIP v1.0 in being primarily based on individual forcing simulations. That said, natural-only (hist-nat) simulations can be equivalently described and used as all-but-anthropogenic forcing simulations, and DAMIP v2.0 links with an AerChemMIP2 simulation with all forcings but aerosols (hist-piAer), which can be used to address particular questions relating to the dependence of the anthropogenic aerosol impact on the climate state and additivity of the aerosol response with the responses to other forcings (Marvel et al., 2015; Shiogama et al., 2013; Simpson et al., 2023).

Recognising the advances in modelling which allow a larger fraction of climate models to be run with interactive CO<sub>2</sub>, the new science questions which may be addressed using interactive CO<sub>2</sub> simulations and the interest in running many CMIP7 simulations with interactive CO<sub>2</sub> (Sanderson et al., 2024), DAMIP v2.0 also proposes a set of individual forcing interactive CO<sub>2</sub> simulations (Sect. 3.3). Such simulations could, for example, be used to evaluate the effects of biogeochemical feedbacks on the responses to particular forcings, such as aerosols (e.g. Szopa et al., 2021), and could be used in studies attributing changes in atmosphere, land or ocean carbon pools to land use change, fossil CO<sub>2</sub> emissions and other GHG changes, and other factors. As in CMIP6, such interactive CO<sub>2</sub> simulations will have distinct experiment names from their corresponding prescribed concentration simulations (because the prognostic CO<sub>2</sub> concentration will generally differ from that prescribed in the corresponding concentration-driven experiments). Interactive CO<sub>2</sub> historical simulations will sample over uncertainties in the understanding and representation of the carbon cycle within ESMs and will allow analysis of the effects of carbon-cycle feedbacks on the responses to individual forcings, e.g. testing the frequently made assumption that the response to historical well-mixed GHG concentration changes is equivalent to the response to historical well-mixed GHG emission changes. However, given that direct and accurate observations of the evolution of atmospheric CO<sub>2</sub> exist, we recommend that prescribed-CO<sub>2</sub> simulations should continue to be

used for detection and attribution studies of changes in the physical climate. For this reason, our highest-priority Assessment Fast Track simulations are concentration-driven, as are the other simulations described in Sect. 3.1 and 3.2 (the Assessment Fast Track simulations are a subset of CMIP7 simulations to be carried out first in support of the IPCC Seventh Assessment Report; Dunne et al., 2024). If modelling centres have the capacity and interest to carry out both prescribed-CO<sub>2</sub> and interactive-CO<sub>2</sub> simulations, we recommend that they carry out both, allowing the effects of carbon-cycle feedbacks on the responses to individual sets of forcings to be isolated in each model.

Like the effects of an interactive carbon cycle, atmospheric chemistry has the potential to modify the simulated response to individual forcings or sets of forcings. In a model with a full representation of atmospheric chemistry, methane and fluorinated gas concentrations influence tropospheric and stratospheric ozone concentrations, solar and volcanic forcings influence stratospheric ozone concentrations and aerosols and aerosol precursors influence ozone and methane concentrations. Such interactions would make DAMIP simulations from models with and without interactive chemistry not fully comparable. Because of such interactions, attribution of physical climate changes to changes in concentrations of radiatively active species is fundamentally different to attribution to emission changes of such species. Typically, detection and attribution studies attribute observed changes in climate to concentration changes (e.g. Eyring et al., 2021), while emissions-driven individual forcing simulations may be used for bottom-up model-based estimates of the climate response to emissions of individual species (e.g. Szopa et al., 2021, Sect. 6.4). Like DAMIP v1.0, our focus here is mainly on supporting attribution to concentration changes, while AerChemMIP2 is proposing simulations to address attribution to emissions. While DAMIP v1.0 did propose an experimental design for hist-GHG and hist-aer for models with interactive chemistry to try to maintain comparability with other models, we note that this was never implemented because no modelling centres actually carried out the DAMIP v1.0 experiments with models with interactive chemistry. To simplify the experimental design and ensure comparability between outputs from different models, we therefore suggest that modelling groups carry out the DAMIP v2.0 experiments using model versions without gas-phase chemistry if possible. While we note that aerosol microphysics and chemistry schemes may make simulated aerosol concentrations sensitive to simulated temperatures, winds and possibly greenhouse gas concentrations and therefore different in the historical and hist-aer simulations, we note that the primary sensitivity of aerosol concentrations is to aerosol and aerosol precursor emissions and that most modelling centres do not have the capacity to run their models with specified aerosol concentrations, and therefore we accept such small differences as a limitation of our experimental design.


**Table 1.** DAMIP v2.0 simulations.

Name	Description of the prescribed concentration simulations (forcing agents perturbed)	Equivalent interactive CO <sub>2</sub> simulation name (modifications)	Modifications for coupled chemistry models	Start year	End year	Minimum ensemble size
historical and Medium scenario	Enlarging the ensemble sizes of the historical simulation (1850–2021) and ScenarioMIP Medium scenario (scen7-mc, 2022–2035) to an ensemble size of at least three (forcings: well-mixed greenhouse gases (WMGHGs), black carbon (BC), organic carbon (OC), SO <sub>2</sub> , SO <sub>4</sub> , NO <sub>x</sub> , NH <sub>3</sub> , CO, NMVOCs, nitrogen deposition, ozone, stratospheric aerosols, solar irradiance and land use)	esm-hist and esm-scen7-m (prescribe fossil CO <sub>2</sub> emissions instead of concentrations)		1850	2035	3
hist-nat	Natural-only historical simulations (solar irradiance and stratospheric aerosol)	esm-hist-nat		1850	2035	3
hist-GHG	Greenhouse-gas-only historical simulations (WMGHGs)	esm-hist-GHG (prescribe fossil CO <sub>2</sub> emissions instead of CO <sub>2</sub> concentrations)	Prescribe emissions instead of the concentrations of those WMGHGs simulated interactively	1850	2035	3
hist-aer	Anthropogenic-aerosol-only historical simulations (BC, OC, SO <sub>2</sub> , SO <sub>4</sub> , NO <sub>x</sub> , NH <sub>3</sub> , CO and NMVOCs)	esm-hist-aer		1850	2035	3
hist-O3	Ozone-only historical simulations (ozone concentration)	esm-hist-O3	n/a	1850	2035	3
hist-lu	Land-use-change-only historical simulations (land use)	esm-hist-lu		1850	2035	3
historical-CMIP6	Historical simulations with forcings from CMIP6 historical and SSP2-4.5 (WMGHGs, BC, OC, SO <sub>2</sub> , SO <sub>4</sub> , NO <sub>x</sub> , NH <sub>3</sub> , CO, NMVOCs, nitrogen deposition, ozone, stratospheric aerosols, solar irradiance and land use)	n/a	n/a	1850	2035	3

Table 1. Continued.

Name	Description of the prescribed concentration simulations (forcing agents perturbed)	Equivalent interactive CO <sub>2</sub> simulation name (modifications)	Modifications for coupled chemistry models	Start year	End year	Minimum ensemble size
hist-volc	Volcanic-aerosol-only historical simulation (stratospheric aerosols)	esm-hist-volc		1850	2035	3
Medium-GHG	Extension of at least one hist-GHG experiment to 2100 using the Medium scenario	esm-Medium-ghg (prescribe fossil CO <sub>2</sub> emissions instead of CO <sub>2</sub> concentrations)	Prescribe emissions instead of the concentrations of those WMGHGs simulated interactively	2036	2100	1
Medium-aer	Extension of at least one hist-aer experiment to 2100 using the Medium scenario	esm-Medium-aer		2036	2100	1
Medium-O3	Extension of at least one hist-O3 experiment to 2100 using the Medium scenario	esm-Medium-O3	n/a	2036	2100	1

n/a – not applicable.



**Figure 2.** Schematic of the experiments proposed in DAMIP v2.0 for CMIP7. The blue boxes are Tier-1 experiments, the yellow boxes are Tier-2 experiments, the green boxes are Tier-3 experiments, and the white boxes contain descriptive text. All historical simulations except for historical-CMIP6 run from 1850 to 2035 using the Medium scenario forcings from 2022, while historical-CMIP6 runs from 1850 to 2035 using the SSP2-4.5 scenario from 2015. The historical and esm-hist simulation shown in the top row uses CMIP7 historical forcings which can be decomposed into the sets of forcings used in each of the simulations in the second row. Three of these simulations are extended using the Medium scenario from 2036 to 2100, as shown in the third row. The fourth row depicts two additional experiments that are complementary to those in the second row. The hatched boxes show simulations which are in other MIPs but which are of particular relevance to DAMIP.

If modelling centres plan to only submit simulations with interactive gas-phase chemistry and are submitting historical simulations including this interactive chemistry, we request that they follow the DAMIP experimental design as indicated but with the following modifications (see also Table 1). For hist-GHG and esm-hist-GHG, they should specify emissions rather than concentrations of all well-mixed non-CO<sub>2</sub> greenhouse gases simulated interactively. hist-O<sub>3</sub> and esm-hist-O<sub>3</sub> simulations should not be carried out using models with interactive gas-phase chemistry. This is because ozone is simulated interactively in response to changes in ozone-depleting substances, methane and other species in such models, and the concentrations of ozone-depleting substances, methane and other species do not change in these simulations. historical-CMIP6 simulations should also not be carried out with such models. All other simulations should be carried out as specified.

While hist-nat simulations in models with gas-phase chemistry will include changes in stratospheric ozone which may modulate the response to solar and volcanic forcing, we expect the effects on the surface climate to be small, and such output could be used together with historical simulations to attribute surface climate change to anthropogenic and natural forcings. Similarly, hist-lu simulations from models with and without interactive chemistry are expected to be comparable. hist-GHG experiments are expected to differ between models with and without gas-phase chemistry, because in models with interactive chemistry, methane emissions will increase tropospheric ozone and halocarbon emissions will decrease stratospheric ozone. hist-aer experiments will also differ because emissions of NO<sub>x</sub>, non-methane volatile organic compounds (NMVOCs) and carbon monoxide (which are aerosol precursors) will all change tropospheric ozone and methane concentrations. Such experiments would not be directly comparable between models with and without gas-phase chemistry, but analysis of the output could inform understanding of atmospheric chemistry feedbacks. To facilitate analysis of the results, we request that modelling centres flag output from models with interactive chemistry with the “f2” flag and indicate in their metadata and documentation that the model included interactive chemistry. We note that AerChemMIP2 also provides a dedicated framework for systematically analysing the effects of such interactions across models with interactive chemistry.

### 3.1 Historical simulations

#### 3.1.1 Increased ensemble size for CMIP7 historical and Medium scenario simulations

While our focus is on historical experiments with single forcings or subsets of forcings, most detection and attribution analyses also use historical simulations with a complete set of forcings, and such analyses generally require an ensemble of such simulations, but only a single historical simulation is

required as part of the DECK (Dunne et al., 2024). Therefore we request at least three ensemble members of the CMIP7 CO<sub>2</sub> concentration-driven historical simulations (historical) and the extension of these simulations to 2035 with the Medium scenario (scen7-mc, intended to represent a frozen policy scenario) proposed by ScenarioMIP for CMIP7 (van Vuuren et al., 2025). Given that actual anthropogenic forcings are expected to diverge only slightly from the Medium emissions scenario over the first decade or so (see Sect. 4), we request that these historical simulations and other DAMIP experiments be extended in this way. This will allow researchers to carry out attribution analyses based on contemporary data over the next decade, at least in the absence of a major volcanic eruption, and will likely ensure an overlap with the next phase of CMIP. DAMIP v1.0 simulations were extended in a similar way from 2015 to 2020, but in hindsight this was not long enough, since at the time of writing CMIP7 simulations are not yet available but observations are available up to the end of 2024, well beyond the end of the DAMIP historical simulations. Such a need is particularly apparent for regularly updated attribution analyses (Forster et al., 2024). Modelling groups should publish the output data as CMIP7 historical simulations (1850–2021) and the Medium scenario simulations of ScenarioMIP (2022–2035). Note that we also request an ensemble size of at least three for all other DAMIP v2.0 historical simulations, though we encourage groups to run larger ensembles if possible. Also note that all DAMIP v2.0 simulations except for historical-CMIP6 use forcings from the CMIP7 historical, esm-hist and Medium simulations described by Dunne et al. (2024). A larger ensemble of CMIP7 historical simulations will also make it easier to evaluate the consistency of simulated historical climate change with that observed, though any such analysis should account for the use of historical temperature change in the tuning of some models (Hourdin et al., 2017).

#### 3.1.2 Simulations with a complete set of forcings

We propose a set of CO<sub>2</sub>-concentration-driven historical simulations driven by subsets of forcings which together add up to give the full set of CMIP7 historical forcings. This set includes the hist-nat, hist-GHG and hist-aer experiments, which are designated as Assessment Fast Track experiments here (Dunne et al., 2024) and also as Tier-1 experiments, based on the extensive use of the corresponding DAMIP v1.0 simulations in the literature and in support of IPCC assessment reports (Fig. 1b). For this reason, we ask modelling groups to prioritise these simulations and to consider running large ensembles of these simulations to support extreme event attribution and other applications (e.g. Smith et al., 2022), if time and resources allow. If the climate response to these forcings is additive, then the sum of the climate responses in this full set of historical simulations (hist-nat, hist-GHG, hist-aer, hist-O<sub>3</sub> and hist-lu) will be equal to the response in the historical experiment. We note that the vari-

ance of this sum will in general be 5 times larger than the variance of the response in historical, which will need to be accounted for when identifying departures from additivity. This represents a minor adjustment relative to the DAMIP v1.0 experimental design in which land use and land cover change and tropospheric ozone changes were not included in any of the original DAMIP v1.0 simulations. All of these simulations should be run using the CMIP7 historical forcings from 1850 to 2021 and the Medium scenario (scen7-mc) from 2022 to 2035, and we request a minimum ensemble size of three for all of the historical experiments.

**hist-nat:** these natural-only simulations parallel the historical and Medium scenario simulations but instead are only forced with solar and volcanic forcings from the historical and Medium scenario simulations, similar to the CMIP6 hist-nat experiment. Such simulations, when compared with the historical and Medium simulations, can be used for attribution of observed changes to anthropogenic influence, as they correspond to the counterfactual world in which human influence is removed. While much of the time evolution of biomass burning emissions has occurred as a result of human activity, the historical simulation includes observed year-to-year variations in biomass burning partly driven by natural variability. However, it is not easy to separate human-induced changes in biomass burning from naturally induced changes. Therefore, we request that modelling centres specify constant biomass burning emissions as in piControl in this hist-nat simulation. In contrast to DAMIP v1.0, and consistent with our aim of ensuring a complete set of forcings across this simulation set and simplifying the experimental design, we propose that no ozone changes be specified in the hist-nat experiment and that all ozone changes be included in the hist-O3 experiment. To test the sensitivity to this change in the experimental design, we carried out a test simulation with CanESM5.0 of the response to the DAMIP v1.0 solar and volcanically induced ozone changes alone. This showed small forced changes in the global mean stratospheric temperature of less than 0.5 °C but no discernible changes in the tropospheric climate.

**hist-GHG:** these greenhouse-gas-only simulations parallel the historical and Medium simulations but are forced by well-mixed greenhouse gas (carbon dioxide, methane, nitrous oxide and fluorinated gas) changes only from the historical and Medium scenario simulations, similarly to the CMIP6 hist-GHG experiment. Both stratospheric and tropospheric ozone should be held constant at piControl levels. Greenhouse gas changes are the dominant anthropogenic forcing, and these simulations will allow the response to this forcing to be quantified. Moreover, historical/Medium, hist-nat and hist-GHG will together allow the attribution of observed climate change to natural, greenhouse gas and other anthropogenic forcings (e.g. Gillett et al., 2021; Ribes et al., 2021).

**hist-aer:** these historical aerosol-only simulations parallel the historical and Medium simulations but are forced by

changes in aerosol and aerosol precursor emission changes only, as in the historical and Medium scenario. This includes changes in sulfur dioxide, sulfate, black carbon, organic carbon, ammonia, NO<sub>x</sub> and volatile organic compounds (VOCs) from biomass burning, industrial emissions and other sources.

**hist-lu:** these simulations parallel the historical and Medium simulations but are forced with prescribed land use and land cover changes only from the historical and Medium scenario simulations, with all other forcings held constant at 1850 values. No such experiments were included in DAMIP v1.0, although they have since been proposed in LESFMIP (Smith et al., 2022), and historical experiments without land use change (hist-noLu) were included in CMIP6 in the Land Use Model Intercomparison Project (LUMIP, Lawrence et al., 2016). These experiments were used to investigate the effects of land use change on surface temperature and other variables in models (Luo et al., 2024; Zhang et al., 2024a) and to diagnose land use change emissions (e.g. Liddicoat et al., 2021). Note that hist-noLu is also a proposed LUMIP experiment for CMIP7, and hence hist-lu and hist-noLu could be used together to evaluate the additivity of the land use change response and the response to other forcings. These experiments could also, for example, be used together to compare and contrast simulated historical land use change emissions with atmospheric CO<sub>2</sub> held constant, with simulated land use change emissions in the presence of CO<sub>2</sub> fertilisation with CO<sub>2</sub> changing through the historical period.

**hist-O3:** these simulations parallel the historical and Medium simulations but are forced by changes in ozone concentrations only from the historical and Medium scenario simulations. They will allow characterisation of the response to combined tropospheric and stratospheric ozone changes, which have played an important role in driving circulation changes in the high latitudes of the Southern Hemisphere and temperature changes in the stratosphere, together with attribution studies of the response to ozone change (e.g. Gillett et al., 2013; Morgenstern, 2021). Since it is increasingly understood that ozone-depleting substances influence ozone concentrations in the troposphere (e.g. Hassler et al., 2022; Li et al., 2024b), in CMIP7 we request simulations forced by changes in ozone concentrations over the whole atmospheric column, as opposed to the stratospheric-ozone-change-only simulations proposed in CMIP6 (Gillett et al., 2016) (although experiments with ozone changes over the whole atmosphere were later proposed and carried out (Liu et al., 2022; Shiogama et al., 2023; Smith et al., 2022)). Moreover, this choice simplifies the experiment design and avoids difficulties associated with specifying stratospheric ozone changes only, given that tropopause height can differ across models (although we acknowledge that inconsistencies between tropopause height and ozone concentrations can still exist when prescribing ozone over the full column, which users should be aware could impact conclusions under certain circumstances; Hardiman et al., 2019).

### 3.1.3 Other historical simulations

**hist-volc:** the hist-volc simulations parallel the hist-nat simulations, except that the hist-volc simulations are driven by stratospheric aerosol changes only. The hist-volc experiments will allow the characterisation of and attribution to volcanic influence, separate from the poorly constrained response to variations in solar forcing. Such analysis can improve our understanding of the response to future volcanic eruptions.

**historical-CMIP6:** these are identical to the CMIP7 historical experiments, but instead of using CMIP7 forcings, they use CMIP6 historical forcings from 1850 to 2014 and ScenarioMIP SSP2-4.5 forcings from 2015 to 2035. CMIP simulations of past and future climate changes are key to attribution and projection of climate change, e.g. in IPCC assessments (Eyring et al., 2021; Lee et al., 2021), and it is important to be able to understand differences in the results based on successive CMIP generations. Comparing historical CMIP6 and CMIP7 simulations will be complicated by the fact that both the models and the forcings will be different between the two CMIP generations, making it difficult to separate the influences of updates to the forcings and changes to the models. By carrying out simulations with a subset of CMIP7 models using CMIP6 forcings, it will be possible to separate the influences of updates to the forcings. A similar approach was used to isolate the influence of updates to forcings between CMIP5 and CMIP6, and the impact of forcing changes was shown to be an important contributor to differences between CMIP6 and CMIP5 simulations (Fyfe et al., 2021; Holland et al., 2024).

### 3.2 Future simulations driven by subsets of forcings

While ScenarioMIP simulations including scenarios of future changes in greenhouse gases, aerosols, ozone and land use change allow the simulation of possible future climate evolution, they do not directly allow the effects of particular forcings, or sets of forcings, to be isolated. Simulations of future climate change with subsets of forcings can help us understand the drivers of future climate change. Moreover, the Allen et al. (2000) and Stott and Kettleborough (2002) approach to constraining climate projections may be used to separately scale the future responses to well-mixed greenhouse gases and aerosols, based on regressions over the historical period, since models may overestimate or underestimate the climate response to greenhouse gases and aerosols by different factors. This kind of analysis requires future scenario simulations with subsets of forcings. DAMIP v1.0 included ssp245-nat, ssp245-GHG, ssp245-aer and ssp245-stratO3 simulations to address such needs (Gillett et al., 2016). Similarly, DAMIP v2.0 also includes future simulations with subsets of forcings following the Medium scenario from ScenarioMIP for CMIP7 (van Vuuren et al., 2025),

which approximately corresponds to a continuation of current climate policies.

**Medium-GHG:** this is an extension of the hist-GHG simulations to 2100 using the Medium scenario's well-mixed greenhouse gas concentrations, with other forcings kept at pre-industrial values.

**Medium-aer:** this is an extension of the hist-aer simulations to 2100 following the Medium scenario's aerosol concentrations and emissions, with other forcings kept at pre-industrial levels. These simulations will also allow a more robust characterisation of the response to future aerosol changes without conflating these changes with the responses to ozone and land use changes, as would occur if the aerosol response were estimated from a difference between the Medium scenario and Medium-GHG.

**Medium-O3:** these simulations are extensions of the hist-O3 simulations to 2100 following the ozone concentrations specified for the Medium scenario. Stratospheric ozone is projected to recover following the successful implementation of the Montreal Protocol and its amendments (e.g. Hassler et al., 2022). These simulations will facilitate a robust multi-model assessment of the climate effects of this recovery on Southern Hemisphere climate and stratospheric temperature.

### 3.3 Interactive CO<sub>2</sub> experiments

Detection and attribution studies have typically attributed changes in the physical climate system to changes in concentrations of greenhouse gases and other species by regressing observed changes onto the simulated response to changes in concentrations of greenhouse gases and other species (e.g. Eyring et al., 2021), but an alternative bottom-up approach uses models directly to simulate the responses to changes in emissions of particular greenhouse gases or other species (e.g. Forster et al., 2021, Sect. 7.3; Szopa et al., 2021, Sect. 6.4). Such approaches can quantify the respective contributions of emissions of various chemical species, such as CO<sub>2</sub>, methane and SO<sub>2</sub>. These approaches can account for feedbacks on the responses to emissions of particular chemical species, based on their representation in models. IPCC (2021, Fig. SPM.2) contrasted observationally constrained attributable warming in response to changes in concentrations of greenhouse gases, among other factors, with estimates of the warming attributable to emissions of CO<sub>2</sub>, methane and other species. These estimates included the simulated effects of tropospheric chemistry (Szopa et al., 2021), which strongly influence the responses to some forcings, but they omitted the effects of carbon-climate feedbacks, which are also expected to influence the responses to such forcings (Szopa et al., 2021). In part to address this knowledge gap, we propose interactive-CO<sub>2</sub> versions of all of the experiments described in Sect. 3.1 and 3.2, with the exception of historical-CMIP6 (where our focus is on understanding differences in the concentration-driven historical experiments between CMIP6 and CMIP7). Such experiments should be

carried out using ESMs with interactive carbon cycles, which can simulate atmospheric concentrations of CO<sub>2</sub> based on prescribed emissions. As noted above, we recommend that modelling centres carrying out these interactive CO<sub>2</sub> experiments also carry out the corresponding prescribed CO<sub>2</sub> experiments, allowing the effects of carbon-climate feedbacks on the response to each set of forcings to be quantified. These simulations should be run from 1850 to 2035 using the esm-hist CO<sub>2</sub> emissions and other forcings for the period 1850–2021 and the Medium scenario's CO<sub>2</sub> emissions and other forcings for the period 2022–2035.

**esm-hist-nat:** these interactive-CO<sub>2</sub> simulations parallel the esm-hist simulations but with only solar and volcanic forcings varying and all other forcings held fixed, including ozone. Such simulations can be used, for example, to evaluate the effects of carbon-climate feedbacks on the response to volcanoes (Kandlbauer et al., 2013; Rothenberg et al., 2012) or as a true counterfactual simulation reflecting the CO<sub>2</sub> evolution in the absence of anthropogenic influence.

**esm-hist-GHG:** these interactive-CO<sub>2</sub> simulations parallel the esm-hist simulations but only include fossil CO<sub>2</sub> emissions and changes in the concentrations of other well-mixed greenhouse gases (methane, nitrous oxide and fluorinated gas). No changes in land use or emissions associated with land use change should be prescribed.

**esm-hist-aer:** these interactive-CO<sub>2</sub> simulations parallel the esm-hist simulations but only include changes in aerosol and aerosol precursor emissions. Changes in aerosols can perturb the carbon cycle, not only through changes in the climate but also through deposition of nutrients such as nitrogen, sulfur, iron and phosphorous and through changes in solar irradiance and diffuse radiation at the surface (Szopa et al., 2021), and the response will depend on the representation of these mechanisms in each ESM.

**esm-hist-lu:** these interactive-CO<sub>2</sub> simulations parallel the esm-hist simulations but are forced with prescribed land use and land cover changes only, with all other forcings held constant at 1850 values. Such experiments will include the simulation of CO<sub>2</sub> emissions based on prescribed land use change and will support the calculation of the net climate influence of land use change (biogeophysical and carbon dioxide effects) (e.g. IPCC, 2019).

**esm-hist-O3:** these interactive-CO<sub>2</sub> simulations parallel the esm-hist simulations but only include prescribed changes in tropospheric and stratospheric ozone concentrations. Tropospheric ozone increases reduce terrestrial plant growth, influencing land carbon uptake (Szopa et al., 2021). Early studies found a substantial role for stratospheric ozone in driving changes in the Southern Ocean carbon sink (Le Quéré et al., 2007), but more recent assessments find a weaker role (Garny et al., 2022). These simulations will support the investigation of the effects of such processes on atmospheric CO<sub>2</sub> concentration.

**esm-hist-volc:** these interactive-CO<sub>2</sub> simulations parallel the esm-hist simulations but are driven by stratospheric aerosol changes only.

**esm-Medium-GHG:** these interactive CO<sub>2</sub> simulations are extensions of the esm-hist-GHG simulations to 2100, with fossil CO<sub>2</sub> emissions and other well-mixed greenhouse gas concentrations following the Medium scenario.


**esm-Medium-aer:** these interactive CO<sub>2</sub> simulations are extensions of the esm-hist-aer simulations to 2100, with aerosol and aerosol precursor emissions following the Medium scenario.

**esm-Medium-O3:** these interactive CO<sub>2</sub> simulations are extensions of the esm-hist-O3 simulations to 2100, with tropospheric and stratospheric ozone concentrations following the Medium scenario.

Figure 3 shows the global mean surface CO<sub>2</sub> concentration and temperature in a subset of such simulations carried out with CanESM5.0 (Swart et al., 2019), compared with the corresponding prescribed concentration simulations for reference. The simulated CO<sub>2</sub> in the esm-hist simulations of this model is within 15 ppm of that observed and that specified in the historical and hist-GHG simulations (compare the solid and dashed black lines in Fig. 3a). As expected, esm-hist-GHG shows lower simulated CO<sub>2</sub> concentrations than esm-hist, because it only includes emissions of fossil CO<sub>2</sub> and omits land use change emissions of CO<sub>2</sub>. Because of this, esm-hist-GHG exhibits less warming than hist-GHG (Fig. 3b). By contrast, esm-hist-lu shows an increase in atmospheric CO<sub>2</sub> due to the interactively simulated effects of land use change, and hence it is warmer than hist-lu, in which a constant pre-industrial CO<sub>2</sub> concentration is specified (Fig. 3b). In this model, atmospheric CO<sub>2</sub> increases slightly in esm-hist-aer. While the ocean takes up carbon in this simulation in response to the simulated cooling, this is more than compensated for by the land giving up carbon, likely due to reduced photosynthesis in response to the reduced solar irradiance. However, this model lacks a representation of the vegetation response to the change in diffuse irradiance associated with the increase in atmospheric aerosols (Szopa et al., 2021), and therefore this response might be different in other models. Because of this increase in atmospheric CO<sub>2</sub>, the cooling in response to aerosols in esm-hist-aer is slightly weaker than in hist-aer, but overall these two experiments exhibit a comparable global surface temperature response in this model.

### 3.4 Updated forcing simulations

As noted in Sect. 3.1, for the purposes of detection and attribution analyses, we recommend extending CMIP7 historical simulations from 2022 to 2035 with the ScenarioMIP Medium scenario and similarly extending DAMIP single-forcing simulations from 2022 to 2035 with the individual forcings prescribed for this scenario. However, observed concentrations and emissions of forcing agents will at some



**Figure 3.** Global mean surface CO<sub>2</sub> concentration (a) and global mean near-surface air temperature (b) in interactive CO<sub>2</sub> experiments performed with CanESM5.0 (Swart et al., 2019). Solid lines show results from interactive CO<sub>2</sub> simulations (esm-hist in black, esm-hist-GHG in red, esm-hist-lu in green, and esm-hist-aer in blue), and dashed lines show results from the corresponding prescribed CO<sub>2</sub> simulations (historical in black, hist-GHG in red, hist-lu in green, and hist-aer in blue). The results shown are the ensemble means of at least five ensemble members. Note that the CO<sub>2</sub> concentration prescribed in hist-GHG is the same as that in historical and is fixed at the pre-industrial level in hist-lu and hist-aer.

point evolve differently from those specified in this scenario. While forcings have evolved broadly consistently with the SSP2-4.5 scenario used to extend DAMIP v1.0 simulations since 2015 (Matthews and Wynes, 2022), there have been some differences. For example, recent aerosol emissions from China have declined more strongly than those specified in the SSP2-4.5 scenario (Wang et al., 2021), a decline in marine sulfur emissions resulting from new International Maritime Organisation regulations was not included in the SSP2-4.5 simulations (Watson-Parris et al., 2022, 2025), the COVID-19 pandemic changed emissions temporarily in ways which were not included in the SSP2-4.5 scenario (e.g. Szopa et al., 2021) and the Hunga Tonga eruption in 2021 injected sulfur dioxide and water vapour into the stratosphere, influencing the climate (e.g. Jenkins et al., 2023). In the past, such differences between forcings used to drive climate model simulations, and those which actually transpired have been cited as reasons for differences between simulated and observed warming trends in the early 21st century (e.g. Eyring et al., 2021; Santer et al., 2014) and in the early 2020s (Rantanen and Laaksonen, 2024). For these reasons, LESFMIP proposed a set of individual forcing simulations using regularly updated forcing estimates as part of CMIP6 (Smith et al., 2022), though at the time of writing such simulations have not yet been carried out and the protocols and processes are not in place for regular updates of forcings which would support such experiments. If this could be achieved in the future, however, such simulations could be used to understand the influence of updates to particular forcing estimates on simulated climate trends, particularly on interannual to decadal timescales, as well as to reduce uncertainties in attribution results associated with forcing changes after 2021. Given the uncertainty in whether future forcings over the next decade or so will diverge strongly

from those specified in the Medium scenario and whether and when updated forcing datasets will be made available to the modelling community, we do not propose additional named experiments here. However, the CMIP community is continuing to explore how forcings can be updated more regularly and how such updates could be used in CMIP7. DAMIP will engage with this discussion to ensure any solutions developed support updates to attribution simulations. Any such effort would require additional community coordination and documentation if and when new forcing datasets are published, and this would be most valuable if carried out in conjunction with updated all-forcing scenario simulations as part of ScenarioMIP or the Decadal Climate Prediction Project (DCPP).

#### 4 Synergies with other MIPs

In CMIP6, several DAMIP v1.0 experiments were coordinated with other MIPs in order to maximise synergies and support research to address a wide range of scientific questions (Gillett et al., 2016). For example, DAMIP v1.0 and the Radiative Forcing Model Intercomparison Project (RFMIP) (Pincus et al., 2016) proposed coordinated coupled and prescribed sea surface temperature simulations with natural-only and well-mixed GHG-only forcings in order to estimate the corresponding effective radiative forcings. For CMIP7, we have again coordinated with a wide range of MIPs to maximise scientific synergies.

As discussed above, all DAMIP v2.0 simulations extend beyond the end of the historical simulation in 2021, and we request that modelling centres extend these simulations with the Medium scenario from ScenarioMIP. Our simulations which are extended to 2100 are also coordinated with

ScenarioMIP to support understanding of the roles of individual forcings in driving future climate change in the Medium scenario. Similarly, we are coordinating our scenario choice with the DCP, such that both DAMIP simulations and DCP simulations will use the same scenario, allowing the contributions of individual forcings to near-future climate change to be calculated and compared with initialised predictions under the same consistent set of forcings.

The AerChemMIP2 Assessment Fast Track hist-piAer experiments parallel the historical and Medium scenario simulations, except that aerosols are kept fixed at pre-industrial levels. Together with the hist-aer and historical experiments, these experiments will allow analysis of the extent to which the climate response to aerosols is additive with the response to other forcings. For this reason we encourage modelling centres participating in DAMIP to also carry out the hist-piAer simulations.

The RFMIP experiment piClim-histaer (which is an atmosphere-only simulation with fixed pre-industrial sea surface temperatures (SSTs), sea ice and transient aerosols) parallels hist-aer, except that it has fixed SSTs. This experiment can be used to diagnose the evolution of the effective radiative forcing of aerosols, while the DAMIP hist-aer experiment can be used to diagnose the climate response to that forcing.

High-resolution climate model simulations are often able to better represent phenomena relevant to climate extremes, such as tropical cyclones, atmospheric rivers, and heat waves and heavy rainfall in mountainous regions (e.g. Roberts et al., 2025). Such events are often the focus of extreme event attribution studies. The HighResMIP2 (Roberts et al., 2025) 1950s control (control-1950), the historical simulation beginning in 1950 (hist-1950) and its extension to 2100 with the Medium scenario will be valuable in supporting extreme event attribution for such variables, with allowance made for the fact that the 1950s control is not a pre-industrial control. We also encourage modelling groups to carry out DAMIP experiments with high-resolution models to the extent possible but realise that computational constraints may prevent this. Finally, we note that the HighResMIP2 choice of the Medium scenario to extend historical simulations to 2100 aligns with our choice of the same scenario in DAMIP v2.0.

## 5 Variables requested by DAMIP

DAMIP has provided input to the harmonised data request for DECK and Assessment Fast Track experiments through the proposal of the “detection and attribution” opportunity. This opportunity contains basic variables that are needed to quantify how the mean climate and its variability change over time, to understand the mechanisms involved and to compare this with the observational record with a view to detecting and attributing climate change signals. Many of these variables overlap with the proposed baseline variable set for

Earth system modelling (Juckes et al., 2025). These proposed variables include fields for assessing the different forcings that the climate system is experiencing, fields for assessing global mean temperature, hydrological, sea level and circulation changes, top-of-atmosphere and surface fluxes for diagnosing energy balances and fields needed to understand the role of clouds in the climate system. With the move towards a greater role for emissions-driven simulations, we also request variables that will help us understand the origins of differences in CO<sub>2</sub> concentrations between simulations when run in emissions-driven mode. A variety of daily fields are also requested to understand the time evolution of extremes and to diagnose synoptic features and daily surface fluxes that could be used to understand the drivers of these extremes. Finally, DAMIP requests zonal mean or 3D atmospheric temperature with a high enough vertical resolution extending deep into the stratosphere to allow comparison with Stratospheric Sounding Unit (SSU) observations (e.g. Mitchell, 2016; Santer et al., 2023). Stratospheric temperature trends are an important part of the climate change signal and are also particularly important for assessing ozone recovery, volcanic aerosol and solar signals (Mitchell, 2016; Santer et al., 2023).

We propose that the “detection and attribution” variable groups be outputted for all DAMIP experiments and related experiments from the DECK and other MIPs shown in Fig. 2. We also suggest that these be outputted for the pre-industrial control experiments to allow the uncertainties due to internal variability alone to be quantified, together with the Atmosphere Model Intercomparison Project (AMIP) experiments, which can be used to explore the role of observed historical trends in SSTs in the evolution of the climate system and in potential differences between coupled models and observations. Since most of the variables proposed as part of the “detection and attribution” opportunity are basic fields needed to characterise the behaviour of the climate system, we also suggest that these variables be outputted for the more idealised experiments that are part of DECK and Assessment Fast Track, which can be used to understand the behaviour of models and to explore the direct effects of radiative forcings and the indirect effects of SST warming on the climate system. These include 1pctCO<sub>2</sub>, 1pctCO<sub>2</sub>-bgc, 1pctCO<sub>2</sub>-rad, abrupt-4xCO<sub>2</sub>, piClim-anthro, hist-piSLCF, amip-p4k and amip-piForcing. We also suggest that these variables be saved for the DCP-initialised predictions from 2025 to 2036 to compare simulated near-term trends in these predictions with those in free-running coupled simulations, allowing for attribution of predicted trends in the climate system to external forcings as well as aspects of variability and change that are imparted to the models through the initial conditions in this particular experiment.

## 6 Summary

This paper describes the Detection and Attribution Model Intercomparison Project (DAMIP v2.0) which will form part of CMIP7 (Dunne et al., 2024) and is intended to underpin detection and attribution analyses informing the IPCC Seventh Assessment Report. DAMIP v1.0 simulations in CMIP6 were used extensively in the fields of attribution of climate trends, extreme event attribution and attribution of climate impacts, and they also underpinned the assessment of human-induced global warming in the IPCC AR6 (IPCC, 2021), which was reported in the Glasgow Climate Pact (UNFCCC, 2022).

DAMIP v2.0 again proposes hist-nat, hist-GHG and hist-aer simulations as high-priority Assessment Fast Track simulations for CMIP7. These simulations were the most heavily used in CMIP6. We propose that they be extended to 2035 using the Medium scenario from ScenarioMIP. However, we also propose a set of historical simulations with forcings which together form the complete set of historical forcings (hist-GHG, hist-aer, hist-nat, hist-lu and hist-O3), allowing additivity to be easily tested and ensuring all forcings are covered. Given that stratospheric and tropospheric ozone changes interact with each other, and to simplify the experimental design, we propose hist-O3 simulations which have prescribed ozone changes in the stratosphere and troposphere rather than just in the stratosphere, as was the case for CMIP6. To evaluate the effects of individual forcings on future climate evolution, we propose an extension of key simulations to 2100 with the Medium emissions scenario. Finally, for CMIP7 we propose a new set of interactive-CO<sub>2</sub> simulations, which will among other things allow the net effect of land use change on the climate to be evaluated and allow the effects of carbon-climate feedbacks on the simulated response to individual forcings to be evaluated.

Updated details on the project and its progress will be available at <https://wcrp-cmip.org/mips/damip/> (last access: 14 July 2025).

**Code availability.** The full CanESM5.0 source code used to run the simulations shown in Fig. 3 is publicly available at <https://gitlab.com/ccma/canesm> (last access: 30 January 2025). The version of the code which can be used to produce all of the simulations described in this paper is tagged as v5.0.3 and has the associated DOI <https://doi.org/10.5281/zenodo.3251113> (Swart et al., 2019).

**Data availability.** The model output from the DAMIP simulations described in this paper will be distributed through the Earth System Grid Federation (ESGF) with DOIs assigned. The model output will be freely accessible through data portals after registration.

**Author contributions.** As members of the Scientific Steering Committee of DAMIP v2.0, NPG, IRS, GH, RK, DM, AR, HS, DS,

CT, PW and WZ together developed the experimental design for DAMIP v2.0. NPG prepared Figs. 1 and 3, and IRS prepared Fig. 2. VKA carried out the interactive-CO<sub>2</sub> simulations shown in Fig. 3. NPG led the writing of the manuscript, and all of the authors contributed to editing and revising the text.

**Competing interests.** The contact author has declared that none of the authors has any competing interests.

**Disclaimer.** Publisher's note: Copernicus Publications remains neutral with regard to jurisdictional claims made in the text, published maps, institutional affiliations, or any other geographical representation in this paper. While Copernicus Publications makes every effort to include appropriate place names, the final responsibility lies with the authors.

**Financial support.** Isla Simpson was supported by the NSF National Center for Atmospheric Research, which is a major facility sponsored by the NSF under Cooperative Agreement No. 1852977. Claudia Tebaldi was supported by the Office of Science, U.S. Department of Energy Biological and Environmental Research, as part of the Water Cycle and Climate Extremes Modeling (WACCEM) project funded by the Regional and Global Model Analysis programme area. Pacific Northwest National Laboratory is operated by Battelle for the U.S. Department of Energy under contract no. DE-AC05-76RL01830. Dáithí Stone was supported by the Ministry of Business, Innovation and Employment (MBIE), Aotearoa New Zealand, through the Whakahura Endeavour Programme.

**Review statement.** This paper was edited by Lele Shu and reviewed by Michael Wehner and one anonymous referee.

## References

- Allen, M. R., Stott, P. A., Mitchell, J. F. B., Schnur, R., and Delworth, T. L.: Quantifying the uncertainty in forecasts of anthropogenic climate change, *Nature*, 407, 617–620, <https://doi.org/10.1038/35036559>, 2000.
- Canadell, J. G., Monteiro, P. M. S., Costa, M. H., Cotrim da Cunha, L., Cox, P. M., Eliseev, A. V., Henson, S., Ishii, M., Jaccard, S., Koven, C., Lohila, A., Patra, P. K., Piao, S., Rogelj, J., Syampungani, S., Zaehle, S., and Zickfeld, K.: Global Carbon and other Biogeochemical Cycles and Feedbacks, edited by: Masson-Delmotte, V., Zhai, P., Pirani, A., Connors, S. L., Péan, C., Berger, S., Caud, N., Chen, Y., Goldfarb, L., Gomis, M. I., Huang, M., Leitzell, K., Lonnoy, E., Matthews, J. B. R., Maycock, T. K., Waterfield, T., Yelekçi, O., Yu, R., and Zhou, B., *Climate Change 2021: The Physical Science Basis. Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change*, Cambridge University Press, Cambridge, UK and New York, NY, USA, 673–816, <https://doi.org/10.1017/9781009157896.007>, 2021.

- Carlson, C. J.: After millions of preventable deaths, climate change must be treated like a health emergency, *Nat. Med.*, 30, 622–622, <https://doi.org/10.1038/s41591-023-02765-y>, 2024.
- Chapman, S., Birch, C. E., Marsham, J. H., Part, C., Hajat, S., Chersich, M. F., Ebi, K. L., Luchters, S., Nakstad, B., and Kovats, S.: Past and projected climate change impacts on heat-related child mortality in Africa, *Environ. Res. Lett.*, 17, 074028, <https://doi.org/10.1088/1748-9326/ac7ac5>, 2022.
- Chiang, F., Mazdiyasi, O., and AghaKouchak, A.: Evidence of anthropogenic impacts on global drought frequency, duration, and intensity, *Nat. Commun.*, 12, 2754, <https://doi.org/10.1038/s41467-021-22314-w>, 2021.
- Christidis, N., Mitchell, D., and Stott, P. A.: Rapidly increasing likelihood of exceeding 50 °C in parts of the Mediterranean and the Middle East due to human influence, *npj Clim. Atmos. Sci.*, 6, 1–12, <https://doi.org/10.1038/s41612-023-00377-4>, 2023.
- Doblas-Reyes, F. J., Sörensson, A. A., Almazroui, M., Dosio, A., Gutowski, W. J., Haarsma, R., Hamdi, R., Hewitson, B., Kwon, W.-T., Lamptey, B. L., Maraun, D., Stephenson, T. S., Takayabu, I., Terray, L., Turner, A., and Zuo, Z.: Linking Global to Regional Climate Change, edited by: Masson-Delmotte, V., Zhai, P., Pirani, A., Connors, S. L., Péan, C., Berger, S., Caud, N., Chen, Y., Goldfarb, L., Gomis, M. I., Huang, M., Leitzell, K., Lonnoy, E., Matthews, J. B. R., Maycock, T. K., Waterfield, T., Yelekçi, O., Yu, R., and Zhou, B., *Climate Change 2021: The Physical Science Basis. Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change*, Cambridge University Press, Cambridge, UK and New York, NY, USA, 1363–1512, <https://doi.org/10.1017/9781009157896.012>, 2021.
- Dong, B., Sutton, R. T., Shaffrey, L., and Harvey, B.: Recent decadal weakening of the summer Eurasian westerly jet attributable to anthropogenic aerosol emissions, *Nat Commun.*, 13, 1148, <https://doi.org/10.1038/s41467-022-28816-5>, 2022.
- Dunne, J. P., Hewitt, H. T., Arblaster, J., Bonou, F., Boucher, O., Cavazos, T., Durack, P. J., Hassler, B., Jukes, M., Miyakawa, T., Mizielinski, M., Naik, V., Nicholls, Z., O'Rourke, E., Pincus, R., Sanderson, B. M., Simpson, I. R., and Taylor, K. E.: An evolving Coupled Model Intercomparison Project phase 7 (CMIP7) and Fast Track in support of future climate assessment, *EGU Sphere* [preprint], <https://doi.org/10.5194/egusphere-2024-3874>, 2024.
- Eyring, V., Bony, S., Meehl, G. A., Senior, C. A., Stevens, B., Stouffer, R. J., and Taylor, K. E.: Overview of the Coupled Model Intercomparison Project Phase 6 (CMIP6) experimental design and organization, *Geosci. Model Dev.*, 9, 1937–1958, <https://doi.org/10.5194/gmd-9-1937-2016>, 2016.
- Eyring, V., Gillett, N. P., Achuta Rao, K. M., Barimalala, R., Barreiro Parrillo, M., Bellouin, N., Cassou, C., Durack, P. J., Kosaka, Y., McGregor, S., Min, S., Morgenstern, O., and Sun, Y.: Human Influence on the Climate System, edited by: Masson-Delmotte, V., Zhai, P., Pirani, A., Connors, S. L., Péan, C., Berger, S., Caud, N., Chen, Y., Goldfarb, L., Gomis, M. I., Huang, M., Leitzell, K., Lonnoy, E., Matthews, J. B. R., Maycock, T. K., Waterfield, T., Yelekçi, O., Yu, R., and Zhou, B., *Climate Change 2021: The Physical Science Basis. Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change*, Cambridge University Press, Cambridge, UK and New York, NY, USA, 423–552, <https://doi.org/10.1017/9781009157896.005>, 2021.
- Fischer, E. M., Beyerle, U., Bloin-Wibe, L., Gessner, C., Humphrey, V., Lehner, F., Pendergrass, A. G., Sippel, S., Zeder, J., and Knutti, R.: Storylines for unprecedented heatwaves based on ensemble boosting, *Nat. Commun.*, 14, 4643, <https://doi.org/10.1038/s41467-023-40112-4>, 2023.
- Forster, P., Storelvmo, T., Armour, K., Collins, W., Dufresne, J.-L., Frame, D., Lunt, D. J., Mauritsen, T., Palmer, M. D., Watanabe, M., Wild, M., and Zhang, H.: *The Earth's Energy Budget, Climate Feedbacks, and Climate Sensitivity*, edited by: Masson-Delmotte, V., Zhai, P., Pirani, A., Connors, S. L., Péan, C., Berger, S., Caud, N., Chen, Y., Goldfarb, L., Gomis, M. I., Huang, M., Leitzell, K., Lonnoy, E., Matthews, J. B. R., Maycock, T. K., Waterfield, T., Yelekçi, O., Yu, R., and Zhou, B., *Climate Change 2021: The Physical Science Basis. Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change*, Cambridge University Press, Cambridge, UK and New York, NY, USA, 923–1054, <https://doi.org/10.1017/9781009157896.009>, 2021.
- Forster, P. M., Smith, C., Walsh, T., Lamb, W. F., Lamboll, R., Hall, B., Hauser, M., Ribes, A., Rosen, D., Gillett, N. P., Palmer, M. D., Rogelj, J., von Schuckmann, K., Trewin, B., Allen, M., Andrew, R., Betts, R. A., Borger, A., Boyer, T., Broersma, J. A., Buontempo, C., Burgess, S., Cagnazzo, C., Cheng, L., Friedlingstein, P., Gettelman, A., Gütschow, J., Ishii, M., Jenkins, S., Lan, X., Morice, C., Mühle, J., Kadow, C., Kennedy, J., Killock, R. E., Krummel, P. B., Minx, J. C., Myhre, G., Naik, V., Peters, G. P., Pirani, A., Pongratz, J., Schleussner, C.-F., Seneviratne, S. I., Szopa, S., Thorne, P., Kovilakam, M. V. M., Majamäki, E., Jalkanen, J.-P., van Marle, M., Hoesly, R. M., Rohde, R., Schumacher, D., van der Werf, G., Vose, R., Zickfeld, K., Zhang, X., Masson-Delmotte, V., and Zhai, P.: Indicators of Global Climate Change 2023: annual update of key indicators of the state of the climate system and human influence, *Earth Syst. Sci. Data*, 16, 2625–2658, <https://doi.org/10.5194/essd-16-2625-2024>, 2024.
- Fox-Kemper, B., Hewitt, H. T., Xiao, C., Aðalgeirsdóttir, G., Drijfhout, S. S., Edwards, T. L., Golledge, N. R., Hemer, M., Kopp, R. E., Krinner, G., Mix, A., Notz, D., Nowicki, S., Nurhati, I. S., Ruiz, L., Sallée, J.-B., Slangen, A. B. A., and Yu, Y.: *Ocean, Cryosphere and Sea Level Change*, edited by: Masson-Delmotte, V., Zhai, P., Pirani, A., Connors, S. L., Péan, C., Berger, S., Caud, N., Chen, Y., Goldfarb, L., Gomis, M. I., Huang, M., Leitzell, K., Lonnoy, E., Matthews, J. B. R., Maycock, T. K., Waterfield, T., Yelekçi, O., Yu, R., and Zhou, B., *Climate Change 2021: The Physical Science Basis. Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change*, Cambridge University Press, Cambridge, UK and New York, NY, USA, 1211–1362, <https://doi.org/10.1017/9781009157896.011>, 2021.
- Fyfe, J. C., Kharin, V. V., Santer, B. D., Cole, J. N. S., and Gillett, N. P.: Significant impact of forcing uncertainty in a large ensemble of climate model simulations, *P. Natl. Acad. Sci. USA*, 118, e2016549118, <https://doi.org/10.1073/pnas.2016549118>, 2021.
- Garny, H., Hendon, H., Abalos, M., Chiodo, G., Purich, A., Randel, W. J., Smith, K., and Thompson, D.: Stratospheric ozone changes and climate, in: *World Meteorological Organisation Scientific assessment of ozone depletion: 2022*, WMO, 978-9914-733-97-6, WMO, Geneva, 271–323, ISBN 978-9914-733-97-6, 2022.
- Gillett, N. P., Fyfe, J. C., and Parker, D. E.: Attribution of observed sea level pressure trends to greenhouse gas, aerosol,

- and ozone changes, *Geophys. Res. Lett.*, 40, 2302–2306, <https://doi.org/10.1002/grl.50500>, 2013.
- Gillett, N. P., Shiogama, H., Funke, B., Hegerl, G., Knutti, R., Matthes, K., Santer, B. D., Stone, D., and Tebaldi, C.: The Detection and Attribution Model Intercomparison Project (DAMIP v1.0) contribution to CMIP6, *Geosci. Model Dev.*, 9, 3685–3697, <https://doi.org/10.5194/gmd-9-3685-2016>, 2016.
- Gillett, N. P., Kirchmeier-Young, M., Ribes, A., Shiogama, H., Hegerl, G. C., Knutti, R., Gastineau, G., John, J. G., Li, L., Nazarenko, L., Rosenbloom, N., Seland, Ø., Wu, T., Yukimoto, S., and Ziehn, T.: Constraining human contributions to observed warming since the pre-industrial period, *Nat. Clim. Change*, 11, 207–212, <https://doi.org/10.1038/s41558-020-00965-9>, 2021.
- Hannart, A., Pearl, J., Otto, F. E. L., Naveau, P., and Ghil, M.: Causal Counterfactual Theory for the Attribution of Weather and Climate-Related Events, *B. Am. Meteorol. Soc.*, 97, 99–110, <https://doi.org/10.1175/BAMS-D-14-00034.1>, 2016.
- Hardiman, S. C., Andrews, M. B., Andrews, T., Bushell, A. C., Dunstone, N. J., Dyson, H., Jones, G. S., Knight, J. R., Neiningner, E., O'Connor, F. M., Ridley, J. K., Ringer, M. A., Scaife, A. A., Senior, C. A., and Wood, R. A.: The impact of prescribed ozone in climate projections run with HadGEM3-GC3.1, *J. Adv. Model Earth Sy.*, 11, 3443–3453, <https://doi.org/10.1029/2019MS001714>, 2019.
- Hassler, B., Young, P. J., Ball, W., Damadeo, R., Keeble, J., Mailard Barras, E., Sofieva, V., and Zeng, G.: Update on global ozone: Past, present and future, in: *World Meteorological Organisation (WMO) Scientific Assessment of Ozone Depletion: 2022*, WMO, 978-9914-733-97-6, WMO, Geneva, 153–214, ISBN 978-9914-733-97-6, 2022.
- Herring, S. C., Christidis, N., Hoell, A., and Stott, P. A.: Explaining Extreme Events of 2020 from a Climate Perspective, *B. Am. Meteorol. Soc.*, 103, S1–S129, <https://doi.org/10.1175/BAMS-ExplainingExtremeEvents2020.1>, 2022.
- Holland, M. M., Hannay, C., Fasullo, J., Jahn, A., Kay, J. E., Mills, M., Simpson, I. R., Wieder, W., Lawrence, P., Kluzek, E., and Bailey, D.: New model ensemble reveals how forcing uncertainty and model structure alter climate simulated across CMIP generations of the Community Earth System Model, *Geosci. Model Dev.*, 17, 1585–1602, <https://doi.org/10.5194/gmd-17-1585-2024>, 2024.
- Hourdin, F., Mauritsen, T., Gettelman, A., Golaz, J.-C., Balaji, V., Duan, Q., Folini, D., Ji, D., Klocke, D., Qian, Y., Rauser, F., Rio, C., Tomassini, L., Watanabe, M., and Williamson, D.: The art and science of climate model tuning, *B. Am. Meteorol. Soc.*, 98, 589–602, <https://doi.org/10.1175/BAMS-D-15-00135.1>, 2017.
- IPCC: Summary for Policymakers, edited by: Shukla, P. R., Skea, J., Calvo Buendia, E., Masson-Delmotte, V., Pörtner, H.-O., Roberts, D. C., Zhai, P., Slade, R., Connors, S., van Diemen, R., Ferrat, M., Haughey, E., Luz, S., Neogi, S., Pathak, M., Petzold, J., Portugal Pereira, J., Vyas, P., Huntley, E., Kissick, K., Belkacemi, M., and Malley, J., *Climate Change and Land: IPCC Special Report on Climate Change, Desertification, Land Degradation, Sustainable Land Management, Food Security, and Greenhouse Gas Fluxes in Terrestrial Ecosystems*, Cambridge University Press, Cambridge, UK and New York, NY, USA, 1–36, <https://doi.org/10.1017/9781009157988>, 2019.
- IPCC: Summary for Policymakers, edited by: Masson-Delmotte, V., Zhai, P., Pirani, A., Connors, S. L., Péan, C., Berger, S., Caud, N., Chen, Y., Goldfarb, L., Gomis, M. I., Huang, M., Leitzell, K., Lonnoy, E., Matthews, J. B. R., Maycock, T. K., Waterfield, T., Yelekçi, O., Yu, R., and Zhou, B., *Climate Change 2021: The Physical Science Basis. Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change*, Cambridge University Press, Cambridge, UK and New York, NY, USA, 3–32, <https://doi.org/10.1017/9781009157896.001>, 2021.
- James, R. A., Jones, R. G., Boyd, E., Young, H. R., Otto, F. E. L., Huggel, C., and Fuglestedt, J. S.: Attribution: How Is It Relevant for Loss and Damage Policy and Practice?, in: *Loss and Damage from Climate Change*, edited by: Mechler, R., Bouwer, L. M., Schinko, T., Surminski, S., and Linnerooth-Bayer, J., Springer International Publishing, Cham, 113–154, [https://doi.org/10.1007/978-3-319-72026-5\\_5](https://doi.org/10.1007/978-3-319-72026-5_5), 2019.
- Jenkins, S., Smith, C., Allen, M., and Grainger, R.: Tonga eruption increases chance of temporary surface temperature anomaly above 1.5 °C, *Nat. Clim. Change*, 13, 127–129, <https://doi.org/10.1038/s41558-022-01568-2>, 2023.
- Jones, C. D., Hickman, J. E., Rumbold, S. T., Walton, J., Lamboll, R. D., Skeie, R. B., Fiedler, S., Forster, P. M., Rogelj, J., Abe, M., Botzet, M., Calvin, K., Cassou, C., Cole, J. N. S., Davini, P., Deushi, M., Dix, M., Fyfe, J. C., Gillett, N. P., Ilyina, T., Kawamiya, M., Kelley, M., Kharin, S., Koshiro, T., Li, H., Mackallah, C., Müller, W. A., Nabat, P., Van Noije, T., Nolan, P., Ohgaito, R., Olivié, D., Oshima, N., Parodi, J., Reerink, T. J., Ren, L., Romanou, A., Séférian, R., Tang, Y., Timmreck, C., Tjiputra, J., Tourigny, E., Tsigaridis, K., Wang, H., Wu, M., Wyser, K., Yang, S., Yang, Y., and Ziehn, T.: The Climate Response to Emissions Reductions Due to COVID-19: Initial Results From CovidMIP, *Geophys. Res. Lett.*, 48, e2020GL091883, <https://doi.org/10.1029/2020GL091883>, 2021.
- Jukes, M., Taylor, K. E., Antonio, F., Brayshaw, D., Buontempo, C., Cao, J., Durack, P. J., Kawamiya, M., Kim, H., Lovato, T., Mackallah, C., Mizieliński, M., Nuzzo, A., Stockhause, M., Visionsi, D., Walton, J., Turner, B., O'Rourke, E., and Dingley, B.: Baseline Climate Variables for Earth System Modelling, *Geosci. Model Dev.*, 18, 2639–2663, <https://doi.org/10.5194/gmd-18-2639-2025>, 2025.
- Kandlbauer, J., Hopcroft, P. O., Valdes, P. J., and Sparks, R. S. J.: Climate and carbon cycle response to the 1815 Tambora volcanic eruption, *J. Geophys. Res.-Atmos.*, 118, 12497–12507, <https://doi.org/10.1002/2013JD019767>, 2013.
- Kang, J. M., Shaw, T. A., Kang, S. M., Simpson, I. R., and Yu, Y.: Revisiting the reanalysis-model discrepancy in Southern Hemisphere winter storm track trends, *npj Clim. Atmos. Sci.*, 7, 1–10, <https://doi.org/10.1038/s41612-024-00801-3>, 2024.
- King, A. D.: Attributing Changing Rates of Temperature Record Breaking to Anthropogenic Influences, *Earth's Future*, 5, 1156–1168, <https://doi.org/10.1002/2017EF000611>, 2017.
- Lanet, M., Li, L., Ehret, A., Turquet, S., and Le Treut, H.: Attribution of summer 2022 extreme wildfire season in Southwest France to anthropogenic climate change, *npj Clim. Atmos. Sci.*, 7, 1–10, <https://doi.org/10.1038/s41612-024-00821-z>, 2024.
- Lawrence, D. M., Hurtt, G. C., Arneth, A., Brovkin, V., Calvin, K. V., Jones, A. D., Jones, C. D., Lawrence, P. J., de Noblet-Ducoudré, N., Pongratz, J., Seneviratne, S. I., and Shevliakova, E.: The Land Use Model Intercomparison Project (LUMIP) contribution to CMIP6: rationale and experimental design, *Geosci.*

- Model Dev., 9, 2973–2998, <https://doi.org/10.5194/gmd-9-2973-2016>, 2016.
- Le Quéré, C., Rödenbeck, C., Buitenhuis, E. T., Conway, T. J., Langenfelds, R., Gomez, A., Labuschagne, C., Ramonet, M., Nakazawa, T., Metz, N., Gillett, N., and Heimann, M.: Saturation of the Southern Ocean CO<sub>2</sub> Sink Due to Recent Climate Change, *Science*, 316, 1735–1738, <https://doi.org/10.1126/science.1136188>, 2007.
- Lee, J.-Y., Marotzke, J., Bala, G., Cao, L., Corti, S., Dunne, J. P., Engelbrecht, F., Fischer, E., Fyfe, J. C., Jones, C., Maycock, A., Mutemi, J., Ndiaye, O., Panickal, S., and Zhou, T.: Future Global Climate: Scenario-Based Projections and Near-Term Information, edited by: Masson-Delmotte, V., Zhai, P., Pirani, A., Connors, S. L., Péan, C., Berger, S., Caud, N., Chen, Y., Goldfarb, L., Gomis, M. I., Huang, M., Leitzell, K., Lonnoy, E., Matthews, J. B. R., Maycock, T. K., Waterfield, T., Yelekçi, O., Yu, R., and Zhou, B., *Climate Change 2021: The Physical Science Basis. Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change*, Cambridge University Press, Cambridge, UK and New York, NY, USA, 553–672, <https://doi.org/10.1017/9781009157896.006>, 2021.
- Li, X., Zhang, L., Wang, G., Cao, H., Zhang, H., Jia, B., Zhou, Z., Liu, L., and Zhang, L.: Anthropogenic forcing decreases the probability of the 2020 Yangtze River extreme flood and future risk, *Atmos. Res.*, 311, 107662, <https://doi.org/10.1016/j.atmosres.2024.107662>, 2024a.
- Li, Y., Xia, Y., Xie, F., and Yan, Y.: Influence of stratosphere-troposphere exchange on long-term trends of surface ozone in CMIP6, *Atmos. Res.*, 297, 107086, <https://doi.org/10.1016/j.atmosres.2023.107086>, 2024b.
- Liddicoat, S. K., Wiltshire, A. J., Jones, C. D., Arora, V. K., Brovkin, V., Cadule, P., Hajima, T., Lawrence, D. M., Pongratz, J., Schwinger, J., Séférian, R., Tjiputra, J. F., and Ziehn, T.: Compatible Fossil Fuel CO<sub>2</sub> Emissions in the CMIP6 Earth System Models' Historical and Shared Socioeconomic Pathway Experiments of the Twenty-First Century, *J. Climate*, 34, 2853–2875, <https://doi.org/10.1175/JCLI-D-19-0991.1>, 2021.
- Liu, W., Hegglin, M. I., Checa-Garcia, R., Li, S., Gillett, N. P., Lyu, K., Zhang, X., and Swart, N. C.: Stratospheric ozone depletion and tropospheric ozone increases drive Southern Ocean interior warming, *Nat. Clim. Change*, 12, 365–372, <https://doi.org/10.1038/s41558-022-01320-w>, 2022.
- Luo, X., Ge, J., Cao, Y., Liu, Y., Yang, L., Wang, S., and Guo, W.: Local and Nonlocal Biophysical Effects of Historical Land Use and Land Cover Changes in CMIP6 Models and the Intermodel Uncertainty, *Earth's Future*, 12, e2023EF004220, <https://doi.org/10.1029/2023EF004220>, 2024.
- Marvel, K., Schmidt, G. A., Shindell, D., Bonfils, C., LeGrande, A. N., Nazarenko, L., and Tsigaridis, K.: Do responses to different anthropogenic forcings add linearly in climate models?, *Environ. Res. Lett.*, 10, 104010, <https://doi.org/10.1088/1748-9326/10/10/104010>, 2015.
- Matthews, H. D. and Wynes, S.: Current global efforts are insufficient to limit warming to 1.5 °C, *Science*, 376, 1404–1409, <https://doi.org/10.1126/science.abo3378>, 2022.
- Menary, M. B., Robson, J., Allan, R. P., Booth, B. B. B., Cassou, C., Gastineau, G., Gregory, J., Hodson, D., Jones, C., Mignot, J., Ringer, M., Sutton, R., Wilcox, L., and Zhang, R.: Aerosol-Forced AMOC Changes in CMIP6 Historical Simulations, *Geophys. Res. Lett.*, 47, e2020GL088166, <https://doi.org/10.1029/2020GL088166>, 2020.
- Mitchell, D. M.: Attributing the forced components of observed stratospheric temperature variability to external drivers, *Q. J. Roy. Meteor. Soc.*, 142, 1041–1047, <https://doi.org/10.1002/qj.2707>, 2016.
- Morgenstern, O.: The Southern Annular Mode in 6th Coupled Model Intercomparison Project Models, *J. Geophys. Res.-Atmos.*, 126, e2020JD034161, <https://doi.org/10.1029/2020JD034161>, 2021.
- Naveau, P., Hannart, A., and Ribes, A.: Statistical Methods for Extreme Event Attribution in Climate Science, *Annu. Rev. Stat. Appl.*, 7, 89–110, <https://doi.org/10.1146/annurev-statistics-031219-041314>, 2020.
- O'Neill, B. C., Tebaldi, C., van Vuuren, D. P., Eyring, V., Friedlingstein, P., Hurtt, G., Knutti, R., Kriegler, E., Lamarque, J.-F., Lowe, J., Meehl, G. A., Moss, R., Riahi, K., and Sanderson, B. M.: The Scenario Model Intercomparison Project (ScenarioMIP) for CMIP6, *Geosci. Model Dev.*, 9, 3461–3482, <https://doi.org/10.5194/gmd-9-3461-2016>, 2016.
- Pincus, R., Forster, P. M., and Stevens, B.: The Radiative Forcing Model Intercomparison Project (RFMIP): experimental protocol for CMIP6, *Geosci. Model Dev.*, 9, 3447–3460, <https://doi.org/10.5194/gmd-9-3447-2016>, 2016.
- Rantanen, M. and Laaksonen, A.: The jump in global temperatures in September 2023 is extremely unlikely due to internal climate variability alone, *npj Clim. Atmos. Sci.*, 7, 1–4, <https://doi.org/10.1038/s41612-024-00582-9>, 2024.
- Ribes, A., Qasmi, S., and Gillett, N. P.: Making climate projections conditional on historical observations, *Sci. Adv.*, 7, eabc0671, <https://doi.org/10.1126/sciadv.abc0671>, 2021.
- Roberts, M. J., Reed, K. A., Bao, Q., Barsugli, J. J., Camargo, S. J., Caron, L.-P., Chang, P., Chen, C.-T., Christensen, H. M., Danabasoglu, G., Frenger, I., Fučkar, N. S., ul Hasson, S., Hewitt, H. T., Huang, H., Kim, D., Kodama, C., Lai, M., Leung, L.-Y. R., Mizuta, R., Nobre, P., Ortega, P., Paquin, D., Roberts, C. D., Scoccimarro, E., Seddon, J., Treguier, A. M., Tu, C.-Y., Ullrich, P. A., Vidale, P. L., Wehner, M. F., Zarzycki, C. M., Zhang, B., Zhang, W., and Zhao, M.: High-Resolution Model Intercomparison Project phase 2 (HighResMIP2) towards CMIP7, *Geosci. Model Dev.*, 18, 1307–1332, <https://doi.org/10.5194/gmd-18-1307-2025>, 2025.
- Rothenberg, D., Mahowald, N., Lindsay, K., Doney, S. C., Moore, J. K., and Thornton, P.: Volcano impacts on climate and biogeochemistry in a coupled carbon–climate model, *Earth Syst. Dynam.*, 3, 121–136, <https://doi.org/10.5194/esd-3-121-2012>, 2012.
- Sanderson, B. M., Booth, B. B. B., Dunne, J., Eyring, V., Fisher, R. A., Friedlingstein, P., Gidden, M. J., Hajima, T., Jones, C. D., Jones, C. G., King, A., Koven, C. D., Lawrence, D. M., Lowe, J., Mengis, N., Peters, G. P., Rogelj, J., Smith, C., Snyder, A. C., Simpson, I. R., Swann, A. L. S., Tebaldi, C., Ilyina, T., Schleussner, C.-F., Séférian, R., Samset, B. H., van Vuuren, D., and Zaehle, S.: The need for carbon-emissions-driven climate projections in CMIP7, *Geosci. Model Dev.*, 17, 8141–8172, <https://doi.org/10.5194/gmd-17-8141-2024>, 2024.
- Santer, B. D., Bonfils, C., Painter, J. F., Zelinka, M. D., Mears, C., Solomon, S., Schmidt, G. A., Fyfe, J. C., Cole, J. N. S., Nazarenko, L., Taylor, K. E., and Wentz, F. J.: Volcanic con-

- tribution to decadal changes in tropospheric temperature, *Nat. Geosci.*, 7, 185–189, <https://doi.org/10.1038/ngeo2098>, 2014.
- Santer, B. D., Po-Chedley, S., Zhao, L., Zou, C.-Z., Fu, Q., Solomon, S., Thompson, D. W. J., Mears, C., and Taylor, K. E.: Exceptional stratospheric contribution to human fingerprints on atmospheric temperature, *P. Natl. Acad. Sci.*, 120, e2300758120, <https://doi.org/10.1073/pnas.2300758120>, 2023.
- Schaller, N., Otto, F., Van Oldenborgh, G. J., Massey, N., Sparrow, S., and Allen, M.: The heavy precipitation event of May–June 2013 in the upper Danube and Elbe basins, *B. Am. Meteorol. Soc.*, 95, S69–S72, 2014.
- Shiogama, H., Stone, D. A., Nagashima, T., Nozawa, T., and Emori, S.: On the linear additivity of climate forcing-response relationships at global and continental scales, *Int. J. Climatol.*, 33, 2542–2550, <https://doi.org/10.1002/joc.3607>, 2013.
- Shiogama, H., Tatebe, H., Hayashi, M., Abe, M., Arai, M., Koyama, H., Imada, Y., Kosaka, Y., Ogura, T., and Watanabe, M.: MIROC6 Large Ensemble (MIROC6-LE): experimental design and initial analyses, *Earth Syst. Dynam.*, 14, 1107–1124, <https://doi.org/10.5194/esd-14-1107-2023>, 2023.
- Simpson, I. R., Rosenbloom, N., Danabasoglu, G., Deser, C., Yeager, S. G., McCluskey, C. S., Yamaguchi, R., Lamarque, J.-F., Tilmes, S., Mills, M. J., and Rodgers, K. B.: The CESM2 Single-Forcing Large Ensemble and Comparison to CESM1: Implications for Experimental Design, *J. Climate*, 36, 5687–5711, <https://doi.org/10.1175/JCLI-D-22-0666.1>, 2023.
- Sippel, S., Mitchell, D., Black, M. T., Dittus, A. J., Harrington, L., Schaller, N., and Otto, F. E. L.: Combining large model ensembles with extreme value statistics to improve attribution statements of rare events, *Weather and Climate Extremes*, 9, 25–35, <https://doi.org/10.1016/j.wace.2015.06.004>, 2015.
- Smith, D. M., Gillett, N. P., Simpson, I. R., Athanasiadis, P. J., Baehr, J., Bethke, I., Bilge, T. A., Bonnet, R., Boucher, O., Findell, K. L., Gastineau, G., Gualdi, S., Hermanson, L., Leung, L. R., Mignot, J., Müller, W. A., Osprey, S., Otterå, O. H., Persad, G. G., Scaife, A. A., Schmidt, G. A., Shiogama, H., Sutton, R. T., Swingedouw, D., Yang, S., Zhou, T., and Ziehn, T.: Attribution of multi-annual to decadal changes in the climate system: The Large Ensemble Single Forcing Model Intercomparison Project (LESFMI), *Front. Clim.*, 4, 955414, <https://doi.org/10.3389/fclim.2022.955414>, 2022.
- Stott, P. A. and Kettleborough, J. A.: Origins and estimates of uncertainty in predictions of twenty-first century temperature rise, *Nature*, 416, 723–726, <https://doi.org/10.1038/416723a>, 2002.
- Stott, P. A., Stone, D. A., and Allen, M. R.: Human contribution to the European heatwave of 2003, *Nature*, 432, 610–614, <https://doi.org/10.1038/nature03089>, 2004.
- Swart, N. C., Cole, J. N. S., Kharin, V. V., Lazare, M., Scinocca, J. F., Gillett, N. P., Anstey, J., Arora, V., Christian, J. R., Hanna, S., Jiao, Y., Lee, W. G., Majaess, F., Saenko, O. A., Seiler, C., Seinen, C., Shao, A., Sigmond, M., Solheim, L., von Salzen, K., Yang, D., and Winter, B.: The Canadian Earth System Model version 5 (CanESM5.0.3), *Geosci. Model Dev.*, 12, 4823–4873, <https://doi.org/10.5194/gmd-12-4823-2019>, 2019.
- Szopa, S., Naik, V., Adhikary, B., Artaxo, P., Berntsen, T., Collins, W. D., Fuzzi, S., Gallardo, L., Kiendler-Scharr, A., Klimont, Z., Liao, H., Unger, N., and Zanis, P.: Short-Lived Climate Forcers, edited by: Masson-Delmotte, V., Zhai, P., Pirani, A., Connors, S. L., Péan, C., Berger, S., Caud, N., Chen, Y., Goldfarb, L., Gomis, M. I., Huang, M., Leitzell, K., Lonnoy, E., Matthews, J. B. R., Maycock, T. K., Waterfield, T., Yelekçi, O., Yu, R., and Zhou, B., *Climate Change 2021: The Physical Science Basis. Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change*, Cambridge University Press, Cambridge, UK and New York, NY, USA, 817–922, <https://doi.org/10.1017/9781009157896.008>, 2021.
- UNFCCC: Report of the Conference of the Parties serving as the meeting of the Parties to the Paris Agreement on its third session, held in Glasgow from 31 October to 13 November 2021, Part two: Action taken by the Conference of the Parties serving as the meeting of the Parties to the Paris Agreement at its third session, *FCCC/PA/CMA/2021/10/Add.1*, 2022.
- van Vuuren, D., O'Neill, B., Tebaldi, C., Chini, L., Friedlingstein, P., Hasegawa, T., Riahi, K., Sanderson, B., Govindasamy, B., Bauer, N., Eyring, V., Fall, C., Frieler, K., Gidden, M., Gohar, L., Jones, A., King, A., Knutti, R., Kriegler, E., Lawrence, P., Lennard, C., Lowe, J., Mathison, C., Mehmood, S., Prado, L., Zhang, Q., Rose, S., Ruane, A., Schleussner, C.-F., Seferian, R., Sillmann, J., Smith, C., Sörensson, A., Panickal, S., Tachiiri, K., Vaughan, N., Vishwanathan, S., Yokohata, T., and Ziehn, T.: The Scenario Model Intercomparison Project for CMIP7 (ScenarioMIP-CMIP7), *EGUsphere* [preprint], <https://doi.org/10.5194/egusphere-2024-3765>, 2025.
- Vicedo-Cabrera, A. M., Scovronick, N., Sera, F., Royé, D., Schneider, R., Tobias, A., Astrom, C., Guo, Y., Honda, Y., Hondula, D. M., Abruzyky, R., Tong, S., Coelho, M. de S. Z. S., Saldiva, P. H. N., Lavigne, E., Correa, P. M., Ortega, N. V., Kan, H., Osorio, S., Kyselý, J., Urban, A., Orru, H., Indermitte, E., Jaakkola, J. J. K., Rytí, N., Pascal, M., Schneider, A., Katsouyanni, K., Samoli, E., Mayvaneh, F., Entezari, A., Goodman, P., Zeka, A., Michelozzi, P., de' Donato, F., Hashizume, M., Alahmad, B., Diaz, M. H., Valencia, C. D. L. C., Overcenco, A., Houthuijs, D., Ameling, C., Rao, S., Di Ruscio, F., Carrasco-Escobar, G., Seposo, X., Silva, S., Madureira, J., Holobaca, I. H., Fratianni, S., Acquafotta, F., Kim, H., Lee, W., Iniguez, C., Forsberg, B., Ragettli, M. S., Guo, Y. L. L., Chen, B. Y., Li, S., Armstrong, B., Aleman, A., Zanobetti, A., Schwartz, J., Dang, T. N., Dung, D. V., Gillett, N., Haines, A., Mengel, M., Huber, V., and Gasparrini, A.: The burden of heat-related mortality attributable to recent human-induced climate change, *Nat. Clim. Change*, 11, 492–500, <https://doi.org/10.1038/s41558-021-01058-x>, 2021.
- Wang, Z., Lin, L., Xu, Y., Che, H., Zhang, X., Zhang, H., Dong, W., Wang, C., Gui, K., and Xie, B.: Incorrect Asian aerosols affecting the attribution and projection of regional climate change in CMIP6 models, *npj Clim. Atmos. Sci.*, 4, 1–8, <https://doi.org/10.1038/s41612-020-00159-2>, 2021.
- Watson-Parris, D., Christensen, M. W., Laursen, A., Clewley, D., Gryspeerdt, E., and Stier, P.: Shipping regulations lead to large reduction in cloud perturbations, *P. Natl. Acad. Sci. USA*, 119, e2206885119, <https://doi.org/10.1073/pnas.2206885119>, 2022.
- Watson-Parris, D., Wilcox, L. J., Stjern, C. W., Allen, R. J., Persad, G., Bollasina, M. A., Ekman, A. M. L., Iles, C. E., Joshi, M., Lund, M. T., McCoy, D., Westervelt, D. M., Williams, A. I. L., and Samset, B. H.: Surface temperature effects of recent reductions in shipping SO<sub>2</sub> emissions are within internal variability, *Atmos. Chem. Phys.*, 25, 4443–4454, <https://doi.org/10.5194/acp-25-4443-2025>, 2025.

- Zhang, M., Gao, Y., Ting, M., Yu, Y., and Wang, G.: Land-use induced changes in extreme temperature predominantly influenced by downward longwave radiation, *Commun. Earth Environ.*, 5, 1–11, <https://doi.org/10.1038/s43247-024-01936-0>, 2024a.
- Zhang, W., Zhou, T., and Wu, P.: Anthropogenic amplification of precipitation variability over the past century, *Science*, 385, 427–432, <https://doi.org/10.1126/science.adp0212>, 2024b.
- Zhang, Y., Hajat, S., Zhao, L., Chen, H., Cheng, L., Ren, M., Gu, K., Ji, J. S., Liang, W., and Huang, C.: The burden of heatwave-related preterm births and associated human capital losses in China, *Nat. Commun.*, 13, 7565, <https://doi.org/10.1038/s41467-022-35008-8>, 2022.
- Zhuang, Y., Fu, R., Santer, B. D., Dickinson, R. E., and Hall, A.: Quantifying contributions of natural variability and anthropogenic forcings on increased fire weather risk over the western United States, *P. Natl. Acad. Sci. USA*, 118, e2111875118, <https://doi.org/10.1073/pnas.2111875118>, 2021.