



Constraining CMIP6 sea ice simulations with ICESat-2

Alek Petty¹, Christopher Cardinale¹, and Madison Smith²

¹Earth System Science Interdisciplinary Center (ESSIC), University of Maryland, College Park, MD, USA

²Woods Hole Oceanographic Institution (WHOI), Falmouth, MA, USA

Correspondence: Alek Petty (akpetty@umd.edu)

Received: 18 February 2025 – Discussion started: 14 March 2025

Revised: 13 June 2025 – Accepted: 18 June 2025 – Published: 25 September 2025

Abstract. This study evaluates sea ice simulations from the Coupled Model Intercomparison Project Phase 6 (CMIP6) using modern-era satellite measurements of sea ice area, total freeboard and thickness. Current global climate models exhibit substantial uncertainties in simulating sea ice, with significant contributions from both model uncertainty and internal variability. In our study, simulated Arctic and Southern Ocean total freeboard and Arctic winter sea ice thickness are assessed with data from NASA's Ice, Cloud and land Elevation Satellite-2 (ICESat-2) mission, providing an additional constraint beyond traditional passive microwave sea ice area comparisons used extensively in previous studies. Freeboard comparisons benefit from accurate observations from satellite laser altimetry but motivate increased focus on bulk sea ice density estimates across models and observations. The short observational time period also increases the role of internal variability. We undertake a plausibility assessment where we account for both observational uncertainty and internal variability across our different metrics for both hemispheres. In general, we see more plausible metrics in the Arctic compared to the Southern Ocean, with important differences when analyzing annual means vs. March and September means. We provide an example of this plausibility assessment by producing constrained estimates of 2015–2035 seasonal sea ice volume, using model subsets constrained using either area metrics or the combined area, freeboard and thickness metrics, with freeboard and thickness providing important additional impacts in terms of the mean seasonal cycle and spread. Finally, we present regional comparisons and a composite analysis, with models showing systematic underestimation of thicker ice in the Western Arctic and clear differences in the simulation of Eastern/Western Arctic sea ice. Overall, our study provides novel insights into sea ice model evaluation and emphasizes the potential benefits of integrat-

ing altimetry data from ICESat-2, as well as providing a discussion on the potential utility of these model constraints and future research priorities.

1 Introduction

Earth's polar sea ice cover is undergoing rapid declines in response to anthropogenic climate change (IPCC, 2023). Coupled global climate models (GCMs) are commonly used to simulate past and future sea ice conditions and disentangle the associated impacts and feedbacks from the rest of the climate system (Notz & SIMIP Community, 2020; Goosse et al., 2018; Jahn et al., 2024; Pithan and Mauritsen, 2014; Smith et al., 2019). Historical GCM outputs are also used to provide important constraints on sea ice mass, energy and freshwater budgets (Holland et al., 2006, 2010; Keen et al., 2021; Massonnet et al., 2018; Meredith et al., 2019; Zanowski et al., 2021) and can provide training input for seasonal/sub-seasonal forecasting models (Andersson et al., 2021). These efforts are often hindered by the large and poorly constrained uncertainties of the sea ice state in current GCMs.

Uncertainty in GCM sea ice conditions arise from the combined impact of model structural uncertainty and internal variability, with additional contributions of forcing uncertainty in future scenario runs. Model uncertainty is typically estimated based on the spread across available models. Model uncertainty has numerous causes, including biases in atmospheric/ocean forcing and errors in model physics (Massonnet et al., 2018). In GCMs, sea ice has historically been considered a simple boundary condition that increases the surface albedo and alters the surface energy balance, rather

than being a crucial climate component in and of itself. Many of the models included in Coupled Model Intercomparison Project 5 (CMIP5) feature only basic parameterizations of sea ice, with only a few models including significant improvements to the underlying sea ice physics schemes in the newly released CMIP6 suite (Notz & SIMIP Community, 2020, referred to as SIMIP2020). However, improvements in sea ice simulation in CMIP6 have been suggested, alluding to improvements in polar atmospheric/ocean forcing and/or model physics (Notz & SIMIP Community, 2020; Roach et al., 2020; Davy and Outten, 2020). Internal variability, which represents the random fluctuations of the climate system, can provide a significant and irreducible source of additional sea ice state uncertainty. The significant uncertainty across CMIP6 sea ice simulations poses important questions about its potential utility for predicting future sea ice conditions, e.g., the potential timing of an ice-free Arctic (Jahn et al., 2024).

There is no agreed upon approach for analyzing multi-model sea ice ensembles. In the simplest approach of full model democracy, all models are considered equally plausible, no exclusion or calibration is employed, and the model uncertainty remains unchanged. This is often the approach taken when observations are too unreliable to provide sufficient constraint or if internal variability estimates are unavailable. Beyond this, model runs can be excluded or weighted based on assessments of the combined observational uncertainty and internal variability. The exclusion approach was adopted in SIMIP2020, whereby models with historical Arctic sea ice area significantly outside a plausible range, calculated from a combination of observational uncertainty and internal variability, were omitted from the final CMIP6 future Arctic sea ice projection analysis. More sophisticated methods for excluding and/or weighting models based on comparisons with observations are available, including methods to recalibrate the models based on their simulated sea ice response to temperature variability (Bonan et al., 2021), atmospheric circulation (Topál and Ding, 2023) and greenhouse gas forcing (Kim et al., 2023). Similarly, emergent constraints, an approach that utilizes statistical relationships between observable quantities and future model diagnostics, have been used to constrain sea ice projections using a variety of metrics (Massonnet et al., 2018; Thackeray and Hall, 2019; Wang et al., 2021). These methods all generally rely on the assumption that model performance is consistent across time periods and depends crucially on the specific research question posed (Notz, 2015). However, recalibration approaches generally do not update associated state variables (e.g., impact on the surface energy budget or freshwater fluxes from the sea ice to the ocean) and can be more challenging to implement, motivating continued attention towards optimal exclusion or weighting approaches.

To date, most of the sea ice model exclusion and calibration efforts have utilized observational estimates of sea ice area or extent from the long-term (> 40 year) passive

microwave record (Lavergne et al., 2019; Parkinson, 2019). Passive microwave sensors measure the brightness temperature at different frequencies and polarities and use this information to infer the concentration of sea ice over the ocean surface, which can then be converted to sea ice areal coverage (by multiplying by the grid-cell area) and extent (grid cells with at least 15 % sea ice concentration). The multi-decadal time period benefits from being long enough to be representative of long-term climate conditions and reduces the role of internal variability. It can also be used to assess the sensitivity of sea ice area to warming or carbon dioxide forcing as an additional observational constraint. Sea ice area uncertainty is often estimated by comparing across different observational products (Notz & SIMIP Community, 2020) or by assuming fixed values/percentages based on community consensus (Massonnet et al., 2012). However, sea ice concentration provides only limited information within the more consolidated ice pack, where large gradients in thickness/volume manifest (Petty et al., 2020).

Dedicated polar-focused satellite altimetry missions have been launched since the early 2000s that can accurately profile sea ice height towards estimates of sea ice freeboard, thickness and, thus, volume. These include the National Aeronautics and Space Administration (NASA) Ice, Cloud and land Elevation Satellite (ICESat) mission (2003–2009, Zwally et al., 2002; Kwok and Cunningham, 2008), the European Space Agency (ESA) CryoSat-2 radar altimetry mission (2010 onwards, Laxon et al., 2013), and, most recently, NASA's ICESat-2 laser altimetry mission (2018 onwards, Neumann et al., 2019; Petty et al., 2020). The altimeters measure the height of sea ice and open water leads between ice floes. The open water height estimates are used to generate an estimate of the local sea surface height. Subtracting the local sea surface height from the sea ice height provides an estimate of freeboard, the extension of sea ice above sea level; see the basic schematic in Fig. 1. Laser altimeters (e.g., NASA's ICESat and ICESat-2) provide estimates of the snow-covered ice surface height (a metric commonly referred to as total freeboard). In contrast, data from radar altimeters (e.g., ESA's CryoSat-2) are typically used to provide an estimate of the less distinct ice–snow interface height and thus an estimate of ice freeboard. The effective radar penetration depth at the Ku band used in CryoSat-2 is generally considered to come from the ice–snow interface, although studies continue to challenge this (Nandan et al., 2017; Willatt et al., 2011). The satellite laser altimeters benefit from higher spatial resolution on the ground than radar (footprints of tens of meters as opposed to hundreds of meters to kilometers). They also profile the upper snow surface, thus providing a useful constraint on total snow loading. However, radar is unaffected by clouds, and CryoSat-2 benefits from continuous data collection and refinement since it launched in 2010. Additional input assumptions regarding snow depth, snow density and ice density, together with an assumption of isostasy, are then typically used to convert measured free-




Figure 1. Schematic showing (a) observations of sea ice total freeboard from laser altimetry towards estimates of sea ice thickness, (b) advanced physics and ice thickness distribution typical of state-of-the-art sea ice models, and (c) fixed ice thickness/snow parameterization used in more basic sea ice models. Note that ice freeboard is the extension of sea ice above sea level, while total freeboard is the extension of the ice and snow layer above freeboard. Sea ice models do not typically simulate freeboard, but this can be estimated based on an assumption of isostasy.

board (ice or total depending on the sensor) into an estimate of sea ice thickness for the Arctic, which introduces significant additional uncertainty to this “observational” estimate (Giles et al., 2007; Kwok and Cunningham, 2008; Petty et al., 2020). Constraining the thickness uncertainty remains challenging due, in part, to the lack of ground-truth data available for validation. For Antarctic sea ice, limited knowledge of the more complex snow loading has generally hindered production of similar snow and sea ice thickness data production efforts to date, although novel approaches show promise (Fons et al., 2021; Garnier et al., 2021).

Due to concerns around accuracy and uncertainty quantification, combined with their more limited temporal coverage, sea ice thickness data have generally been excluded from model assessment efforts to date (Notz & SIMIP Community, 2020; Kay et al., 2022; Roach et al., 2020), despite the fact that the mean thickness has been demonstrated to be the crucial factor controlling sea ice variability and trends (Massonnet et al., 2018). With the recent ICESat-2 period now extending into its seventh year of successful data collection (at the time of writing) and the improved understanding achieved from the joint operation of both ICESat-2 and CryoSat-2, we can begin to reconsider the concerns around accuracy and time period. In addition, comparisons of the direct observations of total freeboard with model estimates of this same quantity may offer another path forward for model assessments, better leveraging the high accuracy of the ICESat-2 laser altimeter observations. Assessments of total freeboard avoid a significant component of the thickness uncertainty that is introduced in the conversion of freeboard to thickness. Laser altimetry total freeboard estimates also avoid the uncertainties associated with radar freeboard profiling (e.g., identifying which interface is dominating the radar return). Total freeboard comparisons are expected to provide significant value in the Southern Ocean, where snow depth and sea ice thickness estimates are less reliable. Southern

Ocean sea ice is also thought to be composed of a higher fraction of thinner first-year ice with higher snow depths compared to the Arctic, such that total freeboard could be considered a better proxy for thickness than in the Arctic (Kurtz and Markus, 2012; Worby et al., 2008). In addition, total freeboard is utilized in some sea ice model parameterizations, e.g., atmospheric form drag (Tsamados et al., 2014) and snow-ice formation (Hunke et al., 2015), providing an additional motivation to assess its representation in models, especially as we prepare for the upcoming release of the CMIP7 output. However, total freeboard integrates information from both ice thickness and snow depth variability concurrently, meaning changes in freeboard can be linked to changes in the underlying ice thickness and/or snow depth, and diagnosing the cause of total freeboard biases is more challenging than the more direct prognostic variables of area and thickness. Additionally, freeboard is not a prognostic variable in sea ice models, and so it is typically calculated within the model as needed assuming hydrostatic equilibrium from the associated ice state variables (see schematic in Fig. 1 and Eq. 2 in Sect. 2.1 below), either in a post-processing step or within the relevant model parameterization scheme.

Following a request from the Sea Ice Model Intercomparison Project (SIMIP) in the lead-up to CMIP6 (Notz et al., 2016), several modeling groups provided direct outputs of ice freeboard, making a comparison effort timely. To our knowledge, we are unaware of any studies that have explored this new model output. In this paper, we thus undertake a first attempt at using satellite altimetry observations of total freeboard from NASA’s ICESat-2 mission to evaluate sea ice output from CMIP6. We compare these with evaluations using the traditional sea ice area and arguably more uncertain (winter Arctic) sea ice thickness metrics derived from satellite observations in order to demonstrate the advantages and disadvantages of these comparisons and look ahead to using this information to better constrain simulations and projections of

sea ice across both poles. Producing a longer-term record of sea ice area, freeboard and thickness through assimilation of sea ice altimetry data into a consistent reanalysis is a current focus of the sea ice community, although challenges unique to sea ice have only recently been assessed (Riedel and Anderson, 2024; Wieringa et al., 2024; Williams et al., 2023). We thus focus only on direct ICESat-2 observations in this study while recognizing the challenges in characterizing internal variability from this short time period.

2 Data and methods

2.1 CMIP6 sea ice model output

We use sea ice model output from the CMIP6 archive (Eyring et al., 2016). CMIP6 data are officially hosted through the Earth System Grid Federation (ESGF), enabling users to directly download all relevant CMIP6 output made available by all contributing model centers using the Open-source Project for a Network Data Access Protocol (OPeN-DAP) system. We additionally make use of the Pangeo Analysis-Ready Cloud-Optimized CMIP6 catalog (<https://pangeo-data.github.io/pangeo-cmip6-cloud/>, last access: 12 September 2025), which is hosted on both the Amazon Web Services (AWS) and Google Cloud Project (GCP) cloud storage services, simplifying the data ingest process (where we use the NASA-funded AWS-hosted CryoCloud platform, <https://book.cryointhecloud.com> (last access: 12 September 2025) and thus the AWS-hosted catalog). Some model outputs are missing from the Pangeo cloud catalogs, so we utilize the ESGF/OPeNDAP data where needed to ensure full model selection. We primarily use data from the SSP2-4.5 future scenario (2015–2100) to ensure full overlap with our ICESat-2 observational period (2018 to 2024). The Code Availability section provides links to all code used to retrieve and analyze these data. We use the monthly mean output across all models except for AWI-CM-1-1-MR, which provides only daily data that we average to a monthly timescale).

As our study builds on the initial SIMIP CMIP6 sea ice evaluation studies, which included only models available in the initial IPCC AR6 analysis time frame (Notz & SIMIP Community, 2020; Roach et al., 2020), we briefly explored differences in Arctic sea ice area from the model runs used in our study compared to the SIMIP2020 study. Overall, our study uses 13 more models compared to the SIMIP2020 study. Despite the significant difference in model subsets, differences in the multi-model CMIP6 mean Arctic sea ice area over the 2015–2035 time period in both March and September were negligible (see Fig. S1 in the Supplement).

The grid-cell mean sea ice variables used in our study include the following (CMIP6 variable names in parentheses): sea ice area (siconc), sea ice thickness (sithick), sea ice freeboard (sifb), snow thickness (sisnthick), sea ice mass

Table 1. Model variables used in our study, its denoted symbol if used in the derivations below, the official CMIP6 variable name and the units. CMIP6 variables denoted NA are not provided directly and are either prescribed or calculated in this study.

Variable name	Symbol	CMIP6 variable	Units
Sea ice area	NA	siconc	m
Sea ice thickness	h_i	sithick	m
Sea ice freeboard	h_{fi}	sifb	m
Total freeboard	h_{ft}	NA	m
Snow thickness	h_s	sisnthick	m
Sea ice mass per unit area	M_i	simass	kg
Sea ice volume per unit area	V_i	sivol	m
Bulk sea ice density	ρ_i	NA	kg m ^{−3}
Bulk snow density	ρ_s	NA	kg m ^{−3}
Seawater density	ρ_ω	NA	kg m ^{−3}

(simass) and sea ice volume (sivol), as summarized in Table 1. Note that the grid-cell mean ice freeboard variable was requested from CMIP6 contributing centers by the Sea-Ice Model Intercomparison Project (SIMIP) consortium at priority level 2 (Notz et al., 2016) and that only 17 of the modeling centers provided this output for the SSP2-4.5 scenario runs. A listing of the CMIP6 models and the relevant variable availability is shown in Table 2. For the models that do not provide ice freeboard output, we can instead estimate this from the provided variables of ice and snow thickness, estimates of ice and snow density, and an assumption of isostasy. This method can also be used to check consistency with the provided ice freeboard output from the models that do provide that output. Starting with the hydrostatic equilibrium equation for ice thickness:

$$h_i = \frac{h_{fi}\rho_\omega + h_s\rho_s}{(\rho_\omega - \rho_i)}, \tag{1}$$

where h_{fi} is sea ice freeboard, h_s is snow thickness, ρ_ω is seawater density (1024 kg m^{−3}), ρ_s is the bulk snow density and ρ_i is the bulk ice density. We can rearrange Eq. (1) to calculate ice freeboard as:

$$h_{fi} = \frac{h_i(\rho_\omega - \rho_i) - h_s\rho_s}{\rho_\omega}. \tag{2}$$

This ice freeboard can be converted into an estimate of total freeboard by simply adding snow thickness (sisnthick) as:

$$h_{ft} = h_{fi} + h_s. \tag{3}$$

Snow thickness was listed as a priority level 1 variable in Notz et al. (2016) and is provided by the 36 models that provide either the grid-cell sea ice thickness or sea ice volume and area. For the models that do provide outputs of ice freeboard, we generally assume that this is calculated in post-processing as in Eq. (2). It is worth noting that differences

Table 2. CMIP6 sea ice model summary and data variable availability. Models are added only if they provide a relevant SSP2-4.5 output. Variables are listed only if used in this study. All data output is monthly and available on the native model grid. There are several exceptions: siconc is daily for AWI-CM-1-1-MR, and data are available only on gr1grids (regridded) for INM-CM4-8, INM-CM5-0 and KIOST-ESM. The first 17 rows indicate the freeboard output subset. Note that in most cases, the underlying sea ice model has been adapted to ensure consistency with other model components. Semtner–Hibler refers to the Semtner zero-layer thermodynamics model (Semtner, 1976) and the Hibler ice dynamics model (Hibler, 1979). Sea ice variables are all grid-cell monthly means and are summarized in Table 1. The final column describes whether any of the model output was used in the Notz & SIMIP Community (2020) study (S), the Roach et al. (2020) study (R), both (B) or neither (N). Sea ice model information taken from https://wcrp-cmip.github.io/CMIP6_CVs/docs/CMIP6_source_id.html (last access: 12 September 2025).

CMIP6 model ID	Sea ice model	Variables	S/R/B/N
ACCESS-CM2	CICE5.1.2	siconc, sithick, sifb, sisnthick, simass, sivol	B
CESM2	CICE5.1.2	siconc, sithick, sifb, sisnthick, simass, sivol	B
CESM2-WACCM	CICE5.1.2	siconc, sithick, sifb, sisnthick, simass, sivol	B
CIESM	CICE4	siconc, sithick, sifb, sisnthick, simass, sivol	N
CMCC-CM2-SR5	CICE4.0	siconc, sithick, sifb, sisnthick, simass, sivol	N
CMCC-ESM2	CICE4.0	siconc, sithick, sifb, sisnthick, simass, sivol	N
CNRM-CM6-1	GELATO6.1	siconc, sithick, sifb, sisnthick, simass, sivol	B
CNRM-CM6-1-HR	GELATO6.1	siconc, sithick, sifb, sisnthick, sivol	B
CNRM-ESM2-1	GELATO6.1	siconc, sithick, sifb, sisnthick, simass, sivol	B
HadGEM3-GC31-LL	CICE5.1.2	siconc, sithick, sifb, sisnthick, simass, sivol	B
IPSL-CM6A-LR	LIM3	siconc, sithick, sifb, sisnthick, simass, sivol	B
MPI-ESM1-2-HR	Semtner-Hibler	siconc, sithick, sifb, sisnthick, simass, sivol	B
MPI-ESM1-2-LR	Semtner-Hibler	siconc, sithick, sifb, sisnthick, simass, sivol	B
MRI-ESM2.0	MRI.COM4.4	siconc, sithick, sifb, sisnthick, simass, sivol	B
NorESM2-LM	CICE5.1.2	siconc, sithick, sifb, sisnthick, simass, sivol	S
NorESM2-MM	CICE5.1.2	siconc, sithick, sifb, sisnthick, simass, sivol	N
UKESM1.0-LL	CICE5.1.2	siconc, sithick, sifb, sisnthick, simass, sivol	B
ACCESS-ESM1-5	CICE4.1	siconc, sithick, sisnthick, sivol	B
AWI-CM-1-1-MR	FESOM1.4	siconc, sithick, sisnthick, sivol	B
BCC-CSM2-MR	SIS2	siconc, sithick, sisnthick, sivol	B
CAMS-CSM1-0	SIS1.0	siconc, sisnthick, sivol	B
CanESM5	LIM2	siconc, sithick, sisnthick	B
CanESM5-1	LIM2	siconc, sithick, sisnthick	N
EC-Earth3	LIM3	siconc, sithick, sisnthick, sivol	B
EC-Earth3-CC	LIM3	siconc, sithick, sisnthick, sivol	N
EC-Earth3-HR	LIM3	siconc, sithick, sivol	N
EC-Earth3-Veg	LIM3	siconc, sithick, sisnthick, sivol	B
EC-Earth3-Veg-LR	LIM3	siconc, sithick, sisnthick	N
FGOALS-f3-L	CICE4.0	siconc, sisnthick, sivol	B
FIO-ESM-2-0	CICE4.0	siconc, sisnthick, sivol	B
GFDL-CM4	GFDL-SIM4p25	siconc, sithick, sisnthick, sivol	B
GFDL-ESM4	GFDL-SIM4p5	siconc, sithick, sisnthick, sivol	B
KIOST-ESM	GFDL-SIS	siconc, sithick, sisnthick	N
MIROC6	COCE4.9	siconc, sithick, sisnthick, sivol	B
MIROC-ES2H	COCO4.9	siconc, sithick, sisnthick	N
MIROC-ES2L	COCO4.9	siconc, sithick, sisnthick	B
NESM3	CICE4.1	siconc, sithick, sisnthick	B
TaiESM1	CICE4	siconc, sithick, sisnthick, sivol	N
CanESM5-CanOE	LIM2	siconc	N
FGOALS-g3	CICE4.0	siconc	N
INM-CM4-8	INM-ICE1	siconc	B
INM-CM5-0	INM-ICE1	siconc	B

could arise both from the calculation of freeboard at sub-monthly time steps before averaging to monthly and from using the categories of ice thickness across the ice thickness distribution (ITD) before averaging across the grid cell (for the models that simulate an ITD).

A crucial additional variable in the conversion between freeboard and thickness is the bulk ice density (ρ_i): the higher the ice and snow density, the lower the freeboard. Neither the bulk ice density nor snow density is provided directly by any of the CMIP6 groups, as generally it is not considered a prognostic variable and instead a prescribed constant. However, for some of the more sophisticated sea ice models, the effective bulk ice density can be considered a function of the variable internal temperature and salinity, which vary based on the internal sea ice physics scheme and need to be calculated during the ice freeboard calculation (e.g., CESM2, D Bailey, personal communication). To our knowledge, all CMIP6 sea ice models currently use a constant snow density of 330 kg m^{-3} (this was also assumed in the CMIP6 sea ice freshwater analysis in Zanowski et al. (2021)). Prescribed (or variable) bulk sea ice densities across CMIP6 were harder to determine from the available documentation. Alternatively, there are two ways in which we can infer the bulk ice density. The first is to infer bulk ice density from provided outputs of total (grid-cell mean) sea ice mass (M_i) and volume (V_i) from the 24 models that provide these outputs (listed in Table 2) as:

$$\rho_i = M_i / V_i. \quad (4)$$

We can also indirectly infer bulk ice density from the 17 models (listed in Table 2) that provide outputs of ice freeboard, ice thickness and snow thickness, together with an estimate of snow density and seawater density, through rearranging Eq. (2) in terms of bulk ice density as:

$$\rho_i = \rho_w - \frac{\rho_w h_{fi} + \rho_s h_s}{h_i}. \quad (5)$$

We set the snow density to 330 kg m^{-3} and the seawater density to 1024 kg m^{-3} based on our review of the default options used across CMIP6 sea ice models.

Model regridding

Model output was regridded to simplify analysis and enable spatial comparisons between the model and observations. We regrid all CMIP6 model data to rectilinear grids depending on the variable and observational comparison. For area, we regrid all data to the EASE 2.0 $25 \text{ km} \times 25 \text{ km}$ grid used by the concentration products described below, while for freeboard and thickness, we use the North Polar Stereographic $25 \text{ km} \times 25 \text{ km}$ grid used by the ATL20/IS2SITMOGR4 datasets, which are described in the following sections. We explored various options to optimize our regridding approach, utilizing the open-source

Python xESMF package (<https://xesmf.readthedocs.io/en/latest/index.html>, last access: 12 September 2025, Zhuang et al., 2024). We primarily utilized the conservative normed regridding method that preserves areal contributions of the input data within each observational grid cell. To prevent unrealistic data interpolation along the coastline, land masks are specified for both the source and destination grids. For native model grids, land masks are calculated using the provided variable sea area percentage (sftof) which, when divided by 100, gives the fraction of the grid cell covered by ocean and used in the conservative normed regridding described above. For the North Polar Stereographic $25 \text{ km} \times 25 \text{ km}$ grid, the NSIDC land mask is used (Meier and Stewart, 2023). For the EASE 2.0 $25 \text{ km} \times 25 \text{ km}$ grid, land is defined as NaN regions in the sea ice concentration data (this method is also used for models with no sftof data). We note that regridding can introduce artificial errors, but we ensured appropriate methods were used to minimize this, and our investigations suggest negligible differences at both the grid-cell and basin scales (see Supplemental Information, Figs. S2 and S3).

2.2 Observational sea ice data

2.2.1 Total freeboard and winter Arctic sea ice thickness from ICESat-2 altimetry

We use monthly gridded total freeboard data from NASA's ICESat-2 ATL20 product (Version 4) disseminated through the National Snow and Ice Data Center (NSIDC) (<https://nsidc.org/data/atl20>, last access: 12 September 2025, Petty et al., 2023b). ATL20 is produced using a simple binning of the along-track freeboard data from the three strong beams of ICESat-2/ATLAS (ATL10, Kwok et al., 2023) on a $25 \text{ km} \times 25 \text{ km}$ North Polar Stereographic grid. Significant data gaps can still exist for various reasons (e.g., cloud attenuation, lack of open water leads, spacecraft issues). Note that the underlying along-track ATL10 freeboard product masks data where the sea ice concentration is less than 50 % and where data are within 25 km of the nearest coastline. ATL20 data are available for both hemispheres across all months (November 2018 onwards). While data are available across the summertime Arctic, these data should be treated with more caution due to the lack of a reliable melt pond classification scheme in the underlying sea ice processing (Tilling et al., 2020, Kwok et al., 2022). No uncertainty term is included in the products that account for the potential misclassification of leads as melt ponds, but the additional height filter in ATL10 (where only leads that pass a strict relative height filter are used to derive the local sea surface height) is expected to mitigate this issue to some degree.

In addition, we use monthly gridded winter Arctic sea ice thickness estimates from ICESat-2 (IS2SITMOGR4, Version 3), also disseminated through the NSIDC (<https://nsidc.org/data/is2sitmogr4>, last access: 12 September 2025, Petty

et al., 2023c). These data use snow loading estimates from the NASA Eulerian Snow On Sea Ice Model (NESOSIM, Petty et al., 2018), now at Version 1.1, a constant bulk ice density of 916 kg m^{-3} and the isostasy assumption to derive estimates of sea ice thickness across the Arctic Ocean between September and April since November 2018 (Petty et al., 2023c). The upgrade of IS2SITMOGR4 from Version 1 to 2 was shown to increase correspondence with ice thickness estimates derived using a similar snow loading approach with CryoSat-2 ice freeboard data (Petty et al., 2023c) and showed good agreement with a product derived directly from ICESat-2 and CryoSat-2 freeboards (Kacimi and Kwok, 2022). Due to the lack of NESOSIM snow loading data in summer, these thickness estimates are currently available only between September and April. To increase data coverage, including the 88°N Arctic pole hole, and to increase consistency across the observational comparisons, we use the linear interpolation/Gaussian smoothing variables from IS2SITMOGR4 as described in Petty et al. (2023c) for the winter Arctic sea ice freeboard/thickness data and apply this method additionally to the Southern Ocean total ATL20 freeboard data. Differences between the raw vs. interpolated/smoothed annual mean ATL20 total freeboards are provided in the Supplementary Information (Fig. S4). We analyze annual means from all months of the year, particularly the months of September and March to capture the peaks and troughs of the seasonal cycle across both hemispheres. For the Arctic thickness data, the annual mean consists only of the January–April and September–December “winter” months, while for total freeboard, we use all months of the year. These data are shown in Figs. 2 and 3.

2.2.2 Bulk ice density estimates

The above ICESat-2 thickness retrievals follow the approach of several other studies in assuming a fixed bulk ice density, in this case 916 kg m^{-3} – the density of pure ice. In reality, sea ice is a complex mixture of pure ice and brine, which increase bulk ice density, but also air pockets that lower bulk density, with their relative contributions varying with the evolving ice state. Various other sea ice remote sensing studies have thus utilized a lower density for multi-year ice (882 kg m^{-3}) based on the analysis of airborne Sever expedition in situ data prior to the 1990s by Alexandrov et al. (2010). However, these bulk ice densities have been challenged in recent studies, using values inferred from multi-sensor airborne profiles (J22, Jutila et al., 2022) and multi-sensor satellite methods (Shi et al., 2023) and from in situ data collected during the Multidisciplinary drifting Observatory for the Study of Arctic Climate (MOSAIC) campaign (Salganik et al., 2025; Zhou et al., 2024), all of which generally show higher densities, linked to the younger ice state and issues with previous ice density measurement approaches.

The IS2SITMOGR4 v3 dataset also includes bulk ice density estimates calculated using the J22 empirical bulk ice

density parameterization, an exponential function of the local ice freeboard derived from coincident laser scanning, snow radar and electromagnetic induction sounding data (calculated using the total freeboard minus snow depth in the ICESat-2 processing). We use these to provide an alternative, and seasonally variable, bulk ice density estimate to compare with the model results and provide added context to both the model and remote-sensing-based estimates. These J22 densities are expected to be higher than the pure ice density approximation of 916 kg m^{-3} , especially for first-year ice regimes. The J22 parameterization has not been validated across different regions and seasons of the Arctic, so we consider these highly experimental and use them here with caution.

2.2.3 Sea ice area from satellite passive microwave

We use sea ice concentration estimates from the European Organisation for the Exploitation of Meteorological Satellites (EUMETSAT) Ocean and Sea Ice Satellite Application Facility (OSI SAF), specifically OSI-450-a, which is the third major version of the OSI SAF Global Sea Ice Concentration Climate Data Record (OSI SAF, 2022a), and OSI-430-a, which is an operational extension of this product with a latency of 16 days, currently for the period 2021 onwards (OSI SAF, 2022b). We use the monthly mean concentration estimates from both datasets across the period November 2018 to April 2024 (the ICESat-2 study period). The data are posted on a $25 \text{ km} \times 25 \text{ km}$ Equal-Area Scalable Earth (EASE) 2.0 grid, meaning all grid cells have a fixed area of 625 km^2 , which we multiply by the grid-cell concentrations to derive sea ice area, before averaging across basins. As in the total freeboard/thickness data, we take annual means (all months of the year) and September and March means across the 2018 to 2024 period.

2.2.4 Observational uncertainties

An important consideration when using observations to evaluate climate models is the observational uncertainty. However, the characterization of uncertainties within sea ice remote sensing products generally focuses on grid-scale uncertainties, estimated primarily using theoretical assessments (e.g., propagation of uncertainties, Giles et al., 2007) or comparisons with the ground truth/imagery (Kern et al., 2022). These uncertainties are generally considered random/uncorrelated at the typical grid scales they are disseminated at ($\sim 10\text{--}100 \text{ km}$) and thus theoretically reduce to zero when averaging at basin scales. Product or algorithm differences that drive regional-scale ($\sim 100\text{--}10\,000 \text{ km}$) systematic uncertainties are rarely accounted for. As such, the approach often used in sea ice climate model diagnostics to estimate observational uncertainty is to calculate differences in hemispheric mean sea ice area/extent across available products or algorithms (Notz & SIMIP Community, 2020; Roach et




Figure 2. (a–c) Total freeboard from ICESat-2 ATL20 v4, and (g–i) sea ice concentration from OSI SAF (bottom) for annual, March and September means in the 2018 to 2024 ICESat-2 period. The sea ice thickness annual mean includes only data between September and April due to data availability. Hatching in the top two rows indicates grid cells not included in the “perennial ice” mask, as data are missing from at least one year in the 2018 to 2024 period. Freeboard and thickness data are the interpolated/smoothed variables.




Figure 3. Same as in Fig. 2 but for the (a–c) Southern Ocean total freeboard and (d–f) sea ice concentration.

al., 2020). For sea ice concentration, multiple well established sea ice concentration/area products exist (e.g., Bootstrap: Comiso et al., 1997, NASA Team: Cavalieri et al., 1996, and OSI SAF: Lavergne et al., 2019), which enables such an approach, although this still has its limitations due to the limited (typically three-product) sample size and the fact that NASA Team data have a well reported low concentration bias, especially in summer (Kern et al., 2019, 2022). Recent efforts in other domains have explored the creation of observational ensembles to better sample the full product spread (Lenssen et al., 2024); however, to our knowledge, no such effort has been undertaken for any of the sea ice metrics used here.

For this study, we instead estimate the basin-scale uncertainty through an evaluation of published values in SIMIP2020 and Roach et al. (2020). We use both a “high” and “low” uncertainty estimate to explore the impact of the observational uncertainty estimate on our model assessments considering the challenge of prescribing observational uncertainty from these limited product “ensembles”. For the passive microwave area, we use the assumption that a 0.5 million km² basin-mean sea ice area uncertainty represents a best-case “low” uncertainty, while 1 million km² represents a less optimistic “high” uncertainty. Uncertainty quantification is more challenging for the ICESat-2 total freeboards due to the lack of alternative freeboard products available to assess the product spread. We instead provide “high” and “low” uncertainty estimates based on a review of the relevant literature. The primary validation of ICESat-2 sea ice height and freeboard with coincident airborne altimetry data from NASA’s Operation IceBridge shows very high accuracies in the sea ice heights and total freeboard errors (10 km along-track means) of less than a few centimeters depending on the methodology used, generally indicative that ICESat-2 was likely to satisfy the mission objectives of < 3 cm freeboard uncertainty at those scales (Kwok et al., 2019). No summer Arctic or Antarctic validation analyses have been completed to date. The summer Arctic is expected to pose more challenges due to the presence of melt ponds but benefits from more openings in the ice cover, reducing the need to interpolate sea surface heights over large distances. In addition, an analysis of monthly mean sea surface height differences between coincident ICESat-2 and CryoSat-2 showed overall mean differences of less than 1 cm (Bagnardi et al., 2021). For our analysis, we proceed with the assumption that a 1.5 cm basin-mean total freeboard uncertainty represents a best-case “low” uncertainty, while 3 cm represents a less optimistic “high” uncertainty.

Winter ICESat-2 Arctic sea ice thickness uncertainty quantification is a less direct measurement than using the total freeboard, so the relative uncertainty increases significantly through the introduction of additional input assumptions related to snow loading and bulk ice density. Inter-comparisons of ICESat-2 and CryoSat-2 winter Arctic sea ice thickness have been undertaken, showing mean differ-

Table 3. Hemispheric monthly mean observational uncertainty estimates applied in our study.

Variable	Low uncertainty estimate	High uncertainty estimate
Sea ice area (million km ²)	0.5	1.0
Total freeboard (cm)	1.5	3.0
Winter Arctic sea ice thickness (cm)	15	30

ences in the mean monthly winter Arctic ice thickness of ~ 10–30 cm between ICESat-2 and the various CryoSat-2 thickness estimates, similar to the comparisons with independent ice draft estimates obtained from upward-looking sonar in the Beaufort Sea (Petty et al., 2023c). Comparisons of monthly winter Arctic mean thickness estimates between ICESat-2 and the Alfred Wegener Institute (AWI) CryoSat-2/SMOS product have also been presented in recent NOAA Arctic report cards (Meier et al., 2023, 2024), showing similar basin-scale monthly mean differences. For our analysis, we proceed with the assumption that a 15 cm basin-mean winter Arctic sea ice thickness uncertainty represents a best-case “low” uncertainty, while 30 cm represents a less optimistic “high” uncertainty estimate. These uncertainty choices are summarized in Table 3.

2.2.5 Ancillary data

We use an Arctic Ocean region mask (Meier and Stewart, 2023) to analyze ICESat-2 data within an Inner Arctic Ocean domain that includes the Central Arctic, Beaufort Sea, Chukchi Sea, East Siberian Sea, Laptev Sea and Kara Sea, as in Petty et al. (2023c, Fig. 5). Focusing on the Inner Arctic avoids challenges of interpretation in the more marginal seas of the Arctic and mitigates issues with more uncertain marginal ice representation in the thickness observations, especially related to snow loading. We refer to these results as “Arctic Ocean” throughout. We do not apply any regional masking to the Southern Ocean analysis. In addition, region masking is applied only to the ICESat-2 total freeboard and thickness data, as the concentration data are considered more reliable in the marginal zones. We discuss the impact of this regional masking in the discussion.

2.3 Methods

2.3.1 Perennial ice masking

Our initial evaluations highlighted the challenge of model-observation assessments related to contrasts in total freeboard and thickness data coverage, especially as the ICESat-2 observations employ various filters, e.g., the 50 % concentration and 25 km coastal filter to improve data quality, and data drop-out due to environmental factors such as

clouds. Spurious model performance was also observed in the more marginal seas and in the comparisons of models with vastly underestimated ice cover in certain seasons. To enhance confidence in our ICESat-2 comparisons, we thus employ a “perennial coverage” masking to both the freeboard and thickness observations from ICESat-2 as follows:

- For the gridded freeboard/thickness observational data, calculate the fractional grid-cell coverage over time for each month across the 2018 to 2024 time period. Arctic data for the area within the Inner Arctic Ocean region described above are used.
- Flag grid cells as “perennial” if they include data every year across our study period.
- Set all monthly model grid-cell freeboards and thicknesses to zero (instead of NaN) across valid regridded ocean/sea ice grid cells.
- For every month, mask all model grid cells (set to NaN) outside the perennial mask for the relevant metric.
- Calculate grid cells of the monthly, hemisphere mean quantities for the observations and the models.
- Compute annual means by averaging the monthly perennial means for each year.
- For the spatial comparisons, compare only data across the “perennial” grid cells.

The impact of this perennial masking is that our annual freeboard/thickness results are likely to be skewed high, as the observations do not include regions of low-concentration (< 50 %) ice. Clouds/data gaps are generally more likely in the more marginal seas also. The freeboard/thickness analysis is thus more of an assessment of ice conditions where we have consistent ICESat-2 data coverage. The assessment of simulated sea ice coverage is addressed more comprehensively with the passive microwave sea ice concentration/area data, which provide full coverage data across both hemispheres.

2.3.2 Internal variability and plausibility estimates

A climate model is not expected to exactly match the time period of our own reality due to the impact of internal variability. To characterize internal variability, we calculate, for each model, the standard deviation of the given metric of interest (e.g., total freeboard) averaged over the given region and the time window of interest (e.g., annual mean or a given month for the 2018 to 2024 ICESat-2 period) across all available ensemble members. As in SIMIP2020, we apply the Bessel correction to estimate an unbiased population standard deviation from a sample, accounting for the variable ensemble size across CMIP6. To explore the sensitivity to ensemble size and our chosen time period, especially for models with

low (or no) ensemble size, we also calculate 6-year running means within a larger time window (starting years of 2015 through 2024, ensuring we analyze only the SSP2-4.5 runs where the ensembles are consistent). We choose this window size as a balance between the benefits of increased sampling and the cost of increasing the likelihood of trend contamination. We additionally calculate a CMIP6 mean internal variability for each given metric by taking the mean internal variability across all models for those with at least five members. We follow SIMIP2020 and define a plausible range as:

$$P = \pm 2\sqrt{(\sigma_{\text{int}}^2 + \sigma_{\text{obs}}^2)}, \quad (6)$$

where σ_{int} is the internal variability and σ_{obs} is the observational uncertainty. The factor 2 effectively provides a 95 % plausibility window based on the observational uncertainty and internal variability. In addition, following earlier CMIP analyses (Olonscheck and Notz, 2017; Santer et al., 2008), we calculate a model plausibility index as:

$$\phi = (\mu_{\text{mod}} + \mu_{\text{obs}}) / \sqrt{(\sigma_{\text{int}}^2 + \sigma_{\text{obs}}^2)}, \quad (7)$$

where μ_{mod} and μ_{obs} are the mean model and observational quantities of interest. A plausibility index of 0 indicates perfect agreement between the model and observation, with higher values in either direction, based on the direction of the model bias, indicating worse agreement and lower plausibility. We use this plausibility index to compare plausibility across metrics.

2.3.3 Spatial assessments

For our freeboard and thickness spatial comparisons, we calculate the mean absolute error (MAE) by comparing the perennial grid-cell means between the observations and models. For area, we choose the sum absolute error (SAE) to express the results in units more consistent with the total area metric. We then additionally estimate spatial internal variability, following the same approach as above, but at the grid-cell level, following the approach of Schaller et al. (2011) for regional precipitation assessments. The regional estimate of internal variability is used to provide context to our regional bias assessments. However, the ability of models to accurately represent the regional manifestation of sea ice internal variability is unclear at best, so we provide this analysis with caution and do not use the spatial comparisons to assess model plausibility explicitly. We use only the perennial masked data in this analysis to increase confidence in our comparisons. Due to the short time period and increasing/uncertain role of internal variability in the spatial assessments, we do not assess pattern correlation, as in other regional assessments (Watts et al., 2021). We instead provide a composite regional bias analysis to provide further insights into typical spatial sea ice bias patterns simulated by CMIP6 models.

3 Results

3.1 Bulk ice density and total freeboard analysis

We first assess bulk ice density and total freeboard across the 17-model subset that provides direct outputs of ice freeboard. As discussed in Sect. 2.1, a key variable in the conversion between ice thickness and freeboard is the bulk ice density, which we estimate using the two methods described in Sect. 2.1.: Method 1 using the provided variables of freeboard, snow depth and ice thickness and a prescribed snow density of 330 kg m^{-3} , and Method 2 using the sea ice mass and volume outputs. Our initial analysis produced notably high bulk ice densities close to that of pure seawater in the ACCESS-CM2 model (see Fig. S5), especially in the Arctic (Method 1/Method 2: $1005/940 \text{ kg m}^{-3}$ for the Arctic and $955/975 \text{ kg m}^{-3}$ for the Southern Ocean, both with significant seasonal variability). We were unable to ascertain the cause of these anomalously high densities and thus dropped this model from the rest of our freeboard subset analysis. We suspect an underlying error in how freeboard was calculated in post-processing, as other key metrics did not display such anomalous behavior.

In Fig. 4, we show estimates of bulk ice density (mean and seasonal variability) for the reduced 16-model subset across both the Arctic Ocean and Southern Ocean over our ICESat-2 study period of 2018–2024. The Method 2 (simass/sivol, Eq. 4) results all show no seasonality and fixed values of 900 kg m^{-3} (MRI-ESM2-0), 910 kg m^{-3} (MPI and CNRM models) and 916 kg m^{-3} (the remaining models). Note that CNRM-CM6-1-HR did not provide the needed output (Table 2), but we expect it to feature the same density as the other CNRM models. For Method 1 (ice freeboard/hydrostatic, Eq. 5), several models show no seasonal variability, as ice density is simply fixed (no variable internal temperature or salinity) in their model setup, and these values are generally consistent with the Method 2 results. However, several Method 1 model estimates produce significant ice density seasonality, which we expect is due to inclusion of a variable internal ice temperature and a salinity physics scheme in the more advanced sea ice models. Interestingly, the bulk ice densities in the seasonally variable Method 1 models are notably higher or lower than the Method 2 results and the pure ice density value. More specifically, the CNRM models produced densities of $898 \pm 8 \text{ kg m}^{-3}$ (Arctic) and $883 \pm 8 \text{ kg m}^{-3}$ (Southern Ocean), whereas the MRI-ESM 2.0 model produced densities of $\sim 891 \pm 2 \text{ kg m}^{-3}$ (both hemispheres, minimal seasonal variability), all consistently lower than the Method 2 results and the pure ice/average bulk ice densities across both hemispheres. In contrast, the CESM2 models produced densities of $\sim 925 \pm 3 \text{ kg m}^{-3}$ (both hemispheres), and the NorESM2 models produced densities of $\sim 925 \pm 3 \text{ kg m}^{-3}$ (Arctic) and $\sim 922 \pm 3 \text{ kg m}^{-3}$ (Southern Ocean), consistently higher than the respective Method 2 results and the pure ice/average bulk ice densities. Cen-

ters with multiple model configurations (CNRM, NorESM2 and CESM2) produced bulk ice densities largely consistent across their respective configurations, as expected. At least one modeling group (CESM2) confirmed there are effectively two bulk ice densities in the model, with the Method 1 density reflecting the internal ice physics and the density used in the freeboard calculation and the Method 2 densities based on fixed salinity/temperature, used in atmospheric coupling assumptions (David Bailey, personal communication, 2025). We expect that similar discrepancies may be the cause of the differences observed across the other models.

Also shown in Fig. 4 is the bulk ice density inferred from the J22 empirical density parameterization applied to ICESat-2 data for the Arctic only (September to April data only due to thickness data availability), showing mean Arctic bulk ice densities of $\sim 931 \pm 3 \text{ kg m}^{-3}$. These densities are notably higher than the other estimates, although within the seasonal range of the CESM2/NorESM2 model Method 1 results and similar to values for bulk density quoted in recent studies (see discussion in Sect. 2.2.2). A more detailed assessment/validation of the bulk ice density estimates is considered beyond the scope of this study considering the significant scale differences between GCMs and field studies. Despite the lack of consensus in methodology, the multi-model annual mean from both methods across the 16-model subset is $912\text{--}914 \text{ kg m}^{-3}$, close to the pure ice bulk ice density assumption (dashed line, discussed in Sect. 2.2).

Next, we briefly explore the impact of differences in bulk ice density on estimates of total freeboard. In Fig. 5, we show estimates of total freeboard from the 17-model subset from both the direct total freeboard output (grid-cell mean snow thickness added to the ice freeboard output) and with total freeboard calculated as in Eqs. (2) and (3) using our own prescribed estimate of bulk ice density (916 kg m^{-3}). We also show observational estimates of total freeboard from ATL20 (using the interpolated/smoothed and perennial analysis) for reference; a more detailed plausible range assessment is provided in the following sections. The results in Fig. 5 show multi-model annual mean total freeboards slightly lower than ATL20 (22 cm vs. 27 cm annual means, respectively), with negligible differences using the prescribed/fixed ice density method compared to the direct output method (differences of $< 1\text{--}2 \text{ cm}$). Note that the ACCESS-CM2 model shows a significant increase in freeboard when using the prescribed ice density method, bringing those freeboards into much better agreement with ATL20, further suggesting issues with the underlying density/freeboard calculation. The MRI-ESM 2.0 model shows the biggest difference in total freeboard from the two methods in the 17-model subset ($\sim 3 \text{ cm}$ difference), with our prescribed density reducing the total freeboard estimate, as this was the model with the lowest estimated bulk ice density ($\sim 890\text{--}895 \text{ kg m}^{-3}$). The strength of the seasonal cycle (the interannual variability contribution to the total monthly variability across this period is low; not shown) appears broadly consistent between the models and ATL20




Figure 4. Derived mean (a) Arctic Ocean and (b) Southern Ocean bulk sea ice density in the 16-model CMIP6 subset (models with available freeboard, snow and ice thickness data, not including ACCESS-CM2). Circles indicate the annual mean for each model, and horizontal lines show the standard deviation across months (a proxy for the seasonality). A 50 % SIC masking is applied before spatial averaging. The dashed vertical black line shows the 916 kg m^{-3} “pure ice” bulk ice density assumption used in the IS2SITMOGR4 dataset. Also shown is the J22 ice density from IS2SITMOGR4, averaged between September and April 2018 to 2024.




Figure 5. Comparison of the mean (a) Arctic Ocean and (b) Southern Ocean total freeboard in the 17-model CMIP6 freeboard subset from the ensemble mean total freeboard output (circles), derived total freeboard assuming a fixed density of 916 kg m^{-3} (triangles), and total freeboard observations from ATL20 v4 (processed as discussed in Sects. 2 and 3) (red) for the ICESat-2 study period (November 2018–11 to April 2024). Horizontal lines show the seasonal variability (monthly standard deviation).

for the Arctic Ocean results (standard deviation of ~ 6 cm for the multi-model mean and ATL20); however, in the Southern Ocean, the ATL20 monthly total freeboard variability is noticeably lower (~ 3 cm) compared to the multi-model mean (~ 6 cm for the multi-model mean). The prescribed vs. variable bulk ice density freeboard estimates do not produce a significant difference in the strength of the freeboard seasonal cycle. A more detailed plausibility assessment accounting for observational uncertainty and internal variability is provided in the following sections. Overall, this bulk ice density and total freeboard analysis suggests the use of a fixed bulk ice density to calculate and analyze total freeboard across the wider 36-model subset that provides the necessary outputs of sea ice thickness and snow depth (Table 2), introduces only small additional uncertainty (< 1 – 2 cm), and ensures consistency in the density approximation used between models and observational freeboard estimates. We discuss the implications of this approach and suggestions for the wider community in the later discussion.

3.2 Plausibility assessments

Next, we explore the plausibility of the full CMIP6 suite, utilizing total freeboard metrics in addition to sea ice area across both hemispheres and winter Arctic sea ice thickness. We undertake this analysis for annual means and the two months that broadly represent the seasonal sea ice maximum and minimum months (September and March), with sea ice thickness data only included between September and April in the annual means due to data availability, as discussed in Sect. 2.2. To assess the plausibility of model output, we need to consider both the internal variability and observational uncertainty of our chosen metrics, which were discussed in detail in Sect. 2. Our estimates of Arctic Ocean mean total freeboard internal variability for the March, September and annual time periods for the 13 models with at least five ensemble members are shown in Fig. 6.

Our results show a significant spread in internal variability estimates across the different models, despite the application of the Bessel correction to account for sample size, with values ranging from ~ 1 to 6 cm depending on the CMIP6 model and time period of interest analyzed. A similar spread was noted in previous studies (Notz & SIMIP Community, 2020; Roach et al., 2020). While the inter-model internal variability spread is high, the differences are largely consistent across the three seasonal time periods. Using the wider time window to increase sampling (2015 to 2029, 10 rolling 6-year time means) provided a moderate impact on the internal variability estimates, with ~ 1 cm higher values on average for the annual, March and September CMIP6 mean internal variability estimates. At the individual model level, impact was highly variable, with some models showing no significant change, most models showing a moderate increase and UKESM1-0-LL especially showing a significant increase. Our multi-model mean estimates of Arc-

tic Ocean total freeboard internal variability using the 2015–2029 window are ~ 2.5 cm (annual), 2.2 cm (March) and 3.6 cm (September). Similar multi-model internal variability differences were found across our other metrics and hemispheres, with the multi-model means for all metrics and time periods summarized in Table 4. For the plausible range analysis below, we proceed with using the multi-model mean internal variability estimates in Table 4 applied to all models. We include a discussion of this assumption in our summary section below.

In Fig. 7, we show comparisons of the annual mean Arctic Ocean total freeboard estimated from the 36-model CMIP6 subset (see Table 2), calculated using our prescribed bulk ice density (916 kg m^{-3}) with observational estimates of total freeboard from ATL20 for our ICESat-2 study period (2018 to 2024). The analysis shows the ensemble mean for each model with a plausibility window for each model calculated following Sect. 2.3.1 based on both the high and low observational uncertainty estimates (Table 3) and multi-model mean estimates of internal variability (Table 4). The models are ranked in order of the ensemble mean differences between the model and observation for the given metric. Overall, the multi-model CMIP6 mean Arctic Ocean total freeboard (26 cm) is similar to ATL20 (27 cm) but with a large multi-model ensemble spread (~ 7 cm). In general, there are more models that are considered plausible (24), compared to implausible (12), at the 2σ , 95 % confidence level. The use of a high uncertainty estimate (3 cm instead of 1.5 cm) impacts the potential plausibility of 4 of the mid-lower-ranked models.

Also shown in Fig. 7 is the mean absolute error (MAE) in the grid-cell annual mean total freeboard for each model vs. ATL20, which captures the regional differences in the time-mean total freeboard between the models and observations. Regional-scale comparisons of climate model outputs are challenging and require deeper consideration of internal variability and model expectations, so we use ensemble mean spatial comparisons for added context of potential offsetting biases rather than as a plausibility constraint. The results show a multi-model MAE of ~ 8 cm compared to the < 1 cm spatial-mean multi-model mean difference, highlighting the significant role of off-setting regional differences. The non-monotonic increase in MAE down the hemisphere-mean difference ranked models alludes to the challenges of hemisphere-scale mean comparisons, although the lowest-ranked models still generally show higher MAE compared to the higher-ranked models, alluding to consistent hemisphere-scale biases in those models. We provide a more detailed plausibility assessment of regional biases in Sect. 3.3.

In Fig. 8, we show the same annual mean total freeboard plausibility analysis but for the Southern Ocean. In this case, the 37-model mean Southern Ocean total freeboard (16 cm) is significantly lower than ATL20 observations (26 cm) and exhibits a more significant multi-model ensemble spread (\sim




Figure 6. Estimates of internal variability (1 ensemble standard deviation, with Bessel correction) of Arctic Ocean total freeboard from ATL20 v4 (processed as discussed in Sects. 2 and 3) for the (a) annual, (b) March and (c) September periods for the 6-year mean over the ICESat-2 study period 2018–2024 (black lines) and running means across the longer 2015–2029 period (red crosses) for all the CMIP6 models with at least five ensemble members. Multi-model CMIP6 mean values are shown on the right.

Table 4. CMIP6 mean internal variability estimates (1 standard deviation) calculated across ensemble members and 10 rolling 6-year means over the period 2015–2029 for the different metrics of interest across the CMIP6 multi-model subset with at least five ensemble members. Note that we do not analyze Southern Ocean sea ice thickness in this study and that the Arctic annual mean thickness estimate does not include data between May and August (based on IS2SITMOGR4 data availability). Total freeboard CMIP6 mean values are shown in Fig. 6.

Metric	Hemisphere	Annual	March	September
Sea ice area (10 ⁶ km ²)	Arctic Ocean/Southern Ocean	0.38/0.31	0.35/0.17	0.54/0.45
Total freeboard (cm)	Arctic Ocean/Southern Ocean	2.4/1.0	2.3/3.3	3.6/1.1
Sea ice thickness (m)	Arctic Ocean	0.16/X	0.15/X	0.29/X




Figure 7. (a) Comparisons of annual mean (2018 to 2024) Arctic Ocean total freeboard estimates from the 37-model CMIP6 subset (ensemble means, using a fixed ice density) and observations from ICESat-2 ATL20 v4 (processed as discussed in Sects. 2 and 3). Horizontal lines on each model show the plausibility window based on internal variability and both the low (bars) and high (whiskers) observational uncertainty estimates. The red circle and vertical red line shows the ATL20 observational mean. ES: ensemble spread. (b) Mean absolute error (MAE) of the spatial differences across the 25 km × 25 km grid cells.




Figure 8. Same as in Fig. 7 but for the Southern Ocean.

9 cm). In general, there are more model estimates of Southern Ocean total freeboard that are considered implausible (29), compared to plausible (7) at the 2σ, 95 % confidence level. The use of the high observational uncertainty estimates results in only three more models being considered plausible at this confidence level. This result was widely expected considering the strong sea ice biases reported in earlier Southern

Ocean CMIP6 analyses (Roach et al., 2020). The Southern Ocean total freeboard MAEs are also higher than the Arctic (~ 14 cm multi-model mean). The strong overall negative bias across most of the CMIP6 models still appears to dominate the MAE contribution but is again explored more in Sect. 3.3.




Figure 9. Plausibility assessment of the full 40-model CMIP6 suite for the individual model ensemble means and the multi-model CMIP6 mean (top row) across all metrics, time periods and hemispheres. The plausibility index (ϕ) is calculated as in Eq. (7) using the 2015–2029 internal variability estimates and observational uncertainties listed in Table 3 (low uncertainty for the Arctic, high for the Southern Ocean). Models are sorted by the average ϕ across all values shown, with the four models missing freeboard/thickness output included at the bottom and the CMIP6 mean included at the top. Lower values are considered more plausible. Values of ϕ greater than 3 are outlined in black.

Figures showing the same plausibility analysis for both hemispheres across all other metrics and selected months (September and March) are provided in the Supplement (Figs. S6 to S18). We instead summarize results from all 15 combinations of metrics, time periods and hemispheres into a single analysis to more efficiently assess CMIP6 model performance. For this analysis, we use the plausibility index described in Sect. 2.3.1. that represents the plausibility of a given model's ensemble mean for the given metric considering internal variability, observational uncertainty and the direction of the overall model observation bias. For the Southern Ocean analysis, we use the “high” observational uncertainty estimate due to added observational complexities discussed earlier and noted in our comparisons and to ensure more models can be included in a constrained subset analysis.

This plausibility index analysis is shown in Fig. 9, with models ranked by the mean plausibility index averaged across all 15 different assessment combinations (columns in Fig. 9). We also highlight the models and metrics where the plausibility index is $> \pm 3$, which is equivalent to a 99 % confidence level that we can consider those models implausible based on our chosen criteria. Four of the models do not provide freeboard/thickness output and are thus included at the bottom of the figure. The CMIP6 model ensemble mean results show some of the best plausibility scores, including plausibility indices < 2 for all Arctic Ocean metrics. This analysis provides further evidence that the common approach of utilizing multi-model means for sea-ice-related projections can provide significant benefits in reducing model biases compared to analyzing any individual model. The results generally show better performance in the Arctic Ocean relative to the Southern Ocean, consistent with the results from

prior foundational CMIP6 analyses (Notz & SIMIP Community, 2020; Roach et al., 2020). We also observe more implausible results in September compared to the March and annual mean results. This seasonal dependence of plausibility is especially strong for total freeboard but also notable for thickness and area, though to a lesser extent. For the Southern Ocean results, we generally observe higher plausibility indices (less plausible) across both area and total freeboard variables as expected. Only one model, ACCESS-ESM1-5, produces a plausibility index < 3 for all 15 of our chosen metrics.

Several models show plausible results for all metrics except one, most frequently the March Southern Ocean total freeboard, and all four of these models show similar negative plausibility indices (-3.1 to -3.5), indicating a slight negative model freeboard bias in austral summer. It is also interesting to note that the March Southern Ocean area results (around the Antarctic sea ice minimum) are typically considered more plausible than the September and annual mean results, but this pattern generally reverses for freeboard. This result is driven partly by the low sea ice coverage in Southern Ocean sea ice in March limiting the magnitude of the bias and also our estimates of internal variability, especially for area – the lower the internal variability, the lower the plausibility threshold. This analysis raises important questions about the ability of current climate models to reasonably constrain more seasonal internal variability estimates, especially for low sea ice months when non-linear/non-Gaussian variability is likely. We discuss these concerns more in Sect. 4.

Overall, this combined plausibility analysis provides what we consider a useful framework for evaluating sea ice model output using a series of metrics and relevant observations. Caution should be used in applying such plausibility results, however, as the results are sensitive to choices regarding internal variability and observational uncertainty, as well as our chosen metrics. Model plausibility and constraint assessments depend ultimately on the overall goal. In the following section, we provide an example of CMIP6 model constraints based on this plausibility analysis, focusing on constraining modern-era CMIP6 sea ice simulations.

3.2.1 Impacts of model plausibility constraints

In Fig. 10, we show the impact of plausibility constraints on CMIP6 estimates of the seasonal cycle in sea ice volume across both the Arctic and Southern Ocean. The same plot but for the individual metrics of sea ice area, total freeboard and thickness is shown in the Supplementary Information (Fig. S19). The constrained model subset is generated independently for each hemisphere, with the assumption being that a model that performs well in one hemisphere should not be used to judge performance in the other hemisphere due to differences in priorities and model development efforts. We produce the constrained subsets in this example using just the annual mean plausibility indices and re-




Figure 10. Seasonal CMIP6 ensemble mean sea ice volume (lines and circles) and ensemble spread (shading, 1 standard deviation) from the unconstrained 36-model subset (black), the annual area constrained subset (blue), and the subset constrained with all considered annual variables, including area and freeboard for both hemispheres and sea ice thickness for the Arctic only (red).

quire a model to have a plausibility index < 3 for all available variables (area, total freeboard and thickness for the Arctic; area and freeboard for the Southern Ocean). We repeat this analysis for just the area metric, then for area, freeboard and thickness (thickness for the Arctic only) to assess the relative benefits of ICESat-2 data in this example constraint analysis. The goal is to explore how our plausibility constraints impact the CMIP6 multi-model mean and spread, and we focus here on the longer 2015 to 2035 time period. For the Arctic Ocean, this results in an area constrained subset of 31 models and an area, freeboard and thickness constrained subset of 27 models. For the Southern Ocean, this results in an area constrained subset of 21 models and an area and freeboard constrained subset of 14 models. We do not include the four models that provide only area (see Table 2) in either subset.

For the Arctic, the area constrained volume seasonal cycle shows a similar inter-model spread (shading in Fig. 10) and a slight increase in sea ice volume ($\sim 1 \times 10^3 \text{ km}^3$) across all months. When constraining also with freeboard and thickness, the inter-model spread reduces considerably (by $\sim 50\%$), and the volume across all months becomes slightly lower ($\sim 0\text{--}1 \times 10^3 \text{ km}^3$) than even the unconstrained volumes, especially in summer. For the Southern Ocean, the

area constrained volume seasonal cycle shows a reduced inter-model spread (again by $\sim 50\%$) and a more significant increase in sea ice volume across all months ($\sim 1\text{--}4 \times 10^3 \text{ km}^3$), especially in austral winter. When constraining also with freeboard, the inter-model spread shows a small extra reduction in some months, and the volume shows a slight additional increase compared to the area constrained results ($0\text{--}1 \times 10^3 \text{ km}^3$). Constraining models also with the March and September Arctic Ocean plausibility results (< 3 , not shown) results in bigger changes in the shape of the inter-model seasonal cycle and closer alignment with the observed seasonal cycles (shown in Fig. S19), including an increase in Arctic total freeboard in early fall, but at the cost of lower model ensemble size and the significant risk of overfitting. For the Southern Ocean, only three models (ACCESS-ESM1.5, AWI-CM-1-1-MR and MRI-ESM2-0) pass all seasonal Southern Ocean metrics, which was not deemed sufficient to determine a constrained model subset.

Overall, this basic analysis highlights the significant additional impact of constraining CMIP6 output with ICESat-2 freeboard and thickness estimates compared to sea ice area.

3.3 Regional plausibility assessments

We primarily explored hemisphere-mean comparisons in the preceding sections, but larger differences can be observed when analyzing model differences at more regional scales. Comparing models and observations at the grid-cell level can be misleading, as fully coupled climate model runs are not intended to perfectly capture our current reality but instead simulate expected climatic conditions within, ideally, a reasonable range of expected internal variability. To explore the ideas of regional plausibility, we follow the approach of other studies (e.g., Schaller et al., 2011) and attempt to characterize internal variability at the grid scale, towards a regional plausibility assessment as described in Sect. 2.3.3. As we assume the observational uncertainties in the hemisphere-mean analysis were driven primarily by biases or systematic error in the observations, we apply these same error estimates at each grid cell as a first-order uncertainty approximation. In reality, we expect a more complex regional combination of different error contributions.

Figure 11 shows maps of the annual Arctic Ocean total freeboard difference between the 36-model CMIP6 subset and ATL20. The areas with hatching indicate regions that are considered implausible based on the regional plausibility index. The regional internal variability estimate calculated from the CMIP6-mean ensemble spread from models with at least five ensemble members is shown also (2 standard deviations), showing a mean value of $\sim 4\text{--}8 \text{ cm}$ across the Arctic, with some small increases along the coast/ice edge. Note again that the 3 cm total freeboard uncertainty is added to this internal variability to produce the regional plausibility threshold. The models are ranked by the overall MAE and help visualize the regional contributions to

the overall bias and regions of implausibility. In the better-performing models, specific regions appear with some consistency across the models, including a few models with negative freeboard anomalies in the thicker Central Arctic regime along the Greenland/Canadian Arctic coast, as well as the Chukchi Sea, and some models with positive anomalies north of Greenland and the Beaufort Sea. Many of these regions represent key dynamical features of the Arctic sea ice cover, e.g., the Beaufort Gyre (Petty et al., 2016) and the Wandel Sea (Schweiger et al., 2021), and allude to deficiencies in atmospheric dynamics (e.g., wind forcing).

For the lower-performing models, the regions of implausible bias extend further across the Arctic and allude to broader thermodynamic drivers of the biases. A full thermodynamic/dynamic accounting for the regional differences is considered beyond the scope of this study but could form the basis of future model development evaluation methods. Difference maps of the remaining Arctic Ocean metrics are shown in the Supplementary Information (Figs. S20–S31). In general, the sea ice area biases are more prevalent in the marginal seas, especially in March, as expected, while the regional distribution of biases and implausibility in thickness align closely with these total freeboard results, further highlighting the potential for total freeboard to provide useful regional diagnostics as a proxy for thickness biases. Similar regional bias patterns in CMIP6 models were observed in Watts et al. (2021), which included comparisons to CryoSat-2 and the earlier ICESat mission.

We next use regional composites, grouping the models based on the biases measured within the Eastern and/or Western Arctic (denoted by the dashed line in Fig. 11 from 60°E to 120°W), to highlight the predominant regional manifestations in freeboard and thickness biases seen in our spatial difference maps. Figure 12 shows composite maps and difference plots based on different combinations of the direction of the total Eastern Arctic (East) vs. Western Arctic (West) bias for both freeboard and thickness (annual means). Note that for this annual mean Arctic Ocean total freeboard comparison, there were no models that showed a $-$ East and $+$ West bias.

For freeboard, nine models feature a $+$ East and $-$ West bias, 11 models feature a $+$ East and $+$ West bias, and 17 models feature a $-$ East and $-$ West bias. The $+$ East and $-$ West bias models show generally consistent freeboard across the Arctic and thus fail to capture the strong gradient from the Western Arctic to the East, while the other model composites show a clear West to East freeboard gradient but with freeboards that are either too high or too low on average. For the $-$ East and $-$ West models, the models show better agreement with observations in the Beaufort Sea region on average but with no clear agreement across the models, again suggestive of unique dynamical challenges for models in this region. The sea ice thickness (September through April) bias composites show virtually identical spatial composite difference patterns and model agreement relative to the freeboard re-




Figure 11. Total freeboard difference of the 2018 to 2024 annual mean for each CMIP6 model (ensemble mean) relative to ATL20 v4 (processed as discussed in Sects. 2 and 3). Areas with hatching indicate regions that are considered implausible based on the regional plausibility index analysis. Models are shown in order of lowest (top left by row) to highest mean absolute error (MAE). The dashed line (50° E and 130° W meridians) is used for the eastern and western sector analysis. Note that only model grid cells within our Inner Arctic Ocean domain and that are “perennially ice-covered” in ATL20 v4 are included here. The final panel shows our estimate of CMIP6 mean regional internal variability (2 standard deviations, Sect. 2.3.3).

sults, with only small differences in the distribution of models across the three composites. Composite maps for both September and March are provided in the Supplementary Information (Figs. S32 and S33), which show similar regional patterns and model composites, with reduced coverage and more models showing a – East and – West bias in September.

In Fig. 13, we show the annual mean total freeboard difference analysis for the Southern Ocean. The mean internal variability is similar to the Arctic (~ 4 – 8 cm), with the high-

est values in the western Weddell Sea, where the ice is generally thickest in the models. As shown in our basin-scale analyses, the Southern Ocean results exhibit larger overall biases compared to the Arctic. However, Southern Ocean sea ice is also generally considered more regionally variable, with significant regional differences in climate driven by its unique geography (where the sea ice pack surrounds the continent of Antarctica and covers three different ocean basins) providing further motivation for such regional assessments. As in the Arctic analysis, there are strong differences across the




Figure 12. (a–d) Annual multi-model CMIP6 ensemble mean total freeboard for all models (a) and then composites based on their East/West bias pattern (b–d). (e–h) Top row minus the ATL20 v4 annual mean total freeboards (processed as in Sects. 2 and 3). (i–p) show the same but for September to April Arctic sea ice thickness from CMIP6 and differences with IS2SITMOGR4 v3 thickness data. Stippling indicates where at least 80 % of the models in the respective composite agree with the sign of the difference.

models where the freeboard biases manifest but also some evidence of problem areas, including the Weddell Sea and other shelf sea regions. It is also interesting to note the significant number of models that show an overall negative bias relative to the observations everywhere except the Ross Sea, a region of thin ice and strong wind-driven polynya activity.

We also observe more consistent negative model biases along the ice edge in the Southern Ocean analysis. These “biases” should be treated with more caution, as wave contamination along the ice edge is a known issue with ATL20 (data are masked below 50 % concentration to try and mitigate this) and is expected to impact the Southern Ocean more than the Arctic due to stronger wave activity (Horvat et al., 2020). In addition, significant implausible negative biases are observed around the Antarctic coastline in several models, which should be a source of future model develop-

ment focus. In an additional analysis (not shown), we confirmed this was present across the native model grids and not introduced in our regridding step. In Fig. 14, we repeat the composite analysis, but as there was no clear longitudinal dependence on the biases, we simply group the models based on the overall direction of the mean bias. For the seven models with positive bias, there is general agreement on the regional pattern of the bias within the ice pack; however, the negative differences around the coast and the ice edge appear more model-specific. The 30 models with negative bias are consistently negative except for the Ross Sea and the eastern Weddell Sea. These again suggest issues with the underlying dynamics and the response to large-scale circulation or the simulated internal variability not fully capturing the regional patterns that would alter the plausibility of the observed biases. September and March composite maps are also shown




Figure 13. Same as in Fig. 11 but for the Southern Ocean.

in the Supplementary Information (Figs. S34 and S35), with the September results showing similar regional distributions, while the limited ice cover in March significantly reduces the value of the regional analysis.

4 Discussion

Sea ice bulk density has long been a source of uncertainty in remote sensing efforts and has arguably not been an explicit focus of global sea ice modeling efforts to date. However, assessments of freeboard and the introduction of variable density schemes provide motivation to increase documentation and focus on the underlying sea ice density assumptions or parameterization schemes used. More direct outputs of freeboard and assessments of these outputs instead of our own derived freeboards could additionally help mitigate the uncertainty introduced by our own density choices. Recent ob-

servational analyses have also alluded to significant increases in bulk ice density compared to historical estimates, linked especially to the thinner and younger Arctic sea ice pack, but also potential issues with prior data collection campaigns and interpretation (Jutila et al., 2022; Salganik et al., 2025; Shi et al., 2023; Zhou et al., 2024). This trend could continue as the Arctic (and perhaps Southern Ocean) ice pack continues to thin and lose older ice, so models with variable and realistic density parameterizations (based on prognostic internal temperature and salinity) calibrated to current-day observations are encouraged.

Uncertainty quantification in remotely sensed sea ice products continues to be a challenge. We provide one basic approach (high vs. low fixed uncertainty estimates based on a literature review) and hope that more community engagement and consensus can be undertaken in the near future. Uncertainty quantification needs to include a full accounting of possible error sources, including sampling/representation




Figure 14. Same as in Fig. 11 but for the Southern Ocean freeboard. No regional bias composite is shown – only composites where basin-averaged differences from ATL20 v4 are either positive or negative.

errors. The ongoing ESA-funded Sea Ice-thickness product iNter-comparison eXerciSe (SIN'XS) initiative (<https://sinxs.noveltis.fr>, last access: 12 September 2025) aims to increase community focus on this issue. In addition, ensemble-based methods would provide a more robust framework for assessing observational uncertainty and provide important insights into the regional uncertainty estimates. Increased development and uncertainty estimates of the freeboard and thickness products in more marginal ice regimes are urgently needed, especially in the Southern Ocean, where some of the model biases were more questionable. Sea ice reanalysis systems or fusion with other sensors is urgently needed.

The current time period of ICESat-2 freeboard and thickness data (2018 to 2024 at the time of writing) is short and not representative of typical climate timescales used in model assessment efforts. The short period increases the contribution of internal variability and the challenge of quantifying internal variability across our chosen metrics, especially from models with lower ensemble member counts and significant model biases. Our utilized method demonstrates a potential approach in the absence of longer time series (an increased window around our current time period), with the benefit of limiting trend contamination. More years of observational data can both increase confidence in our internal variability estimates and reduce its contribution to overall uncertainty, with important implications for our plausibility assessments. We hope future work towards integration of sea ice altimetry data from NASA's ICESat (2003 to 2008) and ESA's CryoSat-2 (2010 to present) mission, together with advances in sea ice reanalyses, will provide important benefits here. Longer records will also enable assessments of thickness and volume trends, especially for the Arctic, where we have more

confidence in the snow loading inputs. Regardless of the time period, different models can produce very different internal variability estimates, which provides a further challenge and source of uncertainty we need to consider more in future work. Utilizing independent large ensembles for constraining internal variability independent of the CMIP6 suite is an alternative approach worth exploring. The ability of models to accurately capture regional internal variability is questionable, which prevents us from more confidently prescribing implausible regional biases in the models.

The summer assessments were more challenging to decipher due to the bigger role of coverage issues and differences between the models and observations. We employed a “perennial” ice masking approach to mitigate coverage issues and improve the robustness of those comparisons, but comparing biases of global climate models in these small regions poses additional questions, e.g., how much should we expect climate models to simulate these more fractional ice packs? Other studies have mitigated this to some degree by focusing more on the strength of the seasonal cycle (Massonnet et al., 2012), which could be worth considering. Similarly, understanding seasonal snow evolution and biases and how they relate to our seasonal biases in freeboard and thickness would provide a logical next phase of this study.

Our regional analysis provided new insights into how total freeboard can be used to diagnose regional sea ice biases in models, with our east–west composite analysis providing a simple framework for assessing and grouping models. Increased focus on regional sea ice internal variability estimates in large ensembles should be explored to enhance confidence in the regional plausibility results. East–West Arctic Ocean sea ice thickness anomalies have been discussed

in previous studies, with links to large-scale atmospheric dynamics including the North Atlantic Oscillation (NAO) (Zhang et al., 2000), which could help diagnose the cause of regional biases across models. Understanding the underlying drivers of bias was considered beyond the scope of this study. The Southern Ocean reanalysis results were more mixed and not as clearly divisible by region or longitude. More sophisticated machine learning tools, e.g., principal component analysis or self-organizing maps, could provide more insight into the dominant regional sea ice bias patterns in the Southern Ocean.

Model calibration efforts depend crucially on the research question posed. Our study focused on plausibility methodologies and the potential benefits of ICESat-2 sea ice altimetry data for evaluating global climate model outputs of sea ice, with a brief demonstration of the impact on constraining seasonal cycles in sea ice volume across both hemispheres. We used the hemisphere-mean plausibility scores in a simple exclusion subset approach, but weighting based on these scores is an alternative option. Additional work could explore the resultant impact of our constrained/plausible subset on associated metrics including surface atmosphere–ice–ocean fluxes and sea ice freshwater fluxes within and out of the polar regions, building on previous CMIP6 studies (Keen et al., 2021; Zanowski et al., 2021). In addition, we hope that these shorter-time-period mean observational data constraints can provide added benefits when combined with more commonly used plausibility metrics (e.g., sea ice sensitivity to temperature over the multi-decadal time period) and/or recalibration approaches (e.g., Bonan et al., 2021) to provide better constrained longer-term CMIP6 sea ice predictions across metrics. Again, this is something we hope to explore in future work with the community.

Finally, assessments of forced sea ice–ocean models, regional models or sea ice/ocean reanalyses could benefit from similar freeboard/thickness diagnostics presented in this study. The increased emphasis on accuracy and agreement with observations would mitigate internal variability considerations and increase focus on the observational uncertainty estimates and associated issues, including representation error. We chose to focus here on fully coupled climate models to explore the various additional issues introduced by the coupled model systems, e.g., the benefit of a large spread in model configurations and outputs, and the challenge of internal variability attribution and contribution. Models with sea ice embedded in the ocean (instead of the levitating assumptions typical of GCMs) may have additional motivation to assess and constrain freeboard.

5 Conclusions

This study provided a comprehensive evaluation of sea ice simulations from the Coupled Model Intercomparison Project Phase 6 (CMIP6) using ICESat-2 altimetry observa-

tions, in addition to sea ice area from passive microwave, introducing new insights into model plausibility and constraints. Freeboard comparisons benefit from accurate observations by satellite laser altimetry and motivate increased focus on bulk sea ice density. The short record increases the challenge in internal variability assessments, which generally contributed more than our observational uncertainty estimates to our plausibility metrics. While CMIP6 Arctic sea ice simulations showed reasonable agreement with ICESat-2 freeboard and thickness data, especially for the multi-model mean, more significant biases were present in the Southern Ocean CMIP6 models, as was largely expected from previous studies. We adopted a similar plausibility approach at the grid scale to highlight the regional manifestation of these model biases and potential regions of implausibility across the models. An east–west composite approach highlighted the consistent model agreement in the types of regional biases observed, often linked to key dynamical features of the ice cover. The regional Southern Ocean analysis was more mixed and could benefit from additional analysis to find consistent patterns of agreement/plausibility.

We demonstrated an example of our plausibility constraints on the seasonal cycles in both Arctic and Southern Ocean sea ice volume, with the freeboard and thickness data providing crucial additional impacts over the standard area constraints in terms of the mean monthly values and inter-model spread, highlighting the role ICESat-2 data can play in CMIP6 model evaluation and constraint.

Future research should prioritize improved uncertainty quantification and expanded assessments of associated metrics in the constrained analysis. More years of data from ICESat-2 and leveraging altimetry data from prior and ongoing satellite missions, e.g., NASA's original ICESat mission and ESA's CryoSat-2, could increase the utility of the freeboard and thickness constraints presented here. The study also emphasizes the importance of addressing challenges in regional sea ice dynamics, which could form the basis of future model development assessments. With the upcoming release of CMIP7, we suggest that these new assessment concepts can be utilized in tandem with more traditional assessment methods to better constrain current and future variability in sea ice and their associated climate impacts.

Code availability. We provide a Jupyter Notebook on GitHub under the *MIT license* (<https://github.com/cjcardinale/CMIP6-ICESat2-Sealce-Petty-2025>, last access: 12 September 2025) and archived on Zenodo (<https://doi.org/10.5281/zenodo.15849468>, Cardinale and Petty, 2025) that showcases the primary aspects of our analysis workflow: code to access our wrangled/regridded CMIP6 data and observations from AWS S3, code to estimate internal variability and model plausibility indices, and code to produce key figures shown in the paper.

Data availability. CMIP6 data in the cloud can be accessed using the `intake.open_esm_datastore()` function from the `intake-esm` Python package, available on GitHub under an Apache-2 license (<https://github.com/intake/intake-esm> (currently at v2025.2.3), last access: 12 September 2025) and archived on Zenodo (<https://doi.org/10.5281/zenodo.14816755>, Banihirwe et al., 2025). The following JSON files served as our data catalogs:

- Sea ice freeboard data: <https://storage.googleapis.com/cmip6/cmip6-pgf-ingestion-test/catalog/catalog.json> (last access: 12 September 2025)
- All other CMIP6 variables: <https://cmip6-pds.s3.amazonaws.com/pangeo-cmip6.json> (last access: 12 September 2025)

We used both the `intake-esgf` and `esgf-pyclient` packages, which provide a Python interface for searching and accessing datasets from ESGF, including CMIP6 data hosted on-prem. The `intake-esgf` package is available on GitHub under a BSD-3-Clause license (<https://github.com/esgf2-us/intake-esgf> (currently at v2025.6.6), last access: 12 September 2025) and archived on Zenodo (Collier et al., 2025). The `esgf-pyclient` package is available on GitHub under a BSD-3-Clause license (<https://github.com/ESGF/esgf-pyclient> (currently at v0.3.1), last access: 12 September 2025). Both packages allow users to query ESGF metadata, filter search results and retrieve URLs for downloading netCDF files. We accessed the CMIP6 data through the Lawrence Livermore National Laboratory (LLNL) ESGF node and the following URL: <https://esgf-node.llnl.gov/esg-search> (last access: 12 September 2025). Data can then be loaded via OPeNDAP and the Xarray Python Library.

We use both final (OSI-450-a, up to 2020, https://doi.org/10.15770/EUM_SAF_OSI_0013, OSI SAF, 2022a) and interim (OSI-430-a, 2021 onwards, https://doi.org/10.15770/EUM_SAF_OSI_0014, OSI SAF, 2022b) OSI SAF sea ice concentration data, which can be accessed from the THREDDS Data Server hosted by the Norwegian Meteorological Institute and loaded via OPeNDAP with the following example URLs:

- OSI-450-a: https://thredds.met.no/thredds/catalog/osisaf/met.no/reprocessed/ice/conc_450a_files/monthly/catalog.html (OSI SAF, 2022a)
- OSI-430-a: https://thredds.met.no/thredds/catalog/osisaf/met.no/reprocessed/ice/conc_cra_files/monthly/catalog.html (OSI SAF, 2022b)

ATL20 total freeboard data (we use Version 4 in this study) are hosted officially through the NSIDC at <https://doi.org/10.5067/ATLAS/ATL20.004> (Petty et al., 2023a).

ICESat-2 IS2SITMOGR4 sea ice thickness data (we use Version 3 in this study) are hosted officially through the NSIDC at <https://doi.org/10.5067/ZCSU8Y5U1BQW> (Petty et al., 2023b).

Supplement. The supplement related to this article is available online at <https://doi.org/10.5194/gmd-18-6313-2025-supplement>.

Author contributions. AP planned the study and drafted the paper with significant input from CC and MS. CC wrangled and analyzed the data and generated all the figures with input from AP/MS. MS produced the schematic and helped plan/draft the paper.

Competing interests. The contact author has declared that none of the authors has any competing interests.

Disclaimer. Publisher's note: Copernicus Publications remains neutral with regard to jurisdictional claims made in the text, published maps, institutional affiliations, or any other geographical representation in this paper. While Copernicus Publications makes every effort to include appropriate place names, the final responsibility lies with the authors.

Acknowledgements. Thanks to the creators and maintainers of the NASA-supported CryoCloud community for providing the AWS cloud-based Jupyter Hub that was used for all of our analysis (CryoCloud is supported by NASA grants 80NSSC22K1877 and 80NSSC23K0002). The platform radically simplifies the process of getting spun-up on AWS resources and enables team members to work collaboratively in the same computational environment. Thanks also to members of Pangeo and the Learning the Earth with Artificial Intelligence and Physics (LEAP) center, who created, maintain and continue to update the Pangeo Analysis-Ready Cloud-Optimized CMIP6 data catalog. Thanks to the various modeling groups who have worked on developing and producing the CMIP6 runs used here, as well as the groups who produced and developed the various observational products used in this study, including members of the ICESat-2 mission and the EUMETSAT Ocean and Sea Ice Satellite Application Facility (OSI SAF). Thanks to the developers and maintainers of the various open-source Python packages we utilized in this study, especially `intake-esm`, `intake-esgf` and `esgf-pyclient`. Thanks also to David Bailey of NCAR for very useful discussions on the internal physics of CESM2/CICE and the calculation of ice freeboard.

Financial support. All team members were supported through a NASA ICESat-2 Science Team award (grant no. 80NSSC23K1253).

Review statement. This paper was edited by Christopher Horvat and reviewed by two anonymous referees.

References

- Alexandrov, V., Sandven, S., Wahlin, J., and Johannessen, O. M.: The relation between sea ice thickness and freeboard in the Arctic, *The Cryosphere*, 4, 373–380, <https://doi.org/10.5194/tc-4-373-2010>, 2010.
- Andersson, T. R., Hosking, J. S., Pérez-Ortiz, M., Paige, B., Elliott, A., Russell, C., Law, S., Jones, D. C., Wilkinson, J., Phillips, T.,

- Byrne, J., Tietsche, S., Sarojini, B. B., Blanchard-Wrigglesworth, E., Aksenov, Y., Downie, R., and Shuckburgh, E.: Seasonal Arctic sea ice forecasting with probabilistic deep learning, *Nat. Commun.*, 12, 5124, <https://doi.org/10.1038/s41467-021-25257-4>, 2021.
- Bagnardi, M., Kurtz, N. T., Petty, A. A., and Kwok, R.: Sea Surface Height Anomalies of the Arctic Ocean From ICESat-2: A First Examination and Comparisons With CryoSat-2, *Geophys. Res. Lett.*, 48, e2021GL093155, <https://doi.org/10.1029/2021GL093155>, 2021.
- Banihirwe, A., Grover, M., bonnland, Kent, J., Bourgault, P., Long, M., Squire, D., Turner, C., Busecke, J., Spring, A., Schulz, H., Lewis, J., Beucher, R., RondeauG, Bartholomew, S. L., Kölling, T., and garciampred: intake/intake-esm: v2025.2.3, Zenodo [data set], <https://doi.org/10.5281/zenodo.14816755>, 2025.
- Bonan, D. B., Schneider, T., Eisenman, I., and Wills, R. C. J.: Constraining the Date of a Seasonally Ice-Free Arctic Using a Simple Model, *Geophys. Res. Lett.*, 48, e2021GL094309, <https://doi.org/10.1029/2021GL094309>, 2021.
- Cardinale, C. and Petty, A. A.: cjc-cardinale/CMIP6-ICESat2-SeaIce-Petty-2025, Zenodo [code], <https://doi.org/10.5281/zenodo.15849468>, 2025.
- Cavalieri, D., Parkinson, C., Gloersen, P., and Zwally, J. H.: Sea Ice Concentrations from Nimbus-7 SMMR and DMSP SSM/I-SSMIS Passive Microwave Data, Version 1. [January 2000–December 2010], NASA DAAC at the National Snow and Ice Data Center [data set], <https://doi.org/10.5067/8GQ8LZQVL0VL>, 1996.
- Collier, N., Grover, M., Stachek, J., and Huard, D.: intake-esgf, Zenodo [code], <https://doi.org/10.5281/zenodo.15609526>, 2025.
- Comiso, J. C., Cavalieri, D. J., Parkinson, C. L., and Gloersen, P.: Passive microwave algorithms for sea ice concentration: A comparison of two techniques, *Remote Sens. Environ.*, 60, 357–384, [https://doi.org/10.1016/S0034-4257\(96\)00220-9](https://doi.org/10.1016/S0034-4257(96)00220-9), 1997.
- Davy, R. and Outten, S.: The Arctic Surface Climate in CMIP6: Status and Developments since CMIP5, *J. Climate*, 33, 8047–8068, <https://doi.org/10.1175/JCLI-D-19-0990.1>, 2020.
- Eyring, V., Bony, S., Meehl, G. A., Senior, C. A., Stevens, B., Stouffer, R. J., and Taylor, K. E.: Overview of the Coupled Model Intercomparison Project Phase 6 (CMIP6) experimental design and organization, *Geosci. Model Dev.*, 9, 1937–1958, <https://doi.org/10.5194/gmd-9-1937-2016>, 2016.
- Fons, S. W., Kurtz, N. T., Bagnardi, M., Petty, A. A., and Tillig, R. L.: Assessing CryoSat-2 Antarctic Snow Freeboard Retrievals Using Data From ICESat-2, *Earth Space Sci.*, 8, e2021EA001728, <https://doi.org/10.1029/2021EA001728>, 2021.
- Garnier, F., Fleury, S., Garric, G., Bouffard, J., Tsamados, M., Laforge, A., Bocquet, M., Fredensborg Hansen, R. M., and Remy, F.: Advances in altimetric snow depth estimates using bi-frequency SARAL and CryoSat-2 Ka–Ku measurements, *The Cryosphere*, 15, 5483–5512, <https://doi.org/10.5194/tc-15-5483-2021>, 2021.
- Giles, K. A., Laxon, S. W., Wingham, D. J., Wallis, D. W., Krabill, W. B., Leuschen, C. J., McAdoo, D., Manizade, S. S., and Raney, R. K.: Combined airborne laser and radar altimeter measurements over the Fram Strait in May 2002, *Remote Sens. Environ.*, 111, 182–194, <https://doi.org/10.1016/j.rse.2007.02.037>, 2007.
- Goosse, H., Kay, J. E., Armour, K. C., Bodas-Salcedo, A., Chepfer, H., Docquier, D., Jonko, A., Kushner, P. J., Lecomte, O., Massonnet, F., Park, H.-S., Pithan, F., Svensson, G., and Vancoppenolle, M.: Quantifying climate feedbacks in polar regions, *Nat. Commun.*, 9, 1919, <https://doi.org/10.1038/s41467-018-04173-0>, 2018.
- Hibler, W. D.: A Dynamic Thermodynamic Sea Ice Model, *J. Phys. Oceanogr.*, 9, 815–846, [https://doi.org/10.1175/1520-0485\(1979\)009<0815:ADTSIM>2.0.CO;2](https://doi.org/10.1175/1520-0485(1979)009<0815:ADTSIM>2.0.CO;2), 1979.
- Holland, M. M., Finnis, J., and Serreze, M. C.: Simulated Arctic Ocean Freshwater Budgets in the Twentieth and Twenty-First Centuries, *J. Climate*, 19, 6221–6242, <https://doi.org/10.1175/JCLI3967.1>, 2006.
- Holland, M. M., Serreze, M. C., and Stroeve, J.: The sea ice mass budget of the Arctic and its future change as simulated by coupled climate models, *Clim. Dynam.*, 34, 185–200, <https://doi.org/10.1007/s00382-008-0493-4>, 2010.
- Horvat, C., Blanchard-Wrigglesworth, E., and Petty, A.: Observing Waves in Sea Ice With ICESat-2, *Geophys. Res. Lett.*, 47, e2020GL087629, <https://doi.org/10.1029/2020GL087629>, 2020.
- Hunke, E. C., Lipscomb, W. H., Turner, A. K., Jeffery, N., and Elliott, S.: CICE: the Los Alamos Sea Ice Model Documentation and Software User's Manual Version 5.1, T-3 Fluid Dynamics Group, Los Alamos National Laboratory, Los Alamos, GitHub [code], <https://github.com/CICE-Consortium/CICE-svn-trunk/blob/main/cicedoc/cicedoc.pdf> (last access: 12 September 2025), 2015.
- IPCC – Intergovernmental Panel on Climate Change (Ed.): Ocean, Cryosphere and Sea Level Change, in: *Climate Change 2021 – The Physical Science Basis: Working Group I Contribution to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change*, Cambridge University Press, Cambridge, 1211–1362, <https://doi.org/10.1017/9781009157896.011>, 2023.
- Jahn, A., Holland, M. M., and Kay, J. E.: Projections of an ice-free Arctic Ocean, *Nat. Rev. Earth Environ.*, 5, 164–176, <https://doi.org/10.1038/s43017-023-00515-9>, 2024.
- Jutila, A., Hendricks, S., Ricker, R., von Albedyll, L., Krumpen, T., and Haas, C.: Retrieval and parameterisation of sea-ice bulk density from airborne multi-sensor measurements, *The Cryosphere*, 16, 259–275, <https://doi.org/10.5194/tc-16-259-2022>, 2022.
- Kacimi, S. and Kwok, R.: Arctic Snow Depth, Ice Thickness, and Volume From ICESat-2 and CryoSat-2: 2018–2021, *Geophys. Res. Lett.*, 49, e2021GL097448, <https://doi.org/10.1029/2021GL097448>, 2022.
- Kay, J. E., DeRepentigny, P., Holland, M. M., Bailey, D. A., DuVivier, A. K., Blanchard-Wrigglesworth, E., Deser, C., Jahn, A., Singh, H., Smith, M. M., Webster, M. A., Edwards, J., Lee, S.-S., Rodgers, K. B., and Rosenbloom, N.: Less Surface Sea Ice Melt in the CESM2 Improves Arctic Sea Ice Simulation With Minimal Non-Polar Climate Impacts, *J. Adv. Model. Earth Syst.*, 14, e2021MS002679, <https://doi.org/10.1029/2021MS002679>, 2022.
- Keen, A., Blockley, E., Bailey, D. A., Boldingh Debernard, J., Bushuk, M., Delhay, S., Docquier, D., Feltham, D., Massonnet, F., O'Farrell, S., Ponsoni, L., Rodriguez, J. M., Schroeder, D., Swart, N., Toyoda, T., Tsujino, H., Vancoppenolle, M., and Wyser, K.: An inter-comparison of the mass budget of the Arctic sea ice in CMIP6 models, *The Cryosphere*, 15, 951–982, <https://doi.org/10.5194/tc-15-951-2021>, 2021.

- Kern, S., Laverne, T., Notz, D., Pedersen, L. T., Tonboe, R. T., Saldo, R., and Sørensen, A. M.: Satellite passive microwave sea-ice concentration data set intercomparison: closed ice and ship-based observations, *The Cryosphere*, 13, 3261–3307, <https://doi.org/10.5194/tc-13-3261-2019>, 2019.
- Kern, S., Laverne, T., Pedersen, L. T., Tonboe, R. T., Bell, L., Meyer, M., and Zeigermann, L.: Satellite passive microwave sea-ice concentration data set intercomparison using Landsat data, *The Cryosphere*, 16, 349–378, <https://doi.org/10.5194/tc-16-349-2022>, 2022.
- Kim, Y.-H., Min, S.-K., Gillett, N. P., Notz, D., and Malinina, E.: Observationally-constrained projections of an ice-free Arctic even under a low emission scenario, *Nat. Commun.*, 14, 3139, <https://doi.org/10.1038/s41467-023-38511-8>, 2023.
- Kurtz, N. T. and Markus, T.: Satellite observations of Antarctic sea ice thickness and volume, *J. Geophys. Res.-Oceans*, 117, C08025, <https://doi.org/10.1029/2012JC008141>, 2012.
- Kwok, R. and Cunningham, G. F.: ICESat over Arctic sea ice: Estimation of snow depth and ice thickness, *J. Geophys. Res.-Oceans*, 113, C08010, <https://doi.org/10.1029/2008JC004753>, 2008.
- Kwok, R., Kacimi, S., Markus, T., Kurtz, N. T., Studinger, M., Sonntag, J. G., Manizade, S. S., Boisvert, L. N., and Harbeck, J. P.: ICESat-2 surface height and sea-ice freeboard assessed with ATM lidar acquisitions from Operation IceBridge, *Geophys. Res. Lett.*, 44, 11228–11236, <https://doi.org/10.1029/2019GL084976>, 2019.
- Kwok, R., Petty, A., Cunningham, G., Markus, T., Hancock, D., Ivanoff, A., Wimert, J., Bagnardi, M., and Kurtz, N.: ATLAS/ICESat-2 L3A Sea Ice Heights, Version 5, USA NSIDC Natl. Snow Ice Data Cent. [data set], <https://doi.org/10.5067/ATLAS/ATL07.005>, 2021.
- Kwok, R., Petty, A., Wimert, J., Bagnardi, M., Cunningham, G., Hancock, D., Ivanoff, A., and Kurtz, N.: Ice, Cloud, and Land Elevation Satellite-2 Project: Algorithm Theoretical Basis Document (ATBD) for Sea Ice Products, Release 006, ICESat-2 Project, <https://doi.org/10.5067/9VT7NJWOTV3I>, 2022.
- Kwok, R., Petty, A., Cunningham, G., Markus, T., Hancock, D., Ivanoff, A., Wimert, J., Bagnardi, M., and Kurtz, N.: ATLAS/ICESat-2 L3A Sea Ice Freeboards, Version 6, USA NSIDC Natl. Snow Ice Data Cent. [data set], <https://doi.org/10.5067/ATLAS/ATL10.006>, 2023.
- Laverne, T., Sørensen, A. M., Kern, S., Tonboe, R., Notz, D., Aaboe, S., Bell, L., Dybkjær, G., Eastwood, S., Gabarro, C., Heygster, G., Killie, M. A., Brandt Kreiner, M., Lavelle, J., Saldo, R., Sandven, S., and Pedersen, L. T.: Version 2 of the EUMETSAT OSI SAF and ESA CCI sea-ice concentration climate data records, *The Cryosphere*, 13, 49–78, <https://doi.org/10.5194/tc-13-49-2019>, 2019.
- Laxon, S. W., Giles, K. A., Ridout, A. L., Wingham, D. J., Willatt, R., Cullen, R., Kwok, R., Schweiger, A., Zhang, J., Haas, C., Hendricks, S., Krishfield, R., Kurtz, N., Farrell, S., and Davidson, M.: CryoSat-2 estimates of Arctic sea ice thickness and volume, *Geophys. Res. Lett.*, 40, 732–737, <https://doi.org/10.1002/grl.50193>, 2013.
- Lenzen, N., Schmidt, G. A., Hendrickson, M., Jacobs, P., Menne, M. J., and Ruedy, R.: A NASA GISTEMPv4 Observational Uncertainty Ensemble, *J. Geophys. Res.-Atmos.*, 129, e2023JD040179, <https://doi.org/10.1029/2023JD040179>, 2024.
- Massonnet, F., Fichefet, T., Goosse, H., Bitz, C. M., Philippon-Berthier, G., Holland, M. M., and Barriat, P.-Y.: Constraining projections of summer Arctic sea ice, *The Cryosphere*, 6, 1383–1394, <https://doi.org/10.5194/tc-6-1383-2012>, 2012.
- Massonnet, F., Vancoppenolle, M., Goosse, H., Docquier, D., Fichefet, T., and Blanchard-Wrigglesworth, E.: Arctic sea-ice change tied to its mean state through thermodynamic processes, *Nat. Clim. Change*, 8, 599–603, <https://doi.org/10.1038/s41558-018-0204-z>, 2018.
- Meier, W. and Stewart, J. S.: Arctic and Antarctic Regional Masks for Sea Ice and Related Data Products, Version 1, NASA National Snow and Ice Data Center Distributed Active Archive Center [data set], <https://doi.org/10.5067/CYW3O8ZUNIWC>, 2023.
- Meier, W., Petty, A. A., Hendricks, S., Kaleschke, L., Divine, D., Farrell, S., Gerland, S., Ricker, R., Tian-Kunze, X., and Webster, M.: Sea Ice, NOAA Arct. Rep. Card 2023, edited by: Moon, T., Druckenmiller, M., and Thoman, R., NOAA, <https://doi.org/10.25923/f5t4-b865>, 2023.
- Meier, W., Petty, A. A., Hendricks, S., Bliss, A., Kaleschke, L., Divine, D., Farrell, S., Gerland, S., Ricker, R., Tian-Kunze, X., and Webster, M.: Sea Ice, NOAA Arct. Rep. Card 2024, edited by: Moon, T., Druckenmiller, M., and Thoman, R., NOAA, <https://doi.org/10.25923/aksk-7p66>, 2024.
- Meredith, M., Sommerkorn, M., Cassotta, S., Derksen, C., Ekaykin, A., Hollowed, A., Kofinas, G., Mackintosh, A., Melbourne-Thomas, J., Muelbert, M. M. C., Ottersen, G., Pritchard, H., and Schuur, E. A. G.: Polar regions, in: IPCC Special Report on the Ocean and Cryosphere in a Changing Climate, edited by: Pörtner, H.-O., Roberts, D. C., Masson-Delmotte, V., Zhai, P., Tignor, M., Poloczanska, E., Mintenbeck, K., Alegría, A., Nicolai, M., Okem, A., Petzold, J., Rama, B., and Weyer, N. M., Cambridge University Press, Cambridge, UK and New York, NY, USA, 203–320, <https://doi.org/10.1017/9781009157964.005>, 2019.
- Nandan, V., Geldsetzer, T., Yackel, J., Mahmud, M., Scharien, R., Howell, S., King, J., Ricker, R., and Else, B.: Effect of Snow Salinity on CryoSat-2 Arctic First-Year Sea Ice Freeboard Measurements, *Geophys. Res. Lett.*, 44, 10419–10426, <https://doi.org/10.1002/2017GL074506>, 2017.
- Neumann, T. A., Martino, A. J., Markus, T., Bae, S., Bock, M. R., Brenner, A. C., Brunt, K. M., Cavanaugh, J., Fernandes, S. T., Hancock, D. W., Harbeck, K., Lee, J., Kurtz, N. T., Luers, P. J., Luthcke, S. B., Magruder, L., Pennington, T. A., Ramos-Izquierdo, L., Rebold, T., Skoog, J., and Thomas, T. C.: The Ice, Cloud, and Land Elevation Satellite – 2 mission: A global geolocated photon product derived from the Advanced Topographic Laser Altimeter System, *Remote Sens. Environ.*, 233, 111325, <https://doi.org/10.1016/j.rse.2019.111325>, 2019.
- Notz, D.: How well must climate models agree with observations?, *Philos. T. Roy. Soc. A*, 373, 20140164, <https://doi.org/10.1098/rsta.2014.0164>, 2015.
- Notz, D., Jahn, A., Holland, M., Hunke, E., Massonnet, F., Stroeve, J., Tremblay, B., and Vancoppenolle, M.: The CMIP6 Sea-Ice Model Intercomparison Project (SIMIP): understanding sea ice through climate-model simulations, *Geosci. Model Dev.*, 9, 3427–3446, <https://doi.org/10.5194/gmd-9-3427-2016>, 2016.
- Notz & SIMIP Community: Arctic Sea Ice in CMIP6, *Geophys. Res. Lett.*, 47, e2019GL086749, <https://doi.org/10.1029/2019GL086749>, 2020.

- Olonscheck, D. and Notz, D.: Consistently Estimating Internal Climate Variability from Climate Model Simulations, *J. Climate*, 30, 9555–9573, <https://doi.org/10.1175/JCLI-D-16-0428.1>, 2017.
- OSI SAF: Global sea ice concentration climate data record 1978–2020, EUMETSAT Ocean Sea Ice Satell. Appl. Facil., OSI-450-a, EUMETSAT [data set], https://doi.org/10.15770/EUM_SAF_OSI_0013, 2022a.
- OSI SAF: Global sea ice concentration interim climate data record, EUMETSAT Ocean Sea Ice Satell. Appl. Facil., OSI-430-a, EUMETSAT [data set], https://doi.org/10.15770/EUM_SAF_OSI_0014, 2022b.
- Parkinson, C. L.: A 40-y record reveals gradual Antarctic sea ice increases followed by decreases at rates far exceeding the rates seen in the Arctic, *P. Natl. Acad. Sci. USA*, 116, 14414–14423, <https://doi.org/10.1073/pnas.1906556116>, 2019.
- Petty, A., Kwok, R., Bagnardi, M., Ivanoff, A., Kurtz, D., Lee, J., and Wimert, J.: ATLAS/ICESat-2 L3B Daily and Monthly Gridded Sea Ice Freeboards, Version 4, NSIDC Natl. Snow Ice Data Cent. [data set], <https://doi.org/10.5067/ATLAS/ATL20.004>, 2023a.
- Petty, A. A., Hutchings, J. K., Richter-Menge, J. A., and Tschudi, M. A.: Sea ice circulation around the Beaufort Gyre: The changing role of wind forcing and the sea ice state, *J. Geophys. Res.-Oceans*, 121, 3278–3296, <https://doi.org/10.1002/2015JC010903>, 2016.
- Petty, A. A., Webster, M., Boisvert, L., and Markus, T.: The NASA Eulerian Snow on Sea Ice Model (NESOSIM) v1.0: initial model development and analysis, *Geosci. Model Dev.*, 11, 4577–4602, <https://doi.org/10.5194/gmd-11-4577-2018>, 2018.
- Petty, A. A., Kurtz, N. T., Kwok, R., Markus, T., and Neumann, T. A.: Winter Arctic Sea Ice Thickness From ICESat-2 Freeboards, *J. Geophys. Res.-Oceans*, 125, e2019JC015764, <https://doi.org/10.1029/2019JC015764>, 2020.
- Petty, A. A., Kurtz, N., Kwok, R., Markus, T., Neumann, T. A., and Keeney, N.: ICESat-2 L4 Monthly Gridded Sea Ice Thickness, Version 3, NASA Natl. Snow Ice Data Cent. Distrib. Act. Arch. Cent. [data set], <https://doi.org/10.5067/ZCSU8Y5U1BQW>, 2023b.
- Petty, A. A., Keeney, N., Cabaj, A., Kushner, P., and Bagnardi, M.: Winter Arctic sea ice thickness from ICESat-2: upgrades to freeboard and snow loading estimates and an assessment of the first three winters of data collection, *The Cryosphere*, 17, 127–156, <https://doi.org/10.5194/tc-17-127-2023>, 2023c.
- Pithan, F. and Mauritsen, T.: Arctic amplification dominated by temperature feedbacks in contemporary climate models, *Nat. Geosci.*, 7, 181–184, <https://doi.org/10.1038/ngeo2071>, 2014.
- Riedel, C. and Anderson, J.: Exploring non-Gaussian sea ice characteristics via observing system simulation experiments, *The Cryosphere*, 18, 2875–2896, <https://doi.org/10.5194/tc-18-2875-2024>, 2024.
- Roach, L. A., Dörr, J., Holmes, C. R., Massonnet, F., Blockley, E. W., Notz, D., Rackow, T., Raphael, M. N., O’Farrell, S. P., Bailey, D. A., and Bitz, C. M.: Antarctic Sea Ice Area in CMIP6, *Geophys. Res. Lett.*, 47, e2019GL086729, <https://doi.org/10.1029/2019GL086729>, 2020.
- Salganik, E., Crabeck, O., Fuchs, N., Hutter, N., Anhaus, P., and Landy, J. C.: Impacts of air fraction increase on Arctic sea ice density, freeboard, and thickness estimation during the melt season, *The Cryosphere*, 19, 1259–1278, <https://doi.org/10.5194/tc-19-1259-2025>, 2025.
- Santer, B. D., Thorne, P. W., Haimberger, L., Taylor, K. E., Wigley, T. M. L., Lanzante, J. R., Solomon, S., Free, M., Gleckler, P. J., Jones, P. D., Karl, T. R., Klein, S. A., Mears, C., Nychka, D., Schmidt, G. A., Sherwood, S. C., and Wentz, F. J.: Consistency of modelled and observed temperature trends in the tropical troposphere, *Int. J. Climatol.*, 28, 1703–1722, <https://doi.org/10.1002/joc.1756>, 2008.
- Schaller, N., Mahlstein, I., Cermak, J., and Knutti, R.: Analyzing precipitation projections: A comparison of different approaches to climate model evaluation, *J. Geophys. Res.-Atmos.*, 116, D10118, <https://doi.org/10.1029/2010JD014963>, 2011.
- Schweiger, A. J., Steele, M., Zhang, J., Moore, G. W. K., and Laidre, K. L.: Accelerated sea ice loss in the Wandel Sea points to a change in the Arctic’s Last Ice Area, *Commun. Earth Environ.*, 2, 1–11, <https://doi.org/10.1038/s43247-021-00197-5>, 2021.
- Semtner, A. J.: A Model for the Thermodynamic Growth of Sea Ice in Numerical Investigations of Climate, *J. Phys. Oceanogr.*, 6, 379–389, 1976.
- Shi, H., Lee, S.-M., Sohn, B.-J., Gasiewski, A. J., Meier, W. N., Dybkjær, G., and Kim, S.-W.: Estimation of Arctic Winter Snow Depth, Sea Ice Thickness and Bulk Density, and Ice Freeboard by Combining CryoSat-2, AVHRR, and AMSR Measurements, *IEEE T. Geosci. Remote*, 61, 1–18, <https://doi.org/10.1109/TGRS.2023.3265274>, 2023.
- Smith, D. M., Screen, J. A., Deser, C., Cohen, J., Fyfe, J. C., García-Serrano, J., Jung, T., Kattsov, V., Matei, D., Msadek, R., Peings, Y., Sigmond, M., Ukita, J., Yoon, J.-H., and Zhang, X.: The Polar Amplification Model Intercomparison Project (PAMIP) contribution to CMIP6: investigating the causes and consequences of polar amplification, *Geosci. Model Dev.*, 12, 1139–1164, <https://doi.org/10.5194/gmd-12-1139-2019>, 2019.
- Thackeray, C. W. and Hall, A.: An emergent constraint on future Arctic sea-ice albedo feedback, *Nat. Clim. Change*, 9, 972–978, <https://doi.org/10.1038/s41558-019-0619-1>, 2019.
- Tilling, R., Kurtz, N. T., Bagnardi, M., Petty, A. A., and Kwok, R.: Detection of Melt Ponds on Arctic Summer Sea Ice From ICESat-2, *Geophys. Res. Lett.*, 47, e2020GL090644, <https://doi.org/10.1029/2020GL090644>, 2020.
- Topál, D. and Ding, Q.: Atmospheric circulation-constrained model sensitivity recalibrates Arctic climate projections, *Nat. Clim. Change*, 13, 710–718, <https://doi.org/10.1038/s41558-023-01698-1>, 2023.
- Tsamados, M., Feltham, D. L., Schroeder, D., Flocco, D., Farrell, S. L., Kurtz, N., Laxon, S. W., and Bacon, S.: Impact of variable atmospheric and oceanic form drag on simulations of Arctic sea ice, *J. Phys. Oceanogr.*, 44, 1329–1353, <https://doi.org/10.1175/JPO-D-13-0215.1>, 2014.
- Wang, B., Zhou, X., Ding, Q., and Liu, J.: Increasing confidence in projecting the Arctic ice-free year with emergent constraints, *Environ. Res. Lett.*, 16, 094016, <https://doi.org/10.1088/1748-9326/ac0b17>, 2021.
- Watts, M., Maslowski, W., Lee, Y. J., Kinney, J. C., and Osinski, R.: A Spatial Evaluation of Arctic Sea Ice and Regional Limitations in CMIP6 Historical Simulations, *J. Climate*, 34, 6399–6420, <https://doi.org/10.1175/JCLI-D-20-0491.1>, 2021.
- Wieringa, M. M., Riedel, C., Anderson, J. L., and Bitz, C. M.: Bounded and categorized: targeting data assimilation for sea

- ice fractional coverage and nonnegative quantities in a single-column multi-category sea ice model, *The Cryosphere*, 18, 5365–5382, <https://doi.org/10.5194/tc-18-5365-2024>, 2024.
- Willatt, R., Laxon, S., Giles, K., Cullen, R., Haas, C., and Helm, V.: Ku-band radar penetration into snow cover on Arctic sea ice using airborne data, *Ann. Glaciol.*, 52, 197–205, <https://doi.org/10.3189/172756411795931589>, 2011.
- Williams, N., Byrne, N., Feltham, D., Van Leeuwen, P. J., Bannister, R., Schroeder, D., Ridout, A., and Nerger, L.: The effects of assimilating a sub-grid-scale sea ice thickness distribution in a new Arctic sea ice data assimilation system, *The Cryosphere*, 17, 2509–2532, <https://doi.org/10.5194/tc-17-2509-2023>, 2023.
- Worby, A. P., Geiger, C. A., Paget, M. J., Van Woert, M. L., Ackley, S. F., and DeLiberty, T. L.: Thickness distribution of Antarctic sea ice, *J. Geophys. Res.*, 113, C05S92, <https://doi.org/10.1029/2007JC004254>, 2008.
- Zanowski, H., Jahn, A., and Holland, M. M.: Arctic Ocean Freshwater in CMIP6 Ensembles: Declining Sea Ice, Increasing Ocean Storage and Export, *J. Geophys. Res.-Oceans*, 126, e2020JC016930, <https://doi.org/10.1029/2020JC016930>, 2021.
- Zhang, J., Rothrock, D., and Steele, M.: Recent Changes in Arctic Sea Ice: The Interplay between Ice Dynamics and Thermodynamics, *Journal of Climate*, 13, 3099–3114, 2000.
- Zhou, Y., Wang, X., Lei, R., von Albedyll, L., Perovich, D. K., Zhang, Y., and Haas, C.: Seasonal evolution and parameterization of Arctic sea ice bulk density: results from the MO-SAiC expedition and ICESat-2/ATLAS, *EGUsphere* [preprint], <https://doi.org/10.5194/egusphere-2024-1240>, 2024.
- Zhuang, J., dussin, raphael, Huard, D., Bourgault, P., Banihirwe, A., Raynaud, S., Malevich, B., Schupfner, M., Filipe, Gauthier, C., Levang, S., Jüling, A., Almansi, M., RichardScottOZ, RondeauG, Rasp, S., Smith, T. J., Mares, B., Stachelek, J., Plough, M., Pierre, Bell, R., Caneill, R., and Li, X.: pangeo-data/xESMF: v0.8.8, Zenodo [code], <https://doi.org/10.5281/zenodo.14025505>, 2024.
- Zwally, H. J., Schutz, B., Abdalati, W., Abshire, J., Bentley, C., Brenner, A., Bufton, J., Dezio, J., Hancock, D., Harding, D., Herring, T., Minster, B., Quinn, K., Palm, S., Spinhirne, J., and Thomas, R.: ICESat's laser measurements of polar ice, atmosphere, ocean, and land, *J. Geodyn.*, 34, 405–445, [https://doi.org/10.1016/S0264-3707\(02\)00042-X](https://doi.org/10.1016/S0264-3707(02)00042-X), 2002.