



Stochastic homogeneous freezing of supercooled droplets in mixed-phase clouds in particle-based idealised microphysics framework (exemplified with PySDM v3.0)

Tim Lüttmer¹, Sylwester Arabas², and Peter Spichtinger¹

¹Institute for Atmospheric Physics, Johannes Gutenberg University Mainz, Mainz, Germany

²Faculty of Physics and Applied Computer Science, AGH University of Krakow, Kraków, Poland

Correspondence: Tim Lüttmer (tluettm@uni-mainz.de)

Received: 17 April 2026 – Discussion started: 27 April 2026

Revised: 31 August 2026 – Accepted: 8 September 2026 – Published: 27 October 2026

Abstract. Homogeneous freezing of supercooled cloud droplets is an important process governing ice formation during the transition from mixed-phase clouds to pure ice-phase clouds in the upper troposphere. In this study, we implement a stochastic representation of homogeneous freezing in the particle-based aerosol–cloud microphysics model PySDM, treating freezing as a Poisson process dependent on droplet volume, time step, and the nucleation rate. We compare two parameterisations of the nucleation rate: a temperature-dependent formulation and a saturation-dependent formulation. Using an idealised adiabatically ascending air-parcel framework, we investigate the distribution of freezing temperatures and the resulting ice number concentrations across ensembles that vary updraft speed, cloud condensation nuclei number concentration, droplet size distribution, and the number of super-particles. Simulations are performed both with and without vapour deposition on ice, enabling assessment of the role of the Wegener-Bergeron-Findeisen process. We find that homogeneous freezing occurs over a broad temperature range rather than at a single threshold, with freezing temperatures strongly controlled by cooling rate and droplet size. When vapour deposition on ice is active, early stochastic freezing events dominate the evolution of the frozen droplet fraction and substantially reduce the fraction of droplets that ultimately freeze. The two nucleation-rate formulations produce similar behaviour near water saturation but diverge significantly when supersaturation or subsaturation with respect to water develops, leading to pronounced differences in freezing temperatures and ice number concentrations. Our results highlight the importance of stochastic freezing for-

mulations and nucleation-rate choice for representing cloud glaciation in models.

1 Introduction

Pure ice-phase clouds at temperatures below $T_{\text{hom}} \sim 235$ K, commonly referred to as cirrus clouds, consist only of ice crystals and snow aggregates. T_{hom} denotes the temperature threshold of homogeneous freezing, below which we can assume that all liquid water droplets of substantial size (e.g., in the μm size range) are frozen. Hence, it separates the pure-ice phase from the mixed-phase thermodynamic regime, where the liquid and ice phase can coexist. Cirrus clouds frequently form in the upper-level outflow of vertically developing dynamical systems such as deep convection and Warm Conveyor Belts (WCBs). Two main formation pathways for cirrus can be distinguished (Krämer et al., 2016; Luebke et al., 2016). In the liquid-origin pathway, cirrus forms from the freezing of supercooled water droplets, hence they stem from glaciation of mixed-phase clouds that were subsequently transported to lower temperatures ($\leq T_{\text{hom}}$). In contrast, in-situ cirrus forms directly from the vapour phase, usually in a thermodynamic regime where (pure) water droplets are already frozen. Observational studies suggest that cirrus formed via the liquid-origin pathway typically exhibit higher ice mass mixing ratios and larger mean ice crystal sizes than cirrus formed via the in-situ-origin pathway (Krämer et al., 2020). These microphysical differences can substantially influence the radiative properties of cirrus clouds and therefore

determine whether their net radiative effect is warming or cooling (Zhang et al., 1999; Fusina et al., 2007; Joos et al., 2014). In addition, they alter the particle size distribution which affects aggregation and sedimentation rates, which can affect precipitation and influence lower-level clouds. In the anvils of deep convective clouds, cirrus forms predominantly through the liquid-origin pathway, as mixed-phase processes dominate in the convective core and are subsequently transported upward into the thermodynamic regime of pure ice-phase clouds (Gasparini et al., 2018; Lüttmer et al., 2025b). In contrast, cirrus clouds in the outflow of WCBs can originate from both liquid-origin and in-situ formation pathways (Krämer et al., 2020). In some cases, the resulting cirrus exhibits a mixed origin, where ice formed through the glaciation of mixed-phase clouds during the WCB ascent coexists with ice produced in-situ above the WCB that subsequently sediments into the ascending air mass (Lüttmer et al., 2025a). Understanding the liquid-origin pathway is essential for characterizing the properties of cirrus clouds in the outflow of WCBs and convective systems. The freezing of supercooled water droplets generally occurs via two nucleation modes: homogeneous nucleation, in which droplets freeze without the presence of foreign substances aiding the nucleation process, and heterogeneous nucleation, in which freezing is initiated by foreign substances, referred to as ice-nucleating particles (INPs), that reduce the temperature and supersaturation required for ice nucleation as compared to homogeneous nucleation (Vali et al., 2015). Heterogeneous freezing plays a key role in mixed-phase cloud microphysics, as it governs the initial formation of ice. Considerable effort has therefore been devoted to improving the representation of INPs in atmospheric models. This includes characterizing their freezing properties (e.g., as a function of temperature) (Phillips et al., 2008; Ullrich et al., 2017), constraining their spatial and temporal variability through observations and climatologies (Hande et al., 2015), and developing physically based aerosol–cloud interaction frameworks (Hoshyari-pour et al., 2026; Costa-Surós et al., 2026). In addition, different conceptual approaches are used to represent INP activation, e.g. “singular” and “time-dependent” parameterizations (Alpert and Knopf, 2016; Arabas et al., 2025). In contrast, homogeneous freezing of micrometre-sized supercooled water droplets has received comparatively less attention in cloud modeling studies. While numerous laboratory experiments provide estimates of homogeneous nucleation rates (see, e.g., Murray et al., 2010; Ickes et al., 2015; Koop and Murray, 2016), their representation in atmospheric models is often simplified. A common approach is to treat homogeneous freezing as a threshold process, whereby all micrometre-sized droplets (“cloud droplets”) instantaneously freeze once temperatures fall below a prescribed threshold T_{hom} . This formulation is widely used in large-scale and cloud-resolving models, such as EC-Earth3-AerChem (Costa-Surós et al., 2026), SCALE-SDM (Shima et al., 2020), IFS (ECMWF, 2023), and the P3 scheme used in WRF (Morrison and Mil-

brandt, 2015). Typical values for T_{hom} are 235 K or -38°C . More advanced approaches attempt to account for the competition between homogeneous ice freezing and droplet evaporation. For example, the microphysics scheme in the Unified Model allows the number of homogeneously frozen droplets to be limited by updraft velocity and environmental conditions, thereby representing the depletion of supersaturation due to vapour deposition onto newly formed ice crystals (“supersaturation quenching”) (Field et al., 2023). In contrast, the microphysics schemes used in ICON and ICON-ART explicitly integrate the homogeneous nucleation rate coefficient J_{hom} over the cloud droplet size distribution, providing a more physically consistent representation of the freezing process (Seifert and Beheng, 2006; Welss et al., 2024; Hoshyari-pour et al., 2026). In this study, we represent the homogeneous freezing of micrometre-sized water droplets as a stochastic process, using classical nucleation theory (Ickes et al., 2015). The super-particle framework is well suited for a physically consistent approach by describing homogeneous freezing as a Poisson process that depends on droplet volume, time step, and the homogeneous nucleation rate. We employ the Python super-particle model PySDM, an open-source Python 3 package designed to simulate populations of super-particles representing atmospheric hydrometeors (Bartman et al., 2022; de Jong et al., 2023). Section 2 introduces the model extensions implemented in this study, namely homogeneous freezing and vapour deposition on ice crystals. In Sect. 3, we present results from idealised simulations of freezing in an adiabatically ascending air parcel. We investigate the sensitivity to updraft velocity, the number of super-particles, and the cloud condensation nuclei (CCN) concentration. In addition, we examine differences arising from formulations of the homogeneous nucleation rate coefficient J_{hom} based on either temperature or water activity. Finally, we highlight the interplay between homogeneous freezing and droplet evaporation driven by the Wegener–Bergeron–Findeisen process.

2 Methods

Hydrometeors (liquid water droplets and ice crystals) are represented as super-particles. Each super-particle represents a number of real particles specified by its multiplicity ξ . In addition to ξ , each super-particle is described by the following attributes: phase state (liquid l or ice i), mass m , radius r , volume V , hygroscopicity parameter κ , and terminal velocity v_t , respectively. The prognostic attributes are the multiplicity ξ , mass m , and phase state. All other attributes are diagnostic, with the exception of κ , which is set to a fixed value (see Sect. 2.2). Attributes of (liquid) droplets are denoted by the subscript l and attributes of ice crystals by the subscript i . Droplets are assumed to be spherical, whereas ice crystals are columnar in shape. All hydrometeors are initially represented as cloud condensation nuclei (CCN) with a dry radius

r_d . Super-particles are sampled uniformly from an analytical size distribution of CCNs. The number of super-particles used for sampling, n_{sd} , determines the accuracy of the simulation.

2.1 Homogeneous freezing

Homogeneous freezing of supercooled liquid droplets can be described using classical nucleation theory (for detailed treatments, see, e.g., Pruppacher et al., 2010; Lamb and Verlinde, 2011; Ickes et al., 2015; Koop and Murray, 2016). Here, we restrict our discussion to the temperature range relevant for ice formation in the troposphere, below the melting temperature of water $T < T_m = 273.13$ K. In a supercooled liquid droplet ($T < T_m$), thermal molecular motion can lead to the spontaneous formation of molecular clusters that serve as ice embryos. Although the ice phase is thermodynamically favored below T_m , nucleation is impeded by a Gibbs free-energy barrier. Formation of a critical embryo is required before further molecular attachment to the crystalline lattice becomes energetically favorable. Once such a critical cluster forms, macroscopic freezing is initiated and the entire droplet freezes. The Gibbs free-energy barrier arises from (i) the change in chemical potential between the liquid and solid phases (a volume term), and (ii) the interfacial free energy associated with the liquid-ice boundary (a surface-area term). On a kinetic level, molecules continuously attach to and detach from the solid embryo, a process that can be described as a diffusive molecular flux. Both the thermodynamic and kinetic effects determine the homogeneous nucleation rate $J_{hom} [\text{m}^{-3} \text{s}^{-1}]$. This rate represents the number of critical ice embryos formed per unit droplet volume and unit time. In general, J_{hom} depends only on the droplet temperature T_d , under the assumption that the droplet is in thermodynamic equilibrium with the ambient air such that $T_d = T$ and the droplet is pure, hence there are no foreign substances present that affect the freezing behaviour. Because the ice embryo forms inside the liquid, it has no interface with the ambient environment, and thus the formulation of J_{hom} should not depend directly on the ambient vapour pressure or saturation ratio. However, as discussed below, the saturation ratio becomes relevant in the parametrisation of homogeneous nucleation for aqueous solution droplets. The value of J_{hom} is sensitive to the parametrisation of the diffusion activation energy, the liquid–ice interfacial energy, and the saturation ratios (Koop and Murray, 2016). Theoretical constraints on these dependencies are limited in the temperature regime relevant for homogeneous nucleation, and they are therefore obtained through fits to nucleation rates measured in laboratory experiments. Significant uncertainties remain, both in the derivation of J_{hom} and in the applicability of classical nucleation theory itself (Ickes et al., 2015). A detailed discussion of these uncertainties is beyond the scope of this study, but they should be kept in mind when evaluating the impact of the choice of J_{hom} parameterization. We define P_{fz} as the

probability of homogeneous freezing of a droplet. First, we formulate this probability as a simple temperature dependent threshold process $P_{fz,thr}$

$$P_{fz,thr}(T) = \begin{cases} 1 & \text{for } T \leq T_{hom} \\ 0 & \text{for } T > T_{hom} \end{cases} \quad (1)$$

where the common choice for $T_{hom} = 235$ K which corresponds to $J_{hom} \sim 10^{15} \text{ m}^{-3} \text{ s}^{-1}$ (see Fig. 1). A more physical approach is to parameterize homogeneous freezing as a Poisson counting process (see, e.g., Bigg, 1953; Pruppacher et al., 2010). Here P_{fz} is the probability that at least one ice embryo of critical size forms inside a droplet of volume V_l within the time interval Δt and thus initiates homogeneous freezing. Assuming that the process rate $r_{proc} = J_{hom} V_l$ remains constant over Δt , the droplet-freezing probability is

$$P_{fz}(J_{hom}, V_l, \Delta t) = 1 - \exp(-J_{hom} V_l \Delta t). \quad (2)$$

P_{fz} therefore depends on the droplet volume ($V_l = 4\pi r_l^3/3$), the thermodynamic state variables that control J_{hom} (primarily T), and the elapsed time t . An important consequence of this time dependence is that even small droplets, or droplets in thermodynamic states with low values of J_{hom} , will eventually freeze if sufficient time is available. The same framework has also been applied to heterogeneous freezing, where the droplet volume is replaced by the surface area of the immersed ice-nucleating particle (Arabas et al., 2025). In the model, freezing is treated using a Monte Carlo approach. At each time step, a uniform random number $x_i \sim \mathcal{U}(0, 1)$ is generated for each super-particle i . Freezing occurs if $x_i < P_{fz}(V_l, J_{hom}, \Delta t)$. Because each super-particle represents ξ real droplets, once freezing is initiated, all real droplets represented by that super-particle freeze. Statistical convergence thus depends not only on the physical time step Δt but also on the sampling of the droplet spectrum – specifically, the number of super-particles n_{sd} and sampling method (e.g., constant multiplicity, uniform in size) used to discretise the size distribution. The formulation of the homogeneous nucleation rate J_{hom} is critical for representing homogeneous freezing in numerical models. The study by Koop et al. (2000) (hereafter KP00) is a foundational contribution in this regard. Solutes inhibit the nucleation ability of droplets, allowing aqueous solution droplets to be supercooled to lower temperatures than pure water droplets before freezing becomes likely. The freezing of submicrometre-sized aqueous solution droplets plays a key role in cirrus formation. KP00 found empirically that J_{hom} for aqueous solution droplets depends solely on thermodynamic state variables and is independent of the type of solute. Moreover, assuming that the droplet is in thermodynamic equilibrium with the ambient environment, J_{hom} can be expressed by shifting the melting curve of water by an offset in water activity, $\Delta a_w = (S_i - 1) \frac{e_{sat,i}}{e_{sat,l}}$, where $e_{sat,i}$ and $e_{sat,l}$ are the saturation vapour pressures over ice and (pure) water, respectively. This formulation extends the dependence of J_{hom} from temperature T

alone to include the saturation ratio with respect to ice S_i . The KP00 rate is formulated as a third-order polynomial in Δa_w . The study of Koop and Murray (2016) (hereafter KM16) revisited the parameterisation of J_{hom} for micrometre-sized pure water droplets. They used new methods to constrain the relevant thermodynamic and kinetic properties controlling J_{hom} and used a selected dataset of nucleation-rate measurements to fit J_{hom} . KM16 use a sixth-order polynomial in T as a fit

$$\log_{10}(J_{\text{hom}})|_{\text{JHOM-T}} = \sum_{k=0}^6 c_k (T - T_m)^k \quad (3)$$

with the melting temperature of ice at standard pressure $T_m = 273.15$ K. The coefficients c_k can be found in Koop and Murray (2016), Table 7. This parameterisation is valid for a temperature regime $245 \leq T \leq 227$ K. The label JHOM-T denotes that J_{hom} depends only on T .

Figure 1a shows three formulations of J_{hom} at water saturation $S_w = 1$. Here we focus first on the parameterisation for JHOM-T (black line), which depends only on T . We note that J_{hom} increases from 240 to 235 K by over 5 orders of magnitude highlighting the sharp increase of J_{hom} with decreasing temperature. Over the same temperature range, the earlier parameterisation of KP00 (blue line), which depends on Δa_w and thus on the saturation ratios (S_w or S_i), follows the same slope as JHOM-T for $S_w = 1$ but shows an offset of about 1.5 orders of magnitude. The slope of the JHOM-T rate diverges from KP00 at $T < 234$ K. While it remains unclear which behaviour is correct, the temperature range below $T = 235$ K is rarely relevant for the homogeneous freezing events investigated in this study (Spichtinger et al., 2023). This highlights the highly uncertain nature of J_{hom} . The JHOM-T parameterisation of KM16 is the latest fit to describe homogeneous freezing of pure water droplets and should therefore be used instead of the earlier KP00 parameterisation in that case. However, from a modeling viewpoint it is beneficial to have a unified, Δa_w -based parameterisation that is valid for both submicrometre-sized aqueous solution droplets (in the pure-ice regime) and micrometre-sized pure water droplets (in the mixed-phase regime). The study of Spichtinger et al. (2023) (hereafter SP23) provides such an option. They reformulated the T -based parameterisation of KM16 as a function of Δa_w

$$\log_{10}(J_{\text{hom}})|_{\text{JHOM-DWA}} = \sum_{k=0}^3 b_k \Delta a_w^k \quad (4)$$

where the b_k coefficients can be found in Spichtinger et al. (2023). The label JHOM-DWA denotes that J_{hom} depends on Δa_w . The black line in Fig. 1a actually shows J_{hom} not only for JHOM-T but also for the JHOM-DWA formulation at $S_w = 1$ as they coincide by construction. So, in principle, the Δa_w -based rate of JHOM-DWA should be valid for both thermodynamic regimes – near water saturation ($S_w \approx$

1) for the homogeneous freezing of micrometre-sized pure droplets, and below water saturation ($S_w < 1$) for the homogeneous freezing of submicrometre-sized aqueous solution droplets. However, at temperatures close to $T = 235$ K Δa_w varies strongly with small changes in S_w . Figure 1b shows J_{hom} for JHOM-DWA at water-saturation ratios ranging from 0.95 to 1.04 (red curves) together with the S_w -independent rate of JHOM-T (black). Here we note that even for small supersaturation or subsaturation, J_{hom} differs strongly from the base JHOM-T rate. As previously discussed, J_{hom} should in general be independent of the vapour pressure in the ambient air (as the ice embryo has no interface with it), at least for the case of pure-water droplets. However, cloud droplets in the atmosphere generally contain foreign substances, e.g., the CCN responsible for droplet activation. In this model, we also assume a soluble CCN. Hence, we should expect some effect on the freezing behaviour from these substances. However, the magnitude of the effect is uncertain, especially since the concentration of foreign substances is only substantial if the droplet is unactivated and thus in the submicrometre size range. Hence, JHOM-T would be the better choice for micrometre-sized cloud droplets, even if the ambient air is super- or subsaturated with respect to water. Note that JHOM-DWA and JHOM-T coincide, as long as water saturation is maintained. In many models featuring bulk schemes, it is a basic assumption that $S_w = 1$ for as long as cloud droplets are present (using saturation adjustment, see Seifert and Beheng, 2006; ECMWF, 2023; Field et al., 2023). However, we will later see that this assumption is often violated during homogeneous freezing events.

2.2 Particle growth by vapour deposition

The growth of droplets by water vapour deposition (condensation) and evaporation is modeled here by κ -Köhler theory (Petters and Kreidenweis, 2007). The hygroscopicity parameter κ is defined by the relation of the water activity a_w to the volume ratio of the solute v_s and water V_l

$$\frac{1}{a_w} = 1 + \kappa \frac{v_s}{v_l} \quad (5)$$

The implementation of condensation in PySDM is described following the formulation in libcloudph++ (Arabas et al., 2015). Following Shima et al. (2020), vapour deposition on ice is implemented in a separate integration step from condensation evaporation. We obtain the mass rate for the growth and sublimation of ice crystals by deposition of water vapour as

$$\frac{dm_i}{dt} = 4\pi C D_v^* G_v (S_i - 1) \quad (6)$$

The capacitance C in m accounts for the non-spherical nature of the ice crystals. Here we use prolate ellipsoids as approximation for columnar ice crystals fitted as in Spichtinger et al. (2023) with the (ice crystal) mass m_i in kg




Figure 1. Homogeneous nucleation rates at (a) water saturation ($S_w = 1$) for the JHOM-T, JHOM-DWA, and KP00 formulations, and (b) for JHOM-DWA at various water saturation ratios, where the rate from JHOM-T is shown as a reference.

$$C = 0.015755 m_i^{0.3} + 0.33565 m_i^{0.43} \quad (7)$$

The diffusion of vapour onto the ice surface is a gas kinetic process. For large ice crystals ($> 1 \mu\text{m}$) we can treat the diffusion of vapour in the continuous regime (Spichtinger and Gierens, 2009; Skrotzki et al., 2013). There the diffusion constant D_v only depends on temperature and pressure (Pruppacher et al., 2010). For larger ice crystals ($> 100 \mu\text{m}$) a ventilation correction must be applied, whereas for smaller ice crystals ($< 1 \mu\text{m}$) a correction for the kinetic regime is required. The latter one is especially relevant in this work, as we investigate the freezing of droplets in the micrometre size range, where kinetic effects dominate. The correction factor for ventilation can be neglected for the ice crystal sizes considered in this study. D_v^* is the diffusion coefficient of water vapour D_v corrected for the kinetic regime

$$D_v^* = \frac{D_v}{\frac{r_i}{r_i + \lambda_a} + \frac{4D_v}{\alpha_d v_{m,v} r_i}} \quad (8)$$

with the equivalent radius of the ice crystal r_i , the mean free path of air λ_a and the mean thermal velocity of water molecules $v_{m,v} = \sqrt{\frac{8R_v T}{\pi}}$. Not every water molecule that impacts the surface of an ice crystal is integrated into the crystal structure. The ratio of colliding to integrated molecules is given by the mass accommodation coefficient α_d . The value of α_d is highly uncertain and might depend on the ice crystal size and temperature (Skrotzki et al., 2013). Here we set $\alpha_d = 0.5$ as in other studies that investigate homogeneous nucleation and which results are compared to this study (Kärcher and Lohmann, 2002; Spichtinger and Gierens, 2009). The phase change of vapour to ice at the surface of the ice crystals results in latent heating. Hence, the surface temperature of the ice crystals is warmer than the ambient air. This changes the local supersaturation experienced by the ice crystal. The growth rate Eq. (6) needs to account for temperature difference between ice crystal and ambient air and the thermal diffusion. This is modeled here by the Howell-factor G_v (Howell, 1949)

$$G_v = \left[\left(\frac{L_s}{R_v T} - 1 \right) \frac{L_s D_v^*}{T K_T^*} + \frac{R_v T}{e_{s,i}} \right]^{-1} \quad (9)$$

with the latent heat of sublimation L_s , the specific gas constant of water vapour R_v and the thermal conductivity of moist air K_T . Similar as for D_v we apply a correction for the kinetic regime to K_T

$$K_T^* = \frac{K_T}{\frac{r_i}{r_i + \lambda_a} + \frac{4K_T}{\alpha_T c_p \rho v_{m,a} r_i}} \quad (10)$$

with the specific heat of dry air at constant pressure c_p , and the mean thermal velocity of air $v_{m,a} = \sqrt{\frac{8R_d T}{\pi}}$. The thermal accommodation coefficient α_T describes the ratio of colliding molecules that attain a thermodynamic equilibrium with the ice crystals during contact. Here we set $\alpha_T = 1$ (Spichtinger and Gierens, 2009). The change of water vapour mixing ratio due to deposition onto ice crystals and evaporation of ice crystals is expressed as

$$\frac{\partial q_v}{\partial t} \Big|_{\text{dep}} = \sum_{k=0}^N \frac{dm_{i,k}}{dt} \frac{\xi_k}{m_{\text{dry}}} \quad (11)$$

where we iterate over all super particles of the ice phase with their mass rate $\frac{dm_{i,k}}{dt}$ and multiplicity ξ_k with the mass of dry air m_{dry} .

2.3 Adiabatic parcel model

We use an adiabatic parcel model to represent an air parcel rising with a prescribed constant velocity w . The thermodynamic state of the air parcel is described by the air pressure p , air temperature T and water vapour mixing ratio q_v . For the (adiabatic) change of pressure we assume hydrostatic balance

$$\frac{dp}{dt} = -g\rho w \quad (12)$$

with the gravitational acceleration g and dry air density ρ . The temperature changes due to adiabatic expansion during

vertical motion and latent heat of vaporization L_v and latent heat of sublimation L_s

$$\frac{dT}{dt} = -\frac{g}{c_p}w - \frac{L_v}{c_p}\frac{\partial q_v}{\partial t}|_{\text{con}} - \frac{L_s}{c_p}\frac{\partial q_v}{\partial t}|_{\text{dep}} \quad (13)$$

with the specific heat of dry air c_p at constant p . Note that we neglect latent heat of freezing of droplets. The change in vapour mass mixing ratio q_v is controlled by phase change due to condensation/evaporation and vapour deposition/sublimation

$$\frac{dq_v}{dt} = \frac{\partial q_v}{\partial t}|_{\text{con}} + \frac{\partial q_v}{\partial t}|_{\text{dep}} \quad (14)$$

The saturation ratios with respect to ice S_i and with respect to water S_w are diagnostic quantities dependent on T , p and q_v

$$S_i = \frac{pq_v}{\epsilon e_{s,i}(T)} S_w = \frac{pq_v}{\epsilon e_{s,w}(T)} \quad (15)$$

with $\epsilon = \frac{R_v}{R_a}$ and the saturation pressure $e_{s,i}$ and $e_{s,w}$ of ice and water, respectively. S_i and S_w control the growth of frozen and liquid super-particles, respectively, and they control J_{hom} for the Δa_w formulation.

2.4 Model setup

All simulations use the simple parcel-model setup described in Sect. 2.3. Unless otherwise specified (e.g., in sensitivity simulations), we use $w = 1 \text{ m s}^{-1}$. The thermodynamic initial conditions for all simulations are temperature $T_0 = 245 \text{ K}$, pressure $p_0 = 500 \text{ hPa}$, and saturation ratio with respect to water $S_w = 0.995$. Super-particles in the considered volume represent particles which are initially unactivated and liquid phase (i.e., CCN), upon their growth and freezing they become cloud droplets or ice crystals. We prescribe a monodisperse CCN size spectrum with a dry aerosol radius of 15 nm and a hygroscopicity parameter $\kappa = 0.65$. Note that these parameter choices affect the simulation outcome only in a minor way. The use of a monodisperse size spectrum ensures that all CCN activate, after which droplet sizes equilibrate to the ambient environment, which is determined by the adiabatic cooling rate (and thus the updraft speed w). Because the simulations start at $T_0 = 245 \text{ K}$, all droplets have sufficient time to activate and equilibrate before thermodynamic conditions favorable for homogeneous freezing are reached. In Sect. 3.3 we will investigate polydisperse CCN size spectra using log-normal distribution with the geometric standard deviation σ as an ensemble parameter. Unless otherwise specified (e.g., in sensitivity simulations), we use a CCN number concentration of $n_{\text{ccn}} = 750 \text{ cm}^{-3}$ at STP (standard temperature and pressure), representative of an average continental air mass (Jurányi et al., 2010; Choudhury and Tesche, 2023). Note, that the local CCN concentration used as input for the simulation is thus lower than the stated CCN

concentration at STP as it has to be scaled by $\rho_{\text{local}}/\rho_{\text{STP}}$. Condensation, homogeneous freezing, and ambient thermodynamics are active for all simulations, whereas ice growth by vapour deposition (DEP-ICE) is only enabled when explicitly stated. The model time step is linked to w such that the parcel's vertical displacement per time step, Δz , is constant. Hence, simulations with stronger updraft speeds employ shorter time steps than those with weaker updrafts. For simulations without DEP-ICE, $\Delta z = 2 \text{ m}$ is sufficient; for simulations with DEP-ICE, we use $\Delta z = 1 \text{ m}$. For simulations with enhanced n_{ccn} and DEP-ICE, $\Delta z = 0.5 \text{ m}$ is used. The simulation is stopped when all super-particles are frozen or evaporated. In this study we investigate the parameter space spanned by these variables: w , n_{ccn} , and n_{sd} . For each of these parameters, ensemble simulations are performed with a choice between the two J_{hom} parametrizations – the T -dependent JHOM-T formulation and the Δa_w -dependent JHOM-DWA formulation – and a choice between DEP-ICE inactive (Sect. 3.1) or active (Sect. 3.2). Additionally, we investigate the impact of choosing a lognormal CCN size distribution instead of a monodisperse distribution, with an ensemble over the width of the distribution, σ , in Sect. 3.3.

3 Results

3.1 Simulations without vapour deposition on ice

First, we investigate the setup where only ambient thermodynamics, condensation, and homogeneous freezing are active. Here, frozen super-particles act as passive tracers, since without vapour deposition on ice (DEP-ICE) or collisions, they do not interact with other super-particles or the ambient air. Without DEP-ICE, the ambient air never becomes subsaturated with respect to water ($S_w < 1$). As a consequence, all droplets eventually freeze, as there is no removal process. Thus, for the bulk quantities – droplet number concentration n_c , CCN number concentration n_{ccn} , and ice number concentration n_i – the following relationship holds: $n_{\text{ccn}} = n_c(t < t_f) = n_i(t_{\text{end}})$, where t_f is the time of the first freezing event, and t_{end} is the end of the simulation when all droplets are frozen. Number concentrations are expressed as number per kg of dry air. Figure 2 shows a sample air parcel ascent for the JHOM-T rate (Fig. 2a–d) and JHOM-DWA rate (Fig. 2e–h) with $n_{\text{sd}} = 1000$ and $w = 2.5 \text{ m s}^{-1}$. We show only a single simulation here, as $n_{\text{sd}} = 1000$ is sufficient to resolve the stochasticity for this setup, and we do not observe significant changes between runs (see Sect. 3.1.1). The dashed line represents the first occurrence of ice ($q_i > 10^{-7} \text{ kg kg}^{-1}$). Since w is constant, there is a constant adiabatic cooling rate applied. Hence, an earlier time step is associated with warmer temperatures. Panels (a) and (e) show the evolution of ambient temperature T in black, as well as the saturation ratios with respect to water (S_w , red) and ice (S_i , blue). T governs J_{hom} for JHOM-T, while the ratio of S_w and S_i governs

J_{hom} for JHOM-DWA. Before homogeneous freezing sets in (dashed line), both simulations show the same evolution of T , S_w , and S_i . After activation of the (monodisperse) CCN population at the beginning, it takes several seconds until the droplets equilibrate with the ambient water vapour and water saturation ($S_w = 1$) is reached. T decreases continuously, mainly driven by adiabatic cooling. This leads to a decrease in $e_{\text{sat},i}$, which in turn increases S_i from 1.3 to 1.5 over the course of the simulation. Panels (b) and (f) show the microphysical bulk quantities for both simulations. The mixing ratio of liquid water (q_w , red) increases steadily due to the constant supply of supersaturation, while the vapour mixing ratio (q_v , black) decreases. Only about $\Delta t \approx 75$ s after the first freezing event does q_w begin to decrease significantly as droplets freeze and contribute to the ice mixing ratio (q_i , blue). At this point, the air parcel becomes supersaturated with respect to water ($S_w > 1$), because too few droplets remain to maintain water saturation through condensation (and DEP-ICE is disabled). Panels c and g show a cumulative histogram of freezing temperatures T_{frz} . Here the cumulative frequency of T_{frz} is the same as the frozen fraction of super-particles. Since all super-particles freeze in this example, the frozen fraction reaches 1 by the end of the simulations. These histograms demonstrate that the stochastic freezing approach used here leads to a gradual freezing event over a range of T_{frz} from ~ 239 to 235 K. The first freezing event, indicated by the dashed line, occurs close to 238 K for both simulations. However, note that the first T_{frz} can vary slightly between simulation runs due to the Monte-Carlo approach. A significant frozen fraction (> 0.01) only occurs at temperatures lower than 237 K. The frozen fraction for JHOM-DWA increases more steeply than for JHOM-T, completing the freezing event at temperatures slightly above 236 K, whereas JHOM-T completes freezing closer to the “classical” homogeneous-freezing limit of 235 K. As discussed in Sect. 2.1, J_{hom} for JHOM-DWA and JHOM-T should be of the same order of magnitude for $S_w = 1$, a condition that is maintained for most of the freezing events. Panels (d) and (h) show J_{hom} for JHOM-T in blue and for JHOM-DWA in red. The line indicates the active parameterisation, while the dashed line shows the other parameterisation under the same thermodynamic conditions for comparison. As freezing begins, the two parameterisations agree closely in order of magnitude. However, plotting the relative error (with JHOM-T considered the reference value) already reveals a divergence during the early onset of freezing. Although such deviations might be considered a “close match” in studies focusing on J_{hom} (see, e.g., Koop and Murray, 2016; Spichtinger et al., 2023), they have a significant impact on the stochastic freezing approach employed here. Since the JHOM-DWA rate is slightly higher than JHOM-T even for $S_w = 1$, this leads to a shift of the histogram toward higher T_{frz} . Once $S_w = 1$ is no longer maintained, the difference between JHOM-DWA and JHOM-T becomes much larger. This is a consequence of

the Δa_w -dependent approach and would also occur even if JHOM-DWA and JHOM-T matched exactly at $S_w = 1$.

3.1.1 n_{sd} ensemble

Figure 3a and b show cumulative density estimates of freezing temperatures T_{frz} for ensemble simulations for a range of super-particle numbers n_{sd} (10, 100 and 1000) used to discretise the initial CCN/droplet size distribution. Panels a and b show the results with the JHOM-T and JHOM-DWA rates, respectively. Each ensemble consists of 5 model runs with the same n_{sd} . Otherwise, all simulations have the same initial conditions with a constant updraft of $w = 1 \text{ m s}^{-1}$. The line is the ensemble mean, and the shaded area indicates the upper and lower bounds of the ensemble. The general impact of J_{hom} rate choice is the same as describe in Sect. 3.1 and holds true for all n_{sd} . JHOM-DWA (Panel b) shows the onset of significant homogeneous freezing at higher T_{frz} . The increase is sharper and full glaciation is achieved at $T_{\text{frz}} > 236 \text{ K}$. JHOM-T (Panel a) on the other hand shows a more gradual increase and full glaciation is achieved at lower T_{frz} than JHOM-DWA but still before the “classical” homogeneous nucleation threshold $T_{\text{frz}} = 235 \text{ K}$. Simulations with only $n_{\text{sd}} = 10$ (grey shaded) show large ensemble spread in the resulting T_{frz} histogram corresponding to a high variance for the small sample size. Simulations with $n_{\text{sd}} = 100$ show a larger spread for JHOM-T than for JHOM-DWA. The higher J_{hom} of JHOM-DWA leads to freezing in a narrower T_{frz} range, and thus to shorter nucleation events (with this setup of constant w). Simulations with $n_{\text{sd}} = 1000$ show a very narrow spread such that there are only negligible differences in the T_{frz} histogram between the 5 model realisations of the ensemble. Hence, for the setup in this section – in particular, with no DEP-ICE and the use of a monodisperse size spectra – just 1000 super-particles are sufficient to resolve homogeneous freezing events using a stochastic approach. Therefore we performed all other simulations that do not include DEP-ICE with $n_{\text{sd}} = 1000$.

3.1.2 w ensemble

Figure 3c and d show histograms of T_{frz} for w ranging from 0.2 to 10 m s^{-1} . The second y axis shows the mean cooling rate of the air parcel. This rate is primarily determined by (constant) adiabatic cooling, but it is modulated over time by latent heating. Consequently, the exact values differ slightly between the two rates, as the timing of freezing events affects the latent heating. Overall, weak updrafts of 0.2 m s^{-1} correspond to $\approx 1.8 \text{ mK s}^{-1}$, whereas strong updrafts of 10 m s^{-1} correspond to $\approx 93 \text{ mK s}^{-1}$.

For the JHOM-T rate (panel c), all T_{frz} values lie between 234.5 and 239 K. We note a shift toward lower T_{frz} for higher w , which is a consequence of the time dependence of the stochastic homogeneous freezing approach: at high w , droplets have less time to trigger a freezing event be-




Figure 2. Air parcel ascent simulations for $n_{sd} = 1000$ using the JHOM-T (a–d) and JHOM-DWA (e–h) J_{hom} formulations, showing (a, e) ambient temperature T and saturation ratios with respect to water S_w and ice S_i , (b, f) bulk microphysical quantities, and (c, g) cumulative histograms of freezing temperatures T_{frz} . Panels (d) and (h) show the J_{hom} rates (solid lines indicate the active formulation, dashed lines the alternative formulation for comparison), as well as the relative difference between the JHOM-T and JHOM-DWA rates (grey). The vertical dashed line marks the first freezing event.

fore reaching lower temperatures. For strong updrafts of 5 and 10 m s^{-1} , representative of conditions in deep convective systems, the T_{frz} histogram extends to temperatures below the classical homogeneous freezing threshold. The marginal plot above the figure shows the T_{frz} histogram aggregated over all w , which for JHOM-T is approximately normally distributed and centered close to 236 K. Panel d shows the T_{frz} histograms of the w ensemble for the JHOM-DWA rate. The overall histogram differs substantially from the ensemble employing JHOM-T. T_{frz} is capped above 236 K for all w , and we do not observe the general shift toward lower T_{frz} at higher w . Instead, T_{frz} remains largely unchanged across most of the w range. Only for convective-like updrafts with $w > 5 \text{ m s}^{-1}$ do we see a notable shift – but unlike JHOM-T, this shift is toward higher T_{frz} . This behavior stems from the Δa_w dependence of JHOM-DWA: for such strong updrafts, water saturation cannot be maintained, and the ambient air becomes slightly supersaturated with respect to water. As discussed in Sect. 2.1, this causes a strong increase in J_{hom} for JHOM-DWA, leading to freezing at higher temperatures.

3.1.3 n_{ccn} ensemble

Figure 3e and f show 2D histograms of T_{frz} for n_{ccn} ranging from 100 to 10000 cm^{-3} (at STP) with $w = 1 \text{ m s}^{-1}$ for the

JHOM-T and JHOM-DWA rates. Since the CCN size distribution is monodisperse, all super-particles always have the same radius. Consequently, the droplet radius r_w decreases with increasing n_{ccn} , as more droplets compete for the available supersaturation. Therefore, the n_{ccn} ensemble can be interpreted as an r_w ensemble, with high n_{ccn} corresponding to small r_w and vice versa. The second y axis shows r_w at the time of the first freezing event, t_f . Since t_f differs between simulations, small differences appear in the labels of the second y axis in panels e and f. As shown in Fig. 2b and f, the droplet mixing ratio q_w increases over time until enough droplets have frozen. r_w also increases over time but changes only weakly after t_f , making it a suitable reference radius for freezing. The process rate of stochastic homogeneous freezing is the product of J_{hom} and the droplet volume $V_l = 4/3\pi r_w^3$. Therefore, larger droplets have a higher probability of freezing. This behavior is evident for both n_{ccn} ensembles using the JHOM-T and JHOM-DWA rates. For higher n_{ccn} (smaller r_w), the T_{frz} histograms shift toward lower temperatures. For JHOM-T, the distribution extends below the classical homogeneous freezing threshold for $n_{ccn} \approx 10000 \text{ cm}^{-3}$. Under the same conditions, for JHOM-DWA all T_{frz} are above the classical homogeneous threshold. Note that the absolute decrease in r_w between very high and very low n_{ccn} is only about $7 \mu\text{m}$. However, this decrease is amplified in the process rates of probabilistic

freezing, since they scale with droplet volume. The process rate for the largest droplets (lowest n_{ccn}) is approximately 90 times larger than that for the smallest droplets (highest n_{ccn}). Comparing JHOM-T and JHOM-DWA, we again observe the same overall trends as before: JHOM-DWA histograms are shifted toward higher T_{frz} by 0.5 to 1 K. Both rates, exhibit long-tailed distributions for each n_{ccn} . In particular, the upper end of the T_{frz} range, where only few freezing events occur, is strongly influenced by the probabilistic formulation of the homogeneous freezing approach.

3.2 Simulations with vapour deposition on ice

We now investigate the setup in which vapour deposition on ice (DEP-ICE) is active in addition to ambient thermodynamics, condensation, and homogeneous freezing. Frozen droplets are no longer passive tracers; instead, they interact with the ambient air through vapour deposition. As vapour is depleted, the air parcel becomes increasingly subsaturated with respect to water ($S_w < 1$). Consequently, once a sufficient number of droplets have frozen, the remaining liquid super-particles begin to evaporate. This marks the onset of the Wegener-Bergeron-Findeisen process. The air parcel is strongly supersaturated with respect to ice ($S_i \sim 1.5$) in the temperature range of homogeneous freezing in this study (239 to 234 K). Thus, the thermodynamic driving force of the Wegener-Bergeron-Findeisen process is exceptionally large, owing to the large difference between S_w and S_i (see also Fig. 2). In this setup, the resulting ice number concentration n_i at the end of the simulation is of particular interest, because not all droplets will freeze: $n_{\text{ccn}} = n_c(t < t_f) \neq n_i(t_{\text{end}})$.

3.2.1 n_{sd} ensemble

Figure 4a and b show cumulative density estimates of freezing temperatures T_{frz} for n_{sd} ensemble simulations using the JHOM-T and JHOM-DWA rates, respectively. In contrast to the n_{sd} ensemble discussed in Sect. 3.1.1, we now consider larger n_{sd} with the range of values – 100, 1000, and 10000, i.e., one order of magnitude higher. As before, five simulations were performed for each n_{sd} , and the shaded areas denote the upper and lower bounds of the frozen fractions (cumulative frequency of T_{frz}) for each ensemble. The solid lines denote the mean of each ensemble. All simulations share the same initial conditions with $w = 1 \text{ m s}^{-1}$. Only those super-particles that actually freeze contribute to the statistics shown in the histograms. The n_{sd} ensemble using the JHOM-T rate (Panel a) shows a substantially larger spread than in the simulations without DEP-ICE. Previously, $n_{\text{sd}} = 1000$ (blue) was sufficient to ensure that the five model realisations produced nearly indistinguishable T_{frz} histograms. With DEP-ICE active, however, $n_{\text{sd}} = 10000$ is required to achieve statistically robust T_{frz} histograms. The freezing histogram is also shifted toward higher temperatures. The onset of sig-

nificant freezing now occurs at $\sim 238 \text{ K}$, and the final homogeneous freezing events occur just below 237 K. Without DEP-ICE, this range extended from 237 to 235.5 K (see Sect. 3.1.1). Consequently, the histograms exhibit a substantially steeper ascent compared to the case without DEP-ICE. The histograms of the JHOM-DWA rate ensemble (Panel b) are not statistically robust even for $n_{\text{sd}} = 10000$. For all values of n_{sd} , the model realisations differ substantially. Freezing also begins at higher temperatures, with $T_{\text{frz}} > 238.5 \text{ K}$. However, the final freezing events occur at slightly lower temperatures than for JHOM-T. In contrast to the simulations without DEP-ICE, the ascent of the JHOM-DWA histogram is thus more gradual than that of JHOM-T. This shift of the T_{frz} histogram toward higher temperatures, together with the reduced statistical robustness, arises from the Wegener-Bergeron-Findeisen process. In Sect. 3.1 we noted that the tail of the T_{frz} histograms is long, meaning that a few droplets occasionally froze at relatively high temperatures, but these rare events had little influence on the overall statistics. With DEP-ICE active, however, such “lucky” early-freezing super-particles encounter high S_i , and still liquid super-particles begin to evaporate before the air parcel cools further. As a result, the model becomes more sensitive to the stochasticity inherent in the homogeneous freezing formulation. Moreover, fewer than 7.5 % of droplets freeze, while the remaining population evaporates. This is shown by the frozen fraction $n_{\text{frz}} = 1 - \frac{n_c(0) - n_i(t_{\text{end}})}{n_c(0)}$ in Fig. 5a, where the red and blue lines indicate the ensemble mean of simulations using the JHOM-DWA and JHOM-T rates, respectively. The shaded area denotes the spread across the five ensemble realisations. Because only a small fraction of super-particles freeze, the shown T_{frz} histograms contain far fewer data points than those presented in the previous sections. For clouds transitioning from a liquid or mixed-phase state to a pure ice-phase (cirrus) cloud, the key quantity of interest is the ice crystal number concentration n_i after all droplets have either frozen or evaporated. This is shown in Fig. 5b, where we observe surprisingly little difference between simulations using $n_{\text{sd}} = 1000$ and $n_{\text{sd}} = 10000$. Moreover, JHOM-DWA generally exhibits a smaller ensemble spread in n_i , despite its broader T_{frz} histogram.

3.2.2 w ensemble

For the case without DEP-ICE active (see Sect. 3.1.1), higher updraft velocities led to lower T_{frz} because, in the stochastic homogeneous-freezing framework, droplets experience less time to freeze before reaching lower temperatures. This behaviour still holds when DEP-ICE is active, but the effect is amplified by the Wegener-Bergeron-Findeisen process. In strong updrafts, droplets are advected to lower temperatures more rapidly, where J_{hom} is larger, increasing the likelihood that they freeze before evaporating. This is reflected in Fig. 5c, which shows the frozen fraction n_{frz} for the w ensemble. n_{frz} increases with increasing w . For weak up-




Figure 3. Ensemble simulations without DEP-ICE. Cumulative histograms of T_{frz} for the n_{sd} ensemble using (a) JHOM-T and (b) JHOM-DWA rates, respectively. Two-dimensional histograms of T_{frz} for the w ensemble using (c) JHOM-T and (d) JHOM-DWA rates, respectively. Two-dimensional histograms of T_{frz} for the n_{ccn} ensemble using (e) JHOM-T and (f) JHOM-DWA rates, respectively.

drafts ($w = 0.2 \text{ m s}^{-1}$), fewer than 0.2 % of droplets freeze for both parameterisations. At $w \approx 1 \text{ m s}^{-1}$, more than 2.5 % freeze for JHOM-T, whereas JHOM-DWA remains below 0.5 %. This difference grows further at stronger updrafts, where up to 20 % and 5.5 % of droplets freeze for JHOM-T and JHOM-DWA, respectively. Figure 4c and d show histograms of T_{frz} for w ranging from 0.2 to 10 m s^{-1} . The second y axis indicates the mean cooling rate of the air parcel. While this rate is still primarily set by (dry) adiabatic cooling, it is now additionally modulated by latent heating from both condensation and DEP-ICE. For the JHOM-T rate, the overall histogram (as illustrated by the marginal distribution above) is centered at 236.75 K, where most freezing events occur, but it lacks a substantial number of freezing events at lower T_{frz} . For the JHOM-DWA rate, the T_{frz} histograms are similarly narrow for $w < 1 \text{ m s}^{-1}$, yet become significantly broader for stronger updrafts. Interestingly, as previously discussed, the frozen fraction n_{frz} is much lower for

JHOM-DWA. Thus, fewer freezing events are spread over a much wider temperature range. This behaviour again reflects the Δa_w -dependence of JHOM-DWA. In the case without DEP-ICE, JHOM-DWA was consistently larger than JHOM-T throughout the simulation because $S_w \geq 1$. While this remains true at the onset of freezing, the Wegener-Bergeron-Findeisen process rapidly drives the air parcel into subsaturation with respect to water ($S_w < 1$). Once sufficient droplets have frozen, the JHOM-DWA rate becomes much smaller for the remaining droplets, effectively delaying further freezing events. This can also be seen in Fig. S1 in the Supplement showing a sample ascent at $w = 5 \text{ m s}^{-1}$. Finally, note that the entire freezing period is considerably shorter for JHOM-T than JHOM-DWA, which is reflected in the narrower range of T_{frz} . Figure 5d shows n_i for the w ensemble for the JHOM-DWA (red) and JHOM-T (blue) rates. For both nucleation rates we find that n_i increases with updraft speed, spanning more than 1.5 orders of magnitude. Across the entire range

of w , the JHOM-T rate produces substantially higher n_i than JHOM-DWA. It is also important to note that the resulting n_i values are much lower than those obtained using a homogeneous freezing threshold (where $n_i = n_{\text{ccn}}$). However, we emphasise that this finding applies only to the idealised setup considered here, which excludes ice-nucleating particles and pre-existing ice. Overall, stronger updrafts lead to higher n_i . We do not show the bulk ice mixing ratio q_i here. Since newly formed ice crystals rapidly equilibrate to $S_i = 1$, q_i is thermodynamically constrained and is therefore mainly controlled by the adiabatic cooling rate, which itself primarily depends on w . For completeness, a corresponding figure showing q_i is provided in the Supplement (see Fig. S4).

3.2.3 n_{ccn} ensemble

For the n_{ccn} ensemble, all simulations use $w = 1 \text{ m s}^{-1}$. For the JHOM-T rate (Fig. 4e), the range of T_{fz} is similar to that obtained in the w ensemble at this updraft for all choices of n_{ccn} . The corresponding overall distribution (shown in the marginal plot above) is approximately a normal distribution and centered at $\sim 237.25 \text{ K}$. The frozen fraction n_{fz} (Fig. 5e) shows that fewer than 0.2 % of droplets freeze at high n_{ccn} , whereas about 10 % freeze at low n_{ccn} . Although n_i still increases with n_{ccn} (Fig. 5f), although a larger fraction of droplets evaporates, the total n_i still benefits from the increased number of available droplets. The histograms for the JHOM-DWA rate (Fig. 4f) show a large spread of T_{fz} for low n_{ccn} (large droplets) and a small spread for high n_{ccn} (small droplets), with T_{fz} generally extending to lower temperatures than for JHOM-T. As before, the Δa_w -dependency makes the freezing behaviour more complex, resulting in increased variability across the ensemble. The behaviour of n_{fz} and n_i is similar to that of JHOM-T, but, consistent with the w ensemble, both quantities take lower values. The size dependency of homogeneous freezing acts as a limitation in this ensemble setup. Larger droplets (low n_{ccn}) freeze earlier during the ascent (at higher T), which subsequently leads to the evaporation of a larger number of droplets. That higher n_{ccn} promotes more pronounced homogeneous freezing is also consistent with previous findings (Lüttmer et al., 2025b). We note that $n_{\text{sd}} = 10\,000$ was kept constant and that all ensemble simulations used monodisperse droplet size distributions. Therefore, increasing n_{ccn} increases the multiplicity ξ , i.e., the number of real droplets represented by a single super-particle. In simulations with DEP-ICE active, this becomes important because both the depletion of ambient water vapour and the associated latent heating are weighted by ξ . Additionally, DEP-ICE depends on the size of the super-particle due to the dependence of the mass rate for the growth on capacitance C and ice crystal radius r_i . Hence, a more rigorous approach would adjust n_{sd} when varying n_{ccn} . Nevertheless, for simplicity we vary only one model parameter at a time. Even so, the overall behaviour resulting from changes in n_{ccn} aligns with our expectations.

3.3 Simulations with polydisperse DSD

So far, we have used only a monodisperse mode to discretise the CCN size distribution. In the atmosphere, however, CCN size spectra are more diverse and often follow log-normal distributions or exhibit multiple modes. In this section, we use a log-normal distribution with a geometric mean radius of $r_0 = 0.05 \mu\text{m}$ and geometric standard deviations in the range $\sigma \in [1.5, 2]$. These values of σ are typical for a variety of aerosol species, including several types of mineral dust, sea salt, sulfate, and ammonium nitrate (Hande et al., 2016). A more comprehensive approach would involve varying not only σ , but also r_0 and the hygroscopicity parameter κ to represent CCN spectra more realistically. However, for simplicity, we again vary only a single parameter at a time. Moreover, in this model setup – with a single CCN mode and idealised parcel physics – the exact values of r_0 and κ (chosen within a reasonable range) do not significantly affect the simulation outcome. This is because, after activation, droplet sizes increase to a radius determined by the ambient water saturation ratio S_w . The use of a polydisperse CCN size spectrum has several important consequences. First, the multiplicity ξ is no longer uniformly distributed across all super-particles. As a result, different super-particles have varying impacts on the ambient thermodynamics, since condensation, latent heating, and vapour deposition on ice are weighted by ξ . Second, super-particles exhibit different droplet radii r because they activate at different times during the ascent, with the largest CCN activating first, and have more time to grow. This affects their freezing probabilities as it is size-dependent (see Sect. 2.1) as well as their growth rates by deposition of vapour. In this section, we exclusively employ the JHOM-T parameterization for the homogeneous freezing rate J_{hom} , which we consider the more accurate representation. We present simulations both with and without vapour deposition on ice. If not otherwise specified, all simulations in this section discretise the CCN size spectrum as a lognormal distribution with $r_0 = 0.05 \mu\text{m}$ and $\sigma = 2$, using $n_{\text{sd}} = 10\,000$. A sample ascent for these initial conditions is shown in the Supplement (see Fig. S2). The CCN size spectrum spans radii from 0.0025 to $1 \mu\text{m}$, with ξ varying by more than three orders of magnitude between the maximum and the tail end of the distribution. This leads to a smoother onset of droplet activation (in terms of saturation adjustment for S_w) and a wider, more gradual histogram of T_{fz} . As before, we choose $w = 1 \text{ m s}^{-1}$ and $n_{\text{ccn}} = 750 \text{ cm}^{-3}$ for the ensemble simulations.

3.3.1 σ ensemble

As for the previous sections, we also performed ensemble simulations for different n_{sd} . With vapour deposition on ice (DEP-ICE) inactive, $n_{\text{sd}} = 1000$ is sufficient to robustly represent the T_{fz} histogram. With DEP-ICE active, however, $n_{\text{sd}} = 10\,000$ is required. Since these results do not introduce




Figure 4. Ensemble simulations with vapour deposition on ice. Cumulative histograms of T_{frz} for the n_{sd} ensemble using (a) JHOM-T and (b) JHOM-DWA rates, respectively. Two-dimensional histograms of T_{frz} for the w ensemble using (c) JHOM-T and (d) JHOM-DWA rates, respectively. Two-dimensional histograms of T_{frz} for the n_{ccn} ensemble using (e) JHOM-T and (f) JHOM-DWA rates, respectively.

new insights beyond our previous findings, the corresponding plots are provided in the Supplement (see Fig. S3). Figure 6a shows the T_{frz} histogram for varying σ with DEP-ICE inactive. The histogram spans a T_{frz} range between 234 and 238 K for all choices of σ . In contrast to the monodisperse CCN spectra (see Sect. 3.1), T_{frz} extends below the classical homogeneous freezing threshold at 235 K. This is caused by small droplets populating the lower end of the CCN spectrum. The T_{frz} histogram does not change substantially with σ . Figure 6b shows the 2-D T_{frz} histogram for varying σ with DEP-ICE active. As in our previous findings, DEP-ICE active substantially narrows the range of T_{frz} , with the lowest values occurring just below 237 K. Because droplets on the upper tail of the CCN spectrum grow to the largest radii, they have the highest freezing probabilities. Once these droplets freeze, the smaller (and far more numerous) droplets evapo-

rate before freezing due to the Wegener-Bergeron-Findeisen process. Overall, the T_{frz} histograms show only minor sensitivity to the choice of σ . This is consistent with the resulting ice number concentration n_i and frozen fraction n_{frz} after glaciation, both of which are largely independent of σ (see Fig. S5). Hence, the freezing behaviour of the droplet spectrum is governed primarily by the ambient thermodynamics (here controlled by w) and by the droplet number concentration (controlled by n_{ccn}), rather than by the width of the CCN size distribution (controlled by σ).

3.3.2 w ensemble

Figure 6c shows the 2-D T_{frz} histogram for varying w with DEP-ICE inactive. Compared to the ensemble with a monodisperse CCN spectrum (Fig. 3c), the T_{frz} range is




Figure 5. Frozen fraction n_{frz} for (a) the n_{sd} ensemble, (c) the w ensemble, (e) and the n_{ccn} ensemble. Ice number concentration n_i for (b) the n_{sd} ensemble, (d) the w ensemble, and (f) the n_{ccn} ensemble. Blue lines indicate simulations using the JHOM-T nucleation rate with monodisperse CCN spectra, teal lines the JHOM-T rate with lognormal CCN spectra, and red lines simulations using the JHOM-DWA rate. All simulations were performed with $n_{\text{sd}} = 10000$ and vapour deposition on ice active.

broad. The histogram still exhibits a systematic shift towards lower temperatures for higher w , caused by the time dependence of the stochastic homogeneous freezing process. However, in contrast to the monodisperse case, two distinct modes emerge in the T_{frz} distribution (as shown by the marginal histogram above), rather than a single dominant mode. With DEP-ICE active, the T_{frz} histograms for lognormal and monodisperse CCN size spectra differ only marginally (compare Figs. 6d and 4c). Small differences remain in the bulk properties after the freezing event, with a lower frozen fraction and slightly lower ice number concentration n_i for the lognormal case (compare the blue and teal lines in Fig. 5c and d). Overall, the Wegener-Bergeron-Findeisen process smooths out differences in freezing behaviour between monodisperse and lognormal droplet spectra. In both cases, early freezing events at higher temperatures dominate the T_{frz} histogram and largely determine the bulk properties of the fully glaciated cloud. The timing of these early freezing events seems largely independent of the initial CCN size spectra.

3.3.3 n_{ccn} ensemble

Figure 6e shows the 2-D T_{frz} histogram for varying n_{ccn} with DEP-ICE inactive. The T_{frz} range is large for all n_{ccn} , spanning more than 5 K. The overall structure of the T_{frz} histogram does not change with n_{ccn} ; instead, it is shifted towards lower temperatures. This results in a broad maximum at ≈ 236 K with a pronounced tail towards lower T_{frz} . Consequently, T_{frz} values are generally lower than in the case of monodisperse CCN spectra (see Fig. 3e). With DEP-ICE active, fewer than 12.5 % of real droplets freeze even for low n_{ccn} (see Fig. 5e, teal line). As a consequence, the T_{frz} histogram becomes much narrower and is shifted towards higher temperatures corresponding to the earliest freezing droplets. As for the w ensemble, the difference in freezing behaviour between monodisperse and lognormal CCN spectra is negligible. Again, only the ice number concentration n_i after completion of the freezing event is slightly lower (compare the blue and teal lines in Fig. 5f).




Figure 6. Ensemble simulations using the JHOM-T parametrisation. Two-dimensional histograms of T_{frz} for the σ ensemble with (a) and without (b) vapour deposition on ice; for the w ensemble with (c) and without (d) deposition on ice; and for the n_{ccn} ensemble with (e) and without (f) deposition on ice.

4 Discussion and summary

In this study, we implement stochastic homogeneous freezing of pure water droplets using a Monte Carlo scheme within the super-particle model PySDM. The freezing probability depends on the droplet volume v , the time interval Δt , and the homogeneous nucleation rate J_{hom} . For J_{hom} , we employ two parametrisations: one depending only on temperature (JHOM-T), and one depending on both temperature and saturation ratio with respect to ice (JHOM-DWA). While JHOM-T represents the state-of-the-art formulation for J_{hom} for pure water droplets in the μm size range, the JHOM-DWA rate is additionally applicable to aqueous solution droplets in the nm size range. Both rates “closely”

agree at water saturation, $S_w = 1$. We use a simple parcel-model framework to investigate the distribution of freezing temperatures T_{frz} and the resulting ice number concentration n_i after the freezing event has concluded. We test both nucleation rates for ensembles varying the number of super-particles n_{sd} used to discretise the initial droplet spectrum, the updraft speed w controlling the adiabatic cooling rate, the cloud condensation nuclei (CCN) number concentration n_{ccn} controlling droplet size and super-particle multiplicity, and the geometric standard deviation σ of the lognormal CCN size spectrum. We also performed simulations with and without vapour deposition on ice (DEP-ICE). When DEP-ICE is inactive, frozen droplets act as passive tracers. When DEP-ICE is active, vapour deposition on ice enables the Wegener-

Bergeron-Findeisen process. All simulations in this paper assume a constant updraft (and thus cooling).

4.1 Fundamental characteristics of homogeneous freezing

Our ensemble simulations confirm the fundamental characteristics of homogeneous freezing events. The shift of freezing temperatures T_{frz} towards lower temperatures for stronger updrafts reflects the time dependency of the stochastic freezing process, as droplets have less time to freeze before reaching lower temperatures. Parameters affecting the CCN size spectrum – the width determined by σ and the multiplicity determined by n_{ccn} – control the droplet size, with larger droplets shifting T_{frz} towards higher temperatures. Repeating the simulations yields different T_{frz} histograms, highlighting the probabilistic formulation of homogeneous freezing. To obtain smooth T_{frz} histograms, $n_{\text{sd}} = 1000$ super-particles were sufficient when DEP-ICE was inactive, whereas $n_{\text{sd}} = 10000$ were required when DEP-ICE was active. However, bulk properties such as the ice number concentration n_i were already robust for $n_{\text{sd}} = 1000$.

4.2 Distribution of freezing temperatures T_{frz}

Most cloud models employ a threshold formulation for homogeneous nucleation of pure water droplets, in which all droplets freeze once the ambient temperature drops below a prescribed threshold T_{hom} . A commonly used value is $T_{\text{hom}} = 235 \text{ K}$ (or -38°C). In contrast, we find that most super-particles freeze at temperatures higher than T_{hom} , with the highest freezing temperatures T_{frz} close to 240 K . Only for strong updrafts ($w > 5 \text{ m s}^{-1}$), high CCN number concentrations ($n_{\text{ccn}} > 1250 \text{ cm}^{-3}$), or lognormal CCN size spectra do some super-particles freeze below T_{hom} . With vapour deposition on ice (DEP-ICE) active, the thermodynamic forcing of the Wegener-Bergeron-Findeisen process is particularly strong in this temperature range. As a result, most droplets ($> 90\%$) evaporate before freezing following the first few freezing events, which typically occur at $T_{\text{frz}} > 237 \text{ K}$. In this sense, our results are consistent with the classical homogeneous nucleation threshold $T_{\text{hom}} = 235 \text{ K}$, in that by this temperature most droplets have indeed been removed or are frozen. However, two important caveats must be considered. First, when ice is already present (e.g. due to immersion freezing), liquid droplets may evaporate before reaching T_{hom} . In a more realistic stochastic freezing framework, these droplets would still have a significant probability of freezing. In this case, a threshold formulation underestimates the resulting ice number concentration n_i . Second, in the absence of pre-existing ice or ice-nucleating particles, the threshold formulation overestimates n_i , since in reality early freezing events trigger the evaporation of a large fraction of the remaining droplets via the Wegener-Bergeron-Findeisen process.

4.3 Choice of J_{hom} parametrisation

The choice of homogeneous nucleation rate parameterisation influences both the distribution of T_{frz} and the resulting n_i after the freezing event. Although the JHOM-T and JHOM-DWA formulations have similar slopes and are often considered to agree closely at water saturation ($S_w = 1$), even small absolute differences between the two can lead to different onsets of freezing. Their behaviour diverges further when water saturation is no longer maintained. This divergence arises from the explicit dependence of the JHOM-DWA formulation on water activity, which makes the nucleation rate sensitive to small deviations of S_w . In our simulations, use of the JHOM-DWA formulation is associated with broader T_{frz} distributions and generally lower n_i compared to JHOM-T, particularly when vapour deposition on ice (DEP-ICE) is active. While JHOM-DWA in principle provides a unified framework applicable to both cloud droplets and aqueous solution droplets, our results suggest that its application to micrometre-sized cloud droplets in mixed-phase conditions requires care, especially in models that explicitly predict S_w (e.g., without saturation adjustment). However, we should not neglect the JHOM-DWA approach for cloud droplets entirely. While these can mostly be considered pure droplets, as the amount of solute from the CCN is low in this size range, these foreign substances should still have some effect on the freezing behaviour. However, constraining the magnitude of this effect is beyond the scope of this study.

4.4 Limitations

In this study, we employ a highly idealised model setup. The simple air-parcel framework does not account for turbulent mixing or interactions with the surrounding flow. In addition, sedimentation is neglected, both the sedimentation of particles into the air parcel and their removal from it. Including sedimentation in a parcel framework is inherently problematic, as it depends on the arbitrary choice of parcel volume and the assumed surrounding cloud environment. Droplet coalescence is also not considered. However, this process is not expected to be relevant for most setups investigated here, as the commonly used geometric collision kernel is not applicable to the monodisperse CCN spectra employed in this study. While coalescence would lead to larger droplets that freeze at higher temperatures, it would also modify the multiplicity distribution ξ , thereby affecting how super-particle–environment interactions such as condensation, latent heating, and DEP-ICE are weighted. The Wegener-Bergeron-Findeisen process plays a key role in stochastic homogeneous freezing events, as early freezing of a small number of “lucky” super-particles can trigger the evaporation of the remaining liquid droplets. In our simulations, typically more than 90% of droplets evaporate before freezing. However, several studies suggest that glaciation driven by the Wegener-Bergeron-Findeisen process may be overestimated in cloud

models, and that mixed-phase conditions can be maintained for longer periods (Abade and Albuquerque, 2024). Improving the representation of this process is therefore critical for accurately modelling homogeneous freezing events. Finally, all simulations were performed without ice-nucleating particles (INPs) or pre-existing ice. Including INPs would introduce additional degrees of freedom, such as the freezing temperature and surface area of INPs. Here, we focus on the fundamental behaviour of homogeneous freezing. We further argue that the INP-free conditions considered here are relevant in specific atmospheric situations, such as at rising cloud tops with high droplet concentrations or in regions where pre-existing ice has already sedimented out of the air parcel.

Code and data availability. Figures 1–6 and S1–S5 can be reproduced using Jupyter notebooks openly available at <https://github.com/open-atmos/PySDM> (last access: 9 October 2026). The exact version of the model used to produce the results used in this paper is archived on Zenodo under <https://doi.org/10.5281/zenodo.22080761> (Lüttmer, 2026), as are scripts to run the model and produce the plots for all the simulations presented in this paper.

Supplement. The supplement related to this article is available online at <https://doi.org/10.5194/gmd-19-10185-2026-supplement>.

Author contributions. TL and SA contributed to the model development; TL designed the study; TL performed the simulations and carried out the data analyses; TL and PS contributed to interpreting the results; TL, SA and PS contributed to writing the paper.

Competing interests. At least one of the (co-)authors is a member of the editorial board of *Geoscientific Model Development*. The peer-review process was guided by an independent editor, and the authors also have no other competing interests to declare.

Disclaimer. Publisher's note: Copernicus Publications remains neutral with regard to jurisdictional claims made in the text, published maps, institutional affiliations, or any other geographical representation in this paper. The authors bear the ultimate responsibility for providing appropriate place names. Views expressed in the text are those of the authors and do not necessarily reflect the views of the publisher.

Acknowledgements. We thank the anonymous reviewers for their insightful comments on the original paper.

Financial support. This research has been supported by the Deutsche Forschungsgemeinschaft (grant no. 428312742), by the Polish National Science Centre (grant no. 2020/39/D/ST10/01220)

and by the Mainz Institute of Multiscale Modeling (M3ODEL Junior Fellowship).

Review statement. This paper was edited by Wojciech W. Grabowski and reviewed by two anonymous referees.

References

- Abade, G. C. and Albuquerque, D. G.: Persistent mixed-phase states in adiabatic cloud parcels under idealised conditions, *Q. J. Roy. Meteor. Soc.*, 150, 3450–3474, <https://doi.org/10.1002/qj.4775>, 2024.
- Alpert, P. A. and Knopf, D. A.: Analysis of isothermal and cooling-rate-dependent immersion freezing by a unifying stochastic ice nucleation model, *Atmos. Chem. Phys.*, 16, 2083–2107, <https://doi.org/10.5194/acp-16-2083-2016>, 2016.
- Arabas, S., Jaruga, A., Pawlowska, H., and Grabowski, W. W.: libcloudph++ 1.0: a single-moment bulk, double-moment bulk, and particle-based warm-rain microphysics library in C++, *Geosci. Model Dev.*, 8, 1677–1707, <https://doi.org/10.5194/gmd-8-1677-2015>, 2015.
- Arabas, S., Curtis, J. H., Silber, I., Fridlind, A. M., Knopf, D. A., West, M., and Riemer, N.: Immersion Freezing in Particle-Based Aerosol-Cloud Microphysics: A Probabilistic Perspective on Singular and Time-Dependent Models, *J. Adv. Model. Earth Sy.*, 17, <https://doi.org/10.1029/2024MS004770>, 2025.
- Bartman, P., Bulenok, O., Górski, K., Jaruga, A., Łazarski, G., Oleśik, M. A., Piasecki, B., Singer, C. E., Talar, A., and Arabas, S.: PySDM v1: particle-based cloud modeling package for warm-rain microphysics and aqueous chemistry, *Journal of Open Source Software*, 7, 3219, <https://doi.org/10.21105/joss.03219>, 2022.
- Bigg, E. K.: The Supercooling of Water, *P. Phys. Soc. Lond. B*, 66, 688, <https://doi.org/10.1088/0370-1301/66/8/309>, 1953.
- Choudhury, G. and Tesche, M.: A first global height-resolved cloud condensation nuclei data set derived from spaceborne lidar measurements, *Earth Syst. Sci. Data*, 15, 3747–3760, <https://doi.org/10.5194/essd-15-3747-2023>, 2023.
- Costa-Surós, M., Gonçalves Ageitos, M., Chatziparaschos, M., Georgakaki, P., Thomas, M. A., Montané Pinto, G., Myriokefalitakis, S., van Noije, T., Le Sager, P., Kanakidou, M., Nenes, A., and Pérez García-Pando, C.: Implementation of primary and secondary ice production in EC-Earth3-AerChem: global impacts and insights, *Atmos. Chem. Phys.*, 26, 2667–2690, <https://doi.org/10.5194/acp-26-2667-2026>, 2026.
- de Jong, E. K., Singer, C. E., Azimi, S., Bartman, P., Bulenok, O., Derlatka, K., Dula, I., Jaruga, A., Mackay, J. B., Ward, R. X., and Arabas, S.: New developments in PySDM and PySDM-examples v2: collisional breakup, immersion freezing, dry aerosol initialization, and adaptive time-stepping, *Journal of Open Source Software*, 8, 4968, <https://doi.org/10.21105/joss.04968>, 2023.
- ECMWF: IFS Documentation CY48R1 – Part IV: Physical Processes, 4, ECMWF, <https://doi.org/10.21957/02054f0fbf>, 2023.
- Field, P. R., Hill, A., Shipway, B., Furtado, K., Wilkinson, J., Miltenberger, A., Gordon, H., Grosvenor, D. P., Stevens, R., and Van Weverberg, K.: Implementation of a double moment cloud microphysics scheme in the UK met office regional numerical

- weather prediction model, *Q. J. Roy. Meteor. Soc.*, 149, 703–739, <https://doi.org/10.1002/qj.4414>, 2023.
- Fusina, F., Spichtinger, P., and Lohmann, U.: Impact of ice supersaturated regions and thin cirrus on radiation in the midlatitudes, *J. Geophys. Res.-Atmos.*, 112, <https://doi.org/10.1029/2007JD008449>, 2007.
- Gasparini, B., Meyer, A., Neubauer, D., Münch, S., and Lohmann, U.: Cirrus Cloud Properties as Seen by the CALIPSO Satellite and ECHAM-HAM Global Climate Model, *J. Climate*, 31, 1983–2003, <https://doi.org/10.1175/JCLI-D-16-0608.1>, 2018.
- Hande, L. B., Engler, C., Hoose, C., and Tegen, I.: Seasonal variability of Saharan desert dust and ice nucleating particles over Europe, *Atmos. Chem. Phys.*, 15, 4389–4397, <https://doi.org/10.5194/acp-15-4389-2015>, 2015.
- Hande, L. B., Engler, C., Hoose, C., and Tegen, I.: Parameterizing cloud condensation nuclei concentrations during HOPE, *Atmos. Chem. Phys.*, 16, 12059–12079, <https://doi.org/10.5194/acp-16-12059-2016>, 2016.
- Hoshyaripour, G. A., Baer, A., Bierbauer, S., Bruckert, J., Brunner, D., Förstner, J., Hamzehloo, A., Hanft, V., Keller, C., Klose, M., Kumar, P., Ludwig, P., Metzner, E., Muth, L., Pauling, A., Porz, N., Ramezani Ziarani, M., Reddmann, T., Reißig, L., Ruhnke, R., Satitkovitchai, K., Seifert, A., Sinnhuber, M., Steiner, M., Versick, S., Vogel, H., Weimer, M., Werchner, S., and Hoose, C.: The atmospheric composition component of the ICON modeling framework: ICON-ART version 2025.10, *Geosci. Model Dev.*, 19, 1645–1681, <https://doi.org/10.5194/gmd-19-1645-2026>, 2026.
- Howell, W. E.: The Growth of Cloud Drops in Uniformly Cooled Air, *J. Atmos. Sci.*, 6, 134–149, [https://doi.org/10.1175/1520-0469\(1949\)006<0134:TGOCDI>2.0.CO;2](https://doi.org/10.1175/1520-0469(1949)006<0134:TGOCDI>2.0.CO;2), 1949.
- Ickes, L., Welti, A., Hoose, C., and Lohmann, U.: Classical nucleation theory of homogeneous freezing of water: thermodynamic and kinetic parameters, *Phys. Chem. Chem. Phys.*, 17, 5514–5537, <https://doi.org/10.1039/C4CP04184D>, 2015.
- Joos, H., Spichtinger, P., Reutter, P., and Fusina, F.: Influence of heterogeneous freezing on the microphysical and radiative properties of orographic cirrus clouds, *Atmos. Chem. Phys.*, 14, 6835–6852, <https://doi.org/10.5194/acp-14-6835-2014>, 2014.
- Jurányi, Z., Gysel, M., Weingartner, E., DeCarlo, P. F., Kammermann, L., and Baltensperger, U.: Measured and modelled cloud condensation nuclei number concentration at the high alpine site Jungfraujoch, *Atmos. Chem. Phys.*, 10, 7891–7906, <https://doi.org/10.5194/acp-10-7891-2010>, 2010.
- Kärcher, B. and Lohmann, U.: A parameterization of cirrus cloud formation: Homogeneous freezing of supercooled aerosols, *J. Geophys. Res.-Atmos.*, 107, <https://doi.org/10.1029/2001JD000470>, 2002.
- Koop, T. and Murray, B. J.: A physically constrained classical description of the homogeneous nucleation of ice in water, *J. Chem. Phys.*, 145, <https://doi.org/10.1063/1.4962355>, 2016.
- Koop, T., Luo, B., Tsias, A., and Peter, T.: Water activity as the determinant for homogeneous ice nucleation in aqueous solutions, *Nature*, 406, 611–614, <https://doi.org/10.1038/35020537>, 2000.
- Krämer, M., Rolf, C., Luebke, A., Afchine, A., Spelten, N., Costa, A., Meyer, J., Zöger, M., Smith, J., Herman, R. L., Buchholz, B., Ebert, V., Baumgardner, D., Borrmann, S., Klingebiel, M., and Avallone, L.: A microphysics guide to cirrus clouds – Part 1: Cirrus types, *Atmos. Chem. Phys.*, 16, 3463–3483, <https://doi.org/10.5194/acp-16-3463-2016>, 2016.
- Krämer, M., Rolf, C., Spelten, N., Afchine, A., Fahey, D., Jensen, E., Khaykin, S., Kuhn, T., Lawson, P., Lykov, A., Pan, L. L., Riese, M., Rollins, A., Stroh, F., Thornberry, T., Wolf, V., Woods, S., Spichtinger, P., Quaas, J., and Sourdeval, O.: A microphysics guide to cirrus – Part 2: Climatologies of clouds and humidity from observations, *Atmos. Chem. Phys.*, 20, 12569–12608, <https://doi.org/10.5194/acp-20-12569-2020>, 2020.
- Lamb, D. and Verlinde, J.: *Physics and chemistry of clouds*, Cambridge University Press, <https://doi.org/10.1017/CBO9780511976377>, 2011.
- Luebke, A. E., Afchine, A., Costa, A., Grooß, J.-U., Meyer, J., Rolf, C., Spelten, N., Avallone, L. M., Baumgardner, D., and Krämer, M.: The origin of midlatitude ice clouds and the resulting influence on their microphysical properties, *Atmos. Chem. Phys.*, 16, 5793–5809, <https://doi.org/10.5194/acp-16-5793-2016>, 2016.
- Lüttmer, T., Miltenberger, A., and Spichtinger, P.: On the impact of ice formation processes and sedimentation on cirrus origin classification in warm conveyor belt outflow, *Atmos. Chem. Phys.*, 25, 10245–10265, <https://doi.org/10.5194/acp-25-10245-2025>, 2025a.
- Lüttmer, T., Spichtinger, P., and Seifert, A.: Investigating ice formation pathways using a novel two-moment multi-class cloud microphysics scheme, *Atmos. Chem. Phys.*, 25, 4505–4529, <https://doi.org/10.5194/acp-25-4505-2025>, 2025b.
- Lüttmer, T.: Stochastic homogeneous freezing of supercooled droplets in mixed-phase clouds in particle-based idealised microphysics framework (exemplified with PySDM v3.0), Zenodo [code, data set], <https://doi.org/10.5281/zenodo.22080761>, 2026.
- Morrison, H. and Milbrandt, J. A.: Parameterization of Cloud Microphysics Based on the Prediction of Bulk Ice Particle Properties. Part I: Scheme Description and Idealized Tests, *J. Atmos. Sci.*, 72, 287–311, <https://doi.org/10.1175/JAS-D-14-0065.1>, 2015.
- Murray, B. J., Broadley, S. L., Wilson, T. W., Bull, S. J., Wills, R. H., Christenson, H. K., and Murray, E. J.: Kinetics of the homogeneous freezing of water, *Phys. Chem. Chem. Phys.*, 12, 10380–10387, <https://doi.org/10.1039/C003297B>, 2010.
- Petters, M. D. and Kreidenweis, S. M.: A single parameter representation of hygroscopic growth and cloud condensation nucleus activity, *Atmos. Chem. Phys.*, 7, 1961–1971, <https://doi.org/10.5194/acp-7-1961-2007>, 2007.
- Phillips, V. T., DeMott, P. J., and Andronache, C.: An empirical parameterization of heterogeneous ice nucleation for multiple chemical species of aerosol, *J. Atmos. Sci.*, 65, 2757–2783, <https://doi.org/10.1175/2007JAS2546.1>, 2008.
- Pruppacher, H., Klett, J., Pruppacher, H., and Klett, J.: *Microstructure of atmospheric clouds and precipitation*, Springer, <https://doi.org/10.1007/978-0-306-48100-0>, 2010.
- Seifert, A. and Beheng, K. D.: A two-moment cloud microphysics parameterization for mixed-phase clouds. Part 1: Model description, *Meteorol. Atmos. Phys.*, 92, 45–66, <https://doi.org/10.1007/s00703-005-0112-4>, 2006.
- Shima, S., Sato, Y., Hashimoto, A., and Misumi, R.: Predicting the morphology of ice particles in deep convection using the super-droplet method: development and evaluation of SCALE-SDM 0.2.5-2.2.0, -2.2.1, and -2.2.2, *Geosci. Model Dev.*, 13, 4107–4157, <https://doi.org/10.5194/gmd-13-4107-2020>, 2020.

- Skrotzki, J., Connolly, P., Schnaiter, M., Saathoff, H., Möhler, O., Wagner, R., Niemand, M., Ebert, V., and Leisner, T.: The accommodation coefficient of water molecules on ice – cirrus cloud studies at the AIDA simulation chamber, *Atmos. Chem. Phys.*, 13, 4451–4466, <https://doi.org/10.5194/acp-13-4451-2013>, 2013.
- Spichtinger, P. and Gierens, K. M.: Modelling of cirrus clouds – Part 1a: Model description and validation, *Atmos. Chem. Phys.*, 9, 685–706, <https://doi.org/10.5194/acp-9-685-2009>, 2009.
- Spichtinger, P., Marschalik, P., and Baumgartner, M.: Impact of formulations of the homogeneous nucleation rate on ice nucleation events in cirrus, *Atmos. Chem. Phys.*, 23, 2035–2060, <https://doi.org/10.5194/acp-23-2035-2023>, 2023.
- Ullrich, R., Hoose, C., Möhler, O., Niemand, M., Wagner, R., Höhler, K., Hiranuma, N., Saathoff, H., and Leisner, T.: A New Ice Nucleation Active Site Parameterization for Desert Dust and Soot, *J. Atmos. Sci.*, 74, 699–717, <https://doi.org/10.1175/JAS-D-16-0074.1>, 2017.
- Vali, G., DeMott, P. J., Möhler, O., and Whale, T. F.: Technical Note: A proposal for ice nucleation terminology, *Atmos. Chem. Phys.*, 15, 10263–10270, <https://doi.org/10.5194/acp-15-10263-2015>, 2015.
- Welss, J.-N., Siewert, C., and Seifert, A.: Explicit Habit-Prediction in the Lagrangian Super-Particle Ice Microphysics Model McSnow, J. *Adv. Model. Earth Sy.*, 16, e2023MS003805, <https://doi.org/10.1029/2023MS003805>, 2024.
- Zhang, Y., Macke, A., and Albers, F.: Effect of crystal size spectrum and crystal shape on stratiform cirrus radiative forcing, *Atmos. Res.*, 52, 59–75, [https://doi.org/10.1016/S0169-8095\(99\)00026-5](https://doi.org/10.1016/S0169-8095(99)00026-5), 1999.