



# Enhancing air–sea CO<sub>2</sub> exchange and modulating seawater carbonate–pH dynamics: the role of wave effect mechanisms in the POP2–waves coupled model

Yung-Yao Lan<sup>1,2</sup>, Huang-Hsiung Hsu<sup>1</sup>, Wei-Liang Lee<sup>1</sup>, and Simon Chou<sup>1</sup>

<sup>1</sup>Research Center for Environmental Changes, Academia Sinica, Taipei 11529, Taiwan

<sup>2</sup>National Land Resources Conservation Center, National Chung Hsing University, Taichung 40227, Taiwan

**Correspondence:** Wei-Liang Lee (leelupin@gate.sinica.edu.tw)

Received: 27 September 2025 – Discussion started: 19 December 2025

Revised: 26 July 2026 – Accepted: 29 July 2026 – Published: 13 August 2026

**Abstract.** In this study, a wave module is online-coupled into the Parallel Ocean Program version 2 (POP2) within the CESM1.2.2 framework (hereafter POP2–waves). Unlike empirical data analyses of observation-based products and offline-forced ocean models that treat wave-induced gas transfer velocity and the air–sea difference in partial pressure of CO<sub>2</sub> ( $\Delta p\text{CO}_2$ ) as decoupled variables, the online-coupled POP2–waves model captures their interactive,  $\Delta p\text{CO}_2$ -mediated negative feedbacks. This setup enables wave properties to dynamically modulate gas transfer velocity and physical mixing through sequential processes. POP2–waves is evaluated alongside a control simulation (B–CTL) against the National Oceanic and Atmospheric Administration (NOAA) CarbonTracker (CT2022) inversion product. The spatial air–sea CO<sub>2</sub> flux in POP2–waves simulation shows generally closer structural agreement with NOAA CT2022 than the control B–CTL, thereby improving performance in most selected regions. Specifically, bubble-mediated transfer accounts for up to 41.3 % of the total flux, consistent with the  $\sim 40\%$  contribution reported in recent research. Although the inclusion of waves (POP2–waves) enhances regional oceanic CO<sub>2</sub> uptake by 11.8 % and outgassing by 41.6 %, these two processes largely offset each other. Consequently, this dual enhancement results in only a slight 1.8 % increase in the global net ocean CO<sub>2</sub> sink compared to the B–CTL simulation. Globally, air–sea  $\Delta p\text{CO}_2$  and pH exhibit the strongest positive and negative regression coefficients with the air–sea CO<sub>2</sub> flux, respectively. Regionally, the gas transfer velocity shows a positive (negative) re-

gression coefficient within oceanic CO<sub>2</sub> outgassing (uptake) regions, whereas SST displays the opposite trend.

## 1 Introduction

The exchange of CO<sub>2</sub> between the air and sea is a crucial component of the global carbon cycle, carrying significant implications for Earth’s climate (Bange et al., 2024; Friedlingstein et al., 2022; McKinley et al., 2020; Müller et al., 2023; Shutler et al., 2019). Traditionally, this air–sea exchange is characterized using the bulk formula (e.g., Wanninkhof, 1992, 2014; McKinley et al., 2020; Dong et al., 2022; Fay et al., 2021; Zhou et al., 2023; Heimdal et al., 2024):

$$F_{\text{CO}_2} = k_{w,660} \cdot k_0 \cdot (p\text{CO}_2^w - p\text{CO}_2^a) \cdot (1 - \text{ice}) \quad (1)$$

where the air–sea CO<sub>2</sub> flux ( $F_{\text{CO}_2}$ , mol m<sup>−2</sup> yr<sup>−1</sup>) is determined by the gas transfer velocity expressed relative to a Schmidt number of 660 ( $k_{w,660}$ , cm h<sup>−1</sup>), the difference in partial pressure of CO<sub>2</sub> ( $\Delta p\text{CO}_2$ ,  $\mu\text{atm}$ ) between the seawater and atmosphere (indicated by superscripts “w” and “a”, respectively), and the solubility constant ( $k_0$ ; mol m<sup>−3</sup> atm<sup>−1</sup>), which varies with salinity and temperature (Weiss, 1974). Note that an appropriate scaling factor ( $8.76 \times 10^{-5}$ ) is implicitly required to convert the dimensions of  $k_{w,660}$  and  $\Delta p\text{CO}_2$  into the final flux unit (mol m<sup>−2</sup> yr<sup>−1</sup>). The sign of air–sea CO<sub>2</sub> flux is determined by  $\Delta p\text{CO}_2$ , where a positive air–sea CO<sub>2</sub> flux indicates ocean outgassing to the atmosphere. Additionally, the calculated fluxes are weighted by

the ice-free fraction ( $1 - \text{ice}$ ), where “ice” represents the sea ice fraction. The estimation of bulk air–sea CO<sub>2</sub> fluxes using the gas transfer velocity is typically based on the Wanninkhof (1992) formulation:

$$k_{w,660} = 0.31 < U_{10}^2 > (Sc/660)^{-0.5} \quad (2)$$

where  $< U_{10}^2 >$  represents the squared neutral mean wind speed at 10 m above the surface ( $\text{m s}^{-1}$ ), and  $Sc$  denotes the Schmidt number. The flux estimate error can reach as high as 34 % due to model limitations and uncertainties surrounding bulk parameters (Signorini and McClain, 2009). Furthermore, substantial uncertainties persist in air–sea CO<sub>2</sub> flux estimates, primarily stemming from an incomplete understanding of the spatiotemporal variability in the governing mechanisms (Shutler et al., 2019).

Some studies have highlighted that estimating air–sea CO<sub>2</sub> fluxes involves complexities well beyond neutral wind speed and solubility factors. Monahan and Spillane (1984) previously regarded whitecaps as “low impedance vents” that effectively “shorten” the water-side transfer resistance. Without wave breaking, air–sea CO<sub>2</sub> exchange occurs via mass transport, which under non-turbulent conditions is governed by molecular diffusion; wave breaking then acts as a transitional process from laminar to turbulent flow (Deike, 2022). Furthermore, bubbles provide additional surface area for gas transfer; as they rise through the aqueous mass boundary layer and burst at the sea surface, they significantly enhance near-surface turbulence (Soloviev and Lukas, 2010; Bell et al., 2017; Deike and Melville, 2018; Krall et al., 2019; Czerski et al., 2022). Gutiérrez-Loza et al. (2022) pointed out that during high and intermediate wind speeds (above 6–8  $\text{m s}^{-1}$ ), enhanced air–sea CO<sub>2</sub> exchange is primarily driven by wave-breaking dynamics and bubble mediation, which occur alongside the generation of sea spray droplets that contribute to atmospheric cloud condensation nucleation. Conversely, under low wind conditions (below 6  $\text{m s}^{-1}$ ), water-side convection becomes the dominant control mechanism. Beyond these calm periods, the critical role of breaking waves and bubble injection mechanisms in facilitating CO<sub>2</sub> transfer has been widely corroborated (e.g., Andreas et al., 2016; Brumer et al., 2017; Blomquist et al., 2017; Reichl and Deike, 2020; Li et al., 2023; Zhou et al., 2023; Rustogi et al., 2025). Specifically, Deike and Melville (2018) demonstrated that the bubble-mediated contribution to CO<sub>2</sub> transfer exceeds 40 % at a neutral 10 m wind speed ( $U_{10} \geq 10 \text{ m s}^{-1}$ ), becomes dominant at  $U_{10} \geq 15 \text{ m s}^{-1}$ , and reaches 60 % at 20  $\text{m s}^{-1}$ .

Monitoring sea surface CO<sub>2</sub> is crucial for understanding Earth system dynamics, as climate change has begun to impact the ocean’s carbon uptake capacity (Behncke et al., 2024). However, air–sea CO<sub>2</sub> flux estimates derived from products based on the partial pressure of CO<sub>2</sub> ( $p\text{CO}_2$ ) often suffer from significant uncertainties, stemming primarily from the empirical parameterization of the gas transfer velocity (e.g., Williams et al., 2017; Gray et al., 2018; Coggins

et al., 2023; Behncke et al., 2024; Fay et al., 2024; Yang et al., 2024). To resolve this issue and independently constrain  $k_{w,660}$ , direct flux measurements using the eddy covariance (EC) method are widely utilized as a benchmark (Dong et al., 2021). By directly acquiring the turbulent flux  $F_{\text{CO}_2}$  via the EC method and pairing it with simultaneous measurements of  $\Delta p\text{CO}_2$ , researchers can inversely calculate the gas transfer velocity ( $k_{w,660} = F_{\text{CO}_2} / (k_0 \times \Delta p\text{CO}_2)$ ). This approach provides a direct observational constraint to evaluate and calibrate these observation-based products under complex marine conditions. However, obtaining reliable direct fluxes from shipborne EC setups presents its own instrumental challenges. Marine EC measurements are generally conducted using either open- or closed-path infrared gas analyzers. A closed-path analyzer (e.g., LI-7000, LI-COR) operates with an enclosed measurement chamber (Sutton et al., 2014, 2021; Bakker et al., 2016; Sabine et al., 2020; Akhand et al., 2021; Wu and Qi, 2023), whereas open-path analyzers (e.g., LI-7500, LI-COR) measure infrared absorption directly in ambient air (Edson et al., 2011; Tokoro et al., 2014; Bell et al., 2017; Dong et al., 2021; Van Dam et al., 2021). While both configurations are inherently susceptible to H<sub>2</sub>O cross-sensitivity, each system possesses distinct advantages and systematic limitations (Honkanen et al., 2018). Evaluating and constraining global Earth system models (ESMs) against direct, long-term observations of air–sea CO<sub>2</sub> flux remains challenging due to the severe scarcity of continuous field measurements. Consequently, alternative validation datasets are typically derived by upscaling sparse in situ  $p\text{CO}_2$  observations through a combination of statistical interpolation, machine learning, atmospheric inversion, and climatological averaging to construct long-term, gridded flux products.

Although several recent studies have estimated the air–sea CO<sub>2</sub> flux using Eq. (1), current parameterizations in oceanic and atmospheric models still rely exclusively on neutral wind speed (Long et al., 2013; Moore et al., 2013; Couldrey et al., 2016; Jin et al., 2017; Lovenduski et al., 2019; Ziehn et al., 2020; Chikamoto and DiNezio, 2021). Similarly, surface ocean observation-based products typically calculate those fluxes using standard parameterizations (Wanninkhof, 1992, 2014; Fay et al., 2021). Importantly, the persistent uptake of anthropogenic CO<sub>2</sub> is lowering seawater pH and altering the carbonate system in nonlinear ways, further reducing the oceans’ capacity to absorb additional CO<sub>2</sub> (Moore et al., 2013). The mechanisms by which breaking waves and bubble injection enhance total gas transfer velocity have been investigated primarily through empirical analyses of observation-based products (e.g., Andreas et al., 2016; Blomquist et al., 2017; Reichl and Deike, 2020; Zhou et al., 2023). Rustogi et al. (2025) employed a fully coupled ocean-biogeochemistry modeling system based on the MOM6 ocean circulation model and the COBALTv2 biogeochemical module. Utilizing this framework, they simulated ocean dynamics, temperature, and carbon cycling processes – including dissolved inorganic carbon (DIC) and  $p\text{CO}_2$  – while explicitly ac-

counting for wave effects on air–sea CO<sub>2</sub> exchange. Furthermore, they identified a nonlinear  $p\text{CO}_2$  feedback mechanism within the coupled ocean–carbon system that regulates air–sea CO<sub>2</sub> flux. However, their configuration relies on ocean and biogeochemical modules that are strictly forced by offline atmospheric reanalysis and wave model outputs. This decoupled or unidirectional forcing approach is limited in capturing high-frequency, transient air–sea interactions (e.g., rapid wind shifts or wave breaking dynamics) that significantly modulate gas exchange. In contrast, our study introduces an online coupling framework that integrates wave effects directly into the flux simulation interface at each model time step. This online approach provides a critical advantage: it enables real-time, interactive feedback between wave dynamics and the seawater carbonate system, thereby capturing the nonlinear high-frequency physical controls on the air–sea CO<sub>2</sub> flux that offline-forced systems typically miss.

In this study, we investigate how breaking waves and bubble injection mechanisms influence air–sea CO<sub>2</sub> flux and the ocean’s buffering capacity. Specifically, we examine the biogeochemical responses driven by these physical mechanisms within the seawater carbonate–pH system. To achieve this, we utilize the Parallel Ocean Program version 2 (POP2; Smith et al., 2010) of the Community Earth System Model version 1.2.2 (CESM1.2.2; Hurrell et al., 2013) online-coupled with the wave module (Mellor et al., 2008) of the Princeton Ocean Model (POM; Blumberg and Mellor, 1987). The coupled configuration is referred to as the POP2–waves model.

The structure of this paper is organized as follows. Section 2 introduces the model, data, methodology, and experiments employed in this study, and defines twelve key regions of high air–sea CO<sub>2</sub> flux variability identified by the NOAA CarbonTracker data assimilation system (CT2022; Jacobson et al., 2023) to validate the model simulation. Section 3 evaluates the performance of the POP2–waves coupled model in simulating air–sea CO<sub>2</sub> flux and its interaction with the carbonate–pH system over the Western Pacific (WP; 160–180° E, 35–40° N) and the Equatorial Pacific (EP; 230–250° E, 0–5° S) regions. Section 4 compares the carbonate–pH dynamics of POP2–waves and B–CTL, examines the primary drivers of air–sea CO<sub>2</sub> exchange, and evaluates the uncertainties arising from the absence of a  $\Delta p\text{CO}_2$ -driven negative feedback. Finally, Sect. 5 presents the conclusions.

## 2 Data, model experiments, and methodology


### 2.1 Description of the model framework and experiments

In this study, we investigate the role of waves and bubble mechanisms in modulating the ocean’s biogeochemical response by comparing the control simulation (B–CTL) with the coupled POP2–waves model within the CESM1.2.2

framework (Fig. 1). The CESM simulation was conducted using a fully coupled component set over a 30-year period, with well-mixed greenhouse gases (CO<sub>2</sub>, CH<sub>4</sub>, N<sub>2</sub>O, etc.), ozone, and aerosols fixed to year-2000 values (the B\_2000\_CAM5 compset). The framework features a horizontal resolution of  $1.9^\circ \times 2.5^\circ$  for both the Community Atmosphere Model 5.3 (CAM5.3) and the Community Land Model version 4 (CLM4). In contrast, the ocean component (POP2; configured with 60 vertical layers) and the prognostic Los Alamos Sea Ice Model (CICE) share a nominal  $1^\circ$  horizontal resolution. State fields and fluxes are exchanged across these differing grids via the coupler (CPL) (Hurrell et al., 2013). POP2 is forced by CAM5.3 through CPL via four primary fields: (1) momentum fluxes (zonal and meridional wind stress and friction velocity), (2) heat fluxes (net shortwave radiation, sensible heat flux, longwave radiation, and heat flux from snow/ice melt), (3) freshwater fluxes (precipitation, evaporation, river runoff, and ice melt), and (4) surface winds (10 m zonal and meridional wind speeds). Furthermore, POP2 interacts with the Biogeochemical Elemental Cycling (BEC; Moore et al., 2013) module to regulate carbonate chemistry and air–sea CO<sub>2</sub> fluxes. This interaction is mediated via ocean dynamics (currents and sea surface height), the  $K$ -profile vertical mixing, and tracer transport, with wave-induced processes providing additional physical constraints. Within the BEC module, variations in DIC and nutrients (NO<sub>3</sub>, PO<sub>4</sub>, and Fe) are primarily controlled by biological processes: photosynthesis consumes DIC and nutrients, whereas respiration and remineralization return organic carbon to the DIC pool. Together with the wave module, these biogeochemical and physical processes jointly regulate the carbon cycle.

The control simulation (B–CTL), which served as the baseline for air–sea CO<sub>2</sub> flux and seawater acidification, utilized Eq. (1) with the gas transfer velocity parameterization from Wanninkhof (1992) (Eq. 2). Both the B–CTL and POP2–waves experiments were integrated under a present-day scenario (starting from the year 2000) to estimate air–sea CO<sub>2</sub> flux and pH variations. This experimental design allows us to isolate the impact of wave effects on carbonate chemistry and directly compare our results with the NOAA CT2022 data assimilation system. For the POP2–waves experiment, the wave module (Mellor et al., 2008) from the POM was online-coupled with POP2. This configuration incorporated bubble-mediated gas transfer – computed from a mechanistic model for air bubble entrainment at the breaking-wave scale (Deike and Melville, 2018) – and was integrated with the POP2 marine biogeochemistry module.

Specifically, the POP2–waves experiment adopts the advanced parameterization framework expanded by Reichl and Deike (2020), and Deike (2022). In this framework, both the non-bubble and bubble-mediated gas transfer velocity are explicitly resolved as a function of the ocean surface friction velocity ( $u^*$ , m s<sup>−1</sup>) and significant wave height ( $H_s$ , m). This parameterization effectively captures the main wave



**Figure 1.** Architecture diagram for POP2–waves experiment in CESM1.2.2 framework, all components adhere to the CESM1.2.2 framework, except for certain parts of the ocean component. The model components including components for the atmosphere [Community Atmosphere Model version 5 (CAM5)], land [Community Land Model version 4 (CLM4)], ocean [Parallel Ocean Program, version 2 (POP2)], along with its associated modules – the waves module (waves) and the Biogeochemical Elemental Cycling (BEC) module – sea ice [prognostic Los Alamos Sea Ice Model (CICE)], and the coupler (CPL). Note: (1) The wave-module variables are communicated between neighboring blocks via MPI, including zonal and meridional 10 m winds, wave radiation stress, subsurface momentum, and wave radiation and energy densities. (2) To transfer variables such as zonal and meridional 10 m winds and friction velocity (including sea surface and ice fraction) from the CPL to POP2. (3) The wave module receives variables from POP2, including the interpolation of depth-varying currents from  $z$ -coordinates (60 levels) to the wave module’s vertical sigma coordinates (21 levels), along with ocean grid information in the  $x$ ,  $y$ , and  $z$  directions. (4) The wave module outputs significant wave height and friction velocity from the CPL for calculating the bubble-mediated gas transfer velocity in Eq. (3).



effect, as well as solubility and diffusivity. To standardize the formulation for CO<sub>2</sub>, the total gas transfer velocity is expressed relative to a Schmidt number of 660. Following Deike and Melville (2018), it is calculated as the sum of the non-bubble ( $k_{wNB,660}$ ) and bubble ( $k_{wB,660}$ ) components:

$$k_{w,660} = k_{wNB,660} + k_{wB,660} = A_{NB} u^* \left( \frac{Sc}{660} \right)^{-1/2} + \frac{A_B}{k_0 R T_0} (u^*)^{5/3} (g H_s)^{2/3} \left( \frac{Sc}{660} \right)^{-1/2} \quad (3)$$

where  $A_{NB}$  is an empirical, nondimensional coefficient set to  $1.55 \times 10^{-4}$ ,  $A_B$  is a dimensional fitting coefficient ( $1.2 \times 10^{-5} \text{ s}^2 \text{ m}^{-2}$ ),  $R$  is the ideal gas constant ( $0.082 \text{ L atm mol}^{-1} \text{ K}^{-1}$ ),  $T_0$  is the sea surface temperature (SST, in K), and  $g$  is gravitational acceleration ( $9.806 \text{ m s}^{-2}$ ). Section 3.3 examines the relative contributions of the non-bubble and bubble components in simulating the air–sea CO<sub>2</sub> flux.

The marine carbonic acid system is a nonlinear, coupled system governed by CO<sub>2</sub> dissolution equilibrium,  $\left( \text{CO}_{2(aq)} + \text{H}_2\text{O} \xrightleftharpoons{K_H} \text{H}_2\text{CO}_{3(aq)} \right)$ , where the Henry’s constant ( $K_H$ ) for CO<sub>2</sub> in water at 25 °C is approximately  $3.4 \times 10^{-2} \text{ (m atm}^{-1}\text{)}$ , two-step dissociation reactions, and the conservation of the total dissolved inorganic carbon (DIC) and total alkalinity (TA), ultimately determining pH and the distribution of carbonate species. Carbonic acid (H<sub>2</sub>CO<sub>3</sub>) has

two hydrogen ions and dissociates in two steps:



The DIC can be expressed as

$$\begin{aligned} \text{DIC} &= [\text{H}_2\text{CO}_3] + [\text{HCO}_3^-] + [\text{CO}_3^{2-}] \\ &= K_H p\text{CO}_2^w + \frac{K_{a1} K_H p\text{CO}_2^w}{[\text{H}^+]} + \frac{K_{a1} K_{a2} K_H p\text{CO}_2^w}{[\text{H}^+]^2} \\ &= K_H p\text{CO}_2^w \left( 1 + \frac{K_{a1}}{[\text{H}^+]} + \frac{K_{a1} K_{a2}}{[\text{H}^+]^2} \right) \\ &= K_H p\text{CO}_2^w \left( [\text{H}^+]^2 + K_{a1} [\text{H}^+] + K_{a1} K_{a2} \right) / [\text{H}^+]^2 \end{aligned} \quad (6)$$

The marine carbonate system describes the distribution of DIC among its chemical species in seawater and is jointly governed by TA and pH. From Eq. (6), if the oceanic  $p\text{CO}_2^w$  and pH are known, the DIC can be determined; conversely,  $p\text{CO}_2^w$  can be derived from DIC. Following Dickson (1981), TA represents the net proton-neutralizing capacity of weak acid anions in seawater and is expressed as  $\text{TA} = [\text{HCO}_3^-] + 2[\text{CO}_3^{2-}] + [\text{B(OH)}_4^-] + [\text{OH}^-] + [\text{HPO}_4^{2-}] + 2[\text{PO}_4^{3-}] + [\text{SiO(OH)}_3^-] + [\text{NH}_3] + [\text{HS}^-] - [\text{H}^+] - [\text{H}_3\text{PO}_4]$ . In the POP2–waves framework, TA, DIC

and biogeochemical processes are used to calculate pH and  $p\text{CO}_2^w$ .

When TA increases, seawater becomes more alkaline, leading to a higher  $\text{CO}_3^{2-}$  concentration and enhanced buffering of hydrogen ions. Consequently,  $[\text{H}^+]$  decreases, causing pH to rise and boosting the ocean's capacity to absorb CO<sub>2</sub>. Conversely, when TA decreases, the buffering capacity weakens,  $[\text{H}^+]$  increases, and pH drops, rendering the ocean more acidic with a reduced capacity for CO<sub>2</sub> uptake.

## 2.2 Coupling POP2–waves Methodology

Except for the ocean component, all other components adhere strictly to the CESM1.2.2 framework. The wave module is forced by variables exchanged through the coupler, POP2, and a dedicated input initialization file. Specifically, in addition to the baseline component interactions in CESM1.2.2, the framework passes the zonal and meridional 10 m winds, along with surface friction velocity ( $u^*$ , accounting for both open-ocean and sea-ice fractions) from the CPL to evaluate the wave properties and the associated gas transfer velocity. To drive the wave module calculations, key physical variables are retrieved from POP2; this involves interpolating depth-variable currents from POP2  $z$ -coordinates (60 levels of zonal and meridional horizontal velocities) onto POM (waves) sigma-coordinates (21 vertical levels). This vertical mapping is required to compute both the depth-dependent wave radiation stresses and the specified spectrum. Additionally, spatial grid configurations in the  $x$ ,  $y$ , and  $z$  directions, along with time-step settings, are exchanged. The wave module further integrates external variables ( $F1$ – $F4$ ), which represent the spectrally averaged wave radiation stress components used to force the momentum equation.

Following Mellor et al. (2008), the significant wave height ( $H_s$ ) used in calculating wave radiation stresses represents the total wave energy integrated over the full spectrum, which inherently encompasses both locally wind-generated seas and remotely generated swells. During the simulation of wave energy and propagation, improper handling of parallel boundary formulations within the domain decomposition can induce artificial wave energy accumulation at the subdomain interfaces. To address this issue, this study mirrors the parallel computing infrastructure of the POP2 Message Passing Interface (MPI) protocol. Key physical and prognostic variables – including the zonal and meridional 10 m winds, wave radiation stress, subsurface momentum, and wave radiation and energy densities – are explicitly exchanged across processor subdomains to establish a robust, scalable online-coupling framework.

## 2.3 Model Validation

We evaluate the simulated spatial patterns of the air–sea CO<sub>2</sub> flux from both the uncoupled control simulation (B–CTL) and the coupled POP2–waves simulation against estimates

from the NOAA CT2022 data assimilation system. Unlike our forward, ocean-centric modeling frameworks, CT2022 optimizes the system from an atmospheric perspective, utilizing a top-down approach to infer net surface CO<sub>2</sub> fluxes by assimilating observed atmospheric CO<sub>2</sub> mole fractions. Consequently, this product captures the integrated signature of both anthropogenic emissions and natural carbon cycle variability to provide continuous global flux estimates spanning from 2000 to 2020.


Following the inversion framework established by Jacobson et al. (2007), NOAA CT2022 partitions the global ocean into 30 distinct basins to capture large-scale circulation and biogeochemical dynamics. This basin-based approach groups regions characterized by coherence in physical processes – such as major current systems, upwelling zones, and boundary mixing layers – as well as shared biogeochemical characteristics, including air–sea CO<sub>2</sub> exchange rates and biological productivity. Moreover, because numerous observational datasets and ocean carbon products are resolved at regional scales, a basin-scale aggregation ensures greater methodological consistency. We evaluate the seasonal variability and mean state of global air–sea CO<sub>2</sub> fluxes, specifically excluding regions characterized by seasonal flux sign reversals – a feature that can introduce substantial artifacts during model–data comparisons. Instead, our analysis focuses primarily on regions exhibiting robust, clear oceanic CO<sub>2</sub> flux signals. Recognizing that inherent uncertainties in atmospheric inversions stem from sub-grid-scale anthropogenic emissions and localized sea-ice melt, we follow the protocols of Fay et al. (2024) and exclude both the coastal ocean and high-latitude marginal ice zones from our model–data evaluation.

Based on the spatial patterns of oceanic CO<sub>2</sub> outgassing, uptake, and low-flux regimes characterized in the Pacific (Fig. 2c), this study delineates 12 key regions exhibiting high air–sea CO<sub>2</sub> flux variability as identified by NOAA CT2022 (Fig. 2d). These are selected from the 30 ocean basins defined in the CT2022 model documentation (Jacobson et al., 2023) for optimizing flux inversions. These oceanic domains are categorized into three major basins: (1) the Pacific Ocean, (2) the Indian Ocean, and (3) the Atlantic Ocean (Table 1). Within the Pacific basin, specific labels denote regions exhibiting the most pronounced oceanic CO<sub>2</sub> outgassing (the Equatorial Eastern Pacific, EEP) and uptake (the Northeastern Pacific, NEP).

## 3 Results

### 3.1 The climatological mean and seasonal variations of air–sea CO<sub>2</sub> flux

We utilize the traditional air–sea CO<sub>2</sub> flux bulk formula (Eq. 1) in CESM1.2.2 while also incorporating wave- and bubble-mediated effects via a gas transfer velocity pa-



**Figure 2.** Climatological mean air–sea CO<sub>2</sub> flux (mol m<sup>−2</sup> yr<sup>−1</sup>; shaded) and monthly standard deviation (SD; contour): (a) B–CTL; (b) POP2–waves; (c) NOAA CT2022: red dashed lines highlight the primary oceanic CO<sub>2</sub> outgassing and uptake regions in the Pacific Ocean, while black dashed lines mark regions with an air–sea CO<sub>2</sub> flux below −2 mol m<sup>−2</sup> yr<sup>−1</sup>; (d) based on NOAA CT2022, 12 key regions exhibiting significant air–sea CO<sub>2</sub> flux variability have been identified among the 30 ocean basins defined in the CT2022 model documentation (Jacobson et al., 2023) for optimizing flux inversions. These include: (1) Pacific Ocean: Northwestern Pacific (NWP), Eastern Equatorial Pacific (EEP), as well as Northeastern Pacific (NEP), South Pacific Ocean 1 (SPO1), and South Pacific Ocean 2 (SPO2); (2) Indian Ocean: North Indian Ocean (NIO), South Indian Ocean 1 (SIO1), and South Indian Ocean 2 (SIO2); (3) Atlantic Ocean: North Atlantic Ocean 1 (NAO1), North Atlantic Ocean 2 (NAO2), Equatorial Atlantic Ocean (EAO), and South Atlantic Ocean (SAO), (e) the air–sea CO<sub>2</sub> flux difference between B–CTL and NOAA CT2022, and (f) same as (e), but between POP2–waves and NOAA CT2022.

**Table 1.** Characteristics of the 12 key regions exhibiting prominent air–sea CO<sub>2</sub> flux variability across the three major ocean basins.

| Basins             | Key regions                                                                                                                                                  |
|--------------------|--------------------------------------------------------------------------------------------------------------------------------------------------------------|
| the Pacific Ocean  | the Northwestern Pacific (NWP), Eastern Equatorial Pacific (EEP), Northeastern Pacific (NEP), South Pacific Ocean 1 (SPO1), and South Pacific Ocean 2 (SPO2) |
| the Indian Ocean   | the North Indian Ocean (NIO), South Indian Ocean 1 (SIO1), and South Indian Ocean 2 (SIO2)                                                                   |
| the Atlantic Ocean | the North Atlantic Ocean 1 (NAO1), North Atlantic Ocean 2 (NAO2), Equatorial Atlantic Ocean (EAO), and South Atlantic Ocean (SAO)                            |

rameterized by  $u^*$ , SST, and  $H_s$  (Eq. 3). A comparison of the climatological means and monthly standard deviations indicates that the spatial distribution of the model-simulated air–sea CO<sub>2</sub> fluxes is broadly consistent with NOAA CT2022, with regional differences generally remain-

ing below 0.5 mol m<sup>−2</sup> yr<sup>−1</sup> across most analyzed sub-basins (Fig. 2a–f). Notably, the Pacific CO<sub>2</sub> outgassing and uptake regions simulated by the POP2–waves experiment show smaller range-normalized relative biases (Table 3) against NOAA CT2022 than the B–CTL baseline across most se-

lected regions. This improvement occurs despite POP2–waves having a higher monthly SD across both the Pacific and Indian Oceans.

To ensure an objective and reproducible classification, the 12 ocean regions were categorized into two distinct groups based on rigid statistical criteria, requiring either seasonal phasing consistency or model–data absolute bias magnitudes relative to the NOAA CT2022 baseline to be met:

1. Seasonal phasing consistency (temporal alignment): Regulated by the Pearson correlation coefficient ( $r$ ) between the simulated monthly flux cycles and NOAA CT2022. Regions assigned to Fig. 3 exhibit consistent seasonal phasing with significantly positive correlations ( $r \geq 0.50$ ), whereas regions in Fig. 4 display prominent phase mismatches or anti-correlated trends ( $r < 0.50$  or statistically non-significant  $r$ ).
2. Relative magnitudes of model–data biases (amplitude deviancy): Defined as the monthly averaged absolute deviation normalized by the overall peak-to-peak seasonal range (95th minus 5th percentile) of the NOAA CT2022 baseline. A threshold of 25 % was applied to further distinguish regions where structural amplitude offsets dominate over temporal phase alignment.


Based on these thresholds, the eight regions in Fig. 3 exhibit coherent seasonal phasing and relatively small monthly model–data biases. In contrast, the four regions in Fig. 4 (including the South Atlantic Ocean with  $r = 0.33$ ) display clear structural discrepancies, such as inverted seasonal trends or weak temporal alignment. Importantly, these temporal phase misalignments tend to amplify the calculated monthly model–data biases, as the model and data fluctuate in opposing directions throughout the annual cycle. This phase mismatch contributes to larger relative monthly mean biases; specifically, the model–data biases averaged across the four regions in Fig. 4 (33.7 % in B–CTL and 32.6 % in POP2–waves) are markedly larger than those in the eight regions of Fig. 3 (26.4 % in B–CTL and 22.0 % in POP2–waves).

In terms of the regional mean values of air–sea CO<sub>2</sub> flux, the POP2–waves simulation generally demonstrates closer structural agreement with NOAA CT2022 compared to the uncoupled B–CTL baseline, showing improvements in seven of the eight analyzed regions (with the exception of the Northwest Pacific (NWP); Fig. 3). Nevertheless, despite this larger mean offset in specific regions such as the NWP, POP2–waves successfully captures the autumn transition of oceanic CO<sub>2</sub> from a sink to a source as depicted in NOAA CT2022 – a feature that remains unrepresented in B–CTL. Table 3 presents the range-normalized relative magnitudes of model–data biases across eight selected regions, providing a robust metric to evaluate the simulated seawater carbonate system, pH and flux field performance against the NOAA CT2022 atmospheric inversion baseline. Overall,


the wave-coupled framework (POP2–waves) demonstrates a widespread reduction in systematic biases across the majority of the analyzed regions compared to the uncoupled control simulation (B–CTL). Specifically, the regional mean absolute deviation dropped noticeably in major ocean basins, with the most pronounced improvement captured in the Equatorial Eastern Pacific (EEP), where the relative bias was drastically attenuated by 25.0 % (from 71.5 % in B–CTL down to 46.5 % in POP2–waves). Notable bias reductions are also observed in the Northwest Pacific (NWP, down by 4.9 %), and South Indian Ocean Region 1 (SIO1, down by 4.1 %), confirming that incorporating wave-effect mechanisms successfully rectifies over- or under-estimations in air–sea carbon exchange dynamics. Conversely, the uncoupled baseline (B–CTL) exhibits slightly smaller biases in the Equatorial Atlantic Ocean (EAO) (11.9 % vs. 15.0 %), suggesting potential localized over-compensations in wave parameterizations within this sub-basin. Nonetheless, the reduction of relative biases across most other regions indicates that the online-coupled framework generally provides an improved representation of air–sea CO<sub>2</sub> fluxes.

The POP2–waves simulation generally exhibits a higher SD than B–CTL, reflecting an amplified seasonal variability (Figs. 3–4). Although these SD values typically remain below  $3 \text{ mol m}^{-2} \text{ yr}^{-1}$ , they peak during boreal winter, driven by intense oceanic CO<sub>2</sub> uptake in high-latitude regions of the Northern Hemisphere, such as the Northwestern and Northeastern Pacific, and the North Atlantic Ocean 1. In contrast, the Southern Hemisphere exhibits pronounced monthly SD along with clear oceanic CO<sub>2</sub> uptake during the austral winter (boreal summer), particularly across regions such as the South Pacific Ocean 1 (SPO1), South Indian Ocean 1 (SIO1), and South Atlantic Ocean (SAO). However, the monthly SD derived from NOAA CT2022 shows notable regional variations. In particular, both models underpredict the magnitude of variability observed in the Northwest Pacific (NWP), Equatorial Eastern Pacific (EEP), and North Atlantic Ocean 1 (NAO1).

Four specific sub-basins – South Atlantic Ocean (SAO), South Pacific Ocean Region 2 (SPO2), South Indian Ocean Region 2 (SIO2), and North Atlantic Ocean Region 2 (NAO2) – exhibit prominent model–data discrepancies when evaluated against NOAA CT2022 (Fig. 4). Notably, these systematic deviations are not attributable to wave-mediated processes or the parameterization of the CO<sub>2</sub> flux bulk formula. Within the NAO2, both the B–CTL and POP2–waves models simulate strong seasonal variations that mirror NAO1, characterized by substantial oceanic CO<sub>2</sub> uptake during the boreal winter; conversely, NOAA CT2022 displays the opposite seasonal trajectory in this sub-basin. A structurally similar mismatch is observed in the mid-latitude South Pacific (SPO2 and SIO2), where the simulated seasonal cycles of both models exhibit close synchronization with their respective low-latitude counterparts (SPO1 and



**Figure 3.** Climatological monthly mean CO<sub>2</sub> flux for eight highly correlated regions defined in Fig. 2d (NWP, EEP, NEP, SPO1, NIO, SIO1, NAO1, and EAO). Lines indicate values for B-CTL (black dashed), POP2-waves (red solid), and NOAA CT2022 (blue solid). Regional-average values are distinguished by a black font for B-CTL, red font for POP2-waves, and blue italics for NOAA CT2022. Monthly standard deviation (SD) is shown as bars: black hollow (B-CTL), red solid (POP2-waves), and light blue hollow (NOAA CT2022).



**Figure 4.** Same as Fig. 3, but for the four low-correlation characteristic regions (SAO, SPO2, SIO2, and NAO2), showing the climatological average of the simulated CO<sub>2</sub> flux compared with NOAA CT2022. Note: SAO, South Atlantic Ocean; SPO2, South Pacific Ocean Region 2; SIO2, South Indian Ocean Region 2; NAO2, North Atlantic Ocean Region 2.

SIO1) but differ noticeably from the inverse trend captured by the NOAA CT2022 atmospheric inversion framework.

### 3.2 The relationship between $U_{10}$ and $k_{w,660}$ in two regions with contrasting wind speed regimes


To demonstrate that the sea-state-dependent gas transfer velocity (Deike and Melville, 2018) provides a more suitable estimate of air–sea CO<sub>2</sub> flux than the wind-only formulation (Wanninkhof, 1992) under high-wind conditions ( $U_{10} > 10 \text{ m s}^{-1}$ ), we selected the Western Pacific (WP) and the Equatorial Pacific (EP) regions due to their contrasting wind regimes. The WP region experiences a high frequency of high-wind conditions (exceeding 30 %), whereas wind speeds in the EP region consistently remain below this threshold. Because both regions exhibit stable and significant oceanic CO<sub>2</sub> outgassing or uptake, they enable a clear comparative analysis of the correlation between  $k_{w,660}$  and  $U_{10}$  (Fig. 5a, b).

Generally, the B–CTL data shows higher  $R^2$  but lower  $k_{w,660}$  values compared to those in POP2–waves across both the WP and EP. The POP2–waves experiment exhibits a slightly lower  $R^2$  than B–CTL due to greater deviations between  $k_{w,660}$  and  $U_{10}$ , particularly at higher surface wind speeds. This pattern is consistent with the findings of Gutiérrez-Loza et al. (2022) under conditions where

$H_s > 1.5 \text{ m}$ . The relationship between  $k_{w,660}$  and  $U_{10}$  was evaluated under two scenarios: baseline monthly climatological means ( $H_s < 1.5 \text{ m}$ ) and high-resolution daily data filtered for  $H_s > 1.5 \text{ m}$  (enhanced conditions) aggregated into 30-year monthly averages. Notably, both approaches substantially diminish the differences in  $k_{w,660}$  at identical  $U_{10}$  values.

However, short-term (30 min) observations indicate that gas transfer velocity can be up to twice as high under similar wind speeds when  $H_s > 1.5 \text{ m}$  (Gutiérrez-Loza et al., 2022). Divergence between the models becomes significant when  $U_{10}$  exceeds  $12 \text{ m s}^{-1}$  over the WP region (Fig. 5a), whereas over the EP region, the convergence of  $k_{w,660}$  estimates is most notable around  $U_{10} = 9 \text{ m s}^{-1}$  (Fig. 5b), suggesting that mid-range wind speeds dominate the EP flux signal.

As shown in Fig. 5c, the enhanced WP cases ( $H_s > 1.5 \text{ m}$ ) have data available for all months except August and October, during these available months, their  $k_{w,660}$  values consistently exceed both the all-data POP2–waves and B–CTL averages. In contrast, the EP region exhibits  $H_s > 1.5 \text{ m}$  data only during the summer months (July to October), where corresponding  $k_{w,660}$  values also exceed both all-data POP2–waves and B–CTL averages. The monthly SD indicates no significant differences between the enhanced ( $H_s > 1.5 \text{ m}$ ) data and the standard POP2–waves simulations over the WP, with the  $k_{w,660}$  discrepancies between the two datasets re-



**Figure 5.** Scatter plots of monthly averaged gas transfer velocity ( $k_{w,660}$ ) versus  $U_{10}$  for (a) the Western Pacific (WP; 160–180°E, 35–40°N) and (b) the Eastern Pacific (EP; 230–250°E, 0–5°S). Gray open circles, green open squares, and blue dots with black edges represent B-CTL, POP2-waves data for  $H_s < 1.5$  m, and  $H_s > 1.5$  m, respectively. Corresponding simple linear regressions (SLRs) and squared correlation coefficients ( $R^2$ ) are shown with black dotted lines/text (B-CTL), orange solid lines/text (POP2-waves,  $H_s < 1.5$  m), and red solid lines/text (POP2-waves,  $H_s > 1.5$  m). Panels (c) and (d) display the monthly mean values of  $k_{w,660}$  for WP and EP, respectively. Red and orange solid lines/text indicate POP2-waves ( $H_s > 1.5$  m) and the average of all POP2-waves data, while the black dotted line/text represents B-CTL. The bar graph illustrates the standard deviation (SD) for each month over the WP and EP, red hollow bars for POP2-waves ( $H_s > 1.5$  m), orange for average POP2-waves, and black for B-CTL.


maintaining minimal from January to May because the average  $H_s$  during this period consistently remains below 1.5 m. However, over the EP, a pronounced discrepancy is observed between the  $H_s > 1.5$  m condition and the all-data POP2-waves average from July to September.

### 3.3 Regression of CO<sub>2</sub> flux against $U_{10}$ under high-wind conditions

The wind-only parameterization of gas transfer velocity tends to underestimate values under high-wind conditions ( $U_{10} > 10$  m s<sup>-1</sup>) or high significant wave heights ( $H_s > 1.5$  m) (e.g., Gutiérrez-Loza et al., 2022; Zhou et al., 2023). Under these high-wind conditions, the monthly mean air–sea CO<sub>2</sub> fluxes derived from the B-CTL and POP2-waves simulations diverge notably within the Western Pacific region (WP) (Fig. 6), highlighting key discrepancies between the wind-only and wave-modified  $k_{w,660}$  formulations. The data were filtered to include only grid points that met this high-wind threshold before being regionally averaged. Because not all months contained grid points satisfying the threshold, a final subset of 154 data points was utilized for

the linear regression analysis. These high-wind conditions over the WP region occur primarily in winter (accounting for 42 % of the valid data), when the concurrent low sea surface temperatures further enhance CO<sub>2</sub> solubility. The slopes of the regression equations ( $-0.77$  mol m<sup>-3</sup> s yr<sup>-1</sup> for B-CTL and  $-1.40$  mol m<sup>-3</sup> s yr<sup>-1</sup> for POP2-waves) alongside the mean CO<sub>2</sub> fluxes ( $Y_{\text{avg}}$ :  $-6.35$  mol m<sup>-2</sup> yr<sup>-1</sup> for B-CTL and  $-6.97$  mol m<sup>-2</sup> yr<sup>-1</sup> for POP2-waves) both indicate that POP2-waves exhibits a higher sensitivity to wind speed, leading to stronger oceanic CO<sub>2</sub> uptake across the WP domain under elevated wind speeds.

The correlation coefficient ( $R$ ) of the POP2-waves simulation is higher than that of B-CTL, indicating that the sea-state-dependent  $k_{w,660}$  formulation provides a more realistic representation of the CO<sub>2</sub> flux than the wind-only  $k_{w,660}$  scheme under high-wind conditions. This finding aligns with the observations of Gutiérrez-Loza et al. (2022) and Zhou et al. (2023). The average air–sea CO<sub>2</sub> flux in B-CTL over the WP region during DJF and JJA is  $-0.021$  and  $-0.004$  mol m<sup>-2</sup> yr<sup>-1</sup>, respectively. Meanwhile, POP2-waves exhibits only slight discrepancies relative to B-CTL during these respective seasons, with differences ranging be-



**Figure 6.** Scatter plots of air–sea CO<sub>2</sub> flux versus  $U_{10}$  with regression lines over the Western Pacific (160–180°E, 35–40°N) under high-wind conditions ( $U_{10} > 10 \text{ m s}^{-1}$ ) for (a) B-CTL and (b) POP2-waves experiments. Black dots indicate monthly mean values from each experiment. Blue lines denote the 95 % and 5 % confidence limits of the mean response, while red lines denote the 95 % and 5 % prediction intervals. The regression equation ( $Y = mx + b$ ), Pearson correlation coefficient ( $r$ ), mean CO<sub>2</sub> flux ( $Y_{\text{avg}}$ ), and standard deviation ( $Y_{\text{SD}}$ ) are shown to the right of the legend.


tween 0.001 and 0.002 mol m<sup>-2</sup> yr<sup>-1</sup> in CO<sub>2</sub> uptake (data not shown). This contrast indicates that under high-wind conditions, the episodic air–sea CO<sub>2</sub> flux is significantly greater than the seasonal baseline averages.

### 3.4 Analysis of surface fields and $k_{\text{w},660}$ component in POP2-waves and B-CTL

The spatial co-variations of sea level pressure (SLP),  $U_{10}$ , and  $H_s$  underscore their combined role as key meteorological and wave drivers steering the regional  $k_{\text{w},660}$  patterns (Fig. 7a, b). The region with  $H_s > 1.5 \text{ m}$  is generally situated where  $U_{10} > 8 \text{ m s}^{-1}$  and lies south of the high-SLP region. The spatial patterns of climatological  $U_{10}$  and  $H_s$  align closely with the findings of Reichl and Deike (2020) (Fig. 7a, b). Furthermore, a dominance of bubble-mediated transfer in the POP2-waves simulation, where the  $k_{\text{wB},660}/k_{\text{w},660}$  ratio exceeds 0.5, primarily occurs in regions characterized by  $H_s > 1.5 \text{ m}$  (Fig. 7b). However, an elevated  $H_s$  does not always correspond to a higher  $k_{\text{wB},660}/k_{\text{w},660}$  ratio in the POP2-waves simulation; this decoupling is explicitly observed in the North Pacific (Fig. 7b).

Deike and Melville (2018) demonstrated that the bubble contribution to CO<sub>2</sub> transfer exceeds 40 % when the  $U_{10} \geq 10 \text{ m s}^{-1}$ . In our POP2-waves simulation, the bubble fraction ( $k_{\text{wB},660}/k_{\text{w},660}$ ) accounts for approximately 38 % of total gas transfer velocity, which is slightly higher than the  $\sim 30 \%$  reported by Reichl and Deike (2020). This discrepancy may be attributed to differences in the  $A_B$  scaling coefficients (Eq. 3), which are derived from field data, where gas transfer velocity is estimated from eddy covariance flux measurements (Reichl and Deike, 2020; Brumer et al., 2017).

The structural differences in  $k_{\text{w},660}$  (POP2-waves – B-CTL) reveal that the most pronounced discrepancies are primarily distributed in regions characterized by elevated  $H_s$ , particularly in the Southern and South Indian Oceans (Fig. 7c). Although  $U_{10}$  in the Indian Ocean and tropical Pacific is relatively low ( $< 5 \text{ m s}^{-1}$ ; Fig. 7a), the  $k_{\text{w},660}$  values simulated by POP2-waves remain approximately 9 cm hr<sup>-1</sup> higher than those in B-CTL. In these regions, this wave-driven enhancement of the total gas transfer velocity is predominantly governed by the non-bubble-mediated component ( $k_{\text{wNB},660}$ ; Fig. 7d). Although  $H_s$  is also elevated in the Southern Ocean (Fig. 7b), it does not contribute substantially to the  $k_{\text{w},660}$  difference between POP2-waves and B-CTL; this is because the high  $U_{10}$  in this region (Fig. 7a) already yields high gas transfer velocities through the empirical formulation of Wanninkhof (1992). The spatial patterns of both the B-CTL  $k_{\text{w},660}$  and the wave-induced  $k_{\text{w},660}$  discrepancies (POP2-waves – B-CTL) closely align with the findings of Reichl and Deike (2020) (Fig. 7c). Overall, the two simulations exhibit consistent geographic distributions, yielding 30-year domain-wide mean gas transfer velocities of 17.0 cm hr<sup>-1</sup> for B-CTL and 22.7 cm hr<sup>-1</sup> for POP2-waves, despite their reliance on distinct input parameters ( $U_{10}$  versus  $u^*$  and  $H_s$ ). Aside from the polar regions, the total gas transfer velocity in the POP2-waves coupled model exceeds that in B-CTL, with the most pronounced differences localized in the tropics (Fig. 7c). To elucidate the driving mechanisms behind this global enhancement, it is necessary to quantify whether the non-bubble ( $k_{\text{wNB},660}$ ) or bubble-mediated ( $k_{\text{wB},660}$ ) component dominates the total gas transfer velocity across the different ocean basins. This partitioning demonstrates that bubble-mediated



**Figure 7.** Effects of atmospheric factors and wave components on average gas transfer velocity: (a) Shaded areas represent 10 m wind speed ( $U_{10}$ ,  $\text{m s}^{-1}$ ) and contours show sea level pressure (SLP, mb); (b) color denotes the ratio of bubble-mediated ( $k_{\text{WB},660}$ ) to POP2-waves  $k_{\text{WB},660}$  components, with contours representing significant wave height  $H_s$  (m) and thick lines marking  $H_s = 1.5$  m; (c) shading represents  $k_{\text{WB},660}$  (B-CTL) and contours represent the difference  $k_{\text{WB},660}$  (POP2-waves – B-CTL) in  $\text{cm hr}^{-1}$ ; (d) shading shows the non-bubble-mediated component ( $k_{\text{WNB},660}$ ) and contours show the difference ( $k_{\text{WNB},660} - k_{\text{WB},660}$ ) in  $\text{cm hr}^{-1}$ , with the thick contour lines at 0.

processes dominate ( $k_{\text{WB},660} > k_{\text{WNB},660}$ ) under high- $H_s$  conditions, whereas non-bubble mechanisms prevail ( $k_{\text{WB},660} < k_{\text{WNB},660}$ ) in light-wind zones where  $U_{10} < 7 \text{ m s}^{-1}$  (Fig. 7).


### 3.5 Mean State and model differences in $p\text{CO}_2^{\text{w}}$ and surface pH

Integrating wave dynamics induces a pronounced spatial reorganization of both  $p\text{CO}_2^{\text{w}}$  and ocean surface pH over the 30-year climatological period (Fig. 8). Specifically, the baseline B-CTL simulation shows that high  $p\text{CO}_2^{\text{w}}$  is concentrated in the equatorial eastern Pacific (EEP), contributing to a larger oceanic CO<sub>2</sub> source (Fig. 8a). In contrast, low  $p\text{CO}_2^{\text{w}}$  is found in high-latitude regions of the Pacific and Atlantic (e.g., NWP, NEP, NAO1, and NAO2), driving strong oceanic CO<sub>2</sub> uptake (Fig. 8a). Over the study period, the climatological standard deviation of  $p\text{CO}_2^{\text{w}}$  in B-CTL ranges between 10 and 30 ppm (Fig. 8a).

In the CESM1 framework of the B\_2000\_CAM5 compsets, atmospheric  $p\text{CO}_2^{\text{a}}$  is held constant (367 ppm). The  $p\text{CO}_2^{\text{w}}$  in the Eastern Pacific exceeds 440 ppm, indicating an air–sea  $\Delta p\text{CO}_2$  greater than 73 ppm in B-CTL (109.8 and 92.7 ppm during DJF and JJA, respectively), whereas the  $p\text{CO}_2^{\text{w}}$  in the Western Pacific is below 320 ppm, indicating an air–sea  $\Delta p\text{CO}_2$  lower than –47 ppm (–53.4 and –31.4 ppm during DJF and JJA, respectively). Notably, POP2-waves re-

duces the high  $p\text{CO}_2^{\text{w}}$  by more than 10 ppm over the Eastern Equatorial Pacific and increases the low  $p\text{CO}_2^{\text{w}}$  in other regions (Fig. 8b). Consequently, in the oceanic CO<sub>2</sub> outgassing (uptake) regions, the POP2-waves coupled model decreases (increases)  $p\text{CO}_2^{\text{w}}$ , thereby reducing the magnitude of the air–sea  $\Delta p\text{CO}_2$ . The climatological SD of the  $p\text{CO}_2^{\text{w}}$  discrepancy between POP2-waves and B-CTL displays marked spatial variability over the Eastern Equatorial Pacific, whereas changes across other regions remain relatively minor.

The biogeochemical module of POP2 (BEC) updates TA in each grid cell by solving a coupled transport–reaction equation (Moore et al., 2013). The average pH of the ocean is currently around 8.1 in oceanic CO<sub>2</sub> uptake regions (e.g., the Western Pacific) and 8.0 in outgassing regions (e.g., the Equatorial Pacific; Fig. 8c), reflecting slightly alkaline conditions. Within the DIC system, bicarbonate ions ( $\text{HCO}_3^-$ ) are the dominant species because background seawater pH falls strictly between  $pK_{\text{a}1}$  and  $pK_{\text{a}2}$ . However, continued uptake of atmospheric CO<sub>2</sub> drives a decline in pH – a process termed ocean acidification. Our study highlights that wave-induced  $k_{\text{WB},660}$  significantly influences the marine CO<sub>2</sub> system, altering alkalinity, pH, and carbonate speciation. These chemical shifts, in turn, provide a feedback mechanism that modulates the overall dynamics of the air–sea CO<sub>2</sub> flux. Con-



**Figure 8.** Spatial distributions of the 30-year climatological averages and model differences for  $\Delta p\text{CO}_2^w$  (ppm; panels **a**, **b**) and ocean surface pH (panels **c**, **d**). Climatological means are represented by shading, and their corresponding standard deviation (SD) is superimposed as contours. Panels (**a**) and (**c**) present baseline results from the B-CTL simulation. Panels (**b**) and (**d**) depict the structural differences between the POP2-waves and B-CTL experiments (POP2-waves minus B-CTL), where stippling indicates statistical significance at the 95 % confidence level based on a Student's *t*-test.

sequently, wave-driven perturbations in  $k_{w,660}$  directly alter both  $\Delta p\text{CO}_2$  and net air–sea CO<sub>2</sub> fluxes within the Earth System Model framework.

Ocean pH and surface ocean  $p\text{CO}_2^w$  are strongly negatively correlated (Macovei et al., 2021). Consequently, low pH values are observed in the Equatorial Eastern Pacific (Fig. 8c), as a direct result of the elevated DIC and  $p\text{CO}_2^w$  concentrations characteristic of that region (Fig. 8a). Conversely, elevated pH conditions persist in domains with lower  $p\text{CO}_2^w$ , a signature most prominent throughout high-latitude waters in both the Pacific and Atlantic Basins. Rustogi et al. (2025) indicate that accelerated equilibration of oceanic  $p\text{CO}_2$  in the wind–wave–bubble simulation, driven by enhanced gas exchange, ultimately reduces the magnitude of the air–sea  $\Delta p\text{CO}_2$ . This  $p\text{CO}_2$  feedback is strongest in the extratropics, where it suppresses CO<sub>2</sub> uptake. Our POP2-waves experiment reveals a consistent response, characterized by increases in both oceanic  $p\text{CO}_2$  (Fig. 8b) and hydrogen ion concentration ( $[\text{H}^+]$ ), which are accompanied by a corresponding decline in pH (Fig. 8d) throughout the extratropical ocean basins. Accounting for wave effects results in simulated pH shifts of approximately +0.01 (−0.01) in oceanic CO<sub>2</sub> outgassing (uptake) regions (Fig. 8d). This suggests that incorporating this mechanism alleviates acidification in the equatorial Pacific, whereas it intensifies ocean acidification in the high-latitude domains of both the Pacific and Atlantic Oceans (Fig. 8d). Rustogi et al. (2025) also

showed that the effects of  $\Delta k_w$  (the difference between wave-induced  $k_{w,660}$  and  $k_{w,660}$  based on  $U_{10}$ ) and  $\Delta p\text{CO}_2$  (POP2-waves − B-CTL) partially offset each other. However, because the  $\Delta k_w$  effect is generally 20 %–30 % stronger, it dominates the counteracting  $\Delta p\text{CO}_2$  effect, explaining the modest changes observed in both oceanic outgassing and up-take CO<sub>2</sub> fluxes within the wave-effect simulation.

## 4 Discussion

### 4.1 Impact of wave-dependent $k_w$ on global air–sea CO<sub>2</sub> flux: a comparison between POP2-waves and standard Earth System Models

Several investigations have parameterized the air–sea CO<sub>2</sub> flux using Eq. (1), yet current  $k_{w,660}$  parameterizations within prominent global frameworks still rely exclusively on the neutral 10 m wind speed ( $U_{10}$ ) as seen in models such as MPI-ESM1.2 (Mauritsen et al., 2019), CESM2 (Danabasoglu et al., 2020), NorESM2 (Seland et al., 2020), UKESM1 (Sellar et al., 2019), MIROC6 (Chikamoto and DiNezio, 2021), CMCC-ESM2 (Lovato et al., 2022), and CanESM5 (Sigmond et al., 2023) (Table 2). Concurrently, pioneering efforts have begun quantifying wave-induced effects on both air–sea CO<sub>2</sub> flux and long-term ocean carbon storage by incorporating coupled wind–wave–bubble gas transfer formulations into ocean general circulation models (OGCMs), such

as MOM6–COBALTv2 (Rustogi et al., 2025) and NEMO–PISCES (Wu et al., 2025). Importantly, Wu et al. (2025) emphasize that integrating these wave-mediated boundary layer dynamics into fully coupled Earth System Models is crucial to reducing systemic uncertainties in global carbon cycle projections.

Several studies have incorporated wave effects following the parameterization of Deike and Melville (2018), utilizing the Surface Ocean CO<sub>2</sub> Atlas (SOCAT) database (Bakker et al., 2016) to calculate diagnostic air–sea CO<sub>2</sub> fluxes. In these offline configurations, however, the computed air–sea CO<sub>2</sub> flux is treated independently of  $p\text{CO}_2^w$  dynamics and surface pH evolution, lacking any interactive feedback mechanisms between the atmosphere and the upper-ocean carbonate system.

Rustogi et al. (2025) utilized an ocean-biogeochemistry system forced by offline atmospheric reanalysis and wave model outputs. While comprehensive, such an offline-forced setup is limited in its ability to capture high-frequency, episodic, and synchronous interactions between physical wave states and surface water chemistry during rapid weather transitions. In contrast, our study implements an online coupling framework. The key added value of our setup is its capacity to simulate step-by-step interactive feedbacks where wave properties dynamically alter the gas transfer velocity ( $k_{w,660}$ ) and physical mixing in real-time. The coupled behavior of POP2–waves includes  $\Delta p\text{CO}_2$ -driven negative feedbacks, which fundamentally differs from traditional obs-based products that treat wave-induced  $k_{w,660}$  and  $\Delta p\text{CO}_2$  as independent variables when estimating air–sea CO<sub>2</sub> flux (e.g., Reichl and Deike, 2020). This online coupling mechanism also contrasts with global ocean models forced offline by atmospheric reanalysis and wave model outputs (e.g., Rustogi et al., 2025), where such interactive feedbacks are often omitted. This approach enables our model to resolve non-linear, high-resolution modulations of air–sea CO<sub>2</sub> exchange during rapid dynamic events – effects that are typically muted in offline configurations.

#### 4.2 Impact of air–sea $\Delta p\text{CO}_2$ , $k_{w,660}$ , SST, and pH on air–sea CO<sub>2</sub> flux

To evaluate which driver exerts the most direct control on air–sea CO<sub>2</sub> flux, Fig. 9 displays the spatial distributions of the linear regression coefficients (LRCs) derived from univariate (one-on-one) linear regressions between the flux and various standardized variables: air–sea  $\Delta p\text{CO}_2$  (Fig. 9a),  $k_{w,660}$  (Fig. 9b), SST (Fig. 9c), and pH (Fig. 9d). To remove the confounding artifacts of differing physical units and enable a direct comparison on a common scale, both the air–sea CO<sub>2</sub> flux and the independent variables were standardized into anomalies (with a mean of 0 and a standard deviation of 1) prior to the regression analysis. The statistical significance of these regressed relationships was subsequently evaluated using a Student's *t*-test at the 95 % confidence level.

Because each standardized regression was performed individually with a single independent variable, the resulting LRCs are equivalent to Pearson correlation coefficients (*r*), constraining their values strictly between −1 and +1. Here, we do not delve into the intricate chemical and biochemical feedback mechanisms governing carbonate species. Nevertheless, it is worth noting that the collective influence of carbonate composition – specifically TA and dissolved inorganic carbon (DIC) – on  $p\text{CO}_2^w$  variability has been shown to outweigh that of solubility changes driven by sea surface salinity and SST (e.g., Koseki et al., 2023).

Among these drivers, air–sea  $\Delta p\text{CO}_2$  and air–sea CO<sub>2</sub> flux exhibit the strongest positive LRCs, with coefficients ranging from 0.6 to 0.8 across the global ocean; this underscores the ocean's role as a net CO<sub>2</sub> source or sink depending on the sign of the air–sea  $\Delta p\text{CO}_2$  gradient. Additionally, wave effects increase the LRCs by approximately 0.1 to 0.2, particularly within regions of intense oceanic CO<sub>2</sub> outgassing or uptake. Clear positive (negative) LRCs exist between  $k_{w,660}$  and the air–sea CO<sub>2</sub> flux in oceanic CO<sub>2</sub> outgassing (uptake) regions. Furthermore, wave effects reduce the absolute LRCs between  $k_{w,660}$  and air–sea CO<sub>2</sub> flux in both oceanic CO<sub>2</sub> outgassing (uptake) regions (Fig. 9b). This indicates that once interactions between the modeled CO<sub>2</sub> flux and the ocean carbonate–pH system are considered, the wave-influenced CO<sub>2</sub> flux becomes less correlated with  $k_{w,660}$  and more closely linked to air–sea  $\Delta p\text{CO}_2$ . This shift can be attributed to the limitations of wind-only parameterizations; as indicated by Zhou et al. (2023), wind-only formulas tend to underestimate gas transfer velocities when  $U_{10}$  exceeds  $10 \text{ m s}^{-1}$ , where intense wave breaking and high significant wave height substantially boost air–sea gas exchange. Furthermore, Johansson et al. (2022) reported that  $H_s$  generally increases with wind speed, but the relationship is strongly modulated by wind direction and regional conditions, leading to spatially variable and nonlinear wind–wave coupling.

In contrast to the former, there are clear positive (negative) LRCs between SST and CO<sub>2</sub> flux in oceanic CO<sub>2</sub> uptake (outgassing) regions (Fig. 9c). A negative correlation exists between  $k_{w,660}$  and SST, reflecting the latitudinal gradient where colder, high-latitude waters are characterized by more intense sea states and elevated  $k_{w,660}$  values. This pattern occurs because lower temperatures at high latitudes increase CO<sub>2</sub> solubility, thereby enhancing oceanic uptake. From a thermodynamic perspective, higher SST enhances molecular diffusion while lowering seawater viscosity, thereby reducing the Schmidt number ( $Sc = \nu/D$ , defined as the ratio of kinematic viscosity ( $\nu$ ) to the molecular diffusion coefficient ( $D$ )). Because  $k_{w,660}$  is parameterized as a function of  $Sc^{-0.5}$ , a decrease in  $Sc$  theoretically elevates  $k_{w,660}$ . Separately, Fay et al. (2024) attributed the regional discrepancies and variations in air–sea CO<sub>2</sub> fluxes to a combination of factors, including choices in wind speed products, SST datasets, biological carbon uptake, and the parameterization of gas transfer velocity. Wave effects only slightly reduce the LRCs

**Table 2.** Intercomparison of Earth System Model for estimating air–sea CO<sub>2</sub> flux

| Earth System Model             | Ocean components            | Wave-informed air–sea CO <sub>2</sub> flux | $k_{w,660}$ parameterization          | Reference                                             |
|--------------------------------|-----------------------------|--------------------------------------------|---------------------------------------|-------------------------------------------------------|
| CESM1 <sup>1</sup>             | POP2 <sup>2</sup>           | no                                         | $U_{10}$ (Wanninkhof, 1992)           | Chikamoto and DiNezio (2021), Chikamoto et al. (2023) |
| CESM2 <sup>3</sup>             | POP2 <sup>2</sup>           | no                                         | $U_{10}$ (Wanninkhof, 2014)           | Danabasoglu et al. (2020)                             |
| NorESM2 <sup>4</sup>           | BLOM <sup>5</sup>           | no                                         | $U_{10}$ (Wanninkhof, 2014)           | Seland et al. (2020), Tjiputra et al. (2020)          |
| MPI-ESM1.2 <sup>6</sup>        | MPIOM1.6 <sup>7</sup>       | no                                         | $U_{10}$ (Wanninkhof, 2014)           | Mauritsen et al. (2019), Liu et al. (2021)            |
| ACCESS-ESM1.5 <sup>8</sup>     | MOM5 <sup>9</sup>           | no                                         | $U_{10}$ (Wanninkhof, 1992)           | Ziehn et al. (2020)                                   |
| CMCC-ESM2 <sup>10</sup>        | NEMO v3.6 <sup>11</sup>     | no                                         | $U_{10}$ (Wanninkhof, 1992)           | Lovato et al. (2022)                                  |
| CanESM5 <sup>12</sup>          | CanNEMO <sup>13</sup>       | no                                         | $U_{10}$ (Wanninkhof, 1992)           | Sigmond et al. (2023)                                 |
| GFDL-ESM4.1 <sup>14</sup>      | MOM6 <sup>15</sup>          | no                                         | $U_{10}$ (Wanninkhof, 2014)           | Stock et al. (2020), Roobaert et al. (2024)           |
| IPSL-CM6A-LR <sup>16</sup>     | NEMO-PISCES <sup>17</sup>   | no                                         | $U_{10}$ (Wanninkhof, 1992)           | Boucher et al. (2020)                                 |
| MIROC-ES2L <sup>18</sup>       | COCO 4.0 <sup>19</sup>      | no                                         | $U_{10}$ (Wanninkhof, 1992)           | Hajima et al. (2020)                                  |
| UKESM1 <sup>20</sup>           | NEMO v3.6 <sup>11</sup>     | no                                         | $U_{10}$ (Wanninkhof, 1992)           | Sellar et al. (2019), Yool et al. (2021)              |
| CNRM-ESM2-1 <sup>21</sup>      | NEMO-PISCES <sup>17</sup>   | no                                         | $U_{10}$ (Wanninkhof, 1992)           | S  f  rian et al. (2019)                              |
| _22                            | NEMO-PISCES <sup>17</sup>   | yes                                        | $H_s, u^*$ (Deike and Melville, 2018) | Wu et al. (2025)                                      |
| _22                            | MOM6-COBALTv2 <sup>23</sup> | yes                                        | $H_s, u^*$ (Deike and Melville, 2018) | Rustogi et al. (2025)                                 |
| CESM1-POP2–waves <sup>24</sup> | POP2–waves coupled model    | yes                                        | $H_s, u^*$ (Deike and Melville, 2018) | This study                                            |

Note: <sup>1</sup> CESM1: the Community Earth System Model version 1 of the National Center for Atmospheric Research (NCAR); <sup>2</sup> POP2: the Parallel Ocean Program version 2 of NCAR; <sup>3</sup> CESM2: CESM version 2; <sup>4</sup> NorESM2: The Norwegian Earth System Model version 2; <sup>5</sup> BLOM: Bergen Layered Ocean Model; <sup>6</sup> MPI-ESM1.2: the Max Planck Institute for Meteorology Earth System Model version 1.2; <sup>7</sup> MPIOM1.6: the Max-Planck Institute Ocean Model version 1.6; <sup>8</sup> ACCESS-ESM1.5: the Australian Community Climate and Earth System Simulator form an Earth System Model version 1.5; <sup>9</sup> MOM5: the GFDL Modular Ocean Model version 5; <sup>10</sup> CMCC-ESM2: the Euro-Mediterranean Centre on Climate Change (CMCC) Earth System Model version 2; <sup>11</sup> NEMO v3.6: Nucleus for European Modelling of the Ocean version 3.6; <sup>12</sup> CanESM5: the Canadian Earth System Model version 5; <sup>13</sup> CanNEMO: NEMO version 3.4 modified for CanESM; <sup>14</sup> GFDL-ESM4.1: the Geophysical Fluid Dynamics Laboratory’s Earth System Model 4.1; <sup>15</sup> MOM6: the GFDL Modular Ocean Model version 6; <sup>16</sup> IPSL-CM6A-LR : version 6 of the Institut Pierre-Simon Laplace (IPSL) climate model; <sup>17</sup> NEMO-PISCES: Nucleus for European Modelling of the Ocean, Pelagic Interaction Scheme for Carbon and Ecosystem Studies ocean general circulation and biogeochemistry model; <sup>18</sup> MIROC-ES2L: the Model for Interdisciplinary Research on Climate, Earth System version 2; <sup>19</sup> COCO 4.0: CCSR (Center for Climate System Research) Ocean Component Model version 4.0; <sup>20</sup> UKESM1: the U.K. Earth System Model; <sup>21</sup> CNRM-ESM2-1: the Earth system (ES) model of second generation developed by the Centre National de Recherches M  t  orologiques (CNRM); <sup>22</sup> The simulations were conducted using global ocean-only models rather than full Earth System Models; <sup>23</sup> MOM6-COBALTv2: Geophysical Fluid Dynamics Laboratory global ocean model (Modular Ocean Model, MOM6) coupled with sea ice and biogeochemistry (Carbon, Ocean Biogeochemistry and Lower Trophics version 2, COBALTv2); <sup>24</sup> CESM1-POP2–waves: POP2–waves coupled model based on CESM1.

between SST and the air–sea CO<sub>2</sub> flux – by approximately 0.1 – within the tropics, with negligible impacts observed in other regions. The strong negative correlation between ocean pH and  $p\text{CO}_2^w$  is dictated by the acid dissociation equilibria of the marine carbonate system (Fig. 8a, c), which govern the fundamental relationship between dissolved CO<sub>2</sub> and hydrogen ion concentration (Williams et al., 2017). As sea-

water pH decreases, the resulting increase in hydrogen ion concentration ( $[\text{H}^+]$ ) reacts with carbonate ions ( $\text{CO}_3^{2-}$ ). This consumption of carbonate ions reduces the ocean’s buffering capacity for CO<sub>2</sub> uptake, causing seawater  $p\text{CO}_2^w$  to increase. Because atmospheric  $p\text{CO}_2^a$  is fixed at 367 ppm in the CESM1.2.2 configuration, this rise in surface ocean  $p\text{CO}_2^w$  narrows the air–sea partial pressure gradient  $\Delta p\text{CO}_2$  in up-


**Table 3.** Relative magnitudes of model–data biases (%), defined as the monthly mean absolute deviation normalized by the model’s corresponding monthly variability range (95th minus 5th percentile).

| Model            | NWP  | EEP  | NEP  | SPO1 | NIO  | SIO1 | NAO1 | EOA  |
|------------------|------|------|------|------|------|------|------|------|
| POP2–waves       | 21.9 | 46.5 | 24.7 | 21.1 | 14.5 | 17.8 | 14.3 | 15.0 |
| B–CTL (baseline) | 26.8 | 71.5 | 25.3 | 22.1 | 16.2 | 21.9 | 15.4 | 11.9 |

Note: Abbreviations: NWP: Northwestern Pacific; EEP: Eastern Equatorial Pacific; NEP: Northeastern Pacific; SPO1: South Pacific Ocean 1; NIO: North Indian Ocean; SIO1: South Indian Ocean 1; NAO1: North Atlantic Ocean 1; EAO: Equatorial Atlantic Ocean. The relative magnitude of model–data bias (%) across the 360-month period is defined and calculated as follows:

$$\text{Model-data biases (\%)} = \frac{1}{12} \sum_{i=1}^{12} \frac{|M_i - O_i|}{P_{95}(M) - P_5(M)} \quad (7)$$

where  $M_i$  and  $O_i$  represent the monthly mean values simulated by (POP2–waves or B–CTL) and reference (the NOAA CT2022 inversion) air–sea CO<sub>2</sub> fluxes for the month mean of month  $i$  ( $i = 1, 2, \dots, 12$ ), respectively.  $P_{95}(M)$  and  $P_5(M)$  denote the 95th and 5th percentiles, respectively, of each calendar month’s values over the 30-year period, representing the corresponding range of monthly variability in the model.

**Figure 9.** Spatial distributions of the 30-year averages of the linear regression coefficients (LRCs) between the CO<sub>2</sub> flux and (a)  $\Delta p\text{CO}_2$ , (b)  $k_{w,660}$ , (c) SST, and (d) pH. Shading represents the baseline LRCs from the B–CTL simulation, while overlaid contours depict the structural differences between the POP2–waves and B–CTL experiments (POP2–waves minus B–CTL). White areas denote regions where the LRCs are statistically insignificant at the 95 % confidence level (based on a Student’s  $t$ -test). All variables were standardized prior to the regression analysis.

take regions (where seawater  $p\text{CO}_2^w$  is below atmospheric levels), thereby suppressing the air–sea CO<sub>2</sub> influx. Reflecting this mechanism, pH and air–sea CO<sub>2</sub> flux exhibit strong negative LRCs (less than  $-0.9$ ) across most of the global ocean, with exceptions confined to the Eastern Equatorial Pacific and along the Arctic margins (Fig. 9d). Wave effects marginally attenuate this relationship in the Eastern Equatorial Pacific (EEP), reducing the LRCs by approximately 0.1.

#### 4.3 Uncertainty arising from the absence of interactions between air–sea CO<sub>2</sub> flux and the ocean carbonate–pH system

Rustogi et al. (2025) indicate that as surface seawater  $p\text{CO}_2^w$  equilibrates with changes in DIC and alkalinity, the air–sea  $\Delta p\text{CO}_2$  gradient is partially attenuated. This induces a negative feedback loop whereby enhanced CO<sub>2</sub> uptake (or outgassing) is systematically damped by chemical re-equilibration within the marine carbonate system, a mechanism that is highly consistent with the findings of this

study. To assess the uncertainty in CO<sub>2</sub> flux resulting from the absence of interactions between the flux and the ocean carbonate–pH system (i.e., the decoupling of the  $\Delta p\text{CO}_2$ -driven negative feedback), we performed a series of sensitivity experiments. The air–sea CO<sub>2</sub> flux was reconstructed using the uncoupled  $\Delta p\text{CO}_2$  fields from the control simulation (B–CTL) and the wave-influenced  $k_{w,660}$  parameterization from the coupled wave simulation (POP2–waves). This experimental setup explicitly isolates the statistical mean and standard deviation of the flux under a scenario defined by the “lack of  $\Delta p\text{CO}_2$ -driven negative feedback”. Only data showing a statistically significant air–sea CO<sub>2</sub> flux difference (at a 95 % confidence level) between “lack of  $\Delta p\text{CO}_2$ -driven negative feedback” scenario and POP2–waves are presented in Fig. 10b. Compared to the baseline (Fig. 2b), the CO<sub>2</sub> flux under the “lack of  $\Delta p\text{CO}_2$ -driven negative feedback” scenario is stronger than that in the POP2–waves simulation across both oceanic CO<sub>2</sub> outgassing and uptake regions, accompanied by a larger SD. The most pronounced discrepancies occur in the Pacific basin across both outgassing and uptake zones. However, certain mid-to-high latitude subregions – specifically the Northeastern Pacific (NEP), South Indian Ocean 2 (SIO2), and South Pacific Ocean 2 (SPO2) – do not reach the 95 % confidence threshold.

Rustogi et al. (2025) demonstrate that while wave-enhanced  $k_{w,660}$  accelerates local air–sea CO<sub>2</sub> exchange, it simultaneously attenuates the air–sea  $\Delta p\text{CO}_2$  gradient. In this study, we conceptualize this process as a manifestation of the ocean’s buffering capacity, which encompasses the dynamic coupling between CO<sub>2</sub> flux (estimated via  $k_{w,660}$ ) and the marine carbonate–pH system. Within the coupled POP2–waves framework, incorporating wave effects alleviates surface acidification and reduces air–sea  $\Delta p\text{CO}_2$  over oceanic CO<sub>2</sub> outgassing regions; conversely, the resulting relatively higher pH levels systematically enhance the ocean’s capacity to absorb atmospheric CO<sub>2</sub> within oceanic CO<sub>2</sub> uptake regions in the coupled POP2–waves model. In contrast, the simulation lacking the  $\Delta p\text{CO}_2$ -driven negative feedback – which artificially omits realistic marine buffering effects – overestimates the air–sea exchange, yielding excess CO<sub>2</sub> outgassing into the atmosphere ( $> 0.4 \text{ mol m}^{-2} \text{ yr}^{-1}$ ) over source regions. Concurrently, it amplifies excess CO<sub>2</sub> uptake ( $< -0.4 \text{ mol m}^{-2} \text{ yr}^{-1}$ ) within regions characterized by oceanic CO<sub>2</sub> uptake regions (Fig. 10b) compared to the coupled POP2–waves model.

## 5 Conclusions

The study advances an Earth System Model (ESM) by developing an online coupling framework (POP2–waves) that incorporates wave effects directly into air–sea CO<sub>2</sub> flux simulations, capturing the  $\Delta p\text{CO}_2$ -driven negative feedback within the marine carbonate–pH system. Diagnostically neglecting this coupling by treating  $\Delta p\text{CO}_2$  and wave-induced


$k_{w,660}$  independently leads to a positive bias in global flux calculations, whereas our online coupled simulation captures a buffering negative feedback that moderates the global net response. Fully integrating these wave effects addresses three core modeling challenges:

1. Technical implementation: Resolving grid interpolation, boundary discontinuities in wave energy, and parallel computation.
2. Interdisciplinary analysis: Harmonizing kinematic, thermodynamic, and biogeochemical frameworks across system components.
3. Validation constraints: Overcoming data sparsity by synthesizing indirect flux estimates from atmospheric inversions (e.g., NOAA CT2022), ships, and buoys.

Within the coupled POP2–waves framework, the gas transfer velocity is resolved using the parameterization of Deike and Melville (2018), which attributes bubble-mediated transfer to the combined effects of friction velocity and significant wave height. This setup extends the framework of Wu et al. (2025) by utilizing a dynamically coupled ocean–wave system. This coupling configuration accounts for how flux-driven alterations in surface  $\Delta p\text{CO}_2$  feedback to influence the global air–sea CO<sub>2</sub> flux. Our regional evaluation reveals a mixed and basin-dependent performance when introducing wave effects. Across the majority of the 12 key regions of high variability defined by NOAA CT2022, the POP2–waves simulation shows generally closer structural agreement with the inversion data compared to the uncoupled B–CTL baseline. However, this improvement is not universal; notable discrepancies persist within specific sectors, including the Equatorial Atlantic Ocean (EAO), South Pacific Ocean 2 (SPO2), and South Indian Ocean 2 (SIO2), where the wave-coupled model does not clearly outperform the baseline.

Significant deviations between POP2–waves and B–CTL simulations emerge when significant wave height exceeds 1.5 m. Consistent with Gutiérrez-Loza et al. (2022), this wave-induced divergence results in a systematically enhanced air–sea CO<sub>2</sub> flux under high 10 m wind speeds and elevated wave height. Across the entire spatiotemporal domain, the bubble-mediated contribution to the total gas transfer velocity ( $k_{wB,660}/k_{w,660}$ ) in POP2–waves averages approximately 38 %. This finding slightly exceeds the  $\sim 30$  % baseline reported by Reichl and Deike (2020). As expected, such elevated ratios are most prominent in these high-wind, rough-sea regions. Concurrently, our results indicate that bubble-mediated processes account for up to 41.3 % of the total air–sea CO<sub>2</sub> flux, which is consistent with the  $\sim 40$  % contribution reported by both Reichl and Deike (2020) and Zhou et al. (2023).

Within oceanic CO<sub>2</sub> outgassing regions, POP2–waves attenuates the large air–sea  $\Delta p\text{CO}_2$  gradient, whereas it slightly amplifies lower gradients elsewhere. This feedback



**Figure 10.** Estimated air–sea CO<sub>2</sub> flux assuming the absence of the  $\Delta p\text{CO}_2$ -driven negative feedback mechanism. (a) Air–sea CO<sub>2</sub> flux calculated using uncoupled, independent  $\Delta p\text{CO}_2$  values from the B–CTL baseline and wave-dependent  $k_{w,660}$  formulations (labeled as the “lack of  $\Delta p\text{CO}_2$ -driven negative feedback” case). Shading represents the climatological mean, and contours indicate the corresponding standard deviation (SD). (b) Spatial difference in CO<sub>2</sub> flux between the decoupled feedback case and the fully coupled POP2–waves simulation (decoupled minus POP2–waves). Shading denotes the mean difference, and overlaid contours indicate regions where the difference is statistically significant at the 95 % confidence level based on a Student’s *t*-test.


drives surface pH shifts of approximately  $\pm 0.01$  that spatially correspond to the sign of regional flux. On a global scale, the air–sea  $\Delta p\text{CO}_2$  (pH) displays the strongest positive (negative) regression coefficients with the flux. Additionally, while gas transfer velocity scales positively with the absolute air–sea CO<sub>2</sub> flux magnitude, SST demonstrates an inverse relationship with this sensitivity.

Compared to the B–CTL baseline, POP2–waves simulations show that oceanic CO<sub>2</sub> uptake and outgassing regions expand by 11.8 % and 41.6 %, respectively; however, these two processes largely offset each other, leading to a simulated slight 1.8 % increase in the global ocean CO<sub>2</sub> sink. Consequently, while wave-induced enhancements in  $k_{w,660}$  boost instantaneous air–sea CO<sub>2</sub> flux, the coupled carbonate–pH system limits the net long-term impact via this  $\Delta p\text{CO}_2$ -driven feedback (Rustogi et al., 2025). This feedback effectively dampens the scaling of local CO<sub>2</sub> flux increases into

proportional changes within the global mean ocean carbon sink.

The impacts of the  $k_{w,660}$  bulk formula (Wanninkhof, 1992) and wave dynamics (Deike and Melville, 2018) on air–sea CO<sub>2</sub> flux, surface pH, and air–sea  $\Delta p\text{CO}_2$  are schematically summarized in Fig. 11. The B–CTL baseline captures a pronounced seasonal contrast in Western Pacific air–sea CO<sub>2</sub> fluxes, characterized by strong uptake during DJF that is heavily suppressed in JJA due to SST-driven reductions in solubility. This seasonal variability in air–sea  $\Delta p\text{CO}_2$  engages a negative feedback loop within the POP2–waves framework, thereby modulating regional flux magnitudes. Conversely, the Equatorial Pacific region exhibits lower pH than the Western Pacific and acts as a persistent CO<sub>2</sub> source across both seasons, showing no obvious seasonal variations – a pattern consistent with Fay et al. (2024).

While the current POP2–waves framework highlights the importance of online coupling and  $\Delta p\text{CO}_2$ -driven feed-



**Figure 11.** Schematic diagrams illustrate B–CTL (gray colors) and the differences between POP2–waves and B–CTL (red colors) in air–sea CO<sub>2</sub> flux (arrows), surface pH (tube colors indicator), and air–sea ΔpCO<sub>2</sub> (circle markers) over the oceanic CO<sub>2</sub> uptake region (WP; 160–180° E, 35–40° N) and the oceanic CO<sub>2</sub> outgassing region (EP; 230–250° E, 0–5° S). Each panel includes an ocean–atmosphere interface, with ocean color shading representing relative SST levels across different seasons. The lower-left corner displays the SST value, (a) WP in DJF seasonal mean, (b) EP in DJF seasonal mean, (c) WP in JJA seasonal mean, and (d) EP in JJA seasonal mean.

backs, several avenues remain to future refine the model. A key priority is the continuous improvement of gas transfer velocity formulations. Future efforts will focus on transitioning to the new generalized formulation proposed by Deike et al. (2025), which incorporates updated coefficients and accounts for an asymmetric bubble flux contribution. While this advancement is expected to have a minor impact on oceanic CO<sub>2</sub> flux estimations, it holds potential significance for constraining global marine O<sub>2</sub> fluxes. Furthermore, moving toward a fully coupled Earth System Model (ESM) – which integrates dynamic atmospheric feedback alongside ocean and wave components – will be pivotal to better unravel the long-term impacts of wave-induced processes on global marine biogeochemical cycles.

**Code and data availability.** The model code of POP2–waves coupled model is available at <https://doi.org/10.5281/zenodo.15795234> (Lan, 2025). Input data of POP2–waves using the climatological Hadley Centre Sea Ice and Sea Surface Temperature dataset, including 30-year numerical experiments, are available at <https://doi.org/10.5281/zenodo.5510795> (Lan et al., 2021). CarbonTracker CT2022 data are provided by the National Oceanic and Atmospheric Administration (NOAA) and available from Global Monitoring Laboratory <https://gml.noaa.gov/aftp/products/carbontracker/co2/CT2022/> (last access: 6 August 2026; Jacobson et al., 2023).

**Author contributions.** YYL is the sole developer of the POP2–waves coupled model and writes the majority part of the paper. HHH provides computational support and analysis suggestions. WLL supports reorganization and offers analytical recommendations and SC offers a non-parallel wave module as part of the POM source code.

**Competing interests.** The contact author has declared that none of the authors has any competing interests.

**Disclaimer.** Publisher’s note: Copernicus Publications remains neutral with regard to jurisdictional claims made in the text, published maps, institutional affiliations, or any other geographical representation in this paper. The authors bear the ultimate responsibility for providing appropriate place names. Views expressed in the text are those of the authors and do not necessarily reflect the views of the publisher.

**Acknowledgements.** Our deepest gratitude goes to the editors and anonymous reviewers for their careful work and thoughtful suggestions that have helped improve this paper substantially. We would like to thank NOAA GML (Boulder, Colorado, USA) for providing the CarbonTracker CT2022 data from the website at <http://carbontracker.noaa.gov> (last access: 20 May 2026). We sincerely thank the National Center for Atmospheric Research (NCAR) and their Atmosphere Model Working Group (AMWG) for releasing CESM1.2.2. We are also grateful to the National Center for High-

performance Computing, Taiwan for providing the facilities for the computational procedures for running POP2–waves simulations. Thanks, Gemini and ChatGPT for correcting the English grammar.

**Financial support.** This research received funding from the National Science and Technology Council of Taiwan (grant nos. NSTC 115-2119-M-001-006-, NSTC 114-2119-M-001-011-, NSTC 114-2111-M-001-007-, and NSTC 113-2111-M-001-009-) as well as support from the Academia Sinica Grand Challenge Program in Taiwan (grant no. AS-GCP-112-M03).

**Review statement.** This paper was edited by Chia-Te Chien and reviewed by two anonymous referees.

## References

- Akhand, A., Chanda, A., Watanabe, K., Das, S., Tokoro, T., Hazra, S., and Kuwae, T.: Reduction in riverine freshwater supply changes inorganic and organic carbon dynamics and air–water CO<sub>2</sub> fluxes in a tropical mangrove dominated estuary, *J. Geophys. Res. G-Biogeophys.*, 126, e2020JG006144, <https://doi.org/10.1029/2020JG006144>, 2021.
- Andreas, E., Vlahos, P., and Monahan, E.: The potential role of sea spray droplets in facilitating air–sea gas transfer, in: IOP conference series: earth and environmental science, IOP Publishing, 35, 012003, <https://doi.org/10.1088/1755-1315/35/1/012003>, 2016.
- Bakker, D. C. E., Pfeil, B., Landa, C. S., Metzl, N., O'Brien, K. M., Olsen, A., Smith, K., Cosca, C., Harasawa, S., Jones, S. D., Nakaoka, S., Nojiri, Y., Schuster, U., Steinhoff, T., Sweeney, C., Takahashi, T., Tilbrook, B., Wada, C., Wanninkhof, R., Alin, S. R., Balestrini, C. F., Barbero, L., Bates, N. R., Bianchi, A. A., Bonou, F., Boutin, J., Bozec, Y., Burger, E. F., Cai, W.-J., Castle, R. D., Chen, L., Chierici, M., Currie, K., Evans, W., Featherstone, C., Feely, R. A., Fransson, A., Goyet, C., Greenwood, N., Gregor, L., Hankin, S., Hardman-Mountford, N. J., Harlay, J., Hauck, J., Hoppema, M., Humphreys, M. P., Hunt, C. W., Huss, B., Ibáñez, J. S. P., Johannessen, T., Keeling, R., Kitidis, V., Körtzinger, A., Kozyr, A., Krasakopoulou, E., Kuwata, A., Landschützer, P., Lauvset, S. K., Lefèvre, N., Lo Monaco, C., Manke, A., Mathis, J. T., Merlivat, L., Millero, F. J., Monteiro, P. M. S., Munro, D. R., Murata, A., Newberger, T., Omar, A. M., Ono, T., Paterson, K., Pearce, D., Pierrot, D., Robbins, L. L., Saito, S., Salisbury, J., Schlitzer, R., Schneider, B., Schweitzer, R., Sieger, R., Skjelvan, I., Sullivan, K. F., Sutherland, S. C., Sutton, A. J., Tadokoro, K., Telszewski, M., Tuma, M., van Heuven, S. M. A. C., Vandemark, D., Ward, B., Watson, A. J., and Xu, S.: A multi-decade record of high-quality *f*CO<sub>2</sub> data in version 3 of the Surface Ocean CO<sub>2</sub> Atlas (SOCAT), *Earth Syst. Sci. Data*, 8, 383–413, <https://doi.org/10.5194/essd-8-383-2016>, 2016.
- Bange, H. W., Mongwe, P., Shutler, J. D., Arévalo-Martínez, D. L., Bianchi, D., Lauvset, S. K., Liu, C., Löscher, C. R., Martins, H., Rosentreter, J. A., Schmale, O., Steinhoff, T., Upstill-Goddard, R. C., Wanninkhof, R., Wilson, S. T., and Xie, H.: Advances in understanding of air–sea exchange and cycling of greenhouse gases in the upper ocean, *Elementa*, 12, <https://doi.org/10.1525/elementa.2023.00044>, 2024.
- Behncke, J., Landschützer, P., and Tanhua, T.: A detectable change in the air–sea CO<sub>2</sub> flux estimate from sailboat measurements, *Sci. Rep.*, 14, 3345, <https://doi.org/10.1038/s41598-024-53159-0>, 2024.
- Bell, T. G., Landwehr, S., Miller, S. D., de Bruyn, W. J., Callaghan, A. H., Scanlon, B., Ward, B., Yang, M., and Saltzman, E. S.: Estimation of bubble-mediated air–sea gas exchange from concurrent DMS and CO<sub>2</sub> transfer velocities at intermediate–high wind speeds, *Atmos. Chem. Phys.*, 17, 9019–9033, <https://doi.org/10.5194/acp-17-9019-2017>, 2017.
- Blomquist, B. W., Brumer, S. E., Fairall, C. W., Huebert, B. J., Zappa, C. J., Brooks, I. M., Yang, M., Bariteau, L., Prytherch, J., Hare, J. E., Czerski, H., Matei, A., and Pascal, R. W.: Wind speed and sea state dependencies of air–sea gas transfer: Results from the High Wind speed Gas exchange Study (HiWinGS), *J. Geophys. Res.-Oceans*, 122, 8034–8062, <https://doi.org/10.1002/2017JC013181>, 2017.
- Blumberg, A. F. and Mellor, G. L.: A Description of a Three-Dimensional Coastal Ocean Circulation Model, in: Coastal and Estuarine Sciences, edited by: Heaps, N. S., Book 4, American Geophysical Union, 1–6, ISBN 978-1-118-66504-6, <https://agupubs.onlinelibrary.wiley.com/doi/10.1029/CO004p0001> (last access: 10 August 2026), 1987.
- Boucher, O., Servonnat, J., Albright, A. L., Aumont, O., Balkanski, Y., Bastrikov, V., Bekki, S., Bonnet, R., Bony, S., Bopp, L., Braconnot, P., Brockmann, P., Cadule, P., Caubel, A., Cheruy, F., Codron, F., Cozic, A., Cugnet, D., D'Andrea, F., Davini, P., de Lavergne, C., Denvil, S., Deshayes, J., Devilliers, M., Ducharme, A., Dufresne, J.-L., Dupont, E., Éthé, C., Fairhead, L., Falletti, L., Flavoni, S., Foujols, M.-A., Gardoll, S., Gastineau, G., Ghattas, J., Grandpeix, J.-Y., Guenet, B., Guez, L. E., Guilyardi, E., Guimberteau, M., Hauglustaine, D., Hourdin, F., Idelkadi, A., Joussaume, S., Kageyama, M., Khodri, M., Krinner, G., Lebas, N., Levavasseur, G., Lévy, C., Li, L., Lott, F., Lurton, T., Luyssaert, S., Madec, G., Madeleine, J.-B., Maignan, F., Marchand, M., Marti, O., Melul, L., Meurdesoif, Y., Mignot, J., Musat, I., Ottlé, C., Peylin, P., Planton, Y., Polcher, J., Rio, C., Rochetin, N., Rousset, C., Sepulchre, P., Sima, A., Swingedouw, D., Thiéblemont, R., Traore, A. K., Vancoppenolle, M., Vial, J., Vialard, J., Viovy, N., and Vuichard, N.: Presentation and Evaluation of the IPSL-CM6A-LR Climate Model, *J. Adv. Model. Earth Sy.*, 12, e2019MS002010, <https://doi.org/10.1029/2019MS002010>, 2020.
- Brumer, S. E., Zappa, C. J., Blomquist, B. W., Fairall, C. W., Cifuentes-Lorenzen, A., Edson, J. B., Brooks, I. M., and Huebert, B. J.: Wave-related Reynolds number parameterizations of CO<sub>2</sub> and DMS transfer velocities, *Geophys. Res. Lett.*, 44, 9865–9875, <https://doi.org/10.1002/2017GL074979>, 2017.
- Chikamoto, M. O. and DiNezio, P.: Multi-century changes in the ocean carbon cycle controlled by the tropical oceans and the Southern Ocean, *Global Biogeochem. Cy.*, 35, e2021GB007090, <https://doi.org/10.1029/2021GB007090>, 2021.
- Chikamoto, M. O., DiNezio, P., and Lovenduski, N.: Long-term slowdown of ocean carbon uptake by alkalinity dynamics, *Geophys. Res. Lett.*, 50, e2022GL101954, <https://doi.org/10.1029/2022GL101954>, 2023.
- Coggins, A., Watson, A. J., Schuster, U., Mackay, N., King, B., McDonagh, E., and Poulton, A. J.: Surface ocean carbon budget in

- the 2017 South Georgia diatom bloom: Observations and validation of profiling biogeochemical argo floats, *Deep-Sea Res. Pt. II*, 209, 105275, <https://doi.org/10.1016/j.dsr2.2023.105275>, 2023.
- Couldrey, M. P., Oliver, K. I. C., Yool, A., Halloran, P. R., and Achterberg, E. P.: On which timescales do gas transfer velocities control North Atlantic CO<sub>2</sub> flux variability?, *Global Biogeochem. Cy.*, 30, 787–802, <https://doi.org/10.1002/2015GB005267>, 2016.
- Czerski, H., Brooks, I. M., Gunn, S., Pascal, R., Matei, A., and Blomquist, B.: Ocean bubbles under high wind conditions – Part 2: Bubble size distributions and implications for models of bubble dynamics, *Ocean Sci.*, 18, 587–608, <https://doi.org/10.5194/os-18-587-2022>, 2022.
- Danabasoglu, G., Lamarque, J.-F., Bacmeister, J., Bailey, D. A., DuVivier, A. K., Edwards, J., Emmons, L. K., Fasullo, J., Garcia, R., Gettelman, A., Hannay, C., Holland, M. M., Large, W. G., Lauritzen, P. H., Lawrence, D. M., Lenaerts, J. T. M., Lindsay, K., Lipscomb, W. H., Mills, M. J., Neale, R., Oleson, K. W., Otto-Bliesner, B., Phillips, A. S., Sacks, W., Tilmes, S., van Kampenhout, L., Vertenstein, M., Bertini, A., Dennis, J., Deser, C., Fischer, C., Fox-Kemper, B., Kay, J. E., Kinnison, D., Kushner, P. J., Larson, V. E., Long, M. C., Mickelson, S., Moore, J. K., Nienhouse, E., Polvani, L., Rasch, P. J., and Strand, W. G.: The Community Earth System Model Version 2 (CESM2), *J. Adv. Model. Earth Sy.*, 12, e2019MS001916, <https://doi.org/10.1029/2019MS001916>, 2020.
- Deike, L.: Mass transfer at the ocean-atmosphere interface: The role of wave breaking, droplets, and bubbles, *Annu. Rev. Fluid Mech.*, 54, 191–224, <https://doi.org/10.1146/annurev-fluid-030121-014132>, 2022.
- Deike, L. and Melville, W. K.: Gas transfer by breaking waves, *Geophys. Res. Lett.*, 45, 10482–10492, <https://doi.org/10.1029/2018GL078758>, 2018.
- Deike, L., Zhou, X., Rustogi, P., Stanley, R. H. R., Reichl, B. G., Bushinsky, S. M., and Resplandy, L.: A universal wind–wave–bubble formulation for air–sea gas exchange and its impact on oxygen fluxes, *P. Natl. Acad. Sci. USA*, 122, e2419319122, <https://doi.org/10.1073/pnas.2419319122>, 2025.
- Dickson, A. G.: An exact definition of total alkalinity and a procedure for the estimation of alkalinity and total inorganic carbon from titration data, *Deep-Sea Res.*, 28A, 609–623, [https://doi.org/10.1016/0198-0149\(81\)90121-7](https://doi.org/10.1016/0198-0149(81)90121-7), 1981.
- Dong, Y., Yang, M., Bakker, D. C. E., Kitidis, V., and Bell, T. G.: Uncertainties in eddy covariance air–sea CO<sub>2</sub> flux measurements and implications for gas transfer velocity parameterisations, *Atmos. Chem. Phys.*, 21, 8089–8110, <https://doi.org/10.5194/acp-21-8089-2021>, 2021.
- Dong, Y., Bakker, D. C. E., Bell, T. G., Huang, B., Landschützer, P., Liss, P. S., and Yang, M.: Update on the temperature corrections of global air–sea CO<sub>2</sub> flux estimates, *Global Biogeochem. Cy.*, 36, e2022GB007360, <https://doi.org/10.1029/2022GB007360>, 2022.
- Edson, J. B., Fairall, C. W., Bariteau, L., Zappa, C. J., Cifuentes-Lorenzen, A., McGillis, W. R., Pezoa, S., Hare, J. E., and Helmig, D.: Direct covariance measurement of CO<sub>2</sub> gas transfer velocity during the 2008 Southern Ocean Gas Exchange Experiment: Wind speed dependency, *J. Geophys. Res.-Oceans*, 116, <https://doi.org/10.1029/2011jc007022>, 2011.
- Fay, A. R., Gregor, L., Landschützer, P., McKinley, G. A., Gruber, N., Gehlen, M., Iida, Y., Laruelle, G. G., Rödenbeck, C., Roobaert, A., and Zeng, J.: SeaFlux: harmonization of air–sea CO<sub>2</sub> fluxes from surface pCO<sub>2</sub> data products using a standardized approach, *Earth Syst. Sci. Data*, 13, 4693–4710, <https://doi.org/10.5194/essd-13-4693-2021>, 2021.
- Fay, A. R., Munro, D. R., McKinley, G. A., Pierrot, D., Sutherland, S. C., Sweeney, C., and Wanninkhof, R.: Updated climatological mean  $\Delta f$ CO<sub>2</sub> and net sea–air CO<sub>2</sub> flux over the global open ocean regions, *Earth Syst. Sci. Data*, 16, 2123–2139, <https://doi.org/10.5194/essd-16-2123-2024>, 2024.
- Friedlingstein, P., Jones, M. W., O’Sullivan, M., Andrew, R. M., Bakker, D. C. E., Hauck, J., Le Quéré, C., Peters, G. P., Peters, W., Pongratz, J., Sitoh, S., Canadell, J. G., Ciais, P., Jackson, R. B., Alin, S. R., Anthoni, P., Bates, N. R., Becker, M., Belouin, N., Bopp, L., Chau, T. T. T., Chevallier, F., Chini, L. P., Cronin, M., Currie, K. I., Decharme, B., Djeutchouang, L. M., Dou, X., Evans, W., Feely, R. A., Feng, L., Gasser, T., Gilfillan, D., Gkritzalis, T., Grassi, G., Gregor, L., Gruber, N., Gürses, Ö., Harris, I., Houghton, R. A., Hurtt, G. C., Iida, Y., Ilyina, T., Luijkx, I. T., Jain, A., Jones, S. D., Kato, E., Kennedy, D., Klein Goldewijk, K., Knauer, J., Korsbakken, J. I., Körtzinger, A., Landschützer, P., Lauvset, S. K., Lefèvre, N., Lienert, S., Liu, J., Marland, G., McGuire, P. C., Melton, J. R., Munro, D. R., Nabel, J. E. M. S., Nakaoka, S.-I., Niwa, Y., Ono, T., Pierrot, D., Poulter, B., Rehder, G., Resplandy, L., Robertson, E., Rödenbeck, C., Rosan, T. M., Schwingner, J., Schwingshackl, C., Séférian, R., Sutton, A. J., Sweeney, C., Tanhua, T., Tans, P. P., Tian, H., Tilbrook, B., Tubiello, F., van der Werf, G. R., Vuichard, N., Wada, C., Wanninkhof, R., Watson, A. J., Willis, D., Wiltshire, A. J., Yuan, W., Yue, C., Yue, X., Zaehle, S., and Zeng, J.: Global Carbon Budget 2021, *Earth Syst. Sci. Data*, 14, 1917–2005, <https://doi.org/10.5194/essd-14-1917-2022>, 2022.
- Gray, A. R., Johnson, K. S., Bushinsky, S. M., Riser, S. C., Russell, J. L., Talley, L. D., Wanninkhof, R., Williams, N. L., and Sarmiento, J. L.: Autonomous biogeochemical floats detect significant carbon dioxide outgassing in the high-latitude Southern Ocean, *Geophys. Res. Lett.*, 45, 9049–9057, <https://doi.org/10.1029/2018GL078013>, 2018.
- Gutiérrez-Loza, L., Nilsson, E., Wallin, M. B., Sahlée, E., and Rutgersson, A.: On physical mechanisms enhancing air–sea CO<sub>2</sub> exchange, *Biogeosciences*, 19, 5645–5665, <https://doi.org/10.5194/bg-19-5645-2022>, 2022.
- Hajima, T., Watanabe, M., Yamamoto, A., Tatebe, H., Noguchi, M. A., Abe, M., Ohgaito, R., Ito, A., Yamazaki, D., Okajima, H., Ito, A., Takata, K., Ogochi, K., Watanabe, S., and Kawamiya, M.: Development of the MIROC-ES2L Earth system model and the evaluation of biogeochemical processes and feedbacks, *Geosci. Model Dev.*, 13, 2197–2244, <https://doi.org/10.5194/gmd-13-2197-2020>, 2020.
- Heimdal, T. H., McKinley, G. A., Sutton, A. J., Fay, A. R., and Gloege, L.: Assessing improvements in global ocean pCO<sub>2</sub> machine learning reconstructions with Southern Ocean autonomous sampling, *Biogeosciences*, 21, 2159–2176, <https://doi.org/10.5194/bg-21-2159-2024>, 2024.
- Honkanen, M., Tuovinen, J.-P., Laurila, T., Mäkelä, T., Hatakka, J., Kielosto, S., and Laakso, L.: Measuring turbulent CO<sub>2</sub> fluxes with a closed-path gas analyzer in a marine environment, *At-*

- mos. Meas. Tech., 11, 5335–5350, <https://doi.org/10.5194/amt-11-5335-2018>, 2018.
- Hurrell, J. W., Holland, M. M., Gent, P. R., Ghan, S., Kay, J. E., Kushner, P. J., Lamarque, J.-F., Large, W. G., Lawrence, D., Lindsay, K., Lipscomb, W. H., Long, M. C., Mahowald, N., Marsh, D. R., Neale, R. B., Rasch, P., Vavrus, S., Vertenstein, M., Bader, D., Collins, W. D., Hack, J. J., Kiehl, J., and Marshall, S.: The Community Earth System Model: A framework for collaborative research, *B. Am. Meteorol. Soc.*, 94, 1339–1360, <https://doi.org/10.1175/BAMS-D-12-00121.1>, 2013.
- Jacobson, A. R., Gruber, N., Sarmiento, J. L., Gloor, M., and Fletcher, S. E. M.: A joint atmosphere–ocean inversion for surface fluxes of carbon dioxide: I. methods and global-scale fluxes, *Global. Biogeochem. Cy.*, 21, GB1020, <https://doi.org/10.1029/2005GB002556>, 2007.
- Jacobson, A. R., Schuldt, K. N., Tans, P., Andrews, A., Miller, J. B., Oda, T., Basu, S., Mund, J., Weir, B., Ott, L., Aalto, T., Abshire, J. B., Aikin, K., Aoki, S., Apadula, F., Arnold, S., Baier, B., Bartyzel, J., Beyersdorf, A., Biermann, T., Biraud, S. C., Boenisch, H., Brailsford, G., Brand, W. A., Chen, G., Chen, H., Chmura, L., Clark, S., Colomb, A., Commene, R., Conil, S., Couret, C., Cox, A., Cristofanelli, P., Cuevas, E., Curcoll, R., Daube, B., Davis, K. J., de Wekker, S., Della Coletta, J., Delmotte, M., DiGangi, E., DiGangi, J. P., di Sarra, A. G., Dlugokencky, E., Elkins, J. W., Emmenegger, L., Fang, S., Fischer, M. L., Forster, G., Frumau, A., Galkowski, M., Gatti, L. V., Gehrlein, T., Gerbig, C., Gheusi, F., Gloor, E., Gomez-Trueba, V., Goto, D., Griffiths, T., Hammer, S., Hanson, C., Haszpra, L., Hatakka, J., Heimann, M., Heliasz, M., Hensen, A., Hermansen, O., Hintsa, E., Holst, J., Ivakhov, V., Jaffe, D. A., Jordan, A., Joubert, W., Karion, A., Kawa, S. R., Kazan, V., Keeling, R. F., Keronen, P., Kneuer, T., Kolari, P., Komínková, K., Kort, E., Kozlova, E., Krummel, P., Kubistin, D., Labuschagne, C., Lam, D. H., Lan, X., Langenfelds, R. L., Laurent, O., Laurila, T., Lauvaux, T., Lavric, J., Law, B. E., Lee, J., Lee, O. S., Lehner, I., Lehtinen, K., Leppert, R., Leskinen, A., Leuenberger, M., Levin, I., Levula, J., Lin, J., Lindauer, M., Loh, Z., Lopez, M., Luijkx, I. T., Lunder, C. R., Machida, T., Mammarella, I., Manca, G., Manning, A., Manning, A., Marek, M. V., Martin, M. Y., Matsueda, H., McKain, K., Meijer, H., Meinhardt, F., Merchant, L., Mihalopoulos, N., Miles, N. L., Miller, C. E., Mitchell, L., Mölder, M., Montzka, S., Moore, F., Moossen, H., Morgan, E., Morgui, J.-A., Morimoto, S., Müller-Williams, J., Munger, J. W., Munro, D., Myhre, C. L., Nakaoka, S.-I., Necki, J., Newman, S., Nichol, S., Niwa, Y., Obersteiner, F., O'Doherty, S., Paplawsky, B., Peischl, J., Peltola, O., Piacentino, S., Pichon, J.-M., Pickers, P., Piper, S., Pitt, J., Plass-Dülmer, C., Platt, S. M., Prinzi-valli, S., Ramonet, M., Ramos, R., Reyes-Sanchez, E., Richardson, S. J., Riris, H., Rivas, P. P., Ryerson, T., Saito, K., Sargent, M., Sasakawa, M., Scheeren, B., Schuck, T., Schumacher, M., Seifert, T., Sha, M. K., Shepson, P., Shook, M., Sloop, C. D., Smith, P., Stanley, K., Steinbacher, M., Stephens, B., Sweeney, C., Thoning, K., Timas, K., Torn, M., Tørseth, K., Trisolino, P., Turnbull, J., van den Bulk, P., van Dinter, D., Vermeulen, A., Viner, B., Vitkova, G., Walker, S., Watson, A., Wofsy, S. C., Worsey, J., Worthy, D., Young, D., Zaehle, S., Zahn, A., and Zimnoch, M.: CarbonTracker CT2022, NOAA Global Monitoring Laboratory, <https://doi.org/10.25925/z1gj-3254>, 2023.
- Jin, C. X., Zhou, T. J., Chen, X. L., and Wu, B.: Seasonally evolving dominant interannual variability mode of air–sea CO<sub>2</sub> flux over the western North Pacific simulated by CESM1-BGC, *Sci. China Earth Sci.*, 60, 1854–1865, <https://doi.org/10.1007/s11430-015-9085-4>, 2017.
- Johansson, M. M., Björkqvist, J.-V., Särkkä, J., Leijala, U., and Kahma, K. K.: Correlation of wind waves and sea level variations on the coast of the seasonally ice-covered Gulf of Finland, *Nat. Hazards Earth Syst. Sci.*, 22, 813–829, <https://doi.org/10.5194/nhess-22-813-2022>, 2022.
- Koseki, S., Tjiputra, J., Fransner, F., Crespo, L. R., and Keenlyside, N. S.: Isentangling the impact of Atlantic Niño on sea–air CO<sub>2</sub> flux, *Nat. Commun.*, 14, 3649, <https://doi.org/10.1038/s41467-023-38718-9>, 2023.
- Krall, K. E., Smith, A. W., Takagaki, N., and Jähne, B.: Air–sea gas exchange at wind speeds up to 85 m s<sup>-1</sup>, *Ocean Sci.*, 15, 1783–1799, <https://doi.org/10.5194/os-15-1783-2019>, 2019.
- Lan, Y.-Y.: POP2–waves coupled model in CESM1.2.2 framework, Zenodo [code], <https://doi.org/10.5281/zenodo.15795234>, 2025.
- Lan, Y.-Y., Hsu, H.-H., Tsuang, B.-J., and Tu, C.-Y.: rceclcr/CAM5\_SIT\_v1.0 (v1.0.0), Zenodo [code], <https://doi.org/10.5281/zenodo.5510795>, 2021.
- Li, S., Babanin, A. V., and Guan, C.: Dimensionless Parameterizations of Air–Sea CO<sub>2</sub> Gas Transfer Velocity on Surface Waves, *Tellus B*, 75, 1–12, <https://doi.org/10.16993/tellusb.1867>, 2023.
- Liu, B., Six, K. D., and Ilyina, T.: Incorporating the stable carbon isotope 13C in the ocean biogeochemical component of the Max Planck Institute Earth System Model, *Biogeosciences*, 18, 4389–4429, <https://doi.org/10.5194/bg-18-4389-2021>, 2021.
- Long, M. C., Lindsay, K., Peacock, S., Moore, J. K., and Doney, S. C.: Twentieth-Century oceanic carbon uptake and storage in CESM1 (BGC), *J. Climate*, 26, 6775–6800, <https://doi.org/10.1175/JCLI-D-12-00184.1>, 2013.
- Lovato, T., Peano, D., Butenschön, M., Materia, S., Iovino, D., Scoccimarro, E., Fogli, P. G., Cherchi, A., Bellucci, A., Gualdi, S., Masina, S., and Navarra, A.: CMIP6 Simulations With the CMCC Earth System Model (CMCC-ESM2), *J. Adv. Model. Earth Sy.*, 14, e2021MS002814, <https://doi.org/10.1029/2021MS002814>, 2022.
- Lovenduski, N. S., Yeager, S. G., Lindsay, K., and Long, M. C.: Predicting near-term variability in ocean carbon uptake, *Earth Syst. Dynam.*, 10, 45–57, <https://doi.org/10.5194/esd-10-45-2019>, 2019.
- Macovei, V. A., Petersen, W., Brix, H., and Voynova, Y. G.: Reduced ocean carbon sink in the south and central North Sea (2014–2018) revealed from FerryBox observations, *Geophys. Res. Lett.*, 48, e2021GL092645, <https://doi.org/10.1029/2021GL092645>, 2021.
- Mauritsen, T., Bader, J., Becker, T., Behrens, J., Bittner, M., Brokopf, R., Brovkin, V., Claussen, M., Crueger, T., Esch, M., Fast, I., Fiedler, S., Flaeschner, D., Gayler, V., Giorgetta, M., Goll, D. S., Haak, H., Hagemann, S., Hedemann, C., Hohenegger, C., Ilyina, T., Jahns, T., Jimenez-de-la Cuesta, D., Jungclaus, J., Kleinen, T., Kloster, S., Kracher, D., Kinne, S., Kleberg, D., Lasslop, G., Kornbluh, L., Marotzke, J., Matei, D., Meraner, K., Mikolajewicz, U., Modali, K., Moebis, B., Mueller, W. A., Nabel, J. E. M. S., Nam, C. C. W., Notz, D., Nyawira, S.-S., Paulsen, H., Peters, K., Pincus, R., Pohlmann, H., Pongratz, J., Popp, M., Raddatz, T. J., Rast, S., Redler, R., Reick,

- C. H., Rohrschneider, T., Schemann, V., Schmidt, H., Schnur, R., Schulzweida, U., Six, K. D., Stein, L., Stemmler, I., Stevens, B., von Storch, J.-S., Tian, F., Voigt, A., Vrese, P., Wieners, K.-H., Wilkenskjaeld, S., Winkler, A., and Roeckner, E.: Developments in the MPI-M Earth System Model version 1.2 (MPI-ESM1.2) and Its Response to Increasing CO<sub>2</sub>, *J. Adv. Model. Earth Sy.*, 11, 998–1038, <https://doi.org/10.1029/2018MS001400>, 2019.
- McKinley, G. A., Fay, A. R., Eddebbbar, Y. A., Gloege, L., and Lovenduski, N. S.: External forcing explains recent-decadal variability of the ocean carbonsink, *AGU Advances*, 1, e2019AV00014, <https://doi.org/10.1029/2019AV000149>, 2020.
- Mellor, G. L., Donelan, M. A., and Oey, L.-Y.: A surface wave model for coupling with numerical ocean circulation models, *J. Atmos. Ocean. Technol.*, 25, 1785–1807, <https://doi.org/10.1175/2008JTECHO573.1>, 2008.
- Monahan, E. C. and Spillane, M. C.: The role of oceanic whitecaps in air–sea gas exchange, in: *Gas transfer at water surfaces*, edited by: Brutsaert, W. and Jirka, G. H., Reidel, Hingham, MA, 495–503, [https://doi.org/10.1007/978-94-017-1660-4\\_45](https://doi.org/10.1007/978-94-017-1660-4_45), 1984.
- Moore, J. K., Lindsay, K., Doney, S. C., Long, M. C., and Misumi, K.: Marine ecosystem dynamics and biogeochemical cycling in the Community Earth System Model [CESM1 (BGC)]: Comparison of the 1990s with the 2090s under the RCP4.5 and RCP8.5 scenarios, *J. Climate*, 26, 9291–9312, <https://doi.org/10.1175/JCLI-D-12-00566.1>, 2013.
- Müller, J. D., Gruber, N., Carter, B., Feely, R., Ishii, M., Lange, N., Lauvset, S. K., Murata, A., Olsen, A., Pérez, F. F., Sabine, C., Tanhua, T., Wanninkhof, R., and Zhu, D.: Decadal trends in the oceanic storage of anthropogenic carbon from 1994 to 2014, *AGU Advances*, 4, e2023AV000875, <https://doi.org/10.1029/2023AV000875>, 2023.
- Reichl, B. G. and Deike, L.: Contribution of Sea-State Dependent Bubbles to Air–sea Carbon Dioxide Fluxes, *Geophys. Res. Lett.*, 47, e2020GL087267, <https://doi.org/10.1029/2020GL087267>, 2020.
- Roobaert, A., Resplandy, L., Laruelle, G. G., Liao, E., and Regnier, P.: Unraveling the physical and biological controls of the global coastal CO<sub>2</sub> sink, *Global Biogeochem. Cy.*, 38, e2023GB007799, <https://doi.org/10.1029/2023GB007799>, 2024.
- Rustogi, P., Resplandy, L., Liao, E., Reichl, B. G., and Deike, L.: Influence of wave-induced variability on ocean carbon uptake, *Global Biogeochem. Cy.*, 39, e2024GB008382, <https://doi.org/10.1029/2024GB008382>, 2025.
- Sabine, C., Sutton, A., McCabe, K., Lawrence-Slavas, N., Alin, S., Feely, R., Jenkins, R., Maenner, S., Meinig, C., Thomas, J., van Ooijen, E., Passmore, A., and Tilbrook, B.: Evaluation of a new carbon dioxide system for autonomous surface vehicles, *J. Atmos. Ocean. Tech.*, 37, 1305–1317, <https://doi.org/10.1175/JTECH-D-20-0010.1>, 2020.
- Séférian, R., Nabat, P., Michou, M., Saint-Martin, D., Voldoire, A., Colin, J., Decharme, B., Delire, C., Berthet, S., Chevallier, M., Sénési, S., Franchisteguy, L., Vial, J., Mallet, M., Joetzjer, E., Geoffroy, O., Guérémy, J.-F., Moine, M.-P., Msadek, R., Ribes, A., Rocher, M., Roebrig, R., Salas-y-Méla, D., Sanchez, E., Terray, L., Valcke, S., Waldman, R., Aumont, O., Bopp, L., Deshayes, J., Éthé, C., and Madec, G.: Evaluation of CNRM Earth-System model, CNRM-ESM2-1: role of Earth system processes in present-day and future climate, *J. Adv. Model. Earth Sy.*, 11, 4182–4227, <https://doi.org/10.1029/2019MS001791>, 2019.
- Seland, Ø., Bentsen, M., Olivie, D., Toniazio, T., Gjermundsen, A., Graff, L. S., Debernard, J. B., Gupta, A. K., He, Y.-C., Kirkevåg, A., Schwinger, J., Tjiputra, J., Aas, K. S., Bethke, I., Fan, Y., Griesfeller, J., Grini, A., Guo, C., Ilicak, M., Karset, I. H. H., Landgren, O., Liakka, J., Moseid, K. O., Nummelin, A., Spensberger, C., Tang, H., Zhang, Z., Heinze, C., Iversen, T., and Schulz, M.: Overview of the Norwegian Earth System Model (NorESM2) and key climate response of CMIP6 DECK, historical, and scenario simulations, *Geosci. Model Dev.*, 13, 6165–6200, <https://doi.org/10.5194/gmd-13-6165-2020>, 2020.
- Sellar, A. A., Jones, C. G., Mulcahy, J. P., Tang, Y., Yool, A., Wiltshire, A., O'Connor, F. M., Stringer, M., Hill, R., Palmieri, J., Woodward, S., de Mora, L., Kuhlbrodt, T., Rumbold, S. T., Kelley, D. I., Ellis, R., Johnson, C. E., Walton, J., Abraham, N. L., Andrews, M. B., Andrews, T., Archibald, A. T., Berthou, S., Burke, E., Blockley, E., Carslaw, K., Dalvi, M., Edwards, J., Folberth, G. A., Gedney, N., Griffiths, P. T., Harper, A. B., Hendry, M. A., Hewitt, A. J., Johnson, B., Jones, A., Jones, C. D., Keeble, J., Liddicoat, S., Morgenstern, O., Parker, R. J., Predoi, V., Robertson, E., Siahann, A., Smith, R. S., Swaminathan, R., Woodhouse, M. T., Zeng, G., and Zerroukat, M.: UKESM1: Description and Evaluation of the U.K. Earth System Model, *J. Adv. Model. Earth Syst.*, 11, 4513–4558, <https://doi.org/10.1029/2019MS001739>, 2019.
- Shutler, J. D., Wanninkhof, R., Nightingale, P. D., Woolf, D. K., Bakker, D. C., Watson, A., Ashton, I., Holding, T., Chapron, B., Quilfen, Y., Fairall, C., Schuster, U., Nakajima, M., and Donlon, C. J.: Satellites will address critical science priorities for quantifying ocean carbon, *Front. Ecol. Environ.*, 18, 27–35, <https://doi.org/10.1002/fee.2129>, 2019.
- Sigmond, M., Anstey, J., Arora, V., Digby, R., Gillett, N., Kharin, V., Merryfield, W., Reader, C., Scinocca, J., Swart, N., Virgin, J., Abraham, C., Cole, J., Lambert, N., Lee, W.-S., Liang, Y., Malinina, E., Rieger, L., von Salzen, K., Seiler, C., Seinen, C., Shao, A., Sospedra-Alfonso, R., Wang, L., and Yang, D.: Improvements in the Canadian Earth System Model (CanESM) through systematic model analysis: CanESM5.0 and CanESM5.1, *Geosci. Model Dev.*, 16, 6553–6591, <https://doi.org/10.5194/gmd-16-6553-2023>, 2023.
- Signorini, S. R. and McClain, C. R.: Effect of uncertainties in climatologic wind, ocean *p*CO<sub>2</sub>, and gas transfer algorithms on the estimate of global sea–air CO<sub>2</sub> flux, *Global Biogeochem. Cy.*, 23, GB2025, <https://doi.org/10.1029/2008GB003246>, 2009.
- Smith, R., Jones, P., Briegleb, B., Bryan, F., Danabasoglu, G., Dennis, J., Dukowicz, J., Eden, C., Fox-Kemper, B., Gent, P., Hecht, M., Jayne, S., Jochum, M., Large, W., Lindsay, K., Maltrud, M., Norton, N., Peacock, S., Vertenstein, M., and Yeager, S.: The Parallel Ocean Program (POP) reference manual ocean component of the Community Climate System Model (CCSM) and Community Earth System Model (CESM), Los Alamos National Laboratory Tech. Rep. LAUR-10-01853, 140 pp., <https://files.cesm.ucar.edu/models/pop/2/POPRefManual.pdf> (last access: 8 August 2026), 2010.
- Soloviev, A. and Lukas, R.: Effects of Bubbles and Sea Spray on Air–Sea Exchange in Hurricane Conditions, *Bound.-Layer Meteorol.*, 136, 365–376, <https://doi.org/10.1007/s10546-010-9505-0>, 2010.

- Stock, C. A., Dunne, J. P., Fan, S., Ginoux, P., John, J., Krasting, J. P., Laufkötter, C., Paulot, F., and Zadeh, N.: Ocean biogeochemistry in GFDL's Earth System Model 4.1 and its response to increasing atmospheric CO<sub>2</sub>, *J. Adv. Model. Earth Sy.*, 12, e2019MS002043, <https://doi.org/10.1029/2019MS002043>, 2020.
- Sutton, A. J., Sabine, C. L., Maenner-Jones, S., Lawrence-Slavas, N., Meinig, C., Feely, R. A., Mathis, J. T., Musielewicz, S., Bott, R., McLain, P. D., Fought, H. J., and Kozyr, A.: A high-frequency atmospheric and seawater pCO<sub>2</sub> data set from 14 open-ocean sites using a moored autonomous system, *Earth Syst. Sci. Data*, 6, 353–366, <https://doi.org/10.5194/essd-6-353-2014>, 2014.
- Sutton, A. J., Williams, N. L., and Tilbrook, B.: Constraining Southern Ocean CO<sub>2</sub> flux uncertainty using uncrewed surface vehicle observations, *Geophys. Res. Lett.*, 48, e2020GL091748, <https://doi.org/10.1029/2020GL091748>, 2021.
- Tjiputra, J. F., Schwinger, J., Bentsen, M., Morée, A. L., Gao, S., Bethke, I., Heinze, C., Goris, N., Gupta, A., He, Y.-C., Olivie, D., Seland, Ø., and Schulz, M.: Ocean biogeochemistry in the Norwegian Earth System Model version 2 (NorESM2), *Geosci. Model Dev.*, 13, 2393–2431, <https://doi.org/10.5194/gmd-13-2393-2020>, 2020.
- Tokoro, T., Hosokawa, S., Miyoshi, E., Tada, K., Watanabe, K., Montani, S., Kayanne, H., and Kuwae, T.: Net uptake of atmospheric CO<sub>2</sub> by coastal submerged aquatic vegetation, *Glob. Change Biol.*, 20, 1873–1884, <https://doi.org/10.1111/gcb.12543>, 2014.
- Van Dam, B., Polsenaere, P., Barreras-Apodaca, A., Lopes, C., Sanchez-Mejia, Z., Tokoro, T., Kuwae, T., Loza, L. G., Rutgersson, A., Fourqurean, J., and Thomas, H.: Global trends in air–water CO<sub>2</sub> exchange over seagrass meadows revealed by atmospheric Eddy Covariance, *Global Biogeochem. Cy.*, 35, e2020GB006848, <https://doi.org/10.1029/2020GB006848>, 2021.
- Wanninkhof, R.: Relationship between wind speed and gas exchange over the ocean, *J. Geophys. Res.-Oceans*, 97, 7373–7382, <https://doi.org/10.1029/92JC00188>, 1992.
- Wanninkhof, R.: Relationship between wind speed and gas exchange over the ocean revisited, *Limnol. Oceanogr.-Meth.*, 12, 351–362, <https://doi.org/10.4319/lom.2014.12.351>, 2014.
- Weiss, R. F.: Carbon dioxide in water and seawater: the solubility of a non-ideal gas, *Mar. Chem.*, 2, 203–215, [https://doi.org/10.1016/0304-4203\(74\)90015-2](https://doi.org/10.1016/0304-4203(74)90015-2), 1974.
- Williams, N. L., Juranek, L. W., Feely, R. A., Johnson, K. S., Sarmiento, J. L., Talley, L. D., Dickson, A. G., Gray, A. R., Wanninkhof, R., Russell, J. L., and Riser, S. C.: Calculating surface ocean pCO<sub>2</sub> from biogeochemical Argo floats equipped with pH: An uncertainty analysis, *Global Biogeochem. Cy.*, 31, 591–604, <https://doi.org/10.1002/2016GB005541>, 2017.
- Wu, L., Cai, Y., and Rutgersson, A.: Ocean surface waves impact on global air–sea CO<sub>2</sub> flux, *Biogeochemistry*, 168, 68, <https://doi.org/10.1007/s10533-025-01267-y>, 2025.
- Wu, Y. and Qi, D.: The controversial Southern Ocean air–sea CO<sub>2</sub> flux in the era of autonomous ocean observations, *Sci. Bull.*, 68, 2519–2522, <https://doi.org/10.1016/j.scib.2023.08.059>, 2023.
- Yang, X., Wynn-Edwards, C. A., Strutton, P. G., and Shadwick, E. H.: Drivers of air–sea CO<sub>2</sub> flux in the subantarctic zone revealed by time series observations, *Global Biogeochem. Cy.*, 38, e2023GB007766, <https://doi.org/10.1029/2023GB007766>, 2024.
- Yool, A., Palmieri, J., Jones, C. G., de Mora, L., Kuhlbrodt, T., Popova, E. E., Nurser, A. J. G., Hirschi, J., Blaker, A. T., Coward, A. C., Blockley, E. W., and Sellar, A. A.: Evaluating the physical and biogeochemical state of the global ocean component of UKESM1 in CMIP6 historical simulations, *Geosci. Model Dev.*, 14, 3437–3472, <https://doi.org/10.5194/gmd-14-3437-2021>, 2021.
- Zhou, X., Reichl, B. G., Romero, L., and Deike, L.: A sea state dependent gas transfer velocity for CO<sub>2</sub> unifying theory, model, and field data, *Earth Space Sci.*, 10, e2023EA003237, <https://doi.org/10.1029/2023EA003237>, 2023.
- Ziehn, T., Chamberlain, M., Law, R., Lenton, A., Bodman, R., Dix, M., Stevens, L., Wang, Y.-P., and Srbinovsky, J.: The Australian Earth System Model: ACCESS-ESM1.5, *J. South. Hemisph. Earth Syst. Sci.*, 70, 193–214, <https://doi.org/10.1071/ES19035>, 2020.