



Implementation of the Generalized Double-Moment scaling Normalization method for raindrop size distribution in a WRF 4.3.1 bulk-type cloud microphysics scheme: a case study over the Korean Peninsula

Joonghyun Jo¹, Kyo Sun Lim², Sun-Young Park³, Juhee Kwon², Wonbae Bang⁴, HyangSuk Park¹, Jae-Young Byon¹, and Gyuwon Lee⁴

¹National Institute of Meteorological Sciences, Korea Meteorological Administration, Seogwipo, Republic of Korea

²School of Earth and Environmental Sciences, Seoul National University, Seoul, Republic of Korea

³Climate Prediction Research Center, Seoul National University, Seoul, Republic of Korea

⁴BK21 Weather Extremes Education & Research Team, Department of Atmospheric Sciences, Center for Atmospheric REMote sensing (CARE), Kyungpook National University, Daegu, Republic of Korea

Correspondence: Kyo Sun Lim (kyosunlim@snu.ac.kr) and Gyuwon Lee (gyuwon@knu.ac.kr)

Received: 16 September 2025 – Discussion started: 20 February 2026

Revised: 16 July 2026 – Accepted: 9 August 2026 – Published: 2 September 2026

Abstract. To our knowledge, this study presents the first implementation of an observationally-constrained Generalized Double-Moment scaling Normalization (GDMN)-based rain Drop Size Distribution (DSD) representation within Weather Research and Forecasting (WRF) Double-Moment 6-class (WDM6) scheme and evaluates its impacts in a convection-permitting real-case simulation. The modified scheme, referred to as WDM6-GDMN, is evaluated through simulations of an isolated summer convection case over the Korean Peninsula, using the universal double-moment normalized DSD function, $h(x)$, derived from rain DSDs observed in the Boseong region during the summers of 2018 and 2019. WDM6-GDMN provides a more realistic spatial distribution of surface precipitation by better simulating convection-cell movement. Although none of the cloud microphysics parameterizations, including the bin-type scheme, reproduce the observed convection that developed in the southeast of the analysis domain, only WDM6-GDMN successfully captures this feature. Microphysical analysis demonstrates that, in WDM6-GDMN, enhanced cloud production due to stronger upward motion leads to the formation of more raindrops and, consequently, greater surface precipitation over southeastern region. Furthermore, the contoured frequency by altitude diagrams of radar reflectivity for the WDM6-GDMN reveals slower particle growth and weaker reflectivity in the lower

atmosphere compared with the original scheme, in better agreement with observations.

1 Introduction

Bulk-type cloud microphysics schemes adopt a specific functional form of the Drop Size Distribution (DSD) for each hydrometeor type (e.g., cloud water, rain, cloud ice, snow, and graupel/hail), rather than calculating the evolution of the DSD at each diameter bin (Lin et al., 1983; Rutledge and Hobbs, 1983; Schoenberg Ferrier, 1994; Meyers et al., 1997; Reisner et al., 1998; Morrison et al., 2005; Morrison and Milbrandt 2015; Thompson et al., 2008; Milbrandt and Yau, 2005). This approach to representing DSD evolution is computationally efficient, meaning that bulk-type cloud microphysics schemes can be applied to operational numerical weather or climate models. Among several microphysics schemes, the Weather Research and Forecasting (WRF) Double-moment 6-class (WDM6) bulk microphysics scheme (Lim and Hong, 2010) has been widely used across regions for research and operational purposes (Byun et al., 2011; Gao et al., 2011; Morrison et al., 2015; Lim et al., 2020).

Previous studies have shown that the WDM6 scheme tends to overestimate reflectivity, as demonstrated by contoured frequency by altitude diagram (CFAD) analyses (Min et al., 2015; Chakraborty et al., 2021). Min et al. (2015) evaluated the performance of the WRF Single-Moment 6-class (WSM6) and WDM6 cloud microphysics schemes using radar observations during summer season. They reported that although the WDM6 agrees better with radar observations than WSM6, both schemes overestimate the height of the melting level and the bright band relative to observations, as revealed by CFAD analysis. Additionally, WDM6 tends to overestimate reflectivity below the melting layer during summer monsoon cases. Chakraborty et al. (2021) compared WDM6 with other microphysics schemes, including the WSM6, Milbrandt, and Thompson Aerosol-aware schemes. They found that WDM6 tends to retain precipitation hydrometeors rather than precipitating them immediately, leading to greater accumulation of frozen hydrometeors in the upper layers. In the same study, CFAD analysis revealed that WDM6 produces the strongest reflectivity in the upper and middle levels among the microphysics schemes. This suggests unrealistic precipitation growth in the ice-phase region above the melting layer.

The WDM6 scheme adopts a rain DSD in the form of a generalized gamma distribution with a fixed shape parameter (μ), following the study by Cohard and Pinty (2000). Although most bulk microphysics schemes employ a fixed shape parameter, several observational studies have reported that the rain DSD shape parameter varies considerably (Yang et al., 2019; Uijlenhoet et al., 2003; Dolan et al., 2018; Cha et al., 2023; Lee et al., 2023). Yang et al. (2019) showed that the rain DSD shape parameter proposed by Testud et al. (2001) ranges from -2.3 to 60 over southern England during 2013–2017. Furthermore, they found that increasing the shape parameter initially leads to an underestimation of rainfall, followed by an overestimation. Uijlenhoet et al. (2003) noted that the rain DSD shape parameter varies depending on the stage of precipitation development and the type of cloud (e.g., stratiform or convective). Dolan et al. (2018) analysed 12 disdrometer datasets spanning three latitude bands – high ($\geq 45^\circ\text{N}$), middle ($23\text{--}45^\circ\text{N}$), and low ($\leq 23^\circ\text{N}$) – and various precipitation types, including light rain, orographic precipitation, deep convection, organized midlatitude systems, and tropical oceanic rainfall. They found that the mean shape parameter values of rain observed using a two-dimensional video disdrometer (2DVD), Joss-Waldvogel disdrometer (JWD), and automated particle size velocity (PARSIVEL) units (APU) vary by region, ranging from -1.20 to 7.26 . Similarly, Cha et al. (2023) compared DSD parameters observed with JWD and PARSIVEL across nine regions in Korea and East Asia and reported mean shape parameter values ranging from -0.158 to 8.37 . These findings indicate that using a fixed shape parameter in the gamma DSD model is not optimal and that more representative values should be applied.

Scaling normalization is commonly employed to represent the hydrometeor DSD and has been extensively studied since the 1990s (Sempere-Torres et al., 1994; Testud et al., 2001; Lee et al., 2004; Szyrmer et al., 2005; Berne et al., 2012; Morrison et al., 2019; Lee et al., 2023). Sempere-Torres et al. (1994) performed the first scaling normalization of DSDs based on a single reference variable, the rainfall rate (R). Later, Testud et al. (2001) extended this single normalization to a double-moment normalization using the third and fourth moments of the DSD. They showed that the mean raindrop volume diameter (D_m) exhibits significant variability, while the intercept parameter (N_0^*) shows no correlation with R . Lee et al. (2004) demonstrated that the normalization approach of Testud et al. (2001) was for the particular cases of the normalization in Lee et al. (2004) using the third and fourth moments. To address this limitation, Lee et al. (2004) generalized the normalization concept and proposed a general form of double-moment scaling normalization, $N(D)/N_0' = h(D/D_m')$, where the generalized characteristic number density (N_0') and generalized characteristic diameter (D_m') can be defined using any combination of two moments. The Generalized Double-Moment scaling Normalization (GDMN) significantly reduces the scatter in the normalized DSD compared to single-moment normalization and decreases the standard deviation of the fractional error in estimations of the other moments not used in the normalization, leading to a more accurate representation of the DSD.

Berne et al. (2012) compared the temporal and spatial variability of DSDs using the single-moment normalization method of Sempere-Torres et al. (1994) and the GDMN method of Lee et al. (2004) over the École Polytechnique Fédérale de Lausanne (EPFL) during March 2009 and July 2010. They showed that GDMN better represents the temporal and spatial variability of DSDs. The GDMN method can be also successfully applied to the retrieval of spatiotemporally varying DSDs from dual-polarimetric radar measurement (Kwon et al., 2020; Lee et al., 2023; Shin et al., 2024). Meanwhile, Morrison et al. (2019) extended the GDMN analysis by developing a generalized n th moment normalization using three or more moments with single- and double-moment normalization using disdrometer data obtained from the U.S. Department of Energy (DOE) Atmospheric Radiation (ARM) program sites worldwide. They noted that increasing the number of reference moments can reduce variability in the DSD parameters.

To evaluate the applicability of the GDMN method for representing the DSD in a bulk-type cloud microphysics scheme, this study develops a new version of WDM6, termed WDM6-GDMN, by incorporating the GDMN approach to represent the rain DSD, following Lee et al. (2004). The performance of WDM6-GDMN is examined through simulations of an isolated summer convection event using observed normalized DSD functions. Section 2 outlines the proposed method, Sect. 3 describes the experimental setup, case study,

and observational data, and Sects. 4 and 5 present the results and the summary and conclusions, respectively.

2 Generalized Double-Moment scaling Normalization (GDMN) method

2.1 Original Rain Drop Size Distribution (DSD) in WDM6

The original WDM6 scheme adopts the gamma form of the DSD for rain with a static shape parameter of c and μ , as shown in Eq. (1):

$$N_R(D_R) = N_{0R}(D_R)^{c\mu-1} \exp\{-(\lambda_R D_R)^c\}. \quad (1)$$

Where c and μ have values of 1 and 2, respectively; $N_R(D_R)$, with units of m^{-4} , is the number concentration corresponding to the rain diameter (D_R), which has units of meters; and N_{0R} and λ_R denote the intercept and slope parameters of the rain DSD, which have units of $\text{m}^{-3-c\mu}$ and m^{-1} , respectively, and can be expressed by the following equations:

$$N_{0R} = N_R \frac{c}{\Gamma(\mu)} (\lambda_R)^{c\mu}, \quad (2)$$

$$\lambda_R = \left[\frac{\pi \rho_R N_R \Gamma(\mu + \frac{d_R}{c})}{6 \rho_a q_R^2 \Gamma(\mu)} \right]^{\frac{1}{d_R}}, \quad (3)$$

where q_R and N_R represent the rain mixing ratio and number concentration, respectively, with units of kg kg^{-1} and m^{-3} ; ρ_R and ρ_a are the densities of the rain and air, respectively, with units of kg m^{-3} ; and d_R is the parameter used in the calculation of the mass (M)–diameter (D) relationship ($M = c_R D^{d_R}$). The diameter of rain, D , is an equivalent sphere diameter, which implies d_R equals 3. It is noteworthy that the units of N_{0R} depend on the two shape parameters, c and μ , expressed as $\text{m}^{-3-c\mu}$.

2.2 New DSD adopting the GDMN method

The DSD can be formulated as the product of the zeroth moment and a probability density function, and the following GDMN is then derived from the assumption of multiple power-law relationships among moments (Lee et al., 2004).

$$N(D) = N'_0 h(D/D'_m), \quad (4)$$

where N'_0 and D'_m are the generalized characteristic number density and the generalized characteristic diameter, with units of m^{-4} and m , respectively. They can be defined using any combination of two moments, as follows:

$$N'_0 = M_i^{(j+1)/(j-i)} M_j^{(i+1)/(i-j)}, \quad (5)$$

$$D'_m = (M_j/M_i)^{\frac{1}{j-i}}, \quad (6)$$

where i and j denote the moment order, and M_i and M_j represent the i th and j th moments of the DSD, respectively,

which are calculated as $M_i = \int N(D) D^i dD$. The $h(x)$ in Eq. (4) is the normalized DSD based on the two moments, M_i and M_j , where $x = D/D'_m$. It should be noted that no assumption on the functional form of $h(x)$ is applied. Thus, this GDMN can be applied to any observed DSD if the multiple power-law relationship is satisfied.

When the GDMN method is applied to the generalized gamma DSD, the normalized DSD takes the non-dimensional form $h(x)$.

$$h_{GG(i,j,\mu,c)}(x) = c \Gamma_i^{(j+c\mu)/(i-j)} \cdot \Gamma_j^{(-i-c\mu)/(i-j)} x^{c\mu-1} \exp \left[- \left(\frac{\Gamma_i}{\Gamma_j} \right)^{c/(i-j)} x^c \right], \quad (7)$$

where $\Gamma_i = \Gamma(\mu + i/c)$ and $\Gamma_j = \Gamma(\mu + j/c)$. The subscript $GG(i, j, \mu, c)$ in $h(x)$ denotes the generalized gamma DSD, and i, j, μ , and c in the parenthesis indicate the i th and j th moments and shape parameters. The advantage of expressing the DSD as $h(x)$ (Eq. 4) is that it provides a comprehensive framework to represent any naturally occurring DSD, while $h(x)$ remains relatively stable and less variable because most of significant variability is explained by the two reference moments (Lee et al., 2004).

2.3 Incorporation of GDMN method for rain DSD in WDM6

The WDM6-GDMN allows the observed shape parameters, c and μ , to be applied to the rain DSD. To calculate c and μ from observed rain DSDs over the Korean Peninsula, data from a 2DVD are utilized. The 2DVD is widely recognized as a reference instrument for the large end of the DSD but has been shown to underestimate the concentrations of small drops (Raupach et al., 2019), which significantly affects M_0 . Consequently, it is not appropriate to derive $h_{GG(0,3,\mu,c)}(x)$ directly from observed DSDs by normalizing with the zeroth and third moments. Therefore, this study follows two steps to obtain $h_{GG(0,3,\mu,c)}(x)$. Normalization is first performed using N'_0 and D'_m , defined with the third and fourth moments, M_3 and M_4 , for the observed rainfall events (threshold: $R \geq 0.1 \text{ mm h}^{-1}$) at the Boseong standard weather observatory during 2018 and 2019 (Fig. 3b). Then, $h_{GG(3,4,\mu,c)}(x)$ is derived from the mean $h(x)$ by minimizing $\sum_{i=20}^{i=1} (\log h_{\text{mean}(3,4,\mu,c)}(x_i) - \log h_{GG(3,4,\mu,c)}(x_i))^2$ (Bang et al., 2020), where $h_{\text{mean}}(x)$ is the bin-wise mean of the normalized DSDs and the bin interval is 0.2. The derived parameters of $h_{GG(3,4,\mu,c)}(x)$ are $c=2.70$ and $\mu=0.24$. Then, the theoretical DSDs, $N_R(D)$, are derived from $h_{GG(3,4,\mu,c)}(x)$, $N'_0(M_3, M_4)$, and $D'_m(M_3, M_4)$:

$$N_R(D) = N'_0(M_3, M_4) h_{GG(3,4,\mu,c)}(D/D'_m(M_3, M_4)). \quad (8)$$

The theoretical $N_R(D)$ is then normalized by $N'_0(M_0, M_3)$ and $D'_m(M_0, M_3)$, where M_0 and M_3 are derived from




Figure 1. Normalized DSDs derived from $N'_0(M_0, M_3)$ and $D'_m(M_0, M_3)$, along with the $h_{\text{mean}(0,3,\mu,c)}(x)$ from the theoretical $N_R(D)$ in Eq. (8) over the Boseong area. Colors indicate the normalized frequency of the normalized DSDs with M_0 and M_3 and the solid line represents $h_{\text{mean}(0,3,\mu,c)}(x)$ while the dotted line represents its best fit obtained using with $h_{\text{GG}(0,3,\mu,c)}(x)$ with $c = 2.60$ and $\mu = 0.29$.

$N_R(D)$. The normalized DSDs, $h_{(0,3,\mu,c)}(x)$, are presented as a frequency distribution in Fig. 1. Finally, $h_{\text{GG}(0,3,\mu,c)}(x)$ is obtained by minimizing the value of $\sum (\log h_{\text{mean}(0,3,\mu,c)}(x) - \log h_{\text{GG}(0,3,\mu,c)}(x))^2$. The derived shape parameters c and μ of $h_{\text{GG}(0,3,\mu,c)}(x)$ are 2.60 and 0.29, respectively. Previous research by Bang et al. (2020) demonstrated that the normalized DSD function $h(x)$, derived from observations over South Korea (2011–2015) and Oklahoma (1998–2006), USA, exhibits no substantial differences despite the distinct geographical characteristics of the two regions. These findings support the robustness of the observation-derived $h(x)$ representation and suggest that it is sufficiently general for the intended modeling application in this study.

Recognizing the importance of rain-related microphysical processes such as collision–coalescence and raindrop breakup in precipitation initiation, storm intensity, and cold-pool dynamics (Lim and Hong, 2012; Morrison, 2012; Hagos et al., 2015), this study introduces the first application of the GDMN method to the rain DSD in the WDM6 scheme. To apply the GDMN method of Lee et al. (2004) in WDM6 using the zeroth ($i = 0$) and third ($j = 3$) moments, which are the prognostic variables of rain in WDM6, the original rain DSD ($N_R(D_R)$) and rain diameter (D_R) must be normalized by $N'_0 = (M_0^4/M_3)^{1/3}$ and $D'_m = (M_3/M_0)^{1/3}$ (Eqs. 5 and 6). With this normalization, the rain DSD can be rewritten as Eq. (9):

$$N_R(D_R) = N'_0 h_{\text{GG}(0,3,\mu,c)}(D_R/D'_m), \quad (9)$$

where $h_{\text{GG}(0,3,\mu,c)}(x_R)$ with $x_R = D_R/D'_m$ can be expressed as Eq. (10):

$$h_{\text{GG}(0,3,\mu,c)}(x_R) = \frac{c}{\Gamma(\mu)} \left(\frac{\Gamma(\mu + \frac{3}{c})}{\Gamma(\mu)} \right)^{\frac{c\mu}{3}} \cdot x_R^{c\mu-1} \exp \left[- \left\{ \left(\frac{\Gamma(\mu + \frac{3}{c})}{\Gamma(\mu)} \right)^{\frac{1}{3}} (x_R) \right\}^c \right]. \quad (10)$$

The original WDM6 scheme adopts a single parameter, $\mu_R = c\mu - 1$, instead of treating c and μ separately. This is equivalent to employing fixed shape parameters, $c = 1$ and $\mu = 2$, for the gamma DSD in Eq. (1), resulting in the form $N(D) = N_0 D \exp\{-\lambda D\}$. This implies that μ_R remains unchanged whether $c = 1$, $\mu = 2$ or $c = 2$, $\mu = 1$. As shown by Lee et al. (2004, 2023), the variability of DSDs, arising from changes in number concentration, mean diameter, and shape, can be substantially reduced through scaling normalization. Variations in the shape parameters c and μ have only a minor effect on the normalized DSD, $h(x)$, which therefore becomes stable and nearly universal. The universal $h(x)$ function provides a more effective representation of DSD variability, as implemented in WDM6-GDMN.

2.4 Microphysics

The WDM6 cloud microphysics parameterization scheme predicts the mixing ratios of six hydrometeor types (water vapor, cloud water, cloud ice, rain, snow, and graupel) as well as the total number concentrations of cloud water, rain, and cloud condensation nuclei. Figure 2 presents a flowchart of the WDM6 scheme, illustrating the various microphysical processes among hydrometeors. By applying the GDMN method, $N_R(D_R)$ is expressed in terms of $h_{\text{GG}(0,3,\mu,c)}(x_R)$, as shown in Eq. (9). Consequently, the parameterization of 16 rain-related microphysical processes must be re-derived: nine associated with mixing-ratio predictions and seven with total number concentration predictions. Fourteen of these processes are indicated by green boxes in Fig. 2, excluding the two processes related to sedimentation of the rain mixing ratio and its number concentration. A list of the symbols used in this study is provided in Appendix A.

3 Experimental setup

3.1 Case description and model setup

Song and Sohn (2015) classified summer rainfall events over the Korean Peninsula into two types: cold-type and warm-type. Cold-type events, driven by convective instability, are typically locally developed and short-lived, making them difficult to accurately simulate in numerical models. Compared with warm-type events, cold-type rainfall exhibits stronger




Figure 2. Flowchart of the microphysical processes for the prediction of (a) mixing ratios and (b) number concentrations in the WDM6 scheme (Lim and Hong, 2010). The terms in red (blue) are activated when the temperature is above (below) 0 °C, whereas the terms in black occur across the entire temperature range. The green rectangles represent newly re-derived microphysics processes.

convective instability, allowing clouds to develop to higher altitudes and producing more intense rainfall. In this study, one cold-type summer rainfall event that occurred on 6 August 2013 is analyzed to test the impact of applying the GDMN method to the rain DSD within WDM6 scheme. The selected event exhibited the highest frequency of lightning strikes over the heavy precipitation area during summer of 2013–2018.

The model forecast spans the period from 23:00 UTC on 5 August 2013 to 17:00 UTC on 6 August 2013, with the analysis period from 02:00–17:00 UTC on 6 August 2013. Version 4.3.1 of the WRF model (Skamarock et al., 2019) is employed, with three nested domains using a horizontal grid spacing of 9, 3, and 1 km, as shown in Fig. 4. All domains have 65 vertical levels, with time steps of 45, 15, and 5 s for domains 1 (d01), 2 (d02), and 3 (d03), respectively. The physics parameterizations include the Kain–Fritsch cumulus scheme for D01 only (Kain and Fritsch, 1990; Kain, 2004), the Revised MM5 Monin–Obukhov surface layer scheme (Jiménez et al., 2012), the Unified Noah land surface scheme (Chen and Dudhia, 2001), the Rapid Radiative Transfer Model for General Circulation Models (RRTMG) for both longwave and shortwave radiation (Iacono et al., 2008; Morcrette et al., 2008), and the Yonsei University planetary boundary layer scheme (Hong et al., 2006). The European Centre for Medium-Range Weather Forecasts (ECMWF) Re-analysis 5 (ERA5) dataset (Hersbach et al., 2020) provides the initial and boundary conditions. The model configuration and physical parameterizations for each domain are summarized in Table 1.

3.2 Observation Data and Numerical Experiments

Radar reflectivity observed from two the Kwanaksan (KWK) and Gwangdeuksan (GDK) sites was analysed using Contoured Frequency by Altitude Diagrams (CFADs; Fig. 3), which illustrate the frequency of occurrence of reflectivity

values at various heights. The CFAD analysis is used to evaluate the impact of incorporating the GDMN method into the WDM6 scheme on the vertical structure of reflectivity. Reflectivity is examined using all model grid points located within a 100 km radius of each radar site, where observational data available. The KWK and GDK radars provide data at 10 min intervals, whereas both the original WDM6 and the modified WDM6-GDMN schemes produce outputs at 1 h intervals. To enable a direct one-to-one temporal comparison between the radar and model results, radar data collected within 30 min before and after each hour were compared with the corresponding hourly model outputs.

To examine the effect of the GDMN method on the rain DSD in simulated convection, two experiments, named WDM6 and GDMN, are conducted for the selected case. WDM6 uses the shape parameters $c = 1$ and $\mu = 2$ in the rain DSD, expressed by the gamma function as in the original WDM6 scheme. GDMN applies the WDM6-GDMN (see Sect. 2) with the observed shape parameters of $c = 2.60$ and $\mu = 0.29$ (Fig. 1). The differences in microphysical processes, particularly rain evaporation and near-surface cooling between WDM6 and GDMN, are not sensitive to the prescribed c and μ parameters in GDMN, as these parameters do not vary substantially within the generalized DSD function. To further evaluate simulated precipitation across different cloud microphysics schemes, additional experiments are conducted using three other double-moment bulk-type microphysics schemes – Morrison, Thompson, Predicted Particle Properties (P3) – as well as the simplified Spectral Bin-type Microphysics (SBM) scheme.

4 Results

Figure 5 shows the observed radar reflectivity at 06:00, 07:00, and 08:00 UTC on 6 August 2013, along with the simulated radar reflectivity from the WDM6 and GDMN ex-




Figure 3. (a) Surface synoptic chart at 03:00 UTC on 6 August 2013, and (b) 15 h accumulated surface precipitation (mm) during the analysis period (02:00–17:00 UTC on 6 August 2013), obtained from the Korea Meteorological Administration (KMA) Automatic Weather Station (AWS) network. The red dots in (b) indicate the locations of the Kwanaksan (KWK) and Gwangdeuksan (GDK) radar sites used to validate simulated radar reflectivity. The red solid line from A to B denotes the transect for cross-sectional analysis, while the red dashed box indicates the area used for microphysical process analysis. The red triangle marks the Boseong standard weather observatory, where a two-dimensional video disdrometer (2DVD) was installed to observe rain drop size distributions (DSDs).

Table 1. A summary of the Weather Research and Forecasting (WRF) model configuration and references for all physics schemes except for the cloud microphysics scheme.

	WRF V4.3.1			References
	Domain 1	Domain 2	Domain 3	
Grid number ($x \times y \times z$)	$180 \times 180 \times 65$	$295 \times 349 \times 65$	$532 \times 691 \times 65$	
Horizontal grid spacing	9 km	3 km	1 km	
Cumulus parameterization	Kain-Fritsch	No cumulus	No cumulus	Kain and Fritsch (1990), Kain (2004)
Surface layer	Revised MM5 Monin–Obukhov scheme			Jiménez et al. (2012)
Land surface	Unified Noah land surface model			Chen and Dudhia (2001)
Longwave and shortwave radiation	Rapid Radiative Transfer Model for General Circulation Models			Iacono et al. (2008)
Planetary Boundary Layer	Yonsei University scheme			Hong et al. (2006)
Initial/boundary data	ERA5 (fifth generation ECMWF reanalysis product)			Hersbach et al. (2020)

periments. The Column Maximum (CMAX) observations at 06:00 UTC reveal two distinct regions of strong reflectivity (Fig. 5a): one over the northwestern Korean Peninsula, which moves south-eastward over time, and another, consisting of a smaller convection cell, in the south of the peninsula, moving eastward (Fig. 5a–c). By 08:00 UTC, these two convection cells converge, producing strong reflectivity over the south-eastern Korean Peninsula (Fig. 5c). The WDM6 scheme does not reproduce this movement accurately, especially for the northwestern system. Although the initial stages of the two precipitation cells in WDM6 are similar to the observations (Fig. 5a and d), the stronger northwestern cell fails to move

south-eastward, shifting mostly eastward instead (Fig. 5d–f). As a result, the two precipitation cells move eastward independently and do not merge, as observed. Conversely, GDMN reproduces the movement of the two precipitation cells more realistically (Fig. 5g–i). Although weak reflectivity signals remain over the western Korean Peninsula, the two cells merge in GDMN, producing intense reflectivity over the southeastern peninsula at 08:00 UTC, consistent with observations (Fig. 5g–i).

The spatial distribution of cumulative surface precipitation on 6 August 2013 is shown in Fig. 6a and b. Compared with AWS observations, WDM6 overestimates precipitation over




Figure 4. The model domain, consisting of three nested domains with resolutions of 9 km (d01), 3 km (d02), and 1 km (d03). Shading indicates the terrain height in meters above sea level. The analysis domain is denoted by the black dotted line within the innermost domain (d03).

Table 2. Statistical skill scores for the simulated precipitation over the analysis domain during the analysis period: Root Mean Square Error (RMSE), Bias, Probability of Detection (POD), False Alarm Ratio (FAR), Equivalent Threat Score (ETS) and Fractions Skill Score (FSS).

	RMSE	BIAS	POD	FAR	ETS	FSS
WDM6	6.72	2.0	0.16	0.70	0.05	0.13
GDMN	5.03	1.39	0.18	0.68	0.07	0.14

the northeastern Korean Peninsula and underestimates it in the west and south (Fig. 6c). In WDM6, precipitation cells approaching from the northwest move eastward, leading to excessive surface precipitation in the northeast. In contrast, GDMN produces more precipitation in the southeast due to the merging of two precipitation cells in that area, thereby reducing the bias of WDM6 relative to AWS (Fig. 6d). This reduction in bias is attributed to GDMN's more accurate simulation of the movement of precipitation cells initially located in the northwest. To quantify the spatial distribution of accumulated surface precipitation, statistical skill scores – including Root Mean Square Error (RMSE), Bias, Probability of Detection (POD), False Alarm Ratio (FAR), Equivalent Threat Score (ETS) and Fractions Skill Score (FSS) – are calculated for WDM6 and GDMN (Table 2). Although GDMN underestimates total surface precipitation relative to observations, all statistical skill scores show improvement compared with WDM6. The calculated FSS also indicates that GDMN outperforms WDM6.

To further evaluate the performance of GDMN method, CFADs from the KWK and GDK sites are analysed for two simulations (Fig. 7). The location of the two sites used in the CFAD analysis are shown in Fig. 3b. The highest precipitation during the analysis period was recorded at KWK, whereas only a small amount was observed at GDK, despite the model simulations indicating substantial precipitation at this site. In Fig. 7, the x axis represents reflectivity, the y axis represents altitude, and the colour scale indicates the frequency ratios (%) at specific altitudes and reflectivity values. The reflectivity values range from 0 to 60 dBZ. Black solid lines denote the cumulative reflectivity frequencies at the 25th, 50th, and 75th percentiles. The simulated radar reflectivity is calculated following the methodologies of Koch et al. (2005), Stoelinga (2005), and Park et al. (2025). The model incorporates a simple water coating method for melting solid-phase particles to represent the bright band.

At KWK, between 10 and 12 km in altitude, weak reflectivity values of approximately 6–7 dBZ are frequently observed, gradually increasing toward 4 km, where precipitation particles grow. The median reflectivity below 4 km is about 20 dBZ. In WDM6, the reflectivity is generally overestimated relative to radar observations. No distinct melting layer is simulated, and the increase in reflectivity with decreasing altitude down to the 4 km is similar to observations. However, between 6 and 10 km, the increase occurs more rapidly than in the observations. WDM6 produces a median reflectivity of 30 dBZ below 4 km, about 10 dBZ higher than radar observations, and frequently reproduces strong reflectivity exceeding 40 dBZ. In contrast, GDMN exhibits a slower increase in reflectivity with decreasing altitude compared with WDM6 above 6 km, making it closer to observations. Although GDMN does not reproduce the narrow, distinct bright band observed near the melting layer likely due to the forward operator within WRF, it well reproduces reflectivity intensity below 4 km, bringing it into better agreement with the observations. GDMN also shows a decrease in reflectivity toward the surface, indicating possible substantial evaporation in the lower layers.

For the GDK site, peak reflectivity is observed at 4 km, indicating the presence of a distinct melting layer at this altitude. Radar observations also show gradually increasing reflectivity from 12 to 4 km, similar to the KWK site, followed by a decrease from 4 km to the surface. In WDM6, reflectivity increases rapidly from the top to 4 km, and the bright band is not distinctly simulated. This rapid increase in reflectivity, inconsistent with observations, was also noted in Min et al. (2015), indicating improper growth of solid particles in the WDM6 scheme. Additionally, WDM6 overestimates reflectivity below 4 km by about 10 dBZ compared with radar observations. Particle growth in GDMN is slower than in WDM6 over the upper layers, consistent with observations. GDMN also reproduces a clear trend of decreasing reflectivity below the melting layer at GDK. Although discrepancies with the observations still remain after apply-




Figure 5. The spatial distribution of maximum reflectivity (dBZ) at 06:00 UTC (left), 07:00 UTC (middle), and 08:00 UTC (right) on 6 August 2013 from (a–c) the Column Maximum (CMAX) radar observations, (d–f) WDM6, and (g–i) GDMN.

ing the GDMN approach, particularly above approximately 6 km, GDMN overall reduces the overestimation of reflectivity compared with WDM6.

Figure 8 shows the 1 h accumulated surface precipitation along the cross-section shown in Fig. 3b from 04:00 to

08:00 UTC on 6 August. The cross-section spans the path of the precipitation cell, which AWS observations show was initially located over the northwestern Korean Peninsula and moved south-eastward over time (Fig. 3b). This movement is well simulated in GDMN but not captured in WDM6




Figure 6. Accumulated surface precipitation (mm) during the analysis period (from 02:00 to 17:00 UTC on 6 August 2013) for (a) WDM6, (b) GDMN, and (c) the difference between WDM6 and Automatic Weather Station (AWS) observations (WDM6 minus AWS). (d) Same as (c), but for the difference between GDMN and WDM6 (GDMN minus WDM6).




Figure 7. Contoured Frequency by Altitude Diagram percentiles at the KWK and GDK radar sites (location shown in Fig. 3b) for (a) radar observations, (b) WDM6, and (c) GDMN at the KWK site during the analysis period. Panels (d)–(f) are the same as (a)–(c) but at the GDK site. Solid lines denote the cumulative reflectivity frequencies at the 25th, 50th, and 75th percentiles.




Figure 8. 1 h accumulated surface precipitation along the cross-sectional area indicated by the red solid line in Fig. 3b for (a) 04:00 to 05:00 UTC, (b) 05:00 to 06:00 UTC, (c) 06:00 to 07:00 UTC, and (d) 07:00 to 08:00 UTC on 6 August 2013. Black and blue lines represent WDM6 and GDMN, respectively.

(Fig. 5d–i). In both WDM6 and GDMN, precipitation begins to develop in the northwest (Location “A” in Fig. 8) after 04:00 UTC (Fig. 8a) and gradually shifts south-eastward (Location “B” in Fig. 8) with time (Fig. 8b–d). WDM6 simulates a persistent precipitation peak near “A” from 04:00 to 06:00 UTC failing to capture the south-eastward movement of precipitation cells (Fig. 5). By contrast, in GDMN, the precipitation peak shifts from Area A to Area B between 04:00 to 08:00 UTC (Fig. 8a–d). Between 05:00 and 06:00 UTC, the maximum precipitation location in GDMN is similar to that in WDM6; however, it subsequently shifts markedly south-eastward. The difference between the two schemes arises from GDMN’s more realistic representation of precipitation cell movement.

To identify the main causes of the differences in simulated precipitation and reflectivity between GDMN and WDM6, thermodynamic fields and mixing ratios are further analysed in Figs. 9 and 10, focusing on 07:00 and 08:00 UTC, when two precipitation cells merged into a single cell. Figure 9 presents the vertical structures of temperature and wind along the cross-sectional area shown in Fig. 3b. In WDM6, a strong updraft develops near 37.3° N at 07:00 UTC and dissipates by 08:00 UTC, producing intense precipitation at the same location during this hour (Figs. 9a and 8d). As the precipitation cells originating in the northwest move eastward, the updraft driving the convection at 07:00 UTC is no longer visible at 08:00 UTC. In contrast, GDMN simulates an updraft farther southeast, near 36.5° N at 07:00 UTC (Fig. 9a and c). The south-eastward-moving precipitation cell merges with the eastward-moving cells and continues moving south-

eastward, maintaining an updraft near 35.5° N at 08:00 UTC (Fig. 9d). Additionally, GDMN produces a colder environment below 1 km compared with WDM6 (Fig. 9). The stronger and more extensive near-surface cooling in GDMN drives the precipitation cells south-eastward, whereas the relatively weak near-surface cold environment in WDM6 limits the south-eastward propagation of the system.

At 07:00 and 08:00 UTC, compared to WDM6, GDMN exhibits stronger cooling in the lower layer behind the precipitation cells (Fig. 10a and b). The differences in cloud and rain mixing ratios between GDMN and WDM6 show a similar trend. At 07:00 UTC, the cloud mixing ratio increases significantly near 36.5° N in GDMN, coinciding with a region of strong updrafts and an enhanced rain mixing ratio (Fig. 10c and e). A comparable pattern occurs around 35.5° N at 08:00 UTC (Fig. 10d and f), where vigorous upward motion generates a supersaturated environment, leading to cloud formation and, consequently, precipitation both aloft and at the surface.

Figure 11 shows the differences (GDMN minus WDM6) in the time-area averaged vertical distributions of prognostic hydrometeor mixing ratios (cloud water (q_C), cloud ice (q_I), rain (q_R), snow (q_S), and graupel (q_G)) and number concentrations (cloud condensation nuclei (N_N), cloud water (N_C), and rain (N_R)). Below 9 km, both the rain mixing ratio and the number concentration are significantly higher in GDMN compared to WDM6. In addition, GDMN exhibits slightly higher cloud water mixing ratios and number concentrations below 6 km, as well as greater graupel mixing ratios above




Figure 9. Temperature with terrain along the cross-section indicated by the red line in Fig. 3b at (a) 07:00 UTC, (b) 08:00 UTC on 6 August 2013 for WDM6. Panels (c) and (d) are the same as (a) and (b), but for GDMN. The wind fields are overlaid at the corresponding times. Red contour lines indicate 20 dBZ reflectivity.

3 km. The enhanced graupel production in GDMN is likely a consequence of stronger convective updrafts in the model.

To identify the microphysical processes responsible for the increased generation of rain and cloud water in GDMN compared to WDM6, the vertical profiles of the time-area averaged major source and sink terms for the rain and cloud mixing ratios are analysed in Fig. 12. In GDMN (solid line), the condensation process of water vapor into cloud water (Pcond) is enhanced below 9 km relative to WDM6 (dashed line) under saturated conditions, serving as the primary source of the cloud mixing ratio (Fig. 12a). The higher cloud mixing ratio in GDMN also intensifies accretion with rain (Pracw) (Fig. 12b), which acts not only the main sink of cloud water but also the main source of rain (blue solid line in Fig. 12c). Among the major sources of rain, Pracw produces a greater rain mixing ratio in GDMN than in WDM6. Moreover, graupel melting (Pgmlt), the second-largest source of the rain mixing ratio, is higher in GDMN owing to greater graupel production. The larger rain content in GDMN further increases ascending rain, which subsequently enhances graupel formation through raindrop freezing (Pgfrz) and accretion of rain by graupel (Pgacr) (Fig. 12d). Notably, rain evaporation (Prevp) is the primary sink of rain in the lower atmosphere, particularly below 5 km. Enhanced Prevp in GDMN compared to WDM6 leads to increased evaporation of rain into water vapor, cooling the lower atmosphere. As a result, precipitation cells in GDMN tend to propagate south-eastward, consistent with the observed behaviour.

5 Summary and Discussion

This study applies the Generalized Double-moment scaling Normalization (GDMN) method to parameterize the rain drop size distribution (DSD) in the WDM6 microphysics scheme. The WDM6 and GDMN method employ the same generalized gamma DSD, and therefore they produce identical simulation results when the same shape parameters are used. The current version of WDM6 adopts fixed shape parameters ($c = 1$ and $\mu = 2$) following the previous study (Lim and Hong, 2010). The key advantage of GDMN method is that it constrains the shape parameters using the normalized DSD, which is considerably more stable than the raw DSD. As a result, GDMN method provides a robust and observationally constrained normalized function, $h(x)$, whereas direct estimation of the generalized gamma shape parameters from raw DSDs, as in WDM6, is subject to substantial variability and often leads to unreliable parameter estimates. This study utilizes the fitted generalized gamma function, $h_{GG}(x)$, rather than the observed mean normalized function, $h_{mean}(x)$, because disdrometer measurements are subject to significant uncertainty for small drops. By adopting the GDMN method, the variability of the DSD is more effectively captured, thereby reducing uncertainties of DSD. The 0th and 3rd moments are used to represent the DSD within GDMN, and sixteen microphysical processes related to rain reformulated. Long-term observed rain shape parameter values ($c = 2.6$, $\mu = 0.29$), calculated from 2DVD data col-




Figure 10. Differences (GDMN – WDM6) in (a) temperature ($^{\circ}C$), (c) cloud mixing ratio ($g\ kg^{-1}$), and (e) rain mixing ratio ($g\ kg^{-1}$), shown with terrain (black color) along the cross-section indicated by the red line in Fig. 3b at 07:00 UTC, 6 August 2013. Panels (b), (d), and (f) show the same fields as (a), (c), and (e), respectively, at 08:00 UTC.




Figure 11. Differences in the time-domain averaged vertical distribution (GDMN minus WDM6) over the region denoted by the red dashed box in Fig. 3b for (a) mixing ratio ($g\ kg^{-1}$) and (b) number concentration (m^{-3}) between 06:00 and 08:00 UTC on 6 August 2013.

lected over the Boseong area during May–October of 2018 and 2019, are applied to the DSD through the GDMN method in WDM6. Numerical experiments are conducted for a cold-type summer rainfall event characterized by strong convection.

The results demonstrate that incorporating GDMN into the WDM6 scheme alleviates some limitations of the original WDM6 scheme, which misrepresents precipitation cell propagation and produces accumulated precipitation patterns that deviate from observations for this case. In the modified scheme, precipitation cells move south-eastward and subsequently collide with another cell generated within the Korean Peninsula, leading to increased precipitation in the southeastern regions. Overall, the modified WDM6 reproduces a more realistic propagation of precipitation cells and a more accurate spatial distribution of accumulated surface precipitation than the original scheme, achieving higher scores across all statistical skill metrics for surface precipitation.

To evaluate the simulated vertical reflectivity structure, CFAD analysis is performed using data from two radar sites,




Figure 12. The vertical profiles of the time-domain averaged (a) sources and (b) sinks of the cloud mixing ratios ($\text{g kg}^{-1} \text{s}^{-1}$) over the region indicated by the red dashed box in Fig. 3b between 06:00 and 08:00 UTC on 6 August 2013. Only the major microphysical processes are represented. The solid and dashed lines represent GDMN and WDM6, respectively. Panels (c) and (d) show the same as (a) and (b) but for the rain mixing ratio ($\text{g kg}^{-1} \text{s}^{-1}$).

KWK and GDK. The original WDM6 scheme exhibits excessively rapid particle growth between the upper and middle layers (4–5 km) and consistently overestimates reflectivity at all altitudes. In contrast, the modified WDM6 scheme produces more gradual particle growth from the top to the middle layers, with significantly reduced reflectivity, resulting in better agreement with observed patterns. These results indicate that the modified rain DSD influences not only warm-rain microphysical processes but also mixed-phase ones, including accretion between liquid- and solid-phase particles. Consequently, the simulated reflectivity in the upper layer is more realistically represented.

Thermodynamic analysis along the path of cell movement reveals that the modified WDM6 scheme successfully triggers convective updrafts over the southeastern Korean Peninsula, which are absent in the original WDM6 scheme. In the modified WDM6, enhanced condensation of water vapor into cloud water over the updraft region increases cloud formation and promotes the production of numerous small raindrops through collision–coalescence and graupel melting. Furthermore, the modified WDM6 simulates a relatively cooler lower environment due to stronger rain evaporation, which enhances near-surface cooling. This intensified cool-

ing drives the south-eastward propagation of precipitation cells, consistent with observations.

None of the bulk or bin microphysics schemes analysed in this study accurately reproduce the observed precipitation distribution across the Korean Peninsula on 6 August 2013 (Figs. 13 and 3b). All simulations employing bulk-type cloud microphysics schemes, which adopt a gamma DSD with constant shape parameters, generate spatial precipitation patterns similar to the original WDM6, with excessive precipitation over the central–eastern Korean Peninsula compared to AWS observations (Fig. 13a–c). The prescribed DSD function and fixed shape parameters in bulk schemes and other parameterization assumption or introduced uncertainties in microphysics schemes constrain their ability to realistically simulate precipitation. Tapiador et al. (2026) emphasized that the DSD strongly influences the physical properties of precipitation systems, including total surface area and scattering characteristics of drops, and highlighted the need for DSD modelling to be grounded in a physically consistent framework in which drop shapes follow an appropriate probability density function (PDF). Even the SBM scheme, which explicitly predicts the particle size distribution, fails to reproduce the observed spatial distribution of accumulated precipitation (Fig. 13d).

To address the limitations of the single-case analysis, we additionally conduct a single month-long simulation over East Asia for July 2023, encompassing a wide range of summertime convective systems. The model configuration, including the physical parameterizations and initial and lateral boundary conditions, is identical to that used in the single-case simulation, except that a horizontal grid spacing of 10 km and a model time step of 60 s are employed. Sea surface temperature (SST) is prescribed using the Optimum Interpolation SST (OISST) dataset (Huang et al., 2021). The performance of the original WDM6 and the modified WDM6 is evaluated against hourly precipitation from the Integrated Multi-satellite Retrievals for GPM (IMERG) product (Huffman et al., 2015) (Fig. 14). Compared with the original WDM6, the modified WDM6 scheme reduces the precipitation over Northeastern China and increases the precipitation over the South Sea of Korea, resulting in better agreement with the observations. The statistical skill scores for surface precipitation also demonstrate the improved performance of the modified WDM6, with the pattern correlation (PC) increasing from 0.47 to 0.52 and the bias decreasing from 1.04 to 1.02. These results indicate that the performance improvements achieved by the GDMN approach are maintained in a one-month simulation encompassing a broad range of summertime convective systems, suggesting that its benefits extend beyond the single convective case examined in this study. Our findings also highlight the potential of the GDMN approach for the examined case under the specific model configuration used in this study, while recognizing that further evaluation is required before its superiority can be generalized to a broader range of convective con-




Figure 13. The accumulated surface precipitation (mm) on 6 August 2013 using (a) Morrison, (b) Thompson, (c) P3, and (d) SBM cloud microphysics schemes.




Figure 14. Spatial distribution of hourly precipitation [mm h^{-1}] from (a) the IMERG observations and model simulations using (b) WDM6 and (c) GDMN for the one-month simulation conducted during July 2023.

ditions. Therefore, our study should be regarded primarily as a proof-of-concept demonstration based on limited cases simulations. Additional multi-case evaluations under diverse meteorological conditions, along with the extension of the GDMN framework to other hydrometeor types, are needed in future work.

Appendix A: List of symbols

Symbol	Meaning	Value	SI unit
c	Shape parameter of the rain		
c_R	Empirical formula of M_R	$\pi \rho_R / 6$	
D	Diameter		m
D'_m	Generalized characteristic diameter	$(M_j / M_i)^{\frac{1}{j-i}}$	m
D_R	Rain diameter		m
d_R	Empirical formula of M_R	3	
I	The number of moment		
j	The number of moment		
M_i	i th moment		
M_j	j th moment		
M_0	0th moment		m^{-3}
M_3	Third moment		
$M_R(D_R)$	Mass of rain of diameter D_R		kg
$N_R(D_R)$	Number concentration corresponding to the diameter of rain (D_R).		m^{-3}
$N(D)$	Number concentration corresponding to the diameter (D).		m^{-3}
N_{0R}	Intercept parameter of rain	8×10^6	m^{-4}
N'_0	Generalized characteristic number concentration	$M_i^{\left(\frac{i+1}{j-i}\right)} M_j^{\left(\frac{i+1}{i-j}\right)}$	m^{-4}
N_R	Number concentration of rain		m^{-3}
q_R	Mixing ratio of rain		
Γ	Gamma function		
λ_R	Slope of rain size distribution		m^{-1}
μ	Shape parameter of the rain		
π	Pi	3.14	
ρ_0	Density of air at reference state	1.28	kg m^{-3}
ρ_a	Air density		kg m^{-3}
ρ_R	Density of rain	1000	kg m^{-3}

Code and data availability. The WRF Model version 4.3.1 and the input files required for model integration are available at <https://doi.org/10.5281/zenodo.18328177> (Jo, 2026a). Initial and boundary conditions are provided from the ERA5 reanalysis data at 6 h intervals, available at <https://doi.org/10.24381/cds.adbb2d47> (Hersbach et al., 2023a) and <https://doi.org/10.24381/cds.bd0915c6> (Hersbach et al., 2020, 2023b). ERA5 single-level and pressure-level fields at 6 h intervals are used, with the initial time set to 23:00 UTC on 5 August 2013. For the pressure-level variables, all vertical levels are included. The model codes for WDM6 and GDMN experiments with the scripts for figures in this manuscript can be found in <https://doi.org/10.5281/zenodo.17194841> (Jo, 2025) and <https://doi.org/10.5281/zenodo.18346989> (Jo, 2026b). The AWS and radar data used for the analysis are available at <https://doi.org/10.5281/zenodo.18328177> (Jo, 2026a). The output data for four additional microphysics schemes (Thompson, Morrison, P3, and SBM) used for comparison with the WDM6 and GDMN experiments are also available at <https://doi.org/10.5281/zenodo.18346989> (Jo, 2026b).

Author contributions. JJ conducted the model simulations and performed the data analysis under the supervision of KL and GL. The manuscript was written by KL, JJ, and GL with substantial input from all co-authors. WB processed the DSD observational dataset. KL and other authors contributed to scientific discussions and provided constructive feedback.

Competing interests. The contact author has declared that none of the authors has any competing interests.

Disclaimer. Publisher's note: Copernicus Publications remains neutral with regard to jurisdictional claims made in the text, published maps, institutional affiliations, or any other geographical representation in this paper. The authors bear the ultimate responsibility for providing appropriate place names. Views expressed in the text are those of the authors and do not necessarily reflect the views of the publisher.

Acknowledgements. The authors would like to thank the participants of the field campaign “Korea Precipitation Observation Program: international collaborative experiments for Mesoscale convective system in Seoul metropolitan area” (KPOP-MS), hosted by the Korea Meteorological Administration (KMA).

Financial support. This work was funded by the Korea Meteorological Administration Research and Development Program “Observing Severe Weather in Seoul Metropolitan Area and Developing Its Application Technology for Forecasts” (grant no. KMA2018-00125).

Review statement. This paper was edited by Simon Unterstrasser and reviewed by two anonymous referees.

References

- Bang, W., Lee, G., Ryzhkov, A., Schuur, T., and Lim, K.-S. S.: Comparison of microphysical characteristics between the Southern Korean Peninsula and Oklahoma using two-dimensional video disdrometer data, *J. Hydrometeorol.*, 21, 2675–2690, <https://doi.org/10.1175/JHM-D-20-0087.1>, 2020.
- Berne, A., Jaffrain, J., and Schleiss, M.: Scaling analysis of the variability of the rain drop size distribution at small scale, *Adv. Water Resour.*, 45, 2–12, <https://doi.org/10.1016/j.advwatres.2011.12.016>, 2012.
- Byun, U.-Y., Hong, S.-Y., Shin, H., Lee, J.-W., Song, J.-I., Hahm, S.-J., Kim, J.-K., Kim, H.-W., and Kim, J.-S.: WRF-based short-range forecast system of the Korea Air Force: Verification of prediction skill in 2009 summer, *Atmosphere*, 21, 197–208, <https://doi.org/10.14191/Atmos.2011.21.2.197>, 2011.
- Cha, J. W., Koo, H. J., Kim, B.-Y., Miloslav, B., Hwang, H. J., Kim, M. H., Chang, K.-H., and Lee, Y. H.: Analysis of rain drop size distribution to elucidate the precipitation process using a cloud microphysics conceptual model and in situ measurement, *Asia-Pacific J. Atmos. Sci.*, 59, 257–269, <https://doi.org/10.1007/s13143-022-00299-w>, 2023.
- Chakraborty, T., Pattnaik, S., Jenamani, R., and Baisya, H.: Evaluating the performances of cloud microphysical parameterizations in WRF for the heavy rainfall event of Kerala (2018), *Meteorol. Atmos. Phys.*, 133, 707–737, <https://doi.org/10.1007/s00703-021-00776-3>, 2021.
- Chen, F. and Dudhia, J.: Coupling an advanced land surface–hydrology model with the Penn State–NCAR MM5 modeling system. Part I: Model implementation and sensitivity, *Mon. Weather Rev.*, 129, 569–585, [https://doi.org/10.1175/1520-0493\(2001\)129<0569:CAALSH>2.0.CO;2](https://doi.org/10.1175/1520-0493(2001)129<0569:CAALSH>2.0.CO;2), 2001.
- Cohard, J. M. and Pinty, J. P.: A comprehensive two-moment warm microphysical bulk scheme. I: Description and tests, *Q. J. Roy. Meteor. Soc.*, 126, 1815–1842, <https://doi.org/10.1002/qj.49712656613>, 2000.
- Dolan, B., Fuchs, B., Rutledge, S., Barnes, E., and Thompson, E.: Primary modes of global drop size distributions, *J. Atmos. Sci.*, 75, 1453–1476, <https://doi.org/10.1175/JAS-D-17-0242.1>, 2018.
- Gao, W., Sui, C. H., Chen Wang, T. C., and Chang, W. Y.: An evaluation and improvement of microphysical parameterization from a two-moment cloud microphysics scheme and the Southwest Monsoon Experiment (SoWMEX)/Terrain-influenced Monsoon Rainfall Experiment (TiMREX) observations, *J. Geophys. Res.-Atmos.*, 116, <https://doi.org/10.1029/2011JD015718>, 2011.
- Hagos, S., Leung, L. R., Yang, Q., Zhao, C., and Lu, J.: Resolution and dynamical core dependence of atmospheric river frequency in global model simulations, *J. Climate*, 28, 2764–2776, <https://doi.org/10.1175/JCLI-D-14-00567.1>, 2015.
- Hersbach, H., Bell, B., Berrisford, P., Hirahara, S., Horányi, A., Muñoz-Sabater, J., Nicolas, J., Peubey, C., Radu, R., and Schepers, D.: The ERA5 global reanalysis, *Q. J. Roy. Meteor. Soc.*, 146, 1999–2049, <https://doi.org/10.1002/qj.3803>, 2020.
- Hersbach, H., Bell, B., Berrisford, P., Biavati, G., Horányi, A., Muñoz Sabater, J., Nicolas, J., Peubey, C., Radu, R., Rozum, I., Schepers, D., Simmons, A., Soci, C., Dee, D., and Thépaut, J.-N.: ERA5 hourly data on single levels from 1940 to present, Copernicus Climate Change Service (C3S) Climate Data Store (CDS) [data set], <https://doi.org/10.24381/cds.adbb2d47>, 2023a.

- Hersbach, H., Bell, B., Berrisford, P., Biavati, G., Horányi, A., Muñoz Sabater, J., Nicolas, J., Peubey, C., Radu, R., Rozum, I., Schepers, D., Simmons, A., Soci, C., Dee, D., and Thépaut, J.-N.: ERA5 hourly data on pressure levels from 1940 to present, Copernicus Climate Change Service (C3S) Climate Data Store (CDS) [data set], <https://doi.org/10.24381/cds.bd0915c6>, 2023b.
- Hong, S.-Y., Noh, Y., and Dudhia, J.: A new vertical diffusion package with an explicit treatment of entrainment processes, *Mon. Weather Rev.*, 134, 2318–2341, <https://doi.org/10.1175/MWR3199.1>, 2006.
- Huang, B., Liu, C., Banzon, V., Freeman, E., Graham, G., Hankins, B., Smith, T., and Zhang, H.-M.: Improvements of the daily optimum interpolation sea surface temperature (DOISST) version 2.1, *J. Climate*, 34, 2923–2939, <https://doi.org/10.1175/jcli-d-20-0166.1>, 2021.
- Huffman, G. J., Bolvin, D. T., Braithwaite, D., Hsu, K., Joyce, R., Kidd, C., Nelkin, E. J., and Xie, P.: NASA global precipitation measurement (GPM) integrated multi-satellite retrievals for GPM (IMERG), Algorithm Theoretical Basis Document (ATBD), version 4.5, National Aeronautics and Space Administration (NASA), 30 pp., https://gpm.nasa.gov/sites/default/files/document_files/IMERG_ATBD_V4.5.pdf (last access: 24 August 2026), 2015.
- Iacono, M. J., Delamere, J. S., Mlawer, E. J., Shephard, M. W., Clough, S. A., and Collins, W. D.: Radiative forcing by long-lived greenhouse gases: Calculations with the AER radiative transfer models, *J. Geophys. Res.-Atmos.*, 113, <https://doi.org/10.1029/2008JD009944>, 2008.
- Jiménez, P. A., Dudhia, J., González-Rouco, J. F., Navarro, J., Montávez, J. P., and García-Bustamante, E.: A revised scheme for the WRF surface layer formulation, *Mon. Weather Rev.*, 140, 898–918, <https://doi.org/10.1175/MWR-D-11-00056.1>, 2012.
- Jo, J.: Code for publication – Implementation of Generalized Double-Moment Scaling Normalization in a Cloud Microphysics Scheme: A Case Study in the Korean Peninsula, Zenodo [code], <https://doi.org/10.5281/zenodo.17194841>, 2025.
- Jo, J.: Additional data for publication – Implementation of Generalized Double-Moment Scaling Normalization in a Cloud Microphysics Scheme: A Case Study in the Korean Peninsula, Zenodo [data set], <https://doi.org/10.5281/zenodo.18328177>, 2026a.
- Jo, J.: Additional scripts and output data for publication – Implementation of Generalized Double-Moment Scaling Normalization in a Cloud Microphysics Scheme: A Case Study in the Korean Peninsula, Zenodo [code and data set], <https://doi.org/10.5281/zenodo.18346988>, 2026b.
- Kain, J. S.: The Kain–Fritsch convective parameterization: an update, *J. Appl. Meteorol.*, 43, 170–181, [https://doi.org/10.1175/1520-0450\(2004\)043<0170:TKCPAU>2.0.CO;2](https://doi.org/10.1175/1520-0450(2004)043<0170:TKCPAU>2.0.CO;2), 2004.
- Kain, J. S. and Fritsch, J. M.: A one-dimensional entraining/detraining plume model and its application in convective parameterization, *J. Atmos. Sci.*, 47, 2784–2802, [https://doi.org/10.1175/1520-0469\(1990\)047<2784:AODEPM>2.0.CO;2](https://doi.org/10.1175/1520-0469(1990)047<2784:AODEPM>2.0.CO;2), 1990.
- Koch, S. E., Ferrier, B., Stoelinga, M. T., Szoke, E., Weiss, S. J., and Kain, J. S.: The use of simulated radar reflectivity fields in the diagnosis of mesoscale phenomena from high-resolution WRF model forecasts, Preprints, 11th Conf. on Mesoscale Processes, 24–28 October 2005, Albuquerque, New Mexico, USA, <https://ams.confex.com/ams/32Rad11Meso/webprogram/Paper97032.html> (last access: 2 September 2026), Amer. Meteor. Soc., J4J, 1–9, 2005.
- Kwon, S., Jung, S.-H., and Lee, G.: A case study on microphysical characteristics of mesoscale convective system using generalized DSD parameters retrieved from dual-polarimetric radar observations, *Remote Sens.*, 12, 1812, <https://doi.org/10.3390/rs12111812>, 2020.
- Lee, G., Brangi, V., and Thurai, M.: The retrieval of drop size distribution parameters using a dual-polarimetric radar, *Remote Sens.*, 15, 1063, <https://doi.org/10.3390/rs15041063>, 2023.
- Lee, G. W., Zawadzki, I., Szyrmer, W., Sempere-Torres, D., and Uijlenhoet, R.: A general approach to double-moment normalization of drop size distributions, *J. Appl. Meteorol.*, 43, 264–281, [https://doi.org/10.1175/1520-0450\(2004\)043<0264:AGATDN>2.0.CO;2](https://doi.org/10.1175/1520-0450(2004)043<0264:AGATDN>2.0.CO;2), 2004.
- Lim, K.-S. S. and Hong, S.-Y.: Development of an effective double-moment cloud microphysics scheme with prognostic cloud condensation nuclei (CCN) for weather and climate models, *Mon. Weather Rev.*, 138, 1587–1612, <https://doi.org/10.1175/2009MWR2968.1>, 2010.
- Lim, K.-S. S. and Hong, S.-Y.: Investigation of aerosol indirect effects on simulated flash-flood heavy rainfall over Korea, *Meteorol. Atmos. Phys.*, 118, 199–214, <https://doi.org/10.1007/s00703-012-0216-6>, 2012.
- Lim, K. S. S., Chang, E.-C., Sun, R., Kim, K., Tapiador, F. J., and Lee, G.: Evaluation of simulated winter precipitation using WRF-ARW during the ICE-POP 2018 field campaign, *Weather Forecast.*, 35, 2199–2213, <https://doi.org/10.1175/WAF-D-19-0236.1>, 2020.
- Lin, Y.-L., Farley, R. D., and Orville, H. D.: Bulk Parameterization of the Snow Field in a Cloud Model, *J. Appl. Meteorol. Climatol.*, 22, 1065–1092, [https://doi.org/10.1175/1520-0450\(1983\)022<1065:BPOTSF>2.0.CO;2](https://doi.org/10.1175/1520-0450(1983)022<1065:BPOTSF>2.0.CO;2), 1983.
- Meyers, M. P., Walko, R. L., Harrington, J. Y., and Cotton, W. R.: New RAMS cloud microphysics parameterization. Part II: The two-moment scheme, *Atmos. Res.*, 45, 3–39, [https://doi.org/10.1016/S0169-8095\(97\)00018-5](https://doi.org/10.1016/S0169-8095(97)00018-5), 1997.
- Milbrandt, J. A. and Yau, M.: A multimoment bulk microphysics parameterization. Part I: Analysis of the role of the spectral shape parameter, *J. Atmos. Sci.*, 62, 3051–3064, 2005.
- Min, K.-H., Choo, S., Lee, D., and Lee, G.: Evaluation of WRF cloud microphysics schemes using radar observations, *Weather Forecast.*, 30, 1571–1589, <https://doi.org/10.1175/WAF-D-14-00095.1>, 2015.
- Morcrette, J., Barker, H. W., Cole, J., Iacono, M. J., and Pincus, R.: Impact of a new radiation package, McRad, in the ECMWF Integrated Forecasting System, *Mon. Weather Rev.*, 136, 4773–4798, <https://doi.org/10.1175/2008MWR2363.1>, 2008.
- Morrison, H.: On the robustness of aerosol effects on an idealized supercell storm simulated with a cloud system-resolving model, *Atmos. Chem. Phys.*, 12, 7689–7705, <https://doi.org/10.5194/acp-12-7689-2012>, 2012.
- Morrison, H., Curry, J., and Khvorostyanov, V.: A new double-moment microphysics parameterization for application in cloud and climate models. Part I: Description, *J. Atmos. Sci.*, 62, 1665–1677, <https://doi.org/10.1175/JAS3446.1>, 2005.
- Morrison, H., Milbrandt, J. A., Bryan, G. H., Ikeda, K., Tessendorf, S. A., and Thompson, G.: Parameterization of

- cloud microphysics based on the prediction of bulk ice particle properties. Part II: Case study comparisons with observations and other schemes, *J. Atmos. Sci.*, 72, 312–339, <https://doi.org/10.1175/JAS-D-14-0066.1>, 2015.
- Morrison, H., Kumjian, M. R., Martinkus, C. P., Prat, O. P., and van Lier-Walqui, M.: A general N-moment normalization method for deriving raindrop size distribution scaling relationships, *J. Appl. Meteorol. Climatol.*, 58, 247–267, <https://doi.org/10.1175/JAMC-D-18-0060.1>, 2019.
- Park, S.-Y., Lim, K.-S. S., and Jo, J.: Assessing the relative importance of the prognostic hail mixing ratio and predicted graupel density on the vertical reflectivity structure in bulk cloud microphysics schemes, *Atmos. Res.*, 328, 108440, <https://doi.org/10.1016/j.atmosres.2025.108440>, 2025.
- Raupach, T. H., Thurai, M., Bringi, V., and Berne, A.: Reconstructing the drizzle mode of the raindrop size distribution using double-moment normalization, *J. Appl. Meteorol. Climatol.*, 58, 145–164, <https://doi.org/10.1175/JAMC-D-18-0156.1>, 2019.
- Reisner, J., Rasmussen, R., and Bruintjes, R.: Explicit forecasting of supercooled liquid water in winter storms using the MM5 mesoscale model, *Q. J. Roy. Meteor. Soc.*, 124, 1071–1107, <https://doi.org/10.1002/qj.49712454804>, 1998.
- Rutledge, S. A. and Hobbs, P.: The mesoscale and microscale structure and organization of clouds and precipitation in midlatitude cyclones. VIII: A model for the “seeder-feeder” process in warm-frontal rainbands, *J. Atmos. Sci.*, 40, 1185–1206, [https://doi.org/10.1175/1520-0469\(1983\)040<1185:TMAMSA>2.0.CO;2](https://doi.org/10.1175/1520-0469(1983)040<1185:TMAMSA>2.0.CO;2), 1983.
- Schoenberg Ferrier, B.: A double-moment multiple-phase four-class bulk ice scheme. Part I: Description, *J. Atmos. Sci.*, 51, 249–280, [https://doi.org/10.1175/1520-0469\(1994\)051<0249:ADMMPF>2.0.CO;2](https://doi.org/10.1175/1520-0469(1994)051<0249:ADMMPF>2.0.CO;2), 1994.
- Sempere-Torres, D. S., Porrà, J. M., and Creutin, J.-D.: A general formulation for raindrop size distribution, *J. Appl. Meteorol. Climatol.*, 33, 1494–1502, [https://doi.org/10.1175/1520-0450\(1994\)033<1494:AGFFRS>2.0.CO;2](https://doi.org/10.1175/1520-0450(1994)033<1494:AGFFRS>2.0.CO;2), 1994.
- Shin, K., Kim, K., Song, J. J., and Lee, G.: Polarimetric Retrieval of Raindrop Size Distribution: Double-Moment Normalization Approach and Machine Learning Techniques, *Geophys. Res. Lett.*, 51, e2023GL106057, <https://doi.org/10.1029/2023GL106057>, 2024.
- Skamarock, W. C., Klemp, J. B., Dudhia, J., Gill, D. O., Liu, Z., Berner, J., Wang, W., Powers, J. G., Duda, M. G., and Barker, D. M.: A description of the advanced research WRF version 4, NCAR tech. note ncar/tn-556+ str, 145, https://www2.mmm.ucar.edu/wrf/users/docs/technote/v4_technote.pdf (last access: 24 August 2026), 2019.
- Song, H.-J. and Sohn, B.-J.: Two heavy rainfall types over the Korean peninsula in the humid East Asian summer environment: A satellite observation study, *Mon. Weather Rev.*, 143, 363–382, <https://doi.org/10.1175/MWR-D-14-00184.1>, 2015.
- Stoelinga, M. T.: Simulated equivalent reflectivity factor as currently formulated in RIP: Description and possible improvements, University of Washington Tech. Rep., 5 pp., <http://citeseerx.ist.psu.edu/viewdoc/download?doi=10.1.1.522.925&rep=rep1&typepdf> (last access: 24 August 2026), 2005.
- Szyrmer, W., Laroche, S., and Zawadzki, I.: A microphysical bulk formulation based on scaling normalization of the particle size distribution. Part I: Description, *J. Atmos. Sci.*, 62, 4206–4221, 2005.
- Tapiador, F. J., Shin, K., Leganés, L. J., Lim, K. S., Juárez, G., Bang, W., Navarro, A., Martín, R., Park, S. Y., and Lee, G.: A physically consistent particle size distribution modeling of the microphysics of precipitation for weather and climate models, *Geophys. Res. Lett.*, 53, e2025GL118545, <https://doi.org/10.1029/2025GL118545>, 2026.
- Testud, J., Oury, S., Black, R. A., Amayenc, P., and Dou, X.: The concept of “normalized” distribution to describe raindrop spectra: A tool for cloud physics and cloud remote sensing, *J. Appl. Meteorol.*, 40, 1118–1140, [https://doi.org/10.1175/1520-0450\(2001\)040<1118:TCOND>2.0.CO;2](https://doi.org/10.1175/1520-0450(2001)040<1118:TCOND>2.0.CO;2), 2001.
- Thompson, G., Field, P. R., Rasmussen, R. M., and Hall, W. D.: Explicit forecasts of winter precipitation using an improved bulk microphysics scheme. Part II: Implementation of a new snow parameterization, *Mon. Weather Rev.*, 136, 5095–5115, <https://doi.org/10.1175/2008MWR2387.1>, 2008.
- Uijlenhoet, R., Steiner, M., and Smith, J. A.: Variability of raindrop size distributions in a squall line and implications for radar rainfall estimation, *J. Hydrometeorol.*, 4, 43–61, [https://doi.org/10.1175/1525-7541\(2003\)004<0043:VORSDI>2.0.CO;2](https://doi.org/10.1175/1525-7541(2003)004<0043:VORSDI>2.0.CO;2), 2003.
- Yang, Q., Dai, Q., Han, D., Chen, Y., and Zhang, S.: Sensitivity analysis of raindrop size distribution parameterizations in WRF rainfall simulation, *Atmos. Res.*, 228, 1–13, <https://doi.org/10.1016/j.atmosres.2019.05.019>, 2019.