

## Supplementary materials

Régis Briant<sup>\*,a</sup>, Christian Seigneur<sup>a</sup>, Marie Gadrat<sup>b</sup>, Christine Bugajny<sup>b</sup>

<sup>a</sup>*CEREA, Joint Research Laboratory École des Ponts ParisTech / EDF R&D, Université Paris-Est, 6–8 avenue Blaise Pascal, Cité Descartes, Champs-sur-Marne, 77455 Marne-la-Vallée Cedex 2, France*

<sup>b</sup>*CETE Nord-Picardie, Equipment Technical Study Center for the Nord-Picardie region in France, 42 bis rue Marais Sequedin - BP 10099, 59482 Haubourdin Cedex, France*

---

---

### 1. General results

Table 1: Performance indicators of Polyphemus, the HV formulation with the "urban" option with Cergy-Pontoise background concentrations.

Performance indicator	Summer campaign		Winter campaign	
	HV	Polyphemus	HV	Polyphemus
Measured monthly mean value of NO <sub>2</sub> (µg m <sup>-3</sup> )	26.0		40.5	
Modeled monthly mean value of NO <sub>2</sub> (µg m <sup>-3</sup> )	17.2	17.3	26.1	26.2
Correlation	0.75	0.75	0.79	0.79
RMSE (µg m <sup>-3</sup> )	14.1	14.0	18.0	17.9
MNE	0.32	0.32	0.33	0.33
MNB	-0.22	-0.22	-0.31	-0.31
NME	0.37	0.37	0.37	0.36
NMB	-0.34	-0.34	-0.36	-0.36
MFE	0.38	0.38	0.41	0.40
MFB	-0.30	-0.30	-0.39	-0.39

---

\*Corresponding author. E-mail: briancr@cerea.enpc.fr

Table 2: Performance indicators of Polyphemus, the HV formulation with the "rural" option with Mantes-la-Jolie background concentrations.

Performance indicator	Summer campaign		Winter campaign	
	HV	Polyphemus	HV	Polyphemus
Measured monthly mean value of $\text{NO}_2$ ( $\mu\text{g m}^{-3}$ )	26.0		40.5	
Modeled monthly mean value of $\text{NO}_2$ ( $\mu\text{g m}^{-3}$ )	22.5	22.6	25.6	25.7
Correlation	0.74	0.74	0.78	0.79
RMSE ( $\mu\text{g m}^{-3}$ )	11.2	11.1	17.8	17.7
MNE	0.31	0.30	0.34	0.34
MNB	0.02	0.02	-0.33	-0.33
NME	0.30	0.29	0.37	0.37
NMB	-0.13	-0.13	-0.37	-0.37
MFE	0.30	0.30	0.42	0.42
MFB	-0.05	-0.05	-0.42	-0.41

Table 3: Performance indicators of Polyphemus, the HV formulation with the "urban" option with Mantes-la-Jolie background concentrations.

Performance indicator	Summer campaign		Winter campaign	
	HV	Polyphemus	HV	Polyphemus
Measured monthly mean value of $\text{NO}_2$ ( $\mu\text{g m}^{-3}$ )	26.0		40.5	
Modeled monthly mean value of $\text{NO}_2$ ( $\mu\text{g m}^{-3}$ )	16.0	16.0	22.0	22.0
Correlation	0.75	0.75	0.79	0.79
RMSE ( $\mu\text{g m}^{-3}$ )	14.9	14.8	21.3	21.2
MNE	0.34	0.34	0.42	0.42
MNB	-0.28	-0.28	-0.42	-0.42
NME	0.40	0.40	0.46	0.46
NMB	-0.38	-0.38	-0.46	-0.46
MFE	0.43	0.43	0.56	0.55
MFB	-0.38	-0.38	-0.55	-0.55

## 2. Comparison to the HV formulation




Figure 1: Comparison between the HV and Polyphemus models of simulated  $\text{NO}_2$  hourly concentrations ( $\mu\text{g m}^{-3}$ ). (a) : Map of the passive diffusion tube locations with respect to the roads (coordinates are in meter). (b) : situations when the wind is parallel to the road ( $\pm 10^\circ$ ). (c) : situations when the wind is not parallel to the road (summer campaign). The road direction is  $111^\circ$  ( $0^\circ$  represents a wind coming from the north and  $90^\circ$  a wind coming from the east).




Figure 2: Comparison between the HV and Polyphemus models of simulated  $\text{NO}_2$  hourly concentrations ( $\mu\text{g m}^{-3}$ ). (a) : Map of the passive diffusion tube locations with respect to the roads (coordinates are in meter). (b) : situations when the wind is parallel to the road ( $\pm 10^\circ$ ). (c) : situations when the wind is not parallel to the road (summer campaign). The road direction is  $16^\circ$  ( $0^\circ$  represents a wind coming from the north and  $90^\circ$  a wind coming from the east).

### 3. Sensitivity to input data

Table 4: Performance indicators of Polyphemus using the "rural" option. In the Monin-Obukhov length column stability classes are based on Monin-Obukhov length, in the GENEMIS column the GENEMIS temporal profile was used and in the NO<sub>2</sub> fraction column a 15% NO<sub>2</sub> fraction was used.

Summer campaign	Monin-Obukhov length	GENEMIS	NO <sub>2</sub> fraction
Measured monthly mean value of NO <sub>2</sub> ( $\mu\text{g m}^{-3}$ )		26.0	
Modeled monthly mean value of NO <sub>2</sub> ( $\mu\text{g m}^{-3}$ )	26.5	21.03	24.0
Correlation	0.74	0.74	0.74
RMSE ( $\mu\text{g m}^{-3}$ )	10.28	11.67	10.67
MNE	0.35	0.29	0.33
MNB	0.19	-0.05	0.09
NME	0.29	0.30	0.29
NMB	0.02	-0.19	-0.07
MFE	0.31	0.31	0.30
MFB	0.10	-0.12	0.02
Winter campaign	Monin-Obukhov length	GENEMIS	NO <sub>2</sub> fraction
Measured mean value ( $\mu\text{g m}^{-3}$ )		40.5	
Modeled mean value ( $\mu\text{g m}^{-3}$ )	31.5	28.4	29.9
Correlation	0.76	0.79	0.79
RMSE ( $\mu\text{g m}^{-3}$ )	13.3	15.79	14.4
MNE	0.22	0.28	0.25
MNB	-0.18	-0.25	-0.22
NME	0.25	0.31	0.28
NMB	-0.22	-0.30	-0.26
MFE	0.25	0.33	0.29
MFB	-0.21	-0.31	-0.26