

Supplement of Geosci. Model Dev., 8, 3179–3198, 2015  
<http://www.geosci-model-dev.net/8/3179/2015/>  
doi:10.5194/gmd-8-3179-2015-supplement  
© Author(s) 2015. CC Attribution 3.0 License.



*Supplement of*

## **Simulation of atmospheric N<sub>2</sub>O with GEOS-Chem and its adjoint: evaluation of observational constraints**

**K. C. Wells et al.**

*Correspondence to:* D. B. Millet (dbm@umn.edu)

The copyright of individual parts of the supplement might differ from the CC-BY 3.0 licence.

## Supplemental Information




Figure S1. Verification of the N<sub>2</sub>O adjoint code. Shown are finite difference sensitivities (FD, upper panels), adjoint sensitivities (middle panels), and comparison between the two for N<sub>2</sub>O terrestrial emissions (left panels), ocean emissions (middle panels), and stratospheric loss frequencies (right panels) for a five-day simulation. Sensitivities for emissions were calculated at the surface level in the model; sensitivities for stratospheric loss were calculated at model level 40 (~25 km altitude).




Figure S2. Resolving seasonal emission biases. Shown are zonally-integrated monthly emissions for May 2010, July 2010, September 2010, November 2010, January 2011, and March 2011 with a seasonally-dependent a priori emission bias applied. Actual model emissions are shown in black, model emissions scaled by the a priori guess are shown in red, a posteriori emissions obtained using surface data, CARIBIC data, and HIPPO data are shown in green, blue, and yellow, respectively.