

Geoscientific Model Development Discussions is the access reviewed
discussion forum of *Geoscientific Model Development*

**MOZART-4
description**

L. K. Emmons et al.

Description and evaluation of the Model for Ozone and Related chemical Tracers, version 4 (MOZART-4)

L. K. Emmons¹, S. Walters¹, P. G. Hess^{1,*}, J.-F. Lamarque¹, G. G. Pfister¹,
D. Fillmore^{1,**}, C. Granier^{2,3}, A. Guenther¹, D. Kinnison¹, T. Laepple^{1,***},
J. Orlando¹, X. Tie¹, G. Tyndall¹, C. Wiedinmyer¹, S. L. Baughcum⁴, and
S. Kloster^{5,*}

¹National Center for Atmospheric Research, Boulder, CO, USA

²Centre National de la Recherche Scientifique, Paris, France

³NOAA, Earth System Research Laboratory, Boulder, CO, USA

⁴Boeing Company, Seattle, WA, USA

⁵Max-Planck-Institute for Meteorology, Hamburg, Germany

* now at: Cornell University, Ithaca, NY, USA

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



** now at: Tech-X Corporation, Boulder, CO, USA

*** now at: Alfred Wegener Inst. for Polar and Marine Research Bremerhaven, Germany

Received: 6 August 2009 – Accepted: 10 August 2009 – Published: 27 August 2009

Correspondence to: L. K. Emmons (emmons@ucar.edu)

Published by Copernicus Publications on behalf of the European Geosciences Union.

GMDD

2, 1157–1213, 2009

**MOZART-4
description**

L. K. Emmons et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Abstract

The Model for Ozone and Related chemical Tracers, version 4 (MOZART-4) is an offline global chemical transport model particularly suited for studies of the troposphere. The updates of the model from its previous version MOZART-2 are described, including an expansion of the chemical mechanism to include more detailed hydrocarbon chemistry and bulk aerosols. Online calculations of a number of processes, such as dry deposition, emissions of isoprene and monoterpenes and photolysis frequencies, are now included. Results from an eight-year simulation (2000–2007) are presented and evaluated. The MOZART-4 source code and standard input files are available for download from the NCAR Community Data Portal (<http://cdp.ucar.edu>).

1 Introduction

MOZART-4 (Model for Ozone and Related chemical Tracers, version 4) is a global chemical transport model for the troposphere and includes a number of updates over the previous tropospheric version MOZART-2 (Horowitz et al., 2003). MOZART-3 is an extension of MOZART-2 with detailed stratospheric chemistry (Kinnison et al., 2007). The MOZART models are offline models, requiring meteorological fields from either climate models or assimilations of meteorological observations. MOZART is built on the framework of the Model of Atmospheric Transport and Chemistry (MATCH) (Rasch et al., 1997). Convective mass fluxes are diagnosed by the model, using the shallow and mid-level convective transport formulation of Hack (1994) and deep convection scheme of Zhang and MacFarlane (1995). Vertical diffusion within the boundary layer is based on the parameterization of Holstlag and Boville (1993). Wet deposition is taken from the formulation of Brasseur et al. (1998). Advective transport incorporates the flux form semi-Lagrangian transport algorithm of Lin and Rood (1996), as described in Horowitz et al. (2003). These physical processes have not been updated from MOZART-2. Details of the chemical solver numerics are given in the Auxiliary Material of Kinnison et al. (2007).

GMDD

2, 1157–1213, 2009

MOZART-4 description

L. K. Emmons et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



MOZART-4 can be driven by essentially any meteorological data set and with any emissions inventory, so there is not a unique standard simulation. The meteorological variables needed to run MOZART-4 are given in Table 1. Whereas the published evaluation of MOZART-2 was for a simulation driven by climate model meteorology (Horowitz et al., 2003), this work presents MOZART-4 driven by meteorological analyses. A MOZART-4 simulation of 2000–2007 is used here to illustrate components of the model and for evaluation. The model was driven by meteorology from the NCAR re-analysis of the National Centers for Environmental Prediction (NCEP) forecasts (Kalnay et al., 1996; Kistler et al., 2001), at a horizontal resolution of approximately $2.8^\circ \times 2.8^\circ$, with 28 vertical levels from the surface to approximately 2 hPa. This resolution is typically used for multi-year simulations. MOZART-4 can be run at essentially any resolution, depending on memory limitations, and typically matching the input meteorological fields. Currently the highest resolution that has been run with full chemistry is $0.7^\circ \times 0.7^\circ$ (Emmons et al., 2009b).

MOZART-4 has been tested on computing platforms ranging from a single CPU on a desktop machine to more than one hundred CPUs on a state-of-the-art supercomputer. The following are required for running MOZART-4: the Linux or Unix operating system (including Apple OS X), the cpp utility (available on any Unix or Linux system), the gnu makefile utility, the netcdf utility, and a Fortran 90 compiler. For desktop systems, MOZART-4 has been compiled and is configured to use either the Portland Group, Lahey/Fujitsu, or Intel Fortran 90 or 95 compilers. All compilers produce similar execution times with normal optimizations. MOZART-4, configured at the standard horizontal resolution of 128×64 grid boxes with 28 vertical levels, requires at least 2 GB main memory per cpu, and can be run on single-, dual-, quad- or 8-CPU workstations. The source code is available for download from the NCAR Community Data Portal (<http://cdp.ucar.edu>), after registration and approval from the authors. Instructions on the building and running of MOZART-4 are included with the source code. Further information is posted on the MOZART-4 website (<http://gctm.acd.ucar.edu/mozart/models/m4/index.shtml>).

MOZART-4 description

L. K. Emmons et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



This paper presents a description of MOZART-4, providing details of the features of the model, and serving as a reference for past and future studies using this model. The following section describes the model updates included in MOZART-4 over MOZART-2. Section 3 summarizes previous applications of MOZART-4. The final sections give a description of the 2000–2007 model simulation presented here, and its evaluation through comparison to global networks and satellite observations.

2 Updates to model processes

Details of the improvements to MOZART-4 over MOZART-2 are described below. The chemical mechanism has been significantly expanded to include an improved representation of non-methane hydrocarbons and the online calculation of aerosols. An online photolysis scheme takes into account the impact of aerosols and clouds on the photolysis rates. Dry deposition velocities can be determined online in the model and an improved scheme for the determination of albedo has been implemented.

2.1 Chemical mechanism

The standard MOZART-4 mechanism includes 85 gas-phase species, 12 bulk aerosol compounds, 39 photolysis and 157 gas-phase reactions. The chemical mechanism includes an updated isoprene oxidation scheme and a better treatment of volatile organic compounds, with three lumped species to represent alkanes and alkenes with four or more carbon atoms and aromatic compounds (called BIGALK, BIGENE and TOLUENE). Oxidation products of these lumped species have also been added. MOZART-2 had a single lumped species (“C₄H₁₀”) representing all hydrocarbons C₄ and larger. Discussion of the isoprene oxidation scheme is given in Pfister et al. (2008a). Reaction rates are based on JPL 2002 (Sander et al., 2003). The full list of species and gas-phase, aerosol and photolytic reactions are given in Tables 2–6. Table 7 gives an approximate mapping of the longer-lived VOCs in MOZART-4 to three

MOZART-4 description

L. K. Emmons et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



other mechanisms commonly used in regional models, SAPRC-99, RADM, CBMZ (e.g., Tang et al., 2007; Grell et al., 2005; Fast et al., 2006). One application of this mapping is the matching of MOZART-4 results to regional model boundary conditions.

The MOZART models make use of a chemical pre-processor, making it relatively easy to modify the chemical mechanism. This facilitates updating reaction rates, adding additional species, or running with simplified mechanisms or a few artificial tracers. MOZART-4 also has the capability of reading offline constituents, such as OH, and prescribed chemical production and loss rates, allowing the simulation of species with relatively simple chemistry (e.g., CO or CH₄) without running the full chemistry.

2.2 Aerosols

The representation of tropospheric aerosols in MOZART-4 has been extended from the work of Tie et al. (2001, 2005), and includes the calculation of sulfate, black carbon, primary organic, secondary organic (SOA), ammonium nitrate, and sea salt (Lamarque et al., 2005b). Sulfate aerosols are determined from emissions of SO₂ and DMS (Barth et al., 2000). Black carbon and organic carbon aerosols are emitted in a combination of hydrophobic and hydrophilic forms (80% and 50% hydrophobic, respectively), and are converted from hydrophobic to hydrophilic with a time constant of 1.6 days. Secondary organic aerosols are linked to the gas-phase chemistry through the oxidation of atmospheric VOCs as in Chung and Seinfeld (2002). The ammonium nitrate distribution is determined from NH₃ emissions and the parameterization of gas/aerosol partitioning by Metzger et al. (2002), which is a set of approximations to the equilibrium constant calculation (Seinfeld, 1986) based on the level of sulfate present.

The uptake of N₂O₅, HO₂, NO₂, and NO₃ on aerosols is included (Jacob, 2000), with details given in Table 5. Because only the bulk mass is calculated, a lognormal number distribution is assumed for all aerosols to calculate the surface area, using a different geometric mean radius and standard deviation for each type of aerosol, as listed in Table 6 (based on Chin et al., 2002). Sea salt aerosols are included in the model with four size bins (0.1–0.5, 0.5–1.5, 1.5–5, 5–10 μm) and emissions are calculated

MOZART-4 description

L. K. Emmons et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



online (Mahowald et al., 2006a). The distributions of four sizes of dust (0.05–0.5, 0.5–1.25, 1.25–2.5, 2.5–5.0 μm) are set from monthly mean distributions taken from online calculations in the Community Atmosphere Model (CAM) (Mahowald et al., 2006b). Hygroscopic growth of the aerosols is determined from the ambient relative humidity, with different rates for each type of aerosol (Chin et al., 2002). Washout of all aerosols, except hydrophobic black carbon and organic carbon, is set to 20% of the washout rate of HNO_3 . This washout rate was selected as it gave the best agreement with observations of aerosol optical depth over ocean.

2.3 Photolysis

Photolysis rates are calculated within MOZART-4, taking into account the impact of aerosols and clouds, using FTUV (fast-TUV), based on the TUV (Tropospheric Ultraviolet-Visible) model (Tie et al., 2003). The treatment of aerosols in determining photolysis frequencies and aerosol optical depth are from a lookup table based on the Mie calculations used in the NCAR Community Atmosphere Model (CAM3): soot, organic carbon and sea salt are from the Optical Properties of Aerosols and Clouds (OPAC) software package (Hess et al., 1998); ammonium sulfate is based on Tang and Munkelwitz (1994), Toon et al. (1976), and the appendix of Gong et al. (2003); dust optics are from Zender et al. (2003). A lookup table, based on 4-stream calculations from TUV and also used in MOZART-3, can be used instead of FTUV (Kinnison et al., 2007). The lookup table includes explicit calculation of photolysis frequencies for most of the MOZART-4 species, whereas FTUV includes mapping to a subset of the species. This configuration includes the influence of clouds (Chang et al., 1987), but does not account for the impact of aerosols.

2.4 Albedo

An improved scheme for the determination of albedo has been implemented, based on satellite observations (Laepfle et al., 2005). Monthly snow and non-snow climatologies

MOZART-4 description

L. K. Emmons et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



have been derived from MODIS observations of albedo, and are combined with snow and ice cover information from the model-driving meteorology.

2.5 Online dry deposition

Dry deposition velocities can be determined online in the model, based on the resistance-based parameterization of Wesely (1989), Walmsley and Wesely (1996), Wesely and Hicks (2000). The calculation of surface resistances uses the vegetation distribution of Bonan et al. (2003). If the online calculation is not selected at run-time, a monthly mean climatology, based on the same parameterizations using 10 years of NCEP meteorology, is used. In both cases, the deposition velocity calculation has been extended to take into account special cases for CO, H₂ and PAN. In the case of CO and H₂, surface uptake caused by the oxidation by soil bacteria or enzymes (Yonemura et al., 2000) is parameterized using the approach of Sanderson et al. (2003); in this case, the deposition velocity is strongly dependent on the soil moisture. In the case of PAN, new laboratory experiments have indicated a strong uptake of PAN by leaves (Teklemariam and Sparks, 2004). Using the results from this study, we have included a leaf uptake of PAN that is vegetation-dependent; results from this parameterization agreed with observations during a field experiment (Turnipseed et al., 2006).

2.6 Online water calculation

In MOZART-2 water vapor concentrations were taken from the meteorological fields. However, better agreement with observations of precipitation were found when H₂O was calculated online from the surface moisture flux and all relevant physics parameters, as implemented in MOZART-3 (Kinnison et al., 2007). This online calculation is a runtime option, and the preferred mode of operation, in MOZART-4.

MOZART-4 description

L. K. Emmons et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[I◀](#)

[▶I](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



2.7 Emissions from vegetation based on MEGAN

Online calculation of biogenic emissions of isoprene and monoterpenes is based on the Model of Emissions of Gases and Aerosols from Nature (MEGAN) (Guenther et al., 2006) using the implementation described by Pfister et al. (2008a), and in more detail below. Since Guenther et al. (2006) gives a comprehensive description of MEGAN, with various options for particular applications, the details of its implementation in MOZART-4, and suggested for other global models, are included here. The emission factor maps used in MEGAN are updated periodically, as more measurements are made and the algorithms are refined (available from <http://bai.acd.ucar.edu/Megan>). For the results shown here, version 2.1 of the emission factor maps are used. In other MOZART-4 studies to date, version 2.0 of the isoprene emission factor maps, and version 1.0 of the monoterpenes emissions, have been used. The MOZART-4 lumped monoterpene (“C10H16”) emissions are the sum of the species alpha-pinene, beta-pinene, limonene, myrcene, ocimene, sabinene, and delta-3-carene in the MEGAN emission factor maps.

The MEGAN formulation also requires global maps of plant functional type (PFT) and the monthly leaf area index (LAI). The PFT and LAI maps used in MOZART-4 are based on AVHRR and MODIS data, as used in the NCAR Community Land Model (CLM) (Lawrence and Chase, 2007). The MEGAN emissions dependency on current and past surface air temperature and solar flux are considered using the model meteorology. Pfister et al. (2008a) illustrate the sensitivity of isoprene emissions to vegetation maps used. It is straight-forward to use other vegetation maps, such as for future climate scenarios, if desired. Using the specified driving variables, the global annual isoprene and monoterpene emissions for the years 2000–2007 are calculated in the range of 530–575 Tg/year and 73–76 Tg/year, respectively. Emissions of isoprene and monoterpenes as calculated in MOZART-4 for January and July 2006 are illustrated in Fig. 1.

GMDD

2, 1157–1213, 2009

MOZART-4 description

L. K. Emmons et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



2.7.1 General formulation

The MEGAN emissions are based on the emission factor EF (the emissions of a compound at standard conditions), the emission activity factor γ (a normalized ratio that accounts for deviations from standard conditions), and a factor ρ accounting for the canopy production and loss (assume=1 because this is representative of typical conditions). All equation numbers below are from Guenther et al. (2006).

Global, spatially varying emission factor (EF) maps for each compound (isoprene and monoterpenes for MOZART-4) are provided on a $0.5^\circ \times 0.5^\circ$ grid for five plant functional types (PFTs): broadleaf trees (btr), needleleaf trees (ntr), shrubs (shr), agricultural crops (crp), and grasses (grs). Landcover maps of PFT fractions per grid box, and leaf area index (LAI) for each month for each PFT are needed. These same maps are used for dry deposition calculations in MOZART-4. 17 PFTs are provided in these maps and must be combined to match the 5 MEGAN PFTs (see Table 8).

The emission activity factor depends on the canopy environment (CE), leaf age and soil moisture:

$$\gamma = \gamma_{CE} \cdot \gamma_{age} \cdot \gamma_{SM} \quad (2)$$

where the canopy environment factor depends on leaf area index (LAI), photosynthetic photon flux density (PPFD) and temperature:

$$\gamma_{CE} = \gamma_{LAI} \cdot \gamma_P \cdot \gamma_T \quad (10)$$

and dependence on soil moisture is ignored ($\gamma_{SM}=1$). By not considering the dependence on soil moisture, we could be overestimating global annual isoprene emissions by $\approx 7\%$ (Guenther et al., 2006).

Thus, for each compound, the emissions ϵ are determined by:

$$\epsilon = \sum_{PFT} EF(PFT) * f_{PFT} * \gamma_{LAI} * \gamma_{age} * \gamma_P * \gamma_T$$

where f_{PFT} is the fraction of each grid box covered by each PFT, and γ_{LAI} , γ_{age} , γ_P and γ_T will vary by compound, and may depend on PFT.

GMDD

2, 1157–1213, 2009

MOZART-4 description

L. K. Emmons et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



2.7.2 Isoprene

The leaf age factor (γ_{age}) for isoprene emissions from evergreen canopies equals 1. For deciduous canopies, γ_{age} is a weighted average of emissions from four ages of foliage (new, growing, mature and old):

$$\gamma_{\text{age}} = A_{\text{new}}F_{\text{new}} + A_{\text{gro}}F_{\text{gro}} + A_{\text{mat}}F_{\text{mat}} + A_{\text{old}}F_{\text{old}} \quad (16)$$

For isoprene, the relative weights of emissions from each canopy type are: $A_{\text{new}}=0.05$, $A_{\text{gro}}=0.6$, $A_{\text{mat}}=1.125$ and $A_{\text{old}}=1$. The leaf age fractions (F_x , for new, growing, mature and old states) are calculated according to Eqs. (17–19), and accompanying text, of Guenther et al. (2006). These factors depend on the change in LAI between the current (LAI_c) and previous (LAI_p) timestep, which for MOZART-4 is a month, based on the time resolution of the LAI maps used. The temperature dependencies of Eq. (18) are based on the average temperature of the previous month (cf., Guenther et al., 2006, and corrigendum).

Emission responses to LAI (γ_{LAI}) variations are estimated according to Eq. (15). The factor accounting for variation in solar radiation (γ_p) is a function of the solar angle, the previous month's average above-canopy PPFD, and the above-canopy PPFD transmission ϕ (given by Eqs. 12 and 13). At low sun angles, ϕ can become greater than 1, so should be limited to a maximum value of 1.

The temperature dependence for isoprene emissions is determined from:

$$\gamma_T = E_{\text{opt}} * C_{T2} * \exp(C_{T1} * x) / (C_{T2} - C_{T1} * (1 - \exp(C_{T2} * x))) \quad (14)$$

where $x = (T_p - T_{\text{opt}}) / (T_p * T_{\text{opt}} * 0.00831)$, $C_{T1} = 80$ and $C_{T2} = 200$, $T_{\text{opt}} = 313 + 0.6 * (T_p - T_{\text{clim}})$, $E_{\text{opt}} = 1.75 * \exp(0.08 * (T_p - T_{\text{clim}}))$, with T_p being the average temperature of the previous month and T_{clim} the daily average climatological value (297 K).

2.7.3 Monoterpenes

MOZART-4 includes a simplified implementation of monoterpene emissions that do not depend on sunlight ($\gamma_p=1$) or leaf age ($\gamma_{\text{age}}=1$), and simpler LAI and temperature

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



factors, with no dependence on past solar radiation:

$$\gamma_{LAI} = 0.2 * LAI$$

$$\gamma_T = \exp(0.09 * (T_p - T_{std}))$$

where T_p is the average temperature of the previous month and $T_{std}=303.15$.

5 2.8 Soil and lightning NO emissions

Soil NO emissions are a combination of interactive natural emissions (Yienger and Levy, 1995) and fertilizer use (Bouwman et al., 2002). The soil NO emissions are highly dependent on the degree of soil dryness; in order to keep track of this quantity, we have added to the model a bucket-style parameterization of soil moisture, taking into account the model-calculated precipitation and input latent heat flux. If the soil moisture is not available in the meteorological input dataset, then the model uses a 10-year climatology of monthly-averaged soil moisture from NCEP/NCAR-reanalysis meteorological fields. On average, for late 20th century conditions, the global annual average of soil NO emissions amount to approximately 7 Tg-N/year, including the effect of fertilization.

The lightning parameterization differs slightly from that used in MOZART-2 (Horowitz et al., 2003). The lightning strength still depends on cloud top height, with a stronger dependence over land than ocean (Price et al., 1997). The definition of ocean grid boxes has been refined to include only boxes surrounded by ocean, so that the land parameterization is extended one grid box beyond the continents (Price and Rind, 1992). Flash frequency is determined by area, not grid box. The vertical distribution of NO emissions has been modified from that given by Pickering et al. (1998), to have a reduced proportion of the emissions emitted near the surface. In addition, the strength of intra-cloud (IC) lightning strikes is assumed to be equal to cloud-to-ground strikes, as recommended by Ridley et al. (2005). The annual lightning emissions for 2006 from MOZART-4 driven by NCEP/GFS are shown in Fig. 2.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



2.9 Surface boundary conditions

For some long-lived species such as CH₄, H₂, and N₂O, their tropospheric concentrations are known more accurately than their emissions. Therefore, MOZART-4 simulations are generally run using fixed lower boundary conditions constrained by observations, instead of direct emissions for these species. CH₄ monthly zonal averages are based on the measurements by NOAA/ESRL Global Monitoring Division Cooperative Air Sampling Network (Dlugokencky et al., 2005, 2008). H₂ is set to 530 ppbv, based on global average observations made by NOAA/ESRL/GMD for 1993–2003 Novelli et al. (1999). N₂O concentrations are taken from IPCC (2000).

2.10 Upper boundary conditions

Mixing ratios of several species (O₃, NO_x, HNO₃, N₂O₅, CO, CH₄) are constrained in the stratosphere since MOZART-4 does not have complete stratospheric chemistry. These mixing ratios have been updated to zonal means from a MOZART-3 simulation. Model concentrations are set to the climatology values above 50 hPa, and relaxed to the climatology with a 10-day time scale down to the tropopause. The primary change from the climatology used in MOZART-2 is a reduction in the NO_x and HNO₃ mixing ratios in the lower stratosphere. O₃ is still constrained to observations (from satellite and ozonesondes), as in MOZART-2 (Horowitz et al., 2003). A stratospheric aerosol surface area density climatology (SPARC, 2006) has been included in MOZART-4 so as to more accurately represent the fast NO_y partitioning in the lower stratosphere.

Reanalysis meteorological datasets generally result in too strong a stratospheric flux in offline chemical transport models (Van Noije et al., 2004, and references therein), resulting in errors in the tropospheric ozone budget and too high ozone mixing ratios in the upper troposphere. This can be the case for MOZART-4 when driven by NCEP/NCAR reanalyses (Kalnay et al., 1996; Kistler et al., 2001) and ECMWF reanalyses (ERA-40) (Simmons and Gibson, 2000). In order to constrain the stratospheric flux of ozone independently of the meteorological dataset used, the synthetic ozone

GMDD

2, 1157–1213, 2009

MOZART-4 description

L. K. Emmons et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



(SYNOZ) representation is used (McLinden et al., 2000). SYNOZ is a tracer with a specified source region (30S–30N, 10–70 hPa) and production rate (400–500 Tg/yr); after production, SYNOZ is advected as a passive tracer. In order to provide a tropospheric sink, the SYNOZ mixing ratio is relaxed to 25 ppbv below 500 hPa. O₃ is set to SYNOZ above the tropopause, if SYNOZ > 100 ppbv; this limits the overwriting of the modeled ozone field by SYNOZ to the lower stratosphere. Because SYNOZ does not provide a realistic distribution of ozone in the stratosphere, an additional tracer, named O3RAD, is relaxed to the ozone climatology (mentioned above) in the stratosphere and to the model calculated ozone in the troposphere; this additional tracer is only used for photolysis calculations. An example of the SYNOZ, O3RAD and O3 variables is shown for a monthly mean profile in Fig. 3. When SYNOZ is used, the O3 variable is appropriate only for the troposphere, and should not be used for studies of the tropopause region. For simulations using the NCEP/GFS analyses it was found not to be necessary to use SYNOZ, probably due to the higher vertical resolution than the NCAR reanalysis (42 levels instead of 28) (Pfister et al., 2008b).

3 Previous MOZART-4 applications

MOZART-4 has already been used in several studies where it has been shown to reproduce observations well. Detailed comparisons have been made of MOZART-4 results to the aircraft, ozonesonde and ground-based observations from the International Consortium for Atmospheric Research on Transport and Transformation (ICARTT) during Summer 2004 (Pfister et al., 2005, 2006; Lapina et al., 2006; Horowitz et al., 2007; Pfister et al., 2008b). MOZART-4 simulations have been used for boundary conditions in regional models (Tang et al., 2008; Fast et al., 2009; Mena-Carrasco et al., 2009). Significant improvement in regional model results have been found when time-varying chemical boundary conditions, such as from MOZART-4, are used (e.g., Tang et al., 2008).

MOZART-4 description

L. K. Emmons et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



MOZART-4 results have also been included in multi-model comparisons, such as the study coordinated by the European Union project Atmospheric Composition Change: the European Network of excellence (ACCENT; <http://www.accent-network.org>) (Dentener et al., 2006; Stevenson et al., 2006; Shindell et al., 2006). Chemical forecasts have been produced with MOZART-4 and used in flight planning activities for aircraft experiments such as the NSF Megacities Impact on Regional and Global Environment (MIRAGE) Mexico City campaign in March 2006 (Fast et al., 2007), and the NASA Intercontinental Chemical Transport Experiment (INTEX-B) in April–May 2006 (Singh et al., 2009). Detailed comparisons of MOZART-4 results to the MIRAGE and INTEX-B experiments are given in Emmons et al. (2009b) and Pfister et al. (2009).

4 MOZART-4 simulation of 2000–2007

A simulation of 2000–2007 is presented here to illustrate the capabilities of MOZART-4 and for its evaluation. This simulation was driven by NCEP/NCAR-reanalysis meteorological fields, at a horizontal resolution of $2.8^{\circ} \times 2.8^{\circ}$ (28 levels). The artificial stratospheric ozone mechanism, SYNOZ, was used. Dry deposition velocities, water vapor concentrations, photolysis rates using FTUV, and biogenic emissions for soil NO, and for isoprene and monoterpenes from MEGAN, were calculated online using the new interactive schemes described above.

4.1 Emissions

A number of emissions inventories are suitable for use in MOZART-4, depending on the application. For this overview, several inventories have been used to cover all of the species and emissions types. These inventories, as used in MOZART-4, are provided with the MOZART-4 source code (<http://cdp.ucar.edu>).

The majority of the anthropogenic emissions used for this simulation came from the POET (Precursors of Ozone and their Effects in the Troposphere) database for 2000

MOZART-4 description

L. K. Emmons et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



(Granier et al., 2005; Olivier et al., 2003), which includes anthropogenic emissions (from fossil fuel and biofuel combustion) based on the EDGAR-3 inventory (Olivier and Berdowski, 2001). The anthropogenic emissions (from fossil fuel and biofuel combustion) of black and organic carbon determined for 1996 are from Bond et al. (2004). For SO₂ and NH₃, anthropogenic emissions are from the EDGAR-FT2000 and EDGAR-2 databases, respectively. For Asia, these inventories have been replaced by the Regional Emission inventory for Asia (REAS) that has annually varying amounts (Ohara et al., 2007). Aircraft emissions of NO, CO and SO₂ from scheduled, charter, general aviation and military traffic for 1999 are also included (Baughcum et al., 1996, 1998; Mortlock and Alstyne, 1998; Sutkus et al., 2001), and have global annual totals of 16.2 Tg/yr for NO, 20.4 Tg/yr for CO, and 2.0 Tg/yr for SO₂.

Monthly average biomass burning emissions for each year are from the Global Fire Emissions Database, version 2 (GFED-v2), which is currently available for 1997–2007 (van der Werf et al., 2006). For species not provided in GFED (e.g., individual volatile organic compounds as specified in MOZART-4, SO₂, and NH₃), emissions are determined by scaling the GFED CO₂ emissions by the emission factors of Andreae and Merlet (2001) and updates (Granier et al., 2005), using the vegetation classification provided with GFED. While MOZART-4 provides the capability to use vertically distributed emissions (such as for biomass burning), for these simulations, all emissions, except aircraft and lightning, are emitted at the surface.

Emissions of isoprene and monoterpenes from vegetation, and NO from soil and lightning, are calculated online, as described above. The DMS emissions are monthly means from the marine biogeochemistry model HAMOCC5, representative of the year 2000 (Kloster et al., 2006). SO₂ emissions from continuously outgassing volcanoes are from the GEIA-v1 inventory (Andres and Kasgnoc, 1998).

The surface emissions totals for 2006, separated by category, are shown in Table 9, and summarized for each year in Table 10. NO emissions from lightning are included in Table 10.

MOZART-4 description

L. K. Emmons et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



5 Model evaluation

To illustrate the capability of MOZART-4 to reproduce the true atmosphere, a few examples of the evaluation of the model through comparisons with a standard OH climatology, network ground sites, ozonesondes and satellite observations of CO and aerosol optical depth are shown.

5.1 Hydroxyl radical

Simulated OH is compared to the climatology of Spivakovsky et al. (2000) in Fig. 4 in terms of the annual airmass-weighted burden, as recommended by Lawrence et al. (2001). For this MOZART-4 simulation, OH is somewhat lower than the climatology, and gives a methane lifetime of about 10.5 years (2000–2007 global average, for the troposphere, altitudes below 100 hPa), which is on the high side of previous model estimates. For example, the MOZART-2 standard simulation had a methane lifetime of 9.4 years (Horowitz et al., 2003), and the MATCH simulations of Lawrence et al. (2001) gave lifetimes between 7.8 and 10.3 years. MOZART-4 configurations with different meteorology and emissions could result in different OH distributions, and consequently different CH₄ lifetimes.

5.2 Carbon monoxide

Comparison of carbon monoxide (CO) in MOZART-4 to the NOAA ESRL Global Monitoring Division surface sites (Novelli et al., 2003) is shown in Fig. 5. The model generally reproduces the seasonal cycle well, and captures much of the interannual variability. The model results are interpolated to the pressure altitude of the observation sites, so the mountain sites of Niwot Ridge (Colorado), Tenerife (Canary Islands), and Mauna Loa (Hawaii) are comparisons of free tropospheric air, generally not directly influenced by surface emissions.

MOZART-4 description

L. K. Emmons et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



MOZART-4
descriptionL. K. Emmons et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[I◀](#)[▶I](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

The mid to high southern latitudes are over-estimated by the model, perhaps due to an over-estimation of biogenic or biomass burning emissions (since anthropogenic emissions are relatively small in the Southern Hemisphere), or too low simulated OH concentrations (as indicated in Fig. 4). The large Australian fires at the start of the year in 2003 and 2007 show up strongly in the model in the Cape Grim comparison, due to the large model grid box that encompasses both Cape Grim and southern Australia.

The global distributions of CO from the Measurements of Pollution In The Troposphere instrument on the Terra satellite are a valuable data set for model evaluation (e.g., Shindell et al., 2006). The recently released MOPITT V4 retrievals (Deeter et al., 2009) are used for comparison to the MOZART-4 simulated CO. Figure 6 shows an example comparison of the total column CO for May 2003. The top panel shows the column average mixing ratio directly from the model output and the middle panel shows the MOPITT column retrieval, expressed as average mixing ratio. To properly compare these two products, the averaging kernel and a priori profile associated with each MOPITT retrieval must be applied to the model profiles, as described in Emmons et al. (2009a). The model appears to overestimate the CO produced by the large fires in Siberia during this month. Other regions of biomass burning, such as the Yucatan peninsula in Mexico, and central Africa, are also a bit high in MOZART-4. This may be due to too high fire emissions from the GFED-v2 inventory, or may be an artifact of having all the emissions released into the model at the surface, instead of vertically distributed, which would be more realistic in many cases. Outside of Siberia, MOZART-4 underestimates the Northern Hemisphere distribution (by 15–30 ppbv).

To evaluate the model in greater detail, comparisons for 2002–2007 are shown in Fig. 7 for various regions. The monthly mean retrievals and monthly mean model results (with MOPITT averaging kernels and a priori applied) have been averaged over various continental regions. The error bars on the MOPITT observations, and the ranges indicated for the model results, indicate the range of the data (10th to 90th percentiles) over the region. In general the agreement between MOZART-4 and MOPITT is quite good, as indicated by the correlation coefficients and mean biases (as indicated

in Fig. 7). In the Northern Hemisphere, the model appears to be a bit low, indicating perhaps the anthropogenic emissions are underestimated. In the regions with significant biomass burning impact (such as Southeast Asia, Indonesia, and S. America), the model has a positive bias of 7–16 ppbv, indicating the CO emissions from biomass burning may be over-estimated. The positive bias over Australia is consistent with that seen in the NOAA surface sites, particularly Cape Grim (Fig. 5).

5.3 Ozone

An evaluation of the distribution of ozone in MOZART-4 is made through comparison with ozonesondes, which are available from many sites over the past several decades. The observations used here are from the World Ozone and Ultraviolet Radiation Data Centre (Environment Canada, retrieved 25 March 2009 from <http://www.woudc.org>). The data originate from many sources, including the SHADOZ network (Thompson et al., 2003). The model-data comparisons are summarized in Fig. 8 for latitude and altitude bins. The mean of all observations, for each month of each year, within each latitude range (90S–30S, 30S–Equator, Equator–30N, 30N–90N) and within 100 hPa of each altitude used (400, 650, 900 hPa), are shown as individual points, colored by year. Monthly mean model results have been extracted for each ozonesonde site, and binned as the observations. The sites of the ozonesonde measurements used in each latitude bin are given in Table 11.

Generally good agreement is seen between MOZART-4 and the observations (Fig. 8), but some systematic differences are apparent. In the lower troposphere of the Southern Hemisphere, MOZART-4 generally underestimates the ozonesondes, whereas it is slightly high in the Northern Tropics. Significant interannual variability is seen in the observations in the Southern Tropics in the mid-troposphere at the end of the year, but MOZART-4 underestimates the concentrations, as well as the variation. MOZART-4 also does not fully reproduce the spring-to-summer increase in ozone in the Northern Hemisphere mid-troposphere, most likely due to the constraint on the stratospheric input through the use of SYNOZ, and the low vertical resolution of this

MOZART-4 description

L. K. Emmons et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



simulation with 28 levels.

5.4 Aerosol optical depth

The aerosol optical depth (AOD) retrievals from MODIS are compared to those calculated by MOZART-4 from the simulated aerosol distributions. Fig. 9 shows averages of MODIS AOD, over oceans only, for various regions, with corresponding averages from MOZART-4. The model agrees very well with the MODIS retrievals over the N. Pacific, which is strongly influenced by the export of pollution from Asia each spring. In May 2003 exceptionally large fires were burning in eastern Russia, resulting in very high AOD values (e.g., Edwards et al., 2004) that the model did not capture. Despite having biomass burning emissions specific for the year (see Sect. 4.1), this indicates the fire emissions inventory underestimated the black carbon or organic carbon emissions. In contrast, the model somewhat over-estimates the AOD over the N. Atlantic Ocean, both in the mean, and in the variability. The AOD from MOZART-4 systematically underestimates the observations in the Southern Hemisphere. The seasonal variation over the S. Pacific and S. Atlantic Oceans is driven by biomass burning in the second half of the year (in S. America, impacting the S. Atlantic, and Africa and Indonesia impacting the S. Pacific). These discrepancies could be due to washout rates that are too strong in the model, an underestimate of the biomass burning emissions of aerosols, or problems in the definition of the physical or radiative properties of aerosols.

6 Conclusions

The offline global chemical transport model for the troposphere MOZART-4 includes a number of updates over the previous version MOZART-2, the most significant being the expansion of the chemical mechanism, with the inclusion of aerosols, and the online calculation of photolysis rates, biogenic emissions and dry deposition. Evaluation with several sets of observations shows MOZART-4 has skill at reproducing tropospheric

MOZART-4 description

L. K. Emmons et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[I◀](#)

[▶I](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



chemical composition. When driven with time varying emission inventories (particularly for biomass burning), MOZART-4 reproduces the spatial and temporal variability in observations, such as the NOAA GMD network and MOPITT CO, ozonesondes and MODIS aerosol optical depth measurements. MOZART-4 is now available to the community and is suitable for many tropospheric investigations on the regional to global scale. The MOZART-4 source code and standard input files are available for download from the NCAR Community Data Portal (<http://cdp.ucar.edu>).

Acknowledgements. We acknowledge the many people who have participated in the development and evaluation of all of the versions of the MOZART model over the years, and in particular the leadership of Guy Brasseur. We thank Mary Barth and Valery Yudin for helpful comments on the manuscript. We also acknowledge the science teams producing the data used for model evaluation, including the NOAA Earth System Research Laboratory Global Monitoring Division for surface CO, the Canadian and NCAR MOPITT Teams for MOPITT retrievals, the World Ozone and Ultraviolet Radiation Data Centre (WOUDC) for ozonesondes, and NASA for MODIS AOD retrievals. The National Center for Atmospheric Research is sponsored by the National Science Foundation.

References

- Andreae, M. and Merlet, P.: Emission of trace gases and aerosols from biomass burning, *Global Biogeochem. Cy.*, 15, 966–995, 2001. 1172
- Andres, R. and Kasgnoc, A.: A time-averaged inventory of subaerial volcanic sulfur emissions, *J. Geophys. Res.*, 103, 25251–25261, 1998. 1172
- Barth, M. C., Rasch, P. J., Kiehl, J. T., Benkovitz, C. M., and Schwartz, S. E.: Sulfur chemistry in the National Center for Atmospheric Research Community Climate Model: Description, evaluation, features, and sensitivity to aqueous chemistry, *J. Geophys. Res.*, 105, 1387–1415, 2000. 1162
- Baughcum, S., Tritz, T., Henderson, S., and Pickett, D.: Scheduled Civil Aircraft Emission Inventories for 1992: Database Development and Analysis, NASA Contractor Report 4700, available at: http://ntrs.nasa.gov/archive/nasa/casi.ntrs.nasa.gov/19960038445_1996060751.pdf, 1996. 1172

MOZART-4 description

L. K. Emmons et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



MOZART-4
descriptionL. K. Emmons et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

- Baughcum, S., Henderson, S., and Sutkus, D.: Scheduled Civil Aircraft Emission Inventories Projected for 2015: Database Development and Analysis, NASA/CR-1998-207638, available at: http://ntrs.nasa.gov/archive/nasa/casi.ntrs.nasa.gov/19980055200_1998120133.pdf, 1998. 1172
- 5 Bonan, G., Levis, S., Sitch, S., Vertenstein, M., and Oleson, K.: A dynamic global vegetation model for use with climate models: concepts and description of simulated vegetation dynamics, *Glob. Change Biol.*, 9, 1543–1566, 2003. 1164
- Bond, T., Streets, D. G., Yarber, K. F., Nelson, S. M., Woo, J.-H., and Klimont, Z.: A technology-based global inventory of black and organic carbon emissions from combustion, *J. Geophys. Res.*, 109, D14203, doi:10.1029/2003JD003697, 2004. 1172
- 10 Bouwman, A. F., Boumans, L. J. M., and Batjes, N. H.: Modeling global annual N₂O and NO emissions from fertilized fields, *Global Biogeochem. Cy.*, 16, 1080, doi:10.1029/2001GB001812, 2002. 1168
- Brasseur, G. P., Hauglustaine, D. A., Walters, S., Rasch, P. J., Muller, J.-F., Granier, C., and Tie, X.-X.: MOZART: A global chemical transport model for ozone and related chemical tracers, Part 1: Model description, *J. Geophys. Res.*, 103, 28265–28289, 1998. 1159
- 15 Chang, J. S., Brost, R. A., Isaksen, I. S. A., Madronich, S., Middleton, P., Stockwell, W. R., and Walcek, C. J.: A three-dimensional Eulerian acid deposition model: Physical concepts and model formulation, *J. Geophys. Res.*, 92, 14681–14700, 1987. 1163
- 20 Chin, M., Ginoux, P., Kinne, S., et al.: Tropospheric aerosol optical thickness from the GOCART model and comparisons with satellite and sun photometer measurements, *J. Atmos. Sci.*, 59, 461–483, 2002. 1162, 1163
- Chung, S. and Seinfeld, J.: Global distribution and climate forcing of carbonaceous aerosols, *J. Geophys. Res.*, 107, 4407, doi:10.1029/2001JD001397, 2002. 1162
- 25 Deeter, M., Edwards, D. P., Gille, J. C., Emmons, L. K., Francis, G., Ho, S.-P., Mao, D., Masters, D., Worden, H., Yudin, V., and Drummond, J. R.: The MOPITT Version 4 CO product: Algorithm enhancements, selected results and bias drift, *J. Geophys. Res.*, in preparation, 2009. 1174
- Dentener, F., Stevenson, D., Ellingsen, K., et al.: Global Atmospheric Environment for the next generation, *Environ. Sci. Technol.*, 40, 3586–3594, 2006. 1171
- 30 Dlugokencky, E., Myers, R., Lang, P., Masarie, K., Crotwell, A., Thoning, K., Hall, B., Elkins, J., and Steele, L.: Conversion of NOAA atmospheric dry air methane mole fractions to a gravimetrically-prepared standard scale, *J. Geophys. Res.*, 110, D18306, doi:

10.1029/2005JD006035, 2005. 1169

Elugokencky, E., Lang, P., and Masarie, K.: Atmospheric Methane Dry Air Mole Fractions from the NOAA ESRL Carbon Cycle Cooperative Global Air Sampling Network, 1983–2007, Version: 2008-07-02, available at: <ftp://ftp.cmdl.noaa.gov/ccg/ch4/flask/>, 2008. 1169

5 Edwards, D., Emmons, L., Hauglustaine, D., et al.: Observations of carbon monoxide and aerosols from the Terra satellite: Northern Hemisphere variability, *J. Geophys. Res.*, 109, D24202, doi:10.1029/2004JD004727, 2004. 1176

Emmons, L. K., Edwards, D. P., Deeter, M. N., Gille, J. C., Campos, T., Ndleci, P., Novelli, P., and Sachse, G.: Measurements of Pollution In The Troposphere (MOPITT) validation through
10 2006, *Atmos. Chem. Phys.*, 9, 1795–1803, 2009a. 1174

Emmons, L. K., Lamarque, J.-F., Apel, E. C., et al.: Impact of Mexico City emissions on regional air quality from MOZART-4 simulations, *Atmos. Chem. Phys.*, in preparation, 2009b. 1160, 1171

15 Fast, J., Gustafson Jr., W., Easter, R., Zaveri, R., Barnard, J., Chapman, E., and Grell, G.: Evolution of ozone, particulates, and aerosol direct forcing in an urban area using a new fully-coupled meteorology, chemistry, and aerosol model, *J. Geophys. Res.*, 111, D21305, doi:10.1029/2005JD006721, 2006. 1162

20 Fast, J. D., Aiken, A. C., Allan, J., Alexander, L., Campos, T., Canagaratna, M. R., Chapman, E., DeCarlo, P. F., de Foy, B., Gaffney, J., de Gouw, J., Doran, J. C., Emmons, L., Hodzic, A., Herndon, S. C., Huey, G., Jayne, J. T., Jimenez, J. L., Kleinman, L., Kuster, W., Marley, N., Russell, L., Ochoa, C., Onasch, T. B., Pekour, M., Song, C., Ulbrich, I. M., Warneke, C., Welsh-Bon, D., Wiedinmyer, C., Worsnop, D. R., Yu, X.-Y., and Zaveri, R.: Evaluating simulated primary anthropogenic and biomass burning organic aerosols during MILAGRO: implications for assessing treatments of secondary organic aerosols, *Atmos. Chem. Phys. Discuss.*, 9, 4805–4871, 2009, <http://www.atmos-chem-phys-discuss.net/9/4805/2009/>. 1170

25 Fast, J. D., de Foy, B., Acevedo Rosas, F., Caetano, E., Carmichael, G., Emmons, L., McKenna, D., Mena, M., Skamarock, W., Tie, X., Coulter, R. L., Barnard, J. C., Wiedinmyer, C., and Madronich, S.: A meteorological overview of the MILAGRO field campaigns, *Atmos. Chem. Phys.*, 7, 2233–2257, 2007, <http://www.atmos-chem-phys.net/7/2233/2007/>. 1171

30 Gong, S. L., Barrie, L. A., Blanchet, J.-P., von Salzen, K., Lohmann, U., Lesins, G., Spacek, L., Zhang, L. M., Girard, E., Lin, H., Leaitch, R., Leighton, H., Chylek, P., and Huang, P.: Canadian Aerosol Module: A size-segregated simulation of atmospheric aerosol processes for climate and air quality models 1. Module development, *J. Geophys. Res.*, 108(D1), 4007,

GMDD

2, 1157–1213, 2009

MOZART-4 description

L. K. Emmons et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



doi:10.1029/2001JD002002, 2003. 1163

Granier, C., Guenther, A., Lamarque, J., Mieville, A., Muller, J., Olivier, J., Orlando, J., Peters, J., Petron, G., Tyndall, G., and Wallens, S.: POET, a database of surface emissions of ozone precursors, available at: <http://www.aero.jussieu.fr/projet/ACCENT/POET.php>, 2005. 1172

5 Grell, G., Peckham, S., Schmitz, R., McKeen, S., Frost, G., Skamarock, W., and Eder, B.: Fully coupled “online” chemistry within the WRF model, *Atmos. Environ.*, 39, 6957–6976, 2005. 1162

Guenther, A., Karl, T., Harley, P., Wiedinmyer, C., Palmer, P. I., and Geron, C.: Estimates of global terrestrial isoprene emissions using MEGAN (Model of Emissions of Gases and Aerosols from Nature), *Atmos. Chem. Phys.*, 6, 3181–3210, 2006, <http://www.atmos-chem-phys.net/6/3181/2006/>. 1165, 1166, 1167

10 Hack, J. J.: Parameterization of moist convection in the NCAR community climate model (CCM2), *J. Geophys. Res.*, 99, 5551–5568, 1994. 1159

Hess, M., Koepke, P., and Schult, I.: Optical Properties of Aerosols and Clouds: The software package OPAC, *B. Am. Meteorol. Soc.*, 79, 831–844, 1998. 1163

15 Holstlag, A. and Boville, B. A.: Local versus nonlocal boundary-layer diffusion in a global climate model, *J. Climate*, 6, 1825–1842, 1993. 1159

Horowitz, L., Walters, S., Mauzerall, D., Emmons, L., Rasch, P., Granier, C., Tie, X., Lamarque, J.-F., Schultz, M., Tyndall, G., Orlando, J., and Brasseur, G.: A global simulation of tropospheric ozone and related tracers: Description and evaluation of MOZART, version 2, *J. Geophys. Res.*, 108, 4784, doi:10.1029/2002JD002853, 2003. 1159, 1160, 1168, 1169, 1173

20 Horowitz, L. W., Fiore, A. M., Milly, G. P., Cohen, R. C., Perring, A., Wooldridge, P. J., Hess, P. G., Emmons, L. K., and Lamarque, J.-F.: Observational constraints on the chemistry of isoprene nitrates over the eastern United States, *J. Geophys. Res.*, 112, D12S08, doi: 10.1029/2006JD007747, 2007. 1170

IPCC: Intergovernmental Panel on Climate Change, Emission Scenarios, A Special Report of IPCC Working Group III, Cambridge University Press, Cambridge, UK, 2000. 1169

Jacob, D. J.: Heterogeneous chemistry and tropospheric ozone, *Atmos. Environ.*, 34, 2131–2159, 2000. 1162

25 Kalnay, E., Kanamitsu, M., Kistler, R., et al.: The NCEP/NCAR 40-Year Reanalysis Project, *B. Amer. Meteorol. Soc.*, 77, 437–471, 1996. 1160, 1169

Kinnison, D., Brasseur, G. P., Walters, S., et al.: Sensitivity of chemical tracers to meteorological

GMDD

2, 1157–1213, 2009

MOZART-4 description

L. K. Emmons et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



parameters in the MOZART-3 chemical transport model, *J. Geophys. Res.*, 112, D20302, doi:10.1029/2006JD007879, 2007. 1159, 1163, 1164

Kistler, R., Kalnay, E., Collins, W., et al.: The NCEP-NCAR 50-Year Reanalysis: Monthly Means CD-ROM and Documentation, *B. Amer. Meteorol. Soc.*, 82, 247–267, 2001. 1160, 1169

5 Kloster, S., Feichter, J., Maier-Reimer, E., Six, K. D., Stier, P., and Wetzell, P.: DMS cycle in the marine ocean-atmosphere system - a global model study, *Biogeosciences*, 3, 29–51, 2006, <http://www.biogeosciences.net/3/29/2006/>. 1172

Laepfle, T., Schultz, M., Lamarque, J.-F., Madronich, S., Shetter, R., Lefer, B., and Atlas, E.: Improved albedo formulation for chemistry transport models based on satellite observations and assimilated snow data and its impact on tropospheric photochemistry, *J. Geophys. Res.*, 110, D11308, doi:10.1029/2004JD005463, 2005. 1163

Lamarque, J.-F., Kiehl, J. T., Hess, P. G., Collins, W. D., Emmons, L. K., Ginoux, P., Luo, C., and Tie, X. X.: Response of a coupled chemistry-climate model to changes in aerosol emissions: Global impact on the hydrological cycle and the tropospheric burdens of OH, ozone and NO_x, *Geophys. Res. Lett.*, 32, L16809, doi:10.1029/2005GL023419, 2005b. 1162

15 Lapina, K., Honrath, R. E., Owen, R. C., Martín, M. V., and Pfister, G.: Evidence of significant large-scale impacts of boreal fires on ozone levels in the midlatitude Northern Hemisphere free troposphere, *Geophys. Res. Lett.*, 33, L10815, doi:10.1029/2006GL025878, 2006. 1170

Lawrence, M. G., Jöckel, P., and von Kuhlmann, R.: What does the global mean OH concentration tell us?, *Atmos. Chem. Phys.*, 1, 37–49, 2001, <http://www.atmos-chem-phys.net/1/37/2001/>. 1173

20 Lawrence, P. J. and Chase, T.: Representing a new MODIS consistent land surface in the Community Land Model (CLM 3.0), *J. Geophys. Res.*, 112, G01023, doi:G0102310.1029/2006JG000168, 2007. 1165

25 Lin, S. J. and Rood, R. B.: Multidimensional flux-form semi-Lagrangian transport scheme, *Mon. Weather Rev.*, 124, 2046–2070, 1996. 1159

Mahowald, N., Lamarque, J.-F., Tie, X., and Wolff, E.: Sea salt aerosol response to climate change: last glacial maximum, pre-industrial and doubled carbon dioxide climates, *J. Geophys. Res.*, 111, D05303, doi:10.1029/2005JD006459, 2006a. 1163

30 Mahowald, N. M., Muhs, D. R., Levis, S., Rasch, P. J., Yoshioka, M., Zender, C. S., and Luo, C.: Change in atmospheric mineral aerosols in response to climate: Last glacial period, preindustrial, modern, and doubled carbon dioxide climates, *J. Geophys. Res.*, 111, D10202, doi:10.1029/2005JD006653, 2006b. 1163

GMDD

2, 1157–1213, 2009

MOZART-4 description

L. K. Emmons et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- McLinden, C., Olsen, S., Hannegan, B., Wild, O., and Prather, M.: Stratospheric ozone in 3-D models: A simple chemistry and the cross-tropopause flux, *J. Geophys. Res.*, 105, 14653–14665, 2000. 1170
- Mena-Carrasco, M., Carmichael, G. R., Campbell, J. E., Zimmerman, D., Tang, Y., Adhikary, B., D'allura, A., Molina, L. T., Zavala, M., García, A., Flocke, F., Campos, T., Weinheimer, A. J., Shetter, R., Apel, E., Montzka, D. D., Knapp, D. J., and Zheng, W.: Assessing the regional impacts of Mexico City emissions on air quality and chemistry, *Atmos. Chem. Phys.*, 9, 3731–3743, 2009, <http://www.atmos-chem-phys.net/9/3731/2009/>. 1170
- Metzger, S., Dentener, F., Pandis, S., and Lelieveld, J.: Gas/aerosol partitioning: 1. A computationally efficient model, *J. Geophys. Res.*, 107(D16), 4312, doi:10.1029/2001JD001102, 2002. 1162
- Mortlock, A. and Alstyne, R. V.: Military, Charter, Unreported Domestic Traffic and General Aviation: 1976, 1984, 1992, and 2015 Emission Scenarios, NASA CR-1998-207639, available at: http://ntrs.nasa.gov/archive/nasa/casi.ntrs.nasa.gov/19980047346_1998120131.pdf, 1998. 1172
- Novelli, P. C., Masarie, K., and Lang, P.: Molecular hydrogen in the troposphere: Global distributions and budget, *J. Geophys. Res.*, 104, 30427–30444, 1999. 1169
- Novelli, P. C., Masarie, K., Lang, P., Hall, B., Myers, R., and Elkins, J.: Re-analysis of tropospheric CO trends: Effects of the 1997–1998 wildfires, *J. Geophys. Res.*, 108, 4464, doi:10.1029/2002JD003031, 2003. 1173
- Ohara, T., Akimoto, H., Kurokawa, J., Horii, N., Yamaji, K., Yan, X., and Hayasaka, T.: An Asian emission inventory of anthropogenic emission sources for the period 1980–2020, *Atmos. Chem. Phys.*, 7, 4419–4444, 2007, <http://www.atmos-chem-phys.net/7/4419/2007/>. 1172
- Olivier, J. and Berdowski, J.: Global emissions sources and sinks, in: *The Climate System*, edited by: Berdowski, J., Guicherit, R., and Heij, B. J., A. A. Balkema Publishers/Swets, Zeitlinger Publishers, Lisse, The Netherlands, 33–78, 2001. 1172
- Olivier, J., J. Peters, J., Granier, C., Petron, G., Müller, J., and Wallens, S.: Present and future surface emissions of atmospheric compounds, POET report #2, EU project EVK2-1999-00011, <http://www.aero.jussieu.fr/projet/ACCENT/POET.php>, 2003. 1172
- Pfister, G., Hess, P., Emmons, L., Lamarque, J.-F., Wiedinmyer, C., Edwards, D., Pétron, G., Gille, J., and Sachse, G.: Quantifying CO emissions from the 2004 Alaskan wildfires using MOPITT CO data, *Geophys. Res. Lett.*, 32, L11809, doi:10.1029/2005GL022995, 2005. 1170

MOZART-4
descriptionL. K. Emmons et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Pfister, G. G., Emmons, L. K., Hess, P. G., Honrath, R., et al.: Ozone production from the 2004 North American boreal fires, *J. Geophys. Res.*, 111, D24S07, doi:10.1029/2006JD007695, 2006. 1170
- 5 Pfister, G. G., Emmons, L. K., Hess, P. G., Lamarque, J.-F., Orlando, J. J., Walters, S., Guenther, A., Palmer, P. I., and Lawrence, P. J.: Contribution of isoprene to chemical budgets: A model tracer study with the NCAR CTM MOZART-4, *J. Geophys. Res.*, 113, D05308, doi:10.1029/2007JD008948, 2008a. 1161, 1165
- 10 Pfister, G. G., Emmons, L. K., Hess, P. G., Lamarque, J.-F., Thompson, A. M., and Yorks, J. E.: Analysis of the Summer 2004 ozone budget over the United States using Intercontinental Transport Experiment Ozone-sonde Network Study (IONS) observations and Model of Ozone and Related Tracers (MOZART-4) simulations, *J. Geophys. Res.*, 113, D23306, doi:10.1029/2008JD010190, 2008b. 1170
- 15 Pfister, G. G., Edwards, D. P., Emmons, L. K., Arellano, A. A., Sachse, G., and Campos, T.: Transpacific pollution transport during INTEX-B: spring 2006 in context to previous years, *Atmos. Chem. Phys. Discuss.*, in press, 2009. 1171
- Pickering, K., Wang, Y., Tao, W.-K., Price, C., and Muller, J.-F.: Vertical distributions of lightning NO_x for use in regional and global transport models, *J. Geophys. Res.*, 103, 31203–31216, 1998. 1168
- 20 Price, C. and Rind, D.: A simple lightning parameterization for calculating global lightning distributions, *J. Geophys. Res.*, 97, 9919–9933, doi:10.1029/92JD00719, 1992. 1168
- Price, C., Penner, J., and Prather, M.: NO_x from lightning 1. Global distribution based on lightning physics, *J. Geophys. Res.*, 102, 5929–5941, 1997. 1168
- Rasch, P. J., Mahowald, N. M., and Eaton, B. E.: Representations of transport, convection, and the hydrologic cycle in chemical transport models: Implications for the modeling of short-lived and soluble species, *J. Geophys. Res.*, 102, 28127–28138, 1997. 1159
- 25 Ridley, B., Pickering, K., and Dye, J.: Comments on the parameterization of lightning-produced NO in global chemistry-transport models, *Atmos. Environ.*, 39, 6184–6187, 2005. 1168
- Sander, S., Friedl, R. R., Ravishankara, A. R., et al.: Chemical Kinetics and Photochemical Data for Use in Atmospheric Studies, Evaluation Number 14, JPL Publication 02-25, NASA, Jet Propulsion Laboratory, California Institute of Technology, Pasadena, CA, 2003. 1161
- 30 Sanderson, M., Collins, W., Derwent, R., and Johnson, C.: Simulation of global hydrogen levels using a lagrangian three-dimensional model, *J. Atmos. Chem.*, 46, 15–28, 2003. 1164
- Seinfeld, J. H.: *Atmospheric Chemistry and Physics of Air Pollution*, Wiley, New York, 1986.

**MOZART-4
description**L. K. Emmons et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

- Shindell, D. T., Faluvegi, G., Stevenson, D., et al.: Multimodel simulations of carbon monoxide: Comparison with observations and projected near-future changes, *J. Geophys. Res.*, 111, D19306, doi:10.1029/2006JD007100, 2006. 1171, 1174
- 5 Simmons, A. and Gibson, J.: The ERA-40 project plan, in: ERA-40 Project Report Series, Tech. Rep. 1, Eur. Cent. For Medium-Range Weather Forecasts, Reading, UK, 2000. 1169
- Singh, H. B., Brune, W. H., Crawford, J. H., Flocke, F., and Jacob, D. J.: Chemistry and transport of pollution over the Gulf of Mexico and the Pacific: spring 2006 INTEX-B campaign overview and first results, *Atmos. Chem. Phys.*, 9, 2301–2318, 2009,
 10 <http://www.atmos-chem-phys.net/9/2301/2009/>. 1171
- SPARC: Report No. 4, Assessment of Stratospheric Aerosol Properties, WCRP-124, WMO/TD-No 1295, 2006. 1169
- Spivakovsky, C. M., Logan, J. A., Montzka, S. A., Balkanski, Y. J., et al.: Three-dimensional climatological distribution of tropospheric OH: Update and evaluation, *J. Geophys. Res.*, 105, 8931–8980, 2000. 1173
- 15 Stevenson, D. S., Dentener, F., Schultz, M., et al.: Multimodel ensemble simulations of present-day and near-future tropospheric ozone, *J. Geophys. Res.*, 111, D08301, doi: 10.1029/2005JD006338, 2006. 1171
- Sutkus, D., Baughcum, S., and DuBois, D.: Scheduled Civil Aircraft Emission Inventories for 1999: Database Development and Analysis, NASA/CR-2001-211216, available at: <http://gltrs.grc.nasa.gov/reports/2001/CR-2001-211216.pdf>, 2001. 1172
- 20 Tang, I. N. and Munkelwitz, H. R.: Water activities, densities and refractive indices of aqueous sulfates and sodium nitrate droplets of atmospheric importance, *J. Geophys. Res.*, 99, 18801–18808, 1994. 1163
- Tang, Y., Lee, P., Tsidulko, M., et al.: The impact of chemical lateral boundary conditions on CMAQ predictions of tropospheric ozone over the continental United States, *Environ. Fluid Mech.*, 9, 43–58, 2009. 1170
- 25 Tang, Y. H., Carmichael, G. R., Thongboonchoo, N., Chai, T. F., Horowitz, L. W., Pierce, R., Al-Saadi, J. A., Pfister, G., Vukovich, J. M., Avery, M. A., Sachse, G. W., Ryerson, T. B., Holloway, J. S., Atlas, E. L., Flocke, F. M., Weber, R. J., Huey, L. G., Dibb, J. E., Streets, D., and Brune, W. H.: Influence of lateral and top boundary conditions on regional air quality prediction: A multiscale study coupling regional and global chemical transport models, *J. Geophys. Res.*, 112, D10S18, doi:10.1029/2006JD007515, 2007. 1162
- 30

**MOZART-4
description**

L. K. Emmons et al.

[Title Page](#)
[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

- Teklemariam, T. and Sparks, J.: Gaseous fluxes of peroxyacetyl nitrate (PAN) into plant leaves, *Plant Cell Environ.*, 27, 1149–1158, 2004. 1164
- Thompson, A., Witte, J., McPeters, R., Oltmans, S., et al.: Southern Hemisphere Additional Ozonesondes (SHADOZ) 1998–2000 tropical ozone climatology 1. Comparison with Total Ozone Mapping Spectrometer (TOMS) and ground-based measurements, *J. Geophys. Res.*, 108, 8238, doi:10.1029/2001JD000967, 2003. 1175
- Tie, X., Brasseur, G., Emmons, L., Horowitz, L., and Kinnison, D.: Effects of aerosols on tropospheric oxidants: A global model study, *J. Geophys. Res.*, 106, 2931–2964, 2001. 1162
- Tie, X., Madronich, S., Walters, S., Rasch, P., and Collins, W.: Effect of clouds on photolysis and oxidants in the troposphere, *J. Geophys. Res.*, 108, 4642, doi:10.1029/2003JD003659, 2003. 1163
- Tie, X., Madronich, S., Walters, S., Edwards, D., Ginoux, P., Mahowald, N., Zhang, R., Luo, C., and Brasseur, G.: Assessment of the global impact of aerosols on tropospheric oxidants, *J. Geophys. Res.*, 110, D03204, doi:10.1029/2004JD005359, 2005. 1162
- Toon, O. B., Pollack, J. B., and Khare, B. N.: The optical constants of several atmospheric aerosol species: ammonium sulfate, aluminum oxide and sodium chloride, *J. Geophys. Res.*, 81, 5733–5744, 1976. 1163
- Turnipseed, A., Huey, G., Nemitz, E., Stickel, R., Higgs, J., Tanner, D. and Slusher, D., Sparks, J., Flocke, F., and Guenther, A.: Eddy covariance fluxes of peroxyacyl nitrates (PANs) and NO_y to a coniferous forest, *J. Geophys. Res.*, 111, D09304, doi:10.1029/2005JD006631, 2006. 1164
- van der Werf, G. R., Randerson, J. T., Giglio, L., Collatz, G. J., Kasibhatla, P. S., and Arellano Jr., A. F.: Interannual variability in global biomass burning emissions from 1997 to 2004, *Atmos. Chem. Phys.*, 6, 3423–3441, 2006, <http://www.atmos-chem-phys.net/6/3423/2006/>. 1172
- Van Noije, T., Eskes, H., van Weele, M., and van Velthoven, P.: Implications of the enhanced Brewer-Dobson circulation in European Centre for Medium-Range Weather Forecasts re-analysis ERA-40 for the stratosphere-troposphere exchange of ozone in global chemistry transport models, *J. Geophys. Res.*, 109, D19308, doi:10.1029/2004JD004586, 2004. 1169
- Walmsley, J. and Wesely, M.: Modification of coded parametrizations of surface resistances to gaseous dry deposition, *Atmos. Environ.*, 30, 1181–1188, 1996. 1164
- Wesely, M.: Parameterization of surface resistances to gaseous dry deposition in regional-scale numerical models, *Atmos. Environ.*, 23, 1293–1304, 1989. 1164

MOZART-4
descriptionL. K. Emmons et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

- Wesely, M. and Hicks, B.: A review of the current status of knowledge on dry deposition, *Atmos. Environ.*, 34, 2261–2282, 2000. 1164
- Yienger, J. and Levy, H.: Empirical model of global soil biogenic NO_x emissions, *J. Geophys. Res.*, 100, 11447–11464, 1995. 1168
- 5 Yonemura, S., Kawashima, S., and Tsuruta, H.: Carbon monoxide, hydrogen, and methane uptake by soils in a temperate arable field and a forest, *J. Geophys. Res.*, 105, 14347–14362, 2000. 1164
- Zender, C. S., Bian, H., and Newman, D.: Mineral Dust Entrainment and Deposition (DEAD) model: Description and 1990s dust climatology, *J. Geophys. Res.*, 108, 4416, doi:10.1029/2002JD002775, 2003. 1163
- 10 Zhang, G. J. and MacFarlane, N. A.: Sensitivity of climate simulations to the parameterization of cumulus convection in the Canadian climate centre general circulation model, *Atmos. Ocean*, 33, 407–446, 1995. 1159

GMDD

2, 1157–1213, 2009

**MOZART-4
description**

L. K. Emmons et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Table 1. Meteorological inputs required for MOZART-4.

Variable	MOZART-4 Name	Dimension	Units
Zonal Winds	U	3-D	m s^{-1}
Meridional Winds	V	3-D	m s^{-1}
Temperature	T	3-D	K
Specific Humidity (optional)	Q	3-D	kg kg^{-1}
Surface Temperature	TS	2-D	K
Surface Pressure	PS	2-D	Pa
Land/Ocean/Sea Ice Flag	ORO	2-D	unitless
Surface Geopotential Height	PHIS	2-D	$\text{m}^2 \text{s}^{-2}$
Surface Zonal Stress	TAUX	2-D	N m^{-2}
Surface Meridional Stress	TAUY	2-D	N m^{-2}
Surface Heat Flux	SHFLX	2-D	W m^{-2}
Surface Moisture Flux	QLFX	2-D	$\text{kg m}^{-2} \text{s}^{-1}$
Solar Flux at Surface	FSDS	2-D	W m^{-2}
Snow Height	SNOWH	2-D	m
Soil Moisture Fraction ¹	SOILW	2-D	unitless
Previous month's average TS ¹	TS _{avg}	2-D	K
Previous month's average FSDS ¹	FSDS _{avg}	2-D	W m^{-2}

¹ If SOILW, TS_{avg} or FSDS_{avg} are not available in met fields, a climatology is used.

MOZART-4 description

L. K. Emmons et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Table 2. Chemical species in MOZART-4.

	Symbolic Name	Atomic Composition	Comments
O, H, N species	O	O(³ P)	
	O1D	O(¹ D)	
	O3	O ₃	ozone
	N2O	N ₂ O	
	NO	NO	
	NO2	NO ₂	
	NO3	NO ₃	
	HNO3	HNO ₃	
	HO2NO2	HNO ₄	
	N2O5	N ₂ O ₅	
	H2	H ₂	
	OH	OH	
	HO2	HO ₂	
	H2O2	H ₂ O ₂	
	C ₁ species	CH4	CH ₄
CO		CO	carbon monoxide
CH3O2		CH ₃ O ₂	
CH3OOH		CH ₃ OOH	
CH2O		CH ₂ O	
CH3OH		CH ₃ OH	methanol
C ₂ species		C2H4	C ₂ H ₄
	C2H6	C ₂ H ₆	ethane
	CH3CHO	CH ₃ CHO	acetaldehyde
	C2H5OH	C ₂ H ₅ OH	ethanol
	EO	HOCH ₂ CH ₂ O	
	EO2	HOCH ₂ CH ₂ O ₂	
	CH3COOH	CH ₃ COOH	acetic acid
	GLYOXAL	HCOCHO	
	GLYALD	HOCH ₂ CHO	glycolaldehyde
	C2H5O2	C ₂ H ₅ O ₂	
	C2H5OOH	C ₂ H ₅ OOH	
	CH3CO3	CH ₃ CO ₃	
	CH3COOOH	CH ₃ CO ₃ H	peracetic acid
	PAN	CH ₃ CO ₃ NO ₂	peroxy acetyl nitrate

**MOZART-4
description**

L. K. Emmons et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

I ◀

▶ I

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Table 2. Continued.

	Symbolic Name	Atomic Composition	Comments
C ₃ species	C3H6		
	C3H8		
	C3H7O2		
	C3H7OOH		
	PO2	e.g., CH ₃ CH(OO)CH ₂ OH	
	POOH	e.g., CH ₃ CH(OOH)CH ₂ OH	
	CH3COCH3		acetone
	HYAC	CH ₃ COCH ₂ OH	hydroxyacetone
	CH3COCHO		methyl glyoxal
	RO2	CH ₃ COCH ₂ O ₂	
	ROOH	CH ₃ COCH ₂ OOH	
	ONIT	CH ₃ COCH ₂ ONO ₂	
	C ₄ species	BIGENE	C ₄ H ₈
ENE02		e.g., CH ₃ CH(OH)CH(OO)CH ₃	
MEK		CH ₃ C(O)CH ₂ CH ₃	methyl ethyl ketone
MEK02		CH ₃ COCH(OO)CH ₃	
MEKOOH		CH ₃ COCH(OOH)CH ₃	
MVK		CH ₂ CHCOCH ₃	methyl vinyl ketone
MACR		CH ₂ CCH ₃ CHO	methacrolein
MPAN		CH ₂ CCH ₃ CO ₃ NO ₂	methacryloyl peroxyxynitrate
MACRO2		e.g., CH ₃ COCH(OO)CH ₂ OH	peroxy radical from OH addition to MVK, MACR
MACROOH		e.g., CH ₃ COCH(OOH)CH ₂ OH	
MCO3		CH ₂ CCH ₃ CO ₃	peroxy radical derived from abstraction reaction of OH with MACR
C ₅ species	BIGALK	C ₅ H ₁₂	lumped alkanes C > 3
	ALKO2	C ₅ H ₁₁ O ₂	
	ALKOOH	C ₅ H ₁₁ OOH	
	ISOP	C ₅ H ₈	isoprene
	ISOPO2	e.g., HOCH ₂ C(OO)CH ₃ CHCH ₂	peroxy radical derived from OH+ISOP
	ISOPOOH	e.g., HOCH ₂ C(OOH)CH ₃ CHCH ₂	unsaturated hydroxyhydroperoxide

GMDD

2, 1157–1213, 2009

**MOZART-4
description**

L. K. Emmons et al.

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[I ◀](#)
[▶ I](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)


Table 2. Continued.

	Symbolic Name	Atomic Composition	Comments
	HYDRALD	e.g., HOCH ₂ CCH ₃ CHCHO	lumped unsaturated hydroxycarbonyl
	XO2	e.g., HOCH ₂ C(OO)CH ₃ CH(OH)CHO	peroxy radical from OH+HYDRALD
	XOOH BIGALD	e.g., HOCH ₂ C(OOH)CH ₃ CH(OH)CHO C ₅ H ₆ O ₂	unsaturated dicarbonyl, oxidation product of toluene
	ISOPNO3 ONITR	CH ₂ CHCCH ₃ OOCH ₂ ONO ₂ CH ₂ CCH ₃ CHONO ₂ CH ₂ OH	peroxy radical from NO ₃ +ISOP lumped isoprene nitrate
C ₇ species	TOLUENE CRESOL TOL02 TOLOOH XOH	C ₆ H ₅ (CH ₃) C ₆ H ₄ (CH ₃)(OH) C ₇ H ₉ O ₅ C ₇ H ₁₀ O ₅ C ₇ H ₁₀ O ₆	lumped aromatics
C ₁₀ species	C10H16	C ₁₀ H ₁₆	lumped monoterpenes, as α -pinene
	TERPO2 TERPOOH	C ₁₀ H ₁₆ (OH)(OO) C ₁₀ H ₁₆ (OH)(OOH)	
Aerosol precursors	SO2 DMS NH3		
Bulk aerosols	SO4 NH4 NH4NO3 SOA CB1 CB2 OC1 OC2 SA1 SA2 SA3 SA4	C ₁₂ C C C C NaCl NaCl NaCl NaCl	sulfate ammonium ammonium nitrate secondary organic aerosol black carbon, hydrophobic black carbon, hydrophylic organic carbon, hydrophobic organic carbon, hydrophylic sea salt, 0.1–0.5 μ m sea salt, 0.5–1.5 μ m sea salt, 1.5–5 μ m sea salt, 5–10 μ m

**MOZART-4
description**

L. K. Emmons et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Table 3. Gas-phase reactions. ({CO2} indicates CO2 is not included in the model solution.)

Reactants	Products	Rate
O+O2+M	→ O3+M	6E-34*(300/T)**2.4
O+O3	→ 2*O2	8.00E-12*exp(-2060./T)
O1D+N2	→ O+N2	2.10E-11*exp(115./T)
O1D+O2	→ O+O2	3.20E-11*exp(70./T)
O1D+H2O	→ 2*OH	2.20E-10
H2+O1D	→ HO2+OH	1.10E-10
H2+OH	→ H2O+HO2	5.50E-12*exp(-2000./T)
O+OH	→ HO2+O2	2.20E-11*exp(120./T)
HO2+O	→ OH+O2	3.00E-11*exp(200./T)
OH+O3	→ HO2+O2	1.70E-12*exp(-940./T)
HO2+O3	→ OH+2*O2	1.00E-14*exp(-490./T)
HO2+HO2	→ H2O2+O2	(2.3E-13*exp(600/T)+1.7E-33*[M]*exp(1000/T))* (1+1.4E-21*[H2O]*exp(2200/T))
H2O2+OH	→ H2O+HO2	2.90E-12*exp(-160./T)
OH+HO2	→ H2O+O2	4.80E-11*exp(250./T)
OH+OH	→ H2O+O	4.20E-12*exp(-240./T)
OH+OH+M	→ H2O2+M	ko=6.90E-31*(300/T)**1.00; ki=2.60E-11; f=0.60
N2O+O1D	→ N2+O2	4.90E-11
N2O+O1D	→ 2*NO	6.70E-11
NO+HO2	→ NO2+OH	3.50E-12*exp(250./T)
NO+O3	→ NO2+O2	3.00E-12*exp(-1500./T)
NO2+O	→ NO+O2	5.60E-12*exp(180./T)
NO2+O3	→ NO3+O2	1.20E-13*exp(-2450./T)
NO3+HO2	→ OH+NO2	2.30E-12*exp(170./T)
NO2+NO3+M	→ N2O5+M	ko=2.00E-30*(300/T)**4.40; ki=1.40E-12*(300/T)**0.70; f=0.60
N2O5+M	→ NO2+NO3+M	k(NO2+NO3+M)*3.333E26*exp(-10990/T)
NO2+OH+M	→ HNO3+M	ko=2.00E-30*(300/T)**3.00; ki=2.50E-1; f=0.60
HNO3+OH	→ NO3+H2O	k0+k3[M]/(1+k3[M]/k2); k0=2.4E-14*exp(460/T); k2=2.7E-17*exp(2199/T); k3=6.5E-34*exp(1335/T)
NO3+NO	→ 2*NO2	1.50E-11*exp(170./T)
NO2+HO2+M	→ HO2NO2+M	ko=1.80E-31*(300/T)**3.20; ki=4.70E-12*(300/T)**1.40; f=0.60
HO2NO2+OH	→ H2O+NO2+O2	1.30E-12*exp(380./T)
HO2NO2+M	→ HO2+NO2+M	k(NO2+HO2+M)*exp(-10900/T)/2.1E-27

MOZART-4 description

L. K. Emmons et al.

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)


Table 3. Continued.

Reactants	Products	Rate
CH ₄ +OH	→ CH ₃ O ₂ +H ₂ O	2.45E-12*exp(-1775./T)
CH ₄ +O ₁ D	→ .75*CH ₃ O ₂ +.75*OH+.25*CH ₂ O +.4*HO ₂ +.05*H ₂	1.50E-10
CH ₃ O ₂ +NO	→ CH ₂ O+NO ₂ +HO ₂	2.80E-12*exp(300./T)
CH ₃ O ₂ +CH ₃ O ₂	→ 2*CH ₂ O+2*HO ₂	5.00E-13*exp(-424./T)
CH ₃ O ₂ +CH ₃ O ₂	→ CH ₂ O+CH ₃ OH	1.90E-14*exp(706./T)
CH ₃ O ₂ +HO ₂	→ CH ₃ OOH+O ₂	4.10E-13*exp(750./T)
CH ₃ OOH+OH	→ .7*CH ₃ O ₂ +.3*OH+.3*CH ₂ O+H ₂ O	3.80E-12*exp(200./T)
CH ₂ O+NO ₃	→ CO+HO ₂ +HNO ₃	6.00E-13*exp(-2058./T)
CH ₂ O+OH	→ CO+H ₂ O+HO ₂	9.00E-12
CO+OH	→ {CO ₂ }+HO ₂	1.5E-13*(1+6E-7*P)
CH ₃ OH+OH	→ HO ₂ +CH ₂ O	7.30E-12*exp(-620./T)
C ₂ H ₄ +OH+M	→ .75*EO ₂ +.5*CH ₂ O+.25*HO ₂ +M	ko=1.00E-28*(300/T)**0.80; ki=8.80E-12; f=0.60
C ₂ H ₄ +O ₃	→ CH ₂ O+.12*HO ₂ +.5*CO +.12*OH+.25*CH ₃ COOH	1.20E-14*exp(-2630./T)
EO ₂ +NO	→ EO+NO ₂	4.20E-12*exp(180./T)
EO+O ₂	→ GLYALD+HO ₂	1.00E-14
EO	→ 2*CH ₂ O+HO ₂	1.60E+11*exp(-4150./T)
C ₂ H ₆ +OH	→ C ₂ H ₅ O ₂ +H ₂ O	8.70E-12*exp(-1070./T)
C ₂ H ₅ O ₂ +NO	→ CH ₃ CHO+HO ₂ +NO ₂	2.60E-12*exp(365./T)
C ₂ H ₅ O ₂ +HO ₂	→ C ₂ H ₅ OOH+O ₂	7.50E-13*exp(700./T)
C ₂ H ₅ O ₂ +CH ₃ O ₂	→ .7*CH ₂ O+.8*CH ₃ CHO+HO ₂ +.3*CH ₃ OH+.2*C ₂ H ₅ OH	2.00E-13
C ₂ H ₅ O ₂ +C ₂ H ₅ O ₂	→ 1.6*CH ₃ CHO+1.2*HO ₂ +.4*C ₂ H ₅ OH	6.80E-14
C ₂ H ₅ OOH+OH	→ .5*C ₂ H ₅ O ₂ +.5*CH ₃ CHO+.5*OH	3.80E-12*exp(200./T)
CH ₃ CHO+OH	→ CH ₃ CO ₃ +H ₂ O	5.60E-12*exp(270./T)
CH ₃ CHO+NO ₃	→ CH ₃ CO ₃ +HNO ₃	1.40E-12*exp(-1900./T)
CH ₃ CO ₃ +NO	→ CH ₃ O ₂ +{CO ₂ }+NO ₂	8.10E-12*exp(270./T)
CH ₃ CO ₃ +NO ₂ +M	→ PAN+M	ko=8.50E-29*(300/T)**6.50; ki=1.10E-11*(300/T); f=0.60
CH ₃ CO ₃ +HO ₂	→ .75*CH ₃ COOOH+.25*CH ₃ COOH+.25*O ₃	4.30E-13*exp(1040./T)
CH ₃ CO ₃ +CH ₃ O ₂	→ .9*CH ₃ O ₂ +CH ₂ O+.9*HO ₂ +.1*CH ₃ COOH+.9*{CO ₂ }	2.00E-12*exp(500./T)
CH ₃ COOOH+OH	→ .5*CH ₃ CO ₃ +.5*CH ₂ O+.5*{CO ₂ }+H ₂ O	1.00E-12
PAN+OH	→ CH ₂ O+NO ₃ +{CO ₂ }	4.00E-14

GMDD

2, 1157–1213, 2009

**MOZART-4
description**

L. K. Emmons et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Table 3. Continued.

Reactants	Products	Rate
PAN+M	→ CH ₃ CO ₃ +NO ₂ +M	k(CH ₃ CO ₃ +NO ₂ +M)*1.111E28 *exp(-14000/T)
CH ₃ CO ₃ +CH ₃ CO ₃	→ 2*CH ₃ O ₂ +2*{CO ₂ }	2.50E-12*exp(500./T)
GLYALD+OH	→ HO ₂ +2*GLYOXAL+.8*CH ₂ O+.8*{CO ₂ }	1.00E-11
GLYOXAL+OH	→ HO ₂ +CO+{CO ₂ }	1.10E-11
CH ₃ COOH+OH	→ CH ₃ O ₂ +{CO ₂ }+H ₂ O	7.00E-13
C ₂ H ₅ OH+OH	→ HO ₂ +CH ₃ CHO	6.90E-12*exp(-230./T)
C ₃ H ₆ +OH+M	→ PO ₂ +M	ko=8.00E-27*(300./T)**3.50; ki=3.00E-11; f=0.50
C ₃ H ₆ +O ₃	→ .54*CH ₂ O+.19*HO ₂ +.33*OH +.5*CH ₃ CHO+.56*CO+.31*CH ₃ O ₂ +.25*CH ₃ COOH+.08*CH ₄	6.50E-15*exp(-1900./T)
C ₃ H ₆ +NO ₃	→ ONIT	4.60E-13*exp(-1156./T)
PO ₂ +NO	→ CH ₃ CHO+CH ₂ O+HO ₂ +NO ₂	4.20E-12*exp(180./T)
PO ₂ +HO ₂	→ POOH+O ₂	7.50E-13*exp(700./T)
POOH+OH	→ .5*PO ₂ +.5*OH+.5*HYAC+H ₂ O	3.80E-12*exp(200./T)
C ₃ H ₈ +OH	→ C ₃ H ₇ O ₂ +H ₂ O	1.00E-11*exp(-660./T)
C ₃ H ₇ O ₂ +NO	→ .82*CH ₃ COCH ₃ +NO ₂ +.27*CH ₃ CHO+HO ₂	4.20E-12*exp(180./T)
C ₃ H ₇ O ₂ +HO ₂	→ C ₃ H ₇ OOH+O ₂	7.50E-13*exp(700./T)
C ₃ H ₇ O ₂ +CH ₃ O ₂	→ CH ₂ O+HO ₂ +.82*CH ₃ COCH ₃	3.75E-13*exp(-40./T)
C ₃ H ₇ OOH+OH	→ H ₂ O+C ₃ H ₇ O ₂	3.80E-12*exp(200./T)
CH ₃ COCH ₃ +OH	→ RO ₂ +H ₂ O	3.82E-11*exp(-2000/T)+1.33E-13
RO ₂ +NO	→ CH ₃ CO ₃ +CH ₂ O+NO ₂	2.90E-12*exp(300./T)
RO ₂ +HO ₂	→ ROOH+O ₂	8.60E-13*exp(700./T)
RO ₂ +CH ₃ O ₂	→ .3*CH ₃ CO ₃ +.2*HYAC+.8*CH ₂ O +.5*CH ₃ OH+.3*HO ₂ +.5*CH ₃ COCHO	2.00E-12*exp(500./T)
ROOH+OH	→ RO ₂ +H ₂ O	3.80E-12*exp(200./T)
ONIT+OH	→ NO ₂ +CH ₃ COCHO	6.80E-13
CH ₃ COCHO+OH	→ CH ₃ CO ₃ +CO+H ₂ O	8.40E-13*exp(830./T)
CH ₃ COCHO+NO ₃	→ HNO ₃ +CO+CH ₃ CO ₃	1.40E-12*exp(-1860./T)
HYAC+OH	→ CH ₃ COCHO+HO ₂	3.00E-12
BIGENE+OH	→ ENEO ₂	5.40E-11
ENE ₂ +NO	→ CH ₃ CHO+.5*CH ₂ O+.5*CH ₃ COCH ₃ +HO ₂ +NO ₂	4.20E-12*exp(180./T)
MEK+OH	→ MEKO ₂	2.30E-12*exp(-170./T)
MEKO ₂ +NO	→ CH ₃ CO ₃ +CH ₃ CHO+NO ₂	4.20E-12*exp(180./T)

GMDD

2, 1157–1213, 2009

MOZART-4
description

L. K. Emmons et al.

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[◀](#)
[▶](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)


Table 3. Continued.

Reactants	Products	Rate
MEKO2+HO2	→ MEKOOH	7.50E-13*exp(700./T)
MEKOOH+OH	→ MEKO2	3.80E-12*exp(200./T)
MPAN+OH	→ .5*HYAC+.5*NO3+.5*CH2O +.5*HO2+.5*{CO2}	ko=8.00E-27*(300/T)**3.50; ki=3.00E-11; f=0.50
BIGALK+OH	→ ALKO2	3.50E-12
ALKO2+NO	→ .4*CH3CHO+.1*CH2O+.25*CH3COCH3 +.9*HO2+.75*MEK+.9*NO2+.1*ONIT	4.20E-12*exp(180./T)
ALKO2+HO2	→ ALKOOH	7.50E-13*exp(700./T)
ALKOOH+OH	→ ALKO2	3.80E-12*exp(200./T)
ISOP+OH	→ ISOPO2	2.54E-11*exp(410./T)
ISOP+O3	→ .4*MACR+.2*MVK+.07*C3H6+.27*OH +.06*HO2+.6*CH2O+.3*CO +.1*O3+.2*MCO3+.2*CH3COOH	1.05E-14*exp(-2000./T)
ISOPO2+NO	→ .08*ONITR+.92*NO2+HO2+.55*CH2O +.23*MACR+.32*MVK+.37*HYDRALD	4.40E-12*exp(180./T)
ISOPO2+NO3	→ HO2+NO2+.6*CH2O+.25*MACR +.35*MVK+.4*HYDRALD	2.40E-12
ISOPO2+HO2	→ ISOPOOH	8.00E-13*exp(700./T)
ISOPOOH+OH	→ .5*XO2+.5*ISOPO2	1.52E-11*exp(200./T)
ISOPO2+CH3O2	→ 1.2*CH2O+.19*MACR+.26*MVK+.3*HYDRALD +.25*CH3OH+HO2	5.00E-13*exp(400./T)
ISOPO2+CH3CO3	→ .6*CH2O+.25*MACR+.35*MVK +.4*HYDRALD+CH3O2+HO2+{CO2}	1.40E-11
ISOP+NO3	→ ISOPNO3	3.03E-12*exp(-446./T)
ISOPNO3+NO	→ 1.206*NO2+.072*CH2O+.167*MACR +.039*MVK+.794*ONITR+.794*HO2	2.70E-12*exp(360./T)
ISOPNO3+NO3	→ 1.206*NO2+.072*CH2O+.167*MACR +.039*MVK+.794*ONITR+.794*HO2	2.40E-12
ISOPNO3+HO2	→ .206*NO2+.008*CH2O+.167*MACR +.039*MVK+.794*ONITR+.794*HO2	8.00E-13*exp(700./T)

GMDD

2, 1157–1213, 2009

**MOZART-4
description**

L. K. Emmons et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Table 3. Continued.

Reactants	Products	Rate
MVK+OH	→ MACRO2	4.13E-12*exp(452./T)
MVK+O3	→ .8*CH2O+.95*CH3COCHO+.08*OH+.2*O3 +.06*HO2+.05*CO+.04*CH3CHO	7.52E-16*exp(-1521./T)
MACR+OH	→ .5*MACRO2+.5*H2O+.5*MCO3	1.86E-11*exp(175./T)
MACR+O3	→ .8*CH3COCHO+.275*HO2+.2*CO+.2*O3 +.7*CH2O+.215*OH	4.40E-15*exp(-2500./T)
MACRO2+NO	→ NO2+.47*HO2+.25*CH2O+.25*CH3COCHO +.53*CH3CO3+.53*GLYALD+.22*HYAC+.22*CO	2.70E-12*exp(360./T)
MACRO2+NO	→ 0.8*ONITR	1.30E-13*exp(360./T)
MACRO2+NO3	→ NO2+.53*GLYALD+.22*HYAC +.53*CH3CO3+.25*CH2O +.22*CO+.25*CH3COCHO+.47*HO2	2.40E-12
MACRO2+HO2	→ MACROOH	8.00E-13*exp(700./T)
MACRO2+CH3O2	→ .73*HO2+.88*CH2O+.11*CO +.24*CH3COCHO+.26*GLYALD +.26*CH3CO3+.25*CH3OH+.23*HYAC	5.00E-13*exp(400./T)
MACRO2+CH3CO3	→ .25*CH3COCHO+CH3O2+.22*CO +.47*HO2+.53*GLYALD+.22*HYAC +.25*CH2O+.53*CH3CO3+{CO2}	1.40E-11
MACROOH+OH	→ .5*MCO3+.2*MACRO2+.1*OH+.2*HO2	2.30E-11*exp(200./T)
MCO3+NO	→ NO2+CH2O+CH3CO3+{CO2}	5.30E-12*exp(360./T)
MCO3+NO3	→ NO2+CH2O+CH3CO3+{CO2}	5.00E-12
MCO3+HO2	→ .25*O3+.25*CH3COOH +.75*CH3COOOH+.75*O2	4.30E-13*exp(1040./T)
MCO3+CH3O2	→ 2*CH2O+HO2+{CO2}+CH3CO3	2.00E-12*exp(500./T)
MCO3+CH3CO3	→ 2*{CO2}+CH3O2+CH2O+CH3CO3	4.60E-12*exp(530./T)
MCO3+MCO3	→ 2*{CO2}+2*CH2O+2*CH3CO3	2.30E-12*exp(530./T)
MCO3+NO2+M	→ MPAN+M	1.1E-11*300/T/[M]
MPAN+M	→ MCO3+NO2+M	k(MCO3+NO2+M)*1.111E28 *exp(-14000/T)
ONITR+OH	→ HYDRALD+.4*NO2+HO2	4.50E-11
ONITR+NO3	→ HYDRALD+NO2+HO2	1.40E-12*exp(-1860./T)
HYDRALD+OH	→ XO2	1.86E-11*exp(175./T)
XO2+NO	→ NO2+1.5*HO2+CO+.25*CH3COCHO +.25*HYAC+.25*GLYALD	2.70E-12*exp(360./T)
XO2+NO3	→ NO2+1.5*HO2+CO+.25*CH3COCHO	2.40E-12

GMDD

2, 1157–1213, 2009

**MOZART-4
description**

L. K. Emmons et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Table 3. Continued.

Reactants	Products	Rate
XO2+HO2	→ XOOH	8.00E-13*exp(700./T)
XO2+CH3O2	→ .3*CH3OH+HO2+.7*CH2O+.4*CO+.1*HYAC +.1*CH3COCHO+.1*GLYALD	5.00E-13*exp(400./T)
XO2+CH3CO3	→ CO+CH3O2+1.5*HO2+{CO2} +.25*HYAC+.25*CH3COCHO +.25*GLYALD	1.30E-12*exp(640./T)
XOOH+OH	→ H2O+XO2	1.90E-12*exp(190./T)
XOOH+OH	→ H2O+OH	T**2 * 7.69E-17* exp(253/T)
TOLUENE+OH	→ .25*CRESOL+.25*HO2+.7*TOLO2	1.70E-12*exp(352./T)
CRESOL+OH	→ XOH	3.00E-12
XOH+NO2	→ .7*NO2+.7*BIGALD+.7*HO2	1.00E-11
TOLO2+NO	→ .45*GLYOXAL+.45*CH3COCHO +.9*BIGALD+.9*NO2+.9*HO2	4.20E-12*exp(180./T)
TOLO2+HO2	→ TOLOOH	7.50E-13*exp(700./T)
TOLOOH+OH	→ TOLO2	3.80E-12*exp(200./T)
C10H16+OH	→ TERPO2	1.20E-11*exp(444./T)
C10H16+O3	→ .7*OH+MVK+MACR+HO2	1.00E-15*exp(-732./T)
C10H16+NO3	→ TERPO2+NO2	1.20E-12*exp(490./T)
TERPO2+NO	→ .1*CH3COCH3+HO2+MVK +MACR+NO2	4.20E-12*exp(180./T)
TERPO2+HO2	→ TERPOOH	7.50E-13*exp(700./T)
TERPOOH+OH	→ TERPO2	3.80E-12*exp(200./T)
SO2+OH	→ SO4	ko=3.0E-31*(300/T)**3.3; ki=1.E-12; f=0.6
DMS+OH	→ SO2	9.60E-12*exp(-234./T)
DMS+OH	→ .5*SO2+.5*HO2	1.7E-42*exp(7810/T)*[M]**0.21/ (1+5.5E-31*exp(7460/T)*[M]**0.21)
DMS+NO3	→ SO2+HNO3	1.90E-13*exp(520./T)
NH3+OH	→ H2O	1.70E-12*exp(-710./T)

MOZART-4 description

L. K. Emmons et al.

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)


Table 4. Photolysis reactions.

Reactant	Products
O ₂ +hν	→ 2*O
O ₃ +hν	→ O1D+O ₂
O ₃ +hν	→ O+O ₂
N ₂ O+hν	→ O1D+N ₂
NO ₂ +hν	→ NO+O
N ₂ O ₅ +hν	→ NO ₂ +NO ₃
HNO ₃ +hν	→ NO ₂ +OH
NO ₃ +hν	→ .89*NO ₂ +.11*NO+.89*O ₃
HO ₂ NO ₂ +hν	→ .33*OH+.33*NO ₃ +.66*NO ₂ +.66*HO ₂
CH ₃ OOH+hν	→ CH ₂ O+HO ₂ +OH
CH ₂ O+hν	→ CO+2*HO ₂
CH ₂ O+hν	→ CO+H ₂
H ₂ O ₂ +hν	→ 2*OH
CH ₃ CHO+hν	→ CH ₃ O ₂ +CO+HO ₂
POOH+hν	→ CH ₃ CHO+CH ₂ O+HO ₂ +OH
CH ₃ COOOH+hν	→ CH ₃ O ₂ +OH+{CO ₂ }
PAN+hν	→ .6*CH ₃ CO ₃ +.6*NO ₂ +.4*CH ₃ O ₂ +.4*NO ₃ +.4*{CO ₂ }
MPAN+hν	→ MCO ₃ +NO ₂
MACR+hν	→ .67*HO ₂ +.33*MCO ₃ +.67*CH ₂ O+.67*CH ₃ CO ₃ +.33*OH+0.67*CO
MVK+hν	→ .7*C ₃ H ₆ +.7*CO+.3*CH ₃ O ₂ +.3*CH ₃ CO ₃
C ₂ H ₅ OOH+hν	→ CH ₃ CHO+HO ₂ +OH
C ₃ H ₇ OOH+hν	→ 0.82*CH ₃ COCH ₃ +OH+HO ₂
ROOH+hν	→ CH ₃ CO ₃ +CH ₂ O+OH
CH ₃ COCH ₃ +hν	→ CH ₃ CO ₃ +CH ₃ O ₂
CH ₃ COCHO+hν	→ CH ₃ CO ₃ +CO+HO ₂
XOOH+hν	→ OH
ONITR+hν	→ HO ₂ +CO+NO ₂ +CH ₂ O
ISOPOOH+hν	→ .402*MVK+.288*MACR+.69*CH ₂ O+HO ₂
HYAC+hν	→ CH ₃ CO ₃ +HO ₂ +CH ₂ O
GLYALD+hν	→ 2*HO ₂ +CO+CH ₂ O
MEK+hν	→ CH ₃ CO ₃ +C ₂ H ₅ O ₂
BIGALD+hν	→ .45*CO+.13*GLYOXAL+.56*HO ₂ +.13*CH ₃ CO ₃ +0.18*CH ₃ COCHO
GLYOXAL+hν	→ 2*CO+2*HO ₂
ALKOOH+hν	→ .4*CH ₃ CHO+.1*CH ₂ O+.25*CH ₃ COCH ₃ +.9*HO ₂ +.8*MEK+OH
MEKOOH+hν	→ OH+CH ₃ CO ₃ +CH ₃ CHO
TOLOOH+hν	→ OH+.45*GLYOXAL+.45*CH ₃ COCHO+.9*BIGALD
TERPOOH+hν	→ OH+.1*CH ₃ COCH ₃ +HO ₂ +MVK+MACR

MOZART-4
description

L. K. Emmons et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



MOZART-4 description

L. K. Emmons et al.

Table 5. Heterogeneous reactions included in MOZART-4, along with the reaction probability (γ) and the type of aerosols on which reactions occur.

Reaction	γ	Aerosol
$\text{N}_2\text{O}_5 + \text{H}_2\text{O} \rightarrow 2 \text{HNO}_3$	0.1	OC, SO ₄ , NH ₄ NO ₃ , SOA
$\text{NO}_3 \rightarrow \text{HNO}_3$	0.001	OC, SO ₄ , NH ₄ NO ₃ , SOA
$\text{NO}_2 \rightarrow 0.5 \text{NO} + 0.5 \text{HNO}_3 + 0.5 \text{OH}$	0.0001	OC, SO ₄ , NH ₄ NO ₃ , SOA
$\text{HO}_2 \rightarrow 0.5 \text{H}_2\text{O}_2$	0.2	OC, SO ₄ , NH ₄ NO ₃ , SOA

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)


MOZART-4 description

L. K. Emmons et al.

Table 6. Bulk aerosol parameters used in calculation of surface area: number distribution mean radius (r_m), geometric standard deviation (σ_g) and density.

Aerosol	r_m (μm)	σ_g (μm)	ρ (g/cm^3)
CB1, CB2	0.0118	2.00	1.0
OC1, OC2	0.0212	2.20	1.8
SO4	0.0695	2.03	1.7
NH4NO3	0.0695	2.03	1.7
SOA	0.0212	2.20	1.8

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[I◀](#)
[▶I](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)


Table 7. Approximate matching of MOZART-4 VOCs to other mechanisms.

MOZART-4	SAPRC-99	RADM2	CBMZ
C2H6	ALK1	ETH	C2H6
C3H8	ALK2	HC3	PAR
BIGALK	ALK3+ALK4+ALK5	HC5	PAR
C2H4	ETHE	OL2	ETH
C3H6	OLE1		PAR
BIGENE	OLE2	OLET+OLEI	OLET+OLEI, PAR
TOLUENE	ARO1+ARO2	TOL+XYL	TOL+XYL
ISOP	ISOPRENE	ISO	ISOP
C10H16			
CH3OH	MEOH		CH3OH
C2H5OH			
CH2O	HCHO	HCHO	HCHO
CH3CHO	CCHO	ALD	ALD2
CH3COOH		ORA2	RCOOH
GLYOXAL		GLY	
GLYALD		ALD	ALD2
CH3OOH		OP1	CH3OOH
C2H5OOH		OP2	ETHOOH
CH3COOOH		PAA	
CH3COCH3	ACET	KET	AONE
HYAC		KET	AONE
CH3COCHO		MGLY	MGLY
ONIT		ONIT	ONIT
MEK	MEK+PRD2	KET	AONE
MVK	MVK		ISOPRD
MACR	METHACRO		ISOPRD
MPAN			
HYDRALD			ISOPRD
BIGALD			OPEN
ISOPNO3			ISOPN
ONITR			ONIT
CRESOL		CSL	CRES

**MOZART-4
description**

L. K. Emmons et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



MOZART-4
description

L. K. Emmons et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[I◀](#)[▶I](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Table 8. Plant functional types in Community Land Model (CLM) landcover files and corresponding MEGAN PFTs.

Index	CLM PFT	MEGAN PFT
1	desert, ice and ocean	–
2	needleleaf evergreen temperate tree	ntr
3	needleleaf evergreen boreal tree	ntr
4	needleleaf deciduous temperate tree	ntr
5	broadleaf evergreen tropical tree	btr
6	broadleaf evergreen temperate tree	btr
7	broadleaf deciduous tropical tree	btr
8	broadleaf deciduous temperate tree	btr
9	broadleaf deciduous boreal tree	btr
10	broadleaf evergreen shrub	shr
11	broadleaf deciduous temperate shrub	shr
12	broadleaf deciduous boreal shrub	shr
13	c3 arctic grass	grs
14	c3 non-arctic grass	grs
15	c4 grass	grs
16	corn	crp
17	wheat	crp

Table 9. Emissions totals by category for 2006 in Tg(species)/year.

Species	Anthro	Fires	Biogenic	Soil	Ocean	Volcano	Total
NO	77.3	10.8	0.0	11.7	0.0	0.0	99.8
CO	642.0	388.9	159.9	0.0	19.9	0.0	1210.7
C ₂ H ₆	7.8	3.1	1.0	0.0	0.0	0.0	11.9
C ₃ H ₈	8.4	0.6	2.0	0.0	0.0	0.0	11.0
C ₂ H ₄	7.3	5.8	5.0	0.0	0.0	0.0	18.1
C ₃ H ₆	3.0	1.8	1.0	0.0	0.0	0.0	5.8
BIGALK	77.6	1.4	0.0	0.0	0.0	0.0	78.9
BIGENE	7.6	1.4	0.0	0.0	0.0	0.0	9.1
TOLUENE	31.5	2.8	0.0	0.0	0.0	0.0	34.3
ISOP	0.0	0.0	555.5	0.0	0.0	0.0	555.5
C ₁₀ H ₁₆	0.0	0.0	75.5	0.0	0.0	0.0	75.5
CH ₃ OH	0.4	7.6	229.6	0.0	0.0	0.0	237.6
C ₂ H ₅ OH	5.4	0.4	0.0	0.0	0.0	0.0	5.8
CH ₂ O	1.1	2.9	0.0	0.0	0.0	0.0	4.0
CH ₃ CHO	2.2	5.4	0.0	0.0	0.0	0.0	7.6
CH ₃ COCH ₃	0.3	2.4	24.4	0.0	0.0	0.0	27.2
MEK	1.4	5.2	0.0	0.0	0.0	0.0	6.6
SO ₂	139.4	2.3	0.0	0.0	0.0	9.6	151.4
DMS	0.0	0.0	0.0	0.0	27.4	0.0	27.4
NH ₃	51.2	5.7	0.0	2.4	8.2	0.0	67.5
CB ₁ +CB ₂	5.0	2.6	0.0	0.0	0.0	0.0	7.6
OC ₁ +OC ₂	16.9	21.3	0.0	0.0	0.0	0.0	38.1

MOZART-4 description

L. K. Emmons et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Table 10. Emissions totals for 2000–2007 in Tg(species)/year.

Species	2000	2001	2002	2003	2004	2005	2006	2007
NO	92.3	94.0	96.5	98.3	98.9	99.8	99.8	101.3
Lght-NO	4.8	5.0	4.6	4.8	4.7	5.3	5.6	5.8
CO	1114.5	1146.8	1211.4	1204.1	1222.1	1225.7	1213.2	1235.1
C2H6	10.9	11.1	11.7	11.4	11.9	12.0	11.9	12.0
C3H8	10.5	10.6	10.8	10.8	10.9	10.9	11.0	11.1
C2H4	16.4	16.9	17.8	17.4	18.2	18.4	18.1	18.4
C3H6	5.3	5.5	5.8	5.8	5.8	5.9	5.8	5.9
BIGALK	75.7	76.2	76.8	77.4	78.0	78.5	79.1	79.7
BIGENE	8.1	8.2	8.7	8.8	8.9	9.0	9.1	9.1
TOLUENE	31.6	32.1	33.0	33.4	33.6	34.0	34.3	34.8
ISOP	450.7	463.7	481.5	472.3	473.3	488.4	469.6	466.9
C10H16	88.1	89.6	90.2	90.0	89.2	91.7	90.7	90.6
CH3OH	236.8	237.2	238.1	237.6	238.1	238.0	237.6	238.0
C2H5OH	5.8	5.8	5.9	5.8	5.8	5.8	5.8	5.8
CH2O	3.2	3.3	4.1	4.1	3.9	3.9	4.0	3.9
CH3CHO	6.9	7.3	8.0	7.8	7.9	7.9	7.7	8.0
CH3COCH3	26.9	27.1	27.4	27.2	27.4	27.4	27.2	27.4
MEK	5.8	6.2	6.9	6.6	6.8	6.8	6.6	6.9
SO2	141.3	143.6	146.9	151.8	151.6	151.5	151.5	151.4
DMS	27.4	27.4	27.4	27.4	27.4	27.4	27.4	27.4
NH3	65.4	65.9	66.6	66.4	67.0	67.5	67.7	68.5
CB1	5.7	5.9	6.1	6.0	6.2	6.2	6.1	6.2
CB2	1.4	1.5	1.5	1.5	1.5	1.5	1.5	1.6
OC1	17.5	18.0	20.1	20.0	19.3	19.2	19.1	19.4
OC2	17.5	18.0	20.1	20.0	19.3	19.2	19.1	19.4
SA1	64.8	65.7	66.0	65.8	65.8	65.5	68.8	82.7
SA2	708.3	718.4	722.0	719.4	719.3	716.2	752.1	904.1
SA3	1667.0	1690.7	1699.2	1693.1	1693.0	1685.8	1770.2	2127.8
SA4	953.3	966.8	971.7	968.2	968.2	964.0	1012.3	1216.8

**MOZART-4
description**

L. K. Emmons et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Table 11. Locations of ozonesondes used in Fig. 7.

Station	Location	Station	Location
90S–30S		30N–90N	
Neumayer	(–70N, –8E)	Alert	(82N, –62E)
Syowa	(–69N, 39E)	Eureka	(79N, –85E)
Marambio	(–64N, –56E)	Ny-Alesund	(78N, 11E)
Macquarie	(–54N, 158E)	Resolute	(74N, –94E)
Broad Meadows	(–37N, 144E)	Lerwick	(60N, –1E)
		Churchill	(58N, –94E)
		Edmonton	(53N, –114E)
		Goose Bay	(53N, –60E)
		Lindenberg	(52N, 14E)
		deBilt	(52N, 5E)
		Uccle	(50N, 4E)
		Praha	(50N, 14E)
		Hohenpeissenberg	(47N, 11E)
		Payerne	(46N, 6E)
		Sapporo	(43N, 141E)
		Trinidad Head	(40N, –124E)
		Madrid	(40N, –3E)
		Wallops	(37N, –75E)
		Tateno	(36N, 140E)
		Huntsville	(34N, –86E)
		Kagoshima	(31N, 130E)
30S–Eq			
Irene	(–25N, 28E)		
Reunion	(–21N, 55E)		
Fiji	(–18N, 178E)		
Samoa	(–14N, –170E)		
Ascension	(–7N, –14E)		
Java	(–7N, 112E)		
Natal	(–5N, –35E)		
Malindi	(–2N, 40E)		
Nairobi	(–1N, 36E)		
San Cristobal	(0N, –89E)		
Eq–30N			
Sepang	(2N, 101E)		
Paramaribo	(5N, –55E)		
Hilo	(19N, –155E)		
Hong Kong	(22N, 114E)		
Naha	(26N, 127E)		
New Delhi	(28N, 77E)		

**MOZART-4
description**

L. K. Emmons et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[I◀](#)[▶I](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

MOZART-4
description

L. K. Emmons et al.




Fig. 1. Emissions of isoprene and monoterpenes for January and July 2006, from online calculation by MEGAN in MOZART-4.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[I◀](#)[▶I](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

**MOZART-4
description**

L. K. Emmons et al.




Fig. 2. Emissions of NO from lightning for 2006. Top panel shows the zonal average vertical distribution for the annual average. Bottom panel shows the total column annual total.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**MOZART-4
description**

L. K. Emmons et al.




Fig. 3. Example profiles of the three ozone variables (O3, O3RAD and SYNOZ) used when including the SYNOZ parameterization. Shown is a monthly average profile for July 2006 at a point over the Pacific Ocean.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[I◀](#)[▶I](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

MOZART-4 description

L. K. Emmons et al.




Fig. 4. OH burden from MOZART-4 compared to the climatology of Spivakovsky et al. (2000).

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



MOZART-4
description

L. K. Emmons et al.




Fig. 5. Comparison of MOZART-4 CO (blue line) to NOAA GMD surface CO measurements (black dots) at selected network sites. MOZART-4 results correspond to the pressure altitude of each observation site. Measurements are monthly averages of approximately weekly samples, while model results are averages of the complete month.

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[◀](#)
[▶](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)


MOZART-4
description

L. K. Emmons et al.




Fig. 6. Evaluation of MOZART-4 CO with MOPITT V4 CO column densities; top: MOZART-4 CO total column average mixing ratio; middle: MOPITT CO column retrieval as average mixing ratio; bottom: difference between MOZART-4 CO, transformed with the MOPITT operator (a priori and averaging kernels), and MOPITT CO, for May 2003.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

MOZART-4
description

L. K. Emmons et al.




Fig. 7. Comparison of MOZART-4 CO (blue solid and dashed lines) with MOPITT V4 CO (red points and error bars) averaged over regions, based on column averages of monthly means. MOZART-4 results have been transformed with the MOPITT averaging kernels and a priori. The error bars (and dashed lines for MOZART-4) indicate the range of data over the region for each month.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



MOZART-4
description

L. K. Emmons et al.




Fig. 8. Evaluation of MOZART-4 with ozonesonde measurements, binned by latitude and altitude, for each year 2000–2007. The number of sites in each region is 90S–30S: 5, 30S–Eq: 10, Eq–30N: 6, 30N–60N: 21. The sites used for each region are given in Table 11. Each dot indicates the average over the month and all sites in the latitude bin, colored by year. The lines indicate MOZART-4 monthly means, averaged over the ozonesonde locations.

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[⏪](#)
[⏩](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)


MOZART-4
description

L. K. Emmons et al.




Fig. 9. Evaluation of MOZART-4 AOD with MODIS AOD retrievals. Ocean-only retrievals from MODIS for five regions are compared to ocean-only grid points from MOZART-4 (latitude-longitude ranges indicated in plot titles). Monthly means are shown, with error bars (dotted lines for MOZART-4) indicating the central 80% of the variation over the region.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)