

Abstract

Chemical data assimilation attempts to optimally use noisy observations along with imperfect model predictions to produce a better estimate of the chemical state of the atmosphere. It is widely accepted that a key ingredient for successful data assimilation is a realistic estimation of the background error distribution. Particularly important is the specification of the background error covariance matrix, which contains information about the magnitude of the background errors and about their correlations. Most models currently use diagonal background covariance matrices. As models evolve toward finer resolutions, the diagonal background covariance matrices become increasingly inaccurate, since they captures less of the spatial error correlations. This paper discusses an efficient computational procedure for constructing non-diagonal background error covariance matrices which account for the spatial correlations of errors. The benefits of using the non-diagonal covariance matrices for variational data assimilation with chemical transport models are illustrated.

1 Introduction

Chemical data assimilation attempts to optimally use noisy observations of reality along with imperfect model predictions to produce a better estimate (in some optimal sense) of the chemical state of the atmosphere. This improved estimate state better defines the spatial and temporal fields of key chemical components in relation to their sources and sinks. This information is critical for improved studies of the atmospheric composition. Chemical data assimilation could also, in principle, improve estimates of emission inventories, of model boundary conditions, or of important model parameters like wet deposition velocities or photolysis rates.

The close integration of observational data is recognized as essential in weather/climate analysis and forecast activities. Consequently, considerable experience with data assimilation have been accumulated in the field of numerical

Non-diagonal background error covariance matrices for data assimilation

K. Singh et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)



[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



**Non-diagonal
background error
covariance matrices
for data assimilation**

K. Singh et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

weather prediction (Daley, 1991; Courtier et al., 1998; Rabier et al., 2000; Kalnay, 2002; Navon, 2009). In this work we focus on chemical data assimilation, i.e., on assimilation of observations of pollutant levels in the atmosphere. Chemical data assimilation poses specific challenges related to the multiphysics nature of the system, the stiffness of chemical kinetic equations, the sparseness of chemical observations, and the uncertainty in the levels of anthropogenic and natural pollutants emitted into the atmosphere.

Previous studies have employed various approaches to assimilating observations of trace gases for improved tropospheric chemistry representations. The base concepts of the variational approach to chemical data assimilation, and the construction of adjoint chemical transport models are discussed in detail in Sandu et al. (2005); Carmichael et al. (2008). Early work in chemical data assimilation using variational techniques has been reported in Khattatov et al. (2000); Elbern and Schmidt (2001). Since then there is a growing body of literature with applications of 4D-Var chemical data assimilation. Adjustment of gas phase chemical tracer initial conditions has been studied in Chai et al. (2007); Zhang et al. (2008). Adjustment of pollutant emissions through 4D-Var chemical data assimilation has been discussed in (Chai et al., 2009). Data assimilation studies involving particle measurements to improve aerosol fields have been performed in Hakami et al. (2005); Henze et al. (2009).

Suboptimal Kalman filters have been employed successfully for chemical data assimilation (Menard et al., 2000; Lamarque et al., 2002; Segers et al., 2005; Clark et al., 2006; Pierce et al., 2007; Parrington et al., 2009). The use of the ensemble Kalman filter (EnKF) in chemical data assimilation has been studied in Constantinescu et al. (2007b,c,d). Data assimilation has been used to improve initial conditions, emissions, and boundary values. Besides the initial conditions, improvements in boundary values lead to improved air quality forecasts. Comparisons of the the performance of different techniques for chemical data assimilation have been performed (Lahoz et al., 2007; Wu et al., 2008).

**Non-diagonal
background error
covariance matrices
for data assimilation**

K. Singh et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

It is widely accepted that a key ingredient of successful data assimilation is a realistic estimation of the background error distribution. Particularly important is the specification of the background error covariance matrix, which contains information about the magnitude of the background errors and about their correlations. Background covariance matrices impact how the information from observations is spread both spatially and among the different types of analysis variables.

The construction of background covariance matrices is challenging due to poorly characterized background errors, and to the very large dimension of the state space of realistic atmospheric models. As a consequence, many chemical data assimilation studies to date have used diagonal background covariance matrices. A popular approach to approximate the background covariance matrix is the NMC method (Parrish and Derber, 1992), in which the differences between several forecasts verifying at the same time are used to approximate the background error. This method has been successfully applied to chemical data assimilation (Chai et al., 2006). An alternative approach constructs autoregressive models of background errors based on the short-term linearized model dynamics (Constantinescu et al., 2007a).

A popular ansatz is that the background error correlations decay exponentially in space. This ansatz allows the construction of simple error correlation models, and is the basis of the covariance localization technique used in ensemble Kalman filtering (Gaspari and Cohn, 1999; Ott et al., 2004; Constantinescu et al., 2007b). Experimental studies with chemical transport models support this assumption; for example, in Chai et al. (2006) it has been shown that ozone error correlations decrease follow, on average, an exponential decay curve. Sub-optimal Kalman filters based on covariance models that impose an exponential decay of correlations with distances have been used in the assimilation of chemical constituents (Khattatov et al., 1999; Pierce et al., 2007).

In the troposphere, ozone is an important greenhouse gas and a major pollutant, which adversely impacts air quality. Its distribution is highly heterogeneous, reflecting the combined influence of atmospheric transport and local chemical sources and sinks. Until recently, observations of the three-dimensional structure of tropospheric

Non-diagonal background error covariance matrices for data assimilation

K. Singh et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[⏪](#)

[⏩](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)

ozone have been limited. The Tropospheric Emission Spectrometer (TES) satellite instrument, launched in 2004, produced the first continuous, global profile retrievals of tropospheric ozone. Similar observations are now available from other satellite instruments, such as the Infrared Atmospheric Sounding Interferometer (IASI).

5 Assimilating these data into atmospheric models provides a powerful means to obtain an improved understanding of the processes controlling tropospheric ozone. Parrington et al. (2008) was the first to assimilate the TES ozone profile retrievals, but they did not account for horizontal correlations in the background error.

We propose here a computationally efficient approach for constructing (background) error covariances that account for spatial correlations in both horizontal and vertical directions, and assess its impact on the assimilation of tropospheric ozone profiles from TES. The construction is based but not restricted to the ansatz of exponential decay of error correlations. The correlation lengths in the latitudinal, longitudinal, and vertical directions can be specified according to the application requirements. Due to the large number of state variables an explicit representation of the full background covariance matrix is impractical. The proposed strategy constructs a multi-dimensional correlation matrix from tensor products of one-dimensional correlation matrices. This avoids the explicit construction and storage of full covariance matrices, and allows the needed linear algebra operations to be performed very efficiently.

20 The paper is organized as follows. Section 2 reviews variational data assimilation techniques, and Sect. 3 presents the GEOS-Chem global chemical transport model. The algorithm for constructing multidimensional covariance matrices is discussed in Sect. 4. Section 5 presents assimilation results of TES ozone profiles with the global chemical transport model GEOS-Chem, and illustrates the benefits of nondiagonal covariances in both three and four dimensional variational data assimilation settings.

2 Variational data assimilation

Variational methods solve the data assimilation problem in an optimal control framework (Sasaki, 1958; Le Dimet and Talagrand, 1986; Courtier and Talagrand, 1987; Lions, 1971). Specifically, they attempt to find the control variable values (e.g., initial conditions) which minimize the discrepancy between the model forecast and observations; the minimization is constrained by the governing dynamic equations. In this discussion, for simplicity of presentation, we focus on discrete models (in time and space) where the initial conditions are the control variables.

Data assimilation combines the following three sources of information.

1. The a priori, or background state \mathbf{x}^b represents the best estimate of the true state \mathbf{x}^t available before any measurements are taken. This estimate is assumed unbiased, and the random background (estimation) errors ε^b are typically assumed to have a normal probability density with a background error covariance matrix \mathbf{B}

$$\varepsilon^b = \mathbf{x}^b - \mathbf{x}^t \in \mathcal{N}(0, \mathbf{B}). \quad (1)$$

2. The model encapsulates our knowledge about physical and chemical laws that govern the evolution of the system. The model evolves an initial state $\mathbf{x}_0 \in \mathbb{R}^n$ at the initial time t_0 to future state values $\mathbf{x}_i \in \mathbb{R}^n$ at future times t_i ,

$$\mathbf{x}_i = \mathcal{M}_{t_0 \rightarrow t_i}(\mathbf{x}_0). \quad (2)$$

The size of the state space in realistic chemical transport models is very large. For example, a global chemical transport model with $2^\circ \times 2.5^\circ$ horizontal resolution has $n \sim 10^8$ variables.

3. Observations $\mathbf{x}_i^{\text{obs}} \in \mathbb{R}^m$ of the state are taken at times t_i , $1 = 1, \dots, N$

$$\mathbf{x}_i^{\text{obs}} = \mathcal{H}(\mathbf{x}_i) + \varepsilon_i^{\text{obs}}. \quad (3)$$

Non-diagonal background error covariance matrices for data assimilation

K. Singh et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Non-diagonal background error covariance matrices for data assimilation

K. Singh et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[⏪](#)

[⏩](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



The observation operator \mathcal{H} maps the state space onto the observation space. In many practical situations \mathcal{H} is a highly nonlinear mapping (e.g., with satellite observation operators). Many times the observations are sparsely distributed, and the number of observations is small compared to the dimension of the state space, $m \ll n$.

The observations are corrupted by measurement and representativeness errors $\varepsilon_i^{\text{obs}}$. The observation errors at each time are assumed to be independent of background errors, and independent of the observation errors at other times. They are typically assumed to have a normal distribution with mean zero and covariance \mathbb{R}_i ,

$$\varepsilon_i^{\text{obs}} \in \mathcal{N}(0, \mathbb{R}_i). \quad (4)$$

Based on these three sources of information data assimilation computes the posterior estimate \mathbf{x}^a of the true state; \mathbf{x}^a is called the “analysis”.

2.1 Three dimensional variational (3D-Var) data assimilation

In the 3D-Var data assimilation the observations (3) are considered successively at times t_1, \dots, t_N . The background state (i.e., the best state estimate at time t_i) is given by the model forecast, starting from the previous analysis (i.e., best estimate at time t_{i-1}):

$$\mathbf{x}_i^b = \mathcal{M}_{t_{i-1} \rightarrow t_i}(\mathbf{x}_{i-1}^a).$$

The discrepancy between the model state \mathbf{x}_i and observations at time t_i , together with the departure of the state from the model forecast \mathbf{x}_i^b , are measured by the 3D-Var cost function:

$$\mathcal{J}(\mathbf{x}_i) = \frac{1}{2} (\mathbf{x}_i - \mathbf{x}_i^b)^T \mathbb{B}^{-1} (\mathbf{x}_i - \mathbf{x}_i^b) + \frac{1}{2} (\mathcal{H}(\mathbf{x}_i) - \mathbf{x}_i^{\text{obs}})^T \mathbb{R}_i^{-1} (\mathcal{H}(\mathbf{x}_i) - \mathbf{x}_i^{\text{obs}}) \quad (5)$$

measured by the 4D-var cost function:

$$\mathcal{J}(\mathbf{x}_0) = \frac{1}{2} (\mathbf{x}_0 - \mathbf{x}_0^b)^T \mathbb{B}^{-1} (\mathbf{x}_0 - \mathbf{x}_0^b) + \frac{1}{2} \sum_{i=1}^N (\mathcal{H}(\mathbf{x}_i) - \mathbf{x}_i^{\text{obs}})^T \mathbb{R}_i^{-1} (\mathcal{H}(\mathbf{x}_i) - \mathbf{x}_i^{\text{obs}}) \quad (9)$$

Note that the departure of the initial conditions from the background is weighted by the inverse background covariance matrix, \mathbb{B}^{-1} , while the differences between the model predictions $\mathcal{H}(\mathbf{x}_i)$ and observations $\mathbf{x}_i^{\text{obs}}$ are weighted by the inverse observation error covariances, \mathbb{R}_i^{-1} .

The 4D-Var analysis is computed as the initial condition which minimizes (9) subject to the model equation constraints (2)

$$\mathbf{x}_0^a = \text{argmin } \mathcal{J}(\mathbf{x}_0) \quad \text{subject to (2)}. \quad (10)$$

- 5 The model (2) propagates the optimal initial condition (9) forward in time to provide the analysis at future times, $\mathbf{x}_i^a = \mathcal{M}_{t_0 \rightarrow t_i}(\mathbf{x}_0^a)$.

The optimization problem (10) is solved numerically using a gradient-based technique. The gradient of (9) reads

$$\nabla \mathcal{J}(\mathbf{x}_0) = \mathbb{B}^{-1} (\mathbf{x}_0 - \mathbf{x}_0^b) + \sum_{i=1}^N \left(\frac{\partial \mathbf{x}_i}{\partial \mathbf{x}_0} \right)^T (\mathcal{H}'(\mathbf{x}_i))^T \mathbb{R}_i^{-1} (\mathcal{H}(\mathbf{x}_i) - \mathbf{x}_i^{\text{obs}}) \quad (11)$$

The 4D-Var gradient requires not only the linearized observation operator $\mathcal{H}' = d\mathcal{H}/d\mathbf{x}$, but also the transposed derivative of future states with respect to the initial conditions. The 4D-Var gradient can be obtained effectively by forcing the adjoint model with observation increments, and running it backwards in time. The construction of an

10 adjoint model requires considerable effort, time, and know-how.

Non-diagonal background error covariance matrices for data assimilation

K. Singh et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



3 GEOS-Chem

In this paper we specifically consider GEOS-Chem (<http://acmg.seas.harvard.edu/geos>), a global three-dimensional chemical transport model (CTM) driven by assimilated meteorological fields from Goddard Earth Observing System (GEOS-4) at the NASA Global Modeling and Assimilation Office (GMAO). The model was first described in (Bey et al., 2001). GEOS-Chem accounts in detail for emissions from both natural and anthropogenic sources, for gas phase chemistry, aerosol processes, long range transport of pollutants, troposphere-stratosphere exchanges, etc. GEOS-Chem is being widely used by research groups world-wide for performing global atmospheric chemistry studies.

The GEOS-Chem Adjoint system (http://wiki.seas.harvard.edu/geos-chem/index.php/GEOS-Chem_Adjoint) has been developed through a joint effort of groups at Virginia Tech, University of Colorado, Caltech, Jet Propulsion Laboratory, and Harvard (Henze et al., 2007; Singh et al., 2009a,b; Eller et al., 2009). The system can perform adjoint sensitivity analyses and 4D-Var chemical data assimilation. Inverse modeling studies with GEOS-Chem adjoint are presented in (Kopacz et al., 2009; Henze et al., 2009).

4 Construction of the background error covariance matrix

A correct characterization of the background errors is necessary for obtaining a meaningful analysis, i.e., for the success of the data assimilation procedure. Under the usual assumption that the background errors are normally distributed their probability density is described by the background state \mathbf{x}^b and the background error covariance matrix \mathbf{B} . In variational data assimilation both \mathbf{x}^b and \mathbf{B} enter directly into the formulation of the cost function; errors in their specification directly impact the analysis result (Daescu, 2008).

GMDD

3, 1783–1827, 2010

Non-diagonal background error covariance matrices for data assimilation

K. Singh et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[⏪](#)

[⏩](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)

Non-diagonal background error covariance matrices for data assimilation

K. Singh et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[⏪](#)

[⏩](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



A non-diagonal background error covariance matrix allows the information from local observations to spread out in space to contribute to corrections of state variables in neighboring locations; similarly, it allows observations of certain components of the state vector to contribute to corrections of other components. This spread of information results in a smooth analysis state, and allows different sets of observations to complement each other.

Despite these advantages, most chemical data assimilation studies to date have employed diagonal background covariances. Little work has been devoted to date to modeling off-diagonal terms (Chai et al., 2006; Constantinescu et al., 2007a). This is due to a number of practical difficulties that arise in the construction of background covariance matrices. The large number of state variables make the construction and storage of full covariance matrices impractical. Ensembles can be employed to estimate error correlations; however, the number of ensemble members is necessarily very small and only low rank approximations of the covariance matrix can be obtained. Localization is often employed to remove spurious correlations and to improve the rank of the resulting matrix (Gaspari and Cohn, 1999).

We next discuss the proposed approach to constructing a background error covariance matrix \mathbf{B} that accounts for both vertical and horizontal correlations without explicitly constructing the full covariance matrix. We explain the construction of the matrix in the two-dimensional case, i.e., for capturing horizontal correlations; the extensions to correlations in three dimensions and to correlations among multiple state variables are immediate. Our target application is global chemical data assimilation using GEOS-Chem.

Consider a uniform latitude-longitude grid and denote by x the longitude and by y the latitude level. A grid point (x_i, y_j) has longitude coordinate x_i , $i = 1, \dots, n_x$, and latitude coordinate y_j , $j = 1, \dots, n_y$. The state vector contains the state values at all gridpoints ordered latitude-first:

$$\text{grid order} = \left\{ (x_1, y_1), \dots, (x_1, y_{n_y}), (x_2, y_1), \dots, (x_2, y_{n_y}), (x_{n_x}, y_1), \dots, (x_{n_x}, y_{n_y}) \right\}. \quad (12)$$

4.1 Directional error correlation matrices

The one-dimensional correlation between errors at two locations (x_i, y_k) and (x_j, y_k) situated at the same latitude y_k is modeled as

$$\left(\tilde{\mathbf{C}}_x^k\right)_{i,j} = \text{corr}\left((x_i, y_k), (x_j, y_k)\right) = e^{-\frac{\text{dist}\left((x_i, y_k), (x_j, y_k)\right)^2}{\ell_x^2}}; \quad i, j = 1, \dots, n_x; \quad k = 1, \dots, n_y; \quad (13)$$

where ℓ_x is the correlation distance in the longitude direction. For a uniform lat-lon grid the distance between x_i and x_j depends only on $\min(|i - j|, n_x - |i - j|)$. This distance also depends on the y_k ; for this reason Eq. (13) defines a different longitudinal correlation matrix $\tilde{\mathbf{C}}_x^k \in \mathbb{R}^{n_x \times n_x}$ for each latitude y_k . Due to the periodicity along each latitude circle the point x_1 is strongly correlated with both x_2 and x_{n_x} , etc. The periodicity is captured by the distance function in (13).

Similarly, the one-dimensional correlation between errors at two locations (x_k, y_i) and (x_k, y_j) situated at the same longitude x_k is modeled as

$$\left(\tilde{\mathbf{C}}_y^k\right)_{i,j} = \text{corr}\left((x_k, y_i), (x_k, y_j)\right) = e^{-\frac{\text{dist}\left((x_k, y_i), (x_k, y_j)\right)^2}{\ell_y^2}}, \quad i, j = 1, \dots, n_y; \quad k = 1, \dots, n_x; \quad (14)$$

where ℓ_y is the correlation distance in the latitude direction. Equation (14) defines a single latitudinal correlation matrix $\tilde{\mathbf{C}}_y \in \mathbb{R}^{n_y \times n_y}$. For a uniform lat-lon grid this correlation matrix is the same for each longitude x_k ; consequently the superscript k is dropped. To simplify the construction the correlations due to the periodicity along a meridional circle are ignored. Otherwise, error correlations across the poles would lead to correlations between errors at all longitudes; such cross-correlations are not captured by (14).

Non-diagonal background error covariance matrices for data assimilation

K. Singh et al.

The cost function and gradient calculations described in Eqs. (5), (7), (9), and (11) require the inverse of the background error covariance matrix; this involves the inverses of the correlation matrices in longitudinal and latitudinal directions. The construction of the directional correlation matrices $\tilde{\mathbf{C}}_x$ and $\tilde{\mathbf{C}}_y$ does not guarantee that they are non-singular. To avoid a possible singularity we take a convex combination between the identity matrix and tensor product correlations as follows:

$$\mathbf{C}_x^k = \theta_x \mathbb{I}_{n_x \times n_x} + (1 - \theta_x) \tilde{\mathbf{C}}_x^k, \tag{15}$$

and

$$\mathbf{C}_y^k = \theta_y \mathbb{I}_{n_y \times n_y} + (1 - \theta_y) \tilde{\mathbf{C}}_y^k. \tag{16}$$

The above procedure brings a shift in the spectrum and ensures the positive definiteness of \mathbf{C}_x^k and \mathbf{C}_y^k . In all our experiments both θ_x and θ_y are chosen to be equal to 0.2.

The longitudinal correlation matrix between all the points on the two-dimensional grid (12) can be constructed from the one-dimensional longitudinal correlation matrices as follows

$$\mathbf{C}_x^{1:n_y} \otimes \mathbb{I}_{n_y \times n_y} = \begin{pmatrix} \begin{pmatrix} (\mathbf{C}_x^1)_{1,1} & 0 & \dots & 0 \\ 0 & (\mathbf{C}_x^2)_{1,1} & \dots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \dots & (\mathbf{C}_x^{n_y})_{1,1} \end{pmatrix} & \dots & \begin{pmatrix} (\mathbf{C}_x^1)_{1,n_x} & 0 & \dots & 0 \\ 0 & (\mathbf{C}_x^2)_{1,n_x} & \dots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \dots & (\mathbf{C}_x^{n_y})_{1,n_x} \end{pmatrix} \\ \vdots & \ddots & \vdots & \vdots \\ \begin{pmatrix} (\mathbf{C}_x^1)_{n_x,1} & 0 & \dots & 0 \\ 0 & (\mathbf{C}_x^2)_{n_x,1} & \dots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \dots & (\mathbf{C}_x^{n_y})_{n_x,1} \end{pmatrix} & \dots & \begin{pmatrix} (\mathbf{C}_x^1)_{n_x,n_x} & 0 & \dots & 0 \\ 0 & (\mathbf{C}_x^2)_{n_x,n_x} & \dots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \dots & (\mathbf{C}_x^{n_y})_{n_x,n_x} \end{pmatrix} \end{pmatrix}$$

Title Page

Abstract Introduction

Conclusions References

Tables Figures

⏪ ⏩

◀ ▶

Back Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



With some abuse of notation we extend the use of the Kronecker product symbol \otimes in the above equation in order to highlight the structure of the two-dimensional longitudinal correlation matrix.

Similarly, the latitudinal correlation matrix between all the points on the two-dimensional grid (12) can be constructed from the one-dimensional latitudinal correlation matrices as follows

$$\|_{n_x \times n_x} \otimes \mathbf{C}_y = \begin{pmatrix} \begin{pmatrix} (\mathbf{C}_y)_{1,1} & (\mathbf{C}_y)_{1,2} & \dots & (\mathbf{C}_y)_{1,n_y} \\ (\mathbf{C}_y)_{2,1} & (\mathbf{C}_y)_{2,2} & \dots & (\mathbf{C}_y)_{2,n_y} \\ \vdots & \vdots & \ddots & \vdots \\ (\mathbf{C}_y)_{n_y,1} & (\mathbf{C}_y)_{n_y,2} & \dots & (\mathbf{C}_y)_{n_y,n_y} \end{pmatrix} & \dots & \begin{pmatrix} 0 & 0 & \dots & 0 \\ 0 & 0 & \dots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \dots & 0 \end{pmatrix} \\ \vdots & \ddots & \vdots \\ \begin{pmatrix} 0 & 0 & \dots & 0 \\ 0 & 0 & \dots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \dots & 0 \end{pmatrix} & \dots & \begin{pmatrix} (\mathbf{C}_y)_{1,1} & (\mathbf{C}_y)_{1,2} & \dots & (\mathbf{C}_y)_{1,n_y} \\ (\mathbf{C}_y)_{2,1} & (\mathbf{C}_y)_{2,2} & \dots & (\mathbf{C}_y)_{2,n_y} \\ \vdots & \vdots & \ddots & \vdots \\ (\mathbf{C}_y)_{n_y,1} & (\mathbf{C}_y)_{n_y,2} & \dots & (\mathbf{C}_y)_{n_y,n_y} \end{pmatrix} \end{pmatrix}$$

The structure of the one-dimensional correlation matrices is represented in Fig. 1. The longitudinal correlation \mathbf{C}_x^k is represented at latitude $y_k = 20^\circ \text{N}$; note that due to the periodicity along each latitude circle not only the elements near the diagonal, but also the elements in the corners of the matrix have non-zero values. The latitudinal correlation \mathbf{C}_y does not account for periodicity (along each meridian the grids 1 and n_y correspond to the South and to the North pole, respectively).

Figure 2 represents contour lines of the longitudinal correlation \mathbf{C}_x for points at different latitudes. The correlation length ℓ_x is short (top panel), medium (middle panel), and large (bottom panel). Note that the same correlation length ℓ_x translates into a larger number of correlated grid points at higher latitudes.

Non-diagonal background error covariance matrices for data assimilation

K. Singh et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



4.2 Two-dimensional covariance matrices

Formally the full background error correlation matrix $\mathbf{C} \in \mathbb{R}^{n_x n_y \times n_x n_y}$ (which accounts for both latitudinal and longitudinal correlations) is constructed via the following relation

$$\mathbf{C} = \left(\mathbf{C}_x^{1:n_y} \otimes \mathbb{1}_{n_y \times n_y} \right) \cdot \left(\mathbb{1}_{n_x \times n_x} \otimes \mathbf{C}_y \right). \quad (17)$$

Note that this (huge) matrix is never explicitly formed. One needs to form and store only n_y one-dimensional longitudinal correlation matrices (13) and a single one-dimensional latitudinal correlation matrix (14). Note that the diagonal entries of the tensor product matrix (17) are all equal to one. The tensor product matrix (17) is not symmetric; with some (more) abuse of notation we will continue to call it a correlation matrix.

A symmetric version of the two-dimensional correlation matrix can be constructed as follows. Any symmetric positive definite matrix \mathbf{C} has a matrix square root $\mathbf{C}^{1/2}$ such that

$$\mathbf{C} = \mathbf{C}^{1/2} \mathbf{C}^{T/2}.$$

The matrix square root is not uniquely defined; in particular it can be symmetric or not depending on the decomposition method used as described in Eqs. (21) and (22). Let $\left(\mathbb{1}_{n_x \times n_x} \otimes \mathbf{C}_y \right)^{1/2}$ be a square root of the longitudinal correlation matrix. The symmetric two-dimensional correlation matrix can be constructed as:

$$\mathbf{C}^{\text{sym}} = \left(\mathbb{1}_{n_x \times n_x} \otimes \mathbf{C}_y \right)^{1/2} \cdot \left(\mathbf{C}_x^{1:n_y} \otimes \mathbb{1}_{n_y \cdot n_y} \right) \cdot \left(\mathbb{1}_{n_x \times n_x} \otimes \mathbf{C}_y \right)^{T/2}. \quad (18)$$

Let $\sigma_{i,j}$ be the standard deviation of the error at location (x_i, y_j) and

$$\mathbf{\Sigma} = \text{diag}_{1 \leq i \leq n_x, 1 \leq j \leq n_y} \sigma_{i,j}$$

the diagonal matrix with all standard deviations at all grid points ordered according to

Non-diagonal background error covariance matrices for data assimilation

K. Singh et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[⏪](#)

[⏩](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



(12). The two-dimensional covariance matrix is constructed from the correlation matrix (17) by scaling it from left and right with the diagonal matrix of standard deviations Σ

$$\mathbb{B} = \Sigma \cdot \mathbf{C} \cdot \Sigma = \Sigma \cdot \left(\mathbf{C}_x^{1:n_y} \otimes \mathbb{1}_{n_y \times n_y} \right) \cdot \left(\mathbb{1}_{n_x \times n_x} \otimes \mathbf{C}_y \right) \cdot \Sigma. \quad (19)$$

With some abuse of the nomenclature we call the non-symmetric \mathbb{B} in (19) a covariance matrix.

Similarly, a symmetric version of the covariance matrix can be constructed from the symmetric correlation (18) as follows:

$$\mathbb{B}^{\text{sym}} = \Sigma \cdot \mathbf{C}^{\text{sym}} \cdot \Sigma = \Sigma \cdot \left(\mathbb{1}_{n_x \times n_x} \otimes \mathbf{C}_y \right)^{1/2} \left(\mathbf{C}_x^{1:n_y} \otimes \mathbb{1}_{n_y \times n_y} \right) \cdot \left(\mathbb{1}_{n_x \times n_x} \otimes \mathbf{C}_y \right)^{T/2} \cdot \Sigma. \quad (20)$$

4.3 Efficient covariance matrix function calculations

The symmetric positive definite one-directional longitudinal correlation matrix has a matrix square root $\mathbf{C}_y^{1/2}$. A symmetric square root, the inverse of the symmetric square root, and the matrix inverse can be obtained via the singular value decomposition (SVD)

$$\mathbf{C}_y = \mathbf{U} \Theta \mathbf{U}^T, \quad (\mathbf{C}_y)^r = \mathbf{U} \Theta^r \mathbf{U}^T, \quad \text{for } r \in \left\{ \frac{1}{2}, -\frac{1}{2}, -1 \right\}. \quad (21)$$

while a nonsymmetric square root can be obtained via a Cholesky decomposition

$$\mathbf{C}_y = \mathbb{L}_y \mathbb{L}_y^T, \quad \mathbf{C}_y^{1/2} = \mathbb{L}_y. \quad (22)$$

By the properties of the Kronecker product we have that the square root, the inverse square root, and the inverse of the two-dimensional longitudinal correlation matrix can be constructed in terms of the same matrix functions applied to the one dimensional longitudinal correlations:

$$\left(\mathbb{1}_{n_x \times n_x} \otimes \mathbf{C}_y \right)^r = \mathbb{1}_{n_x \times n_x} \otimes (\mathbf{C}_y)^r \quad \text{for } r \in \left\{ \frac{1}{2}, -\frac{1}{2}, -1 \right\}. \quad (23)$$

Consequently, the symmetric covariance (20) can be implemented as

$$\mathbb{B}^{\text{sym}} = \Sigma \cdot \left(\mathbb{I}_{n_x \times n_x} \otimes (\mathbf{C}_y)^{1/2} \right) \cdot \left(\mathbf{C}_x^{1:n_y} \otimes \mathbb{I}_{n_y \times n_y} \right) \cdot \left(\mathbb{I}_{n_x \times n_x} \otimes (\mathbf{C}_y)^{T/2} \right) \cdot \Sigma$$

using either of the one-dimensional square roots.

Similarly, different powers of each one-dimensional latitudinal correlation matrix can be obtained via a singular value decomposition:

$$\mathbf{C}_x^{1:n_y} = \mathbb{V} \Gamma \mathbb{V}^T, \quad \left(\mathbf{C}_x^{1:n_y} \right)^r = \mathbb{V} \Gamma^r \mathbb{V}^T, \quad \text{for } r \in \left\{ \frac{1}{2}, -\frac{1}{2}, -1 \right\}.$$

By the properties of the extended Kronecker product we have that the square root, the inverse square root, and the inverse of the two-dimensional latitudinal correlation matrix can be constructed in terms of the same matrix functions applied to the one dimensional latitudinal correlations:

$$\left(\mathbf{C}_x^{1:n_y} \otimes \mathbb{I}_{n_y \times n_y} \right)^r = \left(\mathbf{C}_x^{1:n_y} \right)^r \otimes \mathbb{I}_{n_y \times n_y} \quad \text{for } r \in \left\{ \frac{1}{2}, -\frac{1}{2}, -1 \right\}. \quad (24)$$

We now use these relations to build functions of the covariance matrices. The inverse of the background covariance is needed in the formulation of the variational cost function. The inverse of the non-symmetric covariance (19) is

$$\mathbb{B}^{-1} = \Sigma^{-1} \cdot \left(\mathbb{I}_{n_x \times n_x} \otimes (\mathbf{C}_y)^{-1} \right) \cdot \left(\left(\mathbf{C}_x^{1:n_y} \right)^{-1} \otimes \mathbb{I}_{n_y \times n_y} \right) \cdot \Sigma^{-1}. \quad (25)$$

The inverse of the symmetric covariance (20) matrix is

$$\left(\mathbb{B}^{\text{sym}} \right)^{-1} = \Sigma^{-1} \cdot \left(\mathbb{I}_{n_x \times n_x} \otimes (\mathbf{C}_y)^{-T/2} \right) \left(\left(\mathbf{C}_x^{1:n_y} \right)^{-1} \otimes \mathbb{I}_{n_y \times n_y} \right) \cdot \left(\mathbb{I}_{n_x \times n_x} \otimes (\mathbf{C}_y)^{-1/2} \right) \cdot \Sigma^{-1}. \quad (26)$$

Non-diagonal background error covariance matrices for data assimilation

K. Singh et al.

Title Page	
Abstract	Introduction
Conclusions	References
Tables	Figures
⏪	⏩
◀	▶
Back	Close
Full Screen / Esc	
Printer-friendly Version	
Interactive Discussion	



Non-diagonal background error covariance matrices for data assimilation

K. Singh et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Finally, the symmetric covariance (20) has a (non-symmetric) matrix square root

$$(\mathbb{B}^{\text{sym}})^{1/2} = \Sigma \cdot \left(\mathbb{I}_{n_x \times n_x} \otimes (\mathbf{C}_y)^{1/2} \right) \cdot \left(\left(\mathbf{C}_x^{1:n_y} \right)^{1/2} \otimes \mathbb{I}_{n_y \times n_y} \right). \quad (27)$$

This is built out of tensor products involving the square roots of the one-dimensional correlation matrices. The inverse of the square root matrix (27) is

$$(\mathbb{B}^{\text{sym}})^{-1/2} = \left(\left(\mathbf{C}_x^{1:n_y} \right)^{-1/2} \otimes \mathbb{I}_{n_y \times n_y} \right) \cdot \left(\mathbb{I}_{n_x \times n_x} \otimes (\mathbf{C}_y)^{-1/2} \right) \cdot \Sigma^{-1}. \quad (28)$$

4.4 Efficient linear algebra operations involving the covariance matrix

The computations associated with data assimilation involve linear algebra operations with the covariance \mathbb{B} . Matrix vector operations can be performed effectively by exploiting structure of \mathbb{B} . Consider a vector of concentrations (or concentration errors) $u_{i,j}$ - indexed by latitude and longitude, but stored as a state vector with the convention (12).

Consider the non-symmetric covariance matrix. The covariance matrix-vector product $v = \mathbb{B} \cdot u$ can be computed in stages. Each stage produces a temporary result which is a two-dimensional vector.

Expression	Computation	
$\alpha = \Sigma \cdot u$	$\alpha_{i,j} = \sigma_{i,j} u_{i,j}$,	for $i = 1, \dots, n_x, j = 1, \dots, n_y$.
$\beta = \left(\mathbb{I}_{n_x \times n_x} \otimes \mathbf{C}_y \right) \cdot \alpha$	$\beta_{i,1:n_y} = \mathbf{C}_y \cdot \alpha_{i,1:n_y}$,	for $i = 1, \dots, n_x$.
$\gamma = \left(\mathbf{C}_x^{1:n_y} \otimes \mathbb{I}_{n_y \times n_y} \right) \cdot \beta$	$\gamma_{1:n_x,j} = \mathbf{C}_x^j \cdot \beta_{1:n_x,j}$,	for $j = 1, \dots, n_y$.
$v = \Sigma \cdot \gamma$	$v_{i,j} = \sigma_{i,j} \gamma_{i,j}$,	for $i = 1, \dots, n_x, j = 1, \dots, n_y$.

Similarly the inverse covariance matrix-vector product $v = \mathbb{B}^{-1} \cdot u$ can be computed as follows

Expression	Computation
$\alpha = \Sigma^{-1} \cdot u$	$\alpha_{i,j} = u_{i,j} / \sigma_{i,j},$ for $i = 1, \dots, n_x, j = 1, \dots, n_y.$
$\beta = \left(\mathbb{C}_x^{1:n_y} \otimes \mathbb{I}_{n_y \times n_y} \right)^{-1} \cdot \alpha$	Solve $\mathbb{C}_x^j \cdot \beta_{1:n_x,j} = \alpha_{1:n_x,j},$ for $j = 1, \dots, n_y.$
$\gamma = \left(\mathbb{I}_{n_x \times n_x} \otimes \mathbb{C}_y \right)^{-1} \cdot \beta$	Solve $\mathbb{C}_y \cdot \gamma_{i,1:n_y} = \beta_{i,1:n_y},$ for $i = 1, \dots, n_x.$
$v = \Sigma^{-1} \cdot \gamma$	$v_{i,j} = \gamma_{i,j} / \sigma_{i,j},$ for $i = 1, \dots, n_x, j = 1, \dots, n_y.$

(30)

Similar procedures can be developed for the symmetric covariance time vector products.

The square root (27) times vector product $v = (\mathbb{B}^{\text{sym}})^{1/2} u$ is computed as:

Expression	Computation
$\alpha = \left(\left(\mathbb{C}_x^{1:n_y} \right)^{1/2} \otimes \mathbb{I}_{n_y \times n_y} \right) \cdot u$	$\gamma_{1:n_x,j} = \mathbb{C}_x^j \cdot \beta_{1:n_x,j},$ for $j = 1, \dots, n_y$
$\beta = \left(\mathbb{I}_{n_x \times n_x} \otimes \left(\mathbb{C}_y \right)^{1/2} \right) \cdot \alpha$	$\beta_{i,1:n_y} = \mathbb{C}_y^{1/2} \cdot \alpha_{i,1:n_y},$ for $i = 1, \dots, n_x$
$v = \Sigma \cdot \beta$	$v_{i,j} = \sigma_{i,j} \gamma_{i,j},$ for $i = 1, \dots, n_x, j = 1, \dots, n_y.$

(31)

All the above implementations are based on repeated operations involving the one-dimensional covariance matrices and their square roots. These operations are very efficient since all the linear algebra operations (matrix-vector multiplication, SVD, Cholesky factorization, the solution of linear systems) are performed on small dimensional matrices ($n_x \times n_x$ or $n_y \times n_y$).

Non-diagonal background error covariance matrices for data assimilation

K. Singh et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



5 Numerical experiments

For numerical experiments, we employ GEOS-Chem v7-04-10 adjoint code (Singh et al., 2009b), capable of performing both 3D-Var and 4D-Var data assimilations with real data. We assimilate Tropospheric Emission Spectrometer (TES) satellite ozone profile retrievals into the model and validate the generated analyses through an independent observation dataset provided by direct ozone profile measurements from Ozonesondes. The numerical optimization method used in all experiments is the limited memory bound-constrained BFGS (Zhu et al., 1997). This quasi-Newton approach has become the “gold standard” in solving large scale chemical data assimilation problems (Sandu et al., 2005).

5.1 Experimental setting

Simulations with GEOS-Chem v7 adjoint can be carried out at $4^\circ \times 5^\circ$ and $2^\circ \times 2.5^\circ$ resolutions. We have used $4^\circ \times 5^\circ$ resolution in all our experiments. There are 46×72 latitude-longitude grid boxes at this resolution, and 55 vertical levels; near the equator and at ground level each grid box covers an area of about $400 \text{ km} \times 500 \text{ km}$. The current GEOS-Chem model does not capture well the dynamics of the upper troposphere and of the stratosphere. Therefore, we performed data assimilation for only the first 23 model levels (for up to about 50 hPa). The model has been modified to use the linearized ozone (linoz) scheme for a better estimation of ozone exchanges at troposphere-stratosphere boundary. This scheme has been incorporated in the standard GEOS-Chem model (starting with release v8-02-03).

The 3D-Var data assimilation experiments were performed over the months of July and August 2006, starting at 00:00(GMT) on 1 July. The TES satellite data was read once every 4 h of simulation; the observation operator called at model time t (h) reads in all the measurements collected within the interval $t - 2$ (h) to $t + 2$ (h). 3D-Var data assimilation treats all observations in this interval as instantaneous, and assimilates them in the same optimization run. In all our 3D-Var experiments, we performed

Non-diagonal background error covariance matrices for data assimilation

K. Singh et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[⏪](#)

[⏩](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



8 iterations per analysis since the cost function decreased significantly within the first few iterations. It is important to note that 3D-Var does not involve any model adjoint calculations; gradients require only the adjoint of the observation operator. The optimization adjusts ozone concentrations.

The 4D-Var data assimilation experiments were performed over a 5 day assimilation window starting at 00:00(GMT) on 1 August 2006 and ending at 00:00(GMT) on 6 August of the same year. The background initial conditions were generated through a free GEOS-Chem model run. There were 12 optimization iterations performed in order to improve the ozone initial condition. Each iteration during 4D-Var assimilation includes a forward model and a backward model adjoint run. TES satellite profile retrievals are read every 4 h during the model adjoint run, and the cost function and adjoint gradients accumulate the impact of all 4 h data sets throughout the assimilation window.

NOTE: the TES data can be biased by as much as 10% (Nassar et al., 2008). We removed this bias as estimated by (Nassar et al., 2008) before assimilating the data.

5.1.1 Computational costs

As described in Sect. 1, the construction of the background error covariance matrix \mathbf{B} impacts the result of the data assimilation. If one considers no correlation among different model grid points, or among different chemical species, \mathbf{B} turns out to be diagonal. However, this approximation is inaccurate as the ozone errors are highly correlated spatially (Constantinescu et al., 2007a,c,d) and correlated to errors in other chemical species; this correlation is not discussed in this paper. In Sect. 4, we have introduced an efficient methodology to construct a non-diagonal background error covariance matrix, \mathbf{B} . Its inverse, \mathbf{B}^{-1} , needed in 3D-Var (5) and in 4D-Var (9) cost function formulations, can be obtained either via a Cholesky decomposition or via a singular value decomposition. (Note that by the “computation of the inverse” we mean the solution of a linear system).

Non-diagonal background error covariance matrices for data assimilation

K. Singh et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)



[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)

Non-diagonal background error covariance matrices for data assimilation

K. Singh et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Table 1 illustrates the computational cost of data assimilation compared to the cost of free running model for a 24 h simulation. All the simulations are performed on a Dell Precision T5400 workstation with 2 quadcore Intel(R) Xeon(R) processors with clock speed 2.33 GHz and a RAM of 16 GB shared between the two processors. Performing 3D-Var with a non-diagonal error covariance matrix whose inverse is computed by Cholesky decomposition is only about 1.4% more expensive than the 3D-Var with a diagonal covariance matrix. The calculation of the inverse via the Cholesky decomposition is considerably more efficient than the calculation via a singular value decomposition, as expected.

3D-Var framework is built on top of GEOS-Chem v7 package which uses Sparse Matrix Vectorized GEAR (SMVGEAR) solver for chemistry. However, to construct the adjoint of chemistry required by the 4D-Var, we implemented KPP solver (Damian et al., 2002) into GEOS-Chem which not only provides a suite of high performance chemical solvers to choose from but also generates automatically the continuous and discrete adjoint codes (Daescu, 2000, 2003; Sandu et al., 2003). A detailed discussion on interfacing KPP with GEOS-Chem and comparison with native SMVGEAR solver for accuracy and computational performance is presented in (Eller et al., 2009). As pointed out in (Henze et al., 2007), the computational cost of Rosenbrock solver increases significantly with the tolerance levels; higher tolerances use smaller internal time steps requiring more computation. In our experiments, we have set $RTOL = 10^{-3}$ and $ATOL = 10^{-2}$ to achieve moderate to high accuracy.

The 4D-Var assimilation is considerably more expensive than the 3D-Var. The use of the non-diagonal IB (with Cholesky decomposition) in 4D-Var causes a minimal to zero increase in the computational time when compared to the diagonal IB case.

5.1.2 Tropospheric Emission Spectrometer (TES) observations

The Tropospheric Emission Spectrometer (TES, <http://tes.jpl.nasa.gov>), aboard the NASA Aura spacecraft launched in 2004, measures spectrally resolved outgoing thermal radiation from 5 to 15 microns. These spectra are used to infer the vertical

distribution of atmospheric trace gases through a non-linear optimal estimation procedure Bowman et al. (2006). In this work we assimilate the retrieved ozone vertical profiles. Figure 3 shows the location of TES profiles for two days.

A-priori information about the vertical concentration profile of the species of interest is needed to solve the retrieval inverse problem (the prior information does not come from the measurement). Let $\mathbf{x}^{\text{prior}}$ be the prior vertical ozone concentration profile (in volume mixing ratio units), and let $\mathbf{z}^{\text{prior}} = \log \mathbf{x}^{\text{prior}}$. Let $\mathbf{z}^{\text{radiance}} (= \log \mathbf{x}^{\text{true}})$ be the atmospheric profile as resulting directly from the radiances.

The vertical ozone profile retrieval can be expressed according to the formula

$$\hat{\mathbf{z}} = \mathbf{z}^{\text{prior}} + A \left(\mathbf{z}^{\text{radiance}} - \mathbf{z}^{\text{prior}} \right) + \mathbf{G} \eta, \quad \hat{\mathbf{x}} = \exp(\hat{\mathbf{z}}). \quad (32)$$

Here $A \in \mathbb{R}^{P \times P}$ is the averaging kernel matrix, \mathbf{G} is the gain matrix, and η is the spectral measurement error (assumed to have mean zero and covariance S_η). More details can be found in Jones et al. (2003); Worden et al. (2004); Bowman et al. (2006).

The corresponding TES observation operator \mathcal{H} is linear with respect to the logarithm of the concentrations, but nonlinear with respect to the concentration profile:

$$\mathcal{H}(\mathbf{x}) = \mathbf{z}^{\text{prior}} + A \left(\log(L(\mathbf{x})) - \mathbf{z}^{\text{prior}} \right)$$

where L is an interpolation operator that transforms \mathbf{x} from the GEOS-Chem N -level vertical grid to the TES profile retrieval P -level grid.

For this reason several chemical data assimilation studies based on TES retrieved profiles (Jones et al., 2003; Bowman et al., 2006; Parrington et al., 2009) have opted to perform the suboptimal Kalman filtering step in the logarithm of the concentrations:

$$\log \mathbf{x}^{\text{a}} = \log \mathbf{x}^{\text{f}} + K \left(\hat{\mathbf{z}} - \mathcal{H}(\mathbf{x}^{\text{f}}) \right).$$

For variational data assimilation the forcing calculation is carried out in concentrations. For this reason, an adjoint of the observation operator needs to be

Non-diagonal background error covariance matrices for data assimilation

K. Singh et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



derived to update the gradients as described in Eqs. (7) and (11)

$$(\mathcal{H}'(\mathbf{x}))^T \cdot v = \left(\frac{\partial}{\partial \mathbf{x}} (A \log(L(\mathbf{x}))) \right)^T \cdot v = \left(\frac{\partial L}{\partial \mathbf{x}} \right)^T \cdot \begin{pmatrix} (L\mathbf{x})_0^{-1} & 0 & \dots & 0 \\ 0 & (L\mathbf{x})_1^{-1} & \dots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \dots & (L\mathbf{x})_p^{-1} \end{pmatrix} \cdot A^T \cdot v$$

Here, $(\mathcal{H}'(\mathbf{x}))^T$ is a matrix and $v = \mathbb{R}^{-1} (\mathcal{H}(\mathbf{x}) - \mathbf{x}^{\text{obs}})$. The TES averaging kernel A is usually a non-symmetric matrix, and the result of $A^T \cdot v$ is fed to the interpolation operator to construct the diagonal matrix with the i -th element being $1/(L\mathbf{x})_i$. The term $(\partial L / \partial \mathbf{x})^T$ is the adjoint of the interpolation operator and brings entities from the TES profile retrieval domain back to the GEOS-Chem model domain.

5.1.3 Ozonesonde observations

For validation, we use the ozonesonde profiles measured by the INTEX Ozonesonde Network Study 2006 (IONS-6) (<http://croc.gsfc.nasa.gov/intexb/ions06.html> Thompson et al., 2007a,b) for the month of August, assuming that these measurements provide values close to the true state of the atmosphere. The ozonesonde observations are not used in data assimilation, and therefore provide an independent data set against which the analysis results are validated. There are 418 ozonesondes launched from 22 stations across North America as shown in the Fig. 3. A detailed description of the number of ozonesondes launched per station with longitude and latitude information can be found in (Parrington et al., 2008).

5.2 Impact of non-diagonal background error covariance in 3D-Var assimilation

In order to demonstrate the benefits of including spatial correlations in the background error covariance matrix, we first compare the tropospheric ozone concentrations

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



increments is smaller, and that the corrections are distributed over larger areas. Panels (e)–(f) of Fig. 5 compare directly the 3D-Var analyses obtained using a diagonal IB and a non-diagonal IB with a correlation length of 1000 km. The corrections in the non-diagonal case are spread are less aggressive and smoother.

5.3 Impact of non-diagonal background error covariance in 4D-Var assimilation

We now study the effects of using non-diagonal background error covariance matrix in 4D-Var data assimilation. We compare the analyses ozone concentrations generated by 4D-Var with different background error correlation lengths against the ozonesonde observations. The left panel of Fig. 6 shows the forecast, analysis and ozonesonde measured ozone concentrations averaged over the two months assimilation window. The model ozone fields are interpolated to the space-time location of each ozonesonde launch for comparison. The center panel shows the relative errors of model predictions with respect to ozonesonde data, averaged over all ozonesonde launches. The right panel provides the standard deviations of these errors.

The results indicate that 4D-Var is also sensitive to the structure of the background error covariance matrix. The use of non-diagonal correlations leads to improved analyses. The best analysis is obtained with a correlation length of 500 km (about one grid cell near the equator). Note that 4D-Var accounts for all the data available within the assimilation window.

To better understand the impact of different background error correlation lengths on 4D-Var assimilation, we present in Fig. 7 the differences in ozone concentrations generated by the free model run and by the 4D-Var assimilation using diagonal and non-diagonal IB. The use of a non-diagonal IB with a properly-chosen correlation length not only provides a better estimate but also helps generate a smoother analysis. This is due to the fact that the non-diagonal terms control how the information from observations is distributed spatially. The panels (e)–(f) of Fig. 7 compare directly the 4D-Var analyses obtained using a diagonal IB and a non-diagonal IB with a correlation

Non-diagonal background error covariance matrices for data assimilation

K. Singh et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)



[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



length of 500 km; the large localized corrections over North America provided by the diagonal IB are smoothened out when the non-diagonal IB is employed.

5.4 Determining the correlation length through experiments

The correlation length is a very important parameter that impacts the quality of the assimilation when using non-diagonal error covariance matrix. The value of the correlation length depends on various factors such as the lifetime of the tracer under consideration, the grid resolution, the pressure level, and the wind velocity. We propose a method to determine experimentally a value of the correlation length that is appropriate for the model and data at hand.

Recall the construction of one dimensional correlation matrices in Eqs. (13) and (14). Our aim is to determine the number of grid cells (in each direction) where the errors are correlated. For this we use adjoint sensitivity analysis. Specifically, we initialize the adjoint variable to 1 in a specific cell at the end of a given window (and to zero everywhere else), perform a backwards adjoint simulation, and analyze the adjoint fields at the beginning of the window. The error in the specific cell is correlated with errors in those grid cells where the adjoint values are above $1/e$. The length of the time window depends on the time scale of the model under consideration.

Here we consider a time window of 8 h. We run the forward GEOS-Chem model starting on 1 July, 00:00 GMT for 20 h. The adjoint variable for ozone at 20:00 GMT are initialized to 1 in a subset of the grid points ((i, j)) chosen such that $i \bmod 10 = 1$ and $j \bmod 10 = 1$. Adjoint variables for all other grid points and species are initialized to zero. The gap in the initialization helps avoid the interactions between adjoint “plumes” initialized at different locations. The ozone adjoint variable field is analyzed at 12:00 GMT to find out the number of grid cells where the value is greater than or equal to $1/e$. In the current setup the same correlation length is used for all for all pressure levels.

The procedure can be easily extended to estimate different correlation lengths for different vertical levels and for different geographic areas. This is desirable since the

Non-diagonal background error covariance matrices for data assimilation

K. Singh et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[⏪](#)

[⏩](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



ozone lifetime in the boundary layer is much shorter than it is in the upper troposphere; and the wind velocities are weaker in the boundary layer than in the upper troposphere. The chemical lifetime and the wind velocity both impact the correlation length. Based on the above considerations we expect that the correlation length is shorter near ground level and increases at higher altitudes.

Figure 8 shows that the ozone adjoint variables have spread (on average) over one to two grid cells in both longitudinal and latitudinal direction. For the $4^\circ \times 5^\circ$ model resolution each grid box is of size 400 km \times 500 km near the equator. The adjoint sensitivity analysis indicates that the correlation lengths should be chosen in the ranges of $\ell_x \in [500 \text{ km}, 1000 \text{ km}]$ and $\ell_y \in [400 \text{ km}, 800 \text{ km}]$, respectively. This confirms the best correlation lengths empirically observed in the data assimilation results reported in Figs. 4 and 6.

6 Conclusions

This paper presents an efficient methodology to construct non-diagonal background error covariance matrices for data assimilation. The two- or three-dimensional covariance matrices are not formed explicitly. Rather, multi-dimensional correlations are represented by tensor products of one dimensional correlation matrices along longitudinal and latitudinal directions. The technique can be easily extended to include correlations in the vertical direction as well. Highly efficient linear algebra operations are obtained by performing successive matrix-vector products, Cholesky decompositions, etc. with one-dimensional correlation matrices. The correlation lengths are important parameters that need to be specified for each directional correlation. We propose an adjoint sensitivity analysis approach to guide the choice of proper correlation lengths; the approach implicitly accounts for factors such as chemical activity, grid resolution, etc.

The approach to construct non-diagonal covariance matrices has been tested using the 3D-Var and 4D-Var data assimilation frameworks developed for GEOS-Chem. The

Non-diagonal background error covariance matrices for data assimilation

K. Singh et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)



[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



experiments assimilate observations from TES satellite ozone profile retrievals, and validate the results against an independent data set provided by IONS ozonesondes. The change of the covariance matrix formulation in data assimilation from diagonal to non-diagonal adds only a negligible computational overhead. In the same time, the inclusion of spatial correlations leads consistently to improved analyses in both the 3D-Var and the 4D-Var settings.

Acknowledgements. This work has been supported by NASA through the ROSES-2005 AIST project. The authors would like to thank Mark Parrington and Paul Hamer for providing processed TES data and for their help with the data visualization script.

References

- Beer, R., Glavich, T. A., and Rider, D. M.: Tropospheric emission spectrometer for the Earth Observing System's Aura satellite, *Appl. Optics*, 40(15), 2356–2367, 2001.
- Bey, I., Jacob, D. J., Yantosca, R. M., Logan, J. A., Field, B., Fiore, A. M., Li, Q., Liu, H., Mickley, L. J., and Schultz, M.: Global modeling of tropospheric chemistry with assimilated meteorology: Model description and evaluation, *J. Geophys. Res.*, 106, 23073–23096, 2001. 1792
- Bowman, K. W., Worden, J., Steck, T., Worden, H. M., Clough, S., and Rodgers, C.: Capturing time and vertical variability of tropospheric ozone: A study using TES nadir retrievals, *J. Geophys. Res.*, 107(D23), 4723, doi:10.1029/2002JD002150, 2002.
- Bowman, K. W., Rodgers, C. D., Kulawik, S. S., et al.: Tropospheric Emission Spectrometer: Retrieval method and error analysis, *IEEE T. Geosci. Remote*, 44(5), 1297–1307, May 2006. 1805
- Bowman, K. W., Jones, D. B. A., Logan, J. A., Worden, H., Boersma, F., Chang, R., Kulawik, S., Osterman, G., Hamer, P., and Worden, J.: The zonal structure of tropical O₃ and CO as observed by the Tropospheric Emission Spectrometer in November 2004 - Part 2: Impact of surface emissions on O₃ and its precursors, *Atmos. Chem. Phys.*, 9, 3563–3582, doi:10.5194/acp-9-3563-2009, 2009.
- Carmichael, G. R., Sandu, A., Chai, T., Daescu, D., Constantinescu, E. M., and Tang, Y.:

Non-diagonal background error covariance matrices for data assimilation

K. Singh et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)



[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



Non-diagonal background error covariance matrices for data assimilation

K. Singh et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Predicting air quality: Improvements through advanced methods to integrate models and measurements, *J. Comput. Phys.*, 227(7), 3540–3571, 2008. 1785

Chai, T., Carmichael, G. R., Sandu, A., Tang, Y., and Daescu, D. N.: Chemical data assimilation of transport and chemical evolution over the Pacific (TRACE-P) aircraft measurements, *J. Geophys. Res.*, 111, D02301, doi:10.1029/2005JD005883, 2006. 1786, 1793

Chai, T., Carmichael, G. R., Tang, Y., Sandu, A., Hardesty, M., Pilewskie, P., Whitlow, S., Browell, E. V., Avery, M. A., Thouret, V., Nedelec, P., Merrill, J. T., and Thomson, A. M.: Four dimensional data assimilation experiments with ICARTT (International Consortium for Atmospheric Transport and Transformation) ozone measurements, *J. Geophys. Res.*, 112, D12S15, doi:10.1029/2006JD007763, 2007. 1785

Chai, T., Carmichael, G. R., Tang, Y., Sandu, A., et al.: Regional NO_x emission inversion through a four-dimensional variational approach using SCIAMACHY tropospheric NO_2 column observations, *Atmos. Environ.*, 43(32), 5046–5055, doi:10.1016/j.atmosenv.2009.06.052, 2009. 1785

Clark, H. L., Cathala, M.-L., Teyss re, H., Cammas, J.-P., and Peuch, V.-H.: Cross-tropopause fluxes of ozone using assimilation of MOZAIC observations in a global CTM, *Tellus A and B*, 59, 39–49, 2006. 1785

Constantinescu, E. M., Chai, T., Sandu, A., and Carmichael, G. R.: Autoregressive models of background errors for chemical data assimilation, *J. Geophys. Res.*, 112, D12309, doi:10.1029/2006JD008103, 2007a. 1786, 1793, 1803

Constantinescu, E. M., Sandu, A., Chai, T., and Carmichael, G. R.: Investigation of ensemble-based chemical data assimilation in an idealized setting, *Atmos. Environ.*, 41(1), 18–36, 2007b. 1785, 1786

Constantinescu, E. M., Sandu, A., Chai, T., and Carmichael, G. R.: Ensemble-based chemical data assimilation. I: general approach, *Q. J. Roy. Meteor. Soc.*, 133(626), 1229–1243, Online ISSN:1477-870X, Print ISSN:0035-9009, Part A, July 2007c. 1785, 1803

Constantinescu, E. M., Sandu, A., Chai, T. and Carmichael, G. R.: Ensemble-based chemical data assimilation. II: covariance localization, *Q. J. Roy. Meteor. Soc.*, 133(626), 1245–1256, Online ISSN:1477-870X, Print ISSN:0035-9009, Part A, July 2007d. 1785, 1803

Courtier, P. and Talagrand, O.: Variational assimilation of meteorological observations with the adjoint vorticity equations II: Numerical results, *Q. J. Roy. Meteor. Soc.*, 113, 1329–1347, 1987. 1788

Courtier, P., Andersson, E., Heckley, W., et al.: The ECMWF implementation of three-

Non-diagonal background error covariance matrices for data assimilation

K. Singh et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

dimensional variational assimilation (3D-Var) I: Formulation, Q. J. Roy. Meteor. Soc., 124(550), p. 1783, 1998. 1785

Daescu, D., Carmichael, G. R., and Sandu, A.: Adjoint Implementation of Rosenbrock Methods Applied to Variational Data Assimilation Problems, *Comp. Phys.*, 165, 496–510, 2000. 1804

Daescu, D., Sandu, A., and Carmichael, G. R.: Direct and Adjoint Sensitivity Analysis of Chemical Kinetic Systems with KPP: II – Validation and Numerical Experiments, *Atmos. Environ.*, 37, 5097–5114, 2003. 1804

Daescu, D. N.: On the sensitivity equations of four-dimensional variational (4D-Var) data assimilation, *Mon. Weather Rev.*, 136(8), 3050–3065, 2008. 1792

Daley, R.: *Atmospheric Data Analysis*, Cambridge University Press, 457 pp., 1991. 1785

Damian, V., Sandu, A., Damian, M., Potra, F., and Carmichael, G. R.: The Kinetic PreProcessor KPP – A Software Environment for Solving Chemical Kinetics, *Comput. Chem. Eng.*, 26(11), 1567–1579, 2002. 1804

Elbern, H. and Schmidt, H.: Ozone episode analysis by four dimensional variational chemistry data assimilation, *J. Geophys. Res.*, 106(D4), 3569–3590, 2001. 1785

Eller, P., Singh, K., Sandu, A., Bowman, K., Henze, D. K., and Lee, M.: Implementation and evaluation of an array of chemical solvers in the Global Chemical Transport Model GEOS-Chem, *Geosci. Model Dev.*, 2, 89–96, doi:10.5194/gmd-2-89-2009, 2009. 1792, 1804

Gaspari, G. and Cohn, S. E.: Construction of correlation functions in two and three dimensions, *Q. J. Roy. Meteor. Soc.*, 125(554), 723–757, 1999. 1786, 1793

Hakami, A., Henze, D. K., Seinfeld, J. H., Chai, T., Tang, Y., Carmichael, G. R., and Sandu, A.: Adjoint inverse modeling of black carbon during ACE-Asia, *J. Geophys. Res.*, 110, D14301, 25 pp., doi:10.1029/2004JD005671, 2005. 1785

Henze, D. K., Hakami, A., and Seinfeld, J. H.: Development of the adjoint of GEOS-Chem, *Atmos. Chem. Phys.*, 7, 2413–2433, doi:10.5194/acp-7-2413-2007, 2007. 1792, 1804

Henze, D. K., Seinfeld, J. H., and Shindell, D. T.: Inverse modeling and mapping US air quality influences of inorganic PM_{2.5} precursor emissions using the adjoint of GEOS-Chem, *Atmos. Chem. Phys.*, 9, 5877–5903, doi:10.5194/acp-9-5877-2009, 2009. 1785, 1792

Jones, D. B. A., Bowman, K. W., Logan, J. A., Heald, C. L., Liu, J., Luo, M., Worden, J., and Drummond, J.: The zonal structure of tropical O₃ and CO as observed by the Tropospheric Emission Spectrometer in November 2004 - Part 1: Inverse modeling of CO emissions, *Atmos. Chem. Phys.*, 9, 3547–3562, doi:10.5194/acp-9-3547-2009, 2009.

Jones, D. B. A., Bowman, K. W., Palmer, P. I., Worden, J. R., Jacob, D. J., Hoffman, R. N., Bey, I.,

Non-diagonal background error covariance matrices for data assimilation

K. Singh et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

and Yantosca, R. M.: Potential of observations from the Tropospheric Emission Spectrometer to constrain continental sources of carbon monoxide, *J. Geophys. Res.*, 108(D24), 4789, doi:10.1029/2003JD003702, 2003. 1805

5 Kalnay, E.: Atmospheric modeling, data assimilation and predictability, Cambridge University Press, 2002. 1785

Khattatov, B. V., Gille, J. C., Lyjak, L. V., Brasseur, G. P., Dvortsov, V. L., Roche, A. E., and Walters, J.: Assimilation of photochemically active species and a case analysis of UARS data, *J. Geophys. Res.*, 104, 18715–18737, 1999. 1786

10 Khattatov, B. V., Lamarque, J.-F., Lyjak, L. V., Menard, R., Levelt, P., Tie, X., Brasseur, G. P., and Gille, J. C.: Assimilation of satellite observations of long-lived chemical species in global chemistry transport models, *J. Geophys. Res.*, 105(D23), 29–135, 2000. 1785

Kopacz, M., Jacob, D. J., Henze, D. K., Heald, C. L., Streets, D. G., and Zhang, Q.: A comparison of analytical and adjoint Bayesian inversion methods for constraining Asian sources of CO using satellite (MOPITT) measurements of CO columns, *J. Geophys. Res.*, 114, D04305, doi:10.1029/2007JD009264, 2009. 1792

15 Lahoz, W. A., Geer, A. J., Bekki, S., et al.: The Assimilation of Envisat data (ASSET) project, *Atmos. Chem. Phys.*, 7, 1773–1796, doi:10.5194/acp-7-1773-2007, 2007. 1785

Lamarque, J.-F., Khattatov, B. V., and Gille, J. C.: Constraining tropospheric ozone column through data assimilation, *J. Geophys. Res.*, 107(D22), 4651, doi:10.1029/2001JD001249, 2002. 1785

20 LeDimet, F.-X. and Talagrand, O.: Variational algorithms for analysis and assimilation of meteorological observations: theoretical aspects, *Tellus A*, 38, 97–110, 1986. 1788

Lions, J. L.: Optimal control of systems governed by partial differential equations, Springer-Verlag, 1971. 1788

25 Menard, R., Cohn, S. E., Chang, L.-P., and Lyster, P. M.: Assimilation of stratospheric chemical tracer observations using a Kalman Filter I: Formulation, *Mon. Weather Rev.*, 128, 2654–2671, 2000. 1785

Nassar, R., Logan, J. A., Worden, H. M., et al.: Validation of Tropospheric Emission Spectrometer (TES) nadir ozone profiles using ozonesonde measurements, *J. Geophys. Res.*, 113, D15S17, doi:10.1029/2007JD008819, 2008. 1803

30 Navon, I. M.: Data assimilation for Numerical Weather Prediction: a review, in: *Data Assimilation for Atmospheric, Oceanic, and Hydrologic Applications*, edited by: Park, S. K. and Xu, L., Springer-Verlag Berlin Heidelberg, 21–65, doi:10.1007/978-3-540-71056-1,

Non-diagonal background error covariance matrices for data assimilation

K. Singh et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

2009. 1785

Ott, E., Hunt, B. R., Szunyogh, I., Zimin, A. V., Kostelich, E. J., Kostelich, M., Corazza, M., Sauer, T., Kalnay, E., Patil, D. J., and Yorke, J. A.: A local ensemble Kalman Filter for Atmospheric Data Assimilation, *Tellus A*, 56, 415–428, 2004. 1786

5 Palmer, P. I., Jacob, D. J., Jones, D. B. A., Heald, C. L., Yantosca, R. M., Logan, J. A., Sachse, G. W., and Streets, D. G.: Observations over the western Pacific, *J. Geophys. Res.*, 108(D21), 8825, doi:10.1029/2002JD003176, 2003.

Parrington, M., Jones, D. B. A., Bowman, K. W., Horowitz, L. W., Thompson, A. M., Tarasick, D. W., and Witte, J. C.: Estimating the summertime tropospheric ozone distribution over North America through assimilation of observations from the Tropospheric Emission Spectrometer, *J. Geophys. Res.*, 113, D18307, doi:10.1029/2007JD009341, 2008. 1787, 1806

10 Parrington, M., Jones, D. B. A., Bowman, K. W., Thompson, A. M., Tarasick, D. W., Merrill, J., Oltmans, S. J., Leblanc, T., Witte, J. C. and Millet, D. B.: Impact of the assimilation of ozone from the tropospheric emission spectrometer on surface ozone across North America, *Geophys. Res. Lett.*, 36, L04802, doi:10.1029/2008GL036935, 2009. 1785, 1805

Parrish, D. F. and Derber, J. C.: The national meteorological center's spectral statistical-interpolation analysis system, *Mon. Weather Rev.*, 120, 1747–1763, 1992. 1786

Pierce, R. B., Schaack, T., Al-Saadi, J. A., et al.: Chemical data assimilation estimates of continental U. S. ozone and nitrogen budgets during the Intercontinental Chemical Transport Experiment-North America, *J. Geophys. Res.*, 112, D12S21, doi:10.1029/2006JD007722, 2007. 1785, 1786

Rabier, F., Jarvinen, H., Klinker, E., Mahfouf, J.-F., and Simmons, A.: The ECMWF operational implementation of four-dimensional variational data assimilation I: Experimental results with simplified physics, *Q. J. Roy. Meteor. Soc.*, 126, 1143–1170, 2000. 1785

25 Sandu, A., Daescu, D., and Carmichael, G. R.: Direct and Adjoint Sensitivity Analysis of Chemical Kinetic Systems with KPP: I – Theory and Software Tools, *Atmos. Environ.*, 37, 5083–5096, 2003. 1804

Sandu, A., Daescu, D. N., Carmichael, G. R., and Chai, T.: Adjoint sensitivity analysis of regional air quality models, *J. Comput. Phys.*, 204, 222–252, 2005. 1785, 1802

30 Sandu, A. and Zhang, L.: Discrete second order adjoints in atmospheric chemical transport modeling, *J. Computat. Phys.*, 227(12), 5949–5983, 2008.

Sasaki, Y. K.: An objective analysis based on the variational method, *J. Met. Soc. Jap.*, II(36), 77–88, 1958. 1788

Non-diagonal background error covariance matrices for data assimilation

K. Singh et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Segers, A. J., Eskes, H. J., van der A, R. J., van Oss, R. F., and van Velthoven, P. F. J.: Assimilation of GOME ozone profiles and a global chemistry-transport model, using a Kalman Filter with anisotropic covariance, *Q. J. Roy. Meteor. Soc.*, 131, 477–502, 2005. 1785
- 5 Singh, K., Eller, P., Sandu, A., Bowman, K. W., Jones, D. B. A., and Lee, M.: Improving GEOS-Chem model forecasts through profile retrievals from Tropospheric Emission Spectrometer, in: *Lecture Notes on Computational Science*, 5545, 302–311, International Conference on Computational Science 2009, Baton Rouge, Louisiana, May 25–27, 2009. 1792
- 10 Singh, K., Eller, P., Sandu, A., Henze, D., Bowman, K. W., Kopacz, M., and Lee, M.: Towards the construction of a standard adjoint GEOS-Chem model, High Performance Computing Symposium (HPC 2009) at Spring Simulation Multiconference (SpringSim'09), San Diego, California, 22–27 March, 2009. 1792, 1802
- TES Science Team: TES L2 Data Users Guide, Jet Propulsion Laboratory, California Institute of Technology, Pasadena, California, available at: <http://tes.jpl.nasa.gov/uploadedfiles/TES-DPS.V11.8.pdf> (last access: September 2010), 2008.
- 15 Thompson, A. M., et al.: Intercontinental chemical transport experiment ozonesonde network study (IONS) 2004: 1. Summertime upper troposphere/lower stratosphere ozone over northeastern North America, *J. Geophys. Res.*, 112, D12S12, doi:10.1029/2006JD007441, 2007a. 1806
- 20 Thompson, A. M., Stone, J. B., Witte, J. C., et al.: Intercontinental chemical transport experiment ozonesonde network study (IONS) 2004: 2. Tropospheric ozone budgets and variability over northeastern North America, *J. Geophys. Res.*, 112, D12S13, doi:10.1029/2004JD005359, 2007b. 1806
- Worden, J. R., Bowman, K. W., and Jones, D. B. A.: Characterization of atmospheric profile retrievals from Limb Sounding Observations of an inhomogeneous atmosphere, *J. Quant. Spectrosc. Ra.*, 86, (03)00274-7, 2004. 1805
- 25 Wu, L., Mallet, V., Bocquet, M., and Sportisse, B.: A comparison study of data assimilation algorithms for ozone forecasts, *J. Geophys. Res.*, 113, D20310, 2008. 1785
- Zhang, L., Constantinescu, E. M., Sandu, A., et al.: An adjoint sensitivity analysis and 4D-Var data assimilation study of Texas air quality, *Atmos. Environ.*, 42(23), 5787–5804, 2008. 1785
- 30 Zhu, C., Byrd, R. H., and Nocedal, J.: L-BFGS-B: Algorithm 778: L-BFGS-B, FORTRAN routines for large scale bound constrained optimization, *ACM T. Math. Software*, 23(4), 550–560, 1997. 1802

Non-diagonal background error covariance matrices for data assimilation

K. Singh et al.

Table 1. Timing results for GEOS-Chem free model run, 3D-Var and 4D-Var data assimilations with diagonal and non-diagonal IB for a 24 h simulation starting 1 July 2006.

Experiment Description	CPU Time	Scaled time
Free model run with SMVGEAR chemistry solver	2 min 50 sec	1.00
Free model run with KPP chemistry solver	3 min 18 sec	1.16
3D-Var with diagonal IB	3 min 57 sec	1.39
3D-Var with non-diagonal IB, Cholesky	4 min 00 sec	1.41
3D-Var with non-diagonal IB, SVD	9 min 38 sec	3.40
4D-Var with diagonal IB (per iteration)	16 min 51 sec	5.95
4D-Var with non-diagonal IB, Cholesky (per iteration)	16 min 51 sec	5.95

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶


Back

Close


Full Screen / Esc

Printer-friendly Version

Interactive Discussion



(a) Longitudinal correlation matrix C_x^k at latitude $y_k = 20^\circ$ N.



(b) Latitudinal correlation matrix C_y .

Fig. 1. Mesh representation of the one-dimensional longitudinal and latitudinal correlation matrices. The latitude-longitude model grid resolution is $4^\circ \times 5^\circ$ (about $400 \text{ km} \times 500 \text{ km}$ near the equator) and the correlation lengths are $\ell_x = 1500 \text{ km}$ and $\ell_y = 1200 \text{ km}$.

Non-diagonal background error covariance matrices for data assimilation

K. Singh et al.

Title Page

Abstract	Introduction
Conclusions	References
Tables	Figures

⏪	⏩
◀	▶
Back	Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**Non-diagonal
background error
covariance matrices
for data assimilation**

K. Singh et al.




Fig. 2. Contour lines of the longitudinal correlation C_x for points at different latitudes. The correlation length ℓ_x is short (top panel), medium (middle panel), and large (bottom panel). Note that the same correlation length ℓ_x translates into a different number of correlated grid points depending on the latitude.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

**Non-diagonal
background error
covariance matrices
for data assimilation**

K. Singh et al.




Fig. 3. Ozonesonde sounding stations (triangles) used during IONS06 campaign and AURA/TES satellite trajectory snapshots (dots) plotted over the global ozone distribution on 1 August 2006.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Non-diagonal background error covariance matrices for data assimilation

K. Singh et al.




Fig. 4. The impact of non-diagonal background error covariances in 3D-Var data assimilation. Left panel: mean ozone concentrations at ozonesonde locations for 3D-Var analyses and free model trajectories. Center panel: relative mean errors of predicted ozone concentrations with respect to ozonesonde measurements. Right panel: standard deviation of absolute values of errors with respect to ozonesonde measurements. The data is averaged over all ozonesonde launches. These plots were generated from 2 months simulation from 00:00 GMT 1 July to 23:00 GMT August 2006 and compared against ozonesonde data available for the month of August.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Non-diagonal background error covariance matrices for data assimilation

K. Singh et al.

[Title Page](#)

[Abstract](#) [Introduction](#)

[Conclusions](#) [References](#)

[Tables](#) [Figures](#)

[⏪](#) [⏩](#)

[◀](#) [▶](#)

[Back](#)


[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)


[Interactive Discussion](#)

GEOS-4 O_x Absolute Difference: Diagonal B - Forecast




(a) Absolute difference between the 3D-Var analysis using diagonal IB ($\ell_x = 0$ km) and the free model run.

GEOS-4 O_x Absolute Difference: l_x=500Km - Forecast




(b) Absolute difference between the 3D-Var analysis using non-diagonal IB ($\ell_x = 500$ km) and the free model run.

GEOS-4 O_x Absolute Difference: l_x=1000Km - Forecast



(c) Absolute difference between the 3D-Var analysis using non-diagonal IB ($\ell_x = 1000$ km) and the free model run.

GEOS-4 O_x Absolute Difference: l_x=1500Km - Forecast



(d) Absolute difference between the 3D-Var analysis using non-diagonal IB ($\ell_x = 1500$ km) and the free model run.

Fig. 5. Differences in global ozone concentrations at 23:00 GMT on 31 August 2006 averaged over the first 10 GEOS-Chem vertical levels. Panels **(a)–(d)**: differences between the 3D-Var analysis fields and the model forecast (solution without data assimilation); the analyses use different correlation lengths between 0 km and 1500 km.



Non-diagonal background error covariance matrices for data assimilation

K. Singh et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶


Back

Close


Full Screen / Esc

Printer-friendly Version

Interactive Discussion



(e) Absolute difference between the 3D-Var analyses using diagonal IB and non-diagonal IB ($\ell_x = 1000$ km).



(f) Relative difference between the 3D-Var analyses using diagonal IB and non-diagonal IB ($\ell_x = 1000$ km).

Fig. 5. Panels **(e)–(f)**: absolute and relative differences between 3D-Var analyses using diagonal and non-diagonal background covariance matrices.

Non-diagonal background error covariance matrices for data assimilation

K. Singh et al.




Fig. 6. The impact of non-diagonal background error covariances on 4D-Var data assimilation. The results shown are for a single 5-day assimilation window from 00:00 GMT August 1st to 00:00 GMT August 6th, 2006. Left panel: mean ozone concentrations at ozonesonde locations for 4D-Var analyses and free model trajectories. Center panel: relative mean errors of predicted ozone concentrations with respect to ozonesonde measurements. Right panel: standard deviation of absolute values of errors with respect to ozonesonde measurements.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

⏪

⏩

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)


[Printer-friendly Version](#)

[Interactive Discussion](#)


Non-diagonal background error covariance matrices for data assimilation

K. Singh et al.


Title Page	
Abstract	Introduction
Conclusions	References
Tables	Figures
⏪	⏩
◀	▶
Back	Close
Full Screen / Esc	
Printer-friendly Version	
Interactive Discussion	




(a) Absolute difference between the 4D-Var analysis using diagonal IB ($\ell_x = 0$ km) and the free model run.



(b) Absolute difference between the 4D-Var analysis using non-diagonal IB ($\ell_x = 500$ km) and the free model run.



(c) Absolute difference between the 4D-Var analysis using non-diagonal IB ($\ell_x = 1000$ km) and the free model run.




(d) Absolute difference between the 4D-Var analysis using non-diagonal IB ($\ell_x = 1500$ km) and the free model run.

Fig. 7. Differences in global ozone concentrations at 00:00 GMT on 6 August 2006 (end of assimilation window) averaged over the first 10 GEOS-Chem vertical levels. Panels **(a)–(d)**: differences between the 4D-Var analysis fields and the model forecast (solution without data assimilation); the analyses use different correlation lengths between 0 km and 1500 km.



Non-diagonal background error covariance matrices for data assimilation

K. Singh et al.



(e) Absolute difference between the 4D-Var analyses using diagonal IB and non-diagonal IB ($\ell_x = 500$ km).

(f) Relative difference between the 4D-Var analyses using diagonal IB and non-diagonal IB ($\ell_x = 500$ km).

Fig. 7. Panels **(e)–(f)**: absolute and relative differences between 4D-Var analyses using diagonal and non-diagonal background covariance matrices.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close


Full Screen / Esc

Printer-friendly Version


Interactive Discussion

Non-diagonal background error covariance matrices for data assimilation

K. Singh et al.



(a) Adjoint ozone variables are initialized to one in selected grid cells.



(b) Adjoint ozone variables spread after 8 h of backward sensitivity run.

Fig. 8. Ground level ozone adjoint variable values are initialized to one on 1 July 2006, 20:00 GMT, every tenth grid point in longitudinal and latitudinal directions. An 8 h backward adjoint integration spreads the adjoint fields, and helps identify grid cells where ozone errors are correlated.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion