The Chemical Mechanism of MECCA

KPP version: `2.2.1_rs5`

MECCA version: `3.0gmdd`

Date: December 20, 2010.

Selected reactions:
"`!Ara`"

Number of aerosol phases: 1

Number of species in selected mechanism:

Gas phase:	197
Aqueous phase:	89
All species:	286

Number of reactions in selected mechanism:

Gas phase (Gnnn):	378
Aqueous phase (Annn):	147
Henry (Hnnn):	93
Photolysis (Jnnn):	116
Heterogeneous (HETnnn):	21
Equilibria (EQnn):	68
Isotope exchange (DGnnn):	0
Dummy (Dnn):	1
All equations:	826

This document is part of the electronic supplement to our article
"The atmospheric chemistry box model CAABA/MECCA-3.0gmdd"
in Geosci. Model Dev. Discuss. (2011), available at:
`http://www.geosci-model-dev.net`

Table 1: Gas phase reactions

#	labels	reaction	rate coefficient	reference
G1000	StTrG	$O_2 + O(^1D) \rightarrow O(^3P) + O_2$	3.3E-11*EXP(55./temp)	Sander et al. (2006)
G1001	StTrG	$O_2 + O(^3P) \rightarrow O_3$	6.E-34*((temp/300.)**(-2.4))*cair	Sander et al. (2006)
G1002	StG	$O_3 + O(^1D) \rightarrow 2\ O_2$	1.2E-10	Sander et al. (2006)*
G1003	StG	$O_3 + O(^3P) \rightarrow 2\ O_2$	8.E-12*EXP(-2060./temp)	Sander et al. (2006)
G2100	StTrG	$H + O_2 \rightarrow HO_2$	k_3rd(temp,cair,4.4E-32,1.3,4.7E-11, 0.2,0.6)	Sander et al. (2006)
G2101	StG	$H + O_3 \rightarrow OH + O_2$	1.4E-10*EXP(-470./temp)	Sander et al. (2006)
G2102	StG	$H_2 + O(^1D) \rightarrow H + OH$	1.1E-10	Sander et al. (2006)
G2103	StG	$OH + O(^3P) \rightarrow H + O_2$	2.2E-11*EXP(120./temp)	Sander et al. (2006)
G2104	StTrG	$OH + O_3 \rightarrow HO_2 + O_2$	1.7E-12*EXP(-940./temp)	Sander et al. (2006)
G2105	StTrG	$OH + H_2 \rightarrow H_2O + H$	2.8E-12*EXP(-1800./temp)	Sander et al. (2006)
G2106	StG	$HO_2 + O(^3P) \rightarrow OH + O_2$	3.E-11*EXP(200./temp)	Sander et al. (2006)
G2107	StTrG	$HO_2 + O_3 \rightarrow OH + 2\ O_2$	1.E-14*EXP(-490./temp)	Sander et al. (2006)
G2108a	StG	$HO_2 + H \rightarrow 2\ OH$	7.2E-11	Sander et al. (2006)
G2108b	StG	$HO_2 + H \rightarrow H_2 + O_2$	6.9E-12	Sander et al. (2006)
G2108c	StG	$HO_2 + H \rightarrow O(^3P) + H_2O$	1.6E-12	Sander et al. (2006)
G2109	StTrG	$HO_2 + OH \rightarrow H_2O + O_2$	4.8E-11*EXP(250./temp)	Sander et al. (2006)
G2110	StTrG	$HO_2 + HO_2 \rightarrow H_2O_2 + O_2$	k_HO2_HO2	Christensen et al. (2002), Kircher and Sander (1984)*
G2111	StTrG	$H_2O + O(^1D) \rightarrow 2\ OH$	1.63E-10*EXP(60./temp)	Sander et al. (2006)
G2112	StTrG	$H_2O_2 + OH \rightarrow H_2O + HO_2$	1.8E-12	Sander et al. (2006)
G3100	StGN	$N + O_2 \rightarrow NO + O(^3P)$	1.5E-11*EXP(-3600./temp)	Sander et al. (2006)
G3101	StTrG	$N_2 + O(^1D) \rightarrow O(^3P) + N_2$	2.15E-11*EXP(110./temp)	Sander et al. (2006)
G3102a	StGN	$N_2O + O(^1D) \rightarrow 2\ NO$	6.7E-11*EXP(20./temp)	Sander et al. (2006)
G3102b	StGN	$N_2O + O(^1D) \rightarrow N_2 + O_2$	4.7E-11*EXP(20./temp)	Sander et al. (2006)
G3103	StTrGN	$NO + O_3 \rightarrow NO_2 + O_2$	3.E-12*EXP(-1500./temp)	Sander et al. (2006)
G3104	StGN	$NO + N \rightarrow O(^3P) + N_2$	2.1E-11*EXP(100./temp)	Sander et al. (2006)
G3105	StGN	$NO_2 + O(^3P) \rightarrow NO + O_2$	5.1E-12*EXP(210./temp)	Sander et al. (2006)
G3106	StTrGN	$NO_2 + O_3 \rightarrow NO_3 + O_2$	1.2E-13*EXP(-2450./temp)	Sander et al. (2006)
G3107	StGN	$NO_2 + N \rightarrow N_2O + O(^3P)$	5.8E-12*EXP(220./temp)	Sander et al. (2006)
G3108	StTrGN	$NO_3 + NO \rightarrow 2\ NO_2$	1.5E-11*EXP(170./temp)	Sander et al. (2006)
G3109	StTrGN	$NO_3 + NO_2 \rightarrow N_2O_5$	k_NO3_NO2	Sander et al. (2006)*
G3110	StTrGN	$N_2O_5 \rightarrow NO_2 + NO_3$	k_NO3_NO2/(2.7E-27*EXP(11000./temp))	Sander et al. (2006)*

Table 1: Gas phase reactions (... continued)

#	labels	reaction	rate coefficient	reference
G3200	TrGN	$NO + OH \rightarrow HONO$	k_3rd(temp,cair,7.0E-31,2.6,3.6E-11, 0.1,0.6)	Sander et al. (2006)
G3201	StTrGN	$NO + HO_2 \rightarrow NO_2 + OH$	3.5E-12*EXP(250./temp)	Sander et al. (2006)
G3202	StTrGN	$NO_2 + OH \rightarrow HNO_3$	k_3rd(temp,cair,1.8E-30,3.0,2.8E-11,0., 0.6)	Sander et al. (2006)
G3203	StTrGN	$NO_2 + HO_2 \rightarrow HNO_4$	k_NO2_HO2	Sander et al. (2006)*
G3204	TrGN	$NO_3 + HO_2 \rightarrow NO_2 + OH + O_2$	3.5E-12	Sander et al. (2006)
G3205	TrGN	$HONO + OH \rightarrow NO_2 + H_2O$	1.8E-11*EXP(-390./temp)	Sander et al. (2006)
G3206	StTrGN	$HNO_3 + OH \rightarrow H_2O + NO_3$	k_HNO3_OH	Sander et al. (2006)*
G3207	StTrGN	$HNO_4 \rightarrow NO_2 + HO_2$	k_NO2_HO2/(2.1E-27*EXP(10900./temp))	Sander et al. (2006)*
G3208	StTrGN	$HNO_4 + OH \rightarrow NO_2 + H_2O$	1.3E-12*EXP(380./temp)	Sander et al. (2006)
G3209	TrGN	$NH_3 + OH \rightarrow NH_2 + H_2O$	1.7E-12*EXP(-710./temp)	Kohlmann and Poppe (1999)
G3210	TrGN	$NH_2 + O_3 \rightarrow NH_2O + O_2$	4.3E-12*EXP(-930./temp)	Kohlmann and Poppe (1999)
G3211	TrGN	$NH_2 + HO_2 \rightarrow NH_2O + OH$	4.8E-07*EXP(-628./temp)*temp**(-1.32)	Kohlmann and Poppe (1999)
G3212	TrGN	$NH_2 + HO_2 \rightarrow HNO + H_2O$	9.4E-09*EXP(-356./temp)*temp**(-1.12)	Kohlmann and Poppe (1999)
G3213	TrGN	$NH_2 + NO \rightarrow HO_2 + OH + N_2$	1.92E-12*((temp/298.)**(-1.5))	Kohlmann and Poppe (1999)
G3214	TrGN	$NH_2 + NO \rightarrow N_2 + H_2O$	1.41E-11*((temp/298.)**(-1.5))	Kohlmann and Poppe (1999)
G3215	TrGN	$NH_2 + NO_2 \rightarrow N_2O + H_2O$	1.2E-11*((temp/298.)**(-2.0))	Kohlmann and Poppe (1999)
G3216	TrGN	$NH_2 + NO_2 \rightarrow NH_2O + NO$	0.8E-11*((temp/298.)**(-2.0))	Kohlmann and Poppe (1999)
G3217	TrGN	$NH_2O + O_3 \rightarrow NH_2 + O_2$	1.2E-14	Kohlmann and Poppe (1999)
G3218	TrGN	$NH_2O \rightarrow NHOH$	1.3E3	Kohlmann and Poppe (1999)
G3219	TrGN	$HNO + OH \rightarrow NO + H_2O$	8.0E-11*EXP(-500./temp)	Kohlmann and Poppe (1999)
G3220	TrGN	$HNO + NHOH \rightarrow NH_2OH + NO$	1.66E-12*EXP(-1500./temp)	Kohlmann and Poppe (1999)
G3221	TrGN	$HNO + NO_2 \rightarrow HONO + NO$	1.0E-12*EXP(-1000./temp)	Kohlmann and Poppe (1999)
G3222	TrGN	$NHOH + OH \rightarrow HNO + H_2O$	1.66E-12	Kohlmann and Poppe (1999)
G3223	TrGN	$NH_2OH + OH \rightarrow NHOH + H_2O$	4.13E-11*EXP(-2138./temp)	Kohlmann and Poppe (1999)
G3224	TrGN	$HNO + O_2 \rightarrow HO_2 + NO$	3.65E-14*EXP(-4600./temp)	Kohlmann and Poppe (1999)
G4100	StG	$CH_4 + O(^1D) \rightarrow .75\ CH_3O_2 + .75\ OH + .25\ HCHO + .4\ H + .05\ H_2$	1.5E-10	Sander et al. (2006)
G4101	StTrG	$CH_4 + OH \rightarrow CH_3O_2 + H_2O$	1.85E-20*EXP(2.82*log(temp)-987./temp)	Atkinson (2003)
G4102	TrG	$CH_3OH + OH \rightarrow HCHO + HO_2$	2.9E-12*EXP(-345./temp)	Sander et al. (2006)
G4103	StTrG	$CH_3O_2 + HO_2 \rightarrow CH_3OOH + O_2$	4.1E-13*EXP(750./temp)	Sander et al. (2006)*
G4104	StTrGN	$CH_3O_2 + NO \rightarrow HCHO + NO_2 + HO_2$	2.8E-12*EXP(300./temp)	Sander et al. (2006)
G4105	TrGN	$CH_3O_2 + NO_3 \rightarrow HCHO + HO_2 + NO_2$	1.3E-12	Atkinson et al. (2006)

Table 1: Gas phase reactions (... continued)

#	labels	reaction	rate coefficient	reference
G4106a	StTrG	$CH_3O_2 \rightarrow HCHO + HO_2$	2.*RO2*9.5E-14*EXP(390./temp)/(1.+1./ 26.2*EXP(1130./temp))	Sander et al. (2006)
G4106b	StTrG	$CH_3O_2 \rightarrow .5\ HCHO + .5\ CH_3OH + .5\ O_2$	2.*RO2*9.5E-14*EXP(390./temp)/(1.+ 26.2*EXP(-1130./temp))	Sander et al. (2006)
G4107	StTrG	$CH_3OOH + OH \rightarrow .7\ CH_3O_2 + .3\ HCHO + .3\ OH + H_2O$	k_CH3OOH_OH	Sander et al. (2006)*
G4108	StTrG	$HCHO + OH \rightarrow CO + H_2O + HO_2$	9.52E-18*EXP(2.03*log(temp)+636./temp)	Sivakumaran et al. (2003)
G4109	TrGN	$HCHO + NO_3 \rightarrow HNO_3 + CO + HO_2$	3.4E-13*EXP(-1900./temp)	Sander et al. (2006)*
G4110	StTrG	$CO + OH \rightarrow H + CO_2$	(1.57E-13+cair*3.54E-33)	McCabe et al. (2001)
G4111	TrG	$HCOOH + OH \rightarrow CO_2 + HO_2 + H_2O$	4.0E-13	Sander et al. (2006)
G4200	TrGC	$C_2H_6 + OH \rightarrow C_2H_5O_2 + H_2O$	1.49E-17*temp*temp*EXP(-499./temp)	Atkinson (2003)
G4201	TrGC	$C_2H_4 + O_3 \rightarrow HCHO + .63\ CO + .13\ HO_2 + 0.23125\ HCOOH + 0.13875\ HCHO + 0.13875\ H_2O_2 + .13\ OH$	1.2E-14*EXP(-2630./temp)	Sander et al. (2006)*
G4202	TrGC	$C_2H_4 + OH \rightarrow HOCH_2CH_2O_2$	k_3rd(temp,cair,1.0E-28,4.5,8.8E-12, 0.85,0.6)	Sander et al. (2006)
G4203	TrGC	$C_2H_5O_2 + HO_2 \rightarrow C_2H_5OOH$	7.5E-13*EXP(700./temp)	Sander et al. (2006)
G4204	TrGNC	$C_2H_5O_2 + NO \rightarrow CH_3CHO + HO_2 + NO_2$	2.6E-12*EXP(365./temp)	Sander et al. (2006)
G4205	TrGNC	$C_2H_5O_2 + NO_3 \rightarrow CH_3CHO + HO_2 + NO_2$	2.3E-12	Atkinson et al. (1999)
G4206	TrGC	$C_2H_5O_2 \rightarrow .98\ CH_3CHO + .38\ HO_2 + .02\ HOCH_2CH_2O_2$	3.1E-13*RO2	Rickard and Pascoe (2009)*
G4207	TrGC	$C_2H_5OOH + OH \rightarrow .43\ C_2H_5O_2 + .43\ H_2O + .57\ CH_3CHO + .57\ OH$	0.6*k_CH3OOH_OH + 8.01E-12	see note
G4208	TrGC	$CH_3CHO + OH \rightarrow CH_3C(O)OO + H_2O$	4.4E-12*EXP(365./temp)	Atkinson et al. (2006)
G4209	TrGNC	$CH_3CHO + NO_3 \rightarrow CH_3C(O)OO + HNO_3$	KNO3AL	Sander et al. (2006)
G4210	TrGC	$CH_3COOH + OH \rightarrow CH_3O_2 + CO_2 + H_2O$	4.2E-14*EXP(855./temp)	Atkinson et al. (2006)
G4211a	TrGC	$CH_3C(O)OO + HO_2 \rightarrow CH_3C(O)OOH$	4.3E-13*EXP(1040./temp)/(1.+1./ 37.*EXP(660./temp))	Tyndall et al. (2001)
G4211b	TrGC	$CH_3C(O)OO + HO_2 \rightarrow CH_3COOH + O_3$	4.3E-13*EXP(1040./temp)/(1.+ 37.*EXP(-660./temp))	Tyndall et al. (2001)
G4212	TrGNC	$CH_3C(O)OO + NO \rightarrow CH_3O_2 + CO_2 + NO_2$	8.1E-12*EXP(270./temp)	Tyndall et al. (2001)
G4213	TrGNC	$CH_3C(O)OO + NO_2 \rightarrow PAN$	k_CH3CO3_NO2	Sander et al. (2006)
G4214	TrGNC	$CH_3C(O)OO + NO_3 \rightarrow CH_3O_2 + NO_2 + CO_2$	4.E-12	Canosa-Mas et al. (1996)
G4217	TrGC	$CH_3C(O)OO \rightarrow .7\ CH_3O_2 + .7\ CO_2 + .3\ CH_3COOH$	1.00E-11*RO2	Rickard and Pascoe (2009)
G4218	TrGC	$CH_3C(O)OOH + OH \rightarrow CH_3C(O)OO + H_2O$	0.6*k_CH3OOH_OH	Rickard and Pascoe (2009)*
G4220	TrGNC	$PAN + OH \rightarrow HCHO + CO + NO_2 + H_2O$	9.50E-13*EXP(-650./temp)	Rickard and Pascoe (2009)
G4221	TrGNC	$PAN \rightarrow CH_3C(O)OO + NO_2$	k_PAN_M	Sander et al. (2006)*

Table 1: Gas phase reactions (... continued)

#	labels	reaction	rate coefficient	reference
G4222	TrGC	C_2H_2 + OH → 0.636 GLYOX + 0.636 OH + 0.364 HCOOH + 0.364 CO + 0.364 HO_2	k_3rd(temp,cair,5.5e-30,0.0,8.3e-13, -2.,0.6)	Sander et al. (2006)
G4223	TrGC	$HOCH_2CHO$ + OH → .8 $HOCH_2CO_3$ + .2 GLYOX + .2 HO_2 + H_2O	1.00E-11	Rickard and Pascoe (2009)
G4224	TrGNC	$HOCH_2CHO$ + NO_3 → $HOCH_2CO_3$ + HNO_3	KNO3AL	Rickard and Pascoe (2009)
G4225	TrGC	$HOCH_2CO_3$ → .7 HCHO + .7 CO_2 + .7 HO_2 + .3 $HOCH_2CO_2H$	1.00E-11*RO2	Rickard and Pascoe (2009)
G4226	TrGC	$HOCH_2CO_3$ + HO_2 → .71 $HOCH_2CO_3H$ + .29 $HOCH_2CO_2H$ + .29 O_3	KAPHO2	Rickard and Pascoe (2009)
G4227	TrGNC	$HOCH_2CO_3$ + NO → NO_2 + HO_2 + HCHO + CO_2	KAPNO	Rickard and Pascoe (2009)
G4228	TrGNC	$HOCH_2CO_3$ + NO_2 → PHAN	k_CH3CO3_NO2	Rickard and Pascoe (2009)
G4229	TrGNC	$HOCH_2CO_3$ + NO_3 → NO_2 + HO_2 + HCHO + CO_2	KRO2NO3*1.60	Rickard and Pascoe (2009)
G4230	TrGC	$HOCH_2CO_2H$ + OH → HCHO + HO_2 + CO_2 + H_2O	2.73E-12	Rickard and Pascoe (2009)
G4231	TrGC	$HOCH_2CO_3H$ + OH → $HOCH_2CO_3$ + H_2O	6.19E-12	Rickard and Pascoe (2009)
G4232	TrGNC	PHAN → $HOCH_2CO_3$ + NO_2	k_PAN_M	Rickard and Pascoe (2009)
G4233	TrGNC	PHAN + OH → HCHO + CO + NO_2 + H_2O	1.12E-12	Rickard and Pascoe (2009)
G4234	TrGC	GLYOX + OH → 1.2 CO + .6 HO_2 + .4 $HCOCO_3$ + H_2O	1.14E-11	Rickard and Pascoe (2009)
G4235	TrGNC	GLYOX + NO_3 → 1.2 CO + .6 HO_2 + .4 $HCOCO_3$ + HNO_3	KNO3AL	Rickard and Pascoe (2009)
G4236	TrGC	$HCOCO_3$ → .7 CO + .7 HO_2 + .7 CO_2 + .3 $HCOCO_2H$	1.00E-11*RO2	Rickard and Pascoe (2009)
G4237	TrGC	$HCOCO_3$ + HO_2 → .71 $HCOCO_3H$ + .29 $HCOCO_2H$ + .29 O_3	KAPHO2	Rickard and Pascoe (2009)
G4238	TrGNC	$HCOCO_3$ + NO → HO_2 + CO + NO_2 + CO_2	KAPNO	Rickard and Pascoe (2009)
G4239	TrGNC	$HCOCO_3$ + NO_3 → HO_2 + CO + NO_2 + CO_2	KRO2NO3*1.60	Rickard and Pascoe (2009)
G4240	TrGC	$HCOCO_2H$ + OH → CO + HO_2 + CO_2 + H_2O	1.23E-11	Rickard and Pascoe (2009)
G4241	TrGC	$HCOCO_3H$ + OH → $HCOCO_3$ + H_2O	1.58E-11	Rickard and Pascoe (2009)
G4242	TrGC	$HOCH_2CH_2O_2$ → .6 $HOCH_2CH_2O$ + .2 $HOCH_2CHO$ + .2 ETHGLY	2.00E-12*RO2	Rickard and Pascoe (2009)
G4243	TrGNC	$HOCH_2CH_2O_2$ + NO → .24875 HO_2 + .4975 HCHO + .74625 $HOCH_2CH_2O$ + .995 NO_2 + .005 ETHOHNO3	KRO2NO	Rickard and Pascoe (2009)*
G4244	TrGC	$HOCH_2CH_2O_2$ + HO_2 → HYETHO2H	2.00E-13*EXP(1250./temp)	Rickard and Pascoe (2009)
G4245	TrGNC	ETHOHNO3 + OH → $HOCH_2CHO$ + NO_2 + H_2O	8.40E-13	Rickard and Pascoe (2009)
G4246a	TrGC	HYETHO2H + OH → $HOCH_2CH_2O_2$ + H_2O	0.6*k_CH3OOH_OH	Rickard and Pascoe (2009)*
G4246b	TrGC	HYETHO2H + OH → $HOCH_2CHO$ + OH + H_2O	1.38E-11	Rickard and Pascoe (2009)
G4247a	TrGC	$HOCH_2CH_2O$ → HO_2 + $HOCH_2CHO$	6.00E-14*EXP(-550./temp)*C(ind_O2)	Rickard and Pascoe (2009)

Table 1: Gas phase reactions (... continued)

#	labels	reaction	rate coefficient	reference
G4247b	TrGC	$HOCH_2CH_2O \rightarrow HO_2 + HCHO + HCHO$	9.50E13*EXP(-5988./temp)	Rickard and Pascoe (2009)
G4248	TrGC	$ETHGLY + OH \rightarrow HOCH_2CHO + HO_2 + H_2O$	7.70E-12	Rickard and Pascoe (2009)
G4300	TrGC	$C_3H_8 + OH \rightarrow .736\ iC_3H_7O_2 + .264\ C_2H_5O_2 + .264\ CO_2 + .264\ HO_2 + H_2O$	1.55E-17*temp*temp*EXP(-61./temp)	Rickard and Pascoe (2009)*
G4301	TrGC	$C_3H_6 + O_3 \rightarrow .28\ CH_3O_2 + .1\ CH_4 + .075\ CH_3COOH + .56\ CO + .075\ HCOOH + .09\ H_2O_2 + .28\ HO_2 + .2\ CO_2 + .545\ CH_3CHO + .545\ HCHO + .36\ OH$	6.5E-15*EXP(-1900./temp)	Sander et al. (2006)*
G4302	TrGC	$C_3H_6 + OH \rightarrow HYPROPO2$	k_3rd(temp,cair,8.E-27,3.5,3.E-11,0.,0.5)	Atkinson et al. (1999)
G4303	TrGNC	$C_3H_6 + NO_3 \rightarrow PRONO3BO2$	4.6E-13*EXP(-1155./temp)	Atkinson et al. (1999)
G4304	TrGC	$iC_3H_7O_2 + HO_2 \rightarrow iC_3H_7OOH$	1.9E-13*EXP(1300./temp)	Atkinson (1997)*
G4305	TrGNC	$iC_3H_7O_2 + NO \rightarrow .96\ CH_3COCH_3 + .96\ HO_2 + .96\ NO_2 + .04\ iC_3H_7ONO_2$	2.7E-12*EXP(360./temp)	Atkinson et al. (1999)
G4306	TrGC	$iC_3H_7O_2 \rightarrow CH_3COCH_3 + .8\ HO_2$	4.E-14*RO2	Rickard and Pascoe (2009)*
G4307	TrGC	$iC_3H_7OOH + OH \rightarrow .27\ iC_3H_7O_2 + .73\ CH_3COCH_3 + .73\ OH + H_2O$	1.66E-11 + 0.6*k_CH3OOH_OH	Rickard and Pascoe (2009)*
G4311	TrGC	$CH_3COCH_3 + OH \rightarrow CH_3COCH_2O_2 + H_2O$	(1.33E-13+3.82E-11*EXP(-2000./temp))	Sander et al. (2006)
G4312	TrGC	$CH_3COCH_2O_2 + HO_2 \rightarrow CH_3COCH_2O_2H$	8.6E-13*EXP(700./temp)	Tyndall et al. (2001)
G4313	TrGNC	$CH_3COCH_2O_2 + NO \rightarrow CH_3C(O)OO + HCHO + NO_2$	2.9E-12*EXP(300./temp)	Sander et al. (2006)
G4314	TrGC	$CH_3COCH_2O_2 \rightarrow .6\ CH_3C(O)OO + .6\ HCHO + .2\ MGLYOX + .2\ CH_3COCH_2OH$	7.5E-13*EXP(500./temp)*2.*RO2	Tyndall et al. (2001)
G4315a	TrGC	$CH_3COCH_2O_2H + OH \rightarrow CH_3COCH_2O_2 + H_2O$	0.6*k_CH3OOH_OH	see note
G4315b	TrGC	$CH_3COCH_2O_2H + OH \rightarrow MGLYOX + OH + H_2O$	8.39E-12	Rickard and Pascoe (2009)
G4316	TrGC	$CH_3COCH_2OH + OH \rightarrow MGLYOX + HO_2 + H_2O$	3.E-12	Atkinson et al. (1999)
G4317	TrGC	$MGLYOX + OH \rightarrow CH_3C(O)OO + CO$	8.4E-13*EXP(830./temp)	Tyndall et al. (1995)
G4320	TrGNC	$iC_3H_7ONO_2 + OH \rightarrow CH_3COCH_3 + NO_2$	6.2E-13*EXP(-230./temp)	Atkinson et al. (1999)
G4321	TrGNC	$CH_3COCH_2O_2 + NO_3 \rightarrow CH_3C(O)OO + HCHO + NO_2$	KRO2NO3	Rickard and Pascoe (2009)
G4322	TrGC	$HYPROPO2 \rightarrow CH_3CHO + HCHO + HO_2$	8.80E-13*RO2	Rickard and Pascoe (2009)
G4323	TrGC	$HYPROPO2 + HO_2 \rightarrow HYPROPO2H$	KRO2HO2*0.520	Rickard and Pascoe (2009)
G4324	TrGNC	$HYPROPO2 + NO \rightarrow CH_3CHO + HCHO + HO_2 + NO_2$	KRO2NO	Rickard and Pascoe (2009)
G4325	TrGNC	$HYPROPO2 + NO_3 \rightarrow CH_3CHO + HCHO + HO_2 + NO_2$	KRO2NO3	Rickard and Pascoe (2009)
G4326a	TrGC	$HYPROPO2H + OH \rightarrow HYPROPO2$	1.90E-12*EXP(190./temp)	Rickard and Pascoe (2009)
G4326b	TrGC	$HYPROPO2H + OH \rightarrow CH_3COCH_2OH + OH$	2.44E-11	Rickard and Pascoe (2009)
G4327	TrGNC	$PRONO3BO2 + HO_2 \rightarrow PR2O2HNO3$	KRO2HO2*0.520	Rickard and Pascoe (2009)
G4328	TrGNC	$PRONO3BO2 + NO \rightarrow NOA + HO_2 + NO_2$	KRO2NO	Rickard and Pascoe (2009)

Table 1: Gas phase reactions (... continued)

#	labels	reaction	rate coefficient	reference
G4329	TrGNC	PRONO3BO2 + NO_3 → NOA + HO_2 + NO_2	KRO2NO3	Rickard and Pascoe (2009)
G4330a	TrGNC	PR2O2HNO3 + OH → PRONO3BO2	1.90E-12*EXP(190./temp)	Rickard and Pascoe (2009)
G4330b	TrGNC	PR2O2HNO3 + OH → NOA + OH	3.47E-12	Rickard and Pascoe (2009)
G4331	TrGNC	MGLYOX + NO_3 → $CH_3C(O)OO$ + CO + HNO_3	KNO3AL*2.4	Rickard and Pascoe (2009)
G4332	TrGNC	NOA + OH → MGLYOX + NO_2	1.30E-13	Rickard and Pascoe (2009)
G4333	TrGC	HOCH2COCHO + OH → $HOCH_2CO_3$ + CO	1.44E-11	Rickard and Pascoe (2009)
G4334	TrGNC	HOCH2COCHO + NO_3 → $HOCH_2CO_3$ + CO + HNO_3	KNO3AL*2.4	Rickard and Pascoe (2009)
G4335	TrGC	HOCH2COCO2H + OH → $HOCH_2CO_3$ + CO_2	2.89E-12	Rickard and Pascoe (2009)
G4400	TrGC	nC_4H_{10} + OH → $LC_4H_9O_2$ + H_2O	1.81E-17*temp*temp*EXP(114./temp)	Atkinson (2003)*
G4401	TrGC	$LC_4H_9O_2$ → 0.254 CO_2 + 0.5552 MEK + 0.5552 HO_2 + 0.3178 CH_3CHO + 0.4448 $C_2H_5O_2$	2.5E-13*RO2	Rickard and Pascoe (2009)*
G4402	TrGC	$LC_4H_9O_2$ + HO_2 → LC_4H_9OOH	KRO2HO2*0.625	Rickard and Pascoe (2009)
G4403	TrGNC	$LC_4H_9O_2$ + NO → 0.9172 NO_2 + 0.233 CO_2 + 0.5092 MEK + 0.5092 HO_2 + 0.2915 CH_3CHO + 0.408 $C_2H_5O_2$ + 0.0828 LC4H9NO3	KRO2NO	Rickard and Pascoe (2009)*
G4404	TrGC	LC_4H_9OOH + OH → 0.2285796 $LC_4H_9O_2$ + 0.7117253 MEK + 0.1193902 CO_2 + 0.0596951 $C_2H_5O_2$ + 0.7714204 OH + H_2O	2.636E-11	Rickard and Pascoe (2009)*
G4405	TrGC	MVK + O_3 → 0.28 $CH_3C(O)OO$ + 0.56 CO + 0.225 LCARBON + 0.075 HCOOH + 0.09 H_2O_2 + 0.28 HO_2 + 0.1 CO_2 + 0.1 CH_3CHO + 0.645 HCHO + 0.36 OH + 0.545 MGLYOX	7.51E-16*EXP(-1521./temp)	Rickard and Pascoe (2009)
G4406	TrGC	MVK + OH → LHMVKABO2	4.13E-12*EXP(452./temp)	Rickard and Pascoe (2009)
G4413	TrGC	MEK + OH → LMEKO2 + H_2O	3.24E-18*temp*temp*EXP(414./temp)	Rickard and Pascoe (2009)*
G4414	TrGC	LMEKO2 + HO_2 → LMEKOOH	KRO2HO2*0.625	Rickard and Pascoe (2009)
G4415	TrGNC	LMEKO2 + NO → 0.538 HCHO + 0.538 CO_2 + 0.459 $HOCH_2CH_2O_2$ + 0.079 $C_2H_5O_2$ + 0.462 $CH_3C(O)OO$ + 0.462 CH_3CHO + NO_2	KRO2NO	Rickard and Pascoe (2009)*
G4416	TrGC	LMEKOOH + OH → 0.40851 $CH_3COCH_2O_2$ + 0.350196 BIACET + 0.807212 OH + 0.048506 $C_2H_5O_2$ + 0.505522 CO_2 + 0.192788 LMEKO2 + H_2O	3.786E-11	Rickard and Pascoe (2009)*
G4417	TrGNC	LC4H9NO3 + OH → 0.91423 MEK + 0.08577 $C_2H_5O_2$ + 0.17154 CO_2 + NO_2 + H_2O	9.598E-13	Rickard and Pascoe (2009)*
G4418	TrGNC	MPAN + OH → CH_3COCH_2OH + CO + NO_2	3.2E-11	Orlando et al. (2002)
G4419	TrGNC	MPAN → MACO3 + NO_2	k_PAN_M	see note

Table 1: Gas phase reactions (... continued)

#	labels	reaction	rate coefficient	reference
G4420	TrGC	LMEKO2 → 0.538 HCHO + 0.538 CO_2 + 0.459 $HOCH_2CH_2O_2$ + 0.079 $C_2H_5O_2$ + 0.462 $CH_3C(O)OO$ + 0.462 CH_3CHO	1.483E-12*RO2	Rickard and Pascoe (2009)*
G4421	TrGC	MACR + OH → .57 MACO3 + .43 MACRO2	1.86E-11*EXP(175./temp)	Rickard and Pascoe (2009)
G4422	TrGC	MACR + O_3 → .59 MGLYOX + .41 $CH_3C(O)OO$ + .03375 HCOOH + .55625 HCHO + .82 CO + .12375 H_2O_2 + .41 HO_2 + .82 OH	1.36E-15*EXP(-2112./temp)	Rickard and Pascoe (2009)
G4423	TrGNC	MACR + NO_3 → MACO3 + HNO_3	KNO3AL*2.0	Rickard and Pascoe (2009)
G4424	TrGC	MACO3 → .7 $CH_3C(O)OO$ + .7 HCHO + .7 CO_2 + .3 MACO2H	1.00E-11*RO2	Rickard and Pascoe (2009)
G4425	TrGC	MACO3 + HO_2 → .71 MACO3H + .29 MACO2H + .29 O_3	KAPHO2	Rickard and Pascoe (2009)
G4426	TrGNC	MACO3 + NO → $CH_3C(O)OO$ + HCHO + NO_2 + CO_2	8.70E-12*EXP(290./temp)	Rickard and Pascoe (2009)
G4427	TrGNC	MACO3 + NO_2 → MPAN	k_CH3CO3_NO2	Rickard and Pascoe (2009)
G4428	TrGNC	MACO3 + NO_3 → $CH_3C(O)OO$ + HCHO + NO_2 + CO_2	KRO2NO3*1.60	Rickard and Pascoe (2009)
G4429	TrGC	MACRO2 → .7 CH_3COCH_2OH + .7 HCHO + .7 HO_2 + .3 MACROH	9.20E-14*RO2	Rickard and Pascoe (2009)
G4430	TrGC	MACRO2 + HO_2 → MACROOH	KRO2HO2*0.625	Rickard and Pascoe (2009)
G4431	TrGNC	MACRO2 + NO → CH_3COCH_2OH + HCHO + HO_2 + NO_2	KRO2NO	Rickard and Pascoe (2009)
G4432	TrGNC	MACRO2 + NO_3 → CH_3COCH_2OH + HCHO + HO_2 + NO_2	KRO2NO3	Rickard and Pascoe (2009)
G4433	TrGC	MACROOH + OH → MACRO2	2.82E-11	Rickard and Pascoe (2009)
G4434	TrGC	MACROH + OH → CH_3COCH_2OH + HCHO + HO_2	2.46E-11	Rickard and Pascoe (2009)
G4435	TrGC	MACO2H + OH → $CH_3C(O)OO$ + HCHO + CO_2	1.51E-11	Rickard and Pascoe (2009)
G4436	TrGC	MACO3H + OH → MACO3	1.87E-11	Rickard and Pascoe (2009)
G4437	TrGC	LHMVKABO2 → 0.06 CO2H3CHO + 0.18 HO_2 + 0.18 HCHO + 0.18 MGLYOX + 0.42 $CH_3C(O)OO$ + .42 $HOCH_2CHO$+ .2 HO12CO3C4 + .14 BIACETOH	(.3*2.00E-12 + .7*8.80E-13)*RO2	Rickard and Pascoe (2009)*
G4438	TrGC	LHMVKABO2 + HO_2 → LHMVKABOOH	KRO2HO2*0.625	Rickard and Pascoe (2009)
G4439	TrGNC	LHMVKABO2 + NO → .3 MGLYOX + .7 $HOCH_2CHO$ + .7 $CH_3C(O)OO$ + .3 HCHO + .3 HO_2 + NO_2	KRO2NO	Rickard and Pascoe (2009)*
G4440	TrGNC	LHMVKABO2 + NO_3 → .3 MGLYOX + .7 $HOCH_2CHO$ + .7 $CH_3C(O)OO$ + .3 HCHO + .3 HO_2 + NO_2	KRO2NO3	Rickard and Pascoe (2009)*

Table 1: Gas phase reactions (... continued)

#	labels	reaction	rate coefficient	reference
G4441	TrGC	LHMVKABOOH + OH → .3 CO2H3CHO + .7 BIACETOH + OH	4.496E-11	Rickard and Pascoe (2009)*
G4442	TrGC	MVKOH + OH → LMVKOHABO2	4.60E-12*EXP(452./temp)	Rickard and Pascoe (2009)
G4443	TrGC	MVKOH + O_3 → 0.56 CO + 0.545 HOCH2COCHO + 0.075 HOCH2COCO2H + 0.075 HCOOH + 0.09 H_2O_2 + 0.28 $HOCH_2CO_3$ + 0.28 HO_2 + 0.2 CO_2 + 0.545 HCHO + 0.36 OH + 0.1 $HOCH_2CHO$	7.51E-16*EXP(-1521./temp)	Rickard and Pascoe (2009)
G4444	TrGC	LMVKOHABO2 → .7 $HOCH_2CHO$ + .7 $HOCH_2CO_3$ + .3 HOCH2COCHO + .3 HCHO + .3 HO_2	(0.3*2.00E-12+0.7*8.80E-13)*RO2	Rickard and Pascoe (2009)*
G4445	TrGC	LMVKOHABO2 + HO_2 → LMVKOHABOOH	KRO2HO2*0.625	Rickard and Pascoe (2009)
G4446	TrGNC	LMVKOHABO2 + NO → .3 HOCH2COCHO + .3 HCHO + .3 HO_2 + .7 $HOCH_2CHO$ + .7 $HOCH_2CO_3$ + NO_2	KRO2NO	Rickard and Pascoe (2009)*
G4447	TrGNC	LMVKOHABO2 + NO_3 → .3 HOCH2COCHO + .3 HCHO + .3 HO_2 + .7 $HOCH_2CHO$ + .7 $HOCH_2CO_3$ + NO_2	KRO2NO3	Rickard and Pascoe (2009)*
G4448	TrGC	LMVKOHABOOH + OH → .7 HO12CO3C4 + .3 CO2H3CHO + OH	5.98E-11	Rickard and Pascoe (2009)*
G4449	TrGC	CO2H3CHO + OH → CO2H3CO3	2.45E-11	Rickard and Pascoe (2009)
G4450	TrGNC	CO2H3CHO + NO_3 → CO2H3CO3 + HNO_3	KNO3AL*4.0	Rickard and Pascoe (2009)
G4451	TrGC	CO2H3CO3 → MGLYOX + HO_2 + CO_2	1.00E-11*RO2	Rickard and Pascoe (2009)
G4452	TrGC	CO2H3CO3 + HO_2 → CO2H3CO3H	KAPHO2	Rickard and Pascoe (2009)
G4453	TrGNC	CO2H3CO3 + NO → MGLYOX + HO_2 + NO_2 + CO_2	KAPNO	Rickard and Pascoe (2009)
G4454	TrGNC	CO2H3CO3 + NO_3 → MGLYOX + HO_2 + NO_2 + CO_2	KRO2NO3*1.60	Rickard and Pascoe (2009)
G4455	TrGC	CO2H3CO3H + OH → CO2H3CO3	7.34E-12	Rickard and Pascoe (2009)
G4456	TrGC	HO12CO3C4 + OH → BIACETOH + HO_2	1.88E-11	Rickard and Pascoe (2009)
G4500	TrGC	C_5H_8 + O_3 → .051 CH_3O_2 + .1575 $CH_3C(O)OO$ + .054 LHMVKABO2 + .522 CO + .06875 HCOOH + .11 H_2O_2 + .32475 MACR + .1275 C_3H_6 + .2625 HO_2 + .255 CO_2 + .74975 HCHO + .04125 MACO2H + .27 OH + .244 MVK	7.86E-15*EXP(-1913./temp)	Rickard and Pascoe (2009)
G4501	TrGC	C_5H_8 + OH → .25 LISOPACO2 + .491 ISOPBO2 + .259 ISOPDO2	2.54E-11*EXP(410./temp)	Atkinson (1997)
G4509	TrGNC	C_5H_8 + NO_3 → NISOPO2	3.03E-12*EXP(-446./temp)	Rickard and Pascoe (2009)
G4510	TrGC	LISOPACO2 → .9 LHC4ACCHO + .8 HO_2 + .1 ISOPAOH	2.4E-12*RO2	Rickard and Pascoe (2009)
G4511	TrGC	LISOPACO2 + HO_2 → LISOPACOOH	0.706*KRO2HO2	Rickard and Pascoe (2009)
G4512	TrGNC	LISOPACO2 + NO → .892 LHC4ACCHO + .892 HO_2 + .892 NO_2 + .108 LISOPACNO3	KRO2NO	Rickard and Pascoe (2009)

Table 1: Gas phase reactions (... continued)

#	labels	reaction	rate coefficient	reference
G4513	TrGNC	LISOPACO2 + NO_3 → LHC4ACCHO + HO_2 + NO_2	KRO2NO3	Rickard and Pascoe (2009)
G4514	TrGC	LISOPACOOH + OH → LHC4ACCHO + OH	1.07E-10	Rickard and Pascoe (2009)
G4515	TrGC	ISOPAOH + OH → LHC4ACCHO + HO_2	9.30E-11	Rickard and Pascoe (2009)
G4516	TrGNC	LISOPACNO3 + OH → LHC4ACCHO + NO_2	8.91E-11	Rickard and Pascoe (2009)
G4517	TrGC	ISOPBO2 → .6 MVK + .2 MVKOH + .6 HCHO + .6 HO_2 + .2 CH_3O_2 + .2 ISOPBOH	8.E-13*RO2	Rickard and Pascoe (2009)
G4518	TrGC	ISOPBO2 + HO_2 → ISOPBOOH	0.706*KRO2HO2	Rickard and Pascoe (2009)
G4519	TrGNC	ISOPBO2 + NO → .696 MVK + .232 MVKOH + .696 HCHO + .696 HO_2 + .232 CH_3O_2 + .928 NO_2 + .072 ISOPBNO3	KRO2NO	Rickard and Pascoe (2009)
G4520	TrGNC	ISOPBO2 + NO_3 → .75 MVK + .25 MVKOH + .75 HCHO + .75 HO_2 + .25 CH_3O_2 + NO_2	KRO2NO3	Rickard and Pascoe (2009)
G4521	TrGC	ISOPBOOH + OH → ISOPBO2	4.2E-11	Rickard and Pascoe (2009)
G4522	TrGC	ISOPBOH + OH → .75 MVK + .25 MVKOH + .75 HCHO + .75 HO_2 + .25 CH_3O_2	3.85E-11	Rickard and Pascoe (2009)
G4523	TrGNC	ISOPBNO3 + OH → MVK + HCHO + NO_2	3.55E-11	Rickard and Pascoe (2009)
G4524	TrGC	ISOPDO2 → .8 MACR + .8 HCHO + .8 HO_2 + .1 HCOC5 + .1 ISOPDOH	2.9E-12*RO2	Rickard and Pascoe (2009)
G4525	TrGC	ISOPDO2 + HO_2 → ISOPDOOH	0.706*KRO2HO2	Rickard and Pascoe (2009)
G4526	TrGNC	ISOPDO2 + NO → .855 MACR + .855 HCHO + .855 HO_2 + .855 NO_2 + .145 ISOPDNO3	KRO2NO	Rickard and Pascoe (2009)
G4527	TrGNC	ISOPDO2 + NO_3 → MACR + HCHO + HO_2 + NO_2	KRO2NO3	Rickard and Pascoe (2009)
G4528	TrGC	ISOPDOOH + OH → HCOC5 + OH	1.07E-10	Rickard and Pascoe (2009)
G4529	TrGC	ISOPDOH + OH → HCOC5 + HO_2	7.38E-11	Rickard and Pascoe (2009)
G4530	TrGNC	ISOPDNO3 + OH → HCOC5 + NO_2	6.1E-11	Rickard and Pascoe (2009)
G4531	TrGNC	NISOPO2 → .8 NC4CHO + .6 HO_2 + .2 LISOPACNO3	1.3E-12*RO2	Rickard and Pascoe (2009)
G4532	TrGNC	NISOPO2 + HO_2 → NISOPOOH	.706*KRO2HO2	Rickard and Pascoe (2009)
G4533	TrGNC	NISOPO2 + NO → NC4CHO + HO_2 + NO_2	KRO2NO	Rickard and Pascoe (2009)
G4534	TrGNC	NISOPO2 + NO_3 → NC4CHO + HO_2 + NO_2	KRO2NO3	Rickard and Pascoe (2009)
G4535	TrGNC	NISOPOOH + OH → NC4CHO + OH	1.03E-10	Rickard and Pascoe (2009)
G4536	TrGNC	NC4CHO + OH → LNISO3	4.16E-11	Rickard and Pascoe (2009)
G4537	TrGNC	NC4CHO + O_3 → .445 NO_2 + .89 CO + .075625 H_2O_2 + .034375 $HCOCO_2H$ + .555 NOA + .445 HO_2 + .520625 GLYOX + .89 OH + .445 MGLYOX	2.40E-17	Rickard and Pascoe (2009)
G4538	TrGNC	NC4CHO + NO_3 → LNISO3 + HNO_3	KNO3AL*4.25	Rickard and Pascoe (2009)

Table 1: Gas phase reactions (... continued)

#	labels	reaction	rate coefficient	reference
G4539	TrGNC	LNISO3 + HO_2 → LNISOOH	.5*.706*KRO2HO2 + .5*KAPHO2	Rickard and Pascoe (2009)
G4540	TrGNC	LNISO3 + NO → NOA + .5 GLYOX + .5 CO + HO_2 + NO_2 + .5 CO_2	.5*KAPNO +.5*KRO2NO	Rickard and Pascoe (2009)
G4541	TrGNC	LNISO3 + NO_3 → NOA + .5 GLYOX + .5 CO + HO_2 + NO_2 + .5 CO_2	1.3*KRO2NO3	Rickard and Pascoe (2009)
G4542	TrGNC	LNISOOH + OH → LNISO3	2.65E-11	Rickard and Pascoe (2009)
G4543	TrGC	LHC4ACCHO + OH → .52 LC578O2 + .48 LHC4ACCO3	4.52E-11	Rickard and Pascoe (2009)
G4544	TrGC	LHC4ACCHO + O_3 → .2225 $CH_3C(O)OO$ + .89 CO + .0171875 $HOCH_2CO_2H$ + .075625 H_2O_2 + .0171875 $HCOCO_2H$ + .2775 CH_3COCH_2OH + .6675 HO_2 + .2603125 GLYOX + .2225 HCHO + .89 OH + .2603125 $HOCH_2CHO$ + .5 MGLYOX	2.40E-17	Rickard and Pascoe (2009)
G4545	TrGNC	LHC4ACCHO + NO_3 → LHC4ACCO3 + HNO_3	KNO3AL*4.25	Rickard and Pascoe (2009)
G4546	TrGC	LC578O2 → .5 CH_3COCH_2OH + .5 MGLYOX + .5 GLYOX + .5 $HOCH_2CHO$ + HO_2	9.20E-14*RO2	Rickard and Pascoe (2009)
G4547	TrGC	LC578O2 + HO_2 → LC578OOH	KRO2HO2*0.706	Rickard and Pascoe (2009)
G4548	TrGNC	LC578O2 + NO → .5 CH_3COCH_2OH + .5 MGLYOX + .5 GLYOX + .5 $HOCH_2CHO$ + HO_2 + NO_2	KRO2NO	Rickard and Pascoe (2009)
G4549	TrGNC	LC578O2 + NO_3 → .5 CH_3COCH_2OH + .5 MGLYOX + .5 GLYOX + .5 $HOCH_2CHO$ + HO_2 + NO_2	KRO2NO3	Rickard and Pascoe (2009)
G4550	TrGC	LC578OOH + OH → LC578O2	3.16E-11	Rickard and Pascoe (2009)
G4551	TrGC	LHC4ACCO3 → .3 LHC4ACCO2H + .35 CH_3COCH_2OH + .35 $HOCH_2CHO$ + .35 $CH_3C(O)OO$ + .35 CO + .35 HO_2 + .7 CO_2	1.00E-11*RO2	Rickard and Pascoe (2009)
G4552	TrGC	LHC4ACCO3 + HO_2 → .71 LHC4ACCO3H + .29 LHC4ACCO2H + .29 O_3	KAPHO2	Rickard and Pascoe (2009)
G4553	TrGNC	LHC4ACCO3 + NO → .5 CH_3COCH_2OH + .5 $HOCH_2CHO$ + .5 $CH_3C(O)OO$ + .5 CO + .5 HO_2 + NO_2 + CO_2	KAPNO	Rickard and Pascoe (2009)
G4554	TrGNC	LHC4ACCO3 + NO_2 → LC5PAN1719	k_CH3CO3_NO2	Rickard and Pascoe (2009)
G4555	TrGNC	LHC4ACCO3 + NO_3 → .5 CH_3COCH_2OH + .5 $HOCH_2CHO$ + .5 $CH_3C(O)OO$ + .5 CO + .5 HO_2 + NO_2 + CO_2	1.6*KRO2NO3	Rickard and Pascoe (2009)
G4556	TrGC	LHC4ACCO2H + OH → .5 CH_3COCH_2OH + .5 $HOCH_2CHO$ + .5 $CH_3C(O)OO$ + .5 CO + .5 HO_2 + CO_2	2.52E-11	Rickard and Pascoe (2009)

Table 1: Gas phase reactions (... continued)

#	labels	reaction	rate coefficient	reference
G4557	TrGC	LHC4ACCO3H + OH → LHC4ACCO3	2.88E-11	Rickard and Pascoe (2009)
G4558	TrGNC	LC5PAN1719 → LHC4ACCO3 + NO_2	k_PAN_M	Rickard and Pascoe (2009)
G4559	TrGNC	LC5PAN1719 + OH → .5 MACROH + .5 HO12CO3C4 + CO + NO_2	2.52E-11	Rickard and Pascoe (2009)
G4560	TrGC	HCOC5 + OH → C59O2	3.81E-11	Rickard and Pascoe (2009)
G4561	TrGC	C59O2 → CH_3COCH_2OH + $HOCH_2CO_3$	9.20E-14*RO2	Rickard and Pascoe (2009)
G4562	TrGC	C59O2 + HO_2 → C59OOH	KRO2HO2*0.706	Rickard and Pascoe (2009)
G4563	TrGNC	C59O2 + NO → CH_3COCH_2OH + $HOCH_2CO_3$ + NO_2	KRO2NO	Rickard and Pascoe (2009)
G4564	TrGNC	C59O2 + NO_3 → CH_3COCH_2OH + $HOCH_2CO_3$ + NO_2	KRO2NO3	Rickard and Pascoe (2009)
G4565	TrGC	C59OOH + OH → C59O2	9.7E-12	Rickard and Pascoe (2009)
G6100	StTrGCl	Cl + O_3 → ClO + O_2	2.8E-11*EXP(-250./temp)	Atkinson et al. (2007)
G6101	StGCl	ClO + $O(^3P)$ → Cl + O_2	2.5E-11*EXP(110./temp)	Atkinson et al. (2007)
G6102a	StTrGCl	ClO + ClO → Cl_2 + O_2	1.0E-12*EXP(-1590./temp)	Atkinson et al. (2007)
G6102b	StTrGCl	ClO + ClO → 2 Cl + O_2	3.0E-11*EXP(-2450./temp)	Atkinson et al. (2007)
G6102c	StTrGCl	ClO + ClO → Cl + OClO	3.5E-13*EXP(-1370./temp)	Atkinson et al. (2007)
G6102d	StTrGCl	ClO + ClO → Cl_2O_2	k_ClO_ClO	Atkinson et al. (2007)
G6103	StTrGCl	Cl_2O_2 → ClO + ClO	k_ClO_ClO/(9.3E-28*EXP(8835./temp))	Atkinson et al. (2007), Sander et al. (2006)*
G6200	StGCl	Cl + H_2 → HCl + H	3.9E-11*EXP(-2310./temp)	Atkinson et al. (2007)
G6201a	StGCl	Cl + HO_2 → HCl + O_2	4.4E-11-7.5E-11*EXP(-620./temp)	Atkinson et al. (2007)
G6201b	StGCl	Cl + HO_2 → ClO + OH	7.5E-11*EXP(-620./temp)	Atkinson et al. (2007)
G6202	StTrGCl	Cl + H_2O_2 → HCl + HO_2	1.1E-11*EXP(-980./temp)	Atkinson et al. (2007)
G6203	StGCl	ClO + OH → .94 Cl + .94 HO_2 + .06 HCl + .06 O_2	7.3E-12*EXP(300./temp)	Atkinson et al. (2007)
G6204	StTrGCl	ClO + HO_2 → HOCl + O_2	2.2E-12*EXP(340./temp)	Atkinson et al. (2007)
G6205	StTrGCl	HCl + OH → Cl + H_2O	1.7E-12*EXP(-230./temp)	Atkinson et al. (2007)
G6206	StGCl	HOCl+ OH → ClO + H_2O	3.0E-12*EXP(-500./temp)	Sander et al. (2006)
G6300	StTrGNCl	ClO + NO → NO_2 + Cl	6.2E-12*EXP(295./temp)	Atkinson et al. (2007)
G6301	StTrGNCl	ClO + NO_2 → $ClNO_3$	k_3rd_iupac(temp,cair,1.6E-31,3.4, 7.E-11,0.,0.4)	Atkinson et al. (2007)
G6302	TrGCl	$ClNO_3$ → ClO + NO_2	6.918E-7*exp(-10909./temp)*cair	Anderson and Fahey (1990)
G6303	StGNCl	$ClNO_3$ + $O(^3P)$ → ClO + NO_3	4.5E-12*EXP(-900./temp)	Atkinson et al. (2007)
G6304	StTrGNCl	$ClNO_3$ + Cl → Cl_2 + NO_3	6.2E-12*EXP(145./temp)	Atkinson et al. (2007)
G6400	StTrGCl	Cl + CH_4 → HCl + CH_3O_2	6.6E-12*EXP(-1240./temp)	Atkinson et al. (2006)
G6401	StTrGCl	Cl + HCHO → HCl + CO + HO_2	8.1E-11*EXP(-34./temp)	Atkinson et al. (2006)
G6402	StTrGCl	Cl + CH_3OOH → HCHO + HCl + OH	5.9E-11	Atkinson et al. (2006)*

Table 1: Gas phase reactions (... continued)

#	labels	reaction	rate coefficient	reference
G6403	StTrGCl	$ClO + CH_3O_2 \rightarrow HO_2 + Cl + HCHO$	3.3E-12*EXP(-115./temp)	Sander et al. (2006)
G6404	StGCl	$CCl_4 + O(^1D) \rightarrow ClO + 3\ Cl$	3.3E-10	Sander et al. (2006)
G6405	StGCl	$CH_3Cl + O(^1D) \rightarrow OH + Cl$	1.65E-10	see note
G6406	StGCl	$CH_3Cl + OH \rightarrow H_2O + Cl$	2.4E-12*EXP(-1250./temp)	Sander et al. (2006)
G6407	StGCCl	$CH_3CCl_3 + O(^1D) \rightarrow OH + 3\ Cl$	3.E-10	see note
G6408	StTrGCCl	$CH_3CCl_3 + OH \rightarrow H_2O + 3\ Cl$	1.64E-12*EXP(-1520./temp)	Sander et al. (2006)
G6409	TrGCCl	$Cl + C_2H_4 \rightarrow HOCH_2CH_2O_2 + HCl$	k_3rd_iupac(temp,cair,1.85E-29,3.3, 6.0E-10,0.0,0.4)	Atkinson et al. (2006)*
G6410	TrGCCl	$Cl + CH_3CHO \rightarrow HCl + CH_3C(O)OO$	8.0e-11	Atkinson et al. (2006)
G6411	TrGCCl	$C_2H_2 + Cl \rightarrow CH_3O_2 + HCl$	k_3rd_iupac(temp,cair,6.1e-30,3.0, 2.0e-10,0.,0.6)	Atkinson et al. (2006)
G6412	TrGCCl	$C_2H_6 + Cl \rightarrow CH_3O_2 + HCl$	8.3E-11*EXP(-100./temp)	Atkinson et al. (2006)
G6500	StGFCl	$CF_2Cl_2 + O(^1D) \rightarrow ClO + Cl$	1.4E-10	Sander et al. (2006)
G6501	StGFCl	$CFCl_3 + O(^1D) \rightarrow ClO + 2\ Cl$	2.3E-10	Sander et al. (2006)
G7100	StTrGBr	$Br + O_3 \rightarrow BrO + O_2$	1.7E-11*EXP(-800./temp)	Atkinson et al. (2007)
G7101	StGBr	$BrO + O(^3P) \rightarrow Br + O_2$	1.9E-11*EXP(230./temp)	Atkinson et al. (2007)
G7102a	StTrGBr	$BrO + BrO \rightarrow 2\ Br + O_2$	2.7E-12	Atkinson et al. (2007)
G7102b	StTrGBr	$BrO + BrO \rightarrow Br_2 + O_2$	2.9E-14*EXP(840./temp)	Atkinson et al. (2007)
G7200	StTrGBr	$Br + HO_2 \rightarrow HBr + O_2$	7.7E-12*EXP(-450./temp)	Atkinson et al. (2007)
G7201	StTrGBr	$BrO + HO_2 \rightarrow HOBr + O_2$	4.5E-12*EXP(500./temp)	Atkinson et al. (2007)
G7202	StTrGBr	$HBr + OH \rightarrow Br + H_2O$	6.7E-12*EXP(155./temp)	Atkinson et al. (2007)
G7203	StGBr	$HOBr + O(^3P) \rightarrow OH + BrO$	1.2E-10*EXP(-430./temp)	Atkinson et al. (2007)
G7204	StTrGBr	$Br_2 + OH \rightarrow HOBr + Br$	2.0E-11*EXP(240./temp)	Atkinson et al. (2007)
G7300	TrGBr	$Br + BrNO_3 \rightarrow Br_2 + NO_3$	4.9E-11	Orlando and Tyndall (1996)
G7301	StTrGNBr	$BrO + NO \rightarrow Br + NO_2$	8.7E-12*EXP(260./temp)	Atkinson et al. (2007)
G7302	StTrGNBr	$BrO + NO_2 \rightarrow BrNO_3$	k_BrO_NO2	Atkinson et al. (2007)*
G7303	TrGBr	$BrNO_3 \rightarrow BrO + NO_2$	k_BrO_NO2/(5.44E-9*exp(14192./temp) *1.E6*R_gas*temp/(atm2Pa*N_A))	Orlando and Tyndall (1996), Atkinson et al. (2007)*
G7400	StTrGBr	$Br + HCHO \rightarrow HBr + CO + HO_2$	7.7E-12*EXP(-580./temp)	Atkinson et al. (2006)
G7401	TrGBr	$Br + CH_3OOH \rightarrow CH_3O_2 + HBr$	2.6E-12*EXP(-1600./temp)	Kondo and Benson (1984)
G7402a	TrGBr	$BrO + CH_3O_2 \rightarrow HOBr + HCHO$	G7402a_yield*5.7E-12	Aranda et al. (1997)
G7402b	TrGBr	$BrO + CH_3O_2 \rightarrow Br + HCHO + HO_2$	(1.-G7402a_yield)*5.7E-12	Aranda et al. (1997)
G7403	StTrGBr	$CH_3Br + OH \rightarrow H_2O + Br$	2.35E-12*EXP(-1300./temp)	Sander et al. (2006)
G7404	TrGCBr	$Br + C_2H_4 \rightarrow HOCH_2CH_2O_2 + HBr$	2.8E-13*EXP(224./temp)/(1.+ 1.13E24*EXP(-3200./temp)/C(ind_O2))	Atkinson et al. (2006)*

Table 1: Gas phase reactions (... continued)

#	labels	reaction	rate coefficient	reference
G7405	TrGCBr	$Br + CH_3CHO \rightarrow HBr + CH_3C(O)OO$	1.8e-11*EXP(-460./temp)	Atkinson et al. (2006)
G7406	TrGCBr	$Br + C_2H_2 \rightarrow CH_3O_2 + HBr$	6.35e-15*EXP(440./temp)	Atkinson et al. (2006)
G7407	TrGBr	$CHBr_3 + OH \rightarrow H_2O + 3\ Br$	1.35E-12*EXP(-600./temp)	Sander et al. (2006)*
G7408	TrGBr	$CH_2Br_2 + OH \rightarrow H_2O + 2\ Br$	2.0E-12*EXP(-840./temp)	Sander et al. (2006)*
G7600	TrGClBr	$Br + BrCl \rightarrow Br_2 + Cl$	3.32E-15	Manion et al. (2010)
G7601	TrGClBr	$Br + Cl_2 \rightarrow BrCl + Cl$	1.10E-15	Dolson and Leone (1987)
G7602	TrGClBr	$Br_2 + Cl \rightarrow BrCl + Br$	2.3E-10*EXP(135./temp)	Bedjanian et al. (1998)
G7603a	StTrGClBr	$BrO + ClO \rightarrow Br + OClO$	1.6E-12*EXP(430./temp)	Atkinson et al. (2007)
G7603b	StTrGClBr	$BrO + ClO \rightarrow Br + Cl + O_2$	2.9E-12*EXP(220./temp)	Atkinson et al. (2007)
G7603c	StTrGClBr	$BrO + ClO \rightarrow BrCl + O_2$	5.8E-13*EXP(170./temp)	Atkinson et al. (2007)
G7604	TrGClBr	$BrCl + Cl \rightarrow Br + Cl_2$	1.45E-11	Clyne and Cruse (1972)
G7605	TrGClBr	$CHCl_2Br + OH \rightarrow H_2O + Br$	2.0E-12*EXP(-840./temp)	see note
G7606	TrGClBr	$CHClBr_2 + OH \rightarrow H_2O + 2\ Br$	2.0E-12*EXP(-840./temp)	see note
G7607	TrGClBr	$CH_2ClBr + OH \rightarrow H_2O + Br$	2.4E-12*EXP(-920./temp)	Sander et al. (2006)*
G8100	TrGI	$I + O_3 \rightarrow IO + O_2$	2.1E-11*EXP(-830./temp)	Atkinson et al. (2007)
G8102	TrGI	$OIO + OIO \rightarrow I_{part}$	5.E-11	von Glasow et al. (2002)*
G8103	TrGI	$IO + IO \rightarrow .38\ OIO + 1.62\ I + .62\ O_2$	5.4E-11*EXP(180./temp)	Atkinson et al. (2007)*
G8200	TrGI	$I + HO_2 \rightarrow HI + O_2$	1.5E-11*EXP(-1090./temp)	Atkinson et al. (2007)
G8201	TrGI	$IO + HO_2 \rightarrow HOI + O_2$	1.4E-11*EXP(540./temp)	Atkinson et al. (2007)
G8202	TrGI	$HI + OH \rightarrow I + H_2O$	1.6E-11*EXP(440./temp)	Atkinson et al. (2007)
G8203	TrGI	$OIO + OH \rightarrow HIO_3$	2.2E-10*EXP(243./temp)	Plane et al. (2006)
G8204	TrGI	$I_2 + OH \rightarrow HOI + I$	2.1E-10	Atkinson et al. (2007)
G8300	TrGI	$I + NO_2 \rightarrow INO_2$	k_I_NO2	Atkinson et al. (2007)*
G8301	TrGI	$I + NO_3 \rightarrow IO + NO_2$	1.E-10	Dillon et al. (2008)
G8302	TrGI	$IO + NO \rightarrow I + NO_2$	7.15E-12*EXP(300./temp)	Atkinson et al. (2007)
G8303	TrGI	$IO + NO_2 \rightarrow INO_3$	k_3rd_iupac(temp,cair,7.7E-31,5., 1.6E-11,0.,0.4)	Atkinson et al. (2007)
G8304	TrGI	$OIO + NO \rightarrow NO_2 + IO$	1.1E-12*EXP(542./temp)	Atkinson et al. (2007)
G8305	TrGI	$INO_2 \rightarrow I + NO_2$	k_I_NO2/(3.7E-7*exp(9568./temp) *1.E6*R_gas*temp/(atm2Pa*N_A))	van den Bergh and Troe (1976), Atkinson et al. (2007)*
G8306	TrGI	$INO_3 \rightarrow IO + NO_2$	0.	see note
G8307	TrGI	$I_2 + NO_3 \rightarrow I + INO_3$	1.5E-12	Atkinson et al. (2007)
G8308	TrGI	$IO + NO_3 \rightarrow OIO + NO_2$	9.E-12	Dillon et al. (2008)
G8400	TrGI	$C_3H_7I + OH \rightarrow CH_3O_2 + I$	1.22E-12	Carl and Crowley (2001)

Table 1: Gas phase reactions (... continued)

#	labels	reaction	rate coefficient	reference
G8401	TrGI	CH_3O_2 + IO → .4 I + .6 OIO + HCHO + HO_2	2.E-12	Dillon et al. (2006b), Bale et al. (2005)*
G8402	TrGI	CH_3I + NO_3 → HNO_3 + HCHO + IO	3.4E-17	Wayne et al. (1991)*
G8600	TrGClI	IO + ClO → .2 ICl + .25 Cl + .55 OClO + .8 I + .45 O_2	4.7E-12*EXP(280./temp)	Atkinson et al. (2007)
G8700	TrGBrI	I + BrO → IO + Br	1.2E-11	Sander et al. (2006)
G8701	TrGBrI	IO + BrO → Br + .8 OIO + .2 I + .2 O_2	1.5E-11*EXP(510./temp)	Atkinson et al. (2007)*
G9200	StTrGS	SO_2 + OH → H_2SO_4 + HO_2	k_3rd(temp,cair,3.3E-31,4.3,1.6E-12,0.,0.6)	Sander et al. (2006)
G9400a	TrGS	DMS + OH → CH_3SO_2 + HCHO	1.13E-11*EXP(-253./temp)	Atkinson et al. (2004)*
G9400b	TrGS	DMS + OH → DMSO + HO_2	k_DMS_OH	Atkinson et al. (2004)*
G9401	TrGNS	DMS + NO_3 → CH_3SO_2 + HNO_3 + HCHO	1.9E-13*EXP(520./temp)	Atkinson et al. (2004)
G9402	TrGS	DMSO + OH → .6 SO_2 + HCHO + .6 CH_3O_2 + .4 HO_2 + .4 CH_3SO_3H	1.E-10	Hynes and Wine (1996)
G9403	TrGS	CH_3SO_2 → SO_2 + CH_3O_2	1.8E13*EXP(-8661./temp)	Barone et al. (1995)
G9404	TrGS	CH_3SO_2 + O_3 → CH_3SO_3	3.E-13	Barone et al. (1995)
G9405	TrGS	CH_3SO_3 + HO_2 → CH_3SO_3H	5.E-11	Barone et al. (1995)
G9600	TrGSCl	DMS + Cl → CH_3SO_2 + HCl + HCHO	3.3E-10	Atkinson et al. (2004)
G9700	TrGSBr	DMS + Br → CH_3SO_2 + HBr + HCHO	9.E-11*EXP(-2386./temp)	Jefferson et al. (1994)
G9701	TrGSBr	DMS + BrO → DMSO + Br	4.4E-13	Ingham et al. (1999)
G9800	TrGSI	DMS + IO → DMSO + I	3.2E-13*EXP(-925./temp)	Dillon et al. (2006a)
G10100	TrGHg	Hg + O_3 → HgO + O_2	3.0E-20	Hall (1995)
G10200	TrGHg	Hg + OH → HgO + H	3.55E-14*EXP(294./temp)	Pal and Ariya (2004)
G10201	TrGHg	Hg + H_2O_2 → HgO + H_2O	8.5E-19	Tokos et al. (1998)*
G10600	TrGHgCl	Hg + Cl → HgCl	1.0E-11	Ariya et al. (2002)
G10601	TrGHgCl	Hg + Cl_2 → $HgCl_2$	2.6E-18	Ariya et al. (2002)
G10700	TrGHgBr	Hg + Br → HgBr	3.0E-13	Donohoue et al. (2006)
G10701	TrGHgBr	HgBr + Br → $HgBr_2$	3.0E-12	Calvert and Lindberg (2003)
G10702	TrGHgBr	Hg + Br_2 → $HgBr_2$	9.0E-17	Ariya et al. (2002)
G10703	TrGHgBr	Hg + BrO → HgO + Br	1.0E-15	Raofie and Ariya (2003)
G10704	TrGHgBr	HgBr + BrO → BrHgOBr	3.0E-12	Calvert and Lindberg (2003)
G10705	TrGHgClBr	HgCl + BrO → ClHgOBr	3.0E-12	Calvert and Lindberg (2003)
G10706	TrGHgClBr	HgBr + Cl → ClHgBr	3.0E-12	Calvert and Lindberg (2003)
G10707	TrGHgClBr	HgCl + Br → ClHgBr	3.0E-12	Calvert and Lindberg (2003)

*Notes:

Rate coefficients for three-body reactions are defined via the function `k_3rd`(T, M, k_0^{300}, n, k_{inf}^{300}, m, f_c). In the code, the temperature T is called `temp` and the concentration of "air molecules" M is called `cair`. Using the auxiliary variables $k_0(T)$, $k_{\mathrm{inf}}(T)$, and k_{ratio}, `k_3rd` is defined as:

$$k_0(T) = k_0^{300} \times \left(\frac{300\mathrm{K}}{T}\right)^n \quad (1)$$

$$k_{\mathrm{inf}}(T) = k_{\mathrm{inf}}^{300} \times \left(\frac{300\mathrm{K}}{T}\right)^m \quad (2)$$

$$k_{\mathrm{ratio}} = \frac{k_0(T)M}{k_{\mathrm{inf}}(T)} \quad (3)$$

$$\texttt{k\_3rd} = \frac{k_0(T)M}{1+k_{\mathrm{ratio}}} \times f_c^{\left(\frac{1}{1+(\log_{10}(k_{\mathrm{ratio}}))^2}\right)} \quad (4)$$

A similar function, called `k_3rd_iupac` here, is used by Atkinson et al. (2005) for three-body reactions. It has the same function parameters as `k_3rd` and it is defined as:

$$k_0(T) = k_0^{300} \times \left(\frac{300\mathrm{K}}{T}\right)^n \quad (5)$$

$$k_{\mathrm{inf}}(T) = k_{\mathrm{inf}}^{300} \times \left(\frac{300\mathrm{K}}{T}\right)^m \quad (6)$$

$$k_{\mathrm{ratio}} = \frac{k_0(T)M}{k_{\mathrm{inf}}(T)} \quad (7)$$

$$N = 0.75 - 1.27 \times \log_{10}(f_c) \quad (8)$$

$$\texttt{k\_3rd\_iupac} = \frac{k_0(T)M}{1+k_{\mathrm{ratio}}} \times f_c^{\left(\frac{1}{1+(\log_{10}(k_{\mathrm{ratio}})/N)^2}\right)} \quad (9)$$

G1002: The path leading to 2 $O(^3P)$ + O_2 results in a null cycle regarding odd oxygen and is neglected.

G2110: The rate coefficient is: `k_HO2_HO2 = (1.5E-12*EXP(19./temp)+1.7E-33*EXP(1000./temp) *cair)* (1.+1.4E-21*EXP(2200./temp)*C(ind_ H2O))`. The value for the first (pressure-independent) part is from Christensen et al. (2002), the water term from Kircher and Sander (1984).

G3109: The rate coefficient is: `k_NO3_NO2 = k_ 3rd(temp,cair,2.E-30,4.4,1.4E-12,0.7,0.6)`.

G3110: The rate coefficient is defined as backward reaction divided by equilibrium constant.

G3203: The rate coefficient is: `k_NO2_HO2 = k_ 3rd(temp,cair,1.8E-31,3.2,4.7E-12,1.4,0.6)`.

G3206: The rate coefficient is: `k_HNO3_ OH = 2.4E-14 * EXP(460./temp) + 1./ ( 1./ (6.5E-34 * EXP(1335./temp)*cair) + 1./ (2.7E-17 * EXP(2199./temp)) )`

G3207: The rate coefficient is defined as backward reaction divided by equilibrium constant.

G4103: Sander et al. (2006) recommend a zero product yield for HCHO.

G4107: The rate coefficient is: `k_CH3OOH_OH = 3.8E-12*EXP(200./temp)`.

G4109: The same temperature dependence assumed as for CH_3CHO+NO_3.

G4201: The product distribution is from Rickard and Pascoe (2009), after substitution of the Criegee intermediate by its decomposition products.

G4206: The product C_2H_5OH, which reacts only with OH, is substituted by its degradation products $\approx$ 0.1 $HOCH_2CH_2O_2$ + 0.9 CH_3CHO + 0.9 HO_2.

G4207: The rate constant `8.01E-12` is for the H abstraction in alpha to the $-$OOH group (Rickard and Pascoe, 2009) and `0.6*k_CH3OOH_OH` is for the $C_2H_5O_2$ channel. The branching ratios are calculated from the terms of the rate coefficient at 298 K.

G4218: The rate coefficient is the same as for the CH_3O_2 channel in G4107 (CH_3OOH+OH).

G4221: The rate coefficient is`k_PAN_M = k_CH3CO3_ NO2/9.E-29*EXP(-14000./temp)`, i.e. the rate coefficient is defined as backward reaction divided by equilibrium constant.

G4243: Orlando et al. (1998) estimated that about 25% of the $HOCH_2CH_2O$ in this reaction is produced with sufficient excess energy that it decomposes promptly. The decomposition products are 2 HCHO + HO_2.

G4300: The product NC3H7O2 is substituted with its degradation products $C_2H_5O_2$ + CO_2 + HO_2.

G4301: The product distribution is for terminal olefin carbons from Zaveri and Peters (1999).

G4304: The value for the generic RO_2 + HO_2 reaction from Atkinson (1997) is used here.

G4306: The MCM (Rickard and Pascoe, 2009) products are 0.2 IPROPOL + 0.2 CH_3COCH_3 + 0.6 IC3H7O. IPROPOL and IC3H7O are substituted with their degradation products. We assume IPROPOL to be oxidized entirely to CH_3COCH_3 + HO_2 by OH. IC3H7O + O_2 produces the same products.

G4307: Analogous to G4207 for both rate coefficient and branching ratios.

G4400: $LC_4H_9O_2$ represents 0.127 NC4H9O2 + 0.873 SC4H9O2.

G4401: NC4H9O and SC4H9O are substituted with 2 CO_2 + $C_2H_5O_2$ and 0.636 MEK + HO_2 and 0.364 CH_3CHO + $C_2H_5O_2$, repectively. The stoichiometric coefficients on the right side are weighted averages.

G4403: The alkyl nitrate yield is the weighted average yield for the two isomers forming from NC4H9O2 and SC4H9O2.

G4404: The product distribution is the weighted average of the single isomer hydroperoxides. It is calculated from the rate constants of single channels and the ratio of the isomers NC4H9O2 and SC4H9O2. The

overall rate constant for this reaction is calculated as weighted average of the channels rate constants. The relative weight of the products from NC4H9OOH and SC4H9OOH are then 0.0887 and 0.9113. The channels producing RO_2 are given the rate coefficient `0.6*k_CH3OOH_OH` as for G4107. For NC4H9OOH the products are 0.327 NC4H9O2 + 0.673 C3H7CHO + 0.673 OH. C3H7CHO is then substituted with 2 CO_2 + $C_2H_5O_2$. Hence, 0.327 NC4H9O2 + 1.346 CO_2 + 0.673 $C_2H_5O_2$ + 0.673 OH. For SC4H9OOH the products are 0.219 SC4H9O2 + 0.781 MEK + 0.781 OH.

G4413: LMEKO2 represents 0.459 MEKAO2 + 0.462 MEKBO2 + 0.079 MEKCO2.

G4415: Alkyl nitrate formation is neglected. The products of MEKAO and MEKCO are substituted with HCHO + CO_2 + $HOCH_2CH_2O_2$ and HCHO + CO_2 + $C_2H_5O_2$.

G4416: LMEKOOH is assumed having the composition 0.459 MEKAOOH + 0.462 MEKBOOH + 0.079 MEKCOOH. MEKAOOH + OH gives 0.89 CO2C3CHO + 0.89 OH + 0.11 MEKAO2 + H_2O. CO2C3CHO is substituted with $CH_3COCH_2O_2$ + CO_2 and the products become 0.89 $CH_3COCH_2O_2$ + 0.89 CO_2 + 0.89 OH + 0.11 MEKAO2 + H_2O. MEKBOOH + OH gives 0.758 BIACET + 0.758 OH + 0.242 MEKBO2 + H_2O. MEKCOOH + OH gives 0.614 EGLYOX + 0.614 OH + 0.386 MEKCO2 + H_2O. EGLYOX is substituted with $C_2H_5O_2$ + 2 CO_2 and the products become 0.614 $C_2H_5O_2$ + 1.228 CO_2 + 0.614 OH + 0.386 MEKCO2 + H_2O.

G4417: The rate coefficient is the combination of the ones for the two isomers weighted by the relative abundances for NC4H9NO3 and SC4H9NO3, respectively. Product distribution is calculated accordingly. NC4H9NO3 + OH gives C3H7CHO + NO_2 + H_2O with C3H7CHO being substituted with 2 CO_2 + $C_2H_5O_2$. After substitution is obtained 2 CO_2 + $C_2H_5O_2$ + NO_2 + H_2O. SC4H9NO3 + OH gives MEK + NO_2 + H_2O For the product distribution NC4H9NO3 and SC4H9NO3 account for 0.08577 and 0.91423, respectively.

G4419: The same value as for PAN is assumed.

G4420: Products are as in G4415. Only the main channels for each isomer are considered. Rate constant is the weighted average for the isomers.

G4437: LHMVKABO2 is a lumped species of virtual composition 0.3 HMVKAO2 + 0.7 HMVKBO2. The products are the weighted average for the permutation reactions of each single RO2 in the MCM (Rickard and Pascoe, 2009).

G4439: products are the weighted average for the decomposition of 0.3 HMVKAO + 0.7 HMVKBO.

G4440: as for G4439

G4441: The rate coefficient and products are 30% for HMVKAOOH and 70% for HMVKBOOH.

G4444: LMVKOHABO2 is a lumped species of virtual composition 0.3 MVKOHAO2 + 0.7 MVKOHBO2. The products are the weighted average for the permutation reactions of each single RO2 in the MCM (Rickard and Pascoe, 2009).

G4446: products are the weighted average for the decomposition of 0.3 MVKOHAO + 0.7 MVKOHBO.

G4447: as for G4446

G4448: The rate coefficient and products are 30% for MVKOHAOOH and 70% for MVKOHBOOH.

G6103: The rate coefficient is defined as backward reaction divided by equilibrium constant.

G6402: The initial products are probably HCl and CH_2OOH (Atkinson et al., 2006). It is assumed that CH_2OOH dissociates into HCHO and OH.

G6405: Average of reactions with CH_3Br and CH_3F from Sander et al. (2006) (B. Steil, pers. comm.).

G6407: Rough extrapolation from reactions with CH_3CF_3, CH_3CClF_2, and CH_3CCl_2F from Sander et al. (2006).

G6409: It is assumed that the reaction liberates all Cl atoms in the form of HCl.

G7302: The rate coefficient is: `k_BrO_NO2 = k_3rd(temp,cair,5.2E-31,3.2,6.9E-12,2.9,0.6)`.

G7303: The rate coefficient is defined as backward reaction (Atkinson et al., 2007) divided by equilibrium constant (Orlando and Tyndall, 1996).

G7404: It is assumed that the reaction liberates all Br atoms in the form of HBr.

G7407: It is assumed that the reaction liberates all Br atoms. The fate of the carbon atom is currently not considered.

G7408: It is assumed that the reaction liberates all Br atoms. The fate of the carbon atom is currently not considered.

G7605: Same value as for G7408: CH_2Br_2+OH assumed. It is assumed that the reaction liberates all Br atoms but not Cl. The fate of the carbon atom is currently not considered.

G7606: Same value as for G7408: CH_2Br_2+OH assumed. It is assumed that the reaction liberates all Br atoms but not Cl. The fate of the carbon atom is currently not considered.

G7607: It is assumed that the reaction liberates all Br atoms but not Cl. The fate of the carbon atom is currently not considered.

G8102: It is assumed that the reaction produces new particles.

G8103: The yield of 38 % OIO is from Atkinson et al. (2007). It is assumed here that the remaining 62 % produce 2 I + O_2.

G8300: The rate coefficient is: `k_I_NO2 = k_3rd_iupac(temp,cair,3.E-31,1.,6.6E-11,0.,0.63)`.

G8305: The rate coefficient is defined as backward reaction (Atkinson et al., 2007) divided by equilibrium constant (van den Bergh and Troe, 1976).

G8306: According to John Plane and John Crowley (pers. comm. 2007), the rate coefficient of `1.1E15*EXP(-12060./temp)` suggested by Atkinson et al. (2007) is wrong.

G8401: The rate coefficient is from Dillon et al. (2006b), the yield of I atoms is a lower limit given on page 2170 of Bale et al. (2005).

G8402: The products are from Nakano et al. (2005).

G8701: 80% Br + OIO production is from Atkinson et al. (2007). The remaining channels are assumed to produce Br + I + O_2.

G9400: Addition path. The rate coefficient is: `k_DMS_OH = 1.0E-39*EXP(5820./temp)*C(ind_O2) / (1.+5.0E-30*EXP(6280./temp)*C(ind_O2))`.

G10201: Upper limit.

Table 2: Photolysis reactions

#	labels	reaction	rate coefficient	reference
J1000	StTrGJ	$O_2 + h\nu \rightarrow O(^3P) + O(^3P)$	jx(ip_O2)	see note
J1001a	StTrGJ	$O_3 + h\nu \rightarrow O(^1D)$	jx(ip_O1D)	see note
J1001b	StTrGJ	$O_3 + h\nu \rightarrow O(^3P)$	jx(ip_O3P)	see note
J2100	StGJ	$H_2O + h\nu \rightarrow H + OH$	jx(ip_H2O)	see note
J2101	StTrGJ	$H_2O_2 + h\nu \rightarrow 2\ OH$	jx(ip_H2O2)	see note
J3100	StGNJ	$N_2O + h\nu \rightarrow O(^1D)$	jx(ip_N2O)	see note
J3101	StTrGNJ	$NO_2 + h\nu \rightarrow NO + O(^3P)$	jx(ip_NO2)	see note
J3102	StGNJ	$NO + h\nu \rightarrow N + O(^3P)$	jx(ip_NO)	see note
J3103a	StTrGNJ	$NO_3 + h\nu \rightarrow NO_2 + O(^3P)$	jx(ip_NO2O)	see note
J3103b	StTrGNJ	$NO_3 + h\nu \rightarrow NO$	jx(ip_NOO2)	see note
J3104a	StTrGNJ	$N_2O_5 + h\nu \rightarrow NO_2 + NO_3$	jx(ip_N2O5)	see note
J3104b	StGNJ	$N_2O_5 + h\nu \rightarrow NO + O(^3P) + NO_3$	jx(ip_NO3NOO)	see note
J3200	TrGJ	$HONO + h\nu \rightarrow NO + OH$	jx(ip_HONO)	see note
J3201	StTrGNJ	$HNO_3 + h\nu \rightarrow NO_2 + OH$	jx(ip_HNO3)	see note
J3202	StTrGNJ	$HNO_4 + h\nu \rightarrow .667\ NO_2 + .667\ HO_2 + .333\ NO_3 + .333\ OH$	jx(ip_HNO4)	see note
J4100	StTrGJ	$CH_3OOH + h\nu \rightarrow HCHO + OH + HO_2$	jx(ip_CH3OOH)	see note
J4101a	StTrGJ	$HCHO + h\nu \rightarrow H_2 + CO$	jx(ip_COH2)	see note
J4101b	StTrGJ	$HCHO + h\nu \rightarrow H + CO + HO_2$	jx(ip_CHOH)	see note
J4102	StGJ	$CO_2 + h\nu \rightarrow CO + O(^3P)$	jx(ip_CO2)	see note
J4103	StGJ	$CH_4 + h\nu \rightarrow CO + 0.31\ H + 0.69\ H_2 + 1.155\ H_2O$	jx(ip_CH4)	see note
J4200	TrGCJ	$C_2H_5OOH + h\nu \rightarrow CH_3CHO + HO_2 + OH$	jx(ip_CH3OOH)	von Kuhlmann (2001)*
J4201	TrGCJ	$CH_3CHO + h\nu \rightarrow CH_3O_2 + HO_2 + CO$	jx(ip_CH3CHO)	see note
J4202	TrGCJ	$CH_3C(O)OOH + h\nu \rightarrow CH_3O_2 + OH + CO_2$	jx(ip_CH3CO3H)	see note
J4204	TrGNCJ	$PAN + h\nu \rightarrow CH_3C(O)OO + NO_2$	jx(ip_PAN)	see note
J4205	TrGCJ	$HOCH_2CHO + h\nu \rightarrow HO_2 + HCHO + HO_2 + CO$	jx(ip_HOCH2CHO)	see note
J4206	TrGCJ	$HOCH_2CO_3H + h\nu \rightarrow HCHO + HO_2 + OH + CO_2$	jx(ip_CH3OOH)	Rickard and Pascoe (2009)*
J4207	TrGCJ	$PHAN + h\nu \rightarrow HOCH_2CO_3 + NO_2$	jx(ip_PAN)	see note
J4208	TrGCJ	$GLYOX + h\nu \rightarrow 2\ CO + 2\ HO_2$	jx(ip_GLYOX)	see note
J4209	TrGNCJ	$HCOCO_2H + h\nu \rightarrow 2\ HO_2 + CO + CO_2$	jx(ip_MGLYOX)	Rickard and Pascoe (2009)*
J4210	TrGNCJ	$HCOCO_3H + h\nu \rightarrow HO_2 + CO + OH + CO_2$	(jx(ip_CH3OOH)+jx(ip_HOCH2CHO))	Rickard and Pascoe (2009)*

Table 2: Photolysis reactions (... continued)

#	labels	reaction	rate coefficient	reference
J4211	TrGCJ	HYETHO2H + hν → $HOCH_2CH_2O$ + OH	jx(ip_CH3OOH)	Rickard and Pascoe (2009)*
J4212	TrGCJ	ETHOHNO3 + hν → HO_2 + 2 HCHO + NO_2	J_IC3H7NO3	see note
J4300	TrGCJ	iC_3H_7OOH + hν → CH_3COCH_3 + HO_2 + OH	jx(ip_CH3OOH)	von Kuhlmann (2001)*
J4301	TrGCJ	CH_3COCH_3 + hν → $CH_3C(O)OO$ + CH_3O_2	jx(ip_CH3COCH3)	see note
J4302	TrGCJ	CH_3COCH_2OH + hν → $CH_3C(O)OO$ + HCHO + HO_2	J_ACETOL	see note
J4303	TrGCJ	MGLYOX + hν → $CH_3C(O)OO$ + CO + HO_2	jx(ip_MGLYOX)	see note
J4304	TrGCJ	$CH_3COCH_2O_2H$ + hν → $CH_3C(O)OO$ + HCHO + OH	jx(ip_CH3OOH)+J_ACETOL	Rickard and Pascoe (2009)*
J4306	TrGNCJ	$iC_3H_7ONO_2$ + hν → CH_3COCH_3 + NO_2 + HO_2	J_IC3H7NO3	von Kuhlmann et al. (2003)*
J4307	TrGCJ	NOA + hν → $CH_3C(O)OO$ + HCHO + NO_2	J_IC3H7NO3+jx(ip_CH3COCH3)	see note
J4308	TrGCJ	HOCH2COCO2H + hν → $HOCH_2CO_3$ + HO_2 + CO_2	jx(ip_MGLYOX)	Rickard and Pascoe (2009)*
J4309	TrGCJ	HYPROPO2H + hν → CH_3CHO + HCHO + HO_2 + OH	jx(ip_CH3OOH)	Rickard and Pascoe (2009)*
J4310	TrGNCJ	PR2O2HNO3 + hν → NOA + HO_2 + OH	jx(ip_CH3OOH)	Rickard and Pascoe (2009)*
J4311	TrGCJ	HOCH2COCHO + hν → $HOCH_2CO_3$ + CO + HO_2	jx(ip_MGLYOX)	Rickard and Pascoe (2009)*
J4400	TrGCJ	LC_4H_9OOH + hν → OH + 0.254 CO_2 + 0.5552 MEK + 0.5552 HO_2 + 0.3178 CH_3CHO + 0.4448 $C_2H_5O_2$	jx(ip_CH3OOH)	Rickard and Pascoe (2009)*
J4401	TrGCJ	MVK + hν → .5 C_3H_6 + .5 $CH_3C(O)OO$ + .5 HCHO + CO + .5 HO_2	jx(ip_MVK)	see note
J4403	TrGCJ	MEK + hν → $CH_3C(O)OO$ + $C_2H_5O_2$	0.42*jx(ip_CHOH)	von Kuhlmann et al. (2003)*
J4404	TrGCJ	LMEKOOH + hν → 0.538 HCHO + 0.538 CO_2 + 0.459 $HOCH_2CH_2O_2$ + 0.079 $C_2H_5O_2$ + 0.462 $CH_3C(O)OO$ + 0.462 CH_3CHO + OH	jx(ip_CH3OOH)	Rickard and Pascoe (2009)*
J4405	TrGCJ	BIACET + hν → 2 $CH_3C(O)OO$	2.15*jx(ip_MGLYOX)	see note
J4406	TrGNCJ	LC4H9NO3 + hν → NO_2 + 0.254 CO_2 + 0.5552 MEK + 0.5552 HO_2 + 0.3178 CH_3CHO + 0.4448 $C_2H_5O_2$	J_IC3H7NO3	see note
J4407	TrGNCJ	MPAN + hν → MACO3 + NO_2	jx(ip_PAN)	see note
J4408	TrGCJ	LMVKOHABOOH + hν → .3 HOCH2COCHO + .3 HCHO + .3 HO_2 + .7 $HOCH_2CHO$ + .7 $HOCH_2CO_3$ + OH	J_ACETOL+jx(ip_CH3OOH)	Rickard and Pascoe (2009)*

Table 2: Photolysis reactions (... continued)

#	labels	reaction	rate coefficient	reference
J4409	TrGCJ	CO2H3CO3H + hν → MGLYOX + HO_2 + OH + CO_2	jx(ip_CH3OOH)	Rickard and Pascoe (2009)*
J4410	TrGCJ	CO2H3CO3H + hν → $CH_3C(O)OO$ + HO_2 + $HCOCO_3H$	J_ACETOL	Rickard and Pascoe (2009)*
J4411	TrGCJ	MACR + hν → .5 MACO3 + .5 $CH_3C(O)OO$ + .5 HCHO + .5 CO + HO_2	jx(ip_MACR)	see note
J4412	TrGCJ	MACROOH + hν → CH_3COCH_2OH + HCHO + HO_2 + OH	jx(ip_CH3OOH)	Rickard and Pascoe (2009)*
J4413	TrGCJ	MACROOH + hν → CH_3COCH_2OH + CO + HO_2 + OH	2.77*jx(ip_HOCH2CHO)	see note
J4414	TrGCJ	MACROH + hν → CH_3COCH_2OH + CO + HO_2 + HO_2	2.77*jx(ip_HOCH2CHO)	see note
J4415	TrGCJ	MACO3H + hν → $CH_3C(O)OO$ + HCHO + OH + CO_2	jx(ip_CH3OOH)	Rickard and Pascoe (2009)*
J4416	TrGCJ	LHMVKABOOH + hν → .3 MGLYOX + .7 $CH_3C(O)OO$ + .7 $HOCH_2CHO$ + .3 HCHO + .3 HO_2 + OH	jx(ip_CH3OOH)	Rickard and Pascoe (2009)*
J4417	TrGCJ	MVKOH + hν → .5 HCHO + .5 HO_2 + .5 $HOCH_2CO_3$ + CO + 1.5 LCARBON	jx(ip_MVK)	Rickard and Pascoe (2009)*
J4418	TrGCJ	CO2H3CHO + hν → MGLYOX + CO + HO_2 + HO_2	jx(ip_HOCH2CHO)	Rickard and Pascoe (2009)*
J4419	TrGCJ	HO12CO3C4 + hν → $CH_3C(O)OO$ + $HOCH_2CHO$ + HO_2	J_ACETOL	Rickard and Pascoe (2009)*
J4420	TrGCJ	BIACETOH + hν → $CH_3C(O)OO$ + $HOCH_2CO_3$	2.15*jx(ip_MGLYOX)	see note
J4502	TrGCJ	LISOPACOOH + hν → LHC4ACCHO + HO_2 + OH	jx(ip_CH3OOH)	Rickard and Pascoe (2009)*
J4503	TrGNCJ	LISOPACNO3 + hν → LHC4ACCHO + HO_2 + NO_2	0.59*J_IC3H7NO3	see note
J4504	TrGCJ	ISOPBOOH + hν → .75 MVK + .25 MVKOH + .75 HCHO + .75 HO_2 + .25 CH_3O_2 + OH	jx(ip_CH3OOH)	Rickard and Pascoe (2009)*
J4505	TrGNCJ	ISOPBNO3 + hν → .75 MVK + .25 MVKOH + .75 HCHO + .75 HO_2 + .25 CH_3O_2 + NO_2	2.84*J_IC3H7NO3	see note
J4506	TrGCJ	ISOPDOOH + hν → MACR + HCHO + HO_2 + OH	jx(ip_CH3OOH)	Rickard and Pascoe (2009)*
J4507	TrGNCJ	ISOPDNO3 + hν → MACR + HCHO + HO_2 + NO_2	J_IC3H7NO3	see note
J4508	TrGNCJ	NISOPOOH + hν → NC4CHO + HO_2 + OH	jx(ip_CH3OOH)	Rickard and Pascoe (2009)*
J4509	TrGNCJ	NC4CHO + hν → NOA + 2 CO + 2 HO_2	jx(ip_MACR)	see note

Table 2: Photolysis reactions (... continued)

#	labels	reaction	rate coefficient	reference
J4510	TrGNCJ	LNISOOH + hν $\rightarrow$ NOA + OH + .5 GLYOX + .5 CO + HO_2 + .5 CO_2	jx(ip_CH3OOH)	Taraborrelli et al. (2009)*
J4511	TrGCJ	LHC4ACCHO + hν $\rightarrow$ .5 LHC4ACCO3 + .25 CH_3COCH_2OH + .25 $HOCH_2CHO$ + .25 $CH_3C(O)OO$ + .75 CO + 1.25 HO_2	jx(ip_MACR)	Rickard and Pascoe (2009)*
J4512	TrGCJ	LC578OOH + hν $\rightarrow$ .5 CH_3COCH_2OH + .5 MGLYOX + .5 GLYOX + .5 $HOCH_2CHO$ + HO_2 + OH	jx(ip_CH3OOH)	Taraborrelli et al. (2009)*
J4513	TrGCJ	LHC4ACCO3H + hν $\rightarrow$ .5 CH_3COCH_2OH + .5 $HOCH_2CHO$ + .5 $CH_3C(O)OO$ + .5 CO + .5 HO_2 + OH + CO_2	jx(ip_CH3OOH)	Rickard and Pascoe (2009)*
J4514	TrGNCJ	LC5PAN1719 + hν $\rightarrow$ .5 MACROH + .5 HO12CO3C4 + CO + NO_2	jx(ip_PAN)	see note
J4515	TrGCJ	HCOC5 + hν $\rightarrow$ $CH_3C(O)OO$ + HCHO + $HOCH_2CO_3$	0.5*jx(ip_MVK)	see note
J4516	TrGCJ	C59OOH + hν $\rightarrow$ CH_3COCH_2OH + $HOCH_2CO_3$ + OH	J_ACETOL+jx(ip_CH3OOH)	Rickard and Pascoe (2009)*
J6000	StTrGClJ	Cl_2 + hν $\rightarrow$ Cl + Cl	jx(ip_Cl2)	see note
J6100	StTrGClJ	Cl_2O_2 + hν $\rightarrow$ 2 Cl	1.4*jx(ip_Cl2O2)	see note
J6101	StTrGClJ	OClO + hν $\rightarrow$ ClO + $O(^3P)$	jx(ip_OClO)	see note
J6200	StGClJ	HCl + hν $\rightarrow$ Cl + H	jx(ip_HCl)	see note
J6201	StTrGClJ	HOCl + hν $\rightarrow$ OH + Cl	jx(ip_HOCl)	see note
J6300	TrGNClJ	$ClNO_2$ + hν $\rightarrow$ Cl + NO_2	jx(ip_ClNO2)	see note
J6301a	StTrGNClJ	$ClNO_3$ + hν $\rightarrow$ Cl + NO_3	jx(ip_ClNO3)	see note
J6301b	StTrGNClJ	$ClNO_3$ + hν $\rightarrow$ ClO + NO_2	jx(ip_ClONO2)	see note
J6400	StGClJ	CH_3Cl + hν $\rightarrow$ Cl + CH_3O_2	jx(ip_CH3Cl)	see note
J6401	StGClJ	CCl_4 + hν $\rightarrow$ 4 Cl	jx(ip_CCl4)	see note
J6402	StGCClJ	CH_3CCl_3 + hν $\rightarrow$ 3 Cl	jx(ip_CH3CCl3)	see note
J6500	StGFClJ	$CFCl_3$ + hν $\rightarrow$ 3 Cl	jx(ip_CFCl3)	see note
J6501	StGFClJ	CF_2Cl_2 + hν $\rightarrow$ 2 Cl	jx(ip_CF2Cl2)	see note
J7000	StTrGBrJ	Br_2 + hν $\rightarrow$ Br + Br	jx(ip_Br2)	see note
J7100	StTrGBrJ	BrO + hν $\rightarrow$ Br + $O(^3P)$	jx(ip_BrO)	see note
J7200	StTrGBrJ	HOBr + hν $\rightarrow$ Br + OH	jx(ip_HOBr)	see note
J7300	TrGNBrJ	$BrNO_2$ + hν $\rightarrow$ Br + NO_2	jx(ip_BrNO2)	see note
J7301	StTrGNBrJ	$BrNO_3$ + hν $\rightarrow$ 0.29 Br + 0.29 NO_3 + 0.71 BrO + 0.71 NO_2	jx(ip_BrNO3)	see note
J7400	StGBrJ	CH_3Br + hν $\rightarrow$ Br + CH_3O_2	jx(ip_CH3Br)	see note
J7401	TrGBrJ	CH_2Br_2 + hν $\rightarrow$ 2 Br	jx(ip_CH2Br2)	see note
J7402	TrGBrJ	$CHBr_3$ + hν $\rightarrow$ 3 Br	jx(ip_CHBr3)	see note
J7500	StGFBrJ	CF_3Br + hν $\rightarrow$ Br	jx(ip_CF3Br)	see note

Table 2: Photolysis reactions (... continued)

#	labels	reaction	rate coefficient	reference
J7600	StTrGClBrJ	$BrCl + h\nu \rightarrow Br + Cl$	jx(ip_BrCl)	see note
J7601	StGFBrJ	$CF_2ClBr + h\nu \rightarrow Br + Cl$	jx(ip_CF2ClBr)	see note
J7602	TrGClBrJ	$CH_2ClBr + h\nu \rightarrow Br + Cl$	jx(ip_CH2ClBr)	see note
J7603	TrGClBrJ	$CHCl_2Br + h\nu \rightarrow Br + 2\ Cl$	jx(ip_CHCl2Br)	see note
J7604	TrGClBrJ	$CHClBr_2 + h\nu \rightarrow 2\ Br + Cl$	jx(ip_CHClBr2)	see note
J8000	TrGIJ	$I_2 + h\nu \rightarrow I + I$	jx(ip_I2)	see note
J8100	TrGIJ	$IO + h\nu \rightarrow I + O(^3P)$	jx(ip_IO)	see note
J8200	TrGIJ	$HOI + h\nu \rightarrow I + OH$	jx(ip_HOI)	see note
J8300	TrGNIJ	$INO_2 + h\nu \rightarrow I + NO_2$	jx(ip_INO2)	see note
J8301	TrGNIJ	$INO_3 + h\nu \rightarrow I + NO_3$	jx(ip_INO3)	see note
J8400	TrGIJ	$CH_2I_2 + h\nu \rightarrow 2\ I + 2\ HO_2 + CO$	jx(ip_CH2I2)	see note
J8401	TrGIJ	$CH_3I + h\nu \rightarrow I + CH_3O_2$	jx(ip_CH3I)	see note
J8402	TrGIJ	$C_3H_7I + h\nu \rightarrow I + CH_3O_2$	jx(ip_C3H7I)	see note
J8403	TrGIClJ	$CH_2ClI + h\nu \rightarrow I + Cl + 2\ HO_2 + CO$	jx(ip_CH2ClI)	see note
J8600	TrGClIJ	$ICl + h\nu \rightarrow I + Cl$	jx(ip_ICl)	see note
J8700	TrGBrIJ	$IBr + h\nu \rightarrow I + Br$	jx(ip_IBr)	see note
PH3200_a01	TrAa01NJ	$NO_3^-(aq) + h\nu \rightarrow NO_2(aq) + OH(aq) - H^+(aq)$	xaer(01)*jx(ip_NO2) * 1.4E-4	see note
PH10200_a01	TrAa01HgJ	$Hg(OH)_2(aq) + h\nu \rightarrow Hg(aq)$	xaer(01)*6E-5*jx(ip_NO2)	see note

*Notes:

J-values are calculated with an external module and then supplied to the MECCA chemistry.

Values that originate from the Master Chemical Mechanism (MCM) by Rickard and Pascoe (2009) are translated according in the following way:
J(11) → jx(ip_COH2)
J(12) → jx(ip_CHOH)
J(15) → jx(ip_HOCH2CHO)
J(18) → jx(ip_MACR)
J(22) → jx(ip_ACETOL)
J(23)+J(24) → jx(ip_MVK)
J(31)+J(32)+J(33) → jx(ip_GLYOX)
J(34) → jx(ip_MGLYOX)
J(41) → jx(ip_CH3OOH)
J(53) → J($iC_3H_7ONO_2$)
J(54) → J($iC_3H_7ONO_2$)
J(55) → J($iC_3H_7ONO_2$)
J(56)+J(57) → jx(ip_NOA)

J4207: It is assumed that J(PHAN) is the same as J(PAN).

J4212: It is assumed that J(ETHOHNO3) is the same as J($iC_3H_7ONO_2$).

J4302: Following von Kuhlmann et al. (2003), we use J(CH_3COCH_2OH) = 0.11*jx(ip_CHOH). As an additional factor, the quantum yield of 0.65 is taken from Orlando et al. (1999).

J4306: Following von Kuhlmann et al. (2003), we use J($iC_3H_7ONO_2$) = 3.7*jx(ip_PAN).

J4307: NOA contains the cromophores of both CH_3COCH_3 and a nitrate group. It is assumed here that the J values are additive, i.e.: J(NOA) = J(CH_3COCH_3) + J($iC_3H_7ONO_2$).

J4406: It is assumed that J(LC4H9NO3) is the same as J($iC_3H_7ONO_2$).

J4407: It is assumed that J(MPAN) is the same as J(PAN).

J4405: It is assumed that J(BIACET) is 2.15 times larger than J(MGLYOX), consistent with the photolysis rate coefficients used in the MCM (Rickard and Pascoe, 2009).

J4413: It is assumed that J(MACROOH) is 2.77 times larger than J($HOCH_2CHO$), consistent with the pho-

tolysis rate coefficients used in the MCM (Rickard and Pascoe, 2009).

J4414: It is assumed that J(MACROH) is 2.77 times larger than J($HOCH_2CHO$), consistent with the photolysis rate coefficients used in the MCM (Rickard and Pascoe, 2009).

J4420: It is assumed that J(BIACETOH) is 2.15 times larger than J(MGLYOX), consistent with the photolysis rate coefficients used in the MCM (Rickard and Pascoe, 2009).

J4503: It is assumed that J(LISOPACNO3) = 0.59 × J($iC_3H_7ONO_2$), consistent with the photolysis rate coefficients used in the MCM (Rickard and Pascoe, 2009).

J4505: It is assumed that J(ISOPBNO3) = 2.84 × J($iC_3H_7ONO_2$), consistent with the photolysis rate coefficients used in the MCM (Rickard and Pascoe, 2009).

J4509: It is assumed that J(NC4CHO) is the same as J(MACR).

J4514: It is assumed that J(LC5PAN1719) is the same as J(PAN).

J4515: Consistent with the MCM (Rickard and Pascoe, 2009), we assume that J(HCOC5) is half as large as J(MVK).

J6100: Stimpfle et al. (2004) claim that the combination of absorption cross sections from Burkholder et al. (1990) and the Cl_2O_2 formation rate coefficient by Sander et al. (2003) can approximately reproduce the observed Cl_2O_2/ClO ratios and ozone depletion. They give an almost zenith-angle independent ratio of 1.4 for Burkholder et al. (1990) to Sander et al. (2003) J-values. The IUPAC recommendation for the Cl_2O_2 formation rate is about 5 to 15 % less than the value by Sander et al. (2003) but more than 20 % larger than the value by Sander et al. (2000). The J-values by Burkholder et al. (1990) are within the uncertainty range of the IUPAC recommendation.

J7301: The quantum yields are from Sander et al. (2003).

Table 3: Henry's law coefficients

substance	$\frac{k_{\mathrm{H}}^{\ominus}}{\mathrm{M/atm}}$	$\frac{-\Delta_{\mathrm{soln}}H/R}{\mathrm{K}}$	reference
O_2	1.3×10^{-3}	1500.	Wilhelm et al. (1977)
O_3	1.2×10^{-2}	2560.	Chameides (1984)
OH	3.0×10^{1}	4300.	Hanson et al. (1992)
HO_2	3.9×10^{3}	5900.	Hanson et al. (1992)
H_2O_2	$1.\times10^{5}$	6338.	Lind and Kok (1994)
NH_3	58.	4085.	Chameides (1984)
NO	1.9×10^{-3}	1480.	Schwartz and White (1981)
NO_2	7.0×10^{-3}	2500.	Lee and Schwartz (1981)*
NO_3	2.	2000.	Thomas et al. (1993)
N_2O_5	BIG	0.	see note
HONO	4.9×10^{1}	4780.	Schwartz and White (1981)
HNO_3	$2.45\times10^{6}/1.5\times10^{1}$	8694.	Brimblecombe and Clegg (1989)*
HNO_4	1.2×10^{4}	6900.	Régimbal and Mozurkewich (1997)
CH_3O_2	6.	5600.	Jacob (1986)*
CH_3OOH	3.0×10^{2}	5322.	Lind and Kok (1994)
CO_2	3.1×10^{-2}	2423.	Chameides (1984)
HCHO	7.0×10^{3}	6425.	Chameides (1984)
HCOOH	3.7×10^{3}	5700.	Chameides (1984)
CH_3COOH	4.1×10^{3}	6200.	Sander et al. (2006)
PAN	2.8	5730.	Sander et al. (2006)
$C_2H_5O_2$	6.	5600.	see note
CH_3CHO	1.29×10^{1}	5890.	Sander et al. (2006)
CH_3COCH_3	28.1	5050.	Sander et al. (2006)
Cl_2	9.2×10^{-2}	2081.	Bartlett and Margerum (1999)
HCl	2./1.7	9001.	Brimblecombe and Clegg (1989)
HOCl	6.7×10^{2}	5862.	Huthwelker et al. (1995)
$ClNO_3$	BIG	0.	see note
Br_2	7.7×10^{-1}	3837.	Bartlett and Margerum (1999)
HBr	1.3	10239.	Brimblecombe and Clegg (1989)*
HOBr	9.3×10^{1}	5862.	Vogt et al. (1996)*
$BrNO_3$	BIG	0.	see note
BrCl	9.4×10^{-1}	5600.	Bartlett and Margerum (1999)
I_2	3.	4431.	Palmer et al. (1985)

Table 3: Henry's law coefficients (... continued)

substance	$\frac{k_{\mathrm{H}}^{\ominus}}{\mathrm{M/atm}}$	$\frac{-\Delta_{\mathrm{soln}}H/R}{\mathrm{K}}$	reference
IO	4.5×10^{2}	5862.	see note
OIO	BIG	0.	see note
I_2O_2	BIG	0.	see note
HI	BIG	0.	see note
HOI	4.5×10^{2}	5862.	Chatfield and Crutzen (1990)*
HIO_3	BIG	0.	see note
INO_2	BIG	0.	see note
INO_3	BIG	0.	see note
ICl	1.1×10^{2}	5600.	see note
IBr	2.4×10^{1}	5600.	see note
SO_2	1.2	3120.	Chameides (1984)
H_2SO_4	$1.\times10^{11}$	0.	see note
CH_3SO_3H	BIG	0.	see note
DMS	5.4×10^{-1}	3500.	Staudinger and Roberts (2001)
DMSO	$5.\times10^{4}$	6425.	De Bruyn et al. (1994)*
Hg	0.13	0.	Schroeder and Munthe (1998)
HgO	3.2×10^{6}	0.	Shon et al. (2005)
$HgCl_2$	2.4×10^{7}	0.	Shon et al. (2005)
$HgBr_2$	2.4×10^{7}	0.	see note
ClHgBr	2.4×10^{7}	0.	see note
BrHgOBr	2.4×10^{7}	0.	see note
ClHgOBr	2.4×10^{7}	0.	see note

*Notes:

The value "BIG" corresponds to virtually infinite solubility which is represented in the model using a very large but arbitrary number.

The temperature dependence of the Henry constants is:

$$K_{\mathrm{H}} = K_{\mathrm{H}}^{\ominus} \times \exp\left(\frac{-\Delta_{\mathrm{soln}}H}{R}\left(\frac{1}{T} - \frac{1}{T^{\ominus}}\right)\right)$$

where $\Delta_{\mathrm{soln}}H$ = molar enthalpy of dissolution [J/mol] and $R = 8.314$ J/(mol K).

NO_2: The temperature dependence is from Chameides (1984).

HNO_3: Calculated using the acidity constant from Davis and de Bruin (1964).

CH_3O_2: This value was estimated by Jacob (1986).

$C_2H_5O_2$: Assumed to be the same as $K_{\mathrm{H}}(CH_3O_2)$.

HBr: Calculated using the acidity constant from Lax (1969).

HOBr: This value was estimated by Vogt et al. (1996).

IO: Assumed to be the same as K_{H}(HOI).

HOI: Lower limit.

ICl: Calculated using thermodynamic data from Wagman et al. (1982).

IBr: Calculated using thermodynamic data from Wagman et al. (1982).

H_2SO_4: To account for the very high Henry's law coefficient of H_2SO_4, a very high value was chosen arbitrarily.

DMSO: Lower limit cited from another reference.

$HgBr_2$: Assumed to be the same as for $HgCl_2$.

ClHgBr: Assumed to be the same as for $HgCl_2$.

BrHgOBr: Assumed to be the same as for $HgCl_2$.

ClHgOBr: Assumed to be the same as for $HgCl_2$.

Table 4: Accommodation coefficients

substance	$\alpha^{\ominus}$	$\frac{-\Delta_{\mathrm{obs}}H/R}{\mathrm{K}}$	reference
O_2	0.01	2000.	see note
O_3	0.002	(default)	DeMore et al. (1997)*
OH	0.01	(default)	Takami et al. (1998)*
HO_2	0.5	(default)	Thornton and Abbatt (2005)
H_2O_2	0.077	3127.	Worsnop et al. (1989)
NH_3	0.06	(default)	DeMore et al. (1997)*
NO	5.0×10^{-5}	(default)	Saastad et al. (1993)*
NO_2	0.0015	(default)	Ponche et al. (1993)*
NO_3	0.04	(default)	Rudich et al. (1996)*
N_2O_5	(default)	(default)	DeMore et al. (1997)*
HONO	0.04	(default)	DeMore et al. (1997)*
HNO_3	0.5	(default)	Abbatt and Waschewsky (1998)*
HNO_4	(default)	(default)	DeMore et al. (1997)*
CH_3O_2	0.01	2000.	see note
CH_3OOH	0.0046	3273.	Magi et al. (1997)
CO_2	0.01	2000.	see note
HCHO	0.04	(default)	DeMore et al. (1997)*
HCOOH	0.014	3978.	DeMore et al. (1997)
CH_3COOH	2.0×10^{-2}	4079.	Davidovits et al. (1995)
PAN	(default)	(default)	see note
$C_2H_5O_2$	(default)	(default)	see note
CH_3CHO	3.0×10^{-2}	(default)	see note
CH_3COCH_3	3.72×10^{-3}	6395.	Davidovits et al. (1995)
Cl_2	0.038	6546.	Hu et al. (1995)
HCl	0.074	3072.	Schweitzer et al. (2000)*
HOCl	0.5	(default)	see note
$ClNO_3$	0.108	(default)	Deiber et al. (2004)*
Br_2	0.038	6546.	Hu et al. (1995)
HBr	0.032	3940.	Schweitzer et al. (2000)*
HOBr	0.5	(default)	Abbatt and Waschewsky (1998)*
$BrNO_3$	0.063	(default)	Deiber et al. (2004)*
BrCl	0.38	6546.	see note
I_2	0.01	2000.	see note
IO	0.5	2000.	see note

Table 4: Accommodation coefficients (... continued)

substance	$\alpha^{\ominus}$	$\frac{-\Delta_{\mathrm{obs}}H/R}{\mathrm{K}}$	reference
OIO	0.01	(default)	see note
I_2O_2	(default)	2000.	see note
HI	0.036	4130.	Schweitzer et al. (2000)*
HOI	0.5	(default)	see note
HIO_3	0.01	(default)	see note
INO_2	(default)	2000.	see note
INO_3	(default)	2000.	see note
ICl	0.018	2000.	Braban et al. (2007)
IBr	0.018	2000.	see note
SO_2	0.11	(default)	DeMore et al. (1997)
H_2SO_4	0.65	(default)	Pöschl et al. (1998)*
CH_3SO_3H	0.076	1762.	De Bruyn et al. (1994)
DMS	(default)	(default)	see note
DMSO	0.048	2578.	De Bruyn et al. (1994)
Hg	(default)	(default)	see note
HgO	(default)	(default)	see note
$HgCl_2$	(default)	(default)	see note
$HgBr_2$	(default)	(default)	see note
ClHgBr	(default)	(default)	see note
BrHgOBr	(default)	(default)	see note
ClHgOBr	(default)	(default)	see note

*Notes:

If no data are available, the following default values are used:
$\alpha^{\ominus} = 0.1$
$-\Delta_{\mathrm{obs}}H/R = 0$ K

The temperature dependence of the accommodation coefficients is given by (Jayne et al., 1991):

$$\frac{\alpha}{1-\alpha} = \exp\left(\frac{-\Delta_{\mathrm{obs}}G}{RT}\right) = \exp\left(\frac{-\Delta_{\mathrm{obs}}H}{RT} + \frac{\Delta_{\mathrm{obs}}S}{R}\right)$$

where $\Delta_{\mathrm{obs}}G$ is the Gibbs free energy barrier of the transition state toward solution (Jayne et al., 1991), and $\Delta_{\mathrm{obs}}H$ and $\Delta_{\mathrm{obs}}S$ are the corresponding enthalpy and entropy, respectively. The equation can be rearranged to:

$$\ln\left(\frac{\alpha}{1-\alpha}\right) = \frac{-\Delta_{\mathrm{obs}}H}{R} \times \frac{1}{T} + \frac{-\Delta_{\mathrm{obs}}S}{R}$$

and further:

$$\mathrm{d}\ln\left(\frac{\alpha}{1-\alpha}\right) \Big/ \mathrm{d}\left(\frac{1}{T}\right) = \frac{-\Delta_{\mathrm{obs}}H}{R}$$

O_2: Estimate.
O_3: Value measured at 292 K.
OH: Value measured at 293 K.
NH_3: Value measured at 295 K.
NO: Value measured between 193 and 243 K.
NO_2: Value measured at 298 K.

NO_3: Value is a lower limit, measured at 273 K.

N_2O_5: Value for sulfuric acid, measured between 195 and 300 K.

HONO: Value measured between 247 and 297 K.

HNO_3: Value measured at room temperature. Abbatt and Waschewsky (1998) say $\gamma > 0.2$. Here $\alpha = 0.5$ is used.

HNO_4: Value measured at 200 K for water ice.

CH_3O_2: Estimate.

CO_2: Estimate.

HCHO: Value measured between 260 and 270 K.

PAN: Estimate.

$C_2H_5O_2$: Estimate.

CH_3CHO: Using the same estimate as in the CAPRAM 2.4 model (`http://projects.tropos.de/capram/capram_24.html`).

HCl: Temperature dependence derived from published data at 2 different temperatures

HOCl: Assumed to be the same as α(HOBr).

$ClNO_3$: Value measured at 274.5 K.

HBr: Temperature dependence derived from published data at 2 different temperatures

HOBr: Value measured at room temperature. Abbatt and Waschewsky (1998) say $\gamma > 0.2$. Here $\alpha = 0.5$ is used.

$BrNO_3$: Value measured at 273 K.

BrCl: Assumed to be the same as $\alpha(Cl_2)$.

I_2: Estimate.

IO: Estimate.

OIO: Estimate.

I_2O_2: Estimate.

HI: Temperature dependence derived from published data at 2 different temperatures

HOI: Assumed to be the same as α(HOBr). See also Mössinger and Cox (2001) and Holmes et al. (2001).

HIO_3: Estimate.

INO_2: Estimate.

INO_3: Estimate.

IBr: Assumed to be the same as α(ICl).

H_2SO_4: Value measured at 303 K.

Hg: Estimate.

HgO: Estimate.

$HgCl_2$: Estimate.

$HgBr_2$: Estimate.

ClHgBr: Estimate.

BrHgOBr: Estimate.

ClHgOBr: Estimate.

Table 5: Reversible (Henry's law) equilibria and irreversible ("heterogenous") uptake

#	labels	reaction	rate coefficient	reference
H1000f_a01	TrAa01Sc	$O_2 \rightarrow O_2(aq)$	k_exf(01,ind_O2)	see note
H1000b_a01	TrAa01Sc	$O_2(aq) \rightarrow O_2$	k_exb(01,ind_O2)	see note
H1001f_a01	TrAa01MblScScm	$O_3 \rightarrow O_3(aq)$	k_exf(01,ind_O3)	see note
H1001b_a01	TrAa01MblScScm	$O_3(aq) \rightarrow O_3$	k_exb(01,ind_O3)	see note
H2100f_a01	TrAa01Sc	$OH \rightarrow OH(aq)$	k_exf(01,ind_OH)	see note
H2100b_a01	TrAa01Sc	$OH(aq) \rightarrow OH$	k_exb(01,ind_OH)	see note
H2101f_a01	TrAa01Sc	$HO_2 \rightarrow HO_2(aq)$	k_exf(01,ind_HO2)	see note
H2101b_a01	TrAa01Sc	$HO_2(aq) \rightarrow HO_2$	k_exb(01,ind_HO2)	see note
H2102f_a01	TrAa01MblScScm	$H_2O_2 \rightarrow H_2O_2(aq)$	k_exf(01,ind_H2O2)	see note
H2102b_a01	TrAa01MblScScm	$H_2O_2(aq) \rightarrow H_2O_2$	k_exb(01,ind_H2O2)	see note
H3101f_a01	TrAa01NSc	$NO_2 \rightarrow NO_2(aq)$	k_exf(01,ind_NO2)	see note
H3101b_a01	TrAa01NSc	$NO_2(aq) \rightarrow NO_2$	k_exb(01,ind_NO2)	see note
H3102f_a01	TrAa01NSc	$NO_3 \rightarrow NO_3(aq)$	k_exf(01,ind_NO3)	see note
H3102b_a01	TrAa01NSc	$NO_3(aq) \rightarrow NO_3$	k_exb(01,ind_NO3)	see note
H3200f_a01	TrAa01NMblScScm	$NH_3 \rightarrow NH_3(aq)$	k_exf(01,ind_NH3)	see note
H3200b_a01	TrAa01NMblScScm	$NH_3(aq) \rightarrow NH_3$	k_exb(01,ind_NH3)	see note
H3201_a01	TrAa01MblNScScm	$N_2O_5 \rightarrow HNO_3(aq) + HNO_3(aq)$	k_exf_N2O5(01)*C(ind_H2O_a01)	Behnke et al. (1994), Behnke et al. (1997)*
H3202f_a01	TrAa01NSc	$HONO \rightarrow HONO(aq)$	k_exf(01,ind_HONO)	see note
H3202b_a01	TrAa01NSc	$HONO(aq) \rightarrow HONO$	k_exb(01,ind_HONO)	see note
H3203f_a01	TrAa01MblNScScm	$HNO_3 \rightarrow HNO_3(aq)$	k_exf(01,ind_HNO3)	see note
H3203b_a01	TrAa01MblNScScm	$HNO_3(aq) \rightarrow HNO_3$	k_exb(01,ind_HNO3)	see note
H3204f_a01	TrAa01NSc	$HNO_4 \rightarrow HNO_4(aq)$	k_exf(01,ind_HNO4)	see note
H3204b_a01	TrAa01NSc	$HNO_4(aq) \rightarrow HNO_4$	k_exb(01,ind_HNO4)	see note
H4100f_a01	TrAa01MblScScm	$CO_2 \rightarrow CO_2(aq)$	k_exf(01,ind_CO2)	see note
H4100b_a01	TrAa01MblScScm	$CO_2(aq) \rightarrow CO_2$	k_exb(01,ind_CO2)	see note
H4101f_a01	TrAa01ScScm	$HCHO \rightarrow HCHO(aq)$	k_exf(01,ind_HCHO)	see note
H4101b_a01	TrAa01ScScm	$HCHO(aq) \rightarrow HCHO$	k_exb(01,ind_HCHO)	see note
H4102f_a01	TrAa01Sc	$CH_3O_2 \rightarrow CH_3OO(aq)$	k_exf(01,ind_CH3O2)	see note
H4102b_a01	TrAa01Sc	$CH_3OO(aq) \rightarrow CH_3O_2$	k_exb(01,ind_CH3O2)	see note
H4103f_a01	TrAa01ScScm	$HCOOH \rightarrow HCOOH(aq)$	k_exf(01,ind_HCOOH)	see note
H4103b_a01	TrAa01ScScm	$HCOOH(aq) \rightarrow HCOOH$	k_exb(01,ind_HCOOH)	see note
H4104f_a01	TrAa01ScScm	$CH_3OOH \rightarrow CH_3OOH(aq)$	k_exf(01,ind_CH3OOH)	see note
H4104b_a01	TrAa01ScScm	$CH_3OOH(aq) \rightarrow CH_3OOH$	k_exb(01,ind_CH3OOH)	see note

Table 5: Reversible (Henry's law) equilibria and irreversible ("heterogenous") uptake

#	labels	reaction	rate coefficient	reference
H6000f_a01	TrAa01ClMblSc	$Cl_2 \rightarrow Cl_2(aq)$	k_exf(01,ind_Cl2)	see note
H6000b_a01	TrAa01ClMblSc	$Cl_2(aq) \rightarrow Cl_2$	k_exb(01,ind_Cl2)	see note
H6200f_a01	TrAa01ClMblScScm	$HCl \rightarrow HCl(aq)$	k_exf(01,ind_HCl)	see note
H6200b_a01	TrAa01ClMblScScm	$HCl(aq) \rightarrow HCl$	k_exb(01,ind_HCl)	see note
H6201f_a01	TrAa01ClMblSc	$HOCl \rightarrow HOCl(aq)$	k_exf(01,ind_HOCl)	see note
H6201b_a01	TrAa01ClMblSc	$HOCl(aq) \rightarrow HOCl$	k_exb(01,ind_HOCl)	see note
H6300_a01	TrAa01ClMblN	$N_2O_5 + Cl^-(aq) \rightarrow ClNO_2 + NO_3^-(aq)$	k_exf_N2O5(01) * 5.E2	Behnke et al. (1994), Behnke et al. (1997)*
H6301_a01	TrAa01ClMblN	$ClNO_3 \rightarrow HOCl(aq) + HNO_3(aq)$	k_exf_ClNO3(01) * C(ind_H2O_a01)	see note
H6302_a01	TrAa01ClMblN	$ClNO_3 + Cl^-(aq) \rightarrow Cl_2(aq) + NO_3^-(aq)$	k_exf_ClNO3(01) * 5.E2	see note
H7000f_a01	TrAa01BrMblSc	$Br_2 \rightarrow Br_2(aq)$	k_exf(01,ind_Br2)	see note
H7000b_a01	TrAa01BrMblSc	$Br_2(aq) \rightarrow Br_2$	k_exb(01,ind_Br2)	see note
H7200f_a01	TrAa01BrMblScScm	$HBr \rightarrow HBr(aq)$	k_exf(01,ind_HBr)	see note
H7200b_a01	TrAa01BrMblScScm	$HBr(aq) \rightarrow HBr$	k_exb(01,ind_HBr)	see note
H7201f_a01	TrAa01BrMblSc	$HOBr \rightarrow HOBr(aq)$	k_exf(01,ind_HOBr)	see note
H7201b_a01	TrAa01BrMblSc	$HOBr(aq) \rightarrow HOBr$	k_exb(01,ind_HOBr)	see note
H7300_a01	TrAa01BrMblN	$N_2O_5 + Br^-(aq) \rightarrow BrNO_2 + NO_3^-(aq)$	k_exf_N2O5(01) * 3.E5	Behnke et al. (1994), Behnke et al. (1997)*
H7301_a01	TrAa01BrMblN	$BrNO_3 \rightarrow HOBr(aq) + HNO_3(aq)$	k_exf_BrNO3(01) * C(ind_H2O_a01)	see note
H7302_a01	TrAa01BrMblN	$BrNO_3 + Br^-(aq) \rightarrow Br_2(aq) + NO_3^-(aq)$	k_exf_BrNO3(01) * 3.E5	see note
H7600f_a01	TrAa01ClBrMblSc	$BrCl \rightarrow BrCl(aq)$	k_exf(01,ind_BrCl)	see note
H7600b_a01	TrAa01ClBrMblSc	$BrCl(aq) \rightarrow BrCl$	k_exb(01,ind_BrCl)	see note
H7601_a01	TrAa01ClBrMblN	$ClNO_3 + Br^-(aq) \rightarrow BrCl(aq) + NO_3^-(aq)$	k_exf_ClNO3(01) * 3.E5	see note
H7602_a01	TrAa01ClBrMblN	$BrNO_3 + Cl^-(aq) \rightarrow BrCl(aq) + NO_3^-(aq)$	k_exf_BrNO3(01) * 5.E2	see note
H8000f_a01	TrAa01ISc	$I_2 \rightarrow I_2(aq)$	k_exf(01,ind_I2)	see note
H8000b_a01	TrAa01ISc	$I_2(aq) \rightarrow I_2$	k_exb(01,ind_I2)	see note
H8100f_a01	TrAa01IMblSc	$IO \rightarrow IO(aq)$	k_exf(01,ind_IO)	see note
H8100b_a01	TrAa01IMblSc	$IO(aq) \rightarrow IO$	k_exb(01,ind_IO)	see note
H8101_a01	TrAa01I	$OIO \rightarrow HOI(aq) + HO_2(aq)$	k_exf(01,ind_OIO)	see note
H8102_a01	TrAa01I	$I_2O_2 \rightarrow HOI(aq) + H^+(aq) + IO_2^-(aq)$	k_exf(01,ind_I2O2)	see note
H8200f_a01	TrAa01IMblSc	$HOI \rightarrow HOI(aq)$	k_exf(01,ind_HOI)	see note
H8200b_a01	TrAa01IMblSc	$HOI(aq) \rightarrow HOI$	k_exb(01,ind_HOI)	see note
H8201_a01	TrAa01IMblSc	$HI \rightarrow H^+(aq) + I^-(aq)$	$k_{mt}(HI) \cdot lwc$	see note

Table 5: Reversible (Henry's law) equilibria and irreversible ("heterogenous") uptake

#	labels	reaction	rate coefficient	reference
H8202_a01	TrAa01ISc	$HIO_3 \rightarrow IO_3^-(aq) + H^+(aq)$	$k_{mt}(HIO_3) \cdot lwc$	see note
H8300_a01	TrAa01I	$INO_2 \rightarrow HOI(aq) + HONO(aq)$	k_exf(01,ind_INO2)	see note
H8301_a01	TrAa01IMbl	$INO_3 \rightarrow HOI(aq) + HNO_3(aq)$	k_exf(01,ind_INO3)	see note
H8600f_a01	TrAa01ClIMblSc	$ICl \rightarrow ICl(aq)$	k_exf(01,ind_ICl)	see note
H8600b_a01	TrAa01ClIMblSc	$ICl(aq) \rightarrow ICl$	k_exb(01,ind_ICl)	see note
H8700f_a01	TrAa01BrIMblSc	$IBr \rightarrow IBr(aq)$	k_exf(01,ind_IBr)	see note
H8700b_a01	TrAa01BrIMblSc	$IBr(aq) \rightarrow IBr$	k_exb(01,ind_IBr)	see note
H9100f_a01	TrAa01SMblScScm	$SO_2 \rightarrow SO_2(aq)$	k_exf(01,ind_SO2)	see note
H9100b_a01	TrAa01SMblScScm	$SO_2(aq) \rightarrow SO_2$	k_exb(01,ind_SO2)	see note
H9200_a01	TrAa01SMblScScm	$H_2SO_4 \rightarrow H_2SO_4(aq)$	xnom7sulf*k_exf(01,ind_H2SO4)	see note
H9400f_a01	TrAa01S	$DMSO \rightarrow DMSO(aq)$	k_exf(01,ind_DMSO)	see note
H9400b_a01	TrAa01S	$DMSO(aq) \rightarrow DMSO$	k_exb(01,ind_DMSO)	see note
H9401_a01	TrAa01SMbl	$CH_3SO_3H \rightarrow CH_3SO_3^-(aq) + H^+(aq)$	k_exf(01,ind_CH3SO3H)	see note
H9402f_a01	TrAa01S	$DMS \rightarrow DMS(aq)$	k_exf(01,ind_DMS)	see note
H9402b_a01	TrAa01S	$DMS(aq) \rightarrow DMS$	k_exb(01,ind_DMS)	see note
H10000f_a01	TrAa01Hg	$Hg \rightarrow Hg(aq)$	k_exf(01,ind_Hg)	see note
H10000b_a01	TrAa01Hg	$Hg(aq) \rightarrow Hg$	k_exb(01,ind_Hg)	see note
H10100f_a01	TrAa01Hg	$HgO \rightarrow HgO(aq)$	k_exf(01,ind_HgO)	see note
H10100b_a01	TrAa01Hg	$HgO(aq) \rightarrow HgO$	k_exb(01,ind_HgO)	see note
H10600f_a01	TrAa01HgCl	$HgCl_2 \rightarrow HgCl_2(aq)$	k_exf(01,ind_HgCl2)	see note
H10600b_a01	TrAa01HgCl	$HgCl_2(aq) \rightarrow HgCl_2$	k_exb(01,ind_HgCl2)	see note
H10700f_a01	TrAa01HgBr	$HgBr_2 \rightarrow HgBr_2(aq)$	k_exf(01,ind_HgBr2)	see note
H10700b_a01	TrAa01HgBr	$HgBr_2(aq) \rightarrow HgBr_2$	k_exb(01,ind_HgBr2)	see note
H10701f_a01	TrAa01HgClBr	$ClHgBr \rightarrow ClHgBr(aq)$	k_exf(01,ind_ClHgBr)	see note
H10701b_a01	TrAa01HgClBr	$ClHgBr(aq) \rightarrow ClHgBr$	k_exb(01,ind_ClHgBr)	see note
H10702f_a01	TrAa01HgBr	$BrHgOBr \rightarrow BrHgOBr(aq)$	k_exf(01,ind_BrHgOBr)	see note
H10702b_a01	TrAa01HgBr	$BrHgOBr(aq) \rightarrow BrHgOBr$	k_exb(01,ind_BrHgOBr)	see note
H10703f_a01	TrAa01HgClBr	$ClHgOBr \rightarrow ClHgOBr(aq)$	k_exf(01,ind_ClHgOBr)	see note
H10703b_a01	TrAa01HgClBr	$ClHgOBr(aq) \rightarrow ClHgOBr$	k_exb(01,ind_ClHgOBr)	see note

*Notes:

The forward (k_exf) and backward (k_exb) rate coefficients are calculated in the file messy_mecca_aero.f90 using the accommodation coefficients in subroutine mecca_aero_alpha and Henry's law constants in subroutine mecca_aero_henry.

k_{mt} = mass transfer coefficient

lwc = liquid water content of aerosol mode

H3201, H6300, H6301, H6302, H7300, H7301, H7302,

H7601, H7602: For uptake of X (X = N_2O_5, $ClNO_3$, or $BrNO_3$) and subsequent reaction with H_2O, Cl^-, and Br^-, we define:

$$k_{\mathrm{exf}}(\mathrm{X}) = \frac{k_{\mathrm{mt}}(\mathrm{X}) \times \mathrm{LWC}}{[\mathrm{H_2O}] + 5 \times 10^2 [\mathrm{Cl^-}] + 3 \times 10^5 [\mathrm{Br^-}]}$$

The total uptake rate of X is only determined by k_{mt}. The factors only affect the branching between hydrolysis and the halide reactions. The factor $5{\times}10^2$ was chosen such that the chloride reaction dominates over hydrolysis at about $[Cl^-] > 0.1$ M (see Fig. 3 in Behnke et al. (1997)), i.e. when the ratio $[H_2O]/[Cl^-]$ is less than $5{\times}10^2$. The ratio $5{\times}10^2/3{\times}10^5$ was chosen such that the reactions with chloride and bromide are roughly equal for sea water composition (Behnke et al., 1994).

Table 6: Heterogeneous reactions

#	labels	reaction	rate coefficient	reference
HET200	StHetN	$N_2O_5 + H_2O \rightarrow 2\ HNO_3$	khet_St(ihs_N2O5_H2O)	see note
HET201	TrHetN	$N_2O_5 \rightarrow 2\ NO_3^-(aq) + 2\ H^+(aq)$	khet_Tr(iht_N2O5)	see note
HET410	StHetCl	$HOCl + HCl \rightarrow Cl_2 + H_2O$	khet_St(ihs_HOCl_HCl)	see note
HET420	StHetNCl	$ClNO_3 + HCl \rightarrow Cl_2 + HNO_3$	khet_St(ihs_ClNO3_HCl)	see note
HET421	StHetNCl	$ClNO_3 + H_2O \rightarrow HOCl + HNO_3$	khet_St(ihs_ClNO3_H2O)	see note
HET422	StHetNCl	$N_2O_5 + HCl \rightarrow ClNO_2 + HNO_3$	khet_St(ihs_N2O5_HCl)	see note
HET510	StHetBr	$HOBr + HBr \rightarrow Br_2 + H_2O$	khet_St(ihs_HOBr_HBr)	see note
HET520	StHetNBr	$BrNO_3 + H_2O \rightarrow HOBr + HNO_3$	khet_St(ihs_BrNO3_H2O)	see note
HET540	StHetNClBr	$ClNO_3 + HBr \rightarrow BrCl + HNO_3$	khet_St(ihs_ClNO3_HBr)	see note
HET541	StHetNClBr	$BrNO_3 + HCl \rightarrow BrCl + HNO_3$	khet_St(ihs_BrNO3_HCl)	see note
HET542	StHetClBr	$HOCl + HBr \rightarrow BrCl + H_2O$	khet_St(ihs_HOCl_HBr)	see note
HET543	StHetClBr	$HOBr + HCl \rightarrow BrCl + H_2O$	khet_St(ihs_HOBr_HCl)	see note
HET1001	StTrHetHg	$Hg \rightarrow Hg(aq)$	khet_Tr(iht_Hg) + khet_St(ihs_Hg)	see note
HET1002	StTrHetHg	$HgO \rightarrow Hg(aq)$	khet_Tr(iht_RGM) + khet_St(ihs_RGM)	see note
HET1003	StTrHetHg	$HgCl \rightarrow Hg(aq)$	khet_Tr(iht_RGM) + khet_St(ihs_RGM)	see note
HET1004	StTrHetHg	$HgCl_2 \rightarrow Hg(aq)$	khet_Tr(iht_RGM) + khet_St(ihs_RGM)	see note
HET1005	StTrHetHg	$HgBr \rightarrow Hg(aq)$	khet_Tr(iht_RGM) + khet_St(ihs_RGM)	see note
HET1006	StTrHetHg	$HgBr_2 \rightarrow Hg(aq)$	khet_Tr(iht_RGM) + khet_St(ihs_RGM)	see note
HET1007	StTrHetHg	$ClHgBr \rightarrow Hg(aq)$	khet_Tr(iht_RGM) + khet_St(ihs_RGM)	see note
HET1008	StTrHetHg	$BrHgOBr \rightarrow Hg(aq)$	khet_Tr(iht_RGM) + khet_St(ihs_RGM)	see note
HET1009	StTrHetHg	$ClHgOBr \rightarrow Hg(aq)$	khet_Tr(iht_RGM) + khet_St(ihs_RGM)	see note

*Notes:

Heterogeneous reaction rates are calculated with an external module and then supplied to the MECCA chemistry (see www.messy-interface.org for details)

Table 7: Acid-base and other eqilibria

#	labels	reaction	$K_0[M^{m-n}]$	$-\Delta H/R[K]$	reference
EQ20_a01	TrAa01Sc	$HO_2 \rightleftharpoons O_2^- + H^+$	1.6E-5		Weinstein-Lloyd and Schwartz (1991)
EQ21_a01	TrAa01MblScScm	$H_2O \rightleftharpoons H^+ + OH^-$	1.0E-16	-6716	Chameides (1984)
EQ30_a01	TrAa01MblNScScm	$NH_4^+ \rightleftharpoons H^+ + NH_3$	5.88E-10	-2391	Chameides (1984)
EQ31_a01	TrAa01NSc	$HONO \rightleftharpoons H^+ + NO_2^-$	5.1E-4	-1260	Schwartz and White (1981)
EQ32_a01	TrAa01MblNScScm	$HNO_3 \rightleftharpoons H^+ + NO_3^-$	15	8700	Davis and de Bruin (1964)
EQ33_a01	TrAa01NSc	$HNO_4 \rightleftharpoons NO_4^- + H^+$	1.E-5		Warneck (1999)
EQ40_a01	TrAa01MblScScm	$CO_2 \rightleftharpoons H^+ + HCO_3^-$	4.3E-7	-913	Chameides (1984)*
EQ41_a01	TrAa01ScScm	$HCOOH \rightleftharpoons H^+ + HCOO^-$	1.8E-4		Weast (1980)
EQ60_a01	TrAa01Cl	$Cl_2^- \rightleftharpoons Cl + Cl^-$	7.3E-6		Yu (2004)
EQ61_a01	TrAa01ClMblScScm	$HCl \rightleftharpoons H^+ + Cl^-$	1.7E6	6896	Marsh and McElroy (1985)
EQ62_a01	TrAa01ClSc	$HOCl \rightleftharpoons H^+ + ClO^-$	3.2E-8		Lax (1969)
EQ70_a01	TrAa01Br	$Br_2^- \rightleftharpoons Br + Br^-$	2.54E-6	-2256	Liu et al. (2002)
EQ71_a01	TrAa01BrMblScScm	$HBr \rightleftharpoons H^+ + Br^-$	1.0E9		Lax (1969)
EQ72_a01	TrAa01BrSc	$HOBr \rightleftharpoons H^+ + BrO^-$	2.3E-9	-3091	Kelley and Tartar (1956)*
EQ73_a01	TrAa01ClBrMbl	$BrCl + Cl^- \rightleftharpoons BrCl_2^-$	3.8	1191	Wang et al. (1994)
EQ74_a01	TrAa01ClBrMbl	$BrCl + Br^- \rightleftharpoons Br_2Cl^-$	1.8E4	7457	Wang et al. (1994)
EQ75_a01	TrAa01ClBrMbl	$Br_2 + Cl^- \rightleftharpoons Br_2Cl^-$	1.3	0	Wang et al. (1994)
EQ76_a01	TrAa01ClBrMbl	$Br^- + Cl_2 \rightleftharpoons BrCl_2^-$	4.2E6	14072	Wang et al. (1994)
EQ80_a01	TrAa01ClIMblSc	$ICl + Cl^- \rightleftharpoons ICl_2^-$	7.7E1		Wang et al. (1989)
EQ81_a01	TrAa01BrIMblSc	$IBr + Br^- \rightleftharpoons IBr_2^-$	2.9E2		Troy and Margerum (1991)
EQ82_a01	TrAa01ClBrIMblSc	$ICl + Br^- \rightleftharpoons IBr + Cl^-$	3.3E2		see note
EQ90_a01	TrAa01SMblScScm	$SO_2 \rightleftharpoons H^+ + HSO_3^-$	1.7E-2	2090	Chameides (1984)
EQ91_a01	TrAa01SMblScScm	$HSO_3^- \rightleftharpoons H^+ + SO_3^{2-}$	6.0E-8	1120	Chameides (1984)
EQ92_a01	TrAa01SMblScScm	$HSO_4^- \rightleftharpoons H^+ + SO_4^{2-}$	1.2E-2	2720	Seinfeld and Pandis (1998)
EQ93_a01	TrAa01SMblScScm	$H_2SO_4 \rightleftharpoons H^+ + HSO_4^-$	1.0E3		Seinfeld and Pandis (1998)
EQ100_a01	TrAa01Hg	$Hg^{2+} + OH^- \rightleftharpoons HgOH^+$	4.0E10		Ammann and Pöschl (2007)
EQ101_a01	TrAa01Hg	$HgOH^+ + OH^- \rightleftharpoons Hg(OH)_2$	1.58E11		Ammann and Pöschl (2007)
EQ102_a01	TrAa01HgCl	$Hg^{2+} + Cl^- \rightleftharpoons HgCl^+$	5.8E6		Ammann and Pöschl (2007)
EQ103_a01	TrAa01HgCl	$HgCl^+ + Cl^- \rightleftharpoons HgCl_2$	2.5E6		Ammann and Pöschl (2007)
EQ104_a01	TrAa01HgCl	$HgOH^+ + Cl^- \rightleftharpoons Hg(OH)Cl$	2.69E7		Ammann and Pöschl (2007)
EQ105_a01	TrAa01HgBr	$Hg^{2+} + Br^- \rightleftharpoons HgBr^+$	1.1E9		Raofie and Ariya (2004)
EQ106_a01	TrAa01HgBr	$HgBr^+ + Br^- \rightleftharpoons HgBr_2$	2.5E8		Raofie and Ariya (2004)
EQ107_a01	TrAa01HgS	$Hg^{2+} + SO_3^{2-} \rightleftharpoons HgSO_3$	2.E13		van Loon et al. (2001)
EQ108_a01	TrAa01HgS	$HgSO_3 + SO_3^{2-} \rightleftharpoons Hg(SO_3)_2^{2-}$	1.E10		van Loon et al. (2001)

*Notes:

EQ40: For $pK_a(CO_2)$, see also Dickson and Millero (1987).

EQ72: For pK_a(HOBr), see also Keller-Rudek et al. (1992).

EQ82: Thermodynamic calculations on the IBr/ICl equilibrium according to the data tables from Wagman et al. (1982):

ICl	+	Br^-	$\rightleftharpoons$	IBr	+	Cl^-
-17.1		-103.96	=	-4.2		-131.228

$$\frac{\Delta G}{[\text{kJ/mol}]} = -4.2 - 131.228 - (-17.1 - 103.96) = -14.368$$

$$K = \frac{[\text{IBr}] \times [\text{Cl}^-]}{[\text{ICl}] \times [\text{Br}^-]} = \exp\left(\frac{-\Delta G}{RT}\right) = \exp\left(\frac{14368}{8.314 \times 298}\right) = 330$$

This means we have equal amounts of IBr and ICl when the $[Cl^-]/[Br^-]$ ratio equals 330.

Table 8: Aqueous phase reactions

#	labels	reaction	k_0 $[M^{1-n}s^{-1}]$	$-E_a/R[K]$	reference
A1000_a01	TrAa01Sc	$O_3 + O_2^- \rightarrow OH + OH^-$	1.5E9		Sehested et al. (1983)
A2100_a01	TrAa01Sc	$OH + O_2^- \rightarrow OH^-$	1.0E10		Sehested et al. (1968)
A2101_a01	TrAa01Sc	$OH + OH \rightarrow H_2O_2$	5.5E9		Buxton et al. (1988)
A2102_a01	TrAa01Sc	$HO_2 + O_2^- \rightarrow H_2O_2 + OH^-$	1.0E8	-900	Christensen and Sehested (1988)
A2103_a01	TrAa01Sc	$HO_2 + OH \rightarrow H_2O$	7.1E9		Sehested et al. (1968)
A2104_a01	TrAa01Sc	$HO_2 + HO_2 \rightarrow H_2O_2$	9.7E5	-2500	Christensen and Sehested (1988)
A2105_a01	TrAa01Sc	$H_2O_2 + OH \rightarrow HO_2$	2.7E7	-1684	Christensen et al. (1982)
A3100_a01	TrAa01NSc	$NO_2^- + O_3 \rightarrow NO_3^-$	5.0E5	-6950	Damschen and Martin (1983)
A3101_a01	TrAa01NSc	$NO_2 + NO_2 \rightarrow HNO_3 + HONO$	1.0E8		Lee and Schwartz (1981)
A3102_a01	TrAa01NSc	$NO_4^- \rightarrow NO_2^-$	8.0E1		Warneck (1999)
A3200_a01	TrAa01NSc	$NO_2 + HO_2 \rightarrow HNO_4$	1.8E9		Warneck (1999)
A3201_a01	TrAa01NSc	$NO_2^- + OH \rightarrow NO_2 + OH^-$	1.0E10		Wingenter et al. (1999)
A3202_a01	TrAa01NSc	$NO_3 + OH^- \rightarrow NO_3^- + OH$	8.2E7	-2700	Exner et al. (1992)
A3203_a01	TrAa01NSc	$HONO + OH \rightarrow NO_2$	1.0E10		Barker et al. (1970)
A3204_a01	TrAa01NSc	$HONO + H_2O_2 + H^+ \rightarrow HNO_3 + H^+$	4.6E3	-6800	Damschen and Martin (1983)
A4100_a01	TrAa01Sc	$CO_3^- + O_2^- \rightarrow HCO_3^- + OH^-$	6.5E8		Ross et al. (1992)
A4101_a01	TrAa01Sc	$CO_3^- + H_2O_2 \rightarrow HCO_3^- + HO_2$	4.3E5		Ross et al. (1992)
A4102_a01	TrAa01Sc	$HCOO^- + CO_3^- \rightarrow 2\ HCO_3^- + HO_2$	1.5E5		Ross et al. (1992)
A4103_a01	TrAa01Sc	$HCOO^- + OH \rightarrow OH^- + HO_2 + CO_2$	3.1E9	-1240	Chin and Wine (1994)
A4104_a01	TrAa01Sc	$HCO_3^- + OH \rightarrow CO_3^-$	8.5E6		Ross et al. (1992)
A4105_a01	TrAa01Sc	$HCHO + OH \rightarrow HCOOH + HO_2$	7.7E8	-1020	Chin and Wine (1994)
A4106_a01	TrAa01Sc	$HCOOH + OH \rightarrow HO_2 + CO_2$	1.1E8	-991	Chin and Wine (1994)
A4107_a01	TrAa01Sc	$CH_3OO + O_2^- \rightarrow CH_3OOH + OH^-$	5.0E7		Jacob (1986)
A4108_a01	TrAa01Sc	$CH_3OO + HO_2 \rightarrow CH_3OOH$	4.3E5		Jacob (1986)
A4109_a01	TrAa01Sc	$CH_3OH + OH \rightarrow HCHO + HO_2$	9.7E8		Buxton et al. (1988)
A4110a_a01	TrAa01Sc	$CH_3OOH + OH \rightarrow CH_3OO$	2.7E7	-1715	Jacob (1986)
A4110b_a01	TrAa01Sc	$CH_3OOH + OH \rightarrow HCHO + OH$	1.1E7	-1715	Jacob (1986)
A6000_a01	TrAa01Cl	$Cl + Cl \rightarrow Cl_2$	8.8E7		Wu et al. (1980)
A6001_a01	TrAa01Cl	$Cl_2^- + Cl_2^- \rightarrow Cl_2 + 2\ Cl^-$	3.5E9		Yu (2004)
A6100_a01	TrAa01Cl	$Cl^- + O_3 \rightarrow ClO^-$	3.0E-3		Hoigné et al. (1985)
A6101_a01	TrAa01Cl	$Cl_2 + O_2^- \rightarrow Cl_2^-$	1.0E9		Bjergbakke et al. (1981)
A6102_a01	TrAa01Cl	$Cl_2^- + O_2^- \rightarrow 2\ Cl^-$	1.0E9		Jacobi (1996)*
A6200_a01	TrAa01Cl	$Cl \rightarrow H^+ + ClOH^-$	1.8E5		Yu (2004)
A6201_a01	TrAa01Cl	$Cl + H_2O_2 \rightarrow HO_2 + Cl^- + H^+$	2.7E7	-1684	Christensen et al. (1982)

Table 8: Aqueous phase reactions (...continued)

#	labels	reaction	k_0 $[M^{1-n}s^{-1}]$	$-E_a/R[K]$	reference
A6202_a01	TrAa01Cl	$Cl^- + OH \rightarrow ClOH^-$	4.2E9		Yu (2004)
A6203_a01	TrAa01Cl	$Cl_2 + HO_2 \rightarrow Cl_2^- + H^+$	1.0E9		Bjergbakke et al. (1981)
A6204_a01	TrAa01ClMbl	$Cl_2 \rightarrow Cl^- + HOCl + H^+$	21.8	-8012	Wang and Margerum (1994)
A6205_a01	TrAa01Cl	$Cl_2^- + HO_2 \rightarrow 2\ Cl^- + H^+$	1.3E10		Jacobi (1996)
A6206_a01	TrAa01Cl	$HOCl + O_2^- \rightarrow Cl + OH^-$	7.5E6		Long and Bielski (1980)
A6207_a01	TrAa01Cl	$HOCl + HO_2 \rightarrow Cl$	7.5E6		Long and Bielski (1980)
A6208_a01	TrAa01ClMbl	$HOCl + Cl^- + H^+ \rightarrow Cl_2$	2.2E4	-3508	Wang and Margerum (1994)
A6209_a01	TrAa01Cl	$ClOH^- \rightarrow Cl^- + OH$	6.0E9		Yu (2004)
A6210_a01	TrAa01Cl	$ClOH^- + H^+ \rightarrow Cl$	2.4E10		Yu (2004)
A6300_a01	TrAa01Cl	$Cl + NO_3^- \rightarrow NO_3 + Cl^-$	1.0E8		Buxton et al. (1999b)
A6301_a01	TrAa01Cl	$Cl^- + NO_3 \rightarrow NO_3^- + Cl$	3.4E8		Buxton et al. (1999b)*
A6302_a01	TrAa01Cl	$Cl_2^- + NO_2^- \rightarrow 2\ Cl^- + NO_2$	6.0E7		Jacobi et al. (1996)
A6400_a01	TrAa01Cl	$Cl_2^- + CH_3OOH \rightarrow 2\ Cl^- + H^+ + CH_3OO$	5.0E4		Jacobi et al. (1996)
A7000_a01	TrAa01Br	$Br_2^- + Br_2^- \rightarrow 2\ Br^- + Br_2$	1.9E9		Ross et al. (1992)
A7100_a01	TrAa01Br	$Br^- + O_3 \rightarrow BrO^-$	2.1E2	-4450	Haag and Hoigné (1983)
A7101_a01	TrAa01Br	$Br_2 + O_2^- \rightarrow Br_2^-$	5.6E9		Sutton and Downes (1972)
A7102_a01	TrAa01Br	$Br_2^- + O_2^- \rightarrow 2\ Br^-$	1.7E8		Wagner and Strehlow (1987)
A7200_a01	TrAa01Br	$Br^- + OH \rightarrow BrOH^-$	1.1E10		Zehavi and Rabani (1972)
A7201_a01	TrAa01Br	$Br_2 + HO_2 \rightarrow Br_2^- + H^+$	1.1E8		Sutton and Downes (1972)
A7202_a01	TrAa01BrMbl	$Br_2 \rightarrow Br^- + HOBr + H^+$	9.7E1	-7457	Beckwith et al. (1996)
A7203_a01	TrAa01Br	$Br_2^- + HO_2 \rightarrow Br_2 + H_2O_2 + OH^-$	4.4E9		Matthew et al. (2003)
A7204_a01	TrAa01Br	$Br_2^- + H_2O_2 \rightarrow 2\ Br^- + H^+ + HO_2$	1.0E5		Jacobi (1996)
A7205_a01	TrAa01Br	$HOBr + O_2^- \rightarrow Br + OH^-$	3.5E9		Schwarz and Bielski (1986)
A7206_a01	TrAa01Br	$HOBr + HO_2 \rightarrow Br$	1.0E9		Herrmann et al. (1999)
A7207_a01	TrAa01Br	$HOBr + H_2O_2 \rightarrow Br^- + H^+$	1.2E6		Bichsel and von Gunten (1999)
A7208_a01	TrAa01BrMbl	$HOBr + Br^- + H^+ \rightarrow Br_2$	1.6E10		Beckwith et al. (1996)
A7209a_a01	TrAa01Br	$BrOH^- \rightarrow Br^- + OH$	3.3E7		Zehavi and Rabani (1972)
A7209b_a01	TrAa01Br	$BrOH^- \rightarrow Br + OH^-$	4.2E6		Zehavi and Rabani (1972)
A7210_a01	TrAa01Br	$BrOH^- + H^+ \rightarrow Br$	4.4E10		Zehavi and Rabani (1972)
A7300_a01	TrAa01Br	$Br^- + NO_3 \rightarrow Br + NO_3^-$	4.0E9		Neta and Huie (1986)
A7301_a01	TrAa01Br	$Br_2^- + NO_2^- \rightarrow 2\ Br^- + NO_2$	1.7E7	-1720	Shoute et al. (1991)
A7400_a01	TrAa01Br	$Br_2^- + CH_3OOH \rightarrow 2\ Br^- + H^+ + CH_3OO$	1.0E5		Jacobi (1996)*
A7601_a01	TrAa01Br	$Br^- + ClO^- + H^+ \rightarrow BrCl + OH^-$	3.7E10		Kumar and Margerum (1987)
A7602_a01	TrAa01ClBrMbl	$Br^- + HOCl + H^+ \rightarrow BrCl$	1.32E6		Kumar and Margerum (1987)

Table 8: Aqueous phase reactions (...continued)

#	labels	reaction	k_0 $[M^{1-n}s^{-1}]$	$-E_a/R[K]$	reference
A7603_a01	TrAa01ClBrMbl	$HOBr + Cl^- + H^+ \rightarrow BrCl$	2.3E10		see note
A7604_a01	TrAa01ClBrMbl	$BrCl \rightarrow Cl^- + HOBr + H^+$	3.0E6		Liu and Margerum (2001)
A8100_a01	TrAa01IMbl	$I^- + O_3 \rightarrow HOI + OH^-$	4.2E9	-9311	Magi et al. (1997)
A8101_a01	TrAa01IMbl	$IO + IO \rightarrow HOI + IO_2^- + H^+$	1.5E9		Buxton et al. (1986)
A8200_a01	TrAa01IMbl	$IO_2^- + H_2O_2 \rightarrow IO_3^-$	6.0E1		Furrow (1987)
A8201_a01	TrAa01I	$HOI + IO_2^- \rightarrow IO_3^- + I^- + H^+$	6.0E2		Chinake and Simoyi (1996)
A8202_a01	TrAa01IMbl	$HOI + I^- + H^+ \rightarrow I_2$	4.4E12		Eigen and Kustin (1962)
A8203_a01	TrAa01IMbl	$IO_2^- + I^- + H^+ \rightarrow 2\ HOI + OH^-$	2.0E10		Edblom et al. (1987)
A8600_a01	TrAa01ClIMbl	$ICl \rightarrow HOI + Cl^- + H^+$	2.4E6		Wang et al. (1989)
A8601_a01	TrAa01ClIMbl	$I^- + HOCl + H^+ \rightarrow ICl$	3.5E11		Nagy et al. (1988)
A8602_a01	TrAa01ClI	$IO_2^- + HOCl \rightarrow IO_3^- + Cl^- + H^+$	1.5E3		Lengyel et al. (1996)
A8603_a01	TrAa01ClIMbl	$HOI + Cl^- + H^+ \rightarrow ICl$	2.9E10		Wang et al. (1989)
A8604_a01	TrAa01ClI	$HOI + Cl_2 \rightarrow IO_2^- + 2\ Cl^- + 3H^+$	1.0E6		Lengyel et al. (1996)
A8605_a01	TrAa01ClI	$HOI + HOCl \rightarrow IO_2^- + Cl^- + 2\ H^+$	5.0E5		Citri and Epstein (1988)
A8606_a01	TrAa01ClI	$ICl + I^- \rightarrow I_2 + Cl^-$	1.1E9		Margerum et al. (1986)
A8700_a01	TrAa01BrIMbl	$IBr \rightarrow HOI + H^+ + Br^-$	8.0E5		Troy et al. (1991)
A8701_a01	TrAa01BrIMbl	$I^- + HOBr \rightarrow IBr + OH^-$	5.0E9		Troy and Margerum (1991)
A8702_a01	TrAa01BrI	$IO_2^- + HOBr \rightarrow IO_3^- + Br^- + H^+$	1.0E6		Chinake and Simoyi (1996)
A8703_a01	TrAa01BrIMbl	$HOI + Br^- + H^+ \rightarrow IBr$	3.3E12		Troy et al. (1991)
A8704_a01	TrAa01BrI	$HOI + HOBr \rightarrow IO_2^- + Br^- + 2\ H^+$	1.0E6		Chinake and Simoyi (1996)
A8705_a01	TrAa01ClI	$IBr + I^- \rightarrow I_2 + Br^-$	2.0E9		Faria et al. (1993)
A9100_a01	TrAa01SSc	$SO_3^- + O_2 \rightarrow SO_5^-$	1.5E9		Huie and Neta (1987)
A9101_a01	TrAa01SMblScScm	$SO_3^{2-} + O_3 \rightarrow SO_4^{2-}$	1.5E9	-5300	Hoffmann (1986)
A9102_a01	TrAa01SSc	$SO_4^- + O_2^- \rightarrow SO_4^{2-}$	3.5E9		Jiang et al. (1992)
A9103_a01	TrAa01SSc	$SO_4^- + SO_3^{2-} \rightarrow SO_3^- + SO_4^{2-}$	4.6E8		Huie and Neta (1987)
A9104_a01	TrAa01SSc	$SO_5^- + O_2^- \rightarrow HSO_5^- + OH^-$	2.3E8		Buxton et al. (1996)
A9105_a01	TrAa01S	$SO_5^- + SO_3^{2-} \rightarrow .72\ SO_4^- + .72\ SO_4^{2-} + .28\ SO_3^- + .28\ HSO_5^- + .28\ OH^-$	1.3E7		Huie and Neta (1987), Deister and Warneck (1990)*
A9106_a01	TrAa01S	$SO_5^- + SO_5^- \rightarrow O_2 + SO_4^{2-}$	1.0E8		Ross et al. (1992)*
A9200_a01	TrAa01SSc	$SO_3^{2-} + OH \rightarrow SO_3^- + OH^-$	5.5E9		Buxton et al. (1988)
A9201_a01	TrAa01SSc	$SO_4^- + OH \rightarrow HSO_5^-$	1.0E9		Jiang et al. (1992)
A9202_a01	TrAa01SSc	$SO_4^- + HO_2 \rightarrow SO_4^{2-} + H^+$	3.5E9		Jiang et al. (1992)
A9203_a01	TrAa01SSc	$SO_4^- + H_2O \rightarrow SO_4^{2-} + H^+ + OH$	1.1E1	-1110	Herrmann et al. (1995)
A9204_a01	TrAa01SSc	$SO_4^- + H_2O_2 \rightarrow SO_4^{2-} + H^+ + HO_2$	1.2E7		Wine et al. (1989)

Table 8: Aqueous phase reactions (...continued)

#	labels	reaction	k_0 $[M^{1-n}s^{-1}]$	$-E_a/R[K]$	reference
A9205_a01	TrAa01SSc	$HSO_3^- + O_2^- \rightarrow SO_4^{2-} + OH$	3.0E3		see note
A9206_a01	TrAa01SMblScScm	$HSO_3^- + O_3 \rightarrow SO_4^{2-} + H^+$	3.7E5	-5500	Hoffmann (1986)
A9207_a01	TrAa01SSc	$HSO_3^- + OH \rightarrow SO_3^-$	4.5E9		Buxton et al. (1988)
A9208_a01	TrAa01SSc	$HSO_3^- + HO_2 \rightarrow SO_4^{2-} + OH + H^+$	3.0E3		see note
A9209_a01	TrAa01SMblScScm	$HSO_3^- + H_2O_2 \rightarrow SO_4^{2-} + H^+$	5.2E6	-3650	Martin and Damschen (1981)
A9210_a01	TrAa01SSc	$HSO_3^- + SO_4^- \rightarrow SO_3^- + SO_4^{2-} + H^+$	8.0E8		Huie and Neta (1987)
A9211_a01	TrAa01S	$HSO_3^- + SO_5^- \rightarrow .75\ SO_4^- + .75\ SO_4^{2-} + .75\ H^+ + .25\ SO_3^- + .25\ HSO_5^-$	1.0E5		Huie and Neta (1987)
A9212_a01	TrAa01SSc	$HSO_3^- + HSO_5^- + H^+ \rightarrow 2\ HSO_4^- + H^+$	7.1E6		Betterton and Hoffmann (1988)
A9300_a01	TrAa01SSc	$SO_3^{2-} + NO_2 \rightarrow SO_4^{2-} + 2\ HONO - NO_2$	2.0E7		Clifton et al. (1988)
A9301_a01	TrAa01SSc	$SO_4^- + NO_3^- \rightarrow SO_4^{2-} + NO_3$	5.0E4		Exner et al. (1992)
A9302_a01	TrAa01SSc	$SO_4^{2-} + NO_3 \rightarrow NO_3^- + SO_4^-$	1.0E5		Logager et al. (1993)
A9303_a01	TrAa01SSc	$HSO_3^- + NO_2 \rightarrow HSO_4^- + 2\ HONO - NO_2$	2.0E7		Clifton et al. (1988)
A9304_a01	TrAa01SSc	$HSO_3^- + NO_3 \rightarrow SO_3^- + NO_3^- + H^+$	1.4E9	-2000	Exner et al. (1992)
A9305_a01	TrAa01SSc	$HSO_3^- + HNO_4 \rightarrow HSO_4^- + NO_3^- + H^+$	3.1E5		Warneck (1999)
A9400_a01	TrAa01SSc	$SO_3^{2-} + HCHO \rightarrow CH_2OHSO_3^- + OH^-$	1.4E4		Boyce and Hoffmann (1984)
A9401_a01	TrAa01SSc	$SO_3^{2-} + CH_3OOH + H^+ \rightarrow SO_4^{2-} + H^+ + CH_3OH$	1.6E7	-3800	Lind et al. (1987)
A9402_a01	TrAa01SSc	$HSO_3^- + HCHO \rightarrow CH_2OHSO_3^-$	4.3E-1		Boyce and Hoffmann (1984)
A9403_a01	TrAa01SSc	$HSO_3^- + CH_3OOH + H^+ \rightarrow HSO_4^- + H^+ + CH_3OH$	1.6E7	-3800	Lind et al. (1987)
A9404_a01	TrAa01SSc	$CH_2OHSO_3^- + OH^- \rightarrow SO_3^{2-} + HCHO$	3.6E3		Seinfeld and Pandis (1998)
A9600_a01	TrAa01SCl	$SO_3^{2-} + Cl_2^- \rightarrow SO_3^- + 2\ Cl^-$	6.2E7		Jacobi et al. (1996)
A9601_a01	TrAa01SClMbl	$SO_3^{2-} + HOCl \rightarrow Cl^- + HSO_4^-$	7.6E8		Fogelman et al. (1989)
A9602_a01	TrAa01SCl	$SO_4^- + Cl^- \rightarrow SO_4^{2-} + Cl$	2.5E8		Buxton et al. (1999a)
A9603_a01	TrAa01SCl	$SO_4^{2-} + Cl \rightarrow SO_4^- + Cl^-$	2.1E8		Buxton et al. (1999a)
A9604_a01	TrAa01SCl	$HSO_3^- + Cl_2^- \rightarrow SO_3^- + 2\ Cl^- + H^+$	4.7E8	-1082	Shoute et al. (1991)
A9605_a01	TrAa01SClMbl	$HSO_3^- + HOCl \rightarrow Cl^- + HSO_4^- + H^+$	7.6E8		see note
A9606_a01	TrAa01SCl	$HSO_5^- + Cl^- \rightarrow HOCl + SO_4^{2-}$	1.8E-3	-7352	Fortnum et al. (1960)
A9700_a01	TrAa01SBr	$SO_3^{2-} + Br_2^- \rightarrow 2\ Br^- + SO_3^-$	2.2E8	-649	Shoute et al. (1991)
A9701_a01	TrAa01SBr	$SO_3^{2-} + BrO^- \rightarrow Br^- + SO_4^{2-}$	1.0E8		Troy and Margerum (1991)
A9702_a01	TrAa01SBrMbl	$SO_3^{2-} + HOBr \rightarrow Br^- + HSO_4^-$	5.0E9		Troy and Margerum (1991)
A9703_a01	TrAa01SBr	$SO_4^- + Br^- \rightarrow Br + SO_4^{2-}$	2.1E9		Jacobi (1996)
A9704_a01	TrAa01SBr	$HSO_3^- + Br_2^- \rightarrow 2\ Br^- + H^+ + SO_3^-$	6.3E7	-782	Shoute et al. (1991)
A9705_a01	TrAa01SBrMbl	$HSO_3^- + HOBr \rightarrow Br^- + HSO_4^- + H^+$	5.0E9		see note
A9706_a01	TrAa01SBr	$HSO_5^- + Br^- \rightarrow HOBr + SO_4^{2-}$	1.0E0	-5338	Fogelman et al. (1989)

Table 8: Aqueous phase reactions (...continued)

#	labels	reaction	k_0 $[M^{1-n}s^{-1}]$	$-E_a/R[K]$	reference
A9800_a01	TrAa01SI	$HSO_3^- + I_2 \rightarrow 2\ I^- + HSO_4^- + 2\ H^+$	1.7E9		Yiin and Margerum (1990)
A10100_a01	TrAa01Hg	$Hg + O_3 \rightarrow HgO + O_2$	4.7E7		Munthe (1992)
A10200_a01	TrAa01Hg	$HgO + H^+ \rightarrow Hg^{2+} + OH^-$	1.0E10		Pleijel and Munthe (1995)
A10201_a01	TrAa01Hg	$Hg + OH \rightarrow Hg^+ + OH^-$	2.0E9		Lin and Pehkonen (1997)
A10202_a01	TrAa01Hg	$Hg^+ + OH \rightarrow Hg^{2+} + OH^-$	1.0E10		Lin and Pehkonen (1997)
A10203_a01	TrAa01Hg	$Hg^{2+} + HO_2 \rightarrow Hg^+ + O_2 + H^+$	1.7E4		Enami et al. (2007)
A10204_a01	TrAa01Hg	$Hg^+ + HO_2 \rightarrow Hg + O_2 + H^+$	1.0E10		Lin and Pehkonen (1997)*
A10600_a01	TrAa01HgCl	$Hg + HOCl \rightarrow Hg^{2+} + Cl^- + OH^-$	2.09E6		Lin and Pehkonen (1998)
A10601_a01	TrAa01HgCl	$Hg + ClO^- \rightarrow Hg^{2+} + Cl^- + OH^-$	1.99E6		Lin and Pehkonen (1998)*
A10700_a01	TrAa01HgBr	$Hg + HOBr \rightarrow Hg^{2+} + Br^- + OH^-$	0.279		Wang and Pehkonen (2004)
A10701_a01	TrAa01HgBr	$Hg + BrO^- \rightarrow Hg^{2+} + Br^- + OH^-$	0.273		Wang and Pehkonen (2004)*
A10702_a01	TrAa01HgBr	$Hg + Br_2 \rightarrow Hg^{2+} + 2\ Br^-$	0.196		Wang and Pehkonen (2004)
A10900_a01	TrAa01HgS	$HgSO_3 \rightarrow Hg + HSO_4^- + H^+$	0.0106		van Loon et al. (2000)

*Notes:

A6102: Jacobi (1996) found an upper limit of 6E9 and cite an upper limit from another study of 2E9. Here, we set the rate coefficient to 1E9.

A6301: There is also an earlier study by Exner et al. (1992) which found a smaller rate coefficient but did not consider the back reaction.

A7400: Assumed to be the same as for $Br_2^- + H_2O_2$.

A9105: The rate coefficient for the sum of the paths (leading to either HSO_5^- or SO_4^{2-}) is from Huie and Neta (1987), the ratio 0.28/0.72 is from Deister and Warneck (1990).

A9106: See also: (Huie and Neta, 1987; Warneck, 1991). If this reaction produces a lot of SO_4^-, it will have an effect. However, we currently assume only the stable $S_2O_8^{2-}$ as product. Since $S_2O_8^{2-}$ is not treated explicitly in the mechanism, we use SO_4^{2-} as a proxy. Note that this destroys the mass consistency for sulfur species.

A9205: D. Sedlak, pers. comm. (1993).

A9208: D. Sedlak, pers. comm. (1993).

A9605: assumed to be the same as for $SO_3^{2-} + HOCl$.

A9705: assumed to be the same as for $SO_3^{2-} + HOBr$.

References

Abbatt, J. P. D. and Waschewsky, G. C. G.: Heterogeneous interactions of HOBr, HNO_3, O_3, and NO_2 with deliquescent NaCl aerosols at room temperature, J. Phys. Chem. A, 102, 3719–3725, 1998.

Ammann, M. and Pöschl, U.: Kinetic model framework for aerosol and cloud surface chemistry and gas-particle interactions - Part 2: exemplary practical applications and numerical simulations, Atmos. Chem. Phys., 7, 6025–6045, 2007.

Anderson, L. C. and Fahey, D. W.: Studies with $ClONO_2$: Thermal dissociation rate and catalytic conversion to NO using an NO/O_3 chemiluminescence detector, J. Phys. Chem., 94, 644–652, 1990.

Aranda, A., Le Bras, G., La Verdet, G., and Poulet, G.: The BrO + CH_3O_2 reaction: Kinetics and role in the atmospheric ozone budget, Geophys. Res. Lett., 24, 2745–2748, 1997.

Ariya, P. A., Khalizov, A., and Gidas, A.: Reactions of gaseous mercury with atomic and molecular halogens: Kinetics, product studies, and atmospheric implications, J. Phys. Chem. A, 106, 7310–7320, 2002.

Atkinson, R.: Gas-phase tropospheric chemistry of volatile organic compounds: 1. Alkanes and alkenes, J. Phys. Chem. Ref. Data, 26, 215–290, 1997.

Atkinson, R.: Kinetics of the gas-phase reactions of OH radicals with alkanes and cycloalkanes, Atmos. Chem. Phys., 3, 2233–2307, 2003.

Atkinson, R., Baulch, D. L., Cox, R. A., Hampson, Jr., R. F., Kerr, J. A., Rossi, M. J., and Troe, J.: Summary of evaluated kinetic and photochemical data for atmospheric chemistry: Web version August 1999, `http://www.iupac-kinetic.ch.cam.ac.uk/`, 1999.

Atkinson, R., Baulch, D. L., Cox, R. A., Crowley, J. N., Hampson, R. F., Hynes, R. G., Jenkin, M. E., Rossi, M. J., and Troe, J.: Evaluated kinetic and photochemical data for atmospheric chemistry: Volume I – gas phase reactions of O_x, HO_x, NO_x and SO_x species, Atmos. Chem. Phys., 4, 1461–1738, 2004.

Atkinson, R., Baulch, D. L., Cox, R. A., Crowley, J. N., Hampson, Jr., R. F., Hynes, R. G., Jenkin, M. E., Kerr, J. A., Rossi, M. J., and Troe, J.: Summary of evaluated kinetic and photochemical data for atmospheric chemistry: Web version March 2005, `http://www.iupac-kinetic.ch.cam.ac.uk/`, 2005.

Atkinson, R., Baulch, D. L., Cox, R. A., Crowley, J. N., Hampson, R. F., Hynes, R. G., Jenkin, M. E., Rossi, M. J., and Troe, J.: Evaluated kinetic and photochemical data for atmospheric chemistry: Volume II - gas phase reactions of organic species, Atmos. Chem. Phys., 6, 3625–4055, 2006.

Atkinson, R., Baulch, D. L., Cox, R. A., Crowley, J. N., Hampson, R. F., Hynes, R. G., Jenkin, M. E., Rossi, M. J., and Troe, J.: Evaluated kinetic and photochemical data for atmospheric chemistry: Volume III — gas phase reactions of inorganic halogens, Atmos. Chem. Phys., 7, 981–1191, 2007.

Bale, C. S. E., Canosa-Mas, C. E., Shallcross, D. E., and Wayne, R. P.: A discharge-flow study of the kinetics of the reactions of IO with CH_3O_2 and CF_3O_2, Phys. Chem. Chem. Phys., 7, 2164–2172, 2005.

Barker, G. C., Fowles, P., and Stringer, B.: Pulse radiolytic induced transient electrical conductance in liquid solutions, Trans. Faraday Soc., 66, 1509–1519, 1970.

Barone, S. B., Turnipseed, A. A., and Ravishankara, A. R.: Role of adducts in the atmospheric oxidation of dimethyl sulfide, Faraday Discuss., 100, 39–54, 1995.

Bartlett, W. P. and Margerum, D. W.: Temperature dependencies of the Henry's law constant and the aqueous phase dissociation constant of bromine chloride, Environ. Sci. Technol., 33, 3410–3414, 1999.

Beckwith, R. C., Wang, T. X., and Margerum, D. W.: Equilibrium and kinetics of bromine hydrolysis, Inorg. Chem., 35, 995–1000, 1996.

Bedjanian, Y., Laverdet, G., and Le Bras, G.: Low-pressure study of the reaction of Cl atoms with isoprene, J. Phys. Chem. A, 102, 953–959, 1998.

Behnke, W., Scheer, V., and Zetzsch, C.: Production of $BrNO_2$, Br_2 and $ClNO_2$ from the reaction between sea spray aerosol and N_2O_5, J. Aerosol Sci., 25, S277–S278, 1994.

Behnke, W., George, C., Scheer, V., and Zetzsch, C.: Production and decay of $ClNO_2$ from the reaction of gaseous N_2O_5 with NaCl solution: Bulk and aerosol experiments, J. Geophys. Res., 102D, 3795–3804, 1997.

Betterton, E. A. and Hoffmann, M. R.: Oxidation of aqueous SO_2 by peroxymonosulfate, J. Phys. Chem., 92, 5962–5965, 1988.

Bichsel, Y. and von Gunten, U.: Oxidation of iodide and hypoiodous acid in the disinfection of natural waters, Environ. Sci. Technol., 33, 4040–4045, 1999.

Bjergbakke, E., Navartnam, S., Parsons, B. J., and Swallow, A. J.: Reaction between $HO_2\cdot$ and chlorine in aqueous solution, J. Am. Chem. Soc., 103, 5926–5928, 1981.

Boyce, S. D. and Hoffmann, M. R.: Kinetics and mechanism of the formation of hydroxymethanesulfonic acid at low pH, J. Phys. Chem., 88, 4740–4746, 1984.

Braban, C. F., Adams, J. W., Rodriguez, D., Cox, R. A., Crowley, J. N., and Schuster, G.: Heterogeneous reactions of HOI, ICl and IBr on sea salt and sea salt proxies, Phys. Chem. Chem. Phys., 9, 3136–3148, 2007.

Brimblecombe, P. and Clegg, S. L.: Erratum, J. Atmos. Chem., 8, 95, 1989.

Burkholder, J. B., Orlando, J. J., and Howard, C. J.: Ultraviolet absorption cross sections of Cl_2O_2 between 210 and 410 nm, J. Phys. Chem., 94, 687–695, 1990.

Buxton, G. V., Kilner, C., and Sellers, R. M.: Pulse radiolysis of HOI and IO^- in aqueous solution. Formation and characterization of I(II), Proc. Tihany Symp. Radiat. Chem., 6, 155–159, 1986.

Buxton, G. V., Greenstock, C. L., Helman, W. P., and Ross, A. B.: Critical review of rate constants for reactions of hydrated electrons, hydrogen atoms and hydroxyl radicals ($\cdot OH/\cdot O^-$) in aqueous solution, J. Phys. Chem. Ref. Data, 17, 513–886, 1988.

Buxton, G. V., McGowan, S., Salmon, G. A., Williams, J. E., and Wood, N. D.: A study of the spectra and reactivity of oxysulphur-radical anions involved in the chain oxidation of S(IV): A pulse and γ-radiolysis study, Atmos. Environ., 30, 2483–2493, 1996.

Buxton, G. V., Bydder, M., and Salmon, G. A.: The reactivity of chlorine atoms in aqueous solution: Part II. The equilibrium $SO_4^{\cdot-} + Cl^- \rightleftharpoons Cl\cdot + SO_4^{2-}$, Phys. Chem. Chem. Phys., 1, 269–273, 1999a.

Buxton, G. V., Salmon, G. A., and Wang, J. Q.: The equilibrium $NO_3^{\cdot} + Cl^- \rightleftharpoons NO_3^- + Cl\cdot$: A laser flash photolysis and pulse radiolysis study of the reactivity of $NO_3^{\cdot}$ with chloride ion in aqueous solution, Phys. Chem. Chem. Phys., 1, 3589–3593, 1999b.

Calvert, J. G. and Lindberg, S. E.: A modeling study of the mechanism of the halogen-ozone-mercury homogeneous reactions in the troposphere during the polar spring, Atmos. Environ., 37, 4467–4481, 2003.

Canosa-Mas, C. E., King, M. D., Lopez, R., Percival, C. J., Wayne, R. P., Shallcross, D. E., Pyle, J. A., and Daele, V.: Is the reaction between $CH_3(O)O_2$ and NO_3 important in the night-time troposphere?, J. Chem. Soc. Faraday Trans., 92, 2211–2222, 1996.

Carl, S. A. and Crowley, J. N.: 298 K rate coefficients for the reaction of OH with $i-C_3H_7I$, $n-C_3H_7I$ and C_3H_8, Atmos. Chem. Phys., 1, 1–7, 2001.

Chameides, W. L.: The photochemistry of a remote marine stratiform cloud, J. Geophys. Res., 89D, 4739–4755, 1984.

Chatfield, R. B. and Crutzen, P. J.: Are there interactions of iodine and sulfur species in marine air photochemistry?, J. Geophys. Res., 95D, 22 319–22 341, 1990.

Chin, M. and Wine, P. H.: A temperature-dependent competitive kinetics study of the aqueous-phase reactions of OH radicals with formate, formic acid, acetate, acetic acid, and hydrated formaldehyde, in: Aquatic and Surface Photochemistry, edited by Helz, G. R., Zepp, R. G., and Crosby, D. G., pp. 85–96, A. F. Lewis, NY, 1994.

Chinake, C. R. and Simoyi, R. H.: Kinetics and mechanism of the complex bromate-iodine reaction, J. Phys. Chem., 100, 1643–1656, 1996.

Christensen, H. and Sehested, K.: HO_2 and O_2^- radicals at elevated temperatures, J. Phys. Chem., 92, 3007–3011, 1988.

Christensen, H., Sehested, K., and Corfitzen, H.: Reactions of hydroxyl radicals with hydrogen peroxide at ambient and elevated temperatures, J. Phys. Chem., 86, 1588–1590, 1982.

Christensen, L. E., Okumura, M., Sander, S. P., Salawitch, R. J., Toon, G. C., Sen, B., Blavier, J.-F., and Jucks, K. W.: Kinetics of $HO_2 + HO_2 \rightarrow H_2O_2 + O_2$: Implications for stratospheric H_2O_2, Geophys. Res. Lett., 29, doi:10.1029/2001GL014525, 2002.

Citri, O. and Epstein, I. R.: Mechanistic study of a coupled chemical oscillator: the bromate-chlorite-iodide reaction, J. Phys. Chem., 92, 1865–1871, 1988.

Clifton, C. L., Altstein, N., and Huie, R. E.: Rate constant for the reaction of NO_2 with sulfur(IV) over the pH range 5.3–13, Environ. Sci. Technol., 22, 586–589, 1988.

Clyne, M. A. A. and Cruse, H. W.: Atomic resonance fluorescence spectrometry for the rate constants of rapid bimolecular reactions. Part 2. Reactions Cl + BrCl, Cl + Br_2, Cl + ICl, Br + IBr, Br + ICl, J. Chem. Soc. Faraday Trans. 2, 68, 1377–1387, 1972.

Damschen, D. E. and Martin, L. R.: Aqueous aerosol oxidation of nitrous acid by O_2, O_3 and H_2O_2, Atmos. Environ., 17, 2005–2011, 1983.

Davidovits, P., Hu, J. H., Worsnop, D. R., Zahniser, M. S., and Kolb, C. E.: Entry of gas molecules into liquids, Faraday Discuss., 100, 65–81, 1995.

Davis, Jr., W. and de Bruin, H. J.: New activity coefficients of 0-100 per cent aqueous nitric acid, J. Inorg. Nucl. Chem., 26, 1069–1083, 1964.

De Bruyn, W. J., Shorter, J. A., Davidovits, P., Worsnop, D. R., Zahniser, M. S., and Kolb, C. E.: Uptake of gas-phase sulfur species methanesulfonic acid, dimethylsulfoxide, and dimethyl sulfone by aqueous surfaces, J. Geophys. Res., 99D, 16 927–16 932, 1994.

Deiber, G., George, C., Le Calvé, S., Schweitzer, F., and Mirabel, P.: Uptake study of $ClONO_2$ and $BrONO_2$ by halide containing droplets, Atmos. Chem. Phys., 4, 1291–1299, 2004.

Deister, U. and Warneck, P.: Photooxidation of SO_3^{2-} in aqueous solution, J. Phys. Chem., 94, 2191–2198, 1990.

DeMore, W. B., Sander, S. P., Golden, D. M., Hampson, R. F., Kurylo, M. J., Howard, C. J., Ravishankara, A. R., Kolb, C. E., and Molina, M. J.: Chemical kinetics and photochemical data for use in stratospheric modeling. Evaluation number 12, JPL Publication 97-4, Jet Propulsion Laboratory, Pasadena, CA, 1997.

Dickson, A. G. and Millero, F. J.: A comparison of the equilibrium constants for the dissociation of carbonic acid in seawater media, Deep-Sea Res. A, 34, 1733–1743, 1987.

Dillon, T. J., Karunanandan, R., and Crowley, J. N.: The reaction of IO with CH_3SCH_3: Products and temperature dependent rate coefficients by laser induced fluorescence, Phys. Chem. Chem. Phys., 8, 847–855, 2006a.

Dillon, T. J., Tucceri, M. E., and Crowley, J. N.: Laser induced fluorescence studies of iodine oxide chemistry. Part II. The reactions of IO with CH_3O_2, CF_3O_2 and O_3, Phys. Chem. Chem. Phys., 8, 5185–5198, 2006b.

Dillon, T. J., Tucceri, M. E., Sander, R., and Crowley, J. N.: LIF studies of iodine oxide chemistry, part 3. Reactions IO + NO_3 $\rightarrow$ OIO + NO_2, I + NO_3 $\rightarrow$ IO + NO_2, and CH_2I + O_2 $\rightarrow$ (products): Implications for the chemistry of the marine atmosphere at night., Phys. Chem. Chem. Phys., 10, 1540–1554, 2008.

Dolson, D. A. and Leone, S. R.: A reinvestigation of the laser-initiated chlorine/hydrogen bromide chain reaction: absolute rate constants and the v = 2/v = 1 ratio from chlorine atom + hydrogen bromide $\rightarrow$ hydrogen chloride(v) + bromine atom, J. Phys. Chem., 91, 3543–3550, 1987.

Donohoue, D. L., Bauer, D., Cossairt, B., and Hynes, A. J.: Temperature and pressure dependent rate coefficients for the reaction of Hg with Br and the reaction of Br with Br: a pulsed laser photolysis-pulsed laser induced fluorescence study, J. Phys. Chem. A, 110, 6623–6632, 2006.

Edblom, E. C., Györgyi, L., Orbán, M., and Epstein, I. R.: A mechanism for dynamical behavior in the Landolt reaction with ferrocyanide, J. Am. Chem. Soc., 109, 4876–4880, 1987.

Eigen, M. and Kustin, K.: The kinetics of halogen hydrolysis, J. Am. Chem. Soc., 84, 1355–1361, 1962.

Enami, S., Hoshino, Y., and Kawasaki, M.: A kinetic study of the gas-phase reactions of OIO with NO, NO_2, and Cl_2, Int. J. Chem. Kinetics, 39, 688–693, 2007.

Exner, M., Herrmann, H., and Zellner, R.: Laser-based studies of reactions of the nitrate radical in aqueous solution, Ber. Bunsenges. Phys. Chem., 96, 470–477, 1992.

Faria, R. B., Lengyel, I., Epstein, I. R., and Kustin, K.: Combined mechanism explaining nonlinear dynamics in bromine(III) and bromine(V) oxidations of iodide ion, J. Phys. Chem., 97, 1164–1171, 1993.

Fogelman, K. D., Walker, D. M., and Margerum, D. W.: Non-metal redox kinetics: Hypochlorite and hypochlorous acid reactions with sulfite, Inorg. Chem., 28, 986–993, 1989.

Fortnum, D. H., Battaglia, C. J., Cohen, S. R., and Edwards, J. O.: The kinetics of the oxidation of halide ions by monosubstituted peroxides, J. Am. Chem. Soc., 82, 778–782, 1960.

Furrow, S.: Reactions of iodine intermediates in iodate-hydrogen peroxide oscillators, J. Phys. Chem., 91, 2129–2135, 1987.

Haag, W. R. and Hoigné, J.: Ozonation of bromide-containing waters: Kinetics of formation of hypobromous acid and bromate, Environ. Sci. Technol., 17, 261–267, 1983.

Hall, B.: The gas phase oxidation of elemental mercury by ozone, Water Air Soil Pollut., 80, 301–315, 1995.

Hanson, D. R., Burkholder, J. B., Howard, C. J., and Ravishankara, A. R.: Measurement of OH and HO_2 radical uptake coefficients on water and sulfuric acid surfaces, J. Phys. Chem., 96, 4979–4985, 1992.

Herrmann, H., Reese, A., and Zellner, R.: Time resolved UV/VIS diode array absorption spectroscopy of SO_x^- (x=3, 4, 5) radical anions in aqueous solution, J. Mol. Struct., 348, 183–186, 1995.

Herrmann, H., Ervens, B., Nowacki, P., Wolke, R., and Zellner, R.: A chemical aqueous phase radical mechanism for tropospheric chemistry, Chemosphere, 38, 1223–1232, 1999.

Hoffmann, M. R.: On the kinetics and mechanism of oxidation of aquated sulfur dioxide by ozone, Atmos. Environ., 20, 1145–1154, 1986.

Hoigné, J., Bader, H., Haag, W. R., and Staehelin, J.: Rate constants of reactions of ozone with organic and inorganic compounds in water – III Inorganic compounds and radicals, Wat. Res., 19, 993–1004, 1985.

Holmes, N. S., Adams, J. W., and Crowley, J. N.: Uptake and reaction of HOI and $IONO_2$ on frozen and dry NaCl/NaBr surfaces and H_2SO_4, Phys. Chem. Chem. Phys., 3, 1679–1687, 2001.

Hu, J. H., Shi, Q., Davidovits, P., Worsnop, D. R., Zahniser, M. S., and Kolb, C. E.: Reactive uptake of Cl_2(g) and Br_2(g) by aqueous surfaces as a function of Br^- and I^- ion concentration: The effect of chemical reaction at the interface, J. Phys. Chem., 99, 8768–8776, 1995.

Huie, R. E. and Neta, P.: Rate constants for some oxidations of S(IV) by radicals in aqueous solutions, Atmos. Environ., 21, 1743–1747, 1987.

Huthwelker, T., Clegg, S. L., Peter, T., Carslaw, K., Luo, B. P., and Brimblecombe, P.: Solubility of HOCl in water and aqueous H_2SO_4 to stratospheric temperatures, J. Atmos. Chem., 21, 81–95, 1995.

Hynes, A. J. and Wine, P. H.: The atmospheric chemistry of dimethylsulfoxide (DMSO) kinetics and mechanism of the OH + DMSO reaction, J. Atmos. Chem., 24, 23–37, 1996.

Ingham, T., Bauer, D., Sander, R., Crutzen, P. J., and Crowley, J. N.: Kinetics and products of the reactions BrO + DMS and Br + DMS at 298 K, J. Phys. Chem. A, 103, 7199–7209, 1999.

Jacob, D. J.: Chemistry of OH in remote clouds and its role in the production of formic acid and peroxymonosulfate, J. Geophys. Res., 91D, 9807–9826, 1986.

Jacobi, H.-W.: Kinetische Untersuchungen und Modellrechnungen zur troposphärischen Chemie von Radikalanionen und Ozon in wäßriger Phase, Ph.D. thesis, Universität GH Essen, Germany, 1996.

Jacobi, H.-W., Herrmann, H., and Zellner, R.: Kinetic investigation of the Cl_2^- radical in the aqueous phase, in: Air Pollution Research Report 57: Homogenous and hetrogenous chemical Processes in the Troposphere, edited by Mirabel, P., pp. 172–176, Office for official Publications of the European Communities, Luxembourg, 1996.

Jayne, J. T., Duan, S. X., Davidovits, P., Worsnop, D. R., Zahniser, M. S., and Kolb, C. E.: Uptake of gas-phase alcohol and organic acid molecules by water surfaces, J. Phys. Chem., 95, 6329–6336, 1991.

Jefferson, A., Nicovich, J. M., and Wine, P. H.: Temperature-dependent kinetics studies of the reactions $Br(^2P_{3/2})$ + CH_3SCH_3 ↔ CH_3SCH_2 + HBr. Heat of formation of the CH_3SCH_2 radical, J. Phys. Chem., 98, 7128–7135, 1994.

Jiang, P.-Y., Katsumura, Y., Nagaishi, R., Domae, M., Ishikawa, K., Ishigure, K., and Yoshida, Y.: Pulse radiolysis study of concentrated sulfuric acid solutions. Formation mechanism, yield and reactivity of sulfate radicals, J. Chem. Soc. Faraday Trans., 88, 1653–1658, 1992.

Keller-Rudek, H., Koschel, D., Merlet, P., Ohms-Bredemann, U., Wagner, J., and Wietelmann, A.: Gmelin Handbook of Inorganic and Organometallic Chemistry, 8th Edition, Br, Bromine, Supplement Volume B2, Compounds with Oxygen and Nitrogen, Springer Verlag, Berlin, 1992.

Kelley, C. M. and Tartar, H. V.: On the system: bromine-water, J. Am. Chem. Soc., 78, 5752–5756, 1956.

Kircher, C. C. and Sander, S. P.: Kinetics and mechanism of HO_2 and DO_2 disproportionations, J. Phys. Chem., 88, 2082–2091, 1984.

Kohlmann, J.-P. and Poppe, D.: The tropospheric gas-phase degradation of NH_3 and its impact on the formation of N_2O and NO_x, J. Atmos. Chem., 32, 397–415, 1999.

Kondo, O. and Benson, S. W.: Kinetics and equilibria in the system Br + CH_3OOH ⇌ HBr + CH_3OO. An upper limit for the heat of formation of the methylperoxy radical, J. Phys. Chem., 88, 6675–6680, 1984.

Kumar, K. and Margerum, D. W.: Kinetics and mechanism of general-acid-assisted oxidation of bromide by hypochlorite and hypochlorous acid, Inorg. Chem., 26, 2706–2711, 1987.

Lax, E.: Taschenbuch für Chemiker und Physiker, Springer Verlag, Berlin, 1969.

Lee, Y.-N. and Schwartz, S. E.: Reaction kinetics of nitrogen dioxide with liquid water at low partial pressure, J. Phys. Chem., 85, 840–848, 1981.

Lengyel, I., Li, J., Kustin, K., and Epstein, I. R.: Rate constants for reactions between iodine- and chlorine-containing species: A detailed mechanism of the chlorine dioxine/chlorite reaction, J. Am. Chem. Soc., 118, 3708–3719, 1996.

Lin, C.-J. and Pehkonen, S. O.: Aqueous free radical chemistry of mercury in the presence of iron oxides

and ambient aerosol, Atmos. Environ., 31, 4125–4137, 1997.

Lin, C.-J. and Pehkonen, S. O.: Oxidation of elemental mercury by aqueous chlorine ($HOCl/OCl^-$): Implications for tropospheric mercury chemistry, J. Geophys. Res., 103D, 28 093–28 102, 1998.

Lind, J. A. and Kok, G. L.: Correction to "Henry's law determinations for aqueous solutions of hydrogen peroxide, methylhydroperoxide, and peroxyacetic acid" by John A. Lind and Gregory L. Kok, J. Geophys. Res., 99D, 21 119, 1994.

Lind, J. A., Lazrus, A. L., and Kok, G. L.: Aqueous phase oxidation of sulfur(IV) by hydrogen peroxide, methylhydroperoxide, and peroxyacetic acid, J. Geophys. Res., 92D, 4171–4177, 1987.

Liu, Q. and Margerum, D. W.: Equilibrium and kinetics of bromine chloride hydrolysis, Environ. Sci. Technol., 35, 1127–1133, 2001.

Liu, Y., Pimentel, A. S., Antoku, Y., Giles, B. J., and Barker, J. R.: Temperature-dependent rate and equilibrium constants for $Br \cdot (aq) + Br^-(aq) \rightleftharpoons Br_2^- \cdot (aq)$, J. Phys. Chem. A, 106, 11 075–11 082, 2002.

Logager, T., Sehested, K., and Holcman, J.: Rate constants of the equilibrium reactions $SO_4 + HNO_3 \rightleftharpoons HSO_4^- + NO_3$ and $SO_4 + NO_3 \rightleftharpoons SO_4^{2-} + NO_3$, Radiat. Phys. Chem., 41, 539–543, 1993.

Long, C. A. and Bielski, B. H. J.: Rate of reaction of superoxide radical with chloride-containing species, J. Phys. Chem., 84, 555–557, 1980.

Magi, L., Schweitzer, F., Pallares, C., Cherif, S., , Mirabel, P., and George, C.: Investigation of the uptake rate of ozone and methyl hydroperoxide by water surfaces, J. Phys. Chem. A, 101, 4943–4949, 1997.

Manion, J. A., Huie, R. E., Levin, R. D., Burgess, Jr., D. R., Orkin, V. L., Tsang, W., McGivern, W. S., Hudgens, J. W., Knyazev, V. D., Atkinson, D. B., Chai, E., Tereza, A. M., Lin, C.-Y., Allison, T. C., Mallard, W. G., Westley, F., Herron, J. T., Hampson, R. F., and Frizzell, D. H.: NIST Chemical Kinetics Database (Web Version), `http://kinetics.nist.gov/`, 2010.

Margerum, D. W., Dickson, P. N., Nagy, J. C., Kumar, K., Bowers, C. P., and Fogelman, K. D.: Kinetics of the iodine monochloride reaction with iodide measured by the pulsed-accelerated-flow method, Inorg. Chem., 25, 4900–4904, 1986.

Marsh, A. R. W. and McElroy, W. J.: The dissociation constant and Henry's law constant of HCl in aqueous solution, Atmos. Environ., 19, 1075–1080, 1985.

Martin, L. R. and Damschen, D. E.: Aqueous oxidation of sulfur dioxide by hydrogen peroxide at low pH, Atmos. Environ., 15, 1615–1621, 1981.

Matthew, B. M., George, I., and Anastasio, C.: Hydroperoxyl radical (HO2·) oxidizes dibromide radical anion ($\cdot Br_2^-$) to bromine (Br_2) in aqueous solution: Implications for the formation of Br_2 in the marine boundary layer, Geophys. Res. Lett., 30, doi: 10.1029/2003GL018572, 2003.

McCabe, D. C., Gierczak, T., Talukdar, R. K., and Ravishankara, A. R.: Kinetics of the reaction OH + CO under atmospheric conditions, Geophys. Res. Lett., 28, 3135–3138, 2001.

Mössinger, J. C. and Cox, R. A.: Heterogeneous reaction of HOI with sodium halide salts, J. Phys. Chem. A, 105, 5165–5177, 2001.

Munthe, J.: The aqueous oxidation of elemental mercury by ozone, Atmos. Environ., 26A, 1461–1468, 1992.

Nagy, J. C., Kumar, K., and Margerum, D. W.: Non-metal redox kinetics: Oxidation of iodide by hypochlorous acid and by nitrogen trichloride measured by the pulsed-accelerated-flow method, Inorg. Chem., 27, 2773–2780, 1988.

Nakano, Y., Ishiwata, T., and Kawasaki, M.: Rate constants of the reaction of NO_3 with CH_3I measured with use of cavity ring-down spectroscopy, J. Phys. Chem. A, 109, 6527–6531, 2005.

Neta, P. and Huie, R. E.: Rate constants for reactions of NO_3 radicals in aqueous solutions, J. Phys. Chem., 90, 4644–4648, 1986.

Orlando, J. J. and Tyndall, G. S.: Rate coefficients for the thermal decomposition of $BrONO_2$ and the heat of formation of $BrONO_2$, J. Phys. Chem., 100, 19 398–19 405, 1996.

Orlando, J. J., Tyndall, G. S., Bilde, M., Ferronato, C., Wallington, T. J., Vereecken, L., and Peeters, J.: Laboratory and theoretical study of the oxy radicals in the OH- and Cl-initiated oxidation of ethene, J. Phys. Chem. A, 102, 8116–8123, 1998.

Orlando, J. J., Tyndall, G. S., Fracheboud, J. M., Estupinan, E. G., Haberkorn, S., and Zimmer, A.: The rate and mechanism of the gas-phase oxidation of hydroxyacetone, Atmos. Environ., 33, 1621–1629, 1999.

Orlando, J. J., Tyndall, G. S., Bertman, S. B., Chen, W., and Burkholder, J. B.: Rate coefficient for the reaction of OH with $CH_2{=}C(CH_3)C(O)OONO_2$ (MPAN), Atmos. Environ., 36, 1895–1900, 2002.

Pal, B. and Ariya, P. A.: Gas-phase HO-initiated reactions of elemental mercury: Kinetics, product studies, and atmospheric implications, Environ. Sci. Technol., 38, 5555–5566, 2004.

Palmer, D. A., Ramette, R. W., and Mesmer, R. E.: The hydrolysis of iodine: Equilibria at high temperatures, J. Nucl. Mater., 130, 280–286, 1985.

Plane, J. M. C., Joseph, D. M., Allan, B. J., Ashworth, S. H., and Francisco, J. S.: An experimental and theoretical study of the reactions OIO + NO and OIO + OH, J. Phys. Chem. A, 110, 93–100, 2006.

Pleijel, K. and Munthe, J.: Modelling the atmospheric mercury cycle — Chemistry in fog droplets, Atmos. Environ., 29, 1441–1457, 1995.

Ponche, J. L., George, C., and Mirabel, P.: Mass transfer at the air/water interface: Mass accommodation coefficients of SO_2, HNO_3, NO_2 and NH_3, J. Atmos. Chem., 16, 1–21, 1993.

Pöschl, U., Canagaratna, M., Jayne, J. T., Molina, L. T., Worsnop, D. R., Kolb, C. E., and Molina, M. J.: Mass accommodation coefficient of H_2SO_4 vapor on aqueous sulfuric acid surfaces and gaseous diffusion coefficient of H_2SO_4 in N_2/H_2O, J. Phys. Chem. A, 102, 10 082–10 089, 1998.

Raofie, F. and Ariya, P. A.: Kinetics and products study of the reaction of BrO radicals with gaseous mercury, J. Phys. IV France, 107, 1119–1121, 2003.

Raofie, F. and Ariya, P. A.: Product study of the gas-phase BrO-initiated oxidation of Hg^0: Evidence for stable Hg^{1+} compounds, Environ. Sci. Technol., 38, 4319–4326, 2004.

Régimbal, J.-M. and Mozurkewich, M.: Peroxynitric acid decay mechanisms and kinetics at low pH, J. Phys. Chem. A, 101, 8822–8829, 1997.

Rickard, A. and Pascoe, S.: The Master Chemical Mechanism (MCM), `http://mcm.leeds.ac.uk`, 2009.

Ross, A. B., Mallard, W. G., Helman, W. P., Bielski, B. H. J., Buxton, G. V., Cabelli, D. E., Greenstock, C. L., Huie, R. E., and Neta, P.: NDRL-NIST Solution Kinetics Database: - Ver. 1, National Institute of Standards and Technology, Gaithersburg, MD, 1992.

Rudich, Y., Talukdar, R. K., Imamura, T., Fox, R. W., and Ravishankara, A. R.: Uptake of NO_3 on KI solutions: Rate coefficient for the $NO_3 + I^-$ reaction and gas-phase diffusion coefficients for NO_3, Chem. Phys. Lett., 261, 467–473, 1996.

Saastad, O. W., Ellermann, T., and Nielsen, C. J.: On the adsorption of NO and NO_2 on cold H_2O/H_2SO_4 surfaces, Geophys. Res. Lett., 20, 1191–1193, 1993.

Sander, S. P., Friedl, R. R., DeMore, W. B., Golden, D. M., Kurylo, M. J., Hampson, R. F., Huie, R. E., Moortgat, G. K., Ravishankara, A. R., Kolb, C. E., and Molina, M. J.: Chemical kinetics and photochemical data for use in stratospheric modeling. Supplement to evaluation 12: Update of key reactions. Evaluation number 13, JPL Publication 00-3, Jet Propulsion Laboratory, Pasadena, CA, `http://jpldataeval.jpl.nasa.gov/`, 2000.

Sander, S. P., Finlayson-Pitts, B. J., Friedl, R. R., Golden, D. M., Huie, R. E., Kolb, C. E., Kurylo, M. J., Molina, M. J., Moortgat, G. K., Orkin, V. L., and Ravishankara, A. R.: Chemical Kinetics and Photochemical Data for Use in Atmospheric Studies, Evaluation Number 14, JPL Publication 02-25, Jet Propulsion Laboratory, Pasadena, CA, 2003.

Sander, S. P., Friedl, R. R., Golden, D. M., Kurylo, M. J., Moortgat, G. K., Keller-Rudek, H., Wine, P. H., Ravishankara, A. R., Kolb, C. E., Molina, M. J., Finlayson-Pitts, B. J., Huie, R. E., and Orkin, V. L.: Chemical Kinetics and Photochemical Data for Use in Atmospheric Studies, Evaluation Number 15, JPL Publication 06-2, Jet Propulsion Laboratory, Pasadena, CA, 2006.

Schroeder, W. H. and Munthe, J.: Atmospheric mercury — An overview, Atmos. Environ., 32, 809–822, 1998.

Schwartz, S. E. and White, W. H.: Solubility equilibria of the nitrogen oxides and oxyacids in dilute aqueous solution, in: Advances in Environmental Science and Engineering, edited by Pfafflin, J. R. and Ziegler, E. N., vol. 4, pp. 1–45, Gordon and Breach Science Publishers, NY, 1981.

Schwarz, H. A. and Bielski, B. H. J.: Reactions of HO_2 and O_2^- with iodine and bromine and the I_2^- and I atom reduction potentials, J. Phys. Chem., 90, 1445–1448, 1986.

Schweitzer, F., Mirabel, P., and George, C.: Uptake of hydrogen halides by water droplets, J. Phys. Chem. A, 104, 72–76, 2000.

Sehested, K., Rasmussen, O. L., and Fricke, H.: Rate constants of OH with HO_2, O_2^-, and $H_2O_2^+$ from hydrogen peroxide formation in pulse-irradiated oxygenated water, J. Phys. Chem., 72, 626–631, 1968.

Sehested, K., Holcman, J., and Hart, E. J.: Rate constants and products of the reactions of e_{aq}^-, O_2^- and H with ozone in aqueous solutions, J. Phys. Chem., 87, 1951–1954, 1983.

Seinfeld, J. H. and Pandis, S. N.: Atmospheric Chemistry and Physics, John Wiley & Sons, Inc., 1998.

Shon, Z.-H., Kim, K.-H., Kim, M.-Y., and Lee, M.: Modeling study of reactive gaseous mercury in the urban air, Atmos. Environ., 39, 749–761, 2005.

Shoute, L. C. T., Alfassi, Z. B., Neta, P., and Huie, R. E.: Temperature dependence of the rate constants for reaction of dihalide and azide radicals with inorganic reductants, J. Phys. Chem., 95, 3238–3242, 1991.

Sivakumaran, V., Hölscher, D., Dillon, T. J., and Crowley, J. N.: Reaction between OH and HCHO: temperature dependent rate coefficients (202-399 K) and product pathways (298 K), Phys. Chem. Chem. Phys., 5, 4821–4827, 2003.

Staudinger, J. and Roberts, P. V.: A critical compilation of Henry's law constant temperature dependence relations for organic compounds in dilute aqueous solutions, Chemosphere, 44, 561–576, 2001.

Stimpfle, R. M., Wilmouth, D. M., Salawitch, R. J., and Anderson, J. G.: First measurements of ClOOCl in the stratosphere: The coupling of ClOOCl and ClO in the Arctic polar vortex, J. Geophys. Res., 109, doi:10.1029/2003JD003811, 2004.

Sutton, H. C. and Downes, M. T.: Reactions of the HO_2 radical in aqueous solution with bromine and related compounds, J. Chem. Soc. Faraday Trans. 1, 68, 1498–1507, 1972.

Takami, A., Kato, S., Shimono, A., and Koda, S.: Uptake coefficient of OH radical on aqueous surface, Chem. Phys., 231, 215–227, 1998.

Taraborrelli, D., Lawrence, M. G., Butler, T. M., Sander, R., and Lelieveld, J.: Mainz Isoprene Mechanism 2 (MIM2): an isoprene oxidation mechanism for regional and global atmospheric modelling, Atmos. Chem. Phys., 9, 2751–2777, http://www.atmos-chem-phys.net/9/2751, 2009.

Thomas, K., Volz-Thomas, A., and Kley, D.: Zur Wechselwirkung von NO_3-Radikalen mit wässrigen Lösungen: Bestimmung des Henry- und des Massenakkomodationskoeffizienten, Ph.D. thesis, Institut für Chemie und Dynamik der Geosphäre 2, Forschungszentrum Jülich GmbH, Germany, 1993.

Thornton, J. and Abbatt, J. P. D.: Measurements of HO_2 uptake to aqueous aerosol: Mass accommodation coefficients and net reactive loss, J. Geophys. Res., 110D, doi:10.1029/2004JD005402, 2005.

Tokos, J. J. S., Hall, B., Calhoun, J. A., and Prestbo, E. M.: Homogeneous gas-phase reaction of Hg^0 with H_2O_2, O_3, CH_3I, and $(CH_3)_2S$: Implications for atmospheric Hg cycling, Atmos. Environ., 32, 823–827, 1998.

Troy, R. C. and Margerum, D. W.: Non-metal redox kinetics: Hypobromite and hypobromous acid reactions with iodide and with sulfite and the hydrolysis of bromosulfate, Inorg. Chem., 30, 3538–3543, 1991.

Troy, R. C., Kelley, M. D., Nagy, J. C., and Margerum, D. W.: Non-metal redox kinetics: Iodine monobromide reaction with iodide ion and the hydrolysis of IBr, Inorg. Chem., 30, 4838–4845, 1991.

Tyndall, G. S., Staffelbach, T. A., Orlando, J. J., and Calvert, J. G.: Rate coefficients for the reactions of OH radicals with methylglyoxal and acetaldehyde, Int. J. Chem. Kinetics, 27, 1009–1020, 1995.

Tyndall, G. S., Cox, R. A., Granier, C., Lesclaux, R., Moortgat, G. K., Pilling, M. J., Ravishankara, A. R., and Wallington, T. J.: The atmospheric chemistry of small organic peroxy radicals, J. Geophys. Res., 106D, 12 157–12 182, 2001.

van den Bergh, H. and Troe, J.: Kinetic and thermodynamic properties of INO and INO_2 intermediate complexes in iodine recombination, J. Chem. Phys., 64, 736–742, 1976.

van Loon, L., Mader, E., and Scott, S. L.: Reduction of the aqueous mercuric ion by sulfite: UV spectrum of $HgSO_3$ and its intramolecular redox reaction, J. Phys. Chem. A, 104, 1621–1626, 2000.

van Loon, L. L., Mader, E. A., and Scott, S. L.: Sulfite stabilization and reduction of the aqueous mercuric ion: Kinetic determination of sequential formation constants, J. Phys. Chem. A, 105, 3190–3195, 2001.

Vogt, R., Crutzen, P. J., and Sander, R.: A mechanism for halogen release from sea-salt aerosol in the remote marine boundary layer, Nature, 383, 327–330, doi:10.1038/383327A0, 1996.

von Glasow, R., Sander, R., Bott, A., and Crutzen, P. J.: Modeling halogen chemistry in the marine boundary layer, 1. Cloud-free MBL, J. Geophys. Res., 107D, 4341, doi:10.1029/2001JD000942, 2002.

von Kuhlmann, R.: Tropospheric photochemistry of ozone, its precursors and the hydroxyl radical: A 3D-modeling study considering non-methane hydrocarbons, Ph.D. thesis, Johannes Gutenberg-Universität, Mainz, Germany, 2001.

von Kuhlmann, R., Lawrence, M. G., Crutzen, P. J., and Rasch, P. J.: A model for studies of tropospheric ozone and nonmethane hydrocarbons: Model description and ozone results, J. Geophys. Res., 108D, doi:10.1029/2002JD002893, 2003.

Wagman, D. D., Evans, W. H., Parker, V. B., Schumm, R. H., Halow, I., Bailey, S. M., Churney, K. L., and Nuttall, R. L.: The NBS tables of chemical thermodynamic properties; Selected values for inorganic and

C_1 and C_2 organic substances in SI units, J. Phys. Chem. Ref. Data, 11, suppl. 2, 1982.

Wagner, I. and Strehlow, H.: On the flash photolysis of bromide ions in aqueous solution, Ber. Bunsenges. Phys. Chem., 91, 1317–1321, 1987.

Wang, T. X. and Margerum, D. W.: Kinetics of reversible chlorine hydrolysis: Temperature dependence and general-acid/base-assisted mechanisms, Inorg. Chem., 33, 1050–1055, 1994.

Wang, T. X., Kelley, M. D., Cooper, J. N., Beckwith, R. C., and Margerum, D. W.: Equilibrium, kinetic, and UV-spectral characteristics of aqueous bromine chloride, bromine, and chlorine species, Inorg. Chem., 33, 5872–5878, 1994.

Wang, Y. L., Nagy, J. C., and Margerum, D. W.: Kinetics of hydrolysis of iodine monochloride measured by the pulsed-accelerated-flow method, J. Am. Chem. Soc., 111, 7838–7844, 1989.

Wang, Z. and Pehkonen, S. O.: Oxidation of elemental mercury by aqueous bromine: atmospheric implications, Atmos. Environ., 38, 3675–3688, 2004.

Warneck, P.: Chemical reactions in clouds, Fresenius J. Anal. Chem., 340, 585–590, 1991.

Warneck, P.: The relative importance of various pathways for the oxidation of sulfur dioxide and nitrogen dioxide in sunlit continental fair weather clouds, Phys. Chem. Chem. Phys., 1, 5471–5483, 1999.

Wayne, R. P., Barnes, I., Biggs, P., Burrows, J. P., Canosa-Mas, C. E., Hjorth, J., Le Bras, G., Moortgat, G. K., Perner, D., Poulet, G., Restelli, G., and Sidebottom, H.: The nitrate radical: Physics, chemistry, and the atmosphere, Atmos. Environ., 25A, 1–203, 1991.

Weast, R. C., ed.: CRC Handbook of Chemistry and Physics, 61st Edition, CRC Press, Inc., Boca Raton, FL, 1980.

Weinstein-Lloyd, J. and Schwartz, S. E.: Low-intensity radiolysis study of free-radical reactions in cloudwater: H_2O_2 production and destruction, Environ. Sci. Technol., 25, 791–800, 1991.

Wilhelm, E., Battino, R., and Wilcock, R. J.: Low-pressure solubility of gases in liquid water, Chem. Rev., 77, 219–262, 1977.

Wine, P. H., Tang, Y., Thorn, R. P., Wells, J. R., and Davis, D. D.: Kinetics of aqueous phase reactions of the SO_4^- radical with potential importance in cloud chemistry, J. Geophys. Res., 94D, 1085–1094, 1989.

Wingenter, O. W., Sive, B. C., Blake, N. J., and Rowland, F. S.: Atomic chlorine concentrations determined from ethane and hydroxyl measurements made over the Central Pacific Ocean, Eos, Trans. AGU (Abstract Supplement), 80, F149–F150, 1999.

Worsnop, D. R., Zahniser, M. S., Kolb, C. E., Gardner, J. A., Watson, L. R., van Doren, J. M., Jayne, J. T., and Davidovits, P.: The temperature dependence of mass accommodation of SO_2 and H_2O_2 on aqueous surfaces, J. Phys. Chem., 93, 1159–1172, 1989.

Wu, D., Wong, D., and Di Bartolo, B.: Evolution of Cl_2^- in aqueous NaCl solutions, J. Photochem., 14, 303–310, 1980.

Yiin, B. S. and Margerum, D. W.: Nonmetal redox kinetics: reactions of iodine and triiodide with sulfite and hydrogen sulfite and the hydrolysis of iodosulfate, Inorg. Chem., 29, 1559–1564, 1990.

Yu, X.-Y.: Critical evaluation of rate constants and equilibrium constants of hydrogen peroxide photolysis in acidic aqueous solutions containing chloride ions, J. Phys. Chem. Ref. Data, 33, 747–763, 2004.

Zaveri, R. A. and Peters, L. K.: A new lumped structure photochemical mechanism for large-scale applications, J. Geophys. Res., 104D, 30 387–30 415, 1999.

Zehavi, D. and Rabani, J.: The oxidation of aqueous bromide by hydroxyl radicals. A pulse radiolytic investigation, J. Phys. Chem., 76, 312–319, 1972.