
Geosci. Model Dev. Discuss., 4, C1179–C1180, 2011 www.geosci-model-dev-discuss.net/4/C1179/2011/ © Author(s) 2011. This work is distributed under the Creative Commons Attribute 3.0 License.



**GMDD** 

4, C1179-C1180, 2011

Interactive Comment

## Interactive comment on "Supersaturation calculation in large eddy simulation models for prediction of the droplet number concentration" by O. Thouron et al.

S. Ghan

steve.ghan@pnl.gov

Received and published: 13 December 2011

I enjoyed reading the paper. A few comments:

- 1. Which aerosol activation scheme is used? There are differences, and those differences could explain some of the differences between the results of Scheme A, B and C.
- 2. Why do you treat in-cloud activation with scheme A? It was never designed for that, and is certainly not done in my applications to large-scale models. Why not just treat activation at cloud base? See Ovchinnikov and Ghan, JGR 2005.

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Discussion Paper



3. It is possible to apply a different diagnostic treatment of supersaturation for secondary nucleation within the interior of clouds. See Ghan et al., J. Advances Modeling Earth Systems (2011).

Steve Ghan

Pacific Northwest National Laboratory

Interactive comment on Geosci. Model Dev. Discuss., 4, 3313, 2011.

## **GMDD**

4, C1179-C1180, 2011

Interactive Comment

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Discussion Paper

