



## ***Interactive comment on “Modeling atmospheric ammonia and ammonium using a backward-in-time stochastic Lagrangian air quality model (STILT-Chem v0.7)” by D. Wen et al.***

**D. Wen et al.**

dwen@uwaterloo.ca

Received and published: 23 January 2013

Response: We thank the reviewer for the comment. We have provided detailed, point-by-point responses to the comments and suggestion below.

The overall impression is that that the authors have done a very nice job, but also that the article lacks regarding two important parts: Description of the ammonia emissions is insufficient, and the other part is that this work needs to be compared and discussed with results obtained in work with similar models and methods.

Response: 1) the following description of  $\text{NH}_3$  emission has been added to line 9 on

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)

[Discussion Paper](#)



One of the key objectives of NAESI was to improve the 2002 national inventory on NH<sub>3</sub> emissions, especially from agricultural sources, using updated, Canadian specific, agricultural activity data and emission factors. The updated inventory can therefore account for spatial variation due to regional differences in farming practices and climatic conditions for each livestock category, and temporal variation due to seasonally different agricultural practices or seasonally variable temperatures that have different effects on agricultural NH<sub>3</sub> emissions throughout the year (Ayres et al., 2009). Total Canadian NH<sub>3</sub> emission in 2006 is about 5 Mt, about 90% of which are from agricultural. More information about emissions from other source types can be found at [http://www.ec.gc.ca/pdb/websol/emissions/2006/2006\\_canada\\_e.cfm](http://www.ec.gc.ca/pdb/websol/emissions/2006/2006_canada_e.cfm).

and the following description was added to Line 21 on Page 2761:

Temporal allocations of emissions were performed by SMOKE using predefined temporal profiles, allowing SMOKE-processed emissions to represent diurnal, weekly, and monthly variations.

The following corresponding reference was added to the paper:

Ayres, J., Bittman, S., Sheppard, S., and Girdhar, S.: Sources of ammonia emissions. In: Lillyman, C., and K. Buset.: Canadian Atmospheric Assessment of Agricultural Ammonia, pp. 112–130. National Agri-Environmental Standards Technical Series Report No. 4-1, 2008.

2) The following text about comparison against a similar model's results was added in Line 23 on Page 2763:

In a previous study (Skjøth et al., 2004) using a similar model (ACDEP), correlation coefficients obtained for the years 1999-2001 for three sites in Denmark varied from 0.43 to 0.69 when measured and simulated diurnal mean NH<sub>3</sub> concentration were used. Correlation coefficients increased to a range of 0.83 to 0.93 when measured

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Discussion Paper



and simulated monthly NH<sub>3</sub> concentrations were used.

The following corresponding reference was added to the paper:

Skjøth, C. A., Hertel, O., Gyldenkerne, S., and Ellermann, T.: Implementing a dynamical ammonia emission parameterization in the large-scale air pollution model ACDEP, *J. Geophys. Res.*, D06306, 109, 1-13, 2004., doi:10.1029/2003JD003895

The authors have completely failed to recognize the rather similar work performed of a Danish research group more than 15 years ago. This Danish group used a similar type of model, the ACDEP model, and they even applied a near to identical chemical scheme (Hertel et al, 1995) including a parameterization of the formation of ammonium nitrate, other inorganic as well as organic nitrates. This model has been extensively used for studies to simulate ammonia concentrations (de Leeuw et al., 2003; Skjøth et al., 2002; Skjøth et al., 2004) and nitrogen depositions (Hertel et al., 2002; Hertel et al., 2003) and uses a recently open source model that dynamically handles ammonia emissions (Skjøth et al., 2004; Gyldenkerne et al., 2005; Skjøth et al., 2011). The emission model is already successfully implemented in the DEHM (Brandt et al., 2012) and the Unified EMEP model (Berge, 2010) and underway in the EMEP4UK model (Reis et al, 2011). Furthermore, several reviews have highlighted the applied approach as a needed requirement for many model calculations with respect to ammonia and ammonium containing particles (Hertel et al., 2006; Hertel et al., 2012; Menut and Bessagnet, 2010). Other groups in Europe have also used similar but more simple methods as those that are based on the FRAME model (Fournier et al., 2004; Kryza et al., 2011; Zhang et al., 2011) that were originally developed for the English area (Fournier et al., 2004). Or the TREND model (Asman, 2011), applying a simple seasonal variation of the ammonia emission.

## REFERENCES

Asman, W. A. H. (2001), Modelling the atmospheric transport and deposition of ammo-

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)

[Discussion Paper](#)



nia and ammonium: an overview with special reference to Denmark, *Atmos. Environ.*, 35(11), 1969–1983.

Brandt, J., Silver, J. D., Frohn, L. M., Geels, C., Gross, A., Hansen, A. B., Hansen, K. M., Hedegaard, G. B., Skjøth, C. A., Villadsen, H., Zare, A., and Christensen, J. H., An integrated model study for Europe and North America using the Danish Eulerian Hemispheric Model with focus on intercontinental transport of air pollution, *Atmos. Environ.*, 53, 156-176, 2012.

Berge, H., New temporal variation of ammonia emissions, in Benedictow et al., "Transboundary acidification, eutrophication and ground level ozone in Europe in 2008", Joint MSC-W & CCC & CEIP Report, Norwegian Meteorological Institute, 2010.

de Leeuw, G., Skjøth, C. A., Hertel, O., Jickells, T., Spokes, L., Vignati, E., Frohn, L., Frydendall, J., Schulz, M., Tamm, S., Sørensen, L. L., and Kunz, G. J., Deposition of nitrogen into the North Sea, *Atmos. Environ.*, 37, Supplement 1, 145-165, 2003.

Fournier, N., Dore, A. J., Vieno, M., Weston, K. J., Dragosits, U., and Sutton, M. A., Modelling the deposition of atmospheric oxidised nitrogen and sulphur to the United Kingdom using a multi-layer long-range transport model, *Atmos. Environ.*, 38, 683-694, 2004.

Gyldenkerne, S., Ambelas Skjøth, C., Hertel, O., and Ellermann, T., A dynamical ammonia emission parameterization for use in air pollution models, *J. Geophys. Res.*, [Atmos. ], D07108, 110, 1-14, 2005., doi:10.1029/2004JD005459

Hertel, O., Christensen, J., Runge, E. H., Asman, W. A. H., Berkowicz, R., Hovmand, M. F., and Hov, O., Development and Testing of A New Variable Scale Air-Pollution Model - ACDEP, *Atmos. Environ.*, 29, 1267-1290, 1995.

Please also note the supplement to this comment: <http://www.geosci-model-dev-discuss.net/5/C1132/2012/gmdd-5-C1132-2012-supplement.pdf>

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Discussion Paper

Response: the following was added to Line 23 on Page 2747 to address the issue:

Extensive efforts have been made to modeling studies of atmospheric NH<sub>3</sub> using different models either in Eulerian framework (Brandt et al., 2012; Berge, 2010; Reis et al., 2011; Wu et al., 2008; Makar et al., 2009; Sakurai et al., 2005) or in Lagrangian framework. Although the Eulerian approach is powerful and widely used for elucidating the chemical and physical mechanism in the atmosphere, the Lagrangian approach demonstrates key advantages in presenting sub-grid scale process, minimizing numerical diffusion, artificial dilution and computing resources. The Lagrangian approach has been widely adopted in various models in atmospheric ammonia modeling such as the FRAME model (Singles et al., 1998; Kryza et al., 2011; Zhang et al., 2011), the TREND model (Asman and van Jaarsveld 1992; Asman 2001), the ACDEP model (Hertel et al, 1995; de Leeuw et al., 2003; Skjøth et al., 2002; Skjøth et al., 2004; Hertel et al., 2002; Hertel et al., 2003; Skjøth et al., 2011; Gyldenkerne et al., 2005), The TERN model (ApSimon et al., 1994), and the NAME model (Redington and Derwent, 2002). Most existing Lagrangian models for atmospheric ammonia modeling are either box-based models or use simplified dry chemical scheme. In this study, we attempt to model atmospheric ammonia using a stochastic time-inverted Lagrangian particle model in which a comprehensive dry chemical scheme (CB4) and a back-trajectory method are used. Plumes in Lagrangian particle models are represented by a large number of fictitious particles, which move with random trajectories to represent atmospheric turbulence. Particle models are able to account in detail for three-dimensional variations in the wind field and the effects of turbulent dispersion. High resolution and the improved accuracy of the vertical dispersion parameterization make these particle models particularly useful for simulating highly variable emission rates in complex dispersion scenarios.

The following corresponding references were added to the paper:

ApSimon, H. M., Barker, B. M., Kayin, S.: Modelling studies of the atmospheric release and transport of ammonia in anticyclonic episodes. *Atmos. Environ.*, 28(4), 665-678,

C1354

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Discussion Paper



1994,.

Asman, W. A. H.: Modelling the atmospheric transport and deposition of ammonia and ammonium: an overview with special reference to Denmark, *Atmos. Environ.*, 35(11), 1969–1983, 2001,.

Asman, W.A.H., van Jaarsveld, J.A.: A variable-resolution transport model applied for  $\text{NH}_x$  in Europe. *Atmos. Environ.*, 26A, 445-464, 1992.

Berge, H.: New temporal variation of ammonia emissions, in Benedictow et al.: Transboundary acidification, eutrophication and ground level ozone in Europe in 2008, Joint MSC-W & CCC & CEIP Report, Norwegian Meteorological Institute, 2010.

Brandt, J., Silver, J. D., Frohn, L. M., Geels, C., Gross, A., Hansen, A. B., Hansen, K. M., Hedegaard, G. B., Skjøth, C. A., Villadsen, H., Zare, A., and Christensen, J. H.: An integrated model study for Europe and North America using the Danish Eulerian Hemispheric Model with focus on intercontinental transport of air pollution, *Atmos. Environ.*, 53, 156-176, 2012.

de Leeuw, G., Skjøth, C. A., Hertel, O., Jickells, T., Spokes, L., Vignati, E., Frohn, L., Frydendall, J., Schulz, M., Tamm, S., Sørensen, L. L., and Kunz, G. J.: Deposition of nitrogen into the North Sea, *Atmos. Environ.*, 37, Supplement 1, 145-165, 2003.

Gyldenkerne, S., Ambelas Skjøth, C., Hertel, O., and Ellermann, T.: A dynamical ammonia emission parameterization for use in air pollution models, *J. Geophys. Res.*, D07108, 110, 1-14, 2005, doi:10.1029/2004JD005459.

Hertel, O., Christensen, J., Runge, E. H., Asman, W. A. H., Berkowicz, R., Hovmand, M. F., and Hov, O.: Development and Testing of A New Variable Scale Air-Pollution Model - ACDEP, *Atmos. Environ.*, 29, 1267-1290, 1995.

Hertel, O., Skjøth, C. A., Brandt, J., Christensen, J. H., Frohn, L. M., and Frydendall, J.: Operational mapping of atmospheric nitrogen deposition to the Baltic Sea, *Atmos. Chem. Phys.*, 3, 2083-2099, 2003.

C1355

**GMDD**

5, C1350–C1357, 2013

Interactive  
Comment

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Discussion Paper



Hertel, O., Skjøth, C. A., Frohn, L. M., Vignati, E., Frydendall, J., de Leeuw, G., Schwarz, U., and Reis, S.: Assessment of the atmospheric nitrogen and sulphur inputs into the North Sea using a Lagrangian model, *Physics and Chemistry of the Earth*, 27, 1507-1515, 2002.

Kryza, M., Dore, A. J., Blas, M., and Sobik, M.: Modelling deposition and air concentration of reduced nitrogen in Poland and sensitivity to variability in annual meteorology, *J. Environ. Manage.*, 92, 1225-1236, 2011.

Redington, A. L., and Derwent, R. G.: Calculation of sulphate and nitrate aerosol concentrations over Europe using a Lagrangian dispersion model. *Atmos. Environ.*, 36(28), 4425-4439, 2002.

Reis, S., Skjøth, C. A., Vioenoa, M., Geels, C., Steinle, S., Lang, M., Sutton, M.: Why time and space matters - arguments for the improvement of temporal emission profiles for atmospheric dispersion modeling of air pollutant emissions, MODSIM2011 Congress, 12-16 Dec 2011, Perth, Australia <http://www.mssanz.org.au/modsim2011/E1/reis.pdf>.

Sakurai, T., Fujita, S. I., Hayami, H., Furuhashi, N.: A study of atmospheric ammonia by means of modeling analysis in the Kanto region of Japan. *Atmos. Environ.*, 39(2), 203-210, 2005.

Singles, R.J., Sutton, M.A., Weston, K.J.: A multi-layer model to describe the atmospheric transport and deposition of ammonia in Great Britain. *Atmos. Environ.*, 32, 393-399, 1998.

Skjøth, C. A., Hertel, O., and Ellermann, T.: Use of the ACDEP trajectory model in the Danish nation-wide Background Monitoring Programme, *Physics and Chemistry of the Earth*, 27, 1469-1477, 2002.

Skjøth, C. A., Hertel, O., Gyldenkerne, S., and Ellermann, T.: Implementing a dynamical ammonia emission parameterization in the large-scale air pollution model ACDEP,

[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)[Discussion Paper](#)

J. Geophys. Res., D06306, 109, 1-13, 2004, doi:10.1029/2003JD003895

Skjøth, C. A., Geels, C., Berge, H., Gyldenkerne, S., Fagerli, H., Ellermann, T., Frohn, L. M., Christensen, J., Hansen, K. M., Hansen, K., and Hertel, O.: Spatial and temporal variations in ammonia emissions - a freely accessible model code for Europe, Atmos. Chem. Phys., 11, 5221-5236, 2011.

Wu, S. Y., Hu, J. L., Zhang, Y., Aneja, V. P.: Modeling atmospheric transport and fate of ammonia in North Carolina - Part II: Effect of ammonia emissions on fine particulate matter formation. Atmos. Environ., 42 (14), 3437-3451, 2008.

Zhang, Y., Dore, A. J., Liu, X., and Zhang, F.: Simulation of nitrogen deposition in the North China Plain by the FRAME model, Biogeosciences, 8, 3319-3329, 2011.

---

Interactive comment on Geosci. Model Dev. Discuss., 5, 2745, 2012.

**GMDD**

5, C1350–C1357, 2013

---

Interactive  
Comment

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Discussion Paper

C1357

