



This discussion paper is/has been under review for the journal Geoscientific Model Development (GMD). Please refer to the corresponding final paper in GMD if available.

$\delta^{18}\text{O}$ water isotope in the *iLOVECLIM* model (version 1.0) – Part 3: A paleoperspective based on present-day data-model comparison for oxygen stable isotopes in carbonates

T. Caley¹ and D. M. Roche^{1,2}

¹Earth and Climate Cluster, Faculty of Earth and Life Sciences, Vrije Universiteit Amsterdam, Amsterdam, the Netherlands

²Laboratoire des Sciences du Climat et de l'Environnement (LSCE), CEA/CNRS-INSU/UVSQ, Gif-sur-Yvette Cedex, France

Received: 5 February 2013 – Accepted: 18 February 2013 – Published: 4 March 2013

Correspondence to: T. Caley (t.caley@vu.nl)

Published by Copernicus Publications on behalf of the European Geosciences Union.

GMDD

6, 1527–1558, 2013

$\delta^{18}\text{O}$ water isotope
in the *iLOVECLIM*
model

T. Caley and D. M. Roche

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Abstract

Oxygen stable isotopes ($\delta^{18}\text{O}$) are among the most usual tools in paleoclimatology/paleoceanography. Simulation of oxygen stable isotopes allows testing how the past variability of these isotopes in water can be interpreted. By modelling the proxy directly in the model, the results can also be directly compared with the data. Water isotopes have been implemented in the global three-dimensional model of intermediate complexity *iLOVECLIM* allowing fully coupled atmosphere-ocean simulations. In this study, we present the validation of the model results for present day climate against global database for oxygen stable isotopes in carbonates. The limitation of the model together with the processes operating in the natural environment reveal the complexity of use the continental calcite $\delta^{18}\text{O}$ signal of speleothems for a data-model comparison exercise. On the contrary, the reconstructed surface ocean calcite $\delta^{18}\text{O}$ signal in *iLOVECLIM* does show a very good agreement with late Holocene database (foraminifers) at the global and regional scales. Our results indicate that temperature and the isotopic composition of the seawater are the main control on the fossil $\delta^{18}\text{O}$ signal recorded in foraminifer shells and that depth habitat and seasonality play a role but have secondary importance. We argue that a data-model comparison for surface ocean calcite $\delta^{18}\text{O}$ in past climate, such as the last glacial maximum ($\approx 21\,000$ yr), could constitute an interesting tool for mapping the potential shifts of the frontal systems and circulation changes throughout time. Similarly, the potential changes in intermediate oceanic circulation systems in the past could be documented by a data (benthic foraminifers)-model comparison exercise whereas future investigations are necessary in order to quantitatively compare the results with data for the deep ocean.

1 Introduction

Water isotopes are commonly used as important tracers of the hydrological cycle. Because of small chemical and physical differences between the main isotopic forms of

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

the water molecule (H_2^{16}O , HDO, H_2^{18}O), an isotopic fractionation occurs, principally during phase transitions of water (evaporation and condensation processes). Stable water isotopes have therefore been measured in a large variety of archives to reconstruct regional climate variations. In polar ice cores, the isotopic composition of the ice has long been used to reconstruct past air temperatures (Dansgaard, 1953; Jouzel, 2003, 2007). For tropical ice cores (Thompson et al., 2000), temperature but also precipitation intensity upstream of the air mass trajectories seems to play an important role to explain water isotopes signal (Hoffmann, 2003; Vimeux et al., 2005). Over the continent, water isotope concentration is preserved in carbonate from speleothems or lacustrine organism such as ostracodes, reflecting the isotopic composition of meteoric water and the temperature dependent water–calcite oxygen isotope fractionation (Urey, 1947; Kim and O’Neil, 1997). For speleothems in tropical regions, some studies have suggested that the amount of precipitation, called amount effect, (Dansgaard, 1964; Rozanski et al., 1993) could be the dominant driver of water isotope concentration (Burns et al., 2002; Fleitmann et al., 2003; Wang et al., 2001). In lakes, water isotopes are commonly attributed to changes in water isotope composition which can be dependent on temperature, air mass source area and/or precipitation/evaporation ratio (Stuiver, 1970; Leng and Marshall, 2004).

Similarly to paleo-continental archives, the seawater oxygen isotope concentration ($\delta^{18}\text{O}_{\text{sw}}$) is preserved in carbonates from various organisms such as foraminifers and corals. The carbonate isotopic concentration is mainly controlled by temperature and by the isotopic composition of seawater during their formation (Urey, 1947; Shackleton, 1974).

If water isotopes are one of the most widely used proxy in climate archives from the different realms, processes that control their compositions are various and complex. Models allow testing how the past variability of isotopes in water can be interpreted. For example, the importance of temperature versus precipitation changes for the climatic interpretations of tropical isotopic records (Risi et al., 2010). Models can also help to determine the validity of using modern spatial gradients to derive temporal gradient, i.e.

$\delta^{18}\text{O}$ water isotope in the *iLOVECLIM* model

T. Caley and D. M. Roche

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



the co-variability of climate and water isotopes in a specific location over time (Schmidt et al., 2007; Legrande and Schmidt, 2009). By modelling the proxy (water isotopes) directly in the model, the results can also be directly compared with the data. In addition, simulation of climate and its associated isotopic signal can provide a “transfer function” between isotopic signal and the considered climate variable (Sturm et al., 2010).

General Circulation Models (GCMs) have been frequently used for isotopic studies (Jouzel et al., 1987; Joussame and Jouzel, 1993; Hoffmann et al., 1998; Schmidt, 1998; Delaygue et al., 2000; Paul et al., 1999; Noone and Simmonds, 2002; Mathieu et al., 2002; Lee et al., 2007; Schimdt et al., 2007; Yoshimura et al., 2008; Zhou et al., 2008; Tindall et al., 2009; Risi et al., 2010; LeGrande and Schmidt, 2011; Werner et al., 2011; Xu et al., 2012). Among them, the majority has been used exclusively to simulate water isotopes separately in the atmosphere and ocean components. Oxygen isotopes have been implemented in the three-dimensional model of intermediate complexity called *iLOVECLIM* (version 1.0), allowing fully coupled atmosphere-ocean simulations. The detailed implementations of oxygen isotopes in *iLOVECLIM* can be found in Roche (2013). The present work aims at showing preindustrial experiment results in comparison to different global datasets from marine and continental realms. We mainly focus our discussion on carbonate archives that are extensively used in climate studies. The validation of the model results against water isotopes observations can be found in Roche and Caley (2013). The benefits, caveats, agreements and problems from this data-model comparison for oxygen stable isotopes in carbonates will be discussed to determine the potential and the interest of using *iLOVECLIM* for paleoclimatic studies.

$\delta^{18}\text{O}$ water isotope in the *iLOVECLIM* model

T. Caley and D. M. Roche

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



2 Method

2.1 Water isotopes definition, notation and fractionation

Isotopic values are measured by a mass spectrometer and expressed relative to a standard as (δ) notations (Sharp, 2007):

$$^5 \delta^{18}\text{O} = R(^{18}\text{O})_p / R(^{18}\text{O})_{\text{std}} - 1$$

R is the abundance ratio of the heavy and light isotopes (e.g. $N(^{18}\text{O})_p / N(^{16}\text{O})_p$) for substance P and δ is commonly reported in units of parts per thousand (‰). The standard for carbonate is Pee Dee Belemnite (PDB) (Craig, 1957) and that for waters is Standard Mean Ocean Water (SMOW) (Baertschi, 1976). More recently, standards 10 have been defined with the prefix “Vienna” or VSMOW and VPDB to account for the exhaustion of the original standard materials (Coplen, 1996).

The magnitude of fractionation can be expressed on the δ scale through the logarithmic transformation $10^3 \ln(\alpha_{A-B})$ where $\alpha_{A-B} = R_A / R_B$, α is the equilibrium fractionation and R is the partitioning of isotopes between two substances or two phases of the 15 same substance A and B .

2.2 Oxygen isotopes in *iLOVECLIM*

The *iLOVECLIM* (version 1.0) model is a derivative of the LOVECLIM-1.2 climate model extensively described in Goosse et al. (2010). With regards to water isotopes, the main development lies in the atmospheric component in which evaporation, condensation 20 and existence of different phases (liquid and solid) all affect the isotopic conditions of the water isotopes. In the ocean, the water isotopes are acting as passive tracers ignoring the small fractionation implied by the presence of sea-ice (Craig and Gordon, 1965). For the land surface model, the implementation in the bucket follows the same procedure as for the water except that equilibrium fractionation is assumed during phase

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



changes. A detailed description of the method used to compute the water isotopes in iLOVECLIM can be found in Roche (2013).

2.3 Global datasets

Global oxygen isotopic datasets for the atmosphere and ocean components have been compiled from published Late Holocene measurements to allow comparison and discussion with *iLOVECLIM* results. Table S1 is a compilation of 74 $\delta^{18}\text{O}$ calcite measurements and drip water data of speleothems from the literature. The $\delta^{18}\text{O}$ of the calcite are reported for the last 1000 yr of each record with the associated 2σ error when it was possible to calculate it. Table S1 also include a compilation of 19 $\delta^{18}\text{O}$ data of ice core from the literature. The $\delta^{18}\text{O}$ of the ice are reported for the last 1000 yr of each record with the associated 2σ error when it was possible to calculate it. For the EPICA Dome C, Vostok and Illimani site the δD values were converted in $\delta^{18}\text{O}$ values using the global meteoric water line $\delta\text{D} = 8.2 \cdot \delta^{18}\text{O} + 11.27$ (Rozanski et al., 1993). To compare global oceanic isotopic datasets with *iLOVECLIM* results, we used the late Holocene planktonic foraminifer's $\delta^{18}\text{O}$ database of Waelbroeck et al. (2005) and extend it with 62 data averaging over the last 3000 yr (Table S2). Similarly, for the intermediate and deep ocean, we used the published Atlantic benthic foraminifer's stable isotopic database of Marchal and Curry (2008) and extend it with 73 data covering the global ocean and averaged over the last 3000 yr (Table S2).

20 3 Results and discussion

3.1 Oxygen isotopes: from the atmosphere to continental records

The water evaporated from the surface ocean undergoes a poleward transport over landmasses that results in gradual rainout and a depletion of the remaining moisture in ^{18}O . The $\delta^{18}\text{O}$ in precipitation shows therefore systematic variations with latitude, altitude and distance from vapour source. The $\delta^{18}\text{O}$ in precipitation can then be conserved

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



in continental carbonate archives. Under equilibrium conditions, the $\delta^{18}\text{O}$ of carbonates (speleothem) depends on both the temperature through its control on equilibrium fractionation between water and calcite (Hendy, 1971; Kim and O'Neil, 1997) and the isotopic composition of the drip water from the cave site in which the speleothem grew.

Using the Kim and O'Neil (1997) equation for synthetic calcite and making the approximation that $1000 \cdot \ln \alpha(\text{calcite-water}) \approx \delta_{\text{calcite}} - \delta_{\text{water}}$, it can be demonstrated that $\delta^{18}\text{O}_{\text{calcite(speleothem)}} = \delta^{18}\text{O}_{\text{water}} + (18.03 \cdot (1000/T) - 32.17)$. T is the temperature in Kelvin and the relationship between the two scales (PDB and SMOW) can be expressed as $\delta^{18}\text{O}(\text{PDB}) = 0.97002(\delta^{18}\text{O}(\text{SMOW})) - 29.98$ (Coplen et al., 1983; Sharp, 2007).

Figure 1 shows the results obtain with *iLOVECLIM* for the present day $\delta^{18}\text{O}$ in precipitation in comparison to drip water data from speleothems and ice core data (Table 1). The observed relationship is good ($R^2 = 0.77$). This confirms that the model is capable to represent the global distribution of $\delta^{18}\text{O}$ in precipitation as demonstrated in Roche and Caley (2013) although an underestimation of the fractionation towards lower temperatures/latitudes is observed (Roche and Caley, 2013).

Combining the $\delta^{18}\text{O}$ of precipitation and atmospheric temperature and using the equation of Kim and O'Neil (1997), we are able to reconstruct the calcite $\delta^{18}\text{O}$ in the model and to compare it with late Holocene calcite $\delta^{18}\text{O}$ data (Fig. 2 and Table S1).

We conducted this analysis separately for the European region and for the rest of the world. Indeed, the interpretation of the $\delta^{18}\text{O}$ in precipitation signal over Europe has been found to be complex in *iLOVECLIM*, probably linked to the representation of the Mediterranean region in the model (Roche and Caley, 2013). Nonetheless, a discrepancy between model results and data exist also when the Europe area is excluded (Fig. 2). An insignificant correlation ($R^2 = 0.1$) can be observed between measured calcite $\delta^{18}\text{O}$ and model results (Fig. 2). This weak relationship can be explained by a combination of factors. First, the representation of the $\delta^{18}\text{O}$ in precipitation with the model. In particular the low resolution and simple representation of the atmospheric component that complicates certain aspects of the data-model comparison (Roche

$\delta^{18}\text{O}$ water isotope in the *iLOVECLIM* model

T. Caley and D. M. Roche

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

◀

▶

Back

Close

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



and Caley, 2013). Second, the altitudinal effect and its impact on the $\delta^{18}\text{O}$ in precipitation (around $-2.5\text{‰}/1000\text{ m}$; Lachniet, 2009) is not fully taken into account in different areas due low resolution of the atmospheric model preventing to obtain high elevations in mountain ranges (e.g. the Alps are $\approx 1200\text{ m}$ in height). Nonetheless a correction of this altitudinal effect does not significantly change the results of the correlation for our global database (not shown). Another factor is the use of atmospheric temperatures rather than cave temperatures in the calcite $\delta^{18}\text{O}$ calculation with the modelling results. Indeed, the relationship between measured temperature in the cave and atmospheric temperature in the model is significant but weak ($R^2 = 0.63$, Fig. 3). Taken together, all these limitations lead to a bias of the calcite $\delta^{18}\text{O}$ reconstructed when using the model results. Another source of uncertainties of this data-model comparison correspond to all potential biases which occurs in the natural environment, processes operating in the atmosphere, soil zone, epikarst and cave system (Lachniet, 2009) and affecting the measured calcite $\delta^{18}\text{O}$ signal (Fig. 2). For example, recent studies suggest the precipitation of the calcite in different caves do not occur under equilibrium conditions indicating significant kinetic isotope effects during speleothem calcite growth (Mickler et al., 2006; Daëron et al., 2011). This is an important limitation for a global comparison between model and data.

For a better representation at the local scale of the calcite $\delta^{18}\text{O}$ signal in tropical and mid latitude region, models with higher complexity and resolution are necessary (Wackerbarth et al., 2012). Furthermore, more processes operating between the atmosphere and the cave system need to be integrated, in addition to the knowledge of the isotopic composition of precipitation and temperature, to allow a good reconstruction of the calcite $\delta^{18}\text{O}$ signal.

Another possibility is to use the method recently developed to extract the fossil water of speleothem record (Vonhof et al., 2006; van Breukelen et al., 2008). With this water, it becomes possible to reconstruct hydrogen and oxygen stable isotope analyses of fossil precipitation. Based on our work, it appears that reconstructed fossil water isotopic signals may allow a better comparison with modelling results (Fig. 1). The signal would

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



also be more easily related to tropical precipitation (Roche and Caley, 2013) than the complex calcite $\delta^{18}\text{O}$ signal.

3.2 Oxygen isotopes: from the ocean to the foraminifer shells

The seawater oxygen isotope concentration ($\delta^{18}\text{O}_{\text{sw}}$) tracks regional freshwater sources (evaporation, precipitation and river input) and water mass exchange through surface ocean fluxes and ocean circulation (Jacobs et al., 1985; Skinner et al., 2003). It yields information on changes in the hydrological cycle in the past. The $\delta^{18}\text{O}_{\text{sw}}$ is preserved in ocean carbonates archives as foraminifer shells. $\delta^{18}\text{O}$ of foraminifers calcite is mainly controlled by temperature and by the isotopic composition of seawater in which the shell grew (Urey, 1947; Shackleton, 1974). The temperature dependence of the equilibrium fractionation of inorganic calcite precipitation around 16.9 °C is given in Shackleton (1974) as:

$$T = 16.9 - 4.38(\delta^{18}\text{O}_{\text{carbonate(PDB)}} - \delta^{18}\text{O}_{\text{sw(SMOW)}}) + 0.1(\delta^{18}\text{O}_{\text{carbonate(PDB)}} - \delta^{18}\text{O}_{\text{sw(SMOW)}})^2$$

15

Mathematically, we can write the inverse equation for the $\delta^{18}\text{O}$ of the water:

$$\delta^{18}\text{O}_{\text{sw(SMOW)}} = \delta^{18}\text{O}_{\text{carbonate(PDB)}} + 0.27 - 21.9 + \sqrt{(310.6 + 10T)}$$

The conversion between the two scale (PDB and SMOW) can be expressed as $\delta^{18}\text{O}_{\text{water(VPDB)}} = \delta^{18}\text{O}_{\text{water(VSMOW)}} - 0.27$ (Hut, 1987).

20 3.2.1 Surface ocean

Combining the $\delta^{18}\text{O}_{\text{sw}}$ and oceanic temperature and using the equation of Shackleton (1974), we are able to reconstruct the calcite $\delta^{18}\text{O}$ in the model and to compare it with late Holocene $\delta^{18}\text{O}$ planktonic foraminifers' data (Fig. 4). We choose specifically the

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



**$\delta^{18}\text{O}$ water isotope
in the iLOVECLIM
model**

T. Caley and D. M. Roche

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

et al., 2004) and *N. Pachyderma dextral* calcifies throughout the year (Ostermann et al., 2001).

To take into account the seasonal effect we calculate the amplitude in calcite $\delta^{18}\text{O}$ for the two extreme months in comparison to the mean annual value (Fig. 6). These two end-members represent the extreme configurations in which blooms lead to a 100 % of the carbonate deposit. The range between these two end-members corresponds to all the different possibilities for carbonate deposit. The entire range is from 0–2 % with a weak or null seasonal effect at low and high latitudes whereas we observe the more pronounced seasonal effect at mid latitudes. This seasonal range in calcite $\delta^{18}\text{O}$ values could explain part of the shift and dispersion observed with the equilibrium 1 : 1 line but correspond to a factor of secondary importance (Fig. 5).

We then realized a data-model comparison for the calcite $\delta^{18}\text{O}$ signal species by species (Fig. 7). A significant relationship is found for each species except *N. Pachyderma sinistral*. This foraminifer lives at high latitudes, in polar waters. Some caveats in the model for the Polar Regions as well as complex environment factor driven the $\delta^{18}\text{O}$ signal of this specie are probably responsible for the weak correlation observed (red circle on Fig. 7). For example, *N. Pachyderma* as been found to have a secondary calcification phase at depth which can be as much as 75 % of the individual mass (Kohfeld et al., 1996). Also, the depth habitat of this species has been found to be very variable (Ottens, 1992; Kohfeld et al., 1996; Bauch, 1997; Ostermann et al., 2001; Bauch, 2002; Pak and Kennett, 2002; Simstich, 2003; Pak et al., 2004; Kuroyangi and Kawahata, 2004; von Langen et al., 2005; Came et al., 2007).

Overall, the model is able to well reproduce the data. This implies that temperature and the isotopic composition of the seawater are the main control on the fossil $\delta^{18}\text{O}$ signal recorded in foraminifer's shells at the global scale. Depth life and seasonality must also play a role, but of secondary importance (Figs. 5 and 6). The other biases linked to sedimentation, ecological effects (dissolved carbonate concentration or the photosynthetic activity of algal symbiontes) (Bemis et al., 1998) and post-deposit effects such

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

as bioturbation or dissolution (Waelbroeck et al., 2005) can be considered as weak compared to the other controls cited previously.

To test the ability of the model to represent the data at a regional scale we realized zooms on the North Atlantic Ocean and South Atlantic Ocean (Fig. 8a). We investigate the potential of the calcite $\delta^{18}\text{O}$ signal to track the main hydrographic conditions in these regions. To determine the position of the fronts, we use the actual temperature and salinity changes that characterize these fronts and convert it into calcite $\delta^{18}\text{O}$ values. The Subarctic Front (SAF) has a high range of variability for temperature and salinity values with a mean annual around 13 °C and 35.4 salinity units (psu), respectively (Belkin and Levitus, 1996 and references therein). Using the actual relationship of Legrande and Schmidt (2006) for the North Atlantic Ocean ($\delta^{18}\text{O}=0.55 \cdot \text{sea surface salinity (SSS)} - 18.98$) and the equation of Shackleton (1974), we obtain a value of 1.13 ‰ for the $\delta^{18}\text{O}_{\text{calcite}}$. The Polar Front (PF) is characterized by surface water temperature and salinity lower than 0 °C and 34.4 psu, respectively (Bauch et al., 1999). Using the actual relationship of Legrande and Schmidt (2006) for the GIN seas ($\delta^{18}\text{O}=0.6 \cdot \text{SSS} - 20.71$), we obtain a value of 3.94 ‰ for the $\delta^{18}\text{O}_{\text{calcite}}$. The transition between the Atlantic waters and the arctic domain is through the arctic front (AF). In this area, the Atlantic inflow of water has salinity above 35 psu and a temperature of 5 °C (Skagseth et al., 2008) leading to a $\delta^{18}\text{O}_{\text{calcite}}$ value of 2.93 ‰. These different values of $\delta^{18}\text{O}_{\text{calcite}}$ track faithfully the actual position of the front as described in Dickson et al. (1988) and are well documented by our data-model comparison results (Fig. 8a).

Similarly, the main hydrographic conditions in the South Atlantic Ocean with the hydrographic fronts are well documented by the calcite $\delta^{18}\text{O}$ signal (Fig. 8b). The STF is best defined by salinity (the 34.81 psu halocline in this area) (Edwards and Emery, 1982) or the 11 °C isotherm at 100 m depth. Using the actual relationship of Legrande and Schmidt (2006) for the South Atlantic Ocean ($\delta^{18}\text{O}=0.51 \cdot \text{SSS} - 17.4$) and the equation of Shackleton (1974), we obtain a value of 0.77 ‰ for the $\delta^{18}\text{O}_{\text{calcite}}$. For the SAF, with use the definition of the isotherm 4 °C near 200 m depth, and a salinity of

$\delta^{18}\text{O}$ water isotope in the iLOVECLIM model

T. Caley and D. M. Roche

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



34.5 psu south of the front (Peterson et Stramma, 1991). Using the actual relationship of Legrande and Schmidt (2006) for the Austral Ocean ($\delta^{18}\text{O}=0.24 \cdot \text{SSS} - 8.45$) we can calculate a $\delta^{18}\text{O}_{\text{calcite}}$ value of 2.65 ‰. For the polar front (PF), the definition relies only on the isotherm and not to salinity. Nowlin et al. (1977) used the northern extent of the 0 °C isotherm. We also use a mean value of −0.4 ‰ for the $\delta^{18}\text{O}_{\text{sw}}$ of the austral ocean (Legrande and Schmidt, 2006) leading to a value of 3.6 ‰ for $\delta^{18}\text{O}_{\text{calcite}}$. These different values of $\delta^{18}\text{O}_{\text{calcite}}$ track well the actual position of the front as described in Peterson and Stramma (1991) (Fig. 8b).

Applying this data-model comparison for calcite $\delta^{18}\text{O}$ in past climate could constitute an interesting tool for mapping the potential shifts of the frontal systems and circulation changes through time. Previous data studies on the amplitude of the calcite $\delta^{18}\text{O}$ have documented hydrographic changes during the 8.2 kyr event, the younger dryas event, Heinrich events and the Last Glacial Maximum (LGM) (Cortijo et al., 2005; Eynaud et al., 2009).

15 3.2.2 Intermediate and deep ocean

Since we obtain very good results at the surface of the ocean it is interesting to investigate intermediate and deep ocean results in comparison to available data of benthic foraminifers. To document the intermediate depth circulation we present on Fig. 9a the ocean calcite $\delta^{18}\text{O}$ for a mean depth of 500–1500 m. A good agreement is observed between model results and data. The North Atlantic drift is well represented and the penetration of the Antarctic intermediate waters moving northward is also visible. Any potential changes in these oceanic circulation systems in the past could be documented by a data-model comparison exercise. Outside of the Atlantic Ocean region, the data are too rare to validate the model results. Supplementary data in the Indian Ocean, Pacific Ocean and Southern Ocean would be a major advance in order to validate the observed distribution in calcite $\delta^{18}\text{O}$ at intermediate depth.

Concerning the mean deep ocean (2000–4000 m), Fig. 9b indicates a relatively good agreement between model results and data for the Atlantic Ocean contrary to the other

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

parts of the world ocean. Nonetheless, the calcite $\delta^{18}\text{O}$ simulated with *iLOVECLIM* for the deep ocean is globally too high in comparison to data. The values in the model are actually close to values reached between interglacial and glacial periods (3.8–4.2‰). After investigation, it appears that two factors can explain this abnormally high $\delta^{18}\text{O}$ signal. First the modelled distribution of $\delta^{18}\text{O}_{\text{sw}}$ is 0.25 per mil lower than data between 20° S and 30° N (Roche and Caley, 2013). Second, the modelled temperature of deep water is around 2 °C lower than data, a pattern particularly marked in the austral ocean. The caveat for the deep ocean temperature with the model would be the matter of future investigation. Together, these two biases explain the high $\delta^{18}\text{O}$ calcite signal in the model. These results also elegantly suggest that changes in the deep water temperature by only few degrees can have an important effect of the benthic $\delta^{18}\text{O}$ calcite signal, a pattern that requires further investigation during past climate changes.

4 Conclusion

Concerning the atmospheric component and contrary to the $\delta^{18}\text{O}$ in precipitation, the limitation of the model together with the processes operating in the atmosphere, soil zone, epikarst and cave system hamper a good data model comparison for the calcite $\delta^{18}\text{O}$ signal. It remains to be tested whether the use of more complex isotope – enabled models at higher resolution added to the inclusion of supplementary processes operating between the atmosphere and the cave system would allow a better data – model comparison. It is necessary to assess such improvements in parallel with improvements regarding the potential biases occurring in the natural environment and affecting the measured calcite $\delta^{18}\text{O}$ signal. Based on our work and for simplicity reasons, it appears that reconstructed fossil water isotopic signals by fluid inclusions in speleothems is a more promising way to compare simulation results with climate data, a comparison which is today restricted to ice core records. The relationship between the $\delta^{18}\text{O}$ in precipitation and climate variables such as temperature and precipitation rates could be investigated for past time periods.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

**$\delta^{18}\text{O}$ water isotope
in the iLOVECLIM
model**

T. Caley and D. M. Roche

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Concerning the oceanic component, the simulated calcite $\delta^{18}\text{O}$ compare quantitatively well with global Late Holocene planktonic foraminifer $\delta^{18}\text{O}$ data, also when each species is studied separately. Our results indicate that temperature and the isotopic composition of the seawater are the main control on the fossil $\delta^{18}\text{O}$ signal recorded in foraminifer's shells at the global and regional scale. Depth life and seasonality play also a role but have secondary importance. Nonetheless, further works with more sophisticated ecological models are needed to refine these conclusions. We argue that a data-model comparison exercise for calcite $\delta^{18}\text{O}$ in past climate could constitutes an interesting tool to mapping the potential shifts of the frontal systems and circulation changes throughout time. Of particular interest is the ocean circulation at the LGM. By simulating both the calcite $\delta^{18}\text{O}$ and $\delta^{18}\text{O}_{\text{sw}}$ in the model and by comparison with foraminifer's calcite $\delta^{18}\text{O}$ data it would be possible to address the LGM tropical sea surface temperature controversy (large east-west gradients) (Waelbroeck et al., 2009).

Similarly to surface Ocean, the potential changes in intermediate oceanic circulation systems in the past could be documented by a data-model comparison exercise but an extension of the spatial data coverage is necessary. For deep ocean calcite $\delta^{18}\text{O}$ signal, the abnormally low-modelled deep ocean temperature would be the matter of future investigation in order to quantitatively compare the results with data.

Supplementary material related to this article is available online at:

[http://www.geosci-model-dev-discuss.net/6/1527/2013/
gmdd-6-1527-2013-supplement.pdf](http://www.geosci-model-dev-discuss.net/6/1527/2013/gmdd-6-1527-2013-supplement.pdf).

Acknowledgements. T. Caley is supported by NWO through the VIDI/AC²ME project no. 864.09.013. D. M. Roche is supported by NWO through the VIDI/AC²ME project no. 864.09.013 and by CNRS-INSU. The authors wish to thank D. Genty for useful discussion on equilibrium conditions during speleothem calcite growth. Institut Pierre Simon Laplace is gratefully acknowledged for hosting the iLOVECLIM model code under the LUDUS framework project (<https://forge.ipsl.jussieu.fr/ludus>). This is NWO/AC²ME contribution number 03.

References

- Anand, P., Elderfield, H., and Conte, M. H.: Calibration of Mg/Ca thermometry in planktonic foraminifera from a sediment trap time series, *Paleoceanography*, 18, 1050, doi:10.1029/2002PA000846, 2003.
- 5 Baertschi, P.: Absolute ^{18}O content of standard mean ocean water, *Earth Planet. Sci. Lett.*, 31, 341–344, 1976.
- Bauch, D.: Oxygen isotope composition of living *Neogloboquadrina pachyderma* (sin.) in the Arctic Ocean, *Earth Planet. Sci. Lett.*, 146, 47–58, 1997.
- 10 Bauch, D., Erlenkeuser, H., Winckler, G., Pavlova, G., and Thiede, J.: Carbon isotopes and habitat of polar planktic foraminifera in the Okhotsk Sea: the “carbonate ion effect” under natural conditions, *Mar. Micropaleontol.*, 45, 83–99, 2002.
- Bauch, H. A., Erlenkeuser, H., Fahl, K., Spielhagen, R. F., Weinelt M. S., Andrleit, H., and Heinrich, R.: Evidence for a steeper Eemian than Holocene sea surface temperature gradient between Arctic and sub-Arctic regions, *Palaeogeogr. Palaeoclim. Palaeoecol.*, 144, 95–117, 1999.
- 15 Belkin, I. M. and Levitus, S.: Temporal variability of the Subarctic Front near the Charlie-Gibbs Fracture Zone, *J. Geophys. Res.*, 101, 28317–28324, 1996.
- Bemis, B. E., Spero, H. J., Bijma, J., and Lea, D. W.: Reevaluation of the oxygen isotopic composition of planktonic foraminifera: experimental results and revised paleotemperature equations, *Paleoceanography*, 13, 150–160, 1998.
- 20 Burns, S. J., Fleitmann, D., Mudelsee, M., Neff, U., Matter, A., and Mangini, A.: A 780-year annually resolved record of Indian Ocean monsoon precipitation from a speleothem from South Oman, *J. Geophys. Res.-Atmos.*, 107, 4434, doi:10.1029/2001JD001281, 2002.
- Cane, R. E., Oppo, D. W., and McManus, J. F.: Amplitude and timing of temperature and salinity variability in the subpolar North Atlantic over the past 10 kyr, *Geology*, 35, 315–318, 2007.
- 25 Coplen, T. B.: Editorial: more uncertainty than necessary, *Paleoceanography*, 11, 369–370, 1996.
- Coplen, T. B., Kendall, C., and Hopple, J.: Comparison of stable isotope reference samples, *Nature* 302, 236–238, 1983.
- 30 Cortijo, E., Duplessy, J., Labeyrie, L., Duprat, J., and Paillard, D.: Heinrich events: hydrological impact, *C. R. Geosci.*, 337, 897–907, doi:10.1016/j.crte.2005.04.011, 2005.

$\delta^{18}\text{O}$ water isotope in the iLOVECLIM model

T. Caley and D. M. Roche

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



$\delta^{18}\text{O}$ water isotope in the iLOVECLIM model

T. Caley and D. M. Roche

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



- Craig, H.: Isotopic standards for carbon and oxygen and correction factors for mass-spectrometric analysis of carbon dioxide, *Geochim. Cosmochim. Ac.*, 12, 133–149, 1957.
- Craig, H. and Gordon, L. I.: Deuterium and oxygen 18 variations in the ocean and the marine atmosphere, in: *Stable Isotopes in Oceanographic Studies and Paleotemperatures*, edited by: Tongiorgi, E., Consiglio nazionale delle ricerche, Spoleto, 9–122, 1965.
- Daëron, M., Guo, W., Eiler, J., Genty, D., Blamart, D., Boch, R., Drysdale, R., Maire, K., Wainer, G., and Zanchetta, G.: $^{13}\text{C}^{18}\text{O}$ clumping in speleothems: observations from natural caves and precipitation experiments, *Geochim. Cosmochim. Ac.*, 75, 3303–3317, 2011.
- Dansgaard, W.: The abundance of ^{18}O in atmospheric water and water vapour, *Tellus*, 5, 461–469, 1953.
- Dansgaard, W.: Stable isotopes in precipitation, *Tellus*, 16, 436–468, 1964.
- Delaygue, G., Jouzel, J., and Dutay, J. C.: Oxygen 18-salinity relationship simulated by an oceanic general circulation model, *Earth Planet. Sci. Lett.*, 178, 113–123, doi:10.1016/S0012-821X(00)00073-X, 2000.
- Dickson, R. R., Meincke, J., Malmberg, S. A., and Lee, A. J.: The “great salinity anomaly” in the northern North Atlantic 1968–1982, *Prog. Oceanogr.*, 20, 103–151, doi:10.1016/0079-6611(88)90049-3, 1988.
- Edwards, R. J. and Emery, W. J.: Australasian southern ocean frontal structure during summer 1976–77, *Aust. J. Mar. Freshw. Res.*, 33, 3–22, 1982.
- Eynaud, F., de Abreu, L., Voelker, A., Schönfeld, J., Salgueiro, E., Turon, J. L., Penaud, A., Toucanne, S., Naughton, F., Sanchez-Goñi, M. F., Malaizé, B., and Cacho, I.: Position of the Polar Front along the western Iberian margin during key cold episodes of the last 45 ka, *Geochem. Geophys. Geosyst.*, 10, Q07U05, doi:10.1029/2009GC002398, 2009.
- Farmer, E. C., Kaplan, A., de Menocal, P. B., and Lynch-Stieglitz, J.: Corroborating ecological depth preferences of planktonic foraminifera in the tropical Atlantic with the stable oxygen isotope ratios of core top specimens, *Paleoceanography*, 22, PA3205, doi:10.1029/2006PA001361, 2007.
- Fleitmann, D., Burns, S. J., Mudelsee, M., Neff, U., Kramers, J., Mangini, A., and Matter, A.: Holocene forcing of the Indian monsoon recorded in a stalagmite from southern Oman, *Science*, 300, 1737–1739, 2003.
- Ganssen, G. M. and Kroon, D.: The isotopic signature of planktonic foraminifera from NE Atlantic surface sediments: implications for the reconstruction of past oceanic conditions, *J. Geol. Soc.*, 157, 693–699, 2000.

$\delta^{18}\text{O}$ water isotope in the iLOVECLIM model

T. Caley and D. M. Roche

- Goosse, H., Brovkin, V., Fichefet, T., Haarsma, R., Huybrechts, P., Jongma, J., Mouchet, A.,
 Selten, F., Barriat, P.-Y., Campin, J.-M., Deleersnijder, E., Driesschaert, E., Goelzer, H.,
 Janssens, I., Loutre, M.-F., Morales Maqueda, M. A., Opsteegh, T., Mathieu, P.-P.,
 Munhoven, G., Pettersson, E. J., Renssen, H., Roche, D. M., Schaeffer, M., Tartinville, B.,
 Timmermann, A., and Weber, S. L.: Description of the Earth system model of intermediate
 complexity LOVECLIM version 1.2, *Geosci. Model Dev.*, 3, 603–633, doi:10.5194/gmd-3-
 603-2010, 2010.
- Hendy, C. H.: The isotopic geochemistry of speleothems, 1: The calculation of the effects of different modes of formation on the isotopic composition of speleothems and their applicability as paleoclimatic indicators, *Geochim. Cosmochim. Ac.*, 35, 801–824, 1971.
- Hoffmann, G., Werner, M., and Heimann, M.: Water isotope module of the ECHAM atmospheric general circulation model: a study on timescales from days to several years, *J. Geophys. Res.*, 103, 16871–16896, 1998.
- Hoffmann, G., Ramirez, E., Taupin, J. D., Francou, B., Ribstein, P., Delmas, R., Dürr, H.,
 Gallaire, R., Simões, J., Schotterer, U., Stievenard, M., and Werner, M.: Coherent isotope history of Andean ice cores over the last century, *Geophys. Res. Lett.*, 30, 1179,
 doi:10.1029/2002GL014870, 2003.
- Hut, G.: Stable Isotope Reference Samples for Geochemical and Hydrological Investigations, Consultant Group Meeting IAEA, Vienna, 16–18 September 1985, Report to the Director General, International Atomic Energy Agency, Vienna, 1987.
- Jacobs, S. S., Fairbanks, R. G., and Horibe, Y.: Origin and evolution of water masses near the Antarctic continental margin: evidence from $\text{H}_2^{18}\text{O}/\text{H}_2^{16}\text{O}$ ratios in seawater, in: Oceanology of the Antarctic Continental Shelf, *Antarct. Res. Ser.*, vol. 43, edited by: Jacobs, S. S., AGU, Washington, DC, 59–85, 1985.
- Joussaume, S. and Jouzel, J.: Paleoclimatic tracers: an investigation using an atmospheric general circulation model under ice age conditions: 2. Water isotopes, *J. Geophys. Res.*, 98, 2807–2830, 1993.
- Jouzel, J.: Water stable isotopes: atmospheric composition and applications in polar ice core studies, *Treatise Geochem.*, 4, 213–243, 2003.
- Jouzel, J., Koster, R. D., Suozzo, R. J., Russel, G. L., White, J. W. C., and Broecker, W. S.: Simulations of the HDO and H_2^{18}O atmospheric cycles using the NASA GISS General Circulation Model: the seasonal cycle for present-day conditions, *J. Geophys. Res.*, 92, 14739–14760, 1987.

Title Page	
Abstract	Introduction
Conclusions	References
Tables	Figures
◀	▶
◀	▶
Back	Close
Full Screen / Esc	
Printer-friendly Version	
Interactive Discussion	



- 5 Jouzel, J., Masson-Delmotte, V., Cattani, O., Dreyfus, G., Falourd, S., Hoffmann, G., Minster, B., Nouet, J., Barnola, J. M., Chappellaz, J., Fischer, H., Gallet, J. C., Johnsen, S., Leuenberger, M., Loulergue, L., Luethi, D., Oerter, H., Parrenin, F., Raisbeck, G., Raynaud, D., Schilt, S., Schwander, J., Selmo, E., Souchez, R., Spahni, R., Stauffer, B., Steffensen, J. P., Stenni, B., Stocker, T. F., Tison, J. L., Werner, M., and Wolff, E. W.: Orbital and millennial Antarctic climate variability over the past 800 000 yr, *Science*, 317, 793–796, 2007.
- 10 Kim, S.-T. and O'Neil, J. R.: Equilibrium and nonequilibrium oxygen isotope effects in synthetic carbonates, *Geochim. Cosmochim. Ac.*, 61, 3461–3475, 1997.
- Kohfeld, K. E., Fairbanks, R. G., Smith, S. L., and Walsh, I. D.: *Neogloboquadrina pachyderma* (sinistral coiling) as paleoceanographic tracers in polar oceans: evidence from Northeast Water Polynya plankton tows, sediment traps, and surface sediments, *Paleoceanography*, 11, 679–699, 1996.
- 15 Kuroyanagi, A. and Kawahata, H.: Vertical distribution of living planktonic foraminifera in the seas around Japan, *Mar. Micropaleontol.*, 53, 173–196, 2004.
- Lachniet, M. S.: Climatic and environmental controls on speleothem oxygen isotope values, *Quat. Sci. Rev.*, 28, 412–432, 2009.
- 20 Lee, J.-E., Fung, I., DePaolo, D., and Fennig, C. C.: Analysis of the global distribution of water isotopes using the NCAR atmospheric general circulation model, *J. Geophys. Res.*, 112, D16306, doi:10.1029/2006JD007657, 2007.
- LeGrande, A. N. and Schmidt, G. A.: Global gridded dataset of the oxygen isotopic composition in seawater, *Geophys. Res. Lett.*, 33, L12604, doi:10.1029/2006GL026011, 2006.
- 25 LeGrande, A. N. and Schmidt, G. A.: Sources of Holocene variability of oxygen isotopes in paleoclimate archives, *Clim. Past*, 5, 441–455, doi:10.5194/cp-5-441-2009, 2009.
- LeGrande, A. N. and Schmidt, G. A.: Water isotopologues as a quantitative paleosalinity proxy, *Paleoceanography*, 26, PA3225, doi:10.1029/2010PA002043, 2011.
- 30 Leng, M. J. and Marshall, J. D.: Palaeoclimate interpretation of stable isotope data from lake sediment archives, *Quat. Sci. Rev.*, 23, 811–831, 2004.
- Marchal, O. and Curry, W. B.: On the abyssal circulation in the glacial Atlantic, *J. Phys. Oceanogr.*, 38, 2014–2037, doi:10.1175/2008JPO3895.1, 2008.
- Mathieu, R., Pollard, D., Cole, J., White, J. W. C., Webb, R. S., and Thompson, S. L.: Simulation of stable water isotope variations by the GENESIS GCM for modern conditions, *J. Geophys. Res.*, 107, 4037, doi:10.1029/2001JD900255, 2002.

 $\delta^{18}\text{O}$ water isotope in the iLOVECLIM model

T. Caley and D. M. Roche

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

$\delta^{18}\text{O}$ water isotope in the *iLOVECLIM* model

T. Caley and D. M. Roche

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



- Mickler, P., Stern, L., and Banner, J.: Large kinetic isotope effects in modern speleothems, *Geol. Soc. Am. B.*, 118, 65–81, doi:10.1130/B25698.1, 2006.
- Noone, D. and Simmonds, I.: Associations between $\delta^{18}\text{O}$ of water and climate parameters in a simulation of atmospheric circulation for 1979–1995, *J. Clim.*, 15, 3150–3169, 2002.
- Nowlin, W. D., Whitworth III, T., and Pillsbury, R. D.: Structure and transport of the Antarctic circumpolar current at Drake Passage from short-term measurements, *J. Phys. Oceanogr.*, 7, 788–802, 1977.
- Ostermann, D. R., Olafsson, J., Manganini, S., Curry, W. B., and Honjo, S.: A dramatic increase in particle flux in the Iceland Sea since 1997, results from a 15-year time series, 7th International Conference on Paleoceanography, Program and Abstracts, Hokkaido University, Sapporo, Japan, 169 pp., 2001.
- Ottens, J. J.: Planktic foraminifera as indicators of ocean environments in the northeast Atlantic, PhD thesis, Vrije Universiteit, Amsterdam, 189 pp., 1992.
- Pak, D. K. and Kennett, J. P.: A foraminiferal proxy for upper water mass stratification, *J. Foraminiferal Res.*, 32, 319–327, 2002.
- Pak, D. K., Lea, D. W., and Kennett, J. P.: Seasonal and interannual variation in Santa Barbara Basin water temperatures observed in sediment trap foraminiferal Mg/Ca, *Geochem. Geophys. Geosyst.*, 5, Q12008, doi:10.1029/2004GC000760, 2004.
- Paul, A., Miltz, S., Patzold, J., and Wolff, T.: Simulation of oxygen isotopes in a global ocean model, in: Use of Proxies in Paleoceanography: Examples From the South Atlantic, edited by: Fischer, G. and Wefer, G., Springer, New York, 655–686, 1999.
- Peterson, R. G. and Stramma, L.: Upper-level circulation in the South Atlantic Ocean, *Prog. Oceanogr.*, 26, 1–73, 1991.
- Risi, C., Bony, S., Vimeux, F., and Jouzel, J.: Water stable isotopes in the LMDZ4 general circulation model: model evaluation for present day and past climates and applications to climatic interpretation of tropical isotopic records, *J. Geophys. Res.*, 115, D12118, doi:10.1029/2009JD013255, 2010.
- Roche, D. M.: $\delta^{18}\text{O}$ water isotope in the *iLOVECLIM* model (version 1.0) – Part 1: Implementation and verification, *Geosci. Model Dev. Discuss.*, 6, 1467–1494, doi:10.5194/gmdd-6-1467-2013, 2013.
- Roche, D. M. and Caley, T.: $\delta^{18}\text{O}$ water isotope in the *iLOVECLIM* model (version 1.0) – Part 2: Evaluation of model results against observed $\delta^{18}\text{O}$ in water samples, *Geosci. Model Dev. Discuss.*, 6, 1495–1525, doi:10.5194/gmdd-6-1495-2013, 2013.

$\delta^{18}\text{O}$ water isotope in the iLOVECLIM model

T. Caley and D. M. Roche

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Rozanski, K., Araguas-Araguas, L., and Gonfiantini, R.: Isotopic patterns in modern global precipitation, in: Climate Change in Continental Isotopic Records, Geophys. Monogr. Ser., vol. 78, edited by: Swart, P. K., AGU, Washington, DC, 1–36, 1993.

Schmidt, G. A.: Oxygen-18 variations in a global ocean model, Geophys. Res. Lett., 25, 1201–1204, 1998.

Schmidt, G. A., LeGrande, A. N., and Hoffmann, G.: Water isotope expressions of intrinsic and forced variability in a coupled ocean-atmosphere model, J. Geophys. Res., 112, D10103, doi:10.1029/2006JD007781, 2007.

Shackleton, N. J.: Attainment of isotopic equilibrium between ocean water and benthonic foraminifera genus *Uvigerina*: isotopic changes in the ocean during the last glacial, Les méthodes quantitatives d'étude des variations du climat au cours du Pléistocène, Gif-sur-Yvette, Colloque international du CNRS 219, 203–210, 1974.

Sharp, Z.: Principles of Stable Isotope Geochemistry, Pearson Prentice Hall, Upper Saddle River, NJ, 2007.

Simstich, J., Sarnthein, M., and Erlenkeuser, H.: Paired $\delta^{18}\text{O}$ signals of *Neogloboquadrina pachyderma* (s.) and *Turborotalita quinqueloba* show thermal stratification structure in Nordic Seas, Mar. Micropaleontol., 48, 107–125, 2003.

Skagseth, Ø., Furevik, T., Ingvaldsen, R., Loeng, H., Mork, K., Orvik, K., and Ozhigin, V.: Volume and heat transport to the Arctic Ocean via the Norwegian and Barents Seas, in: Arctic Subarctic Ocean Fluxes, edited by: Dickson, R., Meincke, J., and Rhines, P., Springer, New York, 45–64, 2008.

Skinner, L. C., Shackleton, N. J., and Elderfield, H.: Millennial-scale variability of deep-water temperature and delta $\delta^{18}\text{O}_{\text{dw}}$ indicating deepwater source variations in the Northeast Atlantic, 0–34 cal. ka BP, Geochim. Geophys. Geosyst., 4, 1098, doi:10.1029/2003GC000585, 2003.

Stuiver, M.: Oxygen and carbon isotope ratios of fresh-water carbonates as climatic indicators, J. Geophys. Res., 75, 5247–5257, 1970.

Sturm, C., Zhang, Q., and Noone, D.: An introduction to stable water isotopes in climate models: benefits of forward proxy modelling for paleoclimatology, Clim. Past, 6, 115–129, doi:10.5194/cp-6-115-2010, 2010.

Thompson, L. G., Mosley-Thompson, E., and Henderson, K. A.: Ice-core paleoclimate records in tropical South America since the Last Glacial Maximum, J. Quat. Sci., 15, 1579–1600, 2000.

- Thunnel, R., Tappa, E., Pride, C., and Kincaid, E.: Sea-surface temperature anomalies associated with the 1997–1998 El Niño recorded in the oxygen isotope composition of planktonic foraminifera, *Geology*, 27, 843–846, 1999.
- Tindall, J. C., Valdes, P., and Sime, L. C.: Stable water isotopes in HadCM3: isotopic signature of 5 El Niño–Southern Oscillation and the tropical amount effect, *J. Geophys. Res.*, 114, D04111, doi:10.1029/2008JD010825, 2009.
- Tolderlund, D. S. and Bé, A. W. H.: Seasonal distribution of planktonic foraminifera in the western North Atlantic, *Micropaleontology*, 17, 297–329, 1971.
- Urey, H. C.: Thermodynamic properties of isotopic substances, *J. Chem. Soc.*, 562–581, 1947.
- van Breukelen, M. R., Vonhof, H. B., Hellstrom, J. C., Wester, W. C. G., and Kroon, D.: Fossil 10 dripwater in stalagmites reveals Holocene temperature and rainfall variation in Amazonia, *Earth Planet. Sci. Lett.*, 275, 54–60, 2008.
- Vimeux, F., Gallaire, R., Bony, S., Hoffmann, G., and Chiang, J. C. H.: What are the climate 15 controls on δD in precipitation in the Zongo Valley (Bolivia)? Implications for the Illimani ice core interpretation, *Earth Planet. Sci. Lett.*, 240, 205–220, 2005.
- von Langen, P. J., Pak, D. K., Spero, H. J., and Lea, D. W.: Effects of temperature on Mg/Ca in neogloboquadrinid shells determined by live culturing, *Geochem. Geophys. Geosyst.*, 6, Q10P03, doi:10.1029/2005GC000989, 2005.
- Vonhof, H. B., van Breukelen, M. R., Postma, O., Rowe, P. J., Atkinson, T. C., and Kroon, D.: 20 A continuous-flow crushing device for on-line d2H analysis of fluid inclusion water in speleothems, *Rapid Commun. Mass Sp.*, 20, 2553–2558, 2006.
- Wackerbarth, A., Langebroek, P. M., Werner, M., Lohmann, G., Riechelmann, S., Borsato, A., and Mangini, A.: Simulated oxygen isotopes in cave drip water and speleothem calcite in European caves, *Clim. Past*, 8, 1781–1799, doi:10.5194/cp-8-1781-2012, 2012.
- Waelbroeck, C., Mulitza, S., Spero, H., Dokken, T., Kiefer, T., and Cortijo, E.: A global compilation of Late Holocene planktonic foraminiferal $\delta^{18}\text{O}$: relationship between surface water 25 temperature and $\delta^{18}\text{O}$, *Quat. Sci. Rev.*, 24, 853–878, 2005.
- Waelbroeck, C., Paul, A., Kucera, M., Rosell-Mele, A., Weinelt, M., Schneider, R., Mix, A. C., Abelmann, A., Armand, L., Bard, E., Barker, S., Barrows, T. T., Benway, H., Cacho, I., Chen, M. T., Cortijo, E., Crosta, X., de Vernal, A., Dokken, T., Duprat, J., Elderfield, H., Eynaud, F., Gersonde, R., Hayes, A., Henry, M., Hillaire-Marcel, C., Huang, C. C., Jansen, E., Juggins, S., Kallel, N., Kiefer, T., Kienast, M., Labeyrie, L., Leclaire, H., Londeix, L., Mangin, S., Matthiessen, J., Marret, F., Meland, M., Morey, A. E., Mulitza, S., Pflau- 30

 $\delta^{18}\text{O}$ water isotope in the iLOVECLIM model

T. Caley and D. M. Roche

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

- mann, U., Pisias, N. G., Radi, T., Rochon, A., Rohling, E. J., Sbaffi, L., Schafer-Neth, C., Solignac, S., Spero, H., Tachikawa, K., Turon, J. L., and Members, M. P.: Constraints on the magnitude and patterns of ocean cooling at the Last Glacial Maximum, *Nat. Geosci.*, 2, 127–132, 2009.
- 5 Wang, Y. J., Cheng, H., Edwards, R. L., An, Z. S., Wu, J. Y., Shen, C. C., and Dorale, J. A.: A high-resolution absolute-dated Late Pleistocene monsoon record from Hulu Cave, China, *Science*, 294, 2345–2348, 2001.
- Werner, M., Langebroek, P. M., Carlsen, T., Herold, M., and Lohmann, G.: Stable water isotopes in the ECHAM5 general circulation model: toward high-resolution isotope modeling on global scale, *J. Geophys. Res.*, 116, D15109, doi:10.1029/2011JD015681, 2011.
- 10 Williams, D. F., Bé, A. W., and Fairbanks, R. G.: Seasonal stable isotopic variations in living planktonic foraminifera from Bermuda plankton tows, *Palaeogeogr. Palaeoclim. Palaeoecol.*, 33, 71–102, 1981.
- 15 Wu, G.-P. and Hillaire-Marcel, C.: Oxygen isotope compositions of sinistral *Neogloboquadrina pachyderma* tests in surface sediments: North Atlantic Ocean, *Geochim. Cosmochim. Ac.*, 58, 1303–1312, 1994.
- Xu, X., Werner, M., Butzin, M., and Lohmann, G.: Water isotope variations in the global ocean model MPI-OM, *Geosci. Model Dev.*, 5, 809–818, doi:10.5194/gmd-5-809-2012, 2012.
- Yoshimura, K., Kanamitsu, M., Noone, D., and Oki, T.: Historical isotope simulation using re-analysis atmospheric data, *J. Geophys. Res.*, 113, D19108, doi:10.1029/2008JD010074, 2008.
- Zhou, J., Poulsen, C. J., Pollard, D., and White, T. S.: Simulation of modern and middle Cretaceous marine $\delta^{18}\text{O}$ with an ocean-atmosphere general circulation model, *Paleoceanography*, 23, PA3223, doi:10.1029/2008PA001596, 2008.

$\delta^{18}\text{O}$ water isotope in the iLOVECLIM model

T. Caley and D. M. Roche

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



$\delta^{18}\text{O}$ water isotope in the *iLOVECLIM* model

T. Caley and D. M. Roche




Fig. 1. Simulated $\delta^{18}\text{O}$ in precipitation with *iLOVECLIM* in comparison to measured $\delta^{18}\text{O}$ in precipitation from climate records (speleothem dripwater and ice core) (Table S1).

Discussion Paper | Discussion Paper

- [Title Page](#)
- [Abstract](#) [Introduction](#)
- [Conclusions](#) [References](#)
- [Tables](#) [Figures](#)
- [◀](#) [▶](#)
- [◀](#) [▶](#)
- [Back](#) [Close](#)
- [Full Screen / Esc](#)
- [Printer-friendly Version](#)
- [Interactive Discussion](#)

$\delta^{18}\text{O}$ water isotope
in the *iLOVECLIM*
model

T. Caley and D. M. Roche




Fig. 2. Simulated calcite $\delta^{18}\text{O}$ values based on $\delta^{18}\text{O}$ in precipitation and atmospheric temperature from *iLOVECLIM* results (using Kim and O’Neil (1997) equation) in comparison to measured calcite $\delta^{18}\text{O}$ of late Holocene speleothem records (Table S1).

- [Title Page](#)
- [Abstract](#) [Introduction](#)
- [Conclusions](#) [References](#)
- [Tables](#) [Figures](#)
- [◀](#) [▶](#)
- [◀](#) [▶](#)
- [Back](#) [Close](#)
- [Full Screen / Esc](#)
- [Printer-friendly Version](#)
- [Interactive Discussion](#)

$\delta^{18}\text{O}$ water isotope in the *iLOVECLIM* model

T. Caley and D. M. Roche




Fig. 3. Relationship between atmospheric (*iLOVECLIM* results) and cave temperature (Table S1).

Title Page

Abstract

stract Introduction

Conclusion

References

Table

Tables | Figures



Bac

[Close](#)

Full Screen / Esc

[Printer-friendly Version](#)

Interactive Discussion



**$\delta^{18}\text{O}$ water isotope
in the *iLOVECLIM*
model**

T. Caley and D. M. Roche




Fig. 4. Simulated ocean calcite $\delta^{18}\text{O}$ (0–50 m) for the present day with *iLOVECLIM*. Superimposed dots represent calcite $\delta^{18}\text{O}$ data measured on various shallowest dwelling foraminifer species (Waelbroeck et al., 2005, extended by 62 points: Table S2).

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[|◀](#)[▶|](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

**$\delta^{18}\text{O}$ water isotope
in the *i*LOVECLIM
model**

T. Caley and D. M. Roche




Fig. 5. Relationship between the ocean calcite $\delta^{18}\text{O}$ calculated in the model (0–50 m depth) and calcite $\delta^{18}\text{O}$ measurements on various shallowest dwelling foraminifer species (Waelbroeck et al., 2005 with $1\sigma < 0.3\text{‰}$, extended by 62 points: Table S2).

**$\delta^{18}\text{O}$ water isotope
in the *iLOVECLIM*
model**

T. Caley and D. M. Roche


[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[|◀](#)[▶|](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Fig. 6. Simulated seasonal amplitude for ocean calcite $\delta^{18}\text{O}$ (surface) for present day with *iLOVECLIM*.

$\delta^{18}\text{O}$ water isotope in the *iLOVECLIM* model

T. Caley and D. M. Roche




Fig. 7. Relationship between the ocean calcite $\delta^{18}\text{O}$ calculated in the model (0–25 m for *G. ruber pink*; 0–50 m for *G. ruber white*, *G. sacculifer* and *G. bulloides*; 0–75 m for *N. pachyderma dextral*; 0–150 m for *N. pachyderma sinistral*) and calcite $\delta^{18}\text{O}$ measurements for each shallowest dwelling foraminifer species (Waelbroeck et al., 2005, extended by 62 points: Table S2). The red circle for *N. pachyderma sinistral* indicates points for which the relationship between model and data results is particularly weak. The exclusion of this points leads to a better relationship for the regression ($R^2 = 0.4$).

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)




Fig. 8. Regional comparison between simulated ocean calcite $\delta^{18}\text{O}$ (0–50 m) for the present day with *iLOVECLIM* and calcite $\delta^{18}\text{O}$ measurements on various shallowest dwelling foraminifer species (Waelbroeck et al., 2005, extended by 62 points: Table S2) for (**A**) the North Atlantic Ocean: the North Atlantic Drift (NAD), major boundary return currents (PC, Portugal Current; AC, Azores Current) (Sverdrup et al., 1942; Crowley, 1981; Dickson et al., 1988) and major hydrographic fronts are indicated: PF, Polar Front; AF, Arctic Front; SAF, Subarctic Front (Dickson et al., 1988). (**B**) The South Atlantic Ocean: major hydrographic fronts are indicated: PF, Polar Front; SAF, South Antarctic Front; STF, SubTropical Front (Peterson et Stramma, 1991).

$\delta^{18}\text{O}$ water isotope in the *iLOVECLIM* model

T. Caley and D. M. Roche

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[◀](#)

[▶](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



$\delta^{18}\text{O}$ water isotope in the *iLOVECLIM* model

T. Caley and D. M. Roche




Fig. 9. Simulated ocean calcite $\delta^{18}\text{O}$ (500–1500 m in **A** and 2000–4000 m in **B**) for the present day with *iLOVECLIM*. Superimposed dots represent calcite $\delta^{18}\text{O}$ data measured on benthic foraminifer species (Marchal and Curry (2008), extended by 73 points: Table S2). The north Atlantic drift (red arrow in **A**) is well represented as the penetration of the Antarctic intermediate waters moving northward (black arrow in **A**).