



This discussion paper is/has been under review for the journal Geoscientific Model Development (GMD). Please refer to the corresponding final paper in GMD if available.

The mathematics of the total alkalinity–pH equation: pathway to robust and universal solution algorithms

G. Munhoven

Dépt. d'Astrophysique, Géophysique et Océanographie, Université de Liège,
4000 Liège, Belgium

Received: 21 February 2013 – Accepted: 22 February 2013 – Published: 12 March 2013

Correspondence to: G. Munhoven (guy.munhoven@ulg.ac.be)

Published by Copernicus Publications on behalf of the European Geosciences Union.

GMDD

6, 2087–2136, 2013

Solving the
alkalinity–pH
equation

G. Munhoven

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Abstract

The total alkalinity– pH equation, which relates total alkalinity and pH for a given set of total concentrations of the acid-base systems that contribute to total alkalinity in a given water sample, is reviewed and its mathematical properties established. We prove that the equation function is strictly monotone and always has exactly one positive root. Different commonly used approximations are discussed and compared. An original method to derive appropriate initial values for the iterative solution of the cubic polynomial equation based upon carbonate-borate-alkalinity is presented. We then review different methods that have been used to solve the total alkalinity– pH equation, with a main focus on biogeochemical models. The shortcomings and limitations of these methods are made out and discussed. We then present two variants of a new, robust and universally convergent algorithm to solve the total alkalinity– pH equation. This algorithm does not require any a priori knowledge of the solution. The iterative procedure is shown to converge from any starting value to the physical solution. The extra computational cost for the convergence security is only 10–15 % compared to the fastest algorithm in our test series.

1 Introduction

Biogeochemical models have become indispensable tools to improve our understanding of the cycling of the elements in the Earth system. A central and critical component of almost all biogeochemical models is the pH calculation routine. In ocean carbon cycle models, the air-sea exchange of CO_2 is directly linked to the surface ocean $[CO_2]$; the preservation of biogenic carbonates in the surface sediments at the sea floor is closely linked to the deep-sea $[CO_3^{2-}]$ (Broecker and Peng, 1982). The fractions of CO_2 , HCO_3^- and CO_3^{2-} in the total dissolved inorganic carbon (i.e. the speciation of the carbonate system) are controlled by pH . Hence, pH changes in seawater may directly influence air-sea exchange of CO_2 or the preservation of carbonates in the deep-sea.

Solving the alkalinity– pH equation

G. Munhoven

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Solving the
alkalinity– pH
equation

G. Munhoven

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Conversely, the dissociation of acids, such as carbonic acid, also controls pH : when the ocean takes up or releases CO_2 (e.g. as a result of a rise or a decline of the abundance of CO_2 in the atmosphere), its pH changes. The currently ongoing ocean acidification due to the massive release of CO_2 into the atmosphere by human activity is but one example of such an induced pH change.

The nitrogen cycle is another important biogeochemical cycle where pH plays an important role. The speciation of dissolved ammonium is – as that of any acid-base system – dependent on pH , NH_3 being more abundant than NH_4^+ at high pH , and less abundant at low pH . At $pH > 9$, the concentration of NH_3 in seawater may reach toxic levels.

The realistic modelling of biologically mediated fluxes (e.g. marine primary or export production) requires the co-limitation or even inhibition by different chemical components to be taken into account. The nitrogen and carbon cycles, already mentioned above, strongly interact, both in the ocean and on land. In the ocean, Fe and other metals act as micronutrients and once again, pH plays an important role as the solubility of metals is strongly dependent on pH (Millero et al., 2009). The resulting coupling of the biogeochemical cycles of different elements makes biogeochemical models become more and more complex and pH calculation more and more difficult.

Biogeochemical models are now increasingly used for settings that are strongly different from present-day. Typical applications include future ocean acidification (e.g. Caldeira and Wickett, 2003), the Paleocene-Eocene-Thermal-Maximum (e.g. Ridgwell and Schmidt, 2010), Snowball Earth (e.g. Le Hir et al., 2009) etc. Some commonly used pH solvers may possibly become unstable and produce unreliable results. The convergence properties of currently used solution methods has actually never been systematically tested.

The speciation of any acid system, i.e. the determination of the concentrations of each one of the undissociated and the different dissociated forms of an acid, is an underdetermined problem if only the total concentration and thermodynamic or stoichiometric constants are known. This underdetermination can be lifted if pH is known.

Solving the
alkalinity– pH
equation

G. Munhoven

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Being dependent on temperature and pressure, neither pH nor $[H^+]$ are, however, well suitable for being used in transport equations, and thus in biogeochemical models. In biogeochemical models, the common way to resolve this underdetermination is to consider another conservative quantity: total alkalinity, also called titration alkalinity. Total alkalinity, which is also an experimentally measurable quantity, ties all the different acid systems present in a water sample together and allows us to solve the speciation problem. In comparison to pH , it has the advantage of being a conservative quantity: it is only controlled by its sources and sinks, and it is independent on temperature and pressure (Zeebe and Wolf-Gladrow, 2001).

In the following section, we provide a comprehensive introduction to the concept of alkalinity. In our exploration of the mathematical properties of the equation that relates $[H^+]$ to total alkalinity start with a detailed presentation of various approximations commonly used for present-day seawater. The analysis of the mathematical properties of these approximations will provide useful hints for the characteristics of the general case. In Sect. 3 of this paper, we present solution methods for deriving pH from each of the various approximations to total alkalinity considered. Complications that might possibly arise from the various pH scales that are in use in marine chemistry are elucidated in Sect. 4. In Sect. 5, we then show that there are intrinsic bounds that bracket the root of the total alkalinity– pH equation, and that can be directly derived from the approximation used to represent total alkalinity. The existence of such bounds makes it possible to define a new, universal algorithm to solve the alkalinity– pH equation, that requires no a priori knowledge of the root. A reference implementation of two variants of the new algorithm is presented in Sect. 6. The algorithms are tested for their efficiency and robustness and their performance compared with that of the most common previously published general solution methods.

2 Total alkalinity: general definition and approximations

In the following parts of this section, we review a number of aspects of total alkalinity in natural waters. The main focus will be put onto seawater and on the carbonate system, but all the presented developments can be applied to any natural water sample, provided the required thermodynamic constants are known. We briefly recall the different approximations commonly used for calculating pH and the speciation of acid systems. We will then establish a few basic properties of the expressions that relate the various types of alkalinity to total concentrations and pH . Although simple, these properties do not seem to have been previously explored in detail, nor exploited for designing methods of solution of the alkalinity- pH equation.

Although we primarily focus on modelling in the following developments, the calculation procedures are obviously also applicable in experimental set-ups.

2.1 Total alkalinity: general definition

Total alkalinity, also called titration alkalinity, denoted here Alk_T , reflects the excess of chemical bases of the solution relative to an arbitrary specified zero level of protons, or equivalence point. Ideally, Alk_T represents the amount of bases contained in a sample of seawater that will accept a proton when the sample is titrated with a strong acid (e.g. hydrochloric acid) to the carbonic acid endpoint. That endpoint is located at the pH below which H^+ ions get more abundant in solution than HCO_3^- ions; its value is close to 4.5. H^+ added to water at this pH by adding strong acid will remain as such in solution. Please notice that, for the sake of a simpler notation, we follow here the common usage of denoting protons in solution by H^+ , although free H^+ ions sensu stricto do only exist in insignificantly small amounts in aqueous solutions. Each proton is rather bound to a water molecule to form an H_3O^+ ion, and each of these H_3O^+ in turn is furthermore generally hydrogen bonded to three other H_2O molecules to form an $H_9O_4^+$ ion (Dickson, 1984).

GMDD

6, 2087–2136, 2013

Solving the alkalinity- pH equation

G. Munhoven

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Rigorously speaking, Alk_T is defined as the number of moles of H^+ ions equivalent to the excess of “proton acceptors”, i.e. bases formed from acids characterized by a $pK_A \geq 4.5$ in a solution of zero ionic strength at 25 °C, over “proton donors”, i.e. acids with $pK_A < 4.5$ under the same conditions, per kilogram of sample (Dickson, 1981).

With emphasis on the most important contributors, a rather complete expression for Alk_T in a seawater sample is

$$\begin{aligned} \text{Alk}_T = & [\text{HCO}_3^-] + 2 \times [\text{CO}_3^{2-}] + [\text{B}(\text{OH})_4^-] + [\text{OH}^-] + [\text{HPO}_4^{2-}] + 2 \times [\text{PO}_4^{3-}] + [\text{H}_3\text{SiO}_4^-] \\ & + [\text{NH}_3] + [\text{HS}^-] + 2 \times [\text{S}^{2-}] + \dots - [\text{H}^+]_f - [\text{HSO}_4^-] - [\text{HF}] - [\text{H}_3\text{PO}_4] - \dots \end{aligned} \quad (1)$$

where the ellipses refer to other potential proton donors and acceptors generally present at negligible concentrations only. All of the concentrations are total concentrations (which include free, hydrated and complexed forms of the individual species), except for $[\text{H}^+]_f$, which only includes the free and hydrated forms. There are alternative definitions that can be found in the literature, which lead to similar, although not necessarily exactly the same, expressions. However, the above definition is the one that reflects the titration procedure used to measure alkalinity the most accurately. We will therefore base the following developments upon it.

In other natural water samples (lake, river, or brines) the constituent list in Eq. (1) needs to be adapted: some constituents may be neglected and bases of other acid systems have to be included (e.g. bases derived from organic acids, from dissolved metals, etc.). While total alkalinity in seawater samples typically ranges between about 2 and 2.6 meq kg⁻¹, acid mine drainage samples may even present negative alkalinity, representing the fact that a strong base instead of a strong acid must be added to reach the equivalence pH point of 4.5. Interested readers may refer, e.g. to Kirby and Cravotta III (2005) and references therein for such – from a marine chemist’s point of view – exotic samples.

Total alkalinity as defined above is a conservative quantity with respect to mixing, changes in temperature and pressure (Wolf-Gladrow et al., 2007). It is therefore a cornerstone in biogeochemical cycle models which are most conveniently formulated on

Solving the alkalinity–pH equation

G. Munhoven

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Solving the
alkalinity–pH
equation

G. Munhoven

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



the basis of conservation equations. In such models, definition/Eq. (1) above, or an adequate variant, is used to solve the inverse problem for $[H^+]$. All of the individual species concentrations appearing in Eq. (1) can be expressed in terms of the total concentrations of the acid systems that they respectively belong to and of $[H^+]$. Given the evolutions of the total concentrations of all the acid systems considered (dissociated and non dissociated forms) and of Alk_T – all of which can be derived from appropriate conservation equations – expression (1) is interpreted as an equation for $[H^+]$ or, equivalently, pH. We will therefore call that equation the total alkalinity–pH equation.

2.2 Common approximations for total alkalinity in seawater

Here we first analyse the forward problem for a few specific approximations used for seawater: for given total concentrations of dissolved inorganic carbon, total borate, etc., we analyse how the expressions for the different types of alkalinity change as a function of $[H^+]$. This simple analysis will already provide valuable insight into the overall mathematical properties of the total alkalinity–pH equation and its subcomponents, that we can exploit later for the most general case.

2.2.1 Carbonate alkalinity

The contribution of the carbonic acid system (or carbonate system) to total alkalinity is called carbonate alkalinity and we denote it by Alk_C :

$$Alk_C = [HCO_3^-] + 2[CO_3^{2-}].$$

Upon substitution of the concentrations of the species by their fractional expressions as a function of $[H^+]$,

$$[HCO_3^-] = C_T \frac{K_1[H^+]}{[H^+]^2 + K_1[H^+] + K_1K_2} \quad \text{and} \quad [CO_3^{2-}] = C_T \frac{K_1K_2}{[H^+]^2 + K_1[H^+] + K_1K_2}$$

where C_T is the total concentration of dissolved inorganic carbon ($C_T = [\text{CO}_2] + [\text{HCO}_3^-] + [\text{CO}_3^{2-}]$), K_1 and K_2 are the first and second dissociation constant for carbonic acid, we get

$$\text{Alk}_C = C_T \frac{K_1[\text{H}^+] + 2K_1K_2}{[\text{H}^+]^2 + K_1[\text{H}^+] + K_1K_2}.$$

- 5 For constant C_T , the right-hand side is a strictly decreasing function of $[\text{H}^+]$: its derivative with respect to $[\text{H}^+]$ is strictly negative for positive $[\text{H}^+]$. As a consequence, $0 < \text{Alk}_C < 2C_T$ if $C_T \neq 0$. Both bounds are strict (i.e. they cannot be reached) and represent the limits of $\text{Alk}_C(C_T; [\text{H}^+])$ for $[\text{H}^+] \rightarrow +\infty$ (lower bound) and $[\text{H}^+] \rightarrow 0$ (upper bound), for C_T fixed.

10 2.2.2 Carbonate and borate alkalinity

The second most important component of natural present-day seawater alkalinity is borate alkalinity, Alk_B . Together with the carbonate alkalinity we have

$$\text{Alk}_{CB} = \text{Alk}_C + \text{Alk}_B = [\text{HCO}_3^-] + 2[\text{CO}_3^{2-}] + [\text{B}(\text{OH})_4^-].$$

- 15 Upon substitution of the individual species concentrations by their fractional expressions as a function of $[\text{H}^+]$, we get

$$\text{Alk}_{CB} = C_T \frac{K_1[\text{H}^+] + 2K_1K_2}{[\text{H}^+]^2 + K_1[\text{H}^+] + K_1K_2} + B_T \frac{K_B}{[\text{H}^+] + K_B},$$

- where B_T is the total concentration of dissolved borates. For constant B_T , Alk_B is again a strictly decreasing function with $[\text{H}^+]$, similarly to Alk_C . Hence, for constant C_T and B_T , Alk_{CB} is a strictly decreasing function with $[\text{H}^+]$ and, as a consequence, $0 < \text{Alk}_{CB} < 2C_T + B_T$ as long as $C_T + B_T \neq 0$.

Solving the alkalinity–pH equation

G. Munhoven

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



2.2.3 Carbonate, borate and water self-ionization alkalinity

In a third stage, we may consider the alkalinity that arises from the dissociation of the solvent water itself (by self-ionization) in addition to carbonate and borate alkalinity and get the next important approximation for natural present-day seawater, called *practical alkalinity* by Zeebe and Wolf-Gladrow (2001):

$$\text{Alk}_{\text{CBW}} = \text{Alk}_{\text{CB}} + [\text{OH}^-] - [\text{H}^+] = [\text{HCO}_3^-] + 2[\text{CO}_3^{2-}] + [\text{B}(\text{OH})_4^-] + [\text{OH}^-] - [\text{H}^+].$$

Upon substitution by the respective speciation relationships, we get

$$\text{Alk}_{\text{CBW}} = C_T \frac{K_1[\text{H}^+] + 2K_1K_2}{[\text{H}^+]^2 + K_1[\text{H}^+] + K_1K_2} + B_T \frac{K_B}{[\text{H}^+] + K_B} + \frac{K_W}{[\text{H}^+]} - [\text{H}^+] \quad (2)$$

At this stage, we do not want to insist on subtleties related to $p\text{H}$ scales. Normally, the last term $[\text{H}^+]$ in the two previous equations should actually read $[\text{H}^+]_f$. We will address the difference between $[\text{H}^+]$ and $[\text{H}^+]_f$ in Sect. 4 below.


Since Alk_{CB} is decreasing with $[\text{H}^+]$, for constant C_T and B_T , the same holds for Alk_{CBW} , because $K_W/[\text{H}^+] - [\text{H}^+]$ is again decreasing with $[\text{H}^+]$. However, unlike Alk_{CB} , Alk_{CBW} is unbounded and it can take arbitrarily low values (for $[\text{H}^+] \gg$) and arbitrarily great values (for $[\text{H}^+] \ll$).

2.2.4 Contribution of a generic acid system to total alkalinity


In common seawater, Alk_{CBW} is entirely sufficient even for applications that require high accuracy. However, in some cases other systems than the carbonate and borate systems need to be considered. This is especially the case in suboxic and anoxic waters, such as semi-closed fjords (e.g. Framvaren Fjord in Norway studied by Yao and Millero, 1995) or at a larger scale, the Black Sea (e.g. Dyrssen, 1999), where, e.g. the contribution from sulphides cannot be neglected.

In order to generalize our analysis of the total alkalinity– $p\text{H}$ equation, let us consider a generic acid, denoted by H_nA , that may potentially lead to n successive dissociation

reactions, characterised by stoichiometric dissociation constants K_1, K_2, \dots, K_n , respectively:



\vdots



For simplicity, we omit the “*” superscript commonly used elsewhere to differentiate stoichiometric from thermodynamic dissociation constants (i.e. elsewhere stoichiometric constants generally write K_i^* instead of K_i). Throughout this paper, the constants used will relate concentrations and not activities. As such, they include the effect of activity coefficients that differ from unity. The values of such constants not only depend on temperature and pressure but also on the ionic strength of the solution. Everything developed here furthermore applies to all kinds of acids, be they of Arrhenius, Brønsted-Lowry, Lewis or any other type, even if the adopted notation could possibly suggest that our developments only apply to Arrhenius-type acids.

If we denote the total concentration of dissolved acid H_nA by $[\Sigma\text{A}] = [\text{H}_n\text{A}] + \dots + [\text{A}^{n-}]$, the fractions of undissociated acid and of the various dissociated forms H_{n-1}A^- ,

GMDD

6, 2087–2136, 2013

Solving the alkalinity–pH equation

G. Munhoven

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



$H_{n-2}A^{2-}, \dots, A^{n-}$ are

$$\begin{aligned} \frac{[H_n A]}{[\Sigma A]} &= \frac{[H^+]^n}{[H^+]^n + K_1[H^+]^{n-1} + K_1 K_2[H^+]^{n-2} + \dots + K_1 K_2 \dots K_n} \\ &= \frac{[H^+]^n}{[H^+]^n + \sum_{j=1}^n [H^+]^{n-j} \prod_{i=1}^j K_i} \\ \frac{[H_{n-1} A^-]}{[\Sigma A]} &= \frac{K_1 [H^+]^{n-1}}{[H^+]^n + \sum_{j=1}^n [H^+]^{n-j} \prod_{i=1}^j K_i} \\ &\vdots \\ \frac{[H_{n-j} A^{j-}]}{[\Sigma A]} &= \frac{(\prod_{i=1}^j K_i) [H^+]^{n-j}}{[H^+]^n + \sum_{k=1}^n [H^+]^{n-k} \prod_{i=1}^k K_i} \\ &\vdots \\ \frac{[A^{n-}]}{[\Sigma A]} &= \frac{\prod_{i=1}^n K_i}{[H^+]^n + \sum_{j=1}^n [H^+]^{n-j} \prod_{i=1}^j K_i}. \end{aligned}$$

The joint contribution of all the different dissociated and non dissociated forms of $H_n A$ to alkalinity, proton donors and proton acceptors alike, is then equal to

$$Alk_A = \sum_{j=0}^n (j - m) [H_{n-j} A^{j-}],$$

Solving the alkalinity–pH equation

G. Munhoven

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



where m is an integer constant, which is dependant on the so-called zero proton level of the system under consideration:

- m is such that $pK_m < 4.5 < pK_{m+1}$ if $pK_1 < 4.5$ and $pK_n > 4.5$
- $m = 0$ if $pK_1 > 4.5$
- $m = n$ if $pK_n < 4.5$

Since $pK_m < 4.5$, all of the $H_{n-j}A^{j-}$ in the $H_nA \dots A^{n-}$ system for $j = 0, \dots, m-1$ are proton donors: the last one ($j = m-1$) has a strength of 1 eqmol^{-1} , the second last one ($j = m-2$) of 2 eqmol^{-1} , etc. Since $pK_{m+1} > 4.5$, the dissociation products $H_{n-j}A^{j-}$ for $j = m+1, \dots, n$ are proton acceptors, the first one ($j = m+1$) with a strength of 1 eqmol^{-1} , the second one ($j = m+2$) with a strength of 2 eqmol^{-1} , etc. For the carbonic acid system, e.g. $n = 2$ and $m = 0$; for the boric acid system, $n = 1$ and $m = 0$; for the phosphoric acid system, $n = 3$ and $m = 1$.

From the previous expressions for the species fractions, we then find that

$$\text{Alk}_A([H^+]) = [\Sigma A] \frac{\sum_{j=0}^n (j-m) \Pi_j [H^+]^{n-j}}{\sum_{j=0}^n \Pi_j [H^+]^{n-j}} = [\Sigma A] \left(\frac{\sum_{j=0}^n j \Pi_j [H^+]^{n-j}}{\sum_{j=0}^n \Pi_j [H^+]^{n-j}} - m \right) \quad (3)$$

where we have defined

$$\Pi_j = \prod_{i=1}^j K_i, \quad j = 1, \dots, n \quad \text{and} \quad \Pi_0 = 1 \quad (4)$$

to simplify the notation.

Similar to the carbonate and borate systems above, Alk_A is strictly decreasing with $[H^+]$, for $[\Sigma A]$ fixed. A mathematically rigorous demonstration of this behaviour for the general case is provided in Appendix A.

Solving the alkalinity–pH equation

G. Munhoven

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Solving the
alkalinity–pH
equation

G. Munhoven

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



There are two corollaries of this monotonic behaviour worth emphasizing:

1. For any acid system $H_nA \dots A^{n-}$, Alk_A is bounded: it has a supremum which is equal to $(n - m)[\Sigma A]$ (i.e. the limit for $[H^+] \rightarrow 0$, not actually reachable though), and an infimum, which is equal to $-m[\Sigma A]$ (i.e. its limit for $[H^+] \rightarrow +\infty$, also not actually reachable); both of these could, theoretically, be negative if m is sufficiently large.
2. For a water sample that contains a set of acids $H_{n_i}A_{[i]}$, ($i = 1, \dots$) of respective known total concentrations $[\Sigma A_{[i]}]$ and with zero proton levels respectively characterised by m_i , the total alkalinity–pH equation

$$\sum_i Alk_{A_{[i]}}([H^+]) + \frac{K_w}{[H^+]} - [H^+]_f - Alk_T = 0 \quad (5)$$

has exactly one positive root $[H^+]$, for any given value of Alk_T : the sum of the respective alkalinity contributions over the set $\{H_{n_i}A_{[i]} | i=1, \dots\}$ of all the acid systems active in the sample is a strictly decreasing function of $[H^+]$; the contribution from the dissociation of water is also strictly decreasing with $[H^+]$, and may theoretically take any value between $+\infty$ and $-\infty$.

3 Alkalinity–pH equation in biogeochemical models: approximations and methods of solution

In this section, we are going to review the most common approximations used in ocean carbon and biogeochemical cycle models, focusing on how the corresponding equation is solved.

3.1 Carbonate alkalinity based solutions

The straight approximation $Alk_T \simeq Alk_C$ is often used in textbooks (e.g. Broecker and Peng, 1982). There are only a few models (e.g. Opdyke and Walker, 1992; Walker and

Opdyke, 1995) that use it directly for their carbonate chemistry speciation. For numerical modelling purposes, its usage is indeed somewhat problematic. $[H^+]$ calculated from Alk_T and C_T data, by assuming that $Alk_C = Alk_T$ are typically 30–40 % too low (i.e. 0.15–0.2 pH units too high) for present-day seawater samples. Furthermore, the sensitivity of the C_T - Alk_C system to perturbations is stronger than that of the C_T - Alk_{CBW} system: equilibrium pCO_2 changes, e.g. are of the order of 20 % larger (Munhoven, 1997).

The calculation of $[H^+]$ from C_T - Alk_C remains nevertheless important, as more advanced methods such as those proposed by Bacastow (1981), Peng et al. (1987) or Follows et al. (2006), where Alk_C is iteratively recalculated from more complete approximations to Alk_T (ICAC methods – see below), rely on it.

3.1.1 Fundamental solution

For given Alk_C and C_T ($C_T > 0$), the equation to solve for $[H^+]$ is

$$R_C([H^+]) \equiv C_T \frac{K_1[H^+] + 2K_1K_2}{[H^+]^2 + K_1[H^+] + K_1K_2} - Alk_C = 0. \quad (6)$$

Following our discussion in Sect. 2.2.1, Eq. (6) has a positive root if and only if $0 < Alk_C < 2C_T$; if there is a positive root, it is unique.

Equation (6) can be directly solved after conversion to the quadratic equation

$$P_C([H^+]) \equiv [H^+]^2 + a_1[H^+] + a_0 = 0 \quad (7)$$

where

$$a_1 = K_1 \left(1 - \frac{C_T}{Alk_C} \right) \quad \text{and} \quad a_0 = K_1K_2 \left(1 - \frac{2C_T}{Alk_C} \right)$$

For valid Alk_C values (i.e. for $0 < \text{Alk}_C < 2C_T$), this quadratic equation has two real roots, a positive and a negative one. The positive root is

$$[\text{H}^+] = Q(\text{Alk}_C, C_T) \equiv \frac{K_1}{2} \left(\frac{C_T}{\text{Alk}_C} - 1 + \sqrt{\Delta_C} \right), \quad (8)$$

where

$$\Delta_C = \left(1 - \frac{C_T}{\text{Alk}_C} \right)^2 + 4 \frac{K_2}{K_1} \left(\frac{2C_T}{\text{Alk}_C} - 1 \right). \quad (9)$$

For Alk_C values that are out of range Eq. (7) either has two negative or two complex roots.

3.1.2 Alternative methods

There are other methods to derive $[\text{H}^+]$ from Alk_C and C_T . All of them ultimately seem to rely on the formulae of Park (1969) for deriving the complete speciation of the carbonate system directly from Alk_C and C_T , without explicitly using $[\text{H}^+]$. Antoine and Morel (1995) first calculate $[\text{CO}_2]$ from C_T and Alk_C (which involves the solution of a first parabolic equation), and then derive $[\text{H}^+]$ from the relationship $[\text{CO}_2] = \text{Alk}_C [\text{H}^+]^2 / (K_1 [\text{H}^+] + 2K_1 K_2)$, which requires the solution of a second parabolic equation. Ridgwell (2001) first determines the complete speciation of the carbonate system, referring for the adopted procedure to Millero and Sohn (1992), who actually only report the formulae of Park (1969). He then derives two different estimates for $[\text{H}^+]$, based upon the definitions of the first and second dissociation constants of carbonic acid and finally uses the geometric mean of these two estimates as a solution for Eq. (6).

There are no obvious advantages for calling upon these methods instead of the direct quadratic solution above. Even if carefully implemented, both require a significantly higher number of operations than the solution outlined above. Those methods offer

Solving the alkalinity–pH equation

G. Munhoven

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



a direct access to carbonate speciation (at least in part). That can, however, also be calculated at little extra cost from $[H^+]$.

3.1.3 Iterative carbonate alkalinity correction methods

In most common natural settings, the difference between Alk_C and Alk_T , albeit small, leads to significant errors on $[H^+]$, if Alk_T is used in place of Alk_C and one of the procedures above is used to calculate it from C_T . To overcome this problem, Alk_C can be estimated from Alk_T , and then iteratively corrected until stabilisation occurs. Such a procedure, which we call here Iterative Carbonate Alkalinity Correction (ICAC) can a priori be used with arbitrary chemical compositions, provided Alk_C represents a significant fraction of Alk_T . If Alk_C makes up only a small fraction of Alk_T , the method is likely to exhibit unstable behaviour.

In the most straightforward ICAC method, one starts from a trial value H_0 for $[H^+]$, a first estimate $Alk_{C,0}$ is obtained by subtracting the concentrations of all non-carbonate components from Alk_T . That $Alk_{C,0}$ is then used to calculate a new (improved) estimate H_1 for $[H^+]$ from Eqs. (8) and (9) or one of the alternative methods. H_1 is then used to calculate a new estimate $Alk_{C,1}$ from Alk_T as above and the procedure is iterated until some predefined convergence criterion is fulfilled. This procedure is a classical fixed-point iteration:

$$H_{n+1} = Q(Alk_C(Alk_T, H_n), C_T) \quad (10)$$

In this recurrence, $Alk_C(Alk_T, H_n)$ is the estimate of Alk_C obtained from Alk_T by subtracting all the non-carbonate components estimated by using H_n . Pure fixed point iterative schemes may be prone to convergence problems (slow convergence or no convergence at all). If the procedure is convergent, the rate of convergence is linear.

This plain fixed-point iteration ICAC method was recently made popular again by Follows et al. (2006). These authors argue that in carbon cycle model simulation experiments, where there is little change in pH from one time step to the next, a single

Solving the alkalinity–pH equation

G. Munhoven

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏮

⏭

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Solving the
alkalinity–pH
equation

G. Munhoven

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



iteration may already provide a sufficiently accurate estimate of $[H^+]$ to derive acceptable pCO_2 estimates, for any chosen approximation of total alkalinity. Follows et al. (2006) suggest, if necessary, to repeat the fixed-point iteration until a sufficiently accurate estimate is found.

There are a number of models that rely on the ICAC approach for their pH determination. Peng et al. (1987) consider Alk_{CBW} plus the contributions from silicic and phosphoric acid systems in their representation of total alkalinity.¹ They use an initial value of 10^{-8} and stop their iterations once $|(\Delta H)/H| < 0.005\%$. They report that less than ten iterations are generally sufficient. Antoine and Morel (1995) adopt Alk_{CBW} as an approximation to Alk_T . At each step, they derive $[H^+]$ from C_T and Alk_C by using their special procedure described above. They iterate until two successive Alk_C estimates differ by less than 10^{-8} (no units given). Ridgwell (2001) adopts $Alk_{CB} + [OH^-] + 1.1[PO_4^{3-}]$ as an approximation to total alkalinity. He calculates $[H^+]$ at each step from C_T and Alk_C by using his own procedure described above. GENIE (Ridgwell et al., 2007) initially used the same procedure as Ridgwell (2001); in more recent versions of GENIE, a complete representation of the phosphoric acid component is used (Ridgwell, personal communication, 2012). Arndt et al. (2011) use $Alk_{CBW} + [HS^-]$ as an approximation to total alkalinity in GEOCLIM *reloaded*. They continue to iterate until $|Alk_{CBW} + [HS^-] - Alk_T| < 10^{-6}$ (no units given). The method is further used in LOVECLIM (Anne Mouchet, personal communication, 2012) with Alk_{CBW} as an approximation for total alkalinity (Goosse et al., 2010) and most probably in still some others, that do, unfortunately not provide details about the calculation procedures adopted.

Bacastow (1981) proposed a variant to improve the rate of convergence of fixed-point iterations. That variant only uses the recurrence described above for the first two iterates. From the third iteration on, Bacastow (1981) switches to a secant method to

¹Peng et al. (1987) adopt, however, a slightly different definition of total alkalinity by systematically weighting species by their respective charge. This leads to differences with the phosphoric acid system, e.g.: the definition of Peng et al. (1987) is equivalent to adopting $m = 0$ for the phosphoric acid system.

Solving the
alkalinity–pH
equation

G. Munhoven

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



solve the fixed-point equation $H - Q(\text{Alk}_C(\text{Alk}_T, H)) = 0$.² Fixed-point iterations are thus only used to provide starting values for the solution of the fixed-point equation by the secant method. The rate of convergence of the method is strongly increased by this approach (and the domain of convergence slightly enlarged – see numerical experiments below). However, for some C_T - Alk_T combinations the underlying fixed-point equation may still give rise to convergence problems, even with the secant method. However as will be shown below, the method of Bacastow (1981) is strongly preferable over the pure fixed-point scheme.

The Hadley Centre Ocean Carbon Cycle (HadOCC) model (Palmer and Totterdell, 2001) uses Bacastow's method for its carbonate speciation calculation, with the Alk_{CBW} approximation.

3.2 Carbonate and borate alkalinity based solution

Only a few models appear to use pH calculation routines based upon Alk_{CB} . MBM-MEDUSA (Munhoven and François, 1996; Munhoven, 1997, 2007) is one of them, the model of Marchal et al. (1998) is another one.

3.2.1 Basic formulation and solution methods

The equation to solve for $[\text{H}^+]$ is, for given Alk_{CB} , C_T and B_T ,

$$R_{\text{CB}}([\text{H}^+]) \equiv C_T \frac{K_1[\text{H}^+] + 2K_1K_2}{[\text{H}^+]^2 + K_1[\text{H}^+] + K_1K_2} + B_T \frac{K_B}{[\text{H}^+] + K_B} - \text{Alk}_{\text{CB}} = 0. \quad (11)$$

This equation may be converted into the polynomial equation

$$P_{\text{CB}}([\text{H}^+]) \equiv [\text{H}^+]^3 + c_2[\text{H}^+]^2 + c_1[\text{H}^+] + c_0 = 0 \quad (12)$$

²Bacastow (1981) actually solves the alkalinity equation for the scaled inverse of $[\text{H}^+]$. We provide codes for the two approaches, although we only base our discussions on the version with secant iterations on $[\text{H}^+]$.

$$c_2 = K_B(1 - \frac{B_T}{Alk_{CB}}) + K_1(1 - \frac{C_T}{Alk_{CB}})$$

$$c_1 = K_1 \left(K_B \left(1 - \frac{B_T}{\text{Alk}_{CB}} - \frac{C_T}{\text{Alk}_{CB}} \right) + K_2 \left(1 - 2 \frac{C_T}{\text{Alk}_{CB}} \right) \right)$$

$$c_0 = K_1 K_2 K_B \left(1 - \frac{2C_T + B_T}{\text{Alk}_{CB}} \right).$$

10

The cubic equation could possibly be solved with closed formulae, such as Cardano's formulae (which may, however, suffer from precision problems, require numerically expensive cubic root evaluations or possibly complex arithmetic) or Viète's trigonometric formulae (which require a combination of an arc-cosine, a cosine and a square root). When adopted, the cubic Eq. (12) is therefore generally solved numerically, with a Newton–Raphson scheme. In this case, determining an adequate starting value is the main problem to address in order to design a robust and fast solution algorithm.

15

An excellent initial value for the Newton–Raphson scheme can be found by adopting the following procedure:

- 20

6, 2087–2136, 2013

G. Munhoven

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



▶

▶

[Back](#)

Close

Full Screen / Esc

[Printer-friendly Version](#)

Interactive Discussion



That local minimum, if it exists (i.e. if $c_2^2 - 3c_1 > 0$), is located at

$$H_{\min} = \frac{-c_2 + \sqrt{c_2^2 - 3c_1}}{3} = \frac{-c_1}{c_2 + \sqrt{c_2^2 - 3c_1}}.$$

The Taylor expansion to second order in H_{\min} , thus intersects the H -axis on the right-hand side of H_{\min} at

$$H_0 = H_{\min} + \sqrt{-\frac{P_{CB}(H_{\min})}{\sqrt{c_2^2 - 3c_1}}},$$

provided $P_{CB}(H_{\min}) < 0$. By completing the Taylor expansion to third order, it is straightforward to show that H_0 is greater than the root.

The so-defined H_0 provides an excellent starting value not only for solving the cubic polynomial equation, but also for other iterative methods.

10 3.3 Carbonate, borate and water self-ionization alkalinity

With Alk_{CBW} , C_T and B_T given, the equation to solve is

$$R_{\text{CBW}}([H^+]) \equiv C_T \frac{K_1[H^+] + 2K_1K_2}{[H^+]^2 + K_1[H^+] + K_1K_2} + B_T \frac{K_B}{[H^+] + K_B} + \frac{K_W}{[H^+]} - [H^+] - \text{Alk}_{\text{CBW}} = 0. \quad (13)$$

15 One may either solve this equation in that rational fraction form with some iterative root-finding method or by one of the ICAC methods described above, or one may transform it into a quintic polynomial equation:

$$P_{\text{CBW}}([H^+]) \equiv [H^+]^5 + q_4[H^+]^4 + q_3[H^+]^3 + q_2[H^+]^2 + q_1[H^+] + q_0 = 0 \quad (14)$$

Solving the alkalinity–pH equation

G. Munhoven

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion





3.4 More complete approximations: rational function based solvers

When additional components in total alkalinity need to be considered besides carbonate, borate and water self-ionization, converting the resulting rational function equation to an equivalent polynomial form becomes more and more tedious and the rational function form becomes the preferred basis for finding the solution. ICAC methods are the only ones that we have encountered so far that could possibly be used to address this problem. However, they bear some potential pitfalls: despite having a solution, the underlying fixed-point equation may be difficult to solve numerically; intermediate estimates of Alk_C may go out of bounds (remember that Alk_C may only take values between 0 and $2C_T$). ICAC methods can therefore not be guaranteed to find the solution.

The only commonly used carbonate chemistry routine that directly solves the rational function form of the equation is that from the Ocean Carbon Cycle Model Intercomparison Project (OCMIP). For the purpose of that project, Orr et al. (2000) prepared standard carbonate speciation routines. Total alkalinity is approximated by $\text{Alk}_T \approx [\text{HCO}_3^-] + 2 \times [\text{CO}_3^{2-}] + [\text{B}(\text{OH})_4^-] + [\text{OH}^-] + [\text{HPO}_4^{2-}] + 2 \times [\text{PO}_4^{3-}] + [\text{H}_3\text{SiO}_4^-] - [\text{H}^+]_f - [\text{HSO}_4^-] - [\text{HF}] - [\text{H}_3\text{PO}_4]$. The different species concentrations were, as above, expressed as a function of the total concentrations of their respective acid systems and of $[\text{H}^+]$. The resulting equation was then solved for pH by a hybrid Newton-bisection method, based upon the `rtsafe` solver from Press et al. (1989). All of the models that participated in OCMIP had to use the provided routines for a set of well defined experiments. A number of models still routinely use these OCMIP routines for their pH calculations. These include some versions of the Bern3D model (Müller et al., 2008) and the NCAR global coupled carbon cycle-climate model CSM1.4-carbon (Doney et al., 2006). As mentioned above, PISCES (Aumont and Bopp, 2006) includes a version of the OCMIP solver trimmed down to Alk_{CBW} only. Other models still offer the OCMIP solvers as an option.

GMDD

6, 2087–2136, 2013

Solving the alkalinity–pH equation

G. Munhoven

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



3.5 Other approaches

Luff et al. (2001) have provided a suite of pH calculation routines mainly meant to be used in reactive transport models, but suitable for general speciation calculations as well. The methods proposed by Luff et al. (2001) solve the complete system of equations that control the chemical equilibria between the individual species considered in the total alkalinity approximation. These are required for grid-based reactive transport models where different species are diffusing at different diffusivities. For common applications in biogeochemical carbon cycle models, this approach is nevertheless unnecessarily complex.

There are still some other fine pH solvers, such as CO2SYS of Lewis and Wallace (1998) and derivatives (spreadsheet versions, MATLAB versions, etc. – see <http://cdiac.ornl.gov/oceans/co2rprt.html> for more information), the MATLAB routines from Zeebe and Wolf-Gladrow (2001) or the SEACARB package for R (Lavigne and Gattuso, 2012). These are, however, generally not suitable for inclusion in global biogeochemical models, as they were developed with special programming environments in mind. Their focus is more on data processing. As their names already suggest, they are mainly aimed at carbonate speciation calculations. They also often offer the possibility to choose any two among pH , $[CO_2]$ (or pCO_2), $[HCO_3^-]$, $[CO_3^{2-}]$, C_T , or Alk_T to calculate all the others.

4 pH -scale considerations

As shortly mentioned above, there are a few subtleties related to pH scales that still need to be clarified. In Eq. (18) above, $[H^+]$ may be expressed on any chosen pH scale (free, total, seawater) as long as all of the stoichiometric constants $K_{A_{[i]1}}, \dots, K_{A_{[i]n_i}}$ for all of the acid systems are expressed on this same pH scale. The same holds for $[H^+]_f$, the free concentration of H^+ , which must also be expressed on (or converted to) this same scale. The concentrations of H^+ on the total and on the sea-water scale are, by

GMDD

6, 2087–2136, 2013

Solving the alkalinity- pH equation

G. Munhoven

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



definition, respectively proportional to the concentration on the free scale (Hansson, 1973; Dickson and Riley, 1979):

$$[\text{H}^+]_{\text{T}} = [\text{H}^+]_{\text{f}} (1 + S_{\text{T}}/K_{\text{HSO}_4}) \quad (15)$$

$$[\text{H}^+]_{\text{SWS}} = [\text{H}^+]_{\text{f}} (1 + S_{\text{T}}/K_{\text{HSO}_4} + F_{\text{T}}/K_{\text{HF}}) \quad (16)$$

where S_{T} and F_{T} are, respectively, the total sulphate and fluoride concentrations in solution, while K_{HSO_4} and K_{HF} are the dissociation constants of HSO_4^- and of HF, respectively, here necessarily expressed on the free pH scale. Accordingly, $[\text{H}^+]_{\text{f}}$ can be expressed as a simple function of the adequate $[\text{H}^+]$ in Eq. (18):

$$[\text{H}^+]_{\text{f}} = \frac{[\text{H}^+]}{s} \quad (17)$$

where $[\text{H}^+] = 10^{-\text{pH}}$, for a chosen pH scale, s is the corresponding scale conversion factor derived from either Eq. (15) or (16). Notice that $s \geq 1$ and remind that s is independent of $[\text{H}^+]$.

Hansson's (1973) original definitions of $[\text{H}^+]_{\text{T}}$ (and of $[\text{H}^+]_{\text{SWS}}$ for seawater containing fluoride) were based upon the total analytical concentration of the hydrogen ion in solution:

$$[\text{H}^+]_{\text{SWS(Hansson)}} = [\text{H}^+]_{\text{f}} + [\text{HSO}_4^-] + [\text{HF}].$$

Differences are negligible in common present-day seawater (Munhoven, 1997). However, since our aim here is to set up a generally valid algorithm, we are not adopting the approximation $[\text{H}^+]_{\text{SWS}} = [\text{H}^+]_{\text{SWS(Hansson)}}$ a priori. Instead, we consider that the effects of HSO_4^- and of HF on total alkalinity are taken into account together with the components of all the other acid systems in the sample, with K_{HSO_4} and K_{HF} being expressed on the same pH scale as the constants for those components.

GMDD

6, 2087–2136, 2013

Solving the alkalinity–pH equation

G. Munhoven

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



5 Development of a universal and robust algorithm

Our ultimate goal here is to develop a universal algorithm to solve the equation

$$R_T([H^+]) \equiv \text{Alk}_{nW}([H^+]) + \frac{K_W}{[H^+]} - \frac{[H^+]}{s} - \text{Alk}_T = 0, \quad (18)$$

where

$$\text{Alk}_{nW}([H^+]) = \sum_i \text{Alk}_{A_{[i]}}([H^+])$$

collects the contributions from all the acid systems to total alkalinity, system by system, proton donors and acceptors combined for each one – except for the contribution that results from the self-ionization of water which we keep explicit in Eq. (18) – and where Alk_T and all the total concentrations $[\Sigma A_{[i]}]$ are given.

We will use a hybrid method that combines the speed of convergence of super-linear and higher-order methods (such as the secant or the Newton–Raphson methods) with the global convergence security of the bisection or the regula falsi method. A similar method is used by the OCMIP carbonate speciation routine. Such methods are standard in root-finding for non-linear equations (e.g. Dowell and Jarrett, 1971; Anderson and Björk, 1973; Bus and Dekker, 1975). They are particularly suitable for our problem with its advantageous mathematical characteristics, the more since that problem also has intrinsic a priori root bracketing, as we will show in the next section. Because of the strict monotonicity of the rational function it is sufficient to make sure that iterates remain within bounds. As long as iterates remain within bounds, they will unconditionally improve either one of the two bounds, thus allowing to tighten the bracketing interval at each step. We can therefore use a high-order numerical root-finding method, such as Newton–Raphson or the secant method as the main iterative scheme. In case the main scheme yields an out-of-bounds iterate at some step, that iterate is rejected and a bisection iterate is used instead. Similarly, if an iteration with the main scheme does

not make the absolute value of the equation decrease faster than expected for a bisection step (i.e. by a factor of two) it is replaced by a bisection step. This helps to prevent cyclic iterations.

A bisection step may temporarily slow down the rate of convergence, but it will secure convergence and again unconditionally improve the bracketing. A regula falsi step could be used instead of bisection; bisection has proven to be more economic though.

5.1 Intrinsic bracketing bounds for the root

Our first aim is now to determine $H_{\text{inf}} > 0$ such that $R_T(H_{\text{inf}}) > 0$ and $H_{\text{sup}} > 0$ such that $R_T(H_{\text{sup}}) < 0$. We have previously established that $\text{Alk}_{\text{nW}}([H^+])$ is a strictly decreasing rational function for $[H^+] > 0$ and that it has the infimum $\text{Alk}_{\text{nWinf}} = -\sum_i m_i [\Sigma A_{[i]}]$. It is therefore sufficient to have H_{inf} such that

$$\frac{K_W}{H_{\text{inf}}} - \frac{H_{\text{inf}}}{s} = \text{Alk}_T - \text{Alk}_{\text{nWinf}}$$

as in this case $R_T(H_{\text{inf}}) = \text{Alk}_{\text{nW}}(H_{\text{inf}}) - \text{Alk}_{\text{nWinf}} > 0$. Equivalently, we require that $H_{\text{inf}}^2 + s(\text{Alk}_T - \text{Alk}_{\text{nWinf}})H_{\text{inf}} - sK_W = 0$, and $H_{\text{inf}} > 0$. This problem has the unique solution

$$H_{\text{inf}} = \frac{-s(\text{Alk}_T - \text{Alk}_{\text{nWinf}}) + \sqrt{\Delta_{\text{inf}}}}{2} \quad (19)$$

where $\Delta_{\text{inf}} = s^2(\text{Alk}_T - \text{Alk}_{\text{nWinf}})^2 + 4sK_W > 0$ is the discriminant of the quadratic.

Similarly, because $\text{Alk}_{\text{nW}}([H^+])$ has the supremum $\text{Alk}_{\text{nWsup}} = \sum_i (n_i - m_i) [\Sigma A_{[i]}]$, it is sufficient to chose H_{sup} such that

$$\frac{K_W}{H_{\text{sup}}} - \frac{H_{\text{sup}}}{s} = \text{Alk}_T - \text{Alk}_{\text{nWsup}}$$

which would lead to $R_T(H_{\text{sup}}) = \text{Alk}_{\text{nW}}(H_{\text{sup}}) - \text{Alk}_{\text{nWsup}} < 0$ as requested. Equivalently, we require that $H_{\text{sup}}^2 + s(\text{Alk}_T - \text{Alk}_{\text{nWsup}})H_{\text{sup}} - sK_W = 0$ and $H_{\text{sup}} > 0$. We finally obtain

$$H_{\text{sup}} = \frac{-s(\text{Alk}_T - \text{Alk}_{\text{nWsup}}) + \sqrt{\Delta_{\text{sup}}}}{2}, \quad (20)$$

where $\Delta_{\text{sup}} = s^2(\text{Alk}_T - \text{Alk}_{\text{nWsup}})^2 + 4sK_W > 0$.

H_{inf} and H_{sup} define a universal bracketing interval for the root of Eq. (18). They only require information that can be directly derived from the nature of the acid systems considered for Alk_T . They can theoretically be used with any numerical scheme to keep iterations bracketed right from the start of the scheme, without any need for manually prescribing them.

5.2 Outline of the algorithm

The proposed algorithm is formally set up in the pH-Alk_T space. There are several advantages for rooting the algorithm in the pH-Alk_T space: (1) the equation's overall appearance is closer to linear in the pH-Alk_T space than in the more commonly used $[\text{H}^+]\text{-Alk}_T$ space; (2) physically meaningless negative $[\text{H}^+]$ values cannot be produced by the iterative scheme; this is not warranted with methods that are rooted in the $[\text{H}^+]\text{-Alk}_T$ space. There is nevertheless also a potential disadvantage: passing between the two spaces a priori requires costly power and logarithm evaluations at each step. As shown below, these operations can, however, be avoided to a large extent by transposing the actual calculations into the $[\text{H}^+]\text{-Alk}_T$ space and carrying them out there.

The algorithm comes in two variants: one based upon the Newton–Raphson and bisection methods, and one that is based upon the secant and bisection methods. We will first describe the Newton–Raphson/bisection variant.

Let $R = R(H)$ denote the rational function chosen to approximate Alk_T . Before starting we determine the intrinsic lower and upper bounds H_{inf} and H_{sup} , and constrain the initial value H_0 to be within bounds.

Solving the alkalinity–pH equation

G. Munhoven

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Solving the
alkalinity–pH
equation

G. Munhoven

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Then, at each step $k + 1, k = 0, \dots$:

1. Prepare to carry out a Newton–Raphson step where $pH_{k+1} = pH_k + \Delta pH$, with $\Delta pH = -R|_{pH_k} / (dR/dpH)|_{pH_k}$: $(dR/dpH)|_{pH_k}$ can be calculated from $R(H_k)$ and $dR/dH|_{H_k}$, noticing that $(dR/dpH)|_{pH_k} = -(dR/dH)|_{H_k} \times H_k \times \ln(10)$.
2. Adapt the bracketing interval: if $R|_{pH_k} > 0$ then adjust $H_{inf} := H_k$, if $R|_{pH_k} < 0$ then adjust $H_{sup} := H_k$.
3. Require $|R(H_k)|$ to decrease faster than one would typically expect from bisection under the same conditions: compare it with $\min(|R(H_j)|, \forall j < k)$ and if greater than half that value (bisection halves the bracketing interval at each step and is linearly convergent), do not complete the Newton–Raphson step, but adopt a bisection iterate between the current pH_{inf} and pH_{sup} (updated just before) and return to stage 1 for the next step.
4. Provisionally set $H_{k+1} = 10^{-pH_{k+1}} = H_k \times \exp(-R(H_k)/(H_k dR/dH|_{H_k}))$ to complete the Newton–Raphson step.
5. Constrain H_{k+1} to remain within the current bracketing interval: if $H_{k+1} > H_{sup}$ or $H_{k+1} < H_{inf}$, reject the Newton–Raphson iterate, replace it by the bisection iterate as in stage 4 and return to stage 1 for the next step.
6. Stop the iterations if either the maximum permissible number of iterations is exceeded or if $|(H_{k+1} - H_k)/H_k| < \epsilon$ (ϵ being a pre-set tolerance), else return to stage 1 for another step.

At most one exponential has to be evaluated per step (at stage 4). This is, computationally speaking, the most expensive operation in each step. It can, however, often be avoided: when $|R_T(H_k)/(H_k dR/dH|_{H_k})| \ll 1$, then $H_k \times \exp(-R_T(H_k)/(H_k dR/dH|_{H_k})) \simeq H_k - R_T(H_k)/(dR/dH|_{H_k})$ and the iterate can be assimilated to a plain $[H^+]$ -Alk_T space Newton–Raphson iterate. Once the argument of the exponential becomes sufficiently small (a threshold value of 1 in absolute value has proven efficient) we may

Solving the
alkalinity–pH
equation

G. Munhoven

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏮

⏭

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



switch to the linear approximation, thereby saving the exponential operation. The relative error $|(H_{k+1} - H_k)/H_k|$ from the stopping criterion can be approximated by $|R(H_k)/(H_k \, dR/dH|_{H_k})|$, i.e. we can reuse the argument of the exponential above (no extra operations required). At any stage, bisection between pH_{inf} and pH_{sup} translates

to calculating H_{k+1} as the geometric mean of H_{inf} and H_{sup} : $H_{k+1} = \sqrt{H_{\text{inf}} H_{\text{sup}}}$. By construction, any accepted iterate will thus be strictly between the current H_{inf} and H_{sup} , and because of the strictly decreasing nature of $R(H)$ will always lead to contribute to improve either the lower or the upper bound.

In a variant of the above we replace the Newton–Raphson scheme by a secant scheme. However, rooting a secant scheme in the $pH\text{-Alk}_T$ space while carrying out operations in the $[H^+]\text{-Alk}_T$ space will require non-integer power operations at each step which are even more costly than exponentials. It is therefore preferable to completely root the calculations in the $[H^+]\text{-Alk}_T$ space with the secant method, despite the potential trade-offs for the convergence efficiency. Secant iterations have the advantage of requiring only one evaluation of the equation at each iteration; in addition to the equation evaluation, Newton–Raphson iterations also require the evaluation of the derivative of the equation. The cheaper iterations of the secant method possibly outweigh its lower order of convergence, which is ≈ 1.62 , compared to 2 for the Newton–Raphson method.

5.3 Discussion: comparison with the OCMIP solver

A similar technique was also used in the `drtsafe` routine in the OCMIP suite (Orr et al., 2000), which is fundamentally the `rtsafe` routine of Press et al. (1989) with some essential error trapping removed. That routine also combines the global convergence properties of the bisection with the speed of convergence of the Newton–Raphson method.

The algorithm presented here differs from that used in `drtsafe` in several significant ways. (1) `drtsafe` iterations are rooted in the $[H^+]\text{-Alk}_T$ space. (2) `drtsafe` requires

Solving the
alkalinity–pH
equation

G. Munhoven

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



brackets to be explicitly provided at the subroutine call. In case these are inappropriate (e.g. no sign change of the equation function over the defined bracketing interval), it would simply iterate to the maximum number of iterations allowed because of the absence of validity checks and return some meaningless result (in general one of the two bounds provided). (3) `drtsafe` always starts its iterations from the midpoint of the provided bracketing interval. It is thus critically dependent on a valid interval (no validity checks performed though) and, because of the rooting in the $[H^+]$ - Alk_T space, on a tight bracketing interval for efficiency. The algorithms proposed here only use the bracketing values to secure convergence in case Newton–Raphson iterates are not decreasing fast enough or would go out of bounds and they use an independent initial value. Because we root our iterations in the pH - Alk_T space, even bisection steps may accommodate $[H^+]$ changes over several orders of magnitude during initial steps in case a far off starting value is used.

6 Sample implementation of the new algorithms in Fortran 90

A sample implementation of the algorithms realized in standard Fortran 90 is provided in the Supplement to this paper. Together with the drivers that were used to carry out the experiments described below they make up the SOLVER Suite for Alkalinity-PH Equations (SOLVESAPHE). Parts of the code contain C-preprocessor directives for enabling or disabling specific parts (debugging messages, optional code parts, ...), and to select among the cases treated below. After pre-processing, the source files are strictly standard conforming Fortran 90. The codes are made available under the GNU Lesser Public Licence, version 3.

A complete user manual that covers the technical details of SOLVESAPHE is included in the archive provided in the Supplement. Here we only give a short overview of the two central modules.

6.1 Summary description

The module `mod_chemconst` provides parametric expressions for the stoichiometric constants of the acid systems taken into account (carbonates, borates, hydrogen sulphate, sulphides, phosphates, ...). The module also hosts the Π_j values (Eq. 4) for the various acid systems.

The module `mod_phsolvers` provides six different solvers:

1. the function `solve_at_general`: the new algorithm described above;
2. the function `solve_at_icacfp`: a fixed-point only ICAC method;
3. the function `solve_at_bacastow`: Bacastow's method, an ICAC method with secant iterations (with secant iterations either on $[H^+]$ or on its scaled inverse);
4. the function `solve_at_general_sec`: the variant of `solve_at_general` that uses secant instead of Newton–Raphson iterations;
5. the function `solve_at_ocmip`: a re-implementation of the OCMIP solver with Newton–Raphson/bisection iterations, completed with a bare minimum of error trapping and fitted with the optional initialisation scheme common to all of the solvers (the latter was only adapted to use an interval of ± 0.5 pH unit interval around an optional initial value to emulate the recommended OCMIP setup after startup);
6. the function `solve_at_fast`: a simplified version of `solve_at_general` without the bracketing control (may not always converge).

Each one of the six solvers takes into account all of the constituents that explicitly appear in Eq. (1), except for S^{2-} whose concentration is negligible even at high pH values. `mod_phsolvers_logging.F90` is a special version of `mod_phsolvers.F90` that includes extra bookkeeping regarding the number of iterations required for convergence, the numbers of bisection iterations due to limiting, the initial values

GMDD

6, 2087–2136, 2013

Solving the alkalinity–pH equation

G. Munhoven

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



adopted, the initial bracketing values (if relevant), the intermediate iterates, etc. `mod.phsolvers.logging.F90` does not include `solve_at_fast` though. For more technical details, please refer to the manual that goes with the source codes.

The modules `mod.acb_solvers` and `mod.acbw_solvers` provide simpler and more streamlined solvers, based upon the Alk_{CB} and Alk_{CBW} approximations, respectively.

6.2 Test case definitions

The pH_{SWS} scale was adopted for all of the calculations below. With each method, iterations were stopped once the relative change of an iterate compared to its predecessor felt below 10^{-8} ; the maximum number of iterations was set to 50. For all of the three cases, we adopt a temperature of 275.15 K, a salinity of 35 and an applied pressure of 0 bar. Additional results for a temperature of 298.15 K or a pressure of 300 bar can be found in the Supplement.

The convergence properties for each one of the pH solvers are explored for three different (nested) subsets of the $C_{\text{T}}\text{-Alk}_{\text{T}}$ space:

SW1 – for C_{T} ranging between $1.85 \text{ mmol kg}^{-1}$ and $2.45 \text{ mmol kg}^{-1}$, and Alk_{T} between 2.20 meq kg^{-1} and 2.50 meq kg^{-1} , on a regular 600×300 cell centred grid;

SW2 – for C_{T} ranging between $1.85 \text{ mmol kg}^{-1}$ and $3.35 \text{ mmol kg}^{-1}$, and Alk_{T} between 2.20 meq kg^{-1} and 3.50 meq kg^{-1} , on a regular 1500×1300 cell centred grid;³

SW3 – for C_{T} ranging between 0 mmol kg^{-1} and 6 mmol kg^{-1} , and Alk_{T} between -1 meq kg^{-1} and 5 meq kg^{-1} , on a regular 600×600 cell centred grid.

The other concentrations are set as follows: $P_{\text{T}} = 0.5 \mu\text{mol kg}^{-1}$, $\text{Si}_{\text{T}} = 5 \mu\text{mol kg}^{-1}$, $[\text{NH}_4]_{\text{T}} = 0 \mu\text{mol kg}^{-1}$ and $[\text{H}_2\text{S}]_{\text{T}} = 0 \mu\text{mol kg}^{-1}$. With each test case, three different

³For the graphs shown below, a 150×130 cell centred grid was used.

Solving the
alkalinity–*pH*
equation

G. Munhoven

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

I◀

▶I

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



schemes are considered to start the iterations: (cub) starting values are derived from the scheme designed for the cubic polynomial in Sect. 3.2.2; (pH8) a constant starting value $[H^+] = 10^{-8} \text{ mol kg}^{-1}$ is used, except for `solve_at_ocrmip`, for which the recommended “cold-start” brackets corresponding to $pH = 6$ and $pH = 9$ are used; (safe) the midpoint of the pH interval defined by the intrinsic brackets H_{inf} and H_{sup} (from Sect. 5.1) is used as a starting value, except for `solve_at_ocrmip`, for which H_{inf} and H_{sup} are used as initial brackets. Timing information is based upon `driver_at_general.F90`, all other information (numbers of iterations, of divergences, errors, ...) upon `driver_at_logging.F90`.

SW1 covers the typical range of present-day seawater samples. Every solver should be able to determine the root of the equation without a single failure. SW2 covers the expected range of sea-water samples under the S750 stabilisation scenario over the next 50 000 yr (derived from simulation experiments with the coupled carbon cycle-sediment model MBM-MEDUSA (Munhoven, 2007, 2009)). SW3 is of more theoretical nature and is meant to analyse the performance of the solvers with extremely low alkalinity or C_T values. It will nevertheless also provide important information about the convergence domains of `solve_at_icacfp` and `solve_at_bacastow`, as we will see below.

6.3 Results

We here only present results obtained with the solvers from `mod_phsolvers` (for the timings) and `mod_phsolvers_logging`. To simplify the presentation, we leave out the prefix “`solve_at_`” when referring to the different solver functions below. The testing platform had a Debian 6.0.6 operating system (32-bit kernel 2.6.32-5-686-bigmem) running on a 2.53 GHz Intel Core2Duo T9400 CPU; all of the source codes were compiled with gfortran 4.4.5, without any optimisation flags set.

6.3.1 Comparison of the six solvers

Figure 1 shows the distributions of pH for test cases SW1, SW2 and SW3, as calculated by the new algorithm with Newton–Raphson iterations. Also shown is the equation residual for SW3 (which encompasses the two others). Residual values smaller than $10^{-21} \text{ mol kg}^{-1}$ in absolute value are shown as $10^{-21} \text{ mol kg}^{-1}$. The residual is at least five orders of magnitude smaller than the actual H^+ concentrations, emphasizing that convergence was significantly reached.

Execution times for the SW1, SW2 and SW3 test series are reported in Table 1. The times for each of the three test series have been normalised to the execution time of the `general_sec` routine with cubic initialisation (shown in italics). `general_sec` was the fastest of the routines that successfully passed the complete test series.

For test cases SW1 and SW2, Bacastow’s method is clearly the fastest. It is about 20 % faster than the `general` and `general_sec` routines developed here, and twice or two and a half times as fast as its closest relative `icacfp`. `general_sec` is generally about 5–10 % faster than `general`. The results obtained with `fast` indicate that the overhead required by the safeguard bracketing requires about 5 to 10 % extra computing time, if everything works fine. However, the comparison of the SW2-pH8 results for `general` and `fast` shows that replacing unacceptable Newton–Raphson steps by safe but inherently slower bisection steps may overall even lead to a gain of time in more critical situations. Neither `fast`, nor `bacastow`, nor `icacfp` were able to complete test case SW3; `ocmip` was furthermore not able to complete the SW2-pH8 test, because of invalid initial bracketing over parts of the domain. Convergence failures with `ocmip` can be avoided if we use the intrinsic bracketing bounds obtained in Sect. 5.1, as can be seen from the ‘safe’ initialisation procedure. However, with this safe initialisation procedure, the execution for `ocmip` exceed those of `general_sec` with the cubic initialisation by a factor of 5.6–5.7 in test case SW1 and SW2, and by a factor of 3.7 for case SW3. As can be seen on Fig. 1, SW3 includes a large number of $C_T\text{-Alk}_T$

GMDD

6, 2087–2136, 2013

Solving the alkalinity–pH equation

G. Munhoven

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



combinations that lead to extreme pH values (either lower than 4 or higher than 10), where the intrinsic bounds are comparatively restrictive.

In each test, and with every method used, the initialisation procedure developed above for the cubic polynomial leads to 30–60 % shorter execution times than the constant initialisation (“pH8”) which may even lead to divergence (e.g. `ocmip`).

The reasons for the strong performance loss of `icacfp` in SW1 become obvious from Fig. 2. That figure shows the number of iterations required to trigger the stopping criterion for the SW2 test. The SW1 domain is included at the lower left of the SW2 domain: it ends at $C_T = 2.45 \text{ mmol kg}^{-1}$ and $\text{Alk}_T = 2.50 \text{ meq kg}^{-1}$. In that area, `general` and `bacastow` require at most four iterations, `ocmip` generally six or seven, but `icacfp` often fifteen and more.

6.3.2 Shortcomings of ICAC methods

As we have seen, ICAC methods have divergence problems on the SW3 grid. These problems are inherent to the method and can only be alleviated to a limited extent. It should be noticed that the fixed-point equation $H = Q(H)$ (see Eq. 10) has a solution, i.e. that $Q(H)$ has a fixed-point, for any $C_T - \text{Alk}_T$ pair, since the total alkalinity– pH equation has a solution.

The divergence pattern of the ICAC methods can easily be explained from the derivative of the underlying function Q with respect to H , shown for SW3 on Fig. 3. The values were calculated from the H^+ concentrations shown on Fig. 1. The derivative values are negative over the whole domain and fixed-point iterations thus oscillate around the solution (i.e. the fixed-point of Q). Fixed-point iterations can only converge to the fixed point of Q where the derivative is strictly smaller than 1 in absolute value, i.e. is strictly greater than -1 here. The thick white line indicates where the derivative of Q is equal to -1 . The white areas on the other graphs indicate where the methods did not converge. The white areas for `icacfp` clearly match the areas where the derivative of Q is lower than -1 . They are even somewhat larger.

Solving the alkalinity– pH equation

G. Munhoven

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Solving the
alkalinity–pH
equation

G. Munhoven

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



When the derivative of Q are just slightly greater than -1 , iterations may become “operationally divergent”: the pre-set maximum number of iterations is insufficient to meet the stopping criterion as the generated suite converges too slowly there. Bacastow’s method, on the other hand, has a slightly larger convergence domain than delimited by the thick white line in the graph of the derivative. The fixed-point equation can indeed be solved by the secant method in some instances where straight fixed-point iterations would produce slowly divergent iteration suites. As the derivative of Q is negative, fixed-point iterations oscillate around the root, as long as they can be evaluated, i.e. as long as the Alk_C estimates obtained from Alk_T with the H iterates range between 0 and $2C_T$. If they can be successfully calculated, the two first iterates used to initialize the secant iterations in Bacastow’s method thus bracket the root and we may expect that the first application of the secant method provides an excellent estimate for the root, even if the two first iterates would generate a fixed-point suite that slowly diverges. This is especially obvious inside the white bulge on Fig. 3c at low C_T values and Alk_T values greater than C_T .

6.3.3 Quality of the cubic equation based initialisation

As shown above, the initialisation scheme especially designed for the iterative solution of the cubic Eq. (12) by the Newton–Raphson method is highly attractive even for more complex approximations to total alkalinity than Alk_{CB} . This is quantified on Fig. 4, exemplified by SW2 results. The relative error of H_0 determined as outlined in Sect. 3.2.2 on the actual H^+ concentration (as calculated with `general`) is less than 7 % over the SW2 domain. In comparison, over that same domain, the approximation $\text{Alk}_C \simeq \text{Alk}_T$ and usage of Eq. (7) gives rise to errors that are about ten times as large.

7 Conclusions

We have explored the mathematical properties of the total alkalinity– pH equation, i.e. the equation that relates $[H^+]$ (or equivalently pH) to total alkalinity and the total concentrations of all the acid systems contributing to total alkalinity. We have demonstrated that the rational function expression of that equation is strictly monotone. If water self-ionization is considered, the total alkalinity– pH equation has one and only one positive root, for any given value of total alkalinity, and for any given non negative values for the total concentrations of the acid system components of total alkalinity. All other roots of the equation are either negative or complex with non-zero imaginary parts. We have shown that there are intrinsic upper and lower bounds for the positive root of the equation that only depend on information from the list of included acid systems. These seemingly straightforward mathematical properties have apparently not been published before and currently available solvers do not take advantage of them. They actually enabled us to design a universal and fail-safe algorithm to solve the total alkalinity– pH equation. We propose two variants, one using the Newton–Raphson, the other the secant scheme.

The performances of the two algorithms (plus one simplified version without safeguarded iterations) were compared with some common existing ones: (1) the fixed-point Iterative Carbonate Alkalinity Correction Method (ICAC), a classical method recently made popular again by Follows et al. (2006); (2) the method of Bacastow (1981), which is a variant of the previous one using secant instead of fixed-point iterations and (3) the OCMIP-2 standard protocol routines (Orr et al., 2000), re-implemented here. Source code with a reference implementation of the six algorithms discussed in the text is provided in the SOLver Suite for Alkalinity-PH Equations (SOLVE_{SAPHE}) in the Supplement for use under the GNU Lesser General Public Licence Version 3 (LGPLv3) license.

We have defined three test cases for a comparative analysis of the six methods: one for the typical open-ocean concentrations of total dissolved inorganic carbon and

GMDD

6, 2087–2136, 2013

Solving the alkalinity– pH equation

G. Munhoven

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Solving the
alkalinity– pH
equation

G. Munhoven

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



total alkalinity of the present-day ocean; another one covering the expected future distributions of these concentrations under progressing ocean acidification and subsequent dissolution of deep-sea surface sedimentary carbonates; a third one covering extremely low concentrations of dissolved inorganic carbon and total alkalinity, and even negative values for total alkalinity. Different approaches for starting iterative root-finding methods have been tested as well for their efficiency.

The two new algorithms are the only ones that successfully complete all of the tests. The same convergence security could be achieved with the OCMIP solver as well after a modification of its initialisation scheme, at the price of much longer execution times though (typically by a factor of three to six). Bacastow's method is the fastest of all the tested general methods overall in the common regions of convergence. The two new algorithms are only 10–20 % slower than Bacastow's method and more than 50 % faster than the next best performant ones. The secant variant of our algorithm is about 5–10 % faster than the Newton–Raphson variant. We have developed an original starting scheme for solving the cubic polynomial equation that is to be solved to determine pH from carbonate and borate alkalinity alone. That starting scheme can easily be completed for usage with general total alkalinity– pH equation solvers and we show here that it typically allows to save 30–60 % of calculation time compared to the standard $pH = 8$ initialisation.

The two proposed algorithms are furthermore extremely robust. As documented in the Supplement, the sample implementation has been successfully used with random values (covering up to six orders of magnitude) for the total concentrations of the acid system components to total alkalinity and of total alkalinity itself, with random pH starting values between 1 and 14 and still ensured convergence in 100 % of the cases.

The two proposed new algorithms thus offer almost convergence security over an extremely wide range of total concentrations for the contributions of the various acid systems to total alkalinity, at a marginal additional computational cost only.

Appendix A

Alkalinity components are strictly decreasing with $[H^+]$: demonstration

We are now going to show that

$$\frac{dAlk_A}{d[H^+]} < 0,$$

- 5 for any given acid system $H_nA - H_{n-1}A^- - \dots - A^{n-}$, i.e. for constant $[\Sigma A]$, at equilibrium.
For notational convenience, we write

$$Alk_A = [\Sigma A] \left(\frac{D_1}{D} - m \right),$$

with

$$D = \sum_{j=0}^n \Pi_j [H^+]^{n-j} \quad \text{and} \quad D_1 = \sum_{j=0}^n j \Pi_j [H^+]^{n-j}.$$

- 10 Since $\Pi_j > 0$ for any $j \geq 0$, we know that $D > 0$ and $D_1 > 0$. We may then write

$$\frac{dAlk_A}{d[H^+]} = [\Sigma A] \frac{d}{d[H^+]} \left(\frac{D_1}{D} \right) = [\Sigma A] \frac{D \frac{dD_1}{d[H^+]} - D_1 \frac{dD}{d[H^+]}}{D^2}.$$

It is straightforward to show that

$$\frac{dD}{d[H^+]} = \frac{1}{[H^+]} (nD - D_1) \quad \text{and} \quad \frac{dD_1}{d[H^+]} = \frac{1}{[H^+]} (nD_1 - D_2),$$

where we have further defined

$$15 \quad D_2 = \sum_{j=0}^n j^2 \Pi_j [H^+]^{n-j},$$

Solving the alkalinity–pH equation

G. Munhoven

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



which is positive, similarly to D and D_1 . We hence find that

$$\frac{d\text{Alk}_A}{d[\text{H}^+]} = [\Sigma A] \frac{D(nD_1 - D_2) - D_1(nD - D_1)}{[\text{H}^+]D^2} = -[\Sigma A] \frac{DD_2 - D_1^2}{[\text{H}^+]D^2}.$$

The result now follows from Lagrange's identity:

$$\left(\sum_{j=0}^n x_j^2 \right) \left(\sum_{j=0}^n y_j^2 \right) - \left(\sum_{j=0}^n x_j y_j \right)^2 = \frac{1}{2} \sum_{j=0}^n \sum_{i=0}^n (x_i y_j - x_j y_i)^2.$$

5 With

$$x_j = \sqrt{\prod_i [\text{H}^+]^{n-j}}$$

$$y_j = j \sqrt{\prod_i [\text{H}^+]^{n-j}},$$

we have

10
$$\sum_{j=0}^n x_j^2 = D, \quad \sum_{j=0}^n y_j^2 = D_2 \quad \text{and} \quad \sum_{j=0}^n x_j y_j = D_1.$$

To conclude, it is then sufficient to notice that

$$\sum_{j=0}^n \sum_{i=0}^n (x_i y_j - x_j y_i)^2 = \sum_{j=0}^n \sum_{i=0}^n \prod_i \prod_j (i - j)^2 [\text{H}^+]^{2n-i-j}$$

which is strictly positive if $n > 0$. Alternatively, the result also follows from the Cauchy-Schwarz inequality in $n + 1$ dimensions, noticing that the conditions for equality are not

15 met.
Accordingly, $\text{Alk}_A([\text{H}^+])$ is strictly decreasing as a function of $[\text{H}^+]$.

Solving the alkalinity–pH equation

G. Munhoven

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏮

⏭

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Supplementary material related to this article is available online at:
[http://www.geosci-model-dev-discuss.net/6/2087/2013/
gmdd-6-2087-2013-supplement.zip](http://www.geosci-model-dev-discuss.net/6/2087/2013/gmdd-6-2087-2013-supplement.zip).

Acknowledgements. Guy Munhoven is a Research Associate with the Belgian Fund for Scientific Research – FNRS.

References

- Anderson, N. and Björk, A.: A new high order method of regula falsi type for computing a root of an equation, BIT, 13, 253–264, 1973. 2111
- Antoine, D. and Morel, A.: Modelling the seasonal course of the upper ocean $p\text{CO}_2$ (I). Development of a one-dimensional model, Tellus B, 47, 103–121, doi:10.1034/j.1600-0889.47.issue1.11.x, 1995. 2101, 2103
- Arndt, S., Regnier, P., Goddérès, Y., and Donnadieu, Y.: GEOCLIM *reloaded* (v 1.0): a new coupled earth system model for past climate change, Geosci. Model Dev., 4, 451–481, doi:10.5194/gmd-4-451-2011, 2011. 2103
- Aumont, O. and Bopp, L.: Globalizing results from ocean in situ iron fertilization studies, Global Biogeochem. Cy., 20, GB2017, doi:10.1029/2005GB002591, 2006. 2107, 2108
- Bacastow, R.: Numerical evaluation of the evasion factor, in: Carbon Cycle Modelling, vol. 16 of SCOPE, chap. 3.4, edited by: Bolin, B., John Wiley & Sons, Chichester, NY, 95–98, 1981. 2100, 2103, 2104, 2123
- Bacastow, R. and Keeling, C. D.: Atmospheric carbon dioxide and radiocarbon in the natural carbon cycle. II. Changes from A.D. 1700 to 2070 as deduced from a geochemical model, in: Carbon and the Biosphere, Proceedings of the 24th Brookhaven Symposium in Biology, edited by: Woodwell, G. M. and Pecan, E. V., US Atomic Energy Commission, 86–135, 16–18 May 1972, Upton, NY, 1973. 2107
- Bolin, B., Björkström, A., Holmén, K., and Moore, B.: The simultaneous use of tracers for ocean circulation studies, Tellus B, 35, 206–236, 1983. 2107
- Broecker, W. S. and Peng, T.-H.: Tracers in the Sea, Lamont-Doherty Geological Observatory of Columbia University, Palisades, NY 10964, 1982. 2088, 2099

GMDD

6, 2087–2136, 2013

Solving the
alkalinity–pH
equation

G. Munhoven

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Solving the
alkalinity–pH
equation

G. Munhoven

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Bus, J. C. P. and Dekker, T. J.: Two efficient algorithms with guaranteed convergence for finding a zero of a function, *ACM T. Math. Software*, 1, 330–345, 1975. 2111
- Caldeira, K. and Wickett, M. E.: Oceanography: anthropogenic carbon and ocean pH, *Nature*, 425, 365, doi:10.1038/425365a, 2003. 2089
- 5 Dickson, A. G.: An exact definition of total alkalinity and a procedure for the estimation of alkalinity and total inorganic carbon from titration data, *Deep-Sea Res.*, 28A, 609–623, 1981. 2092
- Dickson, A. G.: pH scales and proton-transfer reactions in saline media such as sea water, *Geochim. Cosmochim. Ac.*, 48, 2299–2308, 1984. 2091
- 10 Dickson, A. G. and Riley, J. P.: The estimation of acid dissolution constants in seawater media from potentiometric titrations with strong base. I. The ionic product of water – K_w , *Mar. Chem.*, 7, 89–99, 1979. 2110
- Doney, S. C., Lindsay, K., Fung, I., and John, J.: Natural variability in a stable, 1000-yr global coupled climate-carbon cycle simulation, *J. Climate*, 19, 3033–3054, doi:10.1175/JCLI3783.1, 2006. 2108
- 15 Dowell, M. and Jarrett, P.: A modified regula falsi method for computing the root of an equation, *BIT*, 11, 168–174, 1971. 2111
- Dyrssen, D. W.: Framvaren and the Black Sea – similarities and differences, *Aquat. Geochem.*, 5, 59–73, doi:10.1023/A:1009663704604, 1999. 2095
- 20 Follows, M. J., Ito, T., and Dutkiewicz, S.: On the solution of the carbonate chemistry system in ocean biogeochemistry models, *Ocean Model.*, 12, 290–301, doi:10.1016/j.ocemod.2005.05.004, 2006. 2100, 2102, 2103, 2123
- Gangstø, R., Joos, F., and Gehlen, M.: Sensitivity of pelagic calcification to ocean acidification, *Biogeosciences*, 8, 433–458, doi:10.5194/bg-8-433-2011, 2011. 2107
- 25 Goosse, H., Brovkin, V., Fichefet, T., Haarsma, R., Huybrechts, P., Jongma, J., Mouchet, A., Selten, F., Barriat, P.-Y., Campin, J.-M., Deleersnijder, E., Driesschaert, E., Goelzer, H., Janssens, I., Loutre, M.-F., Morales Maqueda, M. A., Opsteegh, T., Mathieu, P.-P., Munhoven, G., Pettersson, E. J., Renssen, H., Roche, D. M., Schaeffer, M., Tartinville, B., Timmermann, A., and Weber, S. L.: Description of the Earth system model of intermediate complexity LOVECLIM version 1.2, *Geosci. Model Dev.*, 3, 603–633, doi:10.5194/gmd-3-603-2010, 2010. 2103, 2107
- 30 Hansson, I.: A new set of pH-scales and standard buffers for sea water, *Deep-Sea Res.*, 20, 479–491, 1973. 2110

Solving the
alkalinity–pH
equation

G. Munhoven

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Heinze, C., Maier-Reimer, E., and Winn, K.: Glacial $p\text{CO}_2$ reduction by the World Ocean: experiments with the Hamburg Carbon Cycle Model, *Paleoceanography*, 6, 395–430, 1991. 2107

Hoffert, M. I., Wey, Y.-C., Callegari, A. J., and Broecker, W. S.: Atmospheric response to deep-sea injections of fossil-fuel carbon dioxide, *Climatic Change*, 2, 53–68, doi:10.1007/BF00138226, 1979. 2107

Keeling, C. D.: The carbon dioxide cycle: reservoir models to depict the exchange of atmospheric carbon dioxide with the oceans and land plants, in: *Chemistry of the Lower Atmosphere*, chap. 6, edited by: Rasool, S. I., Plenum Press, New York, NY, 251–329, 1973. 2107

Kirby, C. S. and Cravotta III, C. A.: Net alkalinity and net acidity. 1: Theoretical considerations, *Appl. Geochem.*, 20, 1920–1940, doi:10.1016/j.apgeochem.2005.07.002, 2005. 2092

Lavigne, H. and Gattuso, J.-P.: Seacarb: Seawater Carbonate Chemistry with R, R Package Version 2.4., available at: <http://cran.r-project.org/web/packages/seacarb/index.html>, last access: 13 December 2012, 2012. 2109

Le Hir, G., Donnadieu, Y., Yves Godd  ris, Y., Pierrehumbert, R. T., Halverson, G. T., Macouin, M., N  d  lec, A., and Ramstein, G.: The snowball Earth aftermath: exploring the limits of continental weathering processes, *Earth Planet. Sc. Lett.*, 277, 453–463, doi:10.1016/j.epsl.2008.11.010, 2009. 2089

Lewis, E. and Wallace, D.: Program Developed for CO_2 System Calculations, Tech. Rep. 105, Carbon Dioxide Analysis Center, Oak Ridge National Laboratory, Oak Ridge (TN), 1998. 2109

Luff, R., Haeckel, M., and Wallmann, K.: Robust and fast FORTRAN and MATLAB libraries to calculate pH distributions in marine systems, *Comput. Geosci.*, 27, 157–169, 2001. 2109

Maier-Reimer, E.: Geochemical cycles in an Ocean General Circulation Model. Preindustrial tracer distributions, *Global Biogeochem. Cy.*, 7, 645–677, 1993. 2107

Maier-Reimer, E. and Hasselmann, K.: Transport and storage of CO_2 in the ocean – an inorganic ocean-circulation carbon cycle model, *Clim. Dynam.*, 2, 63–90, 1987. 2107

Maier-Reimer, E., Kriest, I., Segschneider, J., and Wetzol, P.: The HAMBURG Ocean Carbon Cycle Model HAMOCC 5.1 – Technical Description Release 1.1, *Berichte zur Erdsystemforschung*, Reports on Earth System Science 14, Max-Planck-Institut f  r Meteorologie, Hamburg (DE), 2005. 2107

Solving the
alkalinity–pH
equation

G. Munhoven

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Marchal, O., Stocker, T. F., and Joos, F.: A latitude-depth, circulation-biogeochemical ocean model for paleoclimate studies. Development and sensitivities, *Tellus B*, 50, 290–316, doi:10.1034/j.1600-0889.1998.t01-2-00006.x, 1998. 2104

5 Millero, F. J. and Sohn, M. L.: Chemical Oceanography, CRC Press, Boca Raton, Florida, 531 pp., 1992. 2101

Millero, F. J., Woosley, R., DiTrollo, B., and Waters, J.: Effect of ocean acidification on the speciation of metals in seawater, *Oceanography*, 22, 72–85, doi:10.5670/oceanog.2009.98, 2009. 2089

10 Müller, S. A., Joos, F., Plattner, G.-K., Edwards, N. R., and Stocker, T. F.: Modeled natural and excess radiocarbon: sensitivities to the gas exchange formulation and ocean transport strength, *Global Biogeochem. Cy.*, 22, GB3011, doi:10.1029/2007GB003065, 2008. 2108

Munhoven, G.: Modelling Glacial-Interglacial Atmospheric CO₂ Variations: The Role of Continental Weathering, Ph.D. thesis, Université de Liège, Liège, available at: <http://www.astro.ulg.ac.be/~munhoven/en/PhDThesis.pdf> (last access: 21 February 2013), 1997. 2100, 2104, 2110

15 Munhoven, G.: Glacial-interglacial rain ratio changes: implications for atmospheric CO₂ and ocean-sediment interaction, *Deep-Sea Res. Pt. II*, 54, 722–746, doi:10.1016/j.dsr2.2007.01.008, 2007. 2104, 2119

Munhoven, G.: Future CCD and CSH variations: deep-sea impact of ocean acidification, *Geochim. Cosmochim. Ac.*, 73, p. A917, 2009. 2119

20 Munhoven, G. and François, L. M.: Glacial-interglacial variability of atmospheric CO₂ due to changing continental silicate rock weathering: a model study, *J. Geophys. Res.*, 101, 21423–21437, doi:10.1029/96JD01842, 1996. 2104

Opdyke, B. N. and Walker, J. C. G.: Return of the coral reef hypothesis: basin to shelf partitioning of CaCO₃ and its effect on atmospheric CO₂, *Geology*, 20, 733–736, 1992. 2099

25 Orr, J., Najjar, R., Sabine, C., and Joos, F.: Abiotic-HOWTO, available at: <http://ocmip5.ipsl.jussieu.fr/OCMIP/phase2/simulations/Abiotic/HOWTO-Abiotic.html>, (last access: 20 July 2012), 2000. 2108, 2115, 2123

Palmer, J. R. and Totterdell, I. J.: Production and export in a global ocean ecosystem model, *Deep-Sea Res. Pt. I*, 48, 1169–1198, doi:10.1016/S0967-0637(00)00080-7, 2001. 2104, 2107

30 Park, P. K.: Oceanic CO₂ system: an evaluation of ten methods of investigation, *Limnol. Oceanogr.*, 14, 179–186, 1969. 2101

Solving the
alkalinity–pH
equation

G. Munhoven

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Peng, T.-H., Takahashi, T., Broecker, W. S., and Olafsson, J.: Seasonal variability of carbon dioxide, nutrients and oxygen in the northern North Atlantic surface water: observations and a model, *Tellus B*, 39, 439–458, 1987. 2100, 2103

Press, W. H., Flannery, B. P., Teukolsky, S. A., and Vetterling, W. T.: *Numerical Recipes (FORTRAN Version)*, Cambridge University Press, Cambridge, 1989. 2108, 2115

Ridgwell, A. J.: *Glacial-Interglacial Perturbations of the Global Carbon Cycle*, Ph.D. thesis, University of East Anglia, Norwich, 2001. 2101, 2103

Ridgwell, A. and Schmidt, D. N.: Past constraints on the vulnerability of marine calcifiers to massive carbon dioxide release, *Nat. Geosci.*, 3, 196–200, doi:10.1038/ngeo755, 2010. 2089

Ridgwell, A., Hargreaves, J. C., Edwards, N. R., Annan, J. D., Lenton, T. M., Marsh, R., Yool, A., and Watson, A.: Marine geochemical data assimilation in an efficient Earth System Model of global biogeochemical cycling, *Biogeosciences*, 4, 87–104, doi:10.5194/bg-4-87-2007, 2007. 2103

Shaffer, G., Malskær Olsen, S., and Pepke Pedersen, J. O.: Presentation, calibration and validation of the low-order, DCESS Earth System Model (Version 1), *Geosci. Model Dev.*, 1, 17–51, doi:10.5194/gmd-1-17-2008, 2008. 2107

Walker, J. C. G. and Opdyke, B. N.: Influence of variable rates of neritic carbonate deposition on atmospheric carbon dioxide and pelagic sediments, *Paleoceanography*, 10, 415–427, 1995. 2099

Wolf-Gladrow, D. A., Zeebe, R. E., Klaas, C., Körtzinger, A., and Dickson, A. G.: Total alkalinity : the explicit conservative expression and its application to biogeochemical processes, *Mar. Chem.*, 106, 287–300, doi:10.1016/j.marchem.2007.01.006, 2007. 2092

Yao, W. and Millero, F. J.: The chemistry of the anoxic waters in the Framvaren Fjord, Norway, *Aquat. Geochem.*, 1, 53–88, 1995. 2095

Zeebe, R. E. and Wolf-Gladrow, D.: *CO₂ in seawater: equilibrium, kinetics, isotopes*, vol. 65 of *Elsevier Oceanography Series*, Elsevier, Amsterdam (NL), 2001. 2090, 2095, 2109

Solving the
alkalinity–pH
equation

G. Munhoven

Table 1. Execution times for the SW1, SW2 and SW3 tests, each one with the three initialisation types (see text). Crosses (×) indicate test series that were affected by divergences and could not be considered for time measurements (notice one exception); dashes (–) indicate that the experiment was not carried out. Within each one of the groups SW1, SW2 and SW3, figures were normalized to the execution time of the respective “cub” run with `general_sec` and rounded to the nearest multiple of 0.05 (i.e. the order of the the largest standard deviation). “cub” results for `general_sec` are reported in italics as they have been implicitly set to exactly 1 by the normalization procedure and are not affected by the rounding procedure.

Routine	SW1			SW2			SW3		
	cub	pH8	safe	cub	pH8	safe	cub	pH8	safe
general	1.05	1.55	1.65	1.05	1.60	1.65	1.10	1.65	1.75
<code>general_sec</code>	<i>1.00</i>	1.45	1.60	<i>1.00</i>	1.60	1.55	<i>1.00</i>	1.55	1.55
fast	0.95	1.50	1.70	1.00	1.65	1.90*	×	×	×
icacfp	2.35	2.90	–	1.75	2.10	–	×	×	–
bacastow	0.90	1.15	–	0.85	1.10	–	×	×	–
ocmip	1.85	3.15	5.70	1.75	×	5.60	×	×	3.70

* Note: includes one divergent case on 1 950 000 calls.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Solving the
alkalinity–pH
equation

G. Munhoven




Fig. 1. pH_{SWS} values obtained with the new universal algorithm (general) for test cases **(a)** SW1, **(c)** SW2 and **(b)** SW3 – please notice the extended colour scale. **(d)** Absolute value of the equation residual at the adopted root value, derived with that same algorithm started with the cubic-based initialisation to solve test case SW3. Applied convergence criterion: $|H_{n+1} - H_n|/H_n < 10^{-8}$.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Solving the
alkalinity–pH
equation

G. Munhoven




Fig. 2. Number of iterations required by (a) general, (b) icacfp, (c) bacastow with secant iterations on $[H^+]$ and (d) ocmip, each one using the cubic-based initialisation procedure to solve test case SW2. Applied convergence criterion: $|H_{n+1} - H_n|/H_n < 10^{-8}$.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Solving the
alkalinity–pH
equation

G. Munhoven




Fig. 3. (a) Derivative with respect to H of the function underlying the ICAC methods, i.e. of the function Q given by Eqs. (8) and (9) that defines the recurrence $H_{n+1} = Q(Alk_C(Alk_T, H_n), C_T)$. The white line indicates where the derivative is equal to -1 ; in the stippled area, the derivative is strictly lower than -1 . Also shown are the numbers of iterations required to meet the convergence criterion for (b) general, (c) icacfp and (d) bacastow with secant iterations on $[H^+]$. White areas indicate no convergence or an excessive number of iterations ($n > 50$).

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Solving the
alkalinity–pH
equation

G. Munhoven




Fig. 4. (a) Relative deviation (in %) of the solution of the quadratic Eq. (7), calculated by setting $Alk_C = Alk_T$, from the actual root of the complete system; (b) idem for the cubic polynomial based initial $[H^+]$, calculated by setting $Alk_{CB} = Alk_T$. Please notice the strongly different colour scales (underlying data come from test case SW2).

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

