



**The second order
shallow ice
approximation**

J. Ahlkrona et al.

Accuracy of the zeroth and second order shallow ice approximation – numerical and theoretical results

J. Ahlkrona¹, N. Kirchner², and P. Lötstedt¹

¹Division of Scientific Computing, Department of Information Technology, Uppsala University, Uppsala, Sweden

²Bolin Centre for Climate Research and Department of Physical Geography and Quaternary Geology, Stockholm University, Stockholm, Sweden

Received: 20 June 2013 – Accepted: 25 June 2013 – Published: 7 August 2013

Correspondence to: J. Ahlkrona (josefin.ahlkrona@it.uu.se)

Published by Copernicus Publications on behalf of the European Geosciences Union.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Abstract

In ice sheet modelling, the Shallow Ice Approximation (SIA) and Second Order Shallow Ice Approximation (SOSIA) schemes are approaches to approximate the solution of the full Stokes equations governing ice sheet dynamics. This is done by writing the solution to the full Stokes equations as an asymptotic expansion in the aspect ratio ϵ , i.e. the quotient between a characteristic height and a characteristic length of the ice sheet. SIA retains the zeroth order terms and SOSIA the zeroth, first, and second order terms in the expansion. Here, we evaluate the order of accuracy of SIA and SOSIA by numerically solving a two dimensional model problem for different values of ϵ , and comparing the solutions with a finite element solution of the full Stokes equations obtained from Elmer/Ice. The SIA and SOSIA solutions are also derived analytically for the model problem. For decreasing ϵ , the computed errors in SIA and SOSIA decrease, but not always in the expected way. Moreover, they depend critically on a parameter introduced to avoid singularities in Glen's flow law in the ice model. This is because the assumptions behind the SIA and SOSIA neglect a thick, high viscosity boundary layer near the ice surface. The sensitivity to the parameter is explained by the analytical solutions.

1 Introduction

The cryosphere is an important part of the climate system, and features, among others, the Greenland Ice Sheet and the Antarctic Ice Sheet. With a volume of about $3 \times 10^7 \text{ km}^3$, these two ice sheets represent the largest component of the cryosphere, and store about 77% of the global freshwater. Cryospheric research, and specifically, ice sheet modelling, is a vibrant discipline which receives much scientific, political and societal attention because of its relevance for predictions of the future sea level rise in a warming world (Solomon et al., 2007).

GMDD

6, 4281–4325, 2013

The second order shallow ice approximation

J. Ahlkrona et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



The second order shallow ice approximation

J. Ahlkrona et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Increasingly complex numerical ice sheet models are used to simulate the response of the earth's ice sheets to different climate forcings, and internationally coordinated efforts are devoted to ensure inter-comparability of such modelling exercises by means of benchmarking (Pattyn et al., 2008, 2012; Calov et al., 2010). Participating models range from zeroth order shallow ice approximation (SIA) models, to higher order models and full Stokes models; the common understanding being that higher order models are more accurate than zeroth order models, while full Stokes models are the most accurate (but also most costly) ones. However, as increasingly accurate numerical ice sheet models and corresponding powerful hardware become more readily available, overall less research is focused on e.g. analyses devoted to basic foundations of ice sheet modelling, such as validity and limitations of approximation schemes and associated scaling procedures. The latter were especially important when virtually all ice sheet modelling was relying on SIA schemes because computing power was more restricted than it is today. Yet, they have not lost their relevance because of challenging applications such as paleoglacial simulations or uncertainty quantification, and a large portion of ice sheet simulations is still today performed with SIA codes or higher-order approximations.

In this paper we investigate the order of accuracy and validity of a higher order extension to the SIA; the Second Order Shallow Ice Approximation (SOSIA). In the process we also study the accuracy of the classical SIA. The SIA was constructed in the 1970s and 1980s by Fowler and Larson (1978), Hutter (1983), and Morland (1984). The derivation is based on scaling and asymptotic series expansion in terms of the aspect ratio, ϵ , which expresses the shallowness of an ice sheet. In the SIA, only the zeroth order terms are kept. In the end of the 1990's, the Second Order Shallow Ice Approximation (SOSIA) was derived by Baral (1999) and Baral et al. (2001), pushing the series expansion to second order in ϵ with the objective to include dynamics not captured by the SIA. The SOSIA is thus a higher order model based on the same theory as the SIA, and is almost as computationally cheap as the SIA. The scaling assumptions underlying the SOSIA (and the SIA as derived in Baral, 1999; Baral et al., 2001; Greve,

The second order shallow ice approximation

J. Ahlkrone et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



1997) do however neglect a high viscosity boundary layer near the ice surface. This boundary layer is thick (Ahlkrone et al., 2013), and other scaling assumptions by Johnson and McMeeking (1984) and Schoof and Hindmarsh (2010) are more appropriate than the classical SIA scalings (Ahlkrone et al., 2013). This calls for a proper analysis of the validity and accuracy of the SOSIA equations, and also an investigation of the true order of accuracy of the SIA.

Our analysis is based on analytical solutions of the SOSIA (and SIA) as well as numerical solutions of SIA, SOSIA and the full Stokes equations. Ultimately, it is of interest to assess under which circumstances SOSIA models can be regarded as a significant improvement to the SIA at low computational costs. The SOSIA, as described in Baral et al. (2001), has to our knowledge never been implemented before this work, most likely because the second order expressions are long and tedious to code. It was applied by Mangeney and Califano (1998) for Newtonian, anisotropic ice, and recently by Egholm et al. (2011), however not in its pure form but in a depth-averaged, iterative scheme.

The outline of this paper is as follows: Sect. 2 is devoted to a summary of the general equations pertaining to ice dynamics, and to their zeroth, first, and second order approximations. Recent works by Ahlkrone et al. (2013) and Kirchner et al. (2011) allow us to keep the presentation to a minimum. Section 3 describes the model problem which we focus on throughout the paper; a two dimensional flow over a bumpy bed. In Sect. 4 an analytical SOSIA solution for this model problem is presented and discussed, extending previous work by Baral (1999) and Baral et al. (2001). In Sect. 5 we implement the SOSIA numerically and compute the accuracy of both the SIA and the SOSIA by comparing with the solution obtained with Elmer/ICE (Gagliardini et al., 2013), for varying ϵ . In both Sects. 4 and 5 we investigate the effect of an extra parameter, σ_{res} , which is necessary due to the neglect of the boundary layer at the surface. The paper concludes with a discussion in Sect. 6.

2 Derivation of the zeroth and second order shallow ice approximation

2.1 The exact equations

Ice sheet flow is commonly described using concepts from continuum mechanics, materials science, and thermodynamics, which allow to formulate the spatio-temporal evolution of ice masses as an initial boundary value problem, with free boundaries. Ice flow is momentum, mass, and internal energy conserving. As we will only study isothermal flow the equations regarding energy are not described here. The equations for balance of mass and momentum are

$$0 = \text{div} \mathbf{v}, \quad (1)$$

$$\rho \dot{\mathbf{v}} = -\nabla p + \text{div} \mathbf{T}^D + \rho \mathbf{g}, \quad (2)$$

where ρ is the density, \mathbf{v} the velocity field, \mathbf{T}^D the deviatoric stress tensor, \mathbf{g} gravitational acceleration. The deviatoric stress tensor, \mathbf{T}^D , and the Cauchy stress tensor, \mathbf{T} are related by $\mathbf{T} = -p\mathbf{I} + \mathbf{T}^D$, where p is the pressure. The acceleration term, $\dot{\mathbf{v}}$ being the material time derivative of the velocity, in Eq. (2) is very small and is therefore neglected in glaciological applications. The resulting equations are called the Stokes equations, or in glaciology rather the *full* Stokes equations. Velocity and stress are related by the constitutive equation,

$$\mathbf{D} = \mathcal{A}(T') f(\sigma) \mathbf{T}^D, \quad (3)$$

in glaciology called Glen's flow law. The strain rate tensor \mathbf{D} is defined as $(\nabla \mathbf{v} + (\nabla \mathbf{v})^*)/2$, where $*$ denotes transpose and $\mathcal{A}(T')$ accounts for coupling the viscosity, η , to the pressure melting point corrected temperature T' . For isothermal conditions \mathcal{A} is merely a constant. The so-called creep response function, f , is defined by $f(\sigma) = \sigma^{n-1}$, where we let n be equal to the standard value 3. Its argument σ , the effective stress, is

GMDD

6, 4281–4325, 2013

The second order shallow ice approximation

J. Ahlkrone et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



The second order shallow ice approximation

J. Ahlkrone et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



the square root of the second invariant of \mathbf{T}^D , and is defined by

$$\sigma^2 = \frac{1}{2} \text{tr}(\mathbf{T}^D)^2 = (t_{xz}^D)^2 + (t_{yz}^D)^2 + (t_{xy}^D)^2 + \frac{1}{2} \left((t_{xx}^D)^2 + (t_{yy}^D)^2 + (t_{zz}^D)^2 \right). \quad (4)$$

Here, t_{ij}^D ($i, j = x, y, z$) are the components of \mathbf{T}^D in a Cartesian coordinate system where the z axis is pointing in the opposite direction of gravity. We refer to t_{xx}^D , t_{yy}^D and t_{zz}^D as *normal deviatoric stresses*, t_{xz}^D ($= t_{xz}$) and t_{yz}^D ($= t_{yz}$) as *vertical shear stresses* and t_{xy}^D ($= t_{xy}$) as the *horizontal plane shear stress*. In glaciology it is also common to use the term *longitudinal stresses* for t_{xx}^D , t_{yy}^D , t_{zz}^D and t_{xy}^D . As the creep response function depends on the effective stress, ice is a non-Newtonian fluid with viscosity, η , given by $1/\eta = 2A(T')f(\sigma)$. This non-linearity makes the ice flow simulations a computationally heavy task.

To complete the system, boundary conditions are imposed at the ice base and ice surface. At the impermeable ice base the velocity satisfies no slip conditions on a rigid bedrock, giving the condition

$$\mathbf{v} = \mathbf{0}, \quad (5)$$

and the ice surface is assumed to be stress-free,

$$\mathbf{T} \times \mathbf{n} = \mathbf{0}. \quad (6)$$

Here \mathbf{n} is the outward pointing normal vector of the ice surface. In the time-dependent case a transport equation for the ice surface elevation is solved, where the velocity field enters as coefficients and the accumulation or ablation enters as a forcing. The equation for the height of the free ice surface $h(x, y, t)$ is

$$\frac{\partial h}{\partial t} + v_x \frac{\partial h}{\partial x} + v_y \frac{\partial h}{\partial y} - v_z = a_s, \quad (7)$$

where v_x , v_y and v_z are the velocity components and a_s is the accumulation-ablation function.

2.2 Shallow ice approximations

Here we briefly describe how the SIA and the SOSIA are derived from the exact, full Stokes equations by scaling and perturbation expansions. We exemplify the procedure by the momentum balance Eq. (2). We follow the scalings presented in Baral (1999); Baral et al. (2001); Greve (1997), as these are ones most commonly used today or deriving the SIA and also those used to go further and arrive at the SOSIA equations. These scalings are

$$\begin{aligned}
 (x, y) &= [L](\tilde{x}, \tilde{y}), & \rho &= \rho g[H]\tilde{\rho}, \\
 z &= [H]\tilde{z}, & (t_{xz}^D, t_{yz}^D, \sigma) &= \epsilon \rho g[H](\tilde{t}_{xz}^D, \tilde{t}_{yz}^D, \tilde{\sigma}), \\
 (v_x, v_y) &= [V_L](\tilde{v}_x, \tilde{v}_y), & (t_{xx}^D, t_{yy}^D, t_{xy}^D, t_{zz}^D) &= \epsilon^2 \rho g[H](\tilde{t}_{xx}^D, \tilde{t}_{yy}^D, \tilde{t}_{xy}^D, \tilde{t}_{zz}^D), \\
 v_z &= [V_H]\tilde{v}_z, & t &= ([L]/[V_L])\tilde{t}, \\
 \epsilon &= [H]/[L] = [V_H]/[V_L], & F &= [V_L]^2/g[L],
 \end{aligned} \tag{8}$$

where the aspect ratio, ϵ , has been introduced. The dimensionless quantities are denoted by tilde and are assumed to be of the order of magnitude $\mathcal{O}(1)$. They are multiplied by typical values of height, $[H]$, length, $[L]$, and vertical and horizontal velocity, $[V_H]$, $[V_L]$, where $[H] \ll [L]$ so that the aspect ratio, ϵ , is small. The scalings reflects that the vertical shear stresses are assumed to dominate over the normal deviatoric and normal shear stresses, and that this dominance is stronger the more shallow the ice sheet is. Also, the horizontal velocity is assumed to dominate over the vertical velocity, $[V_H] \ll [V_L]$.

The scalings Eq. (8) are inserted into the equations and a perturbation expansion is performed, i.e. the dimensionless variables, \tilde{q} , are expanded in a power series as

$$\tilde{q} = \tilde{q}_{(0)} + \epsilon \tilde{q}_{(1)} + \epsilon^2 \tilde{q}_{(2)} + \dots \tag{9}$$

Collecting terms of equal order in ϵ gives rise to a hierarchy of models, called the SIA for zeroth order in ϵ , the FOSIA (First Order Shallow Ice Approximation) for first order

in ϵ , and SOSIA for second order. For the momentum balance Eq. (2) (in component form), the SIA model is

$$0 = -\frac{\partial \tilde{\rho}_{(0)}}{\partial \tilde{x}} + \frac{\partial \tilde{t}_{xz(0)}^D}{\partial \tilde{z}}, \quad (10a)$$

$$0 = -\frac{\partial \tilde{\rho}_{(0)}}{\partial \tilde{y}} + \frac{\partial \tilde{t}_{yz(0)}^D}{\partial \tilde{z}}, \quad (10b)$$

$$5 \quad 1 = -\frac{\partial \tilde{\rho}_{(0)}}{\partial \tilde{z}}, \quad (10c)$$

the FOSIA is

$$0 = -\frac{\partial \tilde{\rho}_{(1)}}{\partial \tilde{x}} + \frac{\partial \tilde{t}_{xz(1)}^D}{\partial \tilde{z}}, \quad (11a)$$

$$0 = -\frac{\partial \tilde{\rho}_{(1)}}{\partial \tilde{y}} + \frac{\partial \tilde{t}_{yz(1)}^D}{\partial \tilde{z}}, \quad (11b)$$

$$10 \quad 0 = -\frac{\partial \tilde{\rho}_{(1)}}{\partial \tilde{z}}, \quad (11c)$$

and the SOSIA is

$$0 = -\frac{\partial \tilde{\rho}_{(2)}}{\partial \tilde{x}} + \frac{\partial \tilde{t}_{xz(2)}^D}{\partial \tilde{z}} + \frac{\partial \tilde{t}_{xx(0)}^D}{\partial \tilde{x}} + \frac{\partial \tilde{t}_{xy(0)}^D}{\partial \tilde{y}}, \quad (12a)$$

$$0 = -\frac{\partial \tilde{\rho}_{(2)}}{\partial \tilde{y}} + \frac{\partial \tilde{t}_{yz(2)}^D}{\partial \tilde{z}} + \frac{\partial \tilde{t}_{xy(0)}^D}{\partial \tilde{x}} + \frac{\partial \tilde{t}_{yy(0)}^D}{\partial \tilde{y}}, \quad (12b)$$

$$15 \quad 0 = -\frac{\partial \tilde{\rho}_{(2)}}{\partial \tilde{z}} + \frac{\partial \tilde{t}_{xz(0)}^D}{\partial \tilde{x}} + \frac{\partial \tilde{t}_{yz(0)}^D}{\partial \tilde{y}} + \frac{\partial \tilde{t}_{zz(0)}^D}{\partial \tilde{z}}. \quad (12c)$$

The second order shallow ice approximation

J. Ahlkrone et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Note that there are no other stress components besides the vertical shear stress in the zeroth order equations. This is the main reason why higher order models are becoming increasingly popular: longitudinal stress and horizontal plane shear stress are important in the coupling to ice shelves and in ice streams and ice stream shear margins.

Integrating Eq. (10c) in the vertical direction and using the boundary condition that the pressure is zero at the ice surface (cf. Eq. 6) gives an explicit expression for the pressure. Inserting the pressure in Eqs. (10a) and (10b), integrating and using the stress free condition again yields simple expressions for the vertical shear stresses. The shear stresses are sufficient to obtain the zeroth order (SIA) velocities by integration of the stress strain rate relation Eq. (3), see Sect. 4 for the outcome of this standard procedure. Once the zeroth order solution is available, it can be used to solve the FOSIA, and subsequently the SOSIA equations by the same method.

Just like the SIA, the FOSIA only contains the vertical shear stresses, but no normal deviatoric stress or horizontal plane shear stress. However, the FOSIA does account for first order boundary effects and forcings that cannot be represented in SIA (Baral, 1999). For the model problem that we study, the FOSIA solution is trivially zero, and the FOSIA will hence not be further discussed. It is in the SOSIA that the normal deviatoric stresses and the horizontal plane shear stresses are present for the first time, suggesting that more complex ice dynamical behaviour can be captured than with the SIA. The SOSIA is computationally inexpensive compared to many other higher order models, because the equations can be solved without a coupling of variables in a system of linear equations.

2.3 Boundary layer treatment

The SIA and the SOSIA are both based on the same scaling arguments in Eq. (8), where it is assumed, in addition to the ice body being shallow, that the dominating stress component is vertical shear stress and that the velocity components in the horizontal plane dominate over the vertical velocity. These assumptions are not valid in a number of well known situations, including fast sliding at the ice/bedrock interface as

The second order shallow ice approximation

J. Ahlkrone et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



in for example ice streams, or at the ice divide where the ice flows mainly downwards. Where fast sliding occurs it is common to use other scaling arguments, such as e.g. the shelfy stream approximation (MacAyeal, 1992).

Another region where the SIA scalings break down is in a boundary layer near the entire ice surface, developing in the case of a bumpy bed due to the non-linear rheology of ice (Ahlkrone et al., 2013). Johnson and McMeeking (1984) made the first attempt to treat this issue by matched asymptotics, rescaling the pressure and stress components in the boundary layer. This allows for all stress components to influence the dynamics near the ice surface. By theoretical analysis they found the boundary layer thickness to be $O(\epsilon^{1/3})$. Schoof and Hindmarsh (2010) extended the boundary layer theory by including the degree of slip at the bed in the expansion, and pushing it to second order. In the case of slow sliding, the rescaling of the pressure and stress components in the boundary layer is (in two dimensions, as in Johnson and McMeeking, 1984; Schoof and Hindmarsh, 2010; Ahlkrone et al., 2013):

$$\begin{aligned} p &= \epsilon^{1/3} \rho g[H] \tilde{p}, \\ t_{xz}^D &= \epsilon^{4/3} \rho g[H] \tilde{t}_{xz}^D, \\ (t_{xx}^D, t_{zz}^D) &= \epsilon^{4/3} \rho g[H] (\tilde{t}_{xx}^D, \tilde{t}_{zz}^D). \end{aligned} \quad (13)$$

By numerically solving the full Stokes equations, Ahlkrone et al. (2013) essentially confirmed the appropriateness of these rescalings for the problem described in Sect. 3. There seems to be a common perception that the boundary layer is thin, but as found in Ahlkrone et al. (2013) it is thick and indistinct, and needs to be considered in model development.

Matched asymptotics, as used in Johnson and McMeeking (1984) and Schoof and Hindmarsh (2010) is unfortunately rather involved and has, to our knowledge, never been implemented into a numerical code for practical use. Baral (1999) and Baral et al. (2001) indeed dismiss it as too complicated and use the scalings in Eq. (8) throughout the entire ice sheet in the derivation of the SIA and SOSIA. This, in combination with

The second order shallow ice approximation

J. Ahlkrone et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



the fact that the rheology of ice is singular and non-linear, does introduce complications that need remedy. The singularity arises from the fact that the creep response function is zero for zero effective stress, i.e. $f(0) = 0$. This means that the viscosity is infinite where the effective stress is zero. By the shallow ice scalings Eq. (8), the normal and horizontal plane shear stress $t_{xx}^D, t_{yy}^D, t_{zz}^D, t_{xy}^D$ in Eq. (4) are neglected such that $f(\sigma_{(0)}) = \sigma_{(0)}^2 = t_{xz(0)}^2 + t_{yz(0)}^2$. Thus the (zeroth order) effective stress and the creep response function are zero wherever the zeroth order vertical shear stresses are zero, which is at the entire ice surface (Baral, 1999; Baral et al., 2001; Greve, 1997). In reality, normal deviatoric stresses develop (and depending on the situation, horizontal plane shear stresses), implying non-zero effective stress at the ice surface except for a few points.

As SIA is the zeroth order expansion it can be derived using several different scaling arguments. Therefore it is not very sensitive to the neglect of normal and horizontal plane shear stress in the boundary layer. Indeed, the computation of the SIA solution does not include the reciprocal of the creep response function (except for in the normal deviatoric stress), and the fact that the zeroth order effective stress is zero at the entire surface does not imply singularities in the velocity field, pressure, or shear stress. The SOSIA is however bound to be more sensitive. When pushing the asymptotic expansions to second order, the creep response function of the zeroth order effective stress does occur in the denominator. To remedy this, an extra parameter, σ_{res} , is used in Baral (1999) and Baral et al. (2001), to regularize the problem, following earlier suggestions by Lliboutry (1969) and Colbeck and Evans (1973). This parameter, which we will call the *finite viscosity parameter* σ_{res} , is added to f as: $f(\sigma) = \sigma^2 + \sigma_{\text{res}}^2$ (see e.g. Colbeck and Evans, 1973). Non-singular creep functions of non-additive structure have been proposed by e.g. Lliboutry (1969). The question of the appropriateness in modifying the material law instead of rescaling the variables in the boundary layer immediately arises. As already mentioned, the singularities do not affect SIA. Hence, in practice σ_{res} is not needed in the SIA, but the solution will not be accurate to the order predicted by the theory in Baral (1999) and Baral et al. (2001). It is unclear whether the SOSIA will be an improvement of SIA, because of the neglect of the special boundary layer

dynamics. Moreover, there is no obvious way to choose the value of the finite viscosity parameter. In Baral et al. (2001), $\sigma_{\text{res}} = \sqrt{10^9}$ is used for the Greenland ice sheet, but with no explanation of this choice. In Sect. 5, we will investigate how to choose σ_{res} , how accurate the SIA is, and whether the SOSIA really is an improvement of the SIA.

3 Model problem – ice flow over a bumpy bed

Throughout this paper we will consider the model problem described in this section. It is a slight modification of the problem studied in the ISMIP-HOM benchmark experiment B (Pattyn et al., 2008). As in Pattyn et al. (2008), we investigate a diagnostic, isothermal 2-D-problem for ice flow over an inclined, bumpy bed. The ice surface is fixed, periodic boundary conditions are applied and no-slip conditions are imposed at the base, see Fig. 1. The rate factor \mathcal{A} is $10^{-16} \text{ Pa}^{-3} \text{ a}^{-1}$, ice density is the standard value of 910 kg m^{-3} , and accumulation and ablation are neglected. The mean ice thickness, $[H]$, is 1000 m and the ice surface, h , and ice base, b , are given by

$$h(x) = -x \tan(\alpha), \quad b(x) = h - [H] + \mu[H] \sin\left(\frac{2\pi}{L}x\right). \quad (14)$$

The amplitude of the bumps is $\mu[H]$ and the inclination angle of the surface slope is α . The typical horizontal extent of the problem equals the wavelength of the sinusoidal bumps, i.e. $L = [H]/\epsilon$ m. The wavelength L of the bumps is varied while $[H]$ is kept constant, which corresponds to varying ϵ .

As shown in Ahlkrone et al. (2013), and as can be seen for instance by non-dimensionalising the ice surface and ice bed, the surface slope should ideally be $\arctan \epsilon$ instead of $\alpha = 0.5$ which was used in the ISMIP-HOM benchmark. The bump amplitude μ should be independent of ϵ . In our numerical experiments we will use $\mu = 0.5$ and $\mu = 0.1$.

4 Analytical solutions for the SIA and the SOSIA

For a deeper understanding, we now compute analytical solutions for the second order field variables $t_{xz(2)}$, $p_{x(2)}$ and $v_{x(2)}$, and also give the zeroth order expressions for completeness. Note that for the 2-D case $v_y = t_{yy}^D = t_{xy}^D = 0$. The second order z velocity is excluded for brevity of presentation, since it follows from the mass balance in the same way for all orders.

4.1 General solution

The following expressions (Eqs. 15–19) hold, not only for our model problem, but for all isothermal, 2-D problems with no-slip conditions at the base and without higher order boundary terms. Generalising to 3-D, including higher order boundary terms, and sliding is straight forward. The well-known zeroth order expressions, denoted by subscript (0), for shear stress (cf. Eq. 10) and velocity are

$$t_{xz(0)}^D = -\rho g \frac{\partial h_{(0)}}{\partial x} (h_{(0)} - z), \quad v_{x(0)} = 2\rho g \frac{\partial h_{(0)}}{\partial x} \int_{b_{(0)}}^z \mathcal{A}f(\sigma_{(0)}) (h_{(0)} - z') dz'. \quad (15)$$

In order to compute $t_{xz(2)}^D$ from Eq. (12) we need the zeroth order normal deviatoric stresses (and depending on the situation, horizontal plane shear stress), which are not computed when applying only the zeroth order approximation, as they are not needed for the zeroth order velocities. The normal deviatoric stress in the x direction is given directly by the stress–strain relation in Baral et al. (2001) and Greve (1997):

$$t_{xx(0)}^D = \frac{1}{\mathcal{A}f(\sigma_{(0)})} \frac{\partial v_{x(0)}}{\partial x}. \quad (16)$$

Note that $t_{xx}^D = -t_{zz}^D$ in the 2-D case. The shear stress, pressure and the horizontal velocity are obtained from the horizontal momentum balance, vertical momentum balance and stress strain relation, respectively, where they occur in derivatives.

It is customary to vertically integrate the SIA-equations in order to avoid having to solve for the variables numerically. This is also convenient for the SOSIA. In Baral et al. (2001), the vertical integration is carried out for the shear stresses and the pressure only, and the resulting expressions are presented in Eqs. (17) and (18) (with misprints in Baral et al., 2001, corrected). Going beyond the results of Baral et al. (2001), we derive here also the second order velocity, $v_{x(2)}$ in Eq. (19), computed from the stress-strain relation and boundary conditions at the base:

$$\epsilon^2 p_{(2)} = \rho g \frac{1}{2} (h_{(0)} - z)^2 \frac{\partial^2 h_{(0)}}{\partial x^2} + \rho g (h_{(0)} - z) \left(\frac{\partial h_{(0)}}{\partial x} \right)^2 + t_{zz(0)}^D, \quad (17)$$

$$\begin{aligned} \epsilon^2 t_{xz(2)}^D = & -\rho g \frac{\partial}{\partial x} \left(-\frac{1}{6} (z - h_{(0)})^3 \frac{\partial^2 h_{(0)}}{\partial x^2} h_{(0)} + \frac{1}{2} (z - h_{(0)})^2 \|\nabla_H h_{(0)}\|^2 \right) \\ & - \frac{\partial}{\partial x} \int_z^{h_{(0)}} t_{zz(0)}^D dz' + \frac{\partial}{\partial x} \int_z^{h_{(0)}} t_{xx(0)}^D dz', \end{aligned} \quad (18)$$

$$\begin{aligned} \epsilon^2 v_{x(2)} = & -\frac{\partial}{\partial x} \int_{b_{(0)}}^z v_{z(0)} dz + 6\mathcal{A} \int_{b_{(0)}}^z (t_{xz(0)}^D)^2 \epsilon^2 t_{xz(2)}^D dz' \\ & + 2\mathcal{A} \int_{b_{(0)}}^z t_{xz(0)}^D (t_{xx(0)}^D)^2 dz' + 2\sigma_{\text{res}}^2 \mathcal{A} \int_{b_{(0)}}^z \epsilon^2 t_{xz(2)}^D dz'. \end{aligned} \quad (19)$$

Equations (17)–(19) yield explicit expressions for second order variables.

4.2 Solution for the model problem

For the geometry in Sect. 3 we have computed the integrals and derivatives in these expressions and thus obtained analytical solutions to the SOSIA. To avoid infinite viscosity, we have followed the suggested regularisation in Baral et al. (2001), i.e.

- 5 adding a constant to the creep response function, $f(\sigma) = \sigma^2 + \sigma_{\text{res}}^2$. The solutions are expressed in terms of the inclination angle of the ice surface α , relative amplitude μ , finite viscosity parameter, σ_{res} and wavelength L . In the Appendix A the same solutions are expressed in a more general form. By Eq. (15), the zeroth order shear stress for the model problem is

$$10 \quad t_{xz(0)}^D = -\rho g \tan(\alpha) (x \tan(\alpha) + z). \quad (20)$$

The zeroth order velocity in Eq. (15) is given by

$$\begin{aligned} v_{x(0)} = & 2\rho g A \tan(\alpha) \left(\frac{(\rho g)^2 \tan^2(\alpha)}{4} \left([H]^4 (1 - \mu \sin(2\pi x/L))^4 - (x \tan(\alpha) + z)^4 \right) \right. \\ & \left. + \frac{\sigma_{\text{res}}^2}{2} \left([H]^2 (1 - \mu \sin(2\pi x/L))^2 - (x \tan(\alpha) + z)^2 \right) \right). \end{aligned} \quad (21)$$

- 15 The maximum magnitude of the velocity in Eq. (21) does not depend on L , but it does depend on the relative amplitude μ . Also, there is an extra term stemming from the finite viscosity law with $\sigma_{\text{res}} > 0$. As mentioned in Sect. 2.3, a finite viscosity law is technically not needed for the zeroth order model, since the creep response function never appears in the denominator in the calculations of the velocity field or shear stress (see Eqs. 20 and 21), which are usually the variables of interest. However, for reasons
- 20 of consistency, it should be used also in the zeroth order model when continuing to second order. To calculate second order shear stress and velocity, the zeroth order normal deviatoric stress in Eq. (16) is needed, given for the model problem by

$$t_{xx(0)}^D = -2\rho g \tan^2(\alpha) (x \tan(\alpha) + z)$$

$$\begin{aligned}
& - \frac{4\pi}{L} \rho g [H]^2 \mu \tan(\alpha) \frac{\cos(2\pi x/L) (1 - \mu \sin(2\pi x/L))}{(x \tan(\alpha) + z)^2 \tan^2(\alpha) + \left(\frac{\sigma_{\text{res}}}{\rho g}\right)^2} \\
& \cdot \left(\tan^2(\alpha) [H]^2 (1 - \mu \sin(2\pi x/L))^2 + \left(\frac{\sigma_{\text{res}}}{\rho g}\right)^2 \right).
\end{aligned} \tag{22}$$

Having calculated $t_{xx(0)}^D$, the second order shear stress is computed from Eq. (18) as

$$\begin{aligned}
& \epsilon^2 t_{xz(2)}^D = 3\rho g \tan^3(\alpha) (x \tan(\alpha) + z) \\
& + 8\rho g [H]^2 \frac{\pi}{L} \mu \tan^2(\alpha) \frac{(1 - \mu \sin(2\pi x/L)) \cos(2\pi x/L)}{\tan^2(\alpha) (x \tan(\alpha) + z)^2 + \frac{\sigma_{\text{res}}^2}{(\rho g)^2}} \\
& \cdot \left([H]^2 \tan^2(\alpha) (1 - \mu \sin(2\pi x/L))^2 + \frac{\sigma_{\text{res}}^2}{(\rho g)^2} \right) \\
& - \frac{4(\rho g)^2 [H]^2}{\sigma_{\text{res}}} \left(\frac{2\pi}{L} \right)^2 \mu \left(3[H]^2 (1 - \mu \sin(2\pi x/L))^2 \mu \cos^2(2\pi x/L) \tan^2(\alpha) \right. \\
& + [H]^2 (1 - \mu \sin(2\pi x/L))^3 \sin(2\pi x/L) \tan^2(\alpha) \\
& + \frac{\sigma_{\text{res}}^2}{(\rho g)^2} \left(\mu \cos^2(2\pi x/L) + (1 - \mu \sin(2\pi x/L)) \sin(2\pi x/L) \right) \Big) \\
& \cdot \arctan \left(\rho g \frac{\tan(\alpha) (x \tan(\alpha) + z)}{\sigma_{\text{res}}} \right).
\end{aligned} \tag{23}$$

An explicit dependence on L is introduced in Eqs. (22)–(23), and hence on the aspect ratio ϵ . Also, Eqs. (22) and (23) depend on α , μ and σ_{res} . In $t_{xx(0)}^D$, σ_{res} appears in

The second order shallow ice approximation

J. Ahlkrone et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



the denominator, preventing singularities to occur at the ice surface when z is equal to $h = -x \tan(\alpha)$, see Eq. (22). In the second order shear stress in Eq. (23), σ_{res} is both in the numerator and the denominator. The second order shear stress is dominated by the last term, i.e. by the last four lines in Eq. (23).

Having calculated $t_{xx(0)}^D$ and $t_{xz(2)}^D$, the second order velocity can be derived from Eq. (19). The expression for the second order solution is very long and is included in the Appendix A for the interested reader. In fact it behaves similarly to the second order shear stress, where σ_{res} appears both in the numerator and the denominator. Remember that to get the full second order solution, the zeroth and second order contributions should be added together as $v_x = v_{x(0)} + \epsilon^2 v_{x(2)}$ (since the first order solution is zero for our model problem). The extra term arising from the finite viscosity law in $v_{x(0)}$ in Eq. (21) is quite large, and thus the SOSIA velocity does not allow too large a σ_{res} . The SOSIA shear stress is not as sensitive to large σ_{res} , since the zeroth order shear stress in Eq. (20) does not contain any terms from the finite viscosity law. Both the second order shear stress correction and velocity correction are however very sensitive to too small a σ_{res} .

Note that all terms involving σ_{res} in Eqs. (22), (23), and (A6) in the Appendix A are pre-multiplied with μ ; thus, for decreasing μ the importance of σ_{res} decays. This is consistent with the fact that the boundary layer near the ice surface disappears when $\mu = 0$ (Ahlkrone et al., 2013).

4.3 Choices of σ_{res} and impact on scalings

In order for any scaling relations (Eq. 8 or Eq. 13) to hold, $\tan(\alpha)$ should vary linearly with ϵ , ideally $\tan(\alpha) = \epsilon$, and μ should be independent of ϵ (Ahlkrone et al., 2013). Note however that the scaling relations were derived in a context where the creep response function was not modified by an additional finite viscosity parameter. We now discuss now how σ_{res} , introduced in an a posteriori fashion, can be chosen such that compatibility with either the classical SIA-scalings in Eq. (8) or the ones in Eq. (13) is achieved.

4.3.1 Choosing σ_{res} consistent with the classical SIA-scalings

If we choose the finite viscosity parameter, σ_{res} , to vary in the same way as the effective stress is assumed to do (linearly with ϵ), the SIA and SOSIA solutions fulfil the scaling relations Eq. (8) that they are derived from. Inserting $\tan(\alpha) = \epsilon$, $\sigma_{\text{res}} = C_\sigma \rho g[H] \epsilon$ and the scaled variables in Eq. (8) into Eqs. (20)–(23) yields:

$$t_{xz(0)}^D = -\rho g[H] \epsilon (\tilde{x} + \tilde{z}), \quad (24)$$

$$v_{x(0)} = \mathcal{A}[H](\rho g[H])^3 \epsilon^3 \left(\frac{1}{2} (1 - \mu \sin(2\pi \tilde{x}))^4 - \frac{1}{2} (\tilde{x} + \tilde{z})^4 + C_\sigma^2 \left((1 - \mu \sin(2\pi \tilde{x}))^2 - (\tilde{x} + \tilde{z})^2 \right) \right), \quad (25)$$

$$t_{xx(0)}^D = -\rho g[H] \epsilon^2 \frac{2}{(\tilde{x} + \tilde{z})^2 + C_\sigma^2} \left((\tilde{x} + \tilde{z}) \left((\tilde{x} + \tilde{z})^2 + C_\sigma^2 \right) - 2\pi \mu \cos(2\pi \tilde{x}) (1 - \mu \sin(2\pi \tilde{x})) \left((1 - \mu \sin(2\pi \tilde{x})) + C_\sigma^2 \right) \right), \quad (26)$$

$$\begin{aligned} \epsilon^2 t_{xz(2)}^D = & \rho g[H] \epsilon^3 \left(3(\tilde{x} + \tilde{z}) \right. \\ & \left. + 8\pi \mu \frac{(1 - \mu \sin(2\pi \tilde{x})) \cos(2\pi \tilde{x}) \left((1 - \mu \sin(2\pi \tilde{x}))^2 + C_\sigma^2 \right)}{(\tilde{x} + \tilde{z})^2 + C_\sigma^2} \right. \\ & \left. - 16\pi^2 \frac{\mu}{C_\sigma} \left(3\mu (1 - \mu \sin(2\pi \tilde{x}))^2 \cos^2(2\pi \tilde{x}) + (1 - \mu \sin(2\pi \tilde{x}))^3 \sin(2\pi \tilde{x}) \right. \right. \\ & \left. \left. + C_\sigma^2 \left(\mu \cos^2(2\pi \tilde{x}) + (1 - \mu \sin(2\pi \tilde{x})) \sin(2\pi \tilde{x}) \right) \right) \arctan \left(\frac{\tilde{x} + \tilde{z}}{C_\sigma} \right) \right), \quad (27) \end{aligned}$$

GMDD

6, 4281–4325, 2013

The second order
shallow ice
approximation

J. Ahlkrona et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



The second order shallow ice approximation

J. Ahlkrone et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



where C_σ is a constant and \tilde{x} and \tilde{z} are the non-dimensionalised x and z coordinates in Eq. (8). The expressions in Eqs. (24)–(27) only depend on geometry, material constants, and C_σ . In line with Eq. (8), $t_{xz(0)}^D$ is pre-multiplied by $\rho g[H]\epsilon$, $t_{xx(0)}^D$ is pre-multiplied by $\rho g[H]\epsilon^2$ and $\epsilon^2 t_{xz(2)}^D$ is pre-multiplied by $\rho g[H]\epsilon^3$. Note that the multiplying factor in $v_{x(0)}$ is $\mathcal{A}[H](\rho g[H])^3 \epsilon^3$ as in Blatter (1995) and Schoof and Hindmarsh (2010). In the same manner, the analytical solution for $v_{z(0)}$ (which is not given here for brevity of presentation) is multiplied by $\mathcal{A}[H](\rho g[H])^3 \epsilon^4$.

4.3.2 Choosing σ_{res} consistent with boundary layer theory

We know that the scaling relations Eq. (13) from Schoof and Hindmarsh (2010) (slightly rearranged) are more correct than the classical scaling relations in Eq. (8) (Ahlkrone et al., 2013; Schoof and Hindmarsh, 2010; Johnson and McMeeking, 1984), and that σ_{res} is merely a parameter introduced to address this problem when using Eq. (8). However, we show now that instead of setting $\sigma_{\text{res}} = C_\sigma \rho g[H]\epsilon$, with C_σ constant, we can choose C_σ such that the field variables fulfil the scaling relations in Eq. (13) rather than Eq. (8).

Near the ice surface, σ is dominated by t_{xx}^D (Ahlkrone et al., 2013; Schoof and Hindmarsh, 2010) and the creep response function satisfies $f(\sigma) \approx t_{xx}^{D2}$ in 2-D. In the SOSIA model, t_{xx}^D is neglected in the creep response and instead $f(\sigma) = \sigma_{\text{res}}^2$ at the ice surface. Hence, C_σ in $\sigma_{\text{res}} = C_\sigma \rho g[H]\epsilon$ should be such that $\sigma_{\text{res}} = t_{xx}^D$ in the upper boundary layer. Let $C_\sigma = C_\gamma \epsilon^\gamma$ and insert into Eq. (26). Since $\tilde{x} + \tilde{z} = 0$ at the surface, we have that $t_{xx}^D \sim \epsilon^2 / \epsilon^{2\gamma} \sim \sigma_{\text{res}} \sim \epsilon^{1+\gamma}$ and consequently $\gamma = 1/3$. Then $t_{xx}^D \sim \epsilon^{4/3}$ in the boundary layer as derived in Eq. (3.108) in Schoof and Hindmarsh (2010). Outside the boundary layer in Eq. (26) when $\tilde{x} + \tilde{z} = O(1)$, $t_{xx}^D \sim \epsilon^2$ as expected from the SOSIA-equations. By replacing C_σ by $C_\gamma \epsilon^\gamma$ in the expression for t_{xx}^D in Eq. (26), inserting $\sigma_{\text{res}} = t_{xx}^D$ and $\tilde{x} + \tilde{z} = 0$, solving for C_γ and finally ignoring terms depending on ϵ , we find that C_γ is

approximated by

$$C_Y^3 = 4\pi\mu \cos(2\pi\tilde{x})(1 - \mu \sin(2\pi\tilde{x}))^2. \quad (28)$$

Thus, C_Y decreases with decreasing bump amplitude, only depends on the geometry and is of $O(1)$. Due to the assumptions made when obtaining Eq. (28) it can not be used directly to determine C_Y , but gives an understanding of its behaviour which we will recognize in our numerical experiments.

The first correction term to the SIA solution $t_{xz(0)}^D$ in Eq. (24) is $e^2 t_{xz(2)}^D$ in Eq. (27). With $C_\sigma = C_Y e^{1/3}$, at the ice surface and outside the boundary layer

$$e^2 t_{xz(2)}^D \sim e^{7/3}, \quad \tilde{x} + \tilde{z} = 0, \quad (29a)$$

$$e^2 t_{xz(2)}^D \sim e^{8/3}, \quad \tilde{x} + \tilde{z} = O(1), \quad (29b)$$

in agreement with the second terms in the expansions in Eqs. (3.108) and (3.73) in Schoof and Hindmarsh (2010). The velocity component $v_{x(0)}$ in Eq. (4.11) is of $O(e^3)$ everywhere in the ice. The first correction term $v_{x(2)}$ in the Appendix A depends in the same way as $t_{xz(2)}^D$ on σ_{res}^{-1} outside the boundary layer and σ_{res}^{-2} close to the ice surface.

In our numerical experiments we will apply both $\sigma_{\text{res}} = C_\sigma \rho g[H]e$ and $\sigma_{\text{res}} = C_Y \rho g[H]e^{4/3}$, where C_σ and C_Y are constants.

5 Numerical computation of the accuracy of SIA and SOSIA

In this section, we compute the SIA and SOSIA solution for the problem described in Sect. 3, and compute the accuracy by comparing with a full Stokes solution. All our results presented in this section regard the accuracy for the x velocity, v_x , and shear stress, t_{xz}^D . The normal deviatoric stresses t_{xx}^D and t_{zz}^D are not calculated to second order, since this is not necessary to obtain the velocity field. The vertical velocity is

also excluded as it follows directly from the mass balance, which is the same for all orders.

5.1 Method

We are interested in the order of accuracy, i.e. how the error in SIA and SOSIA varies with ϵ . For this purpose we perform repeated simulations using $L = 10, 20, 40, 80, 160, 320, 640, 1280, 2560, 5120$, and 10240 km while keeping H constant, equivalent to varying the aspect ratio between 9.77×10^{-5} and 0.1 . We do this in order to investigate the accuracy of shallow ice approximations in the limit $\epsilon \rightarrow 0$.

The SOSIA Eqs. (15)–(19) are implemented in MATLAB and our implementation follows the standard in SICOPOLIS (Greve, 1995). Finite differences are used on a staggered grid for stability reasons. The velocities, horizontal volume fluxes, vertical shear stresses and the horizontal derivatives of the bedrock topography and the ice-surface topography are defined in between the grid points. All other quantities are defined at grid points. When a quantity is needed in a point where it is not defined, linear interpolation is used. To ensure that the grid points coincide with physical boundaries, a σ -transformation is used (Greve and Blatter, 2009; Greve, 1995). Central differences are applied when possible, otherwise one sided differences are used. Integrals are computed by the trapezoidal method if the integrand and the integral are defined at the same points. Otherwise, the midpoint rule is used.

The full Stokes solution that we compare with is obtained using the finite element code Elmer/ICE (Gagliardini et al., 2013). The same mesh with the nodes on vertical lines was used for SIA, SOSIA and in Elmer/ICE, and we ensured that the mesh was sufficiently fine to make the solutions mesh independent. Even though the singular behaviour of the viscosity does not introduce singularities in the field variables in the full Stokes setting, an extra parameter, the critical shear rate $\dot{\gamma}_o$, is needed in order to treat numerical instabilities at low stress (Råback et al., 2013), occurring e.g. near the ice surface in the shear stress. The critical shear rate is a lower bound for the shear rate $\dot{\gamma}$, which is related to the effective stress by $\dot{\gamma} = \sqrt{2tr(D^2)} = 2d = 2A\sigma^3$. We have

used $\dot{\gamma}_o = 10^{-10}$ throughout the simulations. For large aspect ratios there is no $\dot{\gamma}_o$ which suppresses the numerical instabilities in the shear stress without altering the velocity.

The accuracy is measured in terms of the relative error defined by

$$\frac{\|q_{x,\text{full Stokes}} - q_{x,\text{SIA/SOSIA}}\|_2}{\|q_{x,\text{full Stokes}}\|_2}, \quad (30)$$

5 where q is v_x or t_{xz}^D and $\|\cdot\|_2$ denotes the L_2 -norm defined by

$$\|q\|_2 = \sqrt{\frac{1}{V_\Omega} \int_\Omega q^2 d\Omega}, \quad (31)$$

where V_Ω is the area of Ω . The integral in Eq. (31) is computed on a discrete grid using the trapezoidal rule.

10 Since the assumptions behind the SOSIA are valid only below the boundary layer, we measure the accuracy in horizontal layers, viz. at $0.01[H]$, $0.5[H]$, $0.75[H]$, $0.9[H]$ and $0.95[H]$ mean height above the ice base, to see if the errors due to the boundary layer assumptions in SOSIA spread down into the basal ice.

5.2 Results

15 We know from our analytical solutions in Sect. 4.3 that the SOSIA solution is very sensitive to the parameter σ_{res} . Therefore we experiment with different values of and ways of setting σ_{res} .

20 Our analytical solutions suggest that if the scaling relations used to derive the SOSIA were correct, σ_{res} should vary with ϵ as $C_\sigma \rho g[H]\epsilon$, C_σ being a constant. This is used in Fig. 2a and b, showing the accuracy of the SIA and SOSIA solution for a bumpy bed with relative bump amplitude $\mu = 0.5$. The SIA solution for the horizontal velocity component is $v_{x(0)}$ in Eq. (21) with $C_\sigma = 0$, since a finite viscosity law is unnecessary in this case. The SOSIA solution is computed for C_σ equal to 0.35.

5.2.1 Accuracy of the SIA

We start by analysing the error in SIA. According to classical SIA theory in Baral (1999), Baral et al. (2001), and Greve (1997), the SIA relative error should be of $\mathcal{O}(\epsilon^2)$. However, we know from Ahlkrona et al. (2013) that the scalings used in Schoof and Hindmarsh (2010) are more correct than the classical SIA scalings. The asymptotic expansions in Schoof and Hindmarsh (2010) yield the same zeroth order solution, the SIA, but the correction terms are different. The relative error in the velocity is estimated by the first neglected term in the expansions of v_x in Eqs. (3.73) and (3.108) in Schoof and Hindmarsh (2010) and is of order $\epsilon^{4/3}$ in the boundary layer and $\epsilon^{5/3}$ outside the layer. The estimated slope, $\log(\text{error})/\log(\epsilon)$, of the relative SIA error in Fig. 2a (thick red line with nodes) is 1.43, which is between the theoretical rates in Schoof and Hindmarsh (2010). Similarly, the relative error in the SIA solution of $t_{xz(0)}^D$ should according to Eqs. (3.108) and (3.73) in Schoof and Hindmarsh (2010) and Eqs. (24) and (29) be $\epsilon^{4/3}$ in the boundary layer and $\epsilon^{5/3}$ outside it. This is in good agreement with the slope of the relative error in Fig. 2b which is 1.38.

5.2.2 Accuracy of the SOSIA: $\sigma_{\text{res}} = C_\sigma \rho g[H]\epsilon$

We now move to analysing the SOSIA error in Fig. 2a and b (thick blue line with nodes). According to the classical SIA theory would be correct, SOSIA should in principle be much more accurate than SIA for sufficiently small ϵ . Clearly this is not the case in Fig. 2a and b. In addition to computing the SOSIA solution for $C_\sigma = 0.35$ we also tried $C_\sigma = 0.11$ and $C_\sigma = 1.12$ with no improvement in accuracy.

The SOSIA is thus not a correction to the SIA, for $\sigma_{\text{res}} = C_\sigma \rho g[H]\epsilon$ with C_σ constant. This is because the scaling relations Eq. (8) are not correct in the thick boundary layer near the ice surface which will dominate in the global error. Since the scaling relations do hold below this layer (Ahlkrona et al., 2013; Schoof and Hindmarsh, 2010; Johnson and McMeeking, 1984), one would like to know if the SOSIA solution is more accurate further down in the ice. To investigate this we compute the accuracy of SIA and SOSIA

at horizontal layers at $0.01[H]$, $0.5[H]$, $0.75[H]$, $0.9[H]$ and $0.95[H]$ mean height above the ice base, with the result included in Fig. 2a and b (the thin red and blue lines). The accuracy of the SOSIA velocity is slightly higher deeper in the ice for small ϵ , but even there the SOSIA is less accurate than SIA. The shear stress is more accurate deeper in the ice for both SIA and SOSIA.

5.2.3 Accuracy of the SOSIA: σ_{res} as a lower threshold

The SOSIA solution is very sensitive to σ_{res} , and the different choices of C_σ yield very different results. To limit the sensitivity of σ_{res} we can choose to use it only where the effective stress is too small. We therefore experiment with σ_{res} as a lower bound on the effective stress, viz.:

$$\sigma = \max(\sigma_{(0)}, \sigma_{\text{res}}). \quad (32)$$

Figure 2c and d show the error using this approach for $C_\sigma = 0.11$, 0.35 and 1.12 . The SOSIA velocity error decreases considerably and there are now aspect ratios for which SOSIA is more accurate than SIA when $C_\sigma = 0.35$. The stress t_{xz}^D is not largely affected. We have also computed the accuracy of SOSIA in layers throughout the ice, in the same way as in Fig. 2a and b. For the velocity there is a significant change for small aspect ratios. At $0.01[H]$ mean height over the ice surface the SOSIA velocity solution is more accurate than the SIA solution for all aspect ratios smaller than 10^{-2} . For the deviatoric shear stress the layer-wise error is very similar to what is shown in Fig. 2b.

The error of the velocity for small aspect ratios is mainly due to the term on the second line in Eq. (21). It is an extra term in the zeroth order velocity in SOSIA introduced by the use of a finite viscosity law. This error is reduced by using σ_{res} as a lower threshold and it is the most influential near the ice surface where the scaling relations do not hold and where the zeroth order shear stress and therefore the zeroth order effective stress is zero. This explains why the error decreases further down in the ice. Since there is no such zeroth order term in the vertical shear stress, it is not affected by this type of error. In fact, for small aspect ratios even an extremely large σ_{res} does not cause

a large error in the second order shear stress. The stress is thus less sensitive to the handling of σ_{res} , which can be seen in Figs. 2a–4d. The zeroth order term involving σ_{res} is dominant if C_σ is too large and results in the velocity being overall too high, see Fig. 3a, where $C_\sigma = 1.12$ and $\epsilon = 1/160$.

For larger aspect ratios, the error is of a different character. It arises from an excessive second order correction, resulting in a dip in the velocity and shear stress at $x = 3L/4$, see Fig. 3b for the velocity. This type of error is dominant when ϵ is large and σ_{res} is too small, see Fig. 3a for $C_\sigma = 0.11$ and Fig. 3b, and is difficult to avoid for the largest ϵ . The character of the dip can be understood by studying the dominant term in the second order shear stress, i.e. the last four lines in Eq. (23), or the last two lines in Eq. (27). As this term depends on $(1 - \mu \sin(2\pi x/L))$ it is largest at $x = 3L/4$, and as it is pre-multiplied by μ/C_σ it will in general decrease with increasing C_σ and decrease with decreasing bump amplitude. The error is attributed to the improper handling of the singularity in Glen's flow law, and at the point $x = 3L/4$ the effective stress is zero in both the SOSIA as well as in the full Stokes setting.

The relative error in v_x decays as $\epsilon^{2.81}$ in Fig. 2c for larger ϵ . This is the order of the next term in the ϵ expansion. The reduction of the error ends when the term proportional to the constant σ_{res} with $C_\sigma = 0.35$ in Eq. (21) becomes important. The same observation is valid also for the SOSIA error in the vertical shear stress in Fig. 2d. The decay of the relative error is here $\epsilon^{2.51}$ before the reduction is damped by the constant σ_{res} . With the classical theory in Baral (1999) and Baral et al. (2001), the relative SOSIA error should be $O(\epsilon^3)$. In practice, the larger aspect ratios are of interest. Numerical errors are commonly of the order 10^{-2} , which is higher than the model error in the SIA solution for aspect ratios smaller than 3×10^{-3} . For ice sheet flow, aspect ratios smaller than 10^{-3} are seldom applicable.

GMDD

6, 4281–4325, 2013

The second order shallow ice approximation

J. Ahlkrona et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



5.2.4 Accuracy of the SOSIA: $\sigma_{\text{res}} = C_Y \rho g [H] e^{4/3}$ and further adjustments

Figure 2c and d indicates that the optimal choice of C_σ might not be independent of the aspect ratio. Indeed, we found in Sect. 4 that if we choose σ_{res} as $C_Y \rho g [H] e^{4/3}$ (where $C_\sigma = C_Y e^{1/3}$) the SOSIA correction terms are remedied so that they are consistent with the scalings in Johnson and McMeeking (1984), Schoof and Hindmarsh (2010), and Ahlkrona et al. (2013), see Eq. (29). We can even go further in limiting the influence of σ_{res} by only using it as a lower threshold in the computations where the creep response function is in the denominator (that is in the computation of the normal deviatoric, and normal shear stresses, see Eqs. 26 and 27). The combined results of these two measures are shown in Fig. 4a and b. The SOSIA velocity and shear stress are now more accurate than SIA for all aspect ratios smaller than 10^{-2} with C_Y chosen to be 3. Using for instance $C_Y = 4$ results in SOSIA being more accurate than SIA for even larger aspect ratios, but on the other hand the improvement is no longer as significant. The parameter C_Y depends on the geometry (see Eq. 28) and is thus problem dependent and difficult to determine beforehand. Interesting to note is that there is no difference in accuracy of the velocity in the different layers through the ice. The accuracy of the deviatoric shear stress through the layers is distributed similarly to Fig. 2b.

5.2.5 Low bump amplitude

Lastly, we investigate how the accuracy changes as the bump amplitude is decreased. There is a common perception that shallow ice approximations are more accurate for lower bump amplitudes. Also, we found in Sect. 4 that the terms involving σ_{res} in the stresses and (Eqs. 26 and 27) and velocity Eq. (A7) were pre-multiplied with the relative bump amplitude μ , suggesting that the influence of σ_{res} decreases with decreasing bump amplitude. We have applied the SIA and SOSIA for $\mu = 0.1$ (a bump amplitude of 10 % of mean ice thickness), and the result is illustrated in Fig. 4c and d. We set σ_{res} in the same way as in Fig. 4a and b but with $C_Y = 1.5$ instead of $C_Y = 3$. Indeed, Eq. (28)

GMDD

6, 4281–4325, 2013

The second order shallow ice approximation

J. Ahlkrona et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



The second order shallow ice approximation

J. Ahlkrone et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



shows that C_Y decreases with μ . There is a small improvement of the accuracy of SIA and SOSIA compared to the case when $\mu = 0.5$. For large aspect ratios of almost 0.1 there is a notable improvement in both SIA and SOSIA, but at these large aspect ratios the errors are still very large and neither SIA nor SOSIA is a good model. An explanation of the lack of significant improvement for lower amplitudes is that even if the classical scalings in Eq. (8) do hold for a flat bed, the thick boundary layer where variables rescale develops very rapidly as a small bump is introduced (Ahlkrone et al., 2013). Note that for very small aspect ratios numerical errors are present, so that the error for SIA and SOSIA is not smaller than 10^{-4} . This causes a bend in the curves in the lower left parts of Fig. 4c and d, but also in Figs. 2a–4b.

6 Discussion and conclusions

We have solved the SIA and the SOSIA equations both analytically and numerically, and determined and analysed the accuracy of the velocity, v_x , and the shear stress, t_{xz}^D , compared to the full Stokes equations. This was done for a model problem representing ice flow on a bumpy, sloping bed with no-slip conditions at the base, slightly modified from the ISMIP-HOM B benchmark set-up (Pattyn et al., 2008). The purpose was to determine whether the SOSIA is an improvement on the SIA which can be used in practice. We also wanted to show that a thick, high viscosity boundary layer at the ice surface cannot be overlooked in higher order asymptotic expansions, and that it results in the SIA not being second order accurate.

We know from Ahlkrone et al. (2013), Johnson and McMeeking (1984), and Schoof and Hindmarsh (2010) that the scaling arguments behind the SIA and SOSIA as stated in Baral et al. (2001) are not valid in the boundary layer near the surface. Consequently, the numerically computed accuracy of the SIA is $O(\epsilon^{1.43})$ for the velocity and $O(\epsilon^{1.38})$ for the shear stress instead of $O(\epsilon^2)$, as expected from the classical SIA theory. Our results rather agree with the analysis in Schoof and Hindmarsh (2010), predicting the accuracy of the SIA velocity and shear stress to be $O(\epsilon^{4/3})$ in the boundary layer and

$O(\epsilon^{5/3})$ outside it. Our accuracy was measured over the whole domain including the boundary layer.

The neglect of the special boundary layer dynamics in the classical SIA scalings results in singularities in the SOSIA field variables. To remedy this the finite viscosity parameter, σ_{res} , is added to the effective stress in the creep response function when computing the SOSIA. Both our analytical solutions and numerical experiments show that the SOSIA solution is very sensitive to this parameter. It can neither be chosen too large nor too small, since it appears in both the numerator and the denominator of the expressions. Too large a σ_{res} yields too high a velocity, especially for small aspect ratios. This is due to an extra zeroth order term in the zeroth order velocity. This error is more important near the surface than further down in the ice. If σ_{res} is simply added to the creep response function as $f(\sigma) = \sigma^2 + \sigma_{\text{res}}^2$, the error is so large that the SOSIA velocity is less accurate than the SIA for all aspect ratios. If σ_{res} is instead used as a lower threshold for the effective stress, as in Eq. (32), the error is decreased so that there is a range of (quite small) aspect ratios for which SOSIA is an improvement of SIA. Further reducing the effect of σ_{res} , so that it is only used in the calculations where singularities would arise otherwise, improves the accuracy even more. The shear stress is not as sensitive to too large a σ_{res} as the velocity is. Too small a σ_{res} yields errors in both the velocity and shear stress for large aspect ratios, due to excessive second order corrections caused by the singularity in Glen's flow law.

Depending on how we choose the value of σ_{res} , the analytical expressions for the field variables conform with either the classical shallow ice scalings that were used in the derivation, or with the scaling relations in Schoof and Hindmarsh (2010) which we know are accurate (Ahlkrona et al., 2013). In order to agree with the classical SIA scalings we conclude that σ_{res} should be $C_\sigma \rho g [H] \epsilon$ where C_σ is a constant. In order for the solution to scale with the aspect ratio as in Schoof and Hindmarsh (2010) (both inside and outside the boundary layer), σ_{res} should be $C_\gamma \rho g [H] \epsilon^{4/3}$, where C_γ is a constant. In our numerical experiments we find that $\sigma_{\text{res}} = C_\gamma \rho g [H] \epsilon^{4/3}$ yields more accurate

The second order shallow ice approximation

J. Ahlkrona et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



results than $\sigma_{\text{res}} = C_{\sigma} \rho g [H] \epsilon$, in the sense that the former choice, combined with using σ_{res} as a lower threshold where singularities occur, makes SOSIA more accurate than SIA even for aspect ratios almost as large as 0.1.

Our analytical solutions suggest that the sensitivity to σ_{res} decreases when the amplitude of the bumps at the ice base decreases. In our numerical experiments, lowering the bump amplitude from half of the mean ice thickness to 10 % of the mean thickness results in a slight improvement of the accuracy of both SIA and SOSIA for small aspect ratios. For large aspect ratios around 0.1 the improvement is significant, but for these aspect ratios the relative error is too large, about 0.1, for the models to be applicable.

As SOSIA is almost as computationally cheap as the SIA it could be a convenient tool in e.g. paleoglacial simulations. Since the SIA is accurate for very small aspect ratios, and as the aspect ratio of a real ice sheets would not be less than 10^{-3} , the potential applicability of SOSIA is for aspect ratios larger than that. SOSIA is an improvement of SIA for aspect ratios as large as 5×10^{-2} depending on how the finite viscosity parameter σ_{res} is set and on the amplitude of the bedrock bumps. A relative error at 1 % is quite good and SOSIA achieves that for larger ϵ than SIA does. A crucial issue to overcome is how to determine σ_{res} , or C_{γ} and C_{σ} . These parameters are problem dependent and the accuracy is sensitive to their values. It is unclear how to choose them in practice for a more complicated geometry and forcings than treated in this paper.

Appendix A

Analytical solutions to the SIA and SOSIA

This Appendix contains analytical solutions for $v_{x(0)}$, $v_{z(0)}$, $t_{xx(0)}^D$, $t_{xz(2)}^D$, $p_{(2)}$ and $v_{x(2)}$ for an isothermal, diagnostic, 2-D-problem with no slip conditions at the impermeable ice base, an inclined plane as ice surface, and a zeroth order ice base. A finite viscosity law

GMDD

6, 4281–4325, 2013

The second order shallow ice approximation

J. Ahlkrone et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



of the form $f(\sigma) = \sigma^2 + \sigma_{\text{res}}^2$ is used, as discussed in Sect. 2.3. For the model problem in Sect. 3 we also express the solutions in terms of L , α , μ , and σ_{res} .

A1 Zeroth order x velocity, $v_{x(0)}$

Computing the integral in the second equation in Eq. (15) gives

$$v_{x(0)} = -2\rho g A \frac{\partial h_{(0)}}{\partial x} \left(\frac{\left(\rho g \frac{\partial h_{(0)}}{\partial x} \right)^2}{4} \left(H^4 - (h_{(0)} - z)^4 \right) + \frac{\sigma_{\text{res}}^2}{2} \left(H^2 - (h_{(0)} - z)^2 \right) \right). \quad (\text{A1})$$

Inserting the expressions for h , b (see Eq. 14) and $H(= h - b)$ yields Eq. (21).

A2 Zeroth order z velocity, $v_{z(0)}$

The z velocity $v_{z(0)}$ is given by

$$\begin{aligned} v_{z(0)} &= \int_{b_{(0)}}^z \frac{\partial v_{x(0)}}{\partial x} dz' \\ &= 2A \frac{\partial h_{(0)}}{\partial x} \left((\rho g)^3 \left(\frac{\partial h_{(0)}}{\partial x} \right)^2 \left(H_{(0)}^3 \frac{\partial H_{(0)}}{\partial x} (z - b_{(0)}) + \frac{1}{4} \frac{\partial h_{(0)}}{\partial x} \left((h_{(0)} - z)^4 - H_{(0)}^4 \right) \right) \right. \\ &\quad \left. + \rho g \sigma_{\text{res}}^2 \left(H_{(0)} \frac{\partial H_{(0)}}{\partial x} (z - b_{(0)}) + \frac{1}{2} \frac{\partial h_{(0)}}{\partial x} \left((h_{(0)} - z)^2 - H_{(0)}^2 \right) \right) \right). \end{aligned} \quad (\text{A2})$$

Inserting the expressions for h , b and H (see Eq. 14) yields

$$v_{z(0)} = 2A(\rho g)^3 \tan^3(\alpha) \left([H]^4 (1 - \mu \sin(2\pi x/L))^3 \mu \frac{2\pi}{L} \cos(2\pi x/L) \right)$$

$$\cdot (z + x \tan(\alpha) + [H] (1 - \mu \sin(2\pi x/L))) + \frac{1}{4} \tan(\alpha) (x \tan(\alpha) + z)^4 - [H]^4 (1 - \mu \sin(2\pi x/L))^4 \quad (\text{A3})$$

$$+ 2A\rho g \sigma_{\text{res}}^2 \tan(\alpha) \cdot \left([H]^2 (1 - \mu \sin(2\pi x/L)) \mu \frac{2\pi}{L} \cos(2\pi x/L) (z + x \tan(\alpha) + [H] (1 - \mu \sin(2\pi x/L))) \right).$$

A3 Zeroth order normal deviatoric stress, $t_{xx(0)}^D$

Inserting Eq. (A1) in Eq. (16) gives

$$t_{xx(0)}^D = \frac{-2\rho g \frac{\partial h_{(0)}}{\partial x}}{\left(\left((h_{(0)} - z) \frac{\partial h_{(0)}}{\partial x} \right)^2 + \left(\frac{\sigma_{\text{res}}}{\rho g} \right)^2 \right)} \left(\left(\frac{\partial h_{(0)}}{\partial x} \right)^2 \left(H^3 \frac{\partial H_{(0)}}{\partial x} - \frac{\partial h_{(0)}}{\partial x} (h_{(0)} - z)^3 \right) + \left(\frac{\sigma_{\text{res}}}{\rho g} \right)^2 \left(H \frac{\partial H_{(0)}}{\partial x} - \frac{\partial h_{(0)}}{\partial x} (h_{(0)} - z) \right) \right). \quad (\text{A4})$$

Inserting the expressions for h , b and H (Eq. 14) yields Eq. (22).

GMDD

6, 4281–4325, 2013

The second order shallow ice approximation

J. Ahlkrona et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



A4 Second order vertical shear stress, $t_{xz(2)}^D$

Inserting Eq. (A4) in Eq. (18) knowing that $t_{zz}^D = -t_{xx}^D$ gives

$$\begin{aligned} \epsilon^2 t_{xz(2)}^D = & 3\rho g \left(\frac{\partial h_{(0)}}{\partial x} \right)^3 (h_{(0)} - z) \\ & - 4\rho g \frac{H_{(0)} \frac{\partial H_{(0)}}{\partial x} \left(\frac{\partial h_{(0)}}{\partial x} \right)^2 \left(\left(\frac{\partial h_{(0)}}{\partial x} \right)^2 H_{(0)}^2 + \frac{\sigma_{\text{res}}^2}{(\rho g)^2} \right)}{\left(\frac{\partial h_{(0)}}{\partial x} \right)^2 (h_{(0)} - z)^2 + \frac{\sigma_{\text{res}}^2}{(\rho g)^2}} \\ & - \frac{4(\rho g)^2}{\sigma_{\text{res}}} \left(3H^2 \left(\frac{\partial H_{(0)}}{\partial x} \right)^2 \left(\frac{\partial h_{(0)}}{\partial x} \right)^2 \right. \\ & + H^3 \frac{\partial^2 H}{\partial x^2} \left(\frac{\partial h_{(0)}}{\partial x} \right)^2 + \frac{\sigma_{\text{res}}^2}{(\rho g)^2} \left(\left(\frac{\partial h_{(0)}}{\partial x} \right)^2 + H_{(0)} \frac{\partial^2 H_{(0)}}{\partial x^2} \right) \\ & \left. \cdot \arctan \left(\rho g \frac{\frac{\partial h_{(0)}}{\partial x} (h_{(0)} - z)}{\sigma_{\text{res}}} \right) \right) \end{aligned} \quad (\text{A5})$$

5 Inserting the expressions for h , b and H (Eq. 14) yields Eq. (23).

A5 Second order pressure, $p_{(2)}$

The calculation of the second order pressure from Eq. (17) is straight forward, knowing that $t_{zz}^D = -t_{xx}^D$ and using Eqs. (A4) and (22).

A6 Second order x velocity, $v_{x(2)}$

Inserting the expressions for $t_{xz(0)}^D$, $t_{xx(0)}^D$, and $t_{xz(2)}^D$ in Eq. (19) gives

$$\begin{aligned} \frac{\epsilon^2 v_{x(2)}}{\mathcal{A}} = & -(z - b_{(0)})^2 \left(\left(\rho g \frac{\partial h_{(0)}}{\partial x} \right)^3 \left(3H_{(0)}^2 \left(\frac{\partial H_{(0)}}{\partial x} \right)^2 + H_{(0)}^3 \frac{\partial^2 H_{(0)}}{\partial x^2} \right) \right. \\ & \left. + \rho g \frac{\partial h_{(0)}}{\partial x} \sigma_{\text{res}}^2 \left(\left(\frac{\partial H_{(0)}}{\partial x} \right)^2 + H_{(0)} \frac{\partial^2 H_{(0)}}{\partial x^2} \right) \right) \\ & - 4(h_{(0)} - z)^2 \frac{\partial h_{(0)}}{\partial x} (\rho g)^3 \left(3H_{(0)}^2 \left(\frac{\partial H_{(0)}}{\partial x} \right)^2 \left(\frac{\partial h_{(0)}}{\partial x} \right)^2 + H_{(0)}^3 \frac{\partial^2 H_{(0)}}{\partial x^2} \left(\frac{\partial h_{(0)}}{\partial x} \right)^2 \right. \\ & \left. + \frac{\sigma_{\text{res}}^2}{(\rho g)^2} \left(\left(\frac{\partial H_{(0)}}{\partial x} \right)^2 + H_{(0)} \frac{\partial^2 H_{(0)}}{\partial x^2} \right) \right) + 4(h_{(0)} - z)^2 \sigma_{\text{res}}^2 \rho g \left(\frac{\partial h_{(0)}}{\partial x} \right)^3 \\ & + (b_{(0)} - z) \left(6(\rho g)^3 \left(\frac{\partial h_{(0)}}{\partial x} \right)^4 H_{(0)}^3 \frac{\partial H_{(0)}}{\partial x} + 6\rho g \sigma_{\text{res}}^2 \left(\frac{\partial h_{(0)}}{\partial x} \right)^2 H_{(0)} \frac{\partial H_{(0)}}{\partial x} \right. \\ & \left. - 2 \left(\rho g \frac{\partial h_{(0)}}{\partial x} \right)^3 H_{(0)}^3 \frac{\partial H_{(0)}}{\partial x} \frac{\partial b_{(0)}}{\partial x} \right) \end{aligned}$$

$$\begin{aligned}
& + 8 \frac{(\rho g)^4}{\sigma_{\text{res}}} \left(\frac{\partial h_{(0)}}{\partial x} \right)^2 \left(3H^2 \left(\frac{\partial H_{(0)}}{\partial x} \right)^2 \left(\frac{\partial h_{(0)}}{\partial x} \right)^2 + H^3 \frac{\partial^2 H}{\partial x^2} \left(\frac{\partial h_{(0)}}{\partial x} \right)^2 \right. \\
& + \left. \frac{\sigma_{\text{res}}^2}{(\rho g)^2} \left(\left(\frac{\partial H_{(0)}}{\partial x} \right)^2 + H_{(0)} \frac{\partial^2 H_{(0)}}{\partial x^2} \right) \right) (h_{(0)} - z)^3 \arctan \left(\frac{\frac{\partial h_{(0)}}{\partial x} \rho g}{\sigma_{\text{res}}} (h_{(0)} - z) \right) \\
& - 4(\rho g)^3 \frac{\partial h_{(0)}}{\partial x} \frac{\left(\left(\frac{\partial h_{(0)}}{\partial x} \right)^2 H^3 \frac{\partial H_{(0)}}{\partial x} + \frac{\sigma_{\text{res}}^2}{(\rho g)^2} H \frac{\partial H_{(0)}}{\partial x} \right)^2}{\left(\frac{\partial h_{(0)}}{\partial x} \right)^2 (h_{(0)} - z)^2 + \frac{\sigma_{\text{res}}^2}{(\rho g)^2}} \\
& - 2(\rho g)^3 \left(\frac{\partial h_{(0)}}{\partial x} \right)^5 (h_{(0)} - z)^4 \\
& + 8\sigma_{\text{res}}(\rho g)^2 \left(3H^2 \left(\frac{\partial H_{(0)}}{\partial x} \right)^2 \left(\frac{\partial h_{(0)}}{\partial x} \right)^2 + H^3 \frac{\partial^2 H}{\partial x^2} \left(\frac{\partial h_{(0)}}{\partial x} \right)^2 \right. \\
& + \left. \frac{\sigma_{\text{res}}^2}{(\rho g)^2} \left(\left(\frac{\partial H_{(0)}}{\partial x} \right)^2 + H_{(0)} \frac{\partial^2 H_{(0)}}{\partial x^2} \right) \right) (h_{(0)} - z) \arctan \left(\frac{\frac{\partial h_{(0)}}{\partial x} \rho g}{\sigma_{\text{res}}} (h_{(0)} - z) \right) \\
& + 2\rho g \sigma_{\text{res}}^2 \frac{\partial h_{(0)}}{\partial x} H_{(0)} \frac{\partial H_{(0)}}{\partial x} \frac{\partial b_{(0)}}{\partial x} (z - b_{(0)})
\end{aligned}$$

The second order shallow ice approximation

J. Ahlkrona et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



$$\begin{aligned}
 & -\frac{8(\rho g)^2}{\sigma_{\text{res}}} \left(\left(\frac{\partial h_{(0)}}{\partial x} \right)^2 (\rho g)^2 H_{(0)}^2 + \sigma_{\text{res}}^2 \right) \left(3H^2 \left(\frac{\partial H_{(0)}}{\partial x} \right)^2 \left(\frac{\partial h_{(0)}}{\partial x} \right)^2 \right. \\
 & + H^3 \frac{\partial^2 H}{\partial x^2} \left(\frac{\partial h_{(0)}}{\partial x} \right)^2 + \frac{\sigma_{\text{res}}^2}{(\rho g)^2} \left(\left(\frac{\partial H_{(0)}}{\partial x} \right)^2 + H_{(0)} \frac{\partial^2 H_{(0)}}{\partial x^2} \right) \left. H_{(0)} \arctan \left(\frac{\frac{\partial h_{(0)}}{\partial x} \rho g}{\sigma_{\text{res}}} H_{(0)} \right) \right) \\
 & + 2(\rho g)^3 \left(\frac{\partial h_{(0)}}{\partial x} \right)^5 H_{(0)}^4 + 16(\rho g)^3 H_{(0)}^4 \left(\frac{\partial H_{(0)}}{\partial x} \right)^2 \left(\frac{\partial h_{(0)}}{\partial x} \right)^3 \\
 & + 8\rho g \frac{\partial h_{(0)}}{\partial x} H_{(0)}^2 \left(\frac{\partial H_{(0)}}{\partial x} \right)^2 \sigma_{\text{res}}^2 - 4\sigma_{\text{res}}^2 \rho g \left(\frac{\partial h_{(0)}}{\partial x} \right)^3 H_{(0)}^2 \\
 & + 4(\rho g)^3 H_{(0)}^5 \frac{\partial^2 H}{\partial x^2} \left(\frac{\partial h_{(0)}}{\partial x} \right)^3 + 4\rho g \sigma_{\text{res}}^2 \frac{\partial h_{(0)}}{\partial x} H_{(0)}^3 \frac{\partial^2 H_{(0)}}{\partial x^2}.
 \end{aligned} \tag{A6}$$

Inserting the expressions for h , b and H (i.e. Eq. 14) yields

$$\begin{aligned}
 \frac{\epsilon^2 v_{x(2)}}{A} &= \left(\frac{2\pi}{L} \right)^2 \mu [H]^3 \left(z + x \tan(\alpha) - (\mu \sin(2\pi x/L) - 1) \right)^2 \\
 &\cdot \left((\rho g)^3 \tan^3(\alpha) [H]^2 \left(3\mu (1 - \mu \sin(2\pi x/L))^2 \cos^2(2\pi x/L) \right. \right. \\
 &+ \left. \left. (1 - \mu \sin(2\pi x/L))^3 \sin(2\pi x/L) \right) + \rho g \sigma_{\text{res}}^2 \tan(\alpha) \left(\mu \cos^2(2\pi x/L) \right. \right. \\
 &+ \left. \left. (1 - \mu \sin(2\pi x/L)) \sin(2\pi x/L) \right) \right)
 \end{aligned}$$

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**The second order
shallow ice
approximation**

J. Ahlkrone et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

$$\begin{aligned}
 & + 4(\rho g)^3 [H]^4 \left(\frac{2\pi}{L} \right)^2 \mu \tan(\alpha) (x \tan(\alpha) + z)^2 \left(3(1 - \mu \sin(2\pi x/L))^2 \mu \cos^2(2\pi x/L) \tan^2(\alpha) \right. \\
 & + (1 - \mu \sin(2\pi x/L))^3 \sin(2\pi x/L) \tan^2(\alpha) \\
 & + \left. \frac{\sigma_{\text{res}}^2}{(\rho g)^2} \left(\mu \cos^2(2\pi x/L) + (1 - \mu \sin(2\pi x/L)) \sin(2\pi x/L) \right) \right) \\
 & - 4\sigma_{\text{res}}^2 \rho g \mu \frac{2\pi}{L} \tan^3(\alpha) (x \tan(\alpha) + z)^2 (x \tan(\alpha) - [H] (\mu \sin(2\pi x/L) - 1) + z) \cos(2\pi x/L) \\
 & \cdot \left(6[H]^4 (\rho g)^3 \tan^4(\alpha) (1 - \mu \sin(2\pi x/L))^3 - 6[H]^2 \rho g \sigma_{\text{res}}^2 \tan^2(\alpha) (1 - \mu \sin(2\pi x/L)) \right. \\
 & - \left. 2[H]^4 (\rho g \tan(\alpha))^3 (1 - \mu \sin(2\pi x/L))^3 \left(-\tan(\alpha) + [H] \frac{2\pi}{L} \mu \cos(2\pi x/L) \right) \right) \\
 & - 8(\rho g)^4 \frac{[H]^4}{\sigma_{\text{res}}} \left(\frac{2\pi}{L} \right)^2 \mu \tan^2(\alpha) \left(3(1 - \mu \sin(2\pi x/L))^2 \mu \cos^2(2\pi x/L) \tan^2(\alpha) \right. \\
 & + (1 - \mu \sin(2\pi x/L))^3 \sin(2\pi x/L) \tan^2(\alpha) \\
 & + \left. \frac{\sigma_{\text{res}}^2}{(\rho g)^2} \left(\mu \cos^2(2\pi x/L) + (1 - \mu \sin(2\pi x/L)) \sin(2\pi x/L) \right) \right) \\
 & \cdot (x \tan(\alpha) + z)^3 \arctan \left(\frac{\tan(\alpha) \rho g}{\sigma_{\text{res}}} (x \tan(\alpha) + z) \right)
 \end{aligned}$$

$$\begin{aligned}
& + 4(\rho g)^3 \left(\frac{2\pi}{L} \right)^2 \mu^2 [H]^4 \tan(\alpha) \\
& \cdot \frac{\cos^2(2\pi x/L) (1 - \mu \sin(2\pi x/L))^2 \left(\tan^2(\alpha) [H]^2 (1 - \mu \sin(2\pi x/L))^2 + \frac{\sigma_{\text{res}}^2}{(\rho g)^2} \right)^2}{\tan^2(\alpha) (x \tan(\alpha) + z)^2 + \frac{\sigma_{\text{res}}^2}{(\rho g)^2}} \\
& + 2(\rho g)^3 \tan^5(\alpha) (x \tan(\alpha) + z)^4 \\
& - 8\sigma_{\text{res}} (\rho g)^2 [H]^4 \left(\frac{2\pi}{L} \right)^2 \mu \left(3\mu (1 - \mu \sin(2\pi x/L))^2 \cos^2(2\pi x/L) \tan^2(\alpha) \right. \\
& + (1 - \mu \sin(2\pi x/L))^3 \sin(2\pi x/L) \tan^2(\alpha) \\
& + \left. \frac{\sigma_{\text{res}}^2}{(\rho g)^2} \left(\mu \cos^2(2\pi x/L) + (1 - \mu \sin(2\pi x/L)) \sin(2\pi x/L) \right) \right) (x \tan(\alpha) + z) \\
& \cdot \arctan \left(\frac{\tan(\alpha) \rho g}{\sigma_{\text{res}}} (x \tan(\alpha) + z) \right) \\
& + \frac{4\pi}{L} \mu \rho g \sigma_{\text{res}}^2 [H]^2 \tan(\alpha) (1 - \mu \sin(2\pi x/L)) \cos(2\pi x/L) \\
& \cdot \left(-\tan(\alpha) + [H] \frac{2\pi}{L} \mu \cos(2\pi x/L) \right) \\
& \cdot (z + x \tan(\alpha) - [H](\mu \sin(2\pi x/L) - 1))
\end{aligned}$$

The second order shallow ice approximation

J. Ahlkrona et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



The second order shallow ice approximation

J. Ahlkrona et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



and Thies, J.: Capabilities and performance of Elmer/Ice, a new generation ice-sheet model, Geosci. Model Dev. Discuss., 6, 1689–1741, doi:10.5194/gmdd-6-1689-2013, 2013. 4284, 4301

Greve, R.: Thermomechanisches Verhalten polythermer Eisschilde – Theorie, Analytik, Numerik, Ph. D. thesis, Department of Mechanics (III), Technical University Darmstadt, Germany, 1995. 4301

Greve, R.: A continuum-mechanical formulation for shallow polythermal ice sheets, Philos. T. Roy. Soc. A, 355, 921–974, 1997. 4283, 4287, 4291, 4293, 4303

Greve, R. and Blatter, H.: Dynamics of Ice Sheets and Glaciers, Advances in Geophysical and Environmental Mechanics and Mathematics (AGEM²), Springer, Berlin, 2009. 4301

Hutter, K.: Theoretical Glaciology, D. Reidel Publishing Company, Terra Scientific Publishing Company, Dordrecht, 1983. 4283

Johnson, R. E. and McMeeking, R. M.: Near-surface flow in glaciers obeying Glen's law, Quart. J. Mech. Appl. Math., 7, 273–291, 1984. 4284, 4290, 4299, 4303, 4306, 4307

Kirchner, N., Hutter, K., Jakobsson, M., and Gyllencreutz, R.: Capabilities and limitations of numerical ice sheet models: a discussion for Earth-scientists and modelers, Quaternary Sci. Rev., 30, 3691–3704, 2011. 4284

Lliboutry, L. A.: The dynamics of temperate glaciers from the detailed viewpoint, J. Glaciol., 8, 185–205, 1969. 4291

MacAyeal, D. R.: The basal stress distribution of Ice Stream E, Antarctica, inferred by control methods, J. Geophys. Res., 97, 595–603, 1992. 4290

Mangeney, A. and Califano, F.: The shallow-ice approximation for anisotropic ice – Formulation and limits, J. Geophys. Res., 103, 691–705, 1998. 4284

Morland, L. W.: Thermo-mechanical balances of ice sheet flows, Geophys. Astrophys. Fluid Dynam., 29, 237–266, 1984. 4283

Pattyn, F., Perichon, L., Aschwanden, A., Breuer, B., de Smedt, B., Gagliardini, O., Gudmundsson, G. H., Hindmarsh, R. C. A., Hubbard, A., Johnson, J. V., Kleiner, T., Konovalov, Y., Martin, C., Payne, A. J., Pollard, D., Price, S., Rückamp, M., Saito, F., Souček, O., Sugiyama, S., and Zwinger, T.: Benchmark experiments for higher-order and full-Stokes ice sheet models (ISMIP-HOM), The Cryosphere, 2, 95–108, doi:10.5194/tc-2-95-2008, 2008. 4283, 4292, 4307

Pattyn, F., Schoof, C., Perichon, L., Hindmarsh, R. C. A., Bueler, E., de Fleurian, B., Durand, G., Gagliardini, O., Gladstone, R., Goldberg, D., Gudmundsson, G. H., Huybrechts, P., Lee, V.,

Nick, F. M., Payne, A. J., Pollard, D., Rybak, O., Saito, F., and Vieli, A.: Results of the Marine Ice Sheet Model Intercomparison Project, MISMP, The Cryosphere, 6, 573–588, doi:10.5194/tc-6-573-2012, 2012. 4283

Råback, P., Malinen, M., Ruokalainen, J., Pursula, A., and Zwinger, T.: Elmer Models Manual, CSC – IT Center for Science, Helsinki, Finland, 2013. 4301

Schoof, C. and Hindmarsh, R.: Thin-film flows with wall slip: an asymptotic analysis of higher order glacier flow models, Quart. J. Mech. Appl. Math., 63, 73–114, 2010. 4284, 4290, 4299, 4300, 4303, 4306, 4307, 4308

Solomon, S., Qin, D., Manning, M., Chen, Z., Marquis, M., Averyt, K., Tignor, M., and Miller, H.: The physical science basis, in: Contribution of Working Group I to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change 2007, Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, 2007. 4282

GMDD

6, 4281–4325, 2013

The second order shallow ice approximation

J. Ahlkrona et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion






Fig. 1. Model set-up showing the basal topography and the ice surface. The ice flows down-slope in the positive x direction.




Fig. 2. Relative error of horizontal velocity v_x and vertical shear stress t_{xz}^D with $\sigma_{res} = C_\sigma \rho g [H] \epsilon$ for both SIA (red) and SOSIA (blue). Thick lines show the error measured over the whole domain and the thin lines in the upper two panels show the errors measured over horizontal layers.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



The second order shallow ice approximation

J. Ahlkrona et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion




Fig. 3. Surface x velocity for the full Stokes solution, the SIA and the SOSIA. For SOSIA, $\sigma_{\text{res}} = C_{\sigma} \rho g [H] \epsilon$ is used as a lower threshold for the effective stress.

The second order
shallow ice
approximation

J. Ahlkrona et al.




Fig. 4. Relative error of horizontal velocity v_x and vertical shear stress t_{xz}^D for both SIA (red) and SOSIA (blue) for $\mu = 0.5$ and $\mu = 0.1$. $\sigma_{\text{res}} = C_\gamma \rho g [H] e^{4/3}$ is only used when needed in the denominator of the expressions.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

