



This discussion paper is/has been under review for the journal Geoscientific Model Development (GMD). Please refer to the corresponding final paper in GMD if available.

# Development of two-moment cloud microphysics for liquid and ice within the NASA Goddard earth observing system model (GEOS-5)

D. Barahona<sup>1</sup>, A. Molod<sup>1,2</sup>, J. Bacmeister<sup>3</sup>, A. Nenes<sup>4</sup>, A. Gettelman<sup>3</sup>,  
H. Morrison<sup>3</sup>, V. Phillips<sup>5</sup>, and A. Eichmann<sup>1,6</sup>

<sup>1</sup>Global Modeling and Assimilation Office, NASA Goddard Space Flight Center, Greenbelt, MD, USA

<sup>2</sup>University of Maryland, College Park, MD, USA

<sup>3</sup>National Center for Atmospheric Research, Boulder, CO, USA

<sup>4</sup>School of Earth and Atmospheric Sciences and School of Chemical and Biomolecular Engineering, Georgia Institute of Technology, Atlanta, GA, USA

<sup>5</sup>School of Earth and Environment, University of Leeds, Leeds, UK

<sup>6</sup>Science Systems and Applications, Inc., Lanham, MD, USA

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Received: 6 September 2013 – Accepted: 23 September 2013 – Published: 29 October 2013

Correspondence to: D. Barahona (donifan.o.barahona@nasa.gov)

Published by Copernicus Publications on behalf of the European Geosciences Union.

# GMDD

6, 5289–5373, 2013

## Cloud microphysics in GEOS-5

D. Barahona et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## Abstract

This work presents the development of a two-moment cloud microphysics scheme within the version 5 of the NASA Goddard Earth Observing System (GEOS-5). The scheme includes the implementation of a comprehensive stratiform microphysics module, a new cloud coverage scheme that allows ice supersaturation and a new microphysics module embedded within the moist convection parameterization of GEOS-5. Comprehensive physically-based descriptions of ice nucleation, including homogeneous and heterogeneous freezing, and liquid droplet activation are implemented to describe the formation of cloud particles in stratiform clouds and convective cumulus. The effect of preexisting ice crystals on the formation of cirrus clouds is also accounted for. A new parameterization of the subgrid scale vertical velocity distribution accounting for turbulence and gravity wave motion is developed. The implementation of the new microphysics significantly improves the representation of liquid water and ice in GEOS-5. Evaluation of the model shows agreement of the simulated droplet and ice crystal effective and volumetric radius with satellite retrievals and in situ observations. The simulated global distribution of supersaturation is also in agreement with observations. It was found that when using the new microphysics the fraction of condensate that remains as liquid follows a sigmoidal increase with temperature which differs from the linear increase assumed in most models and is in better agreement with available observations. The performance of the new microphysics in reproducing the observed total cloud fraction, longwave and shortwave cloud forcing, and total precipitation is similar to the operational version of GEOS-5 and in agreement with satellite retrievals. However the new microphysics tends to underestimate the coverage of persistent low level stratocumulus. Sensitivity studies showed that the simulated cloud properties are robust to moderate variation in cloud microphysical parameters. However significant sensitivity in ice cloud properties was found to variation in the dispersion of the ice crystal size distribution and the critical size for ice autoconversion. The implementation of the new microphysics leads to a more

## Cloud microphysics in GEOS-5

D. Barahona et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



realistic representation of cloud processes in GEOS-5 and allows the linkage of cloud properties to aerosol emissions.

## 1 Introduction

Cloud microphysical schemes in global circulation models (GCMs) have evolved from directly prescribing cloud properties (i.e., particle size and number, cloud amount and concentration of condensate) to explicit representation of the formation, evolution, and removal of cloud droplets and ice crystals (e.g., Gettelman et al., 2010; Lohmann, 2008; Sud et al., 2013). The development of sophisticated cloud microphysics schemes allows a more realistic description of the variability and interdependence of cloud properties, and will likely improve model predictions of climate (Lohmann and Feichter, 2005). However their increased complexity has also brought about new challenges in the description of small-scale dynamics, cloud particle nucleation, and the generation of precipitation. Most models rely on simplified representations of such processes.

Current GCMs typically use either single- (e.g., Del Genio et al., 1996; Bacmeister et al., 1999) or two-moment cloud microphysics schemes (e.g., Gettelman et al., 2010; Sud et al., 2013; Lohmann et al., 2008). More detailed schemes have also been developed, however their computational expense make them unsuitable for climate studies (Khain et al., 2000). The advantage of two and higher moment schemes is that cloud particle size is explicitly calculated and allowed to interact with radiation and the formation of precipitation. Some schemes also allow for supersaturation with respect to the ice phase, required to explicitly model ice nucleation (e.g., Gettelman et al., 2010; Wang and Penner, 2010). When coupled to an appropriate aerosol activation parameterization, two-moment microphysics schemes are capable of modeling the modification of cloud properties by aerosol emissions, an effect that has important implications for the evolution of climate (IPCC, 2007; Lohmann and Feichter, 2005).

Mounting evidence suggests that aerosols, both natural and anthropogenic, play a key role in many atmospheric processes. For example, the presence of ice in

## Cloud microphysics in GEOS-5

D. Barahona et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**Cloud microphysics  
in GEOS-5**

D. Barahona et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

clouds at temperatures above 235 K depends on the presence of water-insoluble ice nuclei (IN) (Pruppacher and Klett, 1997). IN in turn act as precipitation-forming agents in convective systems and mixed-phase clouds (Ramanathan et al., 2001; Rosenfeld and Woodley, 2000). Although they originate mostly from natural sources (i.e., dust and biogenic material), anthropogenic IN emissions can modify the natural IN concentration. The effect of aerosols on clouds has also been associated with planetary radiative perturbations from the modification of clouds by anthropogenic aerosol emissions (Twomey, 1977, 1991; Lohmann and Feichter, 2005). Emissions of cloud condensation nuclei (CCN) may also lead to the modification of the precipitation onset in convective cumulus by decreasing the average size of cloud droplets (Rosenfeld et al., 2008). Recent studies suggest that the interplay between CCN and IN plays a significant role in the maintenance of Arctic clouds (Morrison et al., 2012; Lance et al., 2011). Accurate representation of these effects in atmospheric models is critical for reliable climate prediction, yet difficult due to their complexity and gaps on the understanding of CCN and IN activation.

A recent simulation of the non-hydrostatic implementation of the NASA Goddard Earth Observing System at 14 km spatial resolution demonstrated that as the spatial resolution increases the parameterized convective transport of moisture plays a weaker role in the generation of cloud condensate. At high resolution the simulated cloud properties are controlled by the cloud microphysics (Putman and Suarez, 2011). For typical GCM resolutions ( $\sim 2^\circ$ ) the parameterization of the convective generation of precipitation is critical for the correct simulation of the hydrological cycle and the distribution of cloud tracers in the atmosphere (Arakawa, 2004). Most GCMs use single-moment schemes to describe the microphysics of convective systems. Two-moment microphysical schemes have also been developed for convective clouds, although mostly based on ideas originally developed for stratiform clouds (e.g., Lohmann, 2008; Song and Zhang, 2011; Sud et al., 2013).

The NASA Goddard Earth Observing System, Version 5 (GEOS-5) is a system of models integrated using the Earth System Modeling Framework (ESMF) (Rienecker

et al., 2008). The operational version of GEOS-5 is regularly used for decadal predictions of climate, field campaign support, satellite data assimilation, weather forecasts and basic research (Rienecker et al., 2008, 2011; Molod, 2012). GEOS-5 uses a single-moment cloud microphysics scheme to parameterize condensation, sublimation, evaporation, autoconversion and sedimentation of liquid and ice (Bacmeister et al., 2006). This single-moment approach captures the main climatic features related to the formation of stratocumulus decks and tropical storms (Reale et al., 2009; Putman and Suarez, 2011). However the single-moment approach prevents the explicit linkage of aerosol emissions to cloud properties and omits sub-grid variability in cloud properties. In this work we develop a new microphysical package for GEOS-5 that addresses these issues. The new two-moment cloud microphysics scheme explicitly predicts the mass and number of cloud ice and liquid, rain and snow and links the number concentration of ice crystals and cloud droplets to processes of cloud droplet activation and ice crystal nucleation.

## 2 Model description

### 2.1 Operational GEOS-5

The cloud scheme in the operational version of GEOS-5 considers a single phase of condensate, however the removal and evaporation of cloud water from detrained convection and in situ condensation are treated separately. The fraction of condensate existing as ice is assumed to linearly increase between 273 and 235 K. Processes of autoconversion, evaporation/sublimation, and accretion of cloud water and ice are treated explicitly (Bacmeister et al., 2006). Moist convection is parameterized using the Relaxed Arakawa–Schubert (RAS) scheme (Moorthi and Suarez, 1992). Generation and evaporation of convective, anvil and stratiform precipitation are parameterized according to Bacmeister et al. (2006). Longwave radiative interactions with cloud water, water vapor, carbon dioxide, ozone, N<sub>2</sub>O and methane are treated following

## Cloud microphysics in GEOS-5

D. Barahona et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## Cloud microphysics in GEOS-5

D. Barahona et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Chou and Suarez (1994). The Chou et al. (1992) scheme is used to describe shortwave absorption by water vapor, ozone, carbon dioxide, oxygen, cloud water, and aerosols and scattering by cloud particles and aerosols. Cloud particle effective size is prescribed and tuned to adjust the radiative balance at the top of the atmosphere. The current version of GEOS-5 also accounts for the radiative effect of precipitating rain and snow according to Molod et al. (2012). Aerosol transport is calculated interactively using the GOCART aerosol model (Colarco et al., 2010).

The calculation of large scale condensation and cloud coverage in GEOS-5 follows a total-water-PDF approach (Smith, 1990; Rienecker et al., 2008; Molod, 2012). The total water probability distribution function (PDF) is assumed to follow a uniform distribution characterized by the critical relative humidity, which follows the formulation of Slingo (1987). Anvil cloud fraction is parameterized following Tiedtke (1993).

### 2.2 New cloud variables

The cloud microphysical scheme in GEOS-5 was augmented to calculate the evolution of the mass and number of ice crystals and cloud droplets. Four new prognostic variables were added to GEOS-5:  $q_l$ ,  $q_i$ ,  $n_d$  and  $n_c$  representing the grid-average mass and number mixing ratio of liquid and ice, respectively. The evolution of a given tracer,  $\eta$ , is described by

$$\frac{\partial \eta}{\partial t} = \left( \frac{\partial \eta}{\partial t} \right)_{\text{adv}} + \left( \frac{\partial \eta}{\partial t} \right)_{\text{turb}} + \left( \frac{\partial \eta}{\partial t} \right)_{\text{ls}} + \left( \frac{\partial \eta}{\partial t} \right)_{\text{cv}} \quad (1)$$

where the terms on the right hand side of Eq. (1) represent the tendency in  $\eta$  due to advective and turbulent transport and large scale and convective cloud processes, respectively. Advective and turbulent transport in GEOS-5 are described in Rienecker et al. (2008).  $\left( \frac{\partial \eta}{\partial t} \right)_{\text{ls}}$  refers to the change in  $\eta$  from non-convective cloud processes (i.e., anvil and stratus clouds), whereas  $\left( \frac{\partial \eta}{\partial t} \right)_{\text{cv}}$  describes the change in  $\eta$  from processes occurring within convective cumulus.

## 2.3 Microphysics of Stratiform and Anvil clouds

The stratiform cloud microphysics scheme of Morrison and Gettelman (2008, hereafter MG08) was implemented in GEOS-5. The scheme includes prognostic equations for the mass and number mixing ratio of cloud ice and liquid, and diagnostically predicts the vertical profiles of rain and snow. The version of MG08 implemented in GEOS-5 follows closely the description of Gettelman et al. (2010) with a few modifications. The detailed mass and number balances leading to  $\left(\frac{\partial n_d}{\partial t}\right)_{ls}$ ,  $\left(\frac{\partial q_l}{\partial t}\right)_{ls}$ ,  $\left(\frac{\partial q_i}{\partial t}\right)_{ls}$  and  $\left(\frac{\partial n_c}{\partial t}\right)_{ls}$  are presented in Morrison and Gettelman (2008). The MG08 scheme is used to describe the microphysics of convective detrainment and stratiform condensate.

In MG08 the size distribution of cloud droplets, rain, ice and snow is assumed to follow a gamma distribution, i.e.,

$$n_y(D) = N_{0,y} D_y^{\mu_y} e^{-\lambda_y D_y} \quad (2)$$

where the subscript “ $y$ ” is used to represent a hydrometeor species and  $N_{0,y}$  and  $\lambda_y$  are the slope and intercept parameters of  $n_y(D)$ , calculated as in Morrison and Gettelman (2008) (c.f. Eq. 3). A Marshall–Palmer distribution is assumed for rain and snow, i.e.,  $\mu_y = 0$ .

MG08 also uses an exponential approximation to the size distribution of ice crystals i.e.,  $\mu_i = 0$ . Theoretical considerations however suggest that  $n_i(D_i)$  in recently formed clouds is better represented by lognormal and gamma functions in which the concentration of ice crystals decreases steeply for very small sizes (Barahona and Nenes, 2008). Since this behavior cannot be reproduced using an exponential distribution, setting  $\mu_i = 0$  may lead to underestimation of  $\lambda_i$  and overestimation of crystal size. This assumption is relaxed in GEOS-5 and  $\mu_i$  is calculated as a function of  $T$  following the correlation of Heymsfield et al. (2002), obtained from extensive measurements in cirrus clouds. It is assumed that  $0.5 < \mu_i < 2.5$  where the in situ data are better constrained (Morrison and Grabowski, 2008; Heymsfield et al., 2002). The

GMDD

6, 5289–5373, 2013

### Cloud microphysics in GEOS-5

D. Barahona et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



critical size for ice autoconversion was set to  $D_{cs} = 400 \mu\text{m}$ . The sensitivity of cloud ice water to  $\mu_i$  and  $D_{cs}$  is analyzed in Sect. 4.

The autoconversion parameterization in MG08 (Khairoutdinov and Kogan, 2000) was replaced by the formulation of Liu et al. (2006). The latter was preferred because of its greater flexibility in representing the effect of cloud droplet dispersion on the autoconversion rate. The liquid water content exponent in Liu's parameterization was set to 2.0 (Liu et al., 2006). Following Liu et al. (2008) the cloud droplet size dispersion,  $\mu_l$ , was parameterized in terms of the grid-scale mean droplet mass.

Other modifications to MG08 include the calculation of the nucleated droplet number and ice crystal concentration and the parameterization of the subgrid scale vertical velocity (Sects. 2.3.2–2.3.4). Partitioning of total condensate accounts for the Bergeron–Findeisen process following Morrison and Gettelman (2008) and Gettelman et al. (2010). Ice and liquid cloud fraction are however not discriminated and total cloud fraction is calculated using the probability distribution function (PDF) of total water (Sect. 2.3.1).

### 2.3.1 Stratiform condensation and cloud fraction

Cloud fraction,  $f_c$ , plays a crucial role in microphysical processes and is intimately tied to the in-cloud number and mass mixing ratios. In GEOS-5 it is calculated using a prognostic PDF scheme, i.e.,

$$f_c = \frac{\int_{q^*}^{\infty} P_q(q_t) dq_t}{\int_0^{\infty} P_q(q_t) dq_t} \quad (3)$$

where  $P_q(q_t)$  is the total water PDF,  $q_t = q_v + q_c$ , and  $q_v$ ,  $q_c$ , and  $q_t$  are the water vapor, total condensate and total water mixing ratio, respectively, and  $q^*$  is the weighted saturation mixing ratio between liquid and ice, given by

$$q^* = (1 - f_{ice})q_l^* + f_{ice}q_i^* \quad (4)$$

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



where  $f_{\text{ice}}$  is the mass fraction of ice in the total condensate and  $q_l^*$  and  $q_i^*$  are the saturation specific humidities for liquid and ice, respectively. Total condensate is therefore given by

$$q_c = \frac{\int_{q^*}^{\infty} (q_t - q^*) P_q(q_t) dq_t}{\int_0^{\infty} P_q(q_t) dq_t} \quad (5)$$

5 The total water distribution in GEOS-5 is defined as a uniform PDF in non-anvil regions plus a  $\delta$ -function representing the detrained condensate from convective cumulus (Rienecker et al., 2008). The same assumption is used in this work, however the lower limit of integration in Eqs. (3) and (5) is modified to  $q^* S_{\text{crit}}$ , where  $S_{\text{crit}}$  is termed the critical saturation ratio.  $S_{\text{crit}}$  controls the level of supersaturation required for cloud formation within a model grid cell. As in the operational version of GEOS-5, it is assumed that  $S_{\text{crit}} = 1$  for mixed-phase and liquid clouds. However for ice clouds linking  $S_{\text{crit}}$  to ice nucleation processes increases the minimum relative humidity required for cloud formation, allowing for supersaturation with respect to ice. Thus, in cirrus clouds  $S_{\text{crit}}$  is calculated by the ice nucleation parameterization (Sect. 3.5).

15 Solution of Eq. (3) gives (Rienecker et al., 2008),

$$f_c = \frac{q_{\text{mx}} - S_{\text{crit}} q^*}{\Delta q} + f_{\text{cn}} \quad (6)$$

where  $q_{\text{mx}} = q_t + 0.5\Delta q$  is the upper limit of the uniform distribution,  $\Delta q$  is the width of  $P_q(q_t)$  (Slingo, 1987) and  $f_{\text{cn}}$  is the detrained anvil cloud fraction calculated according to Tiedtke (1993).

20 Similarly, solution of Eq. (5) gives for the total condensate (Rienecker et al., 2008),

$$q_c = \frac{1}{2} \frac{(q_{\text{mx}} - S_{\text{crit}} q^*)^2}{\Delta q} + q_{\text{c, det}} \quad (7)$$

where  $q_{\text{c, det}}$  is the mixing ratio of detrained condensate.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Microphysical processes modify  $q_t$  and  $P_q(q_t)$  via the formation of precipitation (Tompkins, 2002). The effect of the microphysics on the cloud fraction is accounted for as follows. Assuming that the total water PDF (i.e., anvil and stratiform) after microphysical processing follows a uniform distribution, an equation similar to Eq. (7)

5 can be written for the total condensate in the form,

$$q_c + \Delta q_c = \frac{1}{2} \frac{(q'_{\text{mx}} - S_{\text{crit}} q^*)^2}{\Delta q'} \quad (8)$$

where  $\Delta q_c = \left( \frac{\partial q_c}{\partial t} \right)_{\text{is}} \Delta t$  is the change in total condensate over the time step  $\Delta t$ , and  $q'_{\text{mx}}$  and  $\Delta q'$  represent the values of  $q_{\text{mx}}$  and  $\Delta q$  after the microphysics. Similarly for cloud fraction,

$$10 \quad f'_c = \frac{q'_{\text{mx}} - S_{\text{crit}} q^*}{\Delta q'} \quad (9)$$

Inverting Eq. (8) to find  $\Delta q'$  gives,

$$\Delta q' = \frac{2q_c + \Delta q_c}{(q_{\text{mx}} - S_{\text{crit}} q^* + \Delta q_c)^2} \quad (10)$$

Combining Eqs. (9) and (10) eliminates  $\Delta q'$  and  $q_{\text{mx}}$ . The cloud fraction modified by the microphysics then becomes,

$$15 \quad f'_c = \left( \sqrt{1 - \frac{q_t + \Delta q_c - S_{\text{crit}} q^*}{q_c + \Delta q_c}} + 1 \right)^{-1} \quad (11)$$

In practice, an initial estimate of  $f_c$  (Eq. 6) is used to calculate  $\Delta q_c$  assuming that microphysical processes proceed at constant cloud fraction. Then  $f'_c$  is calculated from Eq. (11) and used for radiative calculations. This procedure ensures that  $f'_c$  calculated after microphysical processing is consistent with  $P_q(q_t)$  and the amount of condensate present in the grid cell at the end of each time step.

20

## 2.3.2 Cloud droplet activation

CCN activation into cloud droplets is parameterized following the approach of Fountoukis and Nenes (2005) (FN05). FN05 is an analytical solution of the equations of an ascending cloudy parcel using the method of population splitting (Nenes and Seinfeld, 2003). Sulfates, hydrophilic organics and sea salt are considered CCN active species. Aerosol number concentrations were derived from the predicted mass mixing ratio for each species using size distributions obtained from the literature (Table 1). Sulfate and organics are considered internally mixed and five separate bins are used to describe dust. Aerosol composition is parameterized in terms of the hygroscopicity parameter (Petters and Kreidenweis, 2007):  $\kappa$  was set to 0.65, 0.2 and 1.28 for sulfate, hydrophilic organics, and sea salt, respectively. The water uptake coefficient was set 1.0 (Raatikainen et al., 2013). In this work the adiabatic version of the FN05 parameterization is employed. However FN05 can be readily extended to include dust activation (Kumar et al., 2009b), entrainment (Barahona and Nenes, 2007), and giant CCN (Barahona and Nenes, 2009a). The contribution of CCN activation in stratiform clouds to the droplet number concentration is given by

$$\left(\frac{dN_d}{dt}\right)_{ls, act} = \frac{\max(N_{d, act} - N_d, 0)}{\Delta t} \quad (12)$$

where  $N_d$  and  $N_{d, act}$  are the in-cloud preexisting and activated droplet number concentration, respectively.

## 2.3.3 Ice nucleation

The ice nucleation parameterization implemented in GEOS-5 was developed by Barahona and Nenes (2008, 2009a, b) (BN09), and is summarized in Barahona et al. (2010). BN09 is derived from the analytical solution of the governing equations of an ascending cloud parcel, and considers the dependency of the ice crystal concentration,  $N_c$ , on cloud formation conditions, subgrid scale dynamics, and aerosol properties.

GMDD

6, 5289–5373, 2013

## Cloud microphysics in GEOS-5

D. Barahona et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion





the nucleation spectrum in the form (Barahona and Nenes, 2009b),

$$S_{\text{het}} = \max \left[ 1 + S_{i,\text{max}} - \mathcal{N}_{\text{het}} \left( \frac{\partial \mathcal{N}_{\text{het}}}{\partial S_i} \right)^{-1}, 1 \right] \quad (14)$$

where  $S_{i,\text{max}}$  is the maximum saturation ratio reached in a single parcel ascent, calculated according to BN09. If no IN are present then  $S_{\text{het}}$  approaches the saturation threshold for homogeneous freezing,  $S_{\text{hom}}$  (Barahona and Nenes, 2009b).  $S_{\text{het}}$  and  $S_{\text{hom}}$  represent the minimum saturation ratio required for cloud formation by heterogeneous and homogeneous freezing, respectively. Thus they have the same meaning as the critical saturation ratio of Eq. (6).  $S_{\text{crit}}$  is then calculated as,

$$S_{\text{crit}} = f_{\text{het}} S_{\text{het}} + (1 - f_{\text{het}}) S_{\text{hom}} \quad (15)$$

where  $f_{\text{het}}$  is the fraction of ice crystals produced by heterogeneous ice nucleation (given by the BN09 parameterization), and  $S_{\text{hom}}$  is calculated following Koop et al. (2000).

The contribution of ice nucleation in cirrus to the ice crystal number concentration is given by,

$$\left( \frac{dN_c}{dt} \right)_{\text{cirrus, nuc}} = \frac{\max[N_{c, \text{nuc}} P_q(q_t > S_{\text{crit}} q_i^*) - N_c, 0]}{\Delta t} \quad (16)$$

where  $N_{c, \text{nuc}}$  is the nucleated ice crystal concentration. The factor  $P_q(q_t > S_{\text{crit}} q_i^*)$  accounts for the the probability of finding an air mass leading to cloud formation within the grid cell. This term was proposed by Barahona and Nenes (2011) to account for the effect of prior nucleation events on current cloud formation.

For the mixed-phase regime ( $T > 235$  K), Eq. (13) is applied directly to find the contribution of deposition and condensation heterogeneous nucleation to  $N_c$ . In this regime cloud droplet freezing by immersion and contact ice nucleation contribute to the

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



ice crystal population. The tendency in  $N_c$  from immersion freezing of cloud droplets is given by

$$\left(\frac{dN_c}{dt}\right)_{\text{imm}} = \sum_x N_x \bar{s}_{p,x} \gamma_c \frac{dn_{s,x}}{dT} \exp(-\bar{s}_{p,x} n_{s,x}) \quad (17)$$

where  $\gamma_c = -w_{\text{sub}} \frac{dT}{dz}$  is the cooling rate and  $n_{s,x}$  the active site surface density for the species “x”. The latter is calculated according to Niemand et al. (2012) for dust and Murray et al. (2012) for black carbon.

Contact ice nucleation is parameterized as the product of the collection flux of aerosol particles by the cloud droplets and the ice nucleation efficiency in contact mode. Young (1974) suggested that phoretic effects and Brownian motion are responsible for collection scavenging of ice nuclei. Baker (1991) however showed that Brownian motion is the dominant factor. Therefore the contribution of contact ice nucleation to the ice crystal formation tendency can be written as,

$$\left(\frac{dN_c}{dt}\right)_{\text{cont}} = \sum_x \left(\frac{dN_x}{dt}\right)_{\text{Brw}} \{1 - \exp[-\bar{s}_{p,x} n_{s,x}(T_{\text{cont}})]\} \quad (18)$$

where  $\left(\frac{dN_x}{dt}\right)_{\text{Brw}}$  is the Brownian collection flux of the x aerosol species (Young, 1974). Consistent with laboratory studies (e.g., Fornea et al., 2009; Ladino et al., 2011) the active site density in the contact mode is assumed to be the same as for immersion freezing shifted towards higher temperature, i.e.,  $T_{\text{cont}} \approx T - 3\text{K}$ .

The in-cloud contribution of ice nucleation in mixed-phase clouds to the ice crystal number concentration tendency is given by,

$$\left(\frac{dN_c}{dt}\right)_{\text{mixed, nuc}} = \min \left[ \left(\frac{dN_c}{dt}\right)_{\text{cont}} + \left(\frac{dN_c}{dt}\right)_{\text{imm}} + \left(\frac{dN_c}{dt}\right)_{\text{dep}}, \frac{N_d}{\Delta t} \right] \quad (19)$$

where the subscripts cont, imm, and dep, refer to contact, immersion, and deposition/condensation ice nucleation, respectively. The term  $\frac{N_d}{\Delta t}$  is used to limit the nucleated ice crystal concentration to the existing concentration of cloud droplets.

### 2.3.4 Subgrid scale dynamics

Besides information on the aerosol composition and size, parameterization of cloud droplet and ice crystal formation requires the knowledge of the vertical velocity,  $w_{\text{sub}}$ , at the spatial scale of individual parcels (typically under 100 m), which is not resolved by GEOS-5.  $w_{\text{sub}}$  depends on radiative cooling (Morrison et al., 2005), turbulence (Golaz et al., 2010), gravity wave dynamics (e.g., Barahona and Nenes, 2011; Kärcher and Ström, 2003; Jensen et al., 2010; Joos et al., 2008) and local convection. To account for these dependencies we employ a semiempirical formulation as follows.

In situ measurements (e.g., Peng et al., 2005; Bacmeister et al., 1999; Conant et al., 2004) suggest that  $w_{\text{sub}}$  is approximately normally distributed. The mean vertical velocity of the distribution is written as (Morrison et al., 2005)

$$\bar{w} = w_{\text{is}} - \frac{c_p}{g} \left( \frac{\partial T}{\partial t} \right)_{\text{rad}} \quad (20)$$

where  $w_{\text{is}}$  is the grid-scale vertical velocity,  $c_p$  the heat capacity of air,  $g$  is the acceleration of gravity, and  $\left( \frac{\partial T}{\partial t} \right)_{\text{rad}}$  is the diabatic heating due to radiative transfer. Variance in  $w_{\text{sub}}$  for large scale clouds (i.e., stratus and in situ cirrus) results from subgrid scale eddy motion,  $\sigma_{w,\text{turb}}^2$ , and gravity wave dynamics,  $\sigma_{w,\text{gw}}^2$ , i.e.,

$$\sigma_w^2 = \sigma_{w,\text{turb}}^2 + \sigma_{w,\text{gw}}^2 \quad (21)$$

A first order closure is used to diagnose  $\sigma_{w,\text{turb}}^2$  (Morrison and Gettelman, 2008),

$$\sigma_{w,\text{turb}}^2 = \frac{K_T}{I_m} \quad (22)$$

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



where  $K_T$  is the mixing coefficient for heat (Louis et al., 1983) and  $l_m$  is the mixing length. MG08 prescribed a fixed  $l_m = 300$  m (Morrison and Gettelman, 2008). To account for the spatial variation of  $l_m$ , the formulation of Blackadar (1962) is used instead, i.e.,

$$l_m = \frac{kz}{1 + \frac{kz}{\lambda_m}} \quad (23)$$

where  $k$  is the von Kármán constant,  $z$  is the altitude and  $\lambda_m$  is the value of  $l_m$  in the free troposphere (Blackadar, 1962). This approach also takes into account the vertical variation of  $l_m$  within the planetary boundary layer (PBL). The minimum value of  $\sigma_{w,turb}^2$  is set to  $0.01 \text{ m}^2 \text{ s}^{-2}$  within the PBL.

Small-scale gravity waves strongly affect the formation of cirrus and mixed-phase clouds (e.g., Haag and Kärcher, 2004; Jensen et al., 2010; Joos et al., 2008; Barahona and Nenes, 2011; Dean et al., 2007). In situ measurements suggest that the dynamics of the upper troposphere are characterized by the random superposition of gravity waves from different sources (e.g., Jensen and Pfister, 2004; Bacmeister et al., 1999; Sato, 1990; Herzog and Vial, 2001). Random wave superposition results in a Gaussian distribution of vertical velocities (e.g., Bacmeister et al., 1999; Barahona and Nenes, 2011). Using this a semiempirical parameterization for  $\sigma_{w,gw}^2$  is derived in the form (Eq. A5),

$$\sigma_{w,gw}^2 = 0.0169 \min \left[ \frac{4\pi U |\tau_0|}{\rho_a L_c N}, \left( \frac{2\pi U^2}{NL_c} \right)^2 \right] \quad (24)$$

where  $\tau_0$  is the surface stress (Lindzen, 1981),  $U$  the horizontal wind,  $\rho_a$  the air density,  $N$  the Brunt–Väisälä frequency, and  $L_c$  the wave length of the highest frequency waves in the spectrum, also referred to as the characteristic cirrus scale (here assumed to be 100 m). Equation (24) is obtained by relating  $|\tau_0|$  to the equivalent perturbation height at

## Cloud microphysics in GEOS-5

D. Barahona et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



the surface. This is scaled to obtain the maximum wave amplitude at each vertical level (Joos et al., 2008; McFarlane, 1987) and then used to compute  $\sigma_{w, gw}^2$  (Barahona and Nenes, 2011). This approach parameterizes orographically-generated gravity waves. In practice, both the background and the orographic surface stress are used in Eq. (24) to avoid underestimation of  $\sigma_{w, gw}^2$  in marine regions. The second term in brackets on the right hand side of Eq. (24) limits  $\sigma_{w, gw}$  to account for wave saturation and breaking (Eq. A3). The derivation of Eq. (24) is detailed in the Appendix A.

The nucleated ice crystal concentration is obtained by averaging over the positive values of  $w_{sub}$ ,

$$N_{c, nuc} = \frac{\int_0^{w_{max}} N_{c, nuc}(w_{sub}) \phi(\bar{w}, \sigma_w^2) dw_{sub}}{\int_0^{w_{max}} \phi(\bar{w}, \sigma_w^2) dw_{sub}} \quad (25)$$

where  $\phi(\bar{w}, \sigma_w^2)$  is the normal distribution and  $w_{max} = \bar{w} + 4\sigma_w$ . The latter is used as an upper limit to the integral to avoid numerical instability. For liquid droplet activation Eq. (25) is simplified as (Peng et al., 2005; Fountoukis and Nenes, 2005),

$$N_{d, act} = N_{d, act}(\bar{w} + 0.8\sigma_w) \quad (26)$$

This approximation is valid for  $\bar{w} \ll \sigma_w$  and may introduce up to 20% non-systematic discrepancy in  $N_{d, act}$  when compared to the direct solution of the integral in Eq. (25) (Morales and Nenes, 2010), however it is justified on computational efficiency. Notice that the same approximation cannot be used for ice nucleation since the competition between homogeneous and heterogeneous nucleation introduces strong nonlinearity in  $N_{c, nuc}(w_{sub})$  (Barahona and Nenes, 2009a) and therefore the characteristic value of  $w_{sub}$  for  $N_{c, nuc}$  generally differs from the average vertical velocity. PDF-averaging is also applied for  $S_{crit}$ ,  $S_{i, max}$  and  $S_{i, max}$ . Only activation processes are modified by subgrid vertical velocity variability, i.e.,  $\phi(\bar{w}, \sigma_w^2)$  is assumed uncorrelated to the subgrid distribution of condensate.

### 2.3.5 Preexisting ice crystals

Ice nucleation can be inhibited by water vapor deposition onto preexisting ice crystals (i.e., ice crystals present in the grid cell from previous nucleation events). Their impact on cirrus properties may be significant at low temperature where ice crystals are small and have low sedimentation rates (Barahona and Nenes, 2011). This effect can be parameterized by reducing the vertical velocity for ice nucleation in cirrus by a factor dependent on the preexisting ice crystal concentration and size (Eq. B5), i.e.,

$$w_{\text{sub, pre}} = w_{\text{sub}} \max \left( 1 - \frac{N_{i, \text{pre}} \pi \beta c \rho_i A_i (S_{\text{hom}} - 1)}{2 \lambda_{i, \text{pre}} \alpha w_{\text{sub}} S_{\text{hom}}}, 0 \right) \quad (27)$$

where  $N_{i, \text{pre}}$  is the preexisting ice crystal concentration,  $c$  is a shape factor (here assumed equal to 1),  $\rho_i$  is the bulk density of ice, and  $A_i$ ,  $\alpha$  and  $\beta$  are temperature-dependent parameters (Appendix Table A1). Equation (27) indicates that water vapor deposition onto preexisting crystals acts against the increase in supersaturation from expansion cooling. The derivation of Eq. (27) is detailed in the Appendix B. The effect of preexisting ice crystals on cirrus properties is analyzed in Sect. 4.

### 2.4 Microphysics of convective cumulus

While all the main features of RAS are preserved in the new scheme, the removal of condensate is reformulated to account for the effect of IN and CCN emissions on the generation of convective precipitation. RAS calculates the convective cloud condensate and mass flux at each model level by averaging over an ensemble of ascending parcels, each one lifted from the the top of the PBL (Molod et al., 2012; Rienecker et al., 2008). Each ascending parcel is characterized by its detrainment level and entrainment rate (Moorthi and Suarez, 1992) and saturation adjustment is used to find the amount of condensate present in each parcel. In the current RAS implementation in GEOS-5 a single parcel detrains at each model level so that the tendency of the tracer  $\eta$  due to

GMDD

6, 5289–5373, 2013

## Cloud microphysics in GEOS-5

D. Barahona et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



cloud convective processes is given by

$$\left(\frac{\partial \eta}{\partial t}\right)_{cv} = D\eta - gW \frac{\partial \eta}{\partial p} \quad (28)$$

where  $D$  is the detrainment rate and  $W$  the convective mass flux. In the operational GEOS-5, a prescribed fraction of condensate is assumed to precipitate from each parcel before reaching cloud top. The remaining condensate is then linearly partitioned between ice and liquid as a function of  $T$  and detrained at the neutral buoyancy level. In this approximation there is no remaining condensate in the convective cloud at the end of each time step. Each parcel is assumed to develop independently and the detrained condensate from different parcels is weighted by the convective mass flux. The subscript “cp” in the following equations refers to processes occurring within each parcel. A detailed description of the GEOS-5 convective scheme is presented elsewhere (Moorthi and Suarez, 1992; Rienecker et al., 2008).

The balance of a tracer,  $\eta$ , within a convective parcel is written as

$$\frac{1}{W} \frac{d(\eta W)}{dt} = \left(\frac{d\eta}{dt}\right)_{cp} + \lambda w_{cp}(\eta' - \eta) \quad (29)$$

where  $\left(\frac{d\eta}{dt}\right)_{cp}$  is the rate of change in  $\eta$  from microphysical processes occurring within convective parcels,  $w_{cp}$  is the parcel vertical velocity,  $\lambda$  the per-length entrainment rate and  $\eta'$  the value of  $\eta$  in the cloud-free environment. Detrainment of condensate is assumed to occur only at cloud top.

The rate of change in  $\eta$  from microphysical processes occurring within a convective cloud parcel is given by

$$\left(\frac{d\eta}{dt}\right)_{cp} = \left(\frac{d\eta}{dt}\right)_{source} + \left(\frac{d\eta}{dt}\right)_{precip} + \left(\frac{d\eta}{dt}\right)_{freezing} \quad (30)$$

where the subscript “source” refers to nucleation, condensation and deposition processes, “precip” to precipitation and “freezing” to phase transformation.

Cloud microphysics  
in GEOS-5

D. Barahona et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Equation (29) is integrated for each parcel from cloud base to cloud top at which all remaining condensate detrains into the anvil, i.e.,  $\left[ \frac{1}{W} \frac{d(\eta W)}{dt} \right]_{\text{cloud top}} = D\eta$ . The initial condition in Eq. (29) depends on the tracer. At cloud base the concentration of ice crystals and the ice mass mixing ratio are assumed to be zero, whereas the activation of cloud droplets at cloud base is explicitly considered (Sect. 2.4.2).

Solution of Eq. (29) requires the knowledge of the vertical velocity within each parcel,  $w_{\text{cp}}$ , which is also necessary to drive the droplet activation and ice nucleation parameterizations. This is calculated by solving (Frank and Cohen, 1987),

$$\frac{1}{2} \frac{dw_{\text{cp}}^2}{dz} = \frac{g}{1+\gamma} \frac{T_v - T'_v}{T'_v} - \lambda w_{\text{cp}}^2 - gq_{\text{cn}} \quad (31)$$

where  $\gamma = 0.5$  (Sud and Walker, 1999),  $T_v$  and  $T'_v$  the virtual temperature of the cloud and the environment, respectively, and  $q_{\text{cn}}$  is the mixing ratio of total condensate in the convective parcel. Equation (31) is forwardly integrated from the level below cloud base to cloud top using  $w_{\text{cp, in}} = 0.8 \text{ ms}^{-1}$  as initial condition (e.g., Guo et al., 2008; Gregory, 2001); the vertical profile  $w_{\text{cp}}$  is not very sensitive to this assumption (Sud and Walker, 1999). Notice that  $w_{\text{cp, in}}$  differs from the vertical velocity used for cloud droplet activation. The latter depends on the local buoyancy, i.e.,  $w_{\text{cp, cloud base}} = w_{\text{cp, in}} + \frac{dw_{\text{cp}}}{dz} \Delta z_{\text{base}}$  where  $\Delta z_{\text{base}}$  is the model layer thickness at cloud base.

### 2.4.1 Partitioning of convective condensate

Total condensate is partitioned between liquid and ice as follows. Nucleated ice crystals are assumed to grow by accretion of water vapor in an environment saturated with respect to liquid water. That is, the coexistence of liquid water favors a high concentration of water vapor available for deposition onto the ice crystals and the ice and liquid phases remain in quasi-equilibrium within the convective parcel. Hydrometeor species are assumed to follow a gamma distribution (Eq. 2). The growth

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



rate of ice crystals within convective cumulus is given by (Pruppacher and Klett, 1997; Korolev and Mazin, 2003)

$$\left(\frac{dq_i}{dt}\right)_{\text{dep}} = \frac{n_i \pi c \rho_i A_i (S_i - 1)}{2\lambda_i} \quad (32)$$

where  $c$  is a shape factor (assumed equal to 1),  $\rho_i$  the ice crystal density, and  $A_i$  is a temperature-dependent growth factor (Appendix Table A1). Using Eq. (32), and since  $q_{\text{cn}} = q_l + q_i$ , the source term for liquid water within convective cumulus is given by

$$\left(\frac{dq_l}{dt}\right)_{\text{cond}} = \left(\frac{dq_{\text{cn}}}{dt}\right) - \left(\frac{dq_i}{dt}\right)_{\text{dep}} \quad (33)$$

where  $\left(\frac{dq_{\text{cn}}}{dt}\right)$  is the rate of generation of total condensate calculated by the convective parameterization.

#### 2.4.2 Droplet activation and ice crystal nucleation in convective cumulus

Explicit activation of CCN into cloud droplets is only considered at cloud base and used as an initial condition to Eq. (29) (Sect. 2.4). Entrained aerosols (sulfate, sea salt, and organics) are assumed to activate instantaneously as they enter the cloud parcel. Dust and soot IN lead to the heterogeneous freezing of cloud droplets in the immersion and contact modes, described using Eqs. (17) and (18). Since soot and dust particles would likely adsorb water within convective parcels (Wiacek et al., 2010; Kumar et al., 2009a) ice nucleation in the deposition mode within convective cumulus is not considered. Cloud droplets freeze homogeneously at 235 K. Frozen droplets rapidly quench supersaturation within convective cumulus. Thus the homogeneous nucleation of deliquesced sulfate, which requires high supersaturation ( $S_i \sim 145\text{--}170\%$ , Koop et al., 2000), is not likely to occur within convective parcels. Therefore homogeneous freezing of interstitial aerosol is not considered in convective cumulus.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



### 2.4.3 Generation of convective precipitation

The size dispersion of the droplet population,  $\mu_I$ , follows the formulation of Liu et al. (2008). Droplet-to-rain autoconversion is calculated according to Liu et al. (2006) and all autoconverted water is assumed to be lost as surface precipitation within one time step. Evaporation of convective precipitation is parameterized according to Bacmeister et al. (1999).

Ice water in convective cumulus is likely to exist as graupel, snow and ice crystals, with different size distributions and falling velocities. Following Del Genio et al. (2005) a simplified treatment of ice precipitation is implemented as follows. Total ice water within convective parcels is assumed to partition as ice/snow (taken as a single species) and graupel, and differentiated by their terminal velocity (Table 2). The fraction of total ice existing as graupel is approximated by (Del Genio et al., 2005),

$$f_{gr} = 0.25 \{3.0 + \exp[0.1 \min(T - 273, 0)]\} \quad (34)$$

The particle sizes of ice/snow and graupel are assumed to follow an exponential distribution ( $\mu_g = \mu_{i/s} = 0.0$ ) (McFarquhar and Heymsfield, 1997). The number precipitation rate of ice/snow within convective parcels is given by the number flux across a critical size,  $D_{c, i/s}$  (Seinfeld, 1998),

$$\left(\frac{dn_{i/s}}{dt}\right)_{\text{precip, cp}} = \frac{n_{i/s} A_i (S_i - 1)}{D_{c, i/s}^2} [1 - \exp(-\lambda_{i/s} D_{c, i/s})] \quad (35)$$

where  $n_{i/s} = (1 - f_{gr})n_i$ . The mass precipitation rate of ice/snow is calculated as,

$$\left(\frac{dq_{i/s}}{dt}\right)_{\text{precip, cp}} = \frac{q_{i/s} \xi_{i/s}}{n_{i/s}} \left(\frac{dn_{i/s}}{dt}\right)_{\text{precip, cp}} \quad (36)$$

where  $q_{i/s} = (1 - f_{gr})q_i$ , and  $\xi_{i/s} = \frac{1}{6}[(\lambda_{i/s} D_{c, i/s})^3 + 3(\lambda_{i/s} D_{c, i/s})^2 + 6\lambda_{i/s} D_{c, i/s} + 6]$  is the ratio of the volume to number fraction above  $D_{c, i/s}$  in the size distribution of ice/snow. The term

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



$\xi_{i/s}$  is introduced to account for the preferential precipitation of the largest particles of the population, which tends to enhance the mass over the number precipitation rate. The critical size for precipitation,  $D_{c, i/s}$ , is calculated by equating the hydrometeor terminal velocity,  $w_{\text{term}}$ , to  $w_{\text{cp}}$  (Table 2).

Equations (35) and (36) assume that ice and snow grow mainly by diffusion within the convective parcel. The same assumption cannot be applied to graupel since it also grows by collection of cloud droplets. The precipitation rate of graupel is therefore approximated calculated by removing the fraction of the size distribution above  $D_{c,g}$  at each model level (Ferrier, 1994),

$$\left(\frac{dn_{\text{gr}}}{dt}\right)_{\text{precip, cp}} = \frac{n_{\text{gr}} \exp(-\lambda_g D_{c,g})}{\Delta t_L} \quad (37)$$

where  $n_{\text{gr}} = f_{\text{gr}} n_i$  is the graupel number mixing ratio and  $\Delta t_L = \Delta z \bar{w}_{\text{cv}}^{-1}$  is the time spent by the parcel in a given model layer. Similarly for  $q_{\text{gr}}$ ,

$$\left(\frac{dq_{\text{gr}}}{dt}\right)_{\text{precip, cp}} = \frac{q_{\text{gr}} \exp(-\lambda_g D_{c,g}) [(\lambda_g D_{c,g})^3 + 3(\lambda_g D_{c,g})^2 + 6\lambda_g D_{c,g} + 6]}{6\Delta t_L} \quad (38)$$

where  $q_{\text{gr}} = f_{\text{gr}} q_i$  is the graupel mass mixing ratio

The total mass precipitation rate for ice within convective parcels is given by,

$$\left(\frac{dq_i}{dt}\right)_{\text{precip, cp}} = \left(\frac{dq_{i/s}}{dt}\right)_{\text{precip, cp}} + \left(\frac{dq_{\text{gr}}}{dt}\right)_{\text{precip, cp}} \quad (39)$$

Similarly for the ice crystal number concentration,

$$\left(\frac{dn_i}{dt}\right)_{\text{precip, cp}} = \left(\frac{dn_{i/s}}{dt}\right)_{\text{precip, cp}} + \left(\frac{dn_{\text{gr}}}{dt}\right)_{\text{precip, cp}} \quad (40)$$

Equations (39) and (40) are used into Eq. (30), which then is used to solve Eqs. (28) and (29).

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



### 3 Model evaluation

Model evaluation is carried out by comparing cloud properties against satellite retrievals and in situ observations. Satellite data sets included level 3 products from the NASA MODIS (<http://modis.gsfc.nasa.gov/>) combined TERRA and AQUA data product (Platnick et al., 2003), and the ISCCP (<http://isccp.giss.nasa.gov/>) (Rossow and Schiffer, 1999) and CloudSat (Li et al., 2012, 2013) projects. When possible, the CFMIP Observation Simulator Package (COSP) (Bodas-Salcedo et al., 2011) was used to compare model output against satellite retrievals. Global cloud radiative properties were obtained from the CERES Energy Balanced and Filled (EBAF) level 4 data product (Loeb et al., 2009) and the NASA Earth Radiation Experiment (ERBE Barkstrom, 1984). Total precipitation was obtained from the Global Precipitation Climatology Project data set (GPCP) (Huffman et al., 1997) and the CPC merged analysis of precipitation (CMAP) (Xie and Arkin, 1997). Runs were performed for a period of 10 yr starting on 1 January 2001 with a spin-up time of one year using a c48 cubed-sphere grid (about  $\sim 2^\circ$  spatial resolution) and 72 vertical levels. Sensitivity studies (Sect. 4) were performed running the model for two years at the same resolution. Test runs showed that two years were enough to elucidate the first order effect of variation in microphysical parameters on cloud properties. All simulations were forced with observed sea surface temperatures (Reynolds et al., 2002). Initial conditions were obtained from the MERRA reanalysis (Rienecker et al., 2011). The aerosol concentration was calculated interactively using the GOCART model (Colarco et al., 2010) with emissions as described in Diehl et al. (2012). Results obtained with the operational version of GEOS-5 and using the new microphysics are referred to as the CTL and NEW runs, respectively.

#### 3.1 Cloud fraction

The parameterization of  $f_c$  in GEOS-5 was modified to account for the effect of microphysical processing on  $P_q(q_t)$  (Sect. 2.3.1) and allow supersaturation with respect

GMDD

6, 5289–5373, 2013

## Cloud microphysics in GEOS-5

D. Barahona et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## Cloud microphysics in GEOS-5

D. Barahona et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



to the ice phase. Figure 1 shows the effect of these modifications on the low (CLDLO), middle (CLDMD), and high (CLDHI) cloud fraction in GEOS-5. In general the CTL and NEW simulations present similar distributions of cloud fraction. However in NEW,  $f_c$  tends to be higher and in better agreement with ISCCP retrievals. The new cloud fraction scheme resulted in higher CLDLO in the remote Atlantic and Pacific oceans and reduced the cloud bias over South America and Asia. Still CLDLO associated with the low level stratocumulus decks in the west coast of North, South America and South Africa is underpredicted in the NEW simulation. This feature is common in climate models (Kay et al., 2012); in GEOS-5 it is likely caused by the absence of an explicit shallow cumulus parameterization. The overprediction of CLDLO in the high latitudes of NH in CTL is also significantly reduced in the NEW simulation. Overall, the global mean bias in CLDLO is significantly lower in NEW ( $-3\%$ ) than in CTL ( $-5\%$ ).

The global mean bias in CLDMD is also lower in NEW ( $-9\%$ ) than in CTL ( $-15\%$ ). The overestimation of CLDMD in the low and midlatitudes of SH and NH in CTL is largely removed in NEW, which results from a more realistic distribution of ice water content in NEW than in CTL (Sect. 3.6). The underestimation in CLDMD in the high latitudes of SH and NH is also smaller in NEW than in CTL, particularly over land. However CLDMD in these regions is still about 10% lower in NEW than the ISCCP retrieval. The CTL and the NEW simulations present similar distributions of high level clouds (CLDHI). In general CLDHI tends to be overestimated in the marine high latitudes and underestimated over the continents. The NEW simulation also tends to underpredict CLDHI over the Tropical Warm Pool. The global mean bias in CLDHI is about 1% and 4% in the CTL and NEW run, respectively.

### 3.2 Supersaturation over ice

Two mechanisms lead to ice supersaturation in the new microphysics. Both  $f_c$  and  $q_i$  are produced only when  $S_i > S_{\text{crit}}$  (Eqs. 6 and 7). Ice nucleation is also restricted to supersaturated regions (Eq. 16). Both mechanisms are controlled in part by  $S_{\text{crit}}$  which provides an internal link between ice nucleation,  $f_c$  and  $q_i$ .

Cloud microphysics  
in GEOS-5

D. Barahona et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



The global distribution of  $S_{\text{crit}}$  for  $T < 235$  K (Fig. 2, right panel) presents two characteristic modes, showing regions of predominance of heterogeneous ( $S_{\text{crit}} \sim 120\%$ ) and homogeneous ( $S_{\text{crit}} \sim 140\%$ ) ice nucleation. The mean value of  $S_{\text{crit}}$  in the upper troposphere is about 144%, and  $S_{\text{crit}}$  typically ranges between 120% and 150%, which agrees with values commonly used in GCM studies (e.g., Liu et al., 2007; Salzmann et al., 2010). However  $S_{\text{crit}}$  is highly variable around the globe as it depends on  $w_{\text{sub}}$ ,  $T$ , and the concentration of IN in the upper troposphere. Figure 2 (left panel) shows that values of  $S_{\text{crit}}$  as low 105% and as high as 160% are not uncommon. Low  $S_{\text{crit}}$  is associated with regions of high concentration of active IN (e.g., dust). These are often located around  $T \sim 230$ – $240$  K where deposition/condensation IN are active and abundant enough to impact supersaturation (Sect. 3.5). For lower  $T$ , the concentration of active IN is too low to substantially decrease supersaturation, and  $S_{\text{crit}}$  increases towards the homogeneous freezing threshold (Fig. 2). This behavior suggests that no single value of  $S_{\text{crit}}$  can represent all the characteristic values of critical relative humidity for cirrus formation around the globe.

The distribution of clear sky saturation ratio,  $S_{i,c} = (q_v - f_c q^*) / (1.0 - f_c)$ , is shown in Fig. 3. In-cloud  $S_i$  is assumed to be 100%. In reality supersaturation relaxation may be slow in cirrus clouds particularly at low  $T$  (Krämer et al., 2009; Barahona and Nenes, 2011). However it is expected that for the conditions of Fig. 3 most supersaturation is relaxed inside clouds over the time step of the simulation ( $\sim 1800$  s) (Barahona and Nenes, 2008). Figure 3 also shows data from the AIRS (Gettelman et al., 2006) and MOZAIC (Gierens et al., 1999) projects. The uncertainty in the retrieval increases with  $S_{i,c}$ . However both MOZAIC and AIRS data show an exponential decrease in the frequency of supersaturation,  $P(S_{i,c})$ , with increasing  $S_{i,c}$ . GEOS-5 also shows this exponential decrease and is in agreement with AIRS and MOZAIC data. The peak  $P(S_{i,c})$  in the model is shifted towards  $S_{i,c} \sim 100\%$  since retrievals tend to avoid zones with  $S_{i,c} \sim 100\%$  near the cloud edges (Gettelman and Kinnison, 2007). The frequency of  $S_{i,c} > 101\%$  in GEOS-5 distributes almost symmetrically around the Tropics (Fig. 3, right panel), with a slightly higher probability of supersaturation in SH than in NH. This

is in part due to lower IN concentrations in SH (Fig. 7), although differences in the dynamics of SH and NH also play a significant role. In agreement with AIRS data, GEOS-5 predicts about 10% supersaturation frequency in the upper Tropical levels. GEOS-5 seems to slightly overpredict  $P(S_{i,c})$  above 300 hpa in the high latitudes of the NH and SH and near the TTL, however the uncertainty of the retrieval in these regions is also high (Gettelman and Kinnison, 2007).

### 3.3 Subgrid scale vertical velocity

The nucleation of ice crystals and cloud droplets is strongly influenced by  $w_{\text{sub}} \cdot \phi(\bar{w}, \sigma_w^2)$  in stratocumulus and anvils is mainly determined by  $\sigma_w$  whereas  $\bar{w}$  is typically small ( $\sim 10^{-2} \text{ ms}^{-1}$ ). For convective clouds  $w_{\text{cp}}$  is explicitly calculated by solving Eq. (31). In general the eddy contribution to  $\sigma_w^2$  is significant near the surface and negligible above 500 hpa. At 900 hPa, where mostly liquid clouds are formed,  $\sigma_w$  ranges between 0.1 and  $0.7 \text{ ms}^{-1}$  and is typically lower over the ocean than over land (Fig. 4). High  $\sigma_w$  is however found in the storm track regions of the Southern and Northern Hemispheres. At this vertical level  $\sigma_w$  is the lowest in the Arctic region ( $\sim 0.1 \text{ ms}^{-1}$ ). The range of  $\sigma_w$  shown in Fig. 4 is in good agreement with in situ measurements of vertical velocity at cloud base in marine stratocumulus (Peng et al., 2005; Guo et al., 2008), and continental regions (Fountoukis et al., 2007; Tonttila et al., 2011), showing  $\sigma_w$  mostly between 0.2 and  $1 \text{ ms}^{-1}$ . However global measurements of  $\sigma_w$  have not been reported. Compared to similar schemes (e.g., Golaz et al., 2010) Eq. (22) results in higher velocities within the PBL since the characteristic length decreases near the surface, consistent with the vertical momentum balance within the PBL (Blackadar, 1962). Thus,  $\sigma_w^2$  rarely hits the prescribed minimum ( $\sim 0.01 \text{ ms}^{-1}$ ) within the PBL.

Gravity wave motion dominates the global distribution of  $\sigma_w$  at the 500 hPa and 150 hPa vertical levels, being typically larger over land than over ocean (Fig. 4). Air flowing over orographic features produces high frequency waves that propagate to

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

the free troposphere (Bacmeister et al., 1999; Herzog and Vial, 2001). Thus  $\sigma_w$  is the highest over the mountain ranges of Asia, South America, and the Antarctic. At 500 hPa,  $\sigma_w$  is about  $0.1 \text{ m s}^{-1}$  over land and may reach up to  $0.5 \text{ m s}^{-1}$  over mountain ranges. These values are in good agreement with in situ measurements (Gayet et al., 2004). A similar distribution of  $\sigma_w$  is found at 150 hPa, with values over land slightly higher than at 500 hPa. Over the ocean,  $\sigma_w$  is typically larger at 150 hPa than at 500 hPa, particularly over the Tropics, since gravity waves in these regions can reach larger amplitudes before breaking. Figure 4 shows that  $\sigma_w$  in the upper troposphere varies by up to three orders of magnitude around the globe. Such variability has important implications for the effect of IN emissions on cloud formation (Sect. 3.5).

### 3.4 Cloud droplet number concentration

Comparison of cloud droplet number concentration against satellite retrievals is typically challenging. Retrieval algorithms generally introduce assumptions on the droplet size distribution that may bias  $N_d$ . To compare satellite retrievals and model data over the same basis we take advantage of the COSP output to obtain a “model retrieved” column integrated droplet concentration,  $N_{d, \text{cum}}$ , in the form (Han et al., 1998),

$$N_{d, \text{cum}} = \frac{\tau}{2\pi R_{\text{eff, liq}}^2 (1 - b)(2 - b)} \quad (41)$$

where  $\tau$  is the liquid cloud optical depth and  $b = 0.193$  (Han et al., 1998). To apply Eq. (41),  $R_{\text{eff, liq}}$  and  $\tau$  are obtained either from the GEOS-5 COSP output or the MODIS retrieval. This procedure does not aim to produce an accurate retrieval of  $N_{d, \text{cum}}$  but rather to equally compare GEOS-5 and MODIS data. Equation (41) is applied between  $60^\circ \text{ S}$  and  $60^\circ \text{ N}$  where the MODIS retrieval is more reliable (Platnick et al., 2003).

Figure 5 shows the global distribution of  $N_{d, \text{cum}}$  from GEOS-5 and MODIS. GEOS-5 is able to capture the high  $N_{d, \text{cum}}$  found in regions of high sulfate emissions i.e.,

Europe, Central and South East Asia and the East Coast of North America. There is also agreement between MODIS and GEOS-5 in regions with high biomass burning emissions like Subsaharian Africa and South America. However the model tends to slightly underpredict  $N_{d, cum}$  in the remote Atlantic and Pacific Oceans. There is also underprediction of  $N_{d, cum}$  off the west coasts of North and South America and Africa. This is due to underprediction of shallow stratocumulus in GEOS-5 (Fig. 1) and because  $w_{sub}$  tends to be small in these regions (Fig. 4). The global mean  $N_{d, cum}$  in GEOS-5 ( $1.68 \text{ cm}^{-2}$ ) is in agreement with MODIS results ( $1.96 \text{ cm}^{-2}$ ). The influence of the CCN activation parameterization on  $N_{d, cum}$  is studied in Sect. 4.

### 3.5 Ice crystal number concentration

At any given  $T$ ,  $N_c$  varies by up to four orders of magnitude, although mostly within a factor of ten (Fig. 6, a). The mean  $N_c$  peaks around  $200 \text{ L}^{-1}$  at 225 K, decreasing to  $\sim 20 \text{ L}^{-1}$  at 190 K, and below  $\sim 1 \text{ L}^{-1}$  at 180 K. For  $T > 245 \text{ K}$   $N_c$  remains mostly below  $\sim 10 \text{ L}^{-1}$ . Global mean  $N_c$  is around  $66 \text{ L}^{-1}$  for all clouds and around  $166 \text{ L}^{-1}$  for cirrus ( $T < 235 \text{ K}$ ). Figure 6 shows agreement of GEOS-5 values with in situ measurements of  $N_c$  over the whole  $T$  interval (Krämer et al., 2009; Gultepe and Isaac, 1996). There is good agreement of GEOS-5 with field campaign data at  $T < 200 \text{ K}$  where most models show a large positive bias (e.g., Barahona et al., 2010; Salzmann et al., 2010; Gettelman et al., 2012). This results from the proper consideration of the effect of prior nucleation events on ice crystal nucleation (Sect. 3.5).  $N_c$  is also influenced by the presence of preexisting ice crystals; their effect is analyzed in Sect. 4.

The relative contribution of different mechanisms to the source of  $N_c$  is shown in Fig. 6. To facilitate comparison against in situ measurements of IN and  $N_c$ , integrated variables, instead of number tendencies, are used. Thus, the ice crystal concentration from ice nucleation in the deposition and condensation modes,  $N_{dep}$ , is calculated using Eq. (13) and the BN09 parameterization.  $N_c$  from immersion freeezing,  $N_{imm}$ , is calculated by integration of Eq. (17) over the time scale defined by  $\gamma_c$ . The

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



concentration of detrained ice crystals,  $N_{c, cv}$ , is given by the ice crystal concentration at cloud top calculated by Eq. (29).

$N_{dep}$  varies mostly within 0.1 and  $50 L^{-1}$ , and is the largest around 240 K where the aerosol concentration is large enough to result in significant IN concentration (Fig. 6b).

There is however large variability in  $N_{dep}$  around the globe. Most deposition IN come from dust although the concentration of black carbon IN may be significant reaching  $2 L^{-1}$  at  $T \sim 230$  K (not shown). A few deposition IN ( $\sim 1 L^{-1}$ ) are found at  $T$  as high as 260 K mostly in regions of large dust concentration.

$N_{imm}$  reaches up to  $40 L^{-1}$  around 240 K but decreases rapidly for lower  $T$  where it is prevented by the homogeneous freezing of cloud droplets (Fig. 6c). In agreement with in situ observations of mixed-phase clouds (e.g., DeMott et al., 2010) immersion freezing IN are scarce above 250 K, with typical concentrations below  $0.1 L^{-1}$ . Dust is the most important source of immersion IN, whereas black carbon IN typically contribute less than  $2 L^{-1}$  to  $N_c$ . Contact freezing IN are not explicitly shown in Fig. 6 but they follow a similar tendency as immersion freezing IN, although with lower concentration.

$N_{c, cv}$  remains below  $50 L^{-1}$  for  $T > 240$  K, characteristic of heterogeneous ice nucleation. For  $T > 250$  K,  $N_{c, cv}$  reaches up to  $10 L^{-1}$  mostly from immersion and contact freezing of supercooled droplets within the convective cumulus (Fig. 6d). Homogeneous freezing of cloud droplets is evident by the strong increase in  $N_{c, cv}$  around  $T \sim 240$  K which in some instances may reach up to  $10 cm^{-3}$ . Such very high  $N_{c, cv}$  is responsible for the highest values of  $N_c$  in Fig. 6. Along with immersion freezing, detrainment from convective cumulus determines  $N_c$  for  $T > 240$  K.

The predominance of heterogeneous ice nucleation in cirrus is analyzed in Fig. 7. Globally about 70% of the production of ice crystals in cirrus proceeds by homogeneous freezing with a clear contrast between the Northern (NH) and the Southern (SH) Hemispheres. Homogeneous freezing is most prevalent in SH and only leeward of South America and Africa the contribution of heterogeneous freezing is significant ( $\sim 30\%$ ). In contrast, most of NH is influenced by IN emissions which

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



in some cases dominate crystal production. Part of the contrast between NH and SH is explained by the greater abundance of dust in NH. However comparison of Figs. 4 and 7 also reveals a marked effect of  $\sigma_w$  on  $N_c$ . Low  $\sigma_w$  tends to enhance the effect of IN on  $N_c$  because of the greater residence time of the heterogeneously-frozen ice crystals in each parcel and the lower rate of increase of supersaturation (Barahona and Nenes, 2009a). Thus heterogeneous freezing tends to dominate ice crystal production in regions of low  $\sigma_w$  like Sub-Saharan Africa, the Arctic, and the west coast of North America, even though these regions are not characterized by high emission rates of IN. This result is also consistent with the study of Cziczo et al. (2013) who found predominance of heterogeneous ice nucleation in these regions. Globally however homogeneous ice nucleation dominates ice crystal production. This suggests that variability in  $\sigma_w$  plays a significant role in defining the effect of IN emissions on cirrus formation.

### 3.6 Cloud liquid and ice water

The implementation of the new microphysics resulted in significant improvement of the representation of ice and liquid water content in GEOS-5. Figure 8 shows the zonal mean ice mass mixing ratio,  $q_i$ , from the NEW and CTL simulation compared to the CloudSat retrieval for non-convective, non-precipitating ice (Li et al., 2012). The global distribution of  $q_i$  in the NEW simulation is in better agreement with the satellite retrieval than that obtained in CTL. The excessive freezing around  $T = 235$  K, characterized by the bulls-eye pattern around 600 hPa in the CTL run, is not present in the NEW simulation. In absolute terms,  $q_i$  in the NEW and CTL runs is generally lower than CloudSat data although mostly within the intrinsic error of the retrieval, about a factor of two (Li et al., 2012; Eliasson et al., 2011). Including snow in the comparison (Fig. 8) still results in lower ice + snow concentration than in CloudSat, although within the error of the retrieval.

Figure 9 shows the zonal mean liquid mass mixing ratio  $q_l$  from GEOS-5 for the CTL and NEW runs compared against the CloudSat retrieval for non-convective, non-

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion





Central Asia, and over the mountain Ranges of Africa, and North and South America. The high IWP of the latter regions results in part from strong ice crystal production over mountain ranges (Sect. 3.5). GEOS-5 however underestimates IWP in the Tropical Western Pacific Ocean. The spatial distribution of total water path (liquid + ice) is similar as obtained with CloudSat, although the global mean TWP is higher in GEOS-5 ( $\sim 64 \text{ gm}^{-2}$ ) than in the retrieval ( $\sim 49 \text{ gm}^{-2}$ ) due to the larger LWP in GEOS-5.

### 3.7 Supercooled cloud fraction

Figure 11 shows the supercooled cloud fraction (e.g., the fraction of cloud condensate present as liquid,  $\text{SCF} = 1 - f_{\text{ice}}$ ) in mixed-phase clouds for the CTL and NEW simulations. In the CTL simulation the total condensate is linearly partitioned into liquid and ice between 235 K and 270 K (Bacmeister et al., 2006). In the NEW simulation partitioning of the condensate is carried out taking into account the activity and concentration of IN and the Bergeron–Findeisen process. In CTL most values of SCF below 260 K follow the prescribed linear tendency. Variability in SCF increases strongly above 260 K due to the freezing of condensate at 273 K and ice-enhanced precipitation (Fig. 11). The tendency of SCF with  $T$  in NEW shows different features than in CTL following a sigmoidal instead of a linear tendency. This behavior has been observed in satellited retrievals and field campaigns (Choi et al., 2010; Hu et al., 2010) and is characteristic of immersion freezing mediated mainly by dust (e.g., Murray et al., 2011; Marcolli et al., 2007). The region of maximum SCF frequency in Fig. 11 however expands about 10 K, which results from variation in particle size and concentration, the presence of black carbon IN, enhanced precipitation in mixed-phase clouds, and variation in  $\sigma_w$ . There is also a higher frequency of  $\text{SCF} > 0.4$  for  $T < 255 \text{ K}$  in the NEW than in the CTL simulation which results from a higher fraction of supercooled liquid in the convective detrainment in NEW than in CTL.

Compared with CALIOP, SCF in NEW is shifted by about 6 K towards higher  $T$ , which implies that clouds tend to glaciate at higher  $T$  in the model than observed by the satellite. This would indicate higher IN activity (i.e., higher dust concentration or more

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



active dust) in GEOS-5 than implied by the CALIOP data. However CALIOP is sensitive mostly to cloud-top properties and SCF may be biased low in deep convective clouds where most of the supercooled liquid is found below cloud top (Hu et al., 2010). The influence of these factors on SCF requires more investigation and will be undertaken in a future study. Still the sigmoidal increase of SCF with  $T$  in both GEOS-5 and the satellite retrieval indicates that SCF is significantly influenced by the presence of IN.

### 3.8 Cloud droplet and ice crystal effective radii

The annual mean droplet effective radius  $R_{\text{eff, liq}}$  from the NEW simulation ( $14.3\ \mu\text{m}$ ) is in agreement with MODIS retrievals ( $14.8\ \mu\text{m}$ ) (Fig. 12). This is higher than the prescribed mean for the CTL run and simulated by other models also using the MG08 stratiform microphysics ( $\sim 9\text{--}11\ \mu\text{m}$ ) (Gettelman et al., 2008; Salzmänn et al., 2010) but similar to the one obtained in Sud et al. (2013) in GEOS-5. The results presented in Fig. 12 benefit from using the COSP package which accounts for the preferential cloud-top sampling of MODIS (Bodas-Salcedo et al., 2011). Other studies (Gettelman et al., 2008; Salzmänn et al., 2010) however did not use COSP for comparison. In agreement with the MODIS retrieval the spatial distribution of  $R_{\text{eff, liq}}$  in the NEW run shows a clear ocean-land contrast (Fig. 12).  $R_{\text{eff, liq}}$  is overestimated in the west coasts of South America, Africa and to a lesser extent, North America, due to low  $N_d$  over these regions. Over the land  $R_{\text{eff, liq}}$  is underestimated in South Central Asia, Europe and the West Coast of North America, likely due to the high concentration of cloud droplets predicted by GEOS-5 in these regions (Sect. 3.4).

The global distribution of ice effective radius,  $R_{\text{eff, ice}}$ , for the NEW run is presented in Fig. 13 along with MODIS retrievals. The global mean value of  $R_{\text{eff, ice}}$  in the NEW simulation ( $26\ \mu\text{m}$ , from COSP output) is in good agreement with the satellite ( $24.2\ \mu\text{m}$ ). GEOS-5 is able to reproduce the low  $R_{\text{eff, ice}}$  seen by MODIS over most of the large mountain ranges, e.g., over the Andean and Himalayan regions, although it tends to underestimate  $R_{\text{eff, ice}}$  over north east Asia. Low  $R_{\text{eff, ice}}$  is caused by

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Cloud microphysics  
in GEOS-5

D. Barahona et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



strong homogeneous freezing events with  $N_c > 1 \text{ cm}^{-3}$  in high orographic uplift (Fig. 7), although local convection may also have an effect on  $R_{\text{eff, ice}}$  as detrainment from deep convection tends to increase  $N_c$  (Sect. 3.5). There is some contrast in  $R_{\text{eff, ice}}$  between land and ocean in the MODIS retrievals which is captured by GEOS-5. However the model tends to overestimate  $R_{\text{eff, ice}}$  in the subtropical continental regions of NH and SH, which may be caused by underestimation of  $\sigma_w$  leading to low  $N_c$ .

There may be some uncertainty in the retrieval of  $R_{\text{eff, ice}}$ , particularly for optically thick clouds (Chiriaco et al., 2007). To further corroborate the GEOS-5 results, in situ observations of the volumetric ice crystal radius,  $R_{\text{vol, ice}} = \left( \frac{3q_i}{4\pi N_c \rho_i} \right)^{1/3}$ , are used.

Figure 14 shows  $R_{\text{vol, ice}}$  as a function of  $T$  along with a composite of in situ data from several field campaigns (Krämer et al., 2009; McFarquhar and Heymsfield, 1997). There is agreement between the field data and the model, particularly for  $T < 230 \text{ K}$  where both show a decrease in  $R_{\text{vol, ice}}$  with decreasing  $T$ . Around  $T \sim 230 \text{ K}$  the model tends to predict slightly higher  $R_{\text{vol, ice}}$  than the observations, although mostly within the spread of the data. The discrepancy may also be a result of crystal shattering in ice crystal probes which tends to increase measured  $N_c$  decreasing  $R_{\text{vol, ice}}$  (Krämer et al., 2009). The smooth transition in  $R_{\text{vol, ice}}$  at  $235 \text{ K}$  indicates that both homogeneous and heterogeneous ice nucleation significantly contribute to ice crystal formation at this temperature (Sect. 3.5). In agreement with observations (McFarquhar and Heymsfield, 1997)  $R_{\text{vol, ice}}$  increases steadily for  $T > 235 \text{ K}$ , which results from increasing vapor deposition rates and decreasing  $N_c$  as  $T$  increases (Sect. 3.5).

### 3.9 Annual mean diagnostics

Table 4 and Fig. 15 show the summary of the annual mean cloud properties analyzed in this work. Annual mean LWP is  $37.3 \text{ g m}^{-2}$ , and  $60 \text{ g m}^{-2}$  if the MODIS COSP simulator is used. LWP in NEW is higher than the CloudSat retrieval ( $23.0 \text{ g m}^{-2}$ ) (Li et al., 2013) mostly from higher LWP in the midlatitudes of the SH, and lower than MODIS retrievals ( $\sim 100 \text{ g m}^{-2}$ ). Ocean-only LWP is also lower than SSM/I output by about a factor of

## Cloud microphysics in GEOS-5

D. Barahona et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



two (not shown). LWP in GEOS-5 refers only to non-convective (anvil and stratiform) clouds and it is likely that the discrepancy with SSMI and MODIS originates from the consideration of convective clouds in the retrievals. IWP in NEW ( $27.1 \text{ gm}^{-2}$ ) is in better agreement with CloudSat ( $25.8 \text{ gm}^{-2}$ ) (Li et al., 2012) although GEOS-5 tends to overestimate IWP in the midlatitudes of SH and NH. Including snow in the comparison does not affect IWP in the Tropics but results in larger subtropical IWP in NEW than in CloudSat. Global mean LWP in CTL is higher ( $60.0 \text{ gm}^{-2}$ ) and IWP slightly lower ( $19.0 \text{ gm}^{-2}$ ) than in NEW.

The prescribed  $R_{\text{eff, liq}}$  and  $R_{\text{eff, ice}}$  in CTL are generally smaller than those retrieved by MODIS with a global mean bias of about  $-5 \mu\text{m}$  and  $-4 \mu\text{m}$  for  $R_{\text{eff, liq}}$  and  $R_{\text{eff, ice}}$ , respectively.  $R_{\text{eff, liq}}$  and  $R_{\text{eff, ice}}$  in NEW are closer to MODIS with a global bias of about  $-0.5 \mu\text{m}$  and  $2 \mu\text{m}$ , respectively (Table 4), well within the intrinsic error of the retrieval (King et al., 2003). Zonal mean  $R_{\text{eff, liq}}$  is however overestimated in the Northern Hemisphere from underestimation of  $N_d$  in oceanic regions (Sect. 3.4).

Global mean cloud fraction in the NEW simulation is higher than in CTL but still lower than ISCCP retrievals (Rossow and Schiffer, 1999). The higher  $f_c$  in NEW results from higher cloud coverage over continental regions (Sect. 3.1). There is good agreement between NEW and ISCCP cloud fraction in the continental midlatitudes and most of the underestimation in NEW originates in marine regions. However in these regions both the NEW and CTL simulations show agreement with the MODIS retrieval. The reason for the better agreement of GEOS-5 with MODIS than with ISCCP in these regions is however not clear but may be related to differences in the the cloud masks of ISCCP and MODIS (Pincus et al., 2012).

Global annual mean precipitation,  $P_{\text{tot}}$ , is lower in the NEW ( $2.72 \text{ mm d}^{-1}$ ) than in the CTL ( $2.85 \text{ mm d}^{-1}$ ) simulation and in better agreement with GPCP (Huffman et al., 1997) and CMAP (Xie and Arkin, 1997) observations ( $\sim 2.6 \text{ mm d}^{-1}$ ), although both simulations tend to overestimate  $P_{\text{tot}}$  in the Tropics. In SH the NEW simulation tends to predict  $P_{\text{tot}}$  higher than CMAP and lower than GPCP whereas CTL is in better agreement with GPCP data. In NH,  $P_{\text{tot}}$  in the NEW and CTL simulations is closer

to GPCP than to CMAP data, although in NEW it tends to be lower than the GPCP observations.

The global top of the atmosphere (TOA) net radiative balance is about  $+0.95 \text{ W m}^{-2}$  in the NEW simulation. The slight radiative imbalance in NEW results in part from the negative bias in stratocumulus cloud coverage in the NEW simulation (Sect. 3.1). The liquid cloud optical depth in NEW however agrees with MODIS data (Fig. 15) particularly over the Tropics. In CTL liquid clouds tend to be optically much thicker than MODIS observations (Fig. 15) which results from larger LWP and smaller  $R_{\text{eff, liq}}$  than the observations (Sects. 3.6 and 3.8). The higher optical depth in CTL leads to a more negative SWCF ( $-52.1 \text{ W m}^{-2}$ ) than in CERES and to a higher net radiative imbalance  $-2.4 \text{ W m}^{-2}$ . Long wave cloud effect (LWCF) is similar in the CTL and NEW runs ( $\sim 25.0 \text{ W m}^{-2}$ ) and in agreement with CERES data ( $26.2 \text{ W m}^{-2}$ ). Compared to MODIS ice cloud optical depth is however overestimated in CTL and underestimated in NEW. In NEW the low bias in ice optical depth is compensated by a positive bias in the high level cloud fraction (Sect. 3.1).

## 4 Sensitivity studies

Tables 3 and 4 present a summary of the sensitivity of GEOS-5 to different microphysical parameters. To study the sensitivity of cloud properties to the description of CCN activation, the parameterization of Abdul-Razzak and Ghan (2000) (hereafter, ARG) was implemented. ARG is based on a fit to the numerical solution of the equations of an ascending parcel written in terms of dimensionless parameters. Compared to the NEW run, the usage of ARG resulted in slightly higher  $N_d$  than with the FN05 formulation particularly over marine regions (run ARGACT, Fig. 5). The ARG parameterization also predicts higher droplet concentration in regions of high aerosol emissions like South East Asia and Southern Africa. Global mean  $R_{\text{eff, liq}}$  was lower in ARGACT than in NEW by about  $0.7 \mu\text{m}$  leading to about  $2 \text{ W m}^{-2}$  more negative

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



SWCF. LWP and cloud fraction remained almost the same as in NEW suggesting that the change in SWCF was driven by modification of cloud albedo.

The sensitivity of cloud properties to the characteristic cirrus scale,  $L_c$ , was also investigated.  $L_c$  is associated with the wave length of the highest frequency waves leading to cloud formation (Eq. 24), although it is considered a free parameter. Increasing  $L_c$  from 100 m to 400 m reduced global  $N_c$  by about a factor of two (run LC400). The global mean  $R_{\text{eff, ice}}$  increased by about  $3 \mu\text{m}$  and LWCF decreased by  $2 \text{W m}^{-2}$ . The higher  $L_c$  led to smaller  $\sigma_w$  (Eq. 24) decreasing the rate of ice crystal formation. Global mean  $\sigma_w$  for  $L_c = 400 \text{ m}$  is about  $0.07 \text{ ms}^{-1}$  and  $0.11 \text{ ms}^{-1}$  at 500 hPa and 150 hPa, respectively, about half the value obtained in the NEW simulation (Fig. 4). These values are still within the observed values in field campaigns (e.g., Gayet et al., 2004), and more observations are needed to better constraint  $L_c$ . Table 4 however shows that GEOS-5 results are robust to moderate changes in  $\sigma_w$ .

The effect of the dispersion in the ice crystal size distribution,  $\mu_i$ , on ice cloud properties (Table 4) was analyzed by setting  $\mu_i = 0.0$  (run MUIZERO) instead of using a temperature dependent parameterization for  $\mu_i$  (Sect. 2.3). This led to about a factor of two lower IWP and  $R_{\text{eff, ice}}$  than in NEW, which resulted from an increase in autoconversion and accretion of ice by snow at low  $T$  (not shown). Despite the lower IWP, the lower ice crystal size increased the ice cloud optical depth and resulted in slightly higher LWCF and SWCF than in the NEW simulation. Because of this compensating effect the radiative properties of ice clouds are robust to moderate changes in the ice crystal size distribution. Decreasing the critical size for ice autoconversion from  $400 \mu\text{m}$  to  $200 \mu\text{m}$  (run DCS200) also increased ice autoconversion leading to lower IWP than in NEW.  $R_{\text{eff, ice}}$  was also reduced although to a lower extent than in MUIZERO. Thus the net radiative effect of reducing  $D_{\text{cs}}$  was a decrease of about  $\sim 6 \text{W m}^{-2}$  in LWCF.

Several studies were performed to investigate the sensitivity of GEOS-5 to the description of heterogeneous ice nucleation. In NOBC and NOGLASS the effect of black carbon and glassy IN, respectively, was switched off. These runs suggested

GMDD

6, 5289–5373, 2013

## Cloud microphysics in GEOS-5

D. Barahona et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## Cloud microphysics in GEOS-5

D. Barahona et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



that black carbon and glassy IN only have a subtle effect on global climate (Table 4), although their local effects may be significant. In particular black carbon IN tend to increase LWCF in regions of high aerosol emissions like East Asia and the East Coast of North America. In the same regions glassy IN tend to reduce  $N_c$  at low  $T$  (Fig. 16).

5 The global TOA radiative imbalance due to black carbon and glassy IN amounts to  $-0.05 \text{ W m}^{-2}$  and  $-0.18 \text{ W m}^{-2}$ , respectively. Although these values are comparable to other published studies (Gettelman et al., 2012) they must be taken with caution since they are based on limited results. A comprehensive description of the aerosol indirect effect in GEOS-5 will be addressed in future studies.

10 In the PDA08 run the Phillips et al. (2008) (hereafter Ph08) ice nucleation spectrum was used. Ph08 was employed in previous studies to study the effect to the ice nucleation spectrum on  $N_c$  (Barahona et al., 2010; Morales Betancourt et al., 2012; Liu et al., 2012). The main difference between Ph13 and Ph08 is that Ph08 accounts for the effect of organic material acting as IN (although their effect may be overestimated in Ph08, Phillips et al., 2013). Using the Ph08 parameterization reduced  $N_c$  increasing  $R_{\text{eff, ice}}$  by about  $1 \mu\text{m}$ , slightly decreasing LWCF. This resulted in part from the effect of organic IN inhibiting homogeneous freezing in cirrus clouds. Other cloud properties remained similar as in NEW.

20 The effect of preexisting ice crystals on ice crystal formation was analyzed in NOPREEX where it was assumed that  $N_{i, \text{pre}} = 0$ . For this run  $N_c$  was about twice as in NEW, with the greater increase occurring between 200 K and 240 K (Fig. 16), and mostly in the Tropics (not shown) indicating that the presence of ice crystals from convective detrainment tends to inhibit new ice nucleation events. Mean  $R_{\text{eff, ice}}$  was reduced by about  $6 \mu\text{m}$  increasing LWCF by  $5 \text{ W m}^{-2}$ .

25 In NOCNV the generation of precipitation in cumulus convection was described by a single-moment approach (Bacmeister et al., 2006). Some studies (e.g., Gettelman et al., 2008; Salzmann et al., 2010) did not consider explicitly the freezing and activation of aerosol particles in convective cumulus. Thus it is important to study how this assumption would affect GEOS-5 results. In NOCNV the contribution of convective

## Cloud microphysics in GEOS-5

D. Barahona et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



detrainment to ice crystal and droplet number concentration was approximated by assuming a fixed droplet size of  $10\mu\text{m}$  for droplets and using the correlation of McFarquhar and Heymsfield (1997) to obtain the ice crystal size as a function of  $T$ . Compared to NEW, the single-moment approach resulted in enhanced precipitation rates, particularly over the Tropical Warm Pool. SWCF and LWCF were lower than in NEW by about  $3\text{Wm}^{-2}$ , which was in part the result of a lower detrainment flux of condensate in the Tropical upper troposphere.  $R_{\text{eff, liq}}$  decreased by about of  $1\mu\text{m}$  due to an increase in droplet number concentration. Mean  $R_{\text{eff, ice}}$  only changed by about  $0.5\mu\text{m}$ , however  $N_c$  was slightly increased, particularly at low  $T$  (Fig. 16).

Finally it is important to analyze the effect of microphysical parameters on  $N_c$  at low  $T$ . Figure 16 shows the temperature dependency of  $N_c$  for the runs of Table 4. All curves of Fig. 16 show the same characteristics, increasing  $N_c$  with decreasing  $T$  to a maximum around  $210\text{K}$  and then decreasing to values typically below  $10\text{L}^{-1}$  at  $185\text{K}$ . The only exception to the latter is the NOCNV run in which mean  $N_c$  is about  $140\text{L}^{-1}$  at  $185\text{K}$ , resulting from the lower detrained  $N_c$  acting as preexisting ice crystals at low  $T$ . The maximum  $N_c$  is around  $300\text{L}^{-1}$  for most runs and only for the NOPREEX run it increases up to  $800\text{L}^{-1}$ . The fact that in all runs  $N_c$  decreases for  $T$  below  $200\text{K}$  indicates that as the  $T$  decreases  $N_c$  becomes more dependent on  $S_{\text{crit}}$  (Sect. 3.2). This indicates that parcel history plays a primary role in determining  $N_c$  at low  $T$  whereas preexisting ice crystals and IN only play a secondary role.

## 5 Summary and conclusions

A new cloud microphysics scheme was developed for the the NASA GEOS-5 global atmospheric model. The main features of the new microphysics are:

- A comprehensive two-moment microphysics description for stratiform clouds (Morrison and Gettelman, 2008).

## Cloud microphysics in GEOS-5

D. Barahona et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Consistent coupling of the cloud fraction and stratiform condensation with the microphysics. The stratiform condensation scheme was modified to allow supersaturation in ice clouds.
- 5 – A two-moment microphysics scheme embedded within the RAS convective parameterization. The new scheme explicitly treats the formation of droplets and ice crystals, the partitioning of condensate between ice and liquid, and the generation of precipitation within convective cumulus.
- 10 – A comprehensive description of cloud droplet activation and ice nucleation in stratiform and convective clouds, linked to the aerosol physicochemical properties. The description of ice formation considers homogeneous freezing of cloud droplets and interstitial aerosol as well as heterogeneous ice nucleation on ice nuclei. Competition between homogeneous and heterogeneous ice nucleation, and between different ice nuclei is explicitly treated. Immersion, contact, condensation and deposition ice nucleation modes are considered.
- 15 – Explicit calculation of the critical saturation ratio for ice formation considering aerosol properties, temperature and subgrid scale dynamics.
- Explicit parameterization of the effect of preexisting ice crystals on ice nucleation.
- Explicit parameterization of the distribution of subgrid scale vertical velocity in stratiform clouds, accounting for the effect turbulence and gravity wave motion on the vertical velocity variance. A new parameterization in terms of large scale variables was developed for the latter.

25 The new microphysics was evaluated against satellite retrievals and field campaign data. Usage of the COSP satellite simulator greatly facilitated the comparison against satellite observations, reducing the uncertainty in the sampling of the model results. In general cloud microphysical fields like ice water, liquid water content and droplet and ice crystal size were in much better agreement with observations than obtained with the

operational version of GEOS-5. The model performance in reproducing the observed total cloud fraction and longwave and shortwave cloud forcings is also improved and is in reasonable agreement with satellite observations.

In the new microphysics ice and cloud droplet nucleation are tightly linked to the evolution of the cloud properties. Cloud droplet number impacts the formation of precipitation. Precipitation decreases total water which in turn feeds back into the cloud fraction through modification of  $P_q(q_t)$  (Sect. 2.3.1). The link between  $N_c$ ,  $f_c$ , and  $q_i$  is stronger since the production of condensate is controlled in part by  $S_{crit}$  which depends on the presence of IN (Eq. 15). The linkage between cloud micro and macro physical variables in the model emphasizes the internal consistency of the new cloud scheme.

A new cloud coverage scheme was developed to allow supersaturation with respect to the ice phase. The frequency and spatial distribution of supersaturation simulated by the model was in good agreement with satellite and in situ observations. It was shown that supersaturation is controlled in part by ice crystal nucleation and the value of  $S_{crit}$ . The latter dictates the minimum water vapor threshold required for cloud formation.  $S_{crit}$  is highly variable over the globe, and dependent on aerosol concentration and temperature. Thus models that assume a single threshold for ice cloud formation are inherently biased.

The variation of supercooled cloud fraction with temperature in the new microphysics followed a sigmoidal tendency. This is in agreement with CALIOP data (Choi et al., 2010) and differs from the typical linear increase of SCF with  $T$  assumed in most GCMs. There are no temperature-based constraints to the occurrence of the Bergeron–Findeisen process nor to the partition of total condensate between ice and liquid in the new microphysics. The sigmoidal tendency in SCF resulted from explicit consideration of homogeneous, immersion and contact freezing in the model. This suggests that rather than temperature alone, the presence of IN greatly influences the frequency of supercooled liquid in mixed-phase clouds.

A new approach was proposed to parameterize the distribution of subgrid scale vertical velocity in cirrus and stratocumulus which takes into account turbulence and

Cloud microphysics  
in GEOS-5

D. Barahona et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## Cloud microphysics in GEOS-5

D. Barahona et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



gravity wave motion. Although no studies have been reported on the global distribution of  $\sigma_w$  the parameterization results were within reported values in field campaigns. Since the parameterization proposed here focuses on surface and orographic stresses, which are higher over the land,  $\sigma_w$  may be underestimated in the upper troposphere in oceanic regions. Still the ability to predict  $\sigma_w$  as a function of large scale variables points in the right direction to reduce one of the main sources of uncertainty in the modeling of the effect of aerosol emissions on climate. It was also shown that the variability in  $\sigma_w$  is a determining factor defining the effect of IN emissions on cirrus formation.

The simulated ice crystal concentration was in agreement with field campaign data, even at very low  $T$  where most models tend to overestimate  $N_c$  (e.g., Barahona et al., 2010; Salzmann et al., 2010; Hendricks et al., 2011). In GEOS-5 the decrease of  $N_c$  with decreasing  $T$  results from an increase in  $S_{\text{crit}}$  (Fig. 2) which limits  $P_q(q_t > S_{\text{crit}} q_i^*)$  at low  $T$  decreasing the probability of homogeneous freezing events. The term  $P_q(q_t > S_{\text{crit}} q_i^*)$  in Eq. (16) provides a link between current cloud formation and prior ice nucleation events (Barahona and Nenes, 2011). This suggests that a statistical rather than a single-parcel approach (e.g., Jensen et al., 2012; Spichtinger and Cziczo, 2010) is required for the correct modeling of low temperature cirrus.

A new parameterization of the effect of preexisting ice crystals on ice cloud formation was developed. It was shown that their effect is more pronounced for  $T$  around 200 K, typically reducing  $N_c$ . However preexisting ice crystals alone can not explain the low ice crystal concentration at low  $T$ . The effect of organic glassy IN on cloud formation was also analyzed and it was found that it tends to reduce  $N_c$  at low temperature. Although these factors alone cannot explain the tendency of  $N_c$  at  $T < 190$  K, they are still necessary to reproduce the observed  $N_c$  in the upper troposphere. In fact it was found that the observed values of ice crystal concentration in the upper troposphere result from the combination of several factors: parcel history, IN concentration, convective detrainment and subgrid dynamics.

Effective cloud droplet size simulated with GEOS-5 was in agreement with the MODIS retrieval. There was however a slight underestimation in  $R_{\text{eff, liq}}$  over the land

## Cloud microphysics in GEOS-5

D. Barahona et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



and overestimation over the Tropical marine regions. This points to the need for a more sophisticated description of aerosol microphysics in GEOS-5. Sensible assumptions were made regarding the aerosol size distribution, however there is a high variability in the aerosol properties around the globe which may affect CCN activation. The inclusion

5 of a more comprehensive aerosol microphysics in GEOS-5 will be addressed in a future study. The simulated cloud droplet number concentration also showed some sensitivity to the parameterization of CCN activation, which in turn influences the cloud albedo.

There was good agreement in the global mean ice effective radius between GEOS-5 and the MODIS retrieval. The decrease in  $R_{vol, ice}$  as  $T$  decreases, a common feature of in situ observations (Krämer et al., 2009) was also captured by GEOS-5. The model was able to capture key features of the spatial distribution of  $R_{eff, ice}$ , as for example the predominance of low  $R_{eff, ice}$  near mountain ranges. This was a result of the explicit consideration of ice nucleation and of the spatial variation of  $\sigma_{w, gw}$ .  $R_{eff, ice}$  was however overestimated in marine regions, particularly in the Southern Hemisphere.

15 The parameterization of  $\sigma_{w, gw}$  developed in this work may underestimate  $\sigma_w$  over the ocean. Other IN sources like biological particles (Burrows et al., 2013) and sea salt (Wise et al., 2012) were not considered in this study but may enhance ice nucleation in marine environments. Some uncertainty may be introduced by the single-moment approach used for the aerosol microphysics in GEOS-5 ice nucleation, although ice

20 nucleation is less dependent on aerosol size than CCN activation. Mixing of dust with sulfate may lead to IN deactivation and is currently not modeled by GEOS-5. The role of the uncertainty in the satellite retrieval must also be taken into account. All of these effects require further investigation. Nevertheless, the approach proposed here results in a realistic and reasonable spatial distribution of  $R_{eff, ice}$ .

25 It was shown that the cloud radiative fields modeled in GEOS-5 with new microphysics are in good agreement with observations, although local biases may be significant. GEOS-5 tends to underestimate the optical depth of persistent stratocumulus decks which leads to a negative radiative bias in the Western Pacific. Reducing such bias requires an explicit representation of shallow cumulus

condensation in GEOS-5. The long-term and large-scale climatic response of GEOS-5 with the new microphysics will be analyzed in a future study.

A simple approach was assumed to describe the cloud microphysics in convective clouds. The description of precipitation within convective cores is highly complex due to the interplay of several cloud species (e.g., graupel, hail, rain, ice and snow). Some authors have developed more comprehensive microphysical packages for convective clouds including processes of autoconversion, aggregation, collection and accretion (e.g., Song and Zhang, 2011; Sud and Walker, 1999; Lohmann, 2008). To be effective, a detailed description of microphysics in convective clouds requires prognostic prediction of the vertical profiles of rain and snow which is not implemented in most GCMs. Also collection and aggregation rates depend on the vertical profiles of rain and snow which are not known in advance. Thus the advantages of a complex representation of the microphysics of convective cores must be weighted against the uncertainty introduced in accommodating such descriptions within the diagnostic integration schemes of the convective parameterizations in GCMs.

The model results were quite robust to variation in microphysical parameters. The largest differences from the base configuration were found for a decrease in the size dispersion parameter of the ice crystal size distribution and in the critical size for ice autoconversion. Both changes lead to a reduction in  $R_{\text{eff, ice}}$  and IWP and modified the long wave cloud forcing. The high sensitivity of  $R_{\text{eff, ice}}$  and IWP to the value of  $\mu_i$  suggests that more attention must be put on its correct parameterization in GCMs.

The implementation of the comprehensive microphysics developed in this work resulted in a more realistic simulation of cloud properties in GEOS-5. Mounting evidence suggests that the explicit description of processes of droplet and ice crystal nucleation and precipitation is necessary for the correct representation of clouds in Earth system models. The new microphysics would likely result in improved and more realistic climate simulations in GEOS-5. The new parameterizations developed here may also help to improve our understanding of the role of microphysics and aerosol emissions on the evolution of clouds. Within the larger picture, the further development

GMDD

6, 5289–5373, 2013

## Cloud microphysics in GEOS-5

D. Barahona et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



of the microphysics GEOS-5 will help to understand the role of clouds on climate and eventually reduce the uncertainty in their prediction.

## Appendix A

### Parameterization of $\sigma_{w,gw}$

5 Parameterizations of the subgrid vertical velocity from gravity wave motion consider either the displacement of a single wave from orographic uplift (Joos et al., 2008; Dean et al., 2007) or the spectrum of velocities resulting from the superposition of waves from different sources (Barahona and Nenes, 2011; Jensen and Pfister, 2004). The characteristic scale of gravity wave motion leading to the formation of clouds is typically  
 10 smaller than the scale of the GCM grid cell. Thus a spectrum of vertical velocities rather than a single wave may be a more realistic representation of the subgrid dynamics in the upper troposphere. Still surface perturbations are likely to determine the maximum  $w_{sub}$  in the spectrum (Joos et al., 2010; Barahona and Nenes, 2011). Using this concept a semi-empirical parameterization for  $\sigma_{w,gw}$  can be developed as follows.

15 The mean vertical momentum flux at the surface (McFarlane, 1987) is given by,

$$\tau = \frac{1}{2} k \rho_a U_s N_s \delta h_s^2 \quad (A1)$$

where  $\delta h_s$  is the vertical displacement at the surface caused by the orographic perturbation,  $N_s$  the Brunt–Väisälä frequency at the surface and  $U_s$  the surface wind (taken as the geometrical mean between the meridional and zonal components), and  $k$   
 20 is the horizontal wave number. Equating  $\tau$  to the mean surface stress,  $\tau_0$ , and scaling  $\delta h$  according to McFarlane (1987) i.e.,  $\delta h = \delta h_s [\rho_a U_s N_s / \rho_a U N]^{1/2}$ , the mean wave amplitude,  $\delta h$ , at any height can be written as

$$\delta h^2 = \min \left( \frac{2|\tau_0|}{k\rho_a U N}, \frac{U}{N} \right) \quad (A2)$$

## Cloud microphysics in GEOS-5

D. Barahona et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



where  $\frac{U}{N}$  is the saturation wave amplitude (Dean et al., 2007). The maximum vertical velocity in the gravity wave spectrum is related to  $\delta h$  by (Joos et al., 2008)

$$w_{\max} = kU\delta h \quad (\text{A3})$$

In a spectrum of randomly superimposed gravity waves,  $w_{\max}$  can be empirically related to  $\sigma_{w,\text{gw}}$  by (Barahona and Nenes, 2011)

$$\sigma_{w,\text{gw}} \approx 0.133w_{\max} \quad (\text{A4})$$

making  $k = \frac{2\pi}{L_c}$  and combining Eqs. (A2) to (A4), we obtain.

$$\sigma_{w,\text{gw}}^2 = 0.0169 \min \left[ \frac{4\pi U |\tau_0|}{\rho_a L_c N}, \left( \frac{2\pi U^2}{NL_c} \right)^2 \right] \quad (\text{A5})$$

where  $L_c$  is associated with the wave length of the highest frequency waves in the spectrum, typically between 50 m and 500 m (Bacmeister et al., 1999), although considered a free parameter.

## Appendix B

### Parameterization of the effect of preexisting crystals on ice nucleation

Water vapor deposition onto ice crystals left from previous nucleation events decreases supersaturation and may reduce  $N_c$ , particularly at low temperature (Barahona and Nenes, 2011). To account for this effect the local rate of change of  $S_i$  in a cloudy parcel with preexisting crystals is written in the form (Barahona and Nenes, 2011),

$$\frac{dS_i}{dt} = \alpha w_{\text{sub}} S_i - \beta \frac{dq_{i,\text{nuc}}}{dt} - \beta \frac{dq_{i,\text{pre}}}{dt} \quad (\text{B1})$$

where  $\alpha$  and  $\beta$  are temperature-dependent parameters (Appendix Table A1), and  $\frac{dq_{i, \text{nuc}}}{dt}$  and  $\frac{dq_{i, \text{pre}}}{dt}$  are the local rates of ice crystal growth of recently nucleated and preexisting ice crystals, respectively. The latter is given by,

$$\frac{dq_{i, \text{pre}}}{dt} = \frac{N_{i, \text{pre}} \pi \beta c \rho_i A_i (S_i - 1)}{2\lambda_{i, \text{pre}}} \quad (\text{B2})$$

5 where it was assumed that the size of preexisting ice crystal follows a gamma distribution (Eq. 2). Introducing Eq. (B2) into Eq. (B1) we obtain,

$$\frac{dS_i}{dt} = \alpha w_{\text{sub}} S_i - \beta \frac{dq_{i, \text{nuc}}}{dt} - \beta \frac{N_{i, \text{pre}} \pi \beta c \rho_i A_i (S_i - 1)}{2\lambda_{i, \text{pre}}} \quad (\text{B3})$$

Ice crystal nucleation in cirrus occurs over small  $S_i$  intervals (Barahona and Nenes, 2008; Kärcher and Lohmann, 2002). Therefore to a good approximation the size of preexisting ice crystals can be considered constant during ice nucleation. With this  
10 assumption, Eq. (B3) can be reorganized as,

$$\frac{dS_i}{dt} = \alpha w_{\text{sub}} S_i \left[ 1 - \frac{N_{i, \text{pre}} \pi \beta c \rho_i A_i (S_{\text{hom}} - 1)}{2\lambda_{i, \text{pre}} \alpha w_{\text{sub}} S_{\text{hom}}} \right] - \beta \frac{dq_{i, \text{nuc}}}{dt} \quad (\text{B4})$$

where it was assumed that  $\frac{S_i - 1}{S_i} \approx \frac{S_{\text{hom}} - 1}{S_{\text{hom}}}$ . If  $N_{i, \text{pre}} = 0$  then Eq. (B4) reduces to the saturation balance of a parcel with no preexisting crystals present (Barahona  
15 and Nenes, 2008). Thus the effect of preexisting crystals on ice nucleation can be accounted for by redefining the cloud scale vertical velocity in the form,

$$w_{\text{sub, pre}} = w_{\text{sub}} \max \left[ 1 - \frac{N_{i, \text{pre}} \pi \beta c \rho_i A_i (S_{\text{hom}} - 1)}{2\lambda_{i, \text{pre}} \alpha w_{\text{sub}} S_{\text{hom}}}, 0 \right] \quad (\text{B5})$$

Equation (B5) shows that the effect of water vapor deposition onto preexisting crystals can be understood as a reduction in the rate of increase of supersaturation by expansion cooling. Since  $w_{\text{sub}}$  is typically an input to ice cloud formation parameterizations, Eq. (B5) also provides a simple way of accounting for the effect of preexisting ice crystals on ice cloud formation, independently of the ice nucleation parameterization employed.

*Acknowledgements.* Donifan Barahona was supported by the NASA Modeling, Analysis and Prediction program under WBS 802678.02.17.01.07. GPCP Precipitation data provided by the NOAA/OAR/ESRL PSD, Boulder, Colorado, USA, from their website at <http://www.esrl.noaa.gov/psd/>. MODIS data was downloaded from <http://modis.gsfc.nasa.gov/data/>. We thank Frank Li for providing level 3 CloudSat data. The authors thank Yong-Sang Choi for providing CALIOP-derived SCF data, and Minghuai Wang for his help with the analysis of field campaign data.

*Code availability:* The GEOS-5 source code is available under the NASA Open Source Agreement at <http://opensource.gsfc.nasa.gov/projects/GEOS-5/>.

## References

Abdul-Razzak, H. and Ghan, S.: A parameterization of aerosol activation, 2. Multiple aerosol types, *J. Geophys. Res.*, 105, 6837–6844, doi:10.1029/1999JD901161, 2000. 5326, 5354, 5362

Arakawa, A.: The cumulus parameterization problem: past, present, and future, *J. Climate*, 17, 2493–2525, 2004. 5293

Bacmeister, J., Eckermann, S., Tsias, A., Carslaw, K., and Peter, T.: Mesoscale temperature fluctuations induced by a spectrum of gravity waves: a comparison of parameterizations and their impact on stratospheric microphysics, *J. Atmos. Sci.*, 56, 1913–1924, 1999. 5292, 5304, 5305, 5311, 5317, 5336

Bacmeister, J., Suarez, M., and Robertson, F. R.: Rain reevaporation, boundary-layer convection interactions, and Pacific rainfall patterns in an AGCM, *J. Atmos. Sci.*, 63, 3383–3403, doi:10.1175/JAS3791.1, 2006. 5294, 5322, 5328

Baker, B. A.: On the role of phoresis in cloud ice initiation, *J. Atmos. Sci.*, 48, 1545–1548, doi:10.1175/1520-0469(1991)048<1545:OTROPI>2.0.CO;2, 1991. 5303

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



---

**Cloud microphysics  
in GEOS-5**


---

D. Barahona et al.

---

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[◀](#)
[▶](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)


- Barahona, D.: On the ice nucleation spectrum, *Atmos. Chem. Phys.*, 12, 3733–3752, doi:10.5194/acp-12-3733-2012, 2012. 5301
- Barahona, D. and Nenes, A.: Parameterization of cloud droplet formation in large scale models: including effects of entrainment, *J. Geophys. Res.*, 112, D16026, doi:10.1029/2007JD008473, 2007. 5300
- Barahona, D. and Nenes, A.: Parameterization of cirrus formation in large scale models: homogeneous nucleation, *J. Geophys. Res.*, 113, D11211, doi:10.1029/2007JD009355, 2008. 5296, 5300, 5315, 5337
- Barahona, D. and Nenes, A.: Parameterizing the competition between homogeneous and heterogeneous freezing in cirrus cloud formation – monodisperse ice nuclei, *Atmos. Chem. Phys.*, 9, 369–381, doi:10.5194/acp-9-369-2009, 2009a. 5300, 5306, 5320
- Barahona, D. and Nenes, A.: Parameterizing the competition between homogeneous and heterogeneous freezing in ice cloud formation – polydisperse ice nuclei, *Atmos. Chem. Phys.*, 9, 5933–5948, doi:10.5194/acp-9-5933-2009, 2009b. 5300, 5301, 5302
- Barahona, D. and Nenes, A.: Dynamical states of low temperature cirrus, *Atmos. Chem. Phys.*, 11, 3757–3771, doi:10.5194/acp-11-3757-2011, 2011. 5302, 5304, 5305, 5306, 5307, 5315, 5332, 5335, 5336
- Barahona, D., Rodriguez, J., and Nenes, A.: Sensitivity of the global distribution of cirrus ice crystal concentration to heterogeneous freezing, *J. Geophys. Res.*, 15, D23213, doi:10.1029/2010JD014273, 2010. 5300, 5318, 5328, 5332
- Barkstrom, B.: The earth radiation budget experiment (ERBE), *B. Am. Meteorol. Soc.*, 65, 1170–1185, 1984. 5313
- Blackadar, A. K.: The Vertical Distribution of Wind and Turbulent Exchange in a Neutral Atmosphere, *J. Geophys. Res.*, 67, 3095–3102, 1962. 5305, 5316
- Bodas-Salcedo, A., Webb, M. J., Bony, S., Chepfer, H., Dufresne, J.-L., Klein, S. A., Zhang, Y., Marchand, R., Haynes, J. M., Pincus, R., and John, V. O.: COSP: satellite simulation software for model assessment, *B. Am. Meteorol. Soc.*, 92, 1023–1043, doi:10.1175/2011BAMS2856.1, 2011. 5313, 5323
- Burrows, S. M., Hoose, C., Pöschl, U., and Lawrence, M. G.: Ice nuclei in marine air: biogenic particles or dust?, *Atmos. Chem. Phys.*, 13, 245–267, doi:10.5194/acp-13-245-2013, 2013. 5333
- Chiriacco, M., Chepfer, H., Minnis, P., Haeffelin, M., Platnick, S., Baumgardner, D., Dubuisson, P., McGill, M., Noël, V., Pelon, J., Spangenberg, D., Sun-Mack, S., and Wind, G.: Comparison

## Cloud microphysics in GEOS-5

D. Barahona et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- of CALIPSO-like, LaRC, and MODIS retrievals of ice-cloud properties over SIRTa in France and Florida during CRYSTAL-FACE, *J. Appl. Meteorol. Clim.*, 46, 249–272, 2007. 5324
- Choi, Y., Lindzen, R., Ho, C., and Kim, J.: Space observations of cold-cloud phase change, *P. Natl. Acad. Sci. USA*, 107, 11211–11216, 2010. 5322, 5331, 5368
- 5 Chou, M.-D. and Suarez, M.: An Efficient Thermal Infrared Radiation Parameterization for Use in General Circulation Models, vol. 3 of NASA Tech. Memorandum 104606, NASA Goddard Space Flight Center, Greenbelt, MD, USA, 1994. 5295
- Chou, M.-D., Suarez, M., Ho, C.-H., Yan, M.-H., and Lee, K.-T.: A solar radiation model for use in climate studies, *J. Atmos. Sci.*, 49, 762–772, 1992. 5295
- 10 Colarco, P., da Silva, A., Chin, M., and Diehl, T.: Online simulations of global aerosol distributions in the NASA GEOS-4 model and comparisons to satellite and ground-based aerosol optical depth, *J. Geophys. Res.*, 115, D14207, doi:10.1029/2009JD012820, 2010. 5295, 5313
- Conant, W. C., VanReken, T., Rissman, T., Varutbangkul, V., Jonsson, H., Nenes, A., Jimenez, J., Delia, A., Bahreini, R., Roberts, G., Flagan, R., and Seinfeld, J. H.: Aerosol-cloud drop concentration closure in warm clouds, *J. Geophys. Res.*, 109, D13204, doi:10.1029/2003JD004324, 2004. 5304
- 15 Cziczo, D. J., Froyd, K. D., Hoose, C., Jensen, E. J., Diao, M., Zondlo, M. A., Smith, J. B., Twohy, C. H., and Murphy, D. M.: Clarifying the dominant sources and mechanisms of cirrus cloud formation, *Science*, 340, 1320–1324, 2013. 5320
- Dean, S. M., Flowerdew, J., Lawrence, B. N., and Eckermann, S. D.: Parameterisation of orographic cloud dynamics in a GCM, *Clim. Dynam.*, 28, 581–597, doi:10.1007/s00382-006-0202-0, 2007. 5305, 5335, 5336
- Del Genio, A., Yao, M., Kovari, W., and Lo, K.: A prognostic cloud water parameterization for global climate models, *J. Climate*, 9, 270–304, doi:10.1175/1520-0442(1996)009<0270:APCWPF>2.0.CO;2, 1996. 5292
- 25 Del Genio, A. D., Kovari, W., Yao, M.-S., and Jonas, J.: Cumulus microphysics and climate sensitivity, *J. Climate*, 18, 2376–2387, doi:10.1175/JCLI3413.1, 2005. 5311
- DeMott, P. J., Prenni, A. J., Liu, X., Kreidenweis, S. M., Petters, M. D., Twohy, C. H., Richardson, M. S., Eidhammer, T., and Rogers, D. C.: Predicting global atmospheric ice nuclei distributions and their impacts on climate, *P. Natl. Acad. Sci. USA*, 107, 11217–11222, doi:10.1073/pnas.0910818107, 2010. 5319
- 30

Cloud microphysics  
in GEOS-5

D. Barahona et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Devasthale, A. and Thomas, M. A.: Sensitivity of cloud liquid water content estimates to the temperature-dependent thermodynamic phase: a global study using CloudSat data, *J. Climate*, 25, 7297–7307, 2012. 5321
- Diehl, T., Heil, A., Chin, M., Pan, X., Streets, D., Schultz, M., and Kinne, S.: Anthropogenic, biomass burning, and volcanic emissions of black carbon, organic carbon, and SO<sub>2</sub> from 1980 to 2010 for hindcast model experiments, *Atmos. Chem. Phys. Discuss.*, 12, 24895–24954, doi:10.5194/acpd-12-24895-2012, 2012. 5313
- Eliasson, S., Buehler, S. A., Milz, M., Eriksson, P., and John, V. O.: Assessing observed and modelled spatial distributions of ice water path using satellite data, *Atmos. Chem. Phys.*, 11, 375–391, doi:10.5194/acp-11-375-2011, 2011. 5320, 5321
- Ferrier, S. B.: A double-moment multiple-phase four-class bulk ice scheme. Part I: Description, *J. Atmos. Sci.*, 51, 249–280, doi:10.1175/1520-0469(1994)051<0249:ADMMPF>2.0.CO;2, 1994. 5312
- Fornea, A. P., Brooks, S. D., Dooley, J. B., and Saha, A.: Heterogeneous freezing of ice on atmospheric aerosols containing ash, soot, and soil, *J. Geophys. Res.*, 114, D13201, doi:10.1029/2009JD011958, 2009. 5303
- Fountoukis, C. and Nenes, A.: Continued development of a cloud droplet formation parameterization for global climate models, *J. Geophys. Res.*, 110, D11212, doi:10.1029/2004JD005591, 2005. 5300, 5306
- Fountoukis, C., Nenes, A., Meskhidze, N., Bahreini, R., Conant, W. C., Jonsson, H., Murphy, S., Sorooshian, A., Varutbangkul, V., Brechtel, F., Flagan, R., and Seinfeld, J. H.: Aerosol-cloud drop concentration closure for clouds sampled during the International Consortium for Atmospheric Research on Transport and Transformation 2004 campaign, *J. Geophys. Res.*, 112, D10S30, doi:10.1029/2006JD007272, 2007. 5316
- Frank, W. M. and Cohen, C.: Simulation of tropical convective systems. Part I: A cumulus parameterization, *J. Atmos. Sci.*, 44, 3787–3799, doi:10.1175/1520-0469(1987)044<3787:SOTCSP>2.0.CO;2, 1987. 5309
- Gayet, J., Ovarlez, J., Shcherbakov, V., Ström, J., Schumann, U., Minikin, A., Auriol, F., Petzold, A., and Monier, M.: Cirrus cloud microphysical and optical properties at southern and northern midlatitudes during the INCA experiment, *J. Geophys. Res.*, 109, D20206, doi:10.1029/2004JD004803, 2004. 5317, 5327

**Cloud microphysics  
in GEOS-5**

D. Barahona et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Gettelman, A. and Kinnison, D. E.: The global impact of supersaturation in a coupled chemistry-climate model, *Atmos. Chem. Phys.*, 7, 1629–1643, doi:10.5194/acp-7-1629-2007, 2007. 5315, 5316
- 5 Gettelman, A., Fetzer, E. J., Eldering, A., and Irion, F. W.: The global distribution of supersaturation in the upper troposphere from the Atmospheric Infrared Sounder, *J. Climate*, 19, 6089–6103, 2006. 5315, 5360
- Gettelman, A., Morrison, H., and Ghan, S. J.: A new two-moment bulk stratiform cloud microphysics scheme in the Community Atmosphere Model, version 3 (CAM3). Part II: Single-column and global results, *J. Climate*, 21, 3660–3679, doi:10.1175/2008JCLI2116.1, 10 5323, 5328
- Gettelman, A., Liu, X., Ghan, S. J., Morrison, H., Park, S., Conley, A. J., Klein, S. A., Boyle, J., Mitchell, D. L., and Li, J.-L. F.: Global simulations of ice nucleation and ice supersaturation with an improved cloud scheme in the Community Atmosphere Model, *J. Geophys. Res.*, 115, D18216, doi:10.1029/2009JD013797, 2010. 5292, 5296, 5297
- 15 Gettelman, A., Liu, X., Barahona, D., Lohmann, U., and Chen, C.: Climate impacts of ice nucleation, *J. Geophys. Res.*, 117, D20201, doi:10.1029/2012JD017950, 2012. 5318, 5328
- Gierens, K., Schumann, U., Helten, M., Smit, H., and Marengo, A.: A distribution law for relative humidity in the upper troposphere and lower stratosphere derived from three years of MOZAIC measurements, *Ann. Geophys.*, 17, 1218–1226, doi:10.1007/s00585-999-1218-7, 20 5315, 5360
- Golaz, J.-C., Salzmann, M., Donner, L. J., Horowitz, L. W., Ming, Y., and Zhao, M.: Sensitivity of the aerosol indirect effect to subgrid variability in the cloud parameterization of the GFDL Atmosphere General Circulation Model AM3, *J. Climate*, 24, 3145–3160, doi:10.1175/2010JCLI3945.1, 2010. 5304, 5316
- 25 Gregory, D.: Estimation of entrainment rate in simple models of convective clouds, *Q. J. Roy. Meteor. Soc.*, 127, 53–72, 2001. 5309
- Gultepe, I. and Isaac, G.: The relationship between cloud droplet and aerosol number concentrations for climate models, *Int. J. Climatol.*, 16, 941–946, 1996. 5318, 5363
- Guo, H., Liu, Y., Daum, P., Senum, G., and Tao, W.: Characteristics of vertical velocity in marine stratocumulus: comparison of large eddy simulations with observations, *Environ. Res. Lett.*, 3, 045020, doi:10.1088/1748-9326/3/4/045020, 2008. 5309, 5316
- 30 Haag, W. and Kärcher, B.: The impact of aerosols and gravity waves on cirrus at midlatitudes, *J. Geophys. Res.*, 109, D12202, doi:10.1029/2004JD004579, 2004. 5305

Cloud microphysics  
in GEOS-5

D. Barahona et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Han, Q., Rossow, W. B., Chou, J., and Welch, R. M.: Global variation of column droplet concentration in low-level clouds, *Geophys. Res. Lett.*, 25, 1419–1422, doi:10.1029/98GL01095, 1998. 5317
- Hendricks, J., Kärcher, B., and Lohmann, U.: Effects of ice nuclei on cirrus clouds in a global climate model, *J. Geophys. Res.*, 116, D18206, doi:10.1029/2010JD015302, 2011. 5332
- Herzog, A. and Vial, F.: A study of the dynamics of the equatorial lower stratosphere by use of ultra-long-duration balloons, *J. Geophys. Res.*, 106, 22745–22761, 2001. 5305, 5317
- Heymsfield, A. J., Bansemer, A., Field, P. R., Durden, S. L., Stith, J. L., Dye, J. E., Hall, W., and Grainger, C. A.: Observations and parameterizations of particle size distributions in deep tropical cirrus and stratiform precipitating clouds: results from in situ observations in TRMM field campaigns, *J. Atmos. Sci.*, 59, 3457–3491, doi:10.1175/1520-0469(2002)059<3457:OAIPOPS>2.0.CO;2, 2002. 5296
- Heymsfield, A. J., van Zadelhoff, G.-J., Donovan, D. P., Fabry, F., Hogan, R. J., and Illingworth, A. J.: Refinements to ice particle mass dimensional and terminal velocity relationships for ice clouds. Part II: Evaluation and parameterizations of ensemble ice particle sedimentation velocities, *J. Atmos. Sci.*, 64, 1068–1088, doi:10.1175/JAS3900.1, 2007. 5353
- Hu, Y., Rodier, S., Xu, K.-M., Sun, W., Huang, J., Lin, B., Zhai, P., and Josset, D.: Occurrence, liquid water content, and fraction of supercooled water clouds from combined CALIOP/IIR/MODIS measurements, *J. Geophys. Res.*, 115, D00H34, doi:10.1029/2009JD012384, 2010. 5322, 5323
- Huffman, G., Adler, R., Arkin, P., Chang, A., Ferraro, R., Gruber, A., Janowiak, J., McNab, A., Rudolf, B., and Schneider, U.: The global precipitation climatology project (GPCP) combined precipitation dataset, *B. Am. Meteorol. Soc.*, 78, 5–20, 1997. 5313, 5325, 5372
- IPCC: Climate Change 2007: The Physical Basis. Contribution of Working Group I to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change, Cambridge University Press, Cambridge, UK and New York, NY, USA, 2007. 5292
- Jensen, E. and Pfister, L.: Transport and freeze-drying in the tropical tropopause layer, *J. Geophys. Res.*, 109, D02207, doi:10.1029/2003JD004022, 2004. 5305, 5335
- Jensen, E. J., Pfister, L., Bui, T.-P., Lawson, P., and Baumgardner, D.: Ice nucleation and cloud microphysical properties in tropical tropopause layer cirrus, *Atmos. Chem. Phys.*, 10, 1369–1384, doi:10.5194/acp-10-1369-2010, 2010. 5304, 5305

## Cloud microphysics in GEOS-5

D. Barahona et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Jensen, E. J., Pfister, L., and Bui, T. P.: Physical processes controlling ice concentrations in cold cirrus near the tropical tropopause, *J. Geophys. Res.*, 117, D11205, doi:10.1029/2011JD017319, 2012. 5332

5 Joos, H., Spichtinger, P., Gayet, J., and Minikin, A.: Orographic cirrus in the global climate model ECHAM5, *J. Geophys. Res.*, 113, D18205, doi:10.1029/2007JD009605, 2008. 5304, 5305, 5306, 5335, 5336

Joos, H., Spichtinger, P., and Lohmann, U.: Influence of a future climate on the microphysical and optical properties of orographic cirrus clouds in ECHAM5, *J. Geophys. Res.*, 115, D19129, doi:10.1029/2010JD013824, 2010. 5335

10 Kärcher, B. and Lohmann, U.: A parameterization of cirrus cloud formation: homogeneous freezing including effects on aerosol size, *J. Geophys. Res.*, 107, 4698, doi:10.1029/2001JD001429, 2002. 5337

Kärcher, B. and Ström, J.: The roles of dynamical variability and aerosols in cirrus cloud formation, *Atmos. Chem. Phys.*, 3, 823–838, doi:10.5194/acp-3-823-2003, 2003. 5304

15 Kay, J. E., Hillman, B. R., Klein, S. A., Zhang, Y., Medeiros, B., Pincus, R., Gettelman, A., Eaton, B., Boyle, J., Marchand, R., and Ackerman, A.: Exposing global cloud biases in the Community Atmosphere Model (CAM) using satellite observations and their corresponding instrument simulators, *J. Climate*, 25, 5190–5207, 2012. 5314

Khain, A., Ovtchinnikov, M., Pinsky, M., Pokrovsky, A., and Krugliak, H.: Notes on the state-of-the-art numerical modeling of cloud microphysics, *Atmos. Res.*, 55, 159–224, doi:10.1016/S0169-8095(00)00064-8, 2000. 5292

Khairoutdinov, M. and Kogan, Y.: A new cloud physics parameterization in a Large-Eddy simulation model of marine stratocumulus, *Mon. Weather Rev.*, 128, 229–243, 2000. 5297

25 King, M., Menzel, W., Kaufman, Y., Tanre, D., Gao, B.-C., Platnick, S., Ackerman, S., Remer, L., Pincus, R., and Hubanks, P.: Cloud and aerosol properties, precipitable water, and profiles of temperature and water vapor from MODIS, *IEEE T. Geosci. Remote*, 41, 442–458, doi:10.1109/TGRS.2002.808226, 2003. 5325

Koop, T., Luo, B., Tslas, A., and Peter, T.: Water activity as the determinant for homogeneous ice nucleation in aqueous solutions, *Nature*, 406, 611–614, 2000. 5301, 5302, 5310

30 Korolev, A. and Mazin, I.: Supersaturation of water vapor in clouds, *J. Atmos. Sci.*, 60, 2957–2974, 2003. 5310

Krämer, M., Schiller, C., Afchine, A., Bauer, R., Gensch, I., Mangold, A., Schlicht, S., Spelten, N., Sitnikov, N., Borrmann, S., de Reus, M., and Spichtinger, P.: Ice supersaturations

## Cloud microphysics in GEOS-5

D. Barahona et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



and cirrus cloud crystal numbers, *Atmos. Chem. Phys.*, 9, 3505–3522, doi:10.5194/acp-9-3505-2009, 2009. 5315, 5318, 5324, 5333, 5363, 5371

Kumar, P., Nenes, A., and Sokolik, I. N.: Importance of adsorption for CCN activity and hygroscopic properties of mineral dust aerosol, *Geophys. Res. Lett.*, 36, L24804, doi:10.1029/2009GL040827, 2009a. 5310

Kumar, P., Sokolik, I. N., and Nenes, A.: Parameterization of cloud droplet formation for global and regional models: including adsorption activation from insoluble CCN, *Atmos. Chem. Phys.*, 9, 2517–2532, doi:10.5194/acp-9-2517-2009, 2009b. 5300

Ladino, L., Stetzer, O., Lüönd, F., Welti, A., and Lohmann, U.: Contact freezing experiments of kaolinite particles with cloud droplets, *J. Geophys. Res.*, 116, D22202, doi:10.1029/2011JD015727, 2011. 5303

Lance, S., Nenes, A., and Rissman, T.: Chemical and dynamical effects on cloud droplet number: implication for estimates of the aerosol indirect effect, *J. Geophys. Res.*, 109, D22208, doi:10.1029/2004JD004596, 2004. 5352

Lance, S., Shupe, M. D., Feingold, G., Brock, C. A., Cozic, J., Holloway, J. S., Moore, R. H., Nenes, A., Schwarz, J. P., Spackman, J. R., Froyd, K. D., Murphy, D. M., Brioude, J., Cooper, O. R., Stohl, A., and Burkhardt, J. F.: Cloud condensation nuclei as a modulator of ice processes in Arctic mixed-phase clouds, *Atmos. Chem. Phys.*, 11, 8003–8015, doi:10.5194/acp-11-8003-2011, 2011. 5293

Li, J.-L. F., Waliser, D. E., Chen, W.-T., Guan, B., Kubar, T., Stephens, G., Ma, H.-Y., Deng, M., Donner, L., Seman, C., and Horowitz, L.: An observationally based evaluation of cloud ice water in CMIP3 and CMIP5 GCMs and contemporary reanalyses using contemporary satellite data, *J. Geophys. Res.*, 117, D16105, doi:10.1029/2012JD017640, 2012. 5313, 5320, 5321, 5325, 5365, 5367, 5372

Li, J.-L. F., Lee, S., Waliser, D. E., Lee, S., Guan, B., Stephens, G., and Teixeira, J.: Assessment of Cloud liquid water in CMIP3, CMIP5, and contemporary GCMs and reanalyses with observations, *J. Geophys. Res.*, submitted, 2013. 5313, 5321, 5324, 5366, 5367, 5372

Lindzen, R. S.: Turbulence and stress owing to gravity wave and tidal breakdown, *J. Geophys. Res.*, 86, 9707–9714, doi:10.1029/JC086iC10p09707, 1981. 5305

Liu, X., Penner, J., Das, B., Bergmann, D., Rodriguez, J., Strahan, S., Wang, M., and Feng, Y.: Uncertainties in global aerosol simulations: assessment using three meteorological data sets, *J. Geophys. Res.*, 112, D11212, doi:10.1029/2006JD008216, 2007. 5315

## Cloud microphysics in GEOS-5

D. Barahona et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Liu, X., Shi, X., Zhang, K., Jensen, E. J., Gettelman, A., Barahona, D., Nenes, A., and Lawson, P.: Sensitivity studies of dust ice nuclei effect on cirrus clouds with the Community Atmosphere Model CAM5, *Atmos. Chem. Phys.*, 12, 12061–12079, doi:10.5194/acp-12-12061-2012, 2012. 5328
- 5 Liu, Y., Daum, P., and Yum, S.: Analytical expression for the relative dispersion of the cloud droplet size distribution, *Geophys. Res. Lett.*, 33, L02810, doi:10.1029/2005GL024052, 2006. 5297, 5311
- Liu, Y., Daum, P., Guo, H., and Peng, Y.: Dispersion bias, dispersion effect, and the aerosol-cloud conundrum, *Environ. Res. Lett.*, 3, 045021, doi:10.1088/1748-9326/3/4/045021, 2008. 5297, 5311
- 10 Locatelli, J. D. and Hobbs, P. V.: Fall speeds and masses of solid precipitation particles, *J. Geophys. Res.*, 79, 2185–2197, doi:10.1029/JC079i015p02185, 1974. 5353
- Loeb, N., Wielicki, B., Doelling, D., Smith, G., Keyes, D., Kato, S., Manalo-Smith, N., and Wong, T.: Toward optimal closure of the Earth's top-of-atmosphere radiation budget, *J. Climate*, 22, 748–766, 2009. 5313, 5372
- 15 Lohmann, U.: Global anthropogenic aerosol effects on convective clouds in ECHAM5-HAM, *Atmos. Chem. Phys.*, 8, 2115–2131, doi:10.5194/acp-8-2115-2008, 2008. 5292, 5293, 5334
- Lohmann, U. and Feichter, J.: Global indirect aerosol effects: a review, *Atmos. Chem. Phys.*, 5, 715–737, doi:10.5194/acp-5-715-2005, 2005. 5292, 5293
- 20 Lohmann, U., Spichtinger, P., Jess, S., Peter, T., and Smit, H.: Cirrus cloud formation and ice supersaturated regions in a global climate model, *Environ. Res. Lett.*, 3, 045022, doi:10.1088/1748-9326/3/4/045022, 2008. 5292
- Louis, J. F., Weill, A., and Vidal-Madjar, D.: Dissipation length in stable layers, *Bound.-Lay. Meteorol.*, 25, 229–243, doi:10.1007/BF00119538, 1983. 5305
- 25 Marcolli, C., Gedamke, S., Peter, T., and Zobrist, B.: Efficiency of immersion mode ice nucleation on surrogates of mineral dust, *Atmos. Chem. Phys.*, 7, 5081–5091, doi:10.5194/acp-7-5081-2007, 2007. 5322
- McFarlane, N. A.: The effect of orographically excited gravity wave drag on the general circulation of the lower stratosphere and troposphere, *J. Atmos. Sci.*, 44, 1775–1800, doi:10.1175/1520-0469(1987)044<1775:TEOOEG>2.0.CO;2, 1987. 5306, 5335
- 30 McFarquhar, G. and Heymsfield, A.: Parameterization of tropical cirrus ice crystal size distributions and implications for radiative transfer: results from CEPEX, *J. Atmos. Sci.*, 54, 2187–2200, 1997. 5311, 5324, 5329, 5371

---

**Cloud microphysics  
in GEOS-5**


---

D. Barahona et al.

---

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[◀](#)
[▶](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)


- Molod, A.: Constraints on the total water PDF in GCMs from AIRS and a high resolution model, *J. Climate*, 25, 8341–8352, 2012. 5294, 5295
- Molod, A., Takacs, L., Suarez, M., Bacmeister, J., Song, I., and Eichmann, A.: The GEOS-5 Atmospheric General Circulation Model: Mean Climate and Development from MERRA to Fortuna, vol. 28 of Technical Report Series on Global Modeling and Data Assimilation, NASA Goddard Space Flight Center, Greenbelt, MD, USA, 2012. 5295, 5307
- 5 Moorthi, S. and Suarez, M. J.: Relaxed Arakawa–Schubert – a parameterization of moist convection for general circulation models, *Mon. Weather Rev.*, 120, 978–1002, 1992. 5294, 5307, 5308
- 10 Morales, R. and Nenes, A.: Characteristic updrafts for computing distribution-averaged cloud droplet number, and stratocumulus cloud properties, *J. Geophys. Res.*, 115, D18220, doi:10.1029/2009JD013233, 2010. 5306
- Morales Betancourt, R., Lee, D., Oreopoulos, L., Sud, Y. C., Barahona, D., and Nenes, A.: Sensitivity of cirrus and mixed-phase clouds to the ice nuclei spectra in McRAS-AC: single column model simulations, *Atmos. Chem. Phys.*, 12, 10679–10692, doi:10.5194/acp-12-10679-2012, 2012. 5328
- 15 Morrison, H. and Gettelman, A.: A new two-moment bulk stratiform cloud microphysics scheme in the Community Atmosphere Model, version 3 (CAM3). Part I: Description and numerical tests, *J. Climate*, 21, 3642–3659, doi:10.1175/2008JCLI2105.1, 2008. 5296, 5297, 5304, 5305, 5329
- 20 Morrison, H. and Grabowski, W. W.: A novel approach for representing ice microphysics in models: Description and tests using a kinematic framework, *J. Atmos. Sci.*, 65, 1528–1548, 2008. 5296
- Morrison, H., Curry, J. A., and Khvorostyanov, V. I.: A new double-moment microphysics parameterization for application in cloud and climate models. Part I: Description, *J. Atmos. Sci.*, 62, 1665–1677, doi:10.1175/JAS3446.1, 2005. 5304
- 25 Morrison, H., de Boer, G., Feingold, G., Harrington, J., Shupe, M. D., and Sulia, K.: Resilience of persistent Arctic mixed-phase clouds, *Nat. Geosci.*, 5, 11–17, doi:10.1038/ngeo1332, 2012. 5293
- 30 Murray, B., OSullivan, D., Atkinson, J., and Webb, M.: Ice nucleation by particles immersed in supercooled cloud droplets, *Chem. Soc. Rev.*, 41, 6519–6554, 2012. 5303

## Cloud microphysics in GEOS-5

D. Barahona et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Murray, B. J., Broadley, S. L., Wilson, T. W., Atkinson, J. D., and Wills, R. H.: Heterogeneous freezing of water droplets containing kaolinite particles, *Atmos. Chem. Phys.*, 11, 4191–4207, doi:10.5194/acp-11-4191-2011, 2011. 5301, 5322

Nenes, A. and Seinfeld, J. H.: Parameterization of cloud droplet formation in global climate models, *J. Geophys. Res.*, 108, 4415, doi:10.1029/2002JD002911, 2003. 5300

Niemand, M., Möhler, O., Vogel, B., Vogel, H., Hoose, C., Connolly, P., Klein, H., Bingemer, H., DeMott, P., Skrotzki, J., and Leisner, T.: A particle-surface-area-based parameterization of immersion freezing on desert dust particles, *J. Atmos. Sci.*, 69, 3077–3092, doi:10.1175/JAS-D-11-0249.1, 2012. 5303

Peng, Y., Lohmann, U., and Leaitch, W.: Importance of vertical velocity variations in the cloud droplet nucleation process of marine stratocumulus, *J. Geophys. Res.*, 110, D21213, doi:10.1029/2004JD004922, 2005. 5304, 5306, 5316

Petters, M. D. and Kreidenweis, S. M.: A single parameter representation of hygroscopic growth and cloud condensation nucleus activity, *Atmos. Chem. Phys.*, 7, 1961–1971, doi:10.5194/acp-7-1961-2007, 2007. 5300

Phillips, V., DeMott, P., and Andronache, C.: An empirical parameterization of heterogeneous ice nucleation for multiple chemical species of aerosol, *J. Atmos. Sci.*, 65, 2757–2783, doi:10.1175/2007JAS2546.1, 2008. 5301, 5328

Phillips, V. T., Demott, P. J., Andronache, C., Pratt, K. A., Prather, K. A., Subramanian, R., and Twohy, C.: Improvements to an empirical parameterization of heterogeneous ice nucleation and its comparison with observations, *J. Atmos. Sci.*, 70, 378–409, doi:10.1175/JAS-D-12-080.1, 2013. 5301, 5328

Pincus, R., Platnick, S., Ackerman, S., Hemler, R., and Patrick Hofmann, R.: Reconciling simulated and observed views of clouds: MODIS, ISCCP, and the limits of instrument simulators, *J. Climate*, 25, 4699–4720, 2012. 5325

Platnick, S., King, M., Ackerman, S., Menzel, W., Baum, B., Riédi, J., and Frey, R.: The MODIS cloud products: algorithms and examples from Terra, *IEEE T. Geosci. Remote*, 41, 459–473, 2003. 5313, 5317, 5321

Popovitcheva, O., Kireeva, E., Persiantseva, N., Khokhlova, T., Shonija, N., Tishkova, V., and Demirdjian, B.: Effect of soot on immersion freezing of water and possible atmospheric implications, *Atmos. Res.*, 90, 326–337, 2008. 5301

Pruppacher, H. and Klett, J.: *Microphysics of Clouds and Precipitation*, 2nd Edn., Kluwer Academic Publishers, Boston, MA, 1997. 5293, 5310

## Cloud microphysics in GEOS-5

D. Barahona et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Putman, W. and Suarez, M.: Cloud-system resolving simulations with the NASA Goddard Earth Observing System global atmospheric model (GEOS-5), *Geophys. Res. Lett.*, 38, L16809, doi:10.1029/2011GL048438, 2011. 5293, 5294

Raatikainen, T., Nenes, A., Seinfeld, J. H., Morales, R., Moore, R. H., Latham, T. L., Lance, S., Padro, L. T., Lin, J. J., Cerully, K. M., Bougiatioti, A., Cozic, J., Ruehl, C. R., Chuang, P. Y., Anderson, B. E., Flagan, R. C., Jonsson, H., Mihalopoulos, N., and Smith, J. N.: Worldwide data sets constrain the water vapor uptake coefficient in cloud formation, *P. Natl. Acad. Sci. USA*, 110, 3760–3764, doi:10.1073/pnas.1219591110, 2013. 5300

Ramanathan, V., Crutzen, P., Kiehl, J., and Rosenfeld, D.: Aerosols, climate, and the hydrological cycle, *Science*, 294, 2119–2124, 2001. 5293

Reale, O., Lau, W. K., Kim, K.-M., and Brin, E.: Atlantic tropical cyclogenetic processes during SOP-3 NAMMA in the GEOS-5 global data assimilation and forecast system, *J. Atmos. Sci.*, 66, 3563–3578, doi:10.1175/2009JAS3123.1, 2009. 5294

Reynolds, R. W., Rayner, N. A., Smith, T. M., Stokes, D. C., and Wang, W.: An improved in situ and satellite SST analysis for climate, *J. Climate*, 15, 1609–1625, doi:10.1175/1520-0442(2002)015<1609:AIISAS>2.0.CO;2, 2002. 5313

Rienecker, M., Suarez, M., Todling, R., Bacmeister, J., Takacs, L., Liu, H.-C., Gu, W., Sienkiewicz, M., Koster, R., Gelaro, R., Stajner, I., and Nielsen, J.: The GEOS-5 Data Assimilation System – Documentation of Versions 5.0.1, 5.1.0, and 5.2.0., vol. 27 of Technical Report Series on Global Modeling and Data Assimilation, NASA Goddard Space Flight Center, Greenbelt, MD, USA, 2008. 5293, 5294, 5295, 5298, 5307, 5308

Rienecker, M. M., Suarez, M. J., Gelaro, R., Todling, R., Bacmeister, J., Liu, E., Bosilovich, M. G., Schubert, S. D., Takacs, L., Kim, G.-K., Bloom, S., Chen, J., Collins, D., Conaty, A., da Silva, A., Gu, W., Joiner, J., Koster, R. D., Lucchesi, R., Molod, A., Owens, T., Pawson, S., Pegion, P., Redder, C. R., Reichle, R., Robertson, F. R., Ruddick, A. G., Sienkiewicz, M., and Woollen, J.: MERRA: NASA’s modern-era retrospective analysis for research and applications, *J. Climate*, 24, 3624–3648, doi:10.1175/JCLI-D-11-00015.1, 2011. 5294, 5313

Rosenfeld, D. and Woodley, W. L.: Deep convective clouds with sustained supercooled liquid water down to  $-37.5^{\circ}\text{C}$ , *Nature*, 405, 440–442, doi:10.1038/35013030, 2000. 5293

Rosenfeld, D., Lohmann, U., Raga, G. B., O’Dowd, C. D., Kulmala, M., Fuzzi, S., Reissell, A., and Andreae, M. O.: Flood or drought: how do aerosols affect precipitation?, *Science*, 321, 1309–1313, doi:10.1126/science.1160606, 2008. 5293

## Cloud microphysics in GEOS-5

D. Barahona et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Rossow, W. and Schiffer, R.: Advances in understanding clouds from ISCCP, *B. Am. Meteorol. Soc.*, 80, 2266–2288, 1999. 5313, 5325, 5358
- Salzmann, M., Ming, Y., Golaz, J.-C., Ginoux, P. A., Morrison, H., Gettelman, A., Krämer, M., and Donner, L. J.: Two-moment bulk stratiform cloud microphysics in the GFDL AM3 GCM: description, evaluation, and sensitivity tests, *Atmos. Chem. Phys.*, 10, 8037–8064, doi:10.5194/acp-10-8037-2010, 2010. 5315, 5318, 5323, 5328, 5332
- Sato, K.: Vertical wind disturbances in the troposphere and lower stratosphere observed by the MU radar, *J. Atmos. Sci.*, 47, 2803–2817, 1990. 5305
- Seinfeld, J. H.: Clouds, contrails and climate, *Nature*, 391, 837–838, 1998. 5311
- Slingo, J.: The development and verification of a cloud prediction scheme for the ECMWF model, *Q. J. Roy. Meteor. Soc.*, 113, 899–927, 1987. 5295, 5298
- Smith, R. N. B.: A scheme for predicting layer clouds and their water content in a general circulation model, *Q. J. Roy. Meteor. Soc.*, 116, 435–460, 1990. 5295
- Song, X. and Zhang, G. J.: Microphysics parameterization for convective clouds in a global climate model: description and single-column model tests, *J. Geophys. Res.*, 116, D02201, doi:10.1029/2010JD014833, 2011. 5293, 5334
- Spencer, R., Hood, R., and Goodman, H.: Precipitation retrieval over land and ocean with the SSM/I-Identification and characteristics of the scattering signal, *J. Atmos. Ocean. Tech.*, 6, 254–273, 1989. 5321
- Spichtinger, P. and Cziczo, D. J.: Impact of heterogeneous ice nuclei on homogeneous freezing events in cirrus clouds, *J. Geophys. Res.*, 115, D14208, doi:10.1029/2009JD012168, 2010. 5332
- Sud, Y. C. and Walker, G. K.: Microphysics of clouds with the relaxed Arakawa–Schubert scheme (McRAS). Part I: Design and evaluation with GATE Phase III data, *J. Atmos. Sci.*, 56, 3196–3220, doi:10.1175/1520-0469(1999)056<3196:MOCWTR>2.0.CO;2, 1999. 5309, 5334
- Sud, Y. C., Lee, D., Oreopoulos, L., Barahona, D., Nenes, A., and Suarez, M. J.: Performance of McRAS-AC in the GEOS-5 AGCM: aerosol–cloud–microphysics, precipitation, cloud radiative effects, and circulation, *Geosci. Model Dev.*, 6, 57–79, doi:10.5194/gmd-6-57-2013, 2013. 5292, 5293, 5323
- Tiedtke, M.: Representation of clouds in large-scale models, *Mon. Weather Rev.*, 121, 3040, doi:10.1175/1520-0493(1993)121<3040:ROCILS>2.0.CO;2, 1993. 5295, 5298

Cloud microphysics  
in GEOS-5

D. Barahona et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Tompkins, A.: A prognostic parameterization for the subgrid-scale variability of water vapor and clouds in large-scale models and its use to diagnose cloud cover, *J. Atmos. Sci.*, 59, 1917–1942, 2002. 5299

5 Tonttila, J., O'Connor, E. J., Niemelä, S., Räisänen, P., and Järvinen, H.: Cloud base vertical velocity statistics: a comparison between an atmospheric mesoscale model and remote sensing observations, *Atmos. Chem. Phys.*, 11, 9207–9218, doi:10.5194/acp-11-9207-2011, 2011. 5316

Twomey, S.: The influence of pollution on the shortwave cloud albedo of clouds, *J. Atmos. Sci.*, 34, 1149–1152, 1977. 5293

10 Twomey, S.: Aerosols, clouds and radiation, *Atmos. Environ.*, 25A, 2435–2442, 1991. 5293

Wang, M. and Penner, J. E.: Cirrus clouds in a global climate model with a statistical cirrus cloud scheme, *Atmos. Chem. Phys.*, 10, 5449–5474, doi:10.5194/acp-10-5449-2010, 2010. 5292

15 Wiacek, A., Peter, T., and Lohmann, U.: The potential influence of Asian and African mineral dust on ice, mixed-phase and liquid water clouds, *Atmos. Chem. Phys.*, 10, 8649–8667, doi:10.5194/acp-10-8649-2010, 2010. 5310

Wise, M. E., Baustian, K. J., Koop, T., Freedman, M. A., Jensen, E. J., and Tolbert, M. A.: Depositional ice nucleation onto crystalline hydrated NaCl particles: a new mechanism for ice formation in the troposphere, *Atmos. Chem. Phys.*, 12, 1121–1134, doi:10.5194/acp-12-1121-2012, 2012. 5333

20 Xie, P. and Arkin, P.: Global precipitation: a 17-year monthly analysis based on gauge observations, satellite estimates, and numerical model outputs, *B. Am. Meteorol. Soc.*, 78, 2539–2558, 1997. 5313, 5325, 5372

25 Young, K. C.: The role of contact nucleation in ice phase initiation in clouds, *J. Atmos. Sci.*, 31, 768–776, doi:10.1175/1520-0469(1974)031<0768:TROCNI>2.0.CO;2, 1974. 5303

Cloud microphysics  
in GEOS-5

D. Barahona et al.

**Table 1.** Lognormal size distribution parameters used in this study (Lance et al., 2004).  $\bar{D}_g$  ( $\mu\text{m}$ ) and  $\sigma_g$  are the geometric mean diameter and dispersion, respectively.  $N_i/N_a$  is the particle number fraction in mode  $i$ . The “polluted” size distribution parameters for sulfate and organics are used when the total aerosol mass exceeds  $5.0 \mu\text{g m}^{-3}$ .

Aerosol species	$\bar{D}_g$	$\sigma_g$	$N_i/N_a$
Dust1	1.46	2.0	1.0
Dust2	2.8	2.0	1.0
Dust3	4.8	2.0	1.0
Dust4	9.0	2.0	1.0
Dust5	16.0	2.0	1.0
Black carbon	0.024	2.20	1.0
Sea salt	[0.02, 0.092, 0.58]	[1.47, 2.0, 2.0]	[0.56, 0.43, $7.6 \times 10^{-3}$ ]
Sulfate and organics			
– Clean	[0.016, 0.067, 0.93]	[1.6, 2.1, 2.2]	[0.55, 0.44, $4.1 \times 10^{-2}$ ]
– Polluted	[0.014, 0.054, 0.86]	[1.8, 2.16, 2.21]	[0.77, 0.23, $3.6 \times 10^{-3}$ ]

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

I◀

▶I

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## Cloud microphysics in GEOS-5

D. Barahona et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**Table 2.** Parameters of the terminal velocity relation  $w_{\text{term}} = aD_v^b(1000/p)^{0.4}$  (SI units) for convective ice species.

Species	a	b	Reference
Ice	$2 \exp[4 \times 10^{-4}(T - 273.0)]$	$0.244 - 4.9 \times 10^{-3}(T - 273.0)$	Heymsfield et al. (2007)
Graupel	19.3	0.37	Locatelli and Hobbs (1974)

## Cloud microphysics in GEOS-5

D. Barahona et al.

**Table 3.** Description of sensitivity runs performed with GEOS-5 using the new microphysics.

Run	Description
NOCNV	Single moment microphysics within convective clouds
NOBC	Black carbon not acting as IN
LC400	$L_c = 400$ m
PDA08	Usage of the Phillips (2008) heterogeneous ice nucleation spectrum
MUIZERO	Prescribed constant $\mu_i = 0.0$
ARGACT	Usage of the Abdul-Razzak and Ghan (2000) activation parameterization
NOGLASS	Glassy organics not considered as IN
NOPREEX	Preexisting ice crystals not considered
DCS200	$D_{cs} = 200$ $\mu\text{m}$

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Cloud microphysics  
in GEOS-5

D. Barahona et al.

**Table 4.** Annual mean model results and observations. The experimental data sets are described in Sect. 3. CTL and NEW refer to runs with the operational version of GEOS-5 and with the implementation of the new microphysics, respectively. Sensitivity studies are described in Table 3 and Sect. 4.

Simulation	CTL	NEW	ARGACT	NOBC	NOGLASS	PDA08	NOPREEX	LC400	NOCNV	MUIZERO	DCS200	OBS
$P_{\text{tot}}$ (mm d <sup>-1</sup> )	2.85	2.72	2.72	2.71	2.72	2.73	2.66	2.77	2.90	2.70	2.83	2.68 (GPCP) 2.60 (CMAP)
LWP (gm <sup>-2</sup> )	60.0	37.3	38.0	37.6	37.5	37.1	37.3	37.2	36.1	36.5	35.3	23.0 (CloudSat), 109.8 (MODIS), 88.4 (SSM/I, ocean)
IWP (gm <sup>-2</sup> )	19.0	27.1	27.3	27.0	26.9	27.3	32.8	21.3	25.2	10.3	16.1	25.8 (CloudSat)
TWP (gm <sup>-2</sup> )	79.0	64.4	65.3	64.6	64.4	64.4	70.1	58.6	61.3	46.8	51.4	48.8 (CloudSat)
$f_c$ (%)	46.0	56.0	56.8	56.3	55.8	55.2	58.3	54.2	51.0	56.8	50.0	52 (MODIS) 62 (ISCCP)
$N_{d, \text{cum}}$ (cm <sup>-2</sup> )		1.68	1.85	1.67	1.68	1.70	1.66	1.55	2.29	1.65	2.33	1.96 (MODIS)
$N_c$ (L <sup>-1</sup> )		66	65	64	67	55	135	38	74	60	62	
$N_c$ (L <sup>-1</sup> ) (cirrus)		166	163	160	168	139	359	91	183	154	158	
$R_{\text{eff, liq}}$ (μm)	10.2	14.2	13.5	14.3	14.3	14.3	14.6	14.6	13.2	13.7	13.0	14.8 (MODIS)
$R_{\text{eff, ice}}$ (μm)	20.8	26.2	26.0	26.0	26.2	27.2	23.2	29.3	25.5	12.5	23.6	24.2 (MODIS)
-SWCF (W m <sup>-2</sup> )	52.1	49.5	52.0	50.3	49.7	49.5	53.2	46.7	45.0	49.7	44.6	47.2 (CERES) 51.8 (ERBE)
LWCF (W m <sup>-2</sup> )	25.2	26.6	27.3	27.2	26.2	25.8	31.2	23.2	22.2	26.9	20.8	26.2 (CERES) 30.67 (ERBE)
OLR (W m <sup>-2</sup> )	238.9	238.3	237.3	237.5	238.2	238.9	233.3	241.4	243.0	237.0	244.5	239.8 (CERES) 240.2 (ERBE)
OSR (W m <sup>-2</sup> )	236.5	239.3	236.7	238.4	238.9	239.2	235.6	242.0	243.8	239.2	244.2	240.6 (CERES) 255.7 (ERBE)
Net TOA (W m <sup>-2</sup> )	-2.4	0.95	-0.52	0.90	0.77	0.32	2.24	0.58	0.75	2.08	-0.28	0.75 (CERES)

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**Table A1.** List of symbols and acronyms.

$\gamma$	Virtual mass coefficient
$\gamma_c$	Cooling rate
$\eta$	Cloud tracer
$\Delta H_s$	Enthalpy of sublimation of ice
$\Delta q_c$	Change in total condensate due to the cloud microphysics
$\Delta t$	Model time step
$\Delta t_i$	Average time of a convective parcel within a model layer
$\phi(\bar{w}, \sigma_w^2)$	Subgrid distribution of vertical velocity
$\kappa$	Hygroscopicity parameter
$\lambda$	Entrainment rate
$\lambda_m$	Value of $f_m$ in the free troposphere
$\lambda_{0,y}$	Slope parameter of $n_y(D)$
$\mu_y$	Dispersion of $n_y(D)$
$\rho_i$	Ice density
$\sigma_{g,x}$	Geometric size dispersion of the $x$ species
$\sigma_{w,turb}^2$	Variance in $w_{sub}$ due to turbulence
$\sigma_{w,gw}^2$	Variance in $w_{sub}$ due to gravity wave dynamics
$\sigma_w$	Standard deviation of $w_{sub}$
$\tau_0$	Surface stress
$A_i$	$\left[ \frac{\rho_i \Delta H_s}{k_c R_c T_c^2} + \frac{\rho_i R_c T_c}{\rho_{w,c} D_c} \right]^{-1}$
CCN	Cloud condensation nuclei
$c_p$	Specific heat capacity of air
$D$	Convective detrainment rate
$D_{cs}$	Critical size for ice-snow autoconversion
$D_{c,y}$	Critical size for precipitation of the $y$ cloud species
$D_{g,x}$	Geometric mean diameter of the $x$ species
$D_w$	Water vapor diffusivity in air
$f_c$	Total cloud fraction
$f_c^*$	Cloud fraction modified by the cloud microphysics
$f_{gr}$	Fraction of ice existing as graupel
$f_{het}$	Fraction of ice crystals produced by heterogeneous ice nucleation
$f_{ice}$	Mass fraction of ice in the total condensate
$f_{cn}$	Detrained anvil cloud fraction
$g$	Acceleration of gravity
IN	Ice nuclei
IWC	Ice water content
IWP	Ice water path
$k_a$	Thermal conductivity of air
$K_T$	Mixing coefficient for heat
$L_c$	Characteristic wave length in cirrus
$l_m$	Mixing length
LWC	Liquid water content
LWCF	Longwave cloud forcing
LWP	Liquid water path
$M_w, M_a$	Molar masses of water and air, respectively
$N$	Brunt–Väisälä frequency
$N_{c,cv}$	Ice crystal concentration within convective cumulus
$N_{c,nuc}$	Ice crystal concentration nucleated in cirrus
$N_{d,act}$	Activated cloud droplet number concentration
$N_{d,cum}$	Column integrated droplet number concentration
$n_{d_i}, N_{d_i}$	Grid mean and in-cloud droplet number concentration, respectively
$n_{d_i}, N_{d_i}$	Grid mean and in-cloud ice crystal number concentration, respectively

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Table A1. Continued.

$N_{\text{dep}}$	Ice crystal concentration produced by deposition and condensation nucleation
$n_{\text{gr}}$	Graupel number concentration
$N_{\text{het}}$	Ice nucleation spectrum
$N_{\text{imm}}$	Ice crystal concentration produced by immersion freezing
$N_{0,y}$	Intercept parameter of $n_y(D)$
$n_{s,x}$	Immersion active site surface density for the $x$ species
$N_x$	Aerosol number concentration of the $x$ species
$n_y(D)$	Size distribution of the $y$ species
$p$	Pressure
$P_q(q_i)$	Probability distribution of total cloud condensate
$p_{s,w}, p_{s,i}$	Liquid water and ice saturation vapor pressure, respectively
$P_{\text{tot}}$	Total precipitation
$q^*$	Weighted saturation mixing ratio between liquid and ice
$q_c$	Total condensate mixing ratio
$q_{c,\text{det}}$	Detained condensate mixing ratio
$q_{\text{cn}}$	Mixing ratio of total condensate in a convective parcel
$q_{\text{gr}}$	Graupel mass mixing ratio within a convective cumulus
$q_l$	Ice water mixing ratio
$q_l$	Liquid water mixing ratio
$q_{\text{max}}, q_{\text{min}}$	Upper and lower limits of the total water distribution, respectively
$q_l^*, q_i^*$	Saturation specific humidities for liquid and ice, respectively
$q_t$	Total water mixing ratio, ( $q_l + q_i$ )
$q_v$	Water vapor mixing ratio
$R$	Universal gas constant
$R_{\text{eff,liq}}$	Cloud droplet effective radius
$R_{\text{eff,ice}}$	Ice crystal effective radius
RH	Ambient relative humidity
$R_v$	$R/M_a$
$R_{\text{vol,ice}}$	Volumetric ice crystal radius, $\left(\frac{3q}{4\pi N_c \rho_i}\right)^{1/3}$
$S_{i,c}$	Clear sky saturation ratio
SCF	Supercooled cloud fraction
$S_{\text{crit}}$	Critical saturation ratio
$S_i$	Saturation ratio with respect to ice
$S_{i,\text{max}}$	Maximum water vapor supersaturation with respect to ice
$S_{i,\text{max}}$	Maximum water vapor supersaturation with respect to water
$\bar{s}_{p,x}$	Mean particle surface area of the $x$ species
$S_w^{\text{sat}}$	Value of $S_i$ at water saturation
SWCF	Shortwave cloud forcing
$t$	Time
$T$	Temperature
$T_c$ and $T_v$	Virtual temperature of the cloud and the environment, respectively
TWP	Total water path
$U$	Horizontal wind
$\bar{w}$	Mean vertical velocity
$w_s$	Grid-scale vertical velocity
$w_{\text{sub}}$	Subgrid scale vertical velocity
$w_{\text{term}}$	Hydrometeor terminal velocity
$w_{\text{up}}$	Cumulus vertical velocity
$W$	Convective mass flux
$z$	Altitude

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## Cloud microphysics in GEOS-5

D. Barahona et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶


Back

Close

Full Screen / Esc

Printer-friendly Version


Interactive Discussion



**Fig. 1.** Annual mean differences in low (CLDLO), middle (CLDMD) and high (CLDHI) level cloud fraction between GEOS-5 and ISCCP (Rossow and Schiffer, 1999) for the CTL and NEW runs using the COSP simulator.

Cloud microphysics  
in GEOS-5


D. Barahona et al.



**Fig. 2.** Annual global frequency distribution of (left) and zonal mean (right) of the critical saturation ratio,  $S_{\text{crit}}$  (%), for the cirrus regime ( $T < 235$  K). Solid lines (right) represent the annual mean tropopause pressure.

Cloud microphysics  
in GEOS-5

D. Barahona et al.



**Fig. 3.** Global frequency distribution of clear sky saturation ratio with respect to ice from instantaneous GEOS-5 output using the new microphysics (left panel, black dots). Blue and red shades correspond to the frequency distributions from AIRS satellite retrievals (Gettelman et al., 2006) and the MOZIC data set (Gierens et al., 1999), respectively. Uncertainty in the observations was calculated as one standard deviation around the mean value within  $2^\circ \times 2^\circ$  grid cell and introducing a 10 % perturbation in  $S_i$  along the x axis. The center and right panels show the zonal mean frequency (%) of supersaturation from GEOS-5 and AIRS, respectively.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Cloud microphysics  
in GEOS-5

D. Barahona et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion




Fig. 4. Annual mean  $\sigma_w$  ( $\text{m s}^{-1}$ ) from GEOS-5.

Cloud microphysics  
in GEOS-5

D. Barahona et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶


Back

Close


Full Screen / Esc

Printer-friendly Version

Interactive Discussion




**Fig. 5.** Annual vertically integrated droplet number concentration ( $10^6 \text{ cm}^{-2}$ ) from GEOS-5 (NEW) and the MODIS retrieval calculated using Eq. (41). Also shown are results obtained using the Abdul-Razzak and Ghan (2000) CCN activation parameterization (ARGACT).



**Fig. 6.** Global frequency of in-cloud ice crystal number concentration as a function of temperature from instantaneous GEOS-5 output. **(a)** Ice crystal concentration,  $N_c$ . Solid lines represent the 25% and 75% quantiles from the field campaign data analysis of Krämer et al. (2009). Solid-dotted lines represent the typical range of mean  $N_c$  found in mixed-phase clouds (Gultepe and Isaac, 1996). **(b)** Ice crystal concentration from deposition/condensation ice nucleation,  $N_{dep}$ . **(c)** Ice crystal concentration from immersion ice nucleation,  $N_{imm}$ . **(d)** Ice crystal concentration from convective cumulus detrainment,  $N_{cnv}$ .

Cloud microphysics  
in GEOS-5

D. Barahona et al.




**Fig. 7.** Annual mean ice crystal concentration nucleated in cirrus ( $T < 235$  K) weighted by cloud fraction (left panel). Also shown are the weighted average (center panel) and zonal mean (right panel) fraction of ice crystal production by homogeneous freezing in cirrus.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

## Cloud microphysics in GEOS-5

D. Barahona et al.



**Fig. 8.** Zonal mean non-convective ice water mass mixing ratio ( $\text{mg kg}^{-1}$ ) (upper panels) and total ice condensate (ice and snow, bottom panels) for non-convective clouds from the CTL and NEW runs and the CloudSat retrieval (Li et al., 2012).

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc


Printer-friendly Version

Interactive Discussion



## Cloud microphysics in GEOS-5

D. Barahona et al.



**Fig. 9.** Zonal mean non-convective liquid water mass mixing ratio ( $\text{mg kg}^{-1}$ ) (upper panels) and total liquid condensate (water and rain, bottom panels) for non-convective clouds from the CTL and NEW runs and the CloudSat retrieval (Li et al., 2013).

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc


Printer-friendly Version

Interactive Discussion



Cloud microphysics  
in GEOS-5

D. Barahona et al.



**Fig. 10.** Liquid (LWP), ice (IWP), and total (TWP) water path (gm<sup>-2</sup>) for non-convective, non-precipitating clouds from GEOS-5 output using the new microphysics and from the CloudSat retrieval (Li et al., 2012, 2013).

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc


Printer-friendly Version

Interactive Discussion



Cloud microphysics  
in GEOS-5

D. Barahona et al.



**Fig. 11.** Global frequency of supercooled cloud fraction (SCF) from GEOS-5 for the CTL and NEW runs. The most frequent SCF value for each temperature is marked (\*). The solid lines represent the range of SCF (mean plus and less one standard deviation) from CALIOP data (Choi et al., 2010).

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc


Printer-friendly Version

Interactive Discussion



Cloud microphysics  
in GEOS-5

D. Barahona et al.



**Fig. 12.** Liquid cloud effective radius ( $\mu\text{m}$ ) from GEOS-5 using COSP and from the MODIS retrieval.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[I ◀](#)[▶ I](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Cloud microphysics  
in GEOS-5

D. Barahona et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶


Back

Close


Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**Fig. 13.** Ice cloud effective radius ( $\mu\text{m}$ ) from GEOS-5 using COSP and from the MODIS retrieval.



**Fig. 14.** Global frequency of ice volumetric radius as a function of temperature from GEOS-5. Solid lines represent the 25% and 75% quantiles from the field campaign analysis of Krämer et al. (2009). Filled circles were calculated using the correlation obtained by McFarquhar and Heymsfield (1997) from field measurements in mixed-phase and cirrus clouds.

**Cloud microphysics  
in GEOS-5**

D. Barahona et al.


Title Page

Abstract	Introduction
Conclusions	References
Tables	Figures
◀	▶
◀	▶
Back	Close
Full Screen / Esc	
Printer-friendly Version	
Interactive Discussion	



Cloud microphysics  
in GEOS-5

D. Barahona et al.



**Fig. 15.** Annual zonal means from the GEOS-5 model for the CTL (blue lines) and the NEW (red lines) runs compared against different observations (black lines). **(a, b)** Liquid ( $R_{\text{eff, liq}}$ ) and ice ( $R_{\text{eff, ice}}$ ) effective radius from COSP output against MODIS. **(c, d)** Shortwave (SWCF) and longwave (LWCF) cloud forcing against CERES-EBAF retrievals (Loeb et al., 2009). **(e)** Liquid water path against CloudSat (black lines) and MODIS (black circles) retrievals. **(f)** Non-convective, non-precipitable ice water path against CloudSat retrievals (Li et al., 2012, 2013). Also shown is the total (ice and snow) non-convective ice water path (red circles) from GEOS-5 using the new microphysics. **(g)** Total cloud fraction from COSP output against MODIS (black lines) and ISCCP (black circles). **(h)** Total precipitation against GPCP data (Huffman et al., 1997). Also shown are data from the CMAP dataset (Xie and Arkin, 1997) (black circles). **(i, j)** Liquid and ice optical depth (COD) from COSP output against MODIS retrievals.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc


Printer-friendly Version

Interactive Discussion



Cloud microphysics  
in GEOS-5

D. Barahona et al.



**Fig. 16.** Annual mean ice crystal concentration as a function of temperature for the different runs of Table 4.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)