



This discussion paper is/has been under review for the journal Geoscientific Model Development (GMD). Please refer to the corresponding final paper in GMD if available.

# A system of conservative regridding for ice/atmosphere coupling in a GCM

E. Fischer<sup>1,2</sup>, S. Nowicki<sup>3</sup>, M. Kelley<sup>2,4</sup>, and G. A. Schmidt<sup>2</sup>

<sup>1</sup>Center for Climate Systems Research, Columbia University, New York, NY, USA

<sup>2</sup>NASA Goddard Institute of Space Studies, New York, NY, USA

<sup>3</sup>NASA Goddard Space Flight Center, Greenbelt, MD, USA

<sup>4</sup>Trinnovim LLC, 2880 Broadway, New York, New York 10025, USA

Received: 12 November 2013 – Accepted: 26 November 2013 – Published: 6 December 2013

Correspondence to: E. Fischer (elizabeth.fischer@columbia.edu)

Published by Copernicus Publications on behalf of the European Geosciences Union.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

## Abstract

The method of elevation classes has proven to be a useful way for a low-resolution general circulation model (GCM) to produce high-resolution downscaled surface mass balance fields, for use in one-way studies coupling GCMs and ice flow models. Past uses of elevation classes have been a cause of non-conservation of mass and energy, caused by inconsistency in regridding schemes chosen to regrid to the atmosphere vs. downscaling to the ice model. This causes problems for two-way coupling.

A strategy that resolves this conservation issue has been designed and is presented here. The approach identifies three grids between which data must be regridded, and five transformations between those grids required by a typical coupled GCM–ice flow model. This paper shows how each of those transformations may be achieved in a consistent, conservative manner. These transformations are implemented in GLINT2, a library used to couple GCMs with ice models. Source code and documentation are available for download. Confounding real-world issues are discussed, including the use of projections for ice modeling, how to handle dynamically changing ice geometry, and modifications required for finite element ice models.

## 1 Introduction

Many questions still surround how ice sheets respond to climate forcing and how those changes will affect sea level, regional and global climate. Recent observations have shown accelerating mass loss from the Greenland and Antarctic ice sheets, adding urgency to these questions. In the past, climate models have been able to predict changes in ice sheet surface mass balance, but could not simulate rapid changes in ice dynamics and associated changes in mass. The Intergovernmental Panel on Climate Change Fourth Assessment Report (IPCC AR4; Solomon et al., 2007) updated predictions of future sea level change, but explicitly stated that these predictions excluded “future rapid dynamical changes in ice flow.” The IPCC caveats are widely regarded as

# GMDD

6, 6493–6568, 2013

## Conservative regridding for ice/atmosphere coupling

E. Fischer et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



a crippling qualification of the sea level predictions contained in the IPCC AR4 report and a major and significant uncertainty in future projections.

Recently, a number of climate modeling groups have sought to rectify these deficiencies by adding dynamic ice flow effects to existing climate models (GCMs): GISS ModelE (Schmidt et al., 2006), CESM (Hurrell et al., 2013), HadGEM2 (Collins et al., 2011), etc. This is done by coupling the GCM with an existing *ice flow model*, such as Glimmer/CISM (Rutt et al., 2009), BISICLES (Cornford et al., 2013), ISSM (Larour et al., 2012), PISM (Bueler and Brown, 2009), etc. One can distinguish between *one-way* and *two-way* coupling. In one-way coupling, the GCM is used to develop surface mass balance (SMB) and temperature fields, which are then used to drive the ice flow model off-line. This process misses effects caused by feedbacks from the ice sheet to the rest of the earth system: for example, decreased ice sheet albedo (Qu and Hall, 2006) or lowered atmosphere orography (Ridley et al., 2005).

Past studies with one-way coupling have yielded useful insight into the future of present-day ice sheets, examples include Huybrechts (1994), Greve (2000), Stone et al. (2010), Bindschadler et al. (2013), Lipscomb et al. (2013), Nowicki et al. (2013a), Nowicki et al. (2013b) and Goelzer et al. (2013). However, ice sheet feedbacks are expected to be increasingly important for simulations of the long-term evolution of ice sheets and the climate associated with them. Many events in the paleo record (Dansgaard–Oeschger events, Heinrich events, the Younger Dryas) almost certainly arose from this kind of feedback.

Two-way coupling strategies address these issues by allowing the atmosphere to see changes in the ice sheet elevation, extent and albedo over time. We distinguish between *loose* and *tight* two-way coupling. Loose two-way coupling involves running a series of GCM simulations with different ice sheet configurations, each based on the result of the previous. Each GCM run is a separate simulation, without continuity of mass or energy between runs.

Studies with loose two-way coupling have yielded insight into future equilibrium states for ice sheets and climate (Ridley et al., 2005, 2010). However, they are not able

## GMDD

6, 6493–6568, 2013

### Conservative regridding for ice/atmosphere coupling

E. Fischer et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

to track transients in the evolution of ice sheets. Because ice sheets require thousands of years to reach equilibrium, results relevant to human society require an understanding of the transients.

For successful simulation of transients, we turn to tight two-way coupling. It involves running the ice flow model, step by step, along with the rest of the GCM – while conserving mass and energy along the way. Successful tight two-way coupling requires attention to conservation issues, since the GCM is simulating a more nearly closed system that could run for a long time.

When one couples dynamic ice flow models with GCM atmospheres, two models operating on different grids and timescales must communicate: ice flow models operate at low frequency on a high-resolution grid with local projection, while GCM atmospheres operate at high frequency on a low-resolution global grids. A number of issues arise due to this mismatch, including how one creates high-resolution surface mass balance fields from low-resolution GCM input.

Elevation classes address this issue. They were first introduced for precipitation downscaling in a GCM by Leung and Ghan (1998) and later applied to one-way coupling from GCM atmosphere to ice flow models by Lipscomb et al. (2013). The key insight is that surface flux fields vary approximately by elevation within a local region.

When using elevation classes, a third grid is introduced, the *elevation grid*. This allows the GCM to compute surface fields at a variety of elevations within each atmosphere grid cell, not just the elevation seen by the atmosphere. A high-resolution surface mass balance (SMB) is produced on the ice grid by first computing SMB on the elevation grid, and then using a vertical interpolation scheme to produce SMB on the ice grid. This method of interpolation produces surprisingly good results: although it cannot capture certain localized effects (e.g., wind direction), it has been shown to allow GCMs to produce surface mass balance fields approaching the quality of those produced by regional climate models (e.g., RACMO2, Vizcaino et al., 2013).

Inside the GCM, SMBs computed on the elevation grid must be regridded to the atmosphere grid as well as the ice grid. In order to maintain conservation in a tight

## Conservative regridding for ice/atmosphere coupling

E. Fischer et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

## Conservative regridding for ice/atmosphere coupling

E. Fischer et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

two-way coupled system, it is essential that the set of regridding operation chosen is self-consistent: that is, if a flux field on the elevation grid is regridded simultaneously to the atmosphere and ice grid, then the total amount of flux represented by the resulting two fields should be the same. For conservation purposes, the specifics of these two transformations are not important, as long as they are consistent with each other. Past efforts at one-way coupling have defined these two transformations in ways that each make intuitive sense, but are not consistent with each other. This is not a problem for one-way coupling, but it would cause conservation problems for two-way coupling.

This paper develops the concept of the elevation grid on which the ice surface model runs, and then derives a set of conservative regridding transformations between the atmosphere, elevation and ice grids. The coupled processes under consideration along with the transformations required for tight two way coupling are introduced in Sect. 2. Section 3 focuses on the elevation grid, while the grid fundamentals necessary for the two way coupling are presented in Sect. 4. Section 5 deals with the use of projection and associated issues encountered when bridging between the spherical geometry of GCMs and Cartesian ice sheet models. We show how to choose and implement the transformations in Sects. 6 through 9, and work through realistic examples of these transformations in Sects. 10 through 12. We touch on a number of extra “wrinkles” in the real world problem: procedures to use when the elevation grid is based on a horizontal grid other than the atmosphere grid in Sect. 13, and regridding procedures required when ice elevations, ice extent or elevation grid change in the simulation in Sect. 14. Finally, we present in Sect. 15 a library, GLINT2, that can be used to tightly couple GCMs and ice sheet models.

## 2 Coupled processes

The coupled atmosphere–ice system involves three models interacting with each other: an atmosphere model, an ice flow model, and an ice surface model situated between them (Fig. 1). The atmosphere is coupled with the ice surface model, which tracks the

## Conservative regridding for ice/atmosphere coupling

E. Fischer et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

top few meters of ice. Processes modeled here include a full surface mass/energy balance computation, snow/firn compaction, albedo effects, meltwater percolation, runoff, refreezing, etc. The bottom of the ice surface model is coupled to the ice flow model. Each model runs on its own grid: the atmosphere model runs on the atmosphere grid, the ice flow model runs on the ice grid, and the ice surface model runs on the elevation grid.

### 2.1 Ice surface model

The purpose of the ice surface model in a tightly two-way coupled system is to insulate the ice flow model from high-frequency changes happening at the surface. This is because the ice flow model, which uses a long time step, is not prepared to handle the high-frequency surface mass and energy transfer that occurs daily on an ice sheet: ice flow models generally require surface temperature and mass flux boundary conditions, assumed to be constant over the course of the ice coupling time step.

The ice surface model should be thick enough that surface-induced changes do not propagate to the bottom in any significant manner over the course of an ice coupling time step. However, there will be a small residual heat flux, no matter how thick the ice surface model is.

We will capture this physics by calculating a heat flux between the ice surface and ice flow models over the course of a coupling time step, and then “guessing” the temperature of the boundary condition that produce this flux (Schmidt et al., 2004). Since we do not expect our guess to be fully correct (due in part to ice movement), there will be a residual difference between the expected vs. actual heat flux into the ice flow model. We will ensure conservation by pushing this heat back into the ice surface model after the coupling time step. This correction, originally calculated on the ice grid, must be regrided to the elevation grid in order to be applied to the ice surface model.

The use of an ice surface model solves some important problems, but it also introduces non-physical elements into the model. Ice contained in the ice surface model does not advect along with the ice flow model, nor does it contribute to the stress field

## Conservative regridding for ice/atmosphere coupling

E. Fischer et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

of the ice below it in the ice flow model. In both cases, we expect the relative error to be small: the top 15 m of snow/firn contains less than 1 % of the mass of a 1000 m thick ice sheet. There may be ways to fix these problems – however, there is no need to make the ice surface model thicker than it needs to be. Numerous studies have shown that ice sheets below about 15 m are fully insulated from surface weather and seasonal cycles (Zagorodnov et al., 2006).

Therefore, the ice sheet model is ideally 15 m thick. As snow accumulates on top, it “pushes” excess mass down to the ice flow model. As ice melts on top, it “pulls” ice back from the ice flow model.

## 2.2 Three models, three grids

Each of the three models in the coupled system runs on its own grid. The atmosphere is run on the atmosphere grid ( $A$ ) and the ice flow model is run on the ice grid ( $I$ ). The ice surface model is run on the elevation grid ( $E$ ), which is based on the atmosphere grid (see Sect. 3). All three grids are two dimensional, in the sense that they are used to construct two-dimensional functions  $f(x, y)$  over the domain. Regridding operations are needed to pass mass and energy fluxes between the models.

It is important to keep in mind the relative size of the three grids. We set up a test using the GISS  $2^\circ \times 2\frac{1}{2}^\circ$  atmosphere grid (Schmidt et al., 2006), overlapping with the SeaRISE 5 km grid (Bindschadler et al., 2013) and 40 elevation points, spaced every 100 m from 0 m to 4000 m. In that case,  $I$  had 66 906 grid cells,  $A$  has 146 and  $E$  has 1829. These numbers only account for grid cells involved with the Greenland ice sheet. In general, the ice grid will be finest, the atmosphere grid coarsest, and the elevation grid in the middle.

Time frequency mismatches are another issue. The atmosphere runs at high frequency, each time step being typically 1 h. On the other hand, the atmosphere and ice flow models are coupled at much lower frequency, typically 1 month. We call these two time steps the *atmosphere time step* and *ice coupling time step*.

## 2.3 Fully coupled system

Figure 2 shows the data flow of the fully coupled system. Steps of the data flow are organized based on their frequency: the top circle of steps runs at the same frequency at the GCM atmosphere, while the bottom circle runs at the *ice coupling* frequency – typically one month or more. We describe the steps involved in coupling the GISS ModelE (Schmidt et al., 2006) with PISM (Bueler and Brown, 2009); however, these steps are general for any GCM or ice flow model. We now trace through the data flow on a typical coupled run.

## 2.4 Atmosphere time step

When the atmosphere runs, it produces a set of fields on the atmosphere grid that affect the processes in the ice surface model: for example, downwelling longwave radiation, downwelling shortwave radiation, precipitation, etc. These fields must be regridded from  $A$  to  $E$ . We denote the set of fields to be regridded with a capital letter,  $R^A$ ; the result of the regridding is  $R^E$  (see Appendix A for notation conventions).

The ice surface model is run at the same frequency as the atmosphere. Among its outputs are mass and energy fluxes with the atmosphere: evaporation, sublimation, upwelling longwave radiation, latent heat release, etc. These are represented by  $F^E$  and are regridded to  $F^A$  before being passed back to the atmosphere on the next time step.

The ice surface model also produces fluxes into the ice flow model; these are described immediately below.

## 2.5 Coupling time step

On each atmosphere time step, flux outputs from the ice surface model are accumulated as  $\Phi^E$ . Every coupling time step – about once a month – the accumulated  $\Phi^E$  is regridded to the ice grid ( $\Phi^I$ ) and passed to the ice flow model.

# GMDD

6, 6493–6568, 2013

## Conservative regridding for ice/atmosphere coupling

E. Fischer et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

The ice flow model produces changes in ice surface topography and extent, as well as a small energy flux between the ice flow and ice surface models ( $\Delta^I$ ). These changes are fed back into the atmosphere and ice surface model – after being appropriately regrided to  $\Delta^E$  and  $\Delta^A$ .

## 5 2.6 Regridding requirements

We see that the fully coupled system requires the use of five regridding operations at various points in its run (Fig. 2). These regridding operations must be conservative, in the sense that *none* of them can change the integral of the field on the domain as a whole. In fact, we would like to impose a stronger conservation condition, that values are conserved within each atmosphere grid cell.

We develop these five regridding operations in the sections below. We begin by discussing the method of elevation classes in a general manner (Sect. 3), and then move on to developing basics of conservative regridding (Sect. 4). Because conservation is defined in terms of integration over areas, part of our discussion involves a definition of how to integrate functions on the elevation grid  $E$ . Finally, we show how the required set of operations can be constructed in a fully self-consistent manner.

Throughout, we assume that the atmosphere grid  $A$  has L0 parameterization: functions on  $A$  are represented by their mean value within each grid cell. This becomes our basis for conservation: all regridding operations are conservative over every atmosphere grid cell. We do not require any specific parameterization for the ice grid, but we consider cases for L0 (piecewise constant) and L1 (piecewise linear) parameterizations, the latter which are commonly used in finite element ice models.

## 3 Elevation points

The method of elevation classes, when applied to the atmosphere–ice coupling problem, involves running the ice surface model at one or more elevations in each

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



atmosphere grid cell (Lipscomb et al., 2013). Temperature, pressure and precipitation are extrapolated to a set of elevations within the grid cell. They are then used in an ice surface model to compute a full surface mass and energy balance at each elevation. The result is a set of “what-if” scenarios, giving an estimate of what the fluxes *would have been*, had the ice surface of the grid cell been at the given elevation – rather than the elevation seen by the atmosphere model.

The practitioner has wide latitude in the choice of elevations to use for each atmosphere grid cell. The simplest approach is to use a fixed set of elevations across all grid cells – for example, every 100 m from 0 m to 4000 m.

However elevations are chosen, the result is a new “grid” – the *elevation grid* – on which the ice surface model is run and surface fluxes are generated. (See Appendix A for definition of *grid*.) The elevation grid is derived from the atmosphere grid, in the sense that each elevation grid “cell” (or *elevation point*) is associated with one parent atmosphere grid cell. If elevation point  $E_j$  is associated with atmosphere grid cell  $A_j$ , we write  $E_j \in A_j$ .

Once the GCM has computed a conserved quantity on the elevation grid, those values can be used to develop a relation between elevation and SMB within each atmosphere grid cell. Suppose we have computed a flux field  $f^E$  – SMB, for example. For an atmosphere grid cell  $A_j$ , consider the values  $f_j^E$  for all  $j$  such that  $E_j \in A_j$ . We can think of these values as samples of a 1-D function relating elevation to flux (SMB), *within the localized region* of  $A_j$ . By interpolating between those points using standard methods, one can construct a continuous function relating flux to elevation within  $A_j$ . As long as enough points are used, and the function being interpolated is smooth enough, this procedure will yield an arbitrarily accurate representation of the “true” function.

In fact, the functions we typically wish to interpolate – surface mass and energy balance averaged over about a month – are quite smooth as functions of elevation (Fig. 3). In the face of spatially invariant precipitation, one would expect SMB to be constant above the equilibrium line altitude (ELA), and to decrease linearly with elevation below the ELA. Near the ELA, one would expect a smooth transition because the

## Conservative regridding for ice/atmosphere coupling

E. Fischer et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



ELA goes up and down over a month of diurnal cycles. This is in agreement with experimental work, which has shown SMB below the ELA to vary linearly with temperature (Braithwaite, 1981; Box et al., 2013).

How many elevation points are required to properly resolve the elevation–flux relationship? This depends on the interpolation scheme: higher order schemes will require fewer points than piecewise linear interpolation. But the general shape of the function – two straight lines connected by a curve near the ELA – implies that not many points are needed, except for near the ELA.

If one knows where the ELA is on every atmosphere grid cell, then this is a useful guide in selecting elevation points. But if one is studying ice sheets in a changing climate, then the ELA will be expected to move over time. It is possible to adaptively move elevation points as the ELA moves. But it is simpler just to use many points everywhere. In our tests, we have used 40 points at 100 m spacing, which is probably sufficient for coupled ice sheet simulations.

The method of elevation classes – or elevation points – provides a way to construct an interpolated relation between elevation and surface mass/energy flux within each atmosphere grid cell. Additional choices need to be made in order to produce fully downscaled flux fields on the ice grid (Sect. 6). We are now almost ready to define the transformations posited in Sect. 3. But first, we pause for some groundwork on grid fundamentals and projection issues.

## 4 Grid fundamentals

Numerical models represent continuous fields by linear combinations of a finite set  $G$  of basis functions – which we call a *parametrization* (Appendix A describes our mathematical notation). For example, suppose that  $G$  uses the basis functions  $\mathbf{g}(x, y) = [g_1(x, y) \dots g_n(x, y)]$ , where  $\mathbf{g}(x, y)$  is the vector of all the basis functions. Suppose we have an  $n$ -dimensional vector  $\mathbf{f}^G$  with components  $f_i^G$ . That vector represents the function:

### Conservative regridding for ice/atmosphere coupling

E. Fischer et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



$$f^G(x, y) = \mathbf{f}^G \cdot \mathbf{g}(x, y) \quad (1)$$

Most commonly in climate modeling,  $f_i^G$  represents the mean value of  $f^G(x, y)$  within some well-defined bounded region, which we call a *grid cell*. From a conservation point of view, a function  $f^G(x, y)$  with L0 parameterization may be taken to be constant within each grid cell, with discrete “jumps” from one grid cell to the next. In this case, the basis function  $g_i(x, y)$  for grid cell  $i$  is equal to one inside the cell and zero outside.

L0 parameterizations are widely used in climate and finite difference ice flow models. Finite element ice flow models might use L1 or even higher-order parameterizations (Zienkiewicz et al., 2013), and they use the term “mesh” to describe the geometry of their basis functions. In this paper, we use the term “grid” both to refer to the vector space in which  $\mathbf{f}^G$  lives, along with its associated parameterization.

## 4.1 Integration on grids

Because we are working with conserved quantities, it is essential that we can integrate functions over any well-defined region  $B$ . Because a parameterization defines  $f(x, y)$  on every point, this integral is well-defined. Substituting from Eq. (1) and using linearity, we get:

$$\int_B f(x, y) dA = \mathbf{f}^G \cdot \int_B \mathbf{g}(x, y) dA \quad (2)$$

If we wish to integrate over  $B$  repeatedly, we can pre-compute  $\int_B \mathbf{g}(x, y) dA$ , and then take dot products with  $\mathbf{f}^G$  as needed. How one integrates the basis functions depends on the nature of those basis functions. We give specifics for L0 and L1 grids in Appendices B and C, respectively.



Area-weighted remapping is closely related to our definition of equivalence above. If we have two fields on two different grids, equivalence on  $H$  can be tested by regriding both fields to  $H$  and comparing the resulting vectors. The transformation from  $G$  to  $H$  is linear, and thus representable by a matrix. It is variously called *area-weighted remapping* or *conservative regridding* (Ramshaw, 1985). The matrix is computed by integrating the source basis functions  $g_1(x,y) \dots g_n(x,y)$  over every grid cell in the destination grid.

#### 4.4 Interpolation grid

We are now almost ready to discuss our implementations of the transformations  $E \rightarrow I$  and  $E \rightarrow A$  (see Appendix A for mathematical notation). But first, we must introduce the *interpolation grid*  $G$ , which is used to rigorously define basis functions on  $E$ .

In general,  $G$  is defined simply as the ice grid  $I$ . If  $I$  is L0-parameterized, we might instead choose  $G$  to be the *exchange grid* between  $A$  and  $I$  (Fig. 4): the L0 grid whose grid cell outlines are formed by the intersection of grid cells in  $A$  and  $I$  (Balaji et al., 2006). The exchange grid is a useful choice for  $G$  because every exchange grid cell overlaps at most one atmosphere grid cell. This choice has its pros and cons, which we explore in Sect. 9. Either way, the interpolation grid is similar to the ice grid, and can be thought of as an ice grid proxy in most cases.

#### 5 Projection issues

Whether the source grid is L0 or L1, area-weighted remapping algorithms need to find the intersection of grid cell outlines from two grids. Technically, this is only possible if the two grids exist on the same surface. In our case, atmosphere models exist on the surface of a sphere, whereas ice flow models work on a Cartesian plane. The exchange grid between an atmosphere grid and ice grid cannot be directly computed.

## GMDD

6, 6493–6568, 2013

### Conservative regridding for ice/atmosphere coupling

E. Fischer et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

## Conservative regridding for ice/atmosphere coupling

E. Fischer et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

This problem is solved using a map projection (Snyder, 1987). We let  $A'$  be the original atmosphere grid, and then let  $A$  in be the projected atmosphere grid – projected to the Cartesian plane using the chosen projection. Regridding computations described in this paper are made to/from  $A$ , the projected grid.

Another way to solve projection-related issues would be to formulate an ice flow model in spherical coordinates. This would be convenient for ice sheets the size of Greenland or Antarctica. But for larger globe-spanning ice sheets, such as those on a Snowball Earth, it might turn out to be essential.

### 5.1 Projection error

The area of a grid cell can change when it is projected: for example, the stereographic projection used in SeaRISE (Bindschadler et al., 2013) can change the size of a grid cell by up to 10% over Greenland. Even if an area-preserving projection is used, the area can change due to geometric error or numerical artifacts (Sect. 5.2). Therefore, data must be rescaled in order to maintain conservation when transforming a field  $f^{A'}$  to  $f^A$ :

$$f_i^A = \frac{|A'_i|}{|A_i|} f_i^{A'} \quad (5)$$

Note that this transformation is diagonal and invertible (Fig. 14). Because this is simply a rescaling, the regridding schemes presented in this paper are not operationally affected by projection issues.

The distinction between the original atmosphere grid  $A'$  and the projected atmosphere grid  $A$  allows us to maintain conservation, but it introduces unphysical artifacts. The Stereographic projection used in SeaRISE (Bindschadler et al., 2013) can change the area of grid cells by up to 10%. If a projected grid cell is 10% smaller than the original, then that means that 1 m of accumulation in the GCM will turn into 1.1 m in the ice flow model. This could cause significant discrepancies in dynamic ice flow.

One solution to this problem would be to use an equal area projection – the Lambert Equal Area Projection, for example. However, equal area projections do not preserve angles, which distorts the ice dynamics at a local scale. We conclude that non-physical distortions happen whether an area-preserving or angle-preserving projection is used.

5 It is not yet clear which choice gives better results in the end.

One innovative approach to this problem is to use an angle-preserving projection, but to allow ice flow model grid cells to vary in size, based on the local distortion of space introduced by the projection. Parameters  $dx$  and  $dy$  are kept for each grid cell. These grid parameters are included in the ice flow model equations, thereby accounting  
10 for shape and area distortion caused by the projection in a physically meaningful way (Pollard and DeConto, 2012).

## 5.2 Geometric error

Even if an equal area projection is used, in practice grid cells will change area anyway. This is because projected grid cells have complex shapes, but the algorithms in  
15 Appendix D are only able to approximate these shapes with polygons (Lauritzen and Nair, 2008).

Geometric error may be arbitrarily reduced by increasing the number of sides used in the approximating polygons – the parameter  $n$  specifies the number of line segments used to approximate each side of the original grid cell in its projected state. But this  
20 method has its limit in practical terms.

Consider a typical latitude–longitude grid on the sphere with a Lambert Equal Area Projection (Fig. 5). In this case, the area of a polygonal approximation will always be *smaller* than the area of the actual grid cell: we end up inscribing polygons inside of circles. The geometric error will depend on the number of sides of the inscribed  
25 polygon, which in this case depends on the number of grid cells in the circle, and on  $n$ .

This is the method used by Archimedes to approximate the value of  $\pi$  in  $\sim 250$  BCE (Heath and Archimedes, 1897, p. 91). Unfortunately, it converges only quadratically, as  $O(1/n^2)$ . Meanwhile, memory use to store all those polygons goes up by  $O(n)$ . Memory

## Conservative regridding for ice/atmosphere coupling

E. Fischer et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



and time requirements will be exponential in terms of the number of digits of accuracy required: each additional digit of accuracy will require an increase in the number of sides by a factor of  $\sqrt{10}$ .

### 5.3 Geometric error example

5 We tested this theory in our code. Using the GISS  $2^\circ \times 2\frac{1}{2}^\circ$  grid on a spherical grid and polar Lambert Equal Area Project, we computed the geometric error caused by the polygonal approximation, for a variety of  $n$  from 1 through 100. Because the projection is centered on the pole, all grid cells have the same geometric error.

10 In Fig. 6, we plot the observed error as a function of  $n$ . As expected, error decreases linearly on a log–log plot. However, the decrease becomes sub-linear for  $n$  between 10 and 100, with the total error approaching  $10^{-6}$ . Convergence in practice is even worse than in theory, due to limitations in our numerical methods. In practice, we do not expect to use  $n > 10$ , avoiding sub-linear convergence problems for larger  $n$ .

### 6 Interpolation: elevation to ice grid

15 We can now derive the first of the five transformations posited in Fig. 2. We begin with the transformation  $E \rightarrow G$  from the elevation grid to interpolation grid, because this is the transformation that ice modelers are most concerned with when seeking high-quality downscaled flux fields. In various contexts, this transformation might be referred to a an *interpolation*, *downscaling* or *regridding* operation.

20 In Sect. 3, we showed how values  $f^E$  at elevation points may be used to construct a flux–elevation relationship  $f_i(z)$  in each atmosphere grid cell  $A_i$ . These relationships may then be used to construct  $f^G$  on the interpolation grid. The resulting transformation for  $E \rightarrow G$  potentially involves horizontal and vertical interpolation, of which the modeler has considerable choice.

## Conservative regridding for ice/atmosphere coupling

E. Fischer et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



We present here three methods of interpolation from  $E$  to  $G$ : Z Interpolation (Sect. 6.1), Bilinear Interpolation (Sect. 6.2) and Elevation Class Interpolation (Sect. 6.3). All three methods assume that  $G$  is L0 parameterized and can be used for the basis of a conservative coupling. And since they are all linear, they can all be represented by a (sparse) matrix, which we will call  $\mathbf{M}$ . The methods are extended to the L1 case in Sect. 6.4

## 6.1 Z Interpolation

Consider an L0-parameterized interpolation grid cell  $G_j$  with mean elevation  $z_j$ . Suppose  $G_j$  is wholly contained in one atmosphere grid cell  $A_i$ .  $A_i$  has an interpolated flux–elevation relationship  $f_i(z)$ , based on  $f^E$ , and we set  $f_j^G = f_i(z_j)$ .

Suppose that  $G_j$  intersects more than one atmosphere grid cell. The same procedure is used, selecting values of  $f_i(z_{G_j})$  for every  $A_i$  intersected by  $G_j$ . These values are then summed together, weighted by  $|G_j \cap A_i|$ .

We call this *Z Interpolation*. As long as standard interpolation methods are used to construct  $f_i(z)$ , Z Interpolation defines a linear function from  $f^E$  to  $f^G$  and can be represented by a matrix. An example of the result is shown in Fig. 7. In general, Z Interpolation produces SMB fields that vary smoothly within each atmosphere grid cell, but that contain discontinuities between grid cells.

## 6.2 Bilinear interpolation

There is some concern that discontinuities created by Z Interpolation could cause problems as an input to an ice flow model. In that case, a bilinear interpolation step may be used to create a smooth field, as in Lipscomb et al. (2013). This scheme sets the value of interpolation grid cell  $G_j$  equal to a linear combination of  $f_j(z)$ ,  $f_k(z)$ ,  $f_l(z)$  and  $f_m(z)$ , where atmosphere grid cells  $A_j$ ,  $A_k$ ,  $A_l$  and  $A_m$  are the four grid cells with centers closest to the center of  $G_j$  (Fig. 8). Results are shown in Fig. 9.

# GMDD

6, 6493–6568, 2013

## Conservative regridding for ice/atmosphere coupling

E. Fischer et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Bilinear Interpolation has the advantage over Z Interpolation in that it produces smooth fields. However, bilinear interpolation has a number of problems:

- The fields it produces will have a significantly different total mass than the fields produce by Z Interpolation. Our experience with monthly SMB fields over Greenland indicates that storms tend to dump large amounts of snow in a few localized areas. Bilinear Interpolation tends to reduce the total amount of snowfall in these cases, causing potentially significant differences in the GCM model run.

Variations in total mass caused by the choice of interpolation procedure will *not* cause conservation problems: we make this clear in Sect. 7. However, an interpolation scheme that produces a significantly different mass from the apparent “intent” of the GCM could cause biases. In particular, a scheme that makes snowstorms smaller than the GCM “intended” would produce a negative bias in the equilibrium extent of the ice sheet.

- It introduces significant numerical diffusion into the system.
- The  $A \rightarrow E$  transformation derived with bilinear interpolation introduces non-physical artifacts (Sect. 12).
- It is also not always clear how to extend bilinear interpolation to the case of non-rectangular atmosphere grids. This problem can also affect non-regular points in mostly regular grids (e.g., at the poles of a latitude/longitude grid).

### 6.3 Elevation class interpolation

One final choice for regridding is to define elevation *classes* as they were originally formulated (Leung and Ghan, 1998). In this approach, each atmosphere grid cell is grouped into sub-regions based on elevation bands. The grid  $E$  can then be thought of as an L0 grid with irregularly shaped grid cells (Fig. 10). Area-weighted remapping is used to interpolate from the elevation grid to the atmosphere grid.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Elevation class interpolation is equivalent to a specialized form of Z Interpolation, in which the 1-D function  $f_i(z)$  (for grid cell  $A_i$ ) is interpolated as a discontinuous piecewise constant function (Fig. 11), rather than a piecewise polynomial (Fig. 3).

Unless one expects significant discontinuities in topography or the elevation–flux relationship, a zero-order interpolation scheme such as Elevation Class Interpolation would be expected to give a less accurate version of  $f^G$  than a higher order scheme such as Z Interpolation. Since we expect the elevation–flux relationship to be smooth with elevation, we do not recommend the use of Elevation Class Interpolation over Z Interpolation.

## 6.4 Interpolating to L1 grids

We have outlined recipes for the interpolation  $E \rightarrow G$ , as long as  $G$  uses an L0 parameterization. The procedures need to be modified for interpolation on ice grids using L1 parameterization.

In a finite element mesh (“grid”), field values are determined at mesh vertices and linearly interpolated within each triangular element (Zienkiewicz et al., 2013). Any of the interpolation methods above may be used to determine the value of  $f^G(x, y)$  at each vertex. Once vertex values have been interpolated on the vertices of a finite element mesh, the value of  $f^G(x, y)$  at all other points is fully determined.

## 7 Regridding: elevation to atmosphere grid

As shown in Fig. 2, a regrid step from the elevation to the atmosphere grid is required on every GCM time step. Once the modeler has chosen the transformation  $E \rightarrow G$ , we show here how to derive a transformation for  $E \rightarrow A$  that is consistent with it.

To derive this transformation, consider one GCM time step, during which a flux field  $f^E$  between the atmosphere and ice surface is computed on the elevation grid. That field will be regridded both to the interpolation and atmosphere grids. Conservation

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



requires that the resulting fields are equivalent on  $A$ , i.e:

$$f^A \equiv^A f^G \quad (6)$$

If the transformation  $E \rightarrow G$  derived in Sect. 6 is represented by the matrix  $\mathbf{M}$ , then we can write  $f^G = \mathbf{M}f^E$ . Substituting into Eq. (6), we get the requirement:

$$f^A \equiv^A \mathbf{M}f^E \quad (7)$$

Equivalence on  $A$  can be tested by conservatively remapping both sides to  $A$  and then comparing. Denoting  $\mathbf{R}$  as the matrix that conservatively regrids from  $G$  to  $A$ , we have the requirement:

$$f^A = \mathbf{R}\mathbf{M}f^E \quad (8)$$

If we treat this as a definition for the transformation  $E \rightarrow A$ , we have derived a transformation for  $E \rightarrow A$  that is consistent with the transformation we already chose for  $E \rightarrow G$ .

In other words (Fig. 12): we will regrid from  $E$  to  $A$  by first regridding from  $E$  to  $G$  (represented by the matrix  $\mathbf{M}$ ), and then use area-weighted remapping from  $G$  to  $A$  (represented by the matrix  $\mathbf{R}$ ). The GCM, which needs to use the transformation  $E \rightarrow A$ , does not need to know anything about the ice grid used to derive that transformation. It just needs to know the final matrix  $\mathbf{R}\mathbf{M}$ , which can be pre-computed via matrix multiplication.

We have derived a transformation for  $E \rightarrow A$  that *by definition* is consistent with a previously chosen transformation for  $E \rightarrow G$ . Our approach differs from previous efforts, which would start out with  $E \rightarrow A$  and try to find a transformation for  $E \rightarrow G$  that is consistent with it. We believe our approach makes sense because modelers care more about the details of  $E \rightarrow G$  than  $E \rightarrow A$ .

## Conservative regridding for ice/atmosphere coupling

E. Fischer et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## 7.1 A basis for $E$

Every vector space has basis functions, including the elevation grid  $E$ . Our choice for the transformation  $E \rightarrow A$  constrains the basis functions we use on  $E$ . To find these basis functions, we expand on the central idea in the previous section: that we can determine properties of a field on the elevation grid by regriding to the ice grid. In this case, we define  $f^E(x, y)$  to be equal to  $f^G(x, y)$ , where  $f^G = \mathbf{M}f^E$ . In other words, we can evaluate  $f^E(x, y)$  at a point by interpolating  $f^E$  to  $G$  and then evaluating  $f^G(x, y)$ . This definition is consistent with our recipe for  $f^A$  in the previous section.

We now use this principle to obtain a formula for the basis functions of  $E$ . Substituting into Eq. (1), we get:

$$f^E(x, y) = f^G(x, y) = \mathbf{M}f^E \cdot \mathbf{g}(x, y) \quad (9)$$

Rewriting in indicial notation, using the commutative property of multiplication, and swapping index letters, we get:

$$f^E(x, y) = f_i^E M_{ji} g_j(x, y) \quad (10)$$

We can use this, along with Eq. (1), to determine the basis functions for the elevation grid:

$$e_i(x, y) = M_{ji} g_j(x, y) \quad (11)$$

or switching back to vector notation, we have:

$$\mathbf{e}(x, y) = \mathbf{M}^T \mathbf{g}(x, y) \quad (12)$$

Equivalently, the basis function  $e_i(x, y)$  is the function obtained if we set the  $f_i^E = 1$  and all other components of  $f^E = 0$ , regrid to the interpolation grid, and then examine the resulting function  $f^G(x, y)$ .

# GMDD

6, 6493–6568, 2013

## Conservative regriding for ice/atmosphere coupling

E. Fischer et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



In general, these basis functions will be not orthogonal. Their exact form depends on choices made in choosing **M**: vertical and horizontal interpolation choices, as well as grid geometry issues. We have plotted some example basis functions in Fig. 13.

Note that if elevation class interpolation is used (Sect. 6.3) and  $G$  is the exchange grid, then our scheme reduces to a traditional elevation class scheme, and basis functions will represent constant-value sub-grid tiles, which are orthogonal.

## 7.2 Forward transformations: summary

We have now defined three of the five regriding transformations required by Fig. 2:  $E \rightarrow G$ ,  $G \rightarrow A$  and  $E \rightarrow A$ . We call these the *forward transformations* because they are linear and can be represented by matrices. See Fig. 14 for a diagram of how these transformations may be used to regrid fields.

We defined these three transformations in a consistent manner. We began by allowing the user choice in constructing the transformation from the elevation to interpolation grids, represented by matrix **M**: this makes sense because users desire choice for  $E \rightarrow G$ . We then noted that standard area-weighted remapping, represented by matrix **R**, may be used to regrid from the interpolation grid to the atmosphere grid. From these two choices, transitivity requires that the transformation from the elevation to atmosphere grid is represented by the matrix **RM**. Conservation is maintained because  $E \rightarrow G$  and  $E \rightarrow A$  are consistent with each other, producing  $f^G$  and  $f^A$  that are equivalent on every grid cell in  $A$ .

Our choices for these transformations lead directly to a well-defined set of basis functions for the grid  $E$ , examples of which we plotted. Fields on this grid will be represented in terms of these basis functions – for example, surface–atmosphere fluxes and ice surface model state.

# GMDD

6, 6493–6568, 2013

## Conservative regridting for ice/atmosphere coupling

E. Fischer et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

## 8 Reversing the transformations

We have defined three transformations so far, but Fig. 2 indicates that five are necessary for full functioning of the coupled system. We still need to derive appropriate procedures for the “reverse transformations”  $A \rightarrow E$  and  $G \rightarrow E$ , indicated in Fig. 14 as dotted lines. Because of the differences in dimensionality between the three grids (Sect. 2.3),  $\mathbf{M}$  and  $\mathbf{RM}$  are not invertible in a simple linear algebraic sense:  $A \rightarrow E$  is underdetermined, and  $G \rightarrow E$  is overdetermined. We still must find ways to compute these transformations that retain conservation over atmosphere grid cells.

### 8.1 Transformation: atmosphere to elevation

Suppose we have a flux field  $f^A$  on the atmosphere grid  $A$ . We wish to regrid it to an “equivalent” flux field  $f^E$  on the elevation grid  $E$ . In this case,  $f$  might represent precipitation or downwelling radiation from the atmosphere.

The problem is underdetermined. Any solution for which  $f^A \equiv^A f^E$  will be conservative. This is the same as requiring that:

$$\mathbf{RM}f^E = f^A \quad (13)$$

Because of the many-to-one relationship between elevation and atmosphere grid cells, our intuition tells us that  $f^E$  may be constructed simply by repeating values of  $f^A$  within each atmosphere grid cell. This is physically self-consistent. More precisely, if  $E_j \in A_i$ , then we would like to set  $f_j^E = f_i^A$ . We define a simple linear transformation  $\mathbf{\Lambda}$  that does this:  $f^E = \mathbf{\Lambda}f$ .

If the matrix  $\mathbf{RM}$  computes each  $f_j^A$  as a weighted sum *only* of values  $f_j^E$  where  $E_j \in A_i$ , then we say that  $\mathbf{RM}$  is a *local transformation* – it uses only data from “within” an atmosphere grid cell to compute a value on that grid cell. If  $\mathbf{RM}$  is local, then the intuitive notion that  $f^E = \mathbf{\Lambda}f^A$  will be conservative: it is easy to show that  $\mathbf{RM}\mathbf{\Lambda}f^A = f^A$  (see Eq. 13). The weights involved in  $\mathbf{RM}$  do not matter, as long as they sum to one for each component of  $f^A$ .

Therefore, if **RM** is local, we can simply use  **$\Lambda$**  for our transformation  $A \rightarrow E$ . Even better, it is easy to show that there will be no numerical dispersion in the round-trip transformation  $E \rightarrow A \rightarrow E$  because  **$\Lambda \mathbf{R} \mathbf{M}$**  is the identity matrix.

## 8.2 Non-local RM

5 If **RM** is not local, then in general  **$\Lambda \mathbf{f}^A$**  and  **$\mathbf{f}^A$**  will not be equivalent on  $A$ , or even the entire ice sheet. We are forced to trade off between the most physically self-consistent value for  **$\mathbf{f}^E = \Lambda \mathbf{f}^A$** , vs. something that conserves. We can use  **$\Lambda$** , along with quadratic optimization, to guide us to such a compromise.

We seek a vector  **$\mathbf{f}^E$**  such that:

$$10 \quad \mathbf{R} \mathbf{M} \mathbf{f}^E = \mathbf{f}^A \quad (14)$$

We also wish that vector to be as close as possible to our intuition above. That is, we wish to minimize the quantity:

$$15 \quad \|\mathbf{f}^E - \Lambda \mathbf{f}^A\|^2 \quad (15)$$

where our  $L_2$  norm is weighted by the weight of each basis function. That is, we use a weight vector  **$\mathbf{w}$**  where:

$$w_j = \int_{\infty} e_j(x, y) dA \quad (16)$$

20 This is a sparse quadratic optimization problem with equality constraints. A number of numerical packages can solve it, we used GALAHAD (Gould et al., 2003). In our tests, solution typically requires a fraction of a second on a single core. That is so fast that we have found no need to consider sub-optimal solutions to this problem.

25 This procedure introduces some numerical dispersion into the fully coupled system from non-local regridding operations: by numerical dispersion, we mean movement of

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[⏪](#)

[⏩](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



mass between adjacent atmosphere grid cells, thereby violating our desired property of conservation with each atmosphere grid cell. Considering the round-trip transformation  $E \rightarrow A \rightarrow E$ , it is clear that this transformation is not local: both because **RM** is not local, and because the quadratic program set up for  $A \rightarrow E$  will be non-local.

### 8.3 Practical issues for non-local RM

GCMs are well equipped to deal with *sub-grid tiles*, each one occupying a fraction of the overall grid cell. The GCM will typically implement  $E \rightarrow A$  by computing a weighted sum over sub-grid tiles in each grid cell, with weights based on each tile's fractional area. This capability is used to implement traditional elevation class schemes.

If the **RM** matrix is local, then by definition the value on each atmosphere grid cell is a weighted sum of the elevation points in that grid cell. This is compatible with existing GCM practice that assumes sub-grid tiles. To use the methods in this paper, the computation of  $E \rightarrow A$  inside the GCM does not need to be replaced, it need only be fed a new set of weights. Even though elevation points do not have well-defined areas, the weight for elevation point  $j$  can be thought of as the "fractional area" that an elevation point contributes to its containing atmosphere grid cell. The GCM can be coded as if traditional elevation classes are being used, even if the user has chosen a more numerically accurate form of vertical interpolation in the construction of the **M** matrix.

Things get more complicated for the GCM if the **RM** matrix is *not* local. Instead of using a simple set of weights, the GCM will have multiply by a general sparse matrix **RM** when computing  $E \rightarrow A$ . An MPI gather will be required to compute  $A \rightarrow E$ . Finally, some GCMs use implicit schemes for ice surface/atmosphere coupling (Best et al., 2004), requiring a matrix inversion on every time step. If **RM** is non-local, then a large, sparse, global matrix inversion will be required, rather than a number of small, local inversions.

Because of the significant practical complications that arise with the use of a non-local **RM** matrix, most users find it simplest to use a local **RM** matrix if at all possible.

## Conservative regridding for ice/atmosphere coupling

E. Fischer et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

## 8.4 When is RM not local?

The use of a non-local **RM** transformation can add burden to the GCM developer. It is therefore important to delineate cases in which **RM** is not local.

For L1 ice grids, as is used with a finite-element ice flow model, **RM** will be non-local: ice elements that straddle two atmosphere grid cells will by necessity involve elevation point values from two atmosphere grid cells. This will make **R** slightly non-local.

Even for L0 grids, **M** could be non-local, depending on the choice of interpolation schemes for  $E \rightarrow G$ . Bilinear interpolation is inherently non-local, whereas the other interpolation strategies mentioned above are local.

For L0 ice grids with a local interpolation scheme, **RM** can be made local, as long as  $G$  was chosen to be the exchange grid (see Sect. 4.4). In this case, **M** will be local – because each exchange grid cell attains its value from elevation points in just one atmosphere grid cell. **R** will be local as well, for the same reason.

## 8.5 Transformation: interpolation grid to elevation grid

Suppose we have a flux field  $f^G$  on the interpolation grid  $G$ . We wish to regrid it to an “equivalent” flux field  $f^E$  on the elevation grid  $E$ . As before, we will set up a quadratic optimization problem.

It would be nice if we could find  $f^E$  such that  $\mathbf{M}f^E = f^G$ . This will not usually be possible, since  $G$  has far more degrees of freedom than  $E$ . Instead, we will minimize the quantity:

$$\|\mathbf{M}f^E - f^G\|^2 \quad (17)$$

while we maintain conservation on each atmosphere grid cell:

$$\mathbf{R}Mf^E = \mathbf{R}f^G \quad (18)$$

# GMDD

6, 6493–6568, 2013

## Conservative regridding for ice/atmosphere coupling

E. Fischer et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion





If one directly uses the ice grid for  $G$ , then the **RM** matrix will *not* be local, producing numerical dispersion for the transformation  $E \rightarrow A$  and complicating the reverse transformation  $A \rightarrow E$ . However, there will be no numerical dispersion for the transformation  $E \rightarrow I$ .

5 On balance, we recommend keeping the **RM** matrix local if possible. Not only does this simplify implementation, it moves unavoidable numerical dispersion away from the atmosphere time step to the less frequent ice coupling time step.

## 9.1 Exchange grids for finite elements

10 The discussion in Sect. 9 assumes an L0 ice grid. The results can be extended to L1 or higher-order parameterizations: finite element ice models, for example. In this case, the concept of exchange grid is not meaningful, since exchange grids are by definition L0.

15 Instead of the exchange grid, one can choose  $G$  to be just about *any* L0 grid, of resolution at least as high as the ice mesh, where grid cells do not cross atmosphere cell boundaries: we will call this a *generalized exchange grid*. The user would then need to develop an appropriate conservative transformation for  $G \rightarrow I$ , whereas the transformation for  $I \rightarrow G$  would remain an area-weighted remapping (see Appendix C). Details of this scheme for a particular ice mesh parameterization are left to the reader.

20 As in Sect. 9, the transformation  $G \rightarrow I$  would cause some numerical dispersion, due to the geometry of the ice mesh. Additional dispersion would be introduced because of mismatch between the L0 grid  $G$  and the higher-order mesh  $I$  – although this could be reduced by making  $G$  finer. The tradeoffs of choosing  $G$  to be a generalized exchange grid vs. the ice grid are the same as in Sect. 9: numerical dispersion is introduced in the ice coupling timestep, in exchange for simplifying implementation and eliminating  
25 dispersion in the atmosphere timestep.

## Conservative regridding for ice/atmosphere coupling

E. Fischer et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

## 10 Regridding examples: local RM

Having shown how to compute all five required regridding transformations, we now demonstrate them working within a realistic GCM context. We set up a test using the GISS  $2^\circ \times 2\frac{1}{2}^\circ$  atmosphere grid (Schmidt et al., 2006), overlapping with the SeaRISE 5 km L0 grid (Bindschadler et al., 2013) and 40 elevation points, spaced every 100 m from 0 m to 4000 m. We chose the exchange grid as  $G$  (see Appendix A for notation conventions). We ran GISS ModelE in this configuration, using fixed sea surface temperatures corresponding to the years 1996–2005. The ice surface model was run on the elevation grid, producing monthly averages of SMB over Greenland, which we label  $f^E$  (Fig. 15).

As expected, the broad pattern shows strong melting in narrow bands at low elevations, along with weak accumulation at high elevations. Atmosphere grid cell boundaries are prominent because precipitation – the primary source of positive SMB – is not downscaled to sub-grid resolution (Leung and Ghan, 1998). The small oscillations at high elevation are artifacts introduced by ModelE’s ice surface model.

### 10.1 Example: $E \rightarrow A$

Figure 16 shows the original field  $f^E$  computed on the elevation grid and regridded to the atmosphere grid  $A$ , via the transformation  $f^A = \mathbf{RM}f^E$ . Note how low-elevation ablation regions become broader and weaker (compared to Fig. 15), whereas the interior of the ice sheet remains about the same. This demonstrates the benefits obtained through the use of elevation points, as compared with running the ice surface model on the atmosphere grid (van den Broeke et al., 2008).

By definition, the transformation  $\mathbf{RM}$  is conservative on  $A$ : we can evaluate conservation properties of other transformations by comparing to Fig. 16.

## GMDD

6, 6493–6568, 2013

### Conservative regridding for ice/atmosphere coupling

E. Fischer et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

## 10.2 Example: $E \rightarrow I$

Figure 17 shows the original field  $f^E$  regridded to  $I$ , via  $f^I = \mathbf{X}\mathbf{M}f^E$  using Z Interpolation for  $\mathbf{M}$ . This plot looks like a smoothed version of Fig. 15. Atmosphere grid cell boundaries are still visible because Z Interpolation is local.

The transformation  $\mathbf{X} : G \rightarrow I$  (Fig. 14) introduces a small amount of non-locality. And although it is conservative over the ice sheet in general, it is not conservative over  $A$ . This can be seen (Fig. 18) by regridding  $f^I$  to  $A$  and comparing with the results of Fig. 16. This plot quantifies the amount of numerical dispersion the simulation will encounter every month when preparing SMB input for the ice model. This dispersion is caused by ice grid cells that overlap more than one atmosphere grid cell.

In most areas, numerical dispersion is low, less than 1%. That is because most atmosphere grid cells are overlapped by many ice grid cells, with only a few lying on an atmosphere grid cell boundary. However, numerical dispersion can be significant for atmosphere grid cells that just nick the edge of the ice sheet, where a high proportion of their overlapping ice grid cells also overlap with a neighboring atmosphere grid cell.

The numerical dispersion encountered here is inherent in the regridding problem itself, rather than our approach to that problem. Short of using an ice grid whose grid cells do not overlap atmosphere grid cells, we see no obvious way to eliminate this issue. However, we do not believe it to be a serious problem: the total area of ice sheet affected by it will be small.

It is important to point out that although  $\mathbf{X}^I$  is not (quite) conservative over  $A$ , it is still conservative over the ice sheet in general. We have verified this numerically in our examples.

## 10.3 Example: $I \rightarrow E$

Since we have not yet developed the surface boundary condition described in Sect. 2.1, we do not have realistic fields on  $I$  to try regridding to  $E$ . We will therefore test  $I \rightarrow E$  at this point using  $f^I = \mathbf{X}\mathbf{M}f^E$  as input.

We test the transformation in two steps. First, we test  $G \rightarrow E$  using  $f^G = \mathbf{M}f^E$  as input. Then, we test  $I \rightarrow E$  using  $f^I = \mathbf{X}\mathbf{M}f^E$  as input. This allows us to evaluate the  $G \rightarrow E$  transformation separately from confounding dispersion factors involved in  $G \rightarrow I$ .

We computed  $f'^E = [G \rightarrow E](f^G)$  where  $f^G = \mathbf{M}f^E$ . Theoretically,  $f'^E$  and  $f^E$  should be equal. The conservation property imposed by the quadratic program held: we found that  $f'^E \equiv^A f^G \equiv^A f^E$  to machine precision. However, we found measurable differences between  $f'^E$  and  $f^E$  (Fig. 19). Although differences are small in most cases, one area has a difference of more than 8 out of 37  $\text{mm day}^{-1}$ , resulting in no more than one digit of precision. Differences tend to be greatest in sparsely populated atmosphere grid cells on the edge of the ice sheet. We conclude that the answer to the quadratic program posed in Sect. 8.5 is poorly constrained. A different numerical solver than the one we used might in theory result in a better match between  $f'^E$  and  $f^E$ .

But this is not a problem in practice: the goal of  $G \rightarrow E$  is to obtain a physically plausible field on  $E$  with the correct conservation properties. The preliminary tests presented in this section are consistent with that goal.

The transformation  $I \rightarrow E$  is constructed by transforming  $I \rightarrow G \rightarrow E$  (Sect. 9). We used this to compute  $f'^E = [I \rightarrow E](f^I)$  where  $f^I = \mathbf{X}\mathbf{M}f^E$ . The conservation property imposed by the quadratic program held: we found that  $f'^E$  was equivalent to  $f^I$  on  $A$ , to machine precision.

However, differences between  $f'^E$  and  $f^E$  are even greater in this case: compare Fig. 20 to Fig. 19. This is because the transformation  $\mathbf{X}: G \rightarrow I$  is not conservative on  $A$ . The end result of transforming  $E \rightarrow G \rightarrow I \rightarrow G \rightarrow E$  will produce an  $f'^E$  that is significantly different from the original  $f^E$ . Differences are greatest for elevation points near an atmosphere grid cell boundary.

## 11 Regridding examples: almost local RM

In Sects. 8.3 and 9, we discussed the trade-offs between using the ice vs. exchange grid for the interpolation grid  $G$ . Having shown in Sect. 10 an example of our

transformations using the exchange grid as  $G$ , we now show how things change if the ice grid is chosen for  $G$  by re-running the above examples. In this case, the **RM** matrix will be mostly local, except for non-locality caused by ice grid cells that intersect more than one atmosphere grid cell. We say that **RM** is *almost local*.

In general, differences in results here vs. in those Sect. 10 are insignificant. However, the choice of  $I$  as interpolation grid *does* change the basis functions used for  $E$ , yielding a system in which the transformation  $E \rightarrow I$  is now conservative over  $A$ .

With a change in  $E \rightarrow I$ , the transformation  $E \rightarrow A$  is also changed. We calculated the difference between  $f^A$  computed using this method, vs.  $f^A$  computed in Fig. 16: this difference is exactly the same as Fig. 18! In practice, this difference does not present a real problem: it is impossible to say which version of  $f^A$  is more “correct.” Both maintain conservation over the ice sheet. As predicted in Sect. 9, grid-related dispersion in  $I \rightarrow E$  is eliminated. Thus, the grid-related “errors” demonstrated in Fig. 20 are eliminated.

Finally, the use of a non-local **RM** matrix requires use of the algorithm described in Sect. 8.1 for  $A \rightarrow E$ . Figure 21 shows a July precipitation field  $p^A$  produced by ModelE, for the same month as the SMB fields above. ModelE currently does not downscale precipitation to sub-grid resolution (Leung and Ghan, 1998): it therefore yields a precipitation field on  $A$ .

We then computed  $p^E = [E \rightarrow A](p^A)$ , which could serve as input to the land surface model. Recall that  $\Lambda p^A$  is the most physically self-consistent value for  $p^E$ , but that we choose a different value for  $p^E$  in order to maintain conservation.

Figure 22 shows the difference between the two,  $p^E - \Lambda p^A$ . Differences are typically in the range  $[-0.1, 0.1]$  (about 2%). They tend to be relatively constant within each atmosphere grid cell, and show no discernible pattern between grid cells. In particular, differences are not related to elevation. Any errors introduced by this scheme will be dwarfed by other precipitation errors in the model.

In this example, we used an L0 ice grid to generate a prototypical almost-local **RM** matrix. We expect similar results when using an L1 ice grid because the non-locality in

## GMDD

6, 6493–6568, 2013

### Conservative regridding for ice/atmosphere coupling

E. Fischer et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



GCMs use a horizontal layout for  $E$  that is not related to  $A$ . In this section, we extend our methods to address that case.

The first problem is a choice of the interpolation grid  $G$ . The idea of locality in **RM** no longer makes sense when  $E^0 \neq A$ . For that reason,  $G = I$  is the right choice,  $\mathbf{X}$  and  $\mathbf{X}'$  will both become identity transformations. The transformation  $E \rightarrow I$  does not need to change, nor does  $I \rightarrow A$ : none of these rely on any specific relationship between  $E^0$  and  $A$ . Similarly,  $E \rightarrow A = \mathbf{RM}$  can be computed as before.

The only other thing that must change is the construction of  $A \rightarrow E$ : the intuitive definition for  $\mathbf{\Lambda}$  (Sect. 8.1) no longer makes sense. Instead, we construct an intuitive regriding operation  $\Phi$  as follows. Given  $f^A$ , first regrid to  $f^{E^0}$  using area-weighted remapping. Then convert  $f^{E^0}$  to  $f^E$  using a “repeat” operator akin to  $\mathbf{\Lambda}$  above. We can now apply the QP-based regrid operator developed in Sect. 8.2, using  $\Phi$  instead of  $\mathbf{\Lambda}$  in Eq. (15).

The use of  $E^0 \neq A$  introduces numerical dispersion into the system. This can be seen by evaluating the locality of the round-trip transformation  $E \rightarrow A \rightarrow E$ . This numerical dispersion is a fundamental consequence of the use of a different underlying grid for the ice surface and atmosphere models.

## 14 Regriding in elevation space

We have tacitly assumed so far that there is one single fixed elevation grid  $E$  with one fixed set of basis functions. This is not the case in a changing climate, since the basis functions used for  $E$  depend on ice elevation and extent (Sect. 7.1). The user might wish to explicitly move elevation points as well – for example, to track a mountain glacier as it moves up or down. When the vector space  $E$  changes, the ice surface model state must be regridded from the old elevation grid to the new elevation grid. We address that issue in this section.

# GMDD

6, 6493–6568, 2013

## Conservative regridding for ice/atmosphere coupling

E. Fischer et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Assume two elevation grids, an “old” grid  $E$  and a “new” grid  $F$ . We wish to conservatively regrid a field  $f^E$  to  $f^F$ . In this case,  $f$  will not be a flux variable, but rather a conserved state variable of the ice surface model: snow depth, water fraction, etc.

This problem can be approached by examining the system of grids and transformations available in Fig. 14 when one has multiple elevation grids (see Fig. 24). By “following the arrows,” the most direct way to transform  $f^E$  to  $f^F$  is to first regrid to the interpolation grid, computing  $f^G = \mathbf{M}^E f^E$ . Then use the procedure in Sect. 8.5 to compute  $f^F$ . Because all these transformations are conservative on the atmosphere grid  $A$ , the end result  $f^F$  will be equivalent to  $f^E$  on  $A$ .

## 14.1 Conserved model state

In previous sections,  $f$  represented fluxes between models, which are conserved. In this section,  $f$  is a *model state* variable of the ice surface model. In order for this re-gridding procedure to be physically meaningful, the model state must be expressed in terms of conserved quantities.

Not all ice surface models are formulated in terms of conserved quantities. In this case, the re-gridding procedures may still be used, as long as the ice flow model can be converted to/from a form that *is* expressed in conserved quantities. For example, an ice surface model might track the temperature, mass and water fraction of the top layer of ice. Temperature is not conserved, so this re-gridding procedure cannot be used directly. However, model state can be converted to enthalpy and mass alone (Aschwan- den et al., 2012). These quantities are conserved, and can be correctly re-gridded with conservative transformations. After re-gridding to the new elevation grid, model state can then be converted back to the original non-conserved parameterization.

## 14.2 When to regrid

Regridding in elevation space is not just required if one changes elevation points, but also *any time* that the transformation  $\mathbf{M}$  for  $E \rightarrow I$  changes. Any time that happens,

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



the regridding described in Sect. 14 should be used. Relevant events that change **M** include:

1. Changes in ice topography: for example, as an ice sheet inflates or deflates due to changing climate. When an ice grid cell changes elevation, the weights in **M** used to compute it will change. This change will either be continuous (as in Z Interpolation), or will jump at certain thresholds (as in Elevation Class Interpolation).
2. Changes in ice sheet extent, as an ice sheet grows or melts. For example, if an ice sheet shrinks, then some ice grid cells will no longer participate in the regridding process, and their associated coefficients in **M** will turn to zero.
3. Changes in the ice grid. Some ice flow models use adaptive grids, for example, to track grounding line retreat (Cornford et al., 2013).
4. Changes in the elevation or number of elevation points.

## 15 GLINT2

Here we describe GLINT2, an open-source implementation of the transformations developed in this paper. GLINT2 is a GCM–ice flow model coupling library whose core function is to compute the five transformations required for a coupled GCM–ice flow model system (Fig. 2). These transformations are computed based on a variety of factors: atmosphere and ice grid geometry, ice topography and extent, and elevation levels chosen for the elevation grid (Fig. 25).

GLINT2 does not just compute these transformations and provide them as a library, it also provides an API for GCMs to use to couple with ice flow models. As a realization of the mediator design pattern (Gamma et al., 1994), GLINT2 is able to shield the GCM from a number of details of the coupling. Although it uses a different codebase, GLINT2 follows the intent of GLINT (GLIMMER Interface; Rutt et al., 2009) to build a coupling

# GMDD

6, 6493–6568, 2013

## Conservative regridding for ice/atmosphere coupling

E. Fischer et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



that GLINT2 does not really fit into. We do not believe this should be a significant barrier to adoption for two reasons:

- In most cases, the elevation grid will have unusual “customized” basis functions (Fig. 13). Centralized GCM regriding schemes are not typically equipped to regrid to/from the elevation grid. Nor are they equipped with the algorithms required for the reverse transformations.
- GLINT2 hides all details of the ice grid from the GCM, communicating with the GCM with fields on the atmosphere and elevation grids. Because all ice grid-related issues are encapsulated in GLINT2, it should not matter to the GCM *how* or even *whether* regriding to/from the ice grid is accomplished. As far as the GCM is concerned, the ice flow model might as well be running on the elevation grid!

Another barrier to adoption is the fact that GLINT2 is packaged as a library. Many GCM projects are reluctant to add additional library dependencies, due to the complications such dependencies introduce in the build process. We do not believe this should be a serious problem because coupling with an ice model already involves the use of outside libraries. With or without GLINT2, the GCM still must manage additional external dependencies.

## 15.2 Availability

GLINT2 is written C++ and designed to couple with GCMs and ice flow models written in Fortran 90/2003, C or C++. It also comes with a Python interface, making it easy to test and plot sample regriding problems before incorporating a coupling strategy into a GCM. GLINT2 source and documentation are available for download (Fischer, 2013).

# GMDD

6, 6493–6568, 2013

## Conservative regriding for ice/atmosphere coupling

E. Fischer et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## 16 Discussion and conclusions

This paper focuses on a system of conservative regridding strategies needed to support two-way coupling between a GCM and an ice flow model in the context of elevation-based downscaling in the GCM. Past efforts at one-way coupling have produced downscaling methods that provide SMB fields that match remarkably well with observations and with regional models. However, these efforts used an inconsistent set of transformations, which would result in non-conservation of mass and energy in a two-way coupled system.

In order to achieve consistency, we began by recognizing the elevation grid as an integral part of the coupling problem, along with the atmosphere and ice grids that had previously been considered. We observed that the ice surface model runs on the elevation grid, and that the downscaling step (from elevation to ice grid) is actually another form of regridding. We have therefore transformed a coupling problem involving two models and two grids into one with three models and three grids.

We analyzed the regridding transformations needed in a typical two-way coupling, and determined that five such transformations are required: three “forward” transformations and two “reverse” transformations. We then set out to develop a consistent implementation for these five transformations, starting with the forward transformations.

We observe that conservation of mass and energy requires consistency between the transformations  $E \rightarrow I$  and  $E \rightarrow A$ . We achieve consistency by allowing the user to choose  $E \rightarrow I$ , and then constructing a transformation for  $E \rightarrow A$  that is consistent with the user’s choice. This is a good approach because it allows the user freedom in choosing a downscaling transformation for  $E \rightarrow I$ . Our transformations imply a set of basis functions for the elevation grid, which we demonstrated in plots.

We then addressed the reverse transformations, using the notions of consistency developed in the previous sections. Problems of underdeterminism and overdeterminism, caused by mismatches in dimensionality between grids, are addressed by using quadratic optimization for these transformations.

# GMDD

6, 6493–6568, 2013

## Conservative regridding for ice/atmosphere coupling

E. Fischer et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

## Conservative regridding for ice/atmosphere coupling

E. Fischer et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

The result is a set of five conservative transformations need to support a two-way coupling of GCMs and ice models. We defined a property of our  $E \rightarrow A$  transformation called *locality*. We showed a number of theoretical and practical benefits if that transformation is local. We implemented all five transformations and demonstrated that they work in practice on realistic input fields.

Although three of those five transformations are well-known in the literature (Ramshaw, 1985; Lipscomb et al., 2013), the reverse transformations are new. Most importantly, we showed how to choose a set of grids, basis functions and regridding schemes so that *all five* regridding operations are conservative. Note that the elevation grid uses a “custom” set of basis functions. This is a significant step beyond past efforts that have focused on conservative regridding to L0 pre-chosen grids (Ramshaw, 1985).

Our downscaling transformation from the elevation to ice grids is taken from previous one-way studies (Lipscomb et al., 2013). Our results “look” almost identical, except for subtle differences in the other related transformations, needed to ensure conservation. We therefore provide practitioners a way to bring already-proven downscaling techniques into a conservative two-way coupled setting.

We went on to package these transformations in GLINT2, a coupling library. As a piece of software, GLINT2 serves as a mediator (Gamma et al., 1994) between the GCM and ice flow model, insulating each from the specific details of the other. This will make it easy to couple GCMs with multiple ice models, as needed to support future research.

## Appendix A

### Mathematical conventions

We use the following mathematical conventions in this paper:

## Conservative regridding for ice/atmosphere coupling

E. Fischer et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

1. A “grid” or “parameterization” is a set of basis functions that may be linearly combined to produce functions over a 2-D domain. Grids are denoted by non-bold capital letters:  $A$  for the atmosphere grid,  $I$  for the ice grid,  $E$  for the elevation grid and  $G$  for an interpolation grid. Basis functions are denoted by non-bold lower case letters:  $a_i(x, y)$  is a basis function for  $A$ . At times, the set of basis functions may be represented as a bold-face vector:  $\mathbf{a}(x, y) = [a_1(x, y), \dots, a_n(x, y)]$
2. Subscripts are used to indicate elements of a vector. If  $\mathbf{f}$  is a vector, then  $f_i$  is the value of the  $i$ th index of  $\mathbf{f}$ .
3. If  $A$  is an L0-parameterized grid and  $A_i$  one of the grid cells, then  $|A_i|$  is the area of that grid cell.
4. Vector values are indicated in boldface, scalars in non-bold.
5. Vectors are used to construct a 2-D function (“field”) within the context of a particular grid. For example, suppose that the atmosphere grid  $A$  uses the basis functions  $\mathbf{a}(x, y) = [a_1(x, y) \dots a_n(x, y)]$ , and we have an  $n$ -dimensional vector  $\mathbf{f}^A$  with components  $f_i^A$ . That vector represents the function:

$$f^A(x, y) = \mathbf{f}^A \cdot \mathbf{a}(x, y) \quad (\text{A1})$$

6. Since this paper is about regridding, we need to talk about the “same” field in different grids. Superscripts are used for this: if the vector  $\mathbf{f}^I$  denotes the vector representing a field on the ice grid, then  $\mathbf{f}^A$  denotes a vector representing the “same” field on the atmosphere grid.
7. Similarly,  $f^A(x, y)$  denotes the function implied by the vector  $\mathbf{f}^A$  and the vector of basis functions  $\mathbf{a}(x, y)$ .
8. We use arrows to talk about transformations between grids, which transform fields on one grid to fields on another. Since fields on a grid are represented by vectors,

the transformations are functions from one vector space to another. For example,  $E \rightarrow A$  can be read “the transformation from the elevation grid to the atmosphere grid.” Many of these transformations linear, and can be represented by matrices.

9. If a transformation is non-linear, we use functional notation with square brackets to denote an application of that transformation:  $f^E = [A \rightarrow E](f^A)$ .
10. We use Einstein summation notation at times.

## Appendix B

### Integration of L0 grid

Suppose we have a region  $B$  and an L0-parameterized grid  $G$  with grid cells  $G_1 \dots G_n$ . The basis function  $g_i(x, y)$  is equal to 1 inside of grid cell  $G_i$  and zero elsewhere. We wish to compute:

$$\int_B g_i(x, y) dA = |B \cap G_i| \quad (\text{B1})$$

This is simply equal to the area of intersection of  $B$  and  $G_i$ .

If we wish to regrid to an L0 grid  $A$ , we must do the above computation multiple times, setting  $B$  to every grid cell  $A_j \in A$ . This creates an *overlap matrix*  $\mathbf{L}$  where  $L_{ij}$  is equal to the overlap between source grid cell  $G_i$  and destination grid cell  $A_j$ .

The overlap matrix is directly related to the exchange grid between  $G$  and  $A$ : every non-zero element of  $L_{ij}$  is equal to the area of one grid cell in the exchange grid. The exchange grid may be computed as described in Appendix D. The Surveyor’s Formula (Braden, 1986), a special case of Green’s Theorem, can then be used to compute the overlap matrix from the exchange grid.

## Appendix C

### Integration of L1 grid

Appendix B shows how to integrate over areas on an L0 ice grid. But if an L1 finite element ice flow model is used, one will need to do this over an L1 ice grid. Here, we show how to integrate L1 finite element basis functions over an arbitrary area  $B$ .

An L1 finite element mesh is made up of triangular elements, where the value of a function is defined at the vertices of the triangles. Values inside each element are interpolated based on the values at the vertices.

Each basis function  $N_j(x, y)$  in a finite element mesh corresponds to a vertex (Zienkiewicz et al., 2013). This function is non-zero only in the triangular elements in which the vertex participates. We define a sub-basis function  $N_{ij}(x, y)$  to be equal to  $N_j(x, y)$  within triangular element  $j$ , and zero elsewhere. Thus we define:

$$N_i(x, y) = \sum_j N_{ij}(x, y) \quad (C1)$$

Within its element, a sub-basis function takes the shape of a plane, having the functional form:

$$N_{ij}(x, y) = a + bx + cy \quad (C2)$$

The coefficients  $a$ ,  $b$  and  $c$  are functions of the locations of the vertices, NOT of the values assigned to those vertices. Using the techniques from Appendix D, one can compute the polygon  $\beta_j$  as the intersection between element  $j$  and region  $B$ . Green's Theorem may then be used to compute:

$$\int_{\beta_j} N_{ij}(x, y) dA \quad (C3)$$

## GMDD

6, 6493–6568, 2013

### Conservative regridding for ice/atmosphere coupling

E. Fischer et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion







## Conservative regridding for ice/atmosphere coupling

E. Fischer et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

independent grids, in: *Parallel Computational Fluid Dynamics: Theory and Applications, Proceedings of the 2005 International Conference on Parallel Computational Fluid Dynamics, College Park, MD, USA, 24–27 May*, edited by: Deane, A., Brenner, G., Ecer, A., Emerson, D., McDonough, J., Periaux, J., Satofuka, N., and Tromeur-Dervout, D., Elsevier, 2006. 6506, 6537

Best, M. J., Baljaars, A., Polcher, J., and Viterbo, P.: A proposed structure for coupling tiled surfaces with the planetary boundary layer, *J. Hydrometeorol.*, 5, 1271–1278, doi:10.1175/JHM-382.1, 2004. 6518

Bindschadler, R., Nowicki, S., Abe-Ouchi, A., Aschwanden, A., Choi, H., Fastook, J., Granzow, G., Greve, R., Gutowski, G., Herzfeld, U., Jackson, C., Johnson, J., Khroulev, C., Levermann, A., Lipscomb, W., Martin, M., Morlighem, M., Parizek, B., Pollard, D., Price, S., Ren, D., Saito, F., Sato, T., Seddik, H., Seroussi, H., Takahashi, K., Walker, R., and Wang, W.: Ice-sheet model sensitivities to environmental forcing and their use in projecting future sea level (the SeaRISE project), *J. Glaciol.*, 59, 195–224, doi:10.3189/2013JoG12J125, 2013. 6495, 6499, 6507, 6522

Box, J. E., Cappelen, J., Chen, C., Decker, D., Fettweis, X., Mote, T., Tedesco, M., van de Wal, R. S. W., and Wahr, J.: Greenland Ice Sheet [in Arctic Report Card 2012], available at: [http://www.arctic.noaa.gov/reportcard/greenland\\_ice\\_sheet.html](http://www.arctic.noaa.gov/reportcard/greenland_ice_sheet.html) (last access: 3 December 2013), 2013. 6503

Braden, B.: The surveyor's area formula, *Coll. Math. J.*, 17, 326–337, doi:10.2307/2686282, 1986. 6535

Braithwaite, R.: On glacier energy balance, ablation, and air temperature, *J. Glaciol.*, 27, 381–391, 1981. 6503

Bueler, E. and Brown, J.: Shallow shelf approximation as a sliding law in a thermomechanically coupled ice sheet model, *J. Geophys. Res.*, 114, F03008, doi:10.1029/2008JF001179, 2009. 6495, 6500

Cgal: Computational Geometry Algorithms Library, available at: <http://www.cgal.org>, last access: 3 December 2013. 6537

Chin, F. and Wang, C. A.: Optimal algorithms for the intersection and the minimum distance problems between planar polygons, *IEEE T. Comput.*, 32, 1203, doi:10.1109/TC.1983.1676186, 1983. 6537

Collins, W. J., Bellouin, N., Doutriaux-Boucher, M., Gedney, N., Halloran, P., Hinton, T., Hughes, J., Jones, C. D., Joshi, M., Liddicoat, S., Martin, G., O'Connor, F., Rae, J., Senior, C.,

## Conservative regridding for ice/atmosphere coupling

E. Fischer et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Sitch, S., Totterdell, I., Wiltshire, A., and Woodward, S.: Development and evaluation of an Earth-System model – HadGEM2, *Geosci. Model Dev.*, 4, 1051–1075, doi:10.5194/gmd-4-1051-2011, 2011. 6495

Cornford, S. L., Martin, D. F., Graves, D. T., Ranken, D. F., Le Brocq, A. M., Gladstone, R. M., Payne, A. J., Ng, E. G., and Lipscomb, W. H.: Adaptive mesh, finite volume modeling of marine ice sheets, *J. Comput. Phys.*, 232, 529–549, doi:10.1016/j.jcp.2012.08.037, 2013. 6495, 6529

Dukowicz, J. K. and Kodis, J. W.: Accurate conservative remapping (rezoning) for arbitrary Lagrangian–Eulerian computations, *SIAM J. Sci. Stat. Comp.*, 8, 305–321, doi:10.1137/0908037, 1987. 6538

Fischer, R.: GLINT2, available at: <http://citibob.github.io/glint2> (last access: 3 December 2013), 2013. 6531

Gamma, E., Helm, R., Johnson, R., and Vlissides, J.: *Design Patterns: Elements of Reusable Object-Oriented Software*, 1st Edn., Addison-Wesley Professional, 1994. 6529, 6533

Goelzer, H., Huybrechts, P., Fürst, J. J., Nick, F. M., Andersen, M. L., Edwards, T. L., Fettweis, X., Payne, A. J., and Shannon, S.: Sensitivity of Greenland ice sheet projections to model formulations, *J. Glaciol.*, 59, 733–749, doi:10.3189/2013JoG12J182, 2013. 6495

Gould, N., Orban, D., and Toint, P.: GALAHAD, a library of thread-safe Fortran 90 packages for large-scale nonlinear optimization, *ACM T. Math. Software*, 29, 353–372, doi:10.1145/962437.962438, 2003. 6517

Greve, R.: On the response of the Greenland Ice Sheet to greenhouse climate change, *Climatic Change*, 46, 289–303, doi:10.1023/A:1005647226590, 2000. 6495

Guttman, A.: R-trees – a dynamic index structure for spatial searching, in: *SIGMOD '84, Proceedings of the 1984 ACM SIGMOD international conference on management of data*, 14, 47–57, doi:10.1145/971697.602266, 1984. 6538

Heath, T. and Archimedes: *The Works of Archimedes*, C. J. Clay and Sons, Cambridge University Press Warehouse, available at: <https://archive.org/details/worksofarchimede029517mbp> (last access: 3 December 2013), 1897. 6508

Hurrell, J. W., Holland, M. M., Gent, P. R., Ghan, S., Kay, J. E., Kushner, P. J., Lamarque, J. F., Large, W. G., Lawrence, D., Lindsay, K., Lipscomb, W. H., Long, M. C., Mahowald, N., Marsh, D. R., Neale, R. B., Rasch, P., Vavrus, S., Vertenstein, M., Bader, D., Collins, W. D., Hack, J. J., Kiehl, J., and Marshall, S.: The Community Earth System Model: a framework

for collaborative research, B. Am. Meteorol. Soc., 94, 1339–1360, doi:10.1175/BAMS-D-12-00121.1, 2013. 6495

Huybrechts, P.: The present evolution of the Greenland ice sheet: an assessment by modelling, Global Planet. Change, 9, 39–51, doi:10.1016/0921-8181(94)90006-X, 1994. 6495

Jones, P. W.: First- and second-order conservative remapping schemes for grids in spherical coordinates, Mon. Weather Rev., 127, 2204–2210, doi:10.1175/1520-0493(1999)127<2204:FASOCR>2.0.CO;2, 1999. 6538

Larour, E., Seroussi, H., Morlighem, M., and Rignot, E.: Continental scale, high order, high spatial resolution, ice sheet modeling using the Ice Sheet System Model (ISSM), J. Geophys. Res., 117, F01022, doi:10.1029/2011JF002140, 2012. 6495

Lauritzen, P. H. and Nair, R. D.: Monotone and Conservative Cascade Remapping between Spherical Grids (CaRS): Regular Latitude–Longitude and Cubed-Sphere Grids, Mon. Weather Rev., 136, 1416, doi:10.1175/2007MWR2181.1, 2008. 6508

Leung, L. R. and Ghan, S. J.: Parameterizing subgrid orographic precipitation and surface cover in climate models, Mon. Weather Rev., 126, 3271–3291, doi:10.1175/1520-0493(1998)126<3271:PSOPAS>2.0.CO;2, 1998. 6496, 6511, 6522, 6525

Lipscomb, W. H., Fyke, J. G., Vizcaíno, M., Sacks, W. J., Wolfe, J., Vertenstein, M., Craig, A., Kluzek, E., and Lawrence, D. M.: Implementation and initial evaluation of the Glimmer Community Ice Sheet Model in the Community Earth System Model, J. Climate, 26, 7352–7371, doi:10.1175/JCLI-D-12-00557.1, 2013. 6495, 6496, 6502, 6510, 6533, 6552

Nowicki, S., Bindschadler, R., Abe-Ouchi, A., Ashwanden, A., Bueler, E., Choi, H., Fastook, J., Granzow, G., Greve, R., Gutowski, G., Herzfeld, U., Jackson, C., Johnson, J., Kroulev, C., Larour, E., Levermann, A., Lipsomb, W., Martin, M., Morlighem, M., Parizek, B., Pollard, D., Price, S., Rignot, E., Ren, D., Saito, F., Sato, T., Seddik, H., Seroussi, H., Takahashi, K., Walker, R., and Wang, W.: Insights into spatial sensitivities of ice mass response to environmental change from the SeaRISE ice sheet modeling project I: Antarctica, J. Geophys. Res., 118, 1002–1024, doi:10.1002/jgrf.20081, 2013a. 6495

Nowicki, S., Bindschadler, R., Abe-Ouchi, A., Ashwanden, A., Bueler, E., Choi, H., Fastook, J., Granzow, G., Greve, R., Gutowski, G., Herzfeld, U., Jackson, C., Johnson, J., Kroulev, C., Larour, E., Levermann, A., Lipsomb, W., Martin, M., Morlighem, M., Parizek, B., Pollard, D., Price, S., Rignot, E., Ren, D., Saito, F., Sato, T., Seddik, H., Seroussi, H., Takahashi, K., Walker, R., and Wang, W.: Insights into spatial sensitivities of ice mass response to envi-

**Conservative  
regridding for  
ice/atmosphere  
coupling**

E. Fischer et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

## Conservative regridding for ice/atmosphere coupling

E. Fischer et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

ronmental change from the SeaRISE ice sheet modeling project II: Greenland, *J. Geophys. Res.-Earth*, 118, 1, doi:10.1002/jgrf.20076, 2013b. 6495

Pollard, D. and DeConto, R. M.: Description of a hybrid ice sheet–shelf model, and application to Antarctica, *Geosci. Model Dev.*, 5, 1273–1295, doi:10.5194/gmd-5-1273-2012, 2012. 6508

5 Qu, X. and Hall, A.: Assessing snow albedo feedback in simulated climate change, *J. Climate*, 19, 2617–2630, doi:10.1175/JCLI3750.1, 2006. 6495

Ramshaw, J. D.: Conservative rezoning algorithm for generalized two-dimensional meshes, *J. Comput. Phys.*, 59, 193–199, doi:10.1016/0021-9991(85)90141-X, 1985. 6506, 6533, 6538

10 Ridley, J. K., Huybrechts, P., Gregory, J. M., and Lowe, J. A.: Elimination of the Greenland Ice Sheet in a high CO<sub>2</sub> climate, *J. Climate*, 18, 3409–3427, doi:10.1175/JCLI3482.1, 2005. 6495

Ridley, J., Gregory, J. M., Huybrechts, P., and Lowe, J.: Thresholds for irreversible decline of the Greenland ice sheet, *Clim. Dynam.*, 35, 1059–1057, doi:10.1007/s00382-009-0646-0, 2010. 6495

15 Rutt, I. C., Hagdorn, M., Hulton, N. R. J., and Payne, A. J.: The Glimmer community ice sheet model, *J. Geophys. Res.*, 114, F02004, doi:10.1029/2008JF001015, 2009. 6495, 6529

Schmidt, G. A., Bitz, C. M., Mikolajewicz, U., and Tremblay, L. B.: Ice–ocean boundary conditions for coupled models, *Ocean Model.*, 7, 59–74, doi:10.1016/S1463-5003(03)00030-1, 2004. 6498

20 Schmidt, G. A., Ruedy, R., Hansen, J. E., Aleinov, I., Bell, N., Bauer, M., Bauer, S., Cairns, B., Canuto, V., Cheng, Y., Del Genio, A., Faluvegi, G., Friend, A. D., Hall, T. M., Hu, Y., Kelley, M., Kiang, N. Y., Koch, D., Lacis, A. A., Lerner, J., Lo, K. K., Miller, R. L., Nazarenko, L., Oinas, V., Perlwitz, J. P., Perlwitz, J., Rind, D., Romanou, A., Russell, G. L., Sato, M., Shindell, D. T., Stone, P. H., Sun, S., Tausnev, N., Thresher, D., and Yao, M.-S.: Present day atmospheric simulations using GISS ModelE: comparison to in-situ, satellite and reanalysis data, *J. Climate*, 19, 153–192, doi:10.1175/JCLI3612.1, 2006. 6495, 6499, 6500, 6522

Snyder, J. P.: Map projections – a working manual, available at: <http://pubs.er.usgs.gov/publication/pp1395> (last access: 3 December 2013), 1987. 6507

30 Solomon, S., Qin, D., Manning, M., Chen, Z., Marquis, M., Averyt, K. B., Tignor, M., and Miller, H. L.: IPCC: Climate Change (2007), The Physical Sciences Basis, Contribution of Working Group I to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change, Cambridge University Press, 2007. 6494

## Conservative regridding for ice/atmosphere coupling

E. Fischer et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close


Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Stone, E. J., Lunt, D. J., Rutt, I. C., and Hanna, E.: Investigating the sensitivity of numerical model simulations of the modern state of the Greenland ice-sheet and its future response to climate change, *The Cryosphere*, 4, 397–417, doi:10.5194/tc-4-397-2010, 2010. 6495
- van den Broeke, M., Smeets, P., Ettema, J., and Munneke, P. K.: Surface radiation balance in the ablation zone of the west Greenland ice sheet, *J. Geophys. Res.*, 113, D13105, doi:10.1029/2007JD009283, 2008. 6522
- Vizcaino, M., Lipscomb, W., Sacks, W., van Angelen, J. H., Wouters, B., and van den Broeke, M. R.: Greenland surface mass balance as simulated by the Community Earth System Model. Part I: model validation and 1850-2005 results, *J. Climate*, 26, 7793–7812, doi:10.1175/JCLI-D-12-00615.1, 2013. 6496
- Zagorodnov, V., Nagornov, O., and Thompson, L. G.: Influence of air temperature on a glacier's active-layer temperature, *Ann. Glaciol.*, 43, 285–291, doi:10.3189/172756406781812203, 2006. 6499
- Zienkiewicz, O. C., Taylor, R. L., and Nithiarasu, P.: *The Finite Element Method for Fluid Dynamics*, 7th Edn., Butterworth–Heinemann, 2013. 6504, 6512, 6536



**Fig. 1.** Configuration of the three models, and the separate spatial domains they occupy. The atmosphere model (A) is coupled to the ice surface model (E), which tracks the top few meters of ice. The ice surface model (E) is then coupled to the dynamic ice flow model (I). The dynamic ice flow model, operating at long time steps, is insulated from high-frequency surface processes in the ice/atmosphere interaction.

Conservative  
regridding for  
ice/atmosphere  
coupling

E. Fischer et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close


Full Screen / Esc

Printer-friendly Version

Interactive Discussion

## Conservative regridding for ice/atmosphere coupling

E. Fischer et al.



**Fig. 2.** Data flow for the coupling between atmosphere, ice surface and dynamic ice flow models. Outputs from one model are fed as inputs into another. Since the three models run on different grids, regridding is required at each step.

Title Page

Abstract Introduction

Conclusions References

Tables Figures

◀ ▶

◀ ▶

Back Close


Full Screen / Esc

Printer-friendly Version

Interactive Discussion

## Conservative regridding for ice/atmosphere coupling




E. Fischer et al.



**Fig. 3.** Interpolated surface mass balance (SMB) function within one atmosphere grid cell. These values come from one month (July) of a run of GISS ModelE with elevation points. The vertical dashed line indicates the mean elevation of the grid cell, “seen” by the atmosphere. Dots and squares represent the results of extrapolated SMB computations at other elevations: dots represent elevations found within the grid cell, whereas squares represent elevations outside the cell’s range. SMB values at intermediate elevations are interpolated.


**Conservative  
regridding for  
ice/atmosphere  
coupling**

E. Fischer et al.

**Atmosphere  
Grid****Ice  
Grid****Exchange  
Grid**

**Fig. 4.** An exchange grid, obtained by overlapping a sample atmosphere grid and ice grid (not to scale). Each resulting grid cell, irregularly shaped, overlaps with exactly one atmosphere grid cell, and exactly one ice grid cell.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)



**Fig. 5.** An example of geometric error, in which grid cells change size due to polygonal approximation. This Lambert Equal Area Projection shows a set of latitude/longitude grid cells, a kind commonly used in atmosphere models. Solid blue lines show polygonal approximations, while dotted red shows the actual grid cells on the sphere. Note that all grid cells (in this local map) shrink when approximated.

Conservative  
regridding for  
ice/atmosphere  
coupling

E. Fischer et al.

Title Page

Abstract Introduction

Conclusions References

Tables Figures

⏪ ⏩

◀ ▶


Back Close

Full Screen / Esc


Printer-friendly Version

Interactive Discussion



[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

**Fig. 6.** Geometric error in atmosphere grid cells, as a function of the number of line segments used for each side of the grid cell ( $n$ ). Experimental result.



**Fig. 7.** Results of Z Interpolation, to develop a downscaled field on the ice grid from an elevation point field. Note the visible discontinuities at atmosphere grid cell boundaries. This figure is for demonstration purposes only. See Sects. 10 and 11 for thorough regridding examples.

# GMDD

6, 6493–6568, 2013

## Conservative regridding for ice/atmosphere coupling

E. Fischer et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back


Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion





**Fig. 8.** Setup for bilinear interpolation. The value in an ice grid cell (square) will be a sum of the values at the four nearest atmosphere grid cell centers, weighted by  $\Delta$ -longitude and  $\Delta$ -latitude along the axes.

**Conservative  
regridding for  
ice/atmosphere  
coupling**

E. Fischer et al.

Title Page

Abstract Introduction

Conclusions References

Tables Figures

⏪ ⏩

◀ ▶


Back Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion





**Fig. 9.** Results of Bilinear Interpolation, to develop a downscaled field on the ice grid from an elevation point field (Lipscomb et al., 2013). Compare to Fig. 7. The use of horizontal and well as vertical interpolation eliminates discontinuities as compared to Z Interpolation. However, total SMB is changed, particularly for localized events such as snowstorms.

**Conservative  
regridding for  
ice/atmosphere  
coupling**

E. Fischer et al.

Title Page

Abstract	Introduction
Conclusions	References
Tables	Figures


⏪	⏩
◀	▶
Back	Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion





**Fig. 10.** Visualization of grid used in elevation class schemes. Atmosphere grid cells are “carved up” according to topography of the ice sheet. Grid cells are shown with checkerboard coloring for visibility.

# GMDD


6, 6493–6568, 2013

## Conservative regridding for ice/atmosphere coupling

E. Fischer et al.

<a href="#">Title Page</a>	
<a href="#">Abstract</a>	<a href="#">Introduction</a>
<a href="#">Conclusions</a>	<a href="#">References</a>
<a href="#">Tables</a>	<a href="#">Figures</a>
<a href="#">⏪</a>	<a href="#">⏩</a>
<a href="#">◀</a>	<a href="#">▶</a>
<a href="#">Back</a>	<a href="#">Close</a>
<a href="#">Full Screen / Esc</a>	
<a href="#">Printer-friendly Version</a>	
<a href="#">Interactive Discussion</a>	






**Fig. 11.** Interpolated SMB function within one atmosphere grid cell, when using elevation classes. Compare to Fig. 3.

## Conservative regridding for ice/atmosphere coupling

E. Fischer et al.



**Fig. 12.** The transformation from elevation grid to atmosphere grid is derived by first interpolating to the ice grid (represented by matrix  $M$ ), and then using a conservative area-weighted remapping step to the atmosphere grid (matrix  $R$ ). The two steps may be combined by computing that matrix product  $RM$ . This construction ensures that a quantity computed on the elevation grid will have the same total mass when regrided to the ice grid or atmosphere grid. The two transformations from the elevation grid to the ice and atmosphere grids are said to be *consistent*.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close


Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Conservative  
regridding for  
ice/atmosphere  
coupling


E. Fischer et al.




**Fig. 13.** Basis functions for the elevation grid  $E$ , constructed using 20 elevation points and Z Interpolation. The exchange grid was used for the interpolation grid. Basis functions for elevation point within one atmosphere grid cell at 950 m, 1150 m and 1350 m are shown. Contour lines show elevation point levels. Note that basis functions overlap and are not orthogonal. Because of the Z Interpolation, each basis function has maximum value at its corresponding elevation, but it has a non-zero support up to one elevation point away.

## Conservative regridding for ice/atmosphere coupling

E. Fischer et al.



**Fig. 14.** The five grids used in the coupling problem, and transformations between them. The (linear) interpolation step  $M$ , from  $E$  to  $G$ , is chosen by the user. The (linear) transformation  $R$ , from  $G$  to  $A$ , is an area-weighted remapping step. These two fully constrain the transformation from  $E$  to  $A$  as  $RM$ . If  $I$  is L0-parameterized, then  $G$  is the exchange grid between  $A$  and  $I$ , and  $X$  and  $X'$  are area-weighted remapping transformations. If  $I$  is L1-parameterized, then  $G$  is equivalent to  $I$ , making  $X$  and  $X'$  the identity.  $P$  is a diagonal rescaling operation between  $A'$  and  $A$ . Reverse transformations required by the coupled system are shown as dotted lines.



**Fig. 15.** July SMB computed by ModelE on the elevation grid  $E$ . For plotting purposes, each elevation point is assigned to a nearby region of similar elevation; the value of the elevation point is then plotted in its corresponding region. Bold numbers on the color scale indicate the extreme values of the plot. Elevation contours are plotted.

**Conservative  
regridding for  
ice/atmosphere  
coupling**

E. Fischer et al.

Title Page

Abstract Introduction

Conclusions References

Tables Figures

◀ ▶

◀ ▶


Back Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion






**Fig. 16.** July SMB computed by ModelE on the elevation grid *E*, then regrided to the atmosphere grid *A* via the exchange grid. The same color scale is used as in Fig. 15

## Conservative regridding for ice/atmosphere coupling

E. Fischer et al.

Title Page	
Abstract	Introduction
Conclusions	References
Tables	Figures
◀	▶
◀	▶
Back	Close
Full Screen / Esc	
Printer-friendly Version	
Interactive Discussion	





**Fig. 17.** July SMB computed by ModelE on the elevation grid  $E$ , then regridded to the ice grid / using Z Interpolation.

# GMDD

6, 6493–6568, 2013

## Conservative regridding for ice/atmosphere coupling

E. Fischer et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back


Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion





**Fig. 18.** Difference plot. July SMB on elevation grid  $E$  was regridded to the atmosphere grid  $A$  via the ice grid  $I$ . Plot shows the difference between this result and Fig. 16, which regridded via the exchange grid. Differences are due to ice grid cells that overlap more than one atmosphere grid cell.

Conservative  
regridding for  
ice/atmosphere  
coupling

E. Fischer et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back


Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion





**Fig. 20.** July SMB on elevation grid  $E$  was regridded to the ice grid. The reverse transformation was then used to recover the original SMB on  $E$ . Plot shows the difference between the result and the original. Differences are due mainly to ice grid cells that overlap more than one atmosphere grid cell, making it impossible to recover the exact original SMB. The same color scale is used as in Fig. 19.

**Conservative  
regridding for  
ice/atmosphere  
coupling**

E. Fischer et al.

Title Page

Abstract Introduction

Conclusions References

Tables Figures

◀ ▶

◀ ▶


Back Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion





**Fig. 21.** July Precipitation field compute by ModelE. Precipitation is computed on the atmosphere grid, without any additional downscaling algorithms to achieve sub-grid resolution.

## Conservative regridding for ice/atmosphere coupling

E. Fischer et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back


Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion





**Fig. 22.** July precipitation (Fig. 21) was regridder to the elevation grid  $E$  using the  $A \rightarrow E$  reverse transformation where the matrix for  $E \rightarrow A$  is almost local. The minor non-locality of  $E \rightarrow A$  causes the result on  $E$  to be somewhat different from what one would get if values on  $A$  were simply “repeated” for every elevation point in each atmosphere grid cell. Those differences on  $E$  are plotted here.

**Conservative  
regridding for  
ice/atmosphere  
coupling**

E. Fischer et al.

Title Page

Abstract Introduction

Conclusions References

Tables Figures

◀ ▶

◀ ▶


Back Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion





**Fig. 23.** July precipitation (Fig. 21), regrided to the elevation grid  $E$  using the  $A \rightarrow E$  reverse transformation, where the matrix for  $E \rightarrow A$  is constructed using (non-local) bilinear interpolation. Note the unphysical artifacts (negative precipitation) for some elevation points! Compare to Figs. 21 and 22.

Conservative  
regridding for  
ice/atmosphere  
coupling

E. Fischer et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶


Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion




**Fig. 24.** Grid System used to regrid ice surface model state when the elevation grid changes: *E* is the old elevation grid, and *F* is the new one.

Title Page	
Abstract	Introduction
Conclusions	References
Tables	Figures
◀	▶
◀	▶
Back	Close
Full Screen / Esc	
Printer-friendly Version	
Interactive Discussion	



## Conservative regridding for ice/atmosphere coupling

E. Fischer et al.



**Fig. 25.** The GLINT2 process flow. GLINT2 produces regridding operations based on a variety of factors: atmosphere and ice grid geometry, ice topography and extent, and elevation levels chosen for the elevation grid.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion