



# ***Interactive comment on* “The potential of an observational data set for calibration of a computationally expensive computer model” by D. J. McNeall et al.**

**D. J. McNeall et al.**

doug.mcneall@metoffice.gov.uk

Received and published: 4 September 2013

## **Review 2 I. Vernon**

**Comment:** 1. Page 2379, lines 27–28 “we let the model output  $y$  take the place of a theoretical observational dataset  $z$  in our analysis”. Using  $y$  in place of  $z$  is correct for the case of zero model discrepancy  $\delta$  and zero observation errors  $e$ , but when  $\delta$  and  $e$  are not assumed zero then this represents an approximation to the full calculation (a good approximation though!). The full calculation would involve simulating from  $\delta$  and  $e$  to get  $z$ , and doing this many times for each ensemble value of  $y$  considered, applying the implausibility constraints for each simulated value of  $z$ , to get

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Discussion Paper



a distribution of space cut out for each  $y$  value. (Incidentally this process is described in the presentation: Vernon, I., Goldstein, M., Liu, J., Lindsey, K. "Emulation and Efficient History Matching of Stochastic Systems Biology Models" Presentation at UCM 2012. <http://www.mucm.ac.uk/UCM2012/TalksPosters.html>) Now the authors mention several times that their analysis will provide "a maximum bound for our ability to constrain the model inputs" e.g. in the abstract, p2370 lines 7-8. Due to the above, this claim should perhaps be tempered as it relies upon the above approximation to the full calculation. What the authors calculate is perhaps a maximum bound on the expected space cutout in the full calculation, but this depends on the distributional assumptions. For example, the calculation may say 50

**Response:** This comment is well made. We have tempered the statement in the abstract as requested, pointing out that it relies on some assumptions. We have added a paragraph outlining the most important assumptions in section 2.4 Using an ensemble to find an upper bound of potential constraint. We have explained that the ensemble must span the full distribution of model output, plus model discrepancy and observational uncertainty, as pointed out in the comment above. In that case, the combination of simulator output, plus discrepancy, plus observational uncertainty, might plausibly be represented by any of the ensemble members, for which we could then calculate the potential constraint. This point is also clarified in the discussion (ref. comment 11).

**Comment:** 2. Page 2372: in the description of the two approaches of Bayesian calibration and history matching, it is useful to mention the main differences between these approaches. Bayesian calibration assumes a "best input  $x^*$ ", uses a prior for  $x^*$  and updates this to a posterior. History matching does not assume the existence of a best input  $x^*$  and instead just tests points in input space to determine if they are consistent with the specified model and associated uncertainties (observation error, model discrepancy). This is an important difference because in the Bayesian calibration case, if you have explored some parts of the input space (by doing runs of the model there) and have found those parts to be bad in terms of giving outputs far from observed data,

[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)[Discussion Paper](#)

you automatically think other as yet unexplored parts of the input space are good. This is because you have assumed one single  $x^*$ , and it must lie somewhere! This is avoid in history matching, where your assessment of one part of the space does not affect ther parts of the space. A full description of the benefits of history matching and its application to a large model of galaxy formation, along with a discussion comparing it to Bayesian calibration, can be found in: Vernon, I, Goldstein, M. Bower, R. G. 2010. "Galaxy Formation: a Bayesian Uncertainty Analysis". Bayesian Analysis 05(04): 619–670 (with discussion) which it may be suitable for the authors to cite.

**Response:** The benefits of history matching in comparison to probabilistic calibration are outlined in more detail in the next section: “ 2.1 solving the inverse problem”, and the suggested reference is included.

**Comment:** 3. Page 2374, line 19: "where  $e$  represents systematic errors or biases in the observations". Do these errors really have to be "systematic" or be "biases". Surely  $e$  is a random variable just representing measurement error, which can be systematic or otherwise! Technically, if there were biased, this should change the implausibility measures too. Perhaps just saying "where  $e$  represents measurement errors in the observations" would be clearer?

**Response:** Changed as requested.

**Comment:** 4. Page 2376, lines 2–3 The authors use Pukelsheim’s 3 sigma rule, a very powerful and general result, but it only holds true for unimodal distributions. Unimodality of the distribution underlying the implausibility measure is a reasonable assumption here, but should be stated.

**Response:** Edited to make clear that this holds only for unimodal distributions.

**Comment:** 5. Page 2376, lines 4–7 the authors combine implausibilities from different outputs by maximising over the implausibilities. This is a sensible choice, however, it is worth noting that there are other more complex implausibility measures available such

[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)[Discussion Paper](#)

as the multivariate measure described in (Vernon et. al. (2010) (ref given above)).

**Response:** Added the reference as requested, along with a brief explanation of the reason we didn't use the more complex metric.

**Comment:** 6. Page 2376. Defining a suitable metric is no easy task. The authors introduce the marginal range of NROY space. This is a fairly sensible metric, however, the volume of NROY space is, in my opinion, a far superior metric as it is a clearly defined object and represents how much we have learned about the input space, where as the marginal range can be misleading in several situations. In this example, where 3 outputs are used to constrain a 5 dimensional input space, we expect the NROY space to be in the form of hyper-surfaces (specifically 2 dimensional) due to the remaining degrees of freedom (as the authors have noticed at a later point). These hyper-surfaces may stretch across the input space, ensuring that the marginal range is quite misleading. Could the authors, if they agree, perhaps mention that the volume metric is superior or safer in many cases?

**Response:** The paragraph was edited to more clearly indicate that we consider the Volume metric superior, while still outlining situations where the Range metric might be useful.

**Comment:** 7. Page 2376, lines 24–25 "We can define a volume  $V$  of "not implausible" input parameter space, or alternatively that input space "Not Ruled Out Yet" – as the region bounded by the convex hull where  $I < 3$ ". No, the volume  $V$  is not the region bounded by the convex hull, it is simply the region defined by  $I < 3$ , whatever its shape or geometry. The Monte Carlo estimate for this volume  $V$  does not need any complex hull results: it is a direct estimate of  $V$ .

**Response:** Text edited to remove the reference to the convex hull.

**Comment:** 8. Page 2377, line 22, " or the least implausible point". When an implausibility measure gives a high value, it means we can rule out that input, but when it give

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Discussion Paper



a low value it simply means we are still not sure about the input at this stage (hence Not Ruled Out Yet). Further runs of the simulator (waves), or more detailed statistical modelling (using say more advanced implausibility measures) may subsequently rule out this input. Hence the least implausible point has no real importance and should be treated with caution (indeed, this is why implausibility measures are much easier to use than full bayesian posteriors as the former just models where the bad inputs are, where the latter tries to model the much more complex question of where all the good inputs are!). Perhaps the authors could just remove the phrase "or the least implausible point".

**Response:** This section was identified as distracting by referee 1, and removed from the text.

**Comment:** 9. Page 2378, line 12–13 "This is unlikely to be a practical solution, given the possibly complex nature of  $z$ , and conflicting demands on expensive simulator output". I would add the difficulty of searching high dimensional spaces which in general can have large numbers of local minima.

**Response:** Added as requested.

**Comment:** 10. Page 2384, "We fix the standard deviation of the representative observational error as 10% of the maximum simulated value for each of the outputs in the ensemble". It is good to see that the authors do explore the effects of non-zero observational errors. This value does seem quite large though as this means 3 sigma in the implausibility will correspond to greater than 30% of this value. The authors should then not be too discouraged to see that not much space is subsequently ruled out.

**Response:** While this uncertainty might seem relatively large, it is chosen for three reasons. 1) Real world observations within climate science often have relatively large uncertainties. 2) We wished to use a plausible-but-large uncertainty in order to ensure that we could very clearly see the effect that that uncertainty had. 3) As noted in the text, this uncertainty might be partitioned between observations and model discrep-

[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)[Discussion Paper](#)

ancy, in which case 30

**Comment:** 11. Page 2388, "case that there exists a poorly modeled discrepancy (an "unknown unknown"), the ability of data to constrain the simulator will be overconfident." This may need clarification, because as was discussed previously, the unknown unknown could lead to far more space being ruled out than the maximum bound from this calculation, or far less than the worse case scenario given. As can be seen, reasonable modelling of model discrepancy is usually unavoidable when considering any observed results from the real world.

**Response:** The paragraph has been amended for clarity, and to make the language consistent with the changes requested by reviewer comment 1.

**Comment:** 12. The authors cite Craig et al. (2001) when discussing history matching and implausibility. It might be reasonable to cite the first two papers using this method, which also feature emulation and model discrepancy:

Craig, P. S., Goldstein, M., Seheult, A. H., and Smith, J. A. (1996). "Bayes linear strategies for history matching of hydrocarbon reservoirs." In Bernardo, J. M., Berger, J. O., Dawid, A. P., and Smith, A. F. M. (eds.), *Bayesian Statistics 5*, 69–95. Oxford, UK: Clarendon Press.

Craig, P. S., Goldstein, M., Seheult, A. H., and Smith, J. A. (1997). "Pressure matching for hydrocarbon reservoirs: a case study in the use of Bayes linear strategies for large computer experiments." In Gatsonis, C., Hodges, J. S., Kass, R. E., McCulloch, R., Rossi, P., and Singpurwalla, N. D. (eds.), *Case Studies in Bayesian Statistics*, volume 3, 36–93. New York: Springer-Verlag.

**Response:** References added as requested

### Technical Corrections

**Correction:** 1. Page 2374, lines 12–16 the authors write:

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Discussion Paper



"We can represent output  $y$  as an uncertain function of input  $x$  thus:  $y = g(x)$ . (1) The simulator is complex enough that we cannot trivially predict the output  $y$  at a given  $x$  before the simulator is run."

This section seemed a little unclear, surely the sentence after (1) should come before the authors talk about  $y$  being an uncertain function of  $x$ ?

**Response:** Fixed as requested

**Correction:** 2. Page 2375, line 8 "a constraining  $X$ " should read "a constraint on  $X$ " or something similar.

**Response:** Fixed as requested

**Correction:** 3. Page 2376, line 6 "at the point", perhaps should be "at the input point  $x$ " for clarity.

**Response:** Fixed as requested

**Correction:** 4. Page 2380, line 3 "and also check that with the true value" should read "and also check that the true value"

**Response:** Fixed as requested

**Correction:** 5. Page 2380, line 17 "it it" should read "if it".

**Response:** Fixed as requested

**Correction:** 6. Page 2382, lines 5–6 "The emulator is composed of a basic linear statistical model, along with a more flexible part," I know the authors want to avoid too much statistical terminology but I think it would be reasonable to add here "know as a Gaussian process" or some such phrase.

**Response:** Added as requested

**Correction:** 7. Page 2387, line 10 "This could be a powerful in the process of simulator develop<sup>Â</sup>ment." should read "This could be a powerful technique in the process of

simulator development." or something similar.

**Response:** Added as requested

**Correction:** 8. Page 2396 lines 2–3 "Implausibility is calculated the maximum of that from all three summaries" should read "Implausibility is calculated as the maximum of that from all three summaries".

**Response:** Added as requested

**Correction:** 9. Page 2399, Fig 7.a. 3rd panel from left: the blue representing the PDDFI input seems to be missing from this plot, where as the plot directly below includes this blue. As this input (and several other inputs) are not constrained by the maximum ice thickness output this is not so important but for consistency it would be good to have the same background blue colour in both Fig 7.a. 3rd panel and Fig 7.b. 3rd panel, if it is not too much trouble.

**Response:** The colours are ordered according to the total constraint by each parameter, in order that the most constrained parameters always appear at the “front” of the plot (and are therefore visible against the lesser-constrained parameters). In the noted case, the difference between parameters is very small (neither are constrained very much), but is enough to change the ordering. The reason for ordering has been added to the figure caption.

---

Interactive comment on Geosci. Model Dev. Discuss., 6, 2369, 2013.

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Discussion Paper