

## ***Interactive comment on “A novel model evaluation approach focussing on local and advected contributions to urban PM<sub>2.5</sub> levels – application to Paris, France” by H. Petetin et al.***

**H. Petetin et al.**

hervepetetin@gmail.com

Received and published: 27 February 2014

We would like to thank the second referee for the detailed and relevant questions he raised, notably concerning our methodology, which give us the opportunity to probably improve the clearness of our approach.

1. Back trajectories and determination of the upwind site (page 6402-4-11) : The referee asks several questions about the validity of the simplistic method used to estimate advected contributions. Each day, the whole set of 40 back trajectories calculated for one day (10 trajectories each 6 hours) is considered for the determination of the upwind site and there is no separate determination each 6 hours. As the referee supposed,

C2674

for a specific day, all back trajectories rarely indicate only one sector (this occurs only 8% of time), but most of time two (43%) or three (47%) sectors. Actually, even if it is not the only one, the dominant sector usually gathers the majority of back trajectories : in average, 73% of back trajectories belong to this dominant sector, against 22 and 5% for the second and the third sector, respectively. The occurrence of stagnant conditions remains very low, with only 4% of hourly wind speed values below 1 m s<sup>-1</sup> over the whole year. However, the most important aspect of our procedure is the calculation of uncertainties, which keep us from developing a more detailed and maybe more realistic (but still uncertain) procedure for the quantification of advected contributions. Indeed, whatever the selected upwind site (or even if we would have considered a mean concentration between several sites), by assuming an uncertainty defined as the maximal concentration range among the whole set of rural sites, the value of the advected contribution will always be included in the confidence interval. I will add the following sentences in the revised manuscript at 6402-11 : "Due to the complexity of wind fields, this procedure is certainly too simplistic to account for all meteorological situations that may occur over the Paris region (e.g. back-trajectories originating from more than one sector, recirculation). However, all these problems relative to the choice of the appropriate upwind rural site are tackled by the quantification of advected contributions uncertainties in which all the three rural concentration values are included, as described in the next section."

2. Advected regional background (page 6402-13-14) : The referee then asks several question about the representativeness of the quantified advected regional background. Both RUS and RNW stations take part of the AIRPARIF network and are officially classified as rural according to the criteria defined in the Annex III of the 2008/50/EC Directive. Specifically set for the PARTICULES campaign, the RNE station also follows these criteria and is thus classified as rural. All three sites are located in small municipalities with population density below 100 inhab km<sup>-2</sup>. Both RUS and RNW sites are clearly representative of the rural background. Concerning the RNE site, an influence of local woodburning emissions have been noticed from levoglucosan measurements,

C2675

and OM concentrations measured at this site cannot thus be considered as representative (and have been removed). As the other chemical constituents are far less (EC) or not (secondary inorganic compounds) influenced by such emission sources, their measurements remain representative of the rural background. The representativeness scale of a rural station is difficult to assess precisely since it depends on the heterogeneity of the regional background which in turn is determined by various factors (e.g. location and intensity of emission sources, wind field structure). Except when they are impacted by the Paris plume, these rural sites are representative of the regional background within a large extent. The Paris agglomeration is located in a region mostly occupied by agricultural activities with thus limited PM<sub>2.5</sub> emissions, as illustrated by the PM<sub>2.5</sub> emissions map (Fig. 1 in this Author Comment). The region is rather flat and windy (mean wind speed of 3.1 m s<sup>-1</sup> measured at the ground MONTsouris site in the Paris center), and as previously mentioned stagnant conditions are very rare. All these features thus favor a good dispersion of air pollution in the Paris region. Therefore, most of time, the Paris agglomeration cannot influence PM<sub>2.5</sub> levels at all three rural sites, which allows us to assume that the regional background advected toward the city is comprised between the lowest and the highest rural sites concentration. This hypothesis (on which is based the estimation of uncertainties in Sect. 4.2) thus appears quite reasonable. I will add at 6402-20 : "Except when they are impacted by the Paris plume, rural sites can be considered as representative of the regional background. As the Paris region is rather flat and windy, stagnant conditions remains very rare (over the year, only 4% of hourly wind speed values measured at the MONTsouris site are below 1 m s<sup>-1</sup>), which most of time prevents all rural sites to be simultaneously influenced by the Paris pollution". And at 6402-24 : "[...] at this site. The RNE rural site can thus not be considered as representative of the OM rural background, but this local wood burning pollution is not assumed to impact significantly the other species. In order to [...]".

3. Back trajectories 4-hours before Paris (page 6410-1-3) : The referee asks how an continental origin can be deduced from back trajectories only 4 hours before Paris.

C2676

The 4-hour backward time period only allows to assess which rural sites have been potentially overflowed by air masses before reaching Paris, but is of course insufficient to assess the geographic origin at the continental scale. The link established in the paper between some intense episodes (in particular during wintertime) and a continental origin of air masses from the north-east derived from an analysis of back trajectories a few days before reaching Paris. To avoid any ambiguity, I will change the sentence at 6410-1-3 to : "The variability (standard deviation of 8.6  $\mu\text{g m}^{-3}$ ) strongly depends on the wind regime, with large episodes mostly linked to advection of continental air masses from the north-east wind sector as indicated by back trajectories over a few days."

4. Modeled contributions : The referee wonders how modeled contributions are derived. Both the observed and modeled contributions are calculated the same way, after the interpolation of simulated concentrations at the four sites. I will add at 6406-16 : "After interpolation of concentrations at all four sites, simulated contributions are derived in the same way as observed ones."

5. Vertical distribution of emissions (page 6401) : The referee asks how emissions are vertically allocated in the model. Area and line sources are emitted in the first layer, while emissions from point sources are vertically allocated depending on several parameters (e.g. ambient temperature, ejection speed and temperature, smokestack height). I will add at 6401-8 : "Area and line sources are emitted in the first layer, while emissions from point sources are vertically allocated depending on several parameters (e.g. ambient temperature, ejection speed and temperature, smokestack height)."


6. MFB and MFE (page 6406-15) : The referee asks about the absence of MFE and MFB definitions in Sect. 4.3 (and maybe in the general discussion). To our sense, MFB and MFE statistical indicators are easier to interpret on a graphic way with the Boylan and Russel performance goals, as in Fig. 13, while the traditional indicators (bias, RMSE) make the discussion easier to understand. This explains why these two metrics are not mentioned before Sect. 5.8.

C2677

7. OC versus OM : The referee finally asks why comparisons are not performed on OC rather than OM. Actually, while primary organic aerosol emission inventories correspond to OC (due to emission factors mostly measured by thermo-optical methods), the secondary organic aerosol produced in the model cannot be considered as OC but is closer to OM. That is why we apply a conversion to emission input data (1.6 in our case) in order to represent OM concentrations rather OC ones. Additionally, the OM to OC conversion remains mandatory in the comparisons of PM2.5 total mass.

Interactive comment on Geosci. Model Dev. Discuss., 6, 6391, 2013.

C2678



**Fig. 1.** PM2.5 emission density in 2010 over the Paris region (t km<sup>-2</sup> year<sup>-1</sup>).

C2679