

## ***Interactive comment on*** “Evaluation of the new **UKCA climate-composition model** **– Part 2: The Troposphere” by F. M. O’Connor et al.**

### **Anonymous Referee #1**

Received and published: 6 May 2013

Review of O’Connor et al., Evaluation of the new UKCA climate-composition model.  
Part II. The Troposphere

The manuscript details one variant of the UK Chemistry and Aerosols (UKCA) model; presented here is the StdTrop configuration that has been used for CMIP5. The significant components of the model are described and a wide ranging series of model results are presented and compared with observations, from radionuclides to assess transport and deposition, to chemical fields for methane, ozone, CO and NO<sub>x</sub>. While the comparison with observations is broad and generally well presented, there seem to be a number of important comparisons that are omitted from the article, and which are frequently included in similar model description articles, leaving the reader with an incomplete view of the model. The assessment of the model carbon monoxide, pre-

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)

[Discussion Paper](#)

sented in Section 5.2, is an example. The comparison of surface CO to a selection of available observation stations is routinely done for these type of model description papers and is well presented here. To investigate the distribution of CO away from the surface, comparisons are made with observations from a variety of aircraft campaigns performed during the 1990s. From the comparison, the model is found to be biased high or low depending on the field campaign and the comparison as a whole is subject to the caveat that '...the comparison at best is expected to be qualitative.' There are really fantastic satellite observations of CO available and the reader is referred to comparisons with TES to be found in Voulgarakis et al (2011b) and Telford et al. (2013) but it is not readily apparent how applicable these comparisons with TES are since they were conducted with the TropSop version of the model. Since this manuscript is meant to provide the reference for the StdTrop version of UKCA and the comparison with the aircraft observations seems inconclusive, I would strongly suggest replacing the comparison with aircraft observations presented in Section 5.2 with a comparison of CO with satellites observations.

The distribution of ozone for January shown in Figure 22 shows concentrations over northern hemisphere continents almost uniformly below 20 ppbv. Very low concentrations can also be seen near the surface in the northern hemisphere in the cross-section of zonally averaged ozone in Figure 23. Looking back to Hauglestaie et al. (2004), January surface concentrations are only very rarely below 20 ppbv over northern hemisphere continental regions, with large areas showing average values of 25 – 35 ppbv. A similar range of values is shown by Zeng et al. (2008) for January over northern hemisphere continental locations. I would suggest the inclusion of a few continental sites for surface ozone observations. Many of the sites presently used for the comparison are island or coastal sites and while Niwot Ridge is continental, it is also at high altitude. Other papers have frequently compared to Hohenpeissenberg and find pretty good comparisons, so that would be one suggestion. Is there an idea why the UKCA model produces such low values over continental regions?

[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)[Discussion Paper](#)

The absence of a global ozone budget for the model is also of note. Although largely theoretical, an ozone budget is frequently a component of these model development papers and allows the reader to ascertain the size of strat-trop exchange and dry deposition.

While the comparison with the aircraft campaign CO was inconclusive, these campaigns also provide some of the best (only??) observations for reactive nitrogen compounds like HNO<sub>3</sub> and PAN. I note there is a comparison for NO<sub>x</sub> in the paper now, but nothing similar for HNO<sub>3</sub> and PAN. Why have these been omitted? Perhaps PAN will be significantly underestimated without isoprene chemistry, but that fact would be important to establish - particularly in relation to the other articles on different versions of UKCA.

Minor comments on the manuscript.

Page 1749, Lines 1-5; Is the aerodynamic resistance ( $r_a$ ) tied to the model treatment of the surface layer? A sentence or two on how  $r_a$  is derived would help complete the description.

Page 1752, Line 27; Figure 3 shows the model values with a red vertical line. I assume it represents a moment of the distribution of the model values, but what exactly should be specified.

Page 1757, Lines 16-20; Has the mass conservation also been assessed by comparing the <sup>210</sup>Pb deposition (for the tracer that is deposited) against the <sup>222</sup>Rn emission? Analysing the non-deposited <sup>210</sup>Pb would capture conservation errors in the <sup>222</sup>Rn, but the non-deposited <sup>210</sup>Pb would very quickly become quite homogeneous on at least the hemispheric scale.

Page 1760, Line 3; Where it is stated that ' $t_d$  is the e-folding lifetime', does that refer to the lifetime to radioactive decay of 10.76 years given a bit earlier? As it is currently written, the terminology is a bit inconsistent.

[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)[Discussion Paper](#)

Page 1767, Line 24; I don't see the need to include Figure 12, showing the spatial distribution of CO emissions.

---

Interactive comment on Geosci. Model Dev. Discuss., 6, 1743, 2013.

**GMDD**

6, C497–C500, 2013

---

Interactive  
Comment

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Discussion Paper

C500

