

1 **Grassland Production Under Global Change Scenarios for**
2 **New Zealand Pastoral Agriculture**

3
4 **E. D. Keller¹, W. T. Baisden¹, L. Timar^{1,2}, B. Mullan³, and A. Clark^{3,*}**

5 [1]{GNS Science, Lower Hutt, New Zealand}

6 [2]{Motu Economic and Public Policy Research, Wellington, New Zealand}

7 [3]{NIWA, Wellington, New Zealand}

8 [*]{now at: New South Wales Department of Primary Industries, Orange, NSW, Australia}

9 Correspondence to: E. D. Keller (l.keller@gns.cri.nz)

10
11 **Abstract**

12 We adapt and integrate the Biome-BGC and Land Use in Rural New Zealand models to
13 simulate pastoral agriculture and to make land-use change, intensification of agricultural
14 activity and climate change scenario projections of New Zealand's pasture production at time
15 slices centred on 2020, 2050 and 2100, with comparison to a present-day baseline. Biome-
16 BGC model parameters are optimised for pasture production in both dairy and sheep/beef
17 farm systems, representing a new application of the Biome-BGC model. Results show up to a
18 10% increase in New Zealand's national pasture production in 2020 under intensification and
19 a 1-2% increase by 2050 from economic factors driving land-use change. Climate change
20 scenarios using statistically downscaled global climate models (GCMs) from the IPCC Fourth
21 Assessment Report also show national increases of 1-2% in 2050, with significant regional
22 variations. Projected out to 2100, however, these scenarios are more sensitive to the type of
23 pasture system and the severity of warming: dairy systems show an increase in production of
24 4% under mild change but a decline of 1% under a more extreme case, whereas sheep/beef
25 production declines in both cases by 3% and 13%, respectively. Our results suggest that high-
26 fertility systems such as dairying could be more resilient under future change, with dairy
27 production increasing or only slightly declining in all of our scenarios. These are the first
28 national-scale estimates using a model to evaluate the joint effects of climate change, CO₂
29 fertilisation and N-cycle feedbacks on New Zealand's unique pastoral production systems that

1 dominate the nation's agriculture and economy. Model results emphasize that CO₂
2 fertilisation and N-cycle feedback effects are responsible for meaningful differences in
3 agricultural systems. More broadly, we demonstrate that our model output enables analysis of
4 Decoupled Land-Use Change Scenarios: the Biome-BGC data products at a national or
5 regional level can be re-sampled quickly and cost-effectively for specific land-use change
6 scenarios and future projections.

7

8 **1 Introduction**

9 Intensive pasture grazing systems dominate New Zealand's agricultural production, in
10 contrast to the cultivated cropland and animal feeding operations that make up the majority of
11 agriculture in other developed countries worldwide. In this respect New Zealand is unusual,
12 with the national scale and economic importance of its pastoral agriculture system
13 representing an extreme. The dairy and sheep/beef industries in particular are central to the
14 New Zealand economy. Production from primary industries makes up 12% of New Zealand's
15 GDP, of which the dairy industry alone contributes almost 3%. Dairy products comprise over
16 a quarter of New Zealand's total exports, with an export value of NZ\$13.9 billion for the year
17 2011/12. Meat and wool exports are also significant, with beef and lamb export value at
18 NZ\$5.6 billion in 2011/12 (New Zealand Ministry for Primary Industries, 2012; Schilling et
19 al., 2010). While sheep and beef farms still make up the majority of agricultural land, dairy
20 farming is rapidly expanding in size and area, continuing a long-term decadal shift towards
21 more intensive but historically more profitable dairy pasture. New Zealand's future pasture
22 production thus features prominently in many national projections in economics, primary
23 production and the environment, including water quality and climate change impacts and
24 adaptation.

25 Consequently, there is a need to understand possible changes to the productivity of New
26 Zealand's pastoral agriculture systems under a range of future scenarios. Because New
27 Zealand depends almost exclusively on pasture for animal feed, climate change could have
28 considerable effects on the nature and profitability of dairy and sheep/beef farming in the
29 short- and long-term. In addition to economic considerations, pasture production estimates are
30 crucial in addressing questions of environmental sustainability. Unlike most developed
31 countries, whose greenhouse gas (GHG) emissions are largely derived from fossil fuels and
32 carbon dioxide (CO₂), almost half of New Zealand's GHG emissions arise from agriculture

1 and methane (CH_4). Accurate estimates of pasture production are required to understand such
2 questions as whether feed supply can become a limiting factor on the number of grazing
3 animals used to calculate New Zealand's emissions of methane and nitrous oxide (N_2O).

4 Complex environmental and policy questions such as these can be addressed through
5 integrated modelling assessments of the impact of climatic, economic, and land management
6 factors on future agricultural productivity. There are many examples of studies at the global,
7 regional, and national level that couple dynamic biophysical process-based or statistical crop
8 models with climate and land-use datasets to estimate quantities such as carbon flux and
9 storage, net primary productivity (NPP) and water availability for both managed and natural
10 ecosystems under future climate and management scenarios (Beer et al., 2010; Bondeau et al.,
11 2007; Gumpenberger et al., 2010; Rost et al., 2009; Roudier et al., 2011). Coupling models in
12 this way is important in accounting for non-linear feedbacks and interactions between climate,
13 the carbon cycle, and land-use and management decisions, which can be quite significant
14 (Ronneberger et al., 2006). Additionally, it is essential to integrate climate change effects and
15 feedbacks into economic and policy assessments for their impact to be considered in the
16 decision-making process and to affect long-term planning and preparedness.

17 The present study is intended to develop flexible output and data products at a national or
18 regional level that enable the examination and analysis of a range of scenarios and their
19 possible impacts on pasture production in New Zealand at particular time slices over the next
20 100 years. It builds on previous research efforts and existing datasets to understand some of
21 the biophysical, climatic, economic, and land management variations that might affect the
22 future productivity of New Zealand's pastoral agriculture. Like many small countries, New
23 Zealand (population 4.5 million, land area 260,000 km^2) requires cost-effective model
24 infrastructure that is capable of evaluating policy options on timescales of months or weeks.
25 The development of the Land Use in Rural New Zealand (LURNZ) model meets this
26 challenge and, importantly, enables the integration of global change scenarios with
27 contemporary policy choices. Before outlining the rationale for our model infrastructure, it is
28 useful to briefly review the history of integrated model development in New Zealand,
29 primarily aimed at the climate change component of global change.

30 Earlier integrated modelling studies of New Zealand pasture include the 2001 CLIMPACTS
31 study (Warrick et al., 2001), which used a global climate model in combination with New
32 Zealand datasets to produce estimates of pasture production based on scenarios from the

1 Intergovernmental Panel on Climate Change (IPCC) Second Assessment Report. The more
2 recent EcoClimate Report (Stroombergen et al., 2008) estimated productivity for 2030 and
3 2080 for several agricultural sub-sectors, including sheep/beef and dairy pasture systems, and
4 provided a preliminary integrated assessment of possible economic costs and benefits of
5 climate change. National and regional projections were based on a climate-index approach
6 (Baisden, 2006) and statistically downscaled scenarios from the HadCM2 model in the 2001
7 IPCC Third Assessment Report (Mullen et al., 2005). However, the methodology does not
8 account for the potentially important effects of increased carbon dioxide concentrations in the
9 atmosphere (CO₂ fertilisation) and the interaction with progressive nitrogen (N) limitation.
10 The earlier CLIMPACTS methodology accounted for CO₂ fertilisation but not N limitation.
11 These omissions could alter results substantially because studies involving the effect of CO₂
12 fertilisation on plant physiology and growth suggest that plant biomass increases overall
13 under elevated CO₂, but the response depends strongly on nutrient (N and P) availability
14 (Ainsworth and Long, 2005; de Graaff et al., 2006; Newton et al., 2010).

15 To address these concerns about the lack of CO₂-N-climate interactions and feedbacks, and
16 also to fulfil the need for a suitable temporal and spatial resolution, we use the Biome-BGC
17 model. The model provides a level of complexity intermediate between simple climate-index-
18 driven pasture production and a full farm system model (e.g. APSIM, Keating et al., 2003),
19 which can be difficult to extrapolate across space and apply at a national scale. Biome-BGC is
20 able to simulate daily climate variables, water availability and irrigation, CO₂ fertilisation
21 effects, all relevant nitrogen inputs and outputs, and the utilisation of pasture by grazing
22 animals without requiring an overwhelming level of detail about individual farms.

23 We simulate the two dominant types of pasture systems in New Zealand, which we will refer
24 to as 'dairy' and 'sheep/beef'. The main difference between these two systems is the intensity
25 of grazing, dairy being the more intensive of the two. Dairy farming is associated with highly
26 productive pasture and therefore involves higher stocking rates, more nitrogen fertilisation, a
27 larger amount of animal products extracted from the system, and often the addition of
28 irrigation (which we do not model here). We develop model parameterisations for each type
29 of pasture system that enables the simulation of national and regional pasture production for
30 the present baseline and projections for future scenarios. By then sampling the model's output
31 across land-use extents, we introduce analysis of Decoupled Land-Use Change Scenarios
32 (DLUCS): the pasture data products at a national or regional level can be re-sampled quickly

1 and cost-effectively for specific land-use change projections. This is a flexible approach that
2 can be easily applied to many specific scenarios of climate and land-use change, not only the
3 ones presented here. Herein, we describe our process of parameterising Biome-BGC for
4 grazed pasture agro-ecosystems on a ~5km grid covering New Zealand, explain our
5 methodology and choice of model scenarios, and report national production results for dairy
6 and sheep/beef systems.

7

8 **2 Methods**

9 **2.1 Biome-BGC model**

10 The Biome-BGC (Bio-Geochemical Cycles) model v4.2 Final Release (Thornton et al., 2005)
11 is an ecosystem process model that simulates the biological and physical processes controlling
12 cycles of carbon, nitrogen and water of vegetation and soil in terrestrial ecosystems. The
13 model is capable of simulating evergreen, deciduous and broadleaf forests, C3 and C4
14 grasslands, and shrub ecosystems. The primary input consists of weather conditions at a daily
15 time step, as well as site-specific information such as elevation, soil composition and rooting
16 depth. In addition, there is a set of 43 adjustable ecological parameters that can be customized
17 for a particular ecosystem. The model and its parameters are described in detail in Thornton et
18 al. (2002), White et al. (2000), and Thornton (1998). The Biome-BGC model has been
19 extensively tested and validated for North American and European evergreen and deciduous
20 forest, grassland, and mixed ecosystems (Jung et al., 2007; Pietsch et al., 2005; Bond-
21 Lamberty et al., 2005; Wang et al., 2009). There have also been other adaptations of the
22 Biome-BGC model to managed agricultural systems and crops (Hidy et al., 2012; Di Vittorio
23 et al., 2010; Wang et al., 2005) that involve supplementary code and/or that are specific to
24 grasses and crops in other regions. Extension to New Zealand ecosystems and managed
25 pasture systems represents a new application. We adapt the model through parameter
26 adjustments rather than by modifying model code.

27 We used the Biome-BGC model's built-in C3 grasslands mode to simulate our two managed
28 pasture systems, dairy and sheep/beef. While the core model is not currently designed for
29 farm systems or the presence of grazing animals, we can reinterpret or redefine some of the
30 model's ecological parameters and calibrate them to adequately represent grazing and harvest.
31 Specifically, the 'annual whole-plant mortality fraction' parameter can be related via a simple

1 algebraic formula to pasture utilisation (the fraction of above-ground biomass production
2 eaten by grazing animals):

3
$$\text{whole plant mortality} = \frac{\text{pasture utilisation}}{(1 + \text{leaf : fine root C})}. \quad (1)$$

4 where leaf:fine root C is a model parameter representing the relative allocation of carbon
5 above and below ground. Intuitively, plant mortality increases along with the amount of grass
6 eaten. For the sake of simplicity we have assumed a nationwide fixed level of pasture utility
7 of 0.55 and 0.90 in sheep/beef and dairy pasture systems, respectively. This results in a
8 mortality proportion that is much higher than that of a natural grassland (default is 0.1). In the
9 same manner, the removal of meat and milk products from the system is factored into the
10 model's 'annual fire mortality fraction' parameter, which describes the proportion of plants
11 that die due to fire each year and that is effectively removed from the ecosystem. Since fire is
12 not normally a significant occurrence in managed pasture, in our model this fraction
13 represents the approximate proportion of nutrients removed from the ecosystem via milk and
14 meat production. We have set dairy systems to have twice the proportion removed (0.2) as
15 sheep/beef (0.1). (The default for grassland is 0.1).

16 We have also included the combined effects of managed fertiliser application and fertility-
17 driven nitrogen fixation through the model's site-specific nitrogen fixation input parameter.
18 The symbiotic and asymbiotic nitrogen fixation rate is typically on the order of 10^{-4} kg
19 N/m²/year for most naturally occurring ecosystems. To represent fixation under high P
20 fertilisation regimes and urea or other N additions common in New Zealand dairy farming, we
21 have set this rate much higher, on the order of 10^{-2} . In our parameterisation, dairy systems
22 have twice the rate of N input via fixation and fertilisation (0.032 kg N/m²/year) as sheep/beef
23 (0.018 kg N/m²/year).

24 **2.2 LURNZ model**

25 To develop an estimate of actual total national pasture productivity, we combined information
26 from the Land Use in Rural New Zealand (LURNZ) model v2 with the pasture production
27 outputs from the Biome-BGC model. The LURNZ model (Hendy et al., 2007, 2008; Timar
28 2011; Kerr et al., 2012) was developed to explain and simulate changes in four major rural
29 land-use types in New Zealand: dairy, sheep and beef, plantation forestry and regenerating
30 natural forest (henceforth termed scrubland). LURNZ models land use both dynamically,

1 based on national time-series econometric estimates of land-use change, and spatially, based
2 on cross-sectional observations of biophysical and socio-economic land attributes. Of most
3 relevance to modelling future pasture production is the ability of LURNZ to evaluate any
4 scenario that can be expressed as a commodity price change in one of the four sectors. The
5 model provides a baseline of actual land use in 2008 and scenario projections for changes to
6 land use in 2020 and 2050 in response to an imposed price on carbon (and agricultural CH₄
7 and N₂O emissions, converted to CO₂ equivalent terms). In the climate change scenarios that
8 follow, our dairy and sheep/beef regional breakdown and total national estimates are based on
9 the LURNZ model's observed land-use distribution in 2008, and land-use scenarios are based
10 on projected dynamic change in dairy and sheep/beef land uses in 2020 and 2050.

11 **2.3 Climate and input datasets**

12 The New Zealand National Institute of Water and Atmospheric Research (NIWA) Virtual
13 Climate Station Network (VCSN) provides the daily weather input required by the Biome-
14 BGC model. The VCSN is a set of virtual 'weather stations' that uses interpolation techniques
15 to provide detailed weather information at each point on 0.05° grid covering all of New
16 Zealand, approximately 5 x 5 km resolution (Tait et al., 2006). Daily weather data is available
17 for each grid cell from 1972 to the present. Direct and indirect inputs to the model from the
18 network include maximum and minimum temperature, precipitation, solar radiation, relative
19 humidity, vapour pressure deficit, and wind run (available from 1997). Before running each
20 model scenario, the model is first 'spun up' by recycling the input datasets as many times as
21 necessary for the model to reach a steady state over ~1000-2000 years (Thornton and
22 Rosenbloom, 2005).

23 Future climate change scenarios circa 2050 and 2100 were statistically downscaled to the
24 VCSN using three GCMs from the IPCC Fourth Assessment Report (AR4): giss-eh
25 (NASA/Goddard Institute for Space Shuttles, USA, rererred to as 'GIEH'), mpi_echam5
26 (from the Max Planck Institute in Germany, referred to as 'MPI') and cccma_cgcm3_1 (from
27 the Canadian Climate Centre, referred to as 'CCC'). This input nominally refers to the 9-year
28 period from 2046-2054 and the 15-year period from 2097-2111, although it is meant to
29 represent an approximate timeframe of 50 and 100 years from the present day. Further
30 explanation of these scenarios is in section 3 (see also Renwick et al., 2013; Baisden et al.,
31 2010).

1 Although the Biome-BGC model does not have an explicit mechanism to incorporate daily
 2 wind speed, wind does have a significant evaporative effect on pasture growth in many New
 3 Zealand regions. We account for the role of wind in enhancing water loss from pastures by
 4 correcting the daily water vapour pressure deficit (VPD) input data with the predicted FAO
 5 Penman-Monteith effect of wind on evapotranspiration for grasslands. This effect is
 6 particularly important where hot, dry northwesterly winds enhance seasonal drought in the
 7 hill country and plains of New Zealand's east coast regions. The modified VPD (which
 8 becomes the direct daily VPD input to the model) is calculated from the following equation
 9 for evapotranspiration (Allen et al., 1998):

$$10 \quad ET_o = 0.408 * \Delta(R_n - G) + \frac{\gamma \left(\frac{900}{T + 273} \right) u_2 (e_s - e_a)}{\Delta + \gamma(1 + 0.34u_2)}. \quad (2)$$

11 where ET_o = reference evapotranspiration (mm day^{-1}), R_n = net radiation at the crop surface
 12 ($\text{MJ m}^{-2} \text{ day}^{-1}$), G = soil heat flux density ($\text{MJ m}^{-2} \text{ day}^{-1}$), T = mean daily air temperature at 2
 13 m height ($^{\circ}\text{C}$), u_2 = wind speed at 2 m height (m s^{-1}), e_s = saturation vapour pressure (kPa), e_a
 14 = actual vapour pressure (kPa), $(e_s - e_a)$ = saturation vapour pressure deficit (kPa), Δ = slope
 15 vapour pressure curve ($\text{kPa } ^{\circ}\text{C}^{-1}$), and γ is the psychrometric constant ($\text{kPa } ^{\circ}\text{C}^{-1}$). For our
 16 purposes, we take $G = 0$ and $\gamma = 0.054$, and Δ is the following:

$$17 \quad \Delta = 2503.06 \frac{\exp\left(\frac{17.27T}{237.3+T}\right)}{(237.3+T)^2}. \quad (3)$$

18 where T is again the mean daily air temperature.

19 The required soil texture and effective rooting depth for each site was obtained from the New
 20 Zealand Fundamental Soil Layers (FSL) dataset (Landcare Research, 2013), which contains
 21 spatial information for 16 soil attributes, including soil texture classes. The soil texture classes
 22 were matched to the percentages of sand, silt and clay required by the model by visually
 23 identifying modal soil textures present in the National Soils Database.

24 National and regional pasture production is given in terms of both kilograms of dry matter per
 25 hectare and metabolisable energy (MJ per kg dry matter). Remote sensing, augmented by on-
 26 the-ground calibration, was used to estimate the seasonal metabolisable energy for all sites
 27 from model output.

2.4 Model calibration and validation

To adapt the Biome-BGC model for intensive pastoral agriculture, we adjusted key ecological parameters to optimise model output to measured pasture growth data in selected locations across New Zealand. Treating sheep/beef and dairy pasture systems as two different ‘biomes’, we developed a unique parameterisation for each type of system. We used an automated parameter estimation software package, PEST v12.0 (Doherty, 2005), which employs the Gauss-Marquardt-Levenberg inversion method to optimise a model’s output to user-supplied observation data. We fit the model’s net primary production (NPP) output to historical pasture clipping data from six sites spread temporally and geographically across New Zealand (three dairy and three sheep/beef). Pasture data is typically reported in units of dry matter (DM); NPP can be converted to the equivalent amount of dry matter with the following:

$$DM = 2.0 * \left(\frac{r_{ab}}{r_{ab} + 1} \right) * NPP . \quad (4)$$

where r_{ab} is the ratio of above-ground to below-ground allocation (given by the inverse of the new fine root C : new leaf C allocation parameter in the Biome-BGC model).

Calibration sites are depicted in Fig. 1 and additional detail about the sites is in Table 1. Pasture growth data at the three dairy calibration sites was obtained from Landcorp Farming (personal communication, 2011) and publicly available data from DairyNZ (DairyNZ, 2011) and Lincoln University Dairy Farm (South Island Dairying Development Centre, 2014). Data at the three sheep/beef sites was obtained from Beef + Lamb New Zealand (Clarke-Hill and Fraser, 2007) and previously published articles (Rosser and Ross, 2011; Smith et al., 2012). Data consists of monthly, bi-weekly, or weekly measurements of pasture clippings over a period of at least two continuous years and up to seven years in the case of Winchmore Research Station in Canterbury. These sites were chosen on the basis of geographic location and data availability; we attempted to balance the desire to include a range of climates and regions in New Zealand with the need for high-quality and complete datasets. All data used for calibration was taken from non-irrigated pasture, with the exception of Lincoln University Dairy Farm (in this case irrigation was also simulated in the model during calibration by adding additional precipitation to the meteorological data input file when soil moisture deficit was above a threshold). The site at Te Whanga in the Wairarapa region provides three different datasets from hillside landslide scars of varying ages: a slip that occurred in 1961, a slip in 1977, and one uneroded location. This site is useful for model calibration because

1 recent scars have shallower soils, resulting in lower water storage capacity, thus providing
2 pasture production records under identical climate but varying soil properties (Rosser and
3 Ross, 2011). The Biome-BGC rooting depth parameter is adjusted according to scar age,
4 providing a way to calibrate the model against topsoil depth.

5 The parameterisation produced a good fit between model output and the observed annual
6 mean production and seasonal cycle of pasture growth. Comparing daily average growth rates
7 over each observation period, the correlation coefficient R^2 for all dairy and sheep/beef sites is
8 0.64 and 0.70, respectively. Figure 2 displays the pasture clipping data and the optimised
9 model fit for each of the calibration sites on a temporal scale. Figure 3 shows a direct
10 comparison of observed and modelled data from all dairy (left) and sheep/beef (right) sites.
11 Figures 2 and 3 also report major axis regression statistics performed by the lmodel2 v1.7-2
12 package in R v3.0.2 (Legendre, 2014; R Core Team, 2013) to quantify the model-data
13 relationship. A full list of adjusted and default parameters is in Appendix A.

14 To validate the sheep/beef model, we chose an additional 22 sites with pasture clipping data
15 (Clarke-Hill and Fraser, 2007) over a similar time period (2003-2005), selected for data
16 completeness and geographical spread. Figure 1 shows the locations of the validation sites.
17 Overall correlation R^2 for these sites when comparing individual measurement intervals is
18 0.36. The scatter plot shown in Fig. 4 reveals that the model is biased low at higher values,
19 often underestimating the observed peaks in production in spring and summer. The three
20 northern-most sites (Broadwood, Tauranga, and Whakatane) perform very poorly, possibly
21 because the climate is warmer than the rest of New Zealand and C4 grasses (rather than C3)
22 are common. Removing these sites, R^2 rises to 0.41. There is also significant variation in
23 pasture age, hill slope, fertilization and nutrient content among individual sites that are not
24 included in our model and could account for the large difference in model fit at specific
25 locations. Monthly averages over several years generally compare better to observations (Fig.
26 4), but there is still a bias during spring and summer months. The correlation coefficient R^2
27 for monthly averages aggregated over three years of pasture clipping data for all validation
28 sites is 0.48, and without the three northern sites is 0.59. The relationship between the model
29 and measurements in Fig. 4 is quantified wth major axis regression statistics using the same
30 methodology as in Figs. 2 and 3.

31 For the dairy model, no additional spatially varying data was available for validation at the
32 time of our study. Consequently, we use national milk production data as a proxy for pasture

1 growth and an evaluation of model performance. In a separate report, we examined the
2 relation between modelled pasture production and total national milk solids production data
3 for New Zealand (Keller, 2012), finding excellent correlation in the six years from June 2006
4 to May 2012 (annual milk season in New Zealand runs from June to May) and moderate
5 correlation over the last 15 years ($R^2 = 0.86$ and 0.46, respectively). Although indirect, this
6 demonstrated relation to actual milk production data allows us to have reasonable confidence
7 in the national model output.

8 We focus primarily on seasonal and annual averages at a national level in this study. In
9 addition, because we are evaluating future scenarios relative to the baseline and are not
10 concerned with absolute levels of production, our subsequent analysis is minimally affected
11 by model bias.

12 **2.5 Methodology and model scenarios**

13 We introduce DLUCS methodology to construct and analyse model scenarios: we simulate
14 biophysical conditions affecting grass growth with the Biome-BGC model to produce an
15 estimate of pasture production at all locations on the national grid, then sample selectively
16 according to the specific land use or economic situation modelled with LURNZ. Scenarios
17 can be anything that can be modelled through changes in weather or nutrient input and/or
18 economic drivers of land-use change. Land use is decoupled from the biophysical dynamics
19 of plant growth, and the two are integrated at the final stage. By creating a national production
20 dataset with Biome-BGC and then re-sampling it using the output from LURNZ, we are able
21 to quickly examine many different plausible land use and economic scenarios relevant for
22 policy decisions, including the response to climate change.

23 Scenarios for this project were developed in consultation with the New Zealand Ministry for
24 Primary Industries (MPI) and reflect factors that are assumed to have significant impact on
25 pasture productivity in New Zealand. All scenarios are constructed to represent averages over
26 nine-year periods centred on 2005, 2020, or 2050, and in the case of climate change in 2100,
27 the 15-year period from 2097 to 2111. We run the model for each type of pasture for all grid
28 cells, regardless of actual land use; results are mapped for dairy and sheep/beef systems as if
29 all available land (exclusive of conservation land, water, year-round ice cover and urban
30 areas) was devoted to that system. We then calculate regional and national pasture production
31 totals by summing production from each grid cell categorised as either dairy or sheep/beef in

1 LURNZ. Spatial mapping and production summation were performed using ArcGIS 9.3
2 (ESRI, 2009). With the exception of the land-use change scenarios, all land-use
3 categorisations are derived from the LURNZ model's 2008 map, based on actual data, and
4 stay constant at 2008 levels in climate and intensification scenarios at 2020, 2050 and 2100 in
5 order to keep the effects of land-use change separate from the other effects that we simulate.
6 The scenarios chosen are intended, as much as possible, to isolate a single effect, so that the
7 sensitivity of pasture production to that particular effect alone can be estimated relative to the
8 baseline. However, we note that some scenarios are closely linked, and in practice it might not
9 be realistic to consider each one in isolation. We describe each scenario in detail in the
10 following sections.

11 **2.5.1 Baseline**

12 The baseline scenario is the output from the Biome-BGC model run using actual climate data
13 from the VCSN for 2001–2009, averaged over the nine years. This scenario is intended to
14 represent ‘present-day’ climate and to serve as a benchmark for comparisons to scenarios in
15 2020 and 2050. The baseline that is used as comparison for the 2100 climate change scenarios
16 is slightly different, covering the 15 years from 1997 to 2011, to correspond to the timing and
17 length of the 2100 simulations. The small inconsistency in the baselines does not alter the
18 general trend in the final results and was chosen to ensure matching with the statistically
19 downscaled climate datasets and a sensible averaging period for the scenarios studied.

20 **2.5.2 Land use**

21 We simulated dynamic land-use changes using the LURNZ model by assuming the primary
22 drivers behind land-use change are economic factors that influence the monetary returns to
23 land under different uses. We selected three scenarios focused on the importance and
24 associated uncertainties of the phase-in of emissions trading, corresponding to low, best-guess
25 and high carbon prices (NZ\$ 0, \$50 and \$100 per tonne CO₂e, respectively) under the New
26 Zealand Emissions Trading Scheme (ETS). Land-use projections are provided for 2020 and
27 2050 and subsequently combined with baseline dairy and sheep/beef pasture production.

28 Along with land-use change, intensification of current usage is also likely to be a considerable
29 driver of pasture production over the coming decades in the absence of new environmental
30 regulation (Parfitt et al., 2006, 2008). To simulate a representative ‘intensification’ scenario in
31 2020 with Biome-BGC, we increased the nitrogen fixation levels per hectare and the effective

1 utilisation of pasture nationwide. Parameter values used in the baseline and intensification
2 scenarios are compared in Table 2. Intensification here is based on high nutrient inputs that
3 represent roughly what a farmer would do in a ten-year timeframe in response to long-term
4 increases in commodity prices (near doubling). Climate was held constant at present-day
5 inputs, and CO₂ concentrations are increased according to the A2 scenario from the IPCC
6 Special Report on Emissions Scenarios (SRES; Nakicenovic et al., 2000). The combined
7 effects of CO₂ fertilisation and intensification were modelled together in this case to test for
8 strong non-additive interactions between elevated CO₂ and agricultural N cycling.

9 **2.5.3 Climate change**

10 The climate change scenarios we selected provide mid-range and upper-end estimates for the
11 combined impact of climate change and elevated CO₂ on pasture growth circa 2050 and 2100.
12 Scenarios were chosen from the ensemble of IPCC AR4 GCM simulations. The particular
13 models that were chosen are in good agreement with present-day climate in the New Zealand
14 region but forecast significantly different changes in local patterns of precipitation and
15 temperature by 2100. A more complete description of the projected changes for New Zealand
16 and the range of responses in selected AR4 GCMs is in Renwick et al. (2013). Climate
17 projections for New Zealand are based on a downscaling scheme that uses partial least
18 squares regression to statistically downscale rainfall, temperature, and solar radiation from
19 GCMs directly to the VCSN (Clark et al., 2011).

20 The SRES A2 emissions scenario is used to estimate the increase in CO₂ concentration levels
21 in the atmosphere for all climate change scenarios. This scenario results in approximately 4°C
22 of global mean average temperature increase by 2100 (measured since pre-industrial times,
23 nominally 1750). A2 is suitable as a mid-range projection in the shorter term out to 2050. In
24 longer-range climate change projections, it represents an upper-end scenario, which becomes
25 the case by 2080. Atmospheric concentrations start at 375 ppm in 2005 and rise to 827.3 ppm
26 in 2099 (Nakicenovic et al., 2000; ENSEMBLES, 2009). This scenario is increasingly
27 regarded as more likely since recent emissions have closely tracked its projections.

28 We first examined the effect of CO₂ fertilisation alone on pasture production, keeping climate
29 patterns the same as the baseline but increasing atmospheric CO₂ concentrations according to
30 the A2 scenario. This is referred to as 'Elevated CO₂'. This scenario was evaluated on the
31 short timeframe of 2020 to provide a partial derivative of elevated CO₂ effects on a timescale

1 during which the effects of climate change might remain within the bounds of regional and
2 decadal variability (e.g., Deser et al., 2012).

3 The climate data for our 2050 mid-range scenario has been downscaled from one GCM, MPI,
4 and broadly represents a 'mid-range' projection for the time slice 2045-2055. The upper-end
5 scenario represents the high end of temperature response and, for most of the country,
6 provides a sample of a severe rainfall reduction across all 19 GCMs. Climatic input data for
7 this scenario was provided by downscaled simulations from GIEH.

8 The two scenarios provided for 2100 (Renwick et al., 2013) use downscaled simulations from
9 the MPI and CCC models and also represent 'mid-range' and 'upper-end' projections. These
10 models predict an annual mean temperature change in 2090 for New Zealand of 3.0°C and
11 3.9°C, respectively (relative to 1990). Simulations were run for the nominal years 2097-2111
12 but are meant to represent general climate in approximately 100 years. These were compared
13 to a present-day baseline scenario covering the period 1997-2011.

14 Apart from climate and atmospheric CO₂ concentrations, other parameters remained
15 unchanged, thus providing an understanding of possible effects of climate change with little
16 change in agronomic systems.

17

18 **3 Results and discussion**

19 **3.1 Results**

20 Results from Biome-BGC and LURNZ for selected scenarios are shown in Tables 3 and 4 and
21 Figs. 6-9. We map production results from Biome-BGC across all of New Zealand, then
22 combine with land uses from LURNZ according to the scenario and tabulate totals nationally.
23 Appendix B contains more detailed results tabulated by region and season.

24 Our scenario results are reported relative to the baseline scenario to limit the impact of
25 potential biases in the model. The model estimates have absolute uncertainty that has not been
26 determined, and we therefore emphasize that results are expected to be most robust in terms
27 of comparisons between model scenarios. We show here the percentage difference in average
28 annual pasture production for each scenario, as compared to the baseline.

29 With the baseline model, we calculate that New Zealand's average annual pasture production
30 is approximately 800 to 900 Petajoules (PJ) of metabolisable energy available to grazing

1 animals. This number is consistent with the estimates used to compile New Zealand's most
2 recent UNFCCC emissions inventories, which was 970 PJ in 2011 (New Zealand Ministry for
3 the Environment, 2009, 2013).

4 Land-use change projections from all three modelled carbon price scenarios suggest that, in
5 general, economic factors other than carbon price dominate. Therefore, the current trends in
6 land-use change continue irrespective of carbon price: dairy is expanding, and sheep/beef is
7 contracting over time. Figure 5 shows historical land area under each of the four land-use
8 categories in LURNZ up to and including 2008 and projected future land-use area thereafter,
9 out to 2050, with no carbon price imposed. Overall, LURNZ model projections indicate that
10 carbon prices have very limited effects on land use, and current land-use change trends will
11 continue. However, these trends by themselves can be expected to have a significant effect on
12 New Zealand's pasture production, particularly through an 18-19% expansion of dairy area.
13 Figure 6 summarizes LURNZ projections for 2020 and 2050 in terms of percentage change in
14 land area and in pasture production. All land-use scenarios resulted in little change in 2020
15 relative to the baseline. In 2050, the LURNZ model projected a 4% decrease in the area of
16 sheep/beef pasture and an 18-19% increase in the area of dairy pasture, regardless of carbon
17 price. Pasture production changes were proportional to area changes. At 2050, the sheep/beef
18 decline and dairy expansion nearly offset one another, resulting in a small net increase of
19 1.5% in total national pasture production.

20 Under elevated atmospheric CO₂ concentrations in 2020 (Fig. 7), very small increases in
21 production on the order of ~0.5% were projected, which can be compared to a 10% increase
22 over present-day CO₂ concentration levels. (The A2 emissions scenario contains a 37%
23 increase in CO₂ concentration levels in 2050 and a 94% increase by 2100.) The increase in
24 production from enhanced CO₂ was slightly greater for dairy systems. No region recorded a
25 loss of production, while the largest regional increases were on the order of 1%.

26 Results from the intensification scenario (Fig. 7) indicate potential increases in pasture
27 production of 10% nationally by 2020. Results ranged between 8–14% in different regions.
28 Overall, dairy systems showed about 2-3% more of an increase than sheep/beef systems.
29 When compared to the results due to elevated CO₂ alone, the results imply the majority of
30 production increases can be attributed to the change in model parameters due to
31 intensification, with elevated CO₂ explaining only a small portion, 0.5%.

1 The modelled climate change scenarios in 2050 suggest relatively small changes at a national
2 scale but potentially more significant regional effects. Results (Fig. 8) suggest a 2% increase
3 in total national production in 2050 for the mid-range MPI scenario and a 1% increase for the
4 upper-end GIEH scenario. The dairy model shows slightly larger production increases of
5 about 4%, regardless of scenario. Sheep/beef production appears more sensitive to the
6 difference in the two scenarios, in particular to the more extreme precipitation decreases from
7 GIEH. Sheep/beef production increases by 2% with the milder MPI scenario but results in
8 virtually no change in GIEH, with the losses in some regions balancing the gains in others.
9 For comparison, we show the corresponding differences in precipitation from both models in
10 Fig. 10. The decrease in sheep/beef production in the east and north in the GIEH scenario
11 closely follows the pattern of decrease in precipitation in these regions. Despite some strong
12 regional effects, however, the model suggests that overall there will not be a large impact on
13 pasture production in 2050 from climate change within expected bounds. Seasonal results (see
14 Appendix B) indicate that in general winter production will increase and summer production
15 will decline; spring and autumn production trends vary by region.

16 Under the two 2100 scenarios, MPI is again associated with milder climate change than is
17 CCC (Fig. 9). Sheep/beef pasture production declines slightly, at around -3%, and dairy
18 production increases by 4%, resulting in almost no change nationally with current land use.
19 With the CCC model, however, production for both sheep/beef and dairy systems declined,
20 with national sheep/beef production decreasing by -13% and dairy decreasing by about -3%.
21 The decline is especially pronounced in the South Island regions of Canterbury and Otago,
22 where sheep/beef production decreases by -19% and -15% and dairy decreases by -8% and -
23 7%, respectively (see Appendix B). These results are consistent with the patterns of climate
24 change predicted for New Zealand by each climate model, with CCC predicting a larger
25 increase in temperatures and more drastic changes in rainfall (Fig. 10), especially in the spring
26 and summer months when the majority of pasture growth occurs. The eastern regions of the
27 South Island in particular are drier and warmer during these crucial growing seasons.

28 **3.2 Comparison of climatic and land management factors**

29 To better understand the effects of inputs and model modifications on the results, we compare
30 the relative significance of selected inputs and parameters in more detail for our case study.
31 Looking at all model results, climatic factors (the primary input to Biome-BGC) are clearly
32 influential in production trends. Land management factors are important as well, but our

1 analysis is limited somewhat by the fact that we have not modelled spatial variation within
2 each land-use subtype.

3 A general, comprehensive sensitivity analysis of Biome-BGC model parameters has been
4 done by White et al. (2000). The authors found that variations in C:N ratio of leaves, fire
5 mortality, and parameters relating to litter quality have the most impact on NPP in grass
6 biomes, leading to the conclusion that productivity is primarily nitrogen-limited in nonwoody
7 biomes. In comparison, our calibration reveals that the most significant effects on NPP in both
8 sheep/beef and dairy systems come from varying two different parameters, the maximum
9 stomatal conductance and the fraction of leaf N in Rubisco. This suggests that in our model,
10 New Zealand's highly-managed grasslands are primarily water- and photosynthesis-limited
11 rather than nitrogen-limited.

12 The influence of precipitation is especially visible in model results. Changes in pasture
13 production in our climate change scenarios in 2050 and 2100 closely follow the patterns of
14 change in precipitation (Fig. 10). Looking at the upper-end GIEH scenario in 2050, decreases
15 of 5-10% in precipitation along the East Coast of New Zealand correspond to a 2-3% regional
16 decrease sheep/beef pasture production. Dairy pasture appears less sensitive to changes in
17 precipitation; one explanation for this could be that the higher nitrogen status of dairy systems
18 leads to an increase in photosynthetic water use efficiency.

19 Seasonal patterns of growth also play an important role. In our climate change scenarios,
20 winter production tends to increase while summer production decreases, as one would expect
21 from an overall average temperature increase. Spring and autumn production trends vary
22 regionally, with no consistent national pattern. Spring and summer in particular are crucial
23 growing seasons in our model as well as for pasture production historically. A breakdown of
24 seasonal rainfall (not shown) indicates that dry summers might drastically reduce production
25 even if the remaining seasons have normal levels of precipitation.

26 An examination of model results with and without the wind correction factor applied to VPD
27 input indicates that the modified VPD reduces overall national production by 3-4% (not
28 shown), although the exact amount varies by region. Hence the general effect of including
29 wind in our simulations is to decrease plant productivity, as is expected from the elevated
30 water loss that wind induces. Although our climate change scenarios did not modify wind
31 speed from present-day values, global climate models predict that the mean westerly winds
32 over New Zealand will increase, especially over the South Island in winter and spring. If local

1 wind strength does indeed increase in the future, it could cause pasture production to decrease
2 more than we have estimated with our scenarios.

3 While the model appears most sensitive to weather inputs in the scenarios we considered,
4 parameters involved in grazing and land management are also notable. CO₂ fertilisation is, on
5 the whole, a small effect, resulting in only a 0.5% in production from a 10% increase in CO₂
6 atmospheric concentrations. Nevertheless, with concentrations projected to increase by 90%
7 or more by 2100, it will become much more significant when compared to present day. The
8 model does respond strongly to increases in the plant mortality and nitrogen fixation
9 parameters, having a relatively large positive effect on production in our intensification
10 scenario. Increasing pasture utilisation by 9% and 6% and nitrogen fixation (or fertiliser
11 application) by 16% and 19% for sheep/beef and dairy systems, respectively, produce
12 modelled increases in production of 9.5% and 12.5%. This should be interpreted with caution,
13 however; our model is a fairly simplistic representation of intensification, as we have treated
14 all land within each subtype as having equal utilisation and equal potential for production
15 gains. In practice, gains would likely be limited due to resource availability, environmental
16 concerns and the considerable spatial variation in land quality. In addition, we have not
17 explicitly considered changes in carbon-cycle feedbacks and other biophysical effects due to
18 land-use change and intensification. Other studies have demonstrated that land-use change
19 affects characteristics such as albedo and radiative forcing (Kirschbaum et al., 2011), carbon
20 storage (Bala et al., 2007), and water yield (Beets and Oliver, 2007). The simulation of these
21 effects is beyond the scope of this study but could be considered in future work.

22 An advantage of the DLUCS methodology is the ability to rapidly sample results from
23 different models and model updates, assuming that strongly interrelated global change issues
24 such as N status and enhanced pCO₂ can be handled within model projections. With the recent
25 release of the IPCC Fifth Assessment Report (AR5), updated climate projections can be
26 incorporated in our simulations through dynamically downscaled weather input to the Biome-
27 BGC model to more fully explore the trends that we discuss here. The possibility of including
28 downscaled changes in wind strength now exists as well, which is currently lacking in our
29 climate change simulations. Output from other biogeochemistry models (including those
30 coupled to GCMs) and improved versions or alternatives replacing Biome-BGC output can be
31 incorporated in a similar way, as well as new developments in LURNZ that add spatial detail
32 and allow for variations in land productivity and carrying capacity (Timar and Kerr, 2014).

1

2 **4 Conclusions**

3 Examining all scenarios modelled here, our results suggest a slight increase in pasture
4 production by 2020 is likely, and increases of 10–15% are plausible. The outlook for 2050 is
5 also favourable given the scenarios considered. The projected continued conversion of land to
6 high-intensity dairy farming will likely increase total national production. Although climate
7 change could have an adverse effect on particular regions by 2050, our modelling estimates a
8 small overall national increase. CO₂ fertilisation effects could also contribute to a slight
9 increase. Projected pasture production in 2100 shows a much larger range of possible
10 outcomes. We find significant differences in the impact on pasture production using weather
11 input from two different GCMs. The CCC model results in a pronounced decline in both
12 sheep/beef and dairy pasture production, while the MPI model shows only a slight decrease in
13 sheep/beef and an increase in dairy production. This highlights that the severity of warming
14 will determine the degree of impact on pasture production for both sheep/beef and dairy
15 agriculture in New Zealand.

16 Our results demonstrate the capability of the Biome-BGC model to provide useful production
17 data when integrated with global change scenarios, including results from the LURNZ
18 model's estimates of land-use change. With downscaled weather input from GCMs, the model
19 infrastructure enables the investigation of regional effects of projected climate change. The
20 Biome-BGC model also offers the potential to model forest ecosystems (with the model's
21 built-in forest modules) and compare productivity across forest and pasture land uses in future
22 studies of global change. The advantage of our approach is the flexibility of the model
23 components and the variety of future change scenarios that we are able to explore with
24 relative ease: by modelling national production for all locations coupled with climate
25 projections or other initial input, we can quickly re-sample the national grid for any modelled
26 land-use change or economic scenario.

27 Further work to calibrate the Biome-BGC model to New Zealand conditions is needed to
28 refine and confirm results. Iterative improvements in modelling and experimental approaches
29 will be required to provide robust results given the strong interactions and feedbacks in
30 productive ecosystems. One area to investigate is the interaction between climate change and
31 elevated CO₂ and water and nitrogen availability, and the differences between time-dependent
32 and quasi-steady-state model results. These effects can be constrained by data (for example,

1 eddy covariance towers and FACE experiments) using our modelling approach and have
2 substantial implications for the seasonal cycle of pasture supply, responses to drought and our
3 ability to correctly characterize relative impacts of climate on sheep/beef versus dairy
4 systems. This includes correct estimation of the benefits and limitations of irrigation in both
5 sheep/beef and dairy pasture. Additionally, the simulation of extreme events (e.g. droughts
6 and floods) and new scenarios from the IPCC AR5 with the Biome-BGC model will add to
7 our understanding of the impacts of future climate change. There is also potential for
8 integration with other earth system models. For example, integration with hydrology models
9 would allow us to examine the effects of climate and land-use change on water supply in New
10 Zealand (see, for example, Gerten et al., 2008 and Rockstrom et al., 2009). While achieving a
11 fully integrated assessment model remains challenging, the DLUCS linkage of models
12 presented here provides a useful methodology to investigate global change, with the ability to
13 generate realistic scenario results on timescales required for policy formulation.

14

15 **Appendix A: Biome-BGC parameters**

16 Eco-physiological parameters used in the Biome-BGC model for dairy and sheep/beef
17 ecosystems. The parameters that were adjusted are marked with a footnote. All other
18 parameters were set to C3 grasses default values provided with the Biome-BGC v4.2 Final
19 Release. Full descriptions of the parameters are contained in the model documentation.

20 Table A1. Biome-BGC model parameters.

Parameter Description	Sheep/Beef Baseline	Dairy Baseline	C3 grass default	Type
annual whole-plant mortality fraction ^{a,b}	0.226	0.722	0.1	(1/yr)
annual fire mortality fraction ^b	0.1	0.2	0.1	(1/yr)
(ALLOCATION) new fine root C : new leaf C ^b	1.43	0.246	1	(ratio)
(ALLOCATION) current growth proportion ^b	0.84	0.424	0.5	(prop.)
C:N of leaves ^b	24.0	24.1	24.0	(kgC/kgN)
C:N of leaf litter, after retranslocation ^b	49.0	49.0	49.0	(kgC/kgN)

^a Whole-plant mortality was calculated as (Utilisation) * (above-ground growth)/ (above-ground + below-ground growth). The ratio below-ground/above-ground is given by the parameter new fine root C : new leaf C.

^b Parameters adjusted during calibration

C:N of fine roots ^b	42.0	42.9	42.0	(kgC/kgN)
canopy average specific leaf area (projected area basis) ^b	45.0	45.0	45.0	(m ² /kgC)
fraction of leaf N in Rubisco ^b	0.19	0.05	0.15	(DIM)
maximum stomatal conductance (projected area basis) ^b	0.00534	0.00375	0.005	(m/s)
boundary layer conductance (projected area basis) ^b	0.07	0.0202	0.04	(m/s)
1 = WOODY 0 = NON-WOODY	0	0	0	(flag)
1 = EVERGREEN 0 = DECIDUOUS	0	0	0	(flag)
1 = C3 PSN 0 = C4 PSN	1	1	1	(flag)
1 = MODEL PHENOLOGY 0 = USER-SPECIFIED PHENOLOGY	0	0	0	(flag)
yearday to start new growth (phenology flag = 0)	0	0	0	(yday)
yearday to end litterfall (phenology flag = 0)	364	364	364	(yday)
transfer growth period as fraction of growing	1	1	1	(prop.)
litterfall as fraction of growing season	1	1	1	(prop.)
annual leaf and fine root turnover fraction	1	1	1	(1/yr)
annual live wood turnover fraction	0	0	0	(1/yr)
(ALLOCATION) new stem C : new leaf C	0	0	0	(ratio)
(ALLOCATION) new live wood C : new total wood C	0	0	0	(ratio)
(ALLOCATION) new croot C : new stem C	0	0	0	(ratio)
C:N of live wood	0	0	0	(kgC/kgN)
C:N of dead wood	0	0	0	(kgC/kgN)
leaf litter labile proportion	0.39	0.39	0.39	(DIM)
leaf litter cellulose proportion	0.44	0.44	0.44	(DIM)
leaf litter lignin proportion	0.17	0.17	0.17	(DIM)
fine root labile proportion	0.30	0.30	0.30	(DIM)
fine root cellulose proportion	0.45	0.45	0.45	(DIM)
fine root lignin proportion	0.25	0.25	0.25	(DIM)
dead wood cellulose proportion	0.75	0.75	0.75	(DIM)
dead wood lignin proportion	0.25	0.25	0.25	(DIM)
canopy water interception coefficient	0.021	0.021	0.021	(1/LAI/d)
canopy light extinction coefficient	0.6	0.6	0.6	(DIM)
all-sided to projected leaf area ratio	2.0	2.0	2.0	(DIM)

ratio of shaded SLA:sunlit SLA	2.0	2.0	2.0	(DIM)
cuticular conductance (projected area basis)	0.00001	0.00001	0.00001	(m/s)
leaf water potential: start of conductance reduction	-0.6	-0.6	-0.6	(MPa)
leaf water potential: complete conductance reduction	-2.3	-2.3	-2.3	(MPa)
vapor pressure deficit: start of conductance reduction	930	930	930	(Pa)
vapor pressure deficit: complete conductance reduction	4100	4100	4100	(Pa)

1

2 **Appendix B: Pasture Production Results by Region and Season**

3 The following tables show pasture production results organised by New Zealand's 16 regions
 4 and by season, in terms of metabolisable energy (the estimated amount of energy that is
 5 available to grazing animals from pasture).

6

1 Table B1. Baseline average production by region and season over the period 2001-2009 in
 2 units of metabolisable energy.

Region	Area (kha)	Winter	Spring	Summer	Autumn	Average	Total ME (TJ)
Average Production (MJ ha ⁻¹ d ⁻¹)							
Sheep & Beef							
Northland	239	138	423	296	199	264	22963
Auckland	85	129	445	304	168	262	8158
Waikato	473	112	421	368	201	276	47532
Bay of Plenty	66	108	402	366	219	274	6580
Gisborne	327	106	431	321	200	264	31582
Manawatu-Wanganui	936	92	415	373	185	266	90856
Hawkes Bay	586	94	398	212	179	220	47121
Taranaki	125	105	398	412	214	282	12850
Tasman	57	86	406	268	171	233	4841
Marlborough	310	59	310	230	133	183	20703
Westland	32	81	355	378	204	255	2949
Wellington	281	95	440	249	156	235	24064
Nelson City	2	87	417	274	142	230	208
Canterbury	1824	54	307	211	131	176	117080
Otago	1738	46	296	248	134	181	114986
Southland	698	62	376	370	176	246	62653
National	7778	73	356	279	159	217	615126
Dairy							
Northland	172	221	617	464	376	420	26334
Auckland	53	209	629	504	342	421	8104
Waikato	487	184	628	510	364	422	74904
Bay of Plenty	80	181	613	540	400	434	12611
Gisborne	3	176	580	536	392	421	384
Manawatu-Wanganui	127	149	602	538	314	401	18595
Hawkes Bay	16	136	581	454	334	376	2159
Taranaki	215	168	594	615	394	443	34784
Tasman	27	139	582	548	357	406	4043
Marlborough	9	126	549	456	303	358	1210
Westland	52	124	532	582	373	402	7690
Wellington	30	142	612	433	298	371	4016
Nelson City	0	121	517	339	295	318	49
Canterbury	119	103	513	362	281	315	13723
Otago	62	88	523	488	287	346	7799
Southland	106	93	529	564	332	379	14609
National	1557	163	594	515	353	406	231014
Combined national	9335						846140

1 Table B2. Baseline average production by region and season over the period 1997-2011 in
 2 units of metabolisable energy.

Region	Area (kha)	Winter (MJ ha ⁻¹ d ⁻¹)	Spring	Summer	Autumn	Average	Total ME (TJ)
Sheep & Beef							
Northland	239	131	435	247	151	241	20999
Auckland	85	123	452	266	134	244	7601
Waikato	473	105	446	298	143	248	42743
Bay of Plenty	66	102	416	298	174	248	5951
Gisborne	327	103	422	341	196	266	31745
Manawatu-Wanganui	936	87	438	288	137	237	81024
Hawkes Bay	586	88	387	188	153	204	43661
Taranaki	125	101	421	402	177	275	12535
Tasman	57	83	403	231	152	217	4524
Marlborough	310	58	305	253	134	187	21235
Westland	32	79	369	308	170	231	2677
Wellington	281	89	437	216	129	218	22317
Nelson City	2	83	408	258	117	217	196
Canterbury	1824	51	291	176	122	160	106386
Otago	1738	43	272	194	119	157	99507
Southland	698	57	369	268	146	210	53466
National	7778	69	350	230	135	196	556567
Dairy							
Northland	172	213	617	426	325	395	24815
Auckland	53	199	637	477	293	402	7733
Waikato	487	170	627	470	308	394	69953
Bay of Plenty	80	169	607	468	354	399	11617
Gisborne	3	173	577	539	380	418	381
Manawatu-Wanganui	127	137	604	470	261	368	17063
Hawkes Bay	16	126	561	409	303	350	2007
Taranaki	215	168	598	620	393	445	34946
Tasman	27	133	578	510	325	386	3843
Marlborough	9	118	529	422	276	337	1136
Westland	52	123	537	547	348	389	7434
Wellington	30	131	599	369	264	341	3690
Nelson City	0.4	114	497	333	266	302	47
Canterbury	119	89	462	289	246	272	11835
Otago	62	78	485	399	251	304	6833
Southland	106	85	525	497	282	347	13384
National	1557	153	588	472	311	381	216717
Combined national	9335						773284

1 Table B3. Elevated CO₂ 2020.

Region	Area (kha)	Winter	Spring	Summer	Autumn	Average	Total ME (TJ)	Change (%)
Average Production (MJ ha ⁻¹ d ⁻¹)								
Sheep & Beef								
Northland	239	140	423	297	199	265	23035	0.3%
Auckland	85	131	445	305	169	263	8188	0.4%
Waikato	473	113	421	369	201	276	47622	0.2%
Bay of Plenty	66	108	402	367	219	274	6590	0.2%
Gisborne	327	107	431	323	200	265	31680	0.3%
Manawatu-Wanganui	936	92	415	374	185	267	91042	0.2%
Hawkes Bay	586	95	398	213	180	222	47378	0.5%
Taranaki	125	105	397	412	214	282	12859	0.1%
Tasman	57	87	407	269	172	234	4861	0.4%
Marlborough	310	59	310	231	134	184	20807	0.5%
Westland	32	82	355	379	205	255	2955	0.2%
Wellington	281	96	441	250	157	236	24186	0.5%
Nelson City	2	88	418	275	144	231	209	0.5%
Canterbury	1824	55	308	213	133	177	117777	0.6%
Otago	1738	47	297	249	135	182	115538	0.5%
Southland	698	63	376	371	176	247	62803	0.2%
National	7778	74	356	280	160	218	617531	0.4%
Dairy								
Northland	172	225	620	466	377	422	26467	0.5%
Auckland	53	213	632	505	342	423	8143	0.5%
Waikato	487	187	631	512	364	424	75290	0.5%
Bay of Plenty	80	183	616	542	400	435	12664	0.4%
Gisborne	3	178	583	537	392	423	386	0.4%
Manawatu-Wanganui	127	151	607	539	314	403	18695	0.5%
Hawkes Bay	16	138	585	456	335	378	2172	0.6%
Taranaki	215	171	597	615	394	444	34915	0.4%
Tasman	27	141	585	550	357	408	4061	0.4%
Marlborough	9	128	553	458	304	361	1218	0.6%
Westland	52	126	535	583	373	404	7724	0.4%
Wellington	30	145	617	435	299	374	4046	0.8%
Nelson City	0	123	522	342	296	321	50	0.8%
Canterbury	119	105	519	365	283	318	13851	0.9%
Otago	62	90	527	489	288	349	7846	0.6%
Southland	106	94	533	564	332	381	14675	0.5%
National	1557	166	598	517	354	409	232203	0.5%
Combined national	9335						849734	0.4%

2

3

1 Table B4. Intensification 2020.

Region	Area (kha)	Winter Average	Spring Production	Summer (MJ ha-1 d-1)	Autumn	Average	Total ME (TJ)	Change (%)
Sheep & Beef								
Northland	239	153	476	318	217	291	25337	10.3%
Auckland	85	142	499	320	178	285	8878	8.8%
Waikato	473	126	482	406	217	308	53058	11.6%
Bay of Plenty	66	122	461	405	240	307	7381	12.2%
Gisborne	327	117	485	348	218	292	34886	10.5%
Manawatu-Wanganui	936	102	474	411	199	297	101291	11.5%
Hawkes Bay	586	103	438	223	194	240	51212	8.7%
Taranaki	125	119	462	471	235	322	14652	14.0%
Tasman	57	94	451	287	185	254	5286	9.2%
Marlborough	310	64	341	247	143	199	22533	8.8%
Westland	32	90	401	418	223	283	3279	11.2%
Wellington	281	104	488	261	169	255	26180	8.8%
Nelson City	2	96	462	290	149	250	225	8.4%
Canterbury	1824	59	334	224	141	190	126240	7.8%
Otago	1738	51	325	265	144	196	124421	8.2%
Southland	698	68	423	404	189	271	69003	10.1%
National	7778	80	396	301	172	237	673861	9.5%
Dairy								
Northland	172	246	698	520	428	473	29678	12.7%
Auckland	53	233	712	565	387	474	9136	12.7%
Waikato	487	204	708	574	411	474	84222	12.4%
Bay of Plenty	80	202	694	610	457	491	14268	13.1%
Gisborne	3	197	659	612	449	479	437	13.9%
Manawatu-Wanganui	127	164	675	605	353	449	20850	12.1%
Hawkes Bay	16	149	649	505	377	420	2412	11.7%
Taranaki	215	189	673	708	451	505	39681	14.1%
Tasman	27	154	654	621	406	459	4565	12.9%
Marlborough	9	139	614	515	340	402	1357	12.2%
Westland	52	136	595	662	423	454	8675	12.8%
Wellington	30	155	680	478	335	412	4461	11.1%
Nelson City	0	131	571	375	327	351	54	10.4%
Canterbury	119	112	563	400	312	347	15117	10.2%
Otago	62	96	578	543	319	384	8647	10.9%
Southland	106	101	590	639	369	425	16355	12.0%
National	1557	181	668	581	399	457	259915	12.5%
Combined national	9335						933776	10.4%

2

3

1 Table B5. Climate change 2050 (GIEH model).

Region	Area (kha)	Winter Average	Spring Production	Summer (MJ ha-1 d-1)	Autumn	Average	Total ME (TJ)	Change (%)
Sheep & Beef								
Northland	239	148	421	288	204	265	23103	0.6%
Auckland	85	140	442	299	178	265	8262	1.3%
Waikato	473	120	421	369	205	279	48129	1.3%
Bay of Plenty	66	115	405	361	222	276	6637	0.9%
Gisborne	327	116	426	305	221	267	31885	1.0%
Manawatu-Wanganui	936	98	417	372	191	269	91999	1.3%
Hawkes Bay	586	104	392	201	207	226	48335	2.6%
Taranaki	125	111	399	412	214	284	12943	0.7%
Tasman	57	94	413	266	175	237	4929	1.8%
Marlborough	310	65	315	218	138	184	20860	0.8%
Westland	32	87	358	377	203	256	2967	0.6%
Wellington	281	103	442	239	174	239	24545	2.0%
Nelson City	2	94	423	261	149	232	209	0.7%
Canterbury	1824	61	311	209	142	181	120235	2.7%
Otago	1738	52	307	250	139	187	118549	3.1%
Southland	698	67	384	373	177	250	63704	1.7%
National	7778	80	360	276	168	221	627293	2.0%
Dairy								
Northland	172	250	630	456	385	430	26999	2.5%
Auckland	53	237	643	500	351	433	8332	2.8%
Waikato	487	210	648	517	378	438	77837	3.9%
Bay of Plenty	80	205	632	538	412	447	12995	3.0%
Gisborne	3	201	596	529	410	434	396	3.1%
Manawatu-Wanganui	127	169	632	542	321	416	19309	3.8%
Hawkes Bay	16	161	604	438	372	393	2258	4.6%
Taranaki	215	188	618	618	396	455	35757	2.8%
Tasman	27	158	609	552	359	419	4172	3.2%
Marlborough	9	148	575	449	309	370	1250	3.3%
Westland	52	139	556	584	374	413	7899	2.7%
Wellington	30	165	642	429	319	389	4207	4.8%
Nelson City	0	142	545	334	311	333	52	4.7%
Canterbury	119	126	539	369	309	336	14629	6.6%
Otago	62	103	555	491	299	362	8153	4.5%
Southland	106	107	563	569	338	394	15181	3.9%
National	1557	186	617	518	364	421	239427	3.6%
Combined national	9335						866720	2.4%

2

3

1 Table B6. Climate change 2050 (MPI model).

Region	Area (kha)	Winter Average	Spring Production	Summer (MJ ha-1 d-1)	Autumn	Average	Total ME (TJ)	Change (%)
Sheep & Beef								
Northland	239	154	414	266	210	261	22729	-1.0%
Auckland	85	146	441	265	182	259	8063	-1.2%
Waikato	473	128	423	355	204	277	47869	0.7%
Bay of Plenty	66	124	406	345	225	275	6611	0.5%
Gisborne	327	124	422	284	222	263	31424	-0.5%
Manawatu-Wanganui	936	105	420	353	189	267	91174	0.4%
Hawkes Bay	586	111	365	189	213	219	46914	-0.4%
Taranaki	125	118	402	408	211	285	12970	0.9%
Tasman	57	101	407	244	177	232	4835	-0.1%
Marlborough	310	69	305	199	140	178	20180	-2.5%
Westland	32	91	359	372	201	256	2962	0.4%
Wellington	281	110	435	208	178	233	23850	-0.9%
Nelson City	2	101	415	228	152	224	202	-2.6%
Canterbury	1824	65	298	196	145	176	117198	0.1%
Otago	1738	55	305	235	139	184	116504	1.3%
Southland	698	71	389	360	174	249	63274	1.0%
National	7778	85	354	260	169	217	616759	0.3%
Dairy								
Northland	172	267	619	424	401	428	26855	2.0%
Auckland	53	255	645	463	358	430	8284	2.2%
Waikato	487	232	644	495	385	439	78004	4.1%
Bay of Plenty	80	227	630	514	423	449	13044	3.4%
Gisborne	3	220	592	512	415	435	397	3.3%
Manawatu-Wanganui	127	187	639	514	321	415	19268	3.6%
Hawkes Bay	16	182	586	409	388	391	2246	4.0%
Taranaki	215	205	625	615	397	460	36167	4.0%
Tasman	27	175	614	533	366	422	4198	3.8%
Marlborough	9	167	566	428	317	369	1247	3.0%
Westland	52	151	563	583	378	419	8003	4.1%
Wellington	30	183	642	389	328	386	4174	3.9%
Nelson City	0	163	527	319	324	333	52	4.8%
Canterbury	119	144	508	350	329	333	14491	5.6%
Otago	62	115	562	468	302	362	8146	4.4%
Southland	106	119	576	562	341	400	15393	5.4%
National	1557	204	615	498	372	422	239967	3.9%
Combined national	9335						856726	1.3%

2

3

1 Table B7. Climate change 2100 (CCC model).

Region	Area (kha)	Winter	Spring	Summer	Autumn	Average	Total ME (TJ)	Change (%)
Sheep & Beef								
Northland	239	172	390	127	211	225	19603	-6.6%
Auckland	85	156	408	88	179	207	6470	-14.9%
Waikato	473	152	443	113	158	216	37346	-12.6%
Bay of Plenty	66	151	403	112	213	220	5289	-11.1%
Gisborne	327	157	411	216	235	255	30427	-4.2%
Manawatu-Wanganui	936	134	433	102	147	204	69621	-14.1%
Hawkes Bay	586	138	299	93	221	188	40090	-8.2%
Taranaki	125	144	471	257	133	251	11439	-8.7%
Tasman	57	132	390	90	157	192	3999	-11.6%
Marlborough	310	94	310	116	130	162	18406	-13.3%
Westland	32	107	384	205	135	208	2404	-10.2%
Wellington	281	138	388	59	163	187	19165	-14.1%
Nelson City	2	119	384	82	148	183	165	-15.5%
Canterbury	1824	80	243	80	116	130	86316	-18.9%
Otago	1738	68	259	109	98	134	84713	-14.9%
Southland	698	84	375	168	126	188	47947	-10.3%
National	7778	105	326	112	139	170	483402	-13.1%
Dairy								
Northland	172	327	577	275	419	400	25077	1.1%
Auckland	53	311	644	249	367	393	7560	-2.2%
Waikato	487	289	635	249	347	380	67506	-3.5%
Bay of Plenty	80	294	606	259	427	396	11524	-0.8%
Gisborne	3	294	588	435	437	439	400	5.0%
Manawatu-Wanganui	127	242	645	238	266	348	16136	-5.4%
Hawkes Bay	16	248	510	249	409	354	2032	1.3%
Taranaki	215	268	697	580	318	466	36594	4.7%
Tasman	27	241	638	346	318	386	3837	-0.2%
Marlborough	9	219	550	228	294	323	1090	-4.1%
Westland	52	208	617	435	305	391	7470	0.5%
Wellington	30	244	602	173	302	330	3574	-3.1%
Nelson City	0	211	484	160	321	294	46	-2.7%
Canterbury	119	176	387	158	276	249	10859	-8.2%
Otago	62	146	488	261	239	283	6381	-6.6%
Southland	106	153	613	380	257	351	13507	0.9%
National	1557	258	608	306	331	376	213593	-1.4%
Combined national	9335						696995	-9.9%

2

3

1 Table B8. Climate change 2100 (MPI model).

Region	Area (kha)	Winter	Spring	Summer	Autumn	Average	Total ME (TJ)	Change (%)
Sheep & Beef								
Northland	239	168	399	183	180	233	20258	-3.5%
Auckland	85	153	422	172	157	226	7047	-7.3%
Waikato	473	142	446	218	149	239	41195	-3.6%
Bay of Plenty	66	140	410	199	189	234	5636	-5.3%
Gisborne	327	148	406	262	219	259	30890	-2.7%
Manawatu-Wanganui	936	123	440	209	146	229	78330	-3.3%
Hawkes Bay	586	131	320	147	196	199	42439	-2.8%
Taranaki	125	127	439	359	160	271	12350	-1.5%
Tasman	57	120	399	185	159	216	4490	-0.7%
Marlborough	310	87	306	209	142	186	21081	-0.7%
Westland	32	75	363	279	158	219	2532	-5.4%
Wellington	281	132	407	153	158	212	21775	-2.4%
Nelson City	2	112	393	186	134	206	186	-4.7%
Canterbury	1824	78	269	153	131	158	105037	-1.3%
Otago	1738	60	273	165	113	153	97026	-2.5%
Southland	698	68	375	228	137	202	51436	-3.8%
National	7778	97	338	185	143	191	541710	-2.7%
Dairy								
Northland	172	309	582	346	383	405	25409	2.4%
Auckland	53	292	646	363	337	410	7886	2.0%
Waikato	487	262	642	381	334	405	71941	2.8%
Bay of Plenty	80	264	613	359	396	408	11864	2.1%
Gisborne	3	270	581	470	421	435	397	4.3%
Manawatu-Wanganui	127	218	653	378	278	382	17720	3.8%
Hawkes Bay	16	224	534	335	378	368	2111	5.2%
Taranaki	215	242	651	611	382	471	37034	6.0%
Tasman	27	212	629	458	342	410	4080	6.1%
Marlborough	9	204	552	360	307	356	1202	5.8%
Westland	52	161	581	516	346	401	7665	3.1%
Wellington	30	225	621	298	309	363	3932	6.6%
Nelson City	0.4	195	502	286	310	323	50	6.9%
Canterbury	119	174	434	287	302	299	13031	10.1%
Otago	62	132	526	366	267	323	7266	6.3%
Southland	106	126	597	456	279	365	14044	4.9%
National	1557	235	608	408	336	397	225631	4.1%
Combined national		9335					767341	-0.8%

2

3

1 Table B9. Land-use change, 2020 CO₂ = \$0

Region	Area (kha)	Change (%)	Winter Average	Spring Production	Summer (MJ ha-1 d-1)	Autumn	Average	Total ME (TJ)	Change (%)
Sheep & Beef									
Northland	248	3.9%	138	423	295	198	263	23838	3.8%
Auckland	83	-2.4%	129	446	305	168	262	7976	-2.2%
Waikato	479	1.4%	112	421	371	201	276	48332	1.7%
Bay of Plenty	74	12.5%	108	403	371	219	275	7444	13.1%
Gisborne	326	-0.4%	106	431	322	200	265	31487	-0.3%
Manawatu-Wanganui	909	-2.9%	91	414	373	185	266	88262	-2.9%
Hawkes Bay	583	-0.4%	94	397	212	179	221	46935	-0.4%
Taranaki	123	-1.7%	105	397	412	214	282	12618	-1.8%
Tasman	61	7.7%	86	405	274	173	235	5259	8.6%
Marlborough	312	0.4%	59	310	231	133	183	20827	0.6%
Westland	39	23.8%	83	358	381	207	257	3686	25.0%
Wellington	281	0.0%	95	440	249	156	235	24065	0.0%
Nelson City	3	4.0%	87	413	269	142	228	214	3.1%
Canterbury	1820	-0.2%	54	307	211	131	176	116850	-0.2%
Otago	1741	0.1%	46	296	248	134	181	115185	0.2%
Southland	684	-1.9%	62	375	370	175	246	61348	-2.1%
National	7766	-0.2%	73	356	279	159	217	614327	-0.13%
Dairy									
Northland	162	-5.6%	221	617	464	377	420	24875	-5.5%
Auckland	54	2.6%	209	628	502	342	420	8302	2.4%
Waikato	482	-1.0%	186	629	506	363	421	74063	-1.1%
Bay of Plenty	71	-10.3%	183	616	532	402	433	11299	-10.4%
Gisborne	5	95.0%	182	569	472	377	400	712	85.4%
Manawatu-Wanganui	157	23.1%	150	603	541	313	402	22953	23.4%
Hawkes Bay	18	17.3%	136	582	422	330	368	2476	14.7%
Taranaki	217	0.9%	168	595	615	394	443	35106	0.9%
Tasman	22	-18.0%	142	586	549	358	409	3336	-17.5%
Marlborough	7	-20.5%	125	542	428	293	347	930	-23.1%
Westland	42	-20.6%	123	533	584	374	403	6122	-20.4%
Wellington	30	0.3%	143	614	428	296	370	4021	0.1%
Nelson City	0	-41.2%	124	533	350	300	327	30	-39.5%
Canterbury	121	1.4%	104	515	361	282	315	13938	1.6%
Otago	58	-5.3%	89	526	488	287	347	7409	-5.0%
Southland	119	12.3%	93	529	566	334	381	16480	12.8%
National	1566	0.6%	163	595	514	352	406	232053	0.45%
Combined national	9332							846380	0.03%

2

3

1 Table B10. Land-use change, 2020 CO₂ = \$50 per tonne.

Region	Area (kha)	Change (%)	Winter Average	Spring Production	Summer (MJ ha-1 d-1)	Autumn	Average	Total ME (TJ)	Change (%)
Sheep & Beef									
Northland	248	4.0%	138	423	295	198	263	23856	3.9%
Auckland	84	-2.0%	129	446	305	168	262	8007	-1.8%
Waikato	481	1.7%	112	421	371	201	276	48463	2.0%
Bay of Plenty	74	12.7%	108	403	371	219	275	7455	13.3%
Gisborne	327	-0.2%	106	431	322	200	265	31541	-0.1%
Manawatu-Wanganui	913	-2.4%	91	414	373	185	266	88641	-2.4%
Hawkes Bay	583	-0.4%	94	397	212	179	221	46954	-0.4%
Taranaki	123	-1.6%	105	397	412	214	282	12638	-1.6%
Tasman	61	7.7%	86	405	274	173	235	5259	8.6%
Marlborough	311	0.3%	59	310	231	133	183	20820	0.6%
Westland	39	23.8%	83	358	381	207	257	3686	25.0%
Wellington	281	0.0%	95	440	249	156	235	24078	0.1%
Nelson City	3	4.0%	87	413	269	142	228	214	3.1%
Canterbury	1820	-0.2%	54	307	211	131	176	116849	-0.2%
Otago	1741	0.1%	46	296	248	134	181	115195	0.2%
Southland	689	-1.2%	62	375	370	176	246	61805	-1.4%
National	7777	0.0%	73	356	279	159	217	615462	0.05%
Dairy									
Northland	162	-5.8%	221	617	464	377	420	24830	-5.7%
Auckland	54	1.8%	209	628	502	342	420	8237	1.6%
Waikato	480	-1.3%	186	629	506	363	421	73827	-1.4%
Bay of Plenty	71	-10.6%	183	616	532	402	433	11258	-10.7%
Gisborne	4	63.0%	180	563	455	373	393	584	52.0%
Manawatu-Wanganui	152	19.8%	150	603	541	313	402	22322	20.0%
Hawkes Bay	18	16.1%	136	581	421	330	367	2447	13.3%
Taranaki	217	0.8%	168	595	615	394	443	35083	0.9%
Tasman	22	-18.0%	142	586	549	358	409	3336	-17.5%
Marlborough	7	-20.5%	125	542	428	293	347	930	-23.1%
Westland	42	-20.6%	123	533	584	374	403	6122	-20.4%
Wellington	30	-0.3%	143	614	428	296	370	3997	-0.5%
Nelson City	0	-41.2%	124	533	350	300	327	30	-39.5%
Canterbury	120	0.8%	104	515	361	282	315	13850	0.9%
Otago	58	-5.9%	89	526	488	287	347	7361	-5.6%
Southland	114	7.7%	93	530	566	334	381	15811	8.2%
National	1552	-0.3%	163	595	513	352	406	230025	-0.43%
Combined national	9330							845487	-0.08%

2

3

1 Table B11. Land-use change, 2020 CO₂ = \$100 per tonne.

Region	Area (kha)	Change (%)	Winter Average	Spring Production	Summer (MJ ha-1 d-1)	Autumn	Average	Total ME (TJ)	Change (%)
Sheep & Beef									
Northland	249	4.5%	138	423	295	198	263	23974	4.4%
Auckland	84	-1.6%	129	446	305	168	262	8044	-1.4%
Waikato	485	2.7%	112	421	372	201	276	48968	3.0%
Bay of Plenty	76	15.1%	108	403	371	219	275	7621	15.8%
Gisborne	327	-0.2%	106	431	322	200	265	31546	-0.1%
Manawatu-Wanganui	915	-2.2%	91	414	373	185	266	88850	-2.2%
Hawkes Bay	584	-0.3%	94	397	212	179	221	47003	-0.3%
Taranaki	123	-1.2%	105	397	412	214	282	12691	-1.2%
Tasman	62	8.2%	86	405	274	173	235	5288	9.2%
Marlborough	311	0.3%	59	310	231	133	183	20813	0.5%
Westland	40	26.5%	83	358	381	207	257	3766	27.7%
Wellington	281	0.1%	95	440	249	156	235	24094	0.1%
Nelson City	3	4.0%	87	413	269	142	228	214	3.1%
Canterbury	1819	-0.3%	54	307	211	131	176	116801	-0.2%
Otago	1740	0.1%	47	296	248	134	181	115184	0.2%
Southland	689	-1.2%	62	375	370	176	246	61827	-1.3%
National	7789	0.1%	73	356	279	159	217	616684	0.25%
Dairy									
Northland	161	-6.5%	221	617	464	377	420	24633	-6.5%
Auckland	53	1.0%	209	628	502	342	420	8180	0.9%
Waikato	475	-2.3%	186	629	505	363	421	73025	-2.5%
Bay of Plenty	70	-12.7%	184	616	531	402	433	10994	-12.8%
Gisborne	4	61.0%	180	563	452	372	392	576	49.8%
Manawatu-Wanganui	150	18.0%	150	603	541	312	402	21999	18.3%
Hawkes Bay	18	12.6%	137	582	419	330	367	2370	9.8%
Taranaki	216	0.5%	169	595	615	394	443	34961	0.5%
Tasman	22	-19.1%	142	586	549	358	409	3290	-18.6%
Marlborough	7	-21.9%	125	542	428	293	347	915	-24.4%
Westland	41	-22.5%	123	533	585	374	404	5978	-22.3%
Wellington	29	-0.8%	143	615	427	296	370	3973	-1.1%
Nelson City	0	-41.2%	124	533	350	300	327	30	-39.5%
Canterbury	120	0.3%	104	515	361	282	315	13788	0.5%
Otago	58	-6.2%	89	526	488	287	347	7331	-6.0%
Southland	113	7.4%	93	530	566	334	381	15761	7.9%
National	1537	-1.3%	163	595	513	352	406	227802	-1.39%
Combined national	9326							844486	-0.20%

2

3

1 Table B12. Land-use change, 2050 CO₂ = \$0.

Region	Area (kha)	Change (%)	Winter Average	Spring Production	Summer (MJ ha ⁻¹ d ⁻¹)	Autumn	Average	Total ME (TJ)	Change (%)
Sheep & Beef									
Northland	243	1.7%	138	423	295	198	264	23337	1.6%
Auckland	78	-8.2%	129	446	306	168	262	7514	-7.9%
Waikato	468	-1.1%	111	421	373	202	277	47192	-0.7%
Bay of Plenty	72	9.5%	107	403	372	219	275	7246	10.1%
Gisborne	319	-2.4%	105	432	324	200	265	30918	-2.1%
Manawatu-Wanganui	866	-7.4%	91	413	373	186	266	84111	-7.4%
Hawkes Bay	572	-2.3%	94	397	213	179	221	46074	-2.2%
Taranaki	116	-6.9%	104	396	412	214	282	11946	-7.0%
Tasman	59	3.8%	86	405	275	173	235	5070	4.7%
Marlborough	308	-0.9%	59	310	231	133	183	20564	-0.7%
Westland	38	20.6%	83	358	381	207	257	3593	21.8%
Wellington	268	-4.8%	95	439	251	156	235	22960	-4.6%
Nelson City	3	1.0%	87	414	271	142	228	208	0.3%
Canterbury	1711	-6.2%	53	303	212	130	175	109073	-6.8%
Otago	1718	-1.2%	46	295	248	134	181	113356	-1.4%
Southland	627	-10.2%	61	373	367	174	244	55748	-11.0%
National	7465	-4.0%	73	354	279	159	216	588911	-4.26%
Dairy									
Northland	168	-2.5%	222	617	465	377	420	25715	-2.4%
Auckland	59	11.2%	209	628	502	341	420	8991	10.9%
Waikato	495	1.7%	186	629	506	363	421	76051	1.5%
Bay of Plenty	75	-6.5%	183	615	532	401	433	11772	-6.7%
Gisborne	12	390.0%	179	574	440	370	390	1746	354.5%
Manawatu-Wanganui	203	59.6%	149	603	545	313	403	29830	60.4%
Hawkes Bay	30	89.7%	137	578	392	325	358	3899	80.6%
Taranaki	226	4.8%	168	595	614	393	443	36456	4.8%
Tasman	25	-8.3%	142	587	539	353	405	3696	-8.6%
Marlborough	9	-6.5%	123	538	404	283	337	1064	-12.1%
Westland	42	-20.1%	123	533	584	374	404	6164	-19.8%
Wellington	43	46.5%	141	614	417	294	367	5815	44.8%
Nelson City	0	-41.2%	124	533	350	300	327	30	-39.5%
Canterbury	221	85.0%	103	520	361	278	315	25417	85.2%
Otago	78	26.6%	89	526	482	284	345	9838	26.1%
Southland	176	67.1%	94	532	568	334	382	24593	68.3%
National	1861	19.5%	157	589	506	345	399	271076	17.34%
Combined national	9326							859987	1.64%

2

3

1 Table B13. Land-use change, 2050 CO₂ = \$50 per tonne.

Region	Area (kha)	Change (%)	Winter Average	Spring Production (MJ ha ⁻¹ d ⁻¹)	Summer	Autumn	Average	Total ME (TJ)	Change (%)
Sheep & Beef									
Northland	243	1.9%	138	423	295	198	264	23385	1.8%
Auckland	79	-7.5%	129	446	306	168	262	7569	-7.2%
Waikato	469	-0.9%	111	421	372	202	276	47283	-0.5%
Bay of Plenty	72	9.9%	107	403	372	219	275	7272	10.5%
Gisborne	319	-2.4%	105	432	324	200	265	30920	-2.1%
Manawatu-Wanganui	867	-7.3%	91	413	373	186	266	84201	-7.3%
Hawkes Bay	572	-2.4%	94	397	213	179	221	46069	-2.2%
Taranaki	117	-6.4%	104	397	412	215	282	12013	-6.5%
Tasman	59	3.9%	86	405	275	173	235	5075	4.8%
Marlborough	308	-0.8%	59	310	231	133	183	20573	-0.6%
Westland	38	21.0%	83	359	381	207	257	3603	22.2%
Wellington	268	-4.6%	95	439	251	156	235	22986	-4.5%
Nelson City	3	1.0%	87	414	271	142	228	208	0.3%
Canterbury	1716	-5.9%	53	304	211	130	175	109353	-6.6%
Otago	1719	-1.1%	46	295	248	134	181	113412	-1.4%
Southland	627	-10.1%	61	373	367	174	244	55772	-11.0%
National	7475	-3.9%	73	354	279	159	216	589695	-4.13%
Dairy									
Northland	167	-3.1%	222	617	465	377	420	25557	-3.0%
Auckland	58	9.8%	209	628	502	341	420	8878	9.6%
Waikato	494	1.4%	186	629	506	363	421	75862	1.3%
Bay of Plenty	74	-6.8%	183	615	532	401	433	11730	-7.0%
Gisborne	12	389.0%	178	573	439	370	390	1741	353.4%
Manawatu-Wanganui	202	58.7%	149	603	546	313	403	29672	59.6%
Hawkes Bay	30	89.2%	137	578	392	325	358	3888	80.1%
Taranaki	225	4.4%	168	595	614	393	443	36309	4.4%
Tasman	25	-8.9%	142	587	539	353	405	3673	-9.2%
Marlborough	8	-9.2%	123	541	410	286	340	1043	-13.8%
Westland	42	-20.4%	123	533	584	374	403	6133	-20.2%
Wellington	43	45.4%	141	614	417	294	367	5771	43.7%
Nelson City	0	-41.2%	124	533	350	300	327	30	-39.5%
Canterbury	216	80.9%	103	520	363	278	316	24912	81.5%
Otago	77	24.7%	89	527	484	285	346	9721	24.6%
Southland	176	66.8%	94	532	568	334	382	24542	68.0%
National	1848	18.7%	157	589	506	345	399	269462	16.64%
Combined national	9323							859157	1.54%

2

3

1 Table B14. Land-use change, 2050 CO₂ = \$100 per tonne.

Region	Area (kha)	Change (%)	Average Production (MJ ha ⁻¹ d ⁻¹)				Total ME (TJ)	Change (%)	
			Winter	Spring	Summer	Autumn			
Sheep & Beef									
Northland	244	2.3%	138	423	295	198	264	23471	2.2%
Auckland	79	-7.3%	129	446	306	168	262	7585	-7.0%
Waikato	473	0.2%	111	421	373	202	277	47776	0.5%
Bay of Plenty	74	12.3%	107	403	372	219	275	7436	13.0%
Gisborne	319	-2.4%	105	432	324	200	265	30926	-2.1%
Manawatu-Wanganui	870	-7.1%	91	413	373	186	266	84413	-7.1%
Hawkes Bay	572	-2.3%	94	397	213	179	221	46102	-2.2%
Taranaki	117	-6.0%	104	397	412	215	282	12065	-6.1%
Tasman	59	4.3%	86	405	275	174	235	5095	5.2%
Marlborough	308	-0.9%	59	310	231	133	183	20554	-0.7%
Westland	39	24.0%	83	359	381	207	257	3696	25.3%
Wellington	268	-4.7%	95	439	251	156	235	22983	-4.5%
Nelson City	3	1.0%	87	414	271	142	228	208	0.3%
Canterbury	1716	-5.9%	53	304	211	130	175	109353	-6.6%
Otago	1719	-1.1%	46	295	248	134	181	113419	-1.4%
Southland	627	-10.1%	61	373	367	174	244	55811	-10.9%
National	7487	-3.7%	73	354	279	159	216	590893	-3.94%
Dairy									
Northland	165	-3.9%	222	617	465	377	420	25359	-3.7%
Auckland	58	9.1%	209	628	502	341	420	8818	8.8%
Waikato	488	0.3%	186	629	505	363	421	75035	0.2%
Bay of Plenty	73	-8.8%	183	615	530	401	433	11478	-9.0%
Gisborne	12	386.0%	179	573	438	370	390	1729	350.2%
Manawatu-Wanganui	200	57.0%	149	603	545	313	403	29341	57.8%
Hawkes Bay	29	85.7%	137	578	391	325	358	3811	76.6%
Taranaki	224	4.1%	168	595	614	393	443	36218	4.1%
Tasman	25	-10.0%	142	587	538	353	405	3626	-10.3%
Marlborough	8	-10.5%	123	541	410	286	340	1028	-15.1%
Westland	41	-22.3%	123	533	585	374	404	5989	-22.1%
Wellington	43	44.9%	141	614	417	294	367	5750	43.2%
Nelson City	0	-41.2%	124	533	350	300	327	30	-39.5%
Canterbury	215	80.4%	103	520	363	278	316	24844	81.0%
Otago	77	24.3%	89	527	484	285	346	9690	24.2%
Southland	176	66.4%	94	532	568	334	382	24485	67.6%
National	1833	17.7%	157	590	506	345	399	267232	15.68%
Combined national	9320							858125	1.42%

2

3

1 **Acknowledgements**

2 This work was supported by the New Zealand Ministry for Primary Industries (MPI) and
3 public research funding from the Government of New Zealand (GNS-540GCT81). Pasture
4 production data provided by Collier Isaacs (Landcorp), Brenda Rosser (Landcare
5 Research/GNS Science), Beef + Lamb New Zealand, DairyNZ, and AgResearch.
6 Meteorological VCSN data provided by NIWA. We thank Duncan Smeaton (AgResearch),
7 Anne-Gaelle Ausseil (Landcare Research), William Power (GNS Science) and Wei Zhang
8 (Motu/MPI) for their contributions. Discussion with MPI has improved the final presentation
9 of our results. We thank two anonymous referees whose comments have improved this
10 manuscript.

11

1 **References**

2 Ainsworth, E. A. and Long, S. P.: What have we learned from 15 years of free-air CO₂
3 enrichment (FACE)? A meta-analytic review of the responses of photosynthesis, canopy
4 properties and plant production to rising CO₂, *New Phytol.*, 165, 351-71, 2005.

5 Allen, R. G., Pereira, L. S., Raes, D., and Smith M.: Crop evapotranspiration - Guidelines for
6 computing crop water requirements - FAO Irrigation and drainage paper 56, Food and
7 Agriculture Organization of the United Nations, Rome, Italy, 1998.

8 Baisden, W. T.: Agricultural and forest productivity for modelling policy scenarios:
9 evaluating approaches for New Zealand greenhouse gas mitigation, *Journal of the Royal
10 Society of New Zealand*, 36, 1-15, 2006.

11 Baisden, W. T., Keller, E. D., Timar, L., Smeaton, D., Clark, A., Ausseil, A., Power, W. L.,
12 and Zhang, W.: New Zealand's pasture production in 2020 and 2050, GNS Science
13 consultancy report 2010/154, New Zealand Ministry for Primary Industries, Wellington, New
14 Zealand, 2010.

15 Barker, D. J., Lancashire, J. A., Moloney, S. C., Dymock, N., Stevens, D. R., Turner, J. D.,
16 Scott, D., and Archie, W. J.: Introduction, production, and persistence of five grass species in
17 dry hill country: 8. Summary and conclusions, *New Zeal. J. Agr. Res.*, 36, 61-66, 1993.

18 Beer, C., Reichstein, M., Tomelleri, E., Ciais, P., Jung, M., Carvalhais, N., and Papale, D.:
19 Terrestrial gross carbon dioxide uptake: global distribution and covariation with climate,
20 *Science*, 329, 834-838, 2010.

21 Bond-Lamberty, B., Gower, S. T., Ahl, D. E., and Thornton, P. E.: Reimplementation of the
22 Biome-BGC model to simulate successional change, *Tree Physiol.*, 25, 413-424, 2005.

23 Bondeau, A., Smith, P. C., Zaehle, S., Schaphoff, S., Lucht, W., Cramer, W., Gerten, D.,
24 Lotze-Campen, H., Muller, C., Reichstein, M., and Smith, B.: Modelling the role of
25 agriculture for the 20th century global terrestrial carbon balance, *Glob. Change Biol.*, 13, 679-
26 706, doi: 10.1111/j.1365-2486.2006.01305.x, 2007.

27 Clark, A., Mullan, B., and Porteous, A.: Scenarios of regional drought under climate change,
28 NIWA report number WLG2010-32, New Zealand Ministry for Primary Industries,
29 Wellington, New Zealand, 2011.

1 Clarke-Hill, W.J., and Fraser, T.J.: PasturePlan™ : On-farm pasture growth and quality data
2 for sheep and beef farms throughout New Zealand, New Zealand Grassland Association
3 Conference Proceedings, Vol. 69, 2007.

4 DairyNZ: DairyNZ Farmwatch - Waikato, available at:
5 <http://farmwatch.dairynz.co.nz/Home>ShowData/2#table> (last access: 21 August 2014), 2011.

6 de Graaf, M. A., Van Groenigen, K. J., Six, J., Hungate, B., and Van Kessel, C.: Interactions
7 between plant growth and soil nutrient cycling under elevated CO₂: a meta-analysis, Glob.
8 Change Biol., 12, 2077–2091, doi: 10.1111/j.1365-2486.2006.01240.x, 2006.

9 Deser, C., Phillips, A., Bourdette, V., and Teng, H.: Uncertainty in climate change
10 projections: the role of internal variability, Clim Dyn, 38, 527-546, doi:10.1007/s00382-010-
11 0977-x, 2012.

12 Di Vittorio, A. V., Anderson, R. S., White, J. D., Miller, N. L., and Running, S. W.:
13 Development and optimization of an Agro-BGC ecosystem model for C4 perennial grasses,
14 Ecol. Model., 221, 2038-2053, 2010.

15 Doherty, J.: PEST: Model Independent Parameter Estimation User Manual, 5th ed.,
16 Watermark Numerical Computing, Brisbane, Australia, 2005.

17 ENSEMBLES: ENSEMBLES RT2A Results, available at:
18 <http://www.cnrm.meteo.fr/ensembles/public/results/results.html> (last access: 3 April 2014),
19 2009.

20 ESRI: ArcGIS Desktop Release 9.3, ESRI Environmental Systems Research Institute,
21 Redlands, California, USA, 2009.

22 Gerten, D., Rost, S., von Bloh, W., and Lucht, W.: Causes of change in 20th century global
23 river discharge, Geophys. Res. Lett., 35, L20405, doi:10.1029/2008GL035258, 2008.

24 Gumpenberger, M., Vohland, K., Heyder, U., Poulter, B., Macey, K., Rammig, A., and
25 Cramer, W.: Predicting pan-tropical climate change induced forest stock gains and losses—
26 implications for REDD, Environ. Res. Lett., 5, 014013, doi:10.1088/1748-9326/5/1/014013,
27 2010.

28 Hendy, J., Kerr S., and Baisden, W. T.: The Land Use in Rural New Zealand Model Version 1
29 (LURNZv1): Model Description, Motu Working Paper 07-07, Motu Economic and Public
30 Policy Research, Wellington, New Zealand, available at:

1 http://www.motu.org.nz/publications/detail/the_land_use_in_rural_new_zealand_model_version_1 (last access: 7 May 2014), 2007.

3 Hendy, J., Kerr, S., and Baisden, W. T.: Greenhouse gas emissions charges and credits on
4 agricultural land: what can a model tell us?, in: Clean development mechanism and law, first
5 edition, Lakshmi, L. (Ed.), Icfai University Press, Hyderabad, India, 198-209, 2008.

6 Hidy, D., Barcza, Z., Haszpra, L., Churkina, G., Pintér, K., and Nagy, Z.: Development of the
7 Biome-BGC model for simulation of managed herbaceous ecosystems, Ecol. Model., 226, 99-
8 119, doi:10.1016/j.ecolmodel.2011.11.008, 2012.

9 Jung, M., Le Maire, G., Zaehle, S., Luyssaert, S., Vetter, M., Churkina, G., Ciais, P., Viovy,
10 N., and Reichstein, M.: Assessing the ability of three land ecosystem models to simulate groß
11 carbon uptake of forests from boreal to Mediterranean climate in Europe, Biogeosciences, 4,
12 647-656, 2007.

13 Keating, B. A., Carberry, P. S., Hammer, G. L., Probert, M. E., Robertson, M. J., Holzworth,
14 D., Huth, N. I., Hargreaves, J. N. G., Meinke, H., Hochman, Z., McLean, G., Verburg, K.,
15 Snow, V., Dimes, J. P., Silburn, M., Wang, E., Brown, S., Bristow, K. L., Asseng, S.,
16 Chapman, S., McCown, R. L., Freebairn, D. M., and Smith, C. J.: An overview of APSIM, a
17 model designed for farming systems simulation, Eur. J. Agron., 18, 267-288, 2003.

18 Keller, E. D.: Using the Biome-BGC Model to assess New Zealand's production of animal
19 products from 2006/7 to 2011/12 seasons relative to historical averages, GNS Science
20 consultancy letter report 2012/177LR, New Zealand Ministry for Primary Industries,
21 Wellington, New Zealand, 2012.

22 Keller, E. D., Baisden, W. T., and Timar, L.: Adapting the Biome-BGC model to New
23 Zealand pastoral agriculture: climate change and land-use change, AGU Fall Meeting, 5-9
24 December 2011, San Francisco, USA, GC23C-0955, 2011.

25 Kerr, S., Anastasiadis, S., Olssen, A., Power, W., Timar, L., and Zhang, W.: Spatial and
26 Temporal Responses to an Emissions Trading Scheme Covering Agriculture and Forestry:
27 Simulation Results from New Zealand, Forests, 3, 1133-1156, doi:10.3390/f3041133, 2012.

28 Landcare Research: Fundamental Data Layers (FDLs) – Soils, available at:
29 http://soils.landcareresearch.co.nz/contents/SoilData_FSL_About.aspx?currentPage=SoilData_FSL&menuItem=SoilData (last access: 7 May 2014), 2014.

1 Legendre, P.: lmodel2: Model II Regression, R package version 1.7-2, available at:
2 <http://CRAN.R-project.org/package=lmodel2> (last access: 21 August 2014), 2014.

3 Mullan, B., Porteous, A., Wratt, D., and Hollis, M.: Changes in drought risk with climate
4 change, NIWA client report WLG2005-23, Ministry for the Environment, Wellington, New
5 Zealand, available at: <http://www.mfe.govt.nz/publications/climate/drought-risk-may05/index.html> (last access: 7 May 2014), 2005.

6 Nakicenovic, N. and Swart, R. (Eds.): Special Report on Emissions Scenarios, A Special
7 Report of Working Group III of the Intergovernmental Panel on Climate Change, Cambridge
8 University Press, Cambridge, United Kingdom and New York, NY, USA, available at:
9 <http://www.ipcc.ch/ipccreports/sres/emission/index.php?idp=0> (last access: 22 August 2014),
10 2000.

11 New Zealand Ministry for the Environment: Climate change: Impacts on New Zealand, MfE
12 report INFO 95, New Zealand Ministry for the Environment, Wellington, New Zealand,
13 available at: <http://www.mfe.govt.nz/publications/climate/impacts-on-new-zealand/index.html>
14 (last access: 7 May 2014), 2001.

15 New Zealand Ministry for the Environment: New Zealand's Fifth National Communication
16 under the United Nations Framework Convention on Climate Change, New Zealand Ministry
17 for the Environment, Wellington, New Zealand, available at:
18 <http://www.mfe.govt.nz/publications/climate/nz-fifth-national-communication/> (last access: 7
19 May 2014), 2009.

20 New Zealand Ministry for the Environment: New Zealand's Greenhouse Gas Inventory 1990-
21 2011, MfE Publication number ME 1113, Submission to the United Nations Framework
22 Convention on Climate Change, 12 April 2013.

23 New Zealand Ministry for Primary Industries: Situation and Outlook for Primary Industries
24 (SOPI) 2012, New Zealand Ministry for Primary Industries, Wellington, New Zealand,
25 available at: <http://www.mpi.govt.nz/news-resources/publications> (last access: 7 May 2014),
26 2012.

27 Newton, P. C., Lieffering, M., Bowatte, W. S. D., Brock, S. C., Hunt, C. L., Theobald, P. W.,
28 and Ross, D. J.: The rate of progression and stability of progressive nitrogen limitation at
29 elevated atmospheric CO₂ in a grazed grassland over 11 years of Free Air CO₂ enrichment,
30 Plant and Soil, 336, 433-441, 2010.

1 Parfitt, R. L., Baisden, W. T., Schipper, L. A., and Mackay, A. D.: Nitrogen inputs and
2 outputs for New Zealand at national and regional scales: past, present and future, *J. Roy. Soc.*
3 *New Zeal.*, 38, 71-87, 2008.

4 Parfitt, R. L., Schipper, L. A., Baisden, W. T., and Elliott, A. H.: Nitrogen inputs and outputs
5 for New Zealand in 2001 at national and regional scales, *Biogeochemistry*, 80, 71-88, 2006.

6 Pietsch, S. A., Hasenauer, H., and Thornton, P. E.: BGC-model parameters for tree species
7 growing in central European forests, *Forest Ecol. Manag.*, 211, 264-295, 2005.

8 R Core Team: R: A language and environment for statistical computing, R Foundation for
9 Statistical Computing, Vienna, Austria, available at: <http://www.R-project.org/> (last access:
10 21 August 2014), 2013.

11 Renwick, J., Mullan, B., Wilcocks, L., Zammit, C., Baisden, W. T., Keller, E. D.,
12 Kirschbaum, M., Meason, D., Harrison, D., Verkerk, G., Cooke, A., Marshall, P., and Clark,
13 A.: Four Degrees of Global Warming: effects on the New Zealand primary sector, NIWA
14 client report WLG2012-24, New Zealand Ministry for Primary Industries, Wellington, New
15 Zealand, available at: <http://www.mpi.govt.nz/news-resources/publications> (last access: 7
16 May 2014), 2013.

17 Rockström, J., Falkenmark, M., Karlberg, L., Hoff, H., Rost, S., and Gerten, D.: Future water
18 availability for global food production: the potential of green water for increasing resilience to
19 global change, *Water Resour. Res.*, 45, W00A12, doi: 10.1029/2007WR006767, 2009.

20 Ronneberger, K., Criscuolo, L., Knorr, W., and Tol, R. S.: KLUM@ LPJ: Integrating
21 dynamic land-use decisions into a dynamic global vegetation and crop growth model to assess
22 the impacts of a changing climate. A feasibility study for Europe, Working Papers FNU-113,
23 Research unit Sustainability and Global Change, Hamburg University, Germany, 2006.

24 Rosser, B. J. and Ross, C. W.: Recovery of pasture production and soil properties on soil slip
25 scars in erodible siltstone hill country, Wairarapa, New Zealand, *New Zeal. J. Agr. Res.*, 54,
26 23-44, 2011.

27 Rost, S., Gerten, D., Hoff, H., Lucht, W., Falkenmark, M., and Rockström, J.: Global
28 potential to increase crop production through water management in rainfed agriculture,
29 *Environ. Res. Lett.*, 4, 044002, 2009.

1 Roudier, P., Sultan, B., Quirion, P., and Berg, A.: The impact of future climate change on
2 West African crop yields: What does the recent literature say?, *Global Environmental
3 Change*, 21, 1073-1083, 2011.

4 Schilling, C., Zuccollo, J., and Nixon, C.: Dairy's role in sustaining New Zealand: the sector's
5 contribution to the economy, The New Zealand Institute for Economic Research, Wellington,
6 New Zealand, available at: <http://nzier.org.nz/publications/dairys-role-in-sustaining-new-zealand> (last access: 7 May 2014), 2010.

8 Smith, L.C., Moss, R.A., Morton, J.D., Metherell, A.K., and Fraser, T.J.: Pasture production
9 from a long-term fertiliser trial under irrigation, *New Zeal. J. Agr. Res.*, 55, 105-117, 2012

10 South Island Dairying Development Centre: LU Dairy Farm Weekly Data, available at
11 <http://www.siddc.org.nz/lu-dairy-farm/weekly-data/> (last access: 21 August 2014), 2014.

12 Stevens, D., Walcroft, J., Casey, M., Burggraaf, V., Fraser, T., Webby, R., Smeaton, D., and
13 Lambert, G.: Literature review on effects of pasture renewal in New Zealand and simulation
14 modelling of those effects on whole farm profitability, AgResearch Ltd, Hamilton, New
15 Zealand, available at: http://www.pasturerenewal.org.nz/downloads/benefit_analysis.pdf (last
16 access: 7 May 2014), 2007.

17 Stroombergen, A., Stojanovik, A., Wratt, D., Mullan, B., Tait, A., Woods, R., Baisden, W. T.,
18 Giltrap, D., Lock, K., Hendy, J., and Kerr, S.: Costs and benefits of climate change and
19 adaptation to climate change in New Zealand agriculture: what do we know so far?, New
20 Zealand Ministry of Agriculture and Forestry, Wellington, New Zealand, available at:
21 http://www.motu.org.nz/publications/detail/costs_and_benefits_of_climate_change_and_adaptation_to_climate_change_in_new_zealand (last access: 7 May 2014), 2008.

23 Tait, A., Henderson, R., Turner, R., and Zheng, X. G.: Thin plate smoothing spline
24 interpolation of daily rainfall for New Zealand using a climatological rainfall surface, *Int. J.
25 Climatol.*, 26, 2097-2115, 2006.

26 Thornton, P. E.: Description of a numerical simulation model for predicting the dynamics of
27 energy, water, carbon, and nitrogen in a terrestrial ecosystem, Ph.D. thesis, University of
28 Montana, Missoula, USA, 280 pp., 1998.

29 Thornton, P. E., Law, B. E., Gholz, H. L., Clark, K. L., Falge, E., Ellsworth, D. S., Goldstein,
30 A. H., Monson, R. K., Hollinger, D., Falk, M., Chen, J., and Sparks, J. P.: Modeling and

1 measuring the effects of disturbance history and climate on carbon and water budgets in
2 evergreen needleleaf forests, *Agr. Forest Meteorol.*, 113, 185-222, 2002.

3 Thornton, P. E. and Rosenbloom, N. A.: Ecosystem model spin-up: Estimating steady state
4 conditions in a coupled terrestrial carbon and nitrogen cycle model, *Ecol. Model.*, 189, 25-48,
5 2005.

6 Thornton, P. E., Running, S. W., and Hunt, E. R.: Biome-BGC: Terrestrial Ecosystem Process
7 Model, Version 4.2 Final Release, Numerical Terradynamic Simulation Group, School of
8 Forestry, University of Montana, Missoula, USA, available at:
9 <http://www.ntsg.umt.edu/project/biome-bgc> (last access: 7 May 2014), 2005.

10 Timar, Levente: Rural Land Use and Land Tenure in New Zealand, Motu Working Paper 11-
11 13, Motu Economic and Public Policy Research, Wellington, 2011.

12 Timar, L. and Kerr, S.: Land-use Intensity and Greenhouse Gas Emissions in the LURNZ
13 Model, Motu Working Paper 14-03, Motu Economic and Public Policy Research, Wellington,
14 2014.

15 Wang, Q., Watanabe, M., and Ouyang, Z.: Simulation of water and carbon fluxes using
16 Biome-BGC model over crops in China, *Agr. Forest Meteorol.*, 131, 209-224,
17 doi:10.1016/j.agrformet.2005.06.002, 2005.

18 Wang, W., Ichii, K., Hashimoto, H., Michaelis, A. R., Thornton, P. E., Law, B. E., and
19 Nemani, R. R.: A hierarchical analysis of terrestrial ecosystem model Biome-BGC:
20 Equilibrium analysis and model calibration, *Ecol. Model.*, 220, 2009-2023, 2009.

21 Warrick, R. A., Kenny, G. J., and Harman, J. J. (Eds): The Effects of Climate Change and
22 Variation in New Zealand: An Assessment Using the CLIMACTS System, International
23 Global Change Institute, University of Waikato, Hamilton, New Zealand, available at:
24 <http://docs.niwa.co.nz/library/public/0473079887.pdf> (last access: 7 May 2014), 2001.

25 White, M. A., Thornton, P. E., Running, S. W., and Nemani, R. R.: Parameterization and
26 Sensitivity Analysis of the Biome-BGC Terrestrial Ecosystem Model: Net Primary
27 Production Controls, *Earth Interact.*, 4, 4-003, 2000.

28

1 Table 1. Calibration sites, location, description and dates of pasture growth data used for
 2 model calibration.

Site	Location	Description	Data Availability	Mean	Annual	Mean	Daily
				Rainfall (mm)	Max/Min	Temperature	(°C)
Whatawhata	-37.80° S 175.15° E	Sheep/Beef, easy hills	Jan 2003 – Sep 2005 monthly intervals	1607		19 / 9.6	
Waikato							
Te Whanga Station	-41.03° S 175.74° E	Sheep/Beef, hillside landslide	Jun 2007 – Aug 2009 2 month intervals (3 distinct sites)	886		18 / 7.4	
Wairarapa							
Winchmore Irrigation Research Station	-43.83° S 171.71° E	Sheep/Beef, flat	Jan 1997 – Dec 2003 monthly intervals	715		17 / 5.7	
DairyNZ Scott Farm	-37.77° S 175.36° E	Dairy, large scale farm	Aug 2009 – May 2011 weekly intervals	1086		19 / 8.8	
Canterbury							
Hamilton							
Lincoln University Dairy Farm (LUDF)	-43.64° S 172.44° E	Dairy, irrigated	Jan 2005 – Dec 2009 weekly intervals	604		17 / 6.8	
Canterbury							
Landcorp Waitepeka Dairy Farm	-46.29° S 169.67° E	Dairy	Jan 2004 – Dec 2009 monthly intervals	701		15 / 5.7	
Southland							

3
 4

1 Table 2. Comparison of baseline and intensification Biome-BGC model parameters.

Parameter	Baseline	Intense	Baseline	Intense
	Sheep/Beef	Sheep/Beef	Dairy	Dairy
Pasture Utilisation	0.55	0.60	0.90	0.95
Annual Whole Plant mortality fraction	0.226	0.247	0.722	0.762
Symbiotic & Asymbiotic Nitrogen Fixation (kg N/m ² /year)	0.018	0.021	0.032	0.038

2

3

1 Table 3. Summary of model scenario results with constant land-use area, tabulated by
 2 scenario and season. Percent change calculated in reference to the Baseline scenario. Seasonal
 3 numbers are averages over the modelled time slice, and national totals are the annual average
 4 sum of sheep/beef and dairy results. ME=Metabolisable energy, defined as the amount of
 5 energy available to grazing animals.

Scenario	Area (kha)	Winter	Spring	Summer	Autumn	Average	Total ME (TJ)	Change (%)
		Average Production (MJ ha ⁻¹ d ⁻¹)						
Baseline 2001-2009								
Sheep & Beef	7778	73	356	279	159	217	615126	-
Dairy	1557	163	594	515	353	406	231014	-
National Total	9335						846140	-
Elevated CO₂ 2020								
Sheep & Beef	7778	74	356	280	160	218	617531	0.4%
Dairy	1557	166	598	517	354	409	232203	0.5%
National Total	9335						849734	0.4%
Intensification 2020								
Sheep & Beef	7778	80	396	301	172	237	673861	9.5%
Dairy	1557	181	668	581	399	457	259915	12.5%
National Total	9335						933776	10.4%
MPI climate change 2050								
Sheep & Beef	7778	80	360	276	168	221	627293	2.0%
Dairy	1557	186	617	518	364	421	239427	3.6%
National Total	9335						866720	2.4%
GIEH climate change 2050								
Sheep & Beef	7778	85	354	260	169	217	616759	0.3%
Dairy	1557	204	615	498	372	422	239967	3.9%
National Total	9335						856726	1.3%
Baseline 1997-2011								
Sheep & Beef	7778	69	350	230	135	196	556567	-
Dairy	1557	153	588	472	311	381	216717	-
National Total	9335						773284	-

MPI climate change 2100								
Sheep & Beef	7778	97	338	185	143	191	541710	-2.7%
Dairy	1557	235	608	408	336	397	225631	4.1%
National Total	9335						767341	-0.8%
CCC climate change 2100								
Sheep & Beef	7778	105	326	112	139	170	483402	-13.1%
Dairy	1557	258	608	306	331	376	213593	-1.4%
National Total	9335						696995	-9.9%

1

2


1 Table 4. Summary of model scenario results with land-use change. Percent change calculated
 2 in reference to the Baseline scenario in Table 3. Seasonal numbers are averages over the
 3 modelled time slice, and national totals are the annual average sum of sheep/beef and dairy
 4 results. ME=Metabolisable energy, defined as the amount of energy available to grazing
 5 animals.

Region	Area (kha)	Change (%)	Winter	Spring	Summer	Autumn	Avg	Total ME (TJ)	Change (%)
Average Production (MJ ha ⁻¹ d ⁻¹)									
Land-use 2020 \$0									
Sheep & Beef	7766	-0.2%	73	356	279	159	217	614327	-0.13%
Dairy	1566	0.6%	163	595	514	352	406	232053	0.45%
National Total	9332							846380	0.03%
Land-use 2020 \$50									
Sheep & Beef	7777	0.0%	73	356	279	159	217	615462	0.05%
Dairy	1552	-0.3%	163	595	513	352	406	230025	-0.43%
National Total	9330							845487	-0.08%
Land-use 2020 \$100									
Sheep & Beef	7789	0.1%	73	356	279	159	217	616684	0.25%
Dairy	1537	-1.3%	163	595	513	352	406	227802	-1.39%
National Total	9326							844486	-0.20%
Land-use 2050 \$0									
Sheep & Beef	7465	-4.0%	73	354	279	159	216	588911	-4.26%
Dairy	1861	19.5%	157	589	506	345	399	271076	17.3%
National Total	9326							859987	1.64%
Land-use 2050 \$50									
Sheep & Beef	7475	-3.9%	73	354	279	159	216	589695	-4.13%
Dairy	1848	18.7%	157	589	506	345	399	269462	16.6%
National Total	9323							859157	1.54%
Land-use 2050 \$100									
Sheep & Beef	7487	-3.7%	73	354	279	159	216	590893	-3.94%
Dairy	1833	17.7%	157	590	506	345	399	267232	15.7%
National Total	9320							858125	1.42%




1
 2
 3
 4
 5
 6
 7

Figure 1. Calibration and validation sites. Left: dairy and sheep/beef calibration sites. White circles are sheep/beef sites, and red squares are dairy sites. Right: sheep/beef validation sites. The three most northern sites (Broadwood, Tauranga, and Whakatane) are outliers in terms of model/observation fit




1
2

3 Figure 2. Modelled and measured pasture growth at calibration sites. Growth (in average
4 kilograms of dry matter per hectare per day) versus time at all dairy (a-c) and sheep/beef (d-h)
5 calibration sites: a) Waitepeka Dairy Farm (Southland) from January 2004 – December 2009;
6 b) Scott Dairy Farm (Hamilton) from August 2009 – May 2011; c) Lincoln University Dairy
7 Farm (Canterbury) from January 2005 – December 2009; d) Winchmore Research Station
8 (Canterbury) from January 1997 – December 2003 (in total kilograms of dry matter per
9 hectare rather than daily averages); e) Whatawhata (Waikato) from February 2003 – October
10 2005; f) Te Whanga (Wairarapa) uneroded site, g) Te Whanga 1977 slip site, and h) Te
11 Whanga 1961 slip site, from August 2007 – August 2009.




1

2 Figure 3. Scatter plot of modelled vs. measured pasture growth for calibration sites. Measured
 3 and modelled daily growth rates (kilograms of dry matter per hectare) from all calibrations
 4 sites for dairy (left) and sheep/beef (right). Data are daily averages of growth over cutting
 5 intervals (between 1 week and 1 month). Correlation coefficient $R^2 = 0.64$ (RMSE = 16.6) for
 6 dairy and 0.70 (RMSE = 9.88) for sheep/beef. Type 2 linear regression model shown is $y =$
 7 $0.83 \pm 0.060 x + 6.1 \pm 2.9$ (dairy) and $y = 0.83 \pm 0.090 x + 6.1 \pm 2.0$ (sheep/beef). Reported
 8 errors are 2σ . The 1:1 line is drawn for reference.
 9



1
2 Figure 4. Scatter plot of modelled vs. measured pasture growth for validation sites. Measured
3 and modelled daily growth rates (kilograms of dry matter per hectare) for all sheep/beef
4 validation sites compared at individual cutting intervals (left) and monthly averages over three
5 years (right). Correlation coefficient $R^2 = 0.36$ (RMSE = 20.9) and 0.48 (RMSE = 15.9),
6 respectively. Type 2 linear regression model shown is $y = 0.53 \pm 0.040 x + 10.4 \pm 1.2$ (left)
7 and $y = 0.63 \pm 0.085 x + 7.8 \pm 2.5$ (right). Reported errors are 2σ . The 1:1 line is drawn for
8 reference.
9




1


2

3 Figure 5. Historical and projected land-use areas in New Zealand (carbon price = \$0). Data to
4 the left of the dashed line (2008) is historical, and to the right is estimated by LURNZ based
5 on exogenous forecasts of economic variables.

6




1
2
3 Figure 6. Land area and pasture production changes. Percentage change from baseline
4 estimated by combining outputs of LURNZ and Biome-BGC models in 2020 and 2050.
5




1
2
3
4
5
6
7
8

Figure 7. Elevated CO₂ (top) and Intensification (bottom) 2020 model scenarios, average annual total pasture production, percentage change from baseline for Sheep/Beef (left) and Dairy (right). Each map shows national pasture production as if all of the available land (excluding urban and conservation land) was devoted to sheep/beef or dairy agriculture systems and is not an actual representation of current or projected land use.





1
2
3
4
5
6
7
8

Figure 8. Climate Change 2050 MPI (top) and GIEH (bottom) model scenarios, average annual total pasture production, percentage change from baseline for Sheep/Beef (left) and Dairy (right). Each map shows national pasture production as if all of the available land (excluding urban and conservation land) was devoted to sheep/beef or dairy agriculture systems and is not an actual representation of current or projected land use.



1
2
3
4
5
6
7
8

Figure 9. Climate Change 2100 MPI (top) and CCC (bottom) model scenarios, average annual total pasture production, percentage change from baseline for Sheep/Beef (left) and Dairy (right). Each map shows national pasture production as if all of the available land (excluding urban and conservation land) was devoted to sheep/beef or dairy agriculture systems and is not an actual representation of current or projected land use.



1
2

3 Figure 10. Total annual change in precipitation in 2050 (top) and 2100 (bottom), percentage
4 change from baseline, for MPI (left) and GIEH/CCC (right) climate change scenarios.