

1 *Representing Icebergs in the iLOVECLIM Model (version 1.0)- A Sensitivity*

2 *Study*

3 M. Bügelmayer¹, D. M. Roche^{1,2}, H. Renssen¹

4 m.bugelmayer@vu.nl, didier.roche@vu.nl, h.renssen@vu.nl

5 ¹ Earth and Climate Cluster, Faculty of Earth and Life Sciences, Vrije Universiteit Amsterdam, Amsterdam, The
6 Netherlands

7 ² Laboratoire des Sciences du Climat et de l'Environnement (LSCE), CEA/CNRS-INSU/UVSQ, Gif-sur-Yvette Cedex,
8 France

9

10 Recent modelling studies have indicated that icebergs play an active role in the climate system as
11 they interact with the ocean and the atmosphere. The icebergs' impact is due to their slowly
12 released melt water, which freshens and cools the ocean and consequently alters the ocean
13 stratification and the sea ice conditions. The spatial distribution of the icebergs and their melt
14 water depends on the atmospheric and oceanic forces acting on them as well as on the initial
15 icebergs' size. The studies conducted so far have in common that the icebergs were moved by
16 reconstructed or modelled forcing fields and that the initial size distribution of the icebergs was
17 prescribed according to present day observations. To study the sensitivity of the modelled iceberg
18 distribution to initial and boundary conditions, we performed 15 sensitivity experiments using the
19 climate model *iLOVECLIM* that includes actively coupled ice-sheet and iceberg modules, to analyse
20 1) the impact of the atmospheric and oceanic forces on the iceberg transport, mass and melt flux
21 distribution, and 2) the effect of the initial iceberg size on the resulting Northern Hemisphere
22 climate including the Greenland ice sheet, due to feedback mechanisms such as altered
23 atmospheric temperatures, under different climate conditions (pre-industrial, high/low radiative
24 forcing). Our results show that, under equilibrated pre-industrial conditions, the oceanic currents
25 cause the icebergs to stay close to the Greenland and North American coast, whereas the
26 atmospheric forcing quickly distributes them further away from their calving site. Icebergs
27 remaining close to Greenland last up to two years longer as they reside in generally cooler waters.
28 Moreover, we find that local variations in the spatial distribution due to different iceberg sizes do
29 not result in different climate states and Greenland ice sheet volume, independent of the
30 prevailing climate conditions (pre-industrial, warming or cooling climate). Therefore, we conclude

31 that local differences in the distribution of their melt flux do not alter the prevailing Northern
32 Hemisphere climate and ice sheet under equilibrated conditions and continuous supply of
33 icebergs. Furthermore, our results suggest that the applied radiative forcing scenarios have a
34 stronger impact on climate than the initial size distribution of the icebergs.

35 **1 Introduction**

36 Icebergs are an important part of the climate system as they interact with the ocean, atmosphere
37 and cryosphere (e.g. Hemming, 2004; Smith et al., 2011; Tournadre et al., 2012). Most
38 importantly, icebergs play an important part in the global fresh water cycle since currently up to
39 half of the mass loss of the Antarctic (Rignot et al., 2013) and Greenland ice sheets is due to
40 calving (approx. 0.01 Sv, 1 Sv = $10^6 \text{ m}^3 \text{s}^{-1}$, Hooke et al., 2005). As icebergs are melting, they affect
41 the upper ocean by freshening and cooling due to their uptake of latent heat. Several studies have
42 revealed that freshening and cooling have opposing effects on ocean stratification, as cooling
43 enhances the surface density, promoting deep mixing, whereas freshening decreases the water
44 density, stabilizing the water column (Jongma et al., 2009, 2013, Green et al., 2011).

45 Moreover, the implementation of dynamical icebergs in climate models has revealed that icebergs
46 enhance the formation of sea ice (Jongma et al., 2009, 2013; Wiersma and Jongma 2010;
47 Bügelmayer et al., 2015), which forms a barrier between the ocean and the atmosphere. On the
48 one hand sea ice shields the ocean from being stirred by atmospheric winds, and on the other
49 hand from losing heat to the relatively cold atmosphere, thus, reducing mixing of the upper water
50 column. Further, this reduced oceanic heat loss leads, in combination with an increase in surface
51 albedo, to a changed atmospheric state (Bügelmayer et al., 2015). Thus, icebergs indirectly alter
52 the ice sheet's mass balance through their effect on air temperature and precipitation
53 (Bügelmayer et al., 2015).

54 The amount of icebergs calved and their effects on climate depend on the calving flux provided by
55 the ice sheets, which is altered by the prevailing climate conditions. For instance, in the relatively
56 cold climate of the last glacial massive episodic discharges of icebergs into the North Atlantic
57 Ocean, so-called Heinrich events, have been recorded in distinct layers of ice rafted debris
58 (Andrews 1998; Hemming 2004). These periods of enhanced ice discharge have been proposed to
59 be caused by ice shelf collapses (e.g. MacAyeal, 1993; Hulbe et al., 2004; Alvarez-Solas et al., 2011)
60 and happened during periods of a (partial) collapse of the thermohaline circulation (Broecker et

61 al., 1993; McManus et al. 2004; Gherardi et al., 2005; Kageyama et al., 2010). It has been
62 suggested that the collapse was caused by the long duration (Marcott et al., 2011) and the
63 increased amount of freshwater released (0.04 up to 0.4 Sv, Roberts et al., 2014) and coincided
64 with globally altered climate conditions (Hemming, 2004).

65 So far, different approaches have been taken to incorporate icebergs from the Antarctic and
66 Greenland ice sheets into numerical models for different time periods. Bigg et al. (1996, 1997)
67 presented an iceberg module, which was forced with present-day atmospheric and oceanic input
68 fields from uncoupled model simulations. The forcing was provided off-line by atmospheric and
69 oceanic models to investigate the drift patterns of icebergs in the Northern Hemisphere. Their
70 approach was further developed for the Southern Ocean by Gladstone et al. (2001), who used
71 modelled oceanic and modern reconstructed wind fields, as well as observed calving amounts to
72 seed the iceberg module. Subsequently, the same iceberg module was implemented in an earth
73 system model of intermediate complexity (EMIC) by Jongma et al. (2009) to investigate the impact
74 of icebergs on the Southern Ocean under pre-industrial conditions. In the latter study, the icebergs
75 were seeded based on a prescribed constant calving flux from observational estimates, but moved
76 according to the modeled winds and currents and interacted with the model atmosphere and
77 ocean. Martin and Adcroft (2010) then implemented the iceberg model into a coupled global
78 climate model (CGCM) using the model's variable runoff as a calving flux though still lacking an ice
79 sheet component. Most recently, Bügelmayer et al. (2015) took the next step by using an EMIC
80 with both dynamically coupled ice sheet and iceberg model components. In their model setup, the
81 climate – ice-sheet – iceberg system was fully interactive, with the icebergs' calving positions and
82 amounts being determined by the ice sheet model, and with the ice sheet responding to the
83 icebergs' effect on climate.

84 Coupled climate-iceberg models have been used for several specific purposes, such as the
85 investigation of drift patterns of icebergs under present-day (Venkatesh and El-Tahan, 1988; Bigg
86 et al., 1996) and glacial climate conditions (Death et al., 2005). In addition, these models have
87 been utilized to study the effect of icebergs on the climate during present (e.g. Gladstone et al,
88 2001; Martin and Adcroft, 2010), pre-industrial (Jongma et al., 2009 Bügelmayer et al., 2015) and
89 past times (Levine and Bigg, 2008; Wiersma and Jongma, 2010; Green et al., 2011; Jongma et al.,
90 2013; Roberts et al., 2014) using both prescribed and interactively modelled forcing fields, and
91 have shown that icebergs and their melt water have an impact on climate. The spatial distribution

92 of the icebergs' freshwater flux is according to the atmospheric and oceanic forces acting on the
93 icebergs as they determine the icebergs' movement.

94 Computing iceberg melting and tracks is linked to various types of uncertainties. First, the
95 iceberg's drift and melting, as computed in the iceberg module, are based on empirical parameters
96 and simplifications (e.g. Jongma et al., 2009) that would need further observations to be
97 improved. Second, uncertainties in the reconstructed and modelled wind fields and ocean
98 currents, used to force the icebergs, directly affect the distribution of the freshwater. Third, the
99 initial size distribution of the icebergs is prescribed and based on present day observations
100 (Dowdeswell et al., 1992). Yet, this chosen size distribution may not be a valid representation of
101 calving events in past or future climate conditions.

102 We therefore propose in this study to extend the approach of Bügelmayer et al. (2015), evaluating
103 in detail the impact of the modelled forcing fields and iceberg size distributions. We use the same
104 earth system model of intermediate complexity (*iLOVECLIM*) coupled to an ice sheet/ice shelf
105 model (GRISLI) and an iceberg module to answer the following research questions.

106 1. How do atmospheric and oceanic forcing fields affect the icebergs (their lifetime and
107 movement) in the Northern Hemisphere under pre-industrial conditions?

108 2. How sensitive is the pre-industrial Northern Hemisphere climate and Greenland ice sheet
109 to spatial variations in the iceberg melt flux?

110 3. Do the Northern Hemisphere climate and the Greenland ice sheet respond differently to
111 icebergs of different initial size distributions?

112 4. Is the Northern Hemisphere climate and the Greenland ice sheet response to icebergs of
113 different initial size distribution dependent on the prevailing climate conditions (pre-
114 industrial (PI), warmer than PI and colder than PI)?

115 We will address these questions by presenting results from 15 different sensitivity experiments
116 (Table 1) that differ in the applied forcing (atmospheric, oceanic, pre-industrial, warmer, colder
117 climate) and the initial size distribution (CTRL (standard sizes), BIG, SMALL, Table 2) of the
118 icebergs.

119 We will first introduce the model and the experimental set-up, then present the results and the
120 discussion, followed by a conclusion section.

121 **2 Methods**

122 We use the earth system model of intermediate complexity *i*LOVECLIM (version 1.0) which is a
123 code fork of the LOVECLIM climate model version 1.2 (Goosse et al., 2010). *i*LOVECLIM differs in
124 the ice sheet module included (Roche et al., 2014) and the further developed iceberg module
125 (Bügelmayer et al., 2015), but shares some physical climate components (atmosphere, ocean and
126 vegetation) with LOVECLIM.

127 **2.1 Atmosphere – ocean – vegetation model**

128 The climate model *i*LOVECLIM consists of the atmospheric model ECBilt (Opsteegh et al., 1998), a
129 quasi-geostrophic, spectral model with a horizontal resolution of T21 (5.6° in latitude/longitude)
130 and three vertical pressure levels (800, 500, 200hPa). The atmospheric state (including e.g.,
131 temperature, humidity) is calculated every four hours. Precipitation depends on the available
132 humidity in the lowermost atmospheric level and the total solid precipitation is given to the ice-
133 sheet model at the end of one model year, as are the monthly surface temperatures.

134 *i*LOVECLIM includes the sea-ice and ocean model CLIO, which is a 3D ocean general circulation
135 model (Deleersnijder and Campin, 1995; Deleersnijder et al. 1997; Campin and Goosse, 1999)
136 including a dynamic – thermodynamic sea-ice model (Fichefet and Morales Maqueda, 1997, 1999).
137 Due to its free surface, the freshwater fluxes related to iceberg melting can be directly applied to
138 the ocean's surface. The horizontal resolution is 3°x3° in longitude and latitude and the ocean is
139 vertically divided into 20 unevenly spaced layers. CLIO uses a realistic bathymetry. The oceanic
140 variables (e.g., sea surface temperature and salinity) are computed once a day.

141 The vegetation (type and cover) is calculated by the vegetation model VECODE (Brovkin et al.,
142 1997), which runs on the same grid as ECBilt. VECODE accounts for fractional use of one grid cell
143 because of the small spatial changes in vegetation. It simulates the dynamics of two plant
144 functional types (trees and grass) as well as bare soil in response to the temperature and
145 precipitation coming from ECBilt.

146 The Antarctic ice sheet is prescribed according to present-day conditions following the ETOPO1
147 topography (<http://www.ngdc.noaa.gov/mgg/global/global.html>). Icebergs are parameterized in
148 the form of homogenous uptake of latent heat around Antarctica, thereby cooling the ocean

149 without altering the salinity. Ice shelf melting is computed according to the prevailing ocean
150 temperatures. The Greenland ice sheet is coupled actively using the GRISLI ice-sheet model.

151 **2.2 GRISLI – ice sheet model**

152 The ice-sheet model included in *iLOVECLIM* is the Grenoble model for Ice Shelves and Land Ice
153 (GRISLI), which is a three-dimensional thermomechanical model that was first developed for the
154 Antarctic (Ritz et al., 1997, 2001) and was further developed for the Northern Hemisphere (Peyaud
155 et al., 2007). GRISLI consists of a Lambert azimuthal grid with a 40x40km horizontal resolution. In
156 the present study, it computes the evolution of the thickness and extension of the Greenland ice
157 sheet (GrIS) only, as we exclude the Southern Hemisphere grid. GRISLI distinguishes three types of
158 ice flow: inland ice, ice streams and ice shelves. Calving takes place whenever the ice thickness at
159 the border of the ice sheet is less than 150 metres and the points upstream do not provide enough
160 inflow of ice to maintain this thickness. After one model year, the total yearly amount of calving is
161 given to the iceberg module where icebergs are generated daily, as described in detail in Section
162 2.3. The runoff of GRISLI is calculated at the end of the year by computing the difference between
163 the ice sheet thickness at the beginning of the model year and the end of the year, and taking into
164 account the mass loss due to calving. The runoff is then given to ECBilt where it is re-computed to
165 fit its time-step (4 hours) and incorporated into the land routing system. GRISLI is run for one
166 model year and then provides the runoff and calving, as well as the updated albedo- and
167 topography fields to the atmosphere – ocean – vegetation component. A more detailed
168 explanation of the coupling between ECBilt, CLIO and the ice sheet model GRISLI is provided in
169 Roche et al. (2014) and Bügelmayer et al. (2015).

170 **2.3 Iceberg module**

171 As discussed in detail in Bügelmayer et al. (2015), the dynamic – thermodynamic iceberg module
172 (Jongma et al., 2009; Wiersma and Jongma, 2010) included in *iLOVECLIM* is based on the iceberg-
173 drift model of Smith and co-workers (Smith and Banke, 1983; Smith, 1993; Loset, 1993) and on the
174 developments done by Bigg et al. (1996, 1997) and Gladstone et al. (2001). According to the
175 calving mass and locations calculated by GRISLI over one model year, icebergs of up to ten size
176 classes are generated. The provided ice mass is re-computed to fit the daily time-step of the
177 iceberg module, taking into account the seasonal calving cycle, with the maximum calving

178 occurring from April to June and the minimum occurring in late summer (Martin and Adcroft,
179 2010). The control size distribution of the icebergs is according to Bigg et al. (1996) and based on
180 observations of Dowdeswell et al. (1992) that represent the Greenland present day distribution
181 (Table 2). It does not take into account huge tabular icebergs as those calved from Antarctica, but
182 is a valid representation for icebergs calving from the Greenland ice sheet. The thickness and
183 width of the calving front as defined in GRISLI affects the amount of ice mass available to generate
184 icebergs, but not the icebergs' dimensions. Icebergs are moved by the Coriolis force, the air-,
185 water-, and sea-ice drag, the horizontal pressure gradient force and the wave radiation force. The
186 forcing fields are provided by ECBilt (winds) and CLIO (ocean currents) and are linearly
187 interpolated from the surrounding grid corners to the icebergs' positions. The icebergs melt over
188 time due to basal melt, lateral melt and wave erosion and may roll over as their length to height
189 ratio changes. The heat needed to melt the icebergs is taken from the ocean layers corresponding
190 to the icebergs' depth and the freshwater fluxes are put into the ocean surface layer of the current
191 grid cell. The refreezing of melted water and the break-up of icebergs is not included in the iceberg
192 module.

193 **2.4 Experimental set – up**

194 We have performed 15 sensitivity experiments that differ in the initial size distribution (CTRL /
195 SMALL / BIG, Table 2), in the applied CO₂ forcing (pre-industrial =280ppm, 4xCO₂ =1120ppm,
196 1/4xCO₂ =70ppm) or in the forces that move the icebergs (atmosphere and ocean). A summary of
197 the experiments performed is given in Table 1. All runs were started from an equilibrated climate
198 and Greenland ice sheet under pre-industrial conditions that has already been used in the study of
199 Bügelmayer et al. (2015). The initial ice sheet thickness is about 1/3 bigger than the observed one.
200 We consider this bias negligible for the present study because we focus on differences between
201 our sensitivity runs using the same initial state for all experiments. The differences between the
202 individual simulations are therefore independent of the initial conditions and only functions of the
203 different forcing applied. The model runs were conducted for 200 model years (pre-industrial) and
204 1000 model years (4xCO₂, 1/4xCO₂), respectively. The last 100 years are presented in the results.

205 **2.4.1 Iceberg Dynamical Forcing**

206 To differentiate between the impact of the ocean and the atmosphere, we separate the individual
207 forcing terms of the equation of horizontal motion (Eq. 1) of an iceberg:

208 $M \frac{dV_i}{dt} = -MfkxVi + Fa + Fr + Fw + Fp + Fs \quad (1)$

209 with M being the Mass of the iceberg, \mathbf{V} its velocity, the first term ($-Mfkx \mathbf{V}i$) on the right side
 210 corresponds to the Coriolis force, the second and third are the air drag (Fa) and wave radiation
 211 force (Fr) and therefore depend on the atmospheric winds; the last three terms represent the
 212 oceanic forcing namely water drag (Fw), horizontal pressure gradient(Fp) and sea-ice drag (Fs).

213 In the so-called “COM” experiments, the icebergs are moved according to Equation 1, thus by the
 214 combined atmospheric and oceanic forcing. In the so-called “ATM” set-up, all the forcing terms
 215 corresponding to ocean currents are set to zero, thereby ensuring that the icebergs are only
 216 moved by the Coriolis and the atmospheric forcing. In the “OCE” set-up on the contrary, the air
 217 drag and the wave radiation force are defined to be zero, thus only the Coriolis force and the
 218 ocean currents are acting on the icebergs.

219 The differentiation between atmospheric and oceanic forces was only made in the equation of
 220 motion of an iceberg. The mass balance (Jongma et al., 2009), which depends on bottom- and
 221 lateral melt (oceanic forcing) and the wave erosion (atmospheric forcing), is the same in all
 222 experiments. All the experiments are described in Table 1.

223 **2.4.2 *Iceberg Initial Size Distribution***

224 By altering the initial size distribution of the icebergs we are able to investigate the potential
 225 sensitivity of the atmosphere, ocean and ice sheet to iceberg sizes. In the CTRL experiments,
 226 depending on the available mass, icebergs of all 10 size classes can be generated (Bügelmayer et
 227 al., 2015). In the SMALL (BIG) experiments, the available mass is used to generate an equal
 228 amount of the three smallest (biggest) iceberg sizes (Table 2).

229 **2.4.3 *Radiative forcing***

230 Using the three size distributions described in 2.4.2, we performed three sets of experiments. The
 231 first set was done under pre-industrial equilibrium conditions for 200 years. In the second one, a
 232 “high” experiment, we applied a CO₂ concentration four times as strong as the pre-industrial value
 233 (1120 vs 280ppm CO₂) and in the third, a “low” experiment, only a quarter of the pre-industrial
 234 CO₂ concentration is used (70 vs 280ppm CO₂). The “high” and “low” experiments were conducted
 235 to analyse the effect of the size (CTRL/SMALL/BIG) distribution during periods of a strongly
 236 changing ice-sheet under non – equilibrated conditions.

237 **3 Results**238 **3.1 Impact of dynamical forcing and initial iceberg size on the transport**
239 **and lifetime of icebergs under pre-industrial conditions**240 **3.1.1 The CTRL experiments**

241 The distribution of the CTRL-COM's iceberg melt flux displays the general transport of icebergs of
242 all size classes due to atmospheric and oceanic forces (Fig. 1a). We find that most iceberg melt flux
243 is distributed along the eastern and western coast of Greenland, displaying that the icebergs'
244 movement follows the oceanic currents. Further, they are moved southward along the North
245 American coast and spread into the North Atlantic. In the Arctic, most icebergs are found close to
246 Ellesmere Island as indicated by the freshwater flux, due to the calving sites in this region (not
247 shown) and are then widely distributed by the Beaufort Gyre and the prevailing winds.

248 By applying only atmospheric forcing, we find that CTRL-ATM icebergs distribute their meltwater
249 further into the North Atlantic and Arctic Ocean (Fig. 1d) than seen in CTRL-COM. After calving,
250 they are quickly pushed away from the Greenland ice sheet (GrIS) margin. In CTRL-ATM less
251 icebergs than in CTRL-COM melt along the coast of Greenland, highlighting the lack of ocean
252 currents. Overall, the amount of iceberg melt flux released in CTRL-ATM (Northern Hemisphere:
253 $30 \text{ m}^3/\text{s}$, please note that this is an area weighted mean) is of the same magnitude, but distributed
254 over a broader area than in CTRL-COM (Fig. 2a). Yet, the lifetime of CTRL-ATM icebergs, that is the
255 time (in months) it takes to completely melt the icebergs, is up to one year shorter than in CTRL-
256 COM (Fig. 3) because they are transported faster away from the ice sheet and into warmer waters
257 of the North Atlantic.

258 The effect of the oceanic forcing is in strong contrast to the atmospheric one as it causes the CTRL-
259 OCE icebergs to stay closer to the GrIS margin (Fig. 1g). The icebergs melt flux reflects the
260 prevailing ocean currents, mainly the Beaufort Gyre, the East Greenland and the Labrador Current.
261 Much less icebergs are moved from the ice sheet into the Greenland – Iceland – Norwegian (GIN)
262 Seas and the North Atlantic in CTRL-OCE compared to CTRL-COM (Fig. 1a,g) due to the lack of wind
263 forcing, which is also reflected in the area that they cover (Fig. 2c,d). Also In the Arctic Ocean the
264 CTRL-OCE icebergs do not spread as much, but a slightly larger iceberg melt flux (IMF) is released
265 because the icebergs are not transported southwards by the wind, but stay and melt there.
266 Overall, the amount of freshwater flux is comparable to the CTRL-COM experiment, though over a

267 much smaller area (CTRL-COM: $2.4 \times 10^{13} \text{ m}^2$, CTRL-OCE: $1.2 \times 10^{13} \text{ m}^2$, Fig. 2b) and over a longer
268 time period. The CTRL-OCE icebergs melt up to 4 months slower than CTRL-COM icebergs because
269 they stay close to the GrIS margin and thus in colder water (Fig. 3).

270 3.1.2 *The BIG experiments*

271 The spatial distribution of the BIG-COM icebergs displays, first, the effect of the Coriolis force since
272 there is an eastward movement in the North Atlantic (Fig. 1b). The Coriolis force depends on the
273 size and velocity of the icebergs and thus, is acting stronger on big icebergs than on small ones.
274 Second, the area covered by BIG-COM icebergs is larger in the North Atlantic than in CTRL-COM
275 (Fig. 2d). Over the Northern Hemisphere the area covered by more than 10 BIG-COM icebergs is
276 only slightly bigger than the one of CTRL-COM (Fig. 2a), even though their lifetime is up to three
277 years longer (Fig. 3). But in total there are less BIG icebergs generated than in the CTRL experiment
278 because more mass is needed per berg (Table 2).

279 Applying only wind forcing on the BIG icebergs (BIG-ATM) transports less icebergs into the North
280 Atlantic and especially the GIN Seas (Fig. 1e) where they cover about half the area of BIG-COM
281 ($4 \times 10^{12} \text{ m}^2$ compared to $7 \times 10^{12} \text{ m}^2$), but release the same amount of freshwater ($150 \text{ m}^3/\text{s}$, Fig. 2c).
282 The strong southward component of the wind keeps the icebergs from drifting further into the
283 GIN Seas. Similar to the CTRL experiment, the BIG-ATM icebergs melt up to two years faster than
284 the ones of BIG-COM or BIG-OCE (Fig. 3).

285 The impact of oceanic forcing on the iceberg melt flux is simulated in BIG-OCE. Since the big
286 icebergs melt slowly, they are transported further south than CTRL-OCE icebergs (Fig. 1h). In the
287 GIN Seas the BIG-OCE icebergs are spread from the coast and cover almost the same area as the
288 BIG-ATM (Fig. 2c). In the Arctic Ocean the BIG-OCE icebergs release a higher averaged melt flux
289 than BIG-COM and BIG-ATM ($125 \text{ m}^3/\text{s}$ compared to $75 \text{ m}^3/\text{s}$ and $95 \text{ m}^3/\text{s}$, respectively; Fig. 2b), but
290 over a smaller area. This is because of the missing wind forcing which prevents the icebergs from
291 being distributed out of the Arctic Ocean. Instead the icebergs are stuck close to their calving sites.
292 The higher IMF in BIG-OCE does not strongly impact the Arctic climate because of the prevailing
293 cold conditions. Thus, more IMF, which is released to the ocean surface layer at 0°C and
294 consequently cools and freshens it, does not cause noticeable changes. The area covered by BIG
295 icebergs over the Northern Hemisphere is clearly bigger than SMALL-, or CTRL-OCE (Fig. 2a)
296 because of their lifetime, which is about two years longer compared to CTRL-OCE (Fig. 3).

297 **3.1.3 *The SMALL experiments***

298 Generating only SMALL-COM icebergs results in a similar iceberg melt flux distribution as in CTRL-
 299 COM (Fig. 1c), but less widespread. The amount of freshwater that is released by SMALL-COM
 300 icebergs is almost the same over the Northern Hemisphere as CTRL-COM, but over a smaller area
 301 (Fig. 2a) because all the SMALL-COM icebergs are melted within two years, compared to three
 302 years in CTRL-COM (Fig. 3).

303 In the icebergs' distribution of the SMALL-ATM model runs (Fig. 1f), it is clearly visible that the
 304 light, small icebergs are easily pushed away from their calving sites by the atmospheric forcing, but
 305 as in the COM experiments, over a smaller area because they melt faster. In the North Atlantic,
 306 the general pattern is directed westward, in contrast to BIG-ATM icebergs that are strongly
 307 influenced by the Coriolis force.

308 The wide-spread meltwater distribution of SMALL-ATM is in strong contrast to the one of SMALL-
 309 OCE (Fig. 1i). The oceanic forcing restricts the icebergs' transport to the shore and due to their
 310 smaller size SMALL-OCE icebergs melt before being distributed as far as CTRL-OCE and especially
 311 BIG-OCE (Fig 2a).

312 In short, the impact of the forcing fields is clearly seen in the icebergs' meltwater distribution and
 313 especially lifetime since 90% of all the atmospheric forced icebergs (SMALL-, BIG-, and CTRL-ATM)
 314 melt up to two years faster compared to the oceanic forced icebergs and compared to the
 315 icebergs of the SMALL-, BIG-, and CTRL-COM set-up.

316 **3.2 *Impact of dynamical forcing and initial iceberg size on pre-industrial
 317 climate***

318 The resulting sea surface and air temperatures (SST, TAIR) are comparable between the CTRL-
 319 COM, -ATM, and -OCE experiments (Fig. 4a,b), despite the different spatial distribution of the
 320 iceberg melt flux. The biggest spread in IMF is found in the Arctic Ocean (BIG-COM: 75m³/s, CTRL-
 321 OCE: 150m³/s, Fig. 2c), but these differences do not result in an altered climate state due to the
 322 prevailing cold conditions that are less sensitive to the freshening and cooling effect of icebergs
 323 (not shown). Also in the GIN Seas and North Atlantic the difference in SST and TAIR between the
 324 experiments does not significantly differ from internal variability (Fig. 4). In all the pre-industrial
 325 experiments, we find that the differences in air and ocean temperature between the CTRL and the
 326 BIG, SMALL experiments do not significantly exceed the internal variability of the CTRL

327 experiment. This is also the case for sensitive areas such as the GIN Seas or the North Atlantic, due
328 to the located convection sites there. Therefore, the impact of the dynamical forcing and initial
329 iceberg size is smaller than natural climate variability, which is also reflected in the deep ocean
330 circulation (not shown). This indicates that since the amount of freshwater released is comparable
331 in the model runs, the exact location of the release does not have a strong impact on the
332 prevailing climate conditions or the ocean circulation. Further, the shorter lifetime of the
333 atmospheric driven icebergs does not cause differences in the resulting climate and the GrIS
334 because the calving flux provided by GRISLI is almost constant over the years and comparable in all
335 the pre-industrial experiments. Therefore, the same amount of freshwater is supplied to the
336 ocean. Under pre-industrial equilibrium conditions the atmospheric and oceanic forcing do
337 transport the icebergs differently, but the resulting spatial patterns of the iceberg melt flux cause
338 only local differences in the Greenland ice sheet volume (Table 3) that are within the internal
339 variability of the ice sheet.

340 **3.3 Impact of initial iceberg size under a changing climate**

341 To have more confidence in using the present day iceberg distribution also for simulations of past
342 and future climates, we conducted two more sets of experiments with enhanced or reduced
343 radiative forcing to obtain warmer and colder climate states. This change in radiative forcing was
344 applied through adjustment of the atmospheric CO₂ concentration in two experiments, the so-
345 called HIGH = 4xCO₂ (1120ppm) and LOW = 1/4xCO₂ (70ppm), with a duration of 1000 years. For each
346 of these settings, we performed experiments with CTRL, BIG and SMALL icebergs. The HIGH
347 experiments resulted in an up to 4°C warmer global mean temperature and caused the Greenland
348 ice sheet to lose 10% of its volume, whereas the LOW experiments caused the mean global
349 temperatures to decrease about 4°C and an increase of the Greenland ice sheet volume of up to
350 4%, compared to the pre-industrial ice sheet volume (Table 3).

351 **3.3.1 Experiments with high radiative forcing**

352 The impact of the enhanced radiative forcing on the Greenland ice sheet is displayed in Fig. 5,
353 where the resulting CTRL-HIGH ice sheet extensions and thickness are shown (Fig. 5b).

354 As the ice sheet is shrinking and retreating from the coast (Fig. 5b), the calving flux from the GrIS is
355 decaying (0.003 Sv vs 0.02 Sv in the CTRL-COM), which is reflected in the IMF and the area that
356 they cover (Fig. 2b). The strong retreat of the ice sheet in South Greenland has a direct impact on

357 the icebergs melt flux. The released iceberg melt flux in the GIN Seas is in the range of 20 (SMALL-,
358 CTRL-HIGH) to 50m³/s (BIG-HIGH, Fig. 2c), compared to 150m³/s in the CTRL-COM. Moreover,
359 there are hardly any icebergs entering the North Atlantic, independent of the used size
360 distribution (Fig. 2d). In the Arctic Ocean the HIGH experiments result in a bigger spread between
361 the CTRL, BIG and SMALL runs than any other performed set-up (Fig. 2b). The BIG-HIGH icebergs
362 cover the smallest area because of the decreased calving flux much less BIG ones are generated.
363 Further, there are still SMALL icebergs, but due to their size and the warmer conditions they melt
364 faster than seen in the SMALL experiments performed under pre-industrial conditions. The CTRL-
365 HIGH experiment covers a slightly smaller area than the CTRL-COM,-OCE or -ATM, but much
366 bigger than BIG-, and SMALL-HIGH (Fig. 2b). This is because the different iceberg sizes allow for the
367 production of a higher number of icebergs than in BIG and the existence of icebergs bigger than
368 size 3 (as in SMALL) allows for a longer lifetime.

369 Although the size of the icebergs generated varies from the beginning, the resulting climate
370 conditions, such as sea surface or air temperatures do not vary at the end of the 1000 year period
371 between the SMALL-, BIG-, and CTRL-HIGH experiments (Fig. 4a, b), nor does the GrIS (Table 3).
372 During periods of strong background changes, different iceberg distributions do not result in
373 different climate states. This indicates that the applied forcing has a stronger impact than local
374 differences due to the chosen iceberg size.

375 3.3.2 *Experiments with low radiative forcing*

376 In contrast to the experiments with high radiative forcing, the low radiative forcing causes up to
377 4°C lower global mean temperatures and consequently the ice sheet's volume is thickening and
378 extending further down to the coast line (Fig. 5c), especially along the western margin and in
379 South Greenland. Similar to the other experiments performed, the impact of different initial size
380 distributions of the icebergs is negligible on the resulting climate and ice sheet volume (Table 3).

381 Due to the increased ice sheet thickness, more calving flux is released (0.05 Sv in CTRL-LOW
382 compared to 0.02 Sv in CTRL-COM) and so the iceberg melt flux increases to ~40m³/s over the
383 Northern Hemisphere, compared to 15m³/s in the pre-industrial experiments. The increase is seen
384 almost everywhere around Greenland (Fig. 2), except in the Arctic Ocean. In the Arctic Ocean the
385 released IMF is in the same range as in the experiments performed under pre-industrial conditions
386 because the ice sheet's thickness and consequently the calving sites in North Greenland are not
387 strongly altered by the colder climate (Fig. 5c). In the North Atlantic the released iceberg melt flux

388 displays a big spread between the experiments with the BIG-LOW icebergs being spread the
389 furthest and releasing the most IMF (80m³/s in BIG-LOW vs 45m³/s in CTRL-LOW; Fig. 2d). Since
390 the cold conditions prevent the BIG-LOW icebergs from melting quickly, almost all of them are
391 transported into the North Atlantic where they finally melt. This is also partly the case for the
392 CTRL-LOW icebergs thereby resulting in a higher iceberg melt flux than the SMALL-LOW (Fig. 2e).
393 Independent of the chosen size distribution, the resulting temperatures are about 5°C lower than
394 during pre-industrial conditions in the North Atlantic and the GIN Seas (Fig. 4), displaying the
395 strong CO₂ forcing.

396 During a strongly changing climate, the initial size distribution does not alter the climate response
397 (temperatures, ocean circulation) stronger than internal variability. The BIG-LOW set-up causes a
398 slightly larger mean ice sheet volume at the end of the 1000 years (Table 3), which indicates that
399 the extreme case of BIG icebergs impacts the resulting ice sheet thickness, even though the
400 climate conditions are similar to the CTRL- and SMALL-LOW runs.

401 **4 Discussion**

402 By testing the impact of the atmospheric versus the oceanic forcing on the lifetime and motion of
403 icebergs, we find that the atmospheric forcing causes the icebergs to travel further away from
404 their calving sites and into the North Atlantic, whereas the ocean currents lead to iceberg tracks
405 closer to shore. It is difficult to compare our results to previous studies, since the studies that
406 investigated the impact of the background forcing (Smith 1993; Keghouche et al., 2002) focused
407 on observations of single icebergs and the ability of models reproducing their specific tracks. Bigg
408 et al. (1997) noted that the modelling of specific iceberg tracks is very unlikely to be successful and
409 it is important to notice that we do not expect our model to resolve single tracks due to its coarse
410 resolution, but to reflect the wide spread effect of icebergs on climate.

411 In our model, the impact of icebergs on climate does not strongly depend on the two types of
412 forcing (atmospheric and oceanic), yet their lifetime is shortened by up to two years when they
413 are transported by atmospheric forces only. Bigg et al. (1997) showed that about 80% of the small
414 icebergs of up to 200 m diameter (size class 1 to 3, Table 2) melt within the first year, which is
415 higher than in our SMALL-COM set-up where about 60% are melted. Also Venkatesh and El-Tahan
416 (1988) conducted a study to investigate the impact of modelling complete deterioration of
417 icebergs on the prediction of their tracks. In their study they showed that most of the icebergs
418 corresponding to size class 1 to 3 used in this study, disappear within 3 to 22 months, consistent

419 with our results. The maximum lifetime of the BIG icebergs is found to be almost seven years,
420 which is slightly longer than modelled by Bigg et al. (1997). This discrepancy can be due to the pre-
421 industrial climate conditions used in our study that are slightly colder than the present day
422 conditions applied by Bigg et al. (1997).

423 To better understand the response of the modelled climate to the initial size distribution, we
424 performed different sensitivity experiments. First, using pre-industrial conditions we find that
425 independent of the forcing, SMALL icebergs release less freshwater and spread over a smaller area
426 than BIG and CTRL icebergs. In the North Atlantic the impact of the Coriolis force is especially
427 pronounced in the BIG-ATM and BIG-COM runs, confirming the findings of Roberts et al. (2014). In
428 their study they noted that BIG icebergs travel further south than small icebergs due to the
429 stronger impact of the Coriolis force. Even though the SMALL icebergs cause locally different
430 ocean and atmospheric conditions than the BIG icebergs, the overall effect on climate and on the
431 Greenland ice sheet is within the natural climate variability.

432 Second, we repeated the experiments under a strongly increased and decreased radiative forcing
433 for 1000 years. During this time scale changes in the Southern Ocean can impact the Northern
434 Hemisphere. Jongma et al. (2009) showed that including active icebergs increases the net
435 production of Antarctic Bottom Water by 10% under pre-industrial conditions. We do neglect this
436 direct effect of icebergs here since icebergs and Antarctic ice-sheet runoff are computed using
437 parameterizations that depend on the prevailing climate conditions. Concerning the icebergs
438 released from Greenland, we do not expect that the size of the icebergs have an impact on the
439 Southern Hemisphere through altered ocean circulation because the Atlantic Meridional
440 Overturning Circulation is comparable within all the experiments (not shown). Thus, the
441 uncertainty introduced by not actively coupling the Antarctic ice sheet is comparable in all the
442 radiative forcing experiments.

443 There might be different reasons why the climate conditions and the GrIS are not strongly affected
444 by the initial size distribution during strong radiative background conditions. One reason could be
445 that the ice sheet and the climate model are too insensitive to the experienced changes as they
446 have a relatively coarse resolution. Therefore, it would be interesting to repeat this study with a
447 finer model grid. Another reason might be that in the experiments where really strong forcing was
448 applied (HIGH=1120ppm CO₂, LOW= 70ppm CO₂), the feedbacks related to calving have a smaller
449 signal than the forcing and are therefore overruled.

450 **5 Conclusions**

451 Within a fully coupled climate – ice sheet – iceberg model set up, we have performed sensitivity
452 experiments to investigate the effect of the forcing fields such as winds and ocean currents, as
453 well as the prescribed initial size distribution on the icebergs and the climate.

454 We find that, under pre-industrial conditions, the wind forcing pushes the icebergs further away
455 from their calving sites and further into the North Atlantic, whereas the ocean currents transport
456 the icebergs close to Greenland and southward along the North American coast. The combined
457 effect of the forces (control set-up) displays a lesser spread iceberg distribution in the Arctic Ocean
458 and into the North Atlantic than the purely atmospheric driven icebergs due to the restrictive
459 effects of the oceanic forcing. The spread of icebergs depends on both the forcing fields and the
460 icebergs size with the CTRL icebergs being transported the furthest, followed by the BIG icebergs.
461 The amount of released iceberg melt flux is comparable in all the experiments, though locally
462 different. In our model set-up, the biggest impact of the applied forcing (atmospheric or oceanic)
463 is on the icebergs' lifetime which is up to two years shorter if the icebergs are only transported by
464 winds.

465 In the presented model framework, the implementation of icebergs of different size classes under
466 equilibrated pre-industrial conditions reveals that there are local differences in the released
467 freshwater flux. However, these differences do not cause significant changes in the resulting
468 Greenland ice sheet volume and climate conditions.

469 When repeating the experiments with different size distributions with strong radiative cooling or
470 warming (1120 ppm or 70 ppm CO₂, 1000 model years), the response of the climate and the ice
471 sheet volume are within the climate variability.

472 Even though the iceberg and freshwater distribution differ between the conducted experiments
473 (all size classes, only SMALL, less than 200m width, and only BIG icebergs, 600-1000m width,
474 respectively), their impact on the Northern Hemispheric climate does not differ significantly from
475 internal variability. We can therefore conclude that for the resulting climate and ice sheet small
476 spatial differences between the runs do not have a strong impact as long as there is a wide spread
477 impact of icebergs (cooling and freshening) around Greenland. Furthermore, our results show that
478 the response of the climate to the applied radiative forcing is much stronger than its response to
479 the chosen initial size distribution of the icebergs.

480 Code availability

481 The iLOVECLIM source code is based on the LOVECLIM model version 1.2 whose code is accessible
 482 at <http://www.elic.ucl.ac.be/modx/elic/index.php?id=289>. The developments on the iLOVECLIM
 483 source code are hosted at <https://forge.ipsl.jussieu.fr/ludus>, but are not publicly available due to
 484 copyright restrictions. Access can be granted on demand by request to D. M. Roche
 485 (didier.roche@lsce.ipsl.fr). The specific experimental set-up used for this study is available at
 486 <https://forge.ipsl.jussieu.fr/ludus>.

487 Acknowledgements. M. Bügelmayer is supported by NWO through the VIDI/AC2ME project no
 488 864.09.013. D. M. Roche is supported by NWO through the VIDI/AC2ME project no 864.09.013 and
 489 by CNRS-INSU. The authors wish to thank Catherine Ritz for the use of the GRISLI ice sheet model
 490 and all the anonymous reviewers for providing valuable comments. Institut Pierre Simon Laplace is
 491 gratefully acknowledged for hosting the *iLOVECLIM* model code under the LUDUS framework
 492 project (<https://forge.ipsl.jussieu.fr/ludus>). This is NWO/AC2ME contribution number 08.

493 REFERENCES

494 Álvarez-Solas, J., Montoya, M., Ritz, C., Ramstein, G., Charbit, S., Dumas, C., Nisancioglu, K.,
 495 Dokken, T., and Ganopolski, A.: Heinrich event 1: an example of dynamical ice-sheet reaction to
 496 oceanic changes, *Clim. Past*, 7, 1297–1306, doi:10.5194/cp-7-1297-2011, 2011.

497 Andrews, J. T.: Abrupt changes (Heinrich events) in late Quaternary North Atlantic marine
 498 environments: a history and review of data and concepts, *Journal of Quaternary Science*, 13, 3–16,
 499 doi:10.1002/(SICI)1099-1417(199801/02)13:1<3::AID-JQS361>3.0.CO;2-0, 1998.

500 Bigg, G. R., Wadley, M. R., Stevens, D. P. and Johnson, J.V.: Prediction of iceberg trajectories for
 501 the North Atlantic and Arctic Oceans, *Geophysical Research Letters*, 23, 3587–3590, 1996.

502 Bigg, G.R., Wadley, M. R., Stevens, D. P. and Johnson, J. V.: Modelling the dynamics and
 503 thermodynamics of icebergs, *Cold Regions Science and Technology*, 26, 113–135,
 504 doi:10.1016/S0165-232X00012-8, 1997.

505 Box, J. E.: Greenland ice sheet mass balance reconstruction. Part II: surface mass balance (1840–
 506 2010), *J. Clim.*, 26, 6974–6989, doi:10.1175/JCLI-D-12-00518.1, 2013.

507 Broecker, W.S., Bond, G., McManus, J.: Heinrich events: Triggers of ocean circulation change? In:
 508 Peltier WR (ed) *Ice in the climate system: NATO ASI Series*, Vol 12, Springer-Verlag, Berlin, pp 161–
 509 166, doi: 10.1007/978-3-642-85016-5_10, 1993.

510 Brovkin, V., Ganopolski, A., and Svirszhev, Y.: A continuous climate-vegetation classification for use
 511 in climate-biosphere studies, *Ecological Modelling*, 101, 251–261, 1997.

512 Bügelmayer, M., Roche, D. M. and Renssen, H.: How do icebergs affect the Greenland ice sheet
513 under pre-industrial conditions? – A model study with a fully coupled ice sheet–climate model,
514 *Cryosph. Discuss.*, 8(1), 187–228, doi:10.5194/tcd-8-187-2014, 2014.

515 Campin, J. M. and Goosse, H.: A parameterization of density driven downsloping flow for coarse
516 resolution model in z-coordinate, *Tellus*, 51A, 412–430, 1999.

517 Death, R., Siegert, M. J., Bigg, G. R. and Wadley, M. R.: Modelling iceberg trajectories,
518 sedimentation rates and meltwater input to the ocean from the Eurasian Ice Sheet at the Last
519 Glacial Maximum, *Palaeogeography, Palaeoclimatology, Palaeoecology*, 236, 135–150,
520 doi:10.1016/j.palaeo.2005.11.040, 2005.

521 Deleersnijder, E., Beckers, J.-M., Campin, J.-M., El Mohajir, M., Fichefet, T., and Luyten, P.: Some
522 mathematical problems associated with the development and use of marine models, in: *The*
523 *mathematics of model for climatology and environment*, edited by: Diaz, J. I., *NATO ASI Series*, vol.
524 I 48, Springer-Verlag, pp. 39–86, 1997.

525 Deleersnijder, E. and Campin, J.-M.: On the computation of the barotropic mode of a free-surface
526 world ocean model, *Annales Geophysicae*, 13, 675–688, 1995.

527 Dowdeswell, J. A., Whittington, R. J. and Hodgkins, R.: The Sizes, Frequencies, and Freeboards of
528 East Greenland Icebergs Observed Using Ship Radar and Sextant, *J. Geophys. Res.*, 97, 3515–3528,
529 1992.

530 Fichefet, T. and Morales Maqueda, M. A.: Sensitivity of a global sea ice model to the treatment of
531 ice thermodynamics and dynamics, *Journal of Geophysical Research*, 102, 12609–12646, 1997.

532 Fichefet, T. and Morales Maqueda, M. A.: Modelling the influence of snow accumulation and
533 snow-ice formation on the seasonal cycle of the Antarctic sea-ice cover, *Climate Dynamics*, 15,
534 251–268, 1999.

535 Gherardi, J., Labeyrie, L., McManus, J., Francois, R., Skinner, L. and Cortijo, E.: Evidence from the
536 Northeastern Atlantic basin for variability in the rate of the meridional overturning circulation
537 through the last deglaciation, *Earth Planet. Sci. Lett.*, 240, 710–723,
538 doi:10.1016/j.epsl.2005.09.061, 2005.

539 Gladstone, R. M., Bigg, G. R. and Nicholls, K. W.: Iceberg trajectory modeling and meltwater
540 injection in the Southern Ocean, *Journal of Geophysical Research*, 106, 19903–19915,
541 doi:10.1029/2000JC000347, 2001.

542 Goosse, H., Brovkin, V., Fichefet, T., Haarsma, R., Huybrechts, P., Jongma, J., Mouchet, A., Selten,
543 F., Barriat, P.-Y., Campin, J.-M., Deleersnijder, E., et al.: Description of the Earth system model of
544 intermediate complexity LOVECLIM version 1.2, *Geoscientific Model Development*, 3, 603–633,
545 doi:10.5194/gmd-3-603-2010, 2010.

546 Green, C. L., Green, J. A. M. and Bigg, G. R.: Simulating the impact of freshwater inputs and deep-
547 draft icebergs formed during a MIS 6 Barents Ice Sheet collapse, *Paleoceanography*, 26, 1–16,
548 doi:10.1029/2010PA002088, 2011.

549 Hemming, S. R.: Heinrich Events: Massive Late Pleistocene Detritus Layers of the North Atlantic
550 and their global climate imprint, *Marine Geology*, Doi:10.1029/2003rg000128.1, 2004.

551 Hooke, R.L.: *Principles of Glacier Mechanics*, second ed. Cambridge University Press, 2005.

552 Hulbe, C. L., MacAyeal, D. R., Denton, G. H., Kleman, J. and Vowell, T. V.: Catastrophic ice shelf
553 breakup as the source of Heinrich event icebergs, *Paleoceanography*, 19(1), 1–16,
554 doi:10.1029/2003PA000890, 2004.

555 Jongma, J.I., Driesschaert, E., Fichefet, T., Goosse, H. and Renssen, H.: The effect of dynamic–
556 thermodynamic icebergs on the Southern Ocean climate in a three-dimensional model, *Ocean*
557 *Modelling*, 26, 104–113, doi:10.1016/j.ocemod.2008.09.007, 2009.

558 Jongma, J.I., Renssen, H. and Roche, D.M.: Simulating Heinrich event 1 with interactive
559 icebergs, *Climate Dynamics*, 40, 1373–1385. doi:10.1007/s00382-012-1421-1, 2013.

560 Kageyama, M., Paul, A., Roche, D. M. and Van Meerbeeck, C. J.: Modelling glacial climatic
561 millennial-scale variability related to changes in the Atlantic meridional overturning circulation: a
562 review, *Quat. Sci. Rev.*, 29, 2931–2956, doi:10.1016/j.quascirev.2010.05.029, 2010.

563 Keghouche, I., Bertino, L. and Lisæter, K. A.: Parameterization of an Iceberg Drift Model in the
564 Barents Sea, *J. Atmos. Ocean. Technol.*, 26(10), 2216–2227, doi:10.1175/2009JTECHO678.1, 2009.

565 Levine, R. C., and G. R. Bigg, 2008, The sensitivity of the glacial ocean to Heinrich events from
566 different sources, as modeled by a coupled atmosphere-iceberg- ocean model, *Paleoceanogr.*, 23:
567 PA4213, doi:10.1029/2008PA001613

568 Loset, S.: Thermal-Energy Conservation in Icebergs and Tracking by Temperature, *Journal of*
569 *Geophysical Research Oceans*, 98, 10001–10012, 1993.

570 MacAyeal, D.: Binge/purge oscillations of the Laurentide ice sheet as a cause of the North
571 Atlantic's Heinrich events, *Paleoceanography*, 8, 775–784, 1993.

572 Marcott, S. A, Clark, P. U., Padman, L., Klinkhammer, G. P., Springer, S. R., Liu, Z., Otto-Bliesner, B.
573 L., Carlson, A. E., Ungerer, A., Padman, J., He, F., Cheng, J. and Schmittner, A.: Ice-shelf collapse
574 from subsurface warming as a trigger for Heinrich events., *Proc. Natl. Acad. Sci. U. S. A.*, 108,
575 13415–9, doi:10.1073/pnas.1104772108, 2011.

576 Martin, T. and Adcroft, A.: Parameterizing the fresh-water flux from land ice to ocean with
577 interactive icebergs in a coupled climate model, *Ocean Modelling*, 34, 111–124,
578 doi:10.1016/j.ocemod.2010.05.001, 2010.

579 McManus, J. F., Francois, R., Gherardi, J.-M., Keigwin, L. D. and Brown-Leger, S.: Collapse and rapid
580 resumption of Atlantic meridional circulation linked to deglacial climate changes., *Nature*, 428,
581 834–7, doi:10.1038/nature02494, 2004.

582 Opsteegh, J. D., Haarsma, R. J., Selten, F. M., and Kattenberg, A.: ECBilt: A dynamic alternative to
583 mixed boundary conditions in ocean models, *Tellus*, 50A, 348–367, 1998.

584 Peyaud, V., Ritz, C. and Krinner, G.: Modelling the Early Weichselian Eurasian Ice Sheets: role of ice
585 shelves and influence of ice-dammed lakes, *Climate of the Past*, 3, 375–386, doi:10.5194/cp-3-375-
586 2007, 2007.

587 Ritz, C., Fabre, A. and Letréguilly, A.: Sensitivity of a Greenland ice sheet model to ice flow and
588 ablation parameters: consequences for the evolution through the last climatic cycle, *Climate*
589 *Dynamics*, 13, 11–23, doi:10.1007/s003820050149, 1997.

590 Ritz, C., Rommelaere, V. and Dumas, C.: Modeling the evolution of Antarctic ice sheet over the last
 591 420,000 years: Implications for altitude changes in the Vostok region, *Journal of Geophysical*
 592 *Research*, 106, 31943–31964, doi:10.1029/2001JD900232, 2001.

593 Roberts, W. H. G., Valdes, P. J., and Payne, A. J.: A new constraint on the size of Heinrich Events
 594 from an iceberg/sediment model. *Earth and Planetary Science Letters*, 386, 1–9.
 595 doi:10.1016/j.epsl.2013.10.020, 2014.

596 Roche, D.M., Dumas, C., Bügelmayer, M., Charbit, S., Ritz, C.: Adding a dynamical cryosphere into
 597 iLOVECLIM (version 1.0) – Part 1: Coupling with the GRISLI ice-sheet model, *Geoscientific Model*
 598 *Development Discussion*, 6, 5215–5249, doi:10.5194/gmdd-6-5215-201, 2013.

599 Smith, K. L.: Free-drifting icebergs in the Southern Ocean: An overview, *Deep Sea Res. Part II Top.*
 600 *Stud. Oceanogr.*, 58(11–12), 1277–1284, doi:10.1016/j.dsr2.2010.11.003, 2011.

601 Smith, S.D.: Hindcasting iceberg drift using current profiles and winds, *Cold Reg. Sci. Technol.*,
 602 22(1), 33–45, doi:10.1016/0165-232X(93)90044-9, 1993.

603 Smith, S.D., Banke, E.G.: The influence of winds, currents and towing forces on the drift of
 604 icebergs. *Cold Regions Science and Technology*, 6, 241– 255, 1983.

605 Tournadre, J., Girard-Ardhuin, F. and Legrésy, B.: Antarctic icebergs distributions, 2002–2010, *J.*
 606 *Geophys. Res.*, 117(C5), C05004, doi:10.1029/2011JC007441, 2012.

607 Venkatesh, S. and El-Tahan, M.: Iceberg life expectancies in the Grand Banks and Labrador Sea,
 608 *Cold Reg. Sci. Technol.*, 15, 1–11, 1988.

609 Wiersma, A. P. and Jongma, J. I.: A role for icebergs in the 8.2 ka climate event, *Climate Dynamics*,
 610 35, 535–549, doi:10.1007/s00382-009-0645-1, 2010.

611

612 LIST OF TABLES

	PRE- INDUSTRIAL (ATM & OCE FORCING) = 280 ppm	ONLY ATMOS- PHERIC FORCING	ONLY OCEANIC FORCING	4xCO ₂ (ATM & OCE FORCING) = 1120ppm	1/4xCO ₂ (ATM & OCE FORCING) = 70ppm
ALL SIZES	CTRL-COM	CTRL-ATM	CTRL-OCE	CTRL-HIGH	CTRL-LOW
BIG ICEBERGS	BIG-COM	BIG-ATM	BIG-OCE	BIG-HIGH	BIG-LOW
SMALL ICEBERGS	SMALL-COM	SMALL-ATM	SMALL-OCE	SMALL-HIGH	SMALL-LOW

613 Table 1: performed experiments

CLASS	HEIGHT (m)	WIDTH (m)	VOLUME (m ³) 1E+05	FRACTION of total available Volume	EXPERIMENT
1	67	67	5.16	0.15 / 0.33	CTRL / SMALL
2	133	133	4.07	0.15 / 0.33	CTRL / SMALL
3	200	200	138	0.2 / 0.33	CTRL / SMALL

4	267	267	328	0.15	CTRL
5	300	333	574	0.08	CTRL
6	300	400	828	0.07	CTRL
7	300	500	1297	0.05	CTRL
8	300	600	1860	0.05 / 0.33	CTRL / BIG
9	300	800	3310	0.05 / 0.33	CTRL / BIG
10	300	1000	5180	0.05 / 0.33	CTRL / BIG

614 Table 2: used initial iceberg classes

615

ICE SHEET THICKNESS	Mean (1E+15)	STDEV (1E+12)	Difference (1E+12)	Difference in %
CTRL-COM	3.90	4.04*	-	-
BIG-COM	3.91	2.61	-3.50	-0.09
SMALL-COM	3.91	1.96	-2.97	-0.08
CTRL-ATM	3.91	2.79*	-	-
BIG-ATM	3.91	2.14	-2.58	-0.07
SMALL-ATM	3.91	1.99	-0.430	0,01
CTRL-OCE	3.91	3.18*	-	-
BIG-OCE	3.91	1.29	-1.20	-0.03
SMALL-OCE	3.91	2.20	-5.63	-0.14
CTRL-HIGH	3.50	5.03	-	-
BIG-HIGH	3.49	4.40	11.0	0.32
SMALL-HIGH	3.49	5.69	4.82	0.14
CTRL-LOW	4.04	1.90	-	-
BIG-LOW	4.06	2.74	-16.6	-0.41
SMALL-LOW	4.04	3.20	-1.85	-0.05

616

617 Table 3: Ice-sheet Volume (m^3): Mean and Standard deviation of last 100 years, the *corresponds
 618 to the CTRL stdev that was computed over the last 200 years to have a more representative range
 619 of internal variability as a reference; difference between the ice sheet volume of the CTRL
 620 experiment and the BIG/SMALL experiments in absolute numbers, if the value is above $2 \times$ stdev of
 621 the CTRL experiments (*), then the difference is significantly different from internal variability
 622 (none of the experiments); % diff = difference between the ice sheet volume of the CTRL
 623 experiment and the BIG/SMALL experiments in percent.

624 FIGURE CAPTIONS

625 Figure 1: Iceberg melt flux (m^3/s); first row: atmospheric forcing only (CTRL-, BIG-, SMALL-ATM);
 626 second row: oceanic forcing only (CTRL-, BIG-, SMALL-OCE), third row: the default set-up (icebergs
 627 are moved by both, atmospheric and oceanic forcing; CTRL-, BIG-, SMALL-COM)










628 Figure 2: area (m^2) vs area weighted iceberg melt flux (m^3/s); the area is computed by taking into
 629 account all the grid cells that have at least 10 icebergs passing through per year (be aware that the
 630 area is $10^{13} m^2$ in panel a, $10^{12} m^2$ otherwise); a: Northern Hemisphere: mean computed over 0-
 631 $90^\circ N$ and $180^\circ W-180^\circ E$, values of IMF: $0-40 m^3/s$ (area weighted IMF); b: Arctic Ocean: $80-90^\circ N$ and





632 180°W-180°E, values of IMF: 60-140 m³/s; c: Greenland – Iceland – Norwegian (GIN) Seas: 50-85°N
633 and 45°W-15°E, values of IMF: 40-240 m³/s; d: North Atlantic: 45-60°N and 60-20°W, values of
634 IMF: 0-50 m³/s;

635 Figure 3: cumulative iceberg melt distribution normalized to 100% as a function of time (months);
636 x – Axis corresponds to months, y-axis to cumulative percentage

637 Figure 4: Mean + Standard deviation of last 100 years of the performed experiments: Sea Surface
638 Temperature (SST, °C) and air temperature (TAIR, °C); red = BIG icebergs, blue = CTRL, green =
639 SMALL icebergs; a: North Atlantic: mean computed over: 45-60°N and 60-20°W; b: Greenland –
640 Iceland – Norwegian (GIN) Seas: 50-85°N and 45°W-15°E


641 Figure 5: a: CTRL-COM ice sheet thickness at the end of the experiments (m); b: difference in ice
642 sheet thickness at the end of the model runs CTRL-COM minus CTRL-HIGH; c: difference in ice
643 sheet thickness at the end of the model runs CTRL-COM minus CTRL-LOW






◆ ATM
◆ COM
◆ OCE
◆ HIGH
◆ LOW
◆ BIG
▲ CTRL
■ SMALL


Lifetime of Icebergs




a) North Atlantic (last 100years)




b) GIN Seas (last 100years)



a) CTRL-COM



b) CTRL-COM - CTRL-HIGH



c) CTRL-COM - CTRL-LOW

