

1 **Response to referee J.P. Evans**

2 We updated our text and included in the references-list the suggested publications with
3 respect to the importance of the PBL scheme (page 13, l7-11)

4 Technical comments:

5 Minor corrections for comments 3-6; 8-16 in the revised manuscript

6 Comment 7: All figures which were not shown, are now added as a supplementary material.

7

8 **Response to anonymous referee #2**

9 1) The reviewer wishes to discuss more the issue on the relation between radiation and
10 temperature, including maximum and minimum values. In our revised manuscript we
11 added the temperature-maximum and minimum variables in the discussion (page 9,
12 lines 10-19; page 16, lines 1-5) and included two additional figures in the
13 supplementary material Fig S1 (tas-max) and Fig S2 (tas-min). We shortly discuss the
14 potential relationship with short- and long wave radiation.

15 2) We updated our references-list with the missing references.

16 3) Figures 2, 4 and 5 were updated, as requested

17 4) Several technical corrections and the required clarifications were added in the
18 manuscript

19 5) Table 1 was updated with the requested information on relaxation zone and missing
20 acronyms.

21 6) Details of the post-processing of the cloud cover was added in 2.2 section (page 7 of
22 the revised manuscript)

23

1 Hindcast regional climate simulations within EURO- 2 CORDEX: Evaluation of a WRF multi-physics ensemble.

3

4 E. Katragkou^{1*}, M. García-Díez^{2,3}, R. Vautard⁴, S. Sobolowski⁵, P. Zanis¹, G.
5 Alexandri⁶, R. M. Cardoso⁷, A. Colette⁸, J. Fernandez³, A. Gobiet^{9,10}, K.
6 Goergen¹¹ Goergen^{10,12,11,13,12}, T. Karacostas¹, S. Knist¹² Knist¹⁰, S. Mayer⁵, P. M.
7 M. Soares⁷, I. Pytharoulis¹, I. Tegoulis¹, A. Tsikerdekis¹, D. Jacob¹⁴ Jacob¹³

8

9 [1]{Department of Meteorology and Climatology, School of Geology, Aristotle University of
10 Thessaloniki, Thessaloniki, Greece}

11 [2]{Climate Dynamics and Impacts Unit, Institut Català de Ciències del Clima, Barcelona,
12 Catalonia, Spain Instituto de Física de Cantabria, CSIC–UC, Santander, Spain}

13 [3]{Department of Applied Mathematics and Computer Science, Universidad de Cantabria,
14 Santander, Spain}

15 [4]{Laboratoire des Sciences du Climat et de l'Environnement, IPSL, CEA/CNRS/UVSQ, Gif
16 sur Yvette, France}

17 [5]{Uni Research Climate, & Bjerknes Center for Climate Research, Bergen, Norway}

18 [6]{Laboratory of Atmospheric Physics, School of Physics, Aristotle University of
19 Thessaloniki, Thessaloniki, Greece}

20 [7]{Instituto Dom Luiz, Faculdade de Ciências, Universidade de Lisboa, 1749-016 Lisboa,
21 Portugal}

22 [8]{Institut National de l' Environnement industriel et des risques (INERIS), Verneuil en
23 Halatte, France}

24 ~~[9]{Department of Applied Mathematics and Computer Science, Universidad de Cantabria,
25 Santander, Spain}~~

26 [109]{Wegener Center for Climate and Global Change, University of Graz, Austria}

27 [1210]{Meteorological Institute, University of Bonn, Germany}

28 [1311]{Jülich Supercomputing Centre, Forschungszentrum Jülich GmbH, Jülich, Germany}

1 [12]{Luxembourg Institute of Science and Technology (as of 01/2015) - former Centre de
2 Recherche Public – Gabriel Lippmann (CRPGL), Belvaux, Luxembourg }

3 [~~14~~13]{Climate Service Centre 2, Hamburg, Germany }

4 Correspondence to: E. Katragkou (katragou@auth.gr)

5

6 **Abstract**

7 In the current work we present six hindcast WRF simulations for the EURO-CORDEX
8 domain with different configurations in microphysics, convection and radiation for the time
9 period 1990-2008. All regional model simulations are forced by the ERA-Interim reanalysis
10 and have the same spatial resolution (0.44°). These simulations are evaluated for surface
11 temperature, precipitation, short- and longwave downward radiation at the surface and total
12 cloud cover. The analysis of the WRF ensemble indicates systematic biases in both
13 temperature and precipitation linked to different physical mechanisms for the summer and
14 winter season. Overestimation of total cloud cover and underestimation of downward
15 shortwave radiation at the surface, mostly when using Grell-Devenyi convection and the
16 CAM radiation scheme, intensifies the negative summer temperature bias in northern Europe
17 (max -2.5°C). Conversely, a strong positive downward shortwave summer bias in central (40-
18 60%) and southern Europe mitigates the systematic cold bias in WRF over these regions,
19 signifying a typical case of error compensation. Maximum winter cold bias is over north-
20 eastern Europe (-2.8°C); this location is indicative of land-atmosphere rather than cloud-
21 radiation interactions. Precipitation is systematically overestimated in summer by all model
22 configurations, especially the higher quantiles, which are associated with summertime deep
23 cumulus convection. The Kain-Fritsch convection scheme produces the larger summertime
24 precipitation biases over the Mediterranean. Winter precipitation is reproduced with lower
25 biases by all model configurations (15-30%). The results of this study indicate the importance
26 of evaluating not only the basic climatic parameters of interest for climate change applications
27 (temperature-precipitation), but also other components of the energy and water cycle, in order
28 to identify the sources of systematic biases, possible compensatory or masking mechanisms
29 and suggest methodologies for model improvement.

30

1 1 Introduction

2 Climate models are the primary tools for investigating the response of the climate system to
3 various forcings, making climate predictions on seasonal to decadal time scales and
4 projections of future climate. Regional climate models (RCMs) are applied over limited-area
5 domains with boundary conditions either from global reanalysis or global climate model
6 output. The use of RCMs for dynamical downscaling has grown, their resolution has
7 increased, process-descriptions have developed further, new components have been added,
8 and coordinated ensemble experiments have become more widespread (Rummukainen 2010;
9 Flato et al. 2013). A significant constraint in a comprehensive evaluation of regional
10 downscaling is that available studies often employ different methods, regions, periods and
11 observational data for evaluation. Thus, evaluation results are difficult to generalize. The
12 Coordinated Regional Climate Downscaling Experiment (CORDEX) initiative provides a
13 platform for a joint evaluation of model performance, along with a solid scientific basis for
14 impact assessments and other uses of downscaled climate information (Giorgi et al. 2009).

15 Published work within CORDEX focusing on the European domain (EURO-CORDEX) for
16 present climate, indicates strengths and deficiencies of the state-of-the-art modeling tools,
17 already used to downscale the Coupled Model Intercomparison Project Phase 5 (CMIP5)
18 global model results (Taylor et al., 2012). Kotlarski et al. (2014), in a joint evaluation based
19 on the EURO-CORDEX RCM ensemble, reported bias ranges for temperatures and
20 precipitation corresponding to those of the ENSEMBLES simulations (van der Linden et al.
21 2009) with some improvements identified and strong influence of specific choices of model
22 configuration on model performance. Vautard et al. (2013), focusing on the European
23 heatwaves with the EURO-CORDEX ensemble, found that high temperatures are primarily
24 sensitive to convection and micro-physics. ~~Comparison of WRF multi-physics ensemble
25 versus the multi-model EURO-CORDEX ensemble by Garcia-Diez et al. (2014a) suggests
26 that WRF multi-physics and multi-model ensemble spreads within EURO-CORDEX were of
27 the same order.~~ Giorgi et al. (2012) highlighted the significant sensitivity of model
28 performance on different parameterization schemes and parameter settings in a RegCM4
29 model study over different CORDEX domains including Europe.

30 These findings indicate that combining model evaluation with sensitivity studies is necessary
31 in order to investigate recurring and persistent biases, list potential sources of their origin,
32 dissuade/encourage modelers from using specific configurations responsible for systematic

1 errors over specific regions and suggest tracks for model development. Since large model
2 ensemble spreads and present climate biases are potentially linked with future climate
3 uncertainties (Boberg and Christensen., 2012), it is important to understand contributions of
4 individual processes on the present European climate in order to be able to interpret future
5 climate projections with greater confidence and possibly constrain these projections (Hall and
6 Qu 2006; Stegehuis et al., 2013).

7 In the current work we analyze hindcast simulations of the Weather Research and Forecasting
8 model (WRF) multi-physics ensemble performed within the framework of EURO-CORDEX.
9 Recent research has demonstrated the ability to use WRF (Skamarock et al 2008) to refine
10 global climate modeling results to higher spatial resolutions in Europe (e.g. Soares et al, 2012;
11 Cardoso et al., 2013; Warrach-Sagi et al., 2013). The aim of this study is to identify
12 systematic biases and areas of large uncertainties in present European climate and relate them
13 to specific physical processes (e.g. cloud-radiation or land-atmosphere interactions). This
14 analysis contributes towards a better understanding of WRF as a dynamical downscaling tool
15 for RCM modeling studies and its optimization for this specific region.

16

17 **2 Data and methodology**

18 **2.1 Observations**

19 To evaluate the model simulations we use daily mean, **minimum and maximum** temperature
20 and precipitation values from E-OBS version 9.0 (hereafter **EOBSE-OBS9**) covering the area
21 25–75N and 40W–75E, available on a 0.44 degree rotated pole grid (Haylock et al., 2008).
22 The E-OBS dataset is based on the ECA&D (European Climate Assessment and Data) station
23 dataset and other stations from different archives.

24 Short- and longwave downwelling radiation fluxes at the surface and cloud fraction were
25 evaluated with the International Satellite Cloud Climatology Project (ISCCP) Flux Dataset.
26 The ISCCP radiation fluxes comprise a satellite derived product including shortwave (0.2-5
27 μm) and longwave (5.0-200 μm) radiation at the Earth's surface. ~~Radiation is a satellite
28 derived product including shortwave (0.2-5 μm) and longwave (5.0-200 μm) downwelling
29 and upwelling radiation at the Earth's surface.~~ The radiation estimates come from the
30 synergistic use of ISCCP cloud dataset, satellite data (TOMS, TOVS and SAGE-II), models
31 (NCEP reanalysis, GISS climate model) and climatologies of various tropospheric and

1 stratospheric parameters (aerosols, water vapour, etc). The dataset spans from July 1983 to
2 December 2009 having a temporal resolution of 3hr and a spatial resolution 280 km x 280 km
3 (~2.5x2.5°). Zhang et al. (2004) estimated the uncertainty of the dataset at 10-15W/m²
4 compared with the ERBE (Earth Radiation Budget Experiment) and (Clouds and the Earth's
5 Radiant Energy System) CERES datasets. Since the ISCCP radiation data emerge from the
6 use of a complete radiative transfer model from the GISS global climate model with
7 observations of ISCCP surface, atmosphere and cloud physical properties as input, the
8 radiation and cloud datasets are considered fully compatible. For the current analysis,
9 seasonal averages of the ISCCP variables were calculated for the time period 1990-2008 and
10 were compared to the WRF surface downward short- and longwave radiation, after bilinear
11 interpolation to the 2.5x2.5° ISCCP grid.

12 Model cloudiness was validated against the well-established cloud product from ISCCP,
13 obtained from operational sensors aboard geostationary and polar-orbiting satellites (Rossow
14 and Schiffer, 1999). Single pixel observations in the visible (0.6mm and 1km resolution) and
15 infrared (11mm and 1–4-km resolution depending on the instrument) spectral bands are used.
16 Pixels appearing to be colder and/or brighter than clear sky are characterized as cloudy. Pixel-
17 level retrievals are spatially aggregated at an equal area grid with a resolution of 280km x
18 280km, being available 8 times per day. The ISCCP cloud product is in good agreement to
19 the MODIS cloud mask product (Pincus et al., 2012).

20 An additional, higher resolution, satellite dataset was also used for model validation, in order
21 to confirm the robustness of the validation findings with ISCCP. Shortwave downward
22 radiation at the surface was additionally obtained from Satellite Application Facilities for
23 Climate Monitoring (CMSAF), which is part of the European Organization for the
24 Exploitation of Meteorological Satellites (EUMETSAT). The spatial resolution of the data is
25 0.03°x0.03° while the temporal resolution is 15 min. There ~~is-are~~ a total of six MFG satellites
26 (Meteosat-2 to 7), providing SSR data from 1983 to 2005. This dataset has been validated
27 against homogenized ground-based observations from the Global Energy Balance Archive
28 (GEBA) (Sanchez-Lorenzo et al., 2013) and from the Baseline Surface Radiation Network
29 (BSRN) (Posselt et al., 2012). In this study, seasonal mean solar surface radiation data from
30 CMSAF were re-gridded to the ~~EOBSE-OBS 0.44° resolution domain~~—in order to be
31 compared with the WRF simulations for the time period 1990-2005. Since this dataset does
32 not exactly overlap with the hindcast timeslice (1990-2008), we used the higher resolution

1 dataset only as auxiliary material to support the major findings of the model comparison with
2 the coarser ISCCP satellite retrievals.

3 **2.2 Models**

4 In this work we present EURO-CORDEX hindcast climate simulations performed with the
5 WRF/ARW (version 3.3.1) model. The simulations cover the EURO-CORDEX domain with
6 a resolution of 0.44° . Some **options–settings** are common to all the simulations. The Noah
7 Land Surface Model (NOAH) was the commonly selected land surface model (Chen et al.,
8 1996), the Yonsei University scheme (YSU) was the chosen Planetary Boundary Layer (PBL)
9 scheme (Hong et al., 2006) and MM5 similarity the surface layer option. All simulations were
10 forced by the ERA-~~interim~~-Interim reanalysis dataset (Dee et al., 2011) at 6-hourly intervals
11 with a spatial resolution of 0.75° . The way the forcing fields were pre-processed and
12 implemented in the simulations (relaxation zone, method etc), the setting of vertical layering,
13 land use databases, ~~or~~ and sea surface temperatures were decided by each group separately.

14 In the current ensemble, five different WRF configurations are applied (Table 1). Three
15 different convection schemes were used, namely the Kain-Fritsch (KF, Kain 2004), the Grell-
16 Devenyi (GD, Grell and Devenyi, 2002) and the Betts-Miller-Janjic ensemble (BMJ, Janjic,
17 2000). The radiation physics options tested were: the newer version of the Rapid Radiative
18 Transfer Model (RRTMG, Iacono et al. 2008) and the CAM scheme (Collins et al. 2004). The
19 selected microphysics options were the WRF Single-Moment 3 and 5-class schemes
20 (WSM3/WSM5, Hong et al., 2004) and the WRF Single-Moment 6-class schemes 6 (WSM6,
21 Hong and Lim 2006). **The number of points in relaxation zone and type of relaxation are**
22 **provided in the last column of Table 1.** WRF_A configuration is simulated twice with
23 different SSTs (WRF_A and WRF_A_SST). In WRF_A_SST, the SST field was interpolated
24 as provided in the standard 3.3.1 release (METGRID.TBL). This option results in a coarse
25 resolution of the SSTs resulting in a strong temperature perturbation across the European
26 coastline. In other configurations, either a finer interpolation method is used or the SST fields
27 are replaced by skin temperature.

28 Five meteorological variables are evaluated, namely surface temperature, precipitation, total
29 cloud cover, the short- and longwave downward radiation at the surface. Temperature and
30 precipitation fields were interpolated to the 0.44° E-OBS grid and an elevation correction
31 (standard lapse rate of $6^\circ\text{C}/\text{Km}$) was applied to the simulated temperature to account for the

1 difference between ~~EOBSE-OBS~~9 and model orography. Radiation and cloud data were
2 ~~bilinearly~~-interpolated to a common ISCCP 2.5° grid for comparison to the satellite dataset.

3 In the WRF output the fractional cloud cover is available in each hybrid level. To be able to
4 compute the total cloud cover, an assumption about the overlapping of these fractions is
5 needed. In the present study, we post-processed the fractional cloud cover following the
6 algorithm proposed by Sundqvist (1989). This method assumes maximum overlapping inside
7 cloud layers and random overlapping between them, which is usually summarized as
8 maximum/random overlapping. Radiation parameterizations make their own assumptions to
9 compute cloud effects on radiative fluxes. The overlapping methodology of the Community
10 Atmosphere Model (CAM) radiation parameterization is described in Collins, (2001), and it is
11 also maximum/random overlapping. The RRTMG parameterization also uses
12 maximum/random overlapping. Therefore, except small differences in the algorithms, the
13 overlapping assumptions are consistent through the parameterizations used and in the post-
14 processor.

15

16 **2.3 Methodology**

17 Mean surface temperature, precipitation and solar radiation were calculated for the time
18 period of interest (1990-2008). One year (1989) was used by all simulations as ~~spin-up~~spin-up
19 time. In particular, this spin-up allows for adjustment of the soil moisture and temperature.
20 The seasons were averaged from June to August (JJA) and December to February (DJF). All
21 seasonal averages were calculated based on mean monthly values. The analysis is undertaken
22 over the whole European domain and over the following sub-regions: Alps (AL), British Isles
23 (BI), East Europe (EA), France (FR), Mid-Europe (ME), Mediterranean (MD), Iberian
24 Peninsula (IP) and Scandinavian Peninsula (SC). These sub-domains are described in
25 Christensen and Christensen, 2007.

26 Taylor diagrams are used to provide a concise statistical summary of how well observed and
27 simulated patterns match each other in terms of their correlation R and normalized standard
28 deviation (NSD) (Taylor, 2001). On a Taylor diagram, R and NSD are all indicated by a
29 single point on a two-dimensional polar coordinate plot. The radial distance from the origin
30 corresponds to NSD while the azimuthal position corresponds to R. In the Taylor diagrams
31 the reference point is also displayed, which has R and NSD equal to one. Thus it is easy to

1 identify locations and analysis regions for which the model performs relatively well, as they
2 lie close to the reference point. Furthermore, in case of deviations from the reference, it is
3 easy to distinguish between errors due to poor simulation of variance or due to incorrect
4 phasing (low correlation).

5 Q-q plots compare the probability distribution of two variables, by representing on a Cartesian
6 plane some quantiles of a variable against those of another variable or a theoretical
7 distribution. In this work, following the methodology of Garcia-Diez et al. ~~2013~~2012, we
8 compared the distribution of simulated mean temperature and precipitation (y-axis) against
9 the observations (x-axis) dividing the probability range into 19 pieces (i.e. taking a quantile
10 every 5%). These representations allow one to easily identify deviations in the probability
11 distribution (as departures from a straight diagonal line), biases (as shifts), differences in the
12 variability (as straight lines with a different slope) or asymmetries (as curved lines).

13 In order to test the statistical significance of differences between models and observations ~~of~~
14 we calculate the quantity t (two-independent sample t-test):

$$15 \quad t = (X_m - X_o) / \text{SQRT}((\sigma_m^2 + \sigma_o^2) / n)$$

16 where X_m and X_o are the arithmetic means of the $n = 57$ monthly values for one season in the
17 19-year time slice; σ_m and σ_o are the standard deviations of the n values. The modelled and
18 observed values are significantly different at the 95% level if $t > 1.98$.

19 **3 Results**

20 **3.1 Surface temperature**

21 **3.1.1 Bias**

22 The mean climatological patterns and the annual cycle of temperature are captured quite well
23 by all model configurations, following the spatial characteristics of ~~E~~OBSE-OBS9. This
24 supports the view that major processes governing the surface temperature climatology are
25 represented reasonably by all model configurations. Figure 1 shows the summer and winter
26 mean surface 2m temperature bias with respect to ~~E~~OBSE-OBS9 over Europe averaged over
27 the time slice 1990-2008. Stippling indicates areas where the biases are not statistically
28 significant; over all other regions the models and observations are significantly different at the
29 95% level. Table 2 summarizes the ~~E~~OBSE-OBS9 mean seasonal averages of surface
30 temperature over the different subregions, the absolute model bias (model-~~E~~OBSE-OBS9) of

1 all simulations and the ERA-Interim comparison with **EOBSE-OBS** (ERA-Interim minus
2 **EOBSE-OBS9**). The forcing fields (ERAi) are somewhat warmer ($\sim 0.5^{\circ}\text{C}$) over the whole
3 European domain compared to **EOBSE-OBS9** data. Nearly all WRF configurations
4 underestimate surface temperatures over the different European sub-regions for both seasons.
5 Only the upper quantiles of JJA mean-temperature are overestimated mainly in southern
6 Europe (MD,IP), as indicated by the q-q plots (Fig S1a). Otherwise, the bias remains
7 systematically negative for all configurations, with no obvious asymmetries or differences in
8 variability, except for the behaviour of WRF-G in summer and WRF-A_SST in winter, which
9 are discussed thoroughly in the following sections.

10 **A large negative temperature bias over north-east Europe in winter is also indicated for**
11 **maximum temperatures (-9°C) (Fig. S2) in WRF-A_SST and is also apparent, in all other**
12 **configurations. This feature is more persistent in minimum temperatures (Fig. S3) ranging**
13 **from -2°C (WRF-F) to -13°C (WRF-A_SST). In summer, maximum temperatures are**
14 **reasonably reproduced in most configurations with biases becoming positive over central and**
15 **eastern Europe. Only the WRF-G configuration exhibits the same persistent feature of strong**
16 **negative bias over north Europe. Minimum temperatures in summer are relatively well**
17 **reproduced, with some positive bias mostly seen in WRF-F ($<3^{\circ}\text{C}$).**

18 Mooney et al. 2013 in a WRF-multi physics ensemble forced by ERA-Interim, reported that
19 summer surface temperature is mostly controlled by the selection of Land Surface Model
20 (LSMs). In their study the NOAH and Rapid Update Cycle (RUC) LSMs were tested, and the
21 use of NOAH yielded more accurate surface temperatures than the use of RUC, however the
22 temperature distributions were shifted towards lower values, especially when combined with
23 the CAM radiation scheme. Our current findings can neither support nor contradict this
24 finding, since all models are using the NOAH LSM. We could tentatively attribute, however,
25 the combination of the NOAH LSM along with the CAM radiation scheme, as one possible
26 explanation contributing to the general tendency towards cold biases in the WRF-ensemble.

27 Of all our WRF simulations, WRF-G has the largest cold bias in summer (-2.1°C mean over
28 all European sub-regions). WRF-G uses the GD convective scheme, which may explain the
29 larger cold bias, since the other configuration using the same microphysics (WSM6) and
30 radiation (CAM) as WRF-G, with a different convective scheme (WRF-A with KF scheme)
31 has a smaller bias (-0.3°C). Analysis of the short- and longwave radiation components further
32 support this **findinginterpretation**, as shown below.

1 In winter a negative temperature bias is apparent for all model configurations especially over
2 the north-eastern part of Europe and as indicated by the winter mean temperature q-q- plots
3 (Fig S1b), this underestimation mostly concerns the lower quantiles of the distribution. This
4 finding ~~is~~ is not uncommon among different climate simulations including global modelling
5 studies within CMIP5 (e.g. Cattiaux et al. 2013). Mooney et al. (2013) reported that the
6 radiation scheme (especially the long wave component) has a large impact on winter surface
7 temperature, the CAM option being related to greater negative bias over north and east
8 Europe in comparison to RRTMG. Our simulations do support this finding, since WRF-D
9 and WRF-F using the RRTMG radiation scheme exhibit the smallest bias in winter over the
10 EA domain (-0.2 and 0.6°C respectively). The bias in Scandinavia ranges from -1 to -3°C in
11 the current ensemble during winter.

12 Interestingly, the same subregions (SC, EA) apart from exhibiting the largest **winter** bias, are
13 also the areas with the highest spread in temperature, as indicated by the standard deviation
14 contours (~~not shown~~ Fig. S4). Moreover, the differences between the observed and model
15 distributions over this area are statistically significant for all model configurations.
16 Wintertime standard deviations are considerably larger than summertime and mostly located
17 over north-east Europe (3-4°C) with a northeast-southwest gradient. This spatial pattern of
18 higher uncertainty (spread) over north-east Europe has also been reported in future climate
19 projections for winter temperature, and **is** related to the role of snow cover in cooling down
20 the surface through snow albedo and snow emissivity feedbacks (Deque et al. 2007). Another
21 issue for consideration is that the working WRF version has known problems in treating
22 surface temperature in snow covered areas¹. Garcia-Diez et al. (2014b) show also in their 5-
23 years long multi-physics EURO-CORDEX ensemble that the snow-covered European regions
24 (Alps, and north-east Europe) overestimate the surface albedo, which may be among the
25 sources of bias.

26 WRF-A_SST has an even colder bias for both seasons in comparison to WRF-A, despite
27 using the same primary parameterizations. This disagreement can be attributed to the SST
28 implementation (coarse resolution along the coastline). This perturbation of SSTs affects
29 considerably the inner part of the domain in winter, by lowering the surface temperature, as
30 indicated by additional 1-year long sensitivity studies with the WRF-A_SST modelling

¹ www.atmos.washington.edu/~cliff/WRFWorkshop2013.ppt

1 system (~~results not shown here~~). In the 19-years hindcast simulations, this effect is not so
2 pronounced in summer. The southern part of the Scandinavian Peninsula, the UK and Italy are
3 the areas with the highest temperature differences in winter. This increases the spread in these
4 areas even more, and thus uncertainty in winter temperature, which has already been shown to
5 be large above north-east Europe in winter.

6 The causal link between SSTs and land surface temperature is not easy to depict as they both
7 may influence one another and third factors may influence both at the same time. A similar
8 behaviour is also reported by Cattiaux et al (2011) in a North-Atlantic SST sensitivity
9 experiment of the fall and winter 2006/2007 with a climatic (colder) SST dataset. A similar
10 response in land surface temperature above Europe was showcased, in which anomalous SSTs
11 affected land temperature through the upper-air advection of heat and water vapor, interacting
12 with radiative fluxes over the continent. This mechanism was also found to be more
13 pronounced in autumn and winter, when SSTs anomalies and upper air advection is more
14 efficient.

15 3.1.2 Temporal and spatial agreement

16 We use Taylor plots (Taylor ~~2011~~2001) to investigate the temporal agreement between the
17 simulated and observed fields, *i.e. the reproduction of interannual variations*. With area-
18 averaged temperature fields, we compare time-series of spatially averaged quantities. Figure 2
19 (upper panel) depicts model performance averaged over the different European sub-regions,
20 different colours depict the different WRF configurations. The overall model performance
21 based on average monthly values, indicates very high temporal agreement with observations
22 (0.95) and amplitude of variability higher than the ~~observational—observed~~ ($\sigma_{\text{norm}} > 1$).
23 Inspection of Taylor plots for each different European subregion (~~not shown Fig. S5~~), shows
24 that the largest amplitude of variability *in summer* is produced by WRF-F/WRF-G and the
25 lowest (σ_{norm} slightly below unity) for WRF-C. The worst performance with respect to
26 temporal correlations is found over the Alps for the winter and summer season ($0.7 < R < 0.8$)
27 most probably due to the coarse resolution of the model set up which cannot capture
28 accurately the topographic features of the area.

29 The spatial agreement between observations and the models is investigated by comparing the
30 time-averaged spatial fields *i.e.* two maps without a temporally varying component. The
31 spatial agreement over the whole European domain (Figure 2-bottom) is very high (0.97-

1 0.99), confirming that the spatial representation of surface temperature is captured well. The
2 amplitude of normalized standard deviation (σ_{norm}) in winter is somewhat higher than unity
3 for all configurations. In summer results are more dispersed compared to winter, and the
4 WRF-C configuration again gives the lowest and best (unity) σ_{norm} . On a sub-regional level
5 results appear to have greater spread over ~~over~~ inner continental regions (ME,FR, EA) in
6 comparison to coastal areas (IP,SC,MD, IB).

7

8 **3.2 Precipitation**

9 **3.2.1 Bias**

10 All models depict observed climatological features, namely the major precipitation maxima
11 over the Alps (smaller in winter) and western Norway and the dry regions over the
12 Mediterranean in summer (~~not shown~~Fig S6). Precipitation is overestimated for both seasons
13 over all subregions, except for the British Isles in winter (-5 to -15% relative bias depending
14 on the configuration) (Table 3). The precipitation bias is larger in summer, ranging between
15 25 to 55% for the different model configurations, than in winter (15 to 30%).

16 Figure 3 shows the mean bias in precipitation for all model configurations. The difference
17 between modelled and observed values is statistically significant for all configurations over
18 most subregions. The models show the largest deviation from observations for summer
19 precipitation magnitudes in the Mediterranean area, especially if the KF convective scheme is
20 selected. Convective precipitation along the Dinaric Alps is overestimated in the WRF-C and
21 WRF-A configurations such that the model precipitation is almost double that of the
22 observations. The issue of unrealistically high summer convective precipitation over
23 mountainous regions is also discussed by Torma et al., 2011 and Zanis et al., 2014, indicating
24 that the bias improves in higher resolution simulations by optimizing the convection scheme.
25 Higher precipitation rates (upper quantiles) are overestimated over all subregions for all
26 model configurations (Fig. S2a-S7a). Herwehe et al. 2014 in their study over North America,
27 also reported a large overestimation in larger summertime precipitation amounts (>2.54 cm),
28 attributed to deep cumulus convection. This large overestimation was improved considerably
29 when subgrid-scale cloud-radiation interaction were introduced into the WRF model in ~~to~~ the
30 KF convection scheme (Alapaty et al., 2012).

1 The lowest summer precipitation bias is noted when the GD convective scheme is used (about
2 25-30% on average), followed by the BMJ (about 35%). The KF scheme is related to the
3 highest positive precipitation bias over all European sub-regions but the Scandinavian
4 Peninsula (50-55% in summer and 20-30% in winter). Results are more comparable in winter:
5 the most problematic area with respect to bias appears to be Eastern Europe (50-65% for
6 different model options) while for all other European sub-regions the bias is considerably
7 lower (20-30%). A number of WRF ensemble studies (Evans et al., 2012; Ji et al., 2014; Di
8 Luca et al, 2014) have also reported that the cumulus along with the PBL schemes exhibit the
9 strongest influence on precipitation. Evans et al., 2012 in a WRF ensemble study over
10 southeast Australia, reported that the YSU PBL scheme tends to induce more convection in
11 the KF scheme and lead to an overestimation of precipitation.

12 Precipitation overestimation is not an uncommon feature in WRF simulations (Garcia-Diez et
13 al., 2014b), and often becomes more pronounced at higher resolutions. This systematic error
14 may reflect an unbalanced hydrological cycle, returning moisture from land and/or water
15 bodies to the atmosphere too quickly. Kotlarski et al. (2014) suggest that the wintertime wet
16 bias of WRF is closely related to the distinct negative bias of mean sea-level pressure,
17 indicating a too high intensity of low pressure systems passing over the continent. However,
18 some sensitivity studies performed at WRF-F using spectral nudging for upper air winds and
19 thereby avoiding this problem, showed little changes in bias amplitude (Vautard, personal
20 communication). Sensitivity tests conducted to test alternative choices for convective
21 parameterizations and cloud microphysics are also usually not conclusive but generally none
22 of the options decisively improve the general picture at higher resolutions (Bullock et al.,
23 2014).

24 Figure 4 depicts the annual cycles of all model configurations based on mean monthly values,
25 over the selected subregions. The shaded area corresponds to the observational standard
26 deviation. All configurations reproduce reasonably well the basic characteristics of the
27 seasonal cycle, such as the dry summer of southern Europe or the summer maximum over
28 Scandinavia. All simulations have a wet bias, mostly during spring- and summertime and to a
29 lesser extent in autumn and winter. This fact points to smaller-scale circulations and
30 convection being a critical component to the large positive bias in precipitation. Higher
31 correlations of the ~~modeled~~ modelled with observed annual cycles are seen over the
32 Mediterranean, the Iberian and the Scandinavian Peninsulas, despite the large positive bias.

1 Results are more dispersed and less correlated for the Alps and the Mid-European regions. In
2 a few cases the models have ~~difficulties-difficulty to capture~~ correctly capturing the seasonal
3 cycle over France (WRF-C, WRF-G, WRF-F).

4 The perturbed SSTs in the WRF-A_SST simulation result in a drier climate throughout the
5 year. The physical reason of this colder and drier climate, can be traced ~~on~~ to the water
6 holding capacity of the atmosphere ~~limitings~~ precipitation amounts in colder conditions,
7 assuming a small change in the average relative humidity. Depending on the energetic
8 constraints of a region and its water limitations this relation is modulated accordingly for each
9 season and subregion (Trenberth and Shea, 2005). It should be noted, that the reduced
10 precipitation in WRF-A_SST simulations improves considerably the precipitation bias (Table
11 2) to about 15% on average for both seasons. However, ~~that~~ this is just a case of error
12 compensation, based on the basic WRF feature of predominant overestimation in
13 precipitation.

14 3.2.2 Temporal and spatial agreement

15 Following the same methodology described above for temperature, we proceed with the
16 analysis for precipitation. The temporal Taylor plot ~~are~~ based on mean monthly values, ~~-thus~~
17 ~~indicating interannual variability~~, and ~~are~~ averaged over all European subregions (Fig. 5,
18 upper panel) for precipitation shows that the average JJA temporal correlation is 0.8 for all
19 configurations, with amplitudes of variability being close to unity for WRF-F/WRF-G (GD
20 convection) and somewhat higher for all other configurations. The impact of the selection of
21 convective scheme is clearly seen in the summer season but not in winter. For DJF
22 precipitation, the metrics improve somewhat in comparison to those during the warm period
23 ($0.8 < R < 0.9$ and $\sigma_{\text{norm}} \sim 1$), therefore it seems that WRF captures better the temporal variability
24 in winter than summer, apart from having a lower wet bias. The temporal correlation over the
25 Alps is the lowest in the sub-regional analysis ($0.3 < R < 0.6$) and larger over the Scandinavian
26 Peninsula (0.9 in winter and 0.6-0.8 in summer).

27 With respect to precipitation spatial agreement with observations (Fig 5, bottom), it seems
28 that DJF WRF results are coherent, and that the different model parameterizations do not
29 impact much on the average winter spatial pattern. The average spatial correlation is about 0.7
30 and the amplitude of variability 1.1 to 1.2. In summer results are more dispersed with spatial
31 correlations ranging from 0.8 to 0.9 and higher amplitudes of variability (1.2 - 1.5), indicating

1 that the amplitude of JJA spatial variation is overestimated. This is a common finding among
2 regional climate model studies, reporting summer precipitation to be mostly controlled by
3 internal ~~convection-convective~~ processes, and winter patterns most likely linked to the large-
4 scale circulation and thus the forcing fields (e.g. Rauscher et al. 2010). On a subregional level,
5 the highest spatial correlations ~~is-are~~ seen over the Scandinavian Peninsula and the British
6 Isles ($R=0.9$) in winter and the ~~worst-lowest~~ over France and Mid-Europe in summer ($R=0.4$).
7 The amplitude of variability is exaggerated by all model configurations in summer
8 ($1.5 < \sigma_{\text{norm}} < 2$), with the exception of the British Isles (σ_{norm} close to unity).

9 **3.3 Radiation**

10 The primary driver of latitudinal and seasonal variations in temperature is the seasonally
11 varying pattern of incident sunlight, and a fundamental driver of the circulation of the
12 atmosphere are the local-to-planetary scale imbalances between the shortwave (SW) and
13 longwave (LW) radiation. The impact of the distribution of insolation on temperature can be
14 strongly modified by the distribution of clouds and surface characteristics. In the ~~this current~~
15 section we evaluate two radiation components of the WRF model simulations, namely the
16 surface downwelling SW and LW, which are compared to available ISCCP satellite
17 ~~measurementsretrievals~~. The comparison was also performed with the CMSAF satellite
18 dataset, available in a higher spatial resolution, but only between 1997-2003.

19 **3.3.1 Downward shortwave radiation at the surface**

20 Seasonal average 1990-2008 downward SW radiation components from WRF and ISCCP
21 satellite data are compared over the European domain. Satellite observations exhibit a south-
22 north gradient in summer, with a maximum over the Mediterranean (up to 400 W/m^2) and
23 minima over northern Europe (about 200 W/m^2 on average). All model configurations exhibit
24 this south-north gradient, however with different characteristics: in some configurations
25 (WRF-A/WRF-C with KF or WRF-D with BMJ convection) the SW radiation gradient is less
26 steep towards the north compared to the satellite data, leading to a general positive SW bias
27 over Europe except Scandinavia with a maximum over central Europe, within the range of 40-
28 60% (Fig. 6a). For WRF-F and WRF-G (GD convection) the SW radiation decreases very
29 steeply near $40-45^\circ$, leading to negative bias of SW radiation over north Europe. This can
30 explain, at least partially, the larger ~~summer~~ negative ~~mean~~ temperature bias ~~of-summer~~
31 ~~temperature~~ over mid- and north Europe for WRF-G and WRF-F, compared to other

1 configurations. The SW radiation bias pattern resembles also the bias pattern of maximum
2 surface temperature (Fig. S2a), indicating a strong dependence of maximum temperatures on
3 the SW radiation component. For the WRF-G configuration maximum temperatures are
4 underestimated by up to 8°C over northern Europe, while biases in minimum temperatures are
5 generally smaller (Fig. S3a) and less correlated with SW radiation.

6 Interestingly, Garcia-Diez et al (2014b) showed that the negative SW radiation bias over
7 central and north Europe in summer in the WRF-G configuration is not reproduced in a 5-year
8 long simulation, when the model simulation restarts daily from the ERA-interim forcing fields
9 with 12 hours of spin-up. Thus, it appears this radiation bias is related to internal physical
10 mechanisms, and eventually feedbacks, which develop in a years-long climate simulation. As
11 it will be shown later, the underestimation of SW downward radiation at the surface in GD
12 convection can be linked to a 40-50% overestimation of cloudiness.

13 In winter the observational data indicate maxima of the SW radiation values of about 160
14 W/m² over the southern part of the domain that decreases gradually towards the north. The
15 same spatial pattern is reproduced by all model configurations; however, there is mostly a
16 positive SW radiation bias over the domain, except the Iberian Peninsula and north European
17 coasts of France and Benelux (Fig 6b). The positive bias increases towards the northern and
18 eastern parts of the domain, where it reaches up to 70-80%. WRF-C, with different
19 microphysics (WSM3) has an additional feature, exhibiting of a higher positive SW radiation
20 bias over Mid- and East-Europe (~70%).

21 3.3.2 Downward longwave radiation at the surface

22 Downward LW radiation in summer is higher over southern Europe and decreases towards
23 the north. Comparison with the ISCCP satellite data indicates a negative bias over southern
24 Europe of about 20% -more pronounced for the KF convective scheme- becoming positive in
25 northern Europe with larger positive bias with the GD convective scheme (10%) (Fig 7a).
26 Comparison of Fig 6a and 7a (SW and LW components) shows that summer SW and LW
27 biases are generally anti-correlated, in such a way that regions with positive SW bias, exhibit
28 a negative LW bias and vice versa. If the magnitude of biases were the same, then there would
29 be a cancelling in radiation bias and a better agreement with observed temperature would be
30 expected. However, this is not the case.

1 For WRF-A and WRF-C configurations using the KF convection and CAM radiation schemes
2 there is a strong surplus in downward radiation ($SW_{bias}+LW_{bias} > 0$) over central and
3 southern Europe, leading to lower cold bias or even small warm biases in southern Europe in
4 comparison to northern Europe (Fig S3A8a). The BMJ/RRTMG configuration (WRF-D) has
5 the same features with more enhanced and extended radiative balance surplus extended in east
6 Europe. The GD/CAM (WRF-G) configuration has a predominant summer negative SW
7 radiation bias in north Europe, a smaller in magnitude and positive bias in LW, resulting in a
8 deficit downward radiation regime ($SW_{bias}+LW_{bias} < 0$). Over south Europe the signs change
9 (positive SW bias/ small negative LW bias) resulting in a surplus downward radiation regime
10 ($SW_{bias}+LW_{bias} > 0$). This feature explains the pronounced cold bias in north Europe which
11 becomes lower while moving southwards.

12 The winter LW climatology (~~not shown~~ Fig S9) correlates well spatially with the temperature
13 patterns. It is minimized over north-east Europe and increases towards the south- and western
14 parts of Europe. The winter LW bias is for all model configurations negative over almost all
15 Europe (Fig 7b), only smaller or even positive along the north-west European coast (France,
16 Benelux, Denmark, Baltic countries), compensating for the SW radiation surplus discussed
17 previously. Since the SW amounts over north European winter are very small, the radiation
18 regime is regulated by the LW radiation component, exhibiting a deficit ($SW_{bias}+LW_{bias} < 0$)
19 over north and north-east Europe, which decreases or even becomes positive (WRF-G/WRF-
20 F) in south and south west Europe (Fig S38b).

21 3.3.3 Total cloud cover

22 Since cloudiness is a key component in the discussion concerning radiation, we compare our
23 model results with total cloud cover (CC) of the ISCCP satellite retrievals. During the
24 summer season, observations indicate increased CC over the north and west part of the
25 domain ($CC > 0.8$) i.e. the north-east Atlantic, and the lowest CC in southern Europe ($lat < 40^\circ$).
26 All WRF configurations have a similar pattern, underestimating CC in southern Europe (Fig
27 8a), by more than 50%. The ~~GD~~-configurations with the GD convective scheme have an
28 additional positive bias over northeast Europe. This pattern is very well correlated with the
29 SW radiation bias discussed above, indicating that cloudiness and SW radiation biases have
30 opposite signs, as expected. Herwehe et al. (2014) in a climatic application of WRF over
31 North America reported also an underestimation of summertime cloud fraction over the south-
32 eastern part of their domain, which was considerably improved in their modified case

1 including the sub-grid scale correction in the KF convection scheme. The most pronounced
2 improvement was found in the middle cloud layer (700-500 hPa) consistent with the deep
3 convection of summer. The addition of sub-grid scale cloudiness in the modified case had
4 also the anticipated effect of decreasing ~~in~~ the SW downwelling radiation at the surface and a
5 better agreement with satellite data. The impact on the LW radiation component was minor.

6 In winter the observed CC has a more pronounced peak over the north-west part of the
7 domain over the sea, and reduces gradually towards the south with a secondary maximum
8 over the Black Sea and minima over the Iberian Peninsula (~~not shown~~ Fig S10). The bias
9 pattern in winter (Fig 8b) is negative over the Mediterranean (-20 to -30%) (except in
10 configurations with the GD convective scheme) and positive over north and north-eastern
11 parts of Europe (40 to 50%). The higher than observed ~~modeled~~ modelled cloudiness over
12 northern Europe ~~eventually~~ reduces the amounts of SW radiation reaching the surface, but the
13 positive SW bias remains. Note however, that winter SW radiation absolute amounts are very
14 small over north Europe in winter, so that large relative biases (60-70%) over this area
15 correspond to small absolute changes, which lie within the uncertainty of the satellite data
16 (Zhang et al., 2004).

17 The wintertime positive bias of cloud cover over north Europe is accompanied by negative
18 bias in the LW downward radiation at the surface, in all model configurations. There is not a
19 straightforward explanation for this feature, since increased cloudiness should be associated
20 with increased LW radiation. Both model and observational datasets are internally consistent
21 (the cloud and radiation components), since the ISCCP radiation data are derived by the cloud
22 data (see section 2.1), while WRF has its own internally consistent physics. The results are
23 robust, since they are reproduced by Garcia-Diez et al. (2014b) ~~in a 5-years multi-physics~~
24 ~~ensemble with the same parameterizations, in their 5-year simulations of the same ensemble,~~
25 validated with a different satellite dataset.

26 In order to provide ~~satisfying~~ answers ~~to the questions raised by the modelled cloud and~~
27 ~~radiation biases~~, several issues should be investigated, including a more detailed analysis of
28 cloud coverage and the various radiation components i.e. what are the types of clouds and
29 their impacts on the radiation budget. It is well known that low clouds are thick and non-
30 transparent, reflecting too much of SW radiation back to space (high cloud albedo forcing)
31 and –having almost the same temperature as the surface – do not greatly affect the LW
32 radiation. On the other hand, high thin cirrus clouds are highly transparent to SW radiation

1 but they readily absorb LW radiation. Since they are high and therefore cold, they have a
2 large cloud greenhouse forcing. Finally, the deep convective clouds have a neutral effect since
3 the cloud greenhouse and albedo forcings almost balance. It is clear from the current study,
4 that **in depth** further analysis is necessary, including short- and longwave radiation
5 components, both at the surface and at the top of the atmosphere, as well as various cloud
6 properties which are derived by satellites and are available as output variables in WRF
7 (altitude, optical thickness, cloud albedo).

8

9 **4 Conclusions**

10 Analysis of the WRF ensemble within the EURO-CORDEX framework indicates that the
11 model can represent the present climate with a reasonable degree of fidelity. Temperatures are
12 on average underestimated and the largest temperature spread and bias is seen in winter over
13 north east Europe. Precipitation is overestimated in both seasons but mostly in summer. These
14 general conclusions apply to all ensemble members; the biases ~~range~~-vary depending on the
15 model configuration and the physical parameterizations selected. **The configurations**
16 **appearing to have a more balanced overall behaviour for both precipitation and temperature**
17 **are WRF-D and WRF-F.**

18 Summer temperatures are characterized by a cold bias, more pronounced in northern Europe
19 for the CAM radiation scheme, and a less pronounced, or even slight warm bias for south
20 Europe for the RRTMG radiation scheme. The coldest **mean** temperature bias in north Europe
21 is related to an underestimation of SW radiation at the surface and an overestimation of cloud
22 cover, mostly seen in configurations using the GD convective scheme. **The summer cold bias**
23 **is even more pronounced in maximum temperatures, which are largely controlled by cloud**
24 **coverage and SW radiation.** The strong positive SW bias in summer in southern Europe,
25 mostly induced by the KF or BMJ convective schemes, contributes to a lessening of the
26 systematic cold bias of WRF. When a convective scheme does not suffer from a positive SW
27 bias, then temperatures are grossly underestimated (in our case WRF-G configuration with
28 GD convection).

29 Winter surface temperatures are affected in snow-covered areas in north-east Europe, as a
30 result of a too-strong response of temperature to snow cover. **This underestimation is even**
31 **more pronounced in minimum temperatures, exhibiting bias of up to -9°C over north-east**
32 **Europe in winter, and obviously sensitive to land-atmosphere interactions.** The negative sign

1 in the sum of LW+SW bias over north Europe, contributes to the cold bias problem of the
2 region. Winter cold bias reduces with the application of RRTMG versus the CAM radiation
3 scheme. Mind also, that ERA-Interim has a small (0.4°C) positive bias in comparison to our
4 reference ~~EOBSE-OBS9~~ climatology. If the driving fields suffer from persistent cold bias
5 they can deteriorate ~~even further~~-model performance even further.

6 Precipitation overestimation is reported as a typical WRF behaviour, which remains or even
7 worsens ~~in-at~~ higher spatial resolutions (Kotlarski et al., 2014). Our current findings are in the
8 same line, with the KF convective scheme being related to the highest bias over the
9 Mediterranean in summer. All ensemble members better capture winter than summer
10 precipitation, the latter being locally rather than large-scale controlled. There is no specific
11 configuration that totally alleviates the wet bias of WRF ~~both-either~~ here ~~and-or~~ according to
12 literature. This issue points, among other things, towards weaknesses of ~~in the~~ convective
13 schemes. ~~The precipitation spread of the EURO-CORDEX WRF ensemble analyzed by~~
14 ~~Garcia-Diez (2014a), could only be partially attributed to the selected physical~~
15 ~~parameterizations.~~ Different model domain configurations and datasets seemingly contribute
16 to the precipitation spread. Our study identifies the implementation of SSTs as one important
17 contributing factor. Erroneously, a coarser resolution of implemented SSTs (WRF-A_SST)
18 seemingly “corrects” the average WRF wet bias, by shifting the average climatology towards
19 a colder-drier winter climate regime.

20 Concluding, we stress the importance of such coordinated evaluation exercises, which aim to
21 highlight systematic biases in model performance, and identify the underlying physical
22 mechanisms. The current work ~~evaluates-concentrates~~ only the surface components of the
23 radiation balance ~~and without-leaves other component such as those-at-the~~ top of the
24 atmosphere, the sensible and latent heat fluxes ~~or-looking-into-and~~ cloud properties ~~for future~~
25 ~~analysis.~~ Future ~~analyses-analysis~~ including these parameters is necessary for a more ~~thorough~~
26 ~~complete interpretation-understanding~~ of the physical mechanisms involved in the appearance
27 of temperature and precipitation biases. This work is ongoing within the EURO-CORDEX
28 WRF-groups.

29

30 **Acknowledgements**

1 We acknowledge the E-OBS dataset from the EU-FP6 project ENSEMBLES
2 (<http://ensembles-eu.metoffice.com>) and the data providers in the ECA&D project
3 (<http://www.ecad.eu>).

4 Aristotle University of Thessaloniki simulations have been produced using the EGI and
5 HellasGrid infrastructures and supported by the Scientific Computing Center at the Aristotle
6 University of Thessaloniki. Katragkou E. acknowledges financial support from the AUTH-
7 Special Account for Research Funds.

8 The contribution from the Centre de Recherche Public – Gabriel Lippmann (CRPGL) (now
9 Luxembourg Institute of Science and Technology) was funded by the Luxemburg National
10 Research Fund through the Grant FNR C09/SR/16 (CLIMPACT).

11 Alexandri G. acknowledges financial support from the QUADIEEMS project which is co-
12 financed by the European Social Fund (ESF) and national resources under the operational
13 programme Education and Lifelong Learning (EdLL) within the framework of the Action
14 "Supporting Postdoctoral Researchers".

15 The contribution from Universidad de Cantabria was funded by the Spanish R&D programme
16 through projects CORWES (CGL2010-22158-C02-01) and WRF4G (CGL2011-28864),
17 cofunded by the European Regional Development Fund. García-Díez M. acknowledges
18 financial support from the EXTREMBLES (CGL2010-21869) project.

19

1 **References**

- 2 Alapaty, K., Herwehe, J. A., Otte, T. L., Nolte, C. G., Bullock, O. R., Mallard, M. S., Kain, J.
3 S. and Dudhia, J.: Introducing subgrid-scale cloud feedbacks to radiation for regional
4 meteorological and climate modeling, *Geophys. Res. Lett.*, 39(24), n/a–n/a,
5 doi:10.1029/2012GL054031, 2012.
- 6 Besselaar van den, E. J. M., Haylock, M. R., van der Schrier, G. and Klein Tank, A. M. G.: A
7 European daily high-resolution observational gridded data set of sea level pressure, *J.*
8 *Geophys. Res.*, 116(D11), D11110, doi:10.1029/2010JD015468, 2011.
- 9 Boberg, F. and Christensen, J. H.: Overestimation of Mediterranean summer temperature
10 projections due to model deficiencies, *Nat. Clim. Chang.*, 2(6), 433–436,
11 doi:10.1038/nclimate1454, 2012.
- 12 Bullock, O. R., Alapaty, K., Herwehe, J. A., Mallard, M. S., Otte, T. L., Gilliam, R. C. and
13 Nolte, C. G.: An Observation-Based Investigation of Nudging in WRF for Downscaling
14 Surface Climate Information to 12-km Grid Spacing, *J. Appl. Meteorol. Climatol.*, 53(1), 20–
15 33, doi:10.1175/JAMC-D-13-030.1, 2014.
- 16 Cardoso, R. M., Soares, P. M. M., Miranda, P. M. A. and Belo-Pereira, M.: WRF high
17 resolution simulation of Iberian mean and extreme precipitation climate, *Int. J. Climatol.*,
18 33(11), 2591–2608, doi:10.1002/joc.3616, 2013.
- 19 Cattiaux, J., Vautard, R. and Yiou, P.: North-Atlantic SST amplified recent wintertime
20 European land temperature extremes and trends, *Clim. Dyn.*, 36(11-12), 2113–2128,
21 doi:10.1007/s00382-010-0869-0, 2010.
- 22 **Cattiaux, J., Vautard, R., Yiou, P. North-Atlantic SST amplified recent wintertime European**
23 **land temperature extremes and trends (2011) *Climate Dynamics*, 36 (11-12), pp. 2113-2128.**
- 24 Cattiaux, J., Douville, H., Ribes, A., Chauvin, F. and Plante, C.: Towards a better
25 understanding of changes in wintertime cold extremes over Europe: a pilot study with CNRM
26 and IPSL atmospheric models, *Clim. Dyn.*, 40(9-10), 2433–2445, doi:10.1007/s00382-012-
27 1436-7, 2012.
- 28 **Cattiaux, J., Douville, H., Ribes, A., Chauvin, F., Plante, C. Towards a better understanding**
29 **of changes in wintertime cold extremes over Europe: A pilot study with CNRM and IPSL**
30 **atmospheric models (2013) *Climate Dynamics*, 40 (9-10), pp. 2433-2445.**

1 Chen, F., Mitchell, K., Schaake, J., Xue, Y., Pan, H.-L., Koren, V., Duan, Q. Y., Ek, M. and
2 Betts, A.: Modeling of land surface evaporation by four schemes and comparison with FIFE
3 observations, *J. Geophys. Res.*, 101(D3), 7251, doi:10.1029/95JD02165, 1996.

4 Christensen, J. H. and Christensen, O. B.: A summary of the PRUDENCE model projections
5 of changes in European climate by the end of this century, *Clim. Change*, 81(S1), 7–30,
6 doi:10.1007/s10584-006-9210-7, 2007.

7 Collins, W. D., Rasch, P. J., Boville, B. A., Hack, J. J., McCaa, J. R., Williamson, D. L.,
8 Kiehl, J. T. and Briegleb, B.: Description of the NCAR community atmosphere model (CAM
9 3.0) NCAR Technical Note, NCAR/TN-464+STR. [online] Available from:
10 [http://hanson.geog.udel.edu/~hanson/hanson/CLD_GCM_Experiment_S11_files/description.](http://hanson.geog.udel.edu/~hanson/hanson/CLD_GCM_Experiment_S11_files/description.pdf)
11 pdf (Accessed 14 August 2014), 2004.

12 Collins, William D. 'Parameterization of Generalized Cloud Overlap for Radiative
13 Calculations in General Circulation Models'. *Journal of the Atmospheric Sciences* 58, no. 21
14 (2001): 3224–42.

15 Dee, D. P., Uppala, S. M., Simmons, A. J., Berrisford, P., Poli, P., Kobayashi, S., Andrae, U.,
16 Balmaseda, M. A., Balsamo, G., Bauer, P., Bechtold, P., Beljaars, A. C. M., van de Berg, L.,
17 Bidlot, J., Bormann, N., Delsol, C., Dragani, R., Fuentes, M., Geer, A. J., Haimberger, L.,
18 Healy, S. B., Hersbach, H., Hólm, E. V, Isaksen, L., Kållberg, P., Köhler, M., Matricardi, M.,
19 McNally, A. P., Monge-Sanz, B. M., Morcrette, J.-J., Park, B.-K., Peubey, C., de Rosnay, P.,
20 Tavolato, C., Thépaut, J.-N. and Vitart, F.: The ERA-Interim reanalysis: configuration and
21 performance of the data assimilation system, *Q. J. R. Meteorol. Soc.*, 137(656), 553–597,
22 doi:10.1002/qj.828, 2011.

23 Déqué, M., Rowell, D. P., Lüthi, D., Giorgi, F., Christensen, J. H., Rockel, B., Jacob, D.,
24 Kjellström, E., Castro, M. and Hurk, B.: An intercomparison of regional climate simulations
25 for Europe: assessing uncertainties in model projections, *Clim. Change*, 81(S1), 53–70,
26 doi:10.1007/s10584-006-9228-x, 2007.

27 Flato, G., Marotzke, J., Abiodun, B., Braconnot, P., Chou, S. ., Collins, W., Cox, P.,
28 Driouech, F., Emori, S., Eyring, V., Forest, C., Glecker, P., Guilyardi, E., Jacob, C., Kattsov,
29 V., Reason, C. and Rummukainen, M.: Evaluation of Climate Models. In: *Climate Change*
30 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment

1 Report of the Intergovernmental Panel on Climate Change, Cambridge University Press,
2 Cambridge, United Kingdom and New York, NY, USA., 2013.

3 ~~Garcia-Diez, M., Fernandez, J., L. Fita, C. Yague: Seasonal dependence of WRF model biases
4 and sensitivity to PBL schemes over Europe, Q.J. Meteorol. Soc., doi: 10.1002/gj.1976, 2012~~

5 ~~García-Díez, M., Fernández, J., Vautard, R., Gobiet, A., Goergen, K., Güttler, I., Halenka, T.,
6 Katragkou, E., Kotlarski, S., Krüzselyi, I., Soares, P., Sobolowski, S. and Teichmann, C.:
7 Multiphysics vs. multimodel spread in the new EURO-CORDEX RCM ensemble, Geophys.
8 Res. Lett., 2014a.~~

9 Garcia-Diez, M., Fernandez, J. and Vautard, R.: RCM multi-physics ensemble over Europe:
10 Multi-variable evaluation to avoid error compensation, J. Clim., 2014b.

11 Giorgi, F., Coppola, E., Solmon, F., Mariotti, L., Sylla, M., Bi, X., Elguindi, N., Diro, G.,
12 Nair, V., Giuliani, G., Turuncoglu, U., Cozzini, S., Güttler, I., O'Brien, T., Tawfik, A.,
13 Shalaby, A., Zakey, A., Steiner, A., Stordal, F., Sloan, L. and Brankovic, C.: RegCM4: model
14 description and preliminary tests over multiple CORDEX domains, Clim. Res., 52, 7–29,
15 doi:10.3354/cr01018, 2012.

16 Giorgi, F., Jones, C. and Asrar, G. R.: Addressing climate information needs at the regional
17 level : the CORDEX framework, , 58(July), 175–183, 2009.

18 Grell, G. A. and Dévényi, D.: A generalized approach to parameterizing convection
19 combining ensemble and data assimilation techniques, Geophys. Res. Lett., 29(14), 34–38,
20 doi:10.1029/2002GL015311, 2002.

21 Hall, A. and Qu, X.: Using the current seasonal cycle to constrain snow albedo feedback in
22 future climate change, Geophys. Res. Lett., 33(3), L03502, doi:10.1029/2005GL025127,
23 2006.

24 Haylock, M. R., Hofstra, N., Klein Tank, A. M. G., Klok, E. J., Jones, P. D. and New, M.: A
25 European daily high-resolution gridded data set of surface temperature and precipitation for
26 1950–2006, J. Geophys. Res., 113(D20), D20119, doi:10.1029/2008JD010201, 2008.

27 Herwehe, J.A., Alapaty, K., Spero, T.L., Nolte, C.G.: Increasing the credibility of regional
28 climate simulations by introducing subgrid-scale cloud-radiation interactions, J. Geophys.
29 Res., 119 (9), 2014

1 Hong, S. and Lim, J.: The WRF single-moment 6-class microphysics scheme (WSM6), J.
2 Korean Meteor. Soc [online] Available from:
3 http://www.mmm.ucar.edu/wrf/users/docs/WSM6-hong_and_lim_JKMS.pdf (Accessed 14
4 August 2014), 2006.

5 Hong, S.-Y., Dudhia, J. and Chen, S.-H.: A Revised Approach to Ice Microphysical Processes
6 for the Bulk Parameterization of Clouds and Precipitation, *Mon. Weather Rev.*, 132(1), 103–
7 120, doi:10.1175/1520-0493(2004)132<0103:ARATIM>2.0.CO;2, 2004.

8 Hong, S.-Y., Noh, Y. and Dudhia, J.: A New Vertical Diffusion Package with an Explicit
9 Treatment of Entrainment Processes, *Mon. Weather Rev.*, 134(9), 2318–2341,
10 doi:10.1175/MWR3199.1, 2006.

11 Iacono, M. J., Delamere, J. S., Mlawer, E. J., Shephard, M. W., Clough, S. A. and Collins, W.
12 D.: Radiative forcing by long-lived greenhouse gases: Calculations with the AER radiative
13 transfer models, *J. Geophys. Res.*, 113(D13), D13103, doi:10.1029/2008JD009944, 2008.

14 Jacob, D., Petersen, J., Eggert, B., Alias, A., Christensen, O. B., Bouwer, L. M., Braun, A.,
15 Colette, A., Déqué, M., Georgievski, G., Georgopoulou, E., Gobiet, A., Menut, L., Nikulin,
16 G., Haensler, A., Hempelmann, N., Jones, C., Keuler, K., Kovats, S., Kröner, N., Kotlarski,
17 S., Kriegsman, A., Martin, E., Meijgaard, E., Moseley, C., Pfeifer, S., Preuschmann, S.,
18 Radermacher, C., Radtke, K., Rechid, D., Rounsevell, M., Samuelsson, P., Somot, S.,
19 Soussana, J.-F., Teichmann, C., Valentini, R., Vautard, R., Weber, B. and Yiou, P.: EURO-
20 CORDEX: new high-resolution climate change projections for European impact research,
21 *Reg. Environ. Chang.*, 14(2), 563–578, doi:10.1007/s10113-013-0499-2, 2013.

22 Janjic, Z.: Comments on “Development and evaluation of a convection scheme for use in
23 climate models,” *J. Atmos. Sci.*, 20746 [online] Available from:
24 [http://journals.ametsoc.org/doi/pdf/10.1175/1520-](http://journals.ametsoc.org/doi/pdf/10.1175/1520-0469(2000)057%3C3686%3ACODAE0%3E2.0.CO%3B2)
25 [0469\(2000\)057%3C3686%3ACODAE0%3E2.0.CO%3B2](http://journals.ametsoc.org/doi/pdf/10.1175/1520-0469(2000)057%3C3686%3ACODAE0%3E2.0.CO%3B2) (Accessed 14 August 2014),
26 2000.

27 Kain, J. S.: The Kain–Fritsch Convective Parameterization: An Update, *J. Appl. Meteorol.*,
28 43(1), 170–181, doi:10.1175/1520-0450(2004)043<0170:TKCPAU>2.0.CO;2, 2004.

29 Kotlarski, S., Keuler, K., Christensen, O. B., Colette, A., Déqué, M., Gobiet, A., Goergen, K.,
30 Jacob, D., Lüthi, D., van Meijgaard, E., Nikulin, G., Schär, C., Teichmann, C., Vautard, R.,
31 Warrach-Sagi, K. and Wulfmeyer, V.: Regional climate modeling on European scales: a joint

1 standard evaluation of the EURO-CORDEX RCM ensemble, *Geosci. Model Dev. Discuss.*,
2 7(1), 217–293, doi:10.5194/gmdd-7-217-2014, 2014.

3 Van der Linder, P.: ENSEMBLES: Climate Change and Its Impacts: Summary of Research
4 and Results from the ENSEMBLES Project, Exeter., 2009.

5 **Mooney, P.A., Mulligan., F.J., Fealy, R., Evaluation of the sensitivity of the weather research
6 and forecasting model to parameterization schemes for regional climates of europe over the
7 period 1990-95, *Journal of Climate*, 26 (3), pp. 1002-1017., 2013**

8 Pincus, R., Platnick, S., Ackerman, S. A., Hemler, R. S. and Patrick Hofmann, R. J.:
9 Reconciling Simulated and Observed Views of Clouds: MODIS, ISCCP, and the Limits of
10 Instrument Simulators, *J. Clim.*, 25(13), 4699–4720, doi:10.1175/JCLI-D-11-00267.1, 2012.

11 Posselt, R., Mueller, R. W., Stöckli, R. and Trentmann, J.: Remote sensing of solar surface
12 radiation for climate monitoring — the CM-SAF retrieval in international comparison,
13 *Remote Sens. Environ.*, 118, 186–198, doi:10.1016/j.rse.2011.11.016, 2012.

14 Rauscher, S. A., Coppola, E., Piani, C. and Giorgi, F.: Resolution effects on regional climate
15 model simulations of seasonal precipitation over Europe, *Clim. Dyn.*, 35(4), 685–711,
16 doi:10.1007/s00382-009-0607-7, 2009.

17 Rossow, W. B. and Schiffer, R. A.: Advances in Understanding Clouds from ISCCP, *Bull.*
18 *Am. Meteorol. Soc.*, 80(11), 2261–2287, doi:10.1175/1520-
19 0477(1999)080<2261:AIUCFI>2.0.CO;2, 1999.

20 Rummukainen, M.: State-of-the-art with regional climate models, *Wiley Interdiscip. Rev.*
21 *Clim. Chang.*, 1(1), 82–96, doi:10.1002/wcc.8, 2010.

22 Sanchez-Lorenzo, A., Wild, M. and Trentmann, J.: Validation and stability assessment of the
23 monthly mean CM SAF surface solar radiation dataset over Europe against a homogenized
24 surface dataset (1983–2005), *Remote Sens. Environ.*, 134, 355–366,
25 doi:10.1016/j.rse.2013.03.012, 2013.

26 Skamarock, W. C., Klemp, J. B., Dudhia, J., Gill, D. O., Barker, D. M., Duda, M. G., Huang,
27 X.-Y., Wang, W. and Powers, J. G.: A description of the advanced research WRF version 3.
28 [online] Available from:
29 <http://oai.dtic.mil/oai/oai?verb=getRecord&metadataPrefix=html&identifier=ADA487419>
30 (Accessed 14 August 2014), 2008.

1 Skliris, N., Sofianos, S., Gkanasos, A., Mantziafou, A., Vervatis, V., Axaopoulos, P. and
2 Lascaratos, A.: Decadal scale variability of sea surface temperature in the Mediterranean Sea
3 in relation to atmospheric variability, *Ocean Dyn.*, 62(1), 13–30, doi:10.1007/s10236-011-
4 0493-5, 2011.

5 Soares, P. M. M., Cardoso, R. M., Miranda, P. M. A., Medeiros, J., Belo-Pereira, M. and
6 Espirito-Santo, F.: WRF high resolution dynamical downscaling of ERA-Interim for Portugal,
7 *Clim. Dyn.*, 39(9-10), 2497–2522, doi:10.1007/s00382-012-1315-2, 2012.

8 Stegehuis, A. I., Teuling, A. J., Ciais, P., Vautard, R. and Jung, M.: Future European
9 temperature change uncertainties reduced by using land heat flux observations, *Geophys. Res.*
10 *Let.*, 40(10), 2242–2245, doi:10.1002/grl.50404, 2013.

11 Sundqvist, Hilding, Erik Berge, and Jón Egill Kristjánsson. ‘Condensation and Cloud
12 Parameterization Studies with a Mesoscale Numerical Weather Prediction Model’. *Monthly*
13 *Weather Review* 117, no. 8 (1989): 1641–57.

14 Taylor, K. E.: Summarizing multiple aspects of model performance in a single diagram, *J.*
15 *Geophys. Res.*, 106(D7), 7183, doi:10.1029/2000JD900719, 2001.

16 Taylor, K.E., Stouffer, R.J., Meehl, G.A. An overview of CMIP5 and the experiment design
17 (2012) *Bulletin of the American Meteorological Society*, 93 (4), pp. 485-498.

18 Torma C, Coppola E, Giorgi F, Bartholy J, Pongrácz R.: Validation of a high-resolution
19 version of the regional climate model RegCM3 over the Carpathian basin, *J Hydrometeorol*
20 12:84–100. doi:10.1175/2010JHM1234.1, 2011.

21 Trenberth, K.E., Shea, D.J. Relationships between precipitation and surface temperature
22 (2005) *Geophysical Research Letters*, 32 (14), pp. 1-4.

23 Vautard, R., Gobiet, A., Jacob, D., Belda, M., Colette, A., Déqué, M., Fernández, J., García-
24 Díez, M., Goergen, K., Güttler, I., Halenka, T., Karacostas, T., Katragkou, E., Keuler, K.,
25 Kotlarski, S., Mayer, S., Meijgaard, E., Nikulin, G., Patarčić, M., Scinocca, J., Sobolowski,
26 S., Suklitsch, M., Teichmann, C., Warrach-Sagi, K., Wulfmeyer, V. and Yiou, P.: The
27 simulation of European heat waves from an ensemble of regional climate models within the
28 EURO-CORDEX project, *Clim. Dyn.*, 41(9-10), 2555–2575, doi:10.1007/s00382-013-1714-
29 z, 2013.

1 Warrach-Sagi, K., Schwitalla, T., Wulfmeyer, V. and Bauer, H.-S.: Evaluation of a climate
2 simulation in Europe based on the WRF–NOAH model system: precipitation in Germany,
3 *Clim. Dyn.*, 41(3-4), 755–774, doi:10.1007/s00382-013-1727-7, 2013.

4 Zanis P., Katragkou E. , Ntogras C. , Marougianni G., Tsikerdekis A., Feidas H. ,
5 Anadranistakis E., Melas D., A transient high resolution regional climate simulation for
6 Greece for the period 1960-2100: Evaluation and future projections, submitted to *Regional*
7 *Environmental Change*

8 Zhang, Y., Rossow, W. B., Lacis, A. A., Oinas, V. and Mishchenko, M. I.: Calculation of
9 radiative fluxes from the surface to top of atmosphere based on ISCCP and other global data
10 sets: Refinements of the radiative transfer model and the input data, *J. Geophys. Res.*,
11 109(D19), D19105, doi:10.1029/2003JD004457, 2004.

12

1 Table 1. WRF configurations participating in the study.

Label	Institute	Nz / TOA	Microphys.	Cum.	Rad.	Rel Zone
WRF-A	CRPGL	50 / 20hPa	WSM6	KF	CAM3	10/exp
WRF-A_SST	AUTH	30/ 50hPa	WSM6	KF	CAM3	5/linear
WRF-C	BCCR	30m / 50 hPa	WSM3	KF	CAM3	10 /exp
WRF-D	IDL	40/ 50 hPa	WSM6	BMJ	RRTMG	5/exp
WRF-F	IPSL	32 / 50hPa	WSM5	GD	RRTMG	5/linear
WRF-G	UCAN	30m / 50hPa	WSM6	GD	CAM3	10/linear

- 2 WSM3: Single moment 3 class microphysics scheme
3 WSM5: Double moment 5 class microphysics scheme
4 WSM6: Double moment 6 class microphysics scheme
5 KF: Kain-Fritsch cumulus parameterization
6 BMJ: Betts Miller Janjic cumulus parameterization
7 GD: Grell Devenyi cumulus parameterization
8 CAM3: radiation scheme from the CAM 3 climate model
9 RRTMG: new Rapid Radiative Transfer Model
10 exp: exponential
11 AUTH: Aristotle University of Thessaloniki
12 BCCR: Bjerknes Centre for Climate Research
13 CRPGL: Centre de Recherche Public – Gabriel Lippman
14 IDL: Instituto Dom Luiz
15 IPSL: Institut Pierre Simon Laplace
16 UCAN: Universidad de Cantabria

19 Table 2a. Means (Mobs) of summer (JJA) surface temperature for observations (~~EOBSE-~~
20 **OBS9**) over 1990-2008 and the European subregions and model mean seasonal bias (Mmod-
21 Mobs). Unit is degree Celsius.

	EOBSE- OBS9	WRF-A _SST	WRF-C	CRGPL	WRF-D	WRF-F	WRF-G	ERAi
AL	17.1	-1.0	-1.4	-0.4	-0.2	-0.9	-2.1	0.7
BI	14.7	-2.3	-1.2	-0.9	-0.6	-1.2	-2.4	0.3
EA	18.8	-0.1	-0.1	0.3	0.5	-0.1	-2.3	0.4
FR	18.8	-2.1	-1.6	-0.9	-0.3	-1.2	-2.9	0.2
IP	21.8	-0.5	-1.5	0.0	0.9	0.3	-1.0	0.3
MD	21.9	-0.4	-1.1	0.0	0.7	0.5	-1.0	0.9
ME	17.5	-1.6	-0.7	-0.3	-0.2	-1.1	-2.8	0.3
SC	13.6	-2.3	-0.7	-0.5	-0.4	-0.6	-2.6	0.6

1 Table 2b. Same as Table 2a for winter

	EOBSE- OBS9	WRF-A _SST	WRF-C	CRGPL	WRF-D	WRF-F	WRF-G	ERAi
AL	0.5	-3.6	-1.1	-0.3	-0.4	0.3	-0.7	0.0
BI	4.6	-3.2	-0.1	-0.1	0.2	0.7	0.1	0.7
EA	-1.1	-5.2	-2.0	-1.3	-0.2	0.6	-1.9	0.2
FR	5.1	-3.1	-0.5	-0.4	0.0	0.7	-0.6	0.1
IP	7.0	-2.0	-0.9	-0.4	-0.1	0.4	-0.7	0.3
MD	5.0	-5.5	-1.1	-1.0	-0.5	-0.1	-0.9	0.6
ME	1.8	-3.8	-0.9	-0.5	0.2	0.7	-1.0	0.2
SC	-5.3	-7.0	-2.8	-1.8	-1.8	-0.9	-2.2	0.4

2

3 Table 3a. Mean (Mobs) of summer (JJA) precipitation for observations (**EOBSE-OBS9**) over
 4 1990-2008 and the European subregions, units in mm/day. Units of **EOBSE-OBS9** in
 5 mm/day. Model relative bias (%)

	EOBSE- OBS9	WRF-A _SST	WRF-C	CRGPL	WRF-D	WRF-F	WRF-G
AL	3.20	8%	37%	28%	24%	14%	21%
BI	2.45	15%	34%	23%	27%	-1%	6%
EA	2.22	23%	41%	49%	39%	36%	33%
FR	1.75	15%	83%	47%	16%	37%	35%
IP	0.67	-6%	63%	63%	25%	31%	15%
MD	0.83	-1%	102%	94%	64%	40%	59%
ME	2.35	27%	46%	42%	34%	34%	23%
SC	2.46	26%	33%	39%	54%	22%	7%


6

7 Table 3b. Same as Table 3a for winter.


	EOBSE- OBS9	WRF-A _SST	WRF-C	CRGPL	WRF-D	WRF-F	WRF-G
AL	2.53	16%	14%	26%	41%	17%	7%

BI	3.63	-11%	-4%	-4%	-13%	-11%	-5%
EA	1.13	44%	51%	65%	59%	60%	65%
FR	2.15	45%	33%	38%	20%	18%	24%
IP	1.94	7%	10%	11%	-4%	-15%	-9%
MD	1.98	-15%	33%	32%	14%	1%	10%
ME	1.92	42%	23%	38%	28%	30%	31%
SC	2.01	4%	14%	24%	27%	22%	21%


1




1
 2 Figure 1a Mean summer 1990-2008 surface temperature bias (model-~~EOBSE~~-OBS9).
 3 Stippling indicates areas where the biases are not statistically significant.
 4



1
 2 Figure 1b Mean winter 1990-2008 surface temperature bias (model-EOBSE-OBS9). Stippling
 3 indicates areas where the biases are not statistically significant. Mind the differences in colour
 4 scales.
 5




1




2

3 Figure 2 Temporal (upper panel) and spatial (bottom panel) Taylor plots for surface
 4 temperature averaged over Europe for summer and winter 1990-2008. **Updated plot**


5




1
 2 Figure 3a Mean summer 1990-2008 precipitation bias (model-~~EOBSE-OBS9~~) expressed in
 3 mm/day. Stippling indicates areas where the biases are not statistically significant.
 4




1
 2 Figure 3b Mean winter 1990-2008 precipitation bias (model-~~EOBSE-OBS~~9) expressed in
 3 mm/day. Stippling indicates areas where the biases are not statistically significant.
 4



1
 2 Figure 4 Mean precipitation annual cycle. The grey area indicates observational standard
 3 deviation. Updated plot.




1



2

3 Figure 5 Temporal (upper panel) and spatial (bottom panel) Taylor plots for precipitation
 4 averaged over Europe for summer and winter 1990-2008. Updated plot


5



1

2 Figure 6a Mean summer 1990-2008 downward surface shortwave radiation bias (WRF-
 3 ISCCP)


4



1

2 Figure 6b Mean winter 1990-2008 downward surface shortwave radiation bias (WRF-ISCCP)


3



1

2 Figure 7a Mean summer 1990-2008 downward surface longwave radiation bias (WRF-
 3 ISCCP)


4



1

2 Figure 7b Mean winter 1990-2008 downward surface ~~shortwave-longwave~~ radiation bias
 3 (WRF-ISCCP)


4



1

2 Figure 8a Mean summer 1990-2008 total cloud cover bias (WRF-ISCCP)

3



1

2 Figure 8b Mean winter 1990-2008 total cloud cover bias (WRF-ISCCP)