

1 Response to Reviewer 1

2 **General comments**

3 • Land carbon uptake in the model is represented by net primary production and not by  
4 gross primary production. This may have some conceptual and practical problems  
5 because, i) the autotrophic flux of carbon is not included in the calculations of the land-  
6 atmosphere C exchange, and ii) this land-atmosphere exchange can't be compared against  
7 many available data products. For example, soil respiration fluxes, which include both  
8 autotrophic and heterotrophic sources can't be compared with model predictions.

9 Similarly, ecosystem level fluxes can't be compared with eddy-covariance derived fluxes  
10 or GPP estimates from satellite products. Can you explain why the autotrophic  
11 component of the land C cycle is not included in the model? Do you plan to include this  
12 in the future, or is there a particular reason why you believe this should not be included?  
13

14 *As the reviewer notes, we have chosen, in this 1.0 version, to implement the terrestrial-  
15 atmosphere C exchange as the difference between NPP and RH, rather than breaking out  
16 GPP and RA separately. This makes for a simpler model but, again as the reviewer  
17 correctly notes, limits our ability to compare to, for example, remotely-sensed GPP. This  
18 is a choice that could (and probably will) be changed in the future; we've logged it as an  
19 "issue" on the project repository at <https://github.com/JGCRI/hector/issues/53>. We have  
20 added the following to the manuscript under section 7.0: "For example, Hector does not  
21 currently simulate terrestrial gross primary production, a key metric of comparison to  
22 e.g. the FLUXNET database."*

23  
24 • The documentation of the model in GitHub is incomplete and needs to be finished. In  
25 particular, the authors should describe better the steps for compiling and running the  
26 model in different OS. Given that this documentation is written in Markdown language,  
27 the authors should provide a step-by-step procedure for compiling and running a  
28 simulation using syntax highlighting. A demo on how to analyze the results using the R  
29 scripts would be also very useful.  
30

31 *All documentation of the model is now on Github wiki, including how to compile and run  
32 Hector in each OS, how to guides such as; add new components, unitvals, tseries and a  
33 demo of the R backend. All documentation is found at:  
34 <https://github.com/JGCRI/hector/wiki>*

35  
36 • Figure 4 shows a very high sensitivity of Hector for predicting temperature anomalies.  
37 The slope after the 1960s is much larger in Hector than in the other models. Can you  
38 comment on this large sensitivity?  
39

40 *We have addressed and fixed the high temperature sensitivity after 1960 by including a  
41 variable ocean heat flux, as well as lagging the temperature effects from atmospheric  
42 [CO<sub>2</sub>]. There are numerous processes that are not simulated in Hector that buffer the  
43 temperature effects of increasing GHGs. Therefore, we take a simple approach in this  
44 current version and lag our temperature. We have addressed this in the manuscript  
45 section 4.1, "As global temperatures rise, the uptake capacity of the ocean thus  
46 diminishes, simulating both a saturation of heat in the surface and a slowdown in ocean*

47 *circulation with increased temperatures. Finally, the temperature effects from*  
48 *atmospheric [CO<sub>2</sub>] are lagged in time, as there are numerous real-world processes not*  
49 *simulated in Hector buffering the temperature effects of increasing atmospheric [CO<sub>2</sub>].”*  
50 *See figures 4 and 8 for updated global temperature change.*

51

## 52 **Technical and other comments**

53 • Page 7076, lines 22-23. I would say that fully coupled Earth system models  
54 (atmosphere-ocean-land) are at the complexity end, and not just AOGCMs.

55

56 *The authors agree with the reviewer and have edited the title as suggested:*

57 *“To accomplish this, a hierarchy of climate models with differing levels of complexity*  
58 *and resolution are used, ranging from purely statistical or empirical models, to simple*  
59 *energy balance models, to fully-coupled Earth System Models (ESMs) (Stocker, 2011).”*

60

61 •Page 7080, line 28. Change \_ for d. The \_ notation is commonly used for isotopes in C  
62 cycle models.

63 *Equation edited as suggested. “dC/dt < ε”*

64

65 • How do you calculate NPP0 and RH0? I think this formulation of RH is potentially  
66 dangerous because you may respire more C than what is available in the pools of  
67 equation (8) and (9).

68

69 *In regards to NPP0 and RH: NPP0, the global preindustrial NPP flux, is specified a priori,*  
70 *not calculated. The use of RH0 in Equation 5 was a mistake, for which we apologize. At*  
71 *any point in time, model RH is always a function of the current carbon stocks in soil and*  
72 *litter.*

73

74 • Equation (12). What is the last term Fi? It seems to me that this term violates mass  
75 balance. What additional flux, different from all inputs and outputs, can modify the net  
76 change?

77

78 *The last term Fi is now the carbon flux to/from the atmosphere to/from the ocean.*

79 *Equation 12 has been changed accordingly:*

$$\frac{dC_i}{dt} = \sum_{j=1}^{in} F_{j \rightarrow i} - \sum_{j=1}^{out} F_{i \rightarrow j} + F_{atm \rightarrow i}$$

80

81

82 • Page 7058, line 25. Replace ‘model’ for ‘version’.

83

84 *Model has been changed to version as suggested by the reviewer.*

85

86 • Equation (15). Why do you use a difference equation instead of a differential equation?  
87 Is this process discrete in time?

88

89 *The reviewer brings up a good point. We updated equation 16 to reflect changes from a*  
90 *difference solution to an exact solution. Operating on a finite timescale introduces more*  
91 *error than an exact solution.  $C(t) = C_0 * \exp\left(-\frac{1}{T}\right) + E * T * \left(1 - \exp\left(-\frac{1}{T}\right)\right)$*

92 Response to Reviewer 2:

### 93 **General comments**

94  
95 I have two major concerns with the manuscript: 1) experiments to validate Hector are not  
96 well described, and 2) Hector appear to have issues at longer timescales that are not well  
97 described or acknowledged. I recommend the authors include additional material on the  
98 performed experiments and fidelity of Hector at different timescales. The manuscript also  
99 need significant cleanup of typos and grammatical errors, and could benefit from  
100 improvement of figures.

101  
102 *The authors have since restructured the results section to better describe the*  
103 *experimental design. All experiments are run under prescribed emissions scenarios from*  
104 *the Representative Concentration Pathways (RCP 2.6, 4.5, 6.0, and 8.5). However, the*  
105 *CMIP5 data used to compare with Hector are from prescribed concentration scenarios,*  
106 *with the exception of atmospheric [CO<sub>2</sub>]. We acknowledge that this may not be a perfect*  
107 *comparison, but the CMIP5 archive is limited in the number of models that ran scenarios*  
108 *with prescribed emissions.*

109  
110 *As noted in the title, Hector is concerned with policy relevant timescales, notably the next*  
111 *100-300 years. We agree with the reviewer that a more detailed explanation of the*  
112 *timescales is needed in the manuscript and have since updated this.*

113 *“Hector’s strengths lie within policy relevant time scales of decades to centuries. Studies*  
114 *suggest that 80% of the anthropogenic CO<sub>2</sub> emissions have an average atmospheric*  
115 *lifetime of 300-450 years (Archer et al., 1997; Rogner, 1997; Archer, 2005). Hector has*  
116 *all the necessary components to model the climate system from present day through the*  
117 *next approximately 300 years.”*

118  
119 *Lastly, grammatical errors, and figures have been improved in the updated manuscript.*

120

### 121 **Specific comments**

122

123 **Title:** The title of the paper does not appropriately describe the contents of this  
124 manuscript. The title suggests that this manuscript describes a global carbon cycle model,  
125 but Hector is a full climate model and the paper describes all the components of Hector.  
126 A better title might be something like, A simple object-oriented and open source model  
127 for scientific and policy analyses of the global climate system – Hector v0.1. I  
128 recommend that the authors revise the title to better reflect the overall contents of the  
129 paper.

130

131 *The authors agree with the reviewer and have edited the title as suggested:*

132 *“A simple object-oriented and open source model for scientific and policy analyses of the*  
133 *global climate system–Hector v1.0”*

134

135 **Introduction:** The introduction is lacking a description of previous work in the field and  
136 needs to add citations and discuss the novelty of Hector. The authors properly describe  
137 the purpose of simple climate models, their general structure and implementation. But,  
138 the authors should cite previous simple climate models and explicitly explain why the  
139 design of Hector is novel. Relevant citations include but are not limited to Meinshausen  
140 et al. (2011), Joos et al. (2013), Glotter et al. (2014), and models described in van Vuuren  
141 et al. (2009) and Hof et al. (2011).

142

143 *The authors have added significant changes to the introduction to better reflect the*  
144 *current state of simple climate modeling within Integrated Assessment Models.*

145

146 *“Depending on the purpose of the IAMs (economics, cost-benefit analysis, or more*  
147 *physical based processes), the corresponding climate and carbon component varies in*  
148 *complexity and resolution. For example, models like DICE, FUND, and MERGE have a*  
149 *highly simplified carbon/climate system (Nordhaus, 2008; Anthoff and Tol, 2014; Manne*  
150 *and Richels, 2005). IAMs focusing more on the physical processes of the natural system*  
151 *and the economy employ more complex representations of the climate/carbon system.*  
152 *Models like GCAM (Global Change Assessment Model) and MESSAGE use MAGICC as*  
153 *their SCM (Meinshausen et al., 2011a; Riahi et al., 2007; Calvin et al., 2011).*  
154 *Increasing in complexity, some IAMs include the climate/carbon system at gridded scales*  
155 *(e.g., IMAGE), and can be coupled to earth system models of intermediate complexity*  
156 *(e.g., MIT IGSM), or more recently coupled to a full earth system model (the iESM*  
157 *project) (Bouwman et al., 2006; Sokolov et al., 2005; Bond-Lamberty et al., 2014; Di*  
158 *Vittorio et al., 2014; Collins et al., 2015).”*

159

160 **Results:** The experimental design for the tests performed in this manuscript are not well  
161 described. There remain several ambiguities in Section 5 that must be clarified so results  
162 can be properly assessed. In general, figure captions should be expanded to explain the  
163 experimental design used to make that display. The authors may find it beneficial to add a  
164 table that describes all experiments performed, including Hector’s configuration for each  
165 experiment, input data used to drive Hector, and the model output (or data) that Hector is  
166 compared to. Specific examples of ambiguities related to experimental design include:

167

168 • For most figures, it remains unclear precisely when Hector is driven by an emissions  
169 scenario and when atmospheric carbon is prescribed. For example, is Figure 8 made  
170 using fixed exogenous CO<sub>2</sub> concentrations or with emissions scenarios that reproduce  
171 RCPs? The authors should clarify when RCPs are used and when esmRCPs are used. An  
172 experimental design table (as described above) would help clarify here.

173

174 *The authors have clarified this information in the text as well as in the figure caption.*  
175 *“All CMIP5 variables used in this study are from model runs with prescribed*  
176 *atmospheric concentrations, except for comparisons involving atmospheric [CO<sub>2</sub>] which*  
177 *are from the emissions driven scenario (esmHistorical and esmRCP8.5) (Figures 3 and*  
178 *5). We acknowledge that this comparison, between an emissions-forced model (Hector)*  
179 *and concentration-forced models (CMIP5), is not perfect. However, few CMIP5 models*  
180 *were run under prescribed emissions scenarios.”*

181

182 • The paragraph on page 7081 (lines 4-16) that describes how atmospheric concentrations  
183 are prescribed needs to be re-written. If the model simply inverts concentrations to find  
184 emissions, it is not clear why the assumption in lines 14-15 is necessary. I am also not  
185 sure this statement would hold true for large perturbation scenarios, such as an  
186 instantaneous doubling (or more) of CO<sub>2</sub>. If this is how the authors perform the  
187 prescribed-CO<sub>2</sub> experiments, it is vital that it be described carefully else results are not  
188 interpretable.

189

190 *The paragraph on page 7081 explains some of the capabilities built into Hector to force*  
191 *its output to match a user-supplied time series. This is very helpful with testing and*  
192 *debugging the carbon cycle system within Hector. We do not invert concentrations to*  
193 *find emissions; instead for example, atmospheric CO<sub>2</sub> concentrations are read into*  
194 *Hector and over write any calculated [CO<sub>2</sub>] values. The user-supplied time series can be*  
195 *started and stopped at any point. When the model exits the constrained time period,*  
196 *[CO<sub>2</sub>], in this case, becomes fully prognostic. We have updated the manuscript to better*  
197 *reflect this.*

198

199 *“Hector can be forced to match its output to a user-supplied time series. This is helpful*  
200 *to isolate and test different components. Available constraints currently include*  
201 *atmospheric CO<sub>2</sub>, global temperature anomaly, total ocean-atmosphere carbon*  
202 *exchange, total land-atmosphere carbon exchange, and total radiative forcing.”*

203

204 • Which “historical conditions” are used to run Hector (page 7089, lines 17-18)?

205

206 *The text containing ‘historical conditions’ has been rewritten to better reflect the inputs*  
207 *use to run Hector in this study.*

208

209 *“Within this study, Hector is run with prescribed emissions from 1850 to 2300 under all*  
210 *four Representative Concentration Pathways (RCPs), freely available at*  
211 *<http://tntcat.iiasa.ac.at/RcpDb/>”*

212

213 • Which models run esmRCP8.5 (page 7090, lines 11-12)? Are these different than the 11  
214 CMIP5 models?

215

216 *The authors have updated table 3 within the manuscript to reflect the models that ran*  
217 *esmHistorical and esmrcp85 (emissions prescribed scenarios).*

218

219 • RCPs (by definition) are CO<sub>2</sub> concentration pathways. What does it mean for  
220 atmospheric CO<sub>2</sub> in Hector to be highly correlated with MAGICC for the four RCPs  
221 (Page 7091, lines 23-27)? Shouldn't the definition of an RCP necessitate identical  
222 concentration pathways? This confusion also applies to figures 5-6, and is likely related  
223 to the confusion described in the second bullet. Please clarify.

224

225 *The authors apologize for the confusion on the wording of RCPs and how they relate to*  
226 *the Hector output. We have clarified this issue in section 5.0 below:*

227 “RCPs by definition are concentration pathways; however, for all experiments within this  
228 manuscript we use the corresponding emissions trajectories from each RCP as input for  
229 Hector.”

230

231 • I disagree with the statements at the beginning of section 6.2. It is incorrect that models  
232 that  
233 accurately estimate historical climate are simply “assumed” to be reliable for future  
234 scenarios. The credibility of Hector in making future projections of the climate should not  
235 be based solely on the fact that it can reproduce historical trends. In fact, we see that  
236 Hector has problems at long timescales (where short timescales are more accurate–  
237 Figures 8 and 10), and even some errors appear in the historical record itself (Figure 4).  
238 The authors must re-write this paragraph, but more importantly, must be explicit about  
239 issues with the use of Hector over long timescales. There are issues with the fidelity of  
240 Hector at different timescales that are not acknowledged or described. Hector does not  
241 include the dissolution of calcium carbonate in its representation of the carbon cycle (to  
242 my knowledge) and therefore will not be dependable past ~2000 years. But I do not know  
243 whether Hector is dependable up to 2000 years. Potential users of Hector would benefit  
244 greatly from a dedicated discussion of its usefulness at different timescales. Specific  
245 concerns with the fidelity of Hector include:

246

247 *The authors fully agree with the reviewer that accurately simulating historical conditions*  
248 *does not thereby make them reliable for future scenarios. We also agree with the*  
249 *reviewer that a discussion of the timescales in which Hector is useful over is needed*  
250 *within the manuscript (section 6.2).*

251 *“We compare Hector to MAGICC and CMIP5 under differing future climate projections.*  
252 *Hector’s strengths lie within policy relevant time scales of decades to centuries. Studies*  
253 *suggest that 80% of the anthropogenic CO<sub>2</sub> emissions have an average atmospheric*  
254 *lifetime of 300-450 years (Archer et al., 1997; Rogner, 1997; Archer, 2005). Hector has*  
255 *all the necessary components to model the climate system from present day through the*  
256 *next approximately 300 years.”*

257

258 • Hector is unable to reproduce 1970-2010 temperatures (Fig 4). These errors should be  
259 described in the text, including possible explanations linked to underlying physics.

260

261 *We have addressed and fixed the high temperature sensitivity after 1960 by including a*  
262 *variable ocean heat flux, as well as lagging the temperature effects from atmospheric*  
263 *[CO<sub>2</sub>]. There are numerous processes that are not simulated in Hector that buffer the*  
264 *temperature effects of increasing GHGs. Therefore, we take a simple approach in this*  
265 *current version and lag our temperature. We have addressed this in the manuscript*  
266 *section 4.1, “As global temperatures rise, the uptake capacity of the ocean thus*  
267 *diminishes, simulating both a saturation of heat in the surface and a slowdown in ocean*  
268 *circulation with increased temperatures. Finally, the temperature effects from*  
269 *atmospheric [CO<sub>2</sub>] are lagged in time, as there are numerous real-world processes not*  
270 *simulated in Hector buffering the temperature effects of increasing atmospheric [CO<sub>2</sub>].”*  
271 *See figures 4 and 8 for updated global temperature change.*

272

273 • Atmospheric CO<sub>2</sub> concentrations in figure 5 are only shown from 1850-2100. Is there a  
274 reason why this plot isn't extended to 2300 like figures 6-11? If model errors are  
275 prevalent from 2100-2300, it is essential that this plot show the entire time range.

276

277 *The CMIP5 archive of models that ran esmrcp85, do not run out to 2300. Therefore, we*  
278 *can only compare out to 2100. The caption for Figure 5 has been updated:*

279 *“Figure 5: Atmospheric [CO<sub>2</sub>] from 1850 to 2100 under RCP 8.5 for Hector (blue),*  
280 *MAGICC6 (green), Mauna Loa (purple), Law Dome (brown) and esmRCP 8.5*  
281 *(prescribed emissions scenario) CMIP5 median, one standard deviation and model range*  
282 *(pink, n=4 (1850-2000) and n=5 (2001-2100)). Note that the CMIP5 models run under*  
283 *esmrcp85 do not extend to 2300.”*

284

285 • Hector also appears unable to reproduce temperatures in CMIP5 models past year 2100  
286 (Fig 8). This misrepresentation is downplayed in the text (page 7092, lines 11-20). It is  
287 not sufficient to simply state that errors are negligible because correlations are high. It is  
288 unclear whether this is an error in the temperature or the carbon cycle model of Hector  
289 because the experiment is not well described. Please clarify.

290

291 *Since fixing the temperature problem post 1960s, Hector is now closer to the CMIP5*  
292 *median post 2100, than MAGICC6 is. Post 2100, Hector remains within the standard*  
293 *deviation of the CMIP5 models. We have included in the figure captions, the numbers of*  
294 *models for each scenario, for each time period. Post 2100, the number of model run out*  
295 *to 2300 drops off dramatically, which could be responsible for some of the differences*  
296 *between the CMIP5 median and Hector.*

297

298 • The authors do a nice job highlighting deviations in the atmosphere-ocean flux in  
299 Hector from CMIP5 models after 2100 (Fig 10). However, these deviations do not seem  
300 trivial, and may impact long-term projections. If Hector cannot be trusted after 2100,  
301 this should be stated.

302 Until a later version of Hector is released with an updated modeling approach, the authors  
303 should acknowledge these issues and should add discussion on the physical causes that  
304 may produce deviations from observations (or more complex models). The authors do  
305 include some discussion of the underlying physics at the end of section 6, but more  
306 should be included throughout the manuscript.

307

308 *The authors agree with the reviewer and have since updated section 6.2 with more detail:*

309 *“Hector’s calculation of air-sea fluxes is within the large CMIP5 model range up to*  
310 *2100. However, after that Hector peaks close to 2150, while the CMIP5 models are*  
311 *beginning to decline. One potential reason for this discrepancy after 2100 is that in this*  
312 *version of Hector, we do not simulate changes in ocean circulation, potentially biasing*  
313 *fluxes too high after 2100. Most ESMs in CMIP5 show a weakening of the Atlantic*  
314 *meridional overturning circulation by 2100 between 15% and 60% under RCP 8.5*  
315 *(Cheng et al., 2013). A slowdown in ocean circulation may result in less carbon uptake*  
316 *by the oceans, as seen in Figure 9. Another potential reason for this bias is Hector’s*  
317 *constant pole to equator ocean temperature gradient. Studies show that the Arctic is*  
318 *warming faster than the rest of the globe (e.g., Bintanja and van der Linden, 2013;*

319 *Holland and Bitz, 2003; Bekryaev et al., 2010). A warmer high latitude surface ocean in*  
320 *Hector would suppress the uptake of carbon, potentially contributing to higher air-sea*  
321 *fluxes after 2100.”*

322

### 323 **Technical corrections - figures and tables**

324 • All figures: Figure text is too small.

325

326 *Figure fonts and line size have been enlarged for all figures.*

327

328 • Figure 2: Describe (in caption or key) the definitions of variables TT , EIL, EID, etc.

329 *A reference to Table 2 has been included in the figure caption.*

330

331 • Figures 3-5, and 8: use consistent colors for models across figures. It is very hard to  
332 compare across figures when Hector output is shown as yellow in one plot and green in  
333 another.

334

335 *The authors agree with the reviewer and have updated all figures to have the same color*  
336 *scheme.*

337

338 • Figure 8: Label panels a, b, c, and d.

339

340 *Figure 8 has been updated.*

341

342 • Tables 1 and 2: Include references for initial condition values where applicable.  
343 For example, the recent IPCC estimates a pre-industrial total oceanic carbon content of  
344 \_38,000 GtC. Numbers here are closer to 35,000 GtC. This difference is not likely  
345 significant for Hector, but my confidence in the model would be higher with references to  
346 justify these numbers.

347

348 *The authors agree with the reviewer and have added references for initial values where*  
349 *applicable in Table 1 & 2.*

350

### 351 **Technical corrections - text**

352 (Note that I did not provide comments for sections 4.2.1-4.2.6, and suggest a different  
353 reader with expertise in this area to review this material.)

354 • Page 7077, line 5: #4 (modeling the carbon cycle) seems a subset of #1 (calculating  
355 future concentrations of greenhouse gases). Either remove #4 or move it up as an explicit  
356 subset of #1 (or explain what is meant, if I am missing something). The order should  
357 reflect the general order of operations in an SCM.

358

359 *Sentence edited as suggested:*

360 *“Most SCMs have a few key features: 1) calculating future concentrations of greenhouse*  
361 *gases (GHGs) from given emissions while modeling the global carbon cycle; 2)*

362 *calculating global mean radiative forcing from greenhouse gas concentrations; and 3)*

363 *converting the radiative forcing to global mean temperature (e.g., Wigley, 1991;*

364 *Meinshausen et al., 2011a; Tanaka et al., 2007b; Lenton, 2000).”*



365  
366 • Page 7077, line 7: Recommend changing the word “policy” to “decision making”.  
367  
368 *Sentence edited as suggested:*  
369 “With these capabilities, SCMs play an integral role in decision making and scientific  
370 research.”  
371  
372 • Page 7077, lines 12-13: Recommend changing “have a simple representation” to “rely  
373 on simple representations”.  
374  
375 *Sentence edited as suggested:*  
376 “Therefore, all IAMs rely on a simple representation of the global climate system.”  
377  
378 • Page 7077, lines 24-27: Consider re-writing the first sentence of this paragraph. There is  
379 also a grammatical error in this sentence: “therefore are used for run multiple simulations  
380 of future climate change...”  
381  
382 *Sentence edited as suggested:*  
383 “Lastly, SCMs are computationally efficient and inexpensive to run. Therefore, they are  
384 used to run multiple simulations of future climate change emissions scenarios,  
385 parameter sensitivity experiments, perturbed physics experiments, large ensemble runs,  
386 and uncertainty analyses (Senior and Mitchell, 2000; Hoffert et al., 1980; Harvey and  
387 Schneider, 1985; Ricciuto et al., 2008; Sriver et al., 2012; Irvine et al., 2012).”  
388  
389 • Page 7077, line 29: Please be more specific with wording choice for “fast enough”.  
390  
391 *Sentence edited as suggested:*  
392 “Lastly, SCMs are computationally efficient and inexpensive to run.”  
393  
394 • Page 7078, line 5: “This study introduces Hector v0.1, **an** object-oriented, simple...”  
395  
396 *Sentence edited as suggested:*  
397 “This study introduces Hector v1.0, an object-oriented, simple...”  
398  
399 • Page 7078, line 11: Consider changing the word “basic” to “fundamental”.  
400  
401 *Sentence edited as suggested:*  
402 “One of the fundamental questions faced in developing a SCM is how much detail should  
403 be represented in the climate system.”  
404  
405 • Page 7082, line 19: typo– “-political”  
406 *Formatting issues with a ‘-‘ have been corrected.*  
407  
408 • Page 7083, line 6: typo– “NPP is modified by a the use-specified...”  
409 *Formatting issues with a ‘-‘ have been corrected.*  
410

411 • Page 7083, line 7: Does (or can) beta change with time or temperature? If parameter is  
412 fixed, state that explicitly.  
413  
414 *No, beta (the shape of the NPP response to CO2 fertilization) doesn't change with time. It*  
415 *does, optionally, change spatially: users can define separate beta values for different*  
416 *biomes, for example.*  
417 *“These are commonly used formulations: NPP is modified by a user-specified carbon*  
418 *fertilization parameter,  $\beta$  (Piao et al., 2013), that is constant in time but not necessarily*  
419 *in space. For example, users can define separate  $\beta$  values for different biomes.”*  
420  
421 • Page 7083, line 14: Do you mean Eqs. (7)-(9)? Correct if this is a typo.  
422  
423 *The authors corrected this typo.*  
424  
425 • Page 7083, eqs 7-9: Explicitly define all terms and/or refer to Table 1. Terms do not  
426 match those in Table 1 (e.g. FLC).  
427  
428 *The authors have corrected this.*  
429  
430 • Page 7084, lines 10-12: This assumption is essentially a statement of fixed equator-pole  
431 temperature gradient. But when the Earth warms, the poles tend to warm more than the  
432 equator. This assumption should be discussed explicitly, including under what conditions  
433 it would affect the performance of Hector  
434  
435 *Within Hector it is assumed a fixed equator-pole temperature gradient in sea surface*  
436 *temperature. While this may not hold under future warming scenarios, v1.0 of Hector is a*  
437 *simple representation of the climate system and this change in temperature gradient is a*  
438 *major future improvement to the model. A warmer high latitude ocean will potentially*  
439 *result in less CO<sub>2</sub> uptake in the high latitude ocean.*  
440 *“We assume a constant pole to equator temperature gradient, but acknowledge that this*  
441 *assumption may not hold true if the poles warm faster than the equator.”*  
442  
443 • Page 7084, lines 21-23: Carbon cycle description (section 3 up to 3.1) is incomplete.  
444 Presumably the model includes the non-linear effects in oceanic carbon uptake from  
445 changing ocean acidity as atmospheric carbon is transferred to the upper ocean, but these  
446 are not described. The relevant equations should be included here. Some discussion  
447 comes later on page 7093, but the pH dependence is not well described.  
448  
449 *This has been addressed under section 3.0:*  
450 *“We model the nonlinearity of the inorganic carbon cycle, calculating pCO<sub>2</sub>, pH, and*  
451 *carbonate saturations based on equations from Zeebe and Wolf-Gladrow, (2001). The*  
452 *flux of CO<sub>2</sub> for each box i is calculated by:*  

$$F_i(t) = k \alpha \Delta pCO_2 \quad (11)$$
453 *where k is the CO<sub>2</sub> gas-transfer velocity,  $\alpha$  is the solubility of CO<sub>2</sub> in water based on*  
454 *salinity, temperature, and pressure, and  $\Delta pCO_2$  is the atmosphere-ocean gradient of*  
455 *pCO<sub>2</sub> (Takahashi et al., 2009). The calculation of pCO<sub>2</sub> in each surface box is based on*

456 *the concentration of CO<sub>2</sub> in the ocean and its solubility (a function of temperature,*  
457 *salinity, and pressure)."*  
458

- 459 • Page 7089, line 17: Please be more specific with "other models". Do the authors mean  
460 more complex models? Or widely used models? Or both?

461 *Sentence edited as suggested:*  
462 *"A critical test of Hector's performance is to compare the major climatic variables*  
463 *calculated in Hector, e.g., atmospheric [CO<sub>2</sub>], radiative forcing, and atmospheric*  
464 *temperature, to observational records and both simple and complex climate models. "*  
465

- 466 • Page 7090, line 8: Spell out "SD".

467  
468 *Sentence edited as suggested: "standard deviation"*  
469

- 470 • Page 7090, line 24: Remove words "a few".

471  
472 *Sentence edited as suggested: removed "a few"*  
473

- 474 • Page 7090, line 25-26: Consider re-wording sentence.

475  
476 *Sentence edited as suggested:*  
477 *"After spinup is complete in Hector, atmospheric [CO<sub>2</sub>] in 1850 is 286.0 ppmv, which*  
478 *compares well with observations from Law Dome of 285.2 ppmv."*  
479

- 480 • Page 7091, line 19: Is Hector actually perfectly correlated here, or is R=1.0 from  
481 rounding? Please double check.

482  
483 *The authors have since removed the correlation values from the manuscript. We have*  
484 *replaced them with absolute changes over given time periods. We feel that this is a better*  
485 *comparison between all the models, than correlation. 2 models can be well correlated,*  
486 *but that does not necessarily suggest that they are in agreement.*  
487

- 488 • Page 7092, line 23: Grammatical error – "the higher the correlation and low RMSE  
489 between CMIP5 and : : :". Presumably what is intended is "the lower the RMSE".

490  
491 *The authors have since removed figure 9 from the manuscript as well.*  
492

- 493 • Page 7093, line 23: Change "see" to "estimate".

494  
495 *Sentence edited as suggested:*  
496 *"We estimate a significant drop in pH from present day through 2100."*  
497

498 **A simple object-oriented and open source model for scientific and policy analyses of**  
499 **the global ~~carbon cycle~~climate system–Hector v~~1.00.1~~**

500 C.A. Hartin\*, P. Patel, A. Schwarber, R.P. Link, and B.P. Bond-Lamberty

501

502 Pacific Northwest National Laboratory, Joint Global Change Research Institute at the

503 University of Maryland–College Park, 5825 University Research Court, College Park, MD

504 20740, USA

505

506 \*Corresponding author: [corinne.hartin@pnnl.gov](mailto:corinne.hartin@pnnl.gov)

507

508

509 **Abstract**

510 Simple climate models play an integral role in the policy and scientific communities.  
511 They are used for climate mitigation scenarios within integrated assessment models,  
512 complex climate model emulation, and uncertainty analyses. ~~Here we describe~~  
513 Hector v~~1.00-1~~, an open source, object-oriented, simple global climate carbon-cycle  
514 model. This model runs essentially instantaneously while still representing the most  
515 critical global scale earth system processes. Hector has a three-part main carbon  
516 ~~pool~~cycle: an a one-pool atmosphere, land, and ocean. ~~The model's terrestrial~~  
517 carbon cycle includes primary production and respiration ~~and fluxes primary production,~~  
518 accommodating arbitrary geographic divisions into, e.g., ecological biomes or political  
519 units. Hector actively solves the inorganic carbon system in the surface ocean, directly  
520 calculating air-sea fluxes of carbon and ocean pH. Hector reproduces the global  
521 historical trends of atmospheric [CO<sub>2</sub>], radiative forcing, and surface temperatures. The  
522 model simulates all four Representative Concentration Pathways with equivalent  
523 ~~changes~~high correlations (R > 0.7) of key variables over time compared with to current  
524 observations, MAGICC (a well-known simple climate model), and models from the 5<sup>th</sup>  
525 Coupled Model Intercomparison Project ~~version 5~~. Hector's flexibility, is freely available  
526 ~~under an~~ open source licensenature, and ~~its~~ modular design will facilitate a broad range  
527 of research in various areas.

528

529

530 **1.0 Introduction**

531 Projecting future impacts of anthropogenic perturbations on the climate system  
532 relies on understanding the interactions of key earth system processes. To accomplish  
533 this, a hierarchy of climate models with differing levels of complexity and resolution are  
534 used, ranging from purely statistical or empirical models, to simple energy balance  
535 models, ~~to fully-coupled atmosphere-ocean-general-circulation models (AOGCMs)~~Earth  
536 System Models (ESMs) (Stocker, 2011).

537 Simple-Reduced-complexity or simple climate models (SCMs) lie in the middle of  
538 this spectrum, representing only the most critical global scale earth system processes  
539 with low spatial and temporal resolution, e.g., carbon fluxes between the ocean and  
540 atmosphere, primary production and respiration ~~fluxes and primary production~~ on land.  
541 These models are relatively easy to use and understand, and are computationally  
542 inexpensive. -Most SCMs have a few key features: 1) calculating future concentrations of  
543 greenhouse gases (GHGs) from given emissions and while modeling the global carbon  
544 cycle; 2) calculating global mean radiative forcing from greenhouse gas concentrations;  
545 and 3) converting the radiative forcing to global mean temperature, ~~and 4) modeling~~  
546 ~~the carbon cycle, an essential part of the climate system~~ (e.g., Wigley, 1991;  
547 Meinshausen et al., 2011a; Tanaka et al., 2007b; Lenton, 2000).

548 With these capabilities, SCMs play an integral role in policy decision making and  
549 scientific research. For example, energy-economic-climate models or Integrated  
550 Assessment Models (IAMs) are used to address issues on energy system planning,  
551 climate mitigation, ~~and~~ stabilization pathways, and land-use changes, ~~pollution control,~~

552 ~~and population policies~~ (Wigley et al., 1996; Edmonds and Smith, 2006; van Vuuren et  
553 al., 2011). ~~AOGCMs-ESMs~~ are too computationally expensive to use in these analyses.  
554 Therefore, all IAMs ~~rely on~~have a simple representation ~~of~~ the global climate system  
555 ~~in which emissions data from the IAMs are converted to concentrations and then~~  
556 ~~radiative forcing and global temperature are calculated.~~

557 Depending on the purpose of the IAMs (economics, cost-benefit analysis, or more  
558 physical based processes), the corresponding climate and carbon component varies in  
559 complexity and resolution. For example, models like DICE, FUND, and MERGE have a  
560 strongly highly simplified carbon/climate system (Nordhaus, 2008; Anthoff and Tol,  
561 2014; Manne and Richels, 2005). IAMs focusing more on the physical processes of the  
562 natural system and the economy ~~have a~~employ more complex representations of the  
563 climate/carbon system. Models like GCAM (Global Change Assessment Model) and  
564 MESSAGE use MAGICC as their SCM (Meinshausen et al., 2011a; Riahi et al., 2007; Calvin  
565 et al., 2011). Increasing in complexity, some IAMs include the climate/carbon system at  
566 gridded scales (e.g., IMAGE), and can be coupled to earth system models of  
567 intermediate complexity (e.g., MIT IGSM), or more recently coupled to a full earth  
568 system model (the iESM project) (Bouwman et al., 2006; Sokolov et al., 2005; Bond-  
569 Lamberty et al., 2014; Di Vittorio et al., 2014; Collins et al., 2015).

570 SCMs such as MAGICC, GENIE, and the climate emulation tool at RDCEP are also  
571 used as emulators of more complex ~~AOGCMs-ESMs, such as MAGICC, GENIE, and the~~  
572 climate emulation tool at RDCEP (Meinshausen et al., 2011c; Schlesinger and Jiang,  
573 1990; Challenor, 2012; Ratto et al., 2012; Lenton et al., 2009; Castruccio et al., 2014).

574 The ~~components behavior~~ of SCMs can be constrained to replicate the overall behavior  
575 of the more complex ~~model-ESM-components~~. For instance, the climate sensitivity of a  
576 SCM can be made equal to that of an ~~ESM-AOGCM~~ by altering a single model  
577 parameter. ~~In particular, the One-SCM, MAGICC, model~~ has been central to the  
578 analyses presented in the Intergovernmental Panel on Climate Change (IPCC) reports,  
579 ~~and can be parameterized to emulating emulate~~ a large suite of ~~AOGCMs-ESMs~~  
580 (Meinshausen et al., 2011a).

581 Lastly, SCMs are computationally efficient and inexpensive to run, ~~and therefore,~~  
582 ~~they~~ are used ~~for to~~ run multiple simulations of future climate change emissions  
583 scenarios, parameter sensitivity experiments, perturbed physics experiments, large  
584 ensemble runs, and uncertainty analyses (Senior and Mitchell, 2000; Hoffert et al., 1980;  
585 Harvey and Schneider, 1985; Ricciuto et al., 2008; Sriver et al., 2012; Irvine et al., 2012).  
586 ~~SCMs are fast enough computationally efficient in that multiple scenarios can be~~  
587 ~~simulated, and a wide range of parameter values can be tested. MAGICC, the Bern CC~~  
588 ~~model, and SNEASY are examples of a few models used for uncertainly analysis~~  
589 (Meinshausen et al., 2011c; Urban and Keller, 2010; Joos et al., 2001b). ~~Specifically,~~  
590 SCMs have been useful in reducing uncertainties in future CO<sub>2</sub> sinks, quantifying  
591 parametric uncertainties in sea-level rise, ice-sheet modeling, ocean-heat uptake, and  
592 aerosol forcings (Ricciuto et al., 2008; Sriver et al., 2012; Applegate et al., 2012; Urban  
593 and Keller, 2009).

594 This study introduces Hector v~~0.1.0~~, ~~an open source~~, object-oriented, simple climate  
595 carbon-cycle model. ~~Hector was developed with three main goals in mind. First, Hector~~



596 | is an open source model, ~~Hector is open source~~, an important quality given that the  
597 | scientific community, funding agencies, and journals are increasingly emphasizing  
598 | transparency and open source (E.P. White, 2013; Heron et al., 2013), particularly in  
599 | climate change sciences (Wolkovich et al., 2012) ~~(Wolkovich et al. 2012)~~. ~~With an open~~  
600 | ~~source model a~~ large community of scientists can access, use, and enhance open  
601 | source models, with the potential for long-term utilization, improvement, and  
602 | reproducibility (Ince et al., 2012) ~~—~~. Second, a clean design using an object-oriented  
603 | framework is critical for Hector development and future use. This allows for new  
604 | components to easily be added to Hector, i.e. the model’s functionality to be easily  
605 | extended in the future ~~not currently included in the core version~~. ~~More importantly~~ In  
606 | addition, this framework allows for easy coupling into IAMs, in particular GCAM. Lastly,  
607 | Hector is a stand-alone simple climate model used to answer fundamental scientific  
608 | research questions, uncertainty analysis, parameter sensitivities, etc.

609 |       One of the ~~basic-fundamental~~ questions faced in developing a SCM is how much  
610 | detail should be represented in the climate system. Our goal is to introduce complexity  
611 | only where warranted, keeping the representations of the climate system as simple as  
612 | possible. This results in fewer calculations, faster execution times, and easier analysis  
613 | and interpretation of results. Sections 2, 3, and 4 describe the structure and  
614 | components of Hector. Sections 5 and 6 describe the experiments, results and  
615 | comparison of Hector against observational data and other models (MAGICC and  
616 | CMIP5).

617

## 618 **2.0 Model architecture**

### 619 **2.1 Overall structure and design**

620 Hector is written in C++ and uses an object-oriented design that enforces clean  
621 separation between its different parts, which interact via strictly defined interfaces. The  
622 separation keeps each software module self-contained, which makes the code easy for  
623 users to understand, maintain, and enhance. Entities in the model include a command-  
624 line *wrapper*, the model *coupler*, various *components* organized around scientific areas  
625 (carbon cycling, radiative forcing, etc.) and *visitors* responsible for model output. Each of  
626 these is discussed below.

### 627 **2.2 Model Coupler**

628 Hector's control flow starts with the coupler, which is responsible for: 1) parsing  
629 and routing input data to the model components; 2) tracking how the components  
630 depend on each other; 3) passing messages and data between components; 4) providing  
631 facilities for logging, time series interpolation, etc.; and 5) controlling the main model  
632 loop as it progresses through time. Any errors thrown by the model are caught by the  
633 wrapper, which prints a detailed summary of the error.

634 Input data are specified in flat text files, and during startup are routed to the  
635 correct model component for its initialization. Some of the key initial model conditions  
636 are summarized in **Table 1 and Table 2**. For more details of initial model conditions we  
637 | urge the reader to download Hector v~~0~~1.0 (<https://github.com/JGCRI/hector>).

638 Components can send messages to each other during the model run, most often

639 | requesting data. The messaging interface is also available to external subroutines, such  
640 | as components of IAMs or other linked models. The coupler handles message routing  
641 | (via the *capability* mechanism, below) and enforces mandatory type checking: e.g., if a  
642 | component requests mean global temperature in °C but the data are provided in K, an  
643 | error will be thrown (i.e., execution halts) unless the receiving component can handle  
644 | this situation.

645 | Visitor patterns- are units of code that traverse all model components and  
646 | handle model output (Martin et al., 1997). Two visitors currently exist: one saves an  
647 | easily-readable summary table to an output file, while the other writes a stream of  
648 | model data (both standard outputs and internal diagnostics). After the model ~~is~~  
649 | ~~finisheds-running~~, this ‘stream’ file can be parsed and summarized by R scripts (R  
650 | Development Core Team, 2014) included with ~~the code~~Hector-. Log files may also be  
651 | written by any model entity, using facilities provided by the coupler. The full sequence  
652 | of events during a model run is summarized in **Figure 1**.

### 653 | **2.3 Components**

654 | Model components are submodels that communicate with the coupler. From  
655 | the coupler’s point of view, components are fully defined by their *capabilities* and  
656 | *dependencies*. At model startup, before the run begins, components inform the coupler  
657 | of their capabilities, i.e., what data they can provide to or accept from the larger model  
658 | system. The coupler uses this information to route messages, such as requests for data,  
659 | ~~such as requests for data~~. Components also register their  
660 | dependencies, i.e., what data-results they require from other components in order ~~for~~

661 | ~~the~~ to complete their computations. After initialization, but before the model begins to  
662 | run, the coupler uses this dependency information to determine the order in which  
663 | components will be called in the main control loop.

664 |         The model’s modular architecture, and the *capability/dependency* systems  
665 | described above, allows swapping, enabling and disabling of model components directly  
666 | via the input without recompiling. For example, this means that a user can test two  
667 | different ocean submodels and easily compare results without having to rebuild the  
668 | model.

#### 669 |                 **2.4 Time step, spinup, and constraints**

670 |         The model’s fundamental time step is 1 year, although the carbon cycle can  
671 | operate on a finer resolution when necessary (Section [2.6.13.1](#)). When the model is on  
672 | an integer date (e.g. 1997.0) it is considered to be the midpoint of that particular  
673 | calendar year, in accordance with Representative Concentration Pathway (RCP) data  
674 | (Meinshausen et al., 2011b)-.

675 |         Like many models, Hector has an optional ‘spinup’ step, in which the model runs  
676 | to equilibrium in an a\_historical, perturbation-free mode (Pietsch and Hasenauer, 2006).  
677 | This occurs after model initialization, but before the historical run begins, and ensures  
678 | that the model is in steady state when it enters the main simulation. During spinup, the  
679 | coupler repeatedly calls all the model components in their dependency-driven ordering,  
680 | using an annual time step. Each component signals whether it needs further steps to  
681 | stabilize, and this process repeats until all components signal that they are complete.

682 | Currently only the model's carbon cycle makes use of the spinup phase. ~~Spinup~~  
683 | Spinup takes place prior to land use change or industrial emission inputs, ~~and t~~The  
684 | main carbon cycle moves from its initial, user-defined carbon pool values to a steady  
685 | state in which  $\delta dC/dt < \epsilon$  for all pools; ~~the~~The convergence criterion  $\epsilon$  is user-definable;  
686 | ~~and~~ by default  $\epsilon = 1 \text{ Tg C yr}^{-1}$ . From its default values the preindustrial carbon cycle will  
687 | typically stabilize in 300-400 time steps.

688 | ~~Hector can be forced to~~ The model can be constrained, i.e., matching its output  
689 | to a user-supplied time series. ~~This is helpful to, to allow isolation and testing of~~  
690 | different components. Available constraints currently include atmospheric CO<sub>2</sub>, global  
691 | temperature anomaly, total ocean-atmosphere carbon exchange, total land-atmosphere  
692 | carbon exchange, and total radiative forcing. Most constraints operate by overwriting  
693 | model-calculated values with user-supplied time series data during the run. The  
694 | atmospheric [CO<sub>2</sub>] constraint operates slightly differently, as the global carbon cycle is  
695 | subject to a continuous mass-balance check. As a result, when the user supplies a [CO<sub>2</sub>]  
696 | record between arbitrary dates and orders the model to match it, the model *computes*  
697 | [CO<sub>2</sub>] at each time step, and any deficit (surplus) in comparison with the constraint [CO<sub>2</sub>]  
698 | is drawn from (added to) the deep ocean. The deep ocean holds the largest reservoir of  
699 | carbon; therefore, small changes in this large pool have a negligible effect on the carbon  
700 | cycle dynamics. When the model exits the constraint time period, atmospheric [CO<sub>2</sub>]  
701 | again becomes fully prognostic.

## 702 | **2.5 Code availability and dependencies**

703 All Hector code is open source and available at  
704 <https://github.com/JGCRI/hector/>. The repository includes model code that can be  
705 compiled on Mac, Linux, and Windows, input files for the four Representative  
706 Concentration Pathways (RCP) cases discussed in Section 4.5, R scripts to process model  
707 output, and extensive documentation. ~~We kept the~~ Software dependencies are as  
708 limited as possible, with only the GNU Scientific Library (GSL, Gough, 2009) and the  
709 Boost C++ libraries (<http://www.boost.org>) required. An optional unit testing build  
710 target requires the googletest framework (<http://code.google.com/p/googletest>).  
711 However, this is not needed to compile and run Hector. HTML documentation can be  
712 automatically generated from the code using the Doxygen tool  
713 (<http://www.doxygen.org>). All these tools and libraries are free and open source.

714

### 715 **3.0 ~~Main~~ Carbon Cycle Component**

716 In the model's default terrestrial carbon cycle, terrestrial vegetation, detritus,  
717 and soil are linked with each other and the atmosphere by first-order differential  
718 equations (**Figure 2**). Vegetation net primary production is a function of atmospheric  
719 [CO<sub>2</sub>] and temperature. Carbon flows from the vegetation to detritus and then to soil,  
720 losing fractions to heterotrophic respiration on the way. Land-use change emissions are  
721 specified as inputs. An 'earth' pool debits carbon emitted as anthropogenic emissions,  
722 allowing a continual mass-balance check across the entire carbon cycle.

723 More formally, any change in atmospheric carbon, and thus [CO<sub>2</sub>], occurs as a  
724 function of anthropogenic fossil fuel and industrial emissions (F<sub>A</sub>), land-use change

725 | emissions ( $F_{LC}$ ), and the atmosphere-ocean ( $F_O$ ) and atmosphere-land ( $F_L$ ) carbon fluxes.

726 | The atmosphere is treated as a single well-mixed box whose rate of change is:

$$\frac{dC_{atm}}{dt} = F_A(t) + F_{LC}(t) - F_O(t) - F_L(t) \quad (1)$$

727 | ~~where,  $F_A$  is the anthropogenic emissions,  $F_{LC}$  is the land use change emissions~~

728 | ~~and  $F_O$  and  $F_L$  are the atmosphere-ocean and atmosphere-land fluxes. Note, that the~~

729 | ~~carbon cycle is solved under indeterminate time steps (represented in the text by~~

730 | ~~equations with  $d/dt$ ), while most other submodels of Hector are solved under a fixed~~

731 | ~~time step of 1 year (equations with  $\Delta$ ). Future versions of Hector will incorporate~~

732 | ~~indeterminate time steps within all components of the model.~~ The overall terrestrial

733 | carbon balance (Equation 2) excluding user-specified land-use change fluxes at time  $t$  is

734 | the difference between net primary production ( $NPP$ ) and heterotrophic respiration

735 | ( $RH$ ). This is summed over user-specified  $n$  groups (each typically regarded as a latitude

736 | band, biome, or -political units), with  $n \geq 1$ :

$$F_L(t) = \sum_{i=1}^n NPP_i(t) - RH_i(t) \quad (2)$$

737 | Note that  $NPP$  here is assumed to include non-LUC disturbance effects (e.g., fire), for

738 | which there is currently no separate term. For each biome  $i$ ,  $NPP$  and  $RH$  are computed

739 | as a functions of ~~their~~ its preindustrial values  $NPP_0$  and  $RH_0$ , current atmospheric carbon

740 |  $C_{atm}$ , and the biome's temperature anomaly  $T_i$ , while heterotrophic respiration  $RH$

741 | depends upon the pool sizes of detritus ( $C_d$ ) and soil ( $C_s$ ), and global temperatures:

$$NPP_i(t) = NPP_0 * f(C_{atm}, \beta_i) \quad (3)$$

$$f(C_{atm}, \beta_i) = 1 + \beta_i \left( \log \left( \frac{C_{atm}}{C_0} \right) \right) \quad (4)$$

$$RH_{s,d}(t) = C_{s,d} * f_{rs,rd} * Q_{10}^{T_i(t)/10} \quad (5)$$

$$T_i(t) = T_G(t) * \delta_i \quad (6)$$

742 These are commonly used formulations: *NPP* is modified by a ~~the~~ user-specified carbon  
 743 fertilization parameter,  $\beta$  (Piao et al., 2013), that is constant in— time but not  
 744 necessarily in space. ~~Optionally, it can change spatially.~~ For example, users can define  
 745 separate  $\beta$  values for different biomes. *RH* changes are controlled by a biome-specific  
 746  $Q_{10}$  value. Biomes can experience temperature changes at rates that differ from the  
 747 global mean  $T_G$ , controlled by a user specified temperature factor  $\delta_i$ . Note that in  
 748 equation (5), soil *RH* depends on a running mean of past temperatures, ~~an attempt to~~  
 749 representing ~~king~~ the slower propagation of heat through soil strata.

750 Land carbon pools (vegetation, detritus, and soil) change as a result of *NPP*, *RH*,  
 751 and land-use change fluxes, whose effects are partitioned among these carbon pools. In  
 752 addition, carbon flows from vegetation to detritus and to soil (**Figure 2**). Partitioning  
 753 fractions ( $f$ ) control the flux quantities between pools (**Table 2**). For simplicity Equations  
 754 ~~87-109~~ omit the time  $t$  and biome-specific  $i$  notations, but each pool is tracked  
 755 separately for each biome at each time step:

$$\frac{dC_V}{dt} = NPP f_{nv} - C_V(f_{vd} + f_{vs}) - F_{LC} f_{lv} \quad (7)(8)$$

$$\frac{dC_D}{dt} = NPP f_{nd} + C_V f_{vd} - C_D f_{ds} - RH_{det} - F_{LC} f_{ld} \quad (8)(9)$$

$$\frac{dC_S}{dt} = NPP f_{ns} + C_V f_{vs} + C_D f_{ds} - RH_{soil} - F_{LC} f_{ls} \quad (9)(10)$$



756 | The ocean-atmosphere carbon flux is the sum of the ocean's surface fluxes ( $F_i$ )  
757 | (currently  $n=2$ , high and low latitude surface box):

$$F_o(t) = \sum_{i=1}^n F_i(t) - F_a(t) = \sum_{i=1}^n F_i(t) \quad (10)(11)$$

758 | The surface fluxes of each individual box are directly-calculated from an ocean  
759 | chemistry submodel described in detail by Hartin et al. (in prep). We model the  
760 | nonlinearity of the inorganic carbon cycle, calculating pCO<sub>2</sub>, pH, and carbonate  
761 |  saturations based on equations from Zeebe and Wolf-Gladrow, (2001). The flux of CO<sub>2</sub>  
762 | for each box  $i$  is calculated by:

$$F_i(t) = k \alpha \Delta pCO_2 \quad (121)$$

763 | ~~Where~~ where  $k$  is the CO<sub>2</sub> gas-transfer velocity,  $\alpha$  is the solubility of CO<sub>2</sub> in water based  
764 | on salinity, temperature, and pressure, and  $\Delta pCO_2$  is the atmosphere-ocean gradient of  
765 | pCO<sub>2</sub> (Takahashi et al., 2009). The calculation of pCO<sub>2</sub> in each surface box is based on  
766 | the concentration of CO<sub>2</sub> in the ocean and its solubility (a function of temperature,  
767 | salinity, and pressure). At steady state, the cold high latitude surface box (>55°, subpolar  
768 | gyres) acts as a sink of carbon from the atmosphere, while the warm low latitude  
769 | surface box (<55°) off gases carbon back to the atmosphere. Temperatures of the  
770 | surface boxes are linearly related to atmospheric global temperatures (see section 4.1),  
771 |  $T_{HL} = \Delta T - 13$  and  $T_{LL} = \Delta T + 7$  (Lenton, 2000). ~~but~~ The ocean model, modeled after  
772 | Lenton et al., (2000) and Knox and McElroy, (1984), circulates carbon through four  
773 | boxes (two surface, one intermediate depth, one deep), via water mass advection and  
774 | exchange, simulating a simple thermohaline circulation (**Figure 2**). At steady state,

775 approximately 100Pg of carbon are transferred from the high latitude surface box to the  
 776 deep box based on the volume of the box and transport in Sv ( $10^6 \text{ m}^3 \text{ s}^{-1}$ ) between the  
 777 boxes. The change in carbon of any box  $i$  is given by the fluxes in and out, with  $F_{atm \rightarrow i}$  as  
 778 the atmosphere-ocean carbon flux:

$$\frac{dC_i}{dt} = \sum_{j=1}^{in} F_{j \rightarrow i} - \sum_{j=1}^{out} F_{i \rightarrow j} + F_{atm \rightarrow i} \quad (12)(13)$$

779 As the model advances, the carbon values in PgC is converted to or dissolved inorganic  
 780 carbon (DIC) change in each box. The new DIC values are used within the chemistry  
 781 submodel to calculate  $p\text{CO}_2$  values at the next time step.

### 782 3.1 Adaptive-time step solver

783 The fundamental time step in Hector is currently one year, and most model  
 784 components are solved at this resolution. The carbon cycle, however, can operate on a  
 785 variable time step, helping to stabilize it ensuring accurate ODE solutions, even under  
 786 particularly high-emissions scenarios. This will also allow future sub-annual applications  
 787 where desired. The adaptive time step accomplished using the *gsl\_odeiv2\_evolve\_apply*  
 788 solver package of GSL 1.16, which attempts many different step sizes varies the time  
 789 step to reliably (i.e., with acceptable error) keep truncation error within a specific  
 790 tolerance when advancing the model. Thus all the carbon cycle components handle  
 791 indeterminate time steps less than or equal to  $\leq 1$  year, and can signal the solver if a  
 792 too-large time step is leading to instability. The solver then re-retries the solution, using  
 793 a series of smaller steps. From the coupler's point of view, however, the entire model  
 794 continues to advance in annual increments.

795 **4.0 Other Components**

796 **4.1 Global Atmospheric Temperature**

797 Near surface global atmospheric temperature is calculated by:

$$\Delta T(t) = \lambda * RF(t) - F_H(t) \tag{13}(14)$$

798 where, the user-specified  $\lambda$  is the climate feedback parameter, defined as  $\lambda = S'/S$ ,  
799 ~~where~~  $S'$  is the climate sensitivity parameter (3 Kelvin) and  $S$  is the equilibrium climate  
800 sensitivity for a doubling of  $CO_2$  ( $3.7 \text{ Wm}^{-2}$ ) (Knutti and Hegerl, 2008).  $RF$  is the total  
801 radiative forcing and  $F_H$  is the ocean heat flux.  $F_H$  is calculated by a simple sigmoidal  
802 expression of the ocean heat uptake efficiency  $k$  ( $\text{W m}^{-2} \text{K}^{-1}$ ) that decreases with  
803 increasing global temperatures and multiplied by the atmospheric temperature change  
804 prior to the ocean's removal of heat from the atmosphere ( $T_H$ ) (Raper et al.,  
805 ~~2002~~); (Raper et al., 2002).

$$\Delta F_H(t) = k * \Delta T_H(t) \tag{14}$$

806 As global temperatures rise, the uptake capacity of the ocean is thus diminished  
807 slightly, simulating both a saturation of heat in the surface and a slowdown in ocean  
808 circulation with increased temperatures. Finally, in order to better simulate the late  
809 20<sup>th</sup> century rise in global temperature, the temperature effects from atmospheric [CO<sub>2</sub>]  
810 are lagged in time. T, as there are exist are numerous real-world processes not simulated  
811 in Hector that will buffering the temperature effects of increasing atmospheric [CO<sub>2</sub>].  
812 Future versions of Hector will likely address some of these processes for a better  
813 representation of the climate system.

814 **4.2 Radiative Forcing**

815 Radiative forcing is calculated ~~from from~~ a series of atmospheric greenhouse  
816 gases, aerosols, and pollutants (**Eq. ~~15-17, 19-25, 2716, 18-22, 25, 29-30~~**). Radiative  
817 forcing is reported as the relative radiative forcing. The base year user-specified  
818 forcings are subtracted from the total radiative forcing to yield a forcing relative to the  
819 base year (1750). ~~In the current model of Hector, the gases other than CO<sub>2</sub> are only  
820 used for the calculation of radiative forcing.~~

821 **4.2.1. CO<sub>2</sub>**

822 Radiative forcing from atmospheric [CO<sub>2</sub>] in W m<sup>-2</sup> is calculated based on

823 Meinshausen et al. (2011a):

$$RF_{CO_2} = 5.35 * \log \frac{Ca}{C0} \quad (15)$$

824 where, 5.35 W m<sup>-2</sup> is a scaling parameter from Myhre et al. (1998), Ca is the  
825 current atmospheric [CO<sub>2</sub>] in ppmv and C0 is the preindustrial [CO<sub>2</sub>] in ppmv.

826

827 **4.2.12 Halocarbons**

828 The halocarbon component of the model can accept an arbitrary number of gas  
829 species, each characterized by a name, a lifetime  $\tau$  (yr), a radiative forcing efficiency  ~~$\alpha_p$~~   
830 (W m<sup>-2</sup> pptv<sup>-1</sup>), a an optional user-specified preindustrial concentration (pptv), and a  
831 molar mass (g). For each gas, its concentration ( $C_i$ ) at time  $t$  is then computed based on  
832 a specified emissions time series  $E$ , assuming an exponential decay from the  
833 atmosphere:

$$C(t) = C_0 * \exp\left(-\frac{1}{T}\right) + E * T * \left(1 - \exp\left(-\frac{1}{T}\right)\right) \quad (16)$$

834

$$C_t(t) = C_t(t-1) \left(1 - \frac{1}{\tau}\right) + E_t(t) \quad (16)$$

835

E is corrected for atmospheric dry air mole constant (1.8) and the molar mass of each

836

halocarbon. The default model input files include these parameters and a time series of

837

emissions for C2F6, CCl4, CF4, CFC11, CFC12, CFC113, CFC114, CFC115, CH3Br, CH3CCl3,

838

CH3Cl, HCF22, HCF141b, HCF142b, HFC23, HFC32, HFC125, HFC134a, HFC143a,

839

HFC227ea, HFC245ca, HFC245fa, HFC4310, SF6, halon1211, halon1301, and halon2402.

840

Radiative forcing by halocarbons, and other gases controlled under the Montreal

841

Protocol, SF<sub>6</sub>, and ozone are calculated via:

$$RF = \alpha [ C(t) - C(t_0) ] \quad (17)$$

842

where  $\alpha$  is the radiative efficiency (input parameters) in W m<sup>-2</sup> ppbv<sup>-1</sup>, and C is the

843

atmospheric concentration.

844

### 4.2.3 Ozone

845

Tropospheric ozone concentrations are calculated from the CH<sub>4</sub> concentration

846

and the emissions of three primary pollutants: NO<sub>x</sub>, CO, and NMVOCs, modified from

847

Tanaka et al. (2007a):

$$O_{3t} = (5.0 * \ln[CH_4]) + (0.125 * ENO_x) + (0.0011 * ECO) \quad (18)$$

$$+ (0.0033 * EVOC)$$

848 where the constants are the ozone sensitivity factors for each of the precursors (Ehhalt  
 849 et al., 2001). The radiative forcing of tropospheric ozone is calculated from a linear  
 850 relationship using a radiative efficiency factor (Joos et al., 2001a):

$$RF_{O_3} = 0.042 * [O_3] \quad (19)$$

851 **4.2.2 Ozone**

852 ~~Tropospheric ozone concentrations are calculated by the current CH<sub>4</sub>~~  
 853 ~~concentration and the emissions of three primary pollutants: NO<sub>x</sub>, CO, and~~  
 854 ~~NMVOCs(2007a):~~

$$\begin{aligned}
 O_3(t) = & O_3(2000) \\
 & + 5.0 \ln \left[ \frac{CH_4(t)}{CH_4(2000)} \right] \\
 & + 0.125 [eNO_x(t) - eNO_x(2000)] \\
 & + 0.0011 [eCO(t) - eCO(2000)] \\
 & + 0.0033 [eVOC(t) - eVOC(2000)]
 \end{aligned} \quad (18)$$

855 ~~where the constants are the ozone sensitivity factors for each of the~~  
 856 ~~precursors (Ehhalt et al., 2001). The radiative forcing of tropospheric ozone is~~  
 857 ~~calculated from a linear relationship using a radiative efficiency factor (Joos et~~  
 858 ~~al., 2001a) and a pre-industrial value of ozone of 25 DU (IPCC, 2001):~~

$$RF_{O_3} = 0.042 * [O_3] - [O_3]_{pre} \quad (19)$$

859 **4.2.43 BC and OC**

860 The radiative forcing from black and organic carbon is a function ~~of the black carbon and~~  
 861 ~~organic carbon~~their emissions (eEBC and eEOC).

$$RF_{BC} = 0.0743 \cdot 10^{-9} \text{ Wm}^{-2} \text{ Tkg}^{-1} * EeBC \quad (2020)$$

$$RF_{OC} = -0.0128 \cdot 10^{-9} \text{ Wm}^{-2} \text{ Tkg}^{-1} * EeOC \quad (211)$$

862 The coefficients ~~0.0743 \* 10<sup>-9</sup>~~ and ~~0.0128 \* 10<sup>-9</sup>~~ include both indirect and direct forcings  
863 of black and organic carbon (fossil fuel and biomass) (Bond et al., 2013, table C1).

#### 864 **4.2.54 Sulphate Aerosols**

865 The radiative forcing from sulphate aerosols is a combination of the direct and indirect  
866 forcings (Joos et al., 2001a).

$$RF_{SOx\ Direct} = -0.435 \text{ Wm}^{-2} * \frac{ESO_{x_t} eSO_{\bar{x}}(t)}{ESO_{x_{t0}} eSO_{\bar{x}}(2000)} \quad (222)$$

$$RF_{SOx\ Indirect} = -0.68 \text{ Wm}^{-2} * \frac{(\ln(EeSN) + EeSO_{x_t} eSO_{\bar{x}}(t))}{EeSN} \quad (233)$$

$$* \left( \ln \frac{EeSN + EeSO_{x_{t0}} eSO_{\bar{x}}(2000)}{EeSN} \right)^{-1}$$

867 The direct forcing by sulphate aerosols is proportional to the anthropogenic sulphur  
868 emissions (GgS yr<sup>-1</sup>) divided by the sulphate emissions from 2000. The indirect forcing by  
869 sulphate aerosols is a function of the anthropogenic and natural sulphur emissions.

870 Natural sulphur emissions, denoted by EeSN, ~~is-are~~ equal to 42000 Gg<sub>S</sub>. A time series of  
871 annual mean volcanic stratospheric aerosol forcing (W m<sup>-2</sup>) is supplied from  
872 Meinshausen et al. (2011b) and ~~is~~ added to the indirect and direct forcing for a total  
873 sulphate forcing.

#### 4.2.65 ~~N<sub>2</sub>O~~ and Methane (CH<sub>4</sub>)CH<sub>4</sub>

The change in [CH<sub>4</sub>] is calculated directly from CH<sub>4</sub> emissions, and sinks of CH<sub>4</sub> in the the troposphere (based on the lifetime of OH), stratosphere, and soil based on Wigley et al. (2002).

$$\Delta CH_4 = \frac{E(CH_4)}{2.78} - \frac{[CH_4]}{T_{OH}} - \frac{[CH_4]}{T_{strat}} - \frac{[CH_4]}{T_{soil}} \quad (24)$$

where E is total CH<sub>4</sub> emissions (Tg yr<sup>-1</sup>) from both natural and anthropogenic sources, 2.78 (Tg ppb<sup>-1</sup>) is the conversion factor, and T are the lifetimes of the tropospheric sink (T<sub>OH</sub>), the stratospheric sink (T<sub>strat</sub> = 120 year), and the soil sink (T<sub>soil</sub> = 160 year). Note, that within Hector, natural emissions are held at a constant 300 Tg yr<sup>-1</sup>.

The lifetime of OH is a function of [CH<sub>4</sub>], and the emissions of NO<sub>x</sub>, CO and VOC, based on Tanaka et al. (2007a).

$$\begin{aligned} \ln(OH)_t = & -0.32 (\ln[CH_4]_t - \ln[CH_4]_{t_0}) + 0.0042 (E(NO_x)_t \\ & - (E(NO_x)_{t_0}) - 0.000105 (E(CO)_t - (E(CO)_{t_0})) \\ & - 0.00315 (E(VOC)_t - (E(VOC)_{t_0})) \end{aligned} \quad (25)$$

The radiative forcing equation for CH<sub>4</sub> (Joos et al., 2001a) is a function of the concentrations (ppbv) of both CH<sub>4</sub> and N<sub>2</sub>O:

$$\begin{aligned} RF_{CH_4} = & 0.036 \text{ Wm}^{-2} \left[ \sqrt{[CH_4](t)} - \sqrt{[CH_4](t_0)} \right] \\ & - f[CH_4(t), N_2O(t_0)] - f[CH_4(t_0), N_2O(t_0)] \end{aligned} \quad (26)$$

The function *f* accounts for the overlap in CH<sub>4</sub> and N<sub>2</sub>O in their bands is:



$$f(M, N) = 0.47 \quad (27)$$

$$\begin{aligned} & * \ln(1 + (2.01 * 10^{-5}) * (MN)^{0.75} + (5.31 * 10^{-15}) * M \\ & * (MN)^{1.52}) \end{aligned}$$

887

#### 4.2.7 N<sub>2</sub>O

888

The change in [N<sub>2</sub>O] is a function of N<sub>2</sub>O emissions, and the lifetime of N<sub>2</sub>O based on

889

Ward and Mahowald (2014).

$$\Delta N_{2O} = \frac{E(N_{2O})}{4.8} - \frac{[N_{2O}]}{T_{N_{2O}}} \quad (28)$$

890

where E is total N<sub>2</sub>O emissions (Tg N yr<sup>-1</sup>), both natural and anthropogenic, 4.8 (Tg N

891

ppbv<sup>-1</sup>) is the conversion factor, and T<sub>N<sub>2</sub>O</sub> is the lifetime of N<sub>2</sub>O. We set natural

892

emissions of N<sub>2</sub>O to linearly decrease from 11 Tg N yr<sup>-1</sup> in 1765, to 8 Tg N yr<sup>-1</sup> in 2000

893

and are then held constant at 8 Tg N yr<sup>-1</sup> to 2300. The lifetime of N<sub>2</sub>O is a function of its

894

initial lifetime (T<sub>0</sub>) and concentration ([N<sub>2</sub>O]<sub>t0</sub>).

$$T_{N_{2O}} = T_0 * \left( \frac{[N_{2O}]_t}{[N_{2O}]_{t0}} \right)^{-0.05} \quad (29)$$

895

The radiative forcing equation for N<sub>2</sub>O (Joos et al., 2001a) ~~are-is~~ a function of the

896

concentrations (ppbv) of both CH<sub>4</sub> and N<sub>2</sub>O and their radiative efficiency:

(24)

$$RF_{N_{2O}} = 0.12 \text{ Wm}^{-2} [\sqrt{[N_{2O}]_t} - \sqrt{[N_{2O}]_{t0}}] - f[CH_4(t_0), N_{2O}(t)] \quad (30)$$

$$- f[CH_4(t_0), N_{2O}(t_0)]$$

(25)

897 The function  $f$  accounts for the overlap in CH<sub>4</sub> and N<sub>2</sub>O in their bands is the same as  
 898 equation 27:

$$f(M, N) = 0.47 \tag{26}$$

$$* \ln(1 + (2.01 * 10^{-5}) * (MN)^{0.75} + (5.31 * 10^{-15}) * M$$

$$* (MN)^{1.52})$$

899 ~~Note, we are not explicitly calculating concentrations of CH<sub>4</sub> and N<sub>2</sub>O within Hector,~~  
 900 ~~instead we have input files of concentrations.~~

#### 4.2.86 Stratospheric H<sub>2</sub>O from CH<sub>4</sub> oxidation:

902 The radiative forcing from stratospheric H<sub>2</sub>O is a function of the [CH<sub>4</sub>]~~concentrations~~  
 903 (Tanaka et al., 2007a). The coefficient 0.05 is from Joos et al. (2001a) based on the fact  
 904 that the forcing contribution from stratospheric H<sub>2</sub>O is about 5% of the total CH<sub>4</sub> forcing  
 905 (IPCC, 2001). ~~—~~ The 0.036 value of the coefficient corresponds to the same ~~coefficient~~  
 906 value used in the CH<sub>4</sub> radiative forcing equation.

$$RF_{stratH2O} = 0.05 * \left\{ 0.036 \text{ Wm}^{-2} * \left( \sqrt{[CH_4]_t} - \sqrt{[CH_4]_{t0}} \right) \right\} \tag{2731}$$

907

## 5.0 Model ~~e~~Experiments and ~~D~~ata ~~S~~ources

909 A critical test of Hector's performance is to compare the major climatic variables  
 910 calculated in Hector, e.g., atmospheric [CO<sub>2</sub>], radiative forcing, and atmospheric  
 911 temperature, to observational records and ~~other models~~both simple and complex  
 912 climate models. Within this study, Hector is run~~We run Hector~~ under prescribed  
 913 emissions under historical conditions from 1850-2005 to 2300 and then under for all  
 914 four Representative Concentration Pathways (RCPs), ~~out to 2300~~ freely available at

915 <http://tntcat.iiasa.ac.at/RcpDb/dsd?Action=htmlpage&page=about> (Moss et al., 2010;  
916 van Vuuren et al., 2007; Clarke et al., 2007; Wise et al., 2009; Riahi et al., 2007; Fujino et  
917 al., 2006; Hijioka et al., 2008; Smith and Wigley, 2006). ~~(Moss et al., 2010).~~ The RCPs are  
918 plausible future scenarios that ~~are~~ were developed to improve our understanding of the  
919 coupled human climate system. ~~—~~ RCPs by definition are concentration pathways;  
920 however, for all experiments within this manuscript we use the corresponding emissions  
921 trajectories from each RCP as input for Hector. All necessary emission and concentration  
922 inputs are from the four RCPs (RCP 2.6, RCP 4.5, RCP 6.0 and RCP 8.5) freely available at  
923 (Meinshausen et al., 2011b; Riahi et al., 2011; van Vuuren et al., 2011a; van Vuuren et  
924 al., 2011b; Masui et al., 2011; Thomson et al., 2011; van Vuuren et al., 2011d).

925 Comparison data ~~is~~ was obtained from a series of models. We compared Hector  
926 results to MAGICC, a SCM widely used in the scientific and IAM communities, for global  
927 variables such as atmospheric CO<sub>2</sub>, radiative forcing, and temperature (e.g., Raper et al.,  
928 2001; Wigley, 1995; Meinshausen et al., 2011a). ~~—~~ We also compare Hector to a suite of  
929 eleven Earth System Models included in the 5<sup>th</sup> Coupled Model Intercomparison Project  
930 (CMIP5) archive (Taylor et al., 2012). **(Table 3)**. All CMIP5 data ~~are~~ were converted to  
931 yearly global averages from the historical period through the RCPs and their extensions.  
932 One standard deviation of the annual global averages and the CMIP5 model ~~spread~~  
933 range is were calculated for each variable using the RCMIP5  
934 (<http://github.com/JGCRI/RCMIP5>) package in R. All CMIP5 variables used in this study  
935 are from model runs with prescribed atmospheric concentrations, except for  
936 comparisons involving atmospheric [CO<sub>2</sub>] which are from the emissions driven scenario

937 | (esmHistorical and esmRCP8.5) (Figures 3 and 5). We acknowledge that this is not a  
938 | perfect comparison, between an emissions-forced model (Hector) versus, and  
939 | concentration-forced models (CMIP5), is not perfect. However, very there is a  
940 | significant lack of few CMIP5 models that are were run under prescribed emissions  
941 | scenarios available. . The models that run esmRCP8.5 are typically earth system  
942 | models used to investigate the carbon cycle in further detail.  
943 | WLastly, we compare Hector to observations of atmospheric [CO<sub>2</sub>]  
944 | concentrations from Law Dome (1010-1975) and Mauna Loa (1958 – 2008), (Keeling and  
945 | Whorf, 2005; Etheridge et al., 1996) . Global temperature anomalies are from  
946 | HadCRUT4 (Morice et al., 2012). Observations of air-sea and air-land fluxes are from the  
947 | Global Carbon Project (GCP) (Le Quéré et al., 2013). Lastly, observations of surface  
948 | ocean pH are from Bermuda Atlantic Time Series (BATS) and Hawaii Ocean Time Series  
949 | (HOTS) (Bates, 2007; Fujieki et al., 2013).

950

## 951 | **6.0 Results and Discussion**

### 952 | **6.1 Historical**

953 | A critical test of Hector’s performance is how well it compares to historical and  
954 | present day climate from observations, MAGICC, and a suite of CMIP5 models. . –Delta  
955 | changes and root mean square errors were calculated We carried out a few statistical  
956 | tests on Hector (e.g., correlation and root mean square error) for all Hector’s primary  
957 | outputs variables, which are summarized in **Table 4**. After spinup is complete in Hector,  
958 | the atmospheric [CO<sub>2</sub>] in 1850 is 286.0 ppmv, which compares well with observations

959 from Law Dome of 285.2 ppmv. ~~Compared to observations, MAGICC6, and CMIP5 data~~  
960 ~~from 1850 to 2005,~~ Hector captures the global trends in atmospheric [CO<sub>2</sub>] (**Figure 3**)  
961 with ~~correlation coefficients of  $R > 0.99$  and~~ an average root mean square error (RMSE)  
962 of 2.85 ppmv (**Table 4a**), ~~when compared to observations, MAGICC6, and CMIP5 data~~  
963 ~~from 1850-2005.~~ Delta change of atmospheric [CO<sub>2</sub>] from 1850-2005 is slightly lower  
964 than the observations, MAGICC6, and CMIP5. —Hector ~~can be forced~~has the ability to  
965 match atmospheric [CO<sub>2</sub>] records (section 2.4), but we disabled this feature to highlight  
966 the full performance of the model. Note however, that in the MAGICC6 results a similar  
967 feature was used to force the output to match the historical atmospheric [CO<sub>2</sub>] record.

968         Historical global atmospheric temperature anomalies (relative to 1850) are  
969 compared across Hector, MAGICC6, CMIP5, and observations from HadCRUT4 (**Figure**  
970 **4**). ~~A Hector is running without the effects of volcanic forcing, leading to the smoother~~  
971 ~~representation of temperature with time.~~ Atmospheric temperature change from  
972 Hector (0.98 °C) over the period 1850 to 2005 4 closely match the CMIP5 temperature  
973 change (1.01 °C), both slightly higher than the observational record. Over this time  
974 period a Hector has an is well correlated to ( $> 0.8$ ) to observations and models with an  
975 average RMSE of 0.124 °C. — Note that ~~With a simple climate models like Hector, we~~  
976 ~~not intend~~ do not aim to capture temperature variations in temperature due to  
977 interannual/decadal variability found in either in ~~ESMs or the real world;~~ —W instead  
978 they are interested in simulating the overall trends in global mean temperature  
979 change.

980

981 **6.2 Future Projections**

982 Hector’s strengths lie within policy relevant time scales of decades to centuries,  
983 and here we compare Hector to MAGICC and CMIP5 under differing future climate  
984 projections. Results from all four RCPs are broadly similar when comparing Hector, to  
985 MAGICC6, and CMIP5; we display here RCP8.5 results as representative. Within the  
986 modeling community, models that best simulate the historical and present day climate  
987 are assumed to be credible under future projections. We are confident in Hector’s  
988 ability to reproduce historical trends and are therefore confident in its ability to  
989 simulate future climate changes. We compare Hector to MAGICC and CMIP5 under  
990 differing future climate projections. Studies suggest that 80% of the anthropogenic CO<sub>2</sub>  
991 emissions have an average atmospheric lifetime of 300-450 years (Archer et al., 1997;  
992 Rogner, 1997; Archer, 2005). Hector has all the necessary components to model the  
993 climate system from present day through the next approximately 300 years. other  
994 processes become important: , dominant longer term cycle a process

995 **Figure 5** highlights historical trends in atmospheric [CO<sub>2</sub>], along with projections  
996 of atmospheric [CO<sub>2</sub>] under esmRCP8.5 from 1850 to 2100. Note that the emissions  
997 forced scenario only extends to 2100 and not to 2300 like the concentration forced  
998 scenarios (e.g., Figure 8). Both Hector and MAGICC6 are on the low end of the CMIP5  
999 median, but fall within one standard deviation and model range, with a RMSE of 9.0  
1000 ppmv is perfectly correlated with MAGICC and CMIP5 over this period and with a RMSE  
1001 of 9.2 ppmv (Table 4b). Hector and MAGICC6 diverge from the CMIP5 median most  
1002 notably after 2050, but are both still within the low end of the CMIP5 model spread.

1003 The CMIP5 archive does not provide emissions prescribed scenarios for all RCPs;  
1004 we can only compare ~~Figure 6 compares~~ atmospheric [CO<sub>2</sub>] from Hector ~~and with~~  
1005 MAGICC6 under all four RCP scenarios out to 2300 (Figure 6). Hector's change in [CO<sub>2</sub>]  
1006 (1472.13 ppmv) from 1850 to 2300 is slightly lower than MAGICC6 (1600.0 ppmv) for  
1007 RCP 8.5. This is most likely due to different representations of the global carbon cycle.  
1008 ~~Hector is well correlated with MAGICC6 from 1850 out to 2300 for the four RCPs. Under~~  
1009 ~~all of the scenarios except for RCP 8.5, atmospheric [CO<sub>2</sub>] within Hector fluctuates~~  
1010 ~~around the MAGICC6 atmospheric [CO<sub>2</sub>] values, with the most notable fluctuations~~  
1011 ~~under low carbon emissions. This is due to changes in the flux of carbon over the land~~  
1012 ~~as net primary production and respiration change with CO<sub>2</sub> fertilization and temperature~~  
1013 ~~effects.~~

1014 We compare Hector to MAGICC6 for changes in radiative forcing under the four  
1015 RCPs **(Figure 7)**. Radiative forcing ~~wasis~~ not ~~an output provided from within~~ the CMIP5  
1016 ~~models archive~~ and therefore we can only compare Hector and MAGICC6. ~~Hector is~~  
1017 ~~offset slightly lower compared to MAGICC6, which is expected since atmospheric [CO<sub>2</sub>]~~  
1018 ~~is slightly lower.~~ Over the period 1850 to 2300 Hector (12.80 Wm<sup>-2</sup>) and MAGICC6  
1019 (12.24 Wm<sup>-2</sup>) are comparable in their change in radiative forcing, ~~is well correlated (1.0)~~  
1020 ~~with MAGICC6~~ with a RMSE of 0.265 W m<sup>-2</sup>. One noticeable difference between  
1021 MAGICC6 and Hector during the historical period is the decreases in radiative forcing.  
1022 This is due to ~~We acknowledge that t~~The correlation is lower under in the historical  
1023 ~~period (0.79), because as noted above. This s~~are may be due to slight differences in the  
1024 ~~representation of atmospheric gases, pollutants, and aerosols between the two models.~~

1025 the effects of volcanic emissions on radiative forcing. For simplicity, we have chosen to  
1026 run Hector without these effects.

1027 **Figure 8** compares global temperature anomalies from Hector to MAGICC6 and  
1028 CMIP5 over the four RCPs, from 2005 to 2300. Hector simulates the- CMIP5 median  
1029 more closely than MAGICC6 across all four RCPs, with a temperature change under RCP  
1030 8.5 for Hector of 8.59 °C, compared to MAGICC6 of 7.30 °C, while the temperature  
1031 change for CMIP5 is 9.57 °C (Table 4c)and MAGICC6 are comparable in their  
1032 temperature change across the four RCPs. However, both are lower than the CMIP5  
1033 median under RCP 2.6, 4.5 and 8.5, with the largest discrepancy under high CO<sub>2</sub>  
1034 emissions in RCP 8.5-. To highlight this close comparison, temperature change over the  
1035 entire record (1850-2300) for Hector is 9.58 °C, which is within 1.0 °C of the CMIP5  
1036 median, while MAGGIC6's temperature change is greater than 2.5 °C away from the  
1037 CMIP5 median. the median does 66 is1 Regardless, Hector is still highly correlated  
1038 ( $\rightarrow 0.97$ ) to MAGICC6 and CMIP5 for RCP 8.5, with a RMSE of 0.52 °C compared to CMIP5  
1039 (Table 4c). The fluctuations seen in RCP 2.6 within atmospheric [CO<sub>2</sub>] are also  
1040 apparent in the atmospheric temperature trends. However, the general trends of  
1041 temperature change, peaking around 2050 and then slowly declining out to 2300 are  
1042 captured within Hector.

1043 (Cheng et al., 2013)

1044 Another way to visualize model performance is a Taylor diagram (Figure 9)  
1045 compactly summarizes model performance simulating of global temperature change  
1046 relative to 1850, from 1850 to 2300 for RCP 8.5. The In this figure, the closer the points



1047 ~~are to the reference point (Hector) the higher the correlation and low RMSE between~~  
1048 ~~CMIP5 models and MAGICC6. Those points with a standard deviation similar to that of~~  
1049 ~~Hector experience the same amplitude of temperature change over this time period~~  
1050 ~~(MAGICC6). All of the models are highly correlated with Hector, with a large range in~~  
1051 ~~the their standard deviations (1–5 °C).~~

1052 **Figures 910** and **110** present a detailed view of carbon fluxes under RCP 8.5, for  
1053 CMIP5 and observations (negative represents carbon flux to the atmosphere). The  
1054 ocean is a major sink of carbon through 2100, becoming less effective with time in both  
1055 Hector and the CMIP5 models. MAGICC6 does not include air-sea fluxes in its output,  
1056 and because it is not open source we were unable to obtain these values. Therefore, we  
1057 compare air-sea fluxes of CO<sub>2</sub> to MAGICC5.3, ~~the version currently used in the IA model,~~  
1058 ~~Global Change Assessment Model (Calvin et al., 2011),~~ updated with explicit BC and OC  
1059 forcing as described in Smith and Bond (2014). Hector's calculation of air-sea fluxes is  
1060 within the large CMIP5 model range up to 2100. However, after that Hector peaks close  
1061 to 2150, while the CMIP5 models are beginning to decline. The correlation is high  
1062 between Hector and CMIP5 over the historical period (0.95). However, the correlation,  
1063 but drops off significantly between 2005 and 2300 (0.10) (Table 4c). This is an active  
1064 area of research, ~~investigating the differences between Hector and CMIP5 after 2100.~~  
1065 One potential reason for the discrepancy low correlation after 2100 ~~could be due to~~  
1066 ~~the fact that we are only comparing to three models that run the RCP extension to 2300~~  
1067 ~~(bcc-csm1-1, IPSL-CM5A-LR, and MPI-ESM-LR).~~ With more models included after 2100a  
1068 larger spread of fluxes, Hector may be better correlated. is that in this version of

1069 Hector, we do not simulate changes in ocean circulation, potentially biasing fluxes too  
1070 high after 2100. Most ESMs in CMIP5 show a weakening of the Atlantic meridional  
1071 overturning circulation by 2100 between 15% and 60% under RCP 8.5 (Cheng et al.,  
1072 2013). A slowdown in ocean circulation may result in less carbon uptake by the oceans.  
1073 . Another potential reason for this bias is Hector’s constant pole to equator ocean  
1074 temperature gradient. Studies show that the Arctic is warming faster than the rest of the  
1075 globe (e.g., Bintanja and van der Linden, 2013; Holland and Bitz, 2003; Bekryaev et al.,  
1076 2010). A warmer high latitude surface ocean in Hector would suppress the uptake of  
1077 carbon, potentially bringing the air-sea fluxes closer to the CMIP5 median after  
1078 2100.~~The average correlation over the CMIP5 models over 1850–2300 is higher at 0.80,~~  
1079 ~~with a RMSE of 1.45 PgC yr<sup>-1</sup> (Table 4b).~~  
1080 ~~The land fluxes have a large range of uncertainty into the future within the~~  
1081 ~~CMIP5 models. Hector follows the general trends of the land acting as a sink of carbon~~  
1082 ~~initially with a gradual switch to a carbon source after 2150. Fluxes of carbon over the~~  
1083 ~~land are less well correlated to the CMIP5 median compared to the air-sea fluxes, 0.55~~  
1084 ~~(historical) and 0.65 (RCP 8.5), but it is important to note that CMIP models tend to~~  
1085 ~~show huge divergences in their land responses to changing climate~~ (e.g., Friedlingstein  
1086 et al., 2006), which is evident by the large range in CMIP5 models (Figure 10). Hector  
1087 simulates the general trends, of increasing carbon sink and then a gradual decline to a  
1088 carbon source after 2100. Both land and ocean fluxes within Hector agree well the  
1089 observations from Le Quere-Queré et al., (2013) and other sources.

1090 ~~One feature in Hector that is unique amongst SCMs~~ Lastly, a unique feature of  
1091 ~~Hector~~ is its ability to actively solve the carbonate system in the upper ocean (Hartin et  
1092 al, in prep). This feature allows us to predict changes ocean acidification, calcium  
1093 carbonate saturations and other ~~parameters of the~~ carbonate system parameters.  
1094 **Figure 112** shows low latitude (<55°) pH for Hector compared to CMIP5 and  
1095 observations from 1850 to 2100 under RCP 8.5. ~~We~~ The see estimate model projects a  
1096 significant drop in pH from present day through 2100, which may lead to detrimental  
1097 effects on marine ecosystems (e.g., Fabry et al., 2008).

1098

## 1099 **7.0 Conclusions**

1100 Hector reproduces the large-scale couplings and feedbacks on the climate  
1101 system between the atmosphere, ocean, and land, ~~generally. Hector falls~~ falling within  
1102 the range of the CMIP5 model ~~spread~~ and ~~tracks~~ matches well with MAGICC.  
1103 ~~Our goal was not to~~ it does not simulate the fine details or parameterizations typically  
1104 found in large-scale, complex ~~models~~ ESMs, but instead ~~to~~ represents only the most  
1105 critical global processes in a reduced-complexity form. This allows for fast execution  
1106 times, ease of understanding, and straightforward analysis of the model output. ~~To help~~  
1107 ~~with the analysis of Hector we included within the online database of Hector, R scripts~~  
1108 ~~to process Hector's output as well as the comparison data.~~

1109 Two of Hector's ~~two~~ key features are its open source license nature and modular  
1110 design. This allows the user to ~~manipulate~~ edit the input files and code at will, for  
1111 example to enable/disable/replace components, or include components not found

1112 within the core version of Hector. For example, ~~the a~~ user can design a new submodel  
1113 (e.g., sea-ice) to answer specific climate questions relating to that process. Hector is  
1114 hosted on a widely-used open source software repository (Github), and thus changes  
1115 and improvements can be easily shared with the scientific community. Because of these  
1116 critical features, Hector has the potential to be a key analytical tool in both the policy  
1117 and scientific communities. We welcome user input and encourage use, modifications,  
1118 and collaborations with Hector.

1119 While Hector has many strengths, ~~there the current 1.0 version~~ certainly has are  
1120 a few some limitations ~~that later versions of Hector hope to address~~ and weaknesses.  
1121 For example, Hector does not currently simulate terrestrial gross primary production, a  
1122 key metric of comparison to e.g. the FLUXNET database. ~~For example~~ Also, Hector does  
1123 not have differential radiative forcing and atmospheric temperature calculations over  
1124 land and ocean. This is may be a problem, as e-land responds to changes in emissions of  
1125 greenhouse gases, and aerosols much quicker than the ocean, ~~leading to different~~  
1126 ~~temperature responses over the land and ocean (REFERENCE)~~ (Hansen et al., 2005). In  
1127 addition, it does not currently simulate terrestrial gross primary production, a key  
1128 metric of comparison to e.g. the Fluxnet database. ~~Also,~~ Hector does not explicitly deal  
1129 with oceanic heat uptake, except via a simple empirical formula. Surface  
1130 temperatures are calculated based on a linear relationship with atmospheric  
1131 temperature and ~~heat uptake by the ocean is parameterized by a constant heat uptake~~  
1132 ~~efficiency.~~ we assume a constant pole to equator temperature gradient. We  
1133 acknowledge that this assumption may not hold true if the poles warm faster than the

1134 ~~equator. While Hector can reproduce global trends in atmospheric CO<sub>2</sub> and~~  
1135 ~~temperature, we cannot investigate ocean heat uptake in the deep ocean using Hector.~~  
1136 ~~Currently, there is placeholder in Hector for a more sophisticated sea level rise~~  
1137 ~~submodel. The current edition of Hector uses inputs of concentrations of CH<sub>4</sub> and N<sub>2</sub>O~~  
1138 ~~to calculate radiative forcing from CH<sub>4</sub> and N<sub>2</sub>O. Ideally we would like Hector to~~  
1139 ~~calculate concentrations from emissions of CH<sub>4</sub> and N<sub>2</sub>O. This would allow for quick~~  
1140 ~~integration within IAMs.~~

1141 Future plans with Hector include addressing some of the above limitations and  
1142 conducting numerous scientific experiments, using Hector as a stand-alone simple  
1143 climate carbon-cycle model. ~~Also, Hector~~ It is also being ~~will be~~ incorporated into Pacific  
1144 Northwest National Laboratory's Global Change Assessment Model ~~to begin running for~~  
1145 ~~policy policy~~-relevant experiments. — Hector has the ability to be a key analytical tool  
1146 used across many scientific and policy communities due to its modern software  
1147 architecture, open source, and object-oriented structure.

1148

#### 1149 **Code Availability**

1150 Hector is freely available at <https://github.com/JGCRI/hector> . The specific Hector  
1151 ~~v0.1.0~~ referenced in this paper, as well as code to reproduce all figures and results  
1152 shown here, is available at <https://github.com/JGCRI/hector/releases/tag/v0.1.0>

#### 1153 **Author contributions**

1154 C.A.H. and B.P.B.-L. developed the ocean and terrestrial carbon models, respectively,  
1155 and led the overall development of Hector. R.P.L. and P.P. wrote critical code for

1156 Hector’s coupler and carbon cycle solver. A.S. helped with the development of the  
1157 atmospheric forcing components. C.A.H. wrote the manuscript with contributions from  
1158 all co-authors.

1159

1160 **Acknowledgements**

1161 This research is based on work supported by the U.S. Department of Energy, Office of  
1162 Science, Integrated Assessment Research Program. The Pacific Northwest National  
1163 Laboratory is operated for DOE by Battelle Memorial Institute under contract DE-AC05-  
1164 | 76RL01830.

1165 **References:**

- 1166 Anthoff, D., and Tol, R. S. J.: The income elasticity of the impact of climate change, Is  
1167 the Environment a Luxury? An inquiry into the relationship between environment and  
1168 income, edited by: Tiezzi, S., and Martini, C., Routledge, 2014.
- 1169 Applegate, P. J., Kirchner, N., Stone, E. J., Keller, K., and Greve, R.: An assessment of  
1170 key model parametric uncertainties in projections of Greenland Ice Sheet behavior, *The*  
1171 *Cryosphere*, 6, 589-606, 10.5194/tc-6-589-2012, 2012.
- 1172 Archer, D., Kheshgi, H., and Maier-Reimer, E.: Multiple timescales for neutralization of  
1173 fossil fuel CO<sub>2</sub>, *Geophysical Research Letters*, 24, 405-408, 10.1029/97GL00168, 1997.
- 1174 Archer, D.: Fate of fossil fuel CO<sub>2</sub> in geologic time, *Journal of Geophysical Research:*  
1175 *Oceans*, 110, C09S05, 10.1029/2004JC002625, 2005.
- 1176 Bates, N. R.: Interannual variability of the oceanic CO<sub>2</sub> sink in the subtropical gyre of the  
1177 North Atlantic Ocean over the last 2 decades, *Journal of Geophysical Research: Oceans*,  
1178 112, C09013, 10.1029/2006JC003759, 2007.
- 1179 Bekryaev, R. V., Polyakov, I. V., and Alexeev, V. A.: Role of Polar Amplification in  
1180 Long-Term Surface Air Temperature Variations and Modern Arctic Warming, *Journal of*  
1181 *Climate*, 23, 3888-3906, 10.1175/2010JCLI3297.1, 2010.
- 1182 Bintanja, R., and van der Linden, E. C.: The changing seasonal climate in the Arctic, *Sci.*  
1183 *Rep.*, 3,  
1184 [http://www.nature.com/srep/2013/130327/srep01556/abs/srep01556.html#supplementary](http://www.nature.com/srep/2013/130327/srep01556/abs/srep01556.html#supplementary-information)  
1185 [-information](http://www.nature.com/srep/2013/130327/srep01556/abs/srep01556.html#supplementary-information), 2013.
- 1186 Bond-Lamberty, B., Calvin, K., Jones, A. D., Mao, J., Patel, P., Shi, X., Thomson, A.,  
1187 Thornton, P., and Zhou, Y.: Coupling earth system and integrated assessment models: the  
1188 problem of steady state, *Geosci. Model Dev. Discuss.*, 7, 1499-1524, 10.5194/gmdd-7-  
1189 1499-2014, 2014.
- 1190 Bond, T. C., Doherty, S. J., Fahey, D. W., Forster, P. M., Berntsen, T., DeAngelo, B. J.,  
1191 Flanner, M. G., Ghan, S., Kärcher, B., Koch, D., Kinne, S., Kondo, Y., Quinn, P. K.,  
1192 Sarofim, M. C., Schultz, M. G., Schulz, M., Venkataraman, C., Zhang, H., Zhang, S.,  
1193 Bellouin, N., Guttikunda, S. K., Hopke, P. K., Jacobson, M. Z., Kaiser, J. W., Klimont,  
1194 Z., Lohmann, U., Schwarz, J. P., Shindell, D., Storelvmo, T., Warren, S. G., and Zender,  
1195 C. S.: Bounding the role of black carbon in the climate system: A scientific assessment,  
1196 *Journal of Geophysical Research: Atmospheres*, 118, 5380-5552, 10.1002/jgrd.50171,  
1197 2013.
- 1198 Bouwman, A. F., Hoek, K. W. v. d., Drecht, G. V., and Eickhout, B.: World Livestock  
1199 and crop production systems, land use and environment between 1970 and 2030, *Rural*  
1200 *Lands, Africulture and Climate beyond 2015: A new prespective on suture land use*  
1201 *patterns*, edited by: Brouwer, F., and McCarl, B., Springer, Dordrecht, 2006.
- 1202 Calvin, K., Clarke, L., Edmonds, J., Eom, J., Hejazi, M., Kim, S., Kyle, G., Link, R.,  
1203 Patel, P., Smith, S., and Wise, M.: GCAM Wiki Documentation, PNNL-20809, Pacific  
1204 Northwest National Laboratory, Richland WA, 2011.
- 1205 Castruccio, S., McInerney, D. J., Stein, M. L., Crouch, F. L., Jacob, R. L., and Moyer, E.  
1206 J.: Statistical Emulation of Climate Model Projections Based on Precomputed GCM  
1207 Runs, *Journal of Climate*, 27, 2014.

1208 Challenor, P.: Using emulators to estimate uncertainty in complex models, *Uncertainty*  
1209 *Quantification in Scientific Computing*, edited by: Dienstry, A. M., and Boisvert, R. F.,  
1210 Springer, IFIP AICT 377, 151-164 pp., 2012.

1211 Cheng, W., Chiang, J. C. H., and Zhang, D.: Atlantic Meridional Overturning Circulation  
1212 (AMOC) in CMIP5 Models: RCP and Historical Simulations, *Journal of Climate*, 26,  
1213 7187-7197, 10.1175/JCLI-D-12-00496.1, 2013.

1214 Clarke, L., J. Edmonds, H. Jacoby, H. Pitcher, Reilly, J., and Richels, R.: Scenarios of  
1215 Greenhouse Gas Emissions and Atmospheric Concentrations. Sub-report 2.1A of  
1216 Synthesis and Assessment Product 2.1, edited by: Research, U. S. C. C. S. P. a. t. S. o. G.  
1217 C., Department of Energy, Office of Biological & Environmental Research, Washington,  
1218 7 DC., USA, 2007.

1219 Collins, W. D., Craig, A. P., Truesdale, J. E., Di Vittorio, A. V., Jones, A. D., Bond-  
1220 Lamberty, B., Calvin, K. V., Edmonds, J. A., Kim, S. H., Thomson, A. M., Patel, P.,  
1221 Zhou, Y., Mao, J., Shi, X., Thornton, P. E., Chini, L. P., and Hurtt, G. C.: The integrated  
1222 Earth System Model (iESM): formulation and functionality, *Geosci. Model Dev.*  
1223 *Discuss.*, 8, 381-427, 10.5194/gmdd-8-381-2015, 2015.

1224 Denman, K. L., G. Brasseur, A. Chidthaisong, P. Ciais, P.M. Cox, R.E. Dickinson, D.  
1225 Hauglustaine, C. Heinze, E. Holland, D. Jacob, U. Lohmann, S Ramachandran, P.L. da  
1226 Silva Dias, S.C. Wofsy and X. Zhang: *Climat Change 2007: The Physical Science Basis.*,  
1227 edited by: Change, C. o. W. G. I. t. t. F. A. R. o. t. I. P. o. C., Cambridge University  
1228 Press, Cambridge , United Kingdom and New York, USA, 2007.

1229 Di Vittorio, A. V., Chini, L. P., Bond-Lamberty, B., Mao, J., Shi, X., Truesdale, J., Craig,  
1230 A., Calvin, K., Jones, A., Collins, W. D., Edmonds, J., Hurtt, G. C., Thornton, P., and  
1231 Thomson, A.: From land use to land cover: restoring the afforestation signal in a coupled  
1232 integrated assessment–earth system model and the implications for CMIP5 RCP  
1233 simulations, *Biogeosciences*, 11, 6435-6450, 10.5194/bg-11-6435-2014, 2014.

1234 E.P. White, E. B., Z.T. Brym, K.J. Locey, D.J. McGlenn: Nine simple ways to make it  
1235 easier to (re)use your data, *PeerJ PrePrints*, 1:e7v2,  
1236 <http://dx.doi.org/10.7287/peerj.preprints.7v2>, 2013.

1237 Edmonds, J., and Smith, S. J.: *The Technology of Two Degrees. Avoiding Dangerous*  
1238 *Climate Change*, edited by: Schellnhuber, H. J., Cramer, W., Nakicenovic, N., Wigley,  
1239 T., and Yohe, G., Cambridge University Press, Cambridge, UK, 2006.

1240 Ehhalt, D., Prather, M. J., Dentener, F. J., Derwent, R., Dlugokencky, E. J., Holland, E.  
1241 A., Isaksen, I. S., Katima, J., Kirchoff, V., Matson, P. A., and Wang, M.: Atmospheric  
1242 chemistry and greenhouse gases, in: *Climate Change 2001: The Scientific Basis*, edited  
1243 by: Houghton, J. T., Ding, Y., Griggs, D. J., Noguer, M., van der Linden, L., Dai, X.,  
1244 Maskell, K., and Johnson, C. A., Cambridge University Press, Cambridge, UK, 892,  
1245 2001.

1246 Etheridge, D. M., Steele, L. P., Langenfelds, R. L., Francey, R. J., Barnola, J. M., and  
1247 Morgan, V. I.: Natural and anthropogenic changes in atmospheric CO<sub>2</sub> over the last 1000  
1248 years from air in Antarctic ice and firn, *Journal of Geophysical Research: Atmospheres*,  
1249 101, 4115-4128, 10.1029/95JD03410, 1996.

1250 Fabry, V. J., Seibel, B. A., Feely, R. A., and Orr, J. C.: Impacts of ocean acidification on  
1251 marine fauna and ecosystem processes, *ICES Journal of Marine Science: Journal du*  
1252 *Conseil*, 65, 414-432, 10.1093/icesjms/fsn048, 2008.



1253 Friedlingstein, P., Cox, P., Betts, R., Bopp, L., von Bloh, W., Brovkin, V., Cadule, P.,  
 1254 Doney, S., Eby, M., Fung, I., Bala, G., John, J., Jones, C., Joos, F., Kato, T., Kawamiya,  
 1255 M., Knorr, W., Lindsay, K., Matthews, H. D., Raddatz, T., Rayner, P., Reick, C.,  
 1256 Roeckner, E., Schnitzler, K. G., Schnur, R., Strassmann, K., Weaver, A. J., Yoshikawa,  
 1257 C., and Zeng, N.: Climate–Carbon Cycle Feedback Analysis: Results from the C4MIP  
 1258 Model Intercomparison, *Journal of Climate*, 19, 3337-3353, 10.1175/JCLI3800.1, 2006.  
 1259 Friedlingstein, P., Meinshausen, M., Arora, V. K., Jones, C. D., Anav, A., Liddicoat, S.  
 1260 K., and Knutti, R.: Uncertainties in CMIP5 Climate Projections due to Carbon Cycle  
 1261 Feedbacks, *Journal of Climate*, 27, 511-526, 10.1175/JCLI-D-12-00579.1, 2014.  
 1262 Fujieki, L., Santiago-Mandujano, F., Fumar, C., Liukas, R., and Church, M.: Hawaii  
 1263 Ocean Time-series Program Data Report, 2013.  
 1264 Fujino, J., Nair, R., Kainuma, M., Masui, T., and Matsuoka, Y.: Multi-gas mitigation  
 1265 analysis on stabilization scenarios using AIM global model, *Multigas Mitigation and*  
 1266 *Climate Policy*. The Energy Journal, Special Issue, 2006.  
 1267 Hansen, J., Sato, M., Ruedy, R., Nazarenko, L., Lacis, A., Schmidt, G. A., Russell, G.,  
 1268 Aleinov, I., Bauer, M., Bauer, S., Bell, N., Cairns, B., Canuto, V., Chandler, M., Cheng,  
 1269 Y., Del Genio, A., Faluvegi, G., Fleming, E., Friend, A., Hall, T., Jackman, C., Kelley,  
 1270 M., Kiang, N., Koch, D., Lean, J., Lerner, J., Lo, K., Menon, S., Miller, R., Minnis, P.,  
 1271 Novakov, T., Oinas, V., Perlwitz, J., Perlwitz, J., Rind, D., Romanou, A., Shindell, D.,  
 1272 Stone, P., Sun, S., Tausnev, N., Thresher, D., Wielicki, B., Wong, T., Yao, M., and  
 1273 Zhang, S.: Efficacy of climate forcings, *Journal of Geophysical Research: Atmospheres*,  
 1274 110, D18104, 10.1029/2005JD005776, 2005.  
 1275 Hartin, C. A., Bond-Lamberty, B., and Patel, P.: Projections of ocean acidification over  
 1276 three centuries using a carbonate chemistry box model, *Biogeosciences*, in prep.  
 1277 Harvey, L. D. D., and Schneider, S. H.: Transient climate response to external forcing on  
 1278 100–104 year time scales part 1: Experiments with globally averaged, coupled,  
 1279 atmosphere and ocean energy balance models, *Journal of Geophysical Research:*  
 1280 *Atmospheres*, 90, 2191-2205, 10.1029/JD090iD01p02191, 1985.  
 1281 Heron, M., Hanson, V., and Ricketts, I.: Open source and accessibility: advantages and  
 1282 limitations, *Journal of Interaction Science*, 1, 2, 10.1186/2194-0827-1-2, 2013.  
 1283 Hijioka, Y., Matsuoka, Y., Nishimoto, H., Masui, M., and Kainuma, M.: Global GHG  
 1284 emissions scenarios under GHG concentration stabilization targets, *Journal of*  
 1285 *Environmental Engineering*, 13, 97-108, 2008.  
 1286 Hoffert, M. I., Callegari, A. J., and Hsieh, C.-T.: The Role of Deep Sea Heat Storage in  
 1287 the Secular Response to Climatic Forcing, *J. Geophys. Res.*, 85, 6667-6679,  
 1288 10.1029/JC085iC11p06667, 1980.  
 1289 Holland, M. M., and Bitz, C. M.: Polar amplification of climate change in coupled  
 1290 models, *Climate Dynamics*, 21, 221-232, 10.1007/s00382-003-0332-6, 2003.  
 1291 Ince, D. C., Hatton, L., and Graham-Cumming, J.: The case for open computer programs,  
 1292 *Nature*, 482, 485-488, 2012.  
 1293 IPCC: *Climate Change 2001: The Science of Climate Change*. Contribution of Working  
 1294 Group I to the Second Assessment Report of the Intergovernmental Panel on Climate  
 1295 Change, Cambridge University Press, Cambridge, 2001.  
 1296 Irvine, P. J., Sriviver, R. L., and Keller, K.: Tension between reducing sea-level rise and  
 1297 global warming through solar-radiation management, *Nature Clim. Change*, 2, 97-100,

1298 <http://www.nature.com/nclimate/journal/v2/n2/abs/nclimate1351.html#supplementary->  
1299 [information](http://www.nature.com/nclimate/journal/v2/n2/abs/nclimate1351.html#supplementary-), 2012.

1300 Joos, F., Prentice, I. C., Sitch, S., Meyer, R., Hooss, G., Plattner, G.-K., Gerber, S., and  
1301 Hasselmann, K.: Global warming feedbacks on terrestrial carbon uptake under the  
1302 Intergovernmental Panel on Climate Change (IPCC) Emission Scenarios, *Global*  
1303 *Biogeochemical Cycles*, 15, 891-907, 10.1029/2000GB001375, 2001a.

1304 Joos, F., Prentice, I. C., Sitch, S., Meyer, R., Hooss, G., Plattner, G.-K., Gerber, S., and  
1305 Hasselmann, K.: Global warming feedbacks on terrestrial carbon uptake under the  
1306 Intergovernmental Panel on Climate Change (IPCC) emission scenarios, *Global*  
1307 *Biochemical Cycles*, 15, 891-907, 2001b.

1308 Knox, F., and McElroy, M. B.: Changes in Atmospheric CO<sub>2</sub>: Influence of the Marine  
1309 Biota at High Latitude, *J. Geophys. Res.*, 89, 4629-4637, 10.1029/JD089iD03p04629,  
1310 1984.

1311 Knutti, R., and Hegerl, G. C.: The equilibrium sensitivity of the Earth's temperature to  
1312 radiation changes, *Nature Geosci*, 1, 735-743, 2008.

1313 Le Quéré, C., Andres, R. J., Boden, T., Conway, T., Houghton, R. A., House, J. I.,  
1314 Marland, G., Peters, G. P., van der Werf, G. R., Ahlström, A., Andrew, R. M., Bopp, L.,  
1315 Canadell, J. G., Ciais, P., Doney, S. C., Enright, C., Friedlingstein, P., Huntingford, C.,  
1316 Jain, A. K., Jourdain, C., Kato, E., Keeling, R. F., Klein Goldewijk, K., Levis, S., Levy,  
1317 P., Lomas, M., Poulter, B., Raupach, M. R., Schwinger, J., Sitch, S., Stocker, B. D.,  
1318 Viovy, N., Zaehle, S., and Zeng, N.: The global carbon budget 1959–2011, *Earth Syst.*  
1319 *Sci. Data*, 5, 165-185, 10.5194/essd-5-165-2013, 2013.

1320 Lenton, T. M.: Land and ocean carbon cycle feedback effects on global warming in a  
1321 simple Earth system model, *Tellus B*, 52, 1159-1188, 10.1034/j.1600-0889.2000.01104.x,  
1322 2000.

1323 Lenton, T. M., Myerscough, R. J., Marsh, R., Livina, V. N., Price, A. R., and Cox, S. J.:  
1324 Using GENIE to study a tipping point in the climate system, 1890, 871-884 pp., 2009.

1325 Manne, A. S., and Richels, R. G.: Merge: an integrated assessment model for global  
1326 climate change, *Energy and environment*, edited by: Loulou, R., Waub, J.-P., and  
1327 Zaccour, G., Springer, New York, 2005.

1328 Martin, R. C., Riehle, D., and Buschmann, F.: *Pattern Languages of Program Design 3*,  
1329 Addison-Wesley, Boston, MA, 672 pp., 1997.

1330 Meinshausen, M., Raper, S. C. B., and Wigley, T. M. L.: Emulating coupled atmosphere-  
1331 ocean and carbon cycle models with a simpler model, *MAGICC6 – Part 1: Model*  
1332 *description and calibration*, *Atmos. Chem. Phys.*, 11, 1417-1456, 10.5194/acp-11-1417-  
1333 2011, 2011a.

1334 Meinshausen, M., Smith, S. J., Calvin, K., Daniel, J. S., Kainuma, M. L. T., Lamarque, J.  
1335 F., Matsumoto, K., Montzka, S. A., Raper, S. C. B., Riahi, K., Thomson, A., Velders, G.  
1336 J. M., and Vuuren, D. P. P.: The RCP greenhouse gas concentrations and their extensions  
1337 from 1765 to 2300, *Climatic Change*, 109, 213-241, 10.1007/s10584-011-0156-z, 2011b.

1338 Meinshausen, M., Wigley, T. M. L., and Raper, S. C. B.: Emulating atmosphere-ocean  
1339 and carbon cycle models with a simpler model, *MAGICC6 – Part 2: Applications*,  
1340 *Atmos. Chem. Phys.*, 11, 1457-1471, 10.5194/acp-11-1457-2011, 2011c.

1341 Morice, C. P., Kennedy, J. J., Rayner, N. A., and Jones, P. D.: Quantifying uncertainties  
1342 in global and regional temperature change using an ensemble of observational estimates:

1343 The HadCRUT4 data set, *Journal of Geophysical Research: Atmospheres*, 117, D08101,  
1344 10.1029/2011JD017187, 2012.

1345 Moss, R. H., Edmonds, J. A., Hibbard, K. A., Manning, M. R., Rose, S. K., van Vuuren,  
1346 D. P., Carter, T. R., Emori, S., Kainuma, M., Kram, T., Meehl, G. A., Mitchell, J. F. B.,  
1347 Nakicenovic, N., Riahi, K., Smith, S. J., Stouffer, R. J., Thomson, A. M., Weyant, J. P.,  
1348 and Wilbanks, T. J.: The next generation of scenarios for climate change research and  
1349 assessment, *Nature*, 463, 747-756,  
1350 [http://www.nature.com/nature/journal/v463/n7282/supinfo/nature08823\\_S1.html](http://www.nature.com/nature/journal/v463/n7282/supinfo/nature08823_S1.html), 2010.

1351 Murakami, K., Sasai, T., and Yamaguchi, Y.: A new one-dimensional simple energy  
1352 balance and carbon cycle coupled model for global warming simulation, *Theoretical and  
1353 Applied Climatology*, 101, 459-473, 10.1007/s00704-009-0232-8, 2010.

1354 Myhre, G., Highwood, E. J., Shine, K. P., and Stordal, F.: New estimates of radiative  
1355 forcing due to well mixed greenhouse gases, *Geophysical Research Letters*, 25, 2715-  
1356 2718, 10.1029/98GL01908, 1998.

1357 Nemani, R. R., Keeling, C. D., Hashimoto, H., Jolly, W. M., Piper, S. C., Tucker, C. J.,  
1358 Myneni, R. B., and Running, S. W.: Climate-Driven Increases in Global Terrestrial Net  
1359 Primary Production from 1982 to 1999, *Science*, 300, 1560-1563,  
1360 10.1126/science.1082750, 2003.

1361 Nordhaus, W. D.: A question of balance weighing the options on global warming  
1362 policies, Yale University Press, New Haven, 2008.

1363 Piao, S., Sitch, S., Ciais, P., Friedlingstein, P., Peylin, P., Wang, X., Ahlström, A., Anav,  
1364 A., Canadell, J. G., Cong, N., Huntingford, C., Jung, M., Levis, S., Levy, P. E., Li, J.,  
1365 Lin, X., Lomas, M. R., Lu, M., Luo, Y., Ma, Y., Myneni, R. B., Poulter, B., Sun, Z.,  
1366 Wang, T., Viovy, N., Zaehle, S., and Zeng, N.: Evaluation of terrestrial carbon cycle  
1367 models for their response to climate variability and to CO<sub>2</sub> trends, *Global Change  
1368 Biology*, 19, 2117-2132, 10.1111/gcb.12187, 2013.

1369 Pietsch, S. A., and Hasenauer, H.: Evaluating the self-initialization procedure for large-  
1370 scale ecosystem models, *Global Change Biology*, 12, 1-12, 10.1111/j.1365-  
1371 2486.2006.01211.x 2006.

1372 Raper, S. C., Gregory, J. M., and Stouffer, R. J.: The Role of Climate Sensitivity and  
1373 Ocean Heat Uptake on AOGCM Transient Temperature Response, *Journal of Climate*,  
1374 15, 124-130, 2002.

1375 Raper, S. C. B., Gregory, J. M., and Osborn, T. J.: Use of an upwelling-diffusion energy  
1376 balance climate model to simulate and diagnose A/OGCM results, *Climate Dynamics*, 17,  
1377 601-613, 2001.

1378 Ratto, M., Castelletti, A., and Pagano, A.: Emulation techniques for the reduction and  
1379 sensitivity analysis of complex environmental models, *Environmental Modelling and  
1380 Software*, 34, 1-4, 2012.

1381 Riahi, K., Grubler, A., and Nakicenovic, N.: Scenarios of long-term socio-economic and  
1382 environmental development under climate stabilization, *Technological Forecasting and  
1383 Social Change*, 74, 887-935, 2007.

1384 Ricciuto, D. M., Davis, K. J., and Keller, K.: A Bayesian calibration of a simple carbon  
1385 cycle model: The role of observations in estimating and reducing uncertainty, *Global  
1386 Biogeochemical Cycles*, 22, GB2030, 10.1029/2006GB002908, 2008.

1387 Rogner, H. H.: An assessment of world hydrocarbon resources, *Annual Review of  
1388 Energy and the Environment*, 22, 217-262, 10.1146/annurev.energy.22.1.217, 1997.

1389 Schlesinger, M. E., and Jiang, X.: Simple Model Representation of Atmosphere-Ocean  
 1390 GCMs and Estimation of the Time Scale of CO<sub>2</sub>-Induced Climate Change, *Journal of*  
 1391 *Climate*, 3, 1297-1315, 10.1175/1520-0442(1990)003<1297:SMROAO>2.0.CO;2, 1990.  
 1392 Senior, C. A., and Mitchell, J. F. B.: The time-dependence of climate sensitivity,  
 1393 *Geophysical Research Letters*, 27, 2685-2688, 10.1029/2000GL011373, 2000.  
 1394 Smith, S., and Wigley, T.: Multi-Gas Forcing Stabilization with the MiniCAM, *Energy*  
 1395 *Journal Special Issue #3*, 373-391, 2006.  
 1396 Smith, S. J., and Bond, T. C.: Two hundred fifty years of aerosols and climate: the end of  
 1397 the age of aerosols, *Atmos. Chem. Phys. Discuss.*, 13, 6419-6453, 10.5194/acp-14-537-  
 1398 2014, 2014.  
 1399 Sokolov, A. P., CA, S., S, D., S, P., DW, K., HD, J., RG, P., CE, F., JM, R., C, W., B, F.,  
 1400 MC, S., J, S., PH, S., M, J., and J, C.: The MIT Integrated Global System Model (IGSM)  
 1401 version 2: model description and baseline evaluation, MIT, Cambridge, 2005.  
 1402 Sriver, R., Urban, N., Olson, R., and Keller, K.: Toward a physically plausible upper  
 1403 bound of sea-level rise projections, *Climatic Change*, 115, 893-902, 10.1007/s10584-012-  
 1404 0610-6, 2012.  
 1405 Stocker, T.: Model Hierarchy and Simplified Climate Models, in: *Introduction to Climate*  
 1406 *Modelling, Advances in Geophysical and Environmental Mechanics and Mathematics*,  
 1407 Springer Berlin Heidelberg, 25-51, 2011.  
 1408 Takahashi, T., Sutherland, S. C., Wanninkhof, R., Sweeney, C., Feely, R. A., Chipman,  
 1409 D. W., Hales, B., Friederich, G., Chavez, F., Sabine, C., Watson, A., Bakker, D. C. E.,  
 1410 Schuster, U., Metzl, N., Yoshikawa-Inoue, H., Ishii, M., Midorikawa, T., Nojiri, Y.,  
 1411 Körtzinger, A., Steinhoff, T., Hoppema, M., Olafsson, J., Arnarson, T. S., Tilbrook, B.,  
 1412 Johannessen, T., Olsen, A., Bellerby, R., Wong, C. S., Delille, B., Bates, N. R., and de  
 1413 Baar, H. J. W.: Climatological mean and decadal change in surface ocean pCO<sub>2</sub>, and net  
 1414 sea-air CO<sub>2</sub> flux over the global oceans, *Deep Sea Research Part II: Topical Studies in*  
 1415 *Oceanography*, 56, 554-577, <http://dx.doi.org/10.1016/j.dsr2.2008.12.009>, 2009.  
 1416 Tanaka, K., Kriegler, E., Bruckner, T., Hooss, C., Knorr, W., and Raddatz, T.:  
 1417 Aggregated Carbon Cycle, Atmospheric Chemistry, and Climate Model (ACC2) -  
 1418 description of the forward and inverse models, Max Planck Institute for Meteorology,  
 1419 Hamburg, Germany, 188, 2007a.  
 1420 Tanaka, K., Kriegler, E., Bruckner, T., Hooss, G., Knorr, W., Raddatz, T. J., and Tol, R.:  
 1421 Aggregated carbon cycle, atmospheric chemistry, and climate model (ACC2), Hamburg,  
 1422 188, 2007b.  
 1423 Taylor, K. E., Stouffer, R. J., and Meehl, G. A.: An Overview of CMIP5 and the  
 1424 Experiment Design, *Bulletin of the American Meteorological Society*, 93, 485-498,  
 1425 10.1175/BAMS-D-11-00094.1, 2012.  
 1426 Urban, N. M., and Keller, K.: Complementary observational constraints on climate  
 1427 sensitivity, *Geophysical Research Letters*, 36, L04708, 10.1029/2008GL036457, 2009.  
 1428 Urban, N. M., and Keller, K.: Probabilistic hindcasts and projections of the coupled  
 1429 climate, carbon cycle and Atlantic meridional overturning circulation system: a Bayesian  
 1430 fusion of century-scale observations with a simple model, *Tellus A*, 62, 737-750,  
 1431 10.1111/j.1600-0870.2010.00471.x, 2010.  
 1432 van Vuuren, D., Elzen, M. J., Lucas, P., Eickhout, B., Strengers, B., Ruijven, B., Wonink,  
 1433 S., and Houdt, R.: Stabilizing greenhouse gas concentrations at low levels: an assessment

1434 of reduction strategies and costs, *Climatic Change*, 81, 119-159, 10.1007/s10584-006-  
1435 9172-9, 2007.

1436 van Vuuren, D., Lowe, J., Stehfest, E., Gohar, L., Hof, A., Hope, C., Warren, R.,  
1437 Meinshausen, M., and Plattner, G.-K.: How well do integrated assessment models  
1438 simulate climate change?, *Climatic Change*, 104, 255-285, 10.1007/s10584-009-9764-2,  
1439 2011.

1440 Ward, D. S., and Mahowald, N. M.: Contributions of developed and developing countries  
1441 to global climate forcing and surface temperature change, *Environmental Research*  
1442 *Letters*, 9, 074008, 2014.

1443 Wigley, T. M. L.: A simple inverse carbon cycle model, *Global Biogeochemical Cycles*,  
1444 5, 373-382, 10.1029/91GB02279, 1991.

1445 Wigley, T. M. L.: Global-mean temperature and sea level consequences of greenhouse  
1446 gas concentration stabilization, *Geophysical Research Letters*, 22, 45-48,  
1447 10.1029/94GL01011, 1995.

1448 Wigley, T. M. L., Richels, R., and Edmonds, J. A.: Economic and environmental choices  
1449 in the stabilization of atmospheric CO<sub>2</sub> concentrations, *Nature*, 379, 240-243, 1996.

1450 Wigley, T. M. L., Smith, S. J., and Prather, M. J.: Radiative Forcing Due to Reactive Gas  
1451 Emissions, *Journal of Climate*, 15, 2690-2696, 10.1175/1520-  
1452 0442(2002)015<2690:RFDTRG>2.0.CO;2, 2002.

1453 Wise, M., Calvin, K., Thomson, A., Clarke, L., Bond-Lamberty, B., Sands, R., Smith, S.  
1454 J., Janetos, A., and Edmonds, J.: Implications of Limiting CO<sub>2</sub> Concentrations for Land  
1455 Use and Energy, *Science*, 324, 1183-1186, 10.1126/science.1168475, 2009.

1456 Wolkovich, E. M., Regetz, J., and O'Connor, M. I.: Advances in global change research  
1457 require open science by individual researchers, *Global Change Biology*, 18, 2102-2110,  
1458 10.1111/j.1365-2486.2012.02693.x, 2012.

1459 Zeebe, R. E., and Wolf-Gladrow, D.: *CO<sub>2</sub> in Seawater: Equilibrium, Kinetics, Isotopes*,  
1460 Elsevier, 2001.

1461

1462

1463 **Table and Figure Captions:**

1464 **Table 1:** Initial mModel conditions prior to the spinup phase. Carbon values change  
 1465 slightly after spinning up to a steady state., assuming a pre-industrial steady state.  
 1466

Variable	Description	Initial Value	Units	Notes
*C <sub>atm</sub>	Atmospheric Carbon	588.1	PgC	<a href="#">Murakami(2010)</a>
*C <sub>D</sub>	Detritus Carbon	55.0	PgC	<a href="#">Denman et al.,(2007)</a> <u>Land carbon (detritus, soil and vegetation) totaling ~2300PgC</u>
*C <sub>S</sub>	Soil Carbon	1782.0	PgC	
*C <sub>V</sub>	Vegetation Carbon	550.0	PgC	
C <sub>DO</sub>	Deep Ocean	26000.0	PgC	<a href="#">Denman et al.,(2007)</a> <u>Ocean carbon (deep, intermediate and surface) totaling ~3800PgC **</u>
C <sub>HL</sub>	Surface Ocean High Latitude	140.0	PgC	
C <sub>IO</sub>	Intermediate Ocean	8400.0	PgC	
C <sub>LL</sub>	Surface Ocean Low Latitude	770.0	PgC	
F <sub>L</sub>	Atmosphere-Land Carbon Flux	0.0	PgC yr <sup>-1</sup>	
F <sub>O</sub>	Atmosphere-Ocean Carbon Flux	0.0	PgC yr <sup>-1</sup>	
NPP <sub>0</sub>	Net Primary Production	50.0	PgC yr <sup>-1</sup>	<u>Approximate global value.</u> <a href="#">Nemani et al.,(2003)</a>
T <sub>G</sub>	Global Temperature Anomaly	0.0	°C	
T <sub>HL</sub>	Temperature of high latitude surface ocean box	2.0	°C	<a href="#">Lenton.,(2000)</a>
T <sub>LL</sub>	Temperature of low latitude surface ocean box	22.0	°C	<a href="#">Lenton.,(2000)</a>

1467 \* parameters appearing in the input file.

1468 \*\* in order to obtain a steady state in Hector, carbon values in the intermediate box are

1469 less than reported Denman et al.,(2007):

1470 **Table 2:** Model parameters for the land and ocean carbon components.

Variable	Description	Value	Notes
$f_{ds}$	annual fraction of detritus carbon that is transferred to soil	0.60	<u>The following fractions (<math>f</math>) were selected to be generally consistent with previous simple earth system models (e.g., Meinshausen et al., 2011a; Ricciuto et al., 2008; Murakami et al., 2010).</u>
$*f_{ld}$	annual fraction of land use change flux from detritus	0.01	
$f_{ls}$	annual fraction of land use change flux from soil	0.89	
$*f_{lv}$	annual fraction of land use change flux from vegetation	0.10	
$*f_{nd}$	annual fraction of NPP carbon that is transferred to detritus	0.60	
$f_{ns}$	annual fraction of NPP carbon that is transferred to soil	0.05	
$*f_{nv}$	annual fraction of NPP carbon that is transferred to vegetation	0.35	
$f_{rd}$	annual fraction of respiration carbon that is transferred to detritus	0.25	
$f_{rs}$	annual fraction of respiration carbon that is transferred to soil	0.02	
$f_{vd}$	annual fraction of vegetation carbon that is transferred to detritus	0.034	
$f_{vs}$	annual fraction of vegetation carbon that is transferred to soil	0.001	
$*\beta$	Beta	0.36	
$*Q_{10}$	Q10 respiration	2.45	
$*T_H$	High-latitude circulation	$4.9e7 \text{ m}^3 \text{ s}^{-1}$	<u>Tuned to give ~100 PgC from surface to deep</u>
$*T_T$	Thermohaline circulation	$7.2e7 \text{ m}^3 \text{ s}^{-1}$	<u>Tuned to give ~100</u>

			<u>PgC from surface to deep</u>
*E <sub>ID</sub>	Water mass exchange – intermediate <u>to</u> -deep	1.25e7 m <sup>3</sup> s <sup>-1</sup>	<u>Lenton, 2000; Knox and McElroy, 1984</u>
*E <sub>LI</sub>	Water mass exchange – low latitude <u>to</u> -intermediate	2.0e8 m <sup>3</sup> s <sup>-1</sup>	<u>Lenton, 2000; Knox and McElroy, 1984</u>

1471

1472

\* parameters appearing in the input file.



1473 **Table 3:** CMIP5 ESM models used within this study. We use the same suite of models as  
 1474 found in Friedlingstein et al. (2014). Note, not all variables are reported for each model  
 1475 under all scenarios.  
 1476

Model	Model Name	Institute
bcc-csm1-1	Beijing Climate Center, Climate System Model, version 1.1	Beijing Climate Center, China Meteorological Administration, China
CanESM2_*	Second Generation Canadian Earth System Model	Canadian Center for Climate Modeling and Analysis, BC, Canada
CESM1-BGC_*	Community Earth System Model, version 1.0-Biogeochemistry	National Center for Atmospheric Research, United States
GFDL-ESM2G	Geophysical Fluid Dynamic Laboratory Earth System Model with GOLD ocean component	Geophysical Fluid Dynamics Laboratory, United States
HadGEM2-ES	Hadley Centre Global Environmental Model, version 2 (Earth System)	Met Office Hadley Centre, United Kingdom
inmcm4	Institute of Numerical Mathematics Coupled Model, version 4.0	Institute of Numerical Mathematics, Russia
IPSL-CM5A-LR	L'Institut Pierre-Simon Laplace Coupled Model, version 5A, coupled with NEMO, low resolution	Institut Pierre Simon Laplace, France
MIROC-ESM_*	Model for Interdisciplinary Research on Climate, Earth System Model	Atmosphere and Ocean Research Institute; National Institute for Environmental Studies, Japan Agency for Marine-Earth Science and Technology, Japan
MPI-ESM-LR	Max Planck Institute Earth System Model, low resolution	Max Planck Institute for Meteorology, Germany
MRI-ESM1_*	Meteorological Research Institute Earth System Model, version 1	Meteorological Research Institute Earth, Japan
NorESM1-ME_*	Norwegian Earth System Model, version 1, intermediate	Norwegian Climate Center, Norway

resolution

\* Models used in emissions forced scenarios (esmhist and esmrp85).

1477 **Table 4: Skill of Root mean square error (RMSE) for** Hector versus observations, CMIP5,  
 1478 **and MAGICC, correlation coefficients (R) and root mean square error (RMSE) for**  
 1479 **atmospheric [CO<sub>2</sub>], surface temperature anomaly, radiative forcing, fluxes of carbon**  
 1480 **(ocean and land), and low latitude surface ocean pH and delta change (Δ) for**  
 1481 **atmospheric [CO<sub>2</sub>], surface temperature anomaly and radiative forcing for Hector,**  
 1482 **CMIP5, observations, and MAGICC6.**

Historical 1850 - 2005						
Variable	Skill	Hector	Observations	MAGICC	CMIP5	Units
[CO <sub>2</sub> ]*	RMSE	--	2.85	2.95	2.21	ppmv
	Δ	85.78	94.47	95.0	103.30	
temperature	RMSE	--	0.15	0.13	0.15	deg C
	Δ	0.98	0.91	0.76	1.01	
Forcing	RMSE	--	--	0.39	--	W m <sup>-2</sup>
	Δ	2.16	--	1.75	--	
Ocean Flux	RMSE	--	--	--	0.25	PgC yr <sup>-1</sup>
Land Flux	RMSE	--	--	--	1.27	PgC yr <sup>-1</sup>
pH	RMSE	--	--	--	0.004	unitless

\*[CO<sub>2</sub>] observations are an average of Law Dome and Mauna Loa.

RCP 8.5 1850 - 2300					
Variable		Hector	MAGICC	CMIP5	Units
[CO <sub>2</sub> ] *	RMSE	--	10.41	7.54	ppmv
	Δ	1557.91	1695.0	--	
temperature	RMSE	--	0.12	0.52	deg C
	Δ	9.58	8.05	10.57	
Forcing	RMSE	--	0.26	--	W m <sup>-2</sup>
	Δ	12.80	12.24	--	
Ocean Flux	RMSE	--	--	1.39	PgC yr <sup>-1</sup>
Land Flux	RMSE	--	--	3.86	PgC yr <sup>-1</sup>
pH	RMSE	--	--	0.003	unitless

\*CMIP5 [CO<sub>2</sub>] only to 2100.

1483

## RCP 8.5 2005 - 2300

Variable		Hector	MAGICC	CMIP5	Units
[CO <sub>2</sub> ]*	RMSE	--	10.07	7.23	ppmv
	Δ	1472.13	1600.0	--	
temperature	RMSE	--	0.09	0.58	deg C
	Δ	8.59	7.30	9.57	
Forcing	RMSE	--	0.03	--	W m <sup>-2</sup>
	Δ	10.65	10.49	--	
Ocean Flux	RMSE	--	--	1.41	PgC yr <sup>-1</sup>
Land Flux	RMSE	--	--	4.59	PgC yr <sup>-1</sup>
pH	RMSE	--	--	0.001	unitless


\*CMIP5 [CO<sub>2</sub>] only to 2100.

1484

1485



1486

1487 **Figure 1:** Model phases for the coupler (left) and a typical component (right). Arrows  
 1488 show flow of control and data. The greyed spinup step is optional.




1489  
 1490

1491 **Figure 2:** Representation of Hector’s carbon cycle, land, atmosphere, and ocean. The  
1492 atmosphere consists of one well mixed box. The ocean consists of four boxes, with  
1493 advection and water mass exchange simulating thermohaline circulation ([see Table 2 for](#)  
1494 [description of parameters](#)). At steady state, the high latitude surface ocean takes up  
1495 carbon from the atmosphere, while the low latitude surface ocean off gases carbon to  
1496 the atmosphere. The land consists of a user defined number of biomes or regions for  
1497 vegetation, detritus and soil. At steady state the vegetation takes up carbon from the  
1498 atmosphere while the detritus and soil release carbon back into the atmosphere. The  
1499 earth pool is continually debited with each time step to act as a mass balance check on  
1500 the carbon system.



1503  
1504


1505 | **Figure 3:** Historical atmospheric [CO<sub>2</sub>] from 1850 to 2005 for Hector (blue), CMIP5  
1506 | median, standard deviation, and model range (pink, n=4), MAGICC6 (green), Law Dome  
1507 | (teal), and Mauna Loa (purplebrown). Note CMIP5 data are from the prescribed  
1508 | emissions -historical scenario (esmHistorical). ~~Notice that~~ MAGICC6, however, is  
1509 | constrained to ~~match~~ es the observational record. ~~We have the capabilities of running~~  
1510 | ~~Hector under numerous constraints. Within~~ Although Hector can be run with similar  
1511 | constraints, in this study ~~we are running~~ Hector was unconstrained to highlight the full  
1512 | performance of the model. n=4 is the number of CMIP5 models used to produce this  
1513 | figure.



1514  
1515  
1516




1517 **Figure 4:** Historical global temperature anomaly relative to 1850 for Hector (blue),  
1518 MAGICC6 (green), CMIP5 median, standard deviation and model spread-range (pink,  
1519 n=8), and historical observations from HadCRUT4 (purple). Hector is running without the  
1520 effects of volcanic forcing, leading to a smoother representation of temperature with  
1521 time.  
1522




1523

1524 | **Figure 5:** Atmospheric [CO<sub>2</sub>] from 1850 to 2100 under RCP 8.5 for Hector (**blue**),  
1525 | MAGICC6 (**green**), Mauna Loa (**purplebrown**), Law Dome (**teal**) and esmRCP 8.5  
1526 | (**prescribed emissions scenario**) CMIP5 median, one standard deviation and model  
1527 | **spread range** (pink, **n=4 (1850-2000) and n=5 (2001-2100)**). Note that the CMIP5  
1528 | models run under esmrcp85 do not extend to 2300.




1529  
1530

1531 | **Figure 6:** Atmospheric [CO<sub>2</sub>] from 1850 to 2300 for RCP 2.6 (yellowred), RCP 4.5 (green),  
1532 RCP 6.0 (blue), RCP 8.5 (purple), Hector (solid) and MAGICC6 (dashed).  
1533




1534  
1535

1536 **Figure 7:** Relative radiative forcing from 1850 to 2300 for Hector (solid) and MAGICC6  
1537 (dashed) for all four RCP scenarios, 2.6 (red), 4.5 (green), 6.0 (blue), 8.5 (purple). Hector  
1538 has the option to enable or disable radiative forcing from historical volcanic emissions.  
1539 We have opted to disable this for ease of comparison across all RCPs.  
1540



1541  
1542


1543 **Figure 8:** Global temperature anomaly relative to 1850 for (a) RCP 2.6 (b) RCP 4.5 (c) RCP  
 1544 6.0 and (d) RCP 8.5, comparing Hector (blue), MAGICC6 (green), and CMIP5 median,  
 1545 standard deviation and model spread-range (pink). The CMIP5 models under RCP 6.0  
 1546 used in this study do not extend to 2300. Note the change in scales between the four  
 1547 panels. Number of CMIP5 models in a) n=7 (2006-2100) and n=5 (2101-2300), b) n=9  
 1548 (2006-2100) and n=6(2101-2300), c) n=6 (2006-2100), d) n=9 (2006-2100) and n=3  
 1549 (2101-2300).



1550  
 1551


1552 | ~~Figure 9: Taylor diagram of global temperature anomaly relative to 1850, from 1850 to~~  
1553 | ~~2300 for RCP 8.5, Hector (), MAGICC6 (), CMIP5 median (red), and CMIP5 models (grey).~~  
1554 |  
1555 |

1556 | **Figure 109:** Global air-sea fluxes of carbon under RCP 8.5, Hector (blue), MAGICC5.3  
1557 | (purple, note that this is not the current version of MAGICC), CMIP5 median, standard  
1558 | deviation, and model range (~~redpink~~, n=9 (1850-2100) and n=4 (2101-2300)), and  
1559 | observations from GCP (green) (Le Quéré et al., 2013). The break in the graph at 2100  
1560 | signifies a change in the number of models that ran the RCP 8.5 extension.



1561  
1562


1563 | **Figure 104:** Global air-land fluxes of carbon under RCP 8.5, Hector (blue), CMIP5 median,  
1564 | standard deviation, and model range CMIP5-(redpink, n=8 (1850-2100) and n=2 (2101-  
1565 | 2300)), and observations from GCP (green) (Le Quéré et al., 2013). The break in the  
1566 | graph at 2100 signifies a change in the number of models that ran the RCP 8.5  
1567 | extension.  
1568



1569



1570 | **Figure 112:** Low latitude (< 55) ocean pH for RCP 8.5, from 1850 – 2100, Hector (blue),  
1571 | CMIP5 median, standard deviation, and model range (pink, n=6) and observations from  
1572 | BATS (green) and HOTS (purple).  
1573



1574