

### Response to Anonymous Referee #1

The authors appreciate the constructive feedback provided by Reviewer #1. We address their comments point-by-point below. The reviewer's original comments appear in quotations and our responses follow. All page numbers referring to the manuscript in our responses below are valid for the revised manuscript, which is provided as a supplement to Anonymous Referee #1 within the interactive discussion with track-changes used to show revisions (<http://www.geosci-model-dev-discuss.net/7/C3029/2015/gmdd-7-C3029-2015-supplement.pdf>).

**“This paper contains a good discussion of various approaches to simulating lakes within a regional climate model and the results of these approaches. It has strong value as a review paper, in addition to having a modest amount of original results. My comments are minor in nature, but I encourage the authors to heed them.”**

#### Specific comments:

**1. “P. 7123 uses ‘interpolation’ on line 9, but then contradicts on line 17 by saying ‘no interpolation’. Clarify by possibly using other terms, like ‘spatial analysis.’”**

The first use of “interpolation” in the sentence has been deleted (p. 2, line 55). The meaning of the sentence is clear without it.

**2. “P. 7123, lines 9-15—This seems to be trying to compromise between accuracy and simplicity. I suggest just describing the situation that prevails in areas of interest, and the method that ends up being used there.”**

We agree that the paragraph is detailed in its description of some interpolation methods with only a somewhat vague explanation of the others, which is what we believe the reviewer is describing as a “compromise.” The paragraph has been amended (p. 2, line 58-60) to direct interested readers to where a full description of all interpolation methods is available. A detailed description of all interpolation methods WPS utilizes is outside the scope of our study, as most are not relevant in our area of interest. However, we feel it is important to note that various methods are attempted using several points from the driving fields or just one point, and it is only after a series of methods have been attempted that the search method is employed. Otherwise, a reader could infer that the search method is often used, which may not be the case when the WRF Preprocessing System is used for other applications.

**3. “P. 7124, line 25—It seems strange to refer to a reanalysis dataset as a proxy for a GCM. It’s just a dataset that can be used to drive an RCM for different purposes than would be achieved using GCM data.”**

The sentence has been reworded (p. 4, line 102) to clarify why we refer to this dataset as a “proxy.” It is clearly stated that the R2 is a reanalysis dataset. The term “proxy for a GCM” is applied because it describes the purpose of driving with a coarse GCM which, as you mentioned above, is the historical evaluation of a downscaling methodology. Reminding readers of why the R2 is used is important, because the problems we focus on with the initialization of lakes could easily be solved by driving our runs with a higher resolution analysis. However, this would not serve the purpose of the retrospective downscaled runs, which is to establish a credible methodology for downscaling GCMs.

**4. “This citation likely belongs in this page: Lofgren, B. M., 2004: A model for the simulation of climate and hydrology in the Great Lakes basin. J. Geophys. Res., 109, doi:10.1029/2004JD004602”**

We are unsure for which page the reviewer is suggesting this citation. While Lofgren (2004) does discuss an approach applied within a RAMS-based model which is similar to WRF’s “search” methodology, the present work is specifically focused on the representation of lakes within WRF, and it is beyond the scope of our study to describe how lake surface variables are set in RAMS or any other RCM. We do briefly discuss general problems in NARCCAP RCMs on p. 5 (bottom paragraph), but only to bring to the attention of the reader that WRF is not the only RCM that tends to represent lakes poorly in a downscaling application.

**5. “P. 7129, line 24 implies that there is a correction that was applied in version 3.5.1. What is it?”**

The table suggested below by the reviewer has been added to the manuscript and a reference to it is inserted to further clarify this sentence (p. 8, line 240). As stated in the introduction (p. 4, bottom paragraph), the default water surface temperature at which WRF determines ice is formed was lowered from 271 K to 100 K as of version 3.5.1.

**6. “There are a whole lot of references to features of different versions of WRF spread across the manuscript. I suggest summarizing the relevant updates in a table.”**

A new table (Table 1 in the manuscript) has been inserted into the manuscript and several references to the table have been added to the text in order to aid the reader to noting the changes between versions more easily.

**7. “P. 7131, lines 7-14 overstate the uniqueness of ice as a limiting factor in evaporation. An ice-free lake can have cool water overlaid by warm and moist air, resulting in very limited evaporation.”**

The sentence has been revised to say that its stated conclusions about the link between ice and precipitation are valid during the lake unstable season (p. 10, line 282) which is characterized by warm lake temperatures and relatively cool air masses. The role of temperatures, as well as ice, in suppressing evaporation is discussed more generally in Section 2.0 (p. 6-7). The studies being reviewed in this paragraph (Burnett et al., 2003; Kunkel et al., 2009; and Gula and Peltier, 2012) discuss changes in ice cover and its effects on precipitation and evaporation during the lake unstable season.

**8. “P. 7134, last paragraph discusses how well FLake performs during historical periods. Are there any thoughts on how well it might do at climate change scenarios in which lake thermal structure and stability may well change, e.g. the situation illustrated by Austin and Colman (2007)?”**

FLake’s ability to simulate the thermal structure of the lake can be limited by its inability to simulate 2- and 3-D processes and its assumption that the temperature profile consists of a homogeneous mixed layer and stable layer (thermocline) which extends to lake bottom. However, the shape (and therefore, stability) of its thermal profile can vary due to a variety of physical processes, including convective mixing and radiative heating of the water column. We

expect that its performance over historical periods is indicative of its accuracy under a climate change scenario. However, it does not presently account for changes in lake depths which can be anticipated to change over the coming decades and this could negatively affect its performance in making future projections at multi-decadal time scales. This limitation is mentioned on page 15 (lines 443-446) of the present manuscript.

**9. “What is the horizontal structure of both FLake and CLM? Are you using a horizontal array of non-interacting 1-d columns? The caption of Fig. 4 seems to imply yes for CLM, but unless I missed it, this should be stated more explicitly.”**

The reviewer’s understanding is correct. In the case of both models, there is no horizontal interaction between columns. When the CLM lake model is first introduced we state that it is one-dimensional (p. 11, lines 324-325). The text describing the FLake model has been modified to state explicitly that it is a 1-D column model (p.12, line 350).

**10. “P. 7135, lines 10-15–This is probably the most troubling part of this manuscript for me. The time series in Fig. 3 is showing strong evidence of numerical instability–highly unrealistic oscillation between high and low values over very short periods of time. This shouldn’t happen even during a spin-up period. Without knowing the details of the code, it seems miraculous, first, that this instability continues for so long without crashing the model run, and second, that it suddenly stops and remains stable thereafter.”**

We agree that the results are highly unrealistic, which is why we choose to highlight the spin-up issues with these models so that users would be aware of the potential ill effects of running lake models without accounting for needed spin-up time. In the case of FLake, it should be noted that such values only occur during the *first* annual cycle of the ten cycles which are used to complete spin-up and these early results are sensitive to the model’s initial state. The offline FLake run is initialized with surface temperatures uniformly set with a user-defined default (274.15 K) and a mixed layer depth which defaults to zero. As noted in the text, initial LSTs in the CLM simulation shown in Fig. 4 are taken from the interpolation from the R2, which also provides a poor initial lake state. We would expect that these initial conditions would negatively impact both models’ short-term results, which is why we emphasize the need for adequate spin-up time to allow the simulated lake state to achieve equilibrium with the driving conditions and to become insensitive to its initial state.

To put these results in further context, the caption of Fig. 3 has been amended to include more information about the initialized lake state for the offline FLake simulation and to note that the time series also shows ice temperatures as well. Some drops in surface temperature are expected as ice forms and ice-top temperatures cool. However, this abrupt cooling below 200 K is, as we note in the text, highly unrealistic. Since providing adequate spin-up time resulted in realistic values in our FLake simulations, we have not looked further into this aspect of the model’s behavior. Being less experienced with the CLM lake model, we cannot comment on the potential causes of its behavior.

**11. “P. 7135, lines 16-17–To reiterate and clarify the comment by Anne Clites, the dataset described in Wang et al. (2012b) is not simply the NIC analysis, but is a value-added dataset, with additional quality checks and a gridded format.”**

The text (p. 14, lines 399-401) has been modified to properly cite Wang et al. (2012b) without misattributing the work to the NIC. In addition, we have added a sentence to the Acknowledgements to state that this data was obtained from the Great Lakes Environmental Research Laboratory.

Technical corrections:

**1. “P. 7123, line 1–insert ‘spatial’ before ‘interpolation.’”**

The suggested revision has been made (p. 2, line 47).

**2. “P. 7128, line 26–‘result from a downscaled simulation’ seems clearer than ‘result in...’”**

This correction has been made (p. 7, line 214).

**3. “P. 7130, line 28–Especially because winter itself is a limited time period, it seems to make sense to replace ‘at a later time period’ with simply ‘later.’”**

The wording has been changed, as suggested (p. 9, line 272).

**4. “P. 7131, line 1–‘...open (and free of ice)’ is redundant. I suggest just ‘free of ice.’”**

This revision has been made (p. 9, line 273).

**5. “Check a style guide on hyphen usage: Remove hyphens from p. 7135 line 5 ‘spin up’, p. 7135 line 12 ‘time series’, and p. 7136 line 11 ‘spun up’. Add one to p. 7136 line 3 ‘110-year’.”**

Most of the suggested revisions have been made (p. 13, line 395; p. 14, line 416). The use of “spin-up” was vetted by a technical editor when manuscript proofs were made.

## Response to Anonymous Referee #2

The authors are thankful for the constructive criticism and commentary provided by Reviewer #2. We address their comments point-by-point below. The reviewer's original comments appear in quotations below and our responses follow. All page numbers referring to the manuscript in our responses below are valid for the revised manuscript with track-changes used to show corrections. The revised manuscript is provided as a supplement to Anonymous Referee #1 within the interactive discussion.

**“This paper presents a brief overview of several problems in generating lake surface temperatures from global climate models (GCM) with underrepresentation of lakes to be used to run WRF as a regional climate model (RCM). It gives a good deal of information about many of techniques that can be used to achieve this, as well as the drawbacks from these methods. While most things are considered, there are a few areas that could use further explanation. Most of these are minor in nature.”**

### **Overall Comments:**

**“I think some background information on how WRF, being run as a regional climate model, treats lakes would be beneficial for context.”**

We have revised the introduction (see p. 2, lines 35-38, lines 42-46) to provide further understanding of how WRF functions as an RCM in its default configuration without a lake model. We also address your more specific questions below.

**“Are surface properties like lake temperature taken from the GCM at each time step, or does the land surface model within WRF deal with this temperature calculation?”**

A sentence has been added on p. 2, lines 42-46 to clarify this. In its default configuration with no lake model used to prognosticate lake surface temperatures (LSTs), LSTs are calculated in the preprocessing steps before WRF is run and then prescribed to the model at runtime. Therefore, the land surface model plays no role in calculating lake temperature. The frequency with which the prescribed temperatures are read is set by the user, but it is commonly set to daily or sub-daily, depending on the user's application and availability of driving data.

**“When temperatures are initialized or a scenario without a lake model, is it simply a one-layer slab model, or multiple layers similar to land points? Does WRF apply a diurnal cycle?”**

As stated above, when no lake model is used, lake (and ocean) surface temperatures are simply read in from an input file after having been interpolated from the driving dataset. This is further clarified in the manuscript on p. 2, lines 35-38. Although a diurnal cycle should be present in any well-vetted, observationally-based sea surface temperature (SST) data used to drive the WRF model, no adjustment is done by WRF to create a diurnal cycle in water temperatures.

This presence of a diurnal cycle is also sensitive to the frequency with which the user updates the prescribed water temperatures [i.e., whether the user specifies a sub-daily (3- or 6-h) frequency to read SSTs into the model]. Although this aspect of the model settings can be arbitrarily chosen, we briefly mention that the commonly used timescales in the revised text (p. 2, lines 42-46).

**“I think answering some of these questions would help to put some of the methods into better context and show how errors in initialization may propagate in a model, especially when no lake model is being used.”**

The reviewer’s questions indicate that further explanation of how WRF prescribes water temperatures is needed in the introduction in order to avoid confusion later in the manuscript. The revised article has been corrected to include more general description of WRF’s treatment of water temperatures, as discussed above.

### **Specific Comments:**

**1. “Pg. 7124 line 21- Pg. 7125 line 2. I am not sure this paragraph is entirely needed. The previous paragraph describes the same situation with visuals that is shown by M14. Some further explanation is either needed to show how this is a different problem than what is presented in Figures 1 and 2, or this section should be pared down. This section could also be worked into the first paragraph of Page 7126.”**

We feel that that this paragraph illustrates the motivation for the study best in its current position because it provides an example of how lake temperatures are set when the lake is partially represented by the GCM. This contrasts well with what is shown in Figs. 1 and 2, where the Great Lakes are not represented at all. An additional sentence has been added on p. 4, lines 101-104 to state this contrast and put the paragraph in better context.

**2. “Pg. 7130 Section 2.2. Has this method been used by any other study? You state that linearly increasing lake states maybe useful for some lakes, but you give no examples of this approach being employed. It seems like this approach offers very little in terms of realism and upside, so is it necessary to be mentioned?”**

We have not found examples of this approach being used. However, the use of stationarity assumptions is ubiquitous in regional climate modeling, where future simulations are routinely run with the same land-use and vegetation fields that are used to simulate the present-day climate. Therefore, it seemed necessary that we examine how the option of using present-day lake surface temperatures and ice cover could affect a future simulation. Warming lake temperatures by a constant rate or using a linear increase would be a logical next step to improve the accuracy of such a method; however, we do not have a specific example, either in this work or by referencing another, to illustrate this. To put this work section in further context, we have added a sentence on p. 9, line 260-261 to clarify why this option is discussed.

**3. “Page 7131 Section 2.3. In the use of this method, I understand where the land-lake temperature contrast would be lost, at least in the short-term. But given enough spinup time (similar to what is shown in section 2.6), could these contrasts be generated from lake-atmospheric interactions, or is this still a case of poor initialization leading to poor results?”**

In this methodology, as applied by Gao et al. (2012), no lake model is run and WRF uses prescribed water temperatures which are calculated during the preprocessing steps before the WRF simulation begins. Therefore, it can simulate only a one-way interaction between the atmosphere and the lake (i.e., the lake state impacting atmospheric conditions). Because this is a one-way interaction based on prescribed lake temperatures from the driving data, even with a protracted spin-up time, the lake temperature could not be forced to produce a lake-land temperature contrast.

**4. “Page 7135, lines 8-10. What do you mean by ‘looped’ here? Do you mean using the atmospheric conditions from the year 2005, and ran that same data 10 times while allowing the lake conditions to evolve? Some clarification might be beneficial here.”**

The reviewer’s understanding of how we ran the model is correct. We have revised the wording (p. 13, 387-393) to state this more clearly.

### Response to Short Comment

The original comment appears below and our response follows. All page numbers referring to the manuscript are valid for the revised article, which is provided as a supplement to Anonymous Referee #1 within the interactive discussion with track-changes used to show corrections.

**“On page 7135, lines 15-17, you refer to a reference: ‘as shown from the National Ice Center Great Lakes Ice Analysis’ while referring to a NOAA-GLERL Technical Memorandum (TM-155, by Wang et al, 2012).**

**This description of our paper is incorrect. While it does use NIC data, the article referred to here is not a National Ice Center publication. This sentence should be changed accordingly.**

**Thanks,**

**Anne H. Clites, NOAA Great Lakes Environmental Research Laboratory Ann Arbor, MI”**

We thank Anne Clites for clarifying this distinction. The text (p. 14, 1<sup>st</sup> paragraph) has been modified to state that the National Ice Center analysis charts were processed and provided by the Great Lakes Environmental Research Laboratory (GLERL) and to properly cite Wang et al. (2012b). In addition, we have added a sentence to the Acknowledgements to state that this data was obtained from GLERL.



# Technical challenges and solutions in representing lakes when using WRF in downscaling applications

M. S. Mallard<sup>1,\*</sup>, C. G. Nolte<sup>1</sup>, T. L. Spero<sup>1</sup>, O. R. Bullock<sup>1</sup>, K. Alapaty<sup>1</sup>, J. A. Herwehe<sup>1</sup>, J. Gula<sup>2</sup>, and J. H. Bowden<sup>3</sup>

[1]{National Exposure Research Laboratory, U.S. Environmental Protection Agency, Research Triangle Park, North Carolina, USA }

[2]{Department of Atmospheric and Oceanic Sciences, University of California, Los Angeles, California, USA }

[3]{Institute for the Environment, University of North Carolina, Chapel Hill, North Carolina, USA }

\*Currently at: Institute for the Environment, University of North Carolina, Chapel Hill, North Carolina, USA

Correspondence to: M. S. Mallard (mmallard@email.unc.edu)

Revised manuscript: 21 January 2015

## Abstract

The Weather Research and Forecasting (WRF) model is commonly used to make high resolution future projections of regional climate by downscaling global climate model (GCM) outputs. Because the GCM fields are typically at a much coarser spatial resolution than the target regional downscaled fields, inland lakes are often poorly resolved in the driving global fields, if they are resolved at all. In such an application, using WRF's default interpolation methods can result in unrealistic lake temperatures and ice cover at inland water points. Prior studies have shown that lake temperatures and ice cover impact the simulation of other surface variables, such as air temperatures and precipitation, two fields that are often used in regional climate applications to understand the impacts of climate change on human health and the environment. Here, alternative methods for setting lake surface variables in WRF for downscaling simulations are presented and contrasted.

## 1 Introduction

When using global climate model (GCM) fields to drive finer-scale regional climate model (RCM) runs, typically the RCM does not have an oceanic or lake physics component and relies on the GCM output to provide all water surface temperatures and ice cover. Within a downscaling simulation, by design, the GCM is at a coarser spatial resolution than the RCM, so inland water bodies in the region being simulated are either poorly resolved or not resolved by the GCM. ~~Some RCM configurations do include an oceanic component. The~~ Prior to 2013, the Weather Research and Forecasting (WRF) model (Skamarock et al., 2008) required exogenously prescribed water surface temperatures, as there was not capability to prognosticate water temperatures. WRF has included an optional coupled ocean component since ~~WRF~~-version 3.5 was released in April 2013 (~~Skamarock et al., 2008~~; WRF User's Guide, 2014). Other RCMs have been coupled to ocean models in order to simulate regions around the Arctic, Mediterranean Sea, and Indian Ocean (e.g., Rinke et al., 2003; Ratnam et al., 2009; Artale et al., 2010; Gualdi et al., 2013). However, when using WRF's default configuration, the sea surface temperature (SST) fields used during the simulation are calculated from the driving data during the preprocessing steps performed before WRF runs the simulation; during the model run, these prescribed water temperatures are input at a user-specified frequency which is usually daily or sub-daily. Similarly, lake surface temperatures (LSTs) and lake ice cover are prescribed by spatial interpolation from the ~~sea surface temperature (SST)~~ and sea ice fields in the driving data. In this study, we examine the use of the Advanced Research WRF (Skamarock and Klemp, 2008) model applied as an RCM in regions where the driving larger-scale data have a poor representation of lakes.

When the WRF Preprocessing System (WPS) interpolates skin temperatures from the coarser global dataset (where both land and water temperatures are included in a single field), masks are applied such that water temperatures from the GCM are used to set water temperatures on the finer, target grid. Using the standard ~~interpolation~~-methods in WPS, interpolation is first attempted using 16 surrounding grid cells in the coarser grid; if this method fails due to a lack of the requisite 16 valid data points, WPS attempts other interpolation techniques using as many as four grid cells and as few as one. While a full description of all WPS interpolation techniques is beyond the scope of this study, more information is available in the WRF User's Guide (2014, p.

[3-56 to 3-59](#)). When all other methods fail due to the lack of nearby water grid cells, WPS defaults to the “search” approach, in which the nearest water point is used to set LSTs (~~WRF User’s Guide, 2014~~). When employing the search option, water cells in the driving data are often distant from and unrepresentative of the target cell in the WRF domain. The search option in WPS performs no interpolation or averaging, sometimes resulting in abrupt, non-physical temperature discontinuities.

Here we show the result of using this default methodology to downscale 1° Community Earth System Model (CESM) fields to a 36 km WRF domain ( $198 \times 126$ ) covering the continental US, and subsequently similar examples in other downscaling studies are discussed. However, it should be noted that the use of CESM as an example is arbitrary because similar results have been obtained with other global datasets as well. The CESM ocean mask, used to interpolate the GCM’s SST fields to the WRF grid, has no water grid cells over the North American interior (Fig. 1). As a result, water temperatures in Hudson Bay are used to set temperatures over the larger westernmost areas of the Laurentian Great Lakes, while LSTs in the southeastern areas of the Great Lakes are set by Atlantic SSTs (Fig. 2). At the time shown in Fig. 2, the LSTs interpolated from CESM onto the 36 km WRF grid contain discontinuities of approximately 17 K between adjacent grid cells in Lakes Michigan and Huron, while a smaller discontinuity of approximately 3 K is created in Lake Superior. It should be noted that various interpolation options are available in WPS and can be specified by the user. The description in the paragraph above is representative of the interpolation process as defined by WPS’s default settings. Even though this process could be changed by the model user, the key issue remains that when lakes are poorly represented or completely absent, the problem of how to specify the lake state is not amenable to any interpolation method.

The problems of using larger-scale data to define LSTs with the default options in WPS are not limited to the Great Lakes. None of the inland lakes resolved by WRF at 36 km have valid LSTs in the CESM ocean mask (Fig. 1). Using the search option in WPS results in setting the LSTs to unrealistic values throughout the domain. Temperatures in Pyramid Lake, Great Salt Lake, as well as several smaller lakes east of the Rocky Mountains in both Canada and the US are assigned from the Pacific Ocean (Fig. 2), while lake temperatures in the southeastern and central

US are set from SSTs in the Gulf of Mexico and Atlantic Ocean. Two adjacent grid cells representing Lake Sakakawea in North Dakota are assigned LSTs differing by approximately 10 K because the western cell is set from the Pacific while the eastern cell is prescribed from Hudson Bay (Fig. 2). Using any interpolation method to assign LSTs when no suitable data are available will adversely affect the accuracy of downscaled simulations that are based on forcing from those LSTs.

Mallard et al. (2014; hereafter M14) also discuss problems that arise when downscaling coarse global data to a 12 km grid covering the eastern US. In M14, the National Centers for Environmental Prediction (NCEP)–Department of Energy Atmospheric Model Intercomparison Project (AMIP-II) reanalysis (hereafter R2; Kanamitsu et al., 2002) is used to drive historical simulations as a proxy or stand-in for a similarly-coarse GCM. In contrast to the CESM example discussed above, R2 has at least a partial representation of western Great Lakes, but nevertheless has only three inland water points to represent all five of the Great Lakes (Fig. 1 of M14). Therefore, using the standard interpolation methods with R2 results in unrealistically large, abrupt, and non-physical LST discontinuities in eastern Lake Erie and Lake Ontario, where water temperatures are set using Atlantic SSTs, while the LSTs in western Lake Erie and in the three western Great Lakes are interpolated from the three lake cells in R2 (M14).

In WRF, ice cover can either ~~be treated as a binary field, which is set based on whether the water temperature is below a threshold, or it can~~ be interpolated from the driving data and ~~prescribed as assigned to~~ covering some fraction of a grid cell, ~~or it can be treated as a binary field that is set~~. ~~In the former approach, WRF sets ice cover to 100 % at grid points-cells where the water surface temperatureLST drops below a specified threshold.271 K, slightly below the freshwater freezing temperature of approximately 273 K. Note that t~~ The default threshold value was ~~changed from~~ 271 K (slightly below the freshwater freezing temperature of approximately 273 K), but it was ~~changed~~ to 100 K as of version 3.5.1 (September 2013), ~~presumably~~ to avoid the unintended creation of ice by this method when using WRF's default settings (Table 1). When fractional ice values are prescribed from the driving dataset, the WPS methods applied to interpolate sea and lake ice differ from those used for SSTs and LSTs. If there are no surrounding water grid cells in the driving dataset, an ice cover value of zero is assigned rather than employing the search

method. When M14 downscaled ice cover from R2, it was shown ice concentrations of zero were applied to points through Lakes Huron, Erie and Ontario throughout a two-year simulation (Fig. 3 of M14), even though partial ice coverage was observed on all three lakes during that historical period. Moreover, almost complete ice coverage of Lakes Superior and Michigan occurred in a single day (M14). Wang et al. (2012a) conducted a climatology of ice cover in the Great Lakes over the period 1973 to 2010 and showed that, in the average seasonal cycle of ice cover, the maximum fractional coverage of Lake Superior was approximately 50 % (their Fig. 3). Although Wang et al. noted that the standard deviation of ice cover is quite large (exceeding the mean values in some of the Great Lakes), the seasonal cycles in their study showed the accumulation of ice coverage over months, not the abrupt appearance of lake-wide ice over daily periods. Ultimately, M14 improved the representation of the Great Lakes in their downscaled simulations by applying a coupled lake model, which will be discussed further in a subsequent section. Whereas M14 showed the results of using a single lake model, the current work presents a broader range of approaches, recognizing that the most preferable method to represent lake fields may vary between different RCM applications.

Prior studies downscaling other global datasets and GCMs have also noted findings similar to the example shown here (Fig. 2) and the results of M14. Using WRF as an RCM over Eastern Africa, Argent (2014) showed that the use of WPS's default interpolation methods resulted in oceanic temperatures from a global SST dataset applied to set LSTs throughout Lake Victoria. Discontinuities in LSTs with WRF were noted in the Great Lakes basin by Bullock et al. (2014) who downscaled R2 to 12 km, and by Gao et al. (2012) who downscaled the CESM to a 4 km grid. Within the downscaled simulations produced for the North American Regional Climate Change Assessment Program (NARCCAP; Mearns et al., 2012), problems with producing realistic LSTs and ice cover for the Great Lakes region are documented using several approaches with various RCMs, including WRF (NARCCAP, 2014). For some NARCCAP model configurations, caution is recommended when using surface variables in the region surrounding the Great Lakes. Previous work examining the value of dynamical downscaling has noted that downscaled simulations have the most potential to add value relative to GCM simulations in areas of complex topography and along coastlines because of increased resolution in regional models (e.g., Feser et al., 2011). Although RCMs better resolve the coastlines (and therefore, the

presence of lakes) than the driving GCMs, using erroneous LSTs and lake ice cover could impair the simulation of interactions between lakes and overlying air masses. The potential benefits gained by downscaling to a grid spacing that better resolves land–water interfaces may not be realized if the lake state (defined here by LSTs and ice) is unrealistically represented. Even as additional computing resources allow GCMs to increase in resolution and better represent lakes, RCMs will also be run at finer scales; therefore, it can be expected that smaller lakes with important effects on mesoscale and microscale climatology will continue to be unresolved by the driving data sets.

The purposes of this paper are to describe various techniques that can be used to set LSTs and lake ice cover in the WRF model for downscaling, and to discuss the benefits and possible shortcomings of each approach. The effects of these techniques on simulated lake–atmosphere interactions, both in the present climate and in future climate states, are discussed in context with relevant previous literature.

## **2 Comparison of methods**

As will be shown below, choice of the appropriate methodology for representing a lake in a downscaling configuration is dependent on what interactions must be simulated between the atmospheric fields and the lake state and how the lake state is expected to be impacted by climate change when downscaling future GCM projections. In regional climate simulations conducted over the continental US, the Laurentian Great Lakes are a prominent feature, as Lake Superior is the largest freshwater lake in the world (by surface area) at over 82 000 km<sup>2</sup>. Several studies have concluded that the Great Lakes strongly influence the surrounding regional climate, moderating extremes in near-surface temperatures, and affecting precipitation and passing cyclones and anticyclones on an annual cycle (e.g., Wilson, 1977; Bates et al., 1993; Scott and Huff, 1996; Notaro et al., 2013). Climatologically, the greater heat capacity of the lakes serves to enhance precipitation and convection during September to March, when warmer surface water (relative to low-level atmospheric temperatures) reduces atmospheric stability (e.g., Notaro et al., 2013). Conversely, the slower warming of the lakes in boreal spring results in the opposite effect during the April–August period, where the relatively cool lakes enhance atmospheric stability and reduce precipitation and convection. These periods are referred to as the lake unstable and

lake stable seasons, respectively. Lake-effect precipitation has also been documented outside the Great Lakes as well, such as in Lake Champlain (Tardy, 2000; Laird et al., 2009), Lake Tahoe (Cairns et al., 2001), and the Great Salt Lake (Carpenter, 1993; Steenburgh and Onton, 2001). A review by Schultz et al. (2004) states that lake-effect snowfall has been observed to occur over lakes with fetches of only 30 to 50 km, citing prior studies over Bull Shoals Lake of Arkansas (Wilken, 1997) as well as Lake Tahoe and Pyramid Lake in Nevada (Cairns et al., 2001; Huggins et al., 2001). Interactions between the lakes and surrounding regions are also strong in tropical environments as well. For example, the immediate region surrounding Lake Victoria in Africa has the highest recorded frequency of thunderstorms in the world with approximately 300 storm days per year (Asnani, 1993). Overall, while a comprehensive review of the impact of each lake on regional climate is beyond the scope of this study, prior work indicates that even lakes that are smaller than the Great Lakes can be anticipated to have substantial effects on regional climate.

Prior studies have also illustrated that even relatively small errors in prescribed LSTs in a downscaling configuration can adversely affect simulated precipitation in regions surrounding lakes. The sensitivity study of Wright et al. (2013) showed significant changes in lake-effect snowfall over the Great Lakes in idealized simulations where LSTs were uniformly warmed by 3 °C. Anyah and Semazzi (2004) simulated changes in the spatial patterns and intensity of precipitation, as well as the amount of evaporation, over Lake Victoria in a modeling study where LSTs were uniformly changed by only 1.5 °C.

Interactions between the lakes and overlying air masses are also governed by the amount of lake ice in climates that permit lakes to freeze. Previous studies have found the presence of ice suppresses turbulent latent and sensible heat fluxes from the lake to the air mass (e.g., Zulauf and Krueger, 2003; Gerbush et al., 2008). As shown in the lake-effect snow case studies simulated by Wright et al. (2013), the presence of ice coverage over the lake's surface inhibits downstream precipitation. As a result, lake-effect snowfall decreases in some areas surrounding the Great Lakes during the later portion of the lake unstable season, as the water's surface freezes during the winter and early spring months. Overall, past studies indicate that if LSTs and ice are not properly prescribed, inaccurate values of precipitation and temperature in the lee of lakes result in a downscaled simulation.

215

## 216 **2.1 WRF's alternative lake setting**

217 Since the release of WRF version 3.3 in April 2011, an “alternative initialization of lake SSTs”  
218 | option is provided in WPS to set LSTs (WRF User’s Guide, 2014; [Table 1](#)). When employing  
219 this method, LSTs can be set using temporally averaged 2 m air temperatures from the driving  
220 data set, with the averaging period set by the user. Bullock et al. (2014), when downscaling a  
221 proxy GCM (R2) over a 12 km grid covering the Great Lakes, attempted to use the alternative  
222 lake setting to account for the greater thermal inertia of the Great Lakes by incorporating  
223 seasonal temperature changes after a one-month time lag. Following the procedure of Bullock et  
224 al., if a user were to perform a simulation over the month of May, a single LST field would first  
225 be generated by temporally averaging air temperatures during the previous month of April;  
226 subsequently this static LST field would be used to set inland water temperatures throughout the  
227 month of May. Because Bullock et al. (2014) preprocessed the driving data in monthly segments,  
228 the LST field was prescribed to vary with time on a monthly basis. Using this method may  
229 imitate the seasonal changes observed over the Great Lakes, producing a lake stable and unstable  
230 season during the appropriate months. A drawback to this methodology is that the same lag time  
231 is used throughout the model grid, regardless of lake depth. Therefore, in this approach, large,  
232 deep lakes are implied to heat and cool on the same timescale as small, shallow lakes.  
233 Meanwhile, it is expected that observed seasonal temperature changes over smaller and  
234 shallower lakes would more closely follow atmospheric temperature changes than in large, deep  
235 lakes. If employed for simulations outside the Great Lakes, the procedure used by Bullock et al.  
236 (2014) should be modified to imitate the observed relationship between changing air  
237 temperatures and LSTs.

238

239 | In its default configuration used prior to the release of version 3.5.1 (~~September 2013~~), WRF  
240 | prescribes ice cover at grid cells where LST is less than 271 K ([Table 1](#)). This value is applied at  
241 all water points regardless of salinity. As winter 2 m air temperatures are frequently below  
242 freezing in the Great Lakes area, Bullock et al. (2014) found that unrealistically large spatial  
243 coverage of ice occurred when using the alternative lake setting in WRF version 3.4.1, with all  
244 five Great Lakes completely frozen for most of the winter. Such erroneous ice cover would be



expected to negatively impact the simulation of precipitation, 2 m temperatures, and other variables influenced by sensible and latent heat fluxes supplied by the Great Lakes. Therefore, the use of the alternative lake setting in WRF may not be appropriate in some regions where sub-freezing air temperatures would result in unrealistic temporal and spatial coverage of sub-freezing LSTs and ice.

However, this is not a concern for tropical lakes where air temperatures would not be sufficiently low enough to result in frozen lakes. Argent (2014, Sect. 3) demonstrated the utility of the alternative lake setting in WRF simulations over Lake Victoria in Eastern Africa, finding that it improved the accuracy of simulated rainfall relative to the use of the default interpolation in which oceanic SSTs were used to set Lake Victoria's LSTs.

## 2.2 Climatological LSTs and ice

Another approach for setting LSTs and lake ice coverage when downscaling with WRF is to prescribe these variables from higher-resolution data sets of climatologically averaged quantities. This can be viewed as assuming stationarity for the lake state as is frequently done for other input variables in an RCM, such as land-use and vegetation. Even for retrospective climate simulations, using this approach could be detrimental because the interannual variability of LSTs and ice – and its effects on the prediction of extreme events – would not be captured using this method. When making future projections, it must be considered that prior studies have shown that LSTs cannot be assumed to be stationary in future warmer climates; in fact, some studies conclude that non-linear feedbacks exist between regional climate change and LSTs and ice for some lakes. An observational study by Austin and Colman (2007) found that the multi-decadal warming trend in the Great Lakes region was amplified in the lake temperatures, relative to surrounding inland temperatures, because of the earlier break-up of ice and earlier springtime warming of surface water. In the downscaling simulations of Gula and Peltier (2012), increased snowfall was simulated in the lee of the Great Lakes in a warmer, mid-century climate because lake ice forms ~~at a later time period~~ in the winter. Gula and Peltier conclude that the impact of having the lakes remain ~~open (and free of ice)~~ is that increased latent and sensible heat fluxes are present for a longer time period during the lake unstable season, lessening the stability of the

overlying air mass and enhancing precipitation. Magnuson (2000) concluded that observed ice coverage is decreasing in lakes and rivers throughout the Northern Hemisphere. Such a decrease in ice coverage has been linked by observational studies to increases in lake-effect precipitation in the Great Lakes region (Assel and Robertson, 1995; Burnett et al., 2003; Kunkel et al., 2009). Because ice ~~supresses~~suppresses fluxes of latent and sensible heat (e.g., Zulauf and Krueger, 2003; Gerbush et al., 2008), decreasing ice cover in a warmer climate allows larger fluxes of latent and sensible heat to modify the overlying air mass, increasing downstream precipitation during the lake unstable season. None of the impacts on the lake state reviewed here (the warming of LSTs and more open water from which to produce fluxes) would be considered in the WRF model using LSTs and ice based on present-day climatology, and the effects of changing lake conditions on atmospheric stability, humidity, precipitation and convection would not be simulated.

This approach could be improved by adding a linear increase to observed LSTs over time, which may be a valid approximation for the effect of climate change on some lakes. However, such an approach would not capture the non-linear impacts of climate change (as described by Austin and Colman, 2007) on the Great Lakes. Overall, the efficacy of using of a climatologically-based approach is dependent on the amount of interannual variability, as well as the impacts of climate change on the lake state and whether those effects can be accounted for by the inclusion of a linear LST anomaly.

## **2.3 Land mask modification**

To avoid the issues with LSTs discussed in Sect. 1 and illustrated in Fig. 2, Gao et al. (2012) modified the GCM land mask in the Great Lakes area so that skin temperatures from land points in the GCM were used to set LSTs on the WRF grid in their downscaled simulations. This treatment successfully eliminated the abrupt temperature discontinuities (such as those in Fig. 2) produced by interpolating a coarse data set. However, the effects of the lakes themselves are lost if GCM land temperatures are used to prescribe RCM water temperatures and the lake-land temperature contrasts, with their associated mesoscale phenomena such as lake breezes and lake-effect precipitation, are eliminated. Notaro et al. (2013) conducted an idealized modeling

experiment where the Great Lakes were replaced with forest and field land cover types. They found that the presence of the lakes affected precipitation, 2 m air temperatures and their variability, water vapor, cloud cover, incoming shortwave radiation, the hydrological budget and the intensity of passing cyclones and anticyclones. The approach used by Gao et al. (2012), where land surface temperatures from the GCM are used to specify water temperatures, partially accounts for some lake effects (such as changes in surface friction and albedo) because WRF would recognize the presence of a water surface. However, all processes related to the LST (e.g., ice formation, latent and sensible heat flux, 2 m temperature and moisture values, outgoing longwave radiation from the surface) would be negatively impacted by this treatment. Additionally, some impacts of climate change on the future lake state could be lost. For example, the amplification of Great Lakes LSTs, relative to over-land temperatures, observed by Austin and Colman (2007) will not be captured if land temperatures are used to set LSTs.

## 2.4 Use of simulated lake fields from GCM

A more sophisticated class of approaches for better representing the lake state in a downscaling configuration involves the use of a lake model. This can be done either by using outputs from the GCM's lake model (if available), driving a stand-alone lake model offline with GCM fields to simulate LSTs and ice, or by coupling a lake model to the RCM when downscaling. The CESM ~~model~~ has a lake model embedded within ~~the-its~~ land ~~surfacecomponent~~ model (LSM), version 4 of the Community Land Model (CLM4). CLM4 accounts for the presence of subgrid-scale lakes using the one-dimensional lake model described in Oleson et al. (2010). It is a column model partially based on the Hostetler lake model (e.g., Hostetler and Bartlein, 1990; Hostetler et al., 1993, 1994), and it simulates 10 water layers through the depth of the lake, as well as additional layers for thermally-active soil underneath and snow and ice above. However, when producing the downscaled simulation shown in Fig. 2, output from CLM's lake model was not easily accessible with other CESM outputs from the same simulation within archiving systems such as the Earth System Grid Federation. Lake temperatures and ice from CESM, and other GCMs with embedded lake models, could be leveraged by RCMs such as WRF to account for the impact of climate change on the lake state. In areas where lakes are at least partially resolved by the GCM, this approach would be effective at driving the RCM with simulated changes in LSTs and ice

cover consistent with future projections and at keeping the RCM solution in the regions affected by lakes consistent with the GCM simulation. However, some small lakes may remain unrepresented by GCM data.

## 2.5 Use of a stand-alone lake model

If lake model outputs from the GCM are unavailable, one alternative is to use a standalone lake model driven by GCM fields to downscale the lake state in a manner which is consistent with the GCM's atmospheric fields. In the downscaling experiments performed by Gula and Peltier (2012) over the period 2050–2060, the Freshwater Lake (FLake) model was utilized to provide simulated LSTs and lake ice to WRF in the Great Lakes basin. GCM fields from the Community Climate System Model, with a spectral resolution of T85 (~ 1.4° grid spacing), were used to drive a FLake simulation on a 10 km regional grid, and the LSTs and ice cover simulated by FLake were subsequently used to drive the downscaled WRF simulation. In this 1-way WRF-FLake model configuration, changes in LSTs and ice respond to changes in atmospheric variables in the driving GCM, but the lake model output is produced on the higher-resolution regional WRF grid. FLake is a 1-D column model which is highly reliant on empirical relationships and has been used in several studies with other RCMs (e.g., Mironov, 2008; Kourzeneva et al., 2008; Martynov et al., 2008; Mironov et al., 2010; Samuelsson et al., 2010). FLake requires a 2-D field of lake depths and the 1-D column model is called at each point. Therefore, the simulated LSTs are sensitive to lake depth, as well as the driving GCM fields.

## 2.6 Use of a coupled lake model within an RCM

In WRF version 3.6, ~~released April 2014~~, a CLM-based lake model can be utilized with other non-CLM land surface models (WRF User's Guide, 2014; Table 1). This lake model is taken from CLM version 4.5 (Subin et al., 2012; Oleson et al., 2013) with some modifications by Gu et al. (2013) as discussed further below. Although a version of CLM4 was available as an LSM ~~land surface model~~ option within WRF version 3.5 (~~released April 2013~~), the lake model in CLM4 was disabled in WRF (Table 1). In WRF version 3.6, CLM's Hostetler-based lake model can be applied by using horizontally varying lake depths (which are available in WPS version 3.6) or a

uniform lake depth can be assigned to all lakes at runtime. Gu et al. (2013) demonstrated WRF-CLM's performance in the Great Lakes region using a previous version of this model configuration (WRF 3.2 and CLM 3.5) to simulate a 16 month period from 2001 to 2002 at 10 km grid spacing. It was shown that the lake model simulated LSTs well in Lake Erie but generated large biases in LSTs when compared to buoy observations in Lake Superior. However, the LST bias was reduced by reformulating the eddy diffusivity parameter in the CLM lake model, and it was concluded that the updated lake model within WRF-CLM was reasonably able to reproduce observed LSTs. However, no ice was observed during the period and the ability of WRF-CLM to accurately simulate ice cover was not examined in Gu et al. (2013).

In an alternative coupled approach, the prior work of Gula and Peltier (2012) has been updated with the option of using WRF-FLake as a 2-way coupled model, where atmospheric variables simulated by WRF are used by FLake at each time step in the WRF model, and simulated LSTs and ice thicknesses are provided back to WRF by FLake. M14 concluded that the use of WRF-FLake resulted in a more accurate representation of LSTs and lake ice, relative to interpolation from ~~a proxy GCM~~, the R2. Substantial improvements were shown in the simulation of the temporal and spatial variability of ice cover, and errors in LSTs were reduced by the use of the coupled model. Similar to Martynov et al. (2010), M14 found that FLake performed worst in the largest and deepest lake (Lake Superior) and best for the smallest and shallowest (Lake Erie).

When using an embedded lake model within an RCM, it can be anticipated that the period of time needed for spin-up could be larger than it is when all water conditions are simply prescribed. To spin-up the WRF-FLake model in M14, the stand-alone version of the FLake model was driven with atmospheric conditions from the proxy GCM ~~for 10 annual cycles to achieve equilibrium, as adapted from the~~ in a spin-up procedure recommended by Mironov et al. (2010) when using FLake. In this methodology, the initial year of the simulation ~~(2005)~~ is “looped” over 10 annual cycles with meteorological variables from the initial year repeatedly used to force the lake model, and the lake state at the end of each year used to initialize FLake for the start of the next year ~~until~~, ensuring that the simulated lake state converges to equilibrium with these atmospheric conditions ~~is achieved by the end of the 10-cycle simulation~~. Output from the first year of this offline simulation is shown in Fig. 3 illustrating the adverse effects of using

FLake output without adequate spin-up time. A time-series taken from a representative point in Lake Superior shows unrealistically cool LSTs (below 200 K) occurring during the initial months of the simulation. Also during this period, unrealistically large ice coverage formed, freezing over all five Great Lakes. The observed ice cover plotted in Fig. 3 is much more limited in its spatial extent. ~~as shown~~ Observed ice cover is plotted from the National Ice Center (NIC) Great Lakes Ice Analysis charts, which are processed and provided by the Great Lakes Environmental Research Laboratory (GLERL; Wang et al., 2012b). The FLake model results obtained after the spin-up period showed realistic values of LSTs and ice cover (M14).

To examine how WRF-CLM reacts during the initial months of a simulation, without any spin-up time, output from a 12 km WRF-CLM simulation (version 3.6) is shown in Fig. 4. In this simulation, the same methods as in M14 are followed but with the following changes: the model version is updated from 3.4.1 to 3.6, the CLM lake model is used in place of FLake, and no spin-up procedure is employed for initialization of the lake model (initial LSTs are interpolated from R2). As in M14, the Noah ~~LSM land surface model~~ (Chen and Dudhia, 2001) is used. Similar to the example shown in Fig. 3, significant overestimation of ice coverage occurs during the first year (Fig. 4). Although some adverse effects in this simulation are introduced due to the use of LSTs interpolated from the coarse R2 data to provide an initial state, the similarity of these results to FLake's fields in Fig. 3 suggests that the lack of spin-up time is a common problem to both model runs. It is also implied by the methodologies of other CLM-based studies, which do use spin-up or initialization procedures. Previous work by Subin et al. (2012) with the lake model in CLM4 used a 110-year period for the spin-up of their reference simulation. In their experiments with WRF-CLM, Gu et al. (2013) used an observed LST field for initialization. The 9 sub-surface layers in their model were initialized based on the shape of an observed profile of lake temperatures, valid during that period of the year and taken from Lake Superior. Using this initialization methodology for a future downscaled simulation is not possible due to lack of observations, but simulated future lake profiles could possibly be utilized for initialization of downscaled runs. Overall, when using an embedded lake model in a downscaling application, users should consider how the lake model is being initialized or spun-up in order to achieve results with accuracy similar to the prior studies discussed above. If the lake state is initially poorly prescribed from the GCM (with results similar to those shown in Fig. 2), a protracted

spin-up could be required to reach equilibrium with the driving fields in the RCM and obtain more realistic results.

It has been noted previously that both WRF-FLake and WRF-CLM, as well as other 1-D lake models, tend to exhibit difficulty in simulating deep lakes (e.g., Martynov et al., 2010; Stepanenko et al., 2010; Gu et al., 2013; M14). Some model error can be attributed to the fact that one-dimensional column models cannot represent 2- and 3-D processes (e.g., currents, drifting ice, and formation of a thermal bar). While more sophisticated lake models could be coupled with WRF, using computationally efficient 1-D models is advantageous in downscaling applications, where computational resources are taxed by the use of finer resolution. Additionally, Martynov et al. (2010) noted that more complex 3-D lake models are generally run with much finer grid spacing ( $\sim 2$  km) than typical RCMs. Martynov et al. (2010) also compared the simulated water temperatures and ice coverage from the Hostetler and FLake models, finding that FLake generally performed better, but that the Hostetler model provides more opportunity to improve model performance because it utilizes more vertical layers and is less reliant on parameterization. A comparison of 1-D lake models by Thiery et al. (2014) showed favorable results for both FLake and Hostetler-based models (including the lake model found in CLM4) and noted their computational efficiency. When making regional climate projections with these models it should be noted that both WRF-FLake and WRF-CLM assume that lake depths are constant in time, which could be a poor assumption depending on the lake being modeled and the future period. Also, more complex lake models may be appropriate for higher resolution ( $\sim 2$  km grid spacing) RCM simulations focused on regions where lake dynamics are not adequately captured by the column lake models discussed here.

### 3 Conclusion

It has been shown in the present study and in previous work (e.g., Gao et al., 2012; Bullock et al., 2014; M14) that downscaling typically-coarse GCM data, using WRF's default interpolation methods, to finer resolution WRF grids results in LST discontinuities and spurious ice formation in the Great Lakes (Fig. 2). Although the default interpolation methods in WRF can easily be modified to alter the interpolation scheme or to eliminate the search option, none of these simple changes will overcome the challenges of setting the LSTs for inland water bodies that are not



resolved by driving data when WRF is used as a RCM. Various alternate methods have been presented, and a summary of the positives and potential drawbacks to each approach is shown in Table 21. Using WRF's "alternative" lake setting instead of the default interpolation method in WPS eliminates unrealistically large and abrupt spatial discontinuities in temperature, but causes large, deep lakes (such as Lake Superior) to erroneously freeze when ice is set based on an air-temperature threshold. All the other approaches discussed above can simulate more realistic ice cover than the default interpolation. However, the simulation of ice cover is obviously not a factor in downscaling studies where the environment does not become sufficiently cold to produce lake ice, such as those focusing on tropical regions. For example, the alternative lake setting has been used to improve rainfall results (relative to the use of WRF's default interpolation techniques) over Lake Victoria in Eastern Africa by Argent (2014). Using climatological values in a future warmer climate will adversely affect results because LSTs cannot be assumed to be stationary over time. A warming trend could be applied to observed LST fields in order to improve this approach; however, a realistic trend may be complex to derive for some lakes as Austin and Colman (2007) have shown an observed non-linear amplification of warming LSTs relative to inland temperatures in the Great Lakes region. The land mask alteration method of Gao et al. (2012) is effective at preventing discontinuities in surface temperatures, but the use of temperatures from land grid cells in the GCM to set LSTs in the RCM eliminates the presence of land-lake temperature contrasts which impact precipitation, winds (i.e. land-sea breeze), and other near-surface fields. The use of a lake model (either coupling a lake model to the RCM or using outputs from the GCM's lake model to drive the RCM) can improve the representation of the lakes in retrospective simulations and has the ability to simulate non-linear impacts of climate change on LSTs and ice cover (e.g., Gula and Peltier, 2012, M14).

For downscaling applications using WRF, we recommend setting LSTs and ice cover from either a RCM- or GCM-driven lake model, especially when simulating mid-latitude regions. In their studies focused on the Great Lakes, Notaro et al. (2013) and Wright et al. (2013) state that accurate predictions of changes in LSTs and ice cover from lake models are needed when simulating changes in regional climate. Zhao et al. (2012) also recommended the use of a lake model for simulating changes in regional precipitation in the Great Lakes basin. Including



prognostic changes in the lake state is also possible if GCM data sets include predicted lake surface temperatures and ice within their publicly-available outputs. For regional climate modeling efforts in which the RCM data is being archived for various end-user applications, we recommend the use of GCM- or RCM-driven lake modeling approaches. If such an approach is not used, the potential adverse effects of setting LSTs and ice cover using interpolation from the GCM should be documented, as is currently done in NARCCAP (2014).

The accuracy of the various approaches presented here is sensitive to the characteristics of the lakes to which they are being applied. Approaches which set LSTs as a function of over-land temperatures (such as the land mask modification approach or WRF's alternative lake setting) may perform adequately when applied to smaller, shallower lakes where LST changes are more closely coupled to air temperature changes. Investigators performing RCM experiments should consider both the present-day interactions between the lake and overlying air masses as well as the potential climate change impacts on the lakes within their model domain when choosing an approach.

#### **4 Code availability**

WPS and the WRF model can be downloaded from <http://www2.mmm.ucar.edu/wrf/users/downloads.html>. Source code for the FLake model can be obtained at <http://www.flake.igb-berlin.de/sourcecodes.shtml>, and code needed to run the coupled WRF-FLake model is available for download at <http://web.atmos.ucla.edu/~gula/wrfflake>.

#### **Acknowledgements**

The lead author was supported by an appointment to the Research Participation Program at the US Environmental Protection Agency ([EPA](#)) Office of Research and Development, administered by the Oak Ridge Institute for Science and Education (ORISE). The WRF model is made available by NCAR, funded by the National Science Foundation. [Great Lakes ice cover data were obtained from GLERL \(<http://www.glerl.noaa.gov/data/pgs/glice/glice.html>\)](#). This research has been subjected to the US EPA's administrative review and approved for publication. The

517 views expressed and the contents are solely the responsibility of the authors, and do not  
518 necessarily represent the official views of the US EPA.

## References

- Anyah, R. O. and Semazzi, F. H. M.: Simulation of the sensitivity of Lake Victoria basin climate to lake surface temperatures, *Theor. Appl. Climatol.*, 79, 55–69, 2004.
- Argent, R. E.: Customisation of the WRF model over the Lake Victoria basin in east Africa, M.S. thesis, North Carolina State University, Raleigh, NC, 124 pp., 2014.
- Artale, V., Calmanti, S., Carillo, A., Dell’Aquila, A., Herrmann, M., Pisacane, G., Ruti, P. M., Sannino, G., Struglia, M. V., Giorgi, F., Bi, X., Pal, J. S., Rauscher, S., and The PROTHEUS Group: An atmosphere–ocean regional climate model for the Mediterranean area: assessment of a present climate simulation, *Clim. Dynam.*, 35, 721–740, 2010.
- Asnani, G. C.: Tropical Meteorology, Vol. 1 and 2, Indian Institute of Tropical Meteorology Pashan, Pune, 1012 pp., 1993.
- Assel, R. A. and Robertson, D. M.: Changes in winter air temperatures near Lake Michigan, 1851–1993, as determined from regional lake-ice records, *Limnol. Oceanogr.*, 40, 165–176, 1995.
- Austin, J. and Colman, S.: Lake Superior summer water temperatures are increasing more rapidly than regional air temperatures: A positive ice–albedo feedback, *Geophys. Res. Lett.*, 34, L06604, doi:10.1029/2006GL029021, 2007.
- Bates, G. T., Giorgi, F., and Hostetler, S. W.: Toward the simulation of the effects of the Great Lakes on regional climate, *Mon. Weather Rev.*, 123, 1505–1522, 1993.
- Bullock, O. R., Alapaty, K., Herwehe, J. A., Mallard, M. S., Otte, T. L., Gilliam, R. C., and Nolte, C. G.: An observation-based investigation of nudging in WRF for downscaling surface climate information to 12 km grid spacing, *J. Appl. Meteorol. Clim.*, 53, 20–33, 2014.

Burnett, A. W., Kirby, M. E., Mullins, H. T., and Patterson, W. P.: Increasing Great Lake–effect snowfall during the twentieth century: a regional response to global warming?, *J. Climate*, 16, 3535–3542, 2003.

Cairns, M. M., Collins, R., Cylke, T., Deutschendorf, M., and Mercer, D.: A lake effect snowfall in western Nevada. Part I: Synoptic setting and observations, *Preprints, 18th Conf. on Weather Analysis and Forecasting*, Fort Lauderdale, FL, 29 July 2001, *Am. Meteor. Soc.*, 329–332, 2001.

Carpenter, D. M.: The lake effect of the Great Salt Lake: overview and forecast problems, *Weather Forecast.*, 8, 181–193, 1993.

Chen, F. and Dudhia, J.: Coupling and advanced land surface–hydrology model with the Penn State–NCAR MM5 modeling system, Part I: Model implementation and sensitivity, *Mon. Weather Rev.*, 129, 569–585, 2001.

Feser, F., Rockel, B., von Storch, H., Winterfeldt, J., and Zahn, M.: Regional climate models add value to global model data: a review and selected examples. *B. Am. Meteorol. Soc.*, 92, 1181–1192, 2011.

Gao, Y., Fu, J. S., Drake, J. B., Liu, Y., and Lamarque, J.-F.: Projected changes of extreme weather events in the eastern United States based on a high resolution climate modeling system, *Environ. Res. Lett.*, 7, 044025, doi:10.1088/1748-9326/7/4/044025, 2012.

Gerbush, M. R., Kristovich, D. A. R., and Laird, N. F.: Mesoscale boundary layer and heat flux variations over pack ice–covered Lake Erie, *J. Appl. Meteorol. Clim.*, 47, 668–682, 2008.

Gu, H., Jin, J., Wu, Y., Ek, M., and Subin, Z.: Calibration and validation of lake surface temperature simulations with the coupled WRF-lake model, *Clim. Chang.*, doi: 10.1007/s10584-013-0978-y, in press, 2013.

Gualdi, S., Somot, S., Li, L., Artale, V., Adani, M., Bellucci, A., Braun, A., Calmanti, S., Carillo, A., Dell'Aquila, A., Déqué, M., Dubois, C., Elizalde, A., Harzallah, A., Jacob, D., L'Hévéder, B., May, W., Oddo, P., Ruti, P., Sanna, A., Sannino, G., Scoccimarro, E., Sevault, F., and Navarra, A.: The CIRCE simulations: Regional climate change projections with realistic representation of the Mediterranean Sea, *B. Am. Meteorol. Soc.*, 94, 65–81, 2013.

Gula, J. and Peltier, W. R.: Dynamical downscaling over the Great Lakes basin of North America using the WRF regional climate model: the impact of the Great Lakes system on regional greenhouse warming, *J. Climate*, 25, 7723–7742, 2012.

Hostetler, S. W. and Bartlein, P. J.: Simulation of lake evaporation with application to modeling lake level variations of Harney-Malheur Lake, Oregon, *Water Resour. Res.*, 26, 2603–2612, 1990.

Hostetler, S. W., Bates, G. T., and Giorgi, F.: Interactive coupling of a lake thermal model with a regional climate model, *J. Geophys. Res.*, 98, 5045–5057, 1993.

Hostetler, S. W., Giorgi, F., Bates, G. T., and Bartlein, P. J.: Lake-atmosphere feedbacks associated with paleolakes Bonneville and Lahontan, *Science*, 263, 665–668, 1994.

Huggins, A. W., Kingsmill, D. E., and Cairns, M. M.: A lake effect snowfall in western Nevada—Part II: Radar characteristics and quantitative precipitation estimates, *Preprints, 18th Conf. on Weather Analysis and Forecasting*, Fort Lauderdale, FL, 29 July 2001, *Amer. Meteor. Soc.*, 333–337, 2001.

Kanamitsu, M., Ebisuzaki, W., Woollen, J., Yang, S.-K., Hnilo, J. J., Fiorino, M., and Potter, G. L.: NCEP–DOE AMIP-II Reanalysis (R-2). *B. Am. Meteorol. Soc.*, 83, 1631–1643, 2002.

Kourzeneva, E., Samuelsson, P., Ganbat, G., and Mironov, D.: Implementation of Lake Model FLake into HIRLAM, *HIRLAM Newsletter* 54, 54–64, available from HIRLAM-A Programme,

c/o J. Onvlee, KNMI, P. O. Box 201, 3730 AE De Bilt, the Netherlands, available at: <http://hirlam.org> (last access: 20 October 2014), 2008.

Kunkel, K. E., Ensor, L., Palecki, M., Easterling, D., Robinson, D., Hubbard, K. G., and Redmond, K.: A new look at lake-effect snowfall trends in the Laurentian Great Lakes using a temporally homogeneous data set, *J. Great Lakes Res.*, 35, 23–29, 2009.

Laird, N. F., Desrochers, J., and Payer, M.: Climatology of lake-effect precipitation events over Lake Champlain, *J. Appl. Meteorol. Clim.*, 48, 232–250, 2009.

Magnuson, J. J., Robertson, D. M., Benson, B. J., Wynne, R. H., Livingstone, D. M., Arai, T., Assel, R. A., Barry, R. G., Card, V., Kuusisto, E., Granin, N. G., Prowse, T. D., Stewart, K. M., and Vuglinski, V. S.: Historical trends in lake and river ice cover in the Northern Hemisphere, *Science*, 289, 1743–1746, 2000.

Mallard, M. S., Nolte, C. G., Bullock, O. R., ~~Spero~~~~Otte~~, T. ~~L.~~, and Gula, J.: Using a coupled lake model with WRF for dynamical downscaling, *J. Geophys. Res.*, 119, 7193–7208, doi:10.1002/2014JD021785, 2014.

Martynov, A., Laprise, R., and Sushama, L.: Off-Line lake water and ice simulations: a step towards the interactive lake coupling with the Canadian Regional Climate Model, *Geophysical Research Abstracts*, Vol. 10, EGU2008-A-02898, EGU General Assembly 2008, Vienna, Austria, 2008.

Martynov, A., Sushama, L., and Laprise, R.: Simulation of temperate freezing lakes by one-dimensional lake models: performance assessment for interactive coupling with regional climate models, *Boreal Environ. Res.*, 15, 143–164, 2010.

Mearns, L. O., Arritt, R., Biner, S., Bukovsky, M. S., McGinnis, S., Sain, S., Caya, D., Correia Jr., J., Flory, D., Gutowski, W., Takle, E. S., Jones, R., Leung, R., Moufouma-Okia, W., McDaniel, L., Nunes, A. M. B., Qian, Y., Roads, J., Sloan, L., and Snyder, M.: The North

American regional climate change assessment program: overview of phase I results, *B. Am. Meteorol. Soc.*, 93, 1337–1362, 2012.

Mironov, D. V.: Parameterization of lakes in numerical weather prediction. Description of a lake model, COSMO Technical Report, No. 11, Deutscher Wetterdienst, Offenbach am Main, Germany, 41 pp., 2008.

Mironov, D., Heise, E., Kourzeneva, E., Ritter, B., Schneider, N., and Terzhevik, A.: Implementation of the lake parameterization scheme FLake into the numerical weather prediction model COSMO, *Boreal Env. Res.*, 15, 218–230, 2010.

NARCCAP: Caveats for Users, available at: <http://www.narccap.ucar.edu/about/caveats.html>, last access: 11 August 2014.

Notaro, M., Holman, K., Zarrin, A., Fluck, E., Vavrus, S., and Bennington, V.: Influence of the Laurentian Great Lakes on regional climate, *J. Climate*, 26, 789–804, 2013.

Oleson, K. W., Lawrence, D. M., Bonan, G. B., Flanner, M. G., Kluzek, E., Lawrence, P. J., Levis, S., Swenson, S. C., Thornton, P. E., Dai, A., Decker, M., Dickinson, R., Feddema, J., Heald, C. L., Hoffman, F., Lamarque, J.-F., Mahowald, N., Niu, G.-Y., Qian, T., Randerson, J., Running, S., Sakaguchi, K., Slater, A., Stöckli, R., Wang, A., Yang, Z.-L., Zeng, X., and Zeng, X.: Technical description of version 4.0 of the Community Land Model (CLM), National Center for Atmospheric Research, P.O. Box 3000, Boulder, Colorado 80307, USA, NCAR/TN-478+STR, 2010.

Oleson, K. W., Lawrence, D. M., Bonan, G. B., Drewniak, B., Huang, M., Koven, C. D., Levis, S., Li, F., Riley, W. J., Subin, Z. M., Swenson, S. C., Thornton, P. E., Bozbiyik, A., Fisher, R., Heald, C. L., Kluzek, E., Lamarque, J.-F., Lawrence, P. J., Leung, L. R., Lipscomb, W., Muszala, S., Ricciuto, D. M., Sacks, W., Sun, Y., Tang, J., and Yang, Z.-L.: Technical description of version 4.5 of the Community Land Model (CLM), National Center for

Atmospheric Research, P.O. Box 3000, Boulder, Colorado 80307, USA, NCAR Technical Note  
 NCAR/TN-503+STR, 420 pp., doi:10.5065/D6RR1W7M, 2013.

Ratnam, J. V., Filippo, G., Kaginalkar, A., and Cozzini, S.: Simulation of the Indian monsoon  
 using the RegCM3-ROMS regional coupled model, *Clim. Dynam.*, 33, 119–139, 2009. Rinke,  
 A., Gerdes, R., Dethloff, K., Kandlbinder, T., Karcher, M., Kauker, F., Frickenhaus, S., Köberle,  
 C., and Hiller, W.: A case study of the anomalous Arctic sea ice conditions during 1990: insights  
 from coupled and uncoupled regional climate model simulations, *J. Geophys. Res.*, 108, 4275,  
 doi:10.1029/2002JD003146, 2003.

Samuelsson, P., Kourzeneva, E., and Mironov, D.: The impact of lakes on the European climate  
 as simulated by a regional climate model, *Boreal Env. Res.*, 15, 113–129, 2010.

Schultz, D. M., Arndt, D. S., Stensrud, D. J., and Hanna, J. W.: Snowbands during the cold-air  
 outbreak of 23 January 2003, *Mon. Weather Rev.*, 132, 827–842, 2004.

Scott, R. W. and Huff, F. A.: Impacts of the Great Lakes on regional climate conditions, *J. Great  
 Lakes Res.*, 22, 845–863, 1996.

Skamarock, W. C. and Klemp, J. B.: A time-split nonhydrostatic atmospheric model for weather  
 research and forecasting applications, *J. Comp. Phys.*, 227, 3465–3485, 2008.

Skamarock, W. C., Klemp, J. B., Dudhia, J., Gill, D. O., Barker, D. M., Duda, M., Huang, X.-Y.,  
 Wang, W., and Powers, J. G.: A description of the Advanced Research WRF version 3, Tech.  
 Rep., National Center for Atmospheric Research, P.O. Box 3000, Boulder, Colorado 80307,  
 USA, NCAR/TN-475+STR, 113 pp., 2008.

Steenburgh, W. J. and Onton, D. J.: Multiscale analysis of the 7 December 1998 Great Salt  
 Lake–effect snowstorm, *Mon. Weather Rev.*, 129, 1296–1317, 2001.



Stepanenko, V. M., Goyett, S., Martynov, A., Perroud, M., Fang, X., and Mironov, D.: First steps of the Lake Model Intercomparison Project: LakeMIP, *Boreal Env. Res.*, 15, 191–202, 2010.

Subin, Z. M., Riley, W. J., and Mironov, D.: An improved lake model for climate simulations: Model structure, evaluation, and sensitivity analyses in CESM1, *J. Adv. Model. Earth Syst.*, 4, M02001, doi:10.1029/2011MS000072, 2012.

Tardy, A.: Lake-effect and lake-enhanced snow in the Champlain Valley of Vermont, Tech. Memo. 2000-05, NWS Eastern Region, 27 pp., Burlington, Vermont, USA, 2000.

Thiery, W., Stepanenko, V. M., Fang, X., Johnk, K. D., Li, Z., Martynov, A., Perroud, M., Subin, Z. M., Darchambeau, F., Mironov, D. and van Lipzig, N. P. M.: LakeMIP Kivu: evaluating the representation of a large, deep tropical lake by a set of one-dimensional lake models, *Tellus*, 66, 21390, 2014.

Wang, J., Bai, X., Hu, H., Clites, A., Colton, M., and Lofgren, B.: Temporal and spatial variability of Great Lakes ice cover, 1973–2010, *J. Climate*, 25, 1318–1329, 2012a.

Wang, J., Assel, R. A., Walterscheid, S., Clites, A. H., and Bai, X.: Great Lakes ice climatology update: winter 2006–2011 description of the digital ice cover data set, NOAA Technical Memorandum GLERL-155, 37 pp., Ann Arbor, Michigan, USA, 2012b.

Wilken, G. R.: A lake-effect snow in Arkansas, NWS Southern Region, Tech. Attachment SR/SSD 97-21, 5 pp., Little Rock, Arkansas, USA, 1997.

Wilson, J. W.: Effect of Lake Ontario on precipitation, *Mon. Weather Rev.*, 105, 207–214, 1977.

WRF User's Guide: User's Guide for the Advanced Research WRF (ARW) Modeling System Version 3.6, available at:

[http://www2.mmm.ucar.edu/wrf/users/docs/user\\_guide\\_V3/ARWUsersGuideV3.pdf](http://www2.mmm.ucar.edu/wrf/users/docs/user_guide_V3/ARWUsersGuideV3.pdf), last access:  
12 June 2014.

Wright, D. M., Posselt, D. J., and Steiner, A. L.: Sensitivity of lake-effect snowfall to lake ice  
cover and temperature in the Great Lakes region, *Mon. Weather Rev.*, 141, 670–689, 2013.

Zhao, L., Jin, J., Wang, S.-Y., and Ek, M. B.: Integration of remote-sensing data with WRF to  
improve lake-effect precipitation simulations over the Great Lakes region, *J. Geophys. Res.*, 117,  
D09102, doi:10.1029/2011JD016979, 2012.

Zulauf, M. A. and Kreuger, S. K.: Two-dimensional cloud-resolving modeling of the  
atmospheric effects of Arctic leads based upon midwinter conditions at the surface heat budget  
of the Arctic Ocean ice camp, *J. Geophys. Res.*, 108, 4312, doi:10.1029/2002JD002643, 2003.


744 Table 1. List of WRF versions discussed in the text, ordered chronologically by the date of  
 745 release and with relevant model updates summarized.

<u>WRF version</u>	<u>Released</u>	<u>Updates of Interest</u>
<u>3.3</u>	<u>April 2011</u>	<u>“Alternative initialization of lake SSTs” option included in WPS so users can set LSTs from temporally averaged 2 m temperatures.</u>
<u>3.5</u>	<u>April 2013</u>	<u>CLM available as an LSM within WRF, but with its lake model disabled.</u>
<u>3.5.1</u>	<u>September 2013</u>	<u>Default surface water temperature at which WRF prescribes ice (“seaice_threshold”) is lowered from 271 K to 100 K.</u>
<u>3.6</u>	<u>April 2014</u>	<u>CLM lake model available with any choice of LSM. Lake depths can be prescribed as a constant or as a spatially varying 2-D field.</u>

746


Table 21. A summary of the pros and cons of each method of treating lake surface temperatures and ice coverage described in the text. All approaches were found to eliminate unrealistic temperature discontinuities resulting from WRF's default interpolation methods as shown in Fig. 2.

Methodology	Positives	Potential drawbacks
WRF's Alternative Lake Setting	Effective at representing LSTs when lake temperatures are closely coupled with atmospheric temperatures.	Unrealistic ice formation possible when 2 m temperatures are below freezing. Cannot account for varying lake depths and differing timescales of warming and cooling throughout lakes.
Climatological	Observed LSTs and ice taken from high-resolution analyses.	For long-term simulations, user must include temperature trend or LSTs will not be in equilibrium with future climate state. Does not represent interannual variability of lake state.
Land Mask Modification	Future LSTs can be taken from projected GCM temperatures.	Eliminates land-lake temperature contrasts.
Lake Model Component	Models have ability to simulate future changes in LST and ice.	Additional preprocessing needed to provide lake model spin-up for RCM run or to use lake fields simulated by GCM.



752

753 Figure 1. The ocean mask from the 1° CESM data (which is used by WPS to determine the  
754 locations of land and water points from CESM), as shown in the area corresponding to a WRF  
755 36-km continental US domain (left), and the 36 km WRF grid's land-water mask (right). Labels  
756 are placed to indicate the locations of Lakes Superior ("S"), Michigan ("M"), Huron ("H"), Erie  
757 ("E") and Ontario ("O"), as well as Hudson Bay ("HB").



758

759 Figure 2. The skin temperature (K) processed from CESM to the 36-km WRF grid using WPS  
760 and valid at 00 UTC 1 Dec 1994. White circles indicate the locations of Pyramid Lake, Great  
761 Salt Lake, and Lake Sakakawea, from west to east, respectively.




Figure 3. Surface temperature from the initial year of a 10-year FLake spin-up simulation, taken from a point near the north shore of Lake Superior ( $48.47^{\circ}$  N,  $87.54^{\circ}$  W) and shown hourly from 1 January to 31 December 2005 (top). LSTs at all lake cells are initialized with a default value of 274.15 K, and the time series shows either ice or water surface temperatures depending on whether ice is present. Simulated ice thickness (m) taken from day 30 of the same FLake simulation, valid 30 January 2005 (bottom left). Fractional ice values observed on this date plotted from the NIC ice analysis (bottom right).




Figure 4. Simulated ice cover (%) taken from a WRF simulation (valid 2 March 2006, after ~ 4 months of simulation time) with the same model configuration as described in M14, but simulated with WRF version 3.6 and the use of the CLM lake model in place of FLake (left). A 2-D field of lake depths (instead of a single default value) were used from WPS to set the lake depth in this simulation. Ice coverage observed on this date is plotted from the NIC ice analysis (right).