



This discussion paper is/has been under review for the journal Geoscientific Model Development (GMD). Please refer to the corresponding final paper in GMD if available.

# Global mass fixer algorithms for conservative tracer transport in the ECMWF model

M. Diamantakis and J. Flemming

ECMWF, Shinfield Park, Reading RG2 9AX, UK

Received: 30 December 2013 – Accepted: 7 January 2014 – Published: 21 January 2014

Correspondence to: M. Diamantakis (michail.diamantakis@ecmwf.int)

Published by Copernicus Publications on behalf of the European Geosciences Union.

**GMDD**

7, 777–814, 2014

## Mass fixer algorithms for tracer transport

M. Diamantakis and  
J. Flemming

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## Abstract

Various mass fixer algorithms (MFA) have been implemented in the Integrated Forecasting System (IFS) of ECMWF to ensure mass conservation of atmospheric tracers within the Semi-Lagrangian (SL) advection scheme. Emphasis has been placed in implementing schemes that despite being primarily global in nature adjust the solution mostly in regions where the advected field has large gradients and therefore interpolation (transport) error is assumed larger.

The MFA have been tested in weather forecast, idealised and atmospheric dispersion cases. Applying these fixers to specific humidity and cloud fields did not change the accuracy of 10 day forecasts. In other words, global mass tracer conservation is achieved without deteriorating the solution accuracy. However, for longer forecast timescales or for forecasts in which correlated species are transported, experiments suggest that MFA may improve IFS forecasts.

## 1 Introduction

A drawback of semi-Lagrangian (SL) transport schemes, such as the one used by IFS (Ritchie et al., 1995) is that they do not formally conserve mass as the pointwise nature of the SL method does not take into account gridbox size and fluxes. Between the beginning and the end of each timestep, the total model mass can differ by a very small amount. This difference, although is not significant for the timescales of Numerical Weather Prediction (NWP), may accumulate in a long run. A systematic drift in the total mass of air (or a tracer field) will eventually affect the quality of the forecast (Thuburn, 2008).

As NWP models become more complex, the number of tracers increases and therefore the requirement for conservative schemes becomes more important. Furthermore, as the resolution increases towards cloud resolving scales it becomes increasingly

GMDD

7, 777–814, 2014

## Mass fixer algorithms for tracer transport

M. Diamantakis and  
J. Flemming

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



desirable from the parametrisation point of view to have a mass conserving advection scheme as this may improve further the simulation of cloud processes.

SL advection (SLA) consists of two steps which do not – in principle – ensure conservation of mass: (i) finding departure points and (ii) interpolating the advected field to their location. However, the choice of method for (i) and (ii) has a considerable impact for the amount of the mass-non-conservation.

There is a class of SL-schemes, the so called inherently conserving schemes, which are able to achieve global, local and consistent mass conservation for tracer and air-mass fields. Two examples are SLICE (Semi-Lagrangian Inherently Conserving and Efficient) transport scheme (see Zerroukat and Allen, 2012) and CSLAM (Conservative Semi-Lagrangian Multi-tracer) transport scheme (see Lauritzen et al., 2010). These schemes are an application of a finite-volume type discretization approach on the semi-Lagrangian continuity equation. In general, they are complex algorithms difficult to implement efficiently in an existing operational model which uses a “traditional” SL method. Although inherently conserving SL methods are not currently used in weather forecasting operations there are schemes in this family which are competitive or even more efficient than their Eulerian finite-volume conservative counterparts for applications where a large number of tracers is advected (multi-tracer simulations). CSLAM is an example of such method while another example of a recent development based on LMCSL by Kaas (2008) scheme (Locally Mass Conserving semi-Lagrangian) is given by Sørensen et al. (2013).

An alternative low computational cost approach to ensure global mass conservation which can be easily applied on traditional SL methods is the Mass Fixer Algorithms (MFA). The task of a MFA is to change the tracer concentrations after SLA in such way that the mass before and after advection is the same. A general problem of MFA is to identify regions where it is most appropriate to change the solution of the SL scheme.

Different MFAs implement different strategies for distributing the global mass loss or gain. The simplest ones correct the solution uniformly by simply scaling each grid-point value with the ratio of the global mass before and after advection. This approach is

## Mass fixer algorithms for tracer transport

M. Diamantakis and  
J. Flemming

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



currently used in IFS when long time integrations take place in order to correct the total model mass and of long-lived tracers (Flemming and Huijnen, 2011).

More sophisticated MFA attempt to compute a correction which is proportional to the smoothness of the solution. A larger correction is applied in areas where the solution has large gradients and therefore the error is larger and a very small correction where the solution is smooth and the error is small.

The aim of the paper is to present tracer MFA that were recently implemented in IFS in model cycle 39r1. Using this model cycle as the base for our experiments we shall discuss results from NWP forecasts, long range forecasts where the mass fixers are applied to humidity and cloud fields as well as idealised tracer and volcanic plume forecasts. Availability of globally mass conserving schemes for tracers can be an important addition to IFS based prediction systems such as the EC-Earth (Hazeleger et al., 2012) climate model or atmospheric composition forecast systems where aerosols, greenhouse and reactive gases are transported (Hollingsworth et al., 2008).

The paper is structured as follows. The amount of the non-conservation by the SL advection scheme of IFS is demonstrated in Sect. 2. Section 3 describes the implemented MFA. Their impact on the simulated fields in different applications is discussed in Sect. 4. Conclusions are presented in Sect. 5.

## 2 Air and tracer global mass conservation in IFS

In a 10 day IFS forecast, at the current operational T1279L137 (approximately 16 km in grid-point space on 137 levels), the total model air mass increases by less than 0.01 % of its initial value. The formulation of the continuity equation, based on Ritchie and Tanguay (1996) scheme (see also ECMWF, 2012, Sect. 3.6.2), plays an important role into achieving this accuracy. Orography is removed from the advected mass field resulting in a much smoother field which can be accurately interpolated to the Lagrangian-grid (departure points).

### Mass fixer algorithms for tracer transport

M. Diamantakis and  
J. Flemming

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## Mass fixer algorithms for tracer transport

M. Diamantakis and  
J. Flemming

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Global conservation errors in tracer advection are larger and depend on the smoothness of the field. For example, smoother fields such as ozone and specific humidity have smaller conservation errors than fields with sharp features such as cloud fields. This is demonstrated in Fig. 1 where the global mass conservation error is displayed for ozone, specific humidity ( $Q$ ), liquid cloud water content (CLWC), cloud ice water content (CIWC) for the same number of timesteps at different resolutions using two approaches for the interpolation to the departure point. Mass conservation is represented in Fig. 1 by a line identical to the horizontal 0-axis. The global mass conservation error for a tracer  $\phi$  is expressed as a percentage of its initial mass:

$$E_{\phi} = 100 \times \frac{M_t^{\phi} - M_0^{\phi}}{M_0^{\phi}}$$

where,  $M_0^{\phi}$ ,  $M_t^{\phi}$  is the initial and current step global tracer mass.

In the forecast experiments of Fig. 1 all parametrizations of sink and source terms have been switched off. This allows to test the performance of the advection scheme using real orography. In addition, the following two interpolation methods have been used: (i) the quasi-cubic ECMWF interpolation, Ritchie et al. (1995), with a quasi-monotone limiter and (ii) a linear interpolation (indicated with LIN in plots). Method (i) is used in IFS operationally for  $Q$  and ozone while method (ii) is used operationally for the rougher cloud fields. The experiments are run at the following horizontal and vertical resolutions: (i) T159 L60 i.e. T159 in the horizontal (approximately equal to 125 km) with 60 levels in the vertical (ii) T159 L91 (iii) T1279 L91 (approximately 16 km in the horizontal) and (iv) T1279 L137. To allow direct comparisons of the mass conservation error per timestep, the four top forecasts in Fig. 1 have been run for the same number of timesteps. At coarse horizontal resolution (T159) the timestep is 6 times longer (60 min) than the corresponding timestep for high resolution (T1279).

The results shown in Fig. 1 indicate that the global mass conservation error per timestep tends to decrease as the resolution increases. However, when *horizontal*

*resolution* is increased from T159 to T1279, the accumulated error at  $t = 10$  days decreases only for CLWC, CIWC with cubic interpolation while remains roughly the same for the remaining fields. It seems that the opposite is true when vertical resolution increases, the accumulated error at  $t = 10$  days decreases except for CLWC, CIWC with cubic interpolation. So there are differences between interpolation schemes and between fields of different smoothness but the overall indication is that in the IFS system mass conservation of tracers tends to improve globally as resolution increases and the best way to demonstrate this is by comparing Fig. 1a with d.

### 3 Description of the MFAs

The transport problem we consider here is the advection of a scalar field  $\phi_\chi$  which represents the mass mixing ratio of a tracer:

$$\frac{D\phi_\chi}{Dt} = S, \phi_\chi = \rho_\chi / \rho \quad (1)$$

where  $\rho_\chi, \rho$  are the tracer and air density respectively and  $S$  represents sources or sinks that may be present. Consider SL time-stepping from  $t$  to  $t + \Delta t$ :

$$\phi_\chi^{t+\Delta t} = \phi_{\chi,d}^t + \Delta t S$$

where  $d$  denotes the departure point computed by the trajectory algorithm and  $\phi_{\chi,d}^t$  is obtained by interpolating the known field  $\phi_\chi^t$  at the computed departure point. If  $S = 0$  then the global volume integral of  $\rho\phi_\chi$  at  $t$  and  $t + \Delta t$  (on the model grid) should not change as this represents the total mass of  $\chi$  and the only process operating is advection (transport). However, in practice, as the interpolation scheme generates errors this global conservation law is violated.

Global MFAs of different sophistication are described in the published literature for SL transport models. In general, any MFA will compute the global tracer mass immediately

## Mass fixer algorithms for tracer transport

M. Diamantakis and  
J. Flemming

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



before and after the advection step. Then a small correction is computed for each grid-point in such a way that this global error is eliminated. In the simplest version of the proportional or multiplicative fixer of Rasch and Williamson (1990), each grid-point value is multiplied by the ratio of the mass before and after advection. Here, we will focus on the more local algorithms. In particular, the following algorithms will be discussed: (i) the quasi-monotone Bermejo and Conde (2002) scheme, (ii) Zerroukat (2010) scheme, (iii) the quasi-monotone Priestley (1993) scheme, and (iv) MacGregor (2011) scheme. These algorithms have been implemented in IFS and will be summarised in the following paragraphs. It should be noted that their implementation is three-dimensional given that semi-Lagrangian advection in IFS is fully three-dimensional.

To describe these different fixers, as implemented in IFS, we use the following notation:  $K$  is the number of model levels, starting from the top of the atmosphere and ending on the surface. Each model level has  $N$  grid-points. Each grid-box has horizontal surface area  $A_j$  and height  $\Delta z_{jk}$  where  $z_{jk}$  denotes the height of the  $j$ th model grid-point of the  $k$ th level. The total mass of a tracer  $\chi$  with mass mixing ratio  $\phi = \rho_\chi / \rho$  where  $\rho$  is the air-density field is:

$$M = \sum_{j=1}^N A_j \sum_{k=1}^K \rho_{\chi,jk} (-\Delta z_{jk}) = \sum_{j=1}^N A_j \sum_{k=1}^K \phi_{jk} \frac{\Delta \rho_{jk}}{g}, \quad \Delta z_{jk} = z_{jk} - z_{j,k-1} < 0, \quad (2)$$

$$\Delta \rho_{jk} = \rho_{jk} - \rho_{j,k-1} > 0.$$

The hydrostatic approximation (valid in IFS)  $\Delta p = -\rho g \Delta z$  has been used in Eq. (2) to eliminate  $\Delta z$ .

During the advection step, a tracer field  $\phi^0$  (i.e. the field before the advection step takes place) is interpolated to the departure point field (Lagrangian grid) and changes to  $\phi^*$  while its total mass changes from  $M^0$  to  $M^*$ :

$$M^0 = \sum_{j=1}^N A_j \sum_{k=1}^K \phi_{jk}^0 \frac{\Delta \rho_{jk}^0}{g}, \quad M^* = \sum_{j=1}^N A_j \sum_{k=1}^K \phi_{jk}^* \frac{\Delta \rho_{jk}^*}{g}. \quad (3)$$

Mass fixer algorithms for tracer transport

M. Diamantakis and J. Flemming

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Use of  $\Delta p_{jk}^*$  in Eq. (3) reflects the change of the surface pressure field due to advection. MFAs aim correcting  $\phi^*$  so that a new field is derived which has a total mass equal to  $M^0$ .

### 3.1 Bermejo and Conde (BC) scheme

Bermejo and Conde (2002) algorithm is derived by a variational principle. It computes a new quasi-monotone field minimizing its distance from the original one subject to the constraint of global mass conservation. The correction added at each grid point depends on an estimate of the interpolation error. The global norm of this correction field has the smallest possible magnitude that can give mass conservation and monotonicity. In the original publication, the scheme was tested on idealised 2-dimensional cases of advection. Here it has been implemented in IFS in 3-D mode and has been tested on active meteorological fields.

Let  $\phi^1$  be the field which minimizes the square of the weighted norm:

$$\min_{\phi^1} \|\phi^1 - \phi^*\|_w^2 = \frac{1}{2} \sum_{j=1}^N A_j \sum_{k=1}^K \frac{(\phi_{jk}^1 - \phi_{jk}^*)^2}{w_{jk}} \frac{\Delta p_{jk}^*}{g} \quad (4)$$

subject to

$$\sum_{j=1}^N A_j \sum_{k=1}^K \phi_{jk}^1 \frac{\Delta p_{jk}^*}{g} = M^0$$

where  $w_{jk}$  is a non-negative weighting factor. Having  $w_{jk} = 0$  means that the corresponding grid-point value is not altered and is not included in the cost function.

## Mass fixer algorithms for tracer transport

M. Diamantakis and  
J. Flemming

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



A solution to Eq. (4) is found using a Lagrange multiplier approach. The cost function

$$E(\phi^1, \lambda) = \frac{1}{2} \sum_{j=1}^N A_j \sum_{k=1}^K \frac{(\phi_{jk}^1 - \phi_{jk}^*)^2}{w_{jk}} \frac{\Delta p_{jk}^*}{g} - \lambda \left( \sum_{j=1}^N A_j \sum_{k=1}^K \phi_{jk}^1 \frac{\Delta p_{jk}^*}{g} - M^0 \right),$$

is defined seeking a pair of values  $(\phi^1, \lambda)$  such that:

$$\frac{\partial E}{\partial \phi_{jk}^1} = 0, \frac{\partial E}{\partial \lambda} = 0.$$

Solving these two equations we obtain:

$$\phi_{jk}^1 = \phi_{jk}^* - \lambda w_{jk}, \lambda = \frac{\delta M}{\sum_{j=1}^N A_j \sum_{k=1}^K w_{jk} \frac{\Delta p_{kj}^*}{g}}, \delta M = M^* - M^0 \quad (5)$$

where the weight  $w_{jk}$  depends on the solution smoothness. We choose it to be proportional to the difference between the quasi-cubic, quasi-monotone interpolated field  $\phi^*$  and the linear one  $\phi^L$ :

$$w_{jk} = \max \left[ 0, \text{sgn}(\delta M) \left( \phi_{jk}^* - \phi_{jk}^L \right)^p \right]. \quad (6)$$

The above weights are used to compute a “local correction”, i.e. the global mass surplus or deficit is distributed unevenly to different grid-points depending on the smoothness of the solution which is measured by the difference between a high and a low order interpolant. For the IFS implementation,  $p$  was set to 1 as tests showed no benefit from using the recommended value  $p = 3$ . In fact, higher values led to sharper, bigger size increments which may not be desirable for the model stability.

In sections that follow for convenience this scheme will be called *BC* fixer.

**Mass fixer algorithms  
for tracer transport**

M. Diamantakis and  
J. Flemming

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



### 3.2 Zerroukat's (ZE) scheme

The BC fixer in IFS can also be run in a mode that corresponds to a version of the Zerroukat (2010) fixer. This leads to smoother correction fields. The drawback is that quasi-monotonicity or positive-definiteness cannot be guaranteed. Here an implementation of this scheme is presented which uses the same measure to assess the solution smoothness as the BC scheme, i.e. the difference between a high order scheme (cubic Lagrange interpolation) and a low order scheme (linear interpolation). Here, this scheme will be called *ZE* fixer. It corrects each grid-point value as follows:

$$\phi_{jk}^1 = \phi_{jk}^* - \gamma_{jk} \delta M, \quad \delta M = M^* - M^0, \quad \gamma_{jk} = \frac{|\phi_{jk}^* - \phi_{jk}^L|^p}{\sum_{j=1}^N A_j \sum_{k=1}^K |\phi_{jk}^* - \phi_{jk}^L|^p \frac{\Delta \rho_{jk}^*}{g}} \quad (7)$$

where  $M^0$ ,  $M^*$  are defined by Eq. (3) and again  $\rho = 1$  is sufficient for practical purposes. If

$$\sum_{j=1}^N A_j \sum_{k=1}^K \gamma_{jk} \frac{\Delta \rho_{jk}^*}{g} = 1$$

holds then global mass conservation is guaranteed:

$$\sum_{j=1}^N A_j \sum_{k=1}^K \phi_{jk}^1 \frac{\Delta \rho_{jk}^*}{g} = M^0.$$

It is worth noticing that Eq. (7) can be re-written in a form that resembles Eq. (5):

$$\phi_{jk}^1 = \phi_{jk}^* - \lambda w_{jk}, \quad \lambda = \frac{\delta M}{\sum_{j=1}^N A_j \sum_{k=1}^K w_{jk} \frac{\Delta \rho_{jk}^*}{g}}, \quad \delta M = M - M^0, \quad w_{jk} = |\phi_{jk}^* - \phi_{jk}^L|^p. \quad (8)$$

This implies that the derived field  $\phi^1$  is also a solution of the minimization problem Eq. (4). One difference between Eqs. (5) and (8) is the construction of the weights  $w_{jk}$ . Using the unlimited  $w_{jk} = |\phi_{jk}^* - \phi_{jk}^L|^p$  means that all grid-points will be corrected. The sign of the increment is determined by the sign of  $\delta M$  (which determines the sign of  $\lambda$ ): for  $\delta M > 0$  (surplus)  $\phi_{jk}^1 \leq \phi_{jk} \forall j, k$  and for  $\delta M < 0$  (deficit)  $\phi_{jk}^1 \geq \phi_{jk} \forall j, k$ . However, as this one-directional correction is not limited as in the BC case, it is possible that a new minimum or maximum value may be generated. In practice, if a quasi-monotone scheme was used for advection this happened in less than 0.5% for humidity grid-points but sometimes can go up to 5% of grid-points for a non-smooth field.

### 3.3 Priestley's (PR) scheme

Priestley (1993) is a well known mass fixing scheme. Its objective is to compute a globally conserving monotone solution by blending the original high order with a low order solution thereby departing as little as possible from the high order one. This is equivalent to finding the highest possible values for the weights  $\alpha_{jk}$  such that the “blended” field:

$$\phi_{jk}^1 = \alpha_{jk} (\phi_{jk}^* - \phi_{jk}^L) + \phi_{jk}^L, \quad 0 \leq \alpha_{jk} \leq 1$$

satisfies:

$$\min \left( \{\phi^0, j, k\}, \phi_{jk}^L \right) \leq \phi_{jk}^1 \leq \max \left( \{\phi^0, j, k\}, \phi_{jk}^L \right), \quad \sum_j A_{j=1}^N \sum_{k=1}^K \phi_{jk}^1 \frac{\Delta p_{jk}^*}{g} = M^0 \quad (9)$$

where  $\{\phi^0, j, k\}$  denotes the set of  $\phi$ -field values before advection at grid-points surrounding the  $(j, k)$  departure point and  $\phi^*$ ,  $\phi^L$  the cubically and linearly interpolated field at the departure point respectively. The two conditions in Eq. (9) ensure conservation and monotonicity. The requirement for “highest possible”  $\alpha$  values is an accuracy requirement. It ensures that the final solution is as close as possible to the original high



## Mass fixer algorithms for tracer transport

M. Diamantakis and  
J. Flemming

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



The algorithm can be described as follows:

**Step 1** Compute total mass before and after advection  $M^0$ ,  $M^*$  as in Eq. (3).

**Step 2** Let a minimum allowed value  $\phi^{\min}$ . Scan each grid point, compute and store:

$$\Delta\phi_{jk}^+ = \max(0, \Delta\phi_{jk}), \quad \Delta\phi_{jk}^- = \min(0, \Delta\phi_{jk})$$

where

$$\Delta\phi_{jk} = \max(\phi_{jk}^*, \phi_{jk}^{\min}) - \frac{\Delta\rho_{jk}^0}{\Delta\rho_{jk}^*} \phi_{jk}^0$$

**Step 3** Compute total positive and negative increments and their ratio:

$$\Delta M^+ = \sum_{j=1}^N A_j \sum_{k=1}^K \frac{\Delta\rho_{jk}^*}{g} \Delta\phi_{jk}^+, \quad \Delta M^- = \sum_{j=1}^N A_j \sum_{k=1}^K \frac{\Delta\rho_{jk}^*}{g} \Delta\phi_{jk}^-,$$

$$r = -\frac{\Delta M^-}{\Delta M^+}$$

**Step 4** Set  $\alpha_\phi = \min(r, \sqrt{r})$  and update:

$$\phi_{jk}^1 = \frac{\Delta\rho_{jk}^0}{\Delta\rho_{jk}^*} \phi_{jk}^0 + \alpha_\phi \Delta\phi_{jk}^+ + \frac{1}{\max(1, \alpha_\phi)} \Delta\phi_{jk}^-$$

The last step is equivalent to:

$$\phi_{jk}^1 = \begin{cases} \frac{\Delta\rho_{jk}^0}{\Delta\rho_{jk}^*} \phi_{jk}^0 + \alpha_\phi \Delta\phi_{jk}^+ + \Delta\phi_{jk}^-, & r \leq 1 \\ \frac{\Delta\rho_{jk}^0}{\Delta\rho_{jk}^*} \phi_{jk}^0 + \alpha_\phi \Delta\phi_{jk}^+ + \frac{1}{\alpha_\phi} \Delta\phi_{jk}^-, & r > 1 \end{cases}$$

and implies that the increment is scaled by a factor  $\alpha_\phi$  which reduces positive increments when their total mass exceeds the total mass of the negative increments. When the opposite is true then positive increments will be amplified and negatives will reduce in magnitude. The new field satisfies the global mass conservation constraint:

$$5 \quad \sum_{j=1}^N A_j \sum_{k=1}^K \phi_{jk}^1 \frac{\Delta p_{jk}^*}{g} = M_0.$$

### 3.5 The quasi-monotone limiter

The quasi-monotone limiter renders the interpolation locally monotone, i.e. in the vicinity of the departure point the interpolation curve (or multidimensional surface) passing from the departure point field value and the field values of points surrounding the departure point does not generate new minima or maxima. For the tests presented in the following section two forms of the quasi-monotone Bermejo and Staniforth (1992) mini-max limiter for cubic interpolation will be used:

1. “the default” limiter or filter used operationally in IFS: the scheme is applied immediately after each 1-D cubic interpolation (in longitude, latitude, height) takes place. So, the steps taken are: interpolate in longitude and then apply 1-D limiter on the interpolated field. Repeat this action for each of the remaining two interpolations (in latitude, height). For brevity this scheme will be called *DEF* limiter or filter.
2. the standard Bermejo and Staniforth (1992) limiter: this shall be called *BS* limiter or filter. In this case the limiter is applied after all three interpolations have finished, i.e. this is limiting in 3-D at once.

We should also clarify that the term “cubic interpolation” will imply here the quasi-tri-cubic interpolation scheme used by IFS (linear interpolation along the edges of the stencil, fully cubic in the interior, see Ritchie et al., 1995).

**Mass fixer algorithms for tracer transport**

M. Diamantakis and J. Flemming

|                          |              |
|--------------------------|--------------|
| Title Page               |              |
| Abstract                 | Introduction |
| Conclusions              | References   |
| Tables                   | Figures      |
| ◀                        | ▶            |
| ◀                        | ▶            |
| Back                     | Close        |
| Full Screen / Esc        |              |
| Printer-friendly Version |              |
| Interactive Discussion   |              |



## 4 Testing of MFAs in IFS

In Fig. 2 the global conservation error during the advection of  $Q$  and CLWC with and without MFA is displayed. It is shown there that application of a MFA eliminates this error. This forecast run has the operational resolution (T1279 horizontal with 137 levels) and is identical to the one that corresponds to the results of Fig. 1 i.e. there are no sources or sinks of tracer mass. For brevity we display only results from BC and PR schemes but also the other MFAs give a globally mass conserving solution. The mass conservation error before and after the advection was always close to machine precision.

The impact of the BC MFA on  $Q$  is demonstrated in Fig. 3. Cubic interpolation is used for the advection of this field. Here, physical parametrizations have been switched on and the setup is the same as in an operational forecast. A single timestep increment from the fixer, at  $t = 24$  h and at a model level which over flat terrain is near the 700 hPa pressure level, is compared with the field itself. The figure shows that the computed increments are at least 3 orders of magnitude smaller than their corresponding field magnitude. The sign is negative due to the fact that at this stage of the forecast, advection increases mass and the fixer has to remove a global surplus. The fixer is acting mainly on areas where large gradients are present where interpolation is expected to be less accurate. In areas where the field is smooth the correction is very small regardless of the field magnitude. Similar results have been produced from runs with the remaining MFAs. For brevity these will not be displayed here but they are publically available (see Diamantakis and Flemming, 2013, Sect. 4).

A zonally and 24 h time-averaged vertical cross section for  $Q$  is compared with corresponding cross-sections of increment diagnostics in Fig. 4. The average increment is 4–5 orders of magnitude smaller than the magnitude of the field itself. It is concentrated in areas where large amounts of humidity are present. It is interesting to notice how similar the zonally and time averaged increments are for BC, ZE and PRqm. The fact that their difference is small means that the different algorithms converge roughly to the

GMDD

7, 777–814, 2014

### Mass fixer algorithms for tracer transport

M. Diamantakis and  
J. Flemming

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



same solution. Larger differences can be noticed when any of the previous three fixers is compared with MG and even larger with *PR*.

Usually, increments computed by PR differ in sign and magnitude from the other fixers (see also Figs. 4, 5). This is because this algorithm computes a quasi-monotone and conservative solution iteratively starting from a cubic interpolated field. In the tests presented here it usually takes 3–4 iterations for PR to converge. During this iterative process both positive and negative increments will be computed to derive a locally monotone solution. Mass has to be removed from overshooting points (negative increment) and added at undershooting points (positive increment). This is not the case with *PRqm* which starts with a quasi-monotone field having no undershooting or overshooting points and therefore the only action that the algorithm needs to take is to restore global mass conservation. Regarding the remaining fixers it is worth mentioning that: (i) ZE produces the smallest, in magnitude, increments but these are slightly more widespread (ii) BC and *PRqm* are similar and (iii) MG produces slightly different patterns than the previous two fixers. As expected, the quasi-monotone schemes did not produce any overshoots or undershoots. A very small percentage of undershoots ( $< 0.01\%$  of total points) was found with MG but no negative values were created. This percentage was larger in the ZE fixer for the cloud fields, exceeding slightly  $1.5\%$ , while it was of similar magnitude for  $Q$  ( $\approx 0.01\%$ ). Most of these undershoots generated negative values.

In the plots presented here specific humidity was chosen to examine the local behaviour of MFAs. This choice was made due to the meteorological importance of this tracer field and given that it includes regions that are relatively smooth as well as regions with large gradients. The MF applied to the rougher cloud fields CLWC and CLIC resulted in similar local patterns as shown for  $Q$ . The CLWC increments were used as a diagnostic for demonstrating the step by step behaviour of the MFAs. This is shown in Fig. 5 where the scaled global rms and max norms of the of CLWC fixer (absolute) increments are displayed. These are scaled to be the fraction (percentage) of the rms global norm of the advected CLWC field which is representative to its mean value. The

## GMDD

7, 777–814, 2014

### Mass fixer algorithms for tracer transport

M. Diamantakis and  
J. Flemming

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



plot shows that the smallest increments are computed by ZE fixer, followed by BC and *PRqm* while as expected and explained before PR computes the largest increments. MG increments are in the middle between PR and ZE.

Finally, to assess the computational cost of the fixers, 10 day forecast tests with the high resolution control (T1279 L137) have been done applying the fixers on *Q*, CLWC, CIWC, CRWC (cloud rain water content) and CSWC (cloud snow water content). The extra CPU time consumed by these algorithms is: (i) BC: 1 %, (ii) *PRqm*: 2 %, (iii) *PR*: 3.5 %, (iv) *MG*: 0.75 %, (v) *ZE*: 0.85 %.

As expected *PR* is the most expensive and *MG* the cheapest. All algorithms have been parallelized using MPI and openMP directives.

#### 4.1 Impact of humidity MFAs on temperature fields in long runs

As there is a strong interaction between humidity and temperature, typically because of radiative effects and cloud microphysics, we shall test in this section to what extent the mass fixer increments on humidity and cloud fields alter the temperature field. To show the impact we carried out four 12 month forecasts with full physics at T159 L137 resolution. This is a standard test of IFS which is done to evaluate whether a new scheme impacts on model climate. The experiments run are described in Table 1.

In Fig. 6a the temperature bias is plotted, i.e. the difference of the vertical cross section of a zonally averaged annual mean temperature field (averaged across the four forecasts) from its corresponding field from ERA-Interim run. This figure displays a common problem in semi-Lagrangian models, the extra-tropical tropopause/lower stratosphere cold bias (see Stenke et al., 2008). For the remaining plots, the difference of the same field (zonally averaged annual mean temperature) from the control run is used. This is done to clearly demonstrate the impact of the changes. As a general rule, warming around the extra-tropical tropopause (in the region where the blue area in Fig. 6a appears) would indicate an improvement while cooling would indicate further deterioration.

## Mass fixer algorithms for tracer transport

M. Diamantakis and  
J. Flemming

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## Mass fixer algorithms for tracer transport

M. Diamantakis and  
J. Flemming

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Results show that none of these fixers deteriorates the existing cold bias. When the fixers are combined with the DEF limiter the difference is small (results show a marginal improvement and have not been included here). On the contrary a noticeable improvement, i.e. a reduction of the cold bias, can be noticed when they are combined with the BS limiter. This shows in Fig. 6b–f. Good results are obtained with the quasi-monotone algorithms PR and BC. As condition in equation Eq. (9) shows, PR fixer is limiting the solution using a similar scheme to BS limiter. Bigger positive impact is obtained by fixers that do not guarantee quasi-monotonicity: ZE followed by MG. However, the former generates negatives especially in the cloud fields which are rougher (3–5% of grid-points become negative after correction is applied). This is not the case for the latter where a negative fixer is built in.

### 4.2 Impact on NWP scores in 10 day forecasts

The forecast accuracy of 10 forecast is typically assessed using accuracy measures that describe the realism of the global geopotential or temperature fields. The forecast fields are compared against the Analysis of the fields and expressed as Root Mean Square Error (RMSE) or Anomaly Correlation Coefficient (ACC) (Wilks, 2011).

In general, it is not expected that a global MFA will improve forecasting skill in the short or medium range but neither it should deteriorate the skill. To investigate this the MFAs have been tested running 37 forecast cases, each starting 10 days apart from 1 December 2011 until 25 November 2012. The resolution used is T511 L137 and each forecast is run for 10 days using operational options for the model dynamics and physics. All fixers were activated on  $Q$ , CLWC, CIWC, CRWC, CSWC. Although these tests are specific on moist physics tracers, they do have a general value. We can indirectly measure the impact a fixer has on advection by measuring the overall forecast skill of the experiment: forecast skill deterioration would imply that the tested algorithm deteriorates the accuracy of the advection scheme and therefore is deemed not suitable for tracer advection. Neutral scores should indicate that the fixer is making

the interpolation conservative without damaging solution accuracy at least on the large scale.

Overall, geopotential, wind, temperature verification scores in the three global regions (NH, TR, SH) from runs with MFAs are neutral and there is no forecast that is better in terms of ACC and RMSE. An exception is the temperature RMSE in the tropics at upper tropospheric levels which increases up to 0.07 K (from approximately 1.26 K to 1.33 K) at  $t = 10$  days when any MFA is applied for humidity and cloud fields with cubic interpolation options. The fixer contributes further (by a small amount) to the existing cold bias. This happens because a small amount of humidity is removed from the atmosphere as a small humidity surplus is detected by the fixer. Reducing the humidity content of the troposphere has in general a cooling effect while the opposite is true for the stratosphere due to reduction of radiative cooling. However, there is no impact on the corresponding ACC scores which remain neutral.

### 4.3 Simulation of correlated tracers

Mass conservation is an important property for atmospheric applications where chemical species are transported. It is also important that existing functional relationships in their concentration are maintained by the advection scheme (see Lauritzen and Thuburn, 2012). The ability of IFS and the newly developed fixers to preserve such relationships has been tested using case 11 from DCMIP (Dynamical Core Model Intercomparison Project, see Ulrich et al., 2012). This is a three-dimensional passive advection deformational flow idealised test case in which four tracers are transported. The initial concentration of the first two tracer fields  $q_1$ ,  $q_2$  obeys the nonlinear relationship:

$$q_2(\lambda, \theta, z) = 0.9 - 0.8q_1^2(\lambda, \theta, z)$$

where  $\lambda$ ,  $\theta$ ,  $z$  is the longitude, latitude and height of a tracer. The first one ( $q_1$ ) is represented by two cosine bells placed at the same height and latitude but at different longitudes.

GMDD

7, 777–814, 2014

## Mass fixer algorithms for tracer transport

M. Diamantakis and  
J. Flemming

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## Mass fixer algorithms for tracer transport

M. Diamantakis and  
J. Flemming

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Results for this test case from IFS runs at T159 horizontal resolution and 137 levels in the vertical (this is close to the recommended resolution for this problem) are plotted in Fig. 7. These plots are correlation plots for the pair  $(q_1, q_2)$  at  $t = 6$  days after the initial time. This is half the time required for the tracers to return to their original position i.e.

complete one full rotation around the earth. The initial concentration of these tracers is given by the parabolic dash-dotted black curve. Pairs  $(q_1, q_2)$  (red dots) that fall outside the region marked by the dashed-dotted convex shape correspond to unphysical mixing ratios. Real mixing in the atmosphere can only move scatter points to the concave side of the pre-existing functional curve along mixing lines (Lauritzen and Thuburn, 2012). Lack of spread indicates that the scheme is over-diffusive as peak values are damped.

The plots show that semi-Lagrangian transport with linear interpolation is excessively diffusive but does not produce any unphysical mixing. The opposite is true when cubic Lagrange is used. It results in relatively large amount of unphysical mixing and overshoots/undershoots (new maxima/minima are created corresponding to values above 1 and below 0). Significant improvements can be noticed when a quasi-monotone limiter is used. The DEF limiter, being more strict (and damping) has bigger impact as all points stay inside the convex shape. However, maximum field values are damped. The BS limiter reduces but does not eliminate completely the unphysical mixing occurring with cubic interpolation. However, it preserves better the maxima.

When a MFA is combined with the DEF limiter it does not change further the mixing: it preserves equally well existing tracer correlations as shown in Fig. 7 (compare c and d). It also results in a small further reduction of maximum field values (result not included here). When the fixers are combined with the BS limiter we obtain very similar results with respect to tracer correlations compared with corresponding results from DEF limiter but slightly improved results in terms of accuracy (preservation of maxima). In this case BC and PR give the best results. They both preserve reasonably well the initial correlation (better than the corresponding run without fixer) and maximum field values are not too far from the analytical values. ZE and MG fixers are not as effective in preserving the functional relationship (especially the latter) as a small proportion of

points are outside the bounded sector. The former can produce small negatives values in some regions. But they are both better in preserving maxima.

In conclusion, applying any of the MFAs did not deteriorate the mixing properties of the advection scheme and in some occasions improved them (e.g. compare Fig. 7e and f). This is a desirable result and suggests that MFAs can be a beneficial addition for a semi-Lagrangian scheme used for transport of chemical tracers. Combination of a MFA with the BS limiter works better and BC, PR seem to give the best results.

#### 4.4 Volcanic plume case study

MFAs have also been tested on volcanic plume advection cases. Here a test case is presented where a tracer ( $\text{SO}_2$ ) is emitted into the atmosphere by a single point source and then transported by the winds. This case resembles the Grímsvötn volcanic eruption (see Flemming and Inness, 2013). Due to the highly localised nature of the advected plume, this case is a good test for assessing the local behaviour of a global MFA. The striking fact in this simulation is that the plume total mass is largely overestimated. A conservation error of almost 20% of the total mass of the field occurs during the first timesteps which eventually results to more than 50% gain. This is shown in Fig. 8. The greatly improved performance in terms of conservation of the non-limited cubic Lagrange without MF shown in this plot is due to the presence of large negative undershoots which offset the overshoots when the global integral is computed and is therefore misleading.

Applying a MFA results in a globally conserving solution as shown by the 0 residual line in Fig. 8. The MFA applied there is BC but the same result is obtained by any of the other algorithms. It also results in some reduction of the peak values of the field which is evident in Fig. 9. This can be explained if we consider that a MFA diagnoses that the total mass has been largely overestimated by cubic interpolation and has to remove mass to enforce conservation. As the mass is concentrated in a small area, few grid-points across, peak values will be inevitably reduced when the MFA is applied. Large interpolation errors as a result of large gradients and insufficient resolution near

## Mass fixer algorithms for tracer transport

M. Diamantakis and  
J. Flemming

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



the source is the main reason for this mass overestimation. The sensitivity with respect to the specific mass fixer or quasi-monotone filter used was relatively small and all algorithms tested behave in a similar way. The biggest difference was found between MG fixer and the remaining ones and this shows in Fig. 9.

Although it is difficult to obtain accurate results in test cases of advection of small scale point sources with coarse (global) resolution semi-Lagrangian models, useful qualitative results can still be obtained. The MFA may reduce the amplitude of the field but it will correct its total mass which is necessary for emission parameter estimation.

## 5 Conclusions

A MFA is a technique to correct the global mass conservation error that a non-formally conserving advection scheme introduces. It acts a-posteriori to correct the solution after the field has been advected. In the context of a semi-Lagrangian scheme this means to correct the field after it has been interpolated to the departure point and before other source terms due to physical processes are added.

Different MFAs have been implemented (cf. Sect. 3) in IFS based on different strategies for correcting the global mass conservation error. They all follow a weighted approach, i.e. weights are computed which determine how much to adjust each grid-point value. The aim is to correct the advected field in regions where the interpolation error is large. Results show that indeed these methods act in areas of steep gradients where the solution is not smooth while they apply very small corrections elsewhere. They achieve globally mass conserving solutions without deteriorating accuracy at large scales. This has been demonstrated here by a set of 12 month forecast tests verified against ERA-Interim and standard 10 day forecasts at T511 L137 resolution verified against ECMWF operational analysis. A small local degradation of existing biases cannot be completely ruled out since the sign of the global mass error determines the sign of the corrections everywhere. The key results from this work are:

**GMDD**

7, 777–814, 2014

## Mass fixer algorithms for tracer transport

M. Diamantakis and  
J. Flemming

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## Mass fixer algorithms for tracer transport

M. Diamantakis and  
J. Flemming

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

1. No significant differences have been found between the approaches at the hydrostatic scales tested. But there are small differences in cost.
2. Global conservation is achieved without deteriorating the solution. An exception is the volcanic plume case in which peak values are reduced. However, this side-effect is also related to the lack of sufficient resolution. Despite this, global mass conservation is important for emission parametric estimates because the mass conservation error can reach up half of the emitted mass.
3. The impact on forecast skill was neutral.
4. Noticeable impact was found from the quasi-monotone limiter applied. In long integrations BS improves on the standard quasi-monotone scheme used in IFS.

Based on the above findings the recommendations on the use of the newly implemented MFAs in IFS are:

1. For quasi-monotone cubic advection of moist quantities BC is the preferred option as it is shape preserving and one of the cheapest.
2. If quasi-monotonicity is not essential and positive-definiteness is sufficient, the cheapest fixer MG is sufficient. It is also the only one that can be applied for advection with linear interpolation and would be recommended for any model using such mixed-approach.
3. ZE fixer results to an accurate advection scheme and generates small increments. If quasi-monotonicity is not essential should be the best option for fields having background values away from zero.
4. Currently the BC fixer is recommended for simulations with chemical tracers because it is one of the cheapest and performs well in advecting correlated tracers (cf. Fig. 7).
5. For volcanic plumes, BC is also sufficient.

## Mass fixer algorithms for tracer transport

M. Diamantakis and  
J. Flemming

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

MFAs may be inappropriate at non-hydrostatic, cloud-resolving scales. Future tests will include these regimes. On going developments in the PantaRhei project (ECMWF, 2013) will provide opportunities towards a strictly mass-conserving scheme for these regimes. Until such developments materialise, MFAs can provide a practical alternative for the applications supported by IFS and are attractive due to their low computational cost.

*Acknowledgements.* The authors wish to thank S. Malardel for providing her idealised case study code and John McGregor for providing reference material for implementing one of the algorithms examined here.

The authors also gratefully acknowledge the help of E. Källén and N. Wedi who provided constructive review comments on this manuscript.

Johannes Flemming contribution was funded by the MACC-II project of the European Union's Seventh Framework Programme (Grant Agreement no. 283576.)

## References

- Bermejo, R. and Conde, J.: A conservative quasi-monotone semi-Lagrangian scheme, *Mon. Weather Rev.*, 130, 423–430, 2002. 783, 784
- Bermejo, R. and Staniforth, A.: The conversion of semi-Lagrangian advection schemes to quasi-monotone schemes, *Mon. Weather Rev.*, 120, 2622–2631, 1992. 788, 790
- Diamantakis, M. and Flemming, J.: Global Mass Fixer Algorithms for Conservative Tracer Transport in the ECMWF Model, Tech. Rep. 713, ECMWF, 2013. 791
- ECMWF, R.-D.: IFS Documentation, available at: <http://www.ecmwf.int/research/ifsdocs/CY38r1/IFSPart3.pdf> (last access: 20 January 2014), 2012. 780
- ECMWF: EU PantaRhei Project at ECMWF, available at: [http://www.ecmwf.int/research/EU\\_projects/PANTARHEI/index.html](http://www.ecmwf.int/research/EU_projects/PANTARHEI/index.html) (last access: 20 January 2014), 2013. 800
- Flemming, J. and Huijnen, V.: IFS Tracer Transport Study, available at: [http://www.gmes-atmosphere.eu/documents/deliverables/g-rg/ifs\\_transport\\_study.pdf](http://www.gmes-atmosphere.eu/documents/deliverables/g-rg/ifs_transport_study.pdf) (last access: 20 January 2014), 2011. 780

## Mass fixer algorithms for tracer transport

M. Diamantakis and  
J. Flemming

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Flemming, J. and Inness, A.: Volcanic sulphur dioxide plume forecasts based on UV-satellite retrievals for the 2011 Grímsvötn and the 2010 Eyjafjallajökull eruption, *J. Geophys. Res.*, 118, 172–189, 2013. 797
- Gravel, S. and Staniforth, A.: A mass-conserving semi-Lagrangian scheme for the shallow-water equations, *Mon. Weather Rev.*, 122, 243–248, 1994. 788
- Hazeleger, W., Wang, X., Severijns, C., Ştefănescu, S., Bitanja, R., Sterl, A., Wyser, K., Semmler, T., Yang, S., van der Hurk, B., van Noije, T., van der Linden, E., and van der Wiel, K.: EC-Earth V2.2: description and validation of a new seamless earth system prediction model, *Clim. Dynam.*, 39, 2611–2629, 2012. 780
- Hollingsworth, A., Engelen, R. J., Benedetti, A., Dethof, A., Flemming, J., Kaiser, J. W., Morcrette, J.-J., Simmons, A. J., Textor, C., Boucher, O., Chevallier, F., Rayner, P., Elbern, H., Eskes, H., Granier, C., Peuch, V.-H., Rouil, L., and Schultz, M. G.: The Global Earth-system Monitoring using Satellite and in-situ data (GEMS) Project: towards a monitoring and forecasting system for atmospheric composition, *B. Am. Meteorol. Soc.*, 89, 1147–1164, 2008. 780
- Kaas, E.: A simple and efficient locally mass-conserving semi-Lagrangian transport scheme, *Tellus A*, 60, 305–320, 2008. 779
- Lauritzen, P. and Thuburn, J.: Evaluating advection/transport schemes using interrelated tracers, scatter plots and numerical mixing diagnostics, *Q. J. Roy. Meteor. Soc.*, 138, 906–918, 2012. 795, 796
- Lauritzen, P., Nair, R. D., and Ullrich, P. A.: A conservative semi-Lagrangian multi tracer transport scheme (CSLAM) on the cubed-sphere grid, *J. Comput. Phys.*, 229, 1401–1424, 2010. 779
- MacGregor, J.: C-CAM Geometric Aspects and Dynamical Formulation, Tech. Rep. 70, CSIRO Atmospheric Research, Aspendale, Vic., 2011. 783, 788
- Priestley, A.: A quasi-conservative version of the semi-Lagrangian advection scheme, *Mon. Weather Rev.*, 121, 621–629, 1993. 783, 787
- Rasch, P. and Williamson, D.: Computational aspects of moisture transport in global models of the atmosphere, *Q. J. Roy. Meteor. Soc.*, 116, 1071–1090, 1990. 783
- Ritchie, H. and Tanguay, M.: A comparison of spatially averaged Eulerian and semi-Lagrangian treatment of mountains, *Mon. Weather Rev.*, 124, 167–181, 1996. 780

## Mass fixer algorithms for tracer transport

M. Diamantakis and  
J. Flemming

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Ritchie, H., Temperton, C., Simmons, A., Hortal, M., Davies, T., Dent, D., and Hamrud, M.: Implementation of the semi-Lagrangian method in a high-resolution version of the ECMWF Forecast Model, *Mon. Weather Rev.*, 121, 489–514, 1995. 778, 781, 790
- Sørensen, B., Kaas, E., and Korsholm, U. S.: A mass-conserving and multi-tracer efficient transport scheme in the online integrated Enviro-HIRLAM model, *Geosci. Model Dev.*, 6, 1029–1042, doi:10.5194/gmd-6-1029-2013, 2013. 779
- Stenke, A., Grewe, V., and Ponater, M.: Lagrangian transport of water vapor and cloud water in the ECHAM4 GCM and its impact on the cold bias, *Clim. Dynam.*, 31, 491–506, 2008. 793
- Thuburn, J.: Some conservation issues for dynamical cores of NWP and climate models, *J. Comput. Phys.*, 227, 3715–3730, 2008. 778
- Ulrich, P., Jablonowski, C., Kent, J., Lauritzen, P., Nair, R., and Taylor, M.: Dynamical Core Model Intercomparison Project (DCMIP) Test Case Document, Tech. rep., NCAR, 2012. 795
- Wilks, D.: *Statistical Methods in the Atmospheric Sciences*, Vol. 100 of 10, 3 edn., Elsevier, 2011. 794
- Zerroukat, M.: A simple mass conserving semi-Lagrangian scheme for transport problems, *J. Comput. Phys.*, 229, 9011–9019, 2010. 783, 786
- Zerroukat, M. and Allen, T.: A three-dimensional monotone and conservative semi-Lagrangian scheme (SLICE-3D) for transport problems, *Q. J. Roy. Meteor. Soc.*, 138, 1640–1651, 2012. 779

## Mass fixer algorithms for tracer transport

M. Diamantakis and  
J. Flemming

**Table 1.** List of 12 month forecast experiments.

| Experiment             | Description   |
|------------------------|---|
| control                | operational setup: cubic interpolation on $Q$ with DEF limiter, linear interpolation on CLWC, CIWC, CRWC, CSWC (no fixer) |
| (control, MG)          | operational setup adding MG fixer   |
| (cubic qm)             | cubic on $Q$ with DEF limiter, CLWC, CIWC, CRWC, CSWC (no fixer)  |
| (cubic qm, BC)         | cubic qm setup adding BC fixer on above moist fields  |
| (cubic BSqm)           | cubic qm setup using BS limiter instead of DEF  |
| (unfiltered cubic, PR) | pure cubic Lagrange for moist fields, quasi-monotone advection by PR algorithm on moist fields                            |
| (cubic BSqm, BC)       | cubic BSqm setup adding BC fixer on moist fields  |
| (cubic BSqm, MG)       | cubic BSqm setup adding MG fixer on moist fields  |
| (cubic BSqm, ZE)       | cubic BSqm setup adding ZE fixer on moist fields  |

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc


Printer-friendly Version

Interactive Discussion




## Mass fixer algorithms for tracer transport


M. Diamantakis and  
J. Flemming




(a) T159 L60



(b) T159 L91



(c) T1279 L91



(d) T1279 L137

**Fig. 1.** Mass conservation error of the IFS SL advection scheme as a percent of initial global mass for ozone,  $Q$ , CLWC, CIWC at different horizontal and vertical resolutions using a quasi-monotonic bi-cubic or a linear (LIN, CLWC and CLIC only) interpolation scheme.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back


Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion





**Fig. 2.** Mass conservation errors as a percentage of initial global mass for Q, CLWC at T1279 L137 resolution forecast with/without PR and BC MFAs.

## Mass fixer algorithms for tracer transport

M. Diamantakis and J. Flemming

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back



Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion





(b) BC fixer

**Fig. 3.** Specific humidity ( $Q$ ) and BC fixer increment for  $Q$  (in  $\text{kg kg}^{-1}$ ) at  $t + 24$  h and 700 hPa height from a T1279 L137 forecast.

**Mass fixer algorithms for tracer transport**

M. Diamantakis and J. Flemming

Title Page

Abstract Introduction

Conclusions References

Tables Figures

⏪ ⏩

◀ ▶


Back Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion





**Fig. 4.** Zonally-averaged and time-averaged (24 h) vertical cross sections for  $Q$  **(a)** and different MFA increments (in  $\text{kg kg}^{-1}$ ) for  $Q$  **(b–f)**. Vertical axis: model level number.

# GMDD

7, 777–814, 2014

## Mass fixer algorithms for tracer transport

M. Diamantakis and  
J. Flemming




Fig. 4. Continued.

Title Page

Abstract Introduction

Conclusions References

Tables Figures

⏪ ⏩

◀ ▶

Back Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## Mass fixer algorithms for tracer transport

M. Diamantakis and  
J. Flemming

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶


Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**Fig. 5.** 48 h timeseries of global rms-norms (left) and max-norms (right) of MFAs increments for CLWC expressed as a percentage of the rms-norm of the field.

## Mass fluxer algorithms for tracer transport

M. Diamantakis and  
J. Flemming

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶


Back

Close


Full Screen / Esc

Printer-friendly Version


Interactive Discussion




(a) control - ERAI



(b) [cubic BSqm] - control



(c) [unfiltered cubic, PR] - control




(d) [cubic BSqm, BC] - control


**Fig. 6.** Experiments with BS limiter described in Table 1. Difference of vertical cross-sections of zonally-averaged annual mean temperature fields. **(a)** Difference (in Kelvin) of control forecast from ERA-Interim. **(b–f)** Difference (in Kelvin) of experiments from control forecast.

## Mass fluxer algorithms for tracer transport

M. Diamantakis and  
J. Flemming



(e) [cubic BSqm, MG] - control



(f) [cubic BSqm, ZE] - control

Fig. 6. Continued.

Title Page

Abstract Introduction

Conclusions References

Tables Figures

⏪ ⏩

◀ ▶

Back Close

Full Screen / Esc


Printer-friendly Version

Interactive Discussion



## Mass fixer algorithms for tracer transport

M. Diamantakis and  
J. Flemming



**Fig. 7.**  $q_1 - q_2$  ( $xy$  axis) scatter plots for correlated tracers at  $t = 6$  days. Scatter points ( $q_1, q_2$ ) at  $t = 0$  follow the upper (parabolic) black dashed-dotted curve.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back


Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion





**Fig. 8.** Relative mass residual in volcanic plume simulations ( $\text{SO}_2$ ) for different schemes.

## Mass fixer algorithms for tracer transport

M. Diamantakis and  
J. Flemming

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back


Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion





**Fig. 9.** Comparison of volcanic plume simulation with and without mass fixer using quasi-monotone cubic Lagrange at T1279 L91 resolution. The plotted quantity is the total  $\text{SO}_2$  content (in  $\text{kg m}^{-2}$ ) per model grid-point column. Experiments defined as in Table 1.

**Mass fixer algorithms for tracer transport**

M. Diamantakis and J. Flemming

|                          |              |
|--------------------------|--------------|
| Title Page               |              |
| Abstract                 | Introduction |
| Conclusions              | References   |
| Tables                   | Figures      |
| ⏪                        | ⏩            |
| ◀                        | ▶            |
| Back                     | Close        |
| Full Screen / Esc        |              |
| Printer-friendly Version |              |
| Interactive Discussion   |              |

