

We would like to thank both reviewers for their useful comments on the manuscript. Both reviewers touched on several long standing discussions in land surface modelling, i.e., validation, optimization and the use of PFTs. We make use of the open discussion to better justify the choices we made but we also decided to keep those justification as concise as possible in the manuscript itself. The reason for this is that we prefer to focus on the ORCHIDEE-CAN developments rather than providing a more general discussion on the current opinions and issues in land surface modelling. As an outcome of this approach, the review comments are largely addressed through new and revised figures and tables (which are the result of several additional simulations), whereas substantial changes and edits to the text are limited to a few sections. We believe, however, that with this approach we were able to address most of the reviewers' comments.

The page and line numbers where changes have been made to the manuscript are included at the end of each reply and the lines are indicated in green in the revised manuscript.

Response to referee 1

General comments

Comment 1

This manuscript presented a new version of ORCHIDEE including a new canopy scheme (-CAN) applicable to forest management simulations. As claimed by the authors, forest management is one of the most important and uncertain factors in the global terrestrial carbon cycle, affecting the future climate change. For example, recent carbon-related debates on climate change mitigation, such as REDD+ and BECCS, are deeply related to forest management. The development of the ORCHIDEE-CAN seems to be a great progress toward taking account of forest management in global terrestrial models. The manuscript seems a bit lengthy, because model structure, parameterization, and validation were described separately for each process such as allocation and energy budget. I recommend revising the structure of manuscript in a more simplified manner.

We agree that the manuscript is rather lengthy and felt this is partially due to the extensive tables. We therefore moved the tables with the parameter values (Table 3, 4, 5 and 6) to the supplementary material. The approach and data differ substantially between the sections on model structure, parametrization and validation which seems to justify treating these items in separate sections. A structure that repeats the same subsections for model structure, parametrization and validation where possible, enables a detailed description in an easily navigable structure for the reader. In addition, integrating all information in one coherent paper rather than splitting it into several papers, avoids duplication of the introduction, cross references and leaving it up to the reader to search and combine information from different sources.

Comment 2

Another concern on the manuscript is that more validation (e.g., site-level comparison with filed data) would be necessary for this kind of study. Also, an intimate comparison with other models (e.g., ORCHIDEE-trunk and ORCHIDEE-FM by Ballassen et al.) is strongly recommended to clarify characteristics of the new model. Although the authors showed region-level, snap-shot comparisons with data, I could not be convinced that the new model simulates long-term transition of carbon budget including the impacts of human and natural disturbances. In conclusion, this manuscript needs major revision before being accepted for publication.

We would like to thank the reviewer for this comment as it gives us the opportunity to add some

details to the validation but also to discuss some more philosophical issues of model validation.

As stated in the conclusions of the manuscript, we recognize that additional validation (and parametrization) is recommended for other regional applications, higher resolution studies and/or studies with a different objective (for example studies that relate to croplands or grasslands). Additional validation will be the subject of follow-up studies (see for example Ryder et al.2014), and are not the focus of the current manuscript. Nevertheless we added more detail to the large-scale validation by performing additional testing:

- We performed an additional validation at higher resolution by comparing simulated basal area and tree diameter at the species level with observed site-level data (depending on the species 639-8227 plots) over France (Figure 6, section 5.7).*
- To give the reader more insight in the behaviour of the model regarding the impact of human disturbances we added an evaluation of the impact of different forest management strategies at the pixel level (Figure 5, section 5.7).*
- A test was added in which the effect of species-level parametrization versus MTC parametrization was assessed (Figure S2, section 5.1)*
- Maps showing the differences between observed and simulated GPP and basal area are shown (Figure 4)*

Although a comparison between ORCHIDEE-CAN and previous versions of ORCHIDEE seems straightforward, it cannot be performed for many of the new features which were the objective of these developments. This problem is already apparent from the European-wide validation where we could validate seven variables for ORCHIDEE-CAN of which only four exist in ORCHIDEE-trunk (Figure 3). In addition many of the more fundamental changes (i.e. the layering of photosynthesis, the implementation of water stress and the overhaul of the allocation scheme) make it impossible to assign differences in model version to specific changes. Recoding the model in a more modular way where changes can be switched on and off would not overcome this issue because in the case of the new albedo scheme the changes to allocation are essential to the albedo calculations. Similar code dependencies exist for water stress and allocation, albedo-photosynthesis, forestry-allocation, and water stress and photosynthesis (Figure 1).

Given these limitations we deemed it more important to document the performance of the model against observation-based spatially explicit product than against other model versions (Figures 2, 4 and 6). After all, the presented validation over Europe is performed on the last ten years of 160 years of simulation for biogeochemical fluxes, biophysical fluxes and forest management related vegetation characteristics (section 5). The results show the ability of the model to sufficiently capture large-scale spatial patterns and interannual variability over Europe. Where the comparison can be made, ORCHIDEE-CAN outperforms the trunk over Europe for four variables (Figure 2 and 3, Table 3), where at the same time empirical approaches were replaced by more process-based approaches. The fact that a more process-based code could outperform ORCHIDEE-trunk made us conclude that the model performs well enough for a first forest-related application over Europe.

Specific comments

Page 8568 Line 3–4 In addition to Dixon et al. (1994), Pan et al. (2011, Science 333, 988–993) should be cited.

The citation was added.(Line 26)

Page 8568 Line 18 By definition of albedo, “darkness” should be replaced by “reflectance” or “whiteness”.

This was changed.(Line 39)

Page 8569 Line 5–6 The statement “Vegetated surfaces are sub-divided in patches of different plant functional types” would be incorrect. Several DGVMs (e.g., LPJ-Guess and SEIB-DGVM) have individual-based vegetation dynamics.

The statement was adjusted. (Line 51)

Page 8571 Line 11–12 “to confirm convergence” Do you mean stabilization of carbon budget or agreement between the different spin-up methods? This statement is ambiguous and I recommend clarifying.

The statement was clarified. (Line 110)

Page 8578 Line 5 For “sapwood area”, do you mean “sapwood cross-section area”? Please clarify.

The statement was clarified. (Line 290)

Page 8580 Line 10 Eq. (15) Please add a reference for Eq. (15).

A reference was added. (Line 335)

Page 8581 Line 24 A recent study (Sun et al. 2014, PNAS 111, 15774–15779) implied that leaf mesophyll conductance in DGVMs. Your new model, ORCHIDEE-CAN, does not include mesophyll control in the hydraulic architecture. Is it correct? If so, please discuss potential uncertainties due to neglect of the mesophyll conductance.

Although the photosynthesis model and its implementation could easily account for mesophyll conductance (Yin & Struik, 2009) leaf mesophyll conductance is indeed not accounted for in our photosynthesis model in order to maintain compatibility between the model formulation and its parametrization This is explained in the manuscript, and the potential implication of neglecting mesophyll conductance is discussed (Lines 594-603).

Page 8584 Line 14–16 What kind of numerical method was used to get a solution by iteration (e.g., the Newton-Raphson method)?

A stationary iterative method was used. This information was added to the manuscript. (Line 419)

Page 8585 Line 21–22 No symbol in Eq. (29) has overbar. Please check.

The overbar was added to the equation (Equation 29).

Page 8590 Line 16–17 “ $R_{s,i}$... may be calculated using...”. What do you mean for “may be calculated”? Do you mean “was calculated”?

The suggested change was made (Line 558)

Page 8591 Line 20 Remove “)” after “model”.

The additional bracket was removed.

Page 8593 Line 4 No natural disturbance such as wildfire, insect outbreak, and windthrow, was explicitly included in the present model. It is correct?

Yes, this is correct. These natural disturbances are all assumed to be included in natural mortality. It is explained in the manuscript on lines 656-664.

Page 8597 Line 25 Can you give a list of the 126 site observations as a supplementary material?

The list of sites and their meta-data is given in Campioli et al. (in prep).

Page 8597 Line 28 What is the definition of “acceptable GPP”? Please explain. Otherwise, replace “acceptable GPP” by other words such as “GPP sufficiently close to observations”.

This section has been revised to address comment 6 of reviewer 2 and the statement was removed from the text.

Page 8604 Line 11–13 I agree the statement that there is no data-derived net carbon flux data. However, there should be soil carbon stock maps (e.g., Harmonized World Soil Database) that can be used for model validation. In this regard, there is a need for carbon spin-up.

Thank you for this suggestion, Soil carbon is indeed an essential variable when studying the carbon pools, fluxes and their budget. However, the soil carbon module within ORCHIDEE-CAN was not changed compared to ORCHIDEE-trunk (see Table 1). Therefore, an evaluation of this module is not a priority given the many modules that were changed between ORCHIDEE-trunk and ORCHIDEE-CAN. A validation against the proposed soil data will be part of the first application of ORCHIDEE-CAN in which in-situ carbon sequestration or net ecosystem production is simulated,

Recently, an evaluation of the soil carbon module in ORCHIDEE against empirical data from the Harmonized World Soil Database (HWSD) and the Northern Circumpolar Soil Carbon Database (NCSCD) was published by Todd-Brown et al. (2013). This study shows that the global soil carbon stocks as simulated by ORCHIDEE falls within the 95% confidence interval for both the soil carbon stock of HWSD and NCSCD. Regarding the spatial distribution of soil carbon stocks, ORCHIDEE showed good agreement with HWSD at the biome level, but mediocre agreement at the 1° scale. ORCHIDEE generally overestimated tundra but underestimated desert and shrublands (Todd-Brown et al., 2013). Given the same approach and parameters are used in ORCHIDEE-CAN as in ORCHIDEE-trunk, a rather similar behaviour to the one reported by Todd-Brown et al. (2013) is expected for ORCHIDEE-CAN.

Page 8606 Line 20 Additionally, the ORCHIDEE-CAN might have a few tuning parameters (e.g., plant water storage and soil-root resistance, p 8598 line 14–15), leading to higher likelihoods. If correct, this fact should be mentioned.

Thank you for bringing this up. Model tuning is an interesting issues that inherently comes with model development. While developing ORCHIDEE-CAN we aimed for a more consistent approach and tried to reduce the number of tuning parameters compared to the trunk (i.e. ORCHIDEE-CAN no longer has a cap on LAI). We made use of one tuning parameter to represent plant water storage and soil-root resistance pending a more mechanistic model representation. Besides this parameter we tried to avoid hiding model flaws in the parametrization by extracting the parameters from a wide range of sources including original observations, large databases, primary research reports and remote sensing products. By using this transparent parametrization approach we should have reduced the amount of tuning parameters compared to the trunk version. In the absence of new developments the level of tuning typically increases with the number of years that a model is used as more and more applications allow for more in-depth testing which might result in more tuning. From this perspective ORCHIDEE-CAN is at present most likely less tuned than the trunk. It should be noted that as long as the tuning is within the physically observed range of the parameters, it is justifiable and does not necessarily hints at a model deficiency.

Page 8612 Line 7 Can you show “this finding” by some materials (e.g., figure or map)?
We have added a map that shows the spatial variability of simulated and observed basal area over Europe (Fig 4).

Page 8613 Line 14–16 I don’t agree with the statement that “PFTs have no meaning outside the modeling community”, because plant biologists use a similar concept called “guild”. Also, the concept of “functional group” has been widely used in plant ecophysiology (e.g., W. Larcher 1995, Physiological Plant Ecology, ecophysiology and stress physiology of functional groups, 3rd ed., 506

pp., Springer-Verlag). As there are more than 200,000 plant species on the Earth, global models inevitably use some simplifications such as the concept of PFTs, for the time being.

Thank you for this comment which refers to the long standing debate on PFTs. It is beyond doubt that PFTs have a role to play in modelling and other scientific communities that have to deal with the huge plant diversity but at the same time the approach has strong limitations for outreach and communication outside the scientific community. Furthermore, some of the initial PFTs pool such diverse plants, for example Eucalyptus and Mediterranean Oaks are in the same PFT because both are temperate evergreen broadleaved species that the approach hampers parametrization of ecophysiological processes such as plant water conductivity and photosynthesis. Furthermore if wood production is among the output variables (which is the case for ORCHIDEE-CAN) the differences in wood use alone could justify abandoning the PFT concept in favour of a species/genus based parametrization (i.e. ORCHIDEE-CAN) or a hybrid parametrization of PFTs and plant trait relationships (van Bodegom et al., 2014; Reichstein et al., 2014). Taking into account the above and the comment of the reviewer, we changed the wording to “PFTs are of limited use outside the scientific community”. (Line 1214)

Page 8637 Table 3 Several values in Table 3 is “Table 3”. Do you mean Table 4, 5, or 6? Otherwise, please explain.

The references to the tables were corrected.

Page 8646 Figure 2 and Page 8647 Figure 3 Please add units to each y-axis.

Units are now added (Fig 2 and 3).

Response to referee 2

1. This paper is part of a series of papers that are currently under discussion or submitted (e.g. MvGrath et al.; Ryder et al.; . . .), all these papers are related to the different aspects of the developed ORCHIDEE-CAN model. This fact is not a problem at all for me, because I understand that the variety of aspects covered in the series of papers, can never be covered in one manuscript. However I think that the motivation of the Naudts et al. manuscript (and the relation of this paper to the other submitted manuscripts) should be presented better in the introduction of the paper.

The motivation and the relation to the other submitted manuscripts are explained in the introduction (Lines 67-75): “The aim of this study is to describe the model developments and parametrization within ORCHIDEE-CAN and to evaluate its performance. ORCHIDEE-CAN is validated against structural, biophysical and biogeochemical data at the European scale. A more detailed description and evaluation of the new multi-layer energy budget and multilevel radiative transfer scheme is given by Ryder et al. (2014), Chen et al; (In prep.), McGrath et al. (In prep.). A new forest management reconstruction which is needed to drive forest management in ORCHIDEE-CAN is presented in McGrath et al. (In prep.) and the interactions between forest management and the new albedo scheme have been discussed by Otto et al. (2014)”.

2. At the end of the introduction I missed a paragraph that clarifies the specific goals of the paper. I did not read any research questions. The paper appeared to me as a real model description/validation paper. It would be good to state this very clearly at the end of the introduction. In order to tell the reader what she/he can expect. For me this is not a hypothesis-drive research paper, but rather a model description paper. The last paragraph of the current introduction nicely motivates why the model development is needed. Adding 1 paragraph that tells the reader what will exactly be presented in the paper should solve my comment. This paragraph should also

contain a sentence that the manuscript is focusing on temperate and boreal forests in Europe, and that tropical forest are not treated (yet) in this manuscript.

This is indeed a model description/validation paper. We added a paragraph to the introduction to frame the content of the manuscript. See our reply to the previous

3. The validation –which is a very important part for a manuscript of this type– is visually presented in a very condensed way (only 2 figures). I think the validity of the model would be much more convincing for the readers of GMD if some additional graphs would be shown, to illustrate what is discussed in the validation section of the manuscript. Some maps illustrating the spatial variability of some key model outputs would be very informative, both for cases where ORCHIDEE-CAN performs very good or for cases where the model performs not good (e.g. the spatial variability of height and basal area).

We included three additional validation figures:

- 1. a European map which compares simulated and observed GPP (good performance) and basal area (weak performance) and which illustrates the spatial variability of the model output (Figure 4), a figure showing regional validation results of tree diameter over France (Fig 6),*
- 2. a figure showing the impact of the different forest management strategies (Fig 5).*

In combination with the figures that were already available, the new figures give a better insight in the performance of the model and better details the performance of the modules simulating forest management.

4. The assumption on the circumference class distribution, using the “target distribution” (page 8594) worries me. I have the feeling that this kind of assumption can have a large effect on the simulated results. This is an essential point for a model that attempts to simulate the impacts of forest management on the climate. It remains unclear how sensitive the model is for this assumption. The authors need to elaborate more on this point.

A more intuitive numerical approach would set a fixed number of diameter classes with fixed boundaries and would let the distribution to vary. Such an approach would use computer resources as at any point in time several diameter classes would be empty. In addition, for different points in time, a different number of diameter classes would contain trees. This would not be a problem until only one diameter class would be populated. At that point the rule of Deleuze and Dhoté which is required to differentiate tree growth for different diameter classes would collapse and the model would not be able to recover from that situation. To avoid such a collapse and make the best use of the computer resources (both computational and storage) we choose for the counter intuitive solution of fixing the number of diameter classes and the number of trees per diameter class but letting the boundaries of the classes to vary. Given that both alternatives have the same degree of freedom, they should have the same flexibility and capacities. In our approach a normal distribution could be approximated by using three diameter classes with a prescribed number of trees per class but with very narrow and very similar diameter ranges for each class. Initial tests showed a limited effect of the number of diameter classes and the initial distribution on the model behaviour for stand level variables (i.e., stand density, basal area, diameter, height). However, the size and growth of individual trees is very sensitive to these settings. Finding robust initial distributions to correctly simulate tree ring width is under-way but is falls outside the scope of this study and the initial objectives of developing a model code that can simulate the large scale biogeochemical and biophysical effects of forest management over Europe. We added a concise comment on the issue in the manuscript on Lines 696-698. The comment reads “Note that the

boundaries of each diameter class are recalculated at each time step, this approach is a numerically efficient alternative to fixing the boundaries of each diameter class but letting the distribution to vary.”

5. The authors explain in detail how the model parameter values were determined for Europe (section 4.2), but the manuscript is lacking a discussion on how the model can be applied globally. It would be interesting to see how ORCHIDEE-CAN performs over Europe if only the original MTC classification was used. This would allow estimating how important the specific parameter value settings are to get good results over Europe.

Initial tests showed that the forest grows reasonably well across the globe but at present model performance was not tested outside of Europe. Global runs would make use of the MTC parameters which are derived by averaging the parameters of the species within that MTC. If less details are required, the same MTC parameters can be used over Europe to perform simulations for the original 13 MTC classes which are distinguished in ORCHIDEE-trunk. To give an idea of the impact of the species level parametrization we ran a new 160 year simulation with the MTC parameters and validated the simulation in the same manner as the species level simulation (Fig. S2). For the seven parameters under study, the effects of introducing species are almost negligible. This test result is added to the manuscript (Lines 974-978): “In ORCHIDEE-CAN the PFT concept was refined by parametrizing the main European tree species groups (section 4.1). To evaluate the effect of the species parametrization, we performed a companion simulation for the configuration described above, but at the MTC level. Model performance was barely affected by the use of the MTC parameters, compared to the simulation with the species parameters (Fig. S2 for RMSE scores).

6. Probably my most important general remark deals with the “tuning” of some of the parameters of the model. The authors are not explicit about this tuning process. For example (1) the kls parameter was “tuned to match observations of GPP, ET and LAI”; how was this tuning performed? With an optimization procedure? Were the different observation weighed in this process? (2) The basal rate of autotrophic respiration was optimized using “a rigorous statistical framework (Tarantola 2005)”: how was this framework implemented and used for the ORCHIDEE-CAN model? Was this done in a similar way as this has been done in the past for the standard ORCHIDEE model (Santaren et al. 2007 GBC)? (3) the allocation scheme required parameter values from fitting regression models to inventory data. (4) on page 8599 the authors mention posterior and prior standard deviations. Is this “posterior” after optimization? What kind of optimization? (5) page 8602, “kalfa1 and kbeta1 were estimated by fitting”. . . It is really needed that the authors are explicit on how this tuning process was performed, in order to judge the quality of the presented simulation results. An additional question is what the order was of the different tuning steps? If you tune parameter x against datasets B and C and after that you tune parameter y against dataset C, you might get another result compared to a situation where you fit y first and x after that . . . I am aware that a “perfect” tuning procedure does not exist for a very complex model like ORCHIDEE-CAN. However, for me the parameter optimization phase is a very important step in model development. Therefore the authors need to bring the description of their parameter tuning at the same high level as their model description in this paper. The tuning of the parameters should be described in a transparent way. Maybe a scheme describing the process and the used datasets will help to do this. In addition, in the validation section I miss a discussion on the impact of the tuning process on the performance of the model for Europe. An independent validation for North America, without tuning, would shed an interesting light on this issue.

We would like to thank the reviewer for raising this issue. We re-wrote substantial parts of the parametrization section. First we added a paragraph describing the order and steps that were

performed during the parametrization process (Lines 704-722). This paragraph is followed by sections that provide more details. Attention was paid to address the concerns of the reviewer (See lines 756-762, 824-826-, 865, 888-901 and 921-927) by adding more information on the procedures, by better describing the parameters that were parametrized, by better referring to the supplementary information that contains detailed and complete information for each parameter and by better describing the data and simulations that were used in the process.

An independent validation for North America as suggested by the reviewer would require a species or species-group map for North America which we currently do not have access to. However, outside Europe the current model set-up only uses MTC parameters derived from European species because the focus of the manuscript and the first applications of ORCHIDEE-CAN is on Europe. Therefore, we chose to perform the validation only over Europe. To give the reader an idea of the impact of the optimization, we ran the model at the pixel-level with the prior values of k_{cmaint} and k_{lsmin} for Scots pine and compared the results to the simulation with the optimized values (Fig. S1). The text now reads: “Limited tests over a period of 100 years in a Scots pine forest at 51 – 52 ° N, 13 – 14 ° E (Fig. S1) suggested that optimizing k_{cmaint} and k_{ls} had the largest effect on the maximum LAI, which decreased with almost 17% after optimization compared to a simulation with prior parameter values. Mean annual GPP, mean annual transpiration and basal area decreased with, respectively 6, 6 and 7% compared to a simulation with prior parameter values (Fig. S1). ”

Specific comments

1. Page 8574, line 23: I was a bit confused by the statement that the radiation scheme (which is used in this manuscript I assume) is not “available” yet. . .

This statement was rephrased.

2. Page 8574, line 26: the energy budget is calculated “implicitly”: what does this mean?

As outlined in (Polcher et al., 1998), the use of an implicit solution for coupling between the atmospheric model and the surface layer model is the only way to keep profiles of temperature and humidity synchronised across the two models when the coupled-model is run over large time steps. The difference between explicit and implicit schemes is that an explicit scheme will calculate each value of the variable (e.g., temperature and humidity) at the current time step entirely in terms of values from the previous time step. An implicit scheme requires the solution of equations written only in terms of those at the current time step. The term implicit is now explained in the manuscript (Lines 195-209).

3. Page 8574, line 27: you mention the time step for the first time here. I suggest mentioning the time step of the model early in section 2. This is important information for readers not familiar with ORCHIDEE.

The 15 minute time step is the time step of the coupling to the atmospheric model, we only mention it here because it is the first time we write about the coupling in section 2. We rewrote this part of the text to clarify that the 15 minute time step concerns the time step of the coupling (Line 205). The time steps of the other processes were added to figure 1 to avoid confusion.

4. Page 8576, line 6: “biomass and soil carbon pools are mixed”: what does this mean?

Following a land cover change, the biomass, soil and litter pools of the vegetation that undergoes a land cover change is added to the biomass, soil and litter carbon that already existed for that PFT. If a land cover change results in the loss of an entire PFT, biomass, litter and soil carbon need to be properly re-assigned to other carbon pools. This was clarified in the manuscript by changing the wording as “Following a land cover change, biomass and soil carbon pools (but not soil water columns) are either merged or split to represent the various outcomes of a land cover change (Line

239).

5. Page 8576, line 28: what are the two additional pools that you introduced? (are these the branches and coarse roots?, this was not completely clear for me).

Yes, the two additional pools are the branches and the coarse roots. The manuscript now reads (Lines 258-261): “The inclusion of forest management resulted in two additional carbon pools, branches and coarse roots (i.e., aboveground and belowground woody biomass) and therefore required an extension to the semi-analytical spin-up method (see section 2.1). The semi-analytical spin-up is now run for nine C pools.”

6. Page 8577, line 9: maintenance respiration is a function of nitrogen concentration. Does this mean that the ORCHIDEE-CAN model contains the N-cycle of the O-CN model? This is not clear from the current text.

No, the ORCHIDEE-CAN model does not contain the N-cycle of O-CN. However, the maintenance respiration is adjusted for an estimated nitrogen effect by assuming that the nitrogen distribution decreases exponentially with LAI, following the Beer-Lambert law. To avoid confusion on the use of the N-cycle, the reference to Zaehle and Friend (2010) is removed, while the Beer-Lambert assumption is added to the text (Line 268).

7. Page 8579, lines 16-19: “. . . the same parameter values . . .” the same as what? This sentence was confusing for me.

This sentence was rephrased as: “The allometric relationships between the plant components and the hydraulic architecture of the plant are both based on the pipe model theory, hence, both the allocation and the hydraulic architecture module use the same parameter values for root and sapwood conductivity”. (Lines 320-322)

8. Page 8581: line 15-17: This is a very important statement; it should be included at the start of section 3.1.

The statement was moved to the start of section 3.1. (Lines 283-288)

9. Page 8586, title 3.4: here and elsewhere in the text the authors use “albedo” when talking about the entire radiation scheme. I suggest to use the wording “radiation scheme for tall canopies” here. We replaced “albedo” by “radiation scheme for tall canopies” where appropriate.

10. Page 8645, fig 1: I suggest to rename the box “albedo” and make it “radiation scheme”
This was changed in the figure (Fig 1).

11. Page 8587, Line 11: “similar equations” are used for diffuse sources. Are these equations explained in another publication? These equations should be available for the reader, as accounting for diffuse radiation is an important issue which is lacking in the standard ORCHIDEE version.
The equations have been published in Pinty et al. (2006). We added a reference for the equations in the manuscript (Line 483).

12. Page 8587, line 21: what is the impact of the assumption of an extinction coefficient of 0.5? this is an important assumption with probably a strong impact on the results and the model performance. I suggest adding some discussion on this assumption.

In fact we could have used any value for the extinction coefficient to calculate the effective LAI as long as we also would have used this effective LAI with the same extinction coefficient in direct mode (only the effective LAI would be changed, not the transmitted flux). However, we chose to use 0.5 as an extinction coefficient to be able to use the two-stream inversion derived by Pinty et al. (2011), which was derived using 0.5 (mathematically the simplest possible solution). To avoid confusion we shortly mention this in the manuscript. (Lines 491-492)

13. Page 8592. I found the wording to describe the mortality a bit strange and sometimes confusing. “A whole PFT is now killed”: what does this mean? All trees belonging to 1 PFT within 1 grid cell? “when it comes to time to actually kill the trees”: a strange formulation. “We mainly kill the smallest trees” (page 8593): I would reformulate such statements, for example “we assume that mainly the smallest trees die”. In addition, the meaning of the word “mainly” is too vague in this sentence. . .

This section was largely rewritten (Lines 612-636) to better explain how forest management was implemented (a request of reviewer 1). The specific suggestions and concerns of the reviewer were taken into account at lines 639, 646 and 669.

14. Page 8592, line 19: I suppose it should be “leaf, carbohydrate reserve, and labile pools”. I assume that mortality takes place when all these pools are depleted.

The PFT will be killed if, at the end of the day, the labile pools are empty and there is no carbon available in the leaf or carbohydrate reserve pools to refill it. This is clarified in the manuscript (Lines 646-648).

15. Page 8593, line 4, line 6: at this point the actual time step of the mortality processes in the model was not clear for me. This could be resolved by adding a paragraph that clearly explains the adopted time steps for the different processes in the model.

The time steps of the different processes are added to figure 1.

16. Page 8601, line 24-28: this piece of text belongs to the model description section.

This section will be moved to model description.

17. Page 8604, line 10: it was a bit disappointing to read at this point that the multilayer energy budget was not used for validation in this manuscript. I understand why you did not do this. I suggest mentioning earlier in the text (maybe end of intro) that the validation is restricted to the single layer energy budget.

The complexity and implication of developing and parametrizing a multi-layer energy budget justifies several manuscripts in its own. We therefore refer the readers to Ryder et al. (2014), Chen et al. (in prep) and McGrath et al. (in prep). The use of the single layer energy budget for validation will be mentioned in the introduction (Line 69-70).

Technical comments

All technical comments were taken into account in the revised manuscript.

1. Page 8567, line 14: ‘occurred’ is a bit strange wording according to me.

2. Page 8568, line 6-8: add a reference for this statement

3. Page 8570, line 25: 3000 simulation years

4. Page 8571 line 16: “the largest gains were realized . . .”: please add a reference for this statement.

5. Page 8571, line 26: Pretzsch (2009)

6. Page 8572, line 17: it is not clear what “the latter” is pointing at. (“resistance” in general or “sapwood resistance”?).

7. Page 8572, line 28: you mention the kvmax parameter for the first time here, without explaining what it means, and you haven’t referred to the Table with parameters at this point in the text yet.

8. Page 8572, line 28: you mention for the first time the word ‘trunk’. I think you should explain here that the trunk is the “standard” version of ORCHIDEE. Otherwise this might be confusing for the reader.

9. Page 8580, eq 15: kalfa1, kbeta1: these parameters were not explained in the text.

10. Page 8582, eq 20: pdelta is represented differently in the equation compared to the subsequent sentence (where the greek symbol for delta is used).

11. Page 8585, line 21: I do not see an “overbar” in eq 29.
12. Page 8590, line 16: “may be calculated” is it calculated according to Ball et al. Or not?
13. Page 8590, line 18: “transport of sensible heat flux”. In my view this formulation is a bit awkward. I suggest to say “sensible heat flux” or “transport of sensible heat”. A “flux” always implies transport.
14. Page 8591, line 1: “transport of latent heat flux” same remark as above.
15. Page 8596, line 26: “12 forest species (groups)”
16. Page 8598, line 12: “hydrology”
17. Page 8605, line 12: “Deleuze and Dhote” add publication year.
18. Page 8634: there is no reference to Table 2 in the text.
19. Page 8635, dcirc: I suppose this is “stem circumference”? I suggest to add “stem”
20. Page 8637: I suppose reference to other tables are wrong: 3 should be 4, 4 should be 5.
21. Page 8640: something is wrong with the columns in table 3
22. Page 8641, table 4: “kresid” should be “ktresid”.
23. Page 8646, fig 2: mention in the caption that the greyscale is corresponding to the symbols. (same for fig 3)

References that are not in the manuscript:

- Reichstein, M., Bahn, M., Mahecha, M. D., Kattge, J., & Baldocchi, D. D. (2014). Linking plant and ecosystem functional biogeography. *Proceedings of the National Academy of Sciences*, *111*(38). doi:10.1073/pnas.1216065111
- Todd-Brown, K. E. O., Randerson, J. T., Post, W. M., Hoffman, F. M., Tarnocai, C., Schuur, E. a. G., & Allison, S. D. (2013). Causes of variation in soil carbon simulations from CMIP5 Earth system models and comparison with observations. *Biogeosciences*, *10*(3), 1717–1736. doi:10.5194/bg-10-1717-2013
- Van Bodegom, P. M., Douma, J. C., & Verheijen, L. M. (2014). A fully traits-based approach to modeling global vegetation distribution. *Proceedings of the National Academy of Sciences of the United States of America*, *111*(38), 13733–8. doi:10.1073/pnas.1304551110

Manuscript prepared for Geosci. Model Dev.

with version 2014/07/29 7.12 Copernicus papers of the L^AT_EX class copernicus.cls.

Date: 11 March 2015

A vertically discretised canopy description for ORCHIDEE (SVN r2290) and the modifications to the energy, water and carbon fluxes

Kim Naudts^{1,*}, James Ryder^{1,*}, Matthew J. McGrath^{1,*}, Juliane Otto^{1,10,*},
Yiying Chen¹, Aude Valade¹, Valentin Bellasen², Gonzalo Berhongaray³,
Gerhard Bönisch⁴, Matteo Campioli³, Josefina Ghattas¹, Toon De Groote^{3,11},
Vanessa Haverd⁵, Jens Kattge⁴, Natasha MacBean¹, Fabienne Maignan¹,
Päivi Merilä⁶, Josep Penuelas^{7,12}, Philippe Peylin¹, Bernard Pinty⁸,
Hans Pretzsch⁹, Ernst Detlev Schulze⁴, Didier Solyga^{1,13}, Nicolas Vuichard¹,
Yuan Yan³, and Sebastiaan Luyssaert^{1,*}

¹ LSCE, IPSL, CEA-CNRS-UVSQ, 91191 Gif-sur-Yvette, France

² INRA, 21079 Dijon, France

³ University of Antwerp, 2610 Wilrijk, Belgium

⁴ MPI-Biogeochemistry, Jena, Germany

⁵ CSIRO-Ocean and Atmosphere Flagship, 2600 Canberra, Australia

⁶ METLA, Oulu, Finland

⁷ CSIC, Global Ecology Unit CREAM-CSIC-UAB, Cerdanyola del Valles, Spain

⁸ European Commission, Joint Research Centre, Ispra, Italy

⁹ TUM, Munich, Germany

¹⁰ Helmholtz-Zentrum Geesthacht, Climate Service Center 2.0, Hamburg, Germany

¹¹ VITO, 2400 Mol, Belgium

¹² CREAM, Cerdanyola del Vallès, Spain

¹³ CGG, 91341 Massy, France

* equal contribution to model development and parametrization

Correspondence to: Kim Naudts (kim.naudts@lsce.ipsl.fr)

Abstract.

Since 70% of global forests are managed and forests impact the global carbon cycle and the energy exchange with the overlying atmosphere, forest management has the potential to mitigate climate change. Yet, none of the land surface models used in Earth system models, and therefore none of today's predictions of future climate, account for the interactions between climate and forest management. We addressed this gap in modelling capability by developing and parametrizing a version of the land surface model ORCHIDEE to simulate the biogeochemical and biophysical effects of forest management. The most significant changes between the new branch called ORCHIDEE-CAN (SVN r2290) and the trunk version of ORCHIDEE (SVN r2243) are the allometric-based allocation of carbon to leaf, root, wood, fruit and reserve pools; the transmittance, absorbance and reflectance of radiation within the canopy; and the vertical discretisation of the energy budget calculations. In addition, conceptual changes were introduced towards a better process representation for the interaction of radiation with snow, the hydraulic architecture of plants, the representation of forest management and a numerical solution for the photosynthesis formalism of Farquhar, von Caemmerer and Berry. For consistency reasons, these changes were extensively linked throughout the code. Parametrization was revisited after introducing twelve new parameter sets that represent specific tree species or genera rather than a group of often distantly related or even unrelated species, as is the case in widely used plant functional types. Performance of the new model was compared against the trunk and validated against independent spatially explicit data for basal area, tree height, canopy structure, GPP, albedo and evapotranspiration over Europe. For all tested variables ORCHIDEE-CAN outperformed the trunk regarding its ability to reproduce large-scale spatial patterns as well as their inter-annual variability over Europe. Depending on the data stream, ORCHIDEE-CAN had a 67% to 92% chance to reproduce the spatial and temporal variability of the validation data.

1 Introduction

Forests play a particularly important role in the global carbon cycle. Forests store almost 50% of the terrestrial organic carbon and 90% of vegetation biomass (Dixon et al., 1994; Pan et al., 2011). Globally, 70% of the forest is managed and the importance of management is still increasing both in relative and absolute terms. In densely populated regions, such as Europe, almost all forest is intensively managed by humans. Recently, forest management has become a top priority on the agenda of political negotiations to mitigate climate change (Kyoto Protocol, <http://unfccc.int/resource/docs/convkp/kpeng.pdf>). Because forest plantations may remove CO₂ from the atmosphere, if used for energy production, harvested timber is a substitute for fossil fuel. Forest management thus has great potential for mitigating climate change, which was recognized in the United Nations Framework Convention on Climate Change and the Kyoto Protocol.

35 Forests not only influence the global carbon cycle, they also dramatically affect the water vapour
and energy fluxes exchanged with the overlying atmosphere. It has been shown, for example, that the
evapotranspiration of young plantations can be so great that the streamflow of neighbouring creeks
is reduced by 50% (Jackson et al., 2005). Modelling studies on the impact of forest plantations
in regions that are snow-covered in winter suggest that because of their **reflectance** (the so-called
40 albedo), forest could increase regional temperature by up to four degrees (Betts, 2000; Bala et al.,
2007; Davin et al., 2007; Zhao and Jackson, 2014). Management-related changes in the albedo,
energy balance and water cycle of forests (Amiro et al., 2006a, b) are of the same magnitude as the
differences between forests, grasslands and croplands (Luyssaert et al., 2014). Moreover, changes
in the water vapour and the energy exchange may offset the cooling effect obtained by managing
45 forests as stronger sinks for atmospheric CO₂ (Pielke et al., 2002). Despite the key implications of
forest management on the carbon-energy-water exchange there have been no integrated studies on
the effects of forest management on the Earth's climate.

Earth system models are the most advanced tools to predict future climate (Bonan, 2008). These
models represent the interactions between the atmosphere and the surface beneath, with the surface
50 formalized as a combination of open oceans, sea ice and land. For land, five classes are distinguished:
glacier, lake, wetland, urban and vegetated. Vegetation is **typically** represented by different plant
functional types. ORCHIDEE is the land surface component of the IPSL (Institut Pierre Simon
Laplace) Earth System Model. Hence, by design, the ORCHIDEE model can be run coupled to the
global circulation model LMDz. In this coupled set-up, the atmospheric conditions affect the land
55 surface and the land surface, in turn, affects the atmospheric conditions. Coupled land-atmosphere
models thus offer the possibility to quantify both the climatic effects of changes in the land surface
and the effects of climate change on the land surface. The most advanced land-surface models used,
for instance, in Earth System Models to predict climate changes (see the recent CMIP5 exercise),
account for changes in vegetation cover but consider forests to be mature and ageless, e.g., JSBACH
60 (Reick et al., 2013), CLM (Stöckli et al., 2008), MOSES (Cox et al., 1999), ORCHIDEE (Krinner
et al., 2005) and LPJ-DVGM (Bonan et al., 2003). At present, none of the predictions of future
climate thus account for the essential interactions between forest management and climate. This
gap in modelling capability provides the motivation for further development of the land-surface
model ORCHIDEE to realistically simulate both the biophysical and biogeochemical effects of forest
65 management on the climate. The ORCHIDEE-CAN (short for ORCHIDEE-CANOPY) branch of the
land surface model was specifically developed to quantify the climatic effects of forest management.

**The aim of this study is to describe the model developments and parametrization within ORCHIDEE-
CAN and to evaluate its performance. ORCHIDEE-CAN is validated against structural, biophysical
and biogeochemical data at the European scale. To allow comparison with the standard version of
70 ORCHIDEE, ORCHIDEE-CAN was run with a single-layer energy budget. A more detailed descrip-
tion and evaluation of the new multi-layer energy budget and multilevel radiative transfer scheme**

is given by Ryder et al. (2014), Chen et al. (In prep.) and McGrath et al. (In prep.b). A new forest management reconstruction which is needed to drive forest management in ORCHIDEE-CAN is presented in McGrath et al. (In prep.a) and the interactions between forest management and the new albedo scheme have been discussed by Otto et al. (2014).

2 Model overview

2.1 The Starting point: ORCHIDEE SVN r2243

The land surface model used for this study, ORCHIDEE, is based on two different modules (Krinner et al., 2005, their Fig. 2). The first module describes the fast processes such as the soil water budget and the exchanges of energy, water and CO₂ through photosynthesis between the atmosphere and the biosphere (Ducoudré et al., 1993; de Rosnay and Polcher, 1998). The second module simulates the carbon dynamics of the terrestrial biosphere and essentially represents processes as maintenance and growth respiration, carbon allocation, litter decomposition, soil carbon dynamics and phenology (Viovy and de Noblet-Ducoudré, 1997). The trunk version of ORCHIDEE describes global vegetation by 13 metaclasses (MTC) with a specific parameter set (one for bare soil, eight for forests, two for grasslands and two for croplands). Each MTC can be divided into a user-defined number of PFTs which can be characterised by at least one parameter value that differs from the parameter settings of the MTC. Parameters that are not given at the PFT-level are assigned the default value for the MTC to which the PFT belongs. By default none of the parameters are specified at the PFT-level, hence, MTCs and PFTs are the same for the standard ORCHIDEE trunk version. A concise description of the main processes in the ORCHIDEE-trunk version and a short motivation to change these modules in ORCHIDEE-CAN is given in Table 1.

Before running simulations, it is necessary to bring the soil carbon pools into equilibrium due to their slow fill rates, an approach known as model spin-up (Thornton and Rosenbloom, 2005; Xia et al., 2012). For a long time, spin-ups have been performed by brute force, i.e., running the model iteratively over a sufficiently long period which allows even the slowest carbon pool to reach equilibrium. This naïve approach is reliable but slow (in the case of ORCHIDEE it takes 3000 simulation years) and thus comes with a large computational demand, often exceeding the computational cost of the simulation itself. Alternative spin-up methods calling only parts of the model, e.g., subsequent cycles of 10 years of only photosynthesis followed by 100 year cycles of only soil processes, have been used for ORCHIDEE to reduce the computational cost in the past. These approaches, however, tend to lead to instabilities in litter and carbon pools. In recent years, semi-analytical methods have been proposed as a cost-effective solution to the spin-up issue (Martin et al., 2007; Lardy et al., 2011; Xia et al., 2012). A matrix-sequence method has been implemented in ORCHIDEE following the approach used by the PaSim model (Lardy et al., 2011). The semi-analytical spin-up implemented in ORCHIDEE relies on algebraic methods to solve a linear system of equations describing the seven

carbon pools separately for each PFT. Convergence of the method and thus equilibrium of the carbon pools is assumed to be reached when the variation of the passive carbon pool (which is the slowest) drops below a predefined threshold. The net biome production (NBP) is used as a second diagnostic criterion to confirm **equilibrium of the carbon pools**. In order to optimize computing resources, the semi-analytical spin-up will stop before the end of the run once the convergence criteria are met. ORCHIDEE's implementation of the semi-analytical spin-up has been validated at regional and global scales against a naïve spin-up, and has been found to converge 12 to 20 times faster. The largest gains were realised in the tropics and the smallest gains in boreal climate (**not shown**).

2.2 Modifications between ORCHIDEE SVN r2243 and ORCHIDEE-CAN SVN r2290

One major overarching change in the ORCHIDEE-CAN branch is the increase of internal consistency within the model by adding connections between the different processes (Fig. 1, red arrows). A more specific novelty is the introduction of circumference classes within forest PFTs, based on the work of Bellassen et al. (2010). For the temperate and boreal zone, tree height and crown diameter are calculated from allometric relationships of tree diameter that were parametrized based on the French, Spanish, Swedish and German forest inventory data and the observational data from Pretzsch (2009). The circumference classes thus allow calculation of the social position of trees within the canopy which justifies applying an intra-tree competition rule (Deleuze et al., 2004) to account for the fact that trees with a dominant position in the canopy are more likely to intercept light than suppressed trees, and, therefore, contribute more to the stand level photosynthesis and biomass growth. To respect the competition rule of Deleuze et al. (2004), a new allocation scheme was developed based on the pipe model theory (Shinozaki et al., 1964) and its implementation by Sitch et al. (2003). The scheme allocates carbon to different biomass pools (leaves, fine roots, and sapwood) while respecting the differences in longevity and hydraulic conductivity between the pools. In addition to the biomass of the different pools, LAI, crown volume, crown density, stem diameter, stem height and stand density are calculated and now depend on accumulated growth. The new scheme allows for the removal of the parameter that caps the maximum LAI (Table 1).

The calculation of tree dimensions (e.g., sapwood area and tree height) that respect the pipe theory supports making use of the hydraulic architecture of plants to calculate the plant water supply (Fig. 1, arrow 1), which is the amount of water a plant can transport from the soil to its stomata. The representation of the plant hydraulic architecture is based on the scheme of Hickler et al. (2006). The water supply is calculated as the ratio of the pressure difference between soil and leaves, and the total hydraulic resistance of the roots, leaves and sapwood, where **the sapwood resistance** is increased when cavitation occurs. Species-specific parameter values were compiled from the literature. As the scheme makes use of the soil water potential, it requires the use of the 11 layer hydrology scheme of de Rosnay (2002) (Table 1). When transpiration based on energy supply exceeds transpiration based on the water supply, the latter restricts stomatal conductance directly, which is a physiologically more

realistic representation of drought stress than the reduction of the carboxylation capacity (Flexas et al., 2006) done in the standard version of ORCHIDEE (further also referred to as "trunk" version).

145 In line with this approach, the drought stress factor used to trigger phenology and senescence is now calculated as the ratio between the transpiration based on water supply and transpiration based on atmospheric demand (Fig. 1, arrow 2).

The new allocation scheme also drastically changed the way forests are represented in the ORCHIDEE-CAN branch. Although the exact location of the canopies in the stand is not known, individual tree
150 canopies are now spherical elements with their horizontal location following a Poisson distribution across the stand. Each PFT contains a user-defined number of model trees, each one corresponding to a circumference class. Model trees are replicated to give realistic stand densities. Following tree growth, canopy dimensions and stand density are updated (Fig. 1, arrow 3). This formulation results in a dynamic canopy structure that is exploited in other parts of the model, i.e., precipitation
155 interception, transpiration, energy budget calculations, radiation scheme (Fig. 1, arrow 4) and absorbed light for photosynthesis (Fig. 1, arrow 5). In the trunk version these processes are driven by the big-leaf canopy assumption. The introduction of an explicit canopy structure is thought to be a key development with respect to the objectives of the ORCHIDEE-CAN branch, i.e., quantifying the biogeochemical and biophysical effects of forest management on atmospheric climate.

160 The radiation transfer scheme at the land surface benefits from the introduction of canopy structure. The trunk version of ORCHIDEE prescribes the vegetation albedo solely as a function of LAI. In the ORCHIDEE-CAN branch each tree canopy is assumed to be composed of uniformly distributed single scatterers. Following the assumption of a Poisson distribution of the trees on the land surface, the model of Haverd et al. (2012) calculates the transmission probability of light to any
165 given vertical point in the forest. This transmission probability is then used to calculate an effective LAI, which is a statistical description of the vertical distribution of leaf mass that accounts for stand density and horizontal tree distribution. The complexity and computational costs are largely reduced by using the effective LAI in combination with the 1D two stream radiation transfer model of Pinty et al. (2006) rather than resolving a full 3-D canopy model. By using the effective LAI, the 1-D
170 model reproduces the radiative fluxes of the 3-D model. The approach of the two stream radiation transfer model was extended for a multi-layer canopy (McGrath et al., In prep.b) to be consistent with the multi-layer energy budget and to better account for non-linearities in the photosynthesis model. The scattering parameters and the background albedo (i.e. the albedo of the surface below the dominant tree canopy) for the two stream radiation transfer model were extracted from the Joint Research Centre Two-stream Inversion Package (JRC-TIP) remote sensing product (section 4.7). This
175 approach produces fluxes of the light absorbed, transmitted, and reflected by the canopy at vertically discretized levels, which are then used for the energy budget (Fig. 1, arrow 6) and photosynthesis calculations (Fig. 1, arrow 5).

The canopy radiative transfer scheme of (Pinty et al., 2006) separates the calculation of the fluxes
180 resulting from downwelling direct and diffuse light, with different scattering parameters available
for near-infrared (NIR) and visible (VIS) light sources. The snow albedo scheme in the trunk does
not distinguish between these two shortwave bands. Therefore, the snow scheme of the Biosphere-
Atmosphere Transfer Scheme (BATS) for the Community Climate Model (Dickinson et al., 1986)
was incorporated into the ORCHIDEE-CAN branch, since it distinguishes between the NIR and VIS
185 radiation. The radiation scheme of Pinty et al. (2006) requires snow to be put on the soil below the
tree canopy instead of on the canopy itself. The calculation of the snow coverage of a PFT therefore
had to be revised according to the scheme of Yang et al. (1997), which allows for snow to completely
cover the ground at depths greater than 0.2 m. The parameter values of Yang et al. (1997) were used
in the ORCHIDEE-CAN branch.

190 The ORCHIDEE-CAN branch differs from any other land surface model by the inclusion of a
newly developed multi-layer energy budget. There are now subcanopy wind, temperature, humidity,
longwave radiation and aerodynamic resistance profiles, in addition to a check of energy closure at
all levels. The energy budget represents an implementation of some of the characteristics of detailed
single site, iterative canopy models (e.g., Baldocchi, 1988; Ogee et al., 2003) within a system that is
195 coupled implicitly to the atmosphere. As an enhancement to the trunk version of ORCHIDEE (Table
1), the new approach also generates a leaf temperature, using a vegetation profile and a vertical
shortwave and longwave radiation distribution scheme (Ryder et al., 2014), which will be fully
available when parametrisation of the scheme has been completed across test sites corresponding
to the species within the model (Chen et al., In prep.). As with the trunk version, the new energy
200 budget is calculated implicitly (Polcher et al., 1998; Best et al., 2004). An implicit solution is a linear
solution in which the surface temperature and fluxes are calculated in terms of the atmospheric input
at the same time-step, whereas an explicit solution uses atmospheric input from the previous time
step to calculate the surface temperature and fluxes. Although it is less straightforward to derive, the
implicit solution is more computationally efficient and stable, which allows the model to be run over
205 a time-step of 15 minutes when coupled to the atmospheric model LMDz - much longer than would
be the case for an explicit model. Parameters were derived by optimizing the model against the
observations from short-term field campaigns. The new scheme may also be reduced to the existing
single layer case, so as to provide a means of comparison and compatability with the ORCHIDEE
trunk version.

210 The combined use of the new energy budget and the hydraulic architecture of plants required
changes to the calculation of the stomatal conductance and photosynthesis (Fig. 1, arrow 7). When
water supply limits transpiration, stomatal conductance is reduced and photosynthesis needs to be
recalculated. Given that photosynthesis is among the computational bottlenecks of the model, the
semi-analytical procedure as available in previous trunk versions (r2031 and further) is replaced by
215 an adjusted implementation of the analytical photosynthesis scheme of Yin and Struik (2009), which

is also implemented in the latest ORCHIDEE-trunk version. In addition to an analytical solution for photosynthesis the scheme includes a modified Arrhenius function for the temperature dependence that accounts for a decrease of carboxylation capacity ($k_{V_{cmax}}$) and electron transport capacity ($k_{J_{max}}$, see Table 2 for variable explanations) at high temperatures and a temperature dependent $k_{J_{max}}/V_{cmax}$ ratio (Kattge and Knorr, 2007). The temperature response of $k_{V_{cmax}}$ and $k_{J_{max}}$ was parametrized with values from reanalysed data in literature (Kattge and Knorr, 2007), whereas $k_{V_{cmax}}$ and $k_{J_{max}}$ at a reference temperature of 25° C were derived from observed species-specific values in the TRY database (Kattge et al., 2011). As the amount of absorbed light varies with height (or canopy depth), the absorbed light computed from the albedo routines is now directly used in the photosynthesis scheme resulting in full consistency between the top of the canopy albedo and absorption. This new approach replaces the old scheme which used multiple levels based on the leaf area index, not the physical height.

ORCHIDEE-CAN incorporates a systematic mass balance closure for carbon cycling to assure that carbon is not getting created or destroyed during the simulation. Hence, budget closure is now consistently checked for water, carbon and energy throughout the model.

The trunk uses 13 plant functional types (PFT) to represent vegetation globally: one PFT for bare soil, eight for forests, two for grasslands, and two for croplands. The ORCHIDEE-CAN branch makes use of the externalization of the PFT-depended parameters by adding 12 parameter sets that represent the main European tree species. Species parameters were extracted from a wide range of sources including original observations, large databases, primary research and remote sensing products (section 4). The use of age classes is introduced through externalization of the PFT parameters as well. Age classes are used during land cover change and forest management to simulate the re-growth of a forest. Following a land cover change, biomass and soil carbon pools (but not soil water columns) are either merged or split to represent the various outcomes of a land cover change. The number of age classes is user defined. Contrary to typical age classes, the boundaries are determined by the tree diameter rather than the age of the trees.

Finally, the forest management strategies in the ORCHIDEE-CAN branch were refined from the original forest management branch (Bellassen et al., 2010). Self-thinning was activated for all forests regardless of human management, contrary to the original FM branch. The new default management strategy thus has no human intervention but includes self-thinning, which replaces the fixed 40 year turnover time for woody biomass. Three management strategies with human intervention have been implemented: (1) “high stands”, in which human intervention is restricted to thinning operations based on stand density and diameter, with occasional clearcuts. Aboveground stems are harvested during operations, while branches and belowground biomass are left to litter. (2) “coppices” involve two kinds of cuts. The first coppice cut is based on stem diameter and the aboveground woody biomass is harvested whereas the belowground biomass is left living. From this belowground biomass new shoots sprout, which increases the number of aboveground stems. In subsequent cuts

the amount of shoots is not increased, although all aboveground wood biomass is still harvested. (3) "short rotation coppices", where rotation periods are based on age and are generally very short (3-6 years). The different management strategies can occur with or without litter raking, which reduces the litter pools and has a longterm effect on soil carbon (Gimmi et al., 2012). All management types are parametrized based on forest inventory data, yield tables and guidelines for forest management. The inclusion of forest management resulted in two additional carbon pools, branches and coarse roots (i.e., aboveground and belowground woody biomass) and therefore required an extension to the semi-analytical spin-up method (section 2.1). The semi-analytical spin-up is now run for nine C pools.

3 Description of the developments

3.1 Allocation

Following bud burst, photosynthesis produces carbon that is added to the labile carbon pool. Labile carbon is used to sustain the maintenance respiration flux (F_{rm}), which is the carbon cost to keep existing tissue alive (Amthor, 1984). Maintenance respiration for the whole plant is calculated by summing maintenance respiration of the different plant compartments, which is a function of the nitrogen concentration of the tissue following the Beer-Lambert law and subtracted from the whole-plant labile pool (up to a maximum of 80% of the labile pool).

The remaining labile carbon pool is split into an active and none-active pool. The size of the active pool is calculated as a function of plant phenology and temperature and was formalized following Ryan (1991); Sitch et al. (2003); Zaehle and Friend (2010). The remaining non-active pool is used to restore the labile and carbohydrate reserves pools according to the rules proposed in Zaehle and Friend (2010). The labile pool is limited to 1% of the plant biomass or 10 times the actual daily photosynthesis. Any excess carbon is transferred to the non-respiring carbohydrate reserve pool. The carbohydrate reserve pool is capped to reflect limited starch accumulation in plants, but carbon can move freely between the two reserve pools. After accounting for growth respiration (F_{rg}), i.e., the cost for producing new tissue excluding the carbon required to build the tissue itself (Amthor, 1984), the total allocatable C used for plant growth is obtained (M_{totinc}).

New biomass is allocated to leaves, roots, sapwood, heartwood, and fruits. Allocation to leaves, roots and wood respects the pipe model theory (Shinozaki et al., 1964) and thus assumes that producing one unit of leaf mass requires a proportional amount of sapwood to transport water from the roots to the leaves as well as a proportional fraction of roots to take up the water from the soil. The different biomass pools have different turnover times, and therefore at the end of the daily time step, the actual biomass components may no longer respect the allometric relationships. Consequently, at the start of the time step carbon is first allocated to restore the allometric relationships before the

remaining carbon is allocated in the manner described below. The scaling parameter between leaf and sapwood mass is derived from:

$$d_l = k_{ls} \times m_w \times d_s \quad (1)$$

290 Where d_l is the one-sided leaf area of an individual plant, d_s is the sapwood cross-section area of an individual plant, k_{ls} a parameter linking leaf area to sapwood cross-section area and, m_w is the water stress as defined in section 3.2. Alternatively, leaf area can be written as a function of leaf mass (M_l) and the specific leaf area (k_{sla}):

$$d_l = M_l \times k_{sla} \quad (2)$$

295 Sapwood mass M_s can be calculated from the sapwood cross-section area d_s as follows:

$$M_s = d_s \times d_h \times k_{\rho s} \quad (3)$$

Where d_h is the tree height and $k_{\rho s}$ is the sapwood density. Following substitution of equations 2 and 3 into equation 1, leaf mass can be written as a function of sapwood mass:

$$M_l = (M_s \times f_{KF}) / d_h \quad (4)$$

300 where,

$$f_{KF} = (k_{ls} \times m_w) / (k_{sla} \times k_{\rho s}) \quad (5)$$

where, k_{ls} is calculated as a function of the gap fraction as supported by site-level observations (Simonin et al., 2006):

$$k_{ls} = k_{lsmin} + f_{Pgap,trees} \times (k_{lsmax} - k_{lsmin}) \quad (6)$$

305 k_{lsmin} is the minimum observed leaf area to sapwood area ratio, k_{lsmax} is the maximum observed leaf area to sapwood area ratio and $f_{Pgap,trees}$ is the actual gap fraction. By using the gap fraction as a control of k_{ls} more carbon will be allocated to the leaves until canopy closure is reached.

Following Magnani et al. (2000), sapwood mass and root mass (M_r) are related as follows:

$$M_s = k_{sar} \times d_h \times M_r \quad (7)$$

310 where the parameter k_{sar} is calculated according to Magnani et al. (2000) (their equation (17)):

$$k_{sar} = \sqrt{(k_{rcon}/k_{scon}) \times (k_{\tau s}/k_{\tau r}) \times k_{\rho s}} \quad (8)$$

where k_{rcon} is the hydraulic conductivity of roots, k_{scon} is the hydraulic conductivity of sapwood, $k_{\tau s}$ is the longevity of sapwood and $k_{\tau r}$ is the root longevity. Following substitution of equation (4) into (7) and some rearrangement, leaf mass can be written as a function of root mass:

$$315 \quad M_l = f_{LF} \times M_r \quad (9)$$

where,

$$f_{LF} = k_{sar} \times f_{KF} \quad (10)$$

Parameter values used in equations 1 to 9, i.e., k_{lsmax} , k_{lsmin} , k_{sar} , k_{sla} , $k_{\rho s}$, k_{rcon} , k_{scon} , $k_{\tau s}$ and $k_{\tau r}$, are based on literature review (Table S1,S2 and S3). The allometric relationships between
 320 the plant components and the hydraulic architecture of the plant (section 3.2) are both based on the pipe model theory, hence, both the allocation and the hydraulic architecture module use the same parameter values for root and sapwood conductivity.

In this version of ORCHIDEE, forests are modelled to have k_{ncirc} circumference classes with d_{ind} identical trees in each one. Hence, the allocatable biomass (M_{totinc}) needs to be distributed
 325 across l diameter classes:

$$M_{totinc} = \text{sum}(l)[d_{ind(l)} \times M_{inc(l)}] \quad (11)$$

where $M_{inc(l)}$ is the biomass that can be allocated to diameter class l . Mass conservation thus requires:

$$M_{inc(l)} = M_{linc(l)} + M_{rinc(l)} + M_{sinc(l)} \quad (12)$$

330 where $M_{linc(l)}$, $M_{rinc(l)}$ and, $M_{sinc(l)}$ are the increase in leaf, root and wood biomass for a tree in diameter class l , respectively. Equation 4 and 9 can be rewritten as

$$(M_{l(l)} + M_{linc(l)})/(M_{s(l)} + M_{sinc(l)}) = f_{KF}/(d_{h(l)} + d_{hinc(l)}) \quad (13)$$

$$(M_{l(l)} + M_{linc(l)}) = (M_{r(l)} + M_{rinc(l)}) \times f_{LF} \quad (14)$$

An allometric relationship is used to describe the relationship between tree height and basal area
 335 (Pretzsch, 2009):

$$d_{h(l)} = k_{\alpha 1} \times (4/\pi \times d_{ba(l)})^{(k_{\beta 1}/2)} \quad (15)$$

The change in height is then calculated as

$$d_{hinc(l)} = [k_{\alpha 1} \times (4/\pi \times (d_{ba(l)} + d_{bainc(l)}))^{(k_{\beta 1}/2)}] - d_{h(l)} \quad (16)$$

where $d_{ba(l)}$ and $d_{bainc(l)}$ are the basal area and its increment, respectively. $k_{\alpha 1}$ and $k_{\beta 1}$ are
 340 allometric constants relating tree diameter and height. The distribution of C across the l diameter
 classes depends on the basal area of the model tree within each diameter class. Trees with a large
 basal area are assigned more carbon for wood allocation than trees with a small basal area, according
 to the method of Deleuze et al. (2004).

$$d_{bainc(l)} = f_{\gamma} \times \left(d_{circ(l)} - k_m * g_{\sigma} + \sqrt{(k_m \times g_{\sigma} + d_{circ(l)})^2 - (4 \times g_{\sigma} \times d_{circ(l)})} \right) / 2 \quad (17)$$

345 where k_m is a parameter, f_{γ} and g_{σ} are calculated from parameters and $d_{circ(l)}$ is the circumfer-
 ence of the model tree in diameter class l . g_{σ} is a function of the diameter distribution of the stand
 at a given time step.

Equations 10 to 16 need to be simultaneously solved. An iterative scheme was avoided by lin-
 earising equation 15, which was found to be an acceptable numerical approximation as allocation
 350 is calculated at a daily time step, and hence the changes in height are small and the relationship is
 locally linear:

$$d_{hinc(l)} = d_{bainc(l)} / f_s \quad (18)$$

where f_s is the slope of the locally linearised equation 15 and is calculated as:

$$f_s = k_{step} / (k_{\alpha 1} \times (4/\pi * (d_{ba} + k_{step}))^{(k_{\beta 1}/2)} - k_{\alpha 1} \times (4/\pi \times d_{ba})^{(k_{\beta 1}/2)}) \quad (19)$$

355 Equations 10, 11, 12, 13, 14, 16, 17 and 18 are then solved for f_{γ} . f_{γ} distributes photosynthates
 across the different diameter classes and as such controls the intra-species competition within a
 stand. f_{γ} thus depends on the total allocatable carbon and needs to be optimised at every time step.
 Once f_{γ} has been calculated, $M_{linc(l)}$, $M_{rinc(l)}$ and $M_{sinc(l)}$ can be calculated.

3.2 Hydraulic architecture

360 The representation of the impact of soil moisture stress on water, carbon and energy fluxes has
 been identified as one of the major uncertainties in land surface models (De Kauwe et al., 2013).

Neither the empirical functions nor the soil moisture stress functions, which are commonly used in land surface models, fully capture stomatal closure and limitation of C uptake during drought stress (Bonan et al., 2014; Verhoef and Egea, 2014). Therefore, we replaced the soil moisture stress function which limits C assimilation through a constrain on $k_{V_{cmax}}$ in the ORCHIDEE-trunk, by a constrain based on the amount of water plants can transport from the soil to their leaves.

The model calculates plant water supply according to the implementation of hydraulic architecture by Hickler et al. (2006). Plant water supply is the amount of water the plant can transport from the soil to its stomata, accounting for the resistances to water transport in the roots, sapwood and leaves. If transpiration rate exceeds plant water supply, the stomatal conductance is reduced until equilibrium is reached.

The water flow from the soil to the leaves is driven by a gradient of decreasing water potential. Using Darcy's law (Slatyer, 1967; Whitehead, 1998), the supply of water for transpiration through stomata can be described as:

$$F_{Trs} = p_{delta} / (R_r + R_{sap} + R_l) \quad (20)$$

where p_{delta} is the pressure difference between the soil and the leaves; and R_r , R_{sap} and R_l are the hydraulic resistances of fine roots, sapwood and leaves, respectively. p_{delta} is calculated following Whitehead (1998):

$$p_{delta} = p_{\psi sr} - k_{\psi l} - (d_h \times k_{\rho w} \times k_g) \quad (21)$$

where $k_{\psi l}$ is a PFT-specific minimal leaf water potential, which means that plants are assumed to maximise water uptake by lowering their $k_{\psi l}$ to the minimum, if transpiration exceeds F_{Trs} (Tyree and Sperry, 1989). The product of d_h , $k_{\rho w}$ and k_g accounts for the loss in water potential by lifting a mass of water from the soil to the place of transpiration at height d_h , $k_{\rho w}$ is the density of water, and k_g is the gravitational constant. The soil water potential in the rooting zone ($p_{\psi sr}$) was calculated by adding a modulator (m_ψ) to the bulk soil water potential, which was calculated as the sum of the soil water potential in each soil layer weighted by the relative share of roots (d_{rd}) in the individual soil layer:

$$p_{\psi sr} = sum(l)[p_{\psi s} \times d_{rd}] + m_\psi \quad (22)$$

The soil water potential for each layer $p_{\psi sl}$ is calculated from soil water content according to Van Genuchten (1980).

$$p_{\psi s(l)} = \frac{1}{k_{av}} \left(\left(\frac{M_{swc} - k_{swcr}}{k_{swcs} - k_{swcr}} \right)^{-1/k_{mv}} - 1 \right)^{1/k_{nv}} \quad (23)$$

where M_{swc} is the volumetric soil water content, k_{swcr} and k_{swcs} are respectively the residual and saturated soil water content and k_{av} , k_{mv} and k_{nv} are parameters.

Root resistance is related to the root mass and thus can be expressed as (Weatherley, 1982):

$$395 \quad R_r = \frac{1}{(k_{rcon} \times M_r)} \quad (24)$$

where k_{rcon} is the fine root hydraulic conductivity per unit biomass. Sapwood resistance is calculated according to Magnani et al. (2000):

$$R_{sap} = \frac{d_h}{(d_s \times k_{scon})} \quad (25)$$

where k_{scon} is the sapwood specific conductivity, which is decreased when cavitation occurs. The
400 loss of conductance as a result of cavitation is a function of $p_{\psi_{sr}}$ and was implemented by using an s-shaped vulnerability curve:

$$k_{scon} = k_{scon} \times e^{(-p_{\psi_{rs}}/k_{\psi 50})^{k_c}} \quad (26)$$

where $k_{\psi 50}$ is the $p_{\psi_{sr}}$ that causes 50% loss of conductance; and k_c is a shape parameter.

R_l is related to the specific leaf conductivity per unit leaf area (k_l) and the leaf area index :

$$405 \quad R_l = \frac{1}{(k_{lcon} \times d_{LAI})} \quad (27)$$

The response of water viscosity to low temperatures increases the resistance (Cochard et al., 2000). The relationship is described as:

$$R_{temp} = \frac{R}{(k_{\alpha 1v} + k_{\alpha 2v} \times T)} \quad (28)$$

where $k_{\alpha 1v}$ and $k_{\alpha 2v}$ are empirical parameters (Cochard et al., 2000), R_{temp} is the temperature
410 adjusted R_l , R_{sap} or R_r , T is air temperature for R_l and R_{sap} and T is soil temperature for R_r .

If, for any time step, the transpiration calculated by the energy budget exceeds the amount of water the plant can transport from the soil to its stomata, transpiration is limited to the plant water supply. As the transpiration is now reduced, the initial calculations of the energy budget and photosynthesis, solely based on atmospheric information, are no longer valid. As a result the energy
415 budget and photosynthesis must be recalculated for the time step in question. For this recalculation, stomatal conductance at the canopy level is calculated such that transpiration equals the amount of water the plant can transport. Owing to the feedback between stomatal conductance, leaf surface temperature and transpiration, this calculation may require up to 10 iterations to converge, using a

stationary iterative method. When the multi-layer energy budget is reduced to its single-layer im-

420 **plementation, however,** canopy level stomatal conductance is decomposed to obtain the stomatal conductance at each canopy layer assuming that each layer is equally restricted by drought stress. Finally, the restricted stomatal conductance is used to calculate CO₂ assimilation rate according to the photosynthesis model by Farquhar, von Caemmerer and Berry (section 3.6).

3.3 Canopy structure

425 Stand structure controls the amount of light that penetrates to a given depth in the canopy. For example, the amount of light reaching the forest floor will be higher for a stand with few mature trees compared to many young trees even if both stands have the same leaf area index. Where a big leaf approach assumes a homogeneous block shaped canopy (as in the trunk version of ORCHIDEE) and can therefore rely on the law of Beer-Lambert, a geometric approach is required to calculate light
430 penetration through structured canopies. Light penetration needs to be simulated to calculate albedo (section 3.4), photosynthesis (section 3.6), partitioning of energy fluxes (section 3.5) and the amount of light reaching the forest floor (see for example section 3.1). The gap fraction, which is the basic information in calculating light penetration at different depths in the canopy, is calculated following the approach presented by Haverd et al. (2012) and formalized in their semi-analytical model. Rather
435 than a spatially explicit approach, Haverd et al. (2012) follow a statistical approach which reduces the memory requirements for the simulations and limits the space requirements for storing the model output files.

The model of Haverd et al. (2012) represents the canopy by a statistical height distribution with varying crown sizes and stem diameters for each height class. The crown canopies are treated as
440 spheroids containing homogeneously distributed single scatterers. Although this f_{Pgap} model can explicitly include trunks, we made the decision to exclude them, as the spectral parameters for our radiation model (section 3.4) are extracted from remote sensing data (section 4.8) without distinguishing between leafy and woody masses. This gives the gap probability for trees as a function of height (z) and solar zenith angle (θ_z):

$$445 \quad f_{Pgap}^{trees}(\theta_z, z) = e^{-d_\lambda \times \overline{d_c(\theta_z, z) \times (1 - f_{Pwc}(\theta_z, z))}} \quad (29)$$

where d_λ is the inverse of the tree density, d_c is the projected crown area (for an opaque canopy), and f_{Pwc} is the mean crown porosity. The overbar depicts the mean over the tree distribution as a function of tree height or, in our case, the mean over the l circumference classes. Following minor adaptations, the implementation of Haverd and Lovell (Haverd et al., 2012) was incorporated in

450 ORCHIDEE-CAN. As there also exist crops, grasses, and bare soil in the model, f_{Pgap} was adjusted for these situations as well. For grasses and crops, the same formulation is used:

$$f_{Pgap}^{gc}(\theta_z, z) = e^{-0.5 \times d_{LAIabove} \times m_{LAICorr} / \cos(\theta_z)} \quad (30)$$

where $d_{LAIabove}$ is the total amount of LAI above height z , and $m_{LAICorr}$ is a correction factor to account for the fact that grasses and crops are treated as homogeneous blocks of vegetation with
 455 no internal structure and is often referred to as a clumping factor. Here it is treated as a tunable parameter and therefore the term correction factor was used. For bare soil, there is no vegetation to intercept radiation, and therefore $f_{Pgap}^{bs}(\theta_z, z)$ is always unity.

3.4 Multi-layer two-way radiation scheme for tall canopies

Species-specific radiation absorbance, reflectance and transmittance by the forest canopy were calcu-
 460 lated from a radiation transfer model (Pinty et al., 2006) which was parametrized by satellite-derived species-specific scattering values (section 4.8). Given the complexity of radiation transfer, it remains challenging to accurately simulate radiation transfer through structurally and optically complex vegetation canopies without using explicit 3-D models. The applied 1-D model belongs to the family of two-stream models (Meador and Weaver, 1980) and thus calculates transmittance, absorbance and
 465 reflectance of both the incoming and outgoing radiation. The calculation of the reflectance at the top of the canopy due to a collimated source (i.e., the sun) is divided in three components:

1. scattering of radiation between the vegetated elements with a black background

$$f_{Coll,veg}^{fR} = f(\theta_{mu}, f_{rl}, f_{tl}, g_G, d_{LAIeff}) \quad (31)$$

2. scattering of radiation by the background with a black canopy

$$470 \quad f_{UnColl,bgd}^{fR} = f_{Rbgd} \times e^{(-d_{LAIeff} / (2 \times \theta_{mu}))} \times f_{UnColl,veg}^T \quad (32)$$

3. multiple scattering of radiation between the canopy and the background

$$f_{Coll,bgd}^{fR} = f_{Rbgd} \times [f_{Coll,bgd,1}^{fR} + f_{Coll,bgd,n}^{fR}] \quad (33)$$

Term (1) is widely used in cloud reflectance calculations, and depends on the cosine of the solar zenith angle (θ_{mu}), the reflectance and transmittance of the single leaves (f_{rl} and f_{tl} , respectively),
 475 the leaf orientation function (g_G), and the effective LAI (d_{LAIeff}). The exact definition of this term is given in Equation B2 in Pinty et al. (2006). In term (2), f_{Rbgd} is the reflectance of the ground beneath the canopy and $f_{UnColl,veg}^T$ is the transmitted fraction of light to the ground which has not collided with any canopy elements. In term (3), $f_{Coll,bgd,1}^{fR}$ is the fraction of light which has struck

vegetation and collided with the background a single time, while $f_{Coll,bgd,n}^{fR}$ is the fraction which
 480 has collided multiple times (n) with the background.

The sum of the three components results in the canopy albedo (Pinty et al., 2006). Similar equations can be derived for light originating from diffuse sources (e.g., clouds and other atmospheric scattering)(Pinty et al., 2006). Implementations of the calculations of the canopy fluxes for a single level are available from the JRC, and these implementations were used as the basis of the routines
 485 put into ORCHIDEE-CAN for both the single- and multi-level cases (McGrath et al., In prep.b). This implementation relies on the use of the effective LAI, which is the LAI that needs to be used in a 1-D process representation to obtain the same reflectance, absorbance and transmittance as would be obtained by a 3D-canopy representation (Pinty, 2004). In this study, the effective LAI was calculated by first computing the canopy gap probability, i.e. the probability that light is transmitted to
 490 a specified height in the canopy at a given solar angle. The gap probability is then converted into the effective LAI by passing it as an input to the inverted Beer-Lambert's law (with an extinction coefficient of 0.5 to ensure compatibility with the two-stream inversion of Pinty et al. (2011a)).

$$d_{LAIeff} = -2.0 \times \cos(\theta_z) \times \log(f_{Pgap}) \quad (34)$$

where f_{Pgap} can be f_{Pgap}^{trees} , f_{Pgap}^{gc} , f_{Pgap}^{bs} . Following the introduction of multi-layer photosynthesis and energy budget submodels, the approach proposed by Pinty (2004) had to be adjusted such
 495 that it could be applied for every level for which absorbance needs to be known to calculate photosynthesis (section 3.6) and reflectance needs to be known to calculate the net shortwave radiation (section 3.5). The multi-layer approach basically applies the 1D-two stream canopy radiation transfer model by Pinty et al. (2006) to each canopy level where the light transmitted by the overlaying
 500 level becomes the input for the lower level.

As the multi-level approach is built around the solution of the one-level scheme for each canopy level, no new equations are introduced. The method can be summarized by the following algorithm for which the details are given in (McGrath et al., In prep.b). First, three fluxes are calculated for each level independently: the fraction of light transmitted through the layer without striking vegetation,
 505 the fraction of light reflected after striking vegetation, and the fraction of light transmitted through the layer after striking vegetation. These three fluxes represent the only possible fate of light (any light not taking one of these paths must be absorbed for energy conservation). Next, an iterative approach is invoked which follows the path of a single photon entering the top level. Based on the solutions for each single level, probabilities can be calculated that the photon is transmitted to a
 510 lower level or reflected to a higher level. Any fraction which is reflected upwards from the top level is added to the total canopy albedo and not considered further. The fraction which is transmitted through the top level enters the next highest level, and again the single level solutions determine where this light goes. Any fraction reflected upwards is considered in the next iteration as part of the light entering the upper level. The steps continue until the bottom canopy level is reached. Here,

515 any fraction which is transmitted into the soil is removed from consideration and added to the total transmittance through the canopy. The algorithm then proceeds to the above canopy level. Now the “transmitted” fluxes are moving in the upwards direction towards to the sky, while “reflected fluxes” are moving towards the ground. The code continues towards the top level, taking as input from below both the flux reflected by downwelling light from the level below the current level and the
 520 flux transmitted from the lower level by upwelling light. After each iteration (moving from the top of the canopy to the bottom and back to the top), the total amount of light considered “active” has been reduced by light escaping to the sky or being absorbed by the canopy or ground. Eventually, this “active” light falls below a pre-defined threshold and the calculation is considered as converged.

Due to the iterative procedure, energy is not strictly conserved, although we have attempted to
 525 choose a threshold which minimizes this loss. The multilevel albedo calculation is currently the most expensive part of the model, due to the iterations and the fact that it must be performed over all canopy levels (currently set to 10), grid points, and PFTs at every physical time-step. Levels with no LAI are no less expensive to compute, either, although we have arranged our canopy levels to make sure no levels are empty in most cases.

530 3.5 Multi-layer energy budget

The present generation of land surface models have difficulties in reproducing consistently the energy balances that are observed in field studies (Pitman et al., 2009; Jiménez et al., 2011; de Noblet-Ducoudré et al., 2012). The ORCHIDEE-CAN branch implemented an energy budget scheme that represents more than one canopy layer to simulate the effects of scalar gradients within the canopy
 535 for determining more accurately the net sensible and latent heat fluxes that are passed to the atmosphere. As outlined in Polcher et al. (1998), the use of an implicit solution for coupling between the atmospheric model and the surface layer model is the only way to keep profiles of temperature and humidity synchronised across the two models when the coupled-model is run over large time steps (e.g., of 30 minutes). The difference between explicit and implicit schemes is that an explicit
 540 scheme will calculate each value of the variable (e.g., temperature and humidity) at the current time step entirely in terms of values from the previous time step. An implicit scheme requires the solution of equations written only in terms of those at the current time step.

The modelling approach formalises three constraints that ensure energy conservation. The three equations that describe the main interactions are:

- 545 1. The energy balance at each layer is the sum of incoming and outgoing fluxes of latent and sensible heat and of shortwave and longwave radiation:

$$k_{lhc,i} k_{\rho_v,i} \frac{\delta T_{L,i}}{\delta t} = \left(k_{shc} k_{\rho_a} \frac{(T_{L,i} - T_{a,i})}{R_{a,i}} + k_{\lambda,LE} \rho_a \frac{q_{L,i} - q_{a,i}}{R_{s,i}} + F_{SW,i} + F_{LW,i} \right) \left(\frac{1}{\Delta d_{hl,i}} \right) \quad (35)$$

where $F_{LW,i}$ is the sum total of long wave radiation, that is, the net LW radiation absorbed into layer i and $F_{SW,i}$ is the net absorbed short wave radiation as calculated by the radiation scheme in section 3.4. k_{shc} is the specific heat capacity of air. The source sensible heat flux from the leaf at level i is the difference between the leaf temperature ($T_{L,i}$) and the atmospheric temperature at the same level ($T_{a,i}$), divided by $R_{a,i}$, which is the leaf resistance to sensible heat flux (a combination of stomatal and boundary layer resistance). Similarly, the source latent heat flux from the leaf at level i is the difference between the saturated humidity in the leaf ($q_{L,i}$) and that in the atmosphere at level i ($q_{a,i}$), divided by $R_{s,i}$ which is the leaf resistance to latent heat flux. $R_{a,i}$ is calculated based upon the leaf boundary layer resistance, and is described in the present model according to Baldocchi (1988). $R_{s,i}$ is the stomatal resistance of the leaf that is calculated using the model of Ball et al. (1987).

2. The sensible heat flux between the vegetation ('the leaf') and the surrounding atmosphere at each level, and between adjacent atmospheric levels above and below is provided by the following expression:

$$\frac{\delta T_{a,i}}{\delta t} \Delta d_{V,i} = k_{k,i} \frac{\delta^2 T_{a,i}}{\delta z^2} \Delta d_{A,i} + \left(\frac{T_{L,i} - T_{a,i}}{R_{a,i}} \right) \left(\frac{1}{\Delta d_{hl,i}} \right) \Delta d_{V,i} \quad (36)$$

where z denotes the height above the soil surface. We have re-written the scalar conservation equation, as applied to canopies, in terms of the sensible heat flux, temperature and source sensible heat from the vegetation at each layer.

3. The latent heat flux between the vegetation and surrounding atmosphere at each level, and between adjacent atmospheric levels above and below is described in a form that is analogous to equation (36), above:

$$\frac{\delta q_{a,i}}{\delta t} \Delta d_{V,i} = k_{k,i} \frac{\delta^2 q_{a,i}}{\delta z^2} \Delta d_{A,i} + \left(\frac{q_{L,i} - q_{a,i}}{R_{s,i}} \right) \left(\frac{1}{\Delta d_{hl,i}} \right) \Delta d_{V,i} \quad (37)$$

In addition to these three basic equations various terms had to be parameterised. The 1D second-order closure model of Massman and Weil (1999) was used to simulate the vertical transport coefficients $k_{k,i}$ within the canopy while accounting for the vertical and horizontal distribution of LAI (section 3.3). This set of equations were then written in an implicit form and solved by induction. More details on the implicit multi-layer energy budget and a complete mathematical documentation are given in Ryder et al. (2014).

To complete the energy budget calculations, the multi-layer 1D canopy radiation transfer model (section 3.4) was used to calculate the net shortwave radiation at each canopy layer. Further, the canopy radiation scheme makes use of the Longwave Radiation Transfer Matrix (LRTM) (Gu, 1988;

Gu et al., 1999). This approach separates the calculation of the radiation distribution completely from the implicit expression. Instead, a single source term for the long wave radiation is added at each level. This means that the distribution of LW radiation is now explicit (i.e., makes use of information only from the 'previous' and not the 'current' time step) but the changes within the timestep were small enough to not affect the overall stability of the model. However, an advantage of the approach is that it accounts for a higher order of reflections from adjacent levels than the single order assumed in the process above.

3.6 Analytical solution for photosynthesis

The photosynthesis model by Farquhar, von Caemmerer and Berry (Farquhar et al., 1980) predicts net photosynthesis of C3 plants as the minimum of the Rubisco-limited rate of CO₂ assimilation and the electron transport-limited rate of CO₂ assimilation (Farquhar et al., 1980). The ORCHIDEE-CAN branch calculates net photosynthesis following an analytical algorithm as described by Yin and Struik (2009). In addition, the C4 photosynthesis is calculated by an equivalent version of the Farquhar, von Caemmerer and Berry model that was extended to account for noncyclic electron transport (Yin and Struik, 2009). A detailed derivation of the analytical solution of the Farquhar, von Caemmerer and Berry model is given in Yin and Struik (2009). Although the exclusion of mesophyll conductance from the photosynthesis model could lead to an underestimation of the CO₂ fertilization effect in Earth system models (Sun et al., 2014), mesophyll conductance was not included in ORCHIDEE-CAN to maintain compatibility between the model formulation and its parametrization. Because values of $k_{V_{cmax}}$ and $k_{J_{max}}$ differ between different formulations of the photosynthesis model (Kattge and Knorr, 2007; Medlyn et al., 2002) and the parametrization that was used in ORCHIDEE-CAN did not include mesophyll conductance, it was also not accounted for in the model formulation. The analytical photosynthesis model implemented in ORCHIDEE-CAN could be easily extended to include mesophyll conductance but that would require reparameterizing the photosynthesis model.

Owing to the canopy structure simulated in this model version and the layering of the canopy, the amount of absorbed light now varies with canopy depth. This new approach replaces the old scheme which uses multiple levels based on the leaf area index, not the physical height within the canopy. Photosynthesis is now calculated at each vertically resolved canopy level independently, using the total amount of absorbed light calculated by the radiation transfer scheme, which means that radiation transfer inside the canopy and photosynthesis are now fully consistent. In the new photosynthesis scheme, photosynthesis thus indirectly depends on canopy structure.

3.7 Forest management and natural mortality

Although forest management has developed a wide range of locally-appropriate and species-specific strategies (Pretzsch, 2009), the nature of large scale land surfaces models such as ORCHIDEE-CAN,

require only a limited number of contrasting strategies that are expected to be relevant at the spatial scale (e.g., 50x50 km) of global and regional modelling studies. Four management strategies were implemented based on their expected impact on biogeochemical and biophysical processes:

1. In unmanaged stands self-thinning drives stand dynamics and continues until too few trees are left on site. Subsequently, a stand replacing disturbance moves all standing biomass into the appropriate litter pools and a new stand is established.

2. High stand management is characterised by regular thinning and a final harvest cut. Thinning is decided on the basis of the deviation between the actual and potential stand density for any given diameter. This approach relates to the so-called relative density index (Fortin et al., 2012), the land use disturbance index (Luyssaert et al., 2011) or hemeroby and naturalness approaches (Schall and Ammer, 2013). Exceeding a threshold diameter results in a clear cut and the stand is replanted in the next year. For both thinning and harvest, leaves, roots and belowground wood is transferred to the appropriate litter pools whereas the aboveground woody biomass is removed from the site and stored in a product pool. Trees with a diameter below a species-specific threshold are stored in a short-lived product pool which mimics wood uses for fuel, paper and cardboard. Trees with larger dimensions are moved to medium and long-lived product pools which mimic for example, particle boards and timber usages, respectively.

3. Coppicing of the aboveground biomass is decided on stem diameter. At harvest, the root system is left intact and in between coppicing, no wood is harvested. Note that at present it is not possible to simulate coppicing-with-standards in ORCHIDEE-CAN.

4. In ORCHIDEE-CAN, stands under short rotation management are limited to poplar (*Populus spp.*) and willow (*Salix spp.*) forests. Stands are harvested at a prescribed age. Following a set number of harvest cycles, the root system is uprooted and the whole stand is replanted.

Different age classes are distinguished to better account for the structural diversity and its possible effects on the element, energy and water fluxes. A clear hierarchy was established for the mortality processes regarding the actual killing of trees (i.e., move their biomass to the litter or harvest pools). All of the processes determine first how much biomass they would remove in the absence of all the other processes. Afterwards, the killing is arranged in the most realistic way possible. A clear-cut event has the highest priority, followed by human thinning and finally natural mortality including self-thinning. If, for example, a forest is scheduled to be clear-cut, the entire forest biomass is subjected to the rules of the clear-cut and no other mortality occurs in that time-step.

In addition to forest management and natural prescribed mortality, a variety of changes have been made to processes involving vegetation mortality. A whole PFT within a grid-cell is now killed if, at the end of the day, the labile pool is empty and there is no carbon available in the leaf or carbohydrate reserve pool to refill it. In this situation, it will be impossible for the plant to assimilate

new carbon from the atmosphere as it will not be able to grow new leaves and thus initiate plant recovery. Furthermore, a forest can die if the density falls below a certain prescribed value. In the next time step a new young forest will be prescribed.

If a forest is thinned, it is assumed that the weakest trees will be thinned, and therefore human thinning reduces or even eliminates the natural mortality for that time-step. Natural mortality still happens on a daily time-step, while human-induced mortality happens only at the end of the year.

Self-thinning, as described below, takes priority over environmental mortality which is the mortality of individuals by insects, lightening, wind, drought, frost and heart rot. Environmental mortality is calculated by multiplying the stand biomass by an assumed mortality fraction of $1/k_{t_{resid}}$. Where self-thinning is less than this assumed environmental mortality, self-thinning is complemented by additional mortality to reach the set environmental mortality. Where self-thinning mortality exceeds the set environmental mortality, simulated self-thinning is assumed to include environmental mortality. The fire module that is available for the trunk but not for the ORCHIDEE-CAN branch simulates stand replacing fires rather than individual-tree based mortality due to lightening. The approach implemented in the ORCHIDEE-CAN branch could therefore be extended with models that simulate stand replacing mortality from fire, insects and storms.

The use of circumference classes adds a good deal of realism and flexibility to the ORCHIDEE-CAN simulations, but it also raises additional questions. For example, which trees should be targeted by which mortality? Given that self-thinning reflects the outcome of continuous resource competition, the largest trees are expected to be most successful when competing for resources, and therefore we assume that the smallest trees die first to reduce the stand density. Conversely, larger trees are more likely to die because of environmental stress factors, being more prone to cavitation, wind damage, lightening, and, heart rot. Therefore, we select more older trees to die from environmental mortality. While doing this also trees in the other circumference classes were killed based on the following recursive definition (cf Bellassen et al., 2010):

$$f_{death}^{icir} = \frac{f_{death}^{icir-1} \times k_{ddf}^{1-(k_{ncirc}-1)}}{m_{Ndeath}} \quad (38)$$

Where k_{ddf} is the death distribution factor, which is the factor by which the smallest and largest circumference classes differ (e.g., $k_{ddf} = 10$ means that the largest circumference class will lose ten times as much biomass as the smallest as a result of the mortality), m_{Ndeath} is a normalization factor so that sum of f_{death}^{icir} is unity, and f_{death}^1 is set equal to unity before normalization. As the stands are very close to even-aged, we set the factor k_{ddf} to be equal to 1. This means the same number of trees is killed in each circumference class. If, for some reason, there is not enough biomass in a given class to satisfy this distribution, the extra biomass is taken from the next smallest class (in the case the smallest class does not have enough, it is taken from the largest class).

Related to mortality is the question of the circumference class distribution. As mentioned above, trees in different circumference classes are preferentially killed by different processes. If the simulation is long enough (or if the mortality is aggressive enough), eventually the number of trees in some circumference classes may become zero. This would reduce the numerical resolution of the allocation scheme. When only one circumference remains populated, the scheme effectively loses its meaning as all the newly produced biomass is now be allocated to the only remaining circumference class. In order to maintain the same level of detail through the simulation, the distribution of all the circumference classes is recalculated at the end of each day. A normalized target distribution is specified as an input parameter (an exponential distribution is currently used), and this distribution is scaled to produce a target distribution for the current number of individuals. All of the current individuals are placed in these new classes until the target distribution is satisfied. The target distribution now contains, however, trees of multiple sizes, so we need to average them to find the new “model” tree for each class. By changing the size of the model tree in each class, we are able to preserve the total biomass of the stand as well as the total number of individuals. *Note that the boundaries of each diameter class are recalculated at each time step, this approach is a numerically efficient alternative to fixing the boundaries of each diameter class with a varying distribution.*

4 Description of the parametrization

The ORCHIDEE-CAN branch was specifically developed to quantify the climate effects of forest management over Europe. Although the developments are sufficiently general to be applied outside of Europe, the model was initially parametrized for the boreal, temperate and Mediterranean climate zones and validation focused on Europe. Parametrization of the tropical zone is subject of a follow-up study. *The parametrization of the model, including parameter optimization and tuning consisted of five major steps:*

1. Parameters related to carbon allocation (section 4.2), forest management and mortality (section 4.3, hydraulic architecture (section 4.4, canopy structure (section 4.5)), photosynthesis (section 4.6), and canopy radiation transfer (section 4.7) and for which observations exists at the species-level (section 4.1) were extracted from a wide range of sources (Tables S1-S5). Using the extracted species-level parameter values in ORCHIDEE without further processing avoids hidden model-tuning and largely reduces the likelihood that simulation results are biased by hidden calibration owing to a poor taxonomic definition of PFTs (Scheiter et al., 2013).
2. The phenology-related parameters of the deciduous MTCs were optimised by MacBean et al. (In prep.), using MODIS-derived NDVI data normalised to model fAPAR over the 2000 - 2008 time period.

3. The modulator (m_{ψ}) which accounts for processes in the the soil-plant continuum that are currently not modelled, was manually tuned against species distribution maps (section 4.4).
4. The coefficient for maintenance respiration was optimized making use of Bayesian calibration (section 4.8) against a compilation of 100+ observations of biomass production efficiency.
5. The leaf to sapwood area ratio was manually tuned (section 4.9) to match 100+ site-level GPP and LAI observations recorded over Europe.

4.1 Introducing twelve new PFTs

Similar to the ORCHIDEE trunk, the ORCHIDEE-CAN branch distinguishes 13 metaclasses (MTC) for vegetation. Outside Europe the original MTC classification of ORCHIDEE was kept, while inside Europe 12 new parameter sets representing the main European tree species were added. The default vegetation distribution map in ORCHIDEE, i.e., Olson et al. (1983), was replaced by an up-to-date global MTC map which has been produced using the ESA CCI ECV Land Cover map (<http://www.esa-landcover-cci.org/>)(Poulter et al., 2015). The mapping from land cover to MTC basically followed Poulter et al. (2011), although table 5 (the "cross-walking" table) has been updated following discussions with the LC-CCI team at Universite Catholique de Louvain. For the European domain, the global MTC distribution was overlaid by a tree species distribution map (Brus et al., 2012).

This study focusses on tree species with a coverage of more than 2% in Europe, yielding seven species groups covering in total 78.8% of the European forest area: *Betula sp.*, *Fagus sylvatica*, *Pinus sylvestris*, *Picea sp.*, *Pinus pinaster*, *Quercus ilex* and a group combining *Quercus robur* and *Quercus petraea*. For *Pinus sylvestris*, *Picea sp.* and *Betula sp.* An additional distinction between boreal and temperate forest was made for the species map and parametrization: trees located in Norway, Sweden and Finland were considered boreal, while trees growing at lower latitudes were categorized as temperate. Given the potential role of tree species of the Salicaceae genus in short rotation coppice management, a separate PFT was parametrized for *Populus sp.* Furthermore, to improve the parametrization of the MTC of boreal needleleaved deciduous forest, observations from *Larix sp.* were included when possible.

For these 12 forest species, 12 new PFTs were created with each PFT belonging to a single MTC (Table S2, S3 and S4) Almost 79% of the European forest was parametrized at the species level. The remaining 21% was reclassified in four residual groups, i.e., a temperate and boreal needleleaf evergreen and a temperate and boreal broadleaved residual group. For use outside Europe, the original MTC classification of ORCHIDEE was kept. The parameters of the residual groups and MTCs are the mean of the parameters of the species-level PFTs that are in the MTC, with the exception of albedo parameters that could be extracted from remote sensing products. Finally, separate PFTs were introduced for boreal grasses and croplands, which allowed for a boreal parametrization of

phenology, senescence and growth. This approach, which distinguishes a total of 28 PFTs, allows a higher taxonomic resolution over Europe, better defines forest types compared to the more general MTC approach and facilitates the use of observations to derive parameters.

755 4.2 Allocation

The allocation scheme relies on the leaf to sapwood area ratio (section 4.9) and the relationship between diameter and height. Following a logarithmic transformation of the more than 150,000 data points from the the national forest inventory data of Spain, France, Germany and Sweden, the two parameters (i.e. k_{α_1} and k_{β_1}) describing the relationship between diameter and height (equation 15) were fitted at the species-level making use of a least square regression. Parameter values for MTCs were derived by grouping the species into MTCs and fitting the parameters. Data sources and parameter estimates are presented in Tables S2 and S3.

4.3 Forest management and mortality

Forest management and tree mortality are controlled by (section 3.7): (1) maximum tree diameter (no symbolic notation; called `largest_tree_diam` in ORCHIDEE-CAN), (2) minimum stand density (no symbolic notation; called `ntrees_dia_profit` in ORCHIDEE-CAN), (3) environmental mortality (no symbolic notation; called `residence_time` in ORCHIDEE-CAN), (4) self-thinning (k_{α_2} and k_{β_2}) and, (5) anthropogenic thinning (no symbolic notation; called `alpha_RDI_upper`, `alpha_RDI_lower`, `beta_RDI_upper` and `beta_RDI_lower` in ORCHIDEE-CAN) where the parameters depend on the management strategy..

Maximum tree diameter was extracted from the French, Swedish, German and Spanish forest inventories as the observed 50% quantile for diameter at breast height. The 50% quantile rather than the observed maximum was used to account for the fact that large scale land surface models are expected to reproduce large scale patterns rather than local extremes. Minimum stand density was estimated as the expected stand density for the maximum tree diameter for a stand under self-thinning. Although both criteria are related to each other through the observed self-thinning relationship (see below), the minimum number of trees is used to decide when unmanaged forests should be replaced, whereas both the maximum diameter and the minimum number are used for managed sites as criteria to initiate a clear cut. Parameters for anthropogenic thinning are based on the national forest inventory data and checked against the JRC-database of species specific yield tables. Parameter values are presented in Table S5. Resource competition between trees in the same stand has been reported to result in the so-called self-thinning relationship that relates the number of individuals within a stand to the stand biomass (Reineke, 1933; Kira et al., 1953; Yoda et al., 1963):

$$(M_s + M_h) \times k_{\rho s} = k_{\alpha} \times (d_{ind})^{-k_{\beta}} \quad (39)$$

785 Where k_α and k_β are the constants of the self-thinning relationship. Furthermore, stem volume can be written as a function of tree diameter (d_{dbh}), tree height and stem form factor ($k_{\alpha'}$) to account for the fact that the stem shape is not a perfect cylinder:

$$(M_s + M_h) * k_{\rho s} = k_{\alpha'} \times (d_{dbh})^2 * d_h \quad (40)$$

Following the allometric relationship given in equation 15, tree height can be written as a function
790 of tree diameter. Hence, the self-thinning relationship can be re-written to relate stand diameter to stand density:

$$d_{dbh} = k_{\alpha 2} \times (d_{ind})^{-k_{\beta 2}} \quad (41)$$

Where, $k_{\beta 2}$ relates to $k_{\beta 1}$ (as in equation 15) as follows:

$$k_{\beta 2} = -3/2 \times (2 + k_{\beta 1}) \quad (42)$$

795 $k_{\alpha 1}$ and $k_{\beta 1}$ were estimated by fitting equation 15 to observed diameter and height of individual trees from NFI of Sweden, Germany, France and Spain. $k_{\beta 2}$ was calculated from equation 42 and $k_{\alpha 2}$ was estimated by fitting equation 41 to observations of the quadratic mean stand diameter and stand density from NFI data.

4.4 Hydraulic architecture

800 Initial choices of parameters for this scheme were based on the values and parameter sources listed by Hickler et al. (2006). All data sources were revisited and the search was extended to obtain values at the PFT rather than MTC level. Given that plant hydrology is rather well studied, observed parameters were available for most of the species. Data sources are listed in Table S1 whereas the parameter values are shown in Table S3. Our implementation of hydraulic architecture required
805 the introduction of a tuning parameter (m_ψ) to account for processes that are currently absent in the scheme, e.g. plant water storage and soil-root resistance. A process-based description of these processes (i.e., Sperry et al., 1998; Steppe et al., 2006) is being tested and should reduce the effect of the tuning parameter and eventually allow its removal from the model.

For the time being, the modulator m_ψ was tuned manually against the species distribution map
810 to obtain a match between the simulated and observed species distributions. When the modulator is set to zero, all PFTs experience excessive water stress resulting in large-scale plant mortality. The modulator was increased until the prescribed vegetation distribution which was based on remote-sensing observations (section 4.1), survived where it was prescribed. To this aim, the model was run for 50 years, forced with v5.2 of the CRU-NCEP climatology for Europe (Climatic Research Unit,

815 University of East Anglia). Note that the values of the modulator depend on the climate data that are used to force the model. Similarly the modulators may need to be re-tuned when ORCHIDEE-CAN is coupled to an atmospheric model.

4.5 Canopy structure

The relationship between diameter and projected crown surface area follows the model proposed by
820 Pretzsch (2009):

$$d_{csa} = k_{ap} \times d_{dbh}^{k_{bp}} \quad (43)$$

with parameters estimated using the dataset presented in Pretzsch and Dieler (2012). This dataset contains diameter and projected crown surface areas observations for over 37,000 individual trees in Europe covering almost 30 species. Following logarithmic transformation of the observations a
825 linear least square regression was used to fit species-specific parameter values. Parameter values are shown in Table S2. Parameter values for MTCs were derived by grouping the species into MTCs and fitting the parameters. No observations were available for the boreal zone and temperate evergreen deciduous species. For the boreal species a subset of the temperate observations (*Pinus sylvestris*, *Picea abies* and *Betula pendula*) was used, i.e., the relationship between d_{csa} and d_{dbh} was fitted
830 to all available data for *Pinus sylvestris*. Next, all observations with a d_{csa} that falls below the predicted d_{csa} were selected as considered to represent a boreal subset. Given the importance of snow pressure on crown structure, selecting observations with sub average d_{csa} is justifiable as a first approximation. Subsequently, the parameters were fitted to this subset of data. For *Quercus ilex* no data were available and parameters were tuned such that the crown diameter was 0.85 m less than
835 the tree height.

4.6 Analytical solution for photosynthesis

Three originally MTC-specific photosynthetic parameters ($k_{V_{cmax}}$, $k_{J_{max}}$ and k_{sla}) were derived at the species level by obtaining weighted site means for each species from the global leaf trait database TRY (Kattge et al., 2011) and additionally from Medlyn et al. (2002). Only $k_{V_{cmax}}$ and
840 $k_{J_{max}}$ standardized to a common formulation and parametrization of the photosynthesis model by (Farquhar et al., 1980) were used. Most $k_{V_{cmax}}$ and $k_{J_{max}}$ values in the TRY database had already been standardized to a reference temperature of 25°C (Kattge and Knorr, 2007). Subsequently, a species-specific $k_{J_{max,opt}}/k_{V_{cmax,opt}}$ ratio was calculated from the records which included both $k_{V_{cmax,opt}}$ and $k_{J_{max,opt}}$ measurements. From this ratio, which was within a range of 1.91-2.47
845 for each species, $k_{J_{max,opt}}$ was calculated for records which originally only included $k_{V_{cmax}}$. Only geo-referenced observations within Europe were used and the distinction between boreal and temperate forest was made similar to the species map. Depending on the species this resulted in 5 to 183

observations for k_{sla} and 11 to 173 observations for $k_{V_{cmax,opt}}$ and $k_{J_{max,opt}}$. From these observations species-specific means were calculated, weighted for differences in the number of observations per site. The parameter values are shown in Table S3.

4.7 Multi-layer two-way radiation scheme for tall canopies

The radiation transfer scheme makes use of parameters describing leaf and background properties, i.e., leaf single scattering and preferred scattering direction (for both visible (VIS) and near-infrared (NIR) wavelengths) and the so-called background albedo or the albedo of the surface below the dominant tree canopy (VIS and NIR). All parameters were taken from the Joint Research Centre Two-stream Inversion Package (JRC-TIP) (Pinty et al., 2011a, b). This is a software package (Pinty et al., 2007) which inverts a two-stream model (Pinty et al., 2006) to best fit the MODIS broadband visible and near-infrared white sky surface albedo from 2001 to 2010 at 1 km resolution (Pinty et al., 2011a). The inverse procedure implemented in the JRC-TIP is shown to be robust, reliable, and compliant with large-scale processing requirements (Pinty et al., 2011a). Furthermore, this package ensures the physical consistency between sets of observations, the two-stream model parameters, and radiation fluxes.

Only parameter values for which the posterior standard deviation of the probability density functions were significantly smaller than the prior standard deviation were selected from the JRC-TIP optimization (Pinty et al., 2011a), since this condition ensures statistically significant values. Species and MTC specific values were derived from JRC-TIP by performing a multiple regression. This method determines, in an objective way, how the fractions of each MTC or species explain the JRC-TIP parameter. The multiple regression was performed separately for the six parameters: the single scattering of leaves (for both VIS and NIR), the scattering direction of leaves (VIS and NIR) and the background albedo (VIS and NIR). Each JRC-TIP parameter was used as the dependent variable and the independent variables consisted of the fractions of each MTC (Poulter et al., 2015) or species (Brus et al., 2012). These fractions were used to find a linear function that best predicted each JRC-TIP parameter. The corresponding slope of a regression of each MTC or species fraction gives the MTC or species dependent JRC-TIP value. The multiple regression was performed without an intercept. To avoid pollution by the seasonal cycle, the multiple regression was applied only for the pixels of the Northern Hemisphere. Only pixels that were less than 10% covered by non-vegetative fractions were selected for the analysis and only significant results following an F-test and positive r^2 -values were selected. The derived parameter values are shown in Table S4.

4.8 Maintenance respiration

Both the trunk and ORCHIDEE-CAN branch reduce the definition of net primary production to biomass production; hence, carbon leaching from the roots, volatile organic emissions from the leaves, dissolved and particulate carbon losses through water fluxes and carbon subsidies to my-

corryhzae are not accounted for in the model. These fluxes are (incorrectly) accounted for in the modelled autotrophic respiration. Modelled autotrophic respiration should therefore be considered an effective rather than a true value. For this reason, the basal rate of autotrophic respiration was optimized against 126 site observations of the biomass production efficiency (k_{cmaint}) calculated as the ratio between annual biomass production and annual photosynthesis (Vicca et al., 2012; Campioli et al., In prep.), using a Bayesian optimization scheme. The scheme, for which more details are given in Santaren et al. (2007), uses a standard variational method based on the iterative minimization of a cost function that measures both the model data misfit and the parameter deviations from prior knowledge (Tarantola, 2005).

The simulations that were used in the Bayesian optimization prescribed a 20 m tall vegetation for temperate tree species, a 15 m tall vegetation for boreal tree species and a 10 m tall vegetation for Mediterranean tree species as its initial condition. This approach reduced the need for several decades of simulations to a single year to grow a mature forests. In total the simulations were run for 10 years and covered the European domain. The first year was discarded and the ratio between modelled GPP and NPP was averaged over the remaining 9 years. Prior to the optimization the observations were averaged for agricultural PFTs (0.57), and deciduous (0.44) and evergreen (0.53) forest PFTs, the observed uncertainty was 0.03. The parameter values were set to range between 0.0032 and 0.160. The optimization converged within 11 iterations and the optimized parameter values are shown in Table S2.

It remains untested how well the simulated effective autotrophic respiration represents the (rarely) observed autotrophic respiration. Note that in the cases of both the trunk and the ORCHIDEE-CAN branch of ORCHIDEE, a match between effective and observed autotrophic respiration should not be interpreted as evidence of desired model behaviour because several components of net primary production are not modelled yet.

After the optimization of the maintenance respiration coefficient (k_{cmaint}), the model simulates reasonable biomass production efficiency for a unit of photosynthesis. Hence, the final step of the parametrization focussed on optimizing the leaf area as this is one of the main drivers of photosynthesis.

4.9 Sapwood to leaf area ratio

The vegetation structure simulated by the ORCHIDEE-CAN branch is sensitive to the value of k_{ls} which describes the ratio between the leaf and sapwood area of an individual tree. The available observations show a wide range within and across forest species. Dependencies of k_{ls} on tree height (McDowell et al., 2002; Novick et al., 2009), tree diameter following stand thinning (Simonin et al., 2006) and CO₂ (Pataki et al., 2006) have been reported. Most observations, however, come from experiments where time was substituted by space which hampers teasing apart the sources of variability. Given the variation and uncertainty in the observations and the model sensitivity to this

parameter, we manually tuned its value within the observed range, to match European-wide observations of leaf area index as recorded in the Database of Global Forest Ecosystem Structure and Function Luyssaert et al. (2007).

This database was used to calculate a mean and maximum observed leaf area index at the species-level for the temperate and boreal region. Initially 20-year long European-wide simulations were used to simulate leaf area index of a species, when the large-scale leaf area index approached the mean target value and did not exceed the maximum value, the simulations were extended to reach 100 years for checking the temporal evolution of leaf area index. We deliberately optimized the sapwood to leaf area ratio (k_{ls}) by making use of stand-level data to reduce circularity with the model validation (see below).

Limited tests over a period of 100 years in a Scots pine forest at 51 – 52°N, 13 – 14°E (Fig. S1) suggested that optimizing k_{cmaint} and k_{ls} had the largest effect on the maximum LAI, which decreased with almost 17% after optimization compared to a simulation with prior parameter values. Mean annual GPP, mean annual transpiration and basal area decreased with, respectively 6, 6 and 7% compared to a simulation with prior parameter values (Fig. S1).

5 Validation

ORCHIDEE-CAN is designed as the land surface model to be coupled to the atmospheric model LMDz. As such future applications of ORCHIDEE-CAN are expected to be regional to global in the spatial domain and to span several years in the temporal domain. Given its anticipated uses, the ability of the model to reproduce large-scale spatial patterns as well as their inter-annual variability is essential. The first applications of the model, both offline and coupled to the atmosphere, will focus on Europe. The validation, therefore, reports performance indices both over Europe as over eight separate regions within Europe (Bellprat et al., 2012). These eight regions, which partially overlap, are defined after Bellprat et al. (2012). Furthermore, the performance indices are calculated for winter, spring, summer and autumn and thus allow to evaluate the capacity of the model to reproduce observed annual cycles.

In addition to the root mean square error, a land performance index (LPI) based on the principles laid out for the Climate Performance Index (Murphy et al., 2004, their SI) was also calculated. LPI normalizes the root of the squared differences between the simulations and observations by the observed spatial and temporal variance. The LPI was used to estimate the likelihood that the simulated variable belongs to the same population as the observed variable, defined as $\exp(-0.5LPI^2)$. An LPI equal to 1 indicates that the model correctly reproduces the mean observed value and implies a likelihood of 61% (Murphy et al., 2004) that the simulations and observations come from the same population. Similarly, an LPI of 2 reduces this likelihood to 13%. An LPI of less than 0.32 has a likelihood of more than 95% and therefore indicates a statistically significant result.

While developing ORCHIDEE-CAN, the numerical approaches that added functionality to the code were selected on the basis of their performance at the site-level (see below). Rather than running the same site-level tests for our implementation, we performed a complementary large-scale validation. The strength of our approach lies not in the details, as is the case for site-level validation, but in its width by simultaneously testing model performance for structural variables such as basal area (de Rigo et al., 2014), canopy structure (Pinty et al., 2011a) and canopy height (Simard et al., 2011), biogeochemical fluxes such as GPP (Jung et al., 2008), biophysical fluxes such as albedo (Schaaf et al., 2002) and fluxes at the interface of biogeochemistry and biophysics such as evapotranspiration (Jung et al., 2008). The selection of variables was limited by the availability of spatially explicit data-derived products for Europe.

For the validation, both the trunk and ORCHIDEE-CAN branch were run from 1850 to 1900 using CRU-NCEP climate forcing from 1901-1950 at 0.5 degree resolution. From 1901 until 2012, the corresponding CRU-NCEP forcing data for each year were used. Both versions used the 11-layer soil hydrology, the single-layer energy budget and the same land cover map (Poulter et al., 2015). Given that no European-wide, spatially explicit and data-derived products were found for the validation of the net carbon flux, there was no need for a carbon spin-up. For the ORCHIDEE-CAN branch, the observed tree height and basal area were compared against the simulation values at the end of 2010 (the trunk does not simulate these variables). For both the trunk and the ORCHIDEE-CAN branch, the observed GPP, evapotranspiration, effective LAI and VIS and NIR albedos were compared against monthly means between 2001 and 2010.

5.1 Species versus PFTs

In ORCHIDEE-CAN the PFT concept was refined by parametrizing the main European tree species groups (section 4.1). To evaluate the effect of the species parametrization we performed a companion simulation for the configuration described above, but at the MTC level. Model performance was barely affected by the use of the MTC parameters, compared to the simulation with the species parameters (Fig. S2 for RMSE scores).

5.2 Allocation

In ORCHIDEE-CAN, functional relationships which vary by species and light stress are used to allocate carbon among the fine roots, foliage and sapwood. The allocation scheme largely follows Zaehle and Friend (2010), who in turn was inspired by Sitch et al. (2003). Approaches simulating allocation based on functional relationships were found to outcompete allocation schemes based on constant fractions or resource limitation (De Kauwe et al., 2014). The ability of these schemes to reproduce foliage, fine root and sapwood reported in large observational data sets (for example, Luyssaert et al., 2007) demonstrates that these schemes capture the main observed features (Zaehle and Friend, 2010). In addition, allocation schemes making use of functional relationships were also

capable of simulating the observed effect of elevated CO₂ on two mature forest ecosystems (De
990 Kauwe et al., 2014). Despite these successes, the schemes were reported to be sensitive to their
parametrization. Differences in parameters were reported to result in substantial differences in the
simulated allocation. The parameters for the functional relationships used in ORCHIDEE-CAN are
given in Table S2. The main conceptual difference between the allocation scheme by Zaehle and
Friend (2010) and ORCHIDEE-CAN is that the latter was designed to simulate one or more diameter
995 classes.

Given that photosynthesis is still calculated at the stand level (and thus not at the tree level) the
allocation rule of Deleuze et al. (2004) was integrated in the functional allocation scheme to account
for light and resource competition within a stand. Where the functional relationships are used to sim-
ulate carbon allocation within an individual tree of a given diameter, the rule of Deleuze et al. (2004)
1000 allocates carbon across the different diameter classes. The allocation rule which models the radial
increment for individual trees in pure even-aged stands was successfully tested for Norway spruce
and Douglas fir stands in France (Deleuze et al., 2004). A similar approach for modelling radial
increment has already been implemented in a version close to the trunk of ORCHIDEE (Bellassen
et al., 2010) and was able to successfully simulate stand characteristics such as height, basal area
1005 and stand diameter (Bellassen et al., 2011). This previous implementation differs from the current
implementation in its time resolution (which is now daily instead of yearly), its analytical solution
and the underlying allocation scheme (which is now based on functional relationships instead of
resource limitation).

The aforementioned studies performed a detailed validation of the two approaches dealing with
1010 carbon allocation, which were combined in ORCHIDEE-CAN. Complementary to these studies we
performed a European-wide validation of our implementation and parametrization of these well-
tested schemes against a remote-sensing based map of tree height (Simard et al., 2011), upscaled
eddy-covariance observations for GPP (Jung et al., 2008) and a map of basal area based on national
forest inventory data (de Rigo et al., 2014). The model's ability to reproduce GPP is thought to
1015 reflect its capacity to simulate the foliage biomass, a correct simulation of height reflects the model's
capacity to simulate aboveground woody biomass and its capacity to reproduce observed basal areas
suggest that the interaction of stand density and individual tree diameter are well-captured.

The new implementation and parametrization of the within-tree and within-stand allocation schemes
were found to have an 91, 68 and 72% chance that the simulations reproduced the observations for
1020 GPP, tree height and basal area for Europe, respectively (Table 3). Given that basal area and height
are not available from the trunk version of ORCHIDEE, we could not compare the performance of
model versions in this respect. With respect to GPP, the ORCHIDEE-CAN branch was found to out-
perform the trunk by 12% and thus increased the likelihood that ORCHIDEE-CAN is an unbiased
simulator of the spatial and temporal variability of GPP from 79 to 91%. Improved performance of

the ORCHIDEE-CAN branch compared to the trunk is observed for all regions in summer where the RMSE of GPP was halved from 2.5-5 gC m⁻² day⁻¹ to 1-2 gC m⁻² day⁻¹ (Fig. 2, 3 and 4).

Although part of the high likelihood could be due to the fact that the observed GPP was upscaled making use of similar climatologies being used as the forcings of the models, this circularity could neither have contributed to the improved performance between the trunk and the ORCHIDEE-CAN branch nor to the decrease in RMSE. The improvements are thought to be due to structural changes to the model such as allocation, hydraulic architecture and canopy structure as well as to the use of more consistent parametrization.

5.3 Plant water supply

Our implementation of plant hydraulic architecture was largely based on the scheme of Hickler et al. (2006), which was tested globally and at site level. Global simulation results for actual evapotranspiration were found to reproduce available data (Baumgartner and Reichel, 1975; Henning, 1989). At the site level, the model agreed well with the magnitude and seasonality of eddy-covariance measurements of actual evapotranspiration for 15 European forests sites (EUROFLUX), with a tendency to slightly overestimate actual evapotranspiration for 6 sites (Hickler et al., 2006).

The maximum amount of water that can be transported by a tree relies on the hydraulic architecture of the tree and therefore on the capacity of the model to simulate tree and stand dimensions as well as on the model's capacity to simulate soil water content. As an additional test, our implementation of the model was compared against the upscaled eddy-covariance measurements for GPP and actual evapotranspiration (Jung et al., 2008). The capacity to jointly reproduce GPP and actual evapotranspiration is an indicator that the model successfully reproduces the coupling between CO₂ and water exchange. Model validation showed 91 and 87% chance (compared to 79 and 45% for the trunk) that ORCHIDEE-CAN reproduces the upscaled GPP and actual evapotranspiration data (Table 3, Fig.4). The RMSE for actual evapotranspiration during summer dropped well below 1 mm day⁻¹ for most regions (Fig. 2), whereas it never dropped below 1 mm day⁻¹ for the trunk (Fig. 3).

5.4 Canopy structure

The canopy structure model by Haverd et al. (2012) was previously validated against ground-based LIDAR data for several test sites with varying density, structural complexity, layering and clumping (Lovell et al., 2012). Model-derived canopy gap probabilities compared with observations using a one-sample t-test were significant for 11 out of 12 test sites. We considered this result as a sufficient proof to use this canopy structure model in the ORCHIDEE-CAN branch and added to its validation by comparing the simulated canopy structure model over Europe against a remote-sensing based map of tree height (Simard et al., 2011) and the JRC-TIP effective LAI product (Pinty et al., 2011a). The effective LAI value expresses the capability of the canopy to intercept direct radiation, and is thus associated with the probability distribution function of the canopy gaps (Haverd et al., 2012).

Thus the effective LAI contains information about the forest structure and leaf distribution of the canopy. In the ORCHIDEE-CAN branch, canopy structure is used to calculate the albedo, roughness length, absorbed light for photosynthesis and leaf area that is coupled to the atmosphere for, e.g., transpiration and interception of precipitation.

The ORCHIDEE-CAN branch is the first branch of ORCHIDEE that makes use of an effective LAI to calculate the interaction between the canopy and the atmosphere. The LPI and RMSE of the branch, therefore, cannot be compared against the trunk. Overall, the combined implementation of the allocation scheme and the canopy structure model shows a 67% chance to reproduce the satellite-based estimates for effective LAI. Surprisingly, effective LAI is better simulated in spring and autumn when dynamics within the canopy are substantial due to leaf on-set and senescence. For the periods when the effective LAI is expected to be most stable, i.e., summer and winter, LPI approached and frequently exceeded 1 (data not shown). Part of this shortcoming may be due to the lack of shrubs in the land cover classification. In the model, shrublands are replaced by forest and/or grasslands, likely resulting in differences between the observed and simulated canopy structure. This lapse also appears in the RMSE of effective LAI (RMSE higher than 0.8, Fig. 2)

5.5 Top of the canopy albedo

The radiation transfer model (Pinty et al., 2006) has been validated extensively against realistic complex three-dimensional canopy scenarios (Pinty et al., 2006) and as part of the RAdiation transfer Model Intercomparison (RAMI) project. The 1-D canopy radiation transfer model by Pinty et al. (2006) was demonstrated to accurately simulate both the amplitude and the angular variations of all radiant fluxes with respect to the solar zenith angle (Widłowski et al., 2011). In addition, the radiation transfer model and its effective values extracted from the JRC-TIP data set were successfully applied to a single forest site (Pinty et al., 2011c).

Previously we reported on the capacity of the radiation transfer model to simulate the effects of forest management on albedo (Otto et al., 2014). For the latter, forest properties were prescribed and the radiation transfer model was validated against top-of-the-canopy albedo data from five observational sites. Differences in the spatial scales between the observed and simulated albedo values were accounted for by presenting the mean June albedo during 2001-2010 (Otto et al., 2014). The simulated summertime canopy albedo falls within the range of observation. However, there occurs a slight overestimation in the near-infrared wavelength band compared to the single site measurement. Too high near-infrared single scattering albedo values for pine, as obtained from the JRC-TIP product, are the most likely cause. The observed deviation is not due to a shortcoming in the model itself but reflects the difficulties the JRC-TIP has with optimizing parameter values in the absence of field observations in the specific case of sparse canopies (Otto et al., 2014).

For the spatial validation we use the white-sky albedo (VIS and NIR) from Moderate Resolution Imaging Spectroradiometer (MODIS, Schaaf et al., 2002) at 0.5° resolution (distributed in netCDF

format by the Integrated Climate Data Center (ICDC, <http://icdc.zmaw.de>) University of Hamburg, Hamburg, Germany). Over large spatial and temporal domains the ORCHIDEE-CAN branch reproduces the observed VIS and NIR albedo and its variability; LPI for the albedo in the visible light is especially satisfying with a likelihood of 92% for the simulations to come from the same population as the observations (Table 3). This high overall performance index, however, hides performance issues over Scandinavia and the Alps during the snow season. The RMSE for VIS and NIR albedo without snow lies around 0.05, whereas during the snow season the RMSE increases to 0.20 (VIS) and 0.18 (NIR) over these regions (Fig. 2). When the ORCHIDEE-CAN branch is coupled to an atmospheric model, however, these deviations will only have a minor effect on the climate, owing to low incoming radiation during most of the snow season, especially in Scandinavia.

Previous validation of the radiation transfer model showed that the largest discrepancies were occurring in the near-infrared domain with a snow covered background (Pinty et al., 2006). With the exception of the snow-covered season, the new albedo scheme, that relies on the simulated canopy structure, resulted in a substantial improvement of 0.05-0.15 compared to the trunk for the RMSE in both the VIS and NIR range in Scandinavia and the Alps (Fig. 2 and 3). The European LPI-based likelihood that our model simulations come from the same populations as the MODIS albedo increased by a remarkable 11 and 23% for, respectively, NIR and VIS albedo (from 61% and 69% for the trunk to 72 and 92% for the ORCHIDEE-CAN, Table 3).

Given that the parametrization of the canopy radiation transfer model used in ORCHIDEE-CAN relies on MODIS, the high likelihood may not come as a surprise. However, our implementation of the radiation transfer model also relies on the simulated absorbed light, simulated GPP, simulated allocation and simulated canopy structure (which depends on mortality and forest management). In the absence of all these processes our canopy radiation transfer model is expected to reproduce the MODIS data with a probability of 100%. Hence, the likelihood of 72% and 92% (for NIR and VIS, respectively) could also be interpreted as a verification of the aforementioned calculations; all calculations that determine the canopy structure reduce the reproducibility of the data by only 8-28% (100% to 72 or 92%).

5.6 Energy fluxes

The multilayer scheme is in the process of a detailed evaluation across a range of tests conditions (Ryder et al., 2014), and further validation across a range of sites is on-going. The scheme is able to produce within-canopy temperature and humidity profiles, and successfully simulates the in-canopy radiation distribution, as well as the separation of the canopy from the soil surface. However, in order to preserve a measure of continuity with previous evaluations of the model, the multilayer solution is here set to single layer operation mode, which includes the effects of hydraulic limitation (section 3.2) and canopy structure (section 3.3) on the energy budget.

The single-layer set-up of the multi-layer solution makes use of an improved albedo estimation and is therefore expected to better simulate the net radiation that needs to be redistributed in the canopy. This has been confirmed at a single site with a sparse canopy (Ryder et al., 2014). Furthermore, the improvements in actual evapotranspiration in addition to the low RMSE (Fig. 2) are expected to be propagated in the performance of the energy budget.

5.7 Forest management strategies

Model comparison has previously demonstrated that explicitly treating thinning processes is essential to reproduce local and large scale biomass observations (Wolf et al., 2011). This finding justifies the implementation of generic approaches to forest management despite the difficulties associated with defining and quantifying forest management and its intensity (Schall and Ammer, 2013). Although the use of so-called naturalness indices, in which the current state of the forest is referenced against the potential state of the forest, has been criticised because of difficulties in defining the potential state of the forest (Schall and Ammer, 2013), such approaches were demonstrated to correctly rank different management strategies according to their intensity (Luyssaert et al., 2011).

Naturalness indices making use of only diameter and stand density or the so-called Relative Density Index (RDI) have been previously implemented at the stand-level (Fortin et al., 2012) and as well as in large scale models (Bellassen et al., 2010). This approach was shown to successfully reproduce the biomass changes during the life cycle of a forest (Bellassen et al., 2011; Fortin et al., 2012). The implementation of a forestry model based on the relative density index was reported to perform better than simple statistical models for stand-level variables such as stand density, basal area, standing volume and height (Bellassen et al., 2011). Although the performance of the model was reported as less satisfying for tree-level variables, the approach is nevertheless considered reliable to model the effects of forest management on biomass stocks of forests across a range of scales from plot to country (Bellassen et al., 2011).

In the absence of forest management, ORCHIDEE-CAN simulates that the stands develop into tall canopy (Fig 5a), with a high biomass (Fig 5b), a substantial dead wood and litter pool (Fig 5c) and no harvest (Fig 5d). High stand management reduces the height, standing biomass and litter pools (Fig 5a-c) but produces biomass for harvest (Fig 5d). Under coppicing, the reduction in forest age is reflected in a shorter canopy and lower biomass and litter pools (Fig a-c) compared to high stand management. The harvest is more evenly spread in time but falls below the harvest generated by high stand management (Fig 5d). Given the shorter rotations, canopy height, standing biomass and litter pools are lower for short rotation coppicing with poplar and willow compared to all other management strategies applied on oak forest (Fig 5a-c). Short rotation coppice was harvested every 3 years resulting in a quasi-continuous supply of woody biomass (Fig. 5d).

The forestry model implemented in ORCHIDEE-CAN is based on the RDI approach by Bellassen et al. (2010). We complemented earlier validation of such an approach over France (Bellassen et al.,

2011) by a new European-wide validation for basal area. At the European scale we verified the simulated basal area and height against observed basal area from national forest inventories (de Rigo et al., 2014) and height from remote-sensing (Simard et al., 2011). With an RMSE of 3-7 for height and 7-15 for BA, and a chance of, respectively, 68 and 72% to reproduce the data at the European scale (Table 3), our model is capable of correctly simulating the mean height and basal area but fails to capture much of the spatial variability (Fig. 4, temporal variability was not considered because the data products were only available for one time period).

Furthermore, we evaluated basal area and tree diameter at the species-level for 11 regions over France which represents a finer spatial-scale than targeted by the model developments and their parametrization. The data were extracted from the French forest inventory between 2005 and 2010 and we used the same simulations as for the European validation in the previous paragraph. Pixels contained in the French inventory data we selected and for both simulations and observations we calculated a moving average for the diameter and basal area per age class to then calculated the RMSE. To account for intrinsic species differences in diameter and basal area we normalized the RMSE. The RMSE was lower than 30% of the mean tree diameter or mean basal area for each region for *Betula sp.*, *Pinus pinaster* and *Quercus ilex*. For *Fagus sylvatica*, *Pinus sylvestris*, *Picea sp.* and *Quercus robur/petraea* (Fig. 6). The relative RMSE of diameter and basal area exceeded 50% for one to four regions for tree diameter and basal area (not shown).

The inability to fully capture the observed spatial variability in the simulation could be due to the simulation protocol that started in 1850 with 2 to 3 meter tall trees all over Europe. A longer simulation accounting for the major historical changes in forest management such as the reforestation in the 1700s following an all time low in the European forest cover, the start of high stand management at the expense of coppicing in the early 1800s, and the reforestation programs following World War II (Farrell et al., 2000) is expected to improve the spatial variability in tree height and basal area. Regional deviations such as those observed in the Iberian Peninsula or over the entire Mediterranean (thus including part of the Iberian Peninsula) may be due to the lack of shrubs in the land cover map and parametrization of the ORCHIDEE-CAN branch. Therefore the models simulates a higher stand density and higher basal area for regions where in reality shrubs occur (Fig. 4) .

The parametrization of the forestry module strongly depends on the national forest inventories from Spain, France, Germany and Sweden. Therefore verification against the same data contains little information about the model quality. Nevertheless, no time-dependent relationships were used in the ORCHIDEE-CAN branch thus the model's capacity to reproduce the relationship between basal area and stand age, diameter and stand age or wood volume and stand age could be considered as largely independent test of the model quality. These tests were performed over 8 bioclimatic regions of France and the ORCHIDEE-CAN branch was found to largely capture the time dependencies of basal area, diameter and wood volume (not shown).

6 Conclusions

ORCHIDEE-CAN (SVN r2290) differs from the trunk version of ORCHIDEE (SVN r2243) by the
1205 allometric-based allocation of carbon to leaf, root, wood, fruit and reserve pools; the transmittance,
absorbance and reflectance of radiation within the canopy; and the vertical discretisation of the en-
ergy budget calculations. Conceptual changes towards a better process representation were made
for the interaction of radiation with snow, the hydraulic architecture of plants, the representation
of forest management and a numerical solution for the photosynthesis formalism of Farquhar, von
1210 Caemmerer and Berry. Furthermore, these changes were extensively linked throughout the code to
improve the consistency of the model. By making use of observation-based parameters the physio-
logical realism of the model was improved and significant reparametrization was done by introduc-
ing twelve new parameter sets that represent specific tree species or genera rather than a group of
phylogenetically often unrelated species, as is the case in widely used plant functional types (PFT).
1215 As PFTs have no meaning outside the **scientific** community, the species level parametrization of the
ORCHIDEE-CAN branch can deliver actionable information to decision-makers and forest owners
on the implications of management strategies on the climate.

Model performance was tested against spatial explicit or upscaled data for basal area, tree height,
canopy structure, GPP, albedo and evapotranspiration over Europe. The tested data streams repre-
1220 sented biogeochemical fluxes, biophysical fluxes and forest management related vegetation charac-
teristics. Enhanced process representation in ORCHIDEE-CAN compared to the trunk version, was
found to increase model performance regarding its ability to reproduce large-scale spatial patterns of
all tested data streams as well as their inter-annual variability over Europe. Although this validation
approach gives us confidence in the large-scale performance of the model over Europe, additional
1225 validation is recommended for other regional applications or higher resolution studies.

7 Author contribution

- Developed and parametrized the ORCHIDEE-CAN model: Kim Naudts, James Ryder, Matthew
J. McGrath, Julianne Otto, Sebastiaan Luyssaert, Nicolas Vuichard, Didier Solyga
- 1230 – Evaluated the performance of the ORCHIDEE-CAN model: Kim Naudts, James Ryder, Ju-
lianne Otto, Matthew J. McGrath, Sebastiaan Luyssaert, Aude Valade, Yiyang Chen, Fabienne
Maignan
- Contributed Fortran code: Vanessa Haverd (canopy gaps), Bernard Pinty (albedo), Valentin
Bellassen (forestry)

- 1235 – Provided/shared observational data sets or tools for model parametrization: Hans Pretzsch,
Päivi Merilä, Jens Kattge, Gerhard Bönisch, Matteo Campioli, Josep Penuelas, Detlef Schulze,
Toon De Groote, Gonzalo Berhongaray, Yuan Yan, Philippe Peylin
- Developed driver data: Natasha MacBean
- Maintained and developed the run environment: Josefine Ghattas

1240 **8 Code availability**

The code and the run environment are open source (<http://forge.ipsl.jussieu.fr/orchidee>). Nevertheless readers interested in running ORCHIDEE-CAN are encouraged to contact the corresponding author for full details and latest bug fixes.

Acknowledgements. JR, YC, MJM, JO, KN and SL were funded through ERC starting grant 242564 (DO-
1245 FOCO), and AV was funded through ADEME (BiCaFF). ESA ECV landcover also supported this work. The research leading to these results has received funding from the European Community's Seventh Framework Programme (FP7/ 2007-2013) under the Grant Agreement n° 284181-TREES4FUTURE. The authors would like to thank Daniele de Rigo for providing a basal area map for Europe.

References

- 1250 Amiro, B., Barr, A., Black, T., Iwashita, H., Kljun, N., McCaughey, J., Mogenstern, K., Murayama, S., Nesic, Z., and Orchansky, A.: Carbon, energy and water fluxes at mature and disturbed forest sites, Saskatchewan, Canada, *Agricultural and Forest Meteorology*, 136, 237–251, doi:10.1016/j.agrformet.2004.11.012, <http://linkinghub.elsevier.com/retrieve/pii/S0168192305002017>, 2006a.
- Amiro, B., Orchansky, A., Barr, A., Black, T., Chambers, S., Chapin III, F., Goulden, M., Litvak, M., Liu,
1255 H., McCaughey, J., McMillan, A., and Randerson, J.: The effect of post-fire stand age on the boreal forest energy balance, *Agricultural and Forest Meteorology*, 140, 41–50, doi:10.1016/j.agrformet.2006.02.014, <http://linkinghub.elsevier.com/retrieve/pii/S0168192306002267>, 2006b.
- Amthor, J. S.: The role of maintenance respiration in plant growth., *Plant, Cell and Environment*, 7, 561–569, doi:10.1111/1365-3040.ep11591833, <http://doi.wiley.com/10.1111/1365-3040.ep11591833>, 1984.
- 1260 Bala, G., Caldeira, K., Wickett, M., Phillips, T. J., Lobell, D. B., Delire, C., and Mirin, A.: Combined climate and carbon-cycle effects of large-scale deforestation., *Proceedings of the National Academy of Sciences of the United States of America*, 104, 6550–5, doi:10.1073/pnas.0608998104, <http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=1871823&tool=pmcentrez&rendertype=abstract>, 2007.
- Baldocchi, D.: A multi-layer model for estimating sulfur dioxide deposition to a deciduous oak forest
1265 canopy, *Atmospheric Environment*, 22, 869–884, doi:10.1016/0004-6981(88)90264-8, <http://linkinghub.elsevier.com/retrieve/pii/0004698188902648>, 1988.
- Ball, J. T., Woodrow, I. E., and Berry, J. A.: A model predicting stomatal conductance and its contribution to the control of photosynthesis under different environmental conditions , in: *Progress in Photosynthesis Research*, edited by Biggins, J. and Nijhoff, M., pp. 221–224, Martinus-Nijhoff Publishers, Dordrecht, Netherlands,
1270 1987.
- Baumgartner, A. and Reichel, E.: *Die Weltwasserbilanz*, R. Oldenburg Verlag, Munich, 1975.
- Bellassen, V., Le Maire, G., Dhôte, J., Ciais, P., and Viovy, N.: Modelling forest management within a global vegetation model—Part 1: Model structure and general behaviour, *Ecological Modelling*, 221, 2458–2474, doi:10.1016/j.ecolmodel.2010.07.008, <http://linkinghub.elsevier.com/retrieve/pii/S0304380010003480>, 2010.
1275
- Bellassen, V., le Maire, G., Guin, O., Dhôte, J., Ciais, P., and Viovy, N.: Modelling forest management within a global vegetation model—Part 2: Model validation from a tree to a continental scale, *Ecological Modelling*, 222, 57–75, doi:10.1016/j.ecolmodel.2010.08.038, <http://linkinghub.elsevier.com/retrieve/pii/S0304380010004527>, 2011.
- 1280 Bellprat, O., Kotlarski, S., Lüthi, D., and Schär, C.: Objective calibration of regional climate models, *Journal of Geophysical Research*, 117, D23 115, doi:10.1029/2012JD018262, <http://doi.wiley.com/10.1029/2012JD018262>, 2012.
- Best, M. J., Beljaars, A., Polcher, J., and Viterbo, P.: A Proposed Structure for Coupling Tiled Surfaces with the Planetary Boundary Layer, *Journal of Hydrometeorology*, 5, 1271–1278, doi:10.1175/JHM-382.1, <http://journals.ametsoc.org/doi/abs/10.1175/JHM-382.1>, 2004.
1285
- Betts, R. A.: Offset of the potential carbon sink from boreal forestation by decreases in surface albedo., *Nature*, 408, 187–90, doi:10.1038/35041545, <http://www.ncbi.nlm.nih.gov/pubmed/11089969>, 2000.

- Bonan, G. B.: Forests and climate change: forcings, feedbacks, and the climate benefits of forests., *Science* (New York, N.Y.), 320, 1444–9, doi:10.1126/science.1155121, <http://www.ncbi.nlm.nih.gov/pubmed/18556546>, 2008.
- Bonan, G. B., Levis, S., Sitch, S., Vertenstein, M., and Oleson, K. W.: A dynamic global vegetation model for use with climate models: concepts and description of simulated vegetation dynamics, *Global Change Biology*, 9, 1543–1566, doi:10.1046/j.1365-2486.2003.00681.x, <http://doi.wiley.com/10.1046/j.1365-2486.2003.00681.x>, 2003.
- Bonan, G. B., Williams, M., Fisher, R. a., and Oleson, K. W.: Modeling stomatal conductance in the earth system: linking leaf water-use efficiency and water transport along the soil–plant–atmosphere continuum, *Geoscientific Model Development*, 7, 2193–2222, doi:10.5194/gmd-7-2193-2014, <http://www.geosci-model-dev.net/7/2193/2014/>, 2014.
- Botta, A., Viovy, N., Ciais, P., Friedlingstein, P., and Monfray, P.: A global prognostic scheme of leaf onset using satellite data, *Global Change Biology*, 6, 709–725, doi:10.1046/j.1365-2486.2000.00362.x, <http://doi.wiley.com/10.1046/j.1365-2486.2000.00362.x>, 2000.
- Brus, D., Hengeveld, G., Walvoort, J., Goedhart, P., Heidema, A., Nabuurs, G., and Gunia, K.: Statistical mapping of tree species over Europe., *European Journal of Forest Research*, 131, 145–157, 2012.
- Campioli, M., Vicca, S., Luyssaert, S., Bilcke, J., Ceschia, E., Chapin III, F., Ciais, P., Fernández-Martínez, M., Malhi, Y., Obersteiner, M., Olefeldt, D., Papale, D., Piao, S., Peñuelas, J., Sullivan, P., Wang, X., Zenone, T., and Janssens, I.: Management improves the efficiency of biomass production of global terrestrial ecosystems, In prep.
- Chen, Y., Ryder, J., Naudts, K., Bastriko, V., Mcgrath, M. J., Otto, J., Launiainen, S., Ogée, J., Elbers, J. A., Foken, T., Tiedemann, F., Heinesch, B., Black, A., Haverd, V., Loustau, D., Gorsel, E. V., Knohl, A., Moors, E., Vessala, T., Otlé, C., Pelin, P., Polcher, J., and Luyssaert, S.: Improving energy partitioning and the nighttime energy balance by implementation of a multilayer energy budget in ORCHIDEE-CAN, pp. 1–25, In prep.
- Cochard, H., Martin, R., Gross, P., and Bogeat-Triboulot, M.: Temperature effects on hydraulic conductance and water relations of *Quercus robur* L., *Journal of Experimental Botany*, 51, 1255–1259, 2000.
- Collatz, G., Ribas-Carbo, M., and Berry, J.: Coupled photosynthesis-stomatal conductance model for leaves of C4 plants, , 19, 519–538, 1992 Title, *Aust. J. Plant Physiol.*, 19, 519–538, 1992.
- Cox, P. M., Betts, R. A., Bunton, C. B., Essery, R. L. H., Rowntree, P. R., and Smith, J.: The impact of new land surface physics on the GCM simulation of climate and climate sensitivity, *Climate Dynamics*, 15, 183–203, doi:10.1007/s003820050276, <http://link.springer.com/10.1007/s003820050276>, 1999.
- Davin, E. L., de Noblet-Ducoudré, N., and Friedlingstein, P.: Impact of land cover change on surface climate: Relevance of the radiative forcing concept, *Geophysical Research Letters*, 34, n/a–n/a, doi:10.1029/2007GL029678, <http://doi.wiley.com/10.1029/2007GL029678>, 2007.
- De Kauwe, M. G., Medlyn, B. E., Zaehle, S., Walker, A. P., Dietze, M. C., Hickler, T., Jain, A. K., Luo, Y., Parton, W. J., Prentice, I. C., Smith, B., Thornton, P. E., Wang, S., Wang, Y.-P., Wärlind, D., Weng, E., Crous, K. Y., Ellsworth, D. S., Hanson, P. J., Seok Kim, H., Warren, J. M., Oren, R., and Norby, R. J.: Forest water use and water use efficiency at elevated CO₂: a model-data intercomparison at two contrasting

temperate forest FACE sites., *Global change biology*, 19, 1759–79, doi:10.1111/gcb.12164, <http://www.ncbi.nlm.nih.gov/pubmed/23504858>, 2013.

- De Kauwe, M. G., Medlyn, B. E., Zaehle, S., Walker, A. P., Dietze, M. C., Wang, Y.-P., Luo, Y., Jain, A. K., El-Masri, B., Hickler, T., Wårlind, D., Weng, E., Parton, W. J., Thornton, P. E., Wang, S., Prentice, I. C., Asao, S., Smith, B., McCarthy, H. R., Iversen, C. M., Hanson, P. J., Warren, J. M., Oren, R., and Norby, R. J.: Where does the carbon go? A model-data intercomparison of vegetation carbon allocation and turnover processes at two temperate forest free-air CO₂ enrichment sites., *New Phytologist*, 203, 883–99, doi:10.1111/nph.12847, <http://www.ncbi.nlm.nih.gov/pubmed/24844873>, 2014.
- de Noblet-Ducoudré, N., Boisier, J.-P., Pitman, A., Bonan, G. B., Brovkin, V., Cruz, F., Delire, C., Gayler, V., van den Hurk, B. J. J. M., Lawrence, P. J., van der Molen, M. K., Müller, C., Reick, C. H., Strengers, B. J., and Voldoire, a.: Determining Robust Impacts of Land-Use-Induced Land Cover Changes on Surface Climate over North America and Eurasia: Results from the First Set of LUCID Experiments, *Journal of Climate*, 25, 3261–3281, doi:10.1175/JCLI-D-11-00338.1, <http://journals.ametsoc.org/doi/abs/10.1175/JCLI-D-11-00338.1>, 2012.
- de Rigo, D., Caudullo, G., Busetto, L., and San Miguel, J.: Supporting EFSA assessment of the EU environmental suitability for exotic forestry pests : Final Report, Tech. rep., EFSA Supporting publications, 2014.
- de Rosnay, P.: Impact of a physically based soil water flow and soil-plant interaction representation for modeling large-scale land surface processes, *Journal of Geophysical Research*, 107, 4118, doi:10.1029/2001JD000634, <http://doi.wiley.com/10.1029/2001JD000634>, 2002.
- de Rosnay, P. and Polcher, J.: Modelling root water uptake in a complex land surface scheme coupled to a GCM, *Hydrology and Earth System Sciences*, 2, 239–255, doi:10.5194/hess-2-239-1998, <http://www.hydrol-earth-syst-sci.net/2/239/1998/>, 1998.
- Deleuze, C., Pain, O., Dhôte, J.-F., and Hervé, J.-C.: A flexible radial increment model for individual trees in pure even-aged stands, *Annals of Forest Science*, 61, 327–335, doi:10.1051/forest:2004026, <http://www.edpsciences.org/10.1051/forest:2004026>, 2004.
- Dickinson, R., Henderson-Sellers, A., Kennedy, P., and Wilson, M.: Biosphere-Atmosphere Transfer Scheme (BATS) for the NCAR Community Climate Model, Tech. Rep. December, 1986.
- Dixon, R. K., Solomon, A. M., Brown, S., Houghton, R. A., Trexler, M. C., and Wisniewski, J.: Carbon pools and flux of global forest ecosystems, *Science (New York, N.Y.)*, 263, 185–90, doi:10.1126/science.263.5144.185, <http://www.ncbi.nlm.nih.gov/pubmed/17839174>, 1994.
- D’Orgeval, T., Polcher, J., and de Rosnay, P.: Sensitivity of the West African hydrological cycle in ORCHIDEE to infiltration processes, *Hydrology and Earth System Sciences*, 12, 1387–1401, doi:10.5194/hess-12-1387-2008, <http://www.hydrol-earth-syst-sci.net/12/1387/2008/>, 2008.
- Ducoudré, N. I., Laval, K., and Perrier, A.: SECHIBA, a new set of parametrizations of the hydrologic exchanges at the land-atmosphere interface within the LMD atmospheric general circulation model, *Journal of Climate*, 6, 248–273, 1993.
- Dufresne, J. and Ghattas, J.: Description du schéma de la couche limite turbulente et l’ interface avec la surface planétaire dans LMDZ, 2009.

- 1365 Farquhar, G. D., von Caemmerer, S., and Berry, J. A.: A biochemical model of photosynthetic CO₂ assimilation in leaves of C₃ species, *Planta*, 149, 78–90, doi:10.1007/BF00386231, <http://www.ncbi.nlm.nih.gov/pubmed/24306196>, 1980.
- Farrell, E. P., Führer, E., Ryan, D., Andersson, F., Hüttel, R., and Piussi, P.: European forest ecosystems: building the future on the legacy of the past, *Forest Ecology and Management*, 132, 5–20, doi:10.1016/S0378-1127(00)00375-3, <http://linkinghub.elsevier.com/retrieve/pii/S0378112700003753>, 2000.
- 1370 Flexas, J., Bota, J., Galmés, J., Medrano, H., and Ribas-Carbó, M.: Keeping a positive carbon balance under adverse conditions: responses of photosynthesis and respiration to water stress, *Physiologia Plantarum*, 127, 343–352, doi:10.1111/j.1399-3054.2006.00621.x, <http://doi.wiley.com/10.1111/j.1399-3054.2006.00621.x>, 2006.
- 1375 Fortin, M., Ningre, F., Robert, N., and Mothe, F.: Quantifying the impact of forest management on the carbon balance of the forest-wood product chain: A case study applied to even-aged oak stands in France, *Forest Ecology and Management*, 279, 176–188, doi:10.1016/j.foreco.2012.05.031, <http://linkinghub.elsevier.com/retrieve/pii/S0378112712003088>, 2012.
- Friedlingstein, P., Joel, G., Field, C. B., and Fung, I. Y.: Toward an allocation scheme for global terrestrial carbon models, *Global Change Biology*, 5, 755–770, doi:10.1046/j.1365-2486.1999.00269.x, <http://doi.wiley.com/10.1046/j.1365-2486.1999.00269.x>, 1999.
- 1380 Gimmi, U., Poulter, B., Wolf, A., Portner, H., Weber, P., and Bürgi, M.: Soil carbon pools in Swiss forests show legacy effects from historic forest litter raking, *Landscape Ecology*, 28, 835–846, doi:10.1007/s10980-012-9778-4, <http://link.springer.com/10.1007/s10980-012-9778-4>, 2012.
- 1385 Gu, L.: Longwave radiative transfer in plant canopies, Ph.D. thesis, University of Virginia, 1988.
- Gu, L., Shugart, H. H., Fuentes, J. D., Black, T., and Shewchuk, S. R.: Micrometeorology, biophysical exchanges and NEE decomposition in a two-story boreal forest - development and test of an integrated model, *Agricultural and Forest Meteorology*, 94, 123–148, doi:10.1016/S0168-1923(99)00006-4, <http://linkinghub.elsevier.com/retrieve/pii/S0168192399000064>, 1999.
- 1390 Haverd, V., Lovell, J., Cuntz, M., Jupp, D., Newnham, G., and Sea, W.: The Canopy Semi-analytic Pgap And Radiative Transfer (CanSPART) model: Formulation and application, *Agricultural and Forest Meteorology*, 160, 14–35, doi:10.1016/j.agrformet.2012.01.018, <http://linkinghub.elsevier.com/retrieve/pii/S0168192312000494>, 2012.
- Henning, D.: Atlas of the surface heat balance of the continents, Gebrüder Bornträger, Berlin, Stuttgart, 1989.
- 1395 Hickler, T., Prentice, I. C., Smith, B., Sykes, M. T., and Zaehle, S.: Implementing plant hydraulic architecture within the LPJ Dynamic Global Vegetation Model, *Global Ecology and Biogeography*, 15, 060811081017001–???, doi:10.1111/j.1466-822X.2006.00254.x, <http://onlinelibrary.wiley.com/doi/10.1111/j.1466-822X.2006.00254.x/full><http://doi.wiley.com/10.1111/j.1466-822X.2006.00254.x>, 2006.
- Hourdin, F.: A new representation of the absorption by the CO₂ 15- μ m band for a Martian general circulation model, *Journal of Geophysical Research*, 97, 18 319, doi:10.1029/92JE01985, <http://doi.wiley.com/10.1029/92JE01985>, 1992.
- 1400 Jackson, R. B., Jobbágy, E. G., Avissar, R., Roy, S. B., Barrett, D. J., Cook, C. W., Farley, K. A., le Maitre, D. C., McCarl, B. A., and Murray, B. C.: Trading water for carbon with biological carbon sequestration., *Sci-*

- ence (New York, N.Y.), 310, 1944–7, doi:10.1126/science.1119282, <http://www.ncbi.nlm.nih.gov/pubmed/16373572>, 2005.
- Jiménez, C., Prigent, C., Mueller, B., Seneviratne, S. I., McCabe, M. F., Wood, E. F., Rossow, W. B., Balsamo, G., Betts, A. K., Dirmeyer, P. A., Fisher, J. B., Jung, M., Kanamitsu, M., Reichle, R. H., Reichstein, M., Rodell, M., Sheffield, J., Tu, K., and Wang, K.: Global intercomparison of 12 land surface heat flux estimates, *Journal of Geophysical Research*, 116, D02 102, doi:10.1029/2010JD014545, <http://doi.wiley.com/10.1029/2010JD014545>, 2011.
- Jung, M., Verstraete, M., Gobron, N., Reichstein, M., Papale, D., Bondeau, A., Robustelli, M., and Pinty, B.: Diagnostic assessment of European gross primary production, *Global Change Biology*, 14, 2349–2364, doi:10.1111/j.1365-2486.2008.01647.x, <http://doi.wiley.com/10.1111/j.1365-2486.2008.01647.x>, 2008.
- Kattge, J. and Knorr, W.: Temperature acclimation in a biochemical model of photosynthesis: a reanalysis of data from 36 species., *Plant, cell & environment*, 30, 1176–90, doi:10.1111/j.1365-3040.2007.01690.x, <http://www.ncbi.nlm.nih.gov/pubmed/17661754>, 2007.
- Kattge, J., Díaz, S., Lavorel, S., Prentice, I. C., Leadley, P., Bönsch, G., Garnier, E., Westoby, M., Reich, P. B., Wright, I. J., Cornelissen, J. H. C., Violle, C., Harrison, S. P., Van Bodegom, P. M., Reichstein, M., Enquist, B. J., Soudzilovskaia, N. a., Ackerly, D. D., Anand, M., Atkin, O., Bahn, M., Baker, T. R., Baldocchi, D., Bekker, R., Blanco, C. C., Blonder, B., Bond, W. J., Bradstock, R., Bunker, D. E., Casanoves, F., Cavender-Bares, J., Chambers, J. Q., Chapin III, F. S., Chave, J., Coomes, D., Cornwell, W. K., Craine, J. M., Dobrin, B. H., Duarte, L., Durka, W., Elser, J., Esser, G., Estiarte, M., Fagan, W. F., Fang, J., Fernández-Méndez, F., Fidelis, A., Finegan, B., Flores, O., Ford, H., Frank, D., Freschet, G. T., Fyllas, N. M., Gallagher, R. V., Green, W. A., Gutierrez, A. G., Hickler, T., Higgins, S. I., Hodgson, J. G., Jalili, A., Jansen, S., Joly, C. A., Kerkhoff, A. J., Kirkup, D., Kitajima, K., Kleyer, M., Klotz, S., Knops, J. M. H., Kramer, K., Kühn, I., Kurokawa, H., Laughlin, D., Lee, T. D., Leishman, M., Lens, F., Lenz, T., Lewis, S. L., Lloyd, J., Llusià, J., Louault, F., Ma, S., Mahecha, M. D., Manning, P., Massad, T., Medlyn, B. E., Messier, J., Moles, a. T., Müller, S. C., Nadrowski, K., Naeem, S., Niinemets, U., Nöllert, S., Nüske, A., Ogaya, R., Oleksyn, J., Onipchenko, V. G., Onoda, Y., Ordoñez, J., Overbeck, G., Ozinga, W. A., Patiño, S., Paula, S., Pausas, J. G., Peñuelas, J., Phillips, O. L., Pillar, V., Poorter, H., Poorter, L., Poschlod, P., Prinzing, A., Proulx, R., Rammig, A., Reinsch, S., Reu, B., Sack, L., Salgado-Negret, B., Sardans, J., Shiodera, S., Shipley, B., Siefert, A., Sosinski, E., Soussana, J.-F., Swaine, E., Swenson, N., Thompson, K., Thornton, P., Waldram, M., Weiher, E., White, M., White, S., Wright, S. J., Yguel, B., Zaehle, S., Zanne, a. E., and Wirth, C.: TRY - a global database of plant traits, *Global Change Biology*, 17, 2905–2935, doi:10.1111/j.1365-2486.2011.02451.x, <http://doi.wiley.com/10.1111/j.1365-2486.2011.02451.x>, 2011.
- Kira, T., Ogawa, H., and Sakazaki, N.: Intraspecific competition among higher plants. I. Competition-yield-density interrelationship in regularly dispersed populations, *Journal of the Institute of Polytechnics (Osaka University)*, 4, 1–16, 1953.
- Krinner, G., Nicolas, V., de Noblet-Ducoudre, N., Ogée, J., Polcher, J., Friedlingstein, P., Ciais, P., Sitch, S., and Prentice, I.: A dynamic global vegetation model for studies of the coupled atmosphere-biosphere system, *Global Biogeochemical Cycles*, 19, GB1015, doi:10.1029/2003GB002199, <http://www.agu.org/pubs/crossref/2005/2003GB002199.shtml>, 2005.

- Lardy, R., Bellocchi, G., and Soussana, J.-F.: A new method to determine soil organic carbon equilibrium, *Environmental Modelling & Software*, doi:10.1016/j.envsoft.2011.05.016, <http://linkinghub.elsevier.com/retrieve/pii/S1364815211001459>, 2011.
- 1445 Lovell, J., Haverd, V., Jupp, D., and Newnham, G.: The Canopy Semi-analytic Pgap And Radiative Transfer (CanSPART) model: Validation using ground based lidar, *Agricultural and Forest Meteorology*, 158–159, 1–12, doi:10.1016/j.agrformet.2012.01.020, <http://linkinghub.elsevier.com/retrieve/pii/S0168192312000512>, 2012.
- 1450 Luysaert, S., Inglima, I., Jung, M., Richardson, A. D., Reichstein, M., Papale, D., Piao, S. L., Schulze, E. D., Wingate, L., Matteucci, G., Aragao, L., Aubinet, M., Beer, C., Bernhofer, C., Black, K. G., Bonal, D., Bonnefond, J. M., Chambers, J., Ciais, P., Cook, B., Davis, K. J., Dolman, A. J., Gielen, B., Goulden, M., Grace, J., Granier, A., Grelle, A., Griffis, T., Grünwald, T., Guidolotti, G., Hanson, P. J., Harding, R., Hollinger, D. Y., Hutyrá, L. R., Kolari, P., Kruijt, B., Kutsch, W., Lagergren, F., Laurila, T., Law, B., Le Maire, G., Lindroth, A., Loustau, D., Malhi, Y., Mateus, J., Migliavacca, M., Misson, L., Montagnani, L., Moncrieff, J., Moors, E., Munger, J. W., Nikinmaa, E., Ollinger, S. V., Pita, G., Rebmann, C., Rouspard, O., Saigusa, N., Sanz, M. J., Seufert, G., Sierra, C., Smith, M. L., Tang, J., Valentini, R., Vesala, T., and Janssens, I. A.: CO₂ balance of boreal, temperate, and tropical forests derived from a global database, *Global Change Biology*, 13, 2509–2537, doi:10.1111/j.1365-2486.2007.01439.x, <http://doi.wiley.com/10.1111/j.1365-2486.2007.01439.x>, 2007.
- 1455 Luysaert, S., Hessenmöller, D., von Lüpke, N., Kaiser, S., and Schulze, E. D.: Quantifying land use and disturbance intensity in forestry, based on the self-thinning relationship, *Ecological Applications*, 21, 3272–3284, doi:10.1890/10-2395.1, <http://www.esajournals.org/doi/abs/10.1890/10-2395.1>, 2011.
- 1460 Luysaert, S., Jammot, M., Stoy, P. C., Estel, S., Pongratz, J., Ceschia, E., Churkina, G., Don, A., Erb, K., Ferlicoq, M., Gielen, B., Grünwald, T., Houghton, R. A., Klumpp, K., Knohl, A., Kolb, T., Kuemmerle, T., Laurila, T., Lohila, A., Loustau, D., McGrath, M. J., Meyfroidt, P., Moors, E. J., Naudts, K., Novick, K., Otto, J., Pilegaard, K., Pio, C. A., Rambal, S., Rebmann, C., Ryder, J., Suyker, A. E., Varlagin, A., Wattenbach, M., and Dolman, A. J.: Land management and land-cover change have impacts of similar magnitude on surface temperature, *Nature Climate Change*, 4, 389–393, doi:10.1038/nclimate2196, <http://www.nature.com/doi/abs/10.1038/nclimate2196>, 2014.
- 1465 MacBean, N., Maignan, F., Peylin, P., Bacour, C., and Ciais, P.: Using satellite data to improve the leaf phenology of a global Land Surface Model., In prep.
- Magnani, F., Mencuccini, M., and Grace, J.: Age-related decline in stand productivity : the role of structural acclimation under hydraulic constraints, *Plant, cell & environment*, 23, 251–263, 2000.
- Martin, M. P., Cordier, S., Balesdent, J., and Arrouays, D.: Periodic solutions for soil carbon dynamics equilibriums with time-varying forcing variables, *Ecological Modelling*, 204, 523–530, doi:10.1016/j.ecolmodel.2006.12.030, <http://linkinghub.elsevier.com/retrieve/pii/S0304380006006703>, 2007.
- 1475 Massman, W. and Weil, J.: An analytical one-dimensional second-order closure model of turbulence statistics and the Lagrangian time scale within and above plant canopies of arbitrary structure, *Boundary-layer meteorology*, 91, 81–107, 1999.
- 1480 McDowell, N., Barnard, H., Bond, B., Hinckley, T., Hubbard, R., Ishii, H., Köstner, B., Magnani, F., Marshall, J., Meinzer, F., Phillips, N., Ryan, M., and Whitehead, D.: The relationship between tree height and leaf

- area:sapwood area ratio, *Oecologia*, 132, 12–20, doi:10.1007/s00442-002-0904-x, <http://link.springer.com/10.1007/s00442-002-0904-x>, 2002.
- 1485 McGrath, M. J., Luyssaert, S., Meyfroidt, P., Kaplan, J. O., Buergi, M., Chen, Y., Erb, K., Gimmi, U., McInerney, D., Naudts, K., Otto, J., Pasztor, F., Ryder, J., Schelhaas, M.-J., and Valade, A.: Reconstructing European forest management from 1600 to 2010, In prep.a.
- McGrath, M. J., Pinty, B., Ryder, J., Otto, J., and Luyssaert, S.: A multilevel canopy radiative transfer scheme based on a domain-averaged structure factor, In prep.b.
- 1490 Meador, W. E. and Weaver, W. R.: Two-Stream Approximations to Radiative Transfer in Planetary Atmospheres: A Unified Description of Existing Methods and a New Improvement, *Journal of the Atmospheric Sciences*, 37, 630–643, doi:10.1175/1520-0469(1980)037<0630:TSATRT>2.0.CO;2, <http://journals.ametsoc.org/doi/abs/10.1175/1520-0469%281980%29037%3C0630%3ATSATRT%3E2.0.CO%3B2>, 1980.
- Medlyn, B. E., Dreyer, E., Ellsworth, D., Forstreuter, M., Harley, P. C., Kirschbaum, M. U. F., Le Roux, X., 1495 Montpied, P., Strassmeyer, J., Walcroft, A., Wang, K., and Loustau, D.: Temperature response of parameters of a biochemically based model of photosynthesis. II. A review of experimental data, *Plant, Cell and Environment*, 25, 1167–1179, doi:10.1046/j.1365-3040.2002.00891.x, <http://doi.wiley.com/10.1046/j.1365-3040.2002.00891.x>, 2002.
- Murphy, J. M., Sexton, D. M. H., Barnett, D. N., Jones, G. S., Webb, M. J., Collins, M., and Stainforth, D. A.: 1500 Quantification of modelling uncertainties in a large ensemble of climate change simulations., *Nature*, 430, 768–772, doi:10.1038/nature02771, <http://www.ncbi.nlm.nih.gov/pubmed/15306806>, 2004.
- Novick, K., Oren, R., Stoy, P., Juang, J.-Y., Siqueira, M., and Katul, G.: The relationship between reference canopy conductance and simplified hydraulic architecture, *Advances in Water Resources*, 32, 809–819, doi:10.1016/j.advwatres.2009.02.004, <http://linkinghub.elsevier.com/retrieve/pii/S0309170809000232>, 1505 2009.
- Ogee, J., Brunet, Y., Loustau, D., Berbigier, P., and Delzon, S.: MuSICA, a CO₂, water and energy multilayer, multileaf pine forest model: evaluation from hourly to yearly time scales and sensitivity analysis, *Global Change Biology*, 9, 697–717, doi:10.1046/j.1365-2486.2003.00628.x, <http://doi.wiley.com/10.1046/j.1365-2486.2003.00628.x>, 2003.
- 1510 Olson, J., Watts, J., and Allison, L.: Carbon in live vegetation of major world ecosystems, Tech. rep., Oak Ridge National Laboratory, ORNL-82, Oak Ridge TN, 1983.
- Otto, J., Berveiller, D., Bréon, F.-M., Delpierre, N., Geppert, G., Granier, A., Jans, W., Knohl, A., Kuusk, A., Longdoz, B., Moors, E., Mund, M., Pinty, B., Schelhaas, M.-J., and Luyssaert, S.: Forest summer albedo is sensitive to species and thinning: how should we account for this in Earth system models?, *Biogeosciences*, 11, 2411–2427, doi:10.5194/bg-11-2411-2014, <http://www.biogeosciences.net/11/2411/2014/>, 2014.
- 1515 Pan, Y., Birdsey, R. a., Fang, J., Houghton, R., Kauppi, P. E., Kurz, W. a., Phillips, O. L., Shvidenko, A., Lewis, S. L., Canadell, J. G., Ciais, P., Jackson, R. B., Pacala, S. W., McGuire, a. D., Piao, S., Rautiainen, A., Sitch, S., and Hayes, D.: A large and persistent carbon sink in the world's forests., *Science (New York, N.Y.)*, 333, 988–993, doi:10.1126/science.1201609, <http://www.ncbi.nlm.nih.gov/pubmed/21764754>, 2011.
- 1520 Parton, W. J., Stewart, J. W. B., and Cole, C. V.: Dynamics of C, N, P and S in grassland soils: a model, *Biogeochemistry*, 5, 109–131, doi:10.1007/BF02180320, <http://link.springer.com/10.1007/BF02180320>, 1988.

Pataki, D. E., Alig, R. J., Fung, A. S., Golubiewski, N. E., Kennedy, C. A., Mcpherson, E. G., Nowak, D. J., Pouyat, R. V., and Romero Lankao, P.: Urban ecosystems and the North American carbon cycle, *Global Change Biology*, 12, 2092–2102, doi:10.1111/j.1365-2486.2006.01242.x, <http://doi.wiley.com/10.1111/j.1365-2486.2006.01242.x>, 2006.

1525 Pielke, R. A., Marland, G., Betts, R. A., Chase, T. N., Eastman, J. L., Niles, J. O., Niyogi, D. D. S., and Running, S. W.: The influence of land-use change and landscape dynamics on the climate system: relevance to climate-change policy beyond the radiative effect of greenhouse gases, *Philosophical Transactions of the Royal Society A: Mathematical, Physical and Engineering Sciences*, 360, 1705–1719, doi:10.1098/rsta.2002.1027, <http://rsta.royalsocietypublishing.org/cgi/doi/10.1098/rsta.2002.1027>, 2002.

1530 Pinty, B.: Synergy between 1-D and 3-D radiation transfer models to retrieve vegetation canopy properties from remote sensing data, *Journal of Geophysical Research*, 109, D21 205, doi:10.1029/2004JD005214, <http://doi.wiley.com/10.1029/2004JD005214>, 2004.

Pinty, B., Lavergne, T., Dickinson, R. E., Widlowski, J.-L., Gobron, N., and Verstraete, M. M.: Simplifying the interaction of land surfaces with radiation for relating remote sensing products to climate models, *Journal of Geophysical Research*, 111, D02 116, doi:10.1029/2005JD005952, <http://www.agu.org/pubs/crossref/2006/2005JD005952.shtml>, 2006.

1535 Pinty, B., Lavergne, T., Voß beck, M., Kaminski, T., Aussedat, O., Giering, R., Gobron, N., Taberner, M., Verstraete, M. M., and Widlowski, J.-L.: Retrieving surface parameters for climate models from Moderate Resolution Imaging Spectroradiometer (MODIS)-Multiangle Imaging Spectroradiometer (MISR) albedo products, *Journal of Geophysical Research*, 112, D10 116, doi:10.1029/2006JD008105, <http://doi.wiley.com/10.1029/2006JD008105>, 2007.

1540 Pinty, B., Andredakis, I., Clerici, M., Kaminski, T., Taberner, M., Verstraete, M. M., Gobron, N., Plummer, S., and Widlowski, J.-L.: Exploiting the MODIS albedos with the Two-stream Inversion Package (JRC-TIP): 1. Effective leaf area index, vegetation, and soil properties, *Journal of Geophysical Research*, 116, D09 105, doi:10.1029/2010JD015372, <http://doi.wiley.com/10.1029/2010JD015372>, 2011a.

1545 Pinty, B., Clerici, M., Andredakis, I., Kaminski, T., Taberner, M., Verstraete, M. M., Gobron, N., Plummer, S., and Widlowski, J.-L.: Exploiting the MODIS albedos with the Two-stream Inversion Package (JRC-TIP): 2. Fractions of transmitted and absorbed fluxes in the vegetation and soil layers, *Journal of Geophysical Research*, 116, D09 106, doi:10.1029/2010JD015373, <http://doi.wiley.com/10.1029/2010JD015373>, 2011b.

1550 Pinty, B., Jung, M., Kaminski, T., Lavergne, T., Mund, M., Plummer, S., Thomas, E., and Widlowski, J.-L.: Evaluation of the JRC-TIP 0.01° products over a mid-latitude deciduous forest site, *Remote Sensing of Environment*, 115, 3567–3581, doi:10.1016/j.rse.2011.08.018, <http://linkinghub.elsevier.com/retrieve/pii/S0034425711003221>, 2011c.

1555 Pitman, A. J., de Noblet-Ducoudré, N., Cruz, F. T., Davin, E. L., Bonan, G. B., Brovkin, V., Claussen, M., Delire, C., Ganzeveld, L., Gayler, V., van den Hurk, B. J. J. M., Lawrence, P. J., van der Molen, M. K., Müller, C., Reick, C. H., Seneviratne, S. I., Strengers, B. J., and Voldoire, A.: Uncertainties in climate responses to past land cover change: First results from the LUCID intercomparison study, *Geophysical Research Letters*, 36, L14 814, doi:10.1029/2009GL039076, <http://www.agu.org/pubs/crossref/2009/2009GL039076.shtml>, 2009.

1560 Polcher, J., McAvaney, B., Viterbo, P., Gaertner, M.-A., Hahmann, A., Mahfouf, J.-F., Noilhan, J., Phillips, T., Pitman, A., Schlosser, C., Schulz, J.-P., Timbal, B., Verseghy, D., and Xue, Y.: A proposal for a general inter-

- face between land surface schemes and general circulation models, *Global and Planetary Change*, 19, 261–276, doi:10.1016/S0921-8181(98)00052-6, <http://linkinghub.elsevier.com/retrieve/pii/S0921818198000526>, 1998.
- 1565 Poulter, B., Ciais, P., Hodson, E., Lischke, H., Maignan, F., Plummer, S., and Zimmermann, N. E.: Plant functional type mapping for earth system models, *Geoscientific Model Development*, 4, 993–1010, doi:10.5194/gmd-4-993-2011, <http://www.geosci-model-dev.net/4/993/2011/>, 2011.
- Poulter, B., MacBean, N., Hartley, A., Khlystova, I., Betts, R., Bontemps, S., Brockmann, C., Defournay, P., Hagemann, S., Herold, M., Kirches, G., Lamarche, C., Lederer, D., and Peylin, P.: Plant functional type
1570 classification for Earth System Models: Results from the European Space Agency’s Land Cover Climate Change Initiative, 2015.
- Pretzsch: *Forest dynamics, growth and yield*, Springer-Verlag, Berlin, 2009.
- Pretzsch, H. and Dieler, J.: Evidence of variant intra- and interspecific scaling of tree crown structure and relevance for allometric theory, *Oecologia*, 169, 637–49, doi:10.1007/s00442-011-2240-5, [http://www.](http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=3375085&tool=pmcentrez&rendertype=abstract)
1575 [pubmedcentral.nih.gov/articlerender.fcgi?artid=3375085&tool=pmcentrez&rendertype=abstract](http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=3375085&tool=pmcentrez&rendertype=abstract), 2012.
- Reick, C. H., Raddatz, T., Brovkin, V., and Gayler, V.: Representation of natural and anthropogenic land cover change in MPI-ESM, *Journal of Advances in Modeling Earth Systems*, 5, 459–482, doi:10.1002/jame.20022, <http://doi.wiley.com/10.1002/jame.20022>, 2013.
- Reineke, L.: Perfecting a stand-density index for even-aged forests, *Journal of Agricultural Research*, 46, 627–
1580 638, 1933.
- Richards, L. A.: CAPILLARY CONDUCTION OF LIQUIDS THROUGH POROUS MEDIUMS, *Physics*, 1, 318, doi:10.1063/1.1745010, <http://scitation.aip.org/content/aip/journal/jap/1/5/10.1063/1.1745010>, 1931.
- Ruimy, A., Dedieu, G., and Saugier, B.: TURC: A diagnostic model of continental gross primary productivity and net primary productivity, *Global Biogeochemical Cycles*, 10, 269–285, doi:10.1029/96GB00349, <http://doi.wiley.com/10.1029/96GB00349>, 1996.
1585
- Ryan, M.: The effects of climate change on plant respiration, *Ecological Applications*, 1, 157–167, 1991.
- Ryder, J., Polcher, J., Peylin, P., Ottlé, C., Chen, Y., van Gorsel, E., Haverd, V., McGrath, M. J., Naudts, K., Otto, J., Valade, A., and Luyssaert, S.: A multi-layer land surface energy budget model for implicit coupling with global atmospheric simulations, *Geoscientific Model Development Discussions*, 7, 8649–8701, doi:10.5194/gmdd-7-8649-2014, <http://www.geosci-model-dev-discuss.net/7/8649/2014/>, 2014.
1590
- Santaren, D., Peylin, P., Viovy, N., and Ciais, P.: Optimizing a process-based ecosystem model with eddy-covariance flux measurements: A pine forest in southern France, *Global Biogeochemical Cycles*, 21, n/a–n/a, doi:10.1029/2006GB002834, <http://doi.wiley.com/10.1029/2006GB002834>, 2007.
- Schaaf, C. B., Gao, F., Strahler, A. H., Lucht, W., Li, X., Tsang, T., Strugnell, N. C., Zhang, X., Jin, Y., Muller, J.-P., Lewis, P., Barnsley, M., Hobson, P., Disney, M., Roberts, G., Dunderdale, M., Doll, C., D’Entremont, R. P., Hu, B., Liang, S., Privette, J. L., and Roy, D.: First operational BRDF, albedo nadir reflectance products from MODIS, *Remote Sensing of Environment*, 83, 135–148, doi:10.1016/S0034-4257(02)00091-3, <http://linkinghub.elsevier.com/retrieve/pii/S0034425702000913>, 2002.
- Schall, P. and Ammer, C.: Can land use intensity be reliably quantified by using a single self-thinning relationship?, *Ecological Applications*, 23, 675–677, doi:10.1890/12-0847.1, [http://www.esajournals.org/doi/abs/10.](http://www.esajournals.org/doi/abs/10.1890/12-0847.1)
1600 [1890/12-0847.1](http://www.esajournals.org/doi/abs/10.1890/12-0847.1), 2013.

- Scheiter, S., Langan, L., and Higgins, S. I.: Next-generation dynamic global vegetation models: learning from community ecology, *The New Phytologist*, 198, 957–69, doi:10.1111/nph.12210, <http://www.ncbi.nlm.nih.gov/pubmed/23496172>, 2013.
- 1605 Shinozaki, K., Yoda, K., Hozumi, K., and Kira, T.: A quantitative analysis of plant form-the pipe model theory: I. Basic analyses, *Japanese Journal of Ecology*, 14, 97–105, 1964.
- Simard, M., Pinto, N., Fisher, J. B., and Baccini, A.: Mapping forest canopy height globally with spaceborne lidar, *Journal of Geophysical Research*, 116, G04 021, doi:10.1029/2011JG001708, <http://doi.wiley.com/10.1029/2011JG001708>, 2011.
- 1610 Simonin, K., Kolb, T. E., Montes-Helu, M., and Koch, G. W.: Restoration thinning and influence of tree size and leaf area to sapwood area ratio on water relations of *Pinus ponderosa*, *Tree Physiology*, 26, 493–503, doi:10.1093/treephys/26.4.493, <http://treephys.oxfordjournals.org/cgi/doi/10.1093/treephys/26.4.493>, 2006.
- Sitch, S., Smith, B., Prentice, I. C., Arneth, A., Bondeau, A., Cramer, W., Kaplan, J. O., Levis, S., Lucht, W., Sykes, M. T., Thonicke, K., and Venevsky, S.: Evaluation of ecosystem dynamics, plant geography and terrestrial carbon cycling in the LPJ dynamic global vegetation model, *Global Change Biology*, 9, 161–185, doi:10.1046/j.1365-2486.2003.00569.x, <http://doi.wiley.com/10.1046/j.1365-2486.2003.00569.x>, 2003.
- 1615 Slatyer, R.: *Plant-Water Relationships*, vol. 158, Academic Press, New York, 1967.
- Sperry, J. S., Adler, F. R., Campbell, G. S., and Comstock, J. P.: Limitation of plant water use by rhizosphere and xylem conductance: results from a model, *Plant, Cell and Environment*, 21, 347–359, doi:10.1046/j.1365-3040.1998.00287.x, <http://doi.wiley.com/10.1046/j.1365-3040.1998.00287.x>, 1998.
- 1620 Steppe, K., De Pauw, D. J. W., Lemeur, R., and Vanrolleghem, P. A.: A mathematical model linking tree sap flow dynamics to daily stem diameter fluctuations and radial stem growth, *Tree Physiology*, 26, 257–273, doi:10.1093/treephys/26.3.257, <http://treephys.oxfordjournals.org/cgi/doi/10.1093/treephys/26.3.257>, 2006.
- Stöckli, R., Lawrence, D. M., Niu, G.-Y., Oleson, K. W., Thornton, P. E., Yang, Z.-L., Bonan, G. B., Denning, A. S., and Running, S. W.: Use of FLUXNET in the Community Land Model development, *Journal of Geophysical Research*, 113, G01 025, doi:10.1029/2007JG000562, <http://doi.wiley.com/10.1029/2007JG000562>, 2008.
- 1625 Sun, Y., Gu, L., Dickinson, R. E., Norby, R. J., Pallardy, S. G., and Hoffman, F. M.: Impact of mesophyll diffusion on estimated global land CO₂ fertilization, *Proceedings of the National Academy of Sciences*, doi:10.1073/pnas.1418075111, <http://www.pnas.org/cgi/doi/10.1073/pnas.1418075111>, 2014.
- 1630 Tarantola, A.: *Inverse Problem Theory and Methods for Model Parameter Estimation*, SIAM, Philadelphia, 2005.
- Thornton, P. E. and Rosenbloom, N. A.: Ecosystem model spin-up: Estimating steady state conditions in a coupled terrestrial carbon and nitrogen cycle model, *Ecological Modelling*, 189, 25–48, doi:10.1016/j.ecolmodel.2005.04.008, <http://linkinghub.elsevier.com/retrieve/pii/S0304380005001948>, 2005.
- 1635 Tyree, M. T. and Sperry, J. S.: Vulnerability of xylem to cavitation and embolism, *Annual Review of Plant Physiology and Molecular Biology*, 40, 19–38, 1989.
- Van Genuchten, M.: A Closed-form Equation for Predicting the Hydraulic Conductivity of Unsaturated Soils, *Soil Science Society of America*, 44, 892–898, 1980.
- 1640 Verhoef, A. and Egea, G.: Modeling plant transpiration under limited soil water: Comparison of different plant and soil hydraulic parameterizations and preliminary implications for their use in land surface models,

Agricultural and Forest Meteorology, 191, 22–32, doi:10.1016/j.agrformet.2014.02.009, <http://linkinghub.elsevier.com/retrieve/pii/S0168192314000483>, 2014.

- 1645 Vicca, S., Luyssaert, S., Peñuelas, J., Campioli, M., Chapin, F. S., Ciais, P., Heinemeyer, A., Högberg, P., Kutsch, W. L., Law, B. E., Malhi, Y., Papale, D., Piao, S. L., Reichstein, M., Schulze, E. D., and Janssens, I.: Fertile forests produce biomass more efficiently., *Ecology letters*, 15, 520–6, doi:10.1111/j.1461-0248.2012.01775.x, <http://www.ncbi.nlm.nih.gov/pubmed/22472207>, 2012.
- Viovy, N. and de Noblet-Ducoudré, N.: Coupling water and carbon cycle in the biosphere, *Sci. Geol. Bull.*, 50, 109–121, 1997.
- 1650 Weatherley, P. E.: Water Uptake and Flow in Roots, in: *Physiological Plant Ecology II*, pp. 79–109, Springer Berlin Heidelberg, doi:10.1007/978-3-642-68150-9_4, 1982.
- Whitehead, D.: Regulation of stomatal conductance and transpiration in forest canopies, *Tree Physiology*, 18, 633–644, doi:10.1093/treephys/18.8-9.633, <http://treephys.oxfordjournals.org/cgi/doi/10.1093/treephys/18.8-9.633>, 1998.
- 1655 Widlowski, J.-L., Pinty, B., Clerici, M., Dai, Y., De Kauwe, M., de Ridder, K., Kallel, A., Kobayashi, H., Lavergne, T., Ni-Meister, W., Olchev, A., Quaife, T., Wang, S., Yang, W., Yang, Y., and Yuan, H.: RAMI4PILPS: An intercomparison of formulations for the partitioning of solar radiation in land surface models, *Journal of Geophysical Research*, 116, G02 019, doi:10.1029/2010JG001511, <http://doi.wiley.com/10.1029/2010JG001511>, 2011.
- 1660 Wolf, A., Ciais, P., Bellassen, V., Delbart, N., Field, C. B., and Berry, J. A.: Forest biomass allometry in global land surface models, *Global Biogeochemical Cycles*, 25, n/a–n/a, doi:10.1029/2010GB003917, <http://doi.wiley.com/10.1029/2010GB003917>, 2011.
- Xia, J. Y., Luo, Y. Q., Wang, Y.-P., Weng, E. S., and Hararuk, O.: A semi-analytical solution to accelerate spin-up of a coupled carbon and nitrogen land model to steady state, *Geoscientific Model Development*, 5, 1259–1271, doi:10.5194/gmd-5-1259-2012, <http://www.geosci-model-dev.net/5/1259/2012/>, 2012.
- 1665 Yang, Z., Dickinson, R., Robock, A., and Vinnikov, K.: Validation of the snow submodel of the biosphere – atmosphere transfer scheme with Russian snow cover and meteorological observational data, *Journal of climate*, 10, 353–373, 1997.
- Yin, X. and Struik, P.: C3 and C4 photosynthesis models: An overview from the perspective of crop modelling, *NJAS - Wageningen Journal of Life Sciences*, 57, 27–38, doi:10.1016/j.njas.2009.07.001, <http://linkinghub.elsevier.com/retrieve/pii/S1573521409000025>, 2009.
- Yoda, K., Kira, T., Ogawa, H., and Hozumi, K.: Self-thinning in overcrowded pure stands under cultivated and natural conditions, *Journal of the Institute of Polytechnics (Osaka University)*, 14, 107–129, 1963.
- Zahle, S. and Friend, A. D.: Carbon and nitrogen cycle dynamics in the O-CN land surface model: 1. Model description, site-scale evaluation, and sensitivity to parameter estimates, *Global Biogeochemical Cycles*, 24, n/a–n/a, doi:10.1029/2009GB003521, <http://doi.wiley.com/10.1029/2009GB003521>, 2010.
- 1675 Zhao, K. and Jackson, R. B.: Biophysical forcings of land-use changes from potential forestry activities in North America, *Ecological Monographs*, 84, 329–353, doi:10.1890/12-1705.1, <http://www.esajournals.org/doi/abs/10.1890/12-1705.1>, 2014.




Figure 1. Schematic overview of the changes in ORCHIDEE-CAN. For the trunk the most important processes and connections are indicated in black, while the processes and connections that were added or changed in ORCHIDEE-CAN are indicated in red. Numbered arrows are discussed in section 2.2.




Figure 2. Root mean square error of ORCHIDEE-CAN for gross primary production, evapotranspiration, visible and near-infra-red albedo, effective leaf area index, basal area and height for different regions and periods (DJF: December-February, MAM: March-May, JJA: June-August, SON: September-November). The gray-scale of the symbols indicates the number of pixels included in the calculation. The transition from green to white indicates an RMSE of 100%.




Figure 3. Root mean square error of ORCHIDEE-trunk for gross primary production, evapotranspiration and visible and near-infra-red albedo for different regions and periods (DJF: December-February, MAM: March-May, JJA: June-August, SON: September-November). The gray-scale of the symbols indicates the number of pixels included in the calculation. The transition from green to white indicates an RMSE of 100%.




Figure 4. Comparison between observations and simulations of ORCHIDEE-CAN for gross primary production and basal area over Europe. Gross primary production represents the mean for June-August between 2001-2010 and basal area is the value at the end of 2010.




Figure 5. Impact of the different forest management strategies on an oak forest for unmanged (green), high stand (orange) and coppice (blue) compared to a Poplar short rotation coppicing (red) at 48°N , 2°E . The simulation was run without spin-up to better visualize carbon build up in the coarse woody debris (C.W.D.) pool. The simulation cycled of a single year (1990) of climate data to minimize the inter annual variability due to climatic year to year variability




Figure 6. Relative root square error (RMSE) of tree diameter for different species (shown as different markers) for different regions over France (shown as A to K). Open triangle, *Pinus sylvestris*; open circle, *Pinus pinaster*; open square, *Picea Sp.*; filled diamond, *Quercus ilex/suber*; filled triangle, *Betula Sp.*; filled circle, *Fagus sylvatica*; filled square, *Quercus robur/petraea*.

Table 1. Concise description of the modules in the standard ORCHIDEE version with the motivation to change the modules in ORCHIDEE-CAN.

Module	Description	Motivation for change
Albedo	For each PFT the total albedo for the grid square is computed as a weighted average of the vegetation albedo, the soil albedo, and the snow albedo.	The scheme overlooks the effect of vegetation shading bare soil for sparse canopies and gives the ground in all PFTs the same reflectance properties as bare soil.
Soil hydrology	Vertical water flow in the soil is based on the Fokker–Planck equation that resolves water diffusion in non-saturated conditions from the Richards equation (Richards, 1931). The 2 m soil column consists of eleven moisture layers with an exponentially increasing depth (D’Orgeval et al., 2008).	No change
Soil temperature	The soil temperature is computed according to the Fourier equation using a finite difference implicit scheme with seven numerical nodes unevenly distributed between 0 and 5.5 m (Hourdin, 1992).	No change
Energy budget	The coupled energy balance scheme, and its exchange with the atmosphere, is based on that of Dufresne and Ghattas (2009). The surface is described as a single layer that includes both the soil surface and any vegetation.	A big leaf approach does not account for within canopy transport of carbon, water and energy. Further, it is inconsistent with the current multi-layer photosynthesis approach and the new multi-layer albedo approach.
Photosynthesis	C3 and C4 photosynthesis is calculated following Farquhar et al. (1980) and Collatz et al. (1992), respectively. Photosynthesis assigns artificial LAI levels to calculate the carbon assimilation of the canopy. These levels allow for a saturation of photosynthesis with LAI, but have no physical meaning.	The scheme uses a simple Beer’s law transmission of light to each level, which is inconsistent with the new albedo scheme.
Autotrophic respiration	Autotrophic respiration distinguishes maintenance and growth respiration. Maintenance respiration occurs in living plant compartments and is a function of temperature, biomass and, the prescribed carbon/nitrogen ratio of each tissue (Ruimy et al., 1996). A prescribed fraction of 28% of the photosynthates allocated to growth is used in growth respiration (?). The remaining assimilates are distributed among the various plant organs using an allocation scheme based on resource limitations (see allocation).	No change

Table 1. Continuation of Table 1.

Module	Description	Motivation for change
Carbon allocation	Carbon is allocated to the plant following resource limitations Friedlingstein et al. (1999). Plants allocate carbon to their different tissues in response to external limitations of water, light and nitrogen availability. When the ratios of these limitations are out of bounds, prescribed allocation factors are used.	The resource limitation approach requires capping LAI at a predefined value. Due to this cap, the allocation rules are most often not applied, reducing the scheme to prescribing allocation.
Phenology	At the end of each day, the model checks whether the conditions for leaf onset are satisfied. The PFT-specific conditions are based on long and short term warmth and/or moisture conditions (Botta et al., 2000).	No change
Mortality and turnover	All biomass pools have a turnover time. Living biomass is transferred to the litter pool, litter is decomposed or transferred to the soil pool.	This approach is not capable of modelling stand dimensions.
Soil and litter carbon and heterotrophic respiration	Following (Parton et al., 1988), prescribed fractions of the different plant components go to the metabolic and structural litter pools following senescence, turnover or mortality. The decay of metabolic and structural litter is controlled by temperature and soil or litter humidity. For structural litter, its lignin content also influences the decay rate.	No change
Forest management	An explicit distribution of individual trees (Bellassen et al., 2010) is the basis for a process-based simulation of mortality. The aboveground 'stand-scale' wood increment is distributed on a yearly time step among individual trees according to the rule of (Deleuze et al., 2004): the basal area of each individual tree grows proportionally to its circumference.	The concept of the original implementation were retained, however, the implementation was adjusted for consistency with the new allocation scheme and to have a larger diversity of management strategies.

Table 2. Variable description. Variables were grouped as follows: F =flux, f =fraction, M =pool, m =modulator, d =stand dimension, T =temperature, p =pressure, R =resistance, q =humidity, g =function.

Symbol in text	Unit	Symbol in ORCHIDEE-CAN	Description
F_{rm}	$\text{gC m}^{-2}\text{s}^{-1}$	resp_maint	Maintenance respiration
F_{rg}	$\text{gC m}^{-2}\text{s}^{-1}$	resp_growth	Growth respiration
$F_{LW,i}$	W m^2	r_lw	Long wave radiation incident at vegetation level "i"
$F_{SW,i}$	W m^2	r_sw	Short wave radiation incident at vegetation level "i"
$F_{Tr,s}$	m s^{-1}	Transpir_supply	Amount of water that a tree can get up from the soil to its leaves for transpiration
$T_{a,i}$	K	temp_atmos_pres,	Atmospheric temperature at the 'present' and 'next' time step, respectively, at level "i"
		temp_atmos_next	
$T_{L,i}$	K	temp_leaf_pres	Leaf temperature at level "i"
$q_{a,i}$	kg kg^{-1}	q_atmos_pres, q_atmos_next	Specific humidity at the "present" and "next" time step, respectively, at level "i"
$q_{L,i}$	kg kg^{-1}	q_leaf_pres	Leaf specific humidity at level "i"
M_l	gC plant^{-1}	Cl	Leaf mass of an individual plant
M_s	gC plant^{-1}	Cs	Sapwood mass of an individual plant
M_h	gC plant^{-1}	Ch	Heartwood mass of an individual plant
M_r	gC plant^{-1}	Cr	Root mass of an individual plant
M_{linc}	gC plant^{-1}	Cl_inc	Increment in leaf mass of an individual plant
M_{sinc}	gC plant^{-1}	Cs_inc	Increment in sapwood mass of an individual plant
M_{rinc}	gC plant^{-1}	Cr_inc	Increment in root mass of an individual plant
M_{totinc}	gC	b_inc_tot	Total biomass increment
M_{inc}	gC plant^{-1}	b_inc	Increment in plant biomass of an individual plant
M_{swc}	$\text{m}^3 \text{m}^{-3}$	swc	Volumetric soil water content
m_w	-	wstress_fac	Modulator for water stress as experienced by the plants
m_ψ	MPa	psi_soil_tune	Modulator to account for resistance in the soil-root interface
m_{Ndeath}	-	scale_factor	Normalization factor for mortality
$m_{LAIcorr}$	-	lai_correction_factor	Adjustable parameter in the calculation of gap probabilities of grasses and crops
d_h	m	height	Plant height
d_l	m^{-2}	-	One sided leaf area of an individual plant

Table 2. Continuation of Table 2

Symbol in text	Unit	Symbol in ORCHIDEE-CAN	Description
d_s	m^{-2}	-	Sapwood area of an individual plant
d_{hinc}	m	delta_height	Height increment
d_{dbh}	m	dia	Plant diameter
d_{ba}	$m^2 \text{ plant}^{-1}$	ba	Basal area
d_{bainc}	$m^2 \text{ plant}^{-1}$	delta_ba	Basal area increment
d_{circ}	m	circ	Stem circumference of an individual plant
d_{ind}	trees	n_circ_class	Number of trees in diameter class l
d_c	m^2	crown_shadow_h	Projected area of an opaque tree crown
d_{csa}	m^2	csa_sap	Projected crown surface area
d_{LAI}	$m_{leaf}^2 m_{ground}^{-2}$	-	Leaf area index
d_{LAIeff}	-	laieff	Effective leaf area index
$d_{LAIabove}$	-	lai_sum	Sum of the LAI of all levels above the current level
$d_{A,i}$	m^2	-	Cross-sectional area of vegetation level "i"
$d_{hl,i}$	m	delta_h	Vegetation height of level "i"
$d_{V,i}$	m^3	-	Volume of vegetation level "i"
d_{rd}	-	root_dens	Root density
d_λ	$ind \text{ m}^2$	-	Inverse of the individual plant density
p_{delta}	MPa	delta_P	Pressure difference between leaves and soil
$p_{\psi sr}$	MPa	psi_soilroot	Bulk soil water potential in the rooting zone
$p_{\psi s}$	MPa	psi_soil	Soil water potential for each soil layer
R_r	$MPa \text{ s m}^{-3}$	R_root	Hydraulic resistance of roots
R_{sap}	$MPa \text{ s m}^{-3}$	R_sap	Hydraulic resistance of sapwood
R_l	$MPa \text{ s m}^{-3}$	R_leaf	Hydraulic resistance of leaves
R_{temp}	$MPa \text{ s m}^{-3}$	-	Hydraulic resistance of roots, sapwood or leaves adjusted for temperature
$R_{a,i}$	$s \text{ m}^{-1}$	big_r	Aerodynamic resistance of vegetation at level "i" in the canopy
$R_{s,i}$	$s \text{ m}^{-1}$	big_r_prime	Stomatal resistance of vegetation at level "i" in the canopy

Table 2. Continuation of Table 2

Symbol in text	Unit	Symbol in ORCHIDEE-CAN	Description
f_{Pwc}	-	Pwc_h	Porosity of a tree crown
f_{Pgap}^{trees}	-	PgapL	Gap probability for trees
f_{Pgap}^{gc}	-	PgapL	Gap probability for grasses and crops
f_{Pgap}^{bs}	-	PgapL	Gap probability for bare soil
f_{death}^{ieir}	-	mortality	Mortality fraction per circumference class
f_{KF}	-	KF	Leaf allocation factor
f_{LF}	-	LF	Root allocation factor
f_{γ}	-	gamma	Slope of the intra-specific competition
f_s	m	s	Slope of linearised relationship between height and basal area
f_{rl}	-	leaf_reflectance	Reflectance of a single leaf
f_{tl}	-	leaf_transmittance	Transmittance of a single leaf
f_{rbgd}	-	bdg_reflectance	Reflectance of the ground beneath the canopy
$f_{Coll,veg}^{fR}$	-	Collim_alb_BB,	Reflected fraction of light to the atmosphere which has collided with canopy elements, separated for direct and diffuse sources, respectively
$f_{UnColl,bgd}^{fR}$	-	Isotrop_alb_BB	Reflected fraction of light to the atmosphere which has not collided with any canopy elements, separated for direct and diffuse sources, respectively
$f_{UnColl,veg}^T$	-	Collim_alb_BC,	Transmitted fraction of light to the ground which has not collided with any canopy elements
$f_{Coll,bgd,1}^{fR}$	-	Isotrop_alb_BC	Reflected fraction of light which has struck the background a single time and has collided with vegetation
$f_{Coll,bgd,n}^{fR}$	-	Collim_Tran_Uncoll	Reflected fraction of light which has struck the background multiple times and has collided with vegetation
z	m	z_array	Height above the soil
θ_z	radians	solar_angle	Solar zenith angle
θ_{μ}	radians	-	Cosine of the solar zenith angle
gG	-	-	Leaf orientation function
$g\sigma$	-	sigmas	Cut-off circumference of the intra-specific competition, calculated as a function of $k_{p,circ}$

Table 3. Likelihood that the simulated variable comes from the same population as the data. The ORCHIDEE-trunk version does not include effective LAI, basal area and height. Note that the likelihood of Europe cannot be derived from the values of the other regions due to the overlap between regions.

	ORCHIDEE-CAN							ORCHIDEE-TRUNK						
	GPP	EVAPO	ALB_NIR	ALB_VIS	EFFLAI	BA	HEIGHT	GPP	EVAPO	ALBEDO	EFFLAI	BA	HEIGHT	
British Isles	0.91	0.87	0.78	0.45	0.55	0.47	0.13	0.91	0.49	0.74	0.04	-	-	
Iberian Peninsula	0.80	0.80	0.73	0.65	0.60	0.09	0.66	0.65	0.37	0.25	0.04	-	-	
France	0.86	0.90	0.92	0.46	0.60	0.66	0.60	0.69	0.46	0.75	0.02	-	-	
Mid-Europe	0.92	0.93	0.88	0.86	0.68	0.80	0.76	0.81	0.48	0.64	0.46	-	-	
Scandinavia	0.92	0.83	0.47	0.91	0.59	0.62	0.24	0.81	0.31	0.55	0.65	-	-	
Alps	0.92	0.86	0.46	0.83	0.68	0.80	0.47	0.77	0.52	0.25	0.52	-	-	
Mediterranean	0.84	0.77	0.77	0.80	0.65	0.51	0.72	0.54	0.45	0.43	0.45	-	-	
Eastern Europe	0.93	0.94	0.70	0.93	0.73	0.71	0.76	0.84	0.52	0.51	0.75	-	-	
Europe	0.91	0.87	0.71	0.92	0.67	0.72	0.68	0.79	0.45	0.61	0.69	-	-	