Reply to the referees:

Referee Nr. 2: "Dear authors:

I acknowledged that you made a lot of efforts to describe the model in a consistent manner and to validate the model. I appreciated the addition of Figures 4, 5, and 6 for validation. However, Figure 6 shows relative RMSE values, making it difficult for me to judge whether the model worked well or not. Can you show more straightforward data (e.g., absolute RMSE) for clearer validation?"

We have changed figure 6 according to the suggestion of the reviewer: it now shows the absolute RMSE. We also made some changes in the text related to the figure (Line 1180-1184).

Manuscript prepared for Geosci. Model Dev. with version 2014/07/29 7.12 Copernicus papers of the LATEX class copernicus.cls. Date: 4 May 2015

A vertically discretised canopy description for ORCHIDEE (SVN r2290) and the modifications to the energy, water and carbon fluxes

Kim Naudts^{1,*}, James Ryder^{1,*}, Matthew J. McGrath^{1,*}, Juliane Otto^{1,10,*}, Yiying Chen¹, Aude Valade¹, Valentin Bellasen², Gonzalo Berhongaray³, Gerhard Bönisch⁴, Matteo Campioli³, Josefine Ghattas¹, Toon De Groote^{3,11}, Vanessa Haverd⁵, Jens Kattge⁴, Natasha MacBean¹, Fabienne Maignan¹, Päivi Merilä⁶, Josep Penuelas^{7,12}, Philippe Peylin¹, Bernard Pinty⁸, Hans Pretzsch⁹, Ernst Detlev Schulze⁴, Didier Solyga^{1,13}, Nicolas Vuichard¹, Yuan Yan³, and Sebastiaan Luyssaert^{1,*}

¹ LSCE, IPSL, CEA-CNRS-UVSQ, 91191 Gif-sur-Yvette, France

³ University of Antwerp, 2610 Wilrijk, Belgium

⁴ MPI-Biogeochemistry, Jena, Germany

⁵ CSIRO-Ocean and Atmosphere Flagship, 2600 Canberra, Australia

⁶ METLA, Oulu, Finland

⁷ CSIC, Global Ecology Unit CREAF-CSIC-UAB, Cerdanyola del Valles, Spain

⁸ European Commission, Joint Research Centre, Ispra, Italy

⁹ TUM, Munich, Germany

¹⁰ Helmholtz-Zentrum Geesthacht, Climate Service Center 2.0, Hamburg, Germany

¹¹ VITO, 2400 Mol, Belgium

¹² CREAF, Cerdanyola del Vallès, Spain

¹³ CGG, 91341 Massy, France

* equal contribution to model development and parametrization

Correspondence to: Kim Naudts (kim.naudts@lsce.ipsl.fr)

² INRA, 21079 Dijon, France

Abstract.

Since 70% of global forests are managed and forests impact the global carbon cycle and the energy exchange with the overlying atmosphere, forest management has the potential to mitigate climate change. Yet, none of the land surface models used in Earth system models, and therefore none

- 5 of today's predictions of future climate, account for the interactions between climate and forest management. We addressed this gap in modelling capability by developing and parametrizing a version of the land surface model ORCHIDEE to simulate the biogeochemical and biophysical effects of forest management. The most significant changes between the new branch called ORCHIDEE-CAN (SVN r2290) and the trunk version of ORCHIDEE (SVN r2243) are the allometric-based allocation
- 10 of carbon to leaf, root, wood, fruit and reserve pools; the transmittance, absorbance and reflectance of radiation within the canopy; and the vertical discretisation of the energy budget calculations. In addition, conceptual changes were introduced towards a better process representation for the interaction of radiation with snow, the hydraulic architecture of plants, the representation of forest management and a numerical solution for the photosynthesis formalism of Farquhar, von Caemmerer and Berry.
- 15 For consistency reasons, these changes were extensively linked throughout the code. Parametrization was revisited after introducing twelve new parameter sets that represent specific tree species or genera rather than a group of often distantly related or even unrelated species, as is the case in widely used plant functional types. Performance of the new model was compared against the trunk and validated against independent spatially explicit data for basal area, tree height, canopy strucure, GPP,
- 20 albedo and evapotranspiration over Europe. For all tested variables ORCHIDEE-CAN outperformed the trunk regarding its ability to reproduce large-scale spatial patterns as well as their inter-annual variability over Europe. Depending on the data stream, ORCHIDEE-CAN had a 67% to 92% chance to reproduce the spatial and temporal variability of the validation data.

1 Introduction

- 25 Forests play a particularly important role in the global carbon cycle. Forests store almost 50% of the terrestrial organic carbon and 90% of vegetation biomass (??). Globally, 70% of the forest is managed and the importance of management is still increasing both in relative and absolute terms. In densely populated regions, such as Europe, almost all forest is intensively managed by humans. Recently, forest management has become a top priority on the agenda of political negotiations to mit-
- 30 igate climate change (Kyoto Protocol, http://unfccc.int/resource/docs/convkp/kpeng.pdf). Because forest plantations may remove CO₂ from the atmosphere, if used for energy production, harvested timber is a substitute for fossil fuel. Forest management thus has great potential for mitigating climate change, which was recognized in the United Nations Framework Convention on Climate Change and the Kyoto Protocol.

- 35 Forests not only influence the global carbon cycle, they also dramatically affect the water vapour and energy fluxes exchanged with the overlying atmosphere. It has been shown, for example, that the evapotranspiration of young plantations can be so great that the streamflow of neighbouring creeks is reduced by 50% (?). Modelling studies on the impact of forest plantations in regions that are snow-covered in winter suggest that because of their reflectance (the so-called albedo), forest
- 40 could increase regional temperature by up to four degrees (????). Management-related changes in the albedo, energy balance and water cycle of forests (??) are of the same magnitude as the differences between forests, grasslands and croplands (?). Moreover, changes in the water vapour and the energy exchange may offset the cooling effect obtained by managing forests as stronger sinks for atmospheric CO₂ (?). Despite the key implications of forest management on the carbon-
- 45 energy-water exchange there have been no integrated studies on the effects of forest management on the Earth's climate.

Earth system models are the most advanced tools to predict future climate (?). These models represent the interactions between the atmosphere and the surface beneath, with the surface formalized as a combination of open oceans, sea ice and land. For land, five classes are distinguished: glacier,

- 50 lake, wetland, urban and vegetated. Vegetation is typically represented by different plant functional types. ORCHIDEE is the land surface component of the IPSL (Institut Pierre Simon Laplace) Earth System Model. Hence, by design, the ORCHIDEE model can be run coupled to the global circulation model LMDz. In this coupled set-up, the atmospheric conditions affect the land surface and the land surface, in turn, affects the atmospheric conditions. Coupled land-atmosphere models thus offer
- 55 the possibility to quantify both the climatic effects of changes in the land surface and the effects of climate change on the land surface. The most advanced land-surface models used, for instance, in Earth System Models to predict climate changes (see the recent CMIP5 exercise), account for changes in vegetation cover but consider forests to be mature and ageless, e.g., JSBACH (?), CLM (?), MOSES (?), ORCHIDEE (?) and LPJ-DVGM (?). At present, none of the predictions of future
- 60 climate thus account for the essential interactions between forest management and climate. This gap in modelling capability provides the motivation for further development of the land-surface model ORCHIDEE to realistically simulate both the biophysical and biogeochemical effects of forest management on the climate. The ORCHIDEE-CAN (short for ORCHIDEE-CANOPY) branch of the land surface model was specifically developed to quantify the climatic effects of forest management.
- 65 The aim of this study is to describe the model developments and parametrization within ORCHIDEE-CAN and to evaluate its performance. ORCHIDEE-CAN is validated against structural, biophysical and biogeochemical data at the European scale. To allow comparison with the standard version of ORCHIDEE, ORCHIDEE-CAN was run with a single-layer energy budget. A more detailed description and evaluation of the new multi-layer energy budget and multilevel radiative transfer scheme is
- 70 given by ?, ? and ?. A new forest management reconstruction which is needed to drive forest man-

agement in ORCHIDEE-CAN is presented in ? and the interactions between forest management and the new albedo scheme have been discussed by ?.

2 Model overview

2.1 The Starting point: ORCHIDEE SVN r2243

- 75 The land surface model used for this study, ORCHIDEE, is based on two different modules (?, their Fig. 2). The first module describes the fast processes such as the soil water budget and the exchanges of energy, water and CO₂ through photosynthesis between the atmosphere and the biosphere (??). The second module simulates the carbon dynamics of the terrestrial biosphere and essentially represents processes as maintenance and growth respiration, carbon allocation, litter decomposition, soil
- 80 carbon dynamics and phenology (?). The trunk version of ORCHIDEE describes global vegetation by 13 metaclasses (MTC) with a specific parameter set (one for bare soil, eight for forests, two for grasslands and two for croplands). Each MTC can be divided into a user-defined number of PFTs which can be characterised by at least one parameter value that differs from the parameter settings of the MTC. Parameters that are not given at the PFT-level are assigned the default value for the MTC
- 85 to which the PFT belongs. By default none of the parameters are specified at the PFT-level, hence, MTCs and PFTs are the same for the standard ORCHIDEE trunk version. A concise description of the main processes in the ORCHIDEE-trunk version and a short motivation to change these modules in ORCHIDEE-CAN is given in Table ??.
- Before running simulations, it is necessary to bring the soil carbon pools into equilibrium due to 90 their slow fill rates, an approach known as model spin-up (??). For a long time, spin-ups have been performed by brute force, i.e., running the model iteratively over a sufficiently long period which allows even the slowest carbon pool to reach equilibrium. This naïve approach is reliable but slow (in the case of ORCHIDEE it takes 3000 simulation years) and thus comes with a large computational demand, often exceeding the computational cost of the simulation itself. Alternative spin-up
- 95 methods calling only parts of the model, e.g., subsequent cycles of 10 years of only photosynthesis followed by 100 year cycles of only soil processes, have been used for ORCHIDEE to reduce the computational cost in the past. These approaches, however, tend to lead to instabilities in litter and carbon pools. In recent years, semi-analytical methods have been proposed as a cost-effective solution to the spin-up issue (???). A matrix-sequence method has been implemented in ORCHIDEE
- 100 following the approach used by the PaSim model (?). The semi-analytical spin-up implemented in ORCHIDEE relies on algebraic methods to solve a linear system of equations describing the seven carbon pools separately for each PFT. Convergence of the method and thus equilibrium of the carbon pools is assumed to be reached when the variation of the passive carbon pool (which is the slowest) drops below a predefined threshold. The net biome production (NBP) is used as a second diagnostic
- 105 criterion to confirm equilibrium of the carbon pools. In order to optimize computing resources, the

semi-analytical spin-up will stop before the end of the run once the convergence criteria are met. OR-CHIDEE's implementation of the semi-analytical spin-up has been validated at regional and global scales against a naïve spin-up, and has been found to converge 12 to 20 times faster. The largest gains were realised in the tropics and the smallest gains in boreal climate (not shown).

110 2.2 Modifications between ORCHIDEE SVN r2243 and ORCHIDEE-CAN SVN r2290

One major overarching change in the ORCHIDEE-CAN branch is the increase of internal consistency within the model by adding connections between the different processes (Fig. ??, red arrows). A more specific novelty is the introduction of circumference classes within forest PFTs, based on the work of ?. For the temperate and boreal zone, tree height and crown diameter are calculated

- 115 from allometric relationships of tree diameter that were parametrized based on the French, Spanish, Swedish and German forest inventory data and the observational data from ?. The circumference classes thus allow calculation of the social position of trees within the canopy which justifies applying an intra-tree competition rule (?) to account for the fact that trees with a dominant position in the canopy are more likely to intercept light than suppressed trees, and, therefore, contribute more
- 120 to the stand level photosynthesis and biomass growth. To respect the competition rule of ?, a new allocation scheme was developed based on the pipe model theory (?) and its implementation by ?. The scheme allocates carbon to different biomass pools (leaves, fine roots, and sapwood) while respecting the differences in longevity and hydraulic conductivity between the pools. In addition to the biomass of the different pools, LAI, crown volume, crown density, stem diameter, stem height
- 125 and stand density are calculated and now depend on accumulated growth. The new scheme allows for the removal of the parameter that caps the maximum LAI (Table **??**).

The calculation of tree dimensions (e.g., sapwood area and tree height) that respect the pipe theory supports making use of the hydraulic architecture of plants to calculate the plant water supply (Fig. **??**, arrow 1), which is the amount of water a plant can transport from the soil to its stomata.

- 130 The representation of the plant hydraulic architecture is based on the scheme of **?**. The water supply is calculated as the ratio of the pressure difference between soil and leaves, and the total hydraulic resistance of the roots, leaves and sapwood, where the sapwood resistance is increased when cavitation occurs. Species-specific parameter values were compiled from the literature. As the scheme makes use of the soil water potential, it requires the use of the 11 layer hydrology scheme of **?** (Table
- 135 **??**). When transpiration based on energy supply exceeds transpiration based on the water supply, the latter restricts stomatal conductance directly, which is a physiologically more realistic representation of drought stress than the reduction of the carboxylation capacity (?) done in the standard version of ORCHIDEE (further also referred to as "trunk" version). In line with this approach, the drought stress factor used to trigger phenology and senescence is now calculated as the ratio between the
- transpiration based on water supply and transpiration based on atmospheric demand (Fig. ??, arrow 2).

The new allocation scheme also drastically changed the way forests are represented in the ORCHIDEE-CAN branch. Although the exact location of the canopies in the stand is not known, individual tree canopies are now spherical elements with their horizontal location following a Poisson distribution

- 145 across the stand. Each PFT contains a user-defined number of model trees, each one corresponding to a circumference class. Model trees are replicated to give realistic stand densities. Following tree growth, canopy dimensions and stand density are updated (Fig. **??**, arrow 3). This formulation results in a dynamic canopy structure that is exploited in other parts of the model, i.e., precipitation interception, transpiration, energy budget calculations, radiation scheme (Fig. **??**, arrow 4) and ab-
- 150 sorbed light for photosynthesis (Fig. ??, arrow 5). In the trunk version these processes are driven by the big-leaf canopy assumption. The introduction of an explicit canopy structure is thought to be a key development with respect to the objectives of the ORCHIDEE-CAN branch, i.e., quantifying the biogeochemical and biophysical effects of forest management on atmospheric climate.

The radiation transfer scheme at the land surface benefits from the introduction of canopy struc-155 ture. The trunk version of ORCHIDEE prescribes the vegetation albedo solely as a function of LAI. In the ORCHIDEE-CAN branch each tree canopy is assumed to be composed of uniformly distributed single scatterers. Following the assumption of a Poisson distribution of the trees on the land surface, the model of ? calculates the transmission probability of light to any given vertical point in the forest. This transmission probability is then used to calculate an effective LAI, which is a

- 160 statistical description of the vertical distribution of leaf mass that accounts for stand density and horizontal tree distribution. The complexity and computational costs are largely reduced by using the effective LAI in combination with the 1D two stream radiation transfer model of ? rather than resolving a full 3-D canopy model. By using the effective LAI, the 1-D model reproduces the radiative fluxes of the 3-D model. The approach of the two stream radiation transfer model was extended for a
- 165 multi-layer canopy (?) to be consistent with the multi-layer energy budget and to better account for non-linearities in the photosynthesis model. The scattering parameters and the background albedo (i.e. the albedo of the surface below the dominant tree canopy) for the two stream radiation transfer model were extracted from the Joint Research Centre Two-stream Inversion Package (JRC-TIP) remote sensing product (section 4.7). This approach produces fluxes of the light absorbed, transmitted,
- 170 and reflected by the canopy at vertically discretized levels, which are then used for the energy budget (Fig. ??, arrow 6) and photosynthesis calculations (Fig. ??, arrow 5).

The canopy radiative transfer scheme of (?) separates the calculation of the fluxes resulting from downwelling direct and diffuse light, with different scattering parameters available for near-infrared (NIR) and visible (VIS) light sources. The snow albedo scheme in the trunk does not distinguish be-

175 tween these two shortwave bands. Therefore, the snow scheme of the Biosphere-Atmosphere Transfer Scheme (BATS) for the Community Climate Model (?) was incorporated into the ORCHIDEE-CAN branch, since it distinguishes between the NIR and VIS radiation. The radiation scheme of ? requires snow to be put on the soil below the tree canopy instead of on the canopy itself. The calculation of the snow coverage of a PFT therefore had to be revised according to the scheme of ?, which

180 allows for snow to completely cover the ground at depths greater than 0.2 m. The parameter values of ? were used in the ORCHIDEE-CAN branch.

The ORCHIDEE-CAN branch differs from any other land surface model by the inclusion of a newly developed multi-layer energy budget. There are now subcanopy wind, temperature, humidity, longwave radiation and aerodynamic resistance profiles, in addition to a check of energy closure at

- all levels. The energy budget represents an implementation of some of the characteristics of detailed 185 single site, iterative canopy models (e.g., ??) within a system that is coupled implicitly to the atmosphere. As an enhancement to the trunk version of ORCHIDEE (Table ??), the new approach also generates a leaf temperature, using a vegetation profile and a vertical shortwave and longwave radiation distribution scheme (?), which will be fully available when parametrisation of the scheme
- has been completed across test sites corresponding to the species within the model (?). As with the 190 trunk version, the new energy budget is calculated implicitly (??). An implicit solution is a linear solution in which the surface temperature and fluxes are calculated in terms of the atmospheric input at the same time-step, whereas an explicit solution uses atmospheric input from the previous time step to calculate the surface temperature and fluxes. Although it is less straightforward to derive, the
- 195 implicit solution is more computationally efficient and stable, which allows the model to be run over a time-step of 15 minutes when coupled to the atmospheric model LMDz - much longer than would be the case for an explicit model. Parameters were derived by optimizing the model against the observations from short-term field campaigns. The new scheme may also be reduced to the existing single layer case, so as to provide a means of comparison and compatability with the ORCHIDEE
- 200 trunk version.

The combined use of the new energy budget and the hydraulic architecture of plants required changes to the calculation of the stomatal conductance and photosynthesis (Fig. ??, arrow 7). When water supply limits transpiration, stomatal conductance is reduced and photosynthesis needs to be recalculated. Given that photosynthesis is among the computational bottlenecks of the model, the

- semi-analytical procedure as available in previous trunk versions (r2031 and further) is replaced by 205 an adjusted implementation of the analytical photosynthesis scheme of ?, which is also implemented in the latest ORCHIDEE-trunk version. In addition to an analytical solution for photosynthesis the scheme includes a modified Arrhenius function for the temperature dependence that accounts for a decrease of carboxylation capacity (k_{Vcmax}) and electron transport capacity (k_{Jmax} , see Table ??
- 210 for variable explanations) at high temperatures and a temperature dependent $k_{Jmax/Vcmax}$ ratio (?). The temperature response of k_{Vcmax} and k_{Jmax} was parametrized with values from reanalysed data in literature (?), whereas k_{Vcmax} and k_{Jmax} at a reference temperature of 25° C were derived from observed species-specific values in the TRY database (?). As the amount of absorbed light varies with height (or canopy depth), the absorbed light computed from the albedo routines is now directly
- 215 used in the photosynthesis scheme resulting in full consistency between the top of the canopy albedo

and absorption. This new approach replaces the old scheme which used multiple levels based on the leaf area index, not the physical height.

ORCHIDEE-CAN incorporates a systematic mass balance closure for carbon cycling to assure that carbon is not getting created or destroyed during the simulation. Hence, budget closure is now consistently checked for water, carbon and energy throughout the model.

220

The trunk uses 13 plant functional types (PFT) to represent vegetation globally: one PFT for bare soil, eight for forests, two for grasslands, and two for croplands. The ORCHIDEE-CAN branch makes use of the externalization of the PFT-depedent parameters by adding 12 parameter sets that represent the main European tree species. Species parameters were extracted from a wide range of

sources including original observations, large databases, primary research and remote sensing products (section 4). The use of age classes is introduced through externalization of the PFT parameters as well. Age classes are used during land cover change and forest management to simulate the regrowth of a forest. Following a land cover change, biomass and soil carbon pools (but not soil water columns) are either merged or split to represent the various outcomes of a land cover change. The number of age classes is user defined. Contrary to typical age classes, the boundaries are determined by the tree diameter rather than the age of the trees.

Finally, the forest management strategies in the ORCHIDEE-CAN branch were refined from the original forest management branch (?). Self-thinning was activated for all forests regardless of human management, contrary to the original FM branch. The new default management strategy thus

- has no human intervention but includes self-thinning, which replaces the fixed 40 year turnover time for woody biomass. Three management strategies with human intervention have been implemented: (1) "high stands", in which human intervention is restricted to thinning operations based on stand density and diameter, with occasional clearcuts. Aboveground stems are harvested during operations, while branches and belowground biomass are left to litter. (2) "coppices" involve two kinds of cuts.
- 240 The first coppice cut is based on stem diameter and the aboveground woody biomass is harvested whereas the belowground biomass is left living. From this belowground biomass new shoots sprout, which increases the number of aboveground stems. In subsequent cuts the amount of shoots is not increased, although all aboveground wood biomass is still harvested. (3) "short rotation coppices", where rotation periods are based on age and are generally very short (3-6 years). The different man-
- 245 agement strategies can occur with or without litter raking, which reduces the litter pools and has a longterm effect on soil carbon (?). All management types are parametrized based on forest inventory data, yield tables and guidelines for forest management. The inclusion of forest management resulted in two additional carbon pools, branches and coarse roots (i.e., abovegound and belowground woody biomass) and therefore required an extension to the semi-analytical spin-up method (section
- 250 2.1). The semi-analytical spin-up is now run for nine C pools.

3 Description of the developments

3.1 Allocation

Following bud burst, photosynthesis produces carbon that is added to the labile carbon pool. Labile carbon is used to sustain the maintenance respiration flux (F_{rm}) , which is the carbon cost to

255

keep existing tissue alive (?). Maintenance respiration for the whole plant is calculated by summing maintenance respiration of the different plant compartments, which is a function of the nitrogen concentration of the tissue following the Beer-Lambert law and subtracted from the whole-plant labile pool (up to a maximum of 80% of the labile pool).

The remaining labile carbon pool is split into an active and none-active pool. The size of the active pool is calculated as a function of plant phenology and temperature and was formalized following **???**. The remaining non-active pool is used to restore the labile and carbohydrate reserves pools according to the rules proposed in **?**. The labile pool is limited to 1% of the plant biomass or 10 times the actual daily photosynthesis. Any excess carbon is transferred to the non-respiring carbohydrate reserve pool. The carbohydrate reserve pool is capped to reflect limited starch accumulation in plants,

but carbon can move freely between the two reserve pools. After accounting for growth respiration (F_{rg}) , i.e., the cost for producing new tissue excluding the carbon required to build the tissue itself (?), the total allocatable C used for plant growth is obtained (M_{totinc}) .

New biomass is allocated to leaves, roots, sapwood, heartwood, and fruits. Allocation to leaves, roots and wood respects the pipe model theory (?) and thus assumes that producing one unit of

- 270 leaf mass requires a proportional amount of sapwood to transport water from the roots to the leaves as well as a proportional fraction of roots to take up the water from the soil. The different biomass pools have different turnover times, and therefore at the end of the daily time step, the actual biomass components may no longer respect the allometric relationships. Consequently, at the start of the time step carbon is first allocated to restore the allometric relationships before the remaining carbon is
- 275 allocated in the manner described below. The scaling parameter between leaf and sapwood mass is derived from:

$$d_l = k_{ls} \times m_w \times d_s \tag{1}$$

280

Where
$$d_l$$
 is the one-sided leaf area of an individual plant, d_s is the sapwood cross-section area
of an individual plant, k_{ls} a parameter linking leaf area to sapwood cross-section area and, m_w is
the water stress as defined in section 3.2. Alternatively, leaf area can be written as a function of leaf
mass (M_l) and the specific leaf area (k_{sla}) :

$$d_l = M_l \times k_{sla} \tag{2}$$

Sapwood mass M_s can be calculated from the sapwood cross-section area d_s as follows:

$$M_s = d_s \times d_h \times k_{\rho s} \tag{3}$$

285

Where d_h is the tree height and $k_{\rho s}$ is the sapwood density. Following substitution of equations 2 and 3 into equation 1, leaf mass can be written as a function of sapwood mass:

$$M_l = \left(M_s \times f_{KF}\right)/d_h \tag{4}$$

where,

$$f_{KF} = (k_{ls} \times m_w) / (k_{sla} \times k_{\rho s}) \tag{5}$$

where, k_{ls} is calculated as a function of the gap fraction as supported by site-level observations 290 (?):

$$k_{ls} = k_{lsmin} + f_{Pgap,trees} \times (k_{lsmax} - k_{lsmin}) \tag{6}$$

 k_{lsmin} is the minimum observed leaf area to sapwood area ratio, k_{lsmax} is the maximum observed leaf area to sapwood area ratio and $f_{Pgap,trees}$ is the actual gap fraction. By using the gap fraction as a control of k_{ls} more carbon will be allocated to the leaves until canopy closure is reached.

295

Following ?, sapwood mass and root mass (M_r) are related as follows:

$$M_s = k_{sar} \times d_h \times M_r \tag{7}$$

where the parameter k_{sar} is calculated according to ? (their equation (17)):

$$k_{sar} = \sqrt{(k_{rcon}/k_{scon}) \times (k_{\tau s}/k_{\tau r}) \times k_{\rho s}}$$
(8)

300

where k_{rcon} is the hydraulic conductivity of roots, k_{scon} is the hydraulic conductivity of sapwood, $k_{\tau s}$ is the longevity of sapwood and $k_{\tau r}$ is the root longevity. Following substitution of equation (4) into (7) and some rearrangement, leaf mass can be written as a function of root mass:

$$M_l = f_{LF} \times M_r \tag{9}$$

where,

$$305 \quad f_{LF} = k_{sar} \times f_{KF} \tag{10}$$

Parameter values used in equations 1 to 9, i.e., k_{lsmax} , k_{lsmin} , k_{sar} , k_{sla} , $k_{\rho s}$, k_{rcon} , k_{scon} , $k_{\tau s}$ and $k_{\tau r}$, are based on literature review (Table S1,S2 and S3). The allometric relationships between the plant components and the hydraulic architecture of the plant (section 3.2) are both based on the pipe model theory, hence, both the allocation and the hydraulic architecture module use the same parameter values for root and sapwood conductivity.

In this version of ORCHIDEE, forests are modelled to have k_{ncirc} circumference classes with d_{ind} identical trees in each one. Hence, the allocatable biomass (M_{totinc}) needs to be distributed across l diameter classes:

$$M_{totinc} = sum(l)[d_{ind(l)} \times M_{inc(l)}]$$
⁽¹¹⁾

where $M_{inc(l)}$ is the biomass that can be allocated to diameter class *l*. Mass conservation thus requires:

$$M_{inc(l)} = M_{linc(l)} + M_{rinc(l)} + M_{sinc(l)}$$

$$\tag{12}$$

where $M_{linc(l)}$, $M_{rinc(l)}$ and, $M_{sinc(l)}$ are the increase in leaf, root and wood biomass for a tree in diameter class l, respectively. Equation 4 and 9 can be rewritten as

320
$$(M_{l(l)} + M_{linc(l)})/(M_{s(l)} + M_{sinc(l)}) = f_{KF}/(d_{h(l)} + d_{hinc(l)})$$
 (13)

$$(M_{l(l)} + M_{linc(l)}) = (M_{r(l)} + M_{rinc(l)}) \times f_{LF}$$
(14)

An allometric relationship is used to describe the relationship between tree height and basal area (?):

$$d_{h(l)} = k_{\alpha 1} \times (4/\pi \times d_{ba(l)})^{(k_{\beta 1}/2)}$$
(15)

325 The change in height is then calculated as

$$d_{hinc(l)} = [k_{\alpha 1} \times (4/\pi \times (d_{ba(l)} + d_{bainc(l)}))^{(k_{\beta 1}/2)}] - d_{h(l)}$$
(16)

where $d_{ba(l)}$ and $d_{bainc(l)}$ are the basal area and its increment, respectively. $k_{\alpha 1}$ and $k_{\beta 1}$ are allometic constants relating tree diameter and height. The distribution of C across the *l* diameter classes depends on the basal area of the model tree within each diameter class. Trees with a large basal area are assigned more carbon for wood allocation than trees with a small basal area, according to the method of **?**.

330

310

$$d_{bainc(l)} = f_{\gamma} \times \left(d_{circ(l)} - k_m * g_{\sigma} + \sqrt{(k_m \times g_{\sigma} + d_{circ(l)})^2 - (4 \times g_{\sigma} \times d_{circ(l)})} \right) / 2$$
(17)

where k_m is a parameter, f_{γ} and g_{σ} are calculated from parameters and $d_{circ(l)}$ is the circumference of the model tree in diameter class l. g_{σ} is a function of the diameter distribution of the stand at a given time step.

335 at a given time step.

Equations 10 to 16 need to be simultaneously solved. An iterative scheme was avoided by linearising equation 15, which was found to be an acceptable numerical approximation as allocation is calculated at a daily time step, and hence the changes in height are small and the relationship is locally linear:

$$340 \quad d_{hinc(l)} = d_{bainc(l)} / f_s \tag{18}$$

where f_s is the slope of the locally linearised equation 15 and is calculated as:

$$f_s = k_{step} / (k_{\alpha 1} \times (4/\pi * (d_{ba} + k_{step}))^{(k_{\beta 1}/2)} - k_{\alpha 1} \times (4/\pi \times d_{ba})^{(k_{\beta 1}/2)})$$
(19)

Equations 10, 11, 12, 13, 14, 16, 17 and 18 are then solved for f_{γ} . f_{γ} distributes photosynthates across the different diameter classes and as such controls the intra-species competition within a stand. f_{γ} thus depends on the total allocatable carbon and needs to be optimised at every time step.

Once f_{γ} has been calculated, $M_{linc(l)}$, $M_{rinc(l)}$ and $M_{sinc(l)}$ can be calculated.

3.2 Hydraulic architecture

345

The representation of the impact of soil moisture stress on water, carbon and energy fluxes has been identified as one of the major uncertainties in land surface models (?). Neither the empirical functions nor the soil moisture stress functions, which are commonly used in land surface models

- 350 functions nor the soil moisture stress functions, which are commonly used in land surface models, fully capture stomatal closure and limitation of C uptake during drought stress (??). Therefore, we replaced the soil moisture stress function which limits C assimilation through a constrain on k_{Vcmax} in the ORCHIDEE-trunk, by a constrain based on the amount of water plants can transport from the soil to their leaves.
- The model calculates plant water supply according to the implementation of hydraulic architecture by ?. Plant water supply is the amount of water the plant can transport from the soil to its stomata, accounting for the resistances to water transport in the roots, sapwood and leaves. If transpiration rate exceeds plant water supply, the stomatal conductance is reduced until equilibrium is reached.

The water flow from the soil to the leaves is driven by a gradient of decreasing water potential. 360 Using Darcy's law (??), the supply of water for transpiration through stomata can be described as:

$$F_{Trs} = p_{delta} / (R_r + R_{sap} + R_l) \tag{20}$$

where p_{delta} is the pressure difference between the soil and the leaves; and R_r , R_{sap} and R_l are the hydraulic resistances of fine roots, sapwood and leaves, respectively. pdelta is calculated following ?:

$$365 \quad p_{delta} = p_{\psi sr} - k_{\psi l} - (d_h \times k_{\rho w} \times k_g) \tag{21}$$

where $k_{\psi l}$ is a PFT-specific minimal leaf water potential, which means that plants are assumed to maximise water uptake by lowering their $k_{\psi l}$ to the minimum, if transpiration exceeds F_{Trs} (?). The product of d_h , $k_{\rho w}$ and k_g accounts for the loss in water potential by lifting a mass of water from the soil to the place of transpiration at height d_h , $k_{\rho w}$ is the density of water, and k_g is the gravitational constant. The soil water potential in the rooting zone $(p_{\psi sr})$ was calculated by adding a modulator

 (m_{ψ}) to the bulk soil water potential, which was calculated as the sum of the soil water potential in each soil layer weighted by the relative share of roots (d_{rd}) in the individual soil layer:

$$p_{\psi sr} = sum(l)[p_{\psi s} \times d_{rd}] + m_{\psi} \tag{22}$$

The soil water potential for each layer $p_{\psi sl}$ is calculated from soil water content according to ?.

375
$$p_{\psi s(l)} = \frac{1}{k_{av}} \left(\left(\frac{M_{swc} - k_{swcr}}{k_{swcs} - k_{swcr}} \right)^{-1/k_{mv}} - 1 \right)^{1/k_{nv}}$$
(23)

where M_{swc} is the volumetric soil water content, k_{swcr} and k_{swcs} are respectively the residual and saturated soil water content and k_{av} , k_{mv} and k_{nv} are parameters.

Root resistance is related to the root mass and thus can be expressed as (?):

$$R_r = \frac{1}{(k_{rcon} \times M_r)} \tag{24}$$

380

385

where k_{rcon} is the fine root hydraulic conductivity per unit biomass. Sapwood resistance is calculated according to ?:

$$R_{sap} = \frac{d_h}{(d_s \times k_{scon})} \tag{25}$$

where k_{scon} is the sapwood specific conductivity, which is decreased when cavitation occurs. The loss of conductance as a result of cavitation is a function of $p_{\psi sr}$ and was implemented by using an s-shaped vulnerability curve:

$$k_{scon} = k_{scon} \times e^{(-p_{\psi rs}/k_{\psi 50})^{k_c}}$$
(26)

where $k_{\psi 50}$ is the $p_{\psi sr}$ that causes 50% loss of conductance; and k_c is a shape parameter. R_l is related to the specific leaf conductivity per unit leaf area (k_l) and the leaf area index :

$$R_l = \frac{1}{(k_{lcon} \times d_{LAI})} \tag{27}$$

390 The response of water viscosity to low temperatures increases the resistance (?). The relationship is described as:

$$R_{temp} = \frac{R}{(k_{\alpha 1v} + k_{\alpha 2v} \times T)}$$
(28)

where $k_{\alpha 1v}$ and $k_{\alpha 2v}$ are empirical parameters (?), R_{temp} is the temperature adjusted R_l , R_{sap} or R_r , T is air temperature for R_l and R_{sap} and T is soil temperature for R_r .

395

If, for any time step, the transpiration calculated by the energy budget exceeds the amount of water the plant can transport from the soil to its stomata, transpiration is limited to the plant water supply. As the transpiration is now reduced, the initial calculations of the energy budget and photosynthesis, solely based on atmospheric information, are no longer valid. As a result the energy budget and photoget and photosynthesis must be recalculated for the time step in question. For this recalculation,

- 400 stomatal conductance at the canopy level is calculated such that transpiration equals the amount of water the plant can transport. Owing to the feedback between stomatal conductance, leaf surface temperature and transpiration, this calculation may require up to 10 iterations to converge, using a stationary iterative method. When the multi-layer energy budget is reduced to its single-layer implementation, however, canopy level stomatal conductance is decomposed to obtain the stomatal
- 405 conductance at each canopy layer assuming that each layer is equally restricted by drought stress. Finally, the restricted stomatal conductance is used to calculate CO₂ assimilation rate according to the photosynthesis model by Farquhar, von Caemmerer and Berry (section 3.6).

3.3 Canopy structure

- Stand structure controls the amount of light that penetrates to a given depth in the canopy. For
 example, the amount of light reaching the forest floor will be higher for a stand with few mature trees compared to many young trees even if both stands have the same leaf area index. Where a big leaf approach assumes a homogeneous block shaped canopy (as in the trunk version of ORCHIDEE) and can therefore rely on the law of Beer-Lambert, a geometric approach is required to calculate light penetration through structured canopies. Light penetration needs to be simulated to calculate albedo
- 415 (section 3.4), photosynthesis (section 3.6), partitioning of energy fluxes (section 3.5) and the amount of light reaching the forest floor (see for example section 3.1). The gap fraction, which is the basic information in calculating light penetration at different depths in the canopy, is calculated following the approach presented by ? and formalized in their semi-analytical model. Rather than a spatially

explicit approach, ? follow a statistical approach which reduces the memory requirements for the simulations and limits the space requirements for storing the model output files.

The model of ? represents the canopy by a statistical height distribution with varying crown sizes and stem diameters for each height class. The crown canopies are treated as spheroids containing homogeneously distributed single scatterers. Although this f_{Pgap} model can explicitly include trunks, we made the decision to exclude them, as the spectral parameters for our radiation model (section

425 3.4) are extracted from remote sensing data (section 4.8) without distinguishing between leafy and woody masses. This gives the gap probability for trees as a function of height (z) and solar zenith angle (θ_z):

$$f_{Pgap}^{trees}(\theta_z, z) = e^{-d_\lambda \times \overline{d_c(\theta_z, z) \times (1 - f_{Pwc}(\theta_z, z))}}$$
(29)

430

440

where d_{λ} is the inverse of the tree density, d_c is the projected crown area (for an opaque canopy), and f_{Pwc} is the mean crown porosity. The overbar depicts the mean over the tree distribution as a function of tree height or, in our case, the mean over the *l* circumference classes. Following minor adaptations, the implementation of Haverd and Lovell (?) was incorporated in ORCHIDEE-CAN. As there also exist crops, grasses, and bare soil in the model, f_{Pgap} was adjusted for these situations as well. For grasses and crops, the same formulation is used:

$$435 \quad f_{Pgap}^{gc}(\theta_z, z) = e^{-0.5 \times d_{LAIabove} \times m_{LAIcorr}/\cos(\theta_z)} \tag{30}$$

where $d_{LAIabove}$ is the total amount of LAI above height z, and $m_{LAIcorr}$ is a correction factor to account for the fact that grasses and crops are treated as homogeneous blocks of vegetation with no internal structure and is often referred to as a clumping factor. Here it is treated as a tunable parameter and therefore the term correction factor was used. For bare soil, there is no vegetation to intercept radiation, and therefore $f_{Pgap}^{bs}(\theta_z, z)$ is always unity.

3.4 Multi-layer two-way radiation scheme for tall canopies

Species-specific radiation absorbance, reflectance and transmittance by the forest canopy were calculated from a radiation transfer model (?) which was parametrized by satellite-derived species-specific scattering values (section 4.8). Given the complexity of radiation transfer, it remains challenging to

445 accurately simulate radiation transfer through structurally and optically complex vegetation canopies without using explicit 3-D models. The applied 1-D model belongs to the family of two-stream models (?) and thus calculates transmittance, absorbance and reflectance of both the incoming and outgoing radiation. The calculation of the reflectance at the top of the canopy due to a collimated source (i.e., the sun) is divided in three components: 450 1. scattering of radiation between the vegetated elements with a black background

$$f_{Coll,veg}^{f_R} = f(\theta_{mu}, f_{rl}, f_{tl}, g_G, d_{LAIeff})$$
(31)

2. scattering of radiation by the background with a black canopy

$$f_{UnColl,bgd}^{f_R} = f_{Rbgd} \times e^{(-d_{LAIeff}/(2 \times \theta_{mu}))} \times f_{UnColl,veg}^T$$
(32)

3. multiple scattering of radiation between the canopy and the background

455
$$f_{Coll,bgd}^{f_R} = f_{Rbgd} \times [f_{Coll,bgd,1}^{f_R} + f_{Coll,bgd,n}^{f_R}]$$
 (33)

Term (1) is widely used in cloud reflectance calculations, and depends on the cosine of the solar zenith angle (θ_{mu}), the reflectance and transmittance of the single leaves (f_{rl} and f_{tl} , respectively), the leaf orientation function (g_G), and the effective LAI (d_{LAIeff}). The exact definition of this term is given in Equation B2 in ?. In term (2), f_{Rbgd} is the reflectance of the ground beneath the canopy and $f_{UnColl,veg}^T$ is the transmitted fraction of light to the ground which has not collided with any canopy elements. In term (3), $f_{Coll,bgd,1}^{f_R}$ is the fraction of light which has struck vegetation and collided with the background a single time, while $f_{Coll,bgd,n}^{f_R}$ is the fraction which has collided

multiple times (n) with the background.
The sum of the three components results in the canopy albedo (?). Similar equations can be derived for light originating from diffuse sources (e.g., clouds and other atmospheric scattering)(?).
Implementations of the calculations of the canopy fluxes for a single level are available from the JRC, and these implementations were used as the basis of the routines put into ORCHIDEE-CAN for both the single- and multi-level cases (?). This implementation relies on the use of the effective

470 reflectance, absorbance and transmittance as would be obtained by a 3D-canopy representation (?). In this study, the effective LAI was calculated by first computing the canopy gap probability, i.e. the probability that light is transmitted to a specified height in the canopy at a given solar angle. The gap probability is then converted into the effective LAI by passing it as an input to the inverted Beer-Lambert's law (with an extinction coefficient of 0.5 to ensure compatibility with the two-steam

LAI, which is the LAI that needs to be used in a 1-D process representation to obtain the same

475 inversion of ?).

480

460

$$d_{LAIeff} = -2.0 \times \cos(\theta_z) \times \log(f_{Pgap}) \tag{34}$$

where f_{Pgap} can be f_{Pgap}^{trees} , f_{Pgap}^{gc} , f_{Pgap}^{bs} . Following the introduction of multi-layer photosynthesis and energy budget submodels, the approach proposed by ? had to be adjusted such that it could be applied for every level for which absorbance needs to be known to calculate photosynthesis (section 3.6) and reflectance needs to be known to calculate the net shortwave radiation (section 3.5). The

multi-layer approach basically applies the 1D-two stream canopy radiation transfer model by ? to each canopy level where the light transmitted by the overlaying level becomes the input for the lower level.

As the multi-level approach is built around the solution of the one-level scheme for each canopy level, no new equations are introduced. The method can be summarized by the following algorithm for which the details are given in (?). First, three fluxes are calculated for each level independently: the fraction of light transmitted through the layer without striking vegetation, the fraction of light reflected after striking vegetation, and the fraction of light transmitted through the layer after striking vegetation. These three fluxes represent the only possible fate of light (any light not taking one of

- 490 these paths must be absorbed for energy conservation). Next, an iterative approach is invoked which follows the path of a single photon entering the top level. Based on the solutions for each single level, probabilities can be calculated that the photon is transmitted to a lower level or reflected to a higher level. Any fraction which is reflected upwards from the top level is added to the total canopy albedo and not considered further. The fraction which is transmitted through the top level enters the
- 495 next highest level, and again the single level solutions determine where this light goes. Any fraction reflected upwards is considered in the next iteration as part of the light entering the upper level. The steps continue until the bottom canopy level is reached. Here, any fraction which is transmitted into the soil is removed from consideration and added to the total transmittance through the canopy. The algorithm then proceeds to the above canopy level. Now the "transmitted" fluxes are moving in
- 500 the upwards direction towards to the sky, while "reflected fluxes" are moving towards the ground. The code continues towards the top level, taking as input from below both the flux reflected by downwelling light from the level below the current level and the flux transmitted from the lower level by upwelling light. After each iteration (moving from the top of the canopy to the bottom and back to the top), the total amount of light considered "active" has been reduced by light escaping
- 505 to the sky or being absorbed by the canopy or ground. Eventually, this "active" light falls below a pre-defined threshold and the calculation is considered as converged.

Due to the iterative procedure, energy is not strictly conserved, although we have attempted to choose a threshold which minimizes this loss. The multilevel albedo calculation is currently the most expensive part of the model, due to the iterations and the fact that it must be performed over all

510 canopy levels (currently set to 10), grid points, and PFTs at every physical time-step. Levels with no LAI are no less expensive to compute, either, although we have arranged our canopy levels to make sure no levels are empty in most cases.

3.5 Multi-layer energy budget

The present generation of land surface models have difficulties in reproducing consistently the energy balances that are observed in field studies (???). The ORCHIDEE-CAN branch implemented an energy budget scheme that represents more than one canopy layer to simulate the effects of scalar gradients within the canopy for determining more accurately the net sensible and latent heat fluxes that are passed to the atmosphere. As outlined in ?, the use of an implicit solution for coupling between the atmospheric model and the surface layer model is the only way to keep profiles of

temperature and humidity synchronised across the two models when the coupled-model is run over 520 large time steps (e.g., of 30 minutes). The difference between explicit and implicit schemes is that an explicit scheme will calculate each value of the variable (e.g., temperature and humidity) at the current time step entirely in terms of values from the previous time step. An implicit scheme requires the solution of equations written only in terms of those at the current time step.

525

The modelling approach formalises three constraints that ensure energy conservation. The three equations that describe the main interactions are:

1. The energy balance at each layer is the sum of incoming and outgoing fluxes of latent and sensible heat and of shortwave and longwave radiation:

$$k_{lhc,i}k_{\rho_{v},i}\frac{\delta T_{L,i}}{\delta t} = \left(k_{shc}k_{\rho_{a}}\frac{(T_{L,i} - T_{a,i})}{R_{a,i}} + k_{\lambda,LE}\rho_{a}\frac{q_{L,i} - q_{a,i}}{R_{s,i}} + F_{SW,i} + F_{LW,i}\right) \left(\frac{1}{\Delta d_{hl,i}}\right)$$
(35)

- 530 where $F_{LW,i}$ is the sum total of long wave radiation, that is, the net LW radiation absorbed into layer i and $F_{SW,i}$ is the net absorbed short wave radiation as calculated by the radiation scheme in section 3.4. k_{shc} is the specific heat capacity of air. The source sensible heat flux from the leaf at level i is the difference between the leaf temperature $(T_{L,i})$ and the atmospheric temperature at the same level $(T_{a,i})$, divided by $R_{a,i}$, which is the leaf resistance to sensible 535 heat flux (a combination of stomatal and boundary layer resistance). Similarly, the source latent heat flux from the leaf at level i is the difference between the saturated humidity in the leaf $(q_{L,i})$ and that in the atmosphere at level i $(q_{a,i})$, divided by $R_{s,i}$ which is the leaf resistance to latent heat flux. $R_{a,i}$ is calculated based upon the leaf boundary layer resistance, and is described in the present model according to ?. $R_{s,i}$ is the stomatal resistance of the leaf 540 that is calculated using the model of ?.
 - 2. The sensible heat flux between the vegetation ('the leaf') and the surrounding atmosphere at each level, and between adjacent atmospheric levels above and below is provided by the following expression:

$$\frac{\delta T_{a,i}}{\delta t} \Delta d_{V,i} = k_{k,i} \frac{\delta^2 T_{a,i}}{\delta z^2} \Delta d_{A,i} + \left(\frac{T_{L,i} - T_{a,i}}{R_{a,i}}\right) \left(\frac{1}{\Delta d_{hl,i}}\right) \Delta d_{V,i} \tag{36}$$

545

where z denotes the height above the soil suface. We have re-written the scalar conservation equation, as applied to canopies, in terms of the sensible heat flux, temperature and source sensible heat from the vegetation at each layer.

3. The latent heat flux between the vegetation and surrounding atmosphere at each level, and between adjacent atmospheric levels above and below is described in a form that is analogous to equation (36), above:

550

555

$$\frac{\delta q_{a,i}}{\delta t} \Delta d_{V,i} = k_{k,i} \frac{\delta^2 q_{a,i}}{\delta z^2} \Delta d_{A_i} + \left(\frac{q_{L,i} - q_{a,i}}{R_{s,i}}\right) \left(\frac{1}{\Delta d_{hl,i}}\right) \Delta d_{V_i}$$
(37)

In addition to these three basic equations various terms had to be parameterised. The 1D secondorder closure model of ? was used to simulate the vertical transport coefficients $k_{k,i}$ within the canopy while accounting for the vertical and horizontal distribution of LAI (section 3.3). This set of equations were then written in an implicit form and solved by induction. More details on the implicit multi-layer energy budget and a complete mathematical documentation are given in ?.

To complete the energy budget calculations, the multi-layer 1D canopy radiation transfer model (section 3.4) was used to calculate the net shortwave radiation at each canopy layer. Further, the canopy radiation scheme makes use of the Longwave Radiation Transfer Matrix (LRTM) (??). This approach separates the calculation of the radiation distribution completely from the implicit expres-

- 560 approach separates the calculation of the radiation distribution completely from the implicit expression. Instead, a single source term for the long wave radiation is added at each level. This means that the distribution of LW radiation is now explicit (i.e., makes use of information only from the 'previous' and not the 'current' time step) but the changes within the timestep were small enough to not affect the overall stability of the model. However, an advantage of the approach is that it accounts for a higher order of reflections from adjacent levels than the single order assumed in the process
- above.

3.6 Analytical solution for photosynthesis

The photosynthesis model by Farquhar, von Caemmerer and Berry (?) predicts net photosynthesis of C3 plants as the minimum of the Rubisco-limited rate of CO₂ assimilation and the electron transportlimited rate of CO₂ assimilation (?). The ORCHIDEE-CAN branch calculates net photosynthesis following an analytical algorithm as described by ?. In addition, the C4 photosynthesis is calculated by an equivalent version of the Farquhar, von Caemmerer and Berry model that was extended to account for noncyclic electron transport (?). A detailed derivation of the analytical solution of the

Farquhar, von Caemmerer and Berry model is given in ?. Although the exclusion of mesophyll

- 575 conductance from the photosynthesis model could lead to an underestimation of the CO₂ fertilization effect in Earth system models (?), mesophyll conductance was not included in ORCHIDEE-CAN to maintain compatibility between the model formulation and its parametrization. Because values of k_{Vcmax} and k_{Jmax} differ between different formulations of the photosynthesis model (??) and the parametrization that was used in ORCHIDEE-CAN did not include mesophyll conductance, it was
- also not accounted for in the model formulation. The analytical photosynthesis model implemented

in ORCHIDEE-CAN could be easily extended to include mesophyll conductance but that would require reparameterizing the photosynthesis model.

Owing to the canopy structure simulated in this model version and the layering of the canopy, the amount of absorbed light now varies with canopy depth. This new approach replaces the old scheme which uses multiple levels based on the leaf area index, not the physical height within the canopy. Photosynthesis is now calculated at each vertically resolved canopy level independently, using the total amount of absorbed light calculated by the radiation transfer scheme, which means that radiation transfer inside the canopy and photosynthesis are now fully consistent. In the new photosynthesis scheme, photosynthesis thus indirectly depends on canopy structure.

590 3.7 Forest management and natural mortality

595

615

Although forest management has developed a wide range of locally-appropriate and species-specific strategies (?), the nature of large scale land surfaces models such as ORCHIDEE-CAN, require only a limited number of contrasting strategies that are expected to be relevant at the spatial scale (e.g., 50x50 km) of global and regional modelling studies. Four management strategies were implemented based on their expected impact on biogeochemical and biophysical processes:

- 1. In unmanaged stands self-thinning drives stand dynamics and continues until too few trees are left on site. Subsequently, a stand replacing disturbance moves all standing biomass into the appropriate litter pools and a new stand is established.
- 2. High stand management is characterised by regular thinning and a final harvest cut. Thinning is decided on the basis of the deviation between the actual and potential stand density for any given diameter. This approach relates to the so-called relative density index (?), the land use disturbance index (?) or hemeroby and naturalness approaches (?). Exceeding a threshold diameter results in a clear cut and the stand is replanted in the next year. For both thinning and harvest, leaves, roots and belowground wood is transferred to the appropriate litter pools whereas the aboveground woody biomass is removed from the site and stored in a product pool. Trees with a diameter below a species-specific threshold are stored in a short-lived product pool which mimics wood uses for fuel, paper and cardboard. Trees with larger dimensions are moved to medium and long-lived product pools which mimic for example, particle boards and timber usages, respectively.
- 610 3. Coppicing of the aboveground biomass is decided on stem diameter. At harvest, the root system is left intact an in between coppicing, no wood is harvested. Note that at present it is not possible to simulate coppicing-with-standards in ORCHIDEE-CAN.
 - 4. In ORCHIDEE-CAN, stands under short rotation management are limited to poplar (*Populus spp.*) and willow (*Salix spp.*) forests. Stands are harvested at a prescribed age. Following a set number of harvest cycles, the root system is uprooted and the whole stand is replanted.
 - 20

Different age classes are distinguished to better account for the structural diversity and its possible effects on the element, energy and water fluxes. A clear hierarchy was established for the mortality processes regarding the actual killing of trees (i.e., move their biomass to the litter or harvest pools). All of the processes determine first how much biomass they would remove in the absence

620 of all the other processes. Afterwards, the killing is arranged in the most realistic way possible. A clear-cut event has the highest priority, followed by human thinning and finally natural mortality including self-thinning. If, for example, a forest is scheduled to be clear-cut, the entire forest biomass is subjected to the rules of the clear-cut and no other mortality occurs in that time-step.

In addition to forest management and natural prescribed mortality, a variety of changes have been 625 made to processes involving vegetation mortality. A whole PFT within a grid-cell is now killed if, at the end of the day, the labile pool is empty and there is no carbon available in the leaf or carbohydrate reserve pool to refill it. In this situation, it will be impossible for the plant to assimilate new carbon from the atmosphere as it will not be able to grow new leaves and thus initiate plant recovery. Furthermore, a forest can die if the density falls below a certain prescribed value. In the 630 next time step a new young forest will be prescribed.

If a forest is thinned, it is assumed that the weakest trees will be thinned, and therefore human thinning reduces or even eliminates the natural mortality for that time-step. Natural mortality still happens on a daily time-step, while human-induced mortality happens only at the end of the year. Self-thinning, as described below, takes priority over environmental mortality which is the mortality of individuals by insects, lightening, wind, drought, frost and heart rot. Environmental mortality is

- 635 of individuals by insects, lightening, wind, drought, frost and heart rot. Environmental mortality is calculated by multiplying the stand biomass by an assumed mortality fraction of $1/k_{t_{resid}}$. Where self-thinning is less than this assumed environmental mortality, self-thinning is complemented by additional mortality to reach the set environmental mortality. Where self-thinning mortality exceeds the set environmental mortality, simulated self-thinning is assumed to include environmental mortal-
- 640 ity. The fire module that is available for the trunk but not for the ORCHIDEE-CAN branch simulates stand replacing fires rather than individual-tree based mortality due to lightening. The approach implemented in the ORCHIDEE-CAN branch could therefore be extended with models that simulate stand replacing mortality from fire, insects and storms.
- The use of circumference classes adds a good deal of realism and flexibility to the ORCHIDEE-645 CAN simulations, but it also raises additional questions. For example, which trees should be targeted by which mortality? Given that self-thinning reflects the outcome of continuous resource competition, the largest trees are expected to be most successful when competing for resources, and therefore we assume that the smallest trees die first to reduce the stand density. Conversely, larger trees are more likely to die because of environmental stress factors, being more prone to cavitation, wind
- damage, lightening, and, heart rot. Therefore, we select more older trees to die from environmental

mortality. While doing this also trees in the other circumference classes were killed based on the following recursive definition (cf ?):

$$f_{death}^{icir} = \frac{f_{death}^{icir-1} \times k_{ddf}^{1-(k_{ncirc}-1)}}{m_{Ndeath}}$$
(38)

Where k_{ddf} is the death distribution factor, which is the factor by which the smallest and largest circumference classes differ (e.g., $k_{ddf} = 10$ means that the largest circumference class will lose ten times as much biomass as the smallest as a result of the mortality), m_{Ndeath} is a normalization factor so that sum of f_{death}^{icir} is unity, and f_{death}^1 is set equal to unity before normalization. As the stands are very close to even-aged, we set the factor k_{ddf} to be equal to 1. This means the same number of trees is killed in each circumference class. If, for some reason, there is not enough biomass in a 660 given class to satisfy this distribution, the extra biomass is taken from the next smallest class (in the case the smallest class does not have enough, it is taken from the largest class).

Related to mortality is the question of the circumference class distribution. As mentioned above, trees in different circumference classes are preferentially killed by different processes. If the sim-

ulation is long enough (or if the morality is aggressive enough), eventually the number of trees in

- some circumference classes may become zero. This would reduce the numerical resolution of the allocation scheme. When only one circumference remains populated, the scheme effectively loses its meaning as all the newly produced biomass is now be allocated to the only remaining circumference class. In order to maintain the same level of detail through the simulation, the distribution of all the circumference classes is recalculated at the end of each day. A normalized target distribution is
- 670 specified as an input parameter (an exponential distribution is currently used), and this distribution is scaled to produce a target distribution for the current number of individuals. All of the current individuals are placed in these new classes until the target distribution is satisfied. The target distribution now contains, however, trees of multiple sizes, so we need to average them to find the new "model" tree for each class. By changing the size of the model tree in each class, we are able to preserve the
- 675 total biomass of the stand as well as the total number of individuals. Note that the boundaries of each diameter class are recalculated at each time step, this approach is a numerically efficient alternative to fixing the boundaries of each diameter class with a varying distribution.

4 Description of the parametrization

680

The ORCHIDEE-CAN branch was specifically developed to quantify the climate effects of forest management over Europe. Although the developments are sufficiently general to be applied outside of Europe, the model was initially parametrized for the boreal, temperate and Mediterranean climate zones and validation focused on Europe. Parametrization of the tropical zone is subject of a follow-up study. The parametrization of the model, including parameter optimization and tuning consisted of five major steps:

- Parameters related to carbon allocation (section 4.2), forest management and mortality (section 4.3, hydraulic architecture (section 4.4, canopy structure (section 4.5)), photosynthesis (section 4.6), and canopy radiation transfer (section 4.7) and for which observations exists at the species-level (section 4.1) were extracted from a wide range of sources (Tables S1-S5). Using the extracted species-level parameter values in ORCHIDEE without further processing avoids hidden model-tuning and largely reduces the likelihood that simulation results are biased by hidden calibration owing to a poor taxonomic definition of PFTs (?).
 - 2. The phenology-related parameters of the deciduous MTCs were optimised by ?, using MODISderived NDVI data normalised to model fAPAR over the 2000 - 2008 time period.
 - 3. The modulator (m_{ψ}) which accounts for processes in the the soil-plant continuum that are currently not modelled, was manually tuned against species distribution maps (section 4.4).
 - The coefficient for maintenance respiration was optimized making use of Bayesian calibration (section 4.8) against a compilation of 100+ observations of biomass production efficiency.
 - 5. The leaf to sapwood area ratio was manually tuned (section 4.9) to match 100+ site-level GPP and LAI observations recorded over Europe.

700 4.1 Introducing twelve new PFTs

Similar to the ORCHIDEE trunk, the ORCHIDEE-CAN branch distinguishes 13 metaclasses (MTC) for vegetation. Outside Europe the original MTC classification of ORCHIDEE was kept, while inside Europe 12 new parameter sets representing the main European tree species were added. The default vegetation distribution map in ORCHIDEE, i.e., **?**, was replaced by an up-to-date global

- 705 MTC map which has been produced using the ESA CCI ECV Land Cover map (http://www.esalandcover-cci.org/)(?).The mapping from land cover to MTC basically followed ?, although table 5 (the "cross-walking" table) has been updated following discussions with the LC-CCI team at Universite Catholique de Louvain. For the European domain, the global MTC distribution was overlaid by a tree species distribution map (?).
- 710 This study focusses on tree species with a coverage of more than 2% in Europe, yielding seven species groups covering in total 78.8% of the European forest area: *Betula sp.,Fagus sylvatica, Pinus sylvestris, Picea sp., Pinus pinaster, Quercus ilex* and a group combining *Quercus robur* and *Quercus petraea*. For *Pinus sylvestris, Picea sp.* and *Betula sp.* An additional distinction between boreal and temperate forest was made for the species map and parametrization: trees located in Norway,
- 715 Sweden and Finland were considered boreal, while trees growing at lower latitudes were categorized as temperate. Given the potential role of tree species of the Salicacea genus in short rotation coppice management, a separate PFT was parametrized for *Populus sp.* Furthermore, to improve the parametrization of the MTC of boreal needleaved deciduous forest, observations from *Larix sp.* were included when possible.

695

- For these 12 forest species, 12 new PFTs were created with each PFT belonging to a single MTC (Table S2, S3 and S4) Almost 79% of the European forest was parametrized at the species level. The remaining 21% was reclassified in four residual groups, i.e., a temperate and boreal needleleaf evergreen and a temperate and boreal broadleaved residual group. For use outside Europe, the original MTC classification of ORCHIDEE was kept. The parameters of the residual groups and MTCs
- 725 are the mean of the parameters of the species-level PFTs that are in the MTC, with the exception of albedo parameters that could be extracted from remote sensing products. Finally, separate PFTs were introduced for boreal grasses and croplands, which allowed for a boreal parametrization of phenology, senescence and growth. This approach, which distinguishes a total of 28 PFTs, allows a higher taxonomic resolution over Europe, better defines forest types compared to the more general

730 MTC approach and facilitates the use of observations to derive parameters.

4.2 Allocation

The allocation scheme relies on the leaf to sapwood area ratio (section 4.9) and the relationship between diameter and height. Following a logarithmic transformation of the more than 150,000 data points from the the national forest inventory data of Spain, France, Germany and Sweden, the two

parameters (i.e. k_{α_1} and k_{β_1}) describing the relationship between diameter and height (equation 15) were fitted at the species-level making use of a least square regression. Parameter values for MTCs were derived by grouping the species into MTCs and fitting the parameters. Data sources and parameter estimates are presented in Tables S2 and S3.

4.3 Forest management and mortality

- Forest management and tree mortality are controlled by (section 3.7): (1) maximum tree diameter (no symbolic notation; called largest_tree_diam in ORCHIDEE-CAN), (2) minimum stand density (no symbolic notation; called ntrees_dia_profit in ORCHIDEE-CAN), (3) environmental mortality (no symbolic notation; called residence_time in ORCHIDEE-CAN), (4) self-thinning (k_{α_2} and k_{β_2}) and, (5) anthropogenic thinning (no symbolic notation; called alpha_RDI_upper, alpha_RDI_lower,
- 745 beta_RDI_upper and beta_RDI _lower in ORCHIDEE-CAN) where the parameters depend on the management strategy.

Maximum tree diameter was extracted from the French, Swedish, German and Spanish forest inventories as the observed 50% quantile for diameter at breast height. The 50% quantile rather than the observed maximum was used to account for the fact that large scale land surface models are

750 expected to reproduce large scale patterns rather than local extremes. Minimum stand density was estimated as the expected stand density for the maximum tree diameter for a stand under self-thinning. Although both criteria are related to each other through the observed self-thinning relationship (see below), the minimum number of trees is used to decide when unmanaged forests should be replaced, whereas both the maximum diameter and the minimum number are used for managed sites as criteria

755 to initiate a clear cut. Parameters for anthropogenic thinning are based on the national forest inventory data and checked against the JRC-database of species specific yield tables. Parameter values are presented in Table S5. Resource competition between trees in the same stand has been reported to result in the so-called self-thinning relationship that relates the number of individuals within a stand to the stand biomass (???):

760
$$(M_s + M_h) \times k_{\rho s} = k_{\alpha} \times (d_{ind})^{-k_{\beta}}$$
(39)

Where k_{α} and k_{β} are the constants of the self-thinning relationship. Furthermore, stem volume can be written as a function of tree diameter (d_{dbh}) , tree height and stem form factor $(k_{\alpha'})$ to account for the fact that the stem shape is not a perfect cylinder:

$$(M_s + M_h) * k_{\rho s} = k_{\alpha'} \times (d_{dbh})^2 * d_h \tag{40}$$

Following the allometric relationship given in equation 15, tree height can be written as a function of tree diameter. Hence, the self-thinning relationship can be re-written to relate stand diameter to stand density:

$$d_{dbh} = k_{\alpha 2} \times (d_{ind})^{-k_{\beta 2}} \tag{41}$$

Where, $k_{\beta 2}$ relates to $k_{\beta 1}$ (as in equation 15) as follows:

770
$$k_{\beta 2} = -3/2 \times (2 + k_{\beta 1})$$
 (42)

 $k_{\alpha 1}$ and $k_{\beta 1}$ were estimated by fitting equation 15 to observed diameter and height of individual trees from NFI of Sweden, Germany, France and Spain. $k_{\beta 2}$ was calculated from equation 42 and $k_{\alpha 2}$ was estimated by fitting equation 41 to observations of the quadratic mean stand diameter and stand density from NFI data.

775 4.4 Hydraulic architecture

Initial choices of parameters for this scheme were based on the values and parameter sources listed by **?**. All data sources were revisited and the search was extended to obtain values at the PFT rather than MTC level. Given that plant hydrology is rather well studied, observed parameters were available for most of the species. Data sources are listed in Table S1 whereas the parameter values are shown in Table S3. Our implementation of hydraulic architecture required the introduction of a tun-

shown in Table S3. Our implementation of hydraulic architecture required the introduction of a tuning parameter (m_{ψ}) to account for processes that are currently absent in the scheme, e.g. plant water storage and soil-root resistance. A process-based description of these processes (i.e., ??) is being tested and should reduce the effect of the tuning parameter and eventually allow its removal from the model.

- For the time being, the modulator m_{ψ} was tuned manually against the species distribution map to obtain a match between the simulated and observed species distributions. When the modulator is set to zero, all PFTs experience excessive water stress resulting in large-scale plant mortality. The modulator was increased until the prescribed vegetation distribution which was based on remotesensing observations (section 4.1), survived where it was prescribed. To this aim, the model was run
- 790 for 50 years, forced with v5.2 of the CRU-NCEP climatology for Europe (Climatic Research Unit, University of East Anglia). Note that the values of the modulator depend on the climate data that are used to force the model. Similarly the modulators may need to be re-tuned when ORCHIDEE-CAN is coupled to an atmospheric model.

4.5 Canopy structure

795 The relationship between diameter and projected crown surface area follows the model proposed by **?**:

$$d_{csa} = k_{ap} \times d_{dbh}^{\kappa_{bp}} \tag{43}$$

with parameters estimated using the dataset presented in ?. This dataset contains diameter and projected crown surface areas observations for over 37,000 individual trees in Europe covering almost
30 species. Following logarithmic transformation of the observations a linear least square regression was used to fit species-specific parameter values. Parameter values are shown in Table S2. Parameter values for MTCs were derived by grouping the species into MTCs and fitting the parameters. No observations were available for the boreal zone and temperate evergreen deciduous species. For the boreal species a subset of the temperate observations (*Pinus sylvestris*, *Picea abies* and *Betula pendula*) was used, i.e., the relationship between d_{csa} and d_{dbh} was fitted to all available data for *Pinus sylvestris*. Next, all observations with a d_{csa} that falls below the predicted d_{csa} were selected as considered to represent a boreal subset. Given the importance of snow pressure on crown structure,

selecting observations with sub average d_{csa} is justifiable as a first approximation. Subsequently, the parameters were fitted to this subset of data. For *Quercus ilex* no data were available and parameters
were tuned such that the crown diameter was 0.85 m less than the tree height.

4.6 Analytical solution for photosynthesis

815

Three originally MTC-specific photosynthetic parameters (k_{Vcmax} , k_{Jmax} and k_{sla}) were derived at the species level by obtaining weighted site means for each species from the global leaf trait database TRY (?) and additionally from ?. Only k_{Vcmax} and k_{Jmax} standardized to a common formulation and parametrization of the photosynthesis model by (?) were used. Most k_{Vcmax} and k_{Jmax} values in the TRY database had already been standardized to a reference temperature of 25°C (?). Subsequently, a species-specific $k_{Jmax,opt}/k_{Vcmax,opt}$ ratio was calculated from the records which included both $k_{Vcmax,opt}$ and $k_{Jmax,opt}$ measurements. From this ratio, which was within a range of 1.91-2.47 for each species, $k_{Jmax,opt}$ was calculated for records which originally only included

820 k_{Vcmax} . Only geo-referenced observations within Europe were used and the distinction between boreal and temperate forest was made similar to the species map. Depending on the species this resulted in 5 to 183 observations for k_{sla} and 11 to 173 observations for $k_{Vcmax,opt}$ and $k_{Jmax,opt}$. From these observations species-specific means were calculated, weighted for differences in the number of observations per site. The parameter values are shown in Table S3.

825 4.7 Multi-layer two-way radiation scheme for tall canopies

The radiation transfer scheme makes use of parameters describing leaf and background properties, i.e., leaf single scattering and prefered scattering direction (for both visible (VIS) and near-infrared (NIR) wavelengths) and the so-called background albedo or the albedo of the surface below the dominant tree canopy (VIS and NIR). All parameters were taken from the Joint Research Cen-

- 830 tre Two-stream Inversion Package (JRC-TIP) (??). This is a software package (?) which inverts a two-stream model (?) to best fit the MODIS broadband visible and near-infrared white sky surface albedo from 2001 to 2010 at 1 km resolution (?). The inverse procedure implemented in the JRC-TIP is shown to be robust, reliable, and compliant with large-scale processing requirements (?). Furthermore, this package ensures the physical consistency between sets of observations, the two-stream
- 835 model parameters, and radiation fluxes.

Only parameter values for which the posterior standard deviation of the probability density functions were significantly smaller than the prior standard deviation were selected from the JRC-TIP optimization (?), since this condition ensures statistically significant values. Species and MTC specific values were derived from JRC-TIP by performing a multiple regression. This methods determines,

- 840 in an objective way, how the fractions of each MTC or species explain the JRC-TIP parameter. The multiple regression was performed separately for the six parameters: the single scattering of leaves (for both VIS and NIR), the scattering direction of leaves (VIS and NIR) and the background albedo (VIS and NIR). Each JRC-TIP parameter was used as the dependent variable and the independent variables consisted of the fractions of each MTC (?) or species (?). These fractions were used to
- 845 find a linear function that best predicted each JRC-TIP parameter. The corresponding slope of a regression of each MTC or species fraction gives the MTC or species dependent JRC-TIP value. The multiple regression was performed without an intercept. To avoid pollution by the seasonal cycle, the multiple regression was applied only for the pixels of the Northern Hemisphere. Only pixels that were less than 10% covered by non-vegetative fractions where selected for the analysis and only
- 850 significant results following an F-test and positive r²-values were selected. The derived parameter values are shown in Table S4.

4.8 Maintenance respiration

Both the trunk and ORCHIDEE-CAN branch reduce the definition of net primary production to biomass production; hence, carbon leaching from the roots, volatile organic emissions from the

- leaves, dissolved and particulate carbon losses through water fluxes and carbon subsidies to mycorryhzae are not accounted for in the model. These fluxes are (incorrectly) accounted for in the modelled autotrophic respiration. Modelled autotrophic respiration should therefore be considered an effective rather than a true value. For this reason, the basal rate of autotrophic respiration was optimized against 126 site observations of the biomass production efficiency (k_{cmaint}) calculated
- as the ratio between annual biomass production and annual photosynthesis (??), using a Bayesian optimization scheme. The scheme, for which more details are given in ?, uses a standard variational method based on the iterative minimization of a cost function that measures both the model data misfit and the parameter deviations from prior knowledge (?).
- The simulations that were used in the Bayesian optimization prescribed a 20 m tall vegetation for temperate tree species, a 15 m tall vegetation for boreal tree species and a 10 m tall vegetation for Mediterranean tree species as its initial condition. This approach reduced the need for several decades of simulations to a single year to grow a mature forests. In total the simulations were run for 10 years and covered the European domain. The first year was discarded and the ratio between modelled GPP and NPP was averaged over the remaining 9 years. Prior to the optimization the
- observations were averaged for agricultural PFTs (0.57), and deciduous (0.44) and evergreen (0.53) forest PFTs, the observed uncertainty was 0.03. The parameter values were set to range between 0.0032 and 0.160. The optimization converged within 11 iterations and the optimized parameter values are shown in Table S2.

It remains untested how well the simulated effective autotrophic respiration represents the (rarely) observed autotrophic respiration. Note that in the cases of both the trunk and the ORCHIDEE-CAN branch of ORCHIDEE, a match between effective and observed autotrophic respiration should not be interpreted as evidence of desired model behaviour because several components of net primary production are not modelled yet.

After the optimization of the maintenance respiration coefficient (k_{cmaint}) , the model simulates reasonable biomass production efficiency for a unit of photosynthesis. Hence, the final step of the parametrization focussed on optimizing the leaf area as this is one of the main divers of photosynthesis.

4.9 Sapwood to leaf area ratio

The vegetation structure simulated by the ORCHIDEE-CAN branch is sensitive to the value of k_{ls}
which describes the ratio between the leaf and sapwood area of an individual tree. The available observations show a wide range within and across forest species. Dependencies of k_{ls} on tree height

(??), tree diameter following stand thinning (?) and CO_2 (?) have been reported. Most observations, however, come from experiments where time was substituted by space which hampers teasing apart the sources of variability. Given the variation and uncertainty in the observations and the

890 n

model sensitivity to this parameter, we manually tuned its value within the observed range, to match European-wide observations of leaf area index as recorded in the Database of Global Forest Ecosystem Structure and Function **?**.

This database was used to calculate a mean and maximum observed leaf area index at the specieslevel for the temperate and boreal region. Initially 20-year long European-wide simulations were

- used to simulate leaf area index of a species, when the large-scale leaf area index approached the mean target value and did not exceed the maximum value, the simulations were extended to reach 100 years for checking the temporal evolution of leaf area index. We deliberately optimized the sapwood to leaf area ratio (k_{ls}) by making use of stand-level data to reduce circularity with the model validation (see below).
- 200 Limited tests over a period of 100 years in a Scots pine forest at $51 52^{\circ}$ N, $13 14^{\circ}$ E (Fig. S1) suggested that optimizing k_{cmaint} and k_{ls} had the largest effect on the maximum LAI, which decreased with almost 17% after optimization compared to a simulation with prior parameter values. Mean annual GPP, mean annual transpiration and basal area decreased with, respectively 6, 6 and 7% compared to a simulation with prior parameter values (Fig. S1).

905 5 Validation

ORCHIDEE-CAN is designed as the land surface model to be coupled to the atmospheric model LMDz. As such future applications of ORCHIDEE-CAN are expected to be regional to global in the spatial domain and to span several years in the temporal domain. Given its anticipated uses, the ability of the model to reproduce large-scale spatial patterns as well as their inter-annual variability

- 910 is essential. The first applications of the model, both offline and coupled to the atmosphere, will focus on Europe. The validation, therefore, reports performance indices both over Europe as over eight separate regions within Europe (?). These eight regions, which partially overlap, are defined after ?. Furthermore, the performance indices are calculated for winter, spring, summer and autumn and thus allow to evaluate the capacity of the model to reproduce observed annual cycles.
- 915 In addition to the root mean square error, a land performance index (LPI) based on the principles laid out for the Climate Performance Index (?, their SI) was also calculated. LPI normalizes the root of the squared differences between the simulations and observations by the observed spatial and temporal variance. The LPI was used to estimate the likelihood that the simulated variable belongs to the same population as the observed variable, defined as exp(-0.5LPI²). An LPI equal to 1 indicates
- 920 that the model correctly reproduces the mean observed value and implies a likelihood of 61% (?) that the simulations and observations come from the same population. Similarly, an LPI of 2 reduces

this likelihood to 13%. An LPI of less than 0.32 has a likelihood of more than 95% and therefore indicates a statistically significant result.

- While developing ORCHIDEE-CAN, the numerical approaches that added functionality to the code were selected on the basis of their performance at the site-level (see below). Rather than running the same site-level tests for our implementation, we performed a complementary large-scale validation. The strength of our approach lies not in the details, as is the case for site-level validation, but in its width by simultaneously testing model performance for structural variables such as basal area (?), canopy structure (?) and canopy height (?), biogeochemical fluxes such as GPP (?),
- 930 biophysical fluxes such as albedo (?) and fluxes at the interface of biogeochemistry and biophysics such as evapotranspiration (?). The selection of variables was limited by the availability of spatially explicit data-derived products for Europe.

For the validation, both the trunk and ORCHIDEE-CAN branch were run from 1850 to 1900 using CRU-NCEP climate forcing from 1901-1950 at 0.5 degree resolution. From 1901 until 2012,

- 935 the corresponding CRU-NCEP forcing data for each year were used. Both versions used the 11layer soil hydrology, the single-layer energy budget and the same land cover map (?). Given that no European-wide, spatially explicit and data-derived products were found for the validation of the net carbon flux, there was no need for a carbon spin-up. For the ORCHIDEE-CAN branch, the observed tree height and basal area were compared against the simulation values at the end of 2010 (the
- 940 trunk does not simulate these variables). For both the trunk and the ORCHIDEE-CAN branch, the observed GPP, evapotranspiration, effective LAI and VIS and NIR albedos were compared against monthly means between 2001 and 2010.

5.1 Species versus PFTs

In ORCHIDEE-CAN the PFT concept was refined by parametrizing the main European tree species groups (section 4.1). To evaluate the effect of the species parametrization we performed a companion simulation for the configuration described above, but at the MTC level. Model performance was barely affected by the use of the MTC parameters, compared to the simulation with the species parameters (Fig. S2 for RMSE scores).

5.2 Allocation

- 950 In ORCHIDEE-CAN, functional relationships which vary by species and light stress are used to allocate carbon among the fine roots, foliage and sapwood. The allocation scheme largely follows ?, who in turn was inspired by ?. Approaches simulating allocation based on functional relationships were found to outcompete allocation schemes based on constant fractions or resource limitation (?). The ability of these schemes to reproduce foliage, fine root and sapwood reported in large observa-
- tional data sets (for example, ?) demonstrates that these schemes capture the main observed features(?). In addition, allocation schemes making use of functional relationships were also capable of

simulating the observed effect of elevated CO_2 on two mature forest ecosystems (?). Despite these successes, the schemes were reported to be sensitive to their parametrization. Differences in parameters were reported to result in substantial differences in the simulated allocation. The parameters for

960 the functional relationships used in ORCHIDEE-CAN are given in Table S2. The main conceptual difference between the allocation scheme by **?** and ORCHIDEE-CAN is that the latter was designed

to simulate one or more diameter classes. Given that photosynthesis is still calculated at the stand level (and thus not at the tree level)

the allocation rule of **?** was integrated in the functional allocation scheme to account for light and resource competition within a stand. Where the functional relationships are used to simulate carbon allocation within an individual tree of a given diameter, the rule of **?** allocates carbon across the different diameter classes. The allocation rule which models the radial increment for individual trees in pure even-aged stands was successfully tested for Norway spruce and Douglas fir stands in France

(?). A similar approach for modelling radial increment has already been implemented in a version
970 close to the trunk of ORCHIDEE (?) and was able to successfully simulate stand characteristics such as baight basel area and stand diameter (2). This provides implementation differs from the current

- as height, basal area and stand diameter (?). This previous implementation differs from the current implementation in its time resolution (which is now daily instead of yearly), its analytical solution and the underlying allocation scheme (which is now based on functional relationships instead of resource limitation).
- The aforementioned studies performed a detailed validation of the two approaches dealing with carbon allocation, which were combined in ORCHIDEE-CAN. Complementary to these studies we performed a European-wide validation of our implementation and parametrization of these well-tested schemes against a remote-sensing based map of tree height (?), upscaled eddy-covariance observations for GPP (?) and a map of basal area based on national forest inventory data (?). The
- 980 model's ability to reproduce GPP is thought to reflect its capacity to simulate the foliage biomass, a correct simulation of height reflects the model's capacity to simulate aboveground woody biomass and its capacity to reproduce observed basal areas suggest that the interaction of stand density and individual tree diameter are well-captured.

The new implementation and parametrization of the within-tree and within-stand allocation schemes

- 985 were found to have an 91, 68 and 72% chance that the simulations reproduced the observations for GPP, tree height and basal area for Europe, respectively (Table ??). Given that basal area and height are not available from the trunk version of ORCHIDEE, we could not compare the performance of model versions in this respect. With respect to GPP, the ORCHIDEE-CAN branch was found to outperform the trunk by 12% and thus increased the likelihood that ORCHIDEE-CAN is an unbiased
- 990 simulator of the spatial and temporal variability of GPP from 79 to 91%. Improved performance of the ORCHIDEE-CAN branch compared to the trunk is observed for all regions in summer where the RMSE of GPP was halved from 2.5-5 gC m⁻² day⁻¹ to 1-2 gC m⁻² day⁻¹ (Fig. ??, ?? and ??).

Although part of the high likelihood could be due to the fact that the observed GPP was upscaled making use of similar climatologies being used as the forcings of the models, this circularity could

995 neither have contributed to the improved performance between the trunk and the ORCHIDEE-CAN branch nor to the decrease in RMSE. The improvements are thought to be due to structural changes to the model such as allocation, hydraulic architecture and canopy structure as well as to the use of more consistent parametrization.

5.3 Plant water supply

1000 Our implementation of plant hydraulic architecture was largely based on the scheme of ?, which was tested globally and at site level. Global simulation results for actual evapotranspiration were found to reproduce available data (??). At the site level, the model agreed well with the magnitude and seasonality of eddy-covariance measurements of actual evapotranspiration for 15 European forests sites (EUROFLUX), with a tendency to slightly overestimate actual evapotranspiration for 6 sites 1005 (?).

The maximum amount of water that can be transported by a tree relies on the hydraulic architecture of the tree and therefore on the capacity of the model to simulate tree and stand dimensions as well as on the model's capacity to simulate soil water content. As an additional test, our implementation of the model was compared against the upscaled eddy-covariance measurements for GPP

- 1010 and actual evapotranspiration (?). The capacity to jointly reproduce GPP and actual evapotranspiration is an indicator that the model successfully reproduces the coupling between CO₂ and water exchange. Model validation showed 91 and 87% chance (compared to 79 and 45% for the trunk) that ORCHIDEE-CAN reproduces the upscaled GPP and actual evapotranspiration data (Table ??, Fig.??). The RMSE for actual evapotranspiration during summer dropped well below 1 mm day⁻¹
- 1015 for most regions (Fig. ??), whereas it never dropped below 1 mm day⁻¹ for the trunk (Fig. ??).

5.4 Canopy structure

The canopy structure model by **?** was previously validated against ground-based LIDAR data for several test sites with varying density, structural complexity, layering and clumping (**?**). Model-derived canopy gap probabilities compared with observations using a one-sample t-test were significant for

- 1020 11 out of 12 test sites. We considered this result as a sufficient proof to use this canopy structure model in the ORCHIDEE-CAN branch and added to its validation by comparing the simulated canopy structure model over Europe against a remote-sensing based map of tree height (?) and the JRC-TIP effective LAI product (?). The effective LAI value expresses the capability of the canopy to intercept direct radiation, and is thus associated with the probability distribution function of the
- 1025 canopy gaps (?). Thus the effective LAI contains information about the forest structure and leaf distribution of the canopy. In the ORCHIDEE-CAN branch, canopy structure is used to calculate

the albedo, roughness length, absorbed light for photosynthesis and leaf area that is coupled to the atmosphere for, e.g., transpiration and interception of precipitation.

The ORCHIDEE-CAN branch is the first branch of ORCHIDEE that makes use of an effective 1030 LAI to calculate the interaction between the canopy and the atmosphere. The LPI and RMSE of the branch, therefore, cannot be compared against the trunk. Overall, the combined implementation of the allocation scheme and the canopy structure model shows a 67% chance to reproduce the satellite-based estimates for effective LAI. Surprisingly, effective LAI is better simulated in spring and autumn when dynamics within the canopy are substantial due to leaf on-set and senescence.

1035 For the periods when the effective LAI is expected to be most stable, i.e., summer and winter, LPI approached and frequently exceeded 1 (data not shown). Part of this shortcoming may be due to the lack of shrubs in the land cover classification. In the model, shrublands are replaced by forest and/or grasslands, likely resulting in differences between the observed and simulated canopy structure. This lapse also appears in the RMSE of effective LAI (RMSE higher than 0.8, Fig. ??)

1040 5.5 Top of the canopy albedo

The radiation transfer model (?) has been validated extensively against realistic complex threedimensional canopy scenarios (?) and as part of the RAdiation transfer Model Intercomparison (RAMI) project. The 1-D canopy radiation transfer model by ? was demonstrated to accurately simulate both the amplitude and the angular variations of all radiant fluxes with respect to the solar

1045 zenith angle (?). In addition, the radiation transfer model and its effective values extracted from the JRC-TIP data set were successfully applied to a single forest site (?).

Previously we reported on the capacity of the radiation transfer model to simulate the effects of forest management on albedo (?). For the latter, forest properties were prescribed and the radiation transfer model was validated against top-of-the-canopy albedo data from five observational sites.

- 1050 Differences in the spatial scales between the observed and simulated albedo values were accounted for by presenting the mean June albedo during 2001-2010 (?). The simulated summertime canopy albedo falls within the range of observation. However, there occurs a slight overestimation in the near-infrared wavelength band compared to the single site measurement. Too high near-infrared single scattering albedo values for pine, as obtained from the JRC-TIP product, are the most likely
- 1055 cause. The observed deviation is not due to a shortcoming in the model itself but reflects the difficulties the JRC-TIP has with optimizing parameter values in the absence of field observations in the specific case of sparse canopies (?).

For the spatial validation we use the white-sky albedo (VIS and NIR) from Moderate Resolution Imaging Spectroradiometer (MODIS, ?) at 0.5° resolution (distributed in netCDF format by

1060 the Integrated Climate Data Center (ICDC, http://icdc.zmaw.de) University of Hamburg, Hamburg, Germany). Over large spatial and temporal domains the ORCHIDEE-CAN branch reproduces the observed VIS and NIR albedo and its variability; LPI for the albedo in the visible light is especially satisfying with a likelihood of 92% for the simulations to come from the same population as the observations (Table **??**). This high overall performance index, however, hides performance issues

- 1065 over Scandinavia and the Alps during the snow season. The RMSE for VIS and NIR albedo without snow lies around 0.05, whereas during the snow season the RMSE increases to 0.20 (VIS) and 0.18 (NIR) over these regions (Fig. ??). When the ORCHIDEE-CAN branch is coupled to an atmospheric model, however, these deviations will only have a minor effect on the climate, owing to low incoming radiation during most of the snow season, especially in Scandinavia.
- 1070 Previous validation of the radiation transfer model showed that the largest discrepancies were occurring in the near-infrared domain with a snow covered background (?). With the exception of the snow-covered season, the new albedo scheme, that relies on the simulated canopy structure, resulted in a substantial improvement of 0.05-0.15 compared to the trunk for the RMSE in both the VIS and NIR range in Scandinavia and the Alps (Fig. ?? and ??). The European LPI-based likelihood
- 1075 that our model simulations come from the same populations as the MODIS albedo increased by a remarkable 11 and 23% for, respectively, NIR and VIS albedo (from 61% and 69% for the trunk to 72 and 92% for the ORCHIDEE-CAN, Table ??).

Given that the parametrization of the canopy radiation transfer model used in ORCHIDEE-CAN relies on MODIS, the high likelihood may not come as a surprise. However, our implementation of

- 1080 the radiation transfer model also relies on the simulated absorbed light, simulated GPP, simulated allocation and simulated canopy structure (which depends on mortality and forest management). In the absence of all these processes our canopy radiation transfer model is expected to reproduce the MODIS data with a probability of 100%. Hence, the likelihood of 72% and 92% (for NIR and VIS, respectively) could also be interpreted as a verification of the aforementioned calculations; all
- 1085 calculations that determine the canopy structure reduce the reproducibility of the data by only 8-28% (100% to 72 or 92%).

5.6 Energy fluxes

The multilayer scheme is in the process of a detailed evaluation across a range of tests conditions (?), and further validation across a range of sites is on-going. The scheme is able to produce within-1090 canopy temperature and humidity profiles, and successfully simulates the in-canopy radiation distribution, as well as the separation of the canopy from the soil surface. However, in order to preserve a measure of continuity with previous evaluations of the model, the multilayer solution is here set to single layer operation mode, which includes the effects of hydraulic limitation (section 3.2) and canopy structure (section 3.3) on the energy budget.

1095 The single-layer set-up of the multi-layer solution makes use of an improved albedo estimation and is therefore expected to better simulate the net radiation that needs to be redistributed in the canopy. This has been confirmed at a single site with a sparse canopy (?). Furthermore, the improvements in actual evapotranspiration in addition to the low RMSE (Fig. ??) are expected to be propagated in the performance of the energy budget.

1100 5.7 Forest management strategies

Model comparison has previously demonstrated that explicitly treating thinning processes is essential to reproduce local and large scale biomass observations (?). This finding justifies the implementation of generic approaches to forest management despite the difficulties associated with defining and quantifying forest management and its intensity (?). Although the use of so-called naturalness

- 1105 indices, in which the current state of the forest in referenced against the potential state of the forest, has been criticised because of difficulties in defining the potential state of the forest (?), such approaches were demonstrated to correctly rank different management strategies according to their intensity (?).
- Naturalness indices making use of only diameter and stand density or the so-called Relative Den-1110 sity Index (RDI) have been previously implemented at the stand-level (?) and as well as in large scale models (?). This approach was shown to successfully reproduce the biomass changes during the life cycle of a forest (??). The implementation of a forestry model based on the relative density index was reported to perform better than simple statistical models for stand-level variables such as stand density, basal area, standing volume and height (?). Although the performance of the model was
- 1115 reported as less satisfying for tree-level variables, the approach is nevertheless considered reliable to model the effects of forest management on biomass stocks of forests across a range of scales from plot to country (?).

In the absence of forest management, ORCHIDEE-CAN simulates that the stands develop into tall canopy (Fig 5.7a), with a high biomass (Fig 5.7b), a substantial dead wood and litter pool (Fig

- 1120 5.7c) and no harvest (Fig 5.7d). High stand management reduces the height, standing biomass and litter pools (Fig 5.7a-c) but produces biomass for harvest (Fig 5.7d). Under coppicing, the reduction in forest age is reflected in a shorter canopy and lower biomass and litter pools (Fig a-c) compared to high stand management. The harvest is more evenly spread in time but falls below the harvest generated by high stand management (Fig 5.7d). Given the shorter rotations, canopy height, standing
- 1125 biomass and litter pools are lower for short rotation coppicing with poplar and willow compared to all other management strategies applied on oak forest (Fig 5.7a-c). Short rotation coppice was harvested every 3 years resulting in a quasi-continuous supply of woody biomass (Fig. 5.7d).

The forestry model implemented in ORCHIDEE-CAN is based on the RDI approach by ?. We complemented earlier validation of such an approach over France (?) by a new European-wide vali-

1130 dation for basal area. At the European scale we verified the simulated basal area and height against observed basal area from national forest inventories (?) and height from remote-sensing (?). With an RMSE of 3-7 for height and 7-15 for BA, and a chance of, respectively, 68 and 72% to reproduce the data at the European scale (Table ??), our model is capable of correctly simulating the mean height and basal area but fails to capture much of the spatial variability (Fig. **??**, temporal variability was 1135 not considered because the data products were only available for one time period).

Furthermore, we evaluated basal area and tree diameter at the species-level for 11 regions over France which represents a finer spatial-scale than targeted by the model developments and their parametrization. The data were extracted from the French forest inventory between 2005 and 2010 and we used the same simulations as for the European validation in the previous paragraph. We

- 1140 selected pixels included in the French inventory data and for both simulations and observations we calculated a moving average for the diameter and basal area per age class to then calculated the RMSE (Fig. ??). To account for intrinsic species differences in diameter and basal area we normalized the RMSE. The normalized RMSE was lower than 30% of the mean tree diameter or mean basal area for each region for *Betula sp.*, *Pinus pinaster* and *Quercus ilex*. For *Fagus sylvatica*,
- 1145 *Pinus sylvestris, Picea sp.* and *Quercus robur/petraea* the normalised RMSE of diameter and basal area exceeded 50% for one to four regions for tree diameter and basal area (not shown).

The inability to fully capture the observed spatial variability in the simulation could be due to the simulation protocol that started in 1850 with 2 to 3 meter tall trees all over Europe. A longer simulation accounting for the major historical changes in forest management such as the reforestation in

- 1150 the 1700s following an all time low in the European forest cover, the start of high stand management at the expense of coppicing in the early 1800s, and the reforestation programs following World War II (?) is expected to improve the spatial variability in tree height and basal area. Regional deviations such as those observed in the Iberian Peninsula or over the entire Mediterranean (thus including part of the Iberian Peninsula) may be due to the lack of shrubs in the land cover map and parametrization
- 1155 of the ORCHIDEE-CAN branch. Therefore the models simulates a higher stand density and higher basal area for regions where in reality shrubs occur (Fig. ??).

The parametrization of the forestry module strongly depends on the national forest inventories from Spain, France, Germany and Sweden. Therefore verification against the same data contains little information about the model quality. Nevertheless, no time-dependent relationships were used in

1160 the ORCHIDEE-CAN branch thus the model's capacity to reproduce the relationship between basal area and stand age, diameter and stand age or wood volume and stand age could be considered as largely independent test of the model quality. These tests were performed over 8 bioclimatic regions of France and the ORCHIDEE-CAN branch was found to largely capture the time dependencies of basal area, diameter and wood volume (not shown).

1165 6 Conclusions

ORCHIDEE-CAN (SVN r2290) differs from the trunk version of ORCHIDEE (SVN r2243) by the allometric-based allocation of carbon to leaf, root, wood, fruit and reserve pools; the transmittance, absorbance and reflectance of radiation within the canopy; and the vertical discretisation of the en-

ergy budget calculations. Conceptual changes towards a better process representation were made

- 1170 for the interaction of radiation with snow, the hydraulic architecture of plants, the representation of forest management and a numerical solution for the photosynthesis formalism of Farquhar, von Caemmerer and Berry. Furthermore, these changes were extensively linked throughout the code to improve the consistency of the model. By making use of observation-based parameters the physiological realism of the model was improved and significant reparametrization was done by introduc-
- 1175 ing twelve new parameter sets that represent specific tree species or genera rather than a group of phylogenetically often unrelated species, as is the case in widely used plant functional types (PFT). As PFTs have no meaning outside the scientific community, the species level parametrization of the ORCHIDEE-CAN branch can deliver actionable information to decision-makers and forest owners on the implications of management strategies on the climate.
- 1180 Model performance was tested against spatial explicit or upscaled data for basal area, tree height, canopy strucure, GPP, albedo and evapotranspiration over Europe. The tested data streams represented biogeochemical fluxes, biophysical fluxes and forest management related vegetation characteristics. Enhanced process representation in ORCHIDEE-CAN compared to the trunk version, was found to increase model performance regarding its ability to reproduce large-scale spatial patterns of
- 1185 all tested data streams as well as their inter-annual variability over Europe. Although this validation approach gives us confidence in the large-scale performance of the model over Europe, additional validation is recommended for other regional applications or higher resolution studies.

7 Author contribution

- Developed and parametrized the ORCHIDEE-CAN model: Kim Naudts, James Ryder, Matthew
 J. McGrath, Juliane Otto, Sebastiaan Luyssaert, Nicolas Vuichard, Didier Solyga
 - Evaluated the performance of the ORCHIDEE-CAN model: Kim Naudts, James Ryder, Juliane Otto, Matthew J. McGrath, Sebastiaan Luyssaert, Aude Valade, Yiying Chen, Fabienne Maignan
- 1195 Contributed Fortran code: Vanessa Haverd (canopy gaps), Bernard Pinty (albedo), Valentin Bellassen (forestry)
 - Provided/shared observational data sets or tools for model parametrization: Hans Pretzsch, Päivi Merilä, Jens Kattge, Gerhard Bönisch, Matteo Campioli, Josep Penuelas, Detlef Schulze, Toon De Groote, Gonzalo Berhongaray, Yuan Yan, Philippe Peylin
- 1200 Developed driver data: Natasha MacBean
 - Maintained and developed the run environment: Josefine Ghattas

8 Code availability

1205

The code and the run environment are open source (http://forge.ipsl.jussieu.fr/orchidee). Nevertheless readers interested in running ORCHIDEE-CAN are encouraged to contact the corresponding author for full details and latest bug fixes.

Acknowledgements. JR, YC, MJM, JO, KN and SL were funded through ERC starting grant 242564 (DO-FOCO), and AV was funded through ADEME (BiCaFF). ESA ECV landcover also supported this work. The research leading to these results has received funding from the European Community's Seventh Framework Programme (FP7/ 2007-2013) under the Grant Agreement n° 284181-TREES4FUTURE. The authors would like to these Deniels de Director for providing a basel area men for European

1210 like to thank Daniele de Rigo for providing a basal area map for Europe.

(R1)



Figure 1. Schematic overview of the changes in ORCHIDEE-CAN. For the trunk the most important processes and connections are indicated in black, while the processes and connections that were added or changed in ORCHIDEE-CAN are indicated in red. Numbered arrows are discussed in section 2.2.



Figure 2. Root mean square error of ORCHIDEE-CAN for gross primary production, evapotranspiration, visible and near-infra-red albedo, effective leaf area index, basal area and height for different regions and periods (DJF: December-February, MAM: March-May, JJA: June-August, SON: September-November). The grayscale of the symbols indicates the number of pixels included in the calculation. The transition from green to white indicates an RMSE of 100%.



Figure 3. Root mean square error of ORCHIDEE-trunk for gross primary production, evapotranspiration and visible and near-infra-red albedo for different regions and periods (DJF: December-February, MAM: March-May, JJA: June-August, SON: September-November). The gray-scale of the symbols indicates the number of pixels included in the calculation. The transition from green to white indicates an RMSE of 100%.



Figure 4. Comparison between observations and simulations of ORCHIDEE-CAN for gross primary production and basal area over Europe. Gross primary production represents the mean for June-August between 2001-2010 and basal area is the value at the end of 2010.



Figure 5. Impact of the different forest management strategies on an oak forest for unmanged (green), high stand(orange) and coppice (blue) compared to a Poplar short rotation coppicing (red) at 48 °N, 2 °E. The simulation was run without spin-up to better visualize carbon build up in the coarse woody debris (C.W.D.) pool. The simulation cycled of a single year (1990) of climate data to minimize the inter annual variability due to climatic year to year variability



Figure 6. Root mean square error (RMSE) of tree diameter for different species (shown as different markers) for different regions over France (shown as A to K). Open triangle, *Pinus sylvestris*; open circle, *Pinus pinaster*; open square, *Picea Sp.*; filled diamond, *Quercus ilex/suber*; filled triangle, *Betula Sp.*; filled circle, *Fagus sylvatica*; filled square, *Quercus robur/petraea*.

Table 1. Concise description	on of the modules in the standard ORCHIDEE version with the motivation to change the m	odules in ORCHIDEE-CAN.
Module	Description	Motivation for change
Albedo	For each PFT the total albedo for the grid square is computed as a weighted average of the vegetation albedo, the soil albedo, and the snow albedo.	The scheme overlooks the effect of vegetation shading bare soil for sparse canopies and gives the ground in all PFTs the same reflectance properties as bare soil.
Soil hydrology	Vertical water flow in the soil is based on the Fokker–Planck equation that resolves water diffusion in non-saturated conditions from the Richards equation (?). The 2 m soil column consists of eleven moisture layers with an exponentially increasing depth (?).	No change
Soil temperature	The soil temperature is computed according to the Fourier equation using a finite differ- ence implicit scheme with seven numerical nodes unevenly distributed between 0 and 5.5 m (?).	No change
Energy budget	The coupled energy balance scheme, and its exchange with the atmosphere, is based on that of ? . The surface is described as a single layer that includes both the soil surface and any vegetation.	A big leaf approach does not account for within canopy transport of carbon, water and energy. Further, it is incon- sistent with the current multi-layer photosynthesis approach and the new multi-layer albedo approach.
Photosynthesis	C3 and C4 photosynthesis is calculated following ? and ?, respectively. Photosynthesis assigns artificial LAI levels to calculate the carbon assimilation of the canopy. These levels allow for a saturation of photosynthesis with LAI, but have no physical meaning.	The scheme uses a simple Beer's law transmission of light to each level, which is inconsistent with the new albedo scheme.
Autotrophic respiration	Autotrophic respiration distinguishes maintenance and growth respiration. Maintenance respiration occurs in living plant compartments and is a function of temperature, biomass and, the prescribed carbon/nitrogen ratio of each tissue (?). A prescribed fraction of 28% of the photosynthates allocated to growth is used in growth respiration (?). The remaining assimilates are distributed among the various plant organs using an allocation scheme based on resource limitations (see allocation).	No change

	401C 1.	
Module	Description	Motivation for change
Carbon allocation	Carbon is allocated to the plant following resource limitations ?. Plants allocate carbon to their different tissues in response to external limitations of water, light and nitrogen availability. When the ratios of these limitations are out of bounds, prescribed allocation factors are used.	The resource limitation approach requires capping LAI at a predefined value. Due to this cap, the allocation rules are most often not applied, reducing the scheme to prescribing allocation.
Phenology	At the end of each day, the model checks whether the conditions for leaf onset are satisfied. The PFT-specific conditions are based on long and short term warmth and/or moisture conditions (?).	No change
Mortality and turnover	All biomass pools have a turnover time. Living biomass is transferred to the litter pool, litter is decomposed or transferred to the soil pool.	This approach is not capable of modelling stand dimensions.
Soil and litter carbon and heterotrophic respi- ration	Following (?), prescribed fractions of the different plant components go to the metabolic and structural litter pools following senescence, turnover or mortality. The decay of metabolic and structural litter is controlled by temperature and soil or litter humidity. For structural litter, its lignin content also influences the decay rate.	No change
Forest management	An explicit distribution of individual trees (?) is the basis for a process-based simulation of mortality. The aboveground 'stand-scale' wood increment is distributed on a yearly time step among individual trees according to the rule of (?): the basal area of each individual tree grows proportionally to its circumference.	The concept of the original implementation were retained, however, the implementation was adjusted for consistency with the new allocation scheme and to have a larger diversity of management strategies.

÷
f Table
Ö
latior
ontin
-
Table

q=humidity, g=functi	ion.		
Symbol in text	Unit	Symbol in ORCHIDEE-CAN	Description
F_{rm}	${ m gC}~{ m m}^{-2}{ m s}^{-1}$	resp_maint	Maintenance respiration
F_{rg}	${ m gC}~{ m m^{-2}s^{-1}}$	resp_growth	Growth respiration
$F_{LW,i}$	${ m W}{ m m}^2$	r_lw	Long wave radiation incident at vegetation level "i"
$F_{SW,i}$	${ m W}{ m m}^2$	r_sw	Short wave radiation incident at vegetation level "i"
F_{Trs}	${ m ms^{-1}}$	Transpir_supply	Amount of water that a tree can get up from the soil to its leaves for transpiration
$T_{a,i}$	K	temp_atmos_pres,	Atmospheric temperature at the 'present' and 'next' time step, respectively, at level "i"
		temp_atmos_next	
$T_{L,i}$	K	temp_leaf_pres	Leaf temperature at level "i"
$q_{a,i}$	kg kg ⁻¹	q_atmos_pres, q_atmos_next	Specific humidity at the "present" and "next" time step, respectively, at level "i"
$q_{L,i}$	kg kg ⁻¹	q_leaf_pres	Leaf specific humidity at level "i"
M_l	$gC plant^{-1}$	CI	Leaf mass of an individual plant
M_s	$gC plant^{-1}$	Cs	Sapwood mass of an individual plant
M_{h}	$gC plant^{-1}$	Ch	Heartwood mass of an individual plant
M_r	$gC plant^{-1}$	Cr	Root mass of an individual plant
M_{linc}	$gC plant^{-1}$	Cl_inc	Increment in leaf mass of an individual plant
M_{sinc}	$gC plant^{-1}$	Cs_inc	Increment in sapwood mass of an individual plant
M_{rinc}	$gC plant^{-1}$	Cr_inc	Increment in root mass of an individual plant
M_{totinc}	gC	b_inc_tot	Total biomass increment
M_{inc}	$gC plant^{-1}$	b_inc	Increment in plant biomass of an individual plant
M_{swc}	${ m m}^3~{ m m}^{-3}$	SWC	Volumetric soil water content
m_w	ı	wstress_fac	Modulator for water stress as experienced by the plants
m_ψ	MPa	psi_soil_tune	Modulator to account for resistance in the soil-root interface
m_{Ndeath}	ı	scale_factor	Normalization factor for mortality
$m_{LAIcorr}$	ı	lai_correction_factor	Adjustable parameter in the calculation of gap probabilities of grasses and crops
d_h	ш	height	Plant height
d_l	m^{-2}		One sided leaf area of an individual plant

Table 2. Variable description. Variables were grouped as follows: F=flux, f=fraction, M=pool, m=modulator, d=stand dimension, T=temperature, p=pressure, R=resistance,

47

																								temperature	py	
Description	Sapwood area of an individual plant	Height increment	Plant diameter	Basal area	Basal area increment	Stem circumference of an individual plant	Number of trees in diameter class l	Projected area of an opaque tree crown	Projected crown surface area	Leaf area index	Effective leaf area index	Sum of the LAI of all levels above the current level	Cross-sectional area of vegetation level "i"	Vegetation height of level "i"	Volume of vegetation level "i"	Root density	Inverse of the individual plant density	Pressure difference between leaves and soil	Bulk soil water potential in the rooing zone	Soil water potential for each soil layer	Hydraulic resistance of roots	Hydraulic resistance of sapwood	Hydraulic resistance of leaves	Hydraulic resistance of roots, sapwood or leaves adjusted for	Aerodynamic resistance of vegetation at level "i" in the canor	Stomatal resistance of vegetation at level "i" in the canopy
Symbol in ORCHIDEE-CAN		delta_height	dia	ba	delta_ba	circ	n_circ_class	crown_shadow_h	csa_sap	1	laieff	lai_sum	,	delta_h	1	root_dens	ı	delta_P	psi_soilroot	psi_soil	R_root	R_sap	R_leaf	1	big_r	big_r_prime
Unit	m^{-2}	ш	ш	$m^2 plant^{-1}$	$m^2 plant^{-1}$	ш	trees	m^2	m^2	$\mathrm{m}_{leaf}^{2}\mathrm{m}_{ground}^{-2}$	I	I	m^2	ш	m ³	I	ind m^2	MPa	MPa	MPa	${ m MPa~s~m^{-3}}$	${ m MPa~s~m^{-3}}$	${ m MPa~s~m^{-3}}$	${ m MPa~s~m^{-3}}$	$\mathrm{s}\mathrm{m}^{-1}$	${ m s~m^{-1}}$
Symbol in text	d_s	d_{hinc}	d_{dbh}	d_{ba}	d_{bainc}	d_{circ}	d_{ind}	d_c	d_{csa}	d_{LAI}	d_{LAIeff}	$d_{LAIabove}$	$d_{A,i}$	$d_{hl,i}$	$d_{V,i}$	d_{rd}	d_{λ}	p_{delta}	$p_{\psi sr}$	$p\psi_s$	R_r	R_{sap}	R_l	R_{temp}	$R_{a,i}$	$R_{s,i}$

 Table 2. Continuation of Table 2

			sdc		ce class			tion	ween height and basal area			the canopy	nosphere which has collided with canopy elements, separate	ectively	mosphere which has not collided with any canopy element	urces, respectively	ground which has not collided with any canopy elements	as struck the background a single time and has collided wi		is struck the background multiple times and has collided wi						specific competition, calculated as a function of k_{ncirc}
Description	Porosity of a tree crown	Gap probability for trees	Gap probability for grasses and cro	Gap probability for bare soil	Mortality fraction per circumferen	Leaf allocation factor	Root allocation factor	Slope of the intra-specific competi	Slope of linearised relationship bet	Reflectance of a single leaf	Transmittance of a single leaf	Reflectance of the ground beneath	Reflected fraction of light to the atn	for direct and diffuse sources, resp	Reflected fraction of light to the at	separated for direct and diffuse sou	Transmitted fraction of light to the	Reflected fraction of light which h	vegetation	Reflected fraction of light which ha	vegetation	Height above the soil	Solar zenith angle	Cosine of the solar zenith angle	Leaf orientation function	Cut-off circumference of the intra-
Symbol in ORCHIDEE-CAN	Pwc_h	PgapL	PgapL	PgapL	mortality	KF	LF	gamma	S	leaf_reflectance	leaf_transmittance	bdg_reflectance	Collim_alb_BB,	Isotrop_alb_BB	Collim_alb_BC,	Isotrop_alb_BC	Collim_Tran_Uncoll	ı		I		z_array	solar_angle	ı	ı	sigmas
Unit	I	I	I	I	I	I	I	ı	m	I	I	I	I		I		I	ı		ı		ш	radians	radians	I	ı
Symbol in text	f_{Pwc}	f_{Pgap}^{trees}	f_{Pgap}^{gc}	f^{bs}_{Pgap}	f_{death}^{icir}	f_{KF}	f_{LF}	f_γ	f_s	f_{rl}	f_{tl}	f_{Rbgd}	$f_{Coll,veg}^{f_R}$		$f_{UnColl,bgd}^{f_R}$		$f_{UnColl,veg}^{T}$	$f_{Coll,bgd,1}^{f_R}$		$f_{Coll,bgd,n}^{f_R}$		z	θ_z	θ_{μ}	g_G	g_{σ}

not include effective LAI, bas:
not include effectiv
not incl
oes ms.
k version de etween regio
IIDEE-trunl e overlap be
The ORCH ns due to th
as the data. other regio
population alues of the
n the same I from the v
comes fron t be derived
ed variable rrope canno
the simulat lihood of Eu
lihood that hat the likel
able 3. Like eight. Note t