

Response to Referee #1

We thank the reviewer for the detailed review and very helpful comments and address the suggested changes in detail below. Comments by the referee are shown in *italic* font. In addition to the changes described below, note that the figures and Table 1 have changed slightly from the previous version of the manuscript due to having to redo some simulations to correct a minor inconsistency. None of the statements in the text had to be changed because of this correction. The simulations that were redone are CAM5-chem, CAM5-chem*, CAM5-chem MAM4, and SD-CAM4-chem.

In the abstract I am missing important findings from section 4, notably the general performance for tropospheric ozone and other important trace gases and aerosols (this is summarized in Section 6 only). 8877, 10-12: Mention underestimation of observational data for methane lifetime

We agree with the reviewer and will add a summary of the general performance of the model in the Abstract. The abstract is changed to:

“The Community Atmosphere Model (CAM), version 5, is now coupled to extensive tropospheric and stratospheric chemistry, called CAM5-chem, and is available in addition to CAM4-chem in the Community Earth System Model (CESM) version 1.2. The main focus of this paper is to compare the performance of configurations with internally derived “free running” (FR) meteorology and “specified dynamics” (SD) against observations from surface, aircraft, and satellite, as well as understand the origin of the identified differences. We focus on the representation of aerosols and chemistry. All model configurations reproduce tropospheric ozone for most regions based on in-situ and satellite observations. However, shortcomings exist in the representation of ozone precursors and aerosols. Tropospheric ozone in all model configurations agrees for the most part with ozonesondes and satellite observations in the Tropics and the Northern Hemisphere within the variability of the observations. Southern Hemispheric tropospheric ozone is consistently underestimated by up to 25%. Differences in convection and stratosphere to troposphere exchange processes are mostly responsible for differences in ozone in the different model configurations. Carbon monoxide (CO) and other volatile organic compounds are largely underestimated in Northern Hemisphere mid latitudes based on satellite and aircraft observations. Nitrogen oxides (NO_x) are biased low in the free tropical troposphere, whereas peroxyacetyl nitrate (PAN) is overestimated in particular in high northern latitudes. The present-day methane lifetime estimates are compared among the different model configurations. These range between 7.8 years in the SD configuration of CAM5-chem and 8.8 years in the FR configuration of CAM4-chem and are therefore underestimated compared to observational estimations. We find that differences in tropospheric aerosol surface area between CAM4 and CAM5 play an important role in controlling the burden of the tropical tropospheric hydroxyl radical (OH), which causes differences in tropical methane lifetime of about half a year between CAM4-chem and CAM5-chem. In addition, different distributions of NO_x from lightning explain about half of the difference between SD and FR model versions in both CAM4-chem and CAM5-chem. Remaining differences in the tropical OH burden are due to enhanced tropical ozone burden in SD configurations compared to the FR versions, which are not only caused by differences in chemical production or loss, but also by transport and mixing. For future studies, we recommend the use of CAM5-chem configurations, due to improved aerosol description and inclusion of aerosol-cloud interactions. However, smaller tropospheric surface area density in the current version of CAM5-chem compared to CAM4-chem results in larger oxidizing capacity in the troposphere and therefore a shorter methane lifetime.”

Section 1:

8878, 10: *Additional references would be desirable (e.g. Isaksen et al., doi:10.1016/j.atmosenv.2009.08.003).*

We agree and add the suggested reference.

Section 2:

8879, 26-27: *The term “data ocean” is not known to me. Re-formulate the sentence to “All model simulations are performed with prescribed sea surface temperatures and sea-ice distribution data for present-day climatological conditions, : : :”*

We agree with the suggestion and change the sentence accordingly:

“All model simulations are performed with prescribed sea surface temperatures and sea-ice distribution data for present-day climatological conditions, since we focus on the atmospheric component.”

8884, 5: *You may think about a different naming instead of “CAM5-chem*”. This term is also not searchable with Adobe Reader.*

Since the sensitivity experiment is not the focus of the paper, we prefer to continue using this naming.

8884, 10: *Replace “performance” by “setup and global model diagnostics”.*

We agree with the suggestions of the reviewer and changed the phrase accordingly.

Section 3: One could think of adding more surface station data to the evaluation data base, notably Ozone and other species or aerosol parameters from global or regional air quality networks, but this may be beyond the scope of the paper. At least the choice of evaluation data should be discussed for their relevance.

We thank the reviewer for pointing this out and follow the advice and add surface ozone observations to the evaluation in this paper. We perform a comparison of the probability distribution function of surface ozone stations over North America and Western Europe. For this we have added the new Figure 14 (see below) and add a description of the dataset and a discussion, see below.

8884, 22-25: *I found this very confusing. Why using MOPITT here? The sentence seems not to belong here.*

We agree with the review and moved the sentence to the next paragraph.

8886, 25-27: *Please clarify the meaning of a “profile-to-profile” comparison. How are observational data and model results matched in space and time?*

To clarify we change the following paragraph (line 16):

“For the comparison with model results one has to keep in mind that aircraft campaigns often do not sample climatological or background conditions of the atmosphere, since they are designed to target specific atmospheric conditions. Further, monthly-mean model results that are averaged over various years are not able to identify specific pollution plumes or structures of the atmosphere as observed in a particular campaign.”

To

“For each observed regional profile, monthly-mean model results are averaged over the location and months of the observations. It is assumed that these regional profiles represent typical background conditions. However, one has to keep in mind that aircraft campaigns often target specific atmospheric conditions that may not be captured in multi-year average model results.”

We further change “A profile-to-profile comparison between aircraft and model data is performed for ozone (O₃) carbon monoxide (CO), nitrogen oxides (NO_x), and peroxyacetyl nitrate

(CH₃COO₂NO₂ or PAN) and other hydrocarbons.” To
“A comparison is performed for ozone (O₃) carbon monoxide (CO), nitrogen oxides (NO_x), peroxyacetyl nitrate (CH₃COO₂NO₂ or PAN), selected hydrocarbons, SO₂ and sulfate aerosol for selected aircraft campaigns”

8887, 21-27: The first part of the sentence until “: : (Table 2)” is an unnecessary replication and can be skipped. Instead, another sentence to introduce the IMPROVE measurements would be useful.

We agree with the reviewer and change this sentence:

“In addition to a limited set of aircraft observations available for profile-to-profile comparisons to the model output (see Table 2), we use surface observations from the United States Interagency Monitoring of Protected Visual Environments (IMPROVE) dataset (<http://vista.cira.colostate.edu/improve/>), (Malm, 2004), for years 1998–2009, to compare sulfur dioxide and sulfate.”

To

“We use two sets of surface observations in this study. Surface observations from the United States Interagency Monitoring of Protected Visual Environments (IMPROVE) dataset (<http://vista.cira.colostate.edu/improve/>, Malm, 2004), is used for years 1998–2009, to compare sulfur dioxide and sulfate with the model results. The IMPROVE network includes 165 sites in the US. Major fine particles (with diameter < 2.5 micro meter) are monitored including aerosol species, sulfates, nitrates, organics, light-absorbing carbon, and wind-blown dust. IMPROVE sites are located in rural environments and therefore will not describe the conditions found in large urban areas.”

We further add:

“Ozone surface observations are used to evaluate daily ozone concentration in our model configurations. Daily averages from available hourly surface ozone data were derived from the Clean Air Status and Trends Network (CASTNET) (<http://java.epa.gov/castnet/>) and the European Monitoring and Evaluation Programme (EMEP) network in Europe (<http://www.emep.int/>) for years 1995-2010, as shown in Tilmes et al. (2012).”

Section 4: In section 4.2, the evaluation should be ordered strictly by species (groups). In particular, all CO evaluation should be placed after the ozone evaluation.

We agree with the reviewer and restructure the section in the following:

Section 4.2.1 Aerosols and Aerosol Optical Depth (AOD) (not changed)

Section 4.2.2 Ozone (including former section “Ozone column” and “Ozone profiles”)

Section 4.2.3 CO and hydrocarbons

Section 4.2.4 NO_x and PAN

We arrange the text accordingly, and rearrange the Figures as described below.

I cannot follow the argumentation when it comes to NH lower troposphere ozone differences between CAM5-chem and SD-CAM5-chem, as it is done in Sections 4.2.2, 4.2.3, and 5. From what is presented in the paper, the relation of BL tropospheric ozone to STE in the model configurations seems to be weak (or vague). A more stringent argumentation chain will be appreciated by the readers.

We agree with the reviewer that the relation of ozone in the NH boundary layer and STE in the model configuration was not sufficiently supported. We decided to move this discussion to the model-to-model comparison section, and added a new Figure 3 (see below) supporting the

statements. The new figure includes the comparison of ozone, ozone production, lightning NO_x, CO, H₂O, and HNO₃, between different model versions. Based on this comparison, we can make the argument that higher ozone mixing ratios in high northern latitudes are not due to ozone production, but rather caused by stronger STE in CAM5-chem compared to SD-CAM5-chem. In addition to some other changes in Section 4.1 we add:

“The comparison of chemical constituents in the two model configurations further supports a stronger tropical vertical transport in CAM5-chem compared to SD-CAM5-chem and stronger STE in high latitudes (Fig.3). Stronger tropical vertical transport (mostly in deep convection) in CAM5-chem is evident due to higher mixing ratios in CO and lower mixing ratios of nitric acid in the upper tropical troposphere. The resulting higher CO mixing ratios in the upper troposphere together with increased LNO_x in mid-latitudes lead to greater ozone production, while reduced LNO_x in the tropical belt reduces ozone production. Furthermore, increased nitric acid in addition to higher ozone mixing ratios in high northern latitudes point to more STE. Additionally, lower NO_x and CO values in the boundary layer in CAM5-chem indicate that increased STE rather than chemical processing results in larger ozone mixing ratios in CAM5-chem than SD-CAM5-chem. Differences in low clouds between CAM5-chem and SD-CAM5-chem also impact chemistry and result in reduced ozone production in the boundary layer in CAM5-chem. Similar differences are present between CAM4-chem to SD-CAM4-chem, however, with smaller differences in STE in high latitudes compared to the CAM5-chem configurations (not shown).”

8888, 25-26: When TTL is defined by pressure levels 150-70 hPa, you can see from Fig. 1 both higher and lower ozone mixing ratios.

We agree and corrected the sentence to point to just the region around the tropical tropopause.

8890, 1-5: Write out COSMIC. These observational data, together with MLS and AIRS, need to be introduced before. Also, the observations give different cold point altitudes. All in all, Figure 3 is not really exploited. The whole paragraph (8889, 26 – 8890, 9) together with Figure 3 could be skipped.

We agree with the reviewer and remove the figure and corresponding discussion.

8890, 10: Fig.4. goes higher up than 30 hPa.

This was a typo, we changed it to 10hPa.

8890, 20-24: Instead of showing Fig. 5, AOA could be assessed by simply comparing AOA entries for a certain height, which can be given in Table 1.

AOA could be certainly expressed in a table picking a few values for different regions, like the tropics, and lower stratosphere in high latitudes. However, we do think, that details shown in the zonal mean plots are important, since for example they show differences in the shape of the age of air more clearly than what can be explained by a couple numbers.

8892, 13-16: CAM4-chem and CAM5-chem results are not shown in Fig.7.

We agree with the reviewer and added “not shown” at the end of this paragraph.

8892, 25-27: This is only true for free tropospheric SO₄. SO₂ is largely underestimated.

We agree and changed the sentence to: “In comparison to aircraft observations over Central Canada in July 2008, simulated SO₄ values in the free troposphere are in the range of variability of observations.”

8893, 9-13: I cannot see any significant and systematic differences between the model

configurations, at least from Figure 9. I suggest to skip these sentences.

We agree with the reviewer and removed the following sentences: “CAM4-chem and in part CAM5-MAM4-chem represent the influence from high BC plumes over the Pacific somewhat better than CAM5-chem. However, CAM5-MAM4-chem shows a stronger overestimation of background BC than the other models, especially in the upper troposphere.”

8894, 14-22: The whole paragraph would better fit to 4.2.4.

We agree with the reviewer, and have reorganized section 4.2 (as described above).

8894, 15: Is this total column or tropospheric column (as Figure 11 suggests)? In most studies, satellite CO used for evaluation is total column.

Here we have used the column below 100hPa and therefore approximately the tropospheric column. We have added the following information to the text: “CO columns are derived for altitudes between surface and 100 hPa”, and also add these information to the Figure caption.

8894, 17: It seems you were misled by the color coding in Fig. 11. There is no significant high bias for CAM4-chem for most of the year.

We agree with the reviewer and remove the statement and just state: “The tropical CO column agrees within 5% with the observations.”

8894, 26: The better agreement of SD models with observations for high latitudes cannot be derived directly from Figure 11.

We agree and remove this statement. However, we have added the following sentence in the next paragraph referring to stratospheric ozone column:

“SD configurations do not show the low bias in ozone column during the ozone hole season in both hemispheres, but instead slightly overestimate column ozone at that time. The reason for this is that temperatures in the SD configurations temperatures are slightly higher than for the FR versions especially the lower stratosphere in high latitudes.”

8895, 20-21: Be more honest here. Modelled ozone has deviations of up to 25% for larger world regions.

Based on comments by Reviewer 2 and to discuss the deviations between model results and observations with regard to the uncertainties and variability of observations, we have added two Figures, new Figure 13 and 16, showing the seasonal cycle of selected regions and different altitudes between models and observations. The mean deviations between modeled ozone and observations is still up to 25%, however, differences are for the most part within the standard deviations of the observations. We further added a probability distribution function comparison to surface observations, as mentioned above. We changed this section to:

“Ozonesonde observations (Sect.3.2), aircraft data (Sect.3.3), and surface observations (Sect.3.4) are used to evaluate the simulated tropospheric chemical composition in more detail. We use a Taylor-like diagram to illustrate relative differences between model configurations and ozonesonde observations, and correlations of the seasonal cycle for different regions, seasons, and different pressure levels, see Figs.12 and 14. In addition, seasonal cycle comparisons between model results and observations for specific regions are illustrated in Figs.13 and 15. A comparison of surface ozone is performed, showing probability distribution functions between model results and observations for Western and Eastern North America and Western Europe in Fig.14.

Near surface ozone at 900hPa is for the most part within the range of variability of ozonesonde

observations in both SD and FR configurations (Fig.12 top row). The high bias in summer over Eastern US and Western Europe, as reported in earlier studies (e.g., Lamarque et al., 2012), has been significantly reduced, due to an improved calculation of dry deposition velocities (Val Martin et al., 2014). In comparison to surface observations (Fig.14), in winter, FR model configurations slightly overestimate maximum ozone values for North America and Western Europe. SD configurations show a low bias for Eastern North America and Western Europe. In summer, all model configurations show a high bias of about 10-15 ppb. However, maximum ozone mixing ratios do agree with observations, whereas low ozone mixing ratios are overestimated. A high bias of about 10 ppb can be attributed to the coarse model resolution, which leads to an overestimate of ozone production, because of diluted emissions of ozone precursors and therefore an increase in the lower ozone mixing ratios of its distribution (e.g., Pfister et al., 2014). Ozonesondes are not compared to the model configurations at the surface. Those agree well to surface observations, besides they bias high over Eastern US in summer, as discussed in Tilmes et al., (2012).

In the mid-troposphere, model results agree well with ozonesonde observations at 500hPa (Fig.12, bottom row). The seasonal cycle is well produced, in particular for the FR configurations in mid- and high latitudes, with correlations around 0.95 compared to the observations. The somewhat higher bias in winter and spring over Western Europe and high latitudes in CAM5-chem in 500hPa contributes to the high bias in 900hPa, as more ozone is transported downward, discussed in Sect.4.1. The low bias in ozone in the West Pacific / East Indian Ocean is due to the stronger convection in the FR model configurations compared to SD, as also discussed in Sect.4.1. This bias is also shown in the comparisons at 250hPa, (Figs 14 and 15). At 50hPa, all configurations show a high ozone bias by at least 20% in the tropics during winter and spring. Mid- and high latitude ozone in the stratosphere is reproduced well for all configurations within the range of variability.

Comparisons to the aircraft climatology in the free troposphere (2-7km, Fig.17, top row) confirm the high bias of ozone in CAM5-chem and the low bias in the SD configuration at high latitudes, as well as the low bias in the Tropics in fall. Deviations from the aircraft climatology are much larger (up to 40%) compared to the ozonesonde observations (up to 25%).

In comparison to HIPPO aircraft observations over the Pacific, ozone mixing ratios are biased high in mid and high latitudes in both CAM4-chem and CAM5-chem configurations, mainly in fall and winter (Fig.18 first and second column). In addition, in spring CAM5-chem simulates larger ozone in the NH mid and high latitudes than the other models. The high ozone bias in both CAM4-chem and CAM5-chem in the remote region of the Pacific further points to a too strong STE in the FR versions. In the tropical troposphere, CAM5-chem reproduces observed mean ozone mixing ratios very well, while there is also the low biased summer and fall. However, SD configurations simulate larger ozone mixing ratios in winter and spring compared to ozonesondes and HIPPO observations.

The better representation of tropical ozone in the SD configurations in summer and fall may therefore be the result of more realistic convection, or due to a larger production of LNO_x in this region. The observations further confirm that STE in winter and spring in mid- and high latitudes is slightly too strong in CAM5-chem compared to the other configurations.”

8896, 5-10: I am not convinced here. There are regions with reduced cloud fraction over the NH but also regions with larger cloud fraction. Also, the implications for ozone of cloud differences over Africa and the Middle East are not obvious. Wouldn't it be

more illustrative (also for the mid latitude differences) to show differences in modelled ozone production?

We thank the reviewer for the comment. A significant difference in the BL ozone as a result of differences in low clouds between the FR and SD configurations could not be confirmed. Differences in surface ozone production between the different models are not significant. We removed this discussion from the paper, including Figure 14.

8896, 11-15: Again, the model bias is up to 25% in Figure 13 and up to 40% for the aircraft data.

8896, 17-19: "Especially : : : observations". Skip this sentence or re-phrase.

8896, 23: Compliance of model results with aircraft observations over Europe and US is not shown.

8897, 9-10: The 250 hPa level is not stratospheric.

8898, 7-10: Specific campaigns are not shown directly by the figures.

We have changed the text in this section, see comments above.

8898, 19-21: HNO₃ nor NO_y are shown in Fig. 20. Fig. 15 does, but gives no clear answer.

We agree and have removed this sentence.

Section 5: 8899, 13-16: Re-phrase this sentence. "As shown in model intercomparison projects projects like : : :, the reason for differences cannot be easily ascribed : : :". Give references here, e.g. Naik et al. (2013) or Voulgarakis et al., doi:10.5194/acp-13-2563-2013.

We change "The reason for differences cannot be easily ascribed to specific processes in model intercomparison projects like the Atmospheric Chemistry and Climate Model Intercomparison Project (ACCMIP), since various processes in models differ."

To

"The reason for differences cannot be easily ascribed to specific processes in models that contributed to the intercomparison projects such as the Atmospheric Chemistry and Climate Model Intercomparison Project (ACCMIP) (Voulgarakis et al., 2013, Naik et al., 2013)."

8899, 22-25: This has been already described in Section 4.2.5.

We agree and removed this sentence.

8899, 27 – 8900, 4: Be more precise here and in Figure 22. Is it tropical CO burden or tropospheric CO burden or both. Similar for the Methane lifetime: Atmospheric or tropospheric?

We have added the information in the text and figure caption to make it clearer.

8900, 16 - 8902, 9: I guess that all burdens mentioned here are tropospheric burdens. This should be mentioned somewhere.

See comment above.

Section 6: 8902, 15-21: Overestimation of surface SO₂ and SO₄ for polluted regions points to an overestimation of emissions here?

We add to the text: "In this model version, anthropogenic emissions of SO₂ and SO₄ are emitted at the surface, which can lead to underestimated transport into the free troposphere."

8903, 9-11: *As said before, I don't agree with this conclusion: Deviations are up to 25% for ozonesondes (Fig. 13) and up to 40% for aircraft campaigns (Fig. 15).*

8903, 14-15: *Looking at Fig. 11, the bias looks more like 25-50% for the southern extratropics. We changed this bullet point to:*

“Tropospheric ozone in the tropics and the Northern Hemisphere is very well represented in all model configurations and agrees within the variability of ozonesonde observations of about 25%. Surface observations are well reproduced in winter. The summer high bias of all models over Western Europe and Northern America can be for the most part contributed to a high bias in low and medium ozone mixing ratios as a result the coarse resolution of the model configurations. In the free troposphere, FR configurations slightly overestimate ozone in mid and high latitudes and underestimate ozone in the tropical free troposphere in summer and fall, while SD configurations slightly overestimate ozone in the upper tropical troposphere and in part underestimate ozone in high latitudes. Southern Hemisphere tropospheric ozone is underestimated by 10-25% in all model configurations. The comparison to aircraft observations confirms the differences based on ozonesonde observations, but models show a large bias up to 40% compared to observations.

8903, 23-24: *A general underestimation of NO_x is not supported by Figs. 15, 19 or 20.*

Instead, both overestimation and underestimation can be found.

Here, we do not agree with the reviewer. NO_x is for the most part underestimated, in particular in high latitudes. We added this information to the text.

Table 1: Exchange “model performance” by “global model diagnostics”.

Table 1: Add AOA as model parameter. Make sure that all burdens are atmospheric, otherwise say “tropospheric burden”.

We agree with the reviewer and add the information for tropospheric burden and lifetimes. As mentioned earlier, we would like to keep the AOA figure and do not add additional values into the table. The table caption is changed to:

“Overview of model experiments, setup between different simulations, and global model diagnostics. Lifetimes and burdens are calculated for the troposphere defined for regions where ozone is below 150 ppb.”

Table 3: Entries for the following abbreviations are missing: AERONET, AIRS, AMWG, CLM, COSMIC, MACCity, MOPITT, MOZART, WMO.

We have added those abbreviations to the table.

Fig. 1: How is SAD TROP in the stratosphere defined? You could mask this area in the top right panel.

In this version of the model SAD TROP contributes to the amount of SAD in the stratosphere, and therefore impacts the chemistry. More SAD in particular in the lower-most stratosphere in CAM4-chem contributes therefore to chemical ozone depletion. Since the model prescribes stratospheric SAD, this double counting of SAD in the stratosphere will be removed in future model versions.

Fig. 1: I guess differences in RH are relative differences, but this is not clear from the figure caption (also in Fig. 2).

Relative humidity (RH) is added to the figure caption.

Fig. 10: replace “Kinne (2009)” by “(Kinne, 2009)”. Add “(model – observations)” after

“Differences”.

Done

Fig. 11: Are tropospheric columns (as in the figure title) or total columns (as suspected from the text) shown?

We added “zonally averaged CO column below 100hPa”

Fig. 11: From the main text this is an OMI/MLS climatology.

We agree and changed this accordingly.

Fig. 14: Frames for aircraft campaigns are hard to see (also in Fig. 19), those are also not really needed for the paper. Re-Phrase the Figure captions.

We have removed this figure.

Fig. 15: It is not obvious how the temporal and spatial match between campaigns and model results has been achieved for this Figure (and Fig. 18).

We revised the text to explain this more clearly and point to the text in the figure caption for more details.

Technical corrections:

All technical corrections are implemented in the revised version of the manuscript if not stated differently below.

8879, 2: Reference for Liu et al., 2014 is missing.

8879, 9: replace “Strength” by “Strengths”.

8880, 2: Write out “CLM”.

8882, 1-2: Remove “chemistry including”.

8883, 8-10: Add reference: Rienecker et al., <http://dx.doi.org/10.1175/JCLI-D-11-00015.1>.

8883, 15: For clarification say “prescribed chemical fields for longer-lived substances”.

8883, 27: Add “with” before “lower”.

8884, 14: Replace “We are interested in: : :” by “We limit ourselves to: : :”.

Changed by “We use present-day ..”

8884, 22: Write out “MOPITT”.

8885, 15: Replace “TERRA” by “Terra”.

8885, 20: Write out “AERONET”.

8886, 6: Skip “between 1995 and 2010”.

We will keep this information to identify precisely what is included in the dataset.

8887, 10: Replace “: : : similar regions and different seasons : : :” by “: : : similar regions in different seasons : : :”.

8889, 28: Add “as” before “described”.

8890, 10: Replace “: : : are analyzed: : :” by “: : : is analyzed: : :”.

8890, 18: Replace “configuration” by “configurations”.

8892, 19: Add “free” before “troposphere”.

8892, 27: Data is shown in Fig. 8, bottom right panels.

8894, 7: Replace “Column” with “Columns”.

8895, 17: Add “ozonesonde” before “observations”.

8896, 26: Figure 17 must become new Figure 16, as it is introduced first.

8896, 26: Replace “Fig. 17 first and second column” by “Fig. 16, first and second row”.

8897, 5: Add “(Fig. 17)” after “biased high”.

8897, 8: Skip "(Fig. 16)".

8897, 21: Exchange "(Fig. 15)" by "(Fig. 18)". 8898, 7: Exchange "(Fig. 18)" by "(Fig. 15)". Figures have been reorganized as discussed above;

8898, 17: Add "free" before "troposphere".

8902, 2: Add "at" after "pointing".

8909, 3-5: Please use correct doi: "doi:10.1029/2004JD005537".

Table 2: Skip ", starting 1995".

Fig. 1: Exchange "SD-CAM4-cam" by SD-CAM4-chem" in the figure caption.

Fig. 3: Re-phrase figure caption: "Comparison between zonally (20_S-20_N) and annually averaged fields of : : around the tropical tropopause region, derived from : : : this figure was removed

Fig. 7: Replace "SD-CAM5-chem (blue) and SD-CAM5-chem (red)" by "SD-CAM4-chem (red) and SD-CAM5-chem (blue)".

New Figures:




Figure 3: Comparison of ozone, nitric acid, ozone production, lightning NO_x, carbon monoxide, NO_x, hydroxyl radical, and water vapor, between CAM5-chem and SD-CAM5-chem.




Figure 13: Seasonal cycle comparison between observations using a present-day ozonesonde climatology between 1995--2011 (black) and model results: CAM5-chem (cyan) and CAM4-chem (orange), SD-CAM5-chem (blue) and SD-CAM4-chem (red). Model results are interpolated to the same locations as sampled by the observations and for different pressure levels, 900hPa (top panel) and 500hPa (bottom panel) for selected regions. The standard deviation of ozonesonde observations is shown as error bars and the mean and correlation of the seasonal cycle between observations and models are printed on the top of each figure.




Figure 14: Probability distribution function (PDF) of the regionally-aggregated ozone distribution for Western North America, Eastern North America, and Western Europe from surface ozone observations (grey shaded area) in comparison to regionally-aggregated ozone distributions from

model results interpolated to the location of the ozone stations (different colors), for winter (left) and summer (right).




Figure 16:

As Fig.13, but for different pressure levels, 250hPa (top panel) and 50hPa (bottom panel).

Response to Referee 2:

We thank the reviewer for the critical review of the paper, which has contributed to an improved version of this manuscript. Comments by the referee are shown in *italic* font. In addition to the changes described below, note that the figures and Table 1 have changed slightly from the previous version of the manuscript due to having to redo some simulations to correct a minor inconsistency. None of the statements in the text had to be changed because of this correction. The simulations that were redone are CAM5-chem, CAM5-chem*, CAM5-chem MAM4, and SD-CAM4-chem.

This manuscript presents an overview assessment of the performance of tropospheric chemistry within the CESM1.2. The presentation makes extensive comparison to observation for validation and assessment of model results. In the Abstract, the authors state that the system is “well suited as tools for atmospheric-chemistry modeling studies in the troposphere and lower stratosphere, whether with internally derived “free running” (FR) meteorology, or “specified dynamics” (SD)” and follows with “The main focus of this paper is to compare the performance of these configurations against observations from surface, aircraft, and satellite, as well as understand the origin of the identified differences.” Unfortunately, the paper doesn’t seem to sufficiently address the second part of the statement well enough to support the first.

We agree that the model configurations are not “perfect” in the sense that they reproduce available observations within the range of uncertainty. Indeed many shortcomings still exist in the model and further development is required. We drop the sentence in the abstract that is of concern to the reviewer and instead focus on describing the performance of the different configurations in comparison to observations.

We therefore modify the abstract as follows:

“The Community Atmosphere Model (CAM), version 5, is now coupled to extensive tropospheric and stratospheric chemistry, called CAM5-chem, and is available in addition to CAM4-chem in the Community Earth System Model (CESM) version 1.2. The main focus of this paper is to compare the performance of configurations with internally derived “free running” (FR) meteorology and “specified dynamics” (SD) against observations from surface, aircraft, and satellite, as well as understand the origin of the identified differences. We focus on the representation of aerosols and chemistry. All model configurations reproduce tropospheric ozone for most regions based on in-situ and satellite observations. However, shortcomings exist in the representation of ozone precursors and aerosols. Tropospheric ozone in all model configurations agrees for the most part with ozonesondes and satellite observations in the Tropics and the Northern Hemisphere within the variability of the observations. Southern Hemispheric tropospheric ozone is consistently underestimated by up to 25%. Differences in convection and stratosphere to troposphere exchange processes are mostly responsible for differences in ozone in the different model configurations. Carbon monoxide (CO) and other volatile organic compounds are largely underestimated in Northern Hemisphere mid latitudes based on satellite and aircraft observations. Nitrogen oxides (NO_x) are biased low in the free tropical troposphere, whereas peroxyacetyl nitrate (PAN) is overestimated in particular in high northern latitudes. The present-day methane lifetime estimates are compared among the different model configurations. These range between 7.8 years in the SD configuration of CAM5-chem and 8.8 years in the FR configuration of CAM4-chem and are therefore underestimated compared to observational estimations. We find that differences in tropospheric aerosol surface area between CAM4 and CAM5 play an important role in controlling the burden of the tropical tropospheric hydroxyl radical (OH), which causes differences in tropical methane lifetime of about half a year between

CAM4-chem and CAM5-chem. In addition, different distributions of NO_x from lightning explain about half of the difference between SD and FR model versions in both CAM4-chem and CAM5-chem. Remaining differences in the tropical OH burden are due to enhanced tropical ozone burden in SD configurations compared to the FR versions, which are not only caused by differences in chemical production or loss, but also by transport and mixing. For future studies, we recommend the use of CAM5-chem configurations, due to improved aerosol description and inclusion of aerosol-cloud interactions. However, smaller tropospheric surface area density in the current version of CAM5-chem compared to CAM4-chem results in larger oxidizing capacity in the troposphere and therefore a shorter methane lifetime.”

It is recognized that a truly thorough evaluation of a system as complex as this is difficult and time consuming, but the content of this paper is a pretty extensive information dump that for the most part lacks anything more than speculative and unsupported guesses regarding the results, their main drivers and why they are what they are relative to observation.

We agree with the reviewer that the submitted version of the manuscript was lacking in supportive arguments for some of the conclusions drawn. We also agree that a lot of information is given in the paper. To improve the paper, we have now included what we feel are better arguments for the drawn conclusions and removed those conclusions that could not be supported, as discussed in more detail below. We have tried to make the statements more quantitative, and also improved the structure of Section 4.2 as suggested by the other reviewer, and perform the evaluations grouped by chemical species.

- Why and what is the impact of the new reactions added?

The reason for the change in reactions are described in the paper line 8882: “In CESM1.2 CAM4-chem, the lumped aromatic (“TOLUENE”) was replaced with the specific species benzene, xylene and toluene, along with simplified oxidation products for the two new species, to accommodate the 2-product formation of SOA (new reactions listed in Appendix A).”

The 2-product formation of SOA in CAM4-chem requires additional species that were not included in earlier versions of the model. This change has little impact on the chemistry, but allows the new SOA calculation. We have added the sentence:

“These changes do not have an impact on the chemical performance of the model.”

- Can the impacts of clouds, dynamics, aerosols, etc. be more explicitly evaluated to address the inter-model differences and differences between model and observation?

In this paper, we focus on the performance and evaluation of aerosols and chemistry. Differences in clouds and dynamics between the different configurations are discussed briefly in Section 4.1 but were evaluated in detail in earlier studies. In the revised version of the manuscript we try to improve the paper and identify differences in chemistry as a result of convection and transport, and to some degree clouds, by including an additional figure, new Figure 3 (see below) and add the following text:

“The comparison of chemical constituents in the two model configurations further supports a stronger tropical vertical transport in CAM5-chem compared to SD-CAM5-chem and stronger STE in high latitudes (Fig.3). Stronger tropical vertical transport (mostly in deep convection) in CAM5-chem is evident due to higher mixing ratios in CO and lower mixing ratios of nitric acid in the upper tropical troposphere. The resulting higher CO mixing ratios in the upper troposphere together with increased LNO_x in mid-latitudes lead to greater ozone production, while reduced LNO_x in the tropical belt reduces ozone production. Furthermore, increased nitric acid in addition to higher ozone mixing ratios in high northern latitudes point to more STE. Additionally, lower

NO_x and CO values in the boundary layer in CAM5-chem indicate that increased STE rather than chemical processing results in larger ozone mixing ratios in CAM5-chem than SD-CAM5-chem. Differences in low clouds between CAM5-chem and SD-CAM5-chem also impact chemistry and result in reduced ozone production in the boundary layer in CAM5-chem. Similar differences are present between CAM4-chem to SD-CAM4-chem, however, with smaller differences in STE in high latitudes compared to the CAM5-chem configurations (not shown).”

It seems that this paper should go in one of two directions:

1) It should be extensively revised to include a more explicit statistical analysis of the results and their comparison to observation. This would be perhaps be best done with a smaller set of observations that highlight some of the main driving factors that impact results: e.g. tropospheric O₃ is often a clear signal of vertical transport. Methane lifetimes reflect OH abundance and chemistry, but this pits one of the slowest reacting species as an indicator of one of the fastest reacting. Is it possible to more directly assess their interaction?

We have attempted to improve the presentation of the statistical evaluation of tropospheric ozone in the revised version of the paper. In addition to the discussion of the Taylor-like diagrams for ozone evaluation based on ozonesonde observations, we also show seasonal cycle plots, including information of the variability of the observations (adding standard deviations). Further, we included the evaluation of the probability distribution function of surface ozone observations.

In additions, we have discussed the connection between methane lifetime and OH, and correlations with other factors in detail in Section 5.

2) The paper should be retracted and resubmitted as two papers: One as a technical description paper with a lot more technical discussion and evaluation. And a second that evaluates the model's performance against observation, including a proper statistical analysis.

We agree with the reviewer that this paper does not cover the technical aspects of the various model configurations since many of those are available in previous publications. The reviewer might have been misled in the introduction by the statement: “The performance of CAM4-chem and CAM5-chem is discussed in Sect. 4, which includes model-to-model comparisons of chemistry and aerosol budgets”. We have changed this sentence to: “Model-to-model differences in dynamics, chemistry and aerosols, and global budgets are discussed in Sect. 4. A comprehensive evaluation of chemistry and aerosols, based on satellite and in-situ observations is performed in Sect. 4.2.”

The focus of this paper is on evaluating the inter-model performance and the performance against observations. We believe that our statistical analysis is for the most part proper, but we agree that improvements can be made, as described in the comment above.

The authors have clearly done a lot of work preparing and testing the model system, compiling observations, and developing tools for comparison. They are well regarded for their effort, but this paper, unfortunately, is sub-par. I add, lastly, that with the density of acronyms it is somewhat difficult to read.

Unfortunately many acronyms are used, but there is not really a good solution in changing this. We have added a table for the purpose of making the paper more readable due to the acronyms used, which is updated in the revised version of the manuscript.

Please consider my comments below for more detail.

P8878, L1: The intro starts off rather abruptly. Please include a general introduction to the CESM.

We agree with the reviewer and modify the first paragraph of the introduction:

“The Community Earth System Model (CESM) is a comprehensive model that couples different independent models for atmosphere, land, ocean, sea-ice, land ice, and river runoff (e.g., Neale et al., 2013; Lamarque et al., 2012). It can be used in various configurations, depending on the use of different components and the coupling between them. The atmospheric component of CESM, the Community Atmosphere Model (CAM), has the capability of including chemistry of varying complexity. Default CESM configurations used for long-term climate model simulations usually include prescribed chemical fields in the atmosphere using monthly averages. ...”

and the first sentence of the second paragraph:

“In CESM version 1.2, CAM version 5 (CAM5) extensive tropospheric and stratospheric chemistry, referred hereafter to as CAM5-chem, has been successfully implemented.”

P8879, L27: With fixed SST, what impact might this have on truly evaluating the interactions between chem., physics, and dynamics in the results?

We purposely performed the experiments with prescribed present-day SST, to eliminate an additional factor in differences between the various model simulations. For specified dynamics runs, calculated SST would not make a difference since the atmospheric dynamics is nudged. For the free running simulation, calculated SST would add additional variability to the atmospheric fields, which would make it more difficult to identify the reason for the differences in different chemical fields. The larger variability in the specified dynamics runs for example in lightning NO_x compared to the free-running version may be caused by the fact that the SST are prescribed for all model years, which may reduce the variability in the FR version, but not in the SD version where atmospheric dynamics are from analyses. We will change this sentence in Section 5:

“The spread in the annual LNO_x production is much larger in the SD simulations compared to the FR configurations, which is likely introduced by the use of climatological SSTs in the FR configurations.”

P8880 – Since the system uses chemistry coupled to the modal aerosol scheme, comparison to results from Long et al. (2013, GMD; 2014, ACP) with more explicit multiphase chem would perhaps be valuable to this MS.

This study is focused on evaluating the different configurations of CESM with chemistry. We think that it would be beyond the scope of the paper to compare our results to a different model.

P. 8882, L15-17: Small particles DO impact het. chem. Why do the authors believe they don't? This would only be the case if the specific chemical mechanism used specifically neglected the chemistry associated with smaller particles.

We rephrase the sentence to clarify that heterogeneous chemistry on small particles is not included in the current model version, even though it has been shown that those reactions matter. “The contribution of very small particles, such as the Aitken mode in MAM3 and the primary carbon mode in MAM4, are neglected in the model calculation of surface area density. Further, sea-salt and mineral dust aerosols do not contribute to SAD in either model version, as heterogeneous reactions are not assumed to occur on these surfaces. Since reactions on very small particles are important, this may lead to an underestimation of SAD in the model.”

The conclusions also state:

“An underestimation of SAD in the model is possible, because BC plumes are significantly underestimated over source regions. Since background aerosols are in general overestimated, shortcomings may exist in the calculation of SAD. For example, sea-salt and dust provide

surfaces for heterogeneous reactions that have not been taken into account in any of the simulations (Evans et al., 2005)”

P. 8888, L 20-24: At the resolution used, is CAM even able to resolve STE well enough to permit diagnostic analysis of the impact of stratospheric chemistry and exchange on the results?

The purpose of the performed diagnostics is to identify the differences in STE in the models. Comparisons with ozonesonde and aircraft data indicate that differences in STE can lead to a slight overestimation of ozone (in CAM5-chem) or in a slight underestimation (SD-CAM versions) in mid- to high latitudes, but on average to reasonable values of ozone, as described in the text.

P8889, L10: SAD is used here but defined on p. 8890
SAD is now introduced when it is used the first time.

P8891, L13-15: This statement is self-contradicting. It appears you're using the term SAD in place of the more appropriate term surface-to-volume ratio. Perhaps this should be changed.

Surface area density in $\mu\text{m}^2/\text{cm}^3$ is indeed the surface area to volume ratio.

“Heterogeneous reactions on aerosol particles in the model do not directly relate to the aerosol burden, but rather depend on the amount of tropospheric SAD. SAD depends not only on aerosol burden or mass, but also on their size distribution. For the same aerosol burden, smaller particles provide a larger SAD than larger particles.”

P8892 and in general: Since this is a chemistry paper, more or less, it is important that the appropriate symbols and terminology are used. Is SO2 meant to refer to SO2 gas? If so, it should be SO2(g). SO4, as presented, suggests some non-ionic sulfur oxide radical. If it is meant to be particulate or aqueous SO4²⁻, it should be presented as such.

To clarify in the paper we added “SO2 is referred to as gas-phase species throughout the paper” and further, define SO4 as “sulfate aerosol”.

P8892, L17-20: Unfortunately, for a publication seeking to present a proper analysis of a chemistry model system, the terms “reasonable agreement” and “agrees well with” are insufficient. This work should be a proper analysis based on observational and modeled statistics. Avoiding this type of analysis completely undermines the utility and ability to extract meaningful information from this as a scientific work. The work that the authors have done to date compiling data and creating unbelievably useful tools for making comparisons and analysis possible should be leveraged to do at least something to step beyond the current state. IF, on the other hand, it is the authors intention to perform a more complete analysis for presentation in another manuscript for submission elsewhere – which may be the more suitable way forward – then this should be stated and much of this analysis should be removed from the manuscript.

To address the concerns of the reviewer, in the revised version of the manuscript we have added more quantitative statements to describe how “well” the model compares to the observations. We further added two new figures, Figures 13 and 16 (see below) showing seasonal cycle comparisons of the model to ozonesonde observations, additions the statistical comparison in from of a Taylor-like diagram, to visualize the performance of the model with regard to the variability of observations. We also added a comparison of probability distribution functions between surface measurements and model results (new Figure 14).

The sentence pointed out by the reviewer:

“Comparisons to aircraft observations over the US and high northern latitudes (Fig. 8), show a reasonable agreement of SO2 over the US for all model configurations. Further, SO4 agrees well in the troposphere over the US, while boundary layer values are overestimated.”

Is changed to:

“Comparisons to aircraft observations over the US (Fig. 8, left), show very good agreement for SO₂ that are very close to the observed values for two of the campaigns, while simulated values are slightly larger for ARCTAS-CARB. Further, the model configurations reproduce observed sulfate aerosol for some campaigns, with the exception of CAM4-chem, while boundary layer values are more than double to those observed.”

P8896, L6, L8; P8897, L1; P8904, L9: Invoking cloud impacts on O₃ seems speculative here. Further analysis is both possible and necessary.

We agree and have removed the discussion and corresponding Figure.

P8893, L28 – P8894, L1: If the AOD bias is the result of too much seasalt and/or dust, wouldn't this mean that the winds may be too high? This would be a straightforward analysis, wouldn't it? Also, given that AOD is based on the MIRAGE system, wouldn't a sensitivity to RH also be just as likely to affect AOD?

We give some more information on sea salt and dust in the model configuration and experiment section: “The dust emissions are calibrated so that the global dust AOD is between 0.025 and 0.030 (Mahowald et al, 2006). Further, sea salt emissions are calibrated in present-day conditions so that the global mean AOD (for all species) are within the reasonable range. Those values have been evaluated in Liu et al., 2012, who show that the difference between model simulations and observations are generally within a factor of two.

Further we add the following text to Section 4.2.1: Aerosols and Aerosol Optical Depth (AOD):

“The AOD bias in the subtropical ocean (mostly from coarse mode sea salt) can be due to the model deficiency representing the sea salt emission or sedimentation (scavenging) process that requires further investigation. Using reanalysis winds do not reduce this bias (not shown).”

P8904, L3-4: This is improper English.

We change the sentence: “Differences in the H₂O-tape recorder and in AOA point to a too strong Brewer–Dobson circulation in the FR model configurations, while it is reasonably reproduced in the SD configurations.” to

Differences in the H₂O tape recorder and in AOA indicate that the Brewer–Dobson circulation is too strong in the FR model configurations, while both diagnostics are reasonably reproduced in the SD configurations.”

Table 1 (and most of the rest of the data shown): What are the statistics? Are they means? If so, what about std. dev. If they're medians, what about range? In any case, are the statistics appropriate? For example, if the optical depth data are means, it should be pointed out that AOD is not normally distributed and therefore means are meaningless (no pun intended). In fact, most of the quantities presented in this table and elsewhere are known to not only NOT be normally distributed, but the distribution and resulting statistics are known to differ with model horizontal resolution. In most cases, there are clear modes within the distributions that can be attributed to individual geographic regions, features, or locations in the atmosphere. As such, changes in one area may appear to affect the whole from a statistical standpoint, when in fact there is no change at all elsewhere. It is possible that any analysis or comparison based on these data are insupportable.

The purpose of Table 1 is to show global values and budgets that are meant to give an overview of the performance of the different model configuration. The total budgets, lifetimes, and other variables, are usually presented in this way. The values we are showing are averages over all the years of the simulations. We are not discussing variability of the different variables here, but go into the details of the variability of some of the variables in Section 4 and 5, as well as for AOD. We are very aware of the fact that many variables, for example ozone, are not distributed normally depending on regions and altitudes considered, as for example pointed out in Tilmes et al., 2012. This is why this table is not meant to serve as a full evaluation tool, but to give an overview of the global diagnostics, whereas detailed investigations are presented further along in the paper. We are not discussing the impact of different horizontal resolutions, however, in the new version of the paper, we acknowledge that some shortcoming in the model are likely a result of the coarse resolution (see the additional comparison to surface observations). Further case studies would be beneficial to investigate the behavior of different variables in more depth and specific aspects of the model, but this is beyond the scope of this paper.

New Figures:




Figure 3: Comparison of ozone, nitric acid, ozone production, lightning NO_x, carbon monoxide, NO_x, hydroxyl radical, and water vapor, between CAM5-chem and SD-CAM5-chem.




Figure 13: Seasonal cycle comparison between observations using a present-day ozonesonde climatology between 1995--2011 (black) and model results: CAM5-chem (cyan) and CAM4-chem (orange), SD-CAM5-chem (blue) and SD-CAM4-chem (red). Model results are interpolated to the same locations as sampled by the observations and for different pressure levels, 900hPa (top panel) and 500hPa (bottom panel) for selected regions. The standard deviation of ozonesonde observations is shown as error bars and the mean and correlation of the seasonal cycle between observations and model results are printed on the top of each figure.




Figure 14: Probability distribution function (PDF) of the regionally-aggregated ozone distribution for Western North America, Eastern North America, and Western Europe from surface ozone observations (grey shaded area) in comparison to regionally-aggregated ozone distributions from

model results interpolated to the location of the ozone stations (different colors), for winter (left) and summer (right).




Figure 16:

As Fig.13, but for different pressure levels, 250hPa (top panel) and 50hPa (bottom panel).

Description and evaluation of tropospheric chemistry and aerosols in the Community Earth System Model (CESM1.2)

S. Tilmes¹, J.-F. Lamarque¹, L. K. Emmons¹, D. E. Kinnison¹, P.-L. Ma², X. Liu³, S. Ghan², C. Bardeen¹, S. Arnold⁴, M. Deeter¹, F. Vitt¹, T. Ryerson⁵, J. W. Elkins⁵, F. Moore⁵, R. Spackman⁶, and M. Val Martin⁷

¹National Center for Atmospheric Research, Boulder, Colorado, USA

²Pacific Northwest National Laboratory, Richland, Washington, USA

³University of Wyoming, USA

⁴School of Earth and Environment, University of Leeds, Leeds, UK

⁵NOAA ESRL Chemical Sciences Division, Boulder, Colorado, USA

⁶NOAA ESRL Science and Technology Corporation (STC), Boulder, Colorado, USA

⁷Department of Chemical and Biological Engineering, The University of Sheffield, Sheffield, UK

Correspondence to: S. Tilmes (tilmes@ucar.edu)

Abstract

The Community Atmosphere Model (CAM), version 5, is now coupled to extensive tropospheric and stratospheric chemistry, called CAM5-chem, and is available in addition to CAM4-chem in the Community Earth System Model (CESM) version 1.2. ~~Both configurations are well suited as tools for atmospheric chemistry modeling studies in the troposphere and lower stratosphere, whether~~ The main focus of this paper is to compare the performance of configurations with internally derived “free running” (FR) meteorology, ~~or~~ and “specified dynamics” (SD). ~~The main focus of this paper is to compare the performance of these configurations~~ against observations from surface, aircraft, and satellite, as well as understand the origin of the identified differences. We ~~particularly focus on comparing~~ focus on the representation of aerosols and chemistry. All model configurations reproduce tropospheric ozone for most regions based on in-situ and satellite observations. However, shortcomings exist in the representation of ozone precursors and aerosols. Tropospheric ozone in all model configurations agrees for the most part with ozonesondes and satellite observations in the Tropics and the Northern Hemisphere within the variability of the observations. Southern Hemispheric tropospheric ozone is consistently underestimated by up to 25%. Differences in convection and stratosphere to troposphere exchange processes are mostly responsible for differences in ozone in the different model configurations. Carbon monoxide (CO) and other volatile organic compounds are largely underestimated in Northern Hemisphere mid latitudes based on satellite and aircraft observations. Nitrogen oxides (NO_x) are biased low in the free tropical troposphere, whereas peroxyacetyl nitrate (PAN) is overestimated in particular in high northern latitudes. The present-day methane lifetime estimates within are compared among the different model configurations, ~~which~~ . These range between 7.8 years in the SD configuration of CAM5-chem and 8.8 years in the FR configuration of CAM4-chem and are therefore underestimated compared to observational estimations. We find that ~~tropospheric surface area density is an important factor~~ differences in tropospheric aerosol surface area between CAM4 and CAM5 play an important role in controlling the burden of the tropical tropospheric hydroxyl radical

(OH), which causes differences in tropical methane lifetime of about half a year between CAM4-chem and CAM5-chem. In addition, different distributions of ~~nitrogen oxides (NO_x) produced from lightning production from lightning~~ explain about half of the difference between SD and FR model versions in both CAM4-chem and CAM5-chem. Remaining differences in the tropical OH burden are due to enhanced tropical ozone burden in SD configurations compared to the FR versions, which are not only caused by differences in chemical production or loss, but also by transport and mixing. For future studies, we recommend the use of CAM5-chem [configurations](#), due to improved aerosol description and inclusion of aerosol-cloud interactions. However, smaller tropospheric surface area density in the current version of CAM5-chem compared to CAM4-chem results in larger oxidizing capacity in the troposphere and therefore a shorter methane lifetime.

1 Introduction

The Community Earth System Model (CESM) [is a comprehensive model that couples different independent models for atmosphere, land, ocean, sea-ice, land ice, and river runoff](#) (e.g., Neale et al., 2013; Lamarque et al., 2012). [It](#) can be used in various configurations, depending on the use of different components and the coupling between them. [The atmospheric component of CESM, the Community Atmosphere Model \(CAM\), has the capability of including chemistry of varying complexity.](#) Default CESM configurations ~~for example used for simulations participating in~~ [used for](#) long-term climate model ~~assessments, usually prescribe most of the~~ [simulations usually include prescribed](#) chemical fields in the atmosphere using monthly averages. To produce those prescribed input fields, simulations with a detailed representation of chemistry and aerosol processes are required. Furthermore, non-linear interactions between chemistry and aerosols in the atmosphere are important for chemistry-climate interactions (e.g., Lamarque et al., 2005; Isaksen et al., 2009) or for the simulation of air quality.

In CESM version 1.2, ~~the capability of running the Community Atmosphere Model (CAM)~~ [CAM](#) version 5 (CAM5) ~~with~~ extensive tropospheric and stratospheric chemistry, referred

hereafter to as CAM5-chem, has been successfully implemented. The performance of CAM version 4 (CAM4) with interactive chemistry, referred to as CAM4-chem, has been discussed in Lamarque et al. (2012). In this study, a similar setup of both CAM4-chem and CAM5-chem allows the comparison of both versions and their performance in comparison to observations. The two atmospheric configurations CAM4-chem and CAM5-chem differ in various aspects, including the treatment of cloud, convection, turbulent mixing, and aerosol processes (e.g., Neale et al., 2013; Gent et al., 2011; Kay et al., 2012; Liu et al., 2012), whereas the gas-phase chemistry is identical. Resulting differences in dynamics, clouds, precipitation, and radiation, will alter chemical reactions in the gas, aqueous, and aerosol phase, and removal processes, and therefore the chemical composition of the atmosphere in these configurations.

In addition to exploring differences between the two atmospheric model versions using internally produced meteorology, we also perform simulations in which the meteorology (temperature, winds, and surface fluxes) is nudged towards meteorological analysis (or re-analysis) fields to reduce differences in the dynamics of the two configurations. Further, two slightly different aerosol schemes of the modal aerosol model (MAM) are tested in CAM5-chem, the 3-mode version (MAM3) (Liu et al., 2012) and the 4-mode version (MAM4) (Liu et al., 2014). (Liu et al., 2015). In addition, sensitivity studies are performed to explore differences in the oxidizing capacity of the atmosphere and therefore in tropospheric methane lifetime in the different model configurations. In this way, relationships between tropospheric methane lifetime, aerosol and chemistry composition, and meteorological parameters are explored.

A comprehensive evaluation of all configurations is performed, using a set of present-day observational climatologies of different chemistry and aerosol species from ground-based, aircraft and satellite observations. Strengths and weaknesses of the various model configurations are discussed. Evaluation tools for trace gases and aerosols developed in this study are merged to the Atmospheric Model Working Group (AMWG) diagnostics package, and are available to the community on the CESM website (<https://www2.cesm.ucar.edu/working-groups/amwg/amwg-diagnostics-package>).

This paper is structured as follows. Section 2 gives details of the model configurations and experiments performed for this study. Section 3 describes present-day climatological datasets used in this study to evaluate the model. ~~The performance of CAM4-chem Model-to-model differences in dynamics, chemistry and CAM5-chem is discussed in Sect. 4, which includes model-to-model comparisons of chemistry and aerosol budgets (aerosols, and global budgets are discussed in Sect. 4.1) and a.~~ A comprehensive evaluation of chemistry and aerosols, based on satellite and in-situ observations (is performed in Sect. 4.2). We discuss reasons for differences in tropospheric methane lifetime of the different model configurations, an indicator of the oxidizing capacity of the atmosphere in Sect. 5. A summary and discussion of the results is given in Sect. 6.

2 Model configurations and experiments

The presented results are based on output from simulations performed with the NCAR Community Earth System Model (CESM) Version 1.2. (<https://www2.cesm.ucar.edu/models/current>). All model simulations are performed with ~~a data ocean consisting of~~ prescribed sea surface temperatures and sea-ice ~~distributions~~ distribution data for present-day climatological conditions, since we focus on the atmospheric component. Dry deposition of gases and aerosols are implemented in the ~~land model~~ Community Land Model (CLM) (Oleson, 2010) as described in Lamarque et al. (2012). For all experiments CLM Version 4.0 was used. CESM 1.2 can also include online calculation of biogenic emissions in CLM using the Model of Emissions of Gases and Aerosols from Nature (MEGAN) version 2.1 (Guenther et al., 2012). In this study, biogenic emissions are prescribed (see below) to ensure having the same amount of emissions in all configurations, and interactive bio-geo-chemistry was not included.

CAM4-chem uses 26 vertical levels while CAM5-chem uses 30, and they both have a model top around 40 km. The horizontal resolution of performed simulations is $1.9^\circ \times 2.5^\circ$ and we use the finite volume dynamical core. An important difference between the two atmospheric models is the cloud microphysics, which in CAM4-chem predicts only the mass

concentrations of the cloud species, but in CAM5-chem predicts the number as well as mass concentrations. CAM5-chem consequently treats the microphysical effect of aerosols on clouds (Ghan et al., 2012), while in CAM4-chem aerosols impact physics and dynamics only through their interaction with radiation.

CAM4-chem and CAM5-chem further differ in the parameterization of aerosols. CAM4-chem runs with a bulk aerosol model (BAM), which considers a fixed size distribution of externally-mixed sulfate, black carbon (BC), organic carbon (OC), sea-salt and dust (Tie, 2005). Sea-salt and dust are described using four different bins. In CAM4-chem, the formation of secondary organic aerosols (SOA) is coupled to chemistry. SOA are derived using the 2-product model approach using laboratory determined yields for SOA formation from monoterpene oxidation, isoprene and aromatic photooxidation, as described in Heald et al. (2008).

The current standard CAM5 model version, and therefore also CAM5-chem, uses the modal aerosol model with three modes (MAM3) (Liu et al., 2012). The aerosol components, including BC, primary organic matter (POM), SOA, sea-salt, dust, and sulfate, are internally mixed in each lognormal mode, and the aerosol mass and the total number in each mode are predicted. CAM5-chem is also tested with the 4-mode version, MAM4, called CAM5-MAM4-chem from here on. The main difference between these two modal versions used here is the representation of BC and OC. In MAM3 all BC and OC is assumed to be aged and hence is emitted directly into the accumulation mode with other soluble aerosol species, whereas MAM4 emits the BC and OC in the primary carbon mode and represents the aging process of BC and OC from the primary carbon mode to the accumulation mode, as done in BAM. For the SOA production in CAM5-chem, mass yields of several biogenic and anthropogenic Volatile Organic Compounds (VOCs) are prescribed. The resulting condensable secondary organic gas reversibly and kinetically partitions to the aerosol phase, as described in detail in Liu et al. (2012). The different approach in CAM5-chem than CAM4-chem results in much larger burden of SOA, as shown in Tsigaridis et al. (2014). [The dust emissions are calibrated so that the global dust aerosol optical depth \(AOD\) is between 0.025 and 0.030](#) (Mahowald et al., 2006). [Further, sea salt emissions](#)

are calibrated to present-day conditions so that the global mean AOD (for all species) are within the reasonable range. Those values have been evaluated in Liu et al. (2012), who have shown that the difference between model simulations and observations are generally within a factor of two.

The production of sulfate aerosol (SO_4) in CAM4-chem and CAM5-chem is also parameterized differently. In CAM4-chem, SO_4 is produced directly from sulfur dioxide (SO_2) by oxidation through heterogeneous reactions on aerosols. SO_2 is referred to as gas-phase species throughout the paper. In CAM5-chem, sulfate aerosols are assumed to be partially neutralized by ammonia (NH_3), in the form of ammonium hydrogen sulfate (NH_4HSO_4). Sulfates are produced via sulfuric acid (H_2SO_4) condensation on existing aerosols, where H_2SO_4 is formed by the oxidation of SO_2 . Both CAM4-chem and CAM5-chem include aqueous phase production of SO_4 from SO_2 , with more than half formed by the hydroperoxy (HO_2) uptake and subsequent hydrogen peroxide (H_2O_2) oxidation in clouds (Liu et al., 2012). In addition, CAM5-chem includes nucleation of SO_4 , which contributes less than 1 % to the production of SO_4 mass but is an important source of aerosol number. Also, while in CAM4-chem sulfur oxides emissions are in the form of SO_2 only, in CAM5, 2.5 % of SO_2 is emitted in the form of SO_4 .

Furthermore, the representation of removal processes is different in CAM4-chem and CAM5-chem. In CAM4-chem all of the aerosol in the cloudy fraction of the grid cell is assumed to reside within cloud droplets and is removed in proportion to the cloud water removal rate. In CAM5-chem the mass and number fraction of the cloud-borne aerosol is determined from the aerosol activation parameterization (Ghan and Easter, 2006), so that smaller particles are not removed by nucleation scavenging.

CAM4-chem has been run and tested with comprehensive ~~chemistry including~~ tropospheric and stratospheric chemistry (Lamarque et al., 2012). The chemical mechanism is based on the MOZART-4 Model for Ozone and Related chemical Tracers (MOZART), version 4 mechanism for the troposphere (Emmons et al., 2010), extended stratospheric chemistry (Kinnison et al., 2007), further updates as described in Lamarque et al. (2012), and additional reaction rate updates following JPL 2010 recommendations (Sander et al.,

2011). In CESM1.2 CAM4-chem, the lumped aromatic (“TOLUENE”) was replaced with the specific species benzene, xylene and toluene, along with simplified oxidation products for the two new species, to accommodate the 2-product formation of SOA (new reactions listed in Appendix A). [These changes do not have an impact on the chemical performance of the model.](#)

As in CAM4-chem, CAM5-chem couples tropospheric aerosols to chemistry through heterogeneous reactions, as listed in Lamarque et al. (2012, Table 4). Tropospheric heterogeneous reactions of chemical species are parameterized based on aerosol surface area density (SAD) and therefore depend on the overall aerosol loading. The total tropospheric SAD in both model configurations is derived using the mass and size distributions of ammonium sulfates, black carbon, and organic aerosols. The contribution of very small particles, such as the Aitken mode in MAM3 and the primary carbon mode in MAM4, to the SAD are neglected [in the the model calculation of surface area density](#). Further, sea-salt and mineral dust aerosols do not contribute to SAD in ~~both model versions~~ [either model version](#), as heterogeneous reactions are not assumed to occur on these surfaces. [Since reactions on very small particles are important, this may lead to an underestimation of SAD in the model.](#)

For all simulations, model configurations simulate wet deposition of gas species using the Neu and Prather (2012) scheme, including a bug fix to CESM1.2, where the SO₂ Henry’s law coefficient has been updated, resulting in reduced washout rates. This fix resulted in an increased burden of SO₄ in CAM4-chem, which has been adjusted by increasing the in- and below-cloud solubility factor of SO₄ from 0.3 to 0.4. In addition, improved calculations of dry deposition velocities for gas species, as discussed in Val Martin et al. (2014), are added to this study, which results in an improved representation of surface ozone, as discussed below.

Experiments

Two different configurations of both CAM4-chem and CAM5-chem are used in this study. In the free running (FR) version the meteorology and dynamics are internally derived. We

also run CAM4-chem and CAM5-chem in a specified dynamics (SD) version of the model, called SD-CAM4-chem and SD-CAM5-chem, respectively. In this configuration, the internally derived meteorological fields are nudged every time step (30 min) by 10 % towards analysis fields (i.e., a 5 h Newtonian relaxation time scale for nudging) from the Modern-Era Retrospective Analysis For Research And Applications (MERRA) reanalysis product (<http://gmao.gsfc.nasa.gov/merra/>) (Rienecker et al., 2011), regridded to the model horizontal resolution. The SD model version adopts the vertical levels of the analysis data up to the top of the model (around 40 km), resulting in 56 vertical levels for both CAM4-chem and CAM5-chem simulations; see Lamarque et al. (2012) and Ma et al. (2013) for details. For the SD simulations, we use meteorological analysis for the years 2000 to 2010.

Emissions and [chemical fields prescribed chemical fields for longer-lived substances](#) follow the protocol defined by the Chemistry Climate Model Initiative (CCMI) hindcast simulations for the year 2000 (Eyring et al., 2013), which are repeated for all the simulated model years for both FR and SD configurations. In particular, greenhouse gases are from Meinshausen et al. (2011), surface mixing ratios of ozone depleting substances are taken from WMO (2010, Table 5-A3), anthropogenic and biofuel emissions are from the MACCity emission data set (Granier et al., 2011), and biomass burning emissions are taken from the Atmospheric Chemistry and Climate Model Intercomparison Project (ACCMIP) historical emissions dataset (Lamarque et al., 2010). Biogenic emissions are prescribed in this study for all model configurations using a climatology based on MEGAN version 2.1, with the same emissions for all model experiments; [carbon monoxide \(CO\)](#): 1053 Tg yr⁻¹, isoprene: 525 Tg yr⁻¹, monoterpene: 97 Tg yr⁻¹, and methanol: 170 Tg yr⁻¹. All experiments use the same solar forcing, [with](#) lower boundary conditions fixed for the year 2000.

Two additional sensitivity experiments are performed to test differences between CAM4-chem and CAM5-chem that may be caused by differences in the aerosol description in the model, in particular the amount of tropospheric SAD in the different configurations. CAM5-chem simulates significantly lower SAD than CAM4 (as discussed in Sect. 4.1.2). We perform an additional CAM5-chem (CAM5-chem*) simulation where SAD is increased by a factor of 1.5 to match the averaged tropospheric SAD amount that is simulated in CAM4-

chem. We also perform SD-CAM5-chem* that matches averaged tropospheric SAD of the SD-CAM4-chem simulation, requiring SAD to increase by a factor of 1.9. And finally, we perform a simulation that uses the MAM4 modal scheme, CAM5-MAM4-chem, as described above. An overview of the [performance setup and global model diagnostics](#) of the different model configurations is given in Table 1.

3 Present day climatological datasets

To evaluate the performance of the different model configurations, we made use of several satellite and in-situ chemical datasets. We [are interested in use](#) present-day climatological datasets with a focus on the troposphere that have been derived from observations between 1995 and 2012.

3.1 Satellite climatologies

[The comparison of the model simulations to satellite observations provides a global picture on the representation of CO and ozone columns.](#) To evaluate tropospheric and stratospheric column ozone in the model simulations, we compare the model to a present-day column ozone climatology compiled by Ziemke et al. (2011). This climatology was derived by combining retrievals from the Aura Ozone Monitoring Instrument (OMI) and Microwave Limb Sounder (MLS) observations over the period between October 2004 and December 2010. ~~Monthly mean Level 3 MOPITT a priori and averaging kernels are applied to monthly mean model results to account for the a priori dependence and vertical resolution of the MOPITT data.~~ The monthly-mean thermal tropopause is used to separate between tropospheric and stratospheric ozone for the model results and satellite climatology.

For comparison with ~~carbon monoxide (CO)~~CO, a new climatology is compiled based on [MOPITT Measurements of Pollution in The Troposphere \(MOPITT\)](#) Version 6 Level 3 data, using the multispectral (thermal-infrared plus near-infrared) total column product. This monthly mean gridded climatology on a $1^\circ \times 1^\circ$ horizontal resolution includes data between 2003 and 2012. Only daytime MOPITT data were analyzed. The Version 6 MO-

PITT product is similar to the validated Version 5 product (Deeter et al., 2013) with several differences (Deeter et al., 2014). The V5 products relied on a priori CO concentrations based on the MOZART chemistry transport model and National Centers for Environmental Prediction (NCEP) analysis fields. The a priori for V6 products is based on CAM4-chem simulations for the period from 2000–2009 (Lamarque et al., 2012) and the retrieval processing exploits the MERRA reanalysis product. Finally, geolocation (latitude and longitude) data are more accurate for V6 product as the result of a correction for a slight misalignment between the MOPITT instrument and the ~~TERRA~~Terra spacecraft. The V6 product is described in more detail in a User's Guide available on the MOPITT website (<http://www2.acd.ucar.edu/mopitt/publications>). Monthly mean Level 3 MOPITT a priori and averaging kernels are applied to monthly mean model results to account for the a priori dependence and vertical resolution of the MOPITT data. CO columns are derived for altitudes between surface and 100 hPa.

For the comparison of ~~aerosol optical depth (AOD)~~AOD, we use a $1^\circ \times 1^\circ$ monthly averaged climatology for present-day AOD at 550 nm, derived using various satellite data including ~~AERONET observations~~observations from the Aerosol Robotic Network (AERONET) (Kinne, 2009).

3.2 Ozonesonde climatology

For a detailed evaluation of tropospheric ozone profiles and seasonality, a present-day ozonesonde climatology is used (Tilmes et al., 2012). This climatology covers available ozonesonde observations between 1995 and 2011 for 42 stations around the globe. Ozonesonde observations do agree reasonably well with surface and aircraft observations (Tilmes et al., 2012). Maximum summer time ozonesonde data over Eastern US is biased high by about 10 ppb compared to surface observations, but otherwise, the ozone climatology provides reliable ozone vertical profiles for different seasons and regions. In this study, monthly mean model results are interpolated to the locations of the data and aggregated over defined regions, as suggested in Tilmes et al. (2012).

3.3 Aircraft climatologies

For the evaluation of various chemical species, averaged profiles from various aircraft campaigns between 1995 and 2010 were derived for different regions and seasons around the globe. Details of aircraft campaigns included between 1995 and 2010 are given in Table 2. More details, including information of earlier aircraft campaigns, are provided on <https://www2.acd.ucar.edu/gcm/aircraft-climatology>. As discussed in Emmons et al. (2000), for each aircraft campaign, regions with high frequency occurrence of vertical profiles from the aircraft are identified. Mean and median profiles of available species are compiled over these regions, as well as percentiles of the distribution with a 1 km vertical resolution. Profiles that are outliers of the distribution were removed. Following this approach, we extended the existing climatology as described in Emmons et al. (2000), to include additional aircraft campaigns up to 2010.

The largest sampling frequency of aircraft observations included in this study is over Europe and the US during spring and summer. For ~~the comparison with model results each observed regional profile, monthly-mean model results are averaged over the location and months of the observations. It is assumed that these regional profiles represent typical background conditions. However,~~ one has to keep in mind that aircraft campaigns often ~~do not sample climatological or background conditions of the atmosphere, since they are designed to target specific atmospheric conditions. Further, monthly mean model results that are averaged over various years are not able to identify specific pollution plumes or structures of the atmosphere as observed in a particular campaign that may not be captured in multi-year average model results.~~ Nevertheless, the combination of the numerous aircraft campaigns provides a general overview on the behavior of the chemistry in the model. In this way, aircraft data provide a very powerful evaluation tool, because various species were observed at the same time during the flight and can be evaluated side by side. A ~~profile to profile comparison between aircraft and model data comparison~~ is performed for ozone (O_3), ~~carbon monoxide (CO),~~ CO_2 , nitrogen oxides (NO_x), and peroxyacetyl nitrate ($CH_3COO_2NO_2$ or PAN) ~~and other hydrocarbons,~~ selected hydrocarbons, SO_2 and sulfate

aerosol for selected aircraft campaigns. In addition, we averaged profiles over certain altitude intervals and grouped them into four regions and four seasons, to identify systematic differences between models and observations.

A data set derived during the HIAPER (High-Performance Instrumented Airborne Platform for Environmental Research) Pole-to-Pole Observations (HIPPO) campaigns (Wofsy et al., 2011) is available for model evaluation purposes (Wofsy et al., 2012). During the campaigns, profiles from 85° N–65° S over the Pacific Ocean and North America were sampled in January and November 2009, March/April 2010, June/July 2011, and August/September 2011. Each of the campaigns sampled very similar flight tracks over the Pacific and North America, which provides information for comparing similar regions ~~and~~ in different seasons (Wofsy et al., 2011). For this paper, we use O₃, BC, and PAN data (Schwarz et al., 2013; Wofsy et al., 2011). The aircraft profiles sampled during different HIPPO campaigns were averaged over 5° latitude intervals along the flight path over the Pacific Ocean to produce a gridded dataset that can be easily compared to model output. Likewise, model results are binned over the same latitude regions as done for the aircraft observations. Here, we compare the observations to monthly mean model data that are aligned with the months of the corresponding campaign. It has to be kept in mind that the HIPPO dataset, even though observing the background atmosphere over the Pacific, is influenced by the specific situation for the particular year. This climatological comparison has shortcomings, in particular because the emissions of the particular year were not considered.

3.4 ~~IMPROVE-network~~ Surface observations

~~In addition to a limited set of aircraft observations available for profile-to-profile comparisons to the model output (see Table 2), we use surface observations~~ We use two sets of surface observations in this study. Surface observations from the United States Interagency Monitoring of Protected Visual Environments (IMPROVE) dataset (<http://vista.cira.colostate.edu/improve/>), (Malm, 2004), ~~for years 1998–2009~~ is used for years 1998–2009, to compare sulfur dioxide and sulfate ~~.IMPROVE sites are aerosol with~~

the model results. The IMPROVE network includes 165 sites in the US. Major fine particles (with diameter < 2.5 micro meter) are monitored including aerosol species, sulfates, nitrates, organics, light-absorbing carbon, and wind-blown dust. IMPROVE sites are located in rural environments and therefore will not describe the conditions found in large urban areas.

Ozone surface observations are used to evaluate daily ozone concentration in our model configurations. Daily averages from available hourly surface ozone data were derived from the Clean Air Status and Trends Network (CASTNET) (<http://java.epa.gov/castnet/>) and the European Monitoring and Evaluation Programme (EMEP) network in Europe (<http://www.emep.int/>) for years 1995-2010, as shown in Tilmes et al. (2012).

4 Performance for different model configurations

4.1 Model-to-model comparison

Differences in the physics, including cloud and aerosol schemes between CAM4-chem and CAM5-chem (as described above), result in large differences in tropospheric surface area density, temperatures, relative humidity and cloud fraction, with implications for chemistry, in particular ozone (Figs. 1 and 2). Additional differences in the vertical resolution of different model configurations influence ~~tropospheric and stratospheric dynamics and convection~~ and dynamics in troposphere and stratosphere and therefore atmospheric composition.

4.1.1 ~~Dynamics and ozone~~

~~CAM4 chem and CAM5 chem show large differences in~~ The comparison of zonal and annual mean ~~relative humidity (Fig. 1), with significantly larger values in mid and high latitudes in CAM5-chem compared to CAM4-chem. These are likely caused by the differences in the microphysics in the two configurations. The fraction of low clouds in all configurations varies between 34 and about 60 (Table 1) and are caused by the different parameterizations of cloud macrophysics with some contribution from the cloud microphysics, but also by differences in the assumed minimum relative humidity values that allow clouds to form.~~

~~Differences in cloud fraction between different configurations impact photolysis rates in the lower troposphere and therefore ozone photochemistry (discussed below), and also precipitation and removal processes~~meteorological as well as chemical constituents between different model versions helps to explain differences in ozone and other chemical tracers.

4.1.1 Dynamics and chemistry

CAM5-chem simulates more ozone in the stratosphere than CAM4-chem, most pronounced in high latitudes in the lower stratosphere, ~~which likely contributes to the stronger stratosphere to troposphere exchange (STE) in mid and high latitudes (Table 1).~~ This is aligned with lower temperatures in the stratosphere in the tropics and mid-latitudes in CAM5-chem compared to CAM4-chem, resulting in reduced ozone destroying gas-phase chemistry. Further, lower ozone mixing ratios and a cold bias are present in CAM5-chem in right around the tropical tropopause ~~layer (TTL)~~ in comparison to CAM4-chem. Reduced ozone in the TTL around the tropical tropopause can affect temperatures at the cold point and above (Bardeen et al., 2013). ~~The lower ozone in the TTL in CAM5-chem compared to CAM4-chem may be further caused by differences in the upwelling, as discussed below. In addition, differences in~~

Differences in zonal winds point to a weaker polar vortex in CAM5-chem compared to CAM4-chem, whereby zonal winds in CAM5-chem are more aligned with analysis fields than in CAM4-chem (not shown). Corresponding higher temperatures in the polar lowermost stratosphere are consistent with higher ozone mixing ratios in high latitudes due to a reduction in halogen activation.

Differences in the microphysics between CAM4-chem and CAM5-chem result in significantly larger relative humidity in the troposphere in mid and high latitudes in CAM5-chem compared to CAM4-chem (Fig. 1, as discussed in Bardeen et al. (2013)). The fraction of low clouds in all configurations varies between 34 % and about 60 % (Table 1) and are caused by the different parameterizations of cloud macrophysics with some contribution from the cloud microphysics, but also by differences in the assumed

minimum relative humidity values that allow clouds to form. Differences in cloud fraction between different configurations impact photolysis rates in the lower troposphere and therefore ozone photochemistry (discussed below), and also precipitation and removal processes.

Large differences between CAM5-chem and CAM4-chem configurations are present in the tropospheric surface area density (SAD), as further discussed below. Those differences impact tropospheric chemistry, whereby less SAD in CAM5-chem results in the reduction of NO_x , OH, and therefore changes in CO and ozone production, in additions to changes in lightning NO_x (LNO_x), see further discussion in Sect. 5.

However, differences in dynamics between CAM5-chem and CAM4-chem have a stronger impact on ozone than differences in clouds and SAD, as shown in comparing SD-CAM5-chem and SD-CAM4-chem (Fig. 1, bottom row). In these two configurations, winds and temperatures are nudged to meteorological analyzed fields. Similarities in the meteorological fields lead to much smaller differences in ozone than between the FR versions, despite the large differences in relative humidity, clouds fraction, and SAD, which are similar to the differences between two free running model versions.

The importance of impact of differences in dynamics for tropospheric chemistry is further supported in comparing CAM5-chem and SD-CAM5-chem (Fig. 2 and 3). In these two model simulations, differences in clouds and SAD are much smaller than between CAM4-chem and CAM5-chem. However, the FR version produces a significantly stronger polar vortex and lower temperatures in high latitudes than the SD version. Smaller ozone mixing ratios in the TTL and larger ozone mixing ratios especially in the northern polar region point to a stronger Brewer Dobson Circulation (BDC) in CAM5-chem than in SD-CAM5-chem, as further illustrated in comparisons of stratospheric age of air (AOA) in the different configurations (see below). Furthermore, annually averaged temperatures are lower in the FR version throughout the atmosphere.

Dynamical differences in the TTL and the stratosphere are investigated for the different model configurations in comparing temperatures, water vapor (H_2O) and relative humidity (Fig. ??), as well as the tape recorder (Fig. 4) and stratospheric AOA, described in,

~~(Fig. 5). CAM5-chem simulates the coldest temperatures in the TTL compared to the other configurations and observations, as shown in . SD simulations driven by MERRA temperatures are significantly higher than the FR model versions, whereas CAM4-chem simulates the cold point in higher altitudes compared to the GOSMIC observations and SD versions.~~ As shown in Bardeen et al. (2013), differences of the microphysics between different model versions determine the relative humidity in the model, and therefore the relationship between water and temperature. Warmer temperatures in SD-CAM5-chem compared to CAM5-chem therefore ~~caused cause~~ an increase in water vapor in the stratosphere.

Dynamical differences in the tropics and the stratosphere are investigated for the different model configurations in analysing the H₂O tape recorder (Mote et al., 1996) (Fig. 4) and stratospheric AOA, as described in Garcia et al. (2011), (Fig. 5). The tropical vertical transport between 23° S and 23° N and 100 and ~~30~~10 hPa ~~are is~~ analyzed for different model configurations based on the magnitude and slope of the H₂O tape recorder (Fig. 4). The slope and magnitude of the tape recorder, as derived from MLS observations between 2005 and 2011 (Fig. 4, bottom row), is best reproduced by the SD configurations, even though H₂O mixing ratios are too large in SD-CAM5-chem. CAM5-chem reproduces the magnitude of the tape recorder, while minimum H₂O mixing ratios are too low, and shows a reduced slope compared to SD-CAM5-chem. This points to a faster updraft of air masses above the TTL. CAM4-chem poorly simulates the slope compared to other model ~~configuration configurations~~, whereas SD-CAM4-chem shows a reasonable magnitude of the tape recorder in comparison to MLS observations. Consistent with the poor representation of the slope of the tape recorder compared to observations, CAM4-chem and CAM5-chem produce much shorter stratospheric AOA compared to the SD configurations (Fig. 5), ~~which is also~~. This is consistent with a ~~too strong stronger~~ BDC in both free running model configurations ~~compared and stronger stratosphere to troposphere exchange (STE) (Table 1). Slightly larger AOA values in the tropics and high latitudes are simulated in CAM5-chem compared to CAM4-chem configurations.~~

The comparison of chemical constituents in the two model configurations further supports a stronger tropical vertical transport in CAM5-chem compared to SD-CAM5-chem and

stronger STE in high latitudes (Fig. 3). Stronger tropical vertical transport (mostly in deep convection) in CAM5-chem is evident due to higher mixing ratios in CO and lower mixing ratios of nitric acid in the upper tropical troposphere. The resulting higher CO mixing ratios in the upper troposphere together with increased LNO_x in mid-latitudes lead to greater ozone production, while reduced LNO_x in the tropical belt reduces ozone production. Furthermore, increased nitric acid in addition to higher ozone mixing ratios in high northern latitudes point to more STE. Additionally, lower NO_x and CO values in the boundary layer in CAM5-chem indicate that increased STE rather than chemical processing results in larger ozone mixing ratios in CAM5-chem than SD-CAM5-chem. Differences in low clouds between CAM5-chem and SD-CAM5-chem also impact chemistry and result in reduced ozone production in the boundary layer in CAM5-chem. Similar differences are present between CAM4-chem to ~~observations and therefore smaller ozone mixing ratio in the TTL. SD-CAM4-chem, however, with smaller differences in STE in high latitudes compared to the CAM5-chem configurations (not shown).~~

4.1.2 Aerosol burden and Surface Area Density (SAD)

Optical depth and aerosol loading from the different model configurations are listed in Table 1. Total optical depth is somewhat smaller in CAM4-chem than in the CAM5-chem configuration, which is due to different amounts of internally derived sea-salt and dust emissions, but also differences in the sulfate burden in comparison to observations, as discussed in Sect. 4.2.1. The largest differences in aerosol burden between the configurations occur in the burden of SOA, with about 50 % larger values in CAM5-chem compared to CAM4-chem (as discussed above). The burden of organic matter and black carbon is slightly larger in CAM4-chem compared to CAM5-chem using MAM3, due to the different handling of these aerosols in the two configurations. More similar values of BC and OC in CAM4-chem are simulated in CAM5-MAM4-chem. Running 2 modes for BC in CAM5-MAM4-chem compared to CAM5-chem increases the BC burden by 37 % (see Table 1). SO₄ burdens in CAM4-chem are slightly larger than in CAM5-chem. This is because of the different way SO₄ formation and washout is parameterized, as described in Sect. 2.

Heterogeneous reactions on aerosol particles in the model do not directly relate to the aerosol burden, but rather depend on the amount of tropospheric SAD. SAD depends not only on aerosol burden or mass, but also on their size distribution. For the same aerosol burden, smaller particles provide a larger SAD than larger particles. Both the SD and FR version CAM5-chem simulate much smaller SAD than CAM4-chem. This has implications for chemistry and climate (see Sect. 5). The total tropospheric SAD in the model includes SAD from SO_4 , nitrates, POM, SOA, and BC modes.

We compare the burden and SAD between SD-CAM5-chem and SD-CAM4-chem for SO_4 , BC, and SOA (Fig. 6). Both magnitude and sign of the differences in burden do not agree with differences in SAD, which is caused by different description of the size distribution of aerosols in the two model versions. In CAM4-chem, BAM assumes a fixed mean radius of 69.5 nm (Emmons et al., 2010; Lamarque et al., 2012), while in MAM3, the size distribution of aerosols is represented in three different modes. For instance, most of SO_4 in the middle and upper troposphere is in the accumulation mode, with a dry diameter size range of 58–270 nm (Liu et al., 2012). On average, SO_4 particles are larger in CAM5-chem compared to CAM4-chem. Larger particles in CAM5-chem in the upper troposphere result in smaller SAD despite the slightly larger SO_4 burden compared to CAM4-chem. The increase of BC burden in CAM5-MAM4-chem does not result in an increase of SAD in the model, because only the aged mode of BC is considered in the calculation of SAD. Instead SAD in MAM4 is slightly reduced compared to MAM3 (see Sect. 5).

4.2 Evaluation of model results

4.2.1 Aerosols and Aerosol Optical Depth (AOD)

For the evaluation of aerosols, we compare simulated SO_2 and SO_4 at the surface with observations over the US from the IMPROVE network (see Sect. 4.3.4), shown in Fig. 7 for SD-CAM4-chem and SD-CAM5-chem, only. Aircraft observations are considered over the US and high latitudes to evaluate the tropospheric distributions (Fig. 8). All model configurations overestimate SO_2 at the surface, as shown here for the SD configurations (Fig. 7) with

larger values in CAM5-chem than in CAM4-chem. Annual SO_4 concentrations for all model configurations are about twice as large as observations in rural areas over the US suggest, in particular in summer. In winter, median SO_4 values in SD-CAM4-chem are biased low compared to observation while SD-CAM5-chem is biased high, whereas CAM4-chem values are biased high and CAM5-chem are biased low (not shown).

Comparisons to aircraft observations over the US ~~and high northern latitudes~~ (Fig. 8), show ~~a reasonable agreement of over the US for all model configurations~~ very good agreement for SO_2 that are very close to the observed values for two of the campaigns, while simulated values are slightly larger for ARCTAS-CARB. Further, ~~agrees well in the troposphere over the US~~ the model configurations reproduce observed sulfate aerosol for some campaigns, with the exception of CAM4-chem, while boundary layer values are ~~overestimated. CAM4-chem also overestimates values in the troposphere compared to observations, aligned with the largest burden in in comparison to the other configurations~~ more than double to those observed. In high latitudes, all model configurations underestimate SO_2 and SO_4 compared to observations from aircraft campaigns ARCTAS and ARCPAC in spring. Those campaigns in particular sampled highly concentrated fire plumes that are not captured by climatological simulations. In comparison to aircraft observations over Central Canada in July 2008, ~~the model performs more realistically~~ simulated SO_4 values in the free troposphere are in the range of variability of observations (Fig. 8, bottom ~~left right~~ panels).

The evaluation of simulated BC for CAM4-chem, CAM5-chem, and CAM5-MAM4-chem, is performed by comparing to HIPPO aircraft campaigns over the Pacific Ocean (Sect. 3.3), as shown in Fig. 9. All model configurations overestimate background BC, as for other climate models (Schwarz et al., 2010; Wang et al., 2014; Samset et al., 2014). The most realistic representation of background BC is in CAM5-chem, where primary BC is assumed to be immediately transitioned into the aged mode and therefore directly emitted in the aged mode. On the other hand, all configurations largely underestimate BC plumes, especially in ~~Northern Hemisphere (NH)~~ NH mid and high latitudes in winter and spring, and in August in the Southern Hemisphere (SH). ~~CAM4-chem and in part CAM5-MAM4-chem represent~~

~~the influence from high BC plumes over the Pacific somewhat better than CAM5-chem. However, CAM5-MAM4-chem shows a stronger overestimation of background BC than the other models, especially in the upper troposphere.~~ Shortcomings in the simulation of BC plumes are likely caused by a potential underestimate of BC emissions, as well as shortcomings in transport and wet removal by convection (Ma et al., 2013; Wang et al., 2013), while the overestimation of background values may be in part caused by a too long lifetime of BC in the ~~models~~ model configurations (Samset et al., 2014).

More work is also needed to improve the representation of POM and SOA, which are not further discussed in this study but were evaluated in Tsigaridis et al. (2014). Large uncertainties exist in the amount of global SOA distribution from observations, and the representation of these aerosols in models, and more future work is needed for both understanding observational yields in comparison to model results.

An overall comparison of aerosol can be given by comparing Aerosol Optical Depth (AOD) from satellite and AERONET observations (see Sect. 3.1) with model results, as shown for CAM4-chem and CAM5-chem (Fig. 10). AOD derived using CAM5-MAM4-chem (not shown) is very similar to CAM5-chem. The global AOD average in CAM4-chem is slightly lower compared to the observations dataset, while it is higher in CAM5-chem. An overestimation of AOD compared to the climatology occurs in CAM5-chem in Northern Africa and the Middle East, and around 30° N and 30° S over the ocean in CAM5-chem; ~~likely due to excessive dust and sea salt emissions. On the other hand,~~ The AOD bias in the subtropical ocean (mostly from coarse mode sea salt) can be due to the model deficiency representing the sea salt emission or sedimentation (scavenging) process that requires further investigation. Using reanalysis winds do not reduce this bias (not shown). Furthermore, AOD values are underestimated over polluted regions like India, South-East Asia in both ~~models~~ model configurations. CAM5-chem has a tendency towards lower AOD in northern mid and high latitudes, which could be a result of the significant underestimation of high BC plumes in these regions. Larger values than observed in CAM4-chem over Eastern US and Europe may be in part a result of the larger simulated SO₄ burden.

4.2.2 Ozone and CO Column

The comparison of the model simulations to satellite observations provides a global picture on the representation of CO and ozone column in the different model versions. Figure 11 shows differences of the zonal mean seasonal cycle of tropospheric CO column and tropospheric and stratospheric O₃ column between model results and climatologies from satellite observations from MOPITT (for column CO) and is evaluated in comparison to a monthly-mean OMI/MLS for column climatology (Sect. 3.1).

In comparison to the observations, all model configurations show a significant low bias in column CO with a maximum in spring and fall in the NH and a smaller bias in October in the SH (Fig. 11, left column). The tropical CO column is reproduced reasonably well, with exception of a high bias for CAM4 chem for most of the year. Regional differences in column CO between CAM5-chem and MOPITT (Fig. 19) occur over polluted regions, especially in April and July for the NH and over South America and southern Africa in October. This points to a significant underestimation of CO biomass burning emissions over those regions. Further, CO is largely overestimated in January over Central Africa, which points to an overestimation of fire emissions.

Figure 11 (middle and right columns). The tropospheric ozone column in CAM4-chem and CAM5-chem is overestimated between fall and spring in the NH mid-latitudes, while it is slightly underestimated in the tropics. On the other hand, SD configurations overestimate column ozone in the tropics in summer, while showing a better agreement to observations in high latitudes. All configurations underestimate tropospheric O₃ column in the SH, with a largest deviations to the observations between September and December. Differences between the FR and SD configurations in NH mid to high latitudes are aligned with a stronger STE and stronger BDC between fall and spring in the FR versions, as discussed in Sect. 4.1.1. The reason for differences of the different model configurations in tropical tropospheric ozone column are further discussed in Sect. 5. The underestimation of tropospheric ozone in the SH, especially in October in the tropics and mid-latitudes may be caused by an underestimation of biomass burning at this time of the year, which is con-

sistent with the underestimation of CO column at the same season in the SH (~~Fig. 11, left column~~see below).

Stratospheric ozone column is reasonably well reproduced for the tropics and mid-latitudes, showing slightly more ozone in the SD versions compared to the FR versions. In high latitudes, the ozone column is largely overestimated in winter and spring in each hemisphere compared to the climatology, which points to shortcomings in stratospheric transport most pronounced in the FR simulations. On the other hand, the underestimation of column O_3 in the SH in October and December point to the well known cold bias of polar vortex temperatures in the FR model versions (Eyring et al., 2010). SD configurations do not show the low bias in ozone column during the ozone hole season in both hemispheres, but instead slightly overestimate column ozone at that time. The reason for this is that temperatures in the SD configurations temperatures are slightly higher than for the FR versions especially the lower stratosphere in high latitudes.

4.2.3 Ozone profiles

~~Both ozonesonde~~Ozonesonde observations (Sect. 3.2)~~and~~, aircraft data (Sect. 3.3), and surface observations (Sect. 3.4) are used to evaluate the simulated tropospheric chemical composition in more detail. We use a Taylor-like diagram to illustrate relative differences between ~~models and~~model configurations and ozonesonde observations, and correlations of the seasonal cycle for different regions, seasons, and different pressure levels, see Figs. 12 and 15. In addition, seasonal cycle comparisons between model results and observations for specific regions are illustrated in Figs. 13 and 16. A comparison of surface ozone is performed, showing probability distribution functions between model results and observations for Western and Eastern North America and Western Europe in Figure 14.

Near surface ozone at 900 hPa is ~~in general very well reproduced in for the most part within the range of variability of ozonesonde observations in~~ both SD and FR configurations (Fig. 12, top row). The high bias in summer over Eastern US and Western Europe, as reported in earlier studies (e.g., Lamarque et al., 2012), has been ~~mostly removed~~significantly reduced, due to an improved calculation of dry deposition velocities (Val Martin et al., 2014).

~~Larger ozone mixing ratios still exist over Western Europe in the FR model versions compared to ozonesonde observations and over Canada. Ozone in SH mid-latitudes is underestimated especially in the SD configurations. Further~~ In comparison to surface observations (Fig. 14), in winter, FR model configurations slightly overestimate maximum ozone values for North America and Western Europe. SD configurations show a low bias for Eastern North America and Western Europe. In summer, all model configurations underestimate ozone in the West Pacific/East Indian Ocean, with CAM5-chem showing the largest bias. show a high bias of about 10-15 ppb. However, maximum ozone mixing ratios do agree with observations, whereas low ozone mixing ratios are overestimated. A high bias of about 10 ppb can be attributed to the coarse model resolution, which leads to an overestimate of ozone production, because of diluted emissions of ozone precursors and therefore an increase in the lower ozone mixing ratios of its distribution (e.g. Pfister et al., 2014). Ozonesondes are not compared to the model configurations at the surface. Those agree well to surface observations, besides they bias high over Eastern US in summer, as discussed in Tilmes et al. (2012).

~~To understand differences between the different configurations in the boundary layer (0-3~~ In the mid-troposphere, model results agree well with ozonesonde observations at 500 hPa, for example between CAM5-chem and SD-CAM5-chem, we related regional patterns of simulated ozone to the lower cloud fraction in the models (Fig. 22). In the NH in spring, ozone is higher in CAM5-chem compared to SD-CAM5-chem, consistent with the stronger STE in CAM5-chem than in the SD version. In addition, reduced low cloud fractions over Northern Europe and North America (12, bottom row). The seasonal cycle is well produced, in particular for the FR configurations in mid- and high latitudes, with correlations around 0.95 compared to the observations. The somewhat higher bias in winter and spring over Western Europe and high latitudes in CAM5-chem compared to SD-CAM5-chem support stronger ozone production in these regions due to increased ultraviolet (UV) radiation. Larger cloud fractions over middle and south Africa and the Middle East likely results in less ozone in CAM5-chem over these regions in summer, which contributes to the 500 hPa contributes to the high bias in 900 hPa, as more ozone

is transported downward, discussed in Sect. 4.1. The low bias in ozone in the West Pacific / East Indian Ocean –

~~In the mid-troposphere, models agree well with ozonesonde observations at 500 hPa due to the stronger convection in the FR model configurations compared to SD. This bias is also shown in the comparisons at 250 hPa (Fig. 12, bottom row) and aircraft data between 2–7, (Figs. 15 and 16). At 50 (Fig. 17, top row). Deviations from the observations are around 20% for all model configurations. All configurations reproduce mean tropical ozone mixing ratios very well at 500 hPa, but not the seasonality, indicated by a correlation coefficient of around 0.7. Ozone mixing ratios in the SH are in general underestimated in mid and high latitudes, as in the case for the surface values, compared to ozonesonde and aircraft observations. Especially the seasonal maximum is not reached based on ozonesonde observations (not shown) in agreement with comparisons to satellite observations. The SD configurations underestimate tropospheric ozone mixing ratios in high latitudes, while CAM5-chem overestimates high latitude ozone and underestimates ozone in the tropics, as all configurations show a high ozone bias by at least 20% in the tropics during winter and spring. Mid- and high latitude ozone in the stratosphere is reproduced well for all configurations within the range of variability.~~

Comparisons to the aircraft climatology in the free troposphere (2–7 km, (Fig. 17, top row) confirm the high bias of ozone in the case for the surface values. Mid-latitude ozone over Western Europe and Eastern US is best reproduced by CAM5-chem compared to both ozonesonde and aircraft observations. CAM5-chem and the low bias in the SD configuration at high latitudes, as well as the low bias in the Tropics in fall. Deviations from the aircraft climatology are much larger (up to 40%) compared to the ozonesonde observations (up to 25%).

In comparison to HIPPO aircraft observations over the Pacific, ozone mixing ratios are biased high in mid and high latitudes in both CAM4-chem and CAM5-chem configurations, mainly in fall and winter (Fig. 18 first and second column). In addition, in spring CAM5-chem simulates larger ozone in the NH mid and high latitudes than the other models. This is consistent with larger values in CAM5-chem in mid and high latitudes compared to the

~~other configurations (as discussed in Sect. 4.2.4) and fewer lower clouds in CAM5-chem over high latitudes. model configurations.~~ The high ozone bias in both CAM4-chem and CAM5-chem in the remote region of the Pacific further points to a too strong STE in the FR versions, ~~while ozone is well reproduced in the SD configurations in mid- and high latitudes.~~

~~For the TTL.~~ In the tropical troposphere, CAM5-chem reproduces observed mean ozone mixing ratios very well, while ~~the other configurations are biased high. In particular there is also the low biased summer and fall.~~ However, SD configurations simulate larger ozone mixing ratios in winter and spring compared to ~~ozonesonde ozonesondes~~ and HIPPO observations. ~~At 50all configurations have a high ozone bias by at least 20 in the tropics.~~

The better representation of tropical ozone in the SD configurations in summer and fall may therefore be the result of more realistic convection, or due to a larger production of LNO_x in this region. The observations further confirm that STE in winter and spring in mid- and high latitudes is slightly too strong in CAM5-chem compared to the other configurations.

4.2.3 CO and hydrocarbons

In comparison to MOPITT satellite observations (Fig. 15)–11, left column) all model configurations show a significant low bias in column CO with a maximum in spring and fall in the NH and a smaller bias in October in the SH. The tropical CO column agrees within 5% with the observations. Regional differences in column CO between CAM5-chem and MOPITT (Fig. 19) occur over polluted regions, especially in April and July for the NH and over South America and southern Africa in October. This points to a significant underestimation of CO biomass burning emissions over those regions. Further, CO is largely overestimated in January over Central Africa, which points to an overestimation of fire emissions. ~~Mid- and high latitude ozone in the stratosphere is reproduced well for all configurations, besides an underestimation of ozone in high latitudes at 250 for all configurations except for CAM5-chem.~~

4.2.4 CO and hydrocarbons

CO and other hydrocarbons are strongly controlled by emissions, but also directly impacted by the amount of OH in the atmosphere. The comparison of CO between aircraft measurement and CAM5-chem model results, averaged over 2–7 km (Fig. 20), confirms the pronounced underestimation of CO mixing ratios in the NH troposphere for seasons where data are available. Inter-model differences can be explained by differences in the oxidizing capacity of the atmosphere, showing largest values for CAM4-chem, consistent with the longest methane lifetime with that configuration (Table, 1, and further discussed in Sect. 5). Furthermore, in the tropics, in spring, aircraft campaigns show in some regions larger propane (C_3H_8), and to some degree large acetylene (C_2H_2) and CO values (Fig. 17). Too strong convection in the tropics may lead to enhanced mixing ratios of short-lived species, like C_3H_8 (with an approximately 10 day lifetime) in this region, while longer-lived species are still underestimated by the models for the same campaigns.

4.2.4 NO_x and PAN

Differences in the simulation of NO_x and PAN between the configurations will have implications for simulated distributions of tropospheric ozone. As for ozone, in the FR version, especially CAM5-chem, both PAN and NO_x mixing ratios in the NH mid and high latitudes are slightly larger compared to the SD versions (Fig. 17). Model comparisons to aircraft observations, show in general an underestimation of NO_x and PAN ~~show a reasonable agreement in the gradient between low and high latitudes (Fig. 18).~~ up 80%. Some aircraft campaigns observed much higher NO_x and PAN values than simulated, for instance ARC-PAC in 2008 and SOS in 1999. Both of these campaigns targeted regions with a significant contribution of biomass burning pollution and local pollution.

In the tropics, ozone deviations from specific aircraft observations often occur along with biases in ozone precursors, NO_x , PAN, and CO, and C_3H_8 , see Figs. 17 and 20. Variations in biases between observations and model results are expected in comparing to aircraft campaigns that targeted specific conditions. We investigate aircraft profiles from those

campaigns, where the models reproduced ozone and CO mixing ratios reasonably well in the troposphere (Fig. 21). In this way, shortcomings in NO_x and PAN can be identified. In general, PAN is overestimated in the [free](#) tropical troposphere, which can be an indicator of too much convection in the model compared to observations (e.g., Fischer et al., 2014). ~~Further, SD configurations tend to show larger PAN and mixing ratios compared to the FR model version and therefore larger values in the tropics.~~ In comparison to HIPPO observations of PAN (Fig. 22), all model configurations strongly overestimate PAN in the upper troposphere, and in the NH troposphere especially in winter. Values in the lower troposphere in tropics and the SH are reasonably well reproduced.

Sensitivity studies, CAM5-chem* and SD-CAM5-chem* (Sect. 2), where SAD is increased in CAM5-chem configurations to the amount simulated in CAM4-chem simulations (see Table 1), show that only a small fraction of the differences in PAN mixing ratios between the different configurations can be attributed to differences in SAD (Fig. 21). Larger SAD values in CAM4-chem result in a faster transition of NO_x to NO_y and therefore reduced PAN production, as shown in the example in Fig. 21, top left panel, for SD-CAM5-chem. However, in the FR versions and for the other cases shown in Fig. 21, adjustments of the SAD between CAM4-chem and CAM5-chem configurations is less important.

5 Methane lifetime and OH differences in CAM4-chem and CAM5-chem

Tropospheric chemistry is strongly controlled by the oxidizing capacity of the atmosphere. The most abundant oxidants in the troposphere are OH, ozone, and nitrate radical (NO_3). These control the atmospheric lifetimes of trace gases, including methane. Methane lifetime can therefore be considered as an indicator for the performance of the model. Model configurations differ largely in tropospheric methane lifetime and often underestimate recent observational estimates of 10.2 years (Prinn, 2005) and 11.3 years (Prather et al., 2012). The reason for differences cannot be easily ascribed to specific processes in ~~model intercomparison projects like~~ [models that contributed to the intercomparison projects such](#)

as the Atmospheric Chemistry and Climate Model Intercomparison Project (ACCMIP) ~~since various processes in models differ~~ (Voulgarakis et al., 2013; Naik et al., 2013).

In this study, all simulations are based on the same framework and run with the same emissions, the same gas-phase chemistry, and in the case of the SD versions, nudged with the same dynamics. Differences in the oxidizing capacity of the atmosphere can be therefore attributed to model physics, aerosol description, and differences in dynamics between SD and FR versions, caused by differences in vertical resolution and transport processes. ~~For the two sensitivity simulations, CAM5-chem* and SD-CAM5-chem*, average tropical tropospheric SAD burden matches the values in the corresponding CAM4-chem simulations (see Sect. 2), and differences in mean tropical tropospheric SAD are for the most part removed between these configurations.~~

Methane Tropospheric methane lifetime in all model configurations in this study varies between 7.6 to 8.8 years (Table 1), which is significantly lower than observational estimates. Tropospheric methane lifetime and CO burden in the tropics (between 30° S–30° N) are both correlated to the tropical OH burden (e.g., Wang and Jacob, 1998; Murray et al., 2014), with slightly different correlations for different model configurations, Fig. 23, left and middle panel. Since CO and methane are both controlled by OH, all model configurations show a very similar CH₄/CO correlation (see Fig. 23, right panel).

To understand the processes that lead to the spread of tropical OH in different model configurations in this study, we explore relationships between annual averages of tropical OH burden and other variables averaged over 30° S–30° N over the troposphere, including tropospheric SAD, H₂O₂, ~~lightning (LNO_x)~~, HNO₃, tropospheric and stratospheric column ozone, and ozone production (Figs. 24 and 25).

A consistent difference in OH burden exists between CAM5-chem and CAM4-chem in both FR and SD versions, whereby the CH₄ lifetime of CAM4-chem is about half a year longer than in CAM5-chem, see Fig. 23. Based on the sensitivity simulations CAM5-chem* and SD-CAM5-chem*, most of the difference in OH burden can be attributed to the differences in SAD between CAM4-chem and CAM5-chem (Fig. 24, left top panel). The increased SAD results in increased heterogeneous reaction and therefore increased H₂O₂

(Fig. 24, right top), and further reductions in NO_x burden in comparison to LNO_x production (Fig. 25, left panel). This is due to the fact that enhanced tropospheric heterogeneous reactions increase both the uptake of dinitrogen pentoxide (N_2O_5) as well as the uptake of HO_2 on aerosols, which is the major aqueous-phase source of H_2O_2 . The hydrolysis of N_2O_5 on aerosols results in a reduction of NO_x . Increased H_2O_2 further results in increased production of sulfate, since the reaction of H_2O_2 with SO_2 in cloud drops is the most significant contributor to sulfate formation (Seinfeld and Pandis, 2012). For the gas-phase chemistry, the decrease of NO_x leads to a reduction of ozone and, together with the reduction in HO_x , this leads to reduced OH and therefore to an increase in methane lifetime.

However, SAD differences do not explain all the differences in the OH burden, especially between FR and SD configurations. To further analyze factors that control OH burden, we scale OH to a fixed SAD value for all configurations and use the mean tropical tropospheric SAD derived using CAM4-chem results ($\text{SAD}_{\text{cam4chem}}$) as a reference. For this, we use the slope of the line that describes the OH/SAD change between CAM5-chem and CAM5-chem* configurations, S_{SAD} , see blue and cyan line in Fig. 24, left top panel, to adjust the OH burden for all configurations to the SAD reference for SD and FR configurations:

$$\text{OH (adjusted)} = \text{OH} + S_{\text{SAD}} \cdot (\text{SAD}_{\text{cam4chem}} - \text{SAD}_{\text{model}}). \quad (1)$$

As discussed in Murray et al. (2014), OH is strongly correlated to NO_x and CO emissions, as well as to the stratospheric ozone column. Since all the simulations were performed with the same CO and NO_x emissions, differences in NO_x emissions are due to variations in LNO_x . ~~The annual spread~~ Indeed, Fig. 24, middle top panel, shows a strong dependency of the OH burden to LNO_x . The spread the annual in LNO_x production is much larger in the SD simulations compared to the FR configurations. ~~This indicates a strong dependency of the OH burden to LNO_x , which is likely introduced by the use of climatological SSTs in the FR configurations.~~ However, the same LNO_x in FR and SD does not result in the same OH burden, which shows inter-model differences are only in part (about half) a result of differences in LNO_x (Fig. 25, top, middle panel).

On the other hand, variations in OH cannot be explained by differences in stratospheric column ozone between the different model simulations. Stratospheric column ozone in the model increases between FR and SD configurations. One would expect a decrease in OH as a result of reduced photolysis rates with increasing stratospheric ozone.

Tropospheric ozone is an important driver for the OH burden in all the different model configurations. More tropospheric ozone results in higher OH burden. The question remains why tropospheric ozone is larger in the SD than the FR version. Considering ozone production, increased SAD between CAM5-chem and CAM5-chem* reduces ozone production as a result of the reduced NO_x burden. However, the same amount of ozone production in FR and SD versions does not result in the same OH burden (see Fig. 25, bottom, right panel). Therefore, enhanced ozone in the SD versions is not only due to differences in chemical production of ozone, but must be also due to differences in transport processes between SD and FR version. This is further supported by the OH to HNO_3 correlations (Fig. 25, middle panel). Larger HNO_3 burden is simulated in the SD configurations than in the FR versions, which is pointing at less stratospheric contribution in the FR configurations. Another source of HNO_3 in the troposphere is LNO_x . The correlation between HNO_3 and LNO_x clearly supports the conclusion that larger HNO_3 mixing ratios in the SD configuration compared to the FR simulations are not due to differences in HNO_3 production (Fig. 25, right panel). Furthermore, smaller tropical tropospheric ozone burden in CAM5-chem compared to CAM4-chem is not aligned with the larger ozone production in CAM5-chem due to larger LNO_x . Differences are therefore likely a result of differences in transport and mixing processes in the tropics.

6 Conclusions

The evaluation of the different model configurations using various observations of aerosol and chemical species shows a realistic performance of the model versions especially in simulating tropospheric ozone. Agreements and shortcomings of each model version against observations are summarized in the following:

- Surface values of SO_2 and SO_4 over rural areas of the US are largely overestimated in most model configurations, whereas median values of SO_2 are overestimated by at least a factor of four and sulfate aerosol (SO_4) is overestimated by about 100% compared to IMPROVE observations. In this model version, anthropogenic emissions of SO_2 and SO_4 are emitted at the surface, which can lead to an underestimated transport into the free troposphere. Comparisons to aircraft observations in the troposphere show a reasonable agreement between models and observations in SO_2 and SO_4 , besides a high bias in SO_4 in CAM4-chem over the US. Profiles of SO_2 and SO_4 in high latitudes are for the most part underestimated in the model.
- The different representation of BC in CAM4-chem and CAM5-chem results in a larger burden of BC in CAM4-chem, which is due to its consideration of primary and aged BC. A similar description in CAM5-MAM4-chem leads to enhanced BC burden compared to CAM5-chem. BC plumes are in general underestimated in all model configurations while background values over the Pacific Ocean are overestimated, ~~especially in CAM5 MAM4 chem,~~ whereby CAM5-chem agrees best with observations.
- AOD points to a significant underestimation of biomass burning emissions in the model, and some overestimation in CAM4-chem over West Europe and Eastern US that may be due to the overestimation of SO_4 . An overestimation of AOD over the Pacific points to too large background values in aerosols, potentially also from sea-salt, which is more pronounced in CAM5-chem than in CAM4-chem.
- Tropospheric ozone in the tropics and the Northern Hemisphere is ~~represented very well~~ very well represented in all model configurations and agrees within ~~about 20%~~ the variability of ozonesonde observations of about 25% of in-situ observations, including ozonesondes, and aircraft observations. Surface observations are well reproduced in winter. The summer high bias of all models over Western Europe and Northern America can be for the most part contributed to a high bias in low and medium ozone mixing ratios as a result the coarse resolution of the model configurations. In the free troposphere, FR configurations slightly overestimate ozone in mid and high latitudes

and underestimate ozone in the tropical free troposphere in summer and fall, while SD configurations slightly overestimate ozone in the upper tropical troposphere and in part underestimate ozone in high latitudes. Southern Hemisphere tropospheric ozone is underestimated by 10–25 % in all model configurations. The comparison to aircraft observations confirms the differences based on ozonesonde observations, but models show a large bias up to 40 % compared to observations.

- CO is largely underestimated in the Northern Hemisphere, especially in spring, and in the SH in October, pointing to the underestimation of emissions. Other hydrocarbons that are most frequently observed during aircraft campaigns are also significantly underestimated for all seasons. The lowest values of CO and hydrocarbons occur in SD-CAM5-Chem in the tropics. CO is in reasonable agreement with the observations in the tropics.
- PAN is in general overestimated in the upper troposphere in comparison to aircraft observations for all model configurations, while NO_x is **slightly** underestimated in comparison to aircraft observations, in particular in high latitudes. The largest bias of simulated PAN in comparison to HIPPO observations occurs in mid and high northern latitudes throughout the troposphere in winter months.

Differences in CAM4-chem and CAM5-chem, and FR and SD configurations are to a large part driven by differences in dynamics, including temperature, transport, and mixing processes. Differences in the H₂O ~~tape tape~~ recorder and in AOA ~~point to a too strong Brewer–Dobson circulation indicate that the Brewer–Dobson circulation is too strong~~ in the FR model configurations, while it is both diagnostics are reasonably reproduced in the SD configurations. This is consistent with the overestimation of ozone in high latitudes in FR, particularly in winter and spring for CAM5-chem. Further, shortcomings in transport and mixing are likely responsible for slightly larger ozone mixing ratios in the tropical troposphere in SD compared to FR versions of the model. ~~Further, clouds were shown to impact ozone through changes in photolysis rates.~~


Differences in the oxidizing capacity of the atmosphere, which impacts methane and CO lifetime between different model configurations, are largely controlled by tropospheric surface area density, lightning NO_x , and differences in tropospheric ozone. Smaller SAD values in CAM5-chem are responsible for the smaller methane lifetime compared to CAM4-chem. Smaller values in surface area density in CAM5-chem compared to CAM4-chem are a result of different aerosol descriptions in the two model configurations. An underestimation of SAD in the model is possible, because BC plumes are significantly underestimated over source regions. Since background aerosols are in general overestimated, shortcomings may exist in the calculation of SAD. For example, sea-salt and dust provide surfaces for heterogeneous reactions that have not been taken into account in any of the simulations (Evans and Jacob, 2005).

Besides SAD, tropospheric ozone impacts the oxidizing capacity of the model. For the SD configuration, larger ozone mixing ratios in the tropics compared to FR result in reduced methane lifetime. Therefore, variations in transport and mixing is an important driver for differences in ozone and therefore methane lifetime, which is critical for climate simulations.

Methane lifetime is in general underestimated in all model configurations compared to observational estimates, with a difference of about one year between the different configurations. The main reason for the underestimation compared to observations is likely due to shortcomings in CO and other hydrocarbon emissions, as also found in other model studies (Stein et al., 2014; Monks et al., 2014; Emmons et al., 2014). This is supported by the underestimation of CO over source regions, but also by the underestimation of AOD over source regions, pointing to a general underestimation of biomass burning emissions. Also, the underestimation of isoprene emissions can result in a significant underestimation of methane lifetime (Pike and Young, 2006).

In summary, both CAM4-chem and CAM5-chem configurations are well suited tools for atmospheric-chemistry modeling studies, considering the shortcomings discussed in this study. We recommend the use of CAM5-chem in future studies, due to the improved description of aerosol processes and cloud interactions. Ongoing work is contributing to further improving CAM5-chem configurations.

Appendix A: Additional reactions in CAM4-chem



Acknowledgements. We thank the HIPPO team for performing reliable aircraft observations used in this study, in particular Steven Wofsy for leading the campaigns, and Joshua Schwarz and Anne Perring for providing black carbon observations, and Ru-Shan Gao for providing ozone observations. We further thank Kenneth Aikin for providing airborne observations in an unified and user-friendly format. MERRA data used in this study have been provided by the Global Modeling and Assimilation Office (GMAO) at NASA Goddard Space Flight Center through the NASA GES DISC online archive. The CESM project is supported by the National Science Foundation and the Office of Science (BER) of the US Department of Energy. The National Center for Atmospheric Research is funded by the National Science Foundation. S. Ghan and P.-L. Ma were supported by the US Department of Energy, Office of Science, Basic Energy Research as part of the Scientific Discoveries through Advanced Computing program. The Pacific Northwest National Laboratory is operated for DOE by Battelle Memorial Institute under contract DE-AC05-76RL01830.

References

- Bardeen, C. G., Gettelman, A., Jensen, E. J., Heymsfield, A., Conley, A. J., Delanoë, J., Deng, M., and Toon, O. B.: Improved cirrus simulations in a general circulation model using CARMA sectional microphysics, *J. Geophys. Res.-Atmos.*, 118, 11679–11697, doi:10.1002/2013JD020193, 2013.
- Deeter, M. N., Martínez-Alonso, S., Edwards, D. P., Emmons, L. K., Gille, J. C., Worden, H. M., Pittman, J. V., Daube, B. C., and Wofsy, S. C.: Validation of MOPITT Version 5 thermal-infrared, near-infrared, and multispectral carbon monoxide profile retrievals for 2000–2011, *J. Geophys. Res.-Atmos.*, 118, 6710–6725, doi:10.1002/jgrd.50272, 2013.
- Deeter, M. N., Martínez-Alonso, S., Edwards, D. P., Emmons, L. K., Gille, J. C., Worden, H. M., Sweeney, C., Pittman, J. V., Daube, B. C., and Wofsy, S. C.: The MOPITT Version 6 product: algorithm enhancements and validation, *Atmos. Meas. Tech.*, 7, 3623–3632, doi:10.5194/amt-7-3623-2014, 2014.
- Emmons, L. K., Hauglustaine, D. A., Müller, J.-F., Carroll, M. A., Brasseur, G. P., Brunner, D., Staelhelin, J., Thouret, V., and Marenco, A.: Data composites of airborne observations of tropospheric ozone and its precursors, *J. Geophys. Res.*, 105, 20497, 2000.
- Emmons, L. K., Walters, S., Hess, P. G., Lamarque, J.-F., Pfister, G. G., Fillmore, D., Granier, C., Guenther, A., Kinnison, D., Laepple, T., Orlando, J., Tie, X., Tyndall, G., Wiedinmyer, C., Baugh-

- cum, S. L., and Kloster, S.: Description and evaluation of the Model for Ozone and Related chemical Tracers, version 4 (MOZART-4), *Geosci. Model Dev.*, 3, 43–67, doi:10.5194/gmd-3-43-2010, 2010.
- Emmons, L. K., Arnold, S. R., Monks, S. A., Huijnen, V., Tilmes, S., Law, K. S., Thomas, J. L., Raut, J.-C., Bouarar, I., Turquety, S., Long, Y., Duncan, B., Steenrod, S., Strode, S., Flemming, J., Mao, J., Langner, J., Thompson, A. M., Tarasick, D., Apel, E. C., Blake, D. R., Cohen, R. C., Dibb, J., Diskin, G. S., Fried, A., Hall, S. R., Huey, L. G., Weinheimer, A. J., Wisthaler, A., Mikoviny, T., Nowak, J., Peischl, J., Roberts, J. M., Ryerson, T., Warneke, C., and Helmig, D.: The POLARCAT Model Intercomparison Project (POLMIP): overview and evaluation with observations, *Atmos. Chem. Phys. Discuss.*, 14, 29331–29393, doi:10.5194/acpd-14-29331-2014, 2014.
- Evans, M. J. and Jacob, D. J.: Impact of new laboratory studies of N₂O₅ hydrolysis on global model budgets of tropospheric nitrogen oxides, ozone, and OH, *Geophys. Res. Lett.*, 32, L09813, doi:10.1029/2005GL022469, 2005.
- Eyring, V., Shepherd, T. G., and Waugh, D. W. (Eds.): SPARC Report on the Evaluation of Chemistry-Climate Models, {SPARC} Report No. 5, WCRP-132, WMO/TD-No.1526, 2010.
- Eyring, V., Lamarque, J.-F., and Hess, P.: Overview of IGAC/SPARC Chemistry-Climate Model Initiative (CCMI) Community Simulations in Support of Upcoming Ozone and Climate Assessments, Tech. rep., 2013.
- Fischer, E. V., Jacob, D. J., Yantosca, R. M., Sulprizio, M. P., Millet, D. B., Mao, J., Paulot, F., Singh, H. B., Roiger, A., Ries, L., Talbot, R.W., Dzepina, K., and Pandey Deolal, S.: Atmospheric peroxyacetyl nitrate (PAN): a global budget and source attribution, *Atmos. Chem. Phys.*, 14, 2679–2698, doi:10.5194/acp-14-2679-2014, 2014.
- Garcia, R. R., Randel, W. J., and Kinnison, D. E.: On the determination of age of air trends from atmospheric trace species, *J. Atmos. Sci.*, 68, 139–154, doi:10.1175/2010JAS3527.1, 2011.
- Gent, P. R., Danabasoglu, G., Donner, L. J., Holland, M. M., Hunke, E. C., Jayne, S. R., Lawrence, D. M., Neale, R. B., Rasch, P. J., Vertenstein, M., Worley, P. H., Yang, Z.-L., and Zhang, M.: The Community Climate System Model Version 4, *J. Climate*, 24, 4973–4991, doi:10.1175/2011JCLI4083.1, 2011.
- Ghan, S. J. and Easter, R. C.: Impact of cloud-borne aerosol representation on aerosol direct and indirect effects, *Atmos. Chem. Phys.*, 6, 4163–4174, doi:10.5194/acp-6-4163-2006, 2006.
- Ghan, S. J., Liu, X., Easter, R. C., Zaveri, R., Rasch, P. J., Yoon, J.-H., and Eaton, B.: Toward a minimal representation of aerosols in climate models: comparative decomposi-

- tion of aerosol direct, semidirect, and indirect radiative forcing, *J. Climate*, 25, 6461–6476, doi:10.1175/JCLI-D-11-00650.1, 2012.
- Granier, C., Bessagnet, B., Bond, T., D'Angiola, A., Denier van der Gon, H., Frost, G. J., Heil, A., Kaiser, J. W., Kinne, S., Klimont, Z., Kloster, S., Lamarque, J.-F., Lioussé, C., Masui, T., Meleux, F., Mieville, A., Ohara, T., Raut, J.-C., Riahi, K., Schultz, M. G., Smith, S. J., Thompson, A., van Aardenne, J., van der Werf, G. R., and van Vuuren, D. P.: Evolution of anthropogenic and biomass burning emissions of air pollutants at global and regional scales during the 1980–2010 period, *Climatic Change*, 109, 163–190, doi:10.1007/s10584-011-0154-1, 2011.
- Guenther, A. B., Jiang, X., Heald, C. L., Sakulyanontvittaya, T., Duhl, T., Emmons, L. K., and Wang, X.: The Model of Emissions of Gases and Aerosols from Nature version 2.1 (MEGAN2.1): an extended and updated framework for modeling biogenic emissions, *Geosci. Model Dev.*, 5, 1471–1492, doi:10.5194/gmd-5-1471-2012, 2012.
- Heald, C. L., Henze, D. K., Horowitz, L. W., Feddes, J., Lamarque, J.-F., Guenther, A., Hess, P. G., Vitt, F., Seinfeld, J. H., Goldstein, A. H., and Fung, I.: Predicted change in global secondary organic aerosol concentrations in response to future climate, emissions, and land use change, *J. Geophys. Res.-Atmos.*, 113, D05211, doi:10.1029/2007JD009092, 2008.
- [Isaksen, I. S. a., Granier, C., Myhre, G., Berntsen, T. K., Dalsø ren, S. B., Gauss, M., Laj, P., Lohmann, U., Maione, M., Monks, P., Prevo t, a. S. H., Raes, F., Richter, a., Jo, P., Rognerud, B., Schulz, M., Shindell, D., Stevenson, D. S., Storelvmo, T., Wang, W., Weele, M. V., Wild, M., and Wuebbles, D., Atmospheric composition change : Climate – Chemistry interactions, *Atmospheric Environment*, 43, 5138–5192, 2009.](#)
- Kay, J. E., Holland, M. M., Bitz, C. M., Blanchard-Wrigglesworth, E., Gettelman, A., Conley, A., and Bailey, D.: The influence of local feedbacks and northward heat transport on the equilibrium arctic climate response to increased greenhouse gas forcing, *J. Climate*, 25, 5433–5450, doi:10.1175/JCLI-D-11-00622.1, 2012.
- Kinne, S.: Remote sensing data combinations-superior global maps for aerosol optical depth, in: *Satellite Aerosol Remote Sensing over Land*, Springer, Berlin Heidelberg, 361–381, 2009.
- Kinnison, D. E., Brasseur, G. P., Walters, S., Garcia, R. R., Marsch, D. A., Sassi, F., Boville, B. A., Harvey, V. L., Randall, C. E., Emmons, L., Lamarque, J. F., Hess, P., Orlando, J. J., Tie, X. X., Randel, W., Pan, L. L., Gettelman, A., Granier, C., Diehl, T., Niemaier, U., and Simmons, A. J.: Sensitivity of chemical tracers to meteorological parameters in the MOZART-3 chemical transport model, *J. Geophys. Res.*, 112, D20302, 2007.

- Lamarque, J.-F., Hess, P., Emmons, L., Buja, L., Washington, W., and Granier, C.: Tropospheric ozone evolution between 1890 and 1990, *J. Geophys. Res.*, 110, D08304, doi:10.1029/2004JD005537, 2005.
- Lamarque, J.-F., Bond, T. C., Eyring, V., Granier, C., Heil, A., Klimont, Z., Lee, D., Liousse, C., Mieville, A., Owen, B., Schultz, M. G., Shindell, D., Smith, S. J., Stehfest, E., Van Aardenne, J., Cooper, O. R., Kainuma, M., Mahowald, N., McConnell, J. R., Naik, V., Riahi, K., and van Vuuren, D. P.: Historical (1850–2000) gridded anthropogenic and biomass burning emissions of reactive gases and aerosols: methodology and application, *Atmos. Chem. Phys.*, 10, 7017–7039, doi:10.5194/acp-10-7017-2010, 2010.
- Lamarque, J.-F., Emmons, L. K., Hess, P. G., Kinnison, D. E., Tilmes, S., Vitt, F., Heald, C. L., Holland, E. A., Lauritzen, P. H., Neu, J., Orlando, J. J., Rasch, P. J., and Tyndall, G. K.: CAM-chem: description and evaluation of interactive atmospheric chemistry in the Community Earth System Model, *Geosci. Model Dev.*, 5, 369–411, doi:10.5194/gmd-5-369-2012, 2012.
- Liu, X., Easter, R. C., Ghan, S. J., Zaveri, R., Rasch, P., Shi, X., Lamarque, J.-F., Gettelman, A., Morrison, H., Vitt, F., Conley, A., Park, S., Neale, R., Hannay, C., Ekman, A. M. L., Hess, P., Mahowald, N., Collins, W., Iacono, M. J., Bretherton, C. S., Flanner, M. G., and Mitchell, D.: Toward a minimal representation of aerosols in climate models: description and evaluation in the Community Atmosphere Model CAM5, *Geosci. Model Dev.*, 5, 709–739, doi:10.5194/gmd-5-709-2012, 2012.
- [Liu, X., Ma, P.-L., Wang, H., Tilmes, S., Balwinder, S., Easter, R. C., Ghan, S. J., Rasch, P.: Development of a 4-Mode Version of Modal Aerosol Module for the Community Atmospheric Model, *Geosci. Model Dev. Disc.*, to be submitted, 2015.](#)
- Ma, P.-L., Rasch, P. J., Wang, H., Zhang, K., Easter, R. C., Tilmes, S., Fast, J. D., Liu, X., Yoon, J.-H., and Lamarque, J.-F.: The role of circulation features on black carbon transport into the Arctic in the Community Atmosphere Model version 5 (CAM5), *J. Geophys. Res.-Atmos.*, 118, 4657–4669, doi:10.1002/jgrd.50411, 2013.
- [Mahowald, N. M., Yoshioka, M., Collins, W. D., Conley, A. J., Fillmore, D. W., and Coleman, D. B.: Climate response and radiative forcing from mineral aerosols during the last glacial maximum, pre-industrial, current and doubled-carbon dioxide climates, *Geophysical Research Letters*, 33, 2–5, 2006.](#)
- Malm, W. C.: Spatial and monthly trends in speciated fine particle concentration in the United States, *J. Geophys. Res.*, 109, D03306, doi:10.1029/2003JD003739, 2004.

- Meinshausen, M., Smith, S. J., Calvin, K., Daniel, J. S., Kainuma, M. L. T., Lamarque, J.-F., Matsumoto, K., Montzka, S. A., Raper, S. C. B., Riahi, K., Thomson, A., Velders, G. J. M., and Vuuren, D. P.: The RCP greenhouse gas concentrations and their extensions from 1765 to 2300, *Climatic Change*, 109, 213–241, <http://link.springer.com/10.1007/s10584-011-0156-z>, 2011.
- Monks, S. A., Arnold, S. R., Emmons, L. K., Law, K. S., Turquety, S., Duncan, B. N., Flemming, J., Huijnen, V., Tilmes, S., Langner, J., Mao, J., Long, Y., Thomas, J. L., Steenrod, S. D., Raut, J. C., Wilson, C., Chipperfield, M. P., Schlager, H., and Ancellet, G.: Multi-model study of chemical and physical controls on transport of anthropogenic and biomass burning pollution to the Arctic, *Atmos. Chem. Phys. Discuss.*, 14, 25281–25350, 2014.
- Montzka, S. a., Krol, M., Dlugokencky, E., Hall, B., Jöckel, P., and Lelieveld, J.: Small interannual variability of global atmospheric hydroxyl, *Science (New York, NY)*, 331, 67–69, 2011.
- Mote, P. W., Rosenlof, K. H., McIntyre, M. E., Carr, E. S., Gille, J. G., Holton, J. R., Kinnerson, J. S., Pumphrey, H. C., Russell III, J. M., and Waters, J. W.: An atmospheric tape recorder: the imprint of tropical tropopause temperatures on stratospheric water vapor, *J. Geophys. Res.*, 101, 3989–4006, 1996.
- Murray, L. T., Mickley, L. J., Kaplan, J. O., Sofen, E. D., Pfeiffer, M., and Alexander, B.: Factors controlling variability in the oxidative capacity of the troposphere since the Last Glacial Maximum, *Atmos. Chem. Phys.*, 14, 3589–3622, doi:10.5194/acp-14-3589-2014, 2014.
- Naik, V., Voulgarakis, A., Fiore, A. M., Horowitz, L. W., Lamarque, J.-F., Lin, M., Prather, M. J., Young, P. J., Bergmann, D., Cameron-Smith, P. J., Cionni, I., Collins, W. J., Dalsøren, S. B., Doherty, R., Eyring, V., Faluvegi, G., Folberth, G. A., Josse, B., Lee, Y. H., MacKenzie, I. A., Nagashima, T., van Noije, T. P. C., Plummer, D. A., Righi, M., Rumbold, S. T., Skeie, R., Shindell, D. T., Stevenson, D. S., Strode, S., Sudo, K., Szopa, S., and Zeng, G.: Preindustrial to present-day changes in tropospheric hydroxyl radical and methane lifetime from the Atmospheric Chemistry and Climate Model Intercomparison Project (ACCMIP), *Atmos. Chem. Phys.*, 13, 5277–5298, doi:10.5194/acp-13-5277-2013, 2013.
- Neale, R. B., Richter, J., Park, S., Lauritzen, P. H., Vavrus, S. J., Rasch, P. J., and Zhang, M.: The Mean Climate of the Community Atmosphere Model (CAM4) in forced SST and fully coupled experiments, *J. Climate*, 26, 5150–5168, doi:10.1175/JCLI-D-12-00236.1, 2013.
- Neu, J. L. and Prather, M. J.: Toward a more physical representation of precipitation scavenging in global chemistry models: cloud overlap and ice physics and their impact on tropospheric ozone, *Atmos. Chem. Phys.*, 12, 3289–3310, doi:10.5194/acp-12-3289-2012, 2012.

- Oleson, K. W.: Technical Description of version 4.0 of the Community Land Model (CLM), NCAR Technical Note, NCAR/TN-47, 257, 2010.
- [Pfister, G. G., Walters, S., Lamarque, J. F., Fast, J., Barth, M. C., Wong, J., Done, J., Holland, G., and Bruyère, C. L., Projections of future summertime ozone over the U.S., *Journal of Geophysical Research: Atmospheres*, 119, 5559–5582, 2014.](#)
- Pike, R. C. and Young, P. J.: How plants can influence tropospheric chemistry: the role of isoprene emissions from the biosphere, *Weather*, 64, 2006.
- Prather, M. J., Holmes, C. D., and Hsu, J.: Reactive greenhouse gas scenarios: systematic exploration of uncertainties and the role of atmospheric chemistry, *Geophys. Res. Lett.*, 39, L09803, doi:10.1029/2012GL051440, 2012.
- Prinn, R. G.: Evidence for variability of atmospheric hydroxyl radicals over the past quarter century, *Geophys. Res. Lett.*, 32, L07809, doi:10.1029/2004GL022228, 2005.
- [Rienecker, M. M., Suarez, M. J., Gelaro, R., Todling, R., Bacmeister, J., Liu, E., Bosilovich, M. G., Schubert, S. D., Takacs, L., Kim, G.-K., Bloom, S., Chen, J., Collins, D., Conaty, A., and et al. da Silva, MERRA - NASA's Modern-Era Retrospective Analysis for Research and Applications, *J. Climate*, 24, 3624–3648, 2011.](#)
- Samset, B. H., Myhre, G., Herber, A., Kondo, Y., Li, S.-M., Moteki, N., Koike, M., Oshima, N., Schwarz, J. P., Balkanski, Y., Bauer, S. E., Bellouin, N., Bernsten, T. K., Bian, H., Chin, M., Diehl, T., Easter, R. C., Ghan, S. J., Iversen, T., Kirkevåg, A., Lamarque, J.-F., Lin, G., Liu, X., Penner, J. E., Schulz, M., Seland, Ø., Skeie, R. B., Stier, P., Takemura, T., Tsigaridis, K., and Zhang, K.: Modelled black carbon radiative forcing and atmospheric lifetime in AeroCom Phase II constrained by aircraft observations, *Atmos. Chem. Phys.*, 14, 12465–12477, doi:10.5194/acp-14-12465-2014, 2014.
- Sander, S. P., Friedl, R. R., Barker, J. R., Golden, D. M., Kurylo, M. J., Sciences, G. E., Wine, P. H., Abbatt, J. P. D., Burkholder, J. B., Kolb, C. E., Moortgat, G. K., Huie, R. E., and Orkin, V. L.: Chemical Kinetics and Photochemical Data for Use in Atmospheric Studies Evaluation Number 17 NASA Panel for Data Evaluation, *JLP Publ.*, 10–6, 2011.
- Schwarz, J. P., Spackman, J. R., Gao, R. S., Watts, L. a., Stier, P., Schulz, M., Davis, S. M., Wofsy, S. C., and Fahey, D. W.: Global-scale black carbon profiles observed in the remote atmosphere and compared to models, *Geophys. Res. Lett.*, 37, L18812, doi:10.1029/2010GL044372, 2010.

- Schwarz, J. P., Samset, B. H., Perring, A. E., Spackman, J. R., Gao, R. S., Stier, P., Schulz, M., Moore, F. L., Ray, E. A., and Fahey, D. W.: Global-scale seasonally resolved black carbon vertical profiles over the Pacific, *Geophys. Res. Lett.*, 40, 5542–5547, doi:10.1002/2013GL057775, 2013.
- Seinfeld, J. H. and Pandis, S. N.: *Atmospheric Chemistry and Physics: from Air Pollution to Climate Change*, John Wiley and Sons, New York, 2012.
- Stein, O., Schultz, M. G., Bouarar, I., Clark, H., Huijnen, V., Gaudel, A., George, M., and Clerbaux, C.: On the wintertime low bias of Northern Hemisphere carbon monoxide found in global model simulations, *Atmos. Chem. Phys.*, 14, 9295–9316, doi:10.5194/acp-14-9295-2014, 2014.
- Tie, X.: Assessment of the global impact of aerosols on tropospheric oxidants, *J. Geophys. Res.*, 110, D03204, doi:10.1029/2004JD005359, 2005.
- Tilmes, S., Lamarque, J.-F., Emmons, L. K., Conley, A., Schultz, M. G., Saunio, M., Thouret, V., Thompson, A. M., Oltmans, S. J., Johnson, B., and Tarasick, D.: Technical Note: Ozonesonde climatology between 1995 and 2011: description, evaluation and applications, *Atmos. Chem. Phys.*, 12, 7475–7497, doi:10.5194/acp-12-7475-2012, 2012.
- Tsigaridis, K., Daskalakis, N., Kanakidou, M., Adams, P. J., Artaxo, P., Bahadur, R., Balkanski, Y., Bauer, S. E., Bellouin, N., Benedetti, A., Bergman, T., Berntsen, T. K., Beukes, J. P., Bian, H., Carslaw, K. S., Chin, M., Curci, G., Diehl, T., Easter, R. C., Ghan, S. J., Gong, S. L., Hodzic, A., Hoyle, C. R., Iversen, T., Jathar, S., Jimenez, J. L., Kaiser, J. W., Kirkevåg, A., Koch, D., Kokkola, H., Lee, Y. H., Lin, G., Liu, X., Luo, G., Ma, X., Mann, G. W., Mihalopoulos, N., Morcrette, J.-J., Müller, J.-F., Myhre, G., Myriokefalitakis, S., Ng, N. L., O'Donnell, D., Penner, J. E., Pozzoli, L., Pringle, K. J., Russell, L. M., Schulz, M., Sciare, J., Seland, Ø., Shindell, D. T., Sillman, S., Skeie, R. B., Spracklen, D., Stavrou, T., Steenrod, S. D., Takemura, T., Tiitta, P., Tilmes, S., Tost, H., van Noije, T., van Zyl, P. G., von Salzen, K., Yu, F., Wang, Z., Wang, Z., Zaveri, R. A., Zhang, H., Zhang, K., Zhang, Q., and Zhang, X.: The AeroCom evaluation and intercomparison of organic aerosol in global models, *Atmos. Chem. Phys.*, 14, 10845–10895, doi:10.5194/acp-14-10845-2014, 2014.
- Val Martin, M., Heald, C. L., and Arnold, S. R.: Coupling dry deposition to vegetation phenology in the Community Earth System Model: implications for the simulation of surface O₃, *Geophys. Res. Lett.*, 41, 2988–2996, doi:10.1002/2014GL059651, 2014.
- [Voulgarakis, A., Analysis of present day and future OH and methane lifetime in the ACCMIP simulations, *Atmos. Chem. Phys.*, 2013.](http://www.atmos-chem-phys.net/13/2563/2013/acp-13-2563-2013.html)

- Wang, H., Easter, R. C., Rasch, P. J., Wang, M., Liu, X., Ghan, S. J., Qian, Y., Yoon, J.-H., Ma, P.-L., and Vinoj, V.: Sensitivity of remote aerosol distributions to representation of cloud–aerosol interactions in a global climate model, *Geosci. Model Dev.*, 6, 765–782, doi:10.5194/gmd-6-765-2013, 2013.
- Wang, Q., Jacob, D. J., Spackman, J. R., Perring, A. E., Schwarz, J. P., Moteki, N., Marais, E. A., Ge, C., Wang, J., and Barrett, S. R. H.: Global budget and radiative forcing of black carbon aerosol: constraints from pole-to-pole (HIPPO) observations across the Pacific, *J. Geophys. Res.-Atmos.*, 119, 195–206, doi:10.1002/2013JD020824, 2014.
- Wang, Y. and Jacob, J.: Anthropogenic forcing on tropospheric ozone and OH since preindustrial times sphere, *J. Geophys. Res.*, 103, 1998.
- WMO: Scientific Assessment of Ozone Depletion, available at: www.unep.ch/ozone/AssessmentPanels/SAP/ScientificAssessment2010/00-SAP-2010-Assement-report-2010.
- Wofsy, S. C. and the HIPPO team: HIAPER Pole-to-Pole Observations (HIPPO): fine-grained, global-scale measurements of climatically important atmospheric gases and aerosols, *Philos. T. Ser. A*, 369, 2073–86, <http://www.ncbi.nlm.nih.gov/pubmed/21502177>, 2011.
- Wofsy, S., Daube, B. C., Jimenez, J., et al.: HIPPO Merged 10-second Meteorology, Atmospheric Chemistry, Aerosol Data (R_20121129), Tech. rep., Carbon Dioxide Information Analysis Center, Oak Ridge National Laboratory, Oak Ridge, Tennessee, USA, doi:10.3334/CDIAC/hippo_010, 2012.
- Ziemke, J. R., Chandra, S., Labow, G. J., Bhartia, P. K., Froidevaux, L., and Witte, J. C.: A global climatology of tropospheric and stratospheric ozone derived from Aura OMI and MLS measurements, *Atmos. Chem. Phys.*, 11, 9237–9251, doi:10.5194/acp-11-9237-2011, 2011.

Table 1. Overview of model experiments, setup between different simulations, [overview of and global model performance diagnostics](#). Lifetimes and burdens are calculated for the troposphere defined for [regions where ozone is below 150 ppb](#).

CESM 1.2.2	CAM4-Chem	SD CAM4-Chem	CAM5-Chem	CAM5-Chem*	SD CAM5-Chem	SD-CAM5-Chem*	CAM5-Chem MAM4
Sim. Years	20 years	2000–2009	20 years	10 years	2000–2009	2000–2009	20 years
Meteorology	CAM4	MERRA (10%)	CAM5	CAM5	MERRA (10%)	MERRA (10%)	CAM5
Aerosol	BAM	BAM	MAM3	MAM3, 1.5°SAD	MAM3	MAM3, 1.9°SAD	MAM4
Vert. Res.	26L	56L	30L	30L	56L	56L	30L
CH ₄ Burden (Tg)	4153	4074	4103 4102	4106 4098	4064	4067	4100 4103
CH ₄ Lifet. (yr)	8.82	0.95 8.40	0.91 8.24	0.5 8.4	7.83	8.13	0.24 8.18
CO Burden (Tg)	308	299 301	289	294	283	291	287
CO Lifet. (yr)	0.135	0.128 0.129	0.134 0.132	0.130 0.129	0.120	0.125	0.132 0.131
O ₃ Burden (Tg)	310	309	314	310	306 313	306	314 315
O ₃ Lifet. (days)	24	24	22 23	23	24	24	23
O ₃ Net. chem. ^a (Tg yr ⁻¹)	515	474 471	530 507	519 480	480	454	536 518
O ₃ STE (Tg yr ⁻¹)	344	957 356	990 386	962 401	362	362	967 377
LNO _x (Tg N yr ⁻¹)	4.3	4.3 4.2	4.6 4.8	4.6	4.5 4.7	4.3	4.7 4.9
Total Optical Depth	0.126	0.140 0.108	0.145 0.142	0.144 0.142	0.153	0.153	0.146 0.143
SAD trop	0.35	0.43	0.23	0.35	0.24	0.44	0.22
POM Burden (Tg C)	0.72	0.75 0.76	0.56 0.58	0.56 0.59	0.66	0.66	0.69 0.84
SOA Burden (Tg C)	0.97	1.00	1.56 1.63	1.56 1.63	1.92	1.92	1.56 1.63
BC Burden (Tg C)	0.119	0.119 0.121	0.078 0.082	0.078 0.082	0.093	0.093	0.107 0.110
SO _x Burden (Tg S)	0.54	0.50 0.51	0.46 0.48	0.45 0.47	0.51	0.50	0.45 0.48
SO _x Aqu. Prod. (Tg S yr ⁻¹)	42.8	46.2 46.8	60.5 30.2	31.2 30.8	30.2	31.2	30.4 30.0
SO _x Chem. Prod. (Tg S yr ⁻¹)	11.2	9.9 10.3	12.7 13.7	12.2	14.4	13.7	12.0 13.8
SO _x Lifet. (days)	3.6	3.3	3.8	3.7	3.9 3.9	3.5	3.7 3.8
TOA residual ^b	2.88		0.97 1.35	1.09 1.33			0.95 1.36
FSDS ^c (W m ⁻²)	183.4	153.2 153.6	180.5 181.0	180.5 181.0	176.0	176.0	180.2 180.9
FSDSC ^d (W m ⁻²)	246.5	247.3 247.6	244.2	244.2	243.4	243.4	243.0 243.9
high clouds (%)	31.9	29.3	38.5	38.6	40.8 38.5	40.8	38.3
med. clouds (%)	19.0	21.3	27.3	27.4 27.3	27.3	27.3	27.2
low clouds (%)	34.3	59.3 59.1	44.2 43.8	44.3 43.8	49.7	49.7	44.2 43.8
total clouds (%)	53.9	70.0 69.9	64.6 64.4	64.7 64.5	68.3	68.3	64.5 64.3

^a Net chemical tendency of O₃.

^b Top of the atmosphere (TOA) residual.

^c Downwelling solar flux at surface.

^d Clearsky downwelling solar flux at surface.


Table 2. Measurements from aircraft campaigns used in this study, ~~starting 1995.~~

Campaign	Year	Months	Platform	O ₃	CO	NO	NO ₂	NO _y	PAN	HNO ₃	OH	H ₂ O ₂	C ₂ H ₆	C ₃ H ₈	C ₂ H ₄	C ₂ H ₂	SO ₂	SO ₄
TOTE	1995	12	DC-8	x	x	x		x										
VOTE	1996	01	DC-8	x	x	x		x										
STRAT	1995/96	01–12	ER-2	x				x										
PEM-Trop-A	1996	08–10	P3/DC-8	x	x	x	x		x	x	x	x	x	x	x	x	x	x
SONEX	1997	10–11	DC-8	x	x	x		x	x		x	x	x	x	x	x	x	x
POLARIS	1997	04–06, 09	ER-2	x	x								x	x			x	
POLINAT-2	1997	09–10	Falcon	x	x	x	x						x	x			x	
PEM-Trop-B	1999	03–04	P3/DC-8	x	x	x			x	x	x	x	x	x	x	x	x	x
ACCENT	1999	04, 09–10	WB57	x	x													
SOS	1999	06, 07	NOAA WP-3D	x	x	x	x	x		x								x
SOLVE	99/00	12, 03	DC-8	x	x	x		x										
SOLVE	99/00	12–03	ER-2	x	x													
TOPSE	2000	02–05	C130	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x
TRACE-P	2000	02–04	P3/DC8	x	x	x	x		x	x	x	x	x	x	x	x	x	x
TexAQS	2000	08, 09	NOAA WP-3D	x	x	x	x	x	x	x			x	x	x	x	x	x
ITCT	2002	04, 05	NOAA WP-3D	x	x	x	x	x	x	x			x	x	x	x	x	x
Crystal Face	2002	06–07	WB57	x	x					x								
INTEX-A	2004	03–08	DC8	x	x	x	(NO ₂)		x	x	x	x	x	x	x	x	x	x
NEAQS-ITCT	2004	07, 08	NOAA WP-3D	x	x	x	x	x	x	x			x	x	x	x	x	x
Ave Fall	2004	10, 11	WB57	x	x					x								
Ave Houston	2005	06	WB57	x	x					x		x						
Polar Ave	2005	01, 02	WB57	x	x		(NO ₂)			x								
Cr-Ave	2006	01, 02	WB57	x	x					x								
INTEX-B	2006	03–08	DC8	x	x	x	(NO ₂)		x	x	x	x	x	x	x	x	x	x
TexAQS	2006	09, 10	NOAA WP-3D	x	x	x	x	x	x	x			x	x	x	x	x	x
TC4	2007	07	WB57	x	x								x					
ARCPAC	2008	03, 04	NOAA WP-3D	x	x	x	x	x	x	x								x
ARCTAS	2008	04–06	DC-8	x	x	x	(NO ₂)		x	x	x	x	x	x	x	x	x	x
START08	2008	04–06	G5	x	x	x		x									x	x
CalNex	2010	05, 06	NOAA WP-3D	x	x	x	x	x	x	x								x

Table 3. Summary of abbreviations used in this article.

Abbreviation	Definition
AERONET	AErosol RObotic NETwork
ACCMIP	Atmospheric Chemistry and Climate Model Intercomparison Project
AMWG	Atmospheric Model Working Group
AOA	age of air
AOD	aerosol optical depth
BAM	bulk aerosol model
BC	black carbon
BDC	Brewer Dobson Circulation
CAM	Community Atmosphere Model
CCMI	Chemistry Climate Model Initiative
CESM	Community Earth System Model
FR	free running
HIAPER	High-Performance Instrumented Airborne Platform for Environmental Research
HIPPO	HIAPPER HIAPER Pole-to-Pole Observations
IMPROVE	Interagency Monitoring of Protected Visual Environments
MACCcity	MACC / CityZEN EU projects
MAM	modal aerosol model
MEGAN	Model of Emissions of Gases and Aerosols from Nature
MERRA	Modern-Era Retrospective Analysis For Research And Applications
MLS	Microwave Limb Sounder
MOPITT	Measurements of Pollution in The Troposphere
MOZART	odel for Ozone and Related chemical Tracers
NCEP	National Centers for Environmental Prediction
NH	Northern Hemisphere
OC	organic carbon
OMI	Aura Ozone Monitoring Instrument
POM	primary organic matter
SAD	surface area density
SD	specified dynamics
SH	Southern Hemisphere
SOA	secondary organic aerosols
STE	stratosphere to troposphere exchange
TTL	tropical tropopause layer
VOCs	Volatile Organic Compounds
WMO	World Meteorological Organization

CAM5-chem minus CAM4-chem



SD-CAM5-chem minus SD-CAM4-chem




Figure 1. Comparison of ozone, tropospheric surface area density (SAD TROP), temperature, zonal wind, relative humidity (RH), and cloud fraction, between CAM5-chem and CAM4-chem (row 1–3), and between SD-CAM5-chem and ~~SD-CAM4-chem~~ SD-CAM4-chem (row 4).

CAM5-chem minus SD-CAM5-chem




Figure 2. Comparison of ozone, tropospheric surface area density (SAD TROP), temperature, zonal wind, relative humidity (RH), and cloud fraction, between CAM5-chem and SD-CAM5-chem.

CAM5-chem minus SD-CAM5-chem




Figure 3. Comparison of ozone, nitric acid, ozone production, lightning NO_x, carbon monoxide, NO_x, hydroxyl radical, and water vapor, between CAM5-chem and SD-CAM5-chem.




Figure 4. Zonal average water vapor tape recorder (in ppm) of different model configurations, CAM4-chem (top left), CAM5-chem (top right), SD-CAM4-chem (middle left), SD-CAM5-chem (middle right) and MLS satellite observations averaged over year 2005–2011 (bottom panel), composited over 12 months for all simulated years, and repeated over 24 months.




Figure 5. Age of air of different model configurations and simulated years for CAM4-chem (top left), CAM5-chem (top right), SD-CAM4-chem (bottom left), SD-CAM5-chem (bottom right).




Figure 6. Comparison of aerosol burden (left) and surface area density (right) between SD-CAM5-chem and SD-CAM4-chem of SO₄sulfate aerosol (SO₄), SOA, and BC.




Figure 7. Comparison between IMPROVE network observations over the US in winter (December/January/February) in comparison to SD-CAM5-chem (blue) and ~~SD-CAM5-chem~~ SD-CAM4-chem (red) for SO₂ (left) and sulfate aerosol (SO₄) (right) and different seasons, DJF (top) and JJA (right). The median and correlation coefficient (R) between observations and model results are given on the top left of each panel.




Figure 8. Comparison of SO₂ (left) and [sulfate aerosol](#) (SO₄) (right) between different model configurations and aircraft observations over the US (two left columns) and at high latitudes (2 right columns). Black lines show the median of aircraft profiles and error bars indicate describe the range between the 25th and 75th percentile of the distribution. Model results are averaged over the region and months of each campaign.




Figure 9. HIPPO BC observations for different HIPPO aircraft campaigns taken over the Pacific (left column) and differences between the different model configurations and observations, CAM4-chem (second column), CAM5-chem (third column) and CAM5-MAM4-chem (fourth column).




Figure 10. Aerosol optical depth at 550 nm for CAM4-chem (left) and CAM5-chem (right) in comparison to the satellite and AERONET composite Kinne (2009) (middle). Differences [between model and observations](#) are show in the bottom row. Numbers in the parenthesis are the global average AOD over only areas where the satellite composite has a valid value.




Figure 11. Differences between model results and observations of zonally averaged CO column below 100hPa from the present-day MOPITT climatology (left), and OMI/MLS tropospheric and stratospheric column climatology (right).

Comparison to Ozonesondes




Figure 12. Taylor-like diagram comparing the mean and correlation of the seasonal cycle between observations using a present-day ozonesonde climatology between 1995–2011 and model results, interpolated to the same locations as sampled by the observations and for different pressure levels, 900 hPa (top panel) and 500 hPa (bottom panel). Different numbers are correspondent to a specific region, as defined in Tilmes et al. (2012). Left panels: 1 – NH-Subtropics; 2 – W-Pacific/East Indian Ocean; 3 – equat. Americas; 4 – Atlantic/Africa. Middle panels: 1 – Western Europe; 2 – Eastern US; 3 – Japan; 4 – SH Mid-Latitudes. Right panels: 1 – NH Polar West; 2 – NH Polar East; 3 – Canada; 4 – SH Polar.




Figure 13. Seasonal cycle comparison between observations using a present-day ozonesonde climatology between 1995–2011 (black) and model results: CAM5-chem (cyan) and CAM4-chem (orange), SD-CAM5-chem (blue) and SD-CAM4-chem (red). Model results are interpolated to the same locations as sampled by the observations and for different pressure levels, 900 hPa (top panel) and 500 hPa (bottom panel) for selected regions. The standard deviation of ozonesonde observations is shown as error bars and the mean and correlation of the seasonal cycle between observations and model results are printed on the top of each figure.




Figure 14. Probability distribution function (PDF) of the regionally-aggregated ozone distribution for Western North America, Eastern North America, and Western Europe from surface ozone observations (grey shaded area) in comparison to regionally-aggregated ozone distributions from model results interpolated to the location of the ozone stations (different colors), for winter (left) and summer (right).

Comparison to Ozonesondes




Figure 15. As Fig. 12, but for different pressure levels, 250 hPa (top panel) and 50 hPa (bottom panel). Different numbers are correspondent to a specific region, as defined in Tilmes et al. (2012). Left panels: 1 – NH-Subtropics; 2 – W-Pacific/East Indian Ocean; 3 – equat. Americas; 4 – Atlantic/Africa. Middle panels: 1 – Western Europe; 2 – Eastern US; 3 – Japan; 4 – SH Mid-Latitudes. Right panels: 1 – NH Polar West; 2 – NH Polar East; 3 – Canada; 4 – SH Polar.




Figure 16. As Fig. 13, but for different pressure levels, 250 hPa (top panel) and 50 hPa (bottom panel).




Figure 17. Relative differences between aircraft observations and different model configurations (different colors) over different regions and seasons as listed in Table 1 and sorted with regard to season and location (see text for more details), averaged over 2–7 km, for O_3 , NO_x , NO_y , PAN, and HNO_3 .




Figure 18. HIPPO O₃ observations for different HIPPO aircraft campaigns taken over the Pacific, left column, and differences between the different model configurations and observations, CAM4-chem (second column), CAM5-chem (third column) and SD-CAM5-chem (fourth column).




Figure 19. Regional comparison of CO column for different months, between CAM5-chem model results and MOPITT observations. Model results are shown on the left, and differences between CAM5-chem and MOPITT on the right. The MOPITT averaging kernels and a priori are applied to the model results to account for the a priori dependence and vertical resolution of the MOPITT data.




Figure 20. As Fig. 17, but instead for CO, C₂H₆, C₃H₈, and C₂H₂.




Figure 21. Comparisons of vertical profiles of ozone, CO, NO_x and PAN, from different tropical aircraft campaigns and different model configurations. Black lines show the median of aircraft profiles and error bars indicate describe the range between the 25th and 75th percentile of the distribution. Model results are averaged over the region and months of each campaign.




Figure 22. HIPPO PAN observations for different HIPPO aircraft campaigns taken over the Pacific, left column, and differences between the different model configurations and observations, CAM4-chem (second column), CAM5-chem (third column) and SD-CAM5-chem (fourth column).




Figure 23. Correlations between tropospheric OH burden, methane lifetime, and CO, for different simulations. OH and CO burden are column integrated tropical averages (30° S– 30° N). Each symbol of each configuration (see legend) represents an annual average value.




Figure 24. Column integrated [tropospheric and](#) tropical OH burden in (30°S – 30°N), left top panel, and OH burden, adjusted to a reference SAD value (see text) for the other panels, in correlation to different variables [that are integrated over the same region](#). Each symbol of each configuration (see legend) represents an annual average value.




Figure 25. Correlations of tropospheric column integrated NO_x to column integrated lightning NO_x over the tropics (left panel); correlation of OH burden, adjusted to a reference SAD value (see text) to column integrated HNO₃ over the tropics (middle panel); correlations of column integrated HNO₃ to column integrated lightning NO_x over the tropics (right panel).