

1 **Technical challenges and solutions in representing lakes**
2 **when using WRF in downscaling applications**

3 **M. S. Mallard^{1,*}, C. G. Nolte¹, T. L. Spero¹, O. R. Bullock¹, K. Alapaty¹, J. A.**
4 **Herwehe¹, J. Gula², and J. H. Bowden³**

5 [1]{National Exposure Research Laboratory, U.S. Environmental Protection Agency, Research
6 Triangle Park, North Carolina, USA }

7 [2]{Department of Atmospheric and Oceanic Sciences, University of California, Los Angeles,
8 California, USA }

9 [3]{Institute for the Environment, University of North Carolina, Chapel Hill, North Carolina,
10 USA }

11 *Currently at: Institute for the Environment, University of North Carolina, Chapel Hill, North
12 Carolina, USA

13 Correspondence to: M. S. Mallard (mmallard@email.unc.edu)

14 [Revised manuscript: 21 January 2015](#)

15
16 **Abstract**

17 The Weather Research and Forecasting (WRF) model is commonly used to make high resolution
18 future projections of regional climate by downscaling global climate model (GCM) outputs.

19 Because the GCM fields are typically at a much coarser spatial resolution than the target regional
20 downscaled fields, inland lakes are often poorly resolved in the driving global fields, if they are
21 resolved at all. In such an application, using WRF's default interpolation methods can result in
22 unrealistic lake temperatures and ice cover at inland water points. Prior studies have shown that
23 lake temperatures and ice cover impact the simulation of other surface variables, such as air
24 temperatures and precipitation, two fields that are often used in regional climate applications to
25 understand the impacts of climate change on human health and the environment. Here,
26 alternative methods for setting lake surface variables in WRF for downscaling simulations are
27 presented and contrasted.

28

29 1 Introduction

30 When using global climate model (GCM) fields to drive finer-scale regional climate model
31 (RCM) runs, typically the RCM does not have an oceanic or lake physics component and relies
32 on the GCM output to provide all water surface temperatures and ice cover. Within a
33 downscaling simulation, by design, the GCM is at a coarser spatial resolution than the RCM, so
34 inland water bodies in the region being simulated are either poorly resolved or not resolved by
35 the GCM. ~~Some RCM configurations do include an oceanic component. The~~ Prior to 2013, the
36 Weather Research and Forecasting (WRF) model (Skamarock et al., 2008) required exogenously
37 prescribed water surface temperatures, as there was not capability to prognosticate water
38 temperatures. WRF has included an optional coupled ocean component since ~~WRF~~-version 3.5
39 was released in April 2013 (~~Skamarock et al., 2008~~; WRF User's Guide, 2014). Other RCMs
40 have been coupled to ocean models in order to simulate regions around the Arctic,
41 Mediterranean Sea, and Indian Ocean (e.g., Rinke et al., 2003; Ratnam et al., 2009; Artale et al.,
42 2010; Gualdi et al., 2013). However, when using WRF's default configuration, the sea surface
43 temperature (SST) fields used during the simulation are calculated from the driving data during
44 the preprocessing steps performed before WRF runs the simulation; during the model run, these
45 prescribed water temperatures are input at a user-specified frequency which is usually daily or
46 sub-daily. Similarly, lake surface temperatures (LSTs) and lake ice cover are prescribed by
47 spatial interpolation from the ~~sea surface temperature (SST)~~ and sea ice fields in the driving data.
48 In this study, we examine the use of the Advanced Research WRF (Skamarock and Klemp,
49 2008) model applied as an RCM in regions where the driving larger-scale data have a poor
50 representation of lakes.

51
52 When the WRF Preprocessing System (WPS) interpolates skin temperatures from the coarser
53 global dataset (where both land and water temperatures are included in a single field), masks are
54 applied such that water temperatures from the GCM are used to set water temperatures on the
55 finer, target grid. Using the standard ~~interpolation~~-methods in WPS, interpolation is first
56 attempted using 16 surrounding grid cells in the coarser grid; if this method fails due to a lack of
57 the requisite 16 valid data points, WPS attempts other interpolation techniques using as many as
58 four grid cells and as few as one. While a full description of all WPS interpolation techniques is
59 beyond the scope of this study, more information is available in the WRF User's Guide (2014, p.

60 [3-56 to 3-59](#)). When all other methods fail due to the lack of nearby water grid cells, WPS
61 defaults to the “search” approach, in which the nearest water point is used to set LSTs (~~WRF~~
62 ~~User’s Guide, 2014~~). When employing the search option, water cells in the driving data are often
63 distant from and unrepresentative of the target cell in the WRF domain. The search option in
64 WPS performs no interpolation or averaging, sometimes resulting in abrupt, non-physical
65 temperature discontinuities.

66
67 Here we show the result of using this default methodology to downscale 1° Community Earth
68 System Model (CESM) fields to a 36 km WRF domain (198 × 126) covering the continental US,
69 and subsequently similar examples in other downscaling studies are discussed. However, it
70 should be noted that the use of CESM as an example is arbitrary because similar results have
71 been obtained with other global datasets as well. The CESM ocean mask, used to interpolate the
72 GCM’s SST fields to the WRF grid, has no water grid cells over the North American interior
73 (Fig. 1). As a result, water temperatures in Hudson Bay are used to set temperatures over the
74 larger westernmost areas of the Laurentian Great Lakes, while LSTs in the southeastern areas of
75 the Great Lakes are set by Atlantic SSTs (Fig. 2). At the time shown in Fig. 2, the LSTs
76 interpolated from CESM onto the 36 km WRF grid contain discontinuities of approximately 17
77 K between adjacent grid cells in Lakes Michigan and Huron, while a smaller discontinuity of
78 approximately 3 K is created in Lake Superior. It should be noted that various interpolation
79 options are available in WPS and can be specified by the user. The description in the paragraph
80 above is representative of the interpolation process as defined by WPS’s default settings. Even
81 though this process could be changed by the model user, the key issue remains that when lakes
82 are poorly represented or completely absent, the problem of how to specify the lake state is not
83 amenable to any interpolation method.

84
85 The problems of using larger-scale data to define LSTs with the default options in WPS are not
86 limited to the Great Lakes. None of the inland lakes resolved by WRF at 36 km have valid LSTs
87 in the CESM ocean mask (Fig. 1). Using the search option in WPS results in setting the LSTs to
88 unrealistic values throughout the domain. Temperatures in Pyramid Lake, Great Salt Lake, as
89 well as several smaller lakes east of the Rocky Mountains in both Canada and the US are
90 assigned from the Pacific Ocean (Fig. 2), while lake temperatures in the southeastern and central

91 US are set from SSTs in the Gulf of Mexico and Atlantic Ocean. Two adjacent grid cells
92 representing Lake Sakakawea in North Dakota are assigned LSTs differing by approximately 10
93 K because the western cell is set from the Pacific while the eastern cell is prescribed from
94 Hudson Bay (Fig. 2). Using any interpolation method to assign LSTs when no suitable data are
95 available will adversely affect the accuracy of downscaled simulations that are based on forcing
96 from those LSTs.

97
98 Mallard et al. (2014; hereafter M14) also discuss problems that arise when downscaling coarse
99 global data to a 12 km grid covering the eastern US. In M14, the National Centers for
100 Environmental Prediction (NCEP)–Department of Energy Atmospheric Model Intercomparison
101 Project (AMIP-II) reanalysis (hereafter R2; Kanamitsu et al., 2002) is used to drive historical
102 simulations as a proxy or stand-in for a similarly-coarse GCM. In contrast to the CESM example
103 discussed above, R2 has at least a partial representation of western Great Lakes, but nevertheless
104 has only three inland water points to represent all five of the Great Lakes (Fig. 1 of M14).
105 Therefore, using the standard interpolation methods with R2 results in unrealistically large,
106 abrupt, and non-physical LST discontinuities in eastern Lake Erie and Lake Ontario, where water
107 temperatures are set using Atlantic SSTs, while the LSTs in western Lake Erie and in the three
108 western Great Lakes are interpolated from the three lake cells in R2 (M14).

109
110 In WRF, ice cover can either ~~be treated as a binary field, which is set based on whether the water~~
111 ~~temperature is below a threshold, or it can~~ be interpolated from the driving data and ~~prescribed as~~
112 ~~assigned to~~ covering some fraction of a grid cell, ~~or it can be treated as a binary field that is set~~
113 ~~In the former approach, WRF sets ice cover~~ to 100 % at grid ~~points~~ cells where the water surface
114 temperatureLST drops below a specified threshold.~~271 K, slightly below the freshwater freezing~~
115 ~~temperature of approximately 273 K. Note that t~~ The default threshold value was ~~changed from~~
116 ~~271 K (slightly below the freshwater freezing temperature of approximately 273 K), but it was~~
117 ~~changed~~ to 100 K as of version 3.5.1 (~~September 2013~~), ~~presumably~~ to avoid the unintended
118 creation of ice by this method when using WRF's default settings (Table 1). When fractional ice
119 values are prescribed from the driving dataset, the WPS methods applied to interpolate sea and
120 lake ice differ from those used for SSTs and LSTs. If there are no surrounding water grid cells in
121 the driving dataset, an ice cover value of zero is assigned rather than employing the search

122 method. When M14 downscaled ice cover from R2, it was shown ice concentrations of zero were
123 applied to points through Lakes Huron, Erie and Ontario throughout a two-year simulation (Fig.
124 3 of M14), even though partial ice coverage was observed on all three lakes during that historical
125 period. Moreover, almost complete ice coverage of Lakes Superior and Michigan occurred in a
126 single day (M14). Wang et al. (2012a) conducted a climatology of ice cover in the Great Lakes
127 over the period 1973 to 2010 and showed that, in the average seasonal cycle of ice cover, the
128 maximum fractional coverage of Lake Superior was approximately 50 % (their Fig. 3). Although
129 Wang et al. noted that the standard deviation of ice cover is quite large (exceeding the mean
130 values in some of the Great Lakes), the seasonal cycles in their study showed the accumulation
131 of ice coverage over months, not the abrupt appearance of lake-wide ice over daily periods.
132 Ultimately, M14 improved the representation of the Great Lakes in their downscaled simulations
133 by applying a coupled lake model, which will be discussed further in a subsequent section.
134 Whereas M14 showed the results of using a single lake model, the current work presents a
135 broader range of approaches, recognizing that the most preferable method to represent lake fields
136 may vary between different RCM applications.

137
138 Prior studies downscaling other global datasets and GCMs have also noted findings similar to the
139 example shown here (Fig. 2) and the results of M14. Using WRF as an RCM over Eastern
140 Africa, Argent (2014) showed that the use of WPS's default interpolation methods resulted in
141 oceanic temperatures from a global SST dataset applied to set LSTs throughout Lake Victoria.
142 Discontinuities in LSTs with WRF were noted in the Great Lakes basin by Bullock et al. (2014)
143 who downscaled R2 to 12 km, and by Gao et al. (2012) who downscaled ~~the~~ CESM to a 4 km
144 grid. Within the downscaled simulations produced for the North American Regional Climate
145 Change Assessment Program (NARCCAP; Mearns et al., 2012), problems with producing
146 realistic LSTs and ice cover for the Great Lakes region are documented using several approaches
147 with various RCMs, including WRF (NARCCAP, 2014). For some NARCCAP model
148 configurations, caution is recommended when using surface variables in the region surrounding
149 the Great Lakes. Previous work examining the value of dynamical downscaling has noted that
150 downscaled simulations have the most potential to add value relative to GCM simulations in
151 areas of complex topography and along coastlines because of increased resolution in regional
152 models (e.g., Feser et al., 2011). Although RCMs better resolve the coastlines (and therefore, the

153 presence of lakes) than the driving GCMs, using erroneous LSTs and lake ice cover could impair
154 the simulation of interactions between lakes and overlying air masses. The potential benefits
155 gained by downscaling to a grid spacing that better resolves land–water interfaces may not be
156 realized if the lake state (defined here by LSTs and ice) is unrealistically represented. Even as
157 additional computing resources allow GCMs to increase in resolution and better represent lakes,
158 RCMs will also be run at finer scales; therefore, it can be expected that smaller lakes with
159 important effects on mesoscale and microscale climatology will continue to be unresolved by the
160 driving data sets.

161
162 The purposes of this paper are to describe various techniques that can be used to set LSTs and
163 lake ice cover in the WRF model for downscaling, and to discuss the benefits and possible
164 shortcomings of each approach. The effects of these techniques on simulated lake–atmosphere
165 interactions, both in the present climate and in future climate states, are discussed in context with
166 relevant previous literature.

167

168 **2 Comparison of methods**

169 As will be shown below, choice of the appropriate methodology for representing a lake in a
170 downscaling configuration is dependent on what interactions must be simulated between the
171 atmospheric fields and the lake state and how the lake state is expected to be impacted by climate
172 change when downscaling future GCM projections. In regional climate simulations conducted
173 over the continental US, the Laurentian Great Lakes are a prominent feature, as Lake Superior is
174 the largest freshwater lake in the world (by surface area) at over 82 000 km². Several studies
175 have concluded that the Great Lakes strongly influence the surrounding regional climate,
176 moderating extremes in near-surface temperatures, and affecting precipitation and passing
177 cyclones and anticyclones on an annual cycle (e.g., Wilson, 1977; Bates et al., 1993; Scott and
178 Huff, 1996; Notaro et al., 2013). Climatologically, the greater heat capacity of the lakes serves to
179 enhance precipitation and convection during September to March, when warmer surface water
180 (relative to low-level atmospheric temperatures) reduces atmospheric stability (e.g., Notaro et al.,
181 2013). Conversely, the slower warming of the lakes in boreal spring results in the opposite effect
182 during the April–August period, where the relatively cool lakes enhance atmospheric stability
183 and reduce precipitation and convection. These periods are referred to as the lake unstable and

184 lake stable seasons, respectively. Lake-effect precipitation has also been documented outside the
185 Great Lakes as well, such as in Lake Champlain (Tardy, 2000; Laird et al., 2009), Lake Tahoe
186 (Cairns et al., 2001), and the Great Salt Lake (Carpenter, 1993; Steenburgh and Onton, 2001). A
187 review by Schultz et al. (2004) states that lake-effect snowfall has been observed to occur over
188 lakes with fetches of only 30 to 50 km, citing prior studies over Bull Shoals Lake of Arkansas
189 (Wilken, 1997) as well as Lake Tahoe and Pyramid Lake in Nevada (Cairns et al., 2001; Huggins
190 et al., 2001). Interactions between the lakes and surrounding regions are also strong in tropical
191 environments as well. For example, the immediate region surrounding Lake Victoria in Africa
192 has the highest recorded frequency of thunderstorms in the world with approximately 300 storm
193 days per year (Asnani, 1993). Overall, while a comprehensive review of the impact of each lake
194 on regional climate is beyond the scope of this study, prior work indicates that even lakes that are
195 smaller than the Great Lakes can be anticipated to have substantial effects on regional climate.

196

197 Prior studies have also illustrated that even relatively small errors in prescribed LSTs in a
198 downscaling configuration can adversely affect simulated precipitation in regions surrounding
199 lakes. The sensitivity study of Wright et al. (2013) showed significant changes in lake-effect
200 snowfall over the Great Lakes in idealized simulations where LSTs were uniformly warmed by 3
201 °C. Anyah and Semazzi (2004) simulated changes in the spatial patterns and intensity of
202 precipitation, as well as the amount of evaporation, over Lake Victoria in a modeling study
203 where LSTs were uniformly changed by only 1.5 °C.

204

205 Interactions between the lakes and overlying air masses are also governed by the amount of lake
206 ice in climates that permit lakes to freeze. Previous studies have found the presence of ice
207 suppresses turbulent latent and sensible heat fluxes from the lake to the air mass (e.g., Zulauf and
208 Krueger, 2003; Gerbush et al., 2008). As shown in the lake-effect snow case studies simulated by
209 Wright et al. (2013), the presence of ice coverage over the lake's surface inhibits downstream
210 precipitation. As a result, lake-effect snowfall decreases in some areas surrounding the Great
211 Lakes during the later portion of the lake unstable season, as the water's surface freezes during
212 the winter and early spring months. Overall, past studies indicate that if LSTs and ice are not
213 properly prescribed, inaccurate values of precipitation and temperature in the lee of lakes result

214 | ~~in~~-from a downscaled simulation.

215

216 **2.1 WRF's alternative lake setting**

217 Since the release of WRF version 3.3 in April 2011, an “alternative initialization of lake SSTs”
218 option is provided in WPS to set LSTs (WRF User’s Guide, 2014; [Table 1](#)). When employing
219 this method, LSTs can be set using temporally averaged 2 m air temperatures from the driving
220 data set, with the averaging period set by the user. Bullock et al. (2014), when downscaling a
221 proxy GCM (R2) over a 12 km grid covering the Great Lakes, attempted to use the alternative
222 lake setting to account for the greater thermal inertia of the Great Lakes by incorporating
223 seasonal temperature changes after a one-month time lag. Following the procedure of Bullock et
224 al., if a user were to perform a simulation over the month of May, a single LST field would first
225 be generated by temporally averaging air temperatures during the previous month of April;
226 subsequently this static LST field would be used to set inland water temperatures throughout the
227 month of May. Because Bullock et al. (2014) preprocessed the driving data in monthly segments,
228 the LST field was prescribed to vary with time on a monthly basis. Using this method may
229 imitate the seasonal changes observed over the Great Lakes, producing a lake stable and unstable
230 season during the appropriate months. A drawback to this methodology is that the same lag time
231 is used throughout the model grid, regardless of lake depth. Therefore, in this approach, large,
232 deep lakes are implied to heat and cool on the same timescale as small, shallow lakes.
233 Meanwhile, it is expected that observed seasonal temperature changes over smaller and
234 shallower lakes would more closely follow atmospheric temperature changes than in large, deep
235 lakes. If employed for simulations outside the Great Lakes, the procedure used by Bullock et al.
236 (2014) should be modified to imitate the observed relationship between changing air
237 temperatures and LSTs.

238

239 In its default configuration used prior to the release of version 3.5.1 (~~September 2013~~), WRF
240 prescribes ice cover at grid cells where LST is less than 271 K ([Table 1](#)). This value is applied at
241 all water points regardless of salinity. As winter 2 m air temperatures are frequently below
242 freezing in the Great Lakes area, Bullock et al. (2014) found that unrealistically large spatial
243 coverage of ice occurred when using the alternative lake setting in WRF version 3.4.1, with all
244 five Great Lakes completely frozen for most of the winter. Such erroneous ice cover would be

245 expected to negatively impact the simulation of precipitation, 2 m temperatures, and other
246 variables influenced by sensible and latent heat fluxes supplied by the Great Lakes. Therefore,
247 the use of the alternative lake setting in WRF may not be appropriate in some regions where sub-
248 freezing air temperatures would result in unrealistic temporal and spatial coverage of sub-
249 freezing LSTs and ice.

250
251 However, this is not a concern for tropical lakes where air temperatures would not be sufficiently
252 low enough to result in frozen lakes. Argent (2014, Sect. 3) demonstrated the utility of the
253 alternative lake setting in WRF simulations over Lake Victoria in Eastern Africa, finding that it
254 improved the accuracy of simulated rainfall relative to the use of the default interpolation in
255 which oceanic SSTs were used to set Lake Victoria's LSTs.

256

257 **2.2 Climatological LSTs and ice**

258 Another approach for setting LSTs and lake ice coverage when downscaling with WRF is to
259 prescribe these variables from higher-resolution data sets of climatologically averaged quantities.
260 This can be viewed as assuming stationarity for the lake state as is frequently done for other
261 input variables in an RCM, such as land-use and vegetation. Even for retrospective climate
262 simulations, using this approach could be detrimental because the interannual variability of LSTs
263 and ice – and its effects on the prediction of extreme events – would not be captured using this
264 method. When making future projections, it must be considered that prior studies have shown
265 that LSTs cannot be assumed to be stationary in future warmer climates; in fact, some studies
266 conclude that non-linear feedbacks exist between regional climate change and LSTs and ice for
267 some lakes. An observational study by Austin and Colman (2007) found that the multi-decadal
268 warming trend in the Great Lakes region was amplified in the lake temperatures, relative to
269 surrounding inland temperatures, because of the earlier break-up of ice and earlier springtime
270 warming of surface water. In the downscaling simulations of Gula and Peltier (2012), increased
271 snowfall was simulated in the lee of the Great Lakes in a warmer, mid-century climate because
272 lake ice forms ~~at a later time period~~ in the winter. Gula and Peltier conclude that the impact of
273 having the lakes remain ~~open (and free of ice)~~ is that increased latent and sensible heat fluxes are
274 present for a longer time period during the lake unstable season, lessening the stability of the

275 overlying air mass and enhancing precipitation. Magnuson (2000) concluded that observed ice
276 coverage is decreasing in lakes and rivers throughout the Northern Hemisphere. Such a decrease
277 in ice coverage has been linked by observational studies to increases in lake-effect precipitation
278 in the Great Lakes region (Assel and Robertson, 1995; Burnett et al., 2003; Kunkel et al., 2009).
279 Because ice ~~supresses~~suppresses fluxes of latent and sensible heat (e.g., Zulauf and Krueger,
280 2003; Gerbush et al., 2008), decreasing ice cover in a warmer climate allows larger fluxes of
281 latent and sensible heat to modify the overlying air mass, increasing downstream precipitation
282 during the lake unstable season. None of the impacts on the lake state reviewed here (the
283 warming of LSTs and more open water from which to produce fluxes) would be considered in
284 the WRF model using LSTs and ice based on present-day climatology, and the effects of
285 changing lake conditions on atmospheric stability, humidity, precipitation and convection would
286 not be simulated.

287

288 This approach could be improved by adding a linear increase to observed LSTs over time, which
289 may be a valid approximation for the effect of climate change on some lakes. However, such an
290 approach would not capture the non-linear impacts of climate change (as described by Austin
291 and Colman, 2007) on the Great Lakes. Overall, the efficacy of using of a climatologically-based
292 approach is dependent on the amount of interannual variability, as well as the impacts of climate
293 change on the lake state and whether those effects can be accounted for by the inclusion of a
294 linear LST anomaly.

295

296 **2.3 Land mask modification**

297 To avoid the issues with LSTs discussed in Sect. 1 and illustrated in Fig. 2, Gao et al. (2012)
298 modified the GCM land mask in the Great Lakes area so that skin temperatures from land points
299 in the GCM were used to set LSTs on the WRF grid in their downscaled simulations. This
300 treatment successfully eliminated the abrupt temperature discontinuities (such as those in Fig. 2)
301 produced by interpolating a coarse data set. However, the effects of the lakes themselves are lost
302 if GCM land temperatures are used to prescribe RCM water temperatures and the lake-land
303 temperature contrasts, with their associated mesoscale phenomena such as lake breezes and lake-
304 effect precipitation, are eliminated. Notaro et al. (2013) conducted an idealized modeling

305 experiment where the Great Lakes were replaced with forest and field land cover types. They
306 found that the presence of the lakes affected precipitation, 2 m air temperatures and their
307 variability, water vapor, cloud cover, incoming shortwave radiation, the hydrological budget and
308 the intensity of passing cyclones and anticyclones. The approach used by Gao et al. (2012),
309 where land surface temperatures from the GCM are used to specify water temperatures, partially
310 accounts for some lake effects (such as changes in surface friction and albedo) because WRF
311 would recognize the presence of a water surface. However, all processes related to the LST (e.g.,
312 ice formation, latent and sensible heat flux, 2 m temperature and moisture values, outgoing
313 longwave radiation from the surface) would be negatively impacted by this treatment.
314 Additionally, some impacts of climate change on the future lake state could be lost. For example,
315 the amplification of Great Lakes LSTs, relative to over-land temperatures, observed by Austin
316 and Colman (2007) will not be captured if land temperatures are used to set LSTs.

317

318 **2.4 Use of simulated lake fields from GCM**

319 A more sophisticated class of approaches for better representing the lake state in a downscaling
320 configuration involves the use of a lake model. This can be done either by using outputs from the
321 GCM's lake model (if available), driving a stand-alone lake model offline with GCM fields to
322 simulate LSTs and ice, or by coupling a lake model to the RCM when downscaling. The CESM
323 | ~~model~~ has a lake model embedded within ~~the-its~~ land ~~surfacecomponent~~ model (LSM), version 4
324 of the Community Land Model (CLM4). CLM4 accounts for the presence of subgrid-scale lakes
325 using the one-dimensional lake model described in Oleson et al. (2010). It is a column model
326 partially based on the Hostetler lake model (e.g., Hostetler and Bartlein, 1990; Hostetler et al.,
327 1993, 1994), and it simulates 10 water layers through the depth of the lake, as well as additional
328 layers for thermally-active soil underneath and snow and ice above. However, when producing
329 the downscaled simulation shown in Fig. 2, output from CLM's lake model was not easily
330 accessible with other CESM outputs from the same simulation within archiving systems such as
331 the Earth System Grid Federation. Lake temperatures and ice from CESM, and other GCMs with
332 embedded lake models, could be leveraged by RCMs such as WRF to account for the impact of
333 climate change on the lake state. In areas where lakes are at least partially resolved by the GCM,
334 this approach would be effective at driving the RCM with simulated changes in LSTs and ice

335 cover consistent with future projections and at keeping the RCM solution in the regions affected
336 by lakes consistent with the GCM simulation. However, some small lakes may remain
337 unrepresented by GCM data.

338

339 **2.5 Use of a stand-alone lake model**

340 If lake model outputs from the GCM are unavailable, one alternative is to use a standalone lake
341 model driven by GCM fields to downscale the lake state in a manner which is consistent with the
342 GCM's atmospheric fields. In the downscaling experiments performed by Gula and Peltier
343 (2012) over the period 2050–2060, the Freshwater Lake (FLake) model was utilized to provide
344 simulated LSTs and lake ice to WRF in the Great Lakes basin. GCM fields from the Community
345 Climate System Model, with a spectral resolution of T85 (~ 1.4° grid spacing), were used to
346 drive a FLake simulation on a 10 km regional grid, and the LSTs and ice cover simulated by
347 FLake were subsequently used to drive the downscaled WRF simulation. In this 1-way WRF-
348 FLake model configuration, changes in LSTs and ice respond to changes in atmospheric
349 variables in the driving GCM, but the lake model output is produced on the higher-resolution
350 regional WRF grid. FLake is a **1-D** column model which is highly reliant on empirical
351 relationships and has been used in several studies with other RCMs (e.g., Mironov, 2008;
352 Kourzeneva et al., 2008; Martynov et al., 2008; Mironov et al., 2010; Samuelsson et al., 2010).
353 FLake requires a 2-D field of lake depths and the 1-D column model is called at each point.
354 Therefore, the simulated LSTs are sensitive to lake depth, as well as the driving GCM fields.
355

356 **2.6 Use of a coupled lake model within an RCM**

357 In WRF version 3.6, ~~released April 2014~~, a CLM-based lake model can be utilized with other
358 non-CLM land surface models (WRF User's Guide, 2014; [Table 1](#)). This lake model is taken
359 from CLM version 4.5 (Subin et al., 2012; Oleson et al., 2013) with some modifications by Gu et
360 al. (2013) as discussed further below. Although a version of CLM4 was available as an **LSMland**
361 ~~surface model~~ option within WRF version 3.5 (~~released April 2013~~), the lake model in CLM4
362 was disabled in WRF ([Table 1](#)). In WRF version 3.6, CLM's Hostetler-based lake model can be
363 applied by using horizontally varying lake depths (which are available in WPS version 3.6) or a

364 uniform lake depth can be assigned to all lakes at runtime. Gu et al. (2013) demonstrated WRF-
365 CLM's performance in the Great Lakes region using a previous version of this model
366 configuration (WRF 3.2 and CLM 3.5) to simulate a 16 month period from 2001 to 2002 at 10
367 km grid spacing. It was shown that the lake model simulated LSTs well in Lake Erie but
368 generated large biases in LSTs when compared to buoy observations in Lake Superior. However,
369 the LST bias was reduced by reformulating the eddy diffusivity parameter in the CLM lake
370 model, and it was concluded that the updated lake model within WRF-CLM was reasonably able
371 to reproduce observed LSTs. However, no ice was observed during the period and the ability of
372 WRF-CLM to accurately simulate ice cover was not examined in Gu et al. (2013).

373

374 In an alternative coupled approach, the prior work of Gula and Peltier (2012) has been updated
375 with the option of using WRF-FLake as a 2-way coupled model, where atmospheric variables
376 simulated by WRF are used by FLake at each time step in the WRF model, and simulated LSTs
377 and ice thicknesses are provided back to WRF by FLake. M14 concluded that the use of WRF-
378 FLake resulted in a more accurate representation of LSTs and lake ice, relative to interpolation
379 from ~~a proxy GCM~~, the R2. Substantial improvements were shown in the simulation of the
380 temporal and spatial variability of ice cover, and errors in LSTs were reduced by the use of the
381 coupled model. Similar to Martynov et al. (2010), M14 found that FLake performed worst in the
382 largest and deepest lake (Lake Superior) and best for the smallest and shallowest (Lake Erie).

383

384 When using an embedded lake model within an RCM, it can be anticipated that the period of
385 time needed for spin-up could be larger than it is when all water conditions are simply
386 prescribed. To spin-up the WRF-FLake model in M14, the stand-alone version of the FLake
387 model was driven with atmospheric conditions from the proxy GCM ~~for 10 annual cycles to~~
388 ~~achieve equilibrium, as adapted from the~~ in a spin-up procedure recommended by Mironov et al.
389 (2010) when using FLake. In this methodology, the initial year of the simulation ~~(2005)~~ is
390 “looped” over 10 annual cycles with meteorological variables from the initial year repeatedly
391 used to force the lake model, and the lake state at the end of each year used to initialize FLake
392 for the start of the next year until, ensuring that the simulated lake state converges to equilibrium
393 with these atmospheric conditions ~~is achieved by the end of the 10-cycle simulation~~. Output from
394 the first year of this offline simulation is shown in Fig. 3 illustrating the adverse effects of using

395 | FLake output without adequate spin-up time. A time-series taken from a representative point in
396 | Lake Superior shows unrealistically cool LSTs (below 200 K) occurring during the initial
397 | months of the simulation. Also during this period, unrealistically large ice coverage formed,
398 | freezing over all five Great Lakes. The observed ice cover plotted in Fig. 3 is much more limited
399 | in its spatial extent. ~~as shown~~ Observed ice cover is plotted from ~~the~~ National Ice Center (NIC)
400 | Great Lakes Ice Analysis charts, which are processed and provided by the Great Lakes
401 | Environmental Research Laboratory (GLERL; Wang et al., 2012b). The FLake model results
402 | obtained after the spin-up period showed realistic values of LSTs and ice cover (M14).

403
404 | To examine how WRF-CLM reacts during the initial months of a simulation, without any spin-
405 | up time, output from a 12 km WRF-CLM simulation (version 3.6) is shown in Fig. 4. In this
406 | simulation, the same methods as in M14 are followed but with the following changes: the model
407 | version is updated from 3.4.1 to 3.6, the CLM lake model is used in place of FLake, and no spin-
408 | up procedure is employed for initialization of the lake model (initial LSTs are interpolated from
409 | R2). As in M14, the Noah ~~LSM~~ land surface model (Chen and Dudhia, 2001) is used. Similar to
410 | the example shown in Fig. 3, significant overestimation of ice coverage occurs during the first
411 | year (Fig. 4). Although some adverse effects in this simulation are introduced due to the use of
412 | LSTs interpolated from the coarse R2 data to provide an initial state, the similarity of these
413 | results to FLake's fields in Fig. 3 suggests that the lack of spin-up time is a common problem to
414 | both model runs. It is also implied by the methodologies of other CLM-based studies, which do
415 | use spin-up or initialization procedures. Previous work by Subin et al. (2012) with the lake
416 | model in CLM4 used a 110-year period for the spin-up of their reference simulation. In their
417 | experiments with WRF-CLM, Gu et al. (2013) used an observed LST field for initialization. The
418 | 9 sub-surface layers in their model were initialized based on the shape of an observed profile of
419 | lake temperatures, valid during that period of the year and taken from Lake Superior. Using this
420 | initialization methodology for a future downscaled simulation is not possible due to lack of
421 | observations, but simulated future lake profiles could possibly be utilized for initialization of
422 | downscaled runs. Overall, when using an embedded lake model in a downscaling application,
423 | users should consider how the lake model is being initialized or spun-up in order to achieve
424 | results with accuracy similar to the prior studies discussed above. If the lake state is initially
425 | poorly prescribed from the GCM (with results similar to those shown in Fig. 2), a protracted

426 spin-up could be required to reach equilibrium with the driving fields in the RCM and obtain
427 more realistic results.

428
429 It has been noted previously that both WRF-FLake and WRF-CLM, as well as other 1-D lake
430 models, tend to exhibit difficulty in simulating deep lakes (e.g., Martynov et al., 2010;
431 Stepanenko et al., 2010; Gu et al., 2013; M14). Some model error can be attributed to the fact
432 that one-dimensional column models cannot represent 2- and 3-D processes (e.g., currents,
433 drifting ice, and formation of a thermal bar). While more sophisticated lake models could be
434 coupled with WRF, using computationally efficient 1-D models is advantageous in downscaling
435 applications, where computational resources are taxed by the use of finer resolution.
436 Additionally, Martynov et al. (2010) noted that more complex 3-D lake models are generally run
437 with much finer grid spacing (~ 2 km) than typical RCMs. Martynov et al. (2010) also compared
438 the simulated water temperatures and ice coverage from the Hostetler and FLake models, finding
439 that FLake generally performed better, but that the Hostetler model provides more opportunity to
440 improve model performance because it utilizes more vertical layers and is less reliant on
441 parameterization. A comparison of 1-D lake models by Thiery et al. (2014) showed favorable
442 results for both FLake and Hostetler-based models (including the lake model found in CLM4)
443 and noted their computational efficiency. When making regional climate projections with these
444 models it should be noted that both WRF-FLake and WRF-CLM assume that lake depths are
445 constant in time, which could be a poor assumption depending on the lake being modeled and the
446 future period. Also, more complex lake models may be appropriate for higher resolution (~ 2 km
447 grid spacing) RCM simulations focused on regions where lake dynamics are not adequately
448 captured by the column lake models discussed here.

449

450 **3 Conclusion**

451 It has been shown in the present study and in previous work (e.g., Gao et al., 2012; Bullock et al.,
452 2014; M14) that downscaling typically-coarse GCM data, using WRF's default interpolation
453 | methods, to finer resolution WRF grids results in LST discontinuities and spurious ice formation
454 in the Great Lakes (Fig. 2). Although the default interpolation methods in WRF can easily be
455 modified to alter the interpolation scheme or to eliminate the search option, none of these simple
456 changes will overcome the challenges of setting the LSTs for inland water bodies that are not

457 resolved by driving data when WRF is used as a RCM. Various alternate methods have been
458 presented, and a summary of the positives and potential drawbacks to each approach is shown in
459 Table 21. Using WRF's "alternative" lake setting instead of the default interpolation method in
460 WPS eliminates unrealistically large and abrupt spatial discontinuities in temperature, but causes
461 large, deep lakes (such as Lake Superior) to erroneously freeze when ice is set based on an air-
462 temperature threshold. All the other approaches discussed above can simulate more realistic ice
463 cover than the default interpolation. However, the simulation of ice cover is obviously not a
464 factor in downscaling studies where the environment does not become sufficiently cold to
465 produce lake ice, such as those focusing on tropical regions. For example, the alternative lake
466 setting has been used to improve rainfall results (relative to the use of WRF's default
467 interpolation techniques) over Lake Victoria in Eastern Africa by Argent (2014). Using
468 climatological values in a future warmer climate will adversely affect results because LSTs
469 cannot be assumed to be stationary over time. A warming trend could be applied to observed
470 LST fields in order to improve this approach; however, a realistic trend may be complex to
471 derive for some lakes as Austin and Colman (2007) have shown an observed non-linear
472 amplification of warming LSTs relative to inland temperatures in the Great Lakes region. The
473 land mask alteration method of Gao et al. (2012) is effective at preventing discontinuities in
474 surface temperatures, but the use of temperatures from land grid cells in the GCM to set LSTs in
475 the RCM eliminates the presence of land-lake temperature contrasts which impact precipitation,
476 winds (i.e. land-sea breeze), and other near-surface fields. The use of a lake model (either
477 coupling a lake model to the RCM or using outputs from the GCM's lake model to drive the
478 RCM) can improve the representation of the lakes in retrospective simulations and has the ability
479 to simulate non-linear impacts of climate change on LSTs and ice cover (e.g., Gula and Peltier,
480 2012, M14).

481
482 For downscaling applications using WRF, we recommend setting LSTs and ice cover from either
483 a RCM- or GCM-driven lake model, especially when simulating mid-latitude regions. In their
484 studies focused on the Great Lakes, Notaro et al. (2013) and Wright et al. (2013) state that
485 accurate predictions of changes in LSTs and ice cover from lake models are needed when
486 simulating changes in regional climate. Zhao et al. (2012) also recommended the use of a lake
487 model for simulating changes in regional precipitation in the Great Lakes basin. Including

488 prognostic changes in the lake state is also possible if GCM data sets include predicted lake
489 surface temperatures and ice within their publicly-available outputs. For regional climate
490 modeling efforts in which the RCM data is being archived for various end-user applications, we
491 recommend the use of GCM- or RCM-driven lake modeling approaches. If such an approach is
492 not used, the potential adverse effects of setting LSTs and ice cover using interpolation from the
493 GCM should be documented, as is currently done in NARCCAP (2014).

494

495 The accuracy of the various approaches presented here is sensitive to the characteristics of the
496 lakes to which they are being applied. Approaches which set LSTs as a function of over-land
497 temperatures (such as the land mask modification approach or WRF's alternative lake setting)
498 may perform adequately when applied to smaller, shallower lakes where LST changes are more
499 closely coupled to air temperature changes. Investigators performing RCM experiments should
500 consider both the present-day interactions between the lake and overlying air masses as well as
501 the potential climate change impacts on the lakes within their model domain when choosing an
502 approach.

503

504 **4 Code availability**

505 WPS and the WRF model can be downloaded from
506 <http://www2.mmm.ucar.edu/wrf/users/downloads.html>. Source code for the FLake model can be
507 obtained at <http://www.flake.igb-berlin.de/sourcecodes.shtml>, and code needed to run the
508 coupled WRF-FLake model is available for download at
509 <http://web.atmos.ucla.edu/~gula/wrfflake>.

510 **Acknowledgements**

511 The lead author was supported by an appointment to the Research Participation Program at the
512 US Environmental Protection Agency ([EPA](#)) Office of Research and Development, administered
513 by the Oak Ridge Institute for Science and Education (ORISE). The WRF model is made
514 available by NCAR, funded by the National Science Foundation. [Great Lakes ice cover data](#)
515 [were obtained from GLERL \(<http://www.glerl.noaa.gov/data/pgs/glice/glice.html>\)](#). This research
516 has been subjected to the US EPA's administrative review and approved for publication. The

517 views expressed and the contents are solely the responsibility of the authors, and do not
518 necessarily represent the official views of the US EPA.

519 **References**

520 Anyah, R. O. and Semazzi, F. H. M.: Simulation of the sensitivity of Lake Victoria basin climate
521 to lake surface temperatures, *Theor. Appl. Climatol.*, 79, 55–69, 2004.

522

523 Argent, R. E.: Customisation of the WRF model over the Lake Victoria basin in east Africa,
524 M.S. thesis, North Carolina State University, Raleigh, NC, 124 pp., 2014.

525

526 Artale, V., Calmanti, S., Carillo, A., Dell'Aquila, A., Herrmann, M., Pisacane, G., Ruti, P. M.,
527 Sannino, G., Struglia, M. V., Giorgi, F., Bi, X., Pal, J. S., Rauscher, S., and The PROTHEUS
528 Group: An atmosphere–ocean regional climate model for the Mediterranean area: assessment of
529 a present climate simulation, *Clim. Dynam.*, 35, 721–740, 2010.

530

531 Asnani, G. C.: *Tropical Meteorology*, Vol. 1 and 2, Indian Institute of Tropical Meteorology
532 Pashan, Pune, 1012 pp., 1993.

533

534 Assel, R. A. and Robertson, D. M.: Changes in winter air temperatures near Lake Michigan,
535 1851–1993, as determined from regional lake-ice records, *Limnol. Oceanogr.*, 40, 165–176,
536 1995.

537

538 Austin, J. and Colman, S.: Lake Superior summer water temperatures are increasing more
539 rapidly than regional air temperatures: A positive ice–albedo feedback, *Geophys. Res. Lett.*, 34,
540 L06604, doi:10.1029/2006GL029021, 2007.

541

542 Bates, G. T., Giorgi, F., and Hostetler, S. W.: Toward the simulation of the effects of the Great
543 Lakes on regional climate, *Mon. Weather Rev.*, 123, 1505–1522, 1993.

544

545 Bullock, O. R., Alapaty, K., Herwehe, J. A., Mallard, M. S., Otte, T. L., Gilliam, R. C., and
546 Nolte, C. G.: An observation-based investigation of nudging in WRF for downscaling surface
547 climate information to 12 km grid spacing, *J. Appl. Meteorol. Clim.*, 53, 20–33, 2014.

548

549 Burnett, A. W., Kirby, M. E., Mullins, H. T., and Patterson, W. P.: Increasing Great Lake–effect
550 snowfall during the twentieth century: a regional response to global warming?, *J. Climate*, 16,
551 3535–3542, 2003.

552

553 Cairns, M. M., Collins, R., Cylke, T., Deutschendorf, M., and Mercer, D.: A lake effect snowfall
554 in western Nevada. Part I: Synoptic setting and observations, Preprints, 18th Conf. on Weather
555 Analysis and Forecasting, Fort Lauderdale, FL, 29 July 2001, *Am. Meteor. Soc.*, 329–332, 2001.

556

557 Carpenter, D. M.: The lake effect of the Great Salt Lake: overview and forecast problems,
558 *Weather Forecast.*, 8, 181–193, 1993.

559

560 Chen, F. and Dudhia, J.: Coupling and advanced land surface–hydrology model with the Penn
561 State–NCAR MM5 modeling system, Part I: Model implementation and sensitivity, *Mon.*
562 *Weather Rev.*, 129, 569–585, 2001.

563

564 Feser, F., Rockel, B., von Storch, H., Winterfeldt, J., and Zahn, M.: Regional climate models add
565 value to global model data: a review and selected examples. *B. Am. Meteorol. Soc.*, 92, 1181–
566 1192, 2011.

567

568 Gao, Y., Fu, J. S., Drake, J. B., Liu, Y., and Lamarque, J.-F.: Projected changes of extreme
569 weather events in the eastern United States based on a high resolution climate modeling system,
570 *Environ. Res. Lett.*, 7, 044025, doi:10.1088/1748-9326/7/4/044025, 2012.

571

572 Gerbush, M. R., Kristovich, D. A. R., and Laird, N. F.: Mesoscale boundary layer and heat flux
573 variations over pack ice–covered Lake Erie, *J. Appl. Meteorol. Clim.*, 47, 668–682, 2008.

574

575 Gu, H., Jin, J., Wu, Y., Ek, M., and Subin, Z.: Calibration and validation of lake surface
576 temperature simulations with the coupled WRF-lake model, *Clim. Chang.*, doi: 10.1007/s10584-
577 013-0978-y, in press, 2013.

578

579 Gualdi, S., Somot, S., Li, L., Artale, V., Adani, M., Bellucci, A., Braun, A., Calmanti, S., Carillo,
580 A., Dell'Aquila, A., Déqué, M., Dubois, C., Elizalde, A., Harzallah, A., Jacob, D., L'Hévéder,
581 B., May, W., Oddo, P., Ruti, P., Sanna, A., Sannino, G., Scoccimarro, E., Sevault, F., and
582 Navarra, A.: The CIRCE simulations: Regional climate change projections with realistic
583 representation of the Mediterranean Sea, *B. Am. Meteorol. Soc.*, 94, 65–81, 2013.

584

585 Gula, J. and Peltier, W. R.: Dynamical downscaling over the Great Lakes basin of North
586 America using the WRF regional climate model: the impact of the Great Lakes system on
587 regional greenhouse warming, *J. Climate*, 25, 7723–7742, 2012.

588

589 Hostetler, S. W. and Bartlein, P. J.: Simulation of lake evaporation with application to modeling
590 lake level variations of Harney-Malheur Lake, Oregon, *Water Resour. Res.*, 26, 2603–2612,
591 1990.

592

593 Hostetler, S. W., Bates, G. T., and Giorgi, F.: Interactive coupling of a lake thermal model with a
594 regional climate model, *J. Geophys. Res.*, 98, 5045–5057, 1993.

595

596 Hostetler, S. W., Giorgi, F., Bates, G. T., and Bartlein, P. J.: Lake-atmosphere feedbacks
597 associated with paleolakes Bonneville and Lahontan, *Science*, 263, 665–668, 1994.

598

599 Huggins, A. W., Kingsmill, D. E., and Cairns, M. M.: A lake effect snowfall in western Nevada–
600 Part II: Radar characteristics and quantitative precipitation estimates, Preprints, 18th Conf. on
601 Weather Analysis and Forecasting, Fort Lauderdale, FL, 29 July 2001, *Amer. Meteor. Soc.*, 333
602 337, 2001.

603

604 Kanamitsu, M., Ebisuzaki, W., Woollen, J., Yang, S.-K., Hnilo, J. J., Fiorino, M., and Potter, G.
605 L.: NCEP–DOE AMIP-II Reanalysis (R-2). *B. Am. Meteorol. Soc.*, 83, 1631–1643, 2002.

606

607 Kourzeneva, E., Samuelsson, P., Ganbat, G., and Mironov, D.: Implementation of Lake Model
608 FLake into HIRLAM, *HIRLAM Newsletter* 54, 54–64, available from HIRLAM-A Programme,

609 c/o J. Onvlee, KNMI, P. O. Box 201, 3730 AE De Bilt, the Netherlands, available at: <http://hirLAM.org> (last access: 20 October 2014), 2008.

610

611

612 Kunkel, K. E., Ensor, L., Palecki, M., Easterling, D., Robinson, D., Hubbard, K. G., and
613 Redmond, K.: A new look at lake-effect snowfall trends in the Laurentian Great Lakes using a
614 temporally homogeneous data set, *J. Great Lakes Res.*, 35, 23–29, 2009.

615

616 Laird, N. F., Desrochers, J., and Payer, M.: Climatology of lake-effect precipitation events over
617 Lake Champlain, *J. Appl. Meteorol. Clim.*, 48, 232–250, 2009.

618

619 Magnuson, J. J., Robertson, D. M., Benson, B. J., Wynne, R. H., Livingstone, D. M., Arai, T.,
620 Assel, R. A., Barry, R. G., Card, V., Kuusisto, E., Granin, N. G., Prowse, T. D., Stewart, K. M.,
621 and Vuglinski, V. S.: Historical trends in lake and river ice cover in the Northern Hemisphere,
622 *Science*, 289, 1743–1746, 2000.

623

624 | Mallard, M. S., Nolte, C. G., Bullock, O. R., [SperoOtte, T. L.](#), and Gula, J.: Using a coupled lake
625 model with WRF for dynamical downscaling, *J. Geophys. Res.*, 119, 7193–7208,
626 doi:10.1002/2014JD021785, 2014.

627

628 Martynov, A., Laprise, R., and Sushama, L.: Off-Line lake water and ice simulations: a step
629 towards the interactive lake coupling with the Canadian Regional Climate Model, *Geophysical*
630 *Research Abstracts*, Vol. 10, EGU2008-A-02898, EGU General Assembly 2008, Vienna,
631 Austria, 2008.

632

633 Martynov, A., Sushama, L., and Laprise, R.: Simulation of temperate freezing lakes by one-
634 dimensional lake models: performance assessment for interactive coupling with regional climate
635 models, *Boreal Environ. Res.*, 15, 143–164, 2010.

636

637 Mearns, L. O., Arritt, R., Biner, S., Bukovsky, M. S., McGinnis, S., Sain, S., Caya, D., Correia
638 Jr., J., Flory, D., Gutowski, W., Takle, E. S., Jones, R., Leung, R., Moufouma-Okia, W.,
639 McDaniel, L., Nunes, A. M. B., Qian, Y., Roads, J., Sloan, L., and Snyder, M.: The North

640 American regional climate change assessment program: overview of phase I results, B. Am.
641 Meteorol. Soc., 93, 1337–1362, 2012.

642

643 Mironov, D. V.: Parameterization of lakes in numerical weather prediction. Description of a lake
644 model, COSMO Technical Report, No. 11, Deutscher Wetterdienst, Offenbach am Main,
645 Germany, 41 pp., 2008.

646

647 Mironov, D., Heise, E., Kourzeneva, E., Ritter, B., Schneider, N., and Terzhevik, A.:
648 Implementation of the lake parameterization scheme FLake into the numerical weather
649 prediction model COSMO, *Boreal Env. Res.*, 15, 218–230, 2010.

650

651 NARCCAP: Caveats for Users, available at: <http://www.narccap.ucar.edu/about/caveats.html>,
652 last access: 11 August 2014.

653

654 Notaro, M., Holman, K., Zarrin, A., Fluck, E., Vavrus, S., and Bennington, V.: Influence of the
655 Laurentian Great Lakes on regional climate, *J. Climate*, 26, 789–804, 2013.

656

657 Oleson, K. W., Lawrence, D. M., Bonan, G. B., Flanner, M. G., Kluzek, E., Lawrence, P. J.,
658 Levis, S., Swenson, S. C., Thornton, P. E., Dai, A., Decker, M., Dickinson, R., Feddema, J.,
659 Heald, C. L., Hoffman, F., Lamarque, J.-F., Mahowald, N., Niu, G.-Y., Qian, T., Randerson, J.,
660 Running, S., Sakaguchi, K., Slater, A., Stöckli, R., Wang, A., Yang, Z.-L., Zeng, X., and Zeng,
661 X.: Technical description of version 4.0 of the Community Land Model (CLM), National Center
662 for Atmospheric Research, P.O. Box 3000, Boulder, Colorado 80307, USA, NCAR/TN-
663 478+STR, 2010.

664

665 Oleson, K. W., Lawrence, D. M., Bonan, G. B., Drewniak, B., Huang, M., Koven, C. D., Levis,
666 S., Li, F., Riley, W. J., Subin, Z. M., Swenson, S. C., Thornton, P. E., Bozbiyik, A., Fisher, R.,
667 Heald, C. L., Kluzek, E., Lamarque, J.-F., Lawrence, P. J., Leung, L. R., Lipscomb, W.,
668 Muszala, S., Ricciuto, D. M., Sacks, W., Sun, Y., Tang, J., and Yang, Z.-L.: Technical
669 description of version 4.5 of the Community Land Model (CLM), National Center for

670 Atmospheric Research, P.O. Box 3000, Boulder, Colorado 80307, USA, NCAR Technical Note
671 NCAR/TN-503+STR, 420 pp., doi:10.5065/D6RR1W7M, 2013.
672

673 Ratnam, J. V., Filippo, G., Kaginalkar, A., and Cozzini, S.: Simulation of the Indian monsoon
674 using the RegCM3-ROMS regional coupled model, *Clim. Dynam.*, 33, 119–139, 2009. Rinke,
675 A., Gerdes, R., Dethloff, K., Kandlbinder, T., Karcher, M., Kauker, F., Frickenhaus, S., Köberle,
676 C., and Hiller, W.: A case study of the anomalous Arctic sea ice conditions during 1990: insights
677 from coupled and uncoupled regional climate model simulations, *J. Geophys. Res.*, 108, 4275,
678 doi:10.1029/2002JD003146, 2003.
679

680 Samuelsson, P., Kourzeneva, E., and Mironov, D.: The impact of lakes on the European climate
681 as simulated by a regional climate model, *Boreal Env. Res.*, 15, 113–129, 2010.
682

683 Schultz, D. M., Arndt, D. S., Stensrud, D. J., and Hanna, J. W.: Snowbands during the cold-air
684 outbreak of 23 January 2003, *Mon. Weather Rev.*, 132, 827–842, 2004.
685

686 Scott, R. W. and Huff, F. A.: Impacts of the Great Lakes on regional climate conditions, *J. Great
687 Lakes Res.*, 22, 845–863, 1996.
688

689 Skamarock, W. C. and Klemp, J. B.: A time-split nonhydrostatic atmospheric model for weather
690 research and forecasting applications, *J. Comp. Phys.*, 227, 3465–3485, 2008.
691

692 Skamarock, W. C., Klemp, J. B., Dudhia, J., Gill, D. O., Barker, D. M., Duda, M., Huang, X.-Y.,
693 Wang, W., and Powers, J. G.: A description of the Advanced Research WRF version 3, Tech.
694 Rep., National Center for Atmospheric Research, P.O. Box 3000, Boulder, Colorado 80307,
695 USA, NCAR/TN-475+STR, 113 pp., 2008.
696

697 Steenburgh, W. J. and Onton, D. J.: Multiscale analysis of the 7 December 1998 Great Salt
698 Lake–effect snowstorm, *Mon. Weather Rev.*, 129, 1296–1317, 2001.
699

700 Stepanenko, V. M., Goyett, S., Martynov, A., Perroud, M., Fang, X., and Mironov, D.: First
701 steps of the Lake Model Intercomparison Project: LakeMIP, *Boreal Env. Res.*, 15, 191–202,
702 2010.

703

704 Subin, Z. M., Riley, W. J., and Mironov, D.: An improved lake model for climate simulations:
705 Model structure, evaluation, and sensitivity analyses in CESM1, *J. Adv. Model. Earth Syst.*, 4,
706 M02001, doi:10.1029/2011MS000072, 2012.

707

708 Tardy, A.: Lake-effect and lake-enhanced snow in the Champlain Valley of Vermont, Tech.
709 Memo. 2000-05, NWS Eastern Region, 27 pp., Burlington, Vermont, USA, 2000.

710

711 Thiery, W., Stepanenko, V. M., Fang, X., Johnk, K. D., Li, Z., Martynov, A., Perroud, M., Subin,
712 Z. M., Darchambeau, F., Mironov, D. and van Lipzig, N. P. M.: LakeMIP Kivu: evaluating the
713 representation of a large, deep tropical lake by a set of one-dimensional lake models, *Tellus*, 66,
714 21390, 2014.

715

716 Wang, J., Bai, X., Hu, H., Clites, A., Colton, M., and Lofgren, B.: Temporal and spatial
717 variability of Great Lakes ice cover, 1973–2010, *J. Climate*, 25, 1318–1329, 2012a.

718

719 Wang, J., Assel, R. A., Walterscheid, S., Clites, A. H., and Bai, X.: Great Lakes ice climatology
720 update: winter 2006–2011 description of the digital ice cover data set, NOAA Technical
721 Memorandum GLERL-155, 37 pp., Ann Arbor, Michigan, USA, 2012b.

722

723 Wilken, G. R.: A lake-effect snow in Arkansas, NWS Southern Region, Tech. Attachment
724 SR/SSD 97-21, 5 pp., Little Rock, Arkansas, USA, 1997.

725

726 Wilson, J. W.: Effect of Lake Ontario on precipitation, *Mon. Weather Rev.*, 105, 207–214, 1977.

727

728 WRF User’s Guide: User’s Guide for the Advanced Research WRF (ARW) Modeling System
729 Version 3.6, available at:

730 http://www2.mmm.ucar.edu/wrf/users/docs/user_guide_V3/ARWUsersGuideV3.pdf, last access:
731 12 June 2014.
732
733 Wright, D. M., Posselt, D. J., and Steiner, A. L.: Sensitivity of lake-effect snowfall to lake ice
734 cover and temperature in the Great Lakes region, *Mon. Weather Rev.*, 141, 670–689, 2013.
735
736 Zhao, L., Jin, J., Wang, S.-Y., and Ek, M. B.: Integration of remote-sensing data with WRF to
737 improve lake-effect precipitation simulations over the Great Lakes region, *J. Geophys. Res.*, 117,
738 D09102, doi:10.1029/2011JD016979, 2012.
739
740 Zulauf, M. A. and Kreuger, S. K.: Two-dimensional cloud-resolving modeling of the
741 atmospheric effects of Arctic leads based upon midwinter conditions at the surface heat budget
742 of the Arctic Ocean ice camp, *J. Geophys. Res.*, 108, 4312, doi:10.1029/2002JD002643, 2003.
743

744 Table 1. List of WRF versions discussed in the text, ordered chronologically by the date of
 745 release and with relevant model updates summarized.


<u>WRF version</u>	<u>Released</u>	<u>Updates of Interest</u>
<u>3.3</u>	<u>April 2011</u>	<u>“Alternative initialization of lake SSTs” option included in WPS so users can set LSTs from temporally averaged 2 m temperatures.</u>
<u>3.5</u>	<u>April 2013</u>	<u>CLM available as an LSM within WRF, but with its lake model disabled.</u>
<u>3.5.1</u>	<u>September 2013</u>	<u>Default surface water temperature at which WRF prescribes ice (“seaice_threshold”) is lowered from 271 K to 100 K.</u>
<u>3.6</u>	<u>April 2014</u>	<u>CLM lake model available with any choice of LSM. Lake depths can be prescribed as a constant or as a spatially varying 2-D field.</u>

746

747 | Table 21. A summary of the pros and cons of each method of treating lake surface temperatures
 748 | and ice coverage described in the text. All approaches were found to eliminate unrealistic
 749 | temperature discontinuities resulting from WRF's default interpolation methods as shown in Fig.
 750 | 2.

Methodology	Positives	Potential drawbacks
WRF's Alternative Lake Setting	Effective at representing LSTs when lake temperatures are closely coupled with atmospheric temperatures.	Unrealistic ice formation possible when 2 m temperatures are below freezing. Cannot account for varying lake depths and differing timescales of warming and cooling throughout lakes.
Climatological	Observed LSTs and ice taken from high-resolution analyses.	For long-term simulations, user must include temperature trend or LSTs will not be in equilibrium with future climate state. Does not represent interannual variability of lake state.
Land Mask Modification	Future LSTs can be taken from projected GCM temperatures.	Eliminates land-lake temperature contrasts.
Lake Model Component	Models have ability to simulate future changes in LST and ice.	Additional preprocessing needed to provide lake model spin-up for RCM run or to use lake fields simulated by GCM.


751



752

753 Figure 1. The ocean mask from the 1⁰ CESM data (which is used by WPS to determine the
754 locations of land and water points from CESM), as shown in the area corresponding to a WRF
755 36-km continental US domain (left), and the 36 km WRF grid's land-water mask (right). Labels
756 are placed to indicate the locations of Lakes Superior ("S"), Michigan ("M"), Huron ("H"), Erie
757 ("E") and Ontario ("O"), as well as Hudson Bay ("HB").

Skin Temperature [K] 1994-12-01




758


759 Figure 2. The skin temperature (K) processed from CESM to the 36-km WRF grid using WPS

760 and valid at 00 UTC 1 Dec 1994. White circles indicate the locations of Pyramid Lake, Great

761 Salt Lake, and Lake Sakakawea, from west to east, respectively.



762
 763 Figure 3. Surface temperature from the initial year of a 10-year FLake spin-up simulation, taken
 764 from a point near the north shore of Lake Superior (48.47° N, 87.54° W) and shown hourly from
 765 1 January to 31 December 2005 (top). LSTs at all lake cells are initialized with a default value
 766 of 274.15 K, and the time series shows either ice or water surface temperatures depending on
 767 whether ice is present. Simulated ice thickness (m) taken from day 30 of the same FLake
 768 simulation, valid 30 January 2005 (bottom left). Fractional ice values observed on this date
 769 plotted from the NIC ice analysis (bottom right).



770
 771 Figure 4. Simulated ice cover (%) taken from a WRF simulation (valid 2 March 2006, after ~ 4
 772 months of simulation time) with the same model configuration as described in M14, but
 773 simulated with WRF version 3.6 and the use of the CLM lake model in place of FLake (left). A
 774 2-D field of lake depths (instead of a single default value) were used from WPS to set the lake
 775 depth in this simulation. Ice coverage observed on this date is plotted from the NIC ice analysis
 776 (right).