

Interactive comment on “The terminator “toy”-chemistry test: a simple tool to assess errors in transport schemes” by P. H. Lauritzen et al.

P. H. Lauritzen et al.

pel@ucar.edu

Received and published: 3 April 2015

Below the reviewer comments are in black font and author reply in blue font:

This paper presents a simplified chemistry-like interaction between two correlated tracers to test transport schemes. The authors particularly focused on the correlation properties of these tracers and the effect of monotonicity/limiters. Overall the paper is clear and well written and the test could be useful to chemistry transport development. Therefore, I don't have any objection for its publication in GMD.

[We thank the reviewer for reviewing the manuscript and providing very useful comments. The revised manuscript is available in the supplement](#)

C3590

Specific comments:

The mathematical equations are not easy on the eye and a bit confusing whether the l in Cl is L or l . Also I don't see why one wants to use 2 or 3 letters to define a variable while one letter will suffice. The notations would be much neater and clearer if, for example, m_1 , m_2 and m_3 are used for Cl , Cl_2 and Cl_y , respectively.

[Response: Agreed](#)

[Changes to manuscript: Notation has been been changed to \$X\$, \$X_2\$, and \$X_T\$ throughout the manuscript.](#)

Section 2. The physical meanings (i.e., mixing ratios) of the three tracers should be clearly defined in this section. The section shows lot of equations without ever saying what are these Cl s.

[Response: Agreed](#)

[Changes to manuscript: In the beginning of Section 2 the units of \$X\$ has been clearly defined, i.e. nomenclature \$X\$ describes the number density \(number of molecules of compound \$X\$ divided by the number of molecules of air\).](#)

Throughout the text the use of “preserve and conserve” are left somehow ambiguous and confusing as to what exactly we mean (examples: page 8772, “Another inspiration for this test is the atomic concentration is conserved.”; “... conserved the total chlorine”). It would help the reader to define exactly what we mean by these words (i.e., tracer mass conservation, preserving a constant, preserving a relationship, etc...), especially in presence of the sources terms.

[Response: Agreed](#)

[Changes to manuscript: This has been clarified. When talking about preservation or conservation the quantity in question is explicitly stated in the manuscript.](#)

Section 5.3 and Fig. 5. The authors has to explain why the “default limiter” doesn't

C3591

remove the overshoot in CI (bottom-left corner picture). Fig. 7. The authors has to provide an explanation as to why the error in CAM-FV is around 10 times smaller than that of CAM-SE?

Response: We stated in the manuscript that an in-depth analysis of why the limiters do not preserve linear relations is up to the scheme developers. That said, we have performed some additional experiments to shed light on why CAM-SE performs as it does. Please see text below.

Changes to the manuscript: The following paragraphs have been edited/added to section 5.2:

CAM-SE does not preserve linear relations either and the errors in X_T are about an order of magnitude larger than CAM-FV. The CAM-SE limiter is optimization-based (using Least-squares) and guarantees no under- or over-shoots at the element level while maintaining mass-conservation at the element level (Guba et al., 2014). While the optimization-based limiter preserves linear relations with exact arithmetic, its present implementation in CAM-SE does not lead to such preservation (most likely due to iteration thresholds, if-statements, etc. that can be non-linear).

To further understand this behavior we have performed some tests (not shown) turning the chemistry off and advecting linearly correlated Cosine hills and linearly correlated step-functions. The Cosine hills are C^0 continuous (the function is continuous but its derivatives are not) and the limiter exactly preserves the linear relationship. For the step-functions, which are discontinuous distributions, the correlation preservation is only maintained up to $O(10^{-8})$ due to an $O(10^{-8})$ overshoot in one of the tracers. So the advection operator introduces an $O(10^{-8})$ error. The terminator chemistry constantly enforces a discontinuity in the distributions and in combination with the CAM-SE limiter, strong error growth is produced. It is beyond the scope of this paper to trace down exactly where in the implementation this error is introduced and to find a remedy. The terminator test is designed to enable scheme developers to test their scheme in

C3592

setups that are directly relevant to some of the issues seen in chemistry application. In this particular case, the terminator test clearly exacerbates small errors in correlation preservation that may be easily overlooked in inert transport testing (such as the tests in Lauritzen et al., 2012).

Final thought: Let us denote the test's equations as follows: (a) $dm1/dt = S1(m1, m2)$, (b) $dm2/dt = S2(m1, m2)$, (c) $dm3/dt = 0$, and (d) $m3 = f(m1, m2)$. I would like to see a discussion about the over-specification of this problem (i.e., there is an arbitrary choice of how one could solve these over-specified equations set). If (d) is an important property to maintain, then one could simply solve (a) + (c) and deduce $m2$ from (d), instead of the approach used in the paper [i.e., (a) + (b) and check how well we achieve (d)]. I am not asking to redo the experiments with the alternative approach but a brief discussion as to why this way and not the other way?

Response: Agreed!

Changes to the manuscript: The following paragraph has been added to the end of Section 3:

We note that instead of advecting each species separately by solving the advection equations in eq:adveqs or eq:rho-phi with $\phi = X, X_2$, one may also choose to advect the sum X_T and one of the species, e.g. X , and then diagnose the remaining species, X_2 . If the transport operator conserves a constant, which many transport operators do, then the constant sum X_T is trivially conserved if chosen as a prognostic variable. This approach is commonly used in chemistry transport models for some families of species. E.g., Douglass *et al.* (e.g., 2004) advected the sum of total inorganic chlorine and bromine to avoid spurious maxima and minima in their distributions in the stratosphere. Idealized tests to evaluate how well families of species that add up to a constant are preserved see, e.g., Lauritzen and Thuburn (2012).

Minor comments:

C3593

Section 5.1, page 8781: There is a bit of mis-use of the word mass. Most the word “mass” in the 2nd paragraph of section 5.1 refers to “density”. We should be clear that $\rho\phi$ is the tracer density and not mass [if ρ is the density of dry air and ϕ is a tracer mixing ratio (mass = $\rho\phi V$ where V is the volume element)].

Response: Text changed to avoid confusion (refer to $\rho\phi$ as tracer density).

References

- Anderson, J., Toohey, D., and Brune, W.: Free radicals within the Antarctic vortex: the role of CFCs in Antarctic ozone loss, *Science*, 251, 39–46, doi:10.1126/science.251.4989.39, 1991.
- Andrews, D. G., Holton, J. R., and Leovy, C. B.: *Middle atmosphere dynamics*, Academic Press, 1987.
- Barth, T. and Jespersen, D.: The design and application of upwind schemes on unstructured meshes, in: *Proc. AIAA 27th Aerospace Sciences Meeting*, Reno, 1989.
- Blossey, P. N. and Durran, D. R.: Selective monotonicity preservation in scalar advection, *J. Comput. Phys.*, 227, 5160–5183, 2008.
- Brasseur, G. and Solomon, S.: *Aeronomy of the Middle Atmosphere*, Springer, 3rd Edn., 2005.
- Colella, P. and Woodward, P. R.: The Piecewise Parabolic Method (PPM) for gas-dynamical simulations, *J. Comput. Phys.*, 54, 174–201, 1984.
- Dennis, J. M., Edwards, J., Evans, K. J., Guba, O., Lauritzen, P. H., Mirin, A. A., St-Cyr, A., Taylor, M. A., and Worley, P. H.: CAM-SE: A scalable spectral element dynamical core for the Community Atmosphere Model, *Int. J. High Perform. C.*, 26, 74–89, doi:10.1177/1094342011428142, 2012.
- Douglas AR, Stolarski RS, Strahan SE, Connell PS. 2004. Radicals and reservoirs in the GMI chemistry and transport model: Comparison to measurements, *J. Geophys. Res.*, **109**, D16302, DOI:10.1029/2004JD004632
- Durran, D.: *Numerical Methods for Fluid Dynamics: With Applications to Geophysics*, vol. 32 of *Texts in Applied Mathematics*, Springer, 2nd Edn., 516 pp., 2010.
- Edouard, S., Legras, B., Zeitlin, V.: The effect of dynamical mixing in a simple model of the ozone hole. *J. Geophys. Res.*, 101(D11), 2156–2202, doi:, 1996.

C3594

- Gravel, S. and A. Staniforth: A mass-conserving semi-Lagrangian scheme for the shallow-water equations. *Mon. Wea. Rev.*, **122**, 243–248, 1994.
- Guba, O., Taylor, M., and St-Cyr, A.: Optimization-based limiters for the spectral element method, *J. Comput. Phys.*, 267, 176–195, doi:10.1016/j.jcp.2014.02.029, 2014.
- Harris, L. M., Lauritzen, P. H., and Mittal, R.: A flux-form version of the Conservative Semi-Lagrangian Multi-tracer transport scheme (CSLAM) on the cubed sphere grid, *J. Comput. Phys.*, 230, 1215–1237, doi:10.1016/j.jcp.2010.11.001, 2010.
- Kinnmark, I. P. and Gray, W. G.: One step integration methods with maximum stability regions, *Math. Comput. Simulat.*, 26, 87–92, doi:10.1016/0378-4754(84)90039-9, 1984a.
- Kinnmark, I. P. and Gray, W. G.: One step integration methods of third-fourth order accuracy with large hyperbolic stability limits, *Math. Comput. Simulat.*, 26, 181–188, doi:10.1016/0378-4754(84)90056-9, 1984b.
- Kristensen, L., Lenschow, D. H., Gurarie, D., and Jensen, N. O.: A simple model for the vertical transport of reactive species in the convective atmospheric boundary layer, *Bound.-Lay. Meteorol.*, 134, 195–221, doi:10.1007/s10546-009-9443-x, 2010.
- Lamarque, J.-F., Emmons, L. K., Hess, P. G., Kinnison, D. E., Tilmes, S., Vitt, F., Heald, C. L., Holland, E. A., Lauritzen, P. H., Neu, J., Orlando, J. J., Rasch, P. J., and Tyndall, G. K.: CAM-chem: description and evaluation of interactive atmospheric chemistry in the Community Earth System Model, *Geosci. Model Dev.*, 5, 369–411, doi:10.5194/gmd-5-369-2012, 2012.
- Lander, J. and Hoskins, B. J.: Believable scales and parameterizations in a spectral transform model. *Mon. Wea. Rev.*, 125, 292–303. doi: , 1997.
- Lauritzen, P. and Thuburn, J.: Evaluating advection/transport schemes using interrelated tracers, scatter plots and numerical mixing diagnostics, *Q. J. Roy. Meteor. Soc.*, 138, 906–918, doi:10.1002/qj.986, 2012.
- Lauritzen, P. H., Nair, R. D., and Ullrich, P. A.: A conservative semi-Lagrangian multi-tracer transport scheme (CSLAM) on the cubed-sphere grid, *J. Comput. Phys.*, 229, 1401–1424, doi:10.1016/j.jcp.2009.10.036, 2010.
- Lauritzen, P. H., Ullrich, P. A., and Nair, R. D.: Atmospheric transport schemes: desirable properties and a semi-Lagrangian view on finite-volume discretizations, in: *Numerical Techniques for Global Atmospheric Models*, Lecture Notes in Computational Science and Engineering, edited by: Lauritzen, P. H., Nair, R. D., Jablonowski, C., Taylor, M., Springer, 80 pp., 2011.
- Lauritzen, P. H., Skamarock, W. C., Prather, M. J., and Taylor, M. A.: A standard test case suite for two-dimensional linear transport on the sphere, *Geosci. Model Dev.*, 5, 887–901,

C3595

doi:10.5194/gmd-5-887-2012, 2012.

- Lauritzen, P. H., Ullrich, P. A., Jablonowski, C., Bosler, P. A., Calhoun, D., Conley, A. J., Enomoto, T., Dong, L., Dubey, S., Guba, O., Hansen, A. B., Kaas, E., Kent, J., Lamarque, J.-F., Prather, M. J., Reinert, D., Shashkin, V. V., Skamarock, W. C., Sørensen, B., Taylor, M. A., and Tolstykh, M. A.: A standard test case suite for two-dimensional linear transport on the sphere: results from a collection of state-of-the-art schemes, *Geosci. Model Dev.*, 7, 105–145, doi:10.5194/gmd-7-105-2014, 2014.
- Lin, S.-J.: A “vertically Lagrangian” finite-volume dynamical core for global models, *Mon. Weather Rev.*, 132, 2293–2307, 2004.
- Lin, S.-J. and Rood, R. B.: Multidimensional flux-form semi-Lagrangian transport schemes, *Mon. Weather Rev.*, 124, 2046–2070, 1996.
- Lin, S.-J., Chao, W. C., Sud, Y. C., and Walker, G. K.: A class of the van Leer-type transport schemes and its application to the moisture transport in a general circulation model, *Mon. Weather Rev.*, 122, 1575–1593, 1994.
- Nair, R. D. and Jablonowski, C.: Moving vortices on the sphere: a test case for horizontal advection problems, *Mon. Weather Rev.*, 136, 699–711, 2008.
- Nair, R. D. and Lauritzen, P. H.: A class of deformational flow test cases for linear transport problems on the sphere, *J. Comput. Phys.*, 229, 8868–8887, doi:10.1016/j.jcp.2010.08.014, 2010.
- Nair, R. D. and Mächenhauer, B.: The mass-conservative cell-integrated semi-Lagrangian advection scheme on the sphere, *Mon. Weather Rev.*, 130, 649–667, 2002.
- Neale, R. B., Chen, C.-C., Gettelman, A., Lauritzen, P. H., Park, S., Williamson, D. L., Conley, A. J., Garcia, R., Kinnison, D., Lamarque, J.-F., Marsh, D., Mills, M., Smith, A. K., Tilmes, S., Vitt, F., Cameron-Smith, P., Collins, W. D., Iacono, M. J., Easter, R. C., Ghan, S. J., Liu, X., Rasch, P. J., and Taylor, M. A.: Description of the NCAR Community Atmosphere Model (CAM 5.0), NCAR Technical Note, National Center of Atmospheric Research, 2010.
- Prather, M., Jaffe, A. H.: Global impact of the Antarctic ozone hole: Chemical propagation. *J. Geophys. Res.*, 95(D4), 3473. doi:, 1990
- Priestley, A.: A quasi-conservative version of the semi-Lagrangian advection scheme. *Mon. Wea. Rev.*, 121, 621–632, doi:, 1993.
- Prigogine, I.: *From Being to Becoming: Time and Complexity in the Physical Sciences*, W H Freeman & Co, 1981.
- Prigogine, I. and Lefever, R.: Symmetry-breaking instabilities in dissipative systems, *J. Chem.*

C3596

Phys., 48, 1695–1700, doi:10.1063/1.1668896, 1968.

- Pudykiewicz, J. A.: Numerical solution of the reaction-advection–diffusion equation on the sphere, *J. Comput. Phys.*, 213, 358–390, doi:10.1016/j.jcp.2005.08.021, 2006.
- Pudykiewicz, J. A.: On numerical solution of the shallow water equations with chemical reactions on icosahedral geodesic grid, *J. Comput. Phys.*, 230, 1956–1991, doi:10.1016/j.jcp.2010.11.045, 2011.
- Salawitch, R. J., Canty, T., Müller, R., Santee, M. L., Schofield, R., Stimpfle, R. M., Stroh, F., Toohey, D. W., and Urban, J.: Sect. 3. Workshop Report: The Role of Halogen Chemistry in Polar Stratospheric Ozone Depletion, in: *Workshop for an Initiative under the Stratospheric Processes and Their Role in climate (SPARC) Project of the World climate Research Programme*, June 2008, Cambridge, UK, 2009.
- Skamarock, W. C. and Weisman, M. L.: The impact of positive-definite moisture transport on NWP precipitation forecasts, *Mon. Weather Rev.*, 137, 488–494, doi:10.1175/2008MWR2583.1, 2009.
- Spiteri, R. and Ruuth, S.: A new class of optimal high-order strong-stability-preserving time discretization methods, *SIAM J. Numer. Anal.*, 40, 469–491, doi:10.1137/S0036142901389025, 2002.
- Strahan, S. E., Douglass, A. R., Stolarski, R. S., Akiyoshi, H., Bekki, S., Braesicke, P., Butchart, N., Chipperfield, M. P., Cugnet, D., Dhomse, S., Frith, S. M., Gettelman, A., Hardiman, S. C., Kinnison, D. E., Lamarque, J.-F., Mancini, E., Marchand, M., Michou, M., Morgenstern, O., Nakamura, T., Olivi, D., Pawson, S., Pitari, G., Plummer, D. A., Pyle, J. A., Scinocca, J. F., Shepherd, T. G., Shibata, K., Smale, D., Teyssède, H., Tian, W., Yamashita, Y.: Using transport diagnostics to understand chemistry climate model ozone simulations, *J. Geophys. Res.*, 116(D17), 2156–2202, doi:, 2011.
- Thuburn, J. and McIntyre, M.: Numerical advection schemes, cross-isentropic random walks, and correlations between chemical species, *J. Geophys. Res.*, 102, 6775–6797, 1997.
- Turing, A.: The chemical basis of morphogenesis, *Philos. T. R. Soc. Lon. B*, 37–72, 1952.
- White, J. B. and Dongarra, J. J.: High-Performance High-Resolution Semi-Lagrangian Tracer Transport on a Sphere, *J. Comput. Phys.* 230, 6778–6799, 2011.
- Williamson, D. L., Drake, J. B., Hack, J. J., Jakob, R., and Swarztrauber, P. N.: A standard test set for numerical approximations to the shallow water equations in spherical geometry, *J. Comput. Phys.*, 102, 211–224, 1992.
- Xiao, F., Yabe, T., Peng, X., and Kobayashi, H.: Conservative and oscillation-less atmo-

C3597

spheric transport schemes based on rational functions, *J. Geophys. Res.*, 107, 4609, 10.1029/2001JD001532, 2002.

Zalesak, S. T.: Fully multidimensional flux-corrected transport algorithms for fluids, *J. Comput. Phys.*, 31, 335–362, 1979.

Please also note the supplement to this comment:

<http://www.geosci-model-dev-discuss.net/7/C3590/2015/gmdd-7-C3590-2015-supplement.pdf>

Interactive comment on *Geosci. Model Dev. Discuss.*, 7, 8769, 2014.

C3598