



This discussion paper is/has been under review for the journal Geoscientific Model Development (GMD). Please refer to the corresponding final paper in GMD if available.

A high-resolution ocean and sea-ice modelling system for the Arctic and North Atlantic Oceans

F. Dupont¹, S. Higginson³, R. Bourdallé-Badie⁴, Y. Lu³, F. Roy², G. C. Smith², J.-F. Lemieux², G. Garric⁴, and F. Davidson⁵

¹MSC, Environment Canada, Dorval, QC, Canada

²MRD, Environment Canada, Dorval, QC, Canada

³Bedford Institute of Oceanography, Fisheries & Oceans Canada, Dartmouth, NS, Canada

⁴Mercator-Océan, Toulouse, France

⁵Northwest Atlantic Fisheries Centre, Fisheries and Oceans Canada, St. John's, NF, Canada

Received: 18 November 2014 – Accepted: 5 December 2014 – Published: 5 January 2015

Correspondence to: F. Dupont (frederic.dupont@ec.gc.ca)

Published by Copernicus Publications on behalf of the European Geosciences Union.

GMDD

8, 1–52, 2015

**Arctic and N-Atlantic
high-resolution
ocean and sea-ice
model**

F. Dupont et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Abstract

As part of the CONCEPTS (Canadian Operational Network of Coupled Environmental Prediction Systems) initiative, The Government of Canada is developing a high resolution ($1/12^\circ$) ice–ocean regional model covering the North Atlantic and the Arctic oceans. The objective is to provide Canada with short-term ice–ocean predictions and hazard warnings in ice infested regions. To evaluate the modelling component (as opposed to the analysis – or data-assimilation – component), a series of hindcasts for the period 2003–2009 is carried out, forced at the surface by the Canadian Global Re-
Forecasts. These hindcasts test how the model represent upper ocean characteristics and ice cover. Each hindcast implements a new aspect of the modelling or the ice–ocean coupling. Notably, the coupling to the multi-category ice model CICE is tested. The hindcast solutions are then assessed using a validation package under development, including in-situ and satellite ice and ocean observations. The conclusions are: (1) the model reproduces reasonably well the time mean, variance and skewness of sea surface height. (2) The model biases in temperature and salinity show that while the mean properties follow expectations, the Pacific Water signature in the Beaufort Sea is weaker than observed. (3) However, the modelled freshwater content of the Arctic agrees well with observational estimates. (4) The distribution and volume of the sea ice is shown to be improved in the latest hindcast thanks to modifications to the drag coefficients and to some degree as well to the ice thickness distribution available in CICE. (5) On the other hand, the model overestimates the ice drift and ice thickness in the Beaufort Gyre.

1 Introduction

The CONCEPTS (Canadian Operational Network of Coupled Environmental Prediction Systems) initiative has fostered collaborations between different federal departments (Fisheries and Oceans Canada, Environment Canada and the Department of

GMDD

8, 1–52, 2015

Arctic and N-Atlantic high-resolution ocean and sea-ice model

F. Dupont et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Arctic and N-Atlantic high-resolution ocean and sea-ice model

F. Dupont et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



National Defence) that yielded the development of several operational prediction systems. These include a coupled (atmosphere–ice–ocean) Gulf of Saint-Lawrence system (officially operational since June 2011, Smith et al., 2012), the Global Ice–Ocean Prediction System (GIOPS, run in real-time since March 2014, Smith et al., 2014b), a Great Lakes coupled system (still in development, Dupont et al., 2012), a regional ice-only prediction system (run in real-time since July 2013, Lemieux et al., 2014) and a regional Arctic–North Atlantic ice–ocean system based on the CREG12 (Canadian REGional) configuration with a nominal horizontal resolution of $1/12^\circ$. The latter is the focus of this paper. The GIOPS, Great Lakes and CREG12-based systems are based on NEMO (Nucleus for European Modelling of the Ocean, <http://www.nemo-ocean.eu>), while the coupled Gulf of Saint-Lawrence system is transitioning to NEMO for the ice–ocean component. The development of these systems has benefited greatly from a collaboration with Mercator-Océan in France.

The goal of the regional system based on CREG12 is to provide Canada with short-term ice–ocean predictions and analyses covering parts of the North Atlantic and whole Arctic oceans at high resolution. For this purpose, the regional system will eventually be coupled to the regional weather prediction system and wave prediction system of Environment Canada. The coupled system is expected to improve regional weather and marine forecasting services such as issuing bulletins and warnings in ice infested waters for navigation, energy-exploration and northern communities requirements. As such, the system development has benefited from financial support from the Canadian METAREA program and the Beaufort Regional Environmental Assessment (BREA) project. However, before the full system (analysis¹ + forecast) can be approved for operational use, we need to understand how to use the forecasting component to its full potential, following the best practices of the community running at comparable resolutions. Hence, a series of incremental hindcasts was performed using the forecasting component, each implementing and testing a different aspect of ocean–ice modelling.

¹The analysis covers the data-assimilation aspect of any prediction system.

Arctic and N-Atlantic high-resolution ocean and sea-ice model

F. Dupont et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



These hindcasts are not long enough to test the full robustness of the model in preserving observed water and ice properties at climatic scales (i.e. several decades), as the initial conditions still imprint the model state after 8 years. Nevertheless, discrepancies between atmospheric forcing products and differences in upper-ocean and ice physics are sufficient to create diverging upper-ocean and ice states and variabilities in this short period, that are worth investigating. Moreover, recent satellite missions and extensive and automatized observing in-situ programs (ARGO floats and ice-tethered profilers to cite a few) create a wealth of data covering the hindcast period, which we take advantage of in our validation approach. We are therefore testing the mean state of the model using a few variables, sometimes focusing on some integrated indices over time, or more extensively mapping the model-observation discrepancy in space and time.

In this contribution, we describe the model components and the validation strategy, along with results of the validation of the latest hindcast. The objective is to present to the community the progress made and challenges met in developing a high resolution modelling system for the Arctic-Atlantic oceans, in the spirit of Megann et al. (2014). In assessing the performance of the latest hindcast in terms of ice properties (concentration, thickness and velocity), we include comparison with an intermediate hindcast and the 1/12° resolution equivalent global simulation ORCA12-T321 of Mercator-Océan.

More precisely, Sect. 2 is divided into the description of the model (domain, model components and parameters; Sect. 2.1), the input bathymetry and other initial and boundary conditions (Sect. 2.2), and the description of the validation package (Sect. 2.3). Section 3 provides details of the hindcast simulations (Sect. 3.1), then describes the simulation results in terms of the statistics of the sea surface height, the hydrography and the general circulation (Sect. 3.2) and in terms of sea-ice metrics (concentration, thickness, volume and drift; Sect. 3.3). Section 4 concludes.

2 Model setup, input data and validation package

2.1 Model description

2.1.1 Domain configuration

The global ORCA12 domain (ORCA family grid at a nominal horizontal resolution of $1/12^\circ$ in both longitudinal and latitudinal directions, Drakkar Group, 2007) is used to derive a seamless (i.e., the “north-fold” discontinuity of the global grid is removed) regional domain covering the whole Arctic Ocean and parts of the North Atlantic down to 27° N. The horizontal grid consists of 1580×1817 points on which resolution varies from 8 km at the open boundary in the Atlantic Ocean to an average of 5 km in the Arctic, and down to slightly below 2 km in some of the southern channels of the Canadian Arctic Archipelago (Fig. 1).

The spatial variation of the first Rossby radius of deformation is shown in Fig. 2a. From about 40 km along the southern Atlantic boundary down to a few kilometers in the Labrador Sea, the Greenland, Iceland and Norwegian (GIN) seas and continental shelves, the radius starts re-increasing in the deep Arctic Ocean to above 10 km. Relative to the local resolution (Fig. 2b), the model resolves – grossly speaking – baroclinic eddies in the Sargasso Sea and the Azores region where there are at least two grid spacings for resolving the Rossby radius, but becomes eddy-permitting in the Labrador Sea (one grid spacing) and less than permitting in the GIN seas (under one grid spacing). However, the model is again eddy-resolving in the central Arctic Ocean, which is of importance for the present application.

2.1.2 Ocean component

The ocean model is version 3.1 of NEMO with some code additions from Mercator-Océan, the UK Met Office and the DRAKKAR community. NEMO is an ocean and ice model developed originally in Europe (Madec and NEMO team, 2008), that has

GMDD

8, 1–52, 2015

Arctic and N-Atlantic high-resolution ocean and sea-ice model

F. Dupont et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



evolved very substantially since its introduction in the 2000s. The ocean engine of NEMO is the primitive equation model OPA (Océan Parallélisé; Madec et al., 1998) adapted to regional and global ocean circulation problems. It is intended to be a flexible tool for studying the ocean and its interactions with the other components of Earth's climate system over a wide range of space and time scales (Masson-Delmotte et al., 2006; Drillet et al., 2005; Barnier et al., 2006). An advantage of the NEMO model is its widespread use and continuous tuning by the scientific community (Rattan et al., 2010).

Previous versions of NEMO have been extensively tested and applied in Canada for global, basin and regional applications (Holloway and Wang, 2009; Zhu et al., 2009; Wang et al., 2010; Lu et al., 2014). Present applications of NEMO in CONCEPTS are based on Version 3.1.

2.1.3 Ocean model parameters

We started from the configuration and parameters of the $1/12^\circ$ resolution equivalent global simulation, ORCA12-T321 of Mercator-Océan, which are described below and notes will be made when departing. NEMO is run with the implicit free-surface solver and linear free-surface (a version using a time-splitting approach and a non-linear free-surface, including the simulation of the main constituents of the tides, is presently being evaluated). The present version uses the same 50 vertical z levels used in GLOPS, with spacing increasing from 1 m at the surface to 450 at 5000 m. Bottom partial steps are employed for an accurate representation of the varying bathymetry. The tracer advection uses the Total Variance Diminishing (TVD) scheme. The vectorial form for momentum is chosen, allowing conservation of both energy and enstrophy. The lateral diffusion operator is biharmonic for momentum along geopotential surfaces and harmonic for tracers along isopycnal surfaces. The biharmonic viscosity has a nominal value of $-1 \times 10^{10} \text{ m}^4 \text{ s}^{-1}$ at the southernmost point, and is scaled by the third power of the grid spacing over the rest of the computational domain. The harmonic diffusion coefficient for tracers follows the same resolution-dependance principle, with a nomi-

GMDD

8, 1–52, 2015

Arctic and N-Atlantic high-resolution ocean and sea-ice model

F. Dupont et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



nal value of $50 \text{ m}^2 \text{ s}^{-1}$ and a linear scaling. For momentum, we additionally tested the purely free-slip and no-slip lateral boundary dynamic conditions, but retained the former one for most of the hindcasts. The background values for vertical viscosity and diffusivity are 10^{-4} and $10^{-5} \text{ m}^2 \text{ s}^{-1}$ respectively. We have also experimented with the turbulent kinetic energy (TKE; Gaspar et al., 1990; Blanke and Delecluse, 1993) and generic length scale (GLS; Umlauf and Burchard, 2003) closure schemes. The bottom drag is quadratic with a fixed non-dimensional coefficient of 10^{-3} . The model time step is 360 s for all hindcasts (including ORCA12-T321), except for hindcast H05 that requires a decrease to 180 s after July 2007 to ensure stability in Dease Strait (Canadian Arctic Archipelago).

2.1.4 Sea-ice models

Within NEMO3.1 the ocean is interfaced with the Louvain-La-Neuve sea-ice model version 2 (LIM2, Fichefet and Maqueda, 1997), or version 3 (LIM3, not tested here; Vancoppenolle et al., 2009b, a). However here we also use another community sea-ice model, CICE (described below).

LIM2 is a simple one-category ice model based on a Semtner 2-layer thermodynamic model (two layers of ice and one layer of snow). A Viscous-Plastic (VP) constitutive law relates the internal ice stresses to the strain rates and the ice strength. It is based on an elliptical yield curve and a normal flow rule (Hibler III, 1979). The VP solution is approached by iteration of a relaxation scheme to the implicit ice velocity problem. LIM2 was used for the first two hindcasts (details given below in Sect. 3.1 and Table 1) for sanity checks relative to the configuration used in ORCA12-T321. The latter actually used an upgraded dynamic solver based on the Elastic-VP (EVP) approach (Hunke and Dukowicz, 1997, 2002; Bouillon et al., 2009) instead of the VP solver described above.

CICE (Hunke, 2001; Lipscomb et al., 2007; Hunke and Lipscomb, 2010) is a dynamic/thermodynamic sea ice model, which can be used as a stand-alone model or

GMDD

8, 1–52, 2015

Arctic and N-Atlantic high-resolution ocean and sea-ice model

F. Dupont et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



coupled to an ocean model inside a climate modelling system. Herein, it is coupled to NEMO on the same grid as a single executable (interface code provided by C. Harris of the UK Met Office and S. Alderson of the National Oceanography Centre, personal communication, 2010). CICE calculates the evolution of a thickness distribution. The thickness distribution evolves with both thermodynamic (vertical growth/melt, new ice formation and lateral melt) and dynamic processes (advection and redistribution). The momentum equation is solved with the same EVP approach as described above for LIM2, although on a slightly different stencil (Arakawa C-grid in LIM2 and B-grid for CICE).

2.1.5 LIM2 and CICE parameters

LIM2 solves the VP dynamics with prescribed ice–water and air–ice drag coefficients. The momentum stress is expressed using a simple quadratic law (McPhee, 1975) with a 0° turning angle for both air and ocean in contact with ice. In the ORCA12-T321 run of Mercator-Océan, the air–ice drag was reduced to 1.5×10^{-3} , whereas the default value of 1.63×10^{-3} is used in our NEMO-LIM2 runs. The ice–water drag is fixed to 1×10^{-2} in all LIM2 runs (including the Mercator-Océan run). In ORCA12-T321, the ice module is called with a time-step of 720 s (every two ocean model time-steps), the EVP solver uses 400 sub-timesteps and a damping elastic time of 1350 s. In our LIM2 runs, the VP solver performs 20 outer loops (the default is 2) with a linear residual at convergence of 1×10^{-6} or a maximum of 550 iterations.

In CICE, both air–ice and ocean–ice stresses are also expressed using a simple quadratic law with a 0° turning angle. Following Roy et al. (2014) for our last two hindcasts and since our first ocean layer thickness is relatively small, the ice–ocean drag coefficient is computed by a log-layer assumption using the oceanic first layer thickness and a roughness length scale of 0.03 m as suggested by Maykut and MCPhee (1995) which yields a drag coefficient of 2.32×10^{-2} . The air–ice stress involves a more sophisticated formulation that takes into account the stability of the atmospheric boundary layer. Following again Roy et al. (2014), the roughness length scale for ice surface

is set in our latest run to the value used in the Canadian Global Re-Forecast (CGRF, Smith et al., 2014a) for consistency between the ice-air stress computed in CGRF and in CICE. These modifications can be seen as a more objective way of deriving the drag coefficients, as they are not retrieved from a calibration exercise.

Ten thickness categories are defined in CICE (as in Smith et al., 2014b), with specific representation of both thin ice and thick ridged ice. The remapping advection scheme is used and the EVP solver is run with 920 sub-timesteps. The ice strength is computed using the more physically realistic approach of Rothrock (1975). The number of layers is set to the default value (four ice layers and one for the snow). Fresh or briny waters are exchanged between LIM or CICE and NEMO through a virtual salt flux approach (during ice formation or melting), where an equivalent freshwater flux is computed to represent dilution or concentration of salt at fixed water volume.

2.2 Model input data

2.2.1 Atmospheric forcing

The model is forced at the surface using the CGRF product from 2002 (2003 for some other runs) to 2009. This product consists of a series of re-forecasts using available historical operational analyses from the Canadian Meteorological Centre of Environment Canada. As such, it is not a true reanalysis as other centres produce. However, because it uses the global Canadian Numerical Weather Prediction (NWP) model (last updated in 2011), it provides a consistent set of global forecasts at higher resolution (nominally 33 km at 60° N) than typical reanalyses. The only source of variation in the quality of the reforecasts is the quality of the initial state (the analysis), which varies during the historical period with the assimilation method and volume of observations used. The resolution offered by this product allows for better resolution of mesoscale atmospheric features. The short and long wave radiation fields however require some level of correction as the NWP model is unable to simulate with sufficient accuracy the marine clouds. A climatological correction based on the

GMDD

8, 1–52, 2015

Arctic and N-Atlantic high-resolution ocean and sea-ice model

F. Dupont et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



month of interest but also on the forecast hour is derived from the GEWEX (https://eosweb.larc.nasa.gov/project/srb/srb_table) radiation product.

The frequency of the forcing fields is set to 3 h, using hours 6–27 of each CGRF initiated at 00:00 UTC. CGRF is provided on 10 m wind and 2 m thermodynamic levels. Those are not true “prognostic” model levels but since conventions and model output dissemination requires these levels, a “diagnostic” procedure is used to derive quantities there. The first prognostic level for wind and temperature in CGRF is in fact approximately at 40 m, and quantities at this level are also available and are thought to be less dependent on assimilated surface conditions and approximations made during the diagnostic procedure. We have therefore used the product at this level as input to the CORE air–sea exchange bulk formulae and the equivalent in CICE. The only limitation to this approach is in LIM2, where input atmospheric conditions are assumed at 10 m with pre-set constant neutral coefficients, causing an over-estimation of wind-stress by approximately 20 to 50 % (the same overestimation problem likely affects the calculation of turbulent heat exchanges).

2.2.2 Bathymetry, initial and lateral boundary conditions

The bathymetry used in the CREG12 configuration is taken from that used in the ORCA12-T321 run of Mercator-Océan. It is based on ETOPO2 (<http://www.ngdc.noaa.gov/mgg/global>, Amante and Eakins, 2009). The minimum depth is set at 20 m.

Two sets of initial ocean conditions (comprising 3-D velocities, temperature, salinity and sea surface height) have been used. Firstly a reanalysis product, GLORYS2v1 (Ferry et al., 2012) is used. This covers the satellite-altimetry and ARGO period (1993–2010), with assimilation of both of these datasets in the reanalysis as well as other in-situ data. However we found that, although the assimilation of observations leads to a remarkable agreement with observations at lower latitudes, GLORYS2v1 suffers from serious departures relative to observations and to the Polar Science Center Hydrographic Climatology (PHC, http://psc.apl.washington.edu/nonwp_projects/PHC/

GMDD

8, 1–52, 2015

Arctic and N-Atlantic high-resolution ocean and sea-ice model

F. Dupont et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Climatology.html) in the Arctic². The second set of initial conditions used is simply derived from the ORCA12-T321 run of Mercator-Océan, which has better hydrographic properties in the Arctic Ocean but is not as accurate as GLORYS2v1 at lower latitudes.

Sea ice initial conditions are taken from the same initial condition product, that is either GLORYS2v1 or ORCA12-T321, which use the mono-category LIM2 model. The ice concentration and ice thickness of these products are applied to the corresponding ice category in CICE, the other categories remaining empty. It then takes several month of simulations before a realistic ice distribution can be recovered. An initial spread among several categories would therefore be more realistic. For snow, the ice category that receives the ice volume also receives the snow volume present in the initial conditions.

Along the lateral open boundaries, time-evolving monthly conditions (comprising 3-D velocities, temperature and salinity from 2002 to 2009) are taken from the same products as the initial conditions. More specifically, a clamped velocity condition is specified (hence lateral transport) and a radiation scheme following the advective characteristic is applied for temperature and salinity combined with restoring to input values. The restoring time is 15 days when radiating outward and 1 day when inward. A closed wall boundary condition is applied to sea ice in LIM2 and CICE.

2.3 Validation package

Validation of the system is performed by comparing model outputs with ocean observations. Additionally, the model outputs are compared with other model estimates and with climatologies. During the development phase, with the model running in hindcast mode, this validation provides an assessment of the improvements introduced with

²Among other poor characteristics, the doming of sea surface height in the Beaufort Sea is absent and the Atlantic layer apparently spreads anti-cyclonically instead of cyclonically. This is in apparent contrast to studies done using GLORYS1, which were more successful, such as in Lique et al. (2011).

GMDD

8, 1–52, 2015

Arctic and N-Atlantic high-resolution ocean and sea-ice model

F. Dupont et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



each change to the model configuration. Once the forecast system is operational, the validation package will provide an assessment of forecast accuracy.

The CONCEPTS validation strategy defines a set of model output fields, a database of ocean observations from both in-situ and remote sensing measurements, and a suite of metrics for comparing the two. This approach has been designed for the CREG12 configuration, but was developed in such a way that it can easily be transferred to other CONCEPTS systems. The key model outputs for validation are sea surface height, ocean temperature, salinity and velocity, and sea ice thickness, concentration and velocity. Additional derived output fields include transports through sections, freshwater content and mixed layer depth. The observation database incorporates measurements included in existing global databases, combined with data from individual observation missions. These include missions using new technologies developed to provide measurements in the ice-covered regions of the Arctic. The ocean observation database includes traditional ship-deployed and moored in-situ measurements of temperature, salinity and velocity, together with measurements from ARGO drifting profilers, ice-tethered profilers, gliders, mammal-mounted instruments and satellite remote sensing. The sea ice observations include thickness and drift measurements from ice mass balance buoys and upward-looking sonar together with remote sensing from aircraft- and satellite-mounted instruments.

3 Model simulations and validation

3.1 Simulations

Five hindcast simulations, H01 to H05, are carried out covering the years 2003 to 2009, and these are briefly described in Table 1. LIM2 is used in H01 and H02, and CICE in H03 and higher. H01 is initialized from GLORYS2v1, which is found less reliable than ORCA12-T321 in the Arctic Ocean, our focus region. Hence H02 and higher are started instead from ORCA12-T321. Changes related to air-ice and ocean-ice drags based

GMDD

8, 1–52, 2015

Arctic and N-Atlantic high-resolution ocean and sea-ice model

F. Dupont et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



on Roy et al. (2014) were incrementally implemented in H03 to H05. Parameters are defined in Sect. 2.1.5. Hence H02 uses for instance a lower ocean–ice drag coefficient relative to H05 (approximately half). The treatment of the air–ice stress is also noteworthy different in H02 as explained in Sect. 2.2.1, and therefore the magnitude of the stress is over-estimated relative to H05.

3.2 Hydrography and circulation validation

The focus of the validation is the most-recent model run, H05, but there are some brief comparisons with the earlier H02, which incorporates the LIM2 ice model rather than the CICE model. In this sense and in spite of other differences, H02 is the closest simulation to the ORCA12-T321 run. Most of the comparisons presented here are for the mean fields for the period 2003–2009 with additional discussions on time variability.

3.2.1 Sea surface height

Satellite altimeters provide a continuous record of sea surface height (SSH) anomalies since 1993 (Benveniste, 2011). Figure 3 shows the mean (top), standard deviation (middle) and skewness (bottom) of SSH for the North Atlantic for the period 2003–2009 from the model hindcast H05 (left panels) and from the satellite record. The altimeter estimates of the standard deviation and skewness are produced using the gridded $1/4^\circ$ SSH anomaly product distributed by Archiving, Validation and Interpretation of Satellite Oceanographic data (AVISO, <http://www.aviso.oceanobs.com/en/data/products/auxiliary-products/mss/index.html>). The mean altimeter SSH is the sum of the 2003–2009 SSH anomalies and the CNES–CLS09 Mean Dynamic Topography (MDT, Rio et al., 2011).

The mean SSH fields from the model and altimeter record are very similar. The sharp gradient of the Gulf Stream can be seen in both, leaving the coast of North America around 35° N, and following a similar path eastwards. The high SSH of the subtropical gyre can be seen to the south of the Gulf Stream, and the low SSH of the subpolar

gyre to the north. The model estimate shows some sharper gradients, for example along the Labrador coast, but this is likely because of the higher horizontal resolution of the model ($1/12^\circ$) compared with the resolution of the altimeter product ($1/4^\circ$).

The spatial distribution of the magnitude of SSH variability, represented by the standard deviation plots, shows good agreement between the model and the altimeter measurements. The altimeter data shows in general though a broader structure of medium values of standard deviation to the south of Gulf Stream whereas that of the model shows medium values extending along the path of the North Atlantic Current.

Positive and negative skewness corresponds to the meandering of a free jet such as the Gulf Stream or the variability caused by warm- and cold-core eddies (Thompson and Demirov, 2006). Typically, the zero contour of skewness separating strong regions of negative and positive skewness is a good indicator of the centre position of the mean currents. There is again good agreement between the model and the altimeter record in terms of the distribution of skewness for the Gulf Stream area, with the zero contour of the model being positioned slightly more to the north. A broad region of negative skewness in the model is also clearly visible in the mid to eastern Atlantic Ocean which is not seen in the altimeter data. The interpretation of this is more difficult, except to note that the model must be producing more intense cyclonic than anticyclonic deviations in this region. Finally, the zero contour helps to define the position of the Azores Current, which is well reproduced in the model in general but with perhaps a slightly more intense and narrower jet.

Most of the satellite altimeters that contribute to the AVISO record are unable to produce useful estimates of SSH in the Arctic, either because their orbits do not extend far enough north or because sea ice prevents the altimeter signal reaching the sea surface. However Farrell et al. (2012) used measurements from the ICESat and Envisat satellite missions to create an Arctic MDT for the period from 2003 to 2009, corresponding with the period of the hindcast. Figure 4 shows the mean Arctic SSH from H05 and the MDT of Farrell et al. (2012). There is good agreement between the two estimates, both in terms of the patterns of SSH and the SSH gradients. For exam-

GMDD

8, 1–52, 2015

Arctic and N-Atlantic high-resolution ocean and sea-ice model

F. Dupont et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



ple H05 shows a cross-Arctic sea level difference, from the high of the Beaufort Gyre to the low north of Spitzbergen, of approximately 60 cm compared with a difference of about 65 cm in the MDT of Farrell et al. (2012). Kwok and Morison (2011) similarly use ICESat data (winter only) to estimate the MDT of the Arctic, including its variability. The interannual variability of mean SSH in H05 (not shown here) compares well with their estimates, particularly in the Canada Basin.

3.2.2 Surface circulation

Figure 5 compares the mean near-surface current speeds for 2003–2009 from hindcast H05 to a $1/2^\circ$ resolution climatology derived from surface drifter velocity estimates (Lumpkin and Johnson, 2013). The general agreement is remarkable, with the model being able to provide an increased level of detail. The estimates of the speed and the position of the Gulf Stream and the North Atlantic Current are in good agreement. On the north flank of the Gulf Stream, a weak but persistent branching is clearly visible in both plots, east of 70° W, although that of the model detaches from and rejoins the Gulf Stream a little too early. This secondary current system is likely related to the Slope Water Current described in Pickart et al. (1999) and Dupont et al. (2006). The East and West Greenland currents, and the Labrador Current, contain more details in the model than can be captured by the drifter resolution, but the separation of coastal and shelf jets is in good agreement with other observations (e.g., Higginson et al., 2011). Again, the path of the Azores Current is visible in both model and observations.

3.2.3 Temperature and salinity

Quality-controlled measurements of ocean temperature and salinity are available from the global CORA3.4 database distributed by myOcean (www.myOcean.eu). This database includes measurements from ship-based surveys, moorings and the drifting profilers of the ARGO network. In the Arctic there are relatively few observations compared with other ocean basins. Whilst some observation programs have been incor-

GMDD

8, 1–52, 2015

Arctic and N-Atlantic high-resolution ocean and sea-ice model

F. Dupont et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



porated into the CORA3.4 database, others are not yet included. We have undertaken a search of data available from all programs, and combined them with the CORA3.4 observations where they are missing. These observation programs include the Beaufort Gyre Exploration Project (BGEF, <http://www.whoi.edu/page.do?pid=66296>), the Ice-Tethered Profiler project (ITP, <http://www.whoi.edu/page.do?pid=20756>), the Canadian Basin Observational System (CABOS, <http://nabos.iarc.uaf.edu/index.php>), the Switchyard project (<http://psc.apl.washington.edu/switchyard/overview.html>), the North Pole Environmental Observatory (NPEO, <http://psc.apl.washington.edu/northpole/>), and monitoring programs in Davis Strait (e.g., Curry et al., 2013), Barrow Strait (e.g., Hamilton et al., 2013) and Fram Strait (e.g., Schauer et al., 2008).

Figure 6 shows the mean TS bias for hindcast H05 for the period 2003–2009. Model values are extracted at the same time and location as observations, and the bias is calculated as the model estimate minus the observation. The biases are averaged in 1° bins for the top 200 m, and between 200 and 500 m depth. These intervals are chosen to quantify the near-surface (including shelf) and intermediate depth anomalies.

Over large areas of both the North Atlantic and the Arctic oceans the average temperature biases are less than $\pm 1^\circ\text{C}$ and the salinity biases are less than ± 0.5 PSU. However, a warm and salty bias is seen in the central North Atlantic and in the currents that form the subpolar gyre. This bias is strongest in the surface layers, averaging more than 2°C and 0.75 PSU in some places, and extends into the Iceland and Norwegian seas. The temperature bias extends southwards along the path of the Gulf Stream, particularly in the surface layers. A cold bias also extends from the north side of the Gulf Stream toward the coast of the US and Canada. The salinity bias is largest in the Labrador Current. A salty bias is seen in the upper layers of the Beaufort Sea, extending along the coast of Canada toward Fram Strait (see also Sect. 3.2.4). Conversely, the waters in the centre of the Greenland-Iceland-Norwegian Seas are colder and fresher.

The vertical structure of the temperature and salinity biases is shown in Fig. 7. It shows the TS bias profiles for all observations between 2003 and 2009, averaged in each of four domains representing regions of oceanographic interest (the subtropical

GMDD

8, 1–52, 2015

Arctic and N-Atlantic high-resolution ocean and sea-ice model

F. Dupont et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



and subpolar gyres, the Beaufort Gyre and the Nordic seas). These domains (except for the subtropical gyre) correspond with regions of relatively high temperature or salinity anomalies identified in Fig. 6. The profiles for the subtropical gyre domain (box d) show that the model does a good job of representing both temperature and salinity, although the top ocean layers are too fresh by 0.5 PSU. In the subpolar gyre domain (box c) the model bias in salinity is positive, with a maximum of less than 0.5 PSU around 100 m depth. The warm bias has a maximum of around 2 °C at a similar depth. In the Greenland and Norwegian seas (box b), there is fresh and cold bias, restricted to the top 100 m of the ocean. In the Beaufort Sea (box a) the temperature biases are very small (less than 0.5 °C), but the profile shows the bias to be cold in the Atlantic water layer (around 500 m depth) and near the surface, and slightly warm in the Pacific water layer (around 150 m depth, Steele et al., 2004). The vertical temperature structure is not well reproduced by the model. This suggests that there may be problems with the transport and transformation of Pacific waters in the model, and this is an area for further investigation. We can only tell at this point that Pacific water signature weakens with time in H05 (not shown). A salty bias in the Beaufort Sea is restricted to the upper 75 m of the water column. Note that GLORYS2v3 and PHC are in good agreement for temperature but both depart significantly from the observations in this area.

Examining the Beaufort Sea salinity bias in a little more detail, Fig. 8 shows the mean liquid freshwater content equivalent depth for the Arctic from H05 and from PHC. The freshwater content is calculated using the method described in Proshutinsky et al. (2009), with a reference salinity of 34.8. There is good agreement in terms of the distribution of liquid freshwater, with the greatest concentration in the Beaufort Gyre, but the total modelled freshwater content in the gyre is greater than in the climatology. This is likely because the PHC does not incorporate observations beyond 1998 and therefore does not reflect the recent increase in freshwater content estimated by Proshutinsky et al. (2009). In order to investigate whether these recent variations are reproduced in the different simulations, we compare the total modelled (H02, H05 and ORCA12-T321) liquid freshwater content in the Beaufort Gyre to the estimate of Proshutinsky

**Arctic and N-Atlantic
high-resolution
ocean and sea-ice
model**

F. Dupont et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



et al. (2009) in Fig. 9. The two CREG12-based hindcasts reproduce fairly realistically the observed increase in freshwater content (although tapering by the end of the simulation period) whereas the ORCA12-T321 content shows no such increase. We partly attribute this discrepancy to differences in atmospheric forcing products used in our hindcasts and ORCA12-T321. The fact that H05 shows a slightly poorer agreement with the observed freshwater estimates than H02 by the end of the simulation is due to the weaker Ekman pumping in the Beaufort Gyre, the latter explained by the smaller roughness and associated ice-air drag as described in Sect. 2.1.5 and Table 1. This will be illustrated from a different point of view in Sect. 3.3.2.

3.2.4 Sections across Fram Strait and Davis Strait

Arrays of moorings have been deployed across the main pathways for exchange of water between the Arctic and Atlantic, for example in Fram Strait (Schauer et al., 2008) and in Davis Strait (Curry et al., 2013). Figures 10 and 11 show a comparison of mean temperature, salinity and velocity observations with corresponding estimates from hindcast H05 for Fram Strait and Davis Strait respectively. The Fram Strait mooring observations cover the period 2005–2009, and the Davis Strait moorings cover 2004–2009, although not all instruments were deployed for the whole period.

Overall there is very good agreement between the model and observations in Fram Strait. The large velocity of the northward-flowing West Spitzbergen Current and the southward-flowing East Greenland Current are very similar in magnitude and location, and the temperature and salinity gradients across the strait are broadly similar. However the modelled northward-flowing water close to Spitzbergen is saltier than observed. Also the observations show a weaker northward-flowing branch of the West Spitzbergen Current in the central channel, as described by Schauer et al. (2004), but this is absent in the simulation. This may explain a cold bias in the modelled near-surface waters in the centre of the strait.

In Davis Strait the observed and modelled temperatures are in good agreement. The salinity fields are also generally good, and the velocity maximums of the northward-

GMDD

8, 1–52, 2015

Arctic and N-Atlantic high-resolution ocean and sea-ice model

F. Dupont et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



flowing West Greenland Current and the southward-flowing Baffin Island Current (BIC) are similar in magnitude. However, the northward-flowing water on the Greenland shelf is a little too salty, likely related to the salty bias in the subpolar gyre described earlier, and the BIC is displaced further offshore in the model. There does not seem to be

a strong temperature or salinity bias in the Arctic outflows through either Fram Strait or Davis Strait, suggesting that this is not the source of the biases seen in the Atlantic and discussed in the previous section.

The mean net liquid volume transport for 2003–2009 in hindcast H05 for Fram Strait is 2.7 Sv (1 Sv is $10^6 \text{ m}^3 \text{ s}^{-1}$) toward the south, compared with an observational estimate of 2 ± 2.7 Sv (Schauer et al., 2008). For Davis Strait the model mean liquid volume transport is 1.9 Sv toward the south, compared with an observed 1.6 ± 0.5 Sv (Curry et al., 2013).

3.3 Sea-ice validation

3.3.1 Ice concentration, thickness and volume

Estimates of the total ice area have been derived from satellite products at the National Snow and Ice Data Center (NSIDC, Cavalieri et al., 1996), filling the North Pole data hole with 95 % ice concentration. Comparing H02 and H05, the implementation of CICE in H05 is beneficial in terms of reproducing the seasonal cycle (Fig. 12, top panel). The multi-ice category allows for larger rates of melting and growth in the small ice thickness categories, thus enhancing the seasonal cycle of ice area and bringing it closer to observations. A secondary effect is related to the different drag coefficients used in the different CREG12 hindcasts and ORCA12-T321, which impacts the Ekman convergence in the ice over the Beaufort Gyre. In terms of interannual variability, looking at September ice area, H05 ice loss is faster than H02 at the beginning of the simulation (2003–2005), bringing the ice area closer to observations between 2005 and 2009 (Fig. 12, bottom panel). H05 September ice area then starts to depart from observations after 2010 due to an anomalous accumulation of ice in the Beaufort Gyre. H02

GMDD

8, 1–52, 2015

Arctic and N-Atlantic high-resolution ocean and sea-ice model

F. Dupont et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



and T321 have a too large September ice area but the decreasing trend is in general closer to observations than in H05. This can be related to the fact that both H02 and ORCA12-T321 are in better thermodynamic balance with the initial condition, which itself is derived from a simulation using LIM2, than H05 which goes through a two-year adjustment period. The 2007 minimum is well reproduced by H05 in terms of total ice area, although the regional structure shows differences from the observations (Fig. 13). The ice concentration in Beaufort and Chukchi Seas is a little too high and that in the tongue of ice connecting the central pack along the Severnaya Zemlya Archipelago to the mainland is somewhat too low. The ORCA12-T321 and H02 ice concentration fields are very similar in spatial structure, with T321 showing a sharper transition at the ice pack edge. They both overestimate the ice concentration in the Beaufort Sea and in the East Siberian sector, in agreement with the total ice area results.

In-situ ice thickness observations are available from a number of different sources. Ice mass balance buoys (e.g., Polashenski et al., 2011) drift with the ice, measuring the evolution of the ice thickness. Figure 14 shows the mean difference between the model sea ice thickness and the measured thickness, averaged over the duration of the model runs. H05, which uses the CICE ice model, clearly produces a result closer to observations than H02, which uses LIM2, but the ice in the Beaufort Gyre is still too thick. Upward-Looking Sonars (ULS) have been deployed on a number of subsurface moorings, providing high frequency measurement of the ice draft from beneath. This data can be used to produce an estimate of the thickness distribution at the mooring location, which can be compared with the ice thickness distribution from the CICE model. Figure 15 compares the estimates from H05 with observations at one of the BGEP moorings (<http://www.whoi.edu/beaufortgyre/data>) and at the NPEO mooring (<http://psc.apl.washington.edu/northpole/>, Morison et al., 2002). The model reproduces the thickness distribution at both sites quite well, although it tends to overestimate the thicker ice categories in the Beaufort Gyre and underestimate them near the North Pole.

The spatial structure of the ice mean thickness (local total ice volume divided by total ice concentration) is also compared to estimates from the ICESat mission (Kwok et al., 2009). Figure 16 shows that ORCA12-T321 and H02 (both of which use LIM2) overestimate thickness over a large area. The mean ice thickness in H05 is closer to the ICESat observations, but there is a region of overestimated thickness in the Beaufort Gyre and an underestimation elsewhere which is consistent with our findings from the ice mass balance and ULS measurements, and also with results of Roy et al. (2014).

Finally, the domain total ice volume of the different model simulations is compared to the estimate of the data-assimilative model PIOMAS (Zhang and Rothrock, 2003). This model ice volume compares well with estimates from ICESat and CryoSat2 (Laxon et al., 1994) and is therefore deemed a reasonable reference. The error bars are not known but could be up to 25 %. The seasonal cycle (Fig. 17, top panel) for H05 is very close to the PIOMAS value, and a clear improvement over H02 and ORCA12-T321. The September values (Fig. 17, bottom panel) emphasize the discrepancy between the different hindcasts: H05 is close to PIOMAS in magnitude and trend, while ORCA12-T321 and H02 do not have a clear trend and the volume is overestimated by 50 to 100 %. This volume overestimation in ORCA12-T321 and H02 is consistent with the findings from in-situ and satellite thickness measurements. Here too, the different drag coefficients partially explained the convergence and accumulation of ice in the Beaufort Gyre. The higher ice–ocean drag and the lower air–ice drag in H05 both concur to reduce the ice velocity and therefore the Ekman convergence there, relative to H02.

The modification to the surface ice roughness between H04 and H05 has a positive impact, improving the absolute value and trend in the volume (not shown). However, although the total volume of H05 is very much on par with estimates from PIOMAS, this conceals regional errors such as an overestimation of ice thickness in the Beaufort Gyre that were discussed earlier. The source of these errors in the Beaufort Gyre is likely related to the ice drift pattern, discussed in the following section.

GMDD

8, 1–52, 2015

Arctic and N-Atlantic high-resolution ocean and sea-ice model

F. Dupont et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



3.3.2 Ice drift

Ice drifts for March 2003–2008 are compared to satellite estimates from Fowler et al. (2013) (source: NSIDC) in Fig. 18 and observations from ice buoys deployed as part of the International Arctic Buoy Program (IABP; <http://iabp.apl.washington.edu/>). One can see the improvement from hindcast H02 to H05 as the ice–water and air–ice drags are adjusted following a semi-objective approach (Roy et al., 2014). However, the ice drift in H05 is still overestimated. In the ORCA12-T321 solution, the air–ice drag is slightly decreased and actually yields the best modelled velocity fields. This is also evident from the ice drift bias calculated relative to the IABP buoys (Fig. 19). The Pathfinder gridded estimates (Fowler et al., 2013) are the closest to buoy drifts, followed by ORCA12-T321 and H05. The H05 bias is close to that of ORCA12-T321 but starts to deviate in late 2006.

This evidence suggests that Ekman transport is still acting too strongly in H05, driving a convergence of ice and maintenance of multi-year ice in the Beaufort Gyre. An obvious reasoning is that the air–ice stress is too large (either due to too large winds or/and drag coefficient), driving the ice too fast. However, the CGRF surface winds tend to show a weak negative bias compared to observations at Ice Station Tara (not shown). This is in contrast to some reanalysis products compared by Jakobson et al. (2012) such as ERA-INT. Moreover, the surface ice roughness lengthscale in CGRF is actually smaller than the one used in ERA-INT. Hence the air–ice stress is less likely to be overestimated. On the other hand, the freshwater content increase during the period 2003–2009 in H05 is slightly weaker than observed (Fig. 9), which suggests the opposite, that is, the convergence of freshwater due to Ekman transport acting of the ocean may be underestimated. This issue needs to be further studied. Some mechanisms explaining variations in the Arctic freshwater content caused by dependencies on model parameters are also assessed in Roy et al. (2014). Preliminary results point to deficiencies in the vertical mixing scheme used in H05 ($k - \epsilon$) in the Arctic upper ocean

GMDD

8, 1–52, 2015

Arctic and N-Atlantic
high-resolution
ocean and sea-ice
model

F. Dupont et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



which would explain the overly strong ice drift in the Beaufort Gyre by underestimating the shallow convection under the ice.

Additionally, we note that the lack of landfast ice parametrization may explain the over-estimation of the ice drift in all model runs in the East Siberian, Laptev and Kara Seas in Fig. 18.

4 Conclusions

The development of a high resolution ice–ocean modelling system is a challenging task that requires a team effort. In CONCEPTS this is achieved by collaborations among different Canadian government departments and international collaborators such as Mercator Océan. The CREG12-based system consists of state-of-the-art ocean and sea-ice models, a comprehensive validation package, and a data assimilation capability under development. Before proposing the system for operational implementation, the capability of the ice–ocean model to produce high-quality hindcasts must be demonstrated. Hence, the present approach of producing a series of hindcasts and by identifying deficiencies, helps in deciding which aspects of the system need to be improved. For instance, the upper ocean physics and more accurate initialization fields appear as areas of particular concern.

Each multi-year hindcast, driven by the high-resolution CGRF forcing, shows incremental improvements with changes to the initial and boundary conditions, the lateral friction schemes, turbulent mixing parametrizations, and finally the change of sea-ice model from LIM2 to CICE. The validation package includes a variety of ocean and sea-ice observations. It demonstrates the capability of the model in hindcasting the mean, variance and skewness of the SSH, the position and strength of the surface circulation.

In terms of temperature-salinity distributions, the initial conditions (however accurate or poor they can be) still imprint the results after 8 years and therefore only variations in the upper ocean can be analyzed. From this point of view, the $k - \epsilon$ mixing scheme seems adequate in the north Atlantic but likely underestimates the shallow convection

GMDD

8, 1–52, 2015

Arctic and N-Atlantic high-resolution ocean and sea-ice model

F. Dupont et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Arctic and N-Atlantic high-resolution ocean and sea-ice model

F. Dupont et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



future. Second, a landfast ice parametrization should improve the representation of ice dynamics over the shelves, especially on the Siberian side, and we are hopeful for results in the very near future in this area as well. Third, two-way coupling between the wave field, the ocean and the ice are in progress (Dumont et al., 2011) and are expected to improve substantially the upper ocean response (with the addition of Stokes currents and induced mixing), the representation of the ice in the marginal ice zone, and improving the wave field in general.

Additionally, promising advances in the parametrization of form drag (Tsamados et al., 2014) between ice-air and sea-ice, and rheology (Tsamados et al., 2013) need to be implemented and tested, although for the latter, it is not clear how beneficial this new rheology can be at high resolution – which is true of any existing rheology for that matter. The two latter advances are already available in CICE5 (Turner and Hunke, 2014). We also plan to move to NEMO Version 3.6 in the upcoming future, which will offer support for coupling to CICE5. We are finally hopeful to increase the vertical resolution of the ocean component to 75 levels with a limit to 250 m thick layers in the deep ocean, instead of the present 450 m limit. This would put us on par with DRAKKAR and Mercator-Océan's latest standards.

Acknowledgements. This research was made possible thanks to grant from BREA. The Gewex data was obtained from the NASA Langley Research Center Atmospheric Science Data Center. The SSH anomaly data was obtained from AVISO. The reanalysis products (GLORYS) and the observational data CORA3.4 were accessed through myOcean. The BGEP and ITP data was downloaded from Woods Hole Oceanographic Institute. We are grateful for NSIDC for providing ice concentration and velocity products and IABP for arctic buoy data. Finally, the manuscript benefited from the insightful comments of Daniel Deacu.

References

- Amante, C. and Eakins, B. W.: ETOPO1 1-Arc-Minute Global Relief Model: Procedures, Data Sources and Analysis, Tech. rep., NOAA Technical Memorandum NESDIS NGDC-24, National Geophysical Data Center, NOAA, doi:10.7289/V5C8276M, 2009. 10
- 5 Barnier, B., Madec, G., Penduff, T., Molines, J.-M., Tréguier, A.-M., Le Sommer, J., Beckmann, A., Biastoch, A., Böning, C., Dengg, J., Derval, C., Durand, E., Gulev, S., Rémy, E., Talandier, C., Theetten, S., Maltrud, M. E., McClean, J., and De Cuevas, B.: Impact of partial steps and momentum advection schemes in a global ocean circulation model at eddy permitting resolution, *Ocean Dynam.*, 56, 543–567, doi:10.1007/s10236-006-0082-1, 2006. 6
- 10 Benveniste, J.: Radar altimetry: past, present and future, in: *Coastal Altimetry*, edited by: Vignudelli, S., Kostianoy, A., Cipolline, P., and Benveniste, J., Springer-Verlag, 1–17, doi:10.1007/978-3-642-12796-0_1, 2011. 13
- Blanke, B. and Delecluse, P.: Variability of the tropical Atlantic Ocean simulated by a general circulation model with two different mixed-layer physics, *J. Phys. Oceanogr.*, 23, 1363–1388, 1993. 7
- 15 Bouillon, S., Morales-Maqueda, M. A., Legat, V., and Fichet, T.: An elastic-viscous-plastic sea ice model formulated on Arakawa B and C grids, *Ocean Model.*, 27, 174–184, doi:10.1016/j.ocemod.2009.01.004, 2009. 7
- Campin, J.-M., Marshall, J., and Ferreira, D.: Sea ice–ocean coupling using a rescaled vertical coordinate z^* , *Ocean Model.*, 24, 1–14, doi:10.1016/j.ocemod.2008.05.005, 2008. 24
- 20 Cavalieri, D. C., Parkinson, C., Gloersen, P., and Zwally, H. J.: Sea Ice Concentrations from NIMBUS-7 SMMR and DMSP SSM/I Passive Microwave Data, [1979–2006], Tech. Rep., digital media, National Snow and Ice Data Center, Boulder, Colorado USA, 1996 (last updated 2008). 19
- 25 Curry, B., Lee, C., Petrie, B., Moritz, R., and Kwok, R.: Multi-year volume, liquid freshwater, and sea ice transports through Davis Strait, 2004–2010, *J. Phys. Oceanogr.*, 44, 1244–1266, doi:10.1175/JPO-D-13-0177.1, 2013. 16, 18, 19
- Drakkar Group: Eddy permitting ocean circulation hindcasts of past decades, *Clivar Exchanges*, 12, 8–10, 2007. 5
- 30 Drillet, Y., Bourdallé-Badie, R., Siefridt, L., and Provost, C. L.: Meddies in the Mercator North Atlantic and Mediterranean Sea eddy-resolving model, *J. Geophys. Res.*, 110, C03016, doi:10.1029/2003JC002170, 2005. 6

Arctic and N-Atlantic high-resolution ocean and sea-ice model

F. Dupont et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Dumont, D., Kohout, A., and Bertino, L.: A wave-based model for the marginal ice zone including a floe breaking parameterization, *J. Geophys. Res.*, 116, C04001, doi:10.1029/2010JC006682, 2011. 25
- Dupont, F., Hannah, C. G., and Wright, D. G.: Model investigation of the slope water, north of the Gulf Stream, *Geophys. Res. Lett.*, 33, L05604, doi:10.1029/2005GL025321, 2006. 15
- Dupont, F., Chittibabu, P., Fortin, V., Rao, Y. R., and Lu, Y.: Assessment of a nemo-based hydrodynamic modelling system for the great lakes, *Water Qual. Res. J. Can.*, 47, 198–214, doi:10.2166/wqrjc.2012.014, 2012. 3
- Farrell, S. L., McAdoo, D. C., Laxon, S. W., Zwally, H. J., Yi, D., Ridout, A., and Giles, K.: Mean dynamic topography of the Arctic Ocean, *Geophys. Res. Lett.*, 39, L01601, doi:10.1029/2011GL050052, 2012. 14, 15, 37
- Ferry, N., Parent, L., Garric, G., Bricaud, C., Testut, C.-E., Le Galloudec, O., Lellouche, J.-M., Drevillon, M., Greiner, E., Barnier, B., Molines, J.-M., Jourdain, N. C., Guinehut, S., Cabanes, C., and Zawadzki, L.: GLORYS2V1 global ocean reanalysis of the altimetric era (1992–2009) at meso scale, *Mercator Quarterly Newsletter*, 44, 29–39, available at: <http://www.mercator-ocean.fr/eng/actualites-agenda/newsletter>, 2012. 10
- Fichefet, T. and Maqueda, M. A. M.: Sensitivity of a global sea ice model to the treatment of ice thermodynamics and dynamics, *J. Geophys. Res.*, 102, 12609–12646, doi:10.1029/97JC00480, 1997. 7
- Fowler, C., Emery, W., and Tschudi, M.: Polar Pathfinder Daily 25 km EASE-Grid Sea Ice Motion Vectors, Version 2 [2002–2009], Tech. Rep., NASA DAAC at the National Snow and Ice Data Center, Boulder, Colorado USA, 2013. 22
- Gaspar, P., Gregoris, Y., and Lefevre, J. M.: A simple eddy kinetic energy model for simulations of the oceanic vertical mixing: tests at station Papa and long-term upper ocean study site, *J. Geophys. Res.*, 95, 16179–16193, 1990. 7
- Griffies, S. M., Pacanowski, R. C., Schmidt, M., and Balaji, V.: The GFDL CM3 coupled climate model: characteristics of the ocean and sea ice simulations, *J. Climate*, 24, 13520–13544, doi:10.1175/2011JCLI3964.1, 2011. 24
- Hamilton, J. M., Collins, K., and Prinsenber, S. J.: Links between ocean properties, ice cover, and plankton dynamics on interannual time scales in the Canadian Arctic Archipelago, *J. Geophys. Res.*, 118, 5625–5639, doi:10.1002/jgrc.20382, 2013. 16
- Hibler III, W. D.: A dynamic thermodynamic sea ice model, *J. Phys. Oceanogr.*, 9, 815–843, 1979. 7

- Hibler, W. D., Roberts, A., Heil, P., Proshutinsky, A., Simmons, H., and Lovick, J.: Modeling M2 tidal variability in arctic sea-ice drift and deformation, *Ann. Glaciol.*, 44, 418–428, 2006. 24
- Higginson, S., Thompson, K. R., Huang, J., Véronneau, M., and Wright, D. G.: The mean surface circulation of the North Atlantic subpolar gyre: a comparison of estimates derived from new gravity and oceanographic measurements, *J. Geophys. Res.*, 116, C08016, doi:10.1029/2010JC006877, 2011. 15
- Holloway, G. and Wang, Z.: Representing eddy stress in an Arctic Ocean model, *J. Geophys. Res.*, 114, C06020, doi:10.1029/2008JC005169, 2009. 6
- Hunke, E. C.: Viscous-plastic sea ice dynamics with the EVP model: linearization issues, *J. Comput. Phys.*, 170, 18–38, 2001. 7
- Hunke, E. C. and Dukowicz, J. K.: An elastic-viscous-plastic model for sea ice dynamics, *J. Phys. Oceanogr.*, 27, 1849–1867, 1997. 7
- Hunke, E. C. and Dukowicz, J. K.: The elastic-viscous-plastic sea ice dynamics model in general orthogonal curvilinear coordinates on a sphere—incorporation of metric terms, *Mon. Weather Rev.*, 130, 1848–1865, 2002. 7
- Hunke, E. C. and Lipscomb, W. H.: CICE: The Los Alamos Sea Ice Model Documentation and Software User's Manual, Tech. Rep., Los Alamos National Laboratory, 2010. 7
- Jakobson, E., Vihma, T., Palo, T., Jakobson, L., Keernik, H., and Jaagus, J.: Validation of atmospheric reanalyses over the central Arctic Ocean, *Geophys. Res. Lett.*, 39, L10802, doi:10.1029/2012GL051591, 2012. 22
- Kwok, R. and Morison, J.: Dynamic topography of the ice-covered Arctic Ocean from ICESat, *Geophys. Res. Lett.*, 38, L02501, doi:10.1029/2010GL046063, 2011. 15
- Kwok, R., Cunningham, G. F., Wensnahan, M., Rigor, I., Zwally, H. J., and Yi, D.: Thinning and volume loss of Arctic sea ice cover: 2003–2008, *J. Geophys. Res.*, 114, C07005, doi:10.1029/2009JC005312, 2009. 21
- Laxon, S., Giles, K. A., Ridout, A. L., Wingham, D. J., Willatt, R., Cullen, R., Kwok, R., Schweiger, A., Zhang, J., Haas, C., Hendricks, S., Krishfield, R., Kurtz, N., Farrell, S., and Davidson, M.: CryoSat-2 estimates of Arctic sea ice thickness and volume, *Geophys. Res. Lett.*, 40, 1–6, doi:10.1002/GRL.50193, 1994. 21
- Lipscomb, W. H., Hunke, E. C., Maslowski, W., and Jakacki, J.: Ridging, strength, and stability in high-resolution sea ice models, *J. Geophys. Res.*, 112, C03S91, doi:10.1029/2005JC003355, 2007. 7

Arctic and N-Atlantic high-resolution ocean and sea-ice model

F. Dupont et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Arctic and N-Atlantic high-resolution ocean and sea-ice model

F. Dupont et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Lique, C., Garric, G., Treguier, A.-M., Barnier, B., Ferry, N., Testut, C.-E., and Girard-Ardhuin, F.: Evolution of the Arctic Ocean salinity, 2007–2008: contrast between the Canadian and the Eurasian basins, *J. Climate*, 24, 1705–1717, doi:10.1175/2010JCLI3762.1, 2011. 11
- Lu, Y., Higginson, S., Nudds, S., and Prinsenberg, S.: Model simulated volume fluxes through the Canadian Arctic Archipelago and Davis Strait: linking monthly variations to forcing in different seasons, *J. Geophys. Res.-Oceans*, 119, 1927–1942, doi:10.1002/2013JC009408, 2014. 6
- Lumpkin, R. and Johnson, G. C.: Global ocean surface velocities from drifters: mean, variance, El Niño–Southern Oscillation response, and seasonal cycle, *J. Geophys. Res.*, 118, 1–15, doi:10.1002/jgrc.20210, 2013. 15
- Madec, G. and NEMO team: NEMO Ocean Engine, Tech. Rep., Note du Pole de Modelisation, 27, Institut Pierre-Simon Laplace, 11, 367 pp., 2008. 5
- Madec, G., Delecluse, P., Imbard, M., and Lévy, C.: OPA 8.1, Ocean General Circulation Model Reference Manual, Tech. Rep., Note du Pole de Modelisation, Institut Pierre-Simon Laplace, 11, 91 pp., 1998. 6
- Masson-Delmotte, V., Kageyama, M., Braconnot, P., Charbit, S., Krinner, G., Ritz, C., Guilyardi, E., Jouzel, J., Abe-Ouchi, A., Crucifix, M., Gladstone, R. M., Hewitt, C. D., Kitoh, A., LeGrande, A. N., Marti, O., Merkel, U., and Motoi, T., Ohgaito, R., Otto-Bliesner, B., Peltier, W. R., Ross, I., Valdes, P. J., Vettoretti, G., Weber, S. L., Wolk, F., and Yu, Y.: Past and future polar amplification of climate change: climate model intercomparisons and ice-core constraints, *Clim. Dynam.*, 26, 513–529, doi:10.1007/s00382-005-0081-9, 2006. 6
- Maykut, G. A. and McPhee, M. G.: Solar heating of the Arctic mixed layer, *J. Geophys. Res.*, 100, 24691–24703, 1995. 8
- McPhee, M. G.: Ice–ocean momentum transfer for the AIDJEX ice model, *AIDJEX Bull.*, 29, 93–111, 1975. 8
- Megann, A., Storkey, D., Aksenov, Y., Alderson, S., Calvert, D., Graham, T., Hyder, P., Sidorn, J., and Sinha, B.: GO5.0: the joint NERC–Met Office NEMO global ocean model for use in coupled and forced applications, *Geosci. Model Dev.*, 7, 1069–1092, doi:10.5194/gmd-7-1069-2014, 2014. 4
- Morison, J., Aagaard, K., Falkner, K. K., Hatakeyama, K., Moritz, R., Overland, J. E., Perovich, D., Shimada, K., Steele, M., Takizawa, T., and Woodgate, R.: North Pole environmental observatory delivers early results, *Eos. T. Am. Geophys. Un.*, 83, 357–361, doi:10.1029/2002EO000259, 2002. 20

- Pickart, R. S., McKee, T. K., Torres, D. J., and Harrington, S.: Mean structure and interannual variability of the Slope Water System south of Newfoundland, *J. Phys. Oceanogr.*, 29, 2541–2558, 1999. 15
- Polashenski, C., Perovich, D., Richter-Menge, J., and Elder, B.: Seasonal ice mass-balance buoys: adapting tools to the changing Arctic, *Ann. Glaciol.*, 52, 18–26, 2011. 20
- Proshutinsky, A., Krishfield, R., Timmermans, M.-L., Toole, J., Carmack, E., McLaughlin, F., Williams, W. J., Zimmermann, S., Itoh, M., and Shimada, K.: Beaufort Gyre fresh-water reservoir: state and variability from observations, *J. Geophys. Res.*, 114, C00A10, doi:10.1029/2008JC005104, 2009. 17, 42
- Rattan, S., Myers, P. G., Treguier, A.-M., Theetten, S., Biastoch, A., and Böning, C.: Towards an understanding of Labrador Sea salinity drift in eddy-permitting simulations, *Ocean Model.*, 35, 77–88, doi:10.1016/j.ocemod.2010.06.007, 2010. 6
- Rio, M. H., Guinehut, S., and Larnicol, G.: New CNES-CLS09 global mean dynamic topography computed from the combination of GRACE data, altimetry, and in situ measurements, *J. Geophys. Res.*, 116, C07018, doi:10.1029/2010JC006505, 2011. 13
- Rothrock, D. A.: The energetics of the plastic deformation of pack ice by ridging, *J. Geophys. Res.*, 33, 4514–4519, 1975. 9
- Roy, F., Chevallier, M., Smith, G., Dupont, F., Garric, G., and Lemieux, J.-F.: Arctic sensitivity to the atmosphere-ice-ocean boundary layer in global simulations, *J. Geophys. Res.*, submitted, 2014. 8, 13, 21, 22, 24
- Schauer, U., Fahrbach, E., Osterhus, S., and Rohardt, G.: Arctic warming through the Fram Strait: oceanic heat transport from 3 years of measurements, *J. Geophys. Res.*, 109, C06026, doi:10.1029/2003JC001823, 2004. 18
- Schauer, U., Beszczynska-Möller, A., Walczowski, W., Fahrbach, E., Piechura, J., and Hansen, E.: Variation of measured heat flow through the Fram Strait between 1997 and 2006, in: *Arctic-Subarctic Ocean Fluxes*, edited by: Dickson, R., Meincke, J., and Rhines, P., Springer, the Netherlands, 65–85, doi:10.1007/978-1-4020-6774-7_4, 2008. 16, 18, 19
- Smith, G. C., Roy, F., and Brasnett, B.: Evaluation of an operational ice-ocean analysis and forecasting system for the Gulf of St Lawrence, *Q. J. Roy. Meteor. Soc.*, 139, 419–433, doi:10.1002/qj.1982, 2012. 3

Arctic and N-Atlantic high-resolution ocean and sea-ice model

F. Dupont et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Arctic and N-Atlantic high-resolution ocean and sea-ice model

F. Dupont et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Smith, G. C., Roy, F., Mann, P., Dupont, F., Brasnett, B., Lemieux, J.-F., Laroche, S., and Belair, S.: a new atmospheric dataset for forcing ice–ocean models: evaluation of reforecasts using the Canadian global deterministic prediction system, *Q. J. Roy. Meteor. Soc.*, 140, 881–894, doi:10.1002/qj.2194, 2014a. 9
- 5 Smith, G. C., Roy, F., Reszka, M., Colan, D. S., He, Z., Deacu, D., Belanger, J.-M., Skachko, S., Liu, Y., Dupont, F., Lemieux, J.-F., Beaudoin, C., Tranchant, B., Drevillon, M., Garric, G., Testut, C.-E., Lellouche, J.-M., Pellerin, P., Ritchie, H., Lu, Y., and Davidson, F.: Sea ice forecast verification in the Canadian global ice ocean prediction system, *Q. J. Roy. Meteor. Soc.*, under review, 2014b. 3, 9
- 10 Steele, M., Morison, J., Ermold, W., Rigor, I., Ortmeyer, M., and Shimada, K.: Circulation of summer Pacific halocline water in the Arctic Ocean, *J. Geophys. Res.*, 109, C02027, doi:10.1029/2003JC002009, 2004. 17
- Thompson, K. and Demirov, E.: Skewness of sea level variability of the world's oceans, *J. Geophys. Res.*, 111, C05005, doi:10.1029/2004JC002839, 2006. 14
- 15 Tsamados, M., Feltham, D. L., and Wilchinsky, A. V.: Impact of a new anisotropic rheology on simulations of Arctic sea ice, *J. Geophys. Res.*, 118, 91–107, doi:10.1029/2012JC007990, 2013. 25
- Tsamados, M., Feltham, D. L., Schroeder, D., Flocco, D., Farrell, S. L., Kurtz, N., Laxon, S. W., and Bacon, S.: Impact of variable atmospheric and oceanic form drag on simulations of Arctic sea ice, *J. Phys. Oceanogr.*, 44, 1329–1353, doi:10.1175/JPO-D-13-0215.1, 2014. 25
- 20 Turner, A. K. and Hunke, E. C.: Impacts of a mushy-layer thermodynamic approach in global sea-ice simulations using the cice sea-ice model, *J. Geophys. Res.*, submitted, 2014. 25
- Umlauf, L. and Burchard, H.: A generic length-scale equation for geophysical turbulence models, *J. Mar. Res.*, 61, 235–265, 2003. 7
- 25 Vancoppenolle, M., Fichefet, T., and Goosse, H.: Simulating the mass balance and salinity of Arctic and Antarctic sea ice, 2. Importance of sea ice salinity variations, *Ocean Model.*, 27, 54–69, 2009a. 7
- Vancoppenolle, M., Fichefet, T., Goosse, H., Bouillon, S., Madec, G., and Morales-Maqueda, M.: Simulating the mass balance and salinity of Arctic and Antarctic sea ice, 1. Model description and validation, *Ocean Model.*, 27, 33–53, doi:10.1016/j.ocemod.2008.10.005, 2009b. 7
- 30 Wang, Z., Lu, Y., Wright, D., and Dupont, F.: Sea ice sensitivity to the parameterisation of open water area, *J. Oper. Ocean.*, 3, 3–9, 2010. 6

- Zhang, J. and Rothrock, D. A.: Modeling global sea ice with a thickness and enthalpy distribution model in generalized curvilinear coordinates, Mon. Weather Rev., 131, 845–861, 2003. 21
- Zhu, J., Demirov, E., Dupont, F., and Wright, D.: Eddy-permitting simulations of the sub-polar North Atlantic: impact of the model bias on water mass properties and circulation, Ocean Dynam., 60, 1177–1192, 2009. 6

GMDD

8, 1–52, 2015

Arctic and N-Atlantic high-resolution ocean and sea-ice model

F. Dupont et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Arctic and N-Atlantic high-resolution ocean and sea-ice model

F. Dupont et al.

Table 1. Summary description of the different hindcasts produced to date. Dates are given in YYYYMMDD format.

Experiment name	Initial and boundary conditions	Lateral dyn. BC	Ice model	Turbulence scheme	Notes
ORCA12-T321	Levitus	free-slip	LIM2	TKE	Started in 19990101. Air-ice drag of 1.5×10^{-3} ; ice-ocean drag of 1.0×10^{-2}
H01	GLORYS2v1	no-slip	LIM2	TKE	Started in 20020101. Air-ice drag of 1.63×10^{-3} ; same ice-ocean drag
H02	ORCA12-T321	free-slip	LIM2	TKE	Started in 20030101
H03	ORCA12-T321	free-slip	CICE	$k - \epsilon$	Started in 20030101, reduced Bering flow to mean 0.8 Sv. Top ice roughness of 5.0×10^{-4} m; icea-ocean drag of 5.36×10^{-3} .
H04	ORCA12-T321	free-slip	CICE	$k - \epsilon$	Increased ice-ocean drag relative to H03 to 2.32×10^{-2} corresponding to a bottom ice roughness of 3.0×10^{-2} m
H05	ORCA12-T321	free-slip	CICE	$k - \epsilon$	H04 with ice surface roughness as in CGRF (1.0×10^{-4} m)

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**Arctic and N-Atlantic
high-resolution
ocean and sea-ice
model**

F. Dupont et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion


**Resolution (km)**

Figure 1. CREG12 domain and horizontal resolution (in km). The 3000 m contour of the modelled bathymetry is overlaid.

**Arctic and N-Atlantic
high-resolution
ocean and sea-ice
model**

F. Dupont et al.




Figure 2. First Rossby radius of deformation (left, in km) and Rossby radius relative to the local resolution in log 2 (right). Grossly speaking, the right panel shows where model is eddy-resolving (values above 1, that is 2 model points to resolve a baroclinic eddy), eddy-permitting (between 0 and 1), or does not resolved eddies (values below 0).

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Arctic and N-Atlantic high-resolution ocean and sea-ice model

F. Dupont et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion




Figure 3. The mean (top), standard deviation (middle) and skewness (bottom) of sea surface height (in m) in the North Atlantic from the model hindcast H05 (left) and from satellite altimeter measurements (right) for the period 2003–2009.

**Arctic and N-Atlantic
high-resolution
ocean and sea-ice
model**

F. Dupont et al.




Figure 4. (Left) Modeled sea surface height (in m) in the Arctic for the period 2003–2009 from hindcast H05. (Right) The estimated Arctic mean dynamic topography for the period 2003–2009, reproduced from Farrell et al. (2012).

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)




Figure 5. Mean current speed at 15 m depth, (top) from a drifter climatology, and (bottom) from hindcast H05 for the period 2003–2009.




Figure 6. The mean model bias for temperature (left) and salinity (right), calculated as the model hindcast H05 estimate minus the observed value, averaged in 1° bins for the top 200 m (left) and the 200–500 m layer (right) for the period 2003–2009.




Figure 7. Average temperature and salinity profiles within the boxes shown in the top panel. The mean observed values are compared with profiles from hindcast H05, the Glorys2v3 ocean reanalysis, and the Polar Science Center Hydrographic Climatology (PHC), calculated in each case by averaging profiles at the same times and locations as the observations. The boxes represent **(a)** the Beaufort Sea, 73–78° N, 152–132° W, **(b)** the Greenland and Norwegian seas, 70–75° N, 10° W–10° E, **(c)** the subpolar gyre, 50–55° N, 50–40° W, and **(d)** the subtropical gyre, 35–40° N, 49–41° W. Note the different scaling on the horizontal axis for each panel.

**Arctic and N-Atlantic
high-resolution
ocean and sea-ice
model**

F. Dupont et al.




Figure 8. Mean liquid freshwater content (in m) from the PHC climatology (left) and from hind-cast H05 (right) for the period 2003–2009.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Arctic and N-Atlantic high-resolution ocean and sea-ice model

F. Dupont et al.




Figure 9. Time series of the estimated liquid freshwater content, averaged over the Beaufort Gyre, from Proshutinsky et al. (2009) and updates (black with uncertainties overlaid as grey area) compared with estimates from the ORCA12-T321 run from Mercator Océan (blue), and CREG12 hindcasts H02 (red) and H05 (magenta).

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Arctic and N-Atlantic high-resolution ocean and sea-ice model

F. Dupont et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion




Figure 10. The mean modeled (hindcast H05, top), observed (middle) and difference (modeled minus observed, bottom) temperature (left), salinity (middle) and northward velocity (right) in Fram Strait.




Figure 11. As for Fig. 10, but for Davis Strait.




Figure 12. Monthly time series of total ice area in the Arctic obtained from satellite observations (black, described as SMMR+SSM/I), the ORCA12 T321 run from Mercator Océan (blue), and CREG12 hindcasts H02 (green) and H05 (red). The top panel shows all months, the bottom panel retains only September from each year.

Arctic and N-Atlantic high-resolution ocean and sea-ice model

F. Dupont et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion




Figure 13. Ice concentration for September 2007 from NSIDC, the ORCA12 T321 run from Mercator Océan and CREG12 hindcasts H02 and H05.




Figure 15. Average ice thickness distributions from ULS measurements (blue) and hindcast H05 (green) at the Beaufort Gyre Exploration Project mooring A (top) and at the North Pole Environmental Observatory mooring (bottom).




Figure 16. The mean ice thickness (in m) for October-November 2007 from ICESat, and the difference between ORCA12 T321 and CREG12 hindcasts H02 and H05 and the ICESat estimate.




Figure 17. Monthly time series of total ice volume in the Arctic obtained from PIOMAS (black), the ORCA12 T321 run from Mercator Océan (blue), and CREG12 hindcasts H02 (green), and H05 (red). The top panel shows all months, the bottom panel retains only September from each year.




Figure 18. Average ice velocity (in m s^{-1}) for March 2003-2008 from NSIDC, the ORCA12 T321 run from Mercator Océan, and CREG12 hindcasts H02 and H05.

Arctic and N-Atlantic high-resolution ocean and sea-ice model

F. Dupont et al.




Figure 19. Monthly timeseries of average bias in monthly ice speed (in ms^{-1}) relative to IABP buoys for NSIDC (black dashed), the ORCA12 T321 run (blue), and CREG12 hindcasts H02 (green) and H05 (red).

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[◀](#)
[▶](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)
