

Coupling aerosol optics to the chemical transport model MATCH (v5.5.0) and aerosol microphysics module SALSA (v1)

E. Andersson¹ and M. Kahnert^{1,2}

¹Department of Earth and Space Sciences, Chalmers University of Technology, 41296 Gothenburg, Sweden

²Swedish Meteorological and Hydrological Institute, 60176 Norrköping, Sweden

Correspondence to: M. Kahnert (michael.kahnert@smhi.se)

Abstract. Modelling aerosol optical properties is a notoriously difficult task due to the particles' complex morphologies and compositions. Yet aerosol particles and their optical properties are important for chemistry-climate modelling and remote sensing applications. Operational optics models often make drastic and non-realistic approximations regarding morphological properties, which can

5 introduce errors. In this study a new aerosol-optics model is implemented, in which more realistic morphologies and mixing states are assumed, especially for black carbon particles. The model includes both external and internal mixing of all chemical species, it treats externally-mixed black carbon as fractal aggregates, and it accounts for inhomogeneous internal mixing of black carbon by use of a novel “core-grey-shell” model. Simulated results of aerosol optical properties, such as aerosol

10 optical depth, backscattering coefficients and Ångström exponent, as well as radiative fluxes computed with the new optics model are compared with results from an older optics-model version that treats all particles as externally mixed homogeneous spheres. For comparison, we perform computations with two different model-versions, one that accounts for aerosol-microphysical processes, and another one that entirely neglects these processes. Since it is well understood that aerosol microphysics

15 has a profound impact on aerosol mass- and number-concentrations, their size-distribution, and their size-dependent chemical composition (which, in turn, strongly impact their optical properties), these additional model-runs can serve as a reference against which we can gauge the significance of the morphological assumptions in the optics model. The results show that using a more detailed description of particle morphology and mixing states influences the optical properties to a degree that is

20 on the same order of magnitude as the corresponding effects of aerosol-microphysical processes. For instance, the aerosol optical depth computed with the two optics models differs over the optical spectrum by -25 – 18 %, while corresponding differences caused by the inclusion or omission of

aerosol microphysics range between -50 – 37 %. The corresponding differences in the backscattering coefficient are -8 – 99 % and -47 – 28 %, respectively. This is an important finding suggesting that simple optics models coupled to a chemical transport model can introduce considerable errors; this can strongly affect simulations of radiative fluxes in chemistry-climate models, and it can compromise the use of remote sensing observations of aerosol particles in model evaluations and chemical data assimilation.

1 Introduction

Aerosol-optics models are employed in large-scale chemical transport models (CTMs) in mainly two contexts, namely, in chemistry-climate modelling (CCM), and in conjunction with remote sensing observations. In a CCM one couples a CTM to an atmosphere-ocean general circulation model (GCM). One purpose is to account for the dynamic effects of aerosol particles on cloud microphysics. Another is to obtain a better description of the direct effect of aerosol particles and radiatively active trace gases on the radiative balance. The aerosol-optics model provides a link that converts the aerosol fields delivered by the CTM to the aerosol optical properties that are required as input to the radiative transfer model, with which one computes the radiative energy budget. In remote sensing applications one is faced with the obstacle that the aerosol concentration fields computed with a CTM are not directly comparable to the radiometric quantities that are observed with remote sensing instruments. The aerosol-optics model provides the observation operator that maps the CTM output to radiometric variables that can be compared to satellite observations or satellite retrieval products. This allows us to either employ satellite observations for evaluating CTM model results, or to assimilate satellite data into a CTM-based air-quality forecasting system. It is clear that the aerosol-optics model has a pivotal role in these kinds of applications. It may constitute an additional source of error that could compromise the reliability of CCMs, impair the reliability of CTM evaluations, or degrade chemical data assimilation results. It is, therefore, important to better understand this potential source of error, quantify its possible impact on model predictions of aerosol radiometric quantities, and assess the level of morphological detail that might be required in aerosol-optics models coupled to CTMs.

A main difficulty is that aerosol particles in nature can have a high degree of morphological complexity. For instance, mineral dust particles can have irregular shape, small-scale surface roughness, and inhomogeneous mineralogical composition (e.g. Nousiainen, 2009). Black carbon particles suspended in air have fractal-aggregate shapes (e.g. Jones, 2006) that can be coated by weakly absorbing liquid-phase components that condense onto the aggregates as they age in the atmosphere (e.g. Adachi and Buseck, 2008). Volcanic ash particles are composed of crustal material in which multiple air vesicles may have been trapped during the generation of the particles. In aerosol-optics models

one has to make a choice what level of morphological detail is necessary and affordable. A detailed discussion of this question can be found in Kahnert et al. (2014).

In environmental modelling practical and computational constraints often force us to invoke drastically simplifying assumptions about aerosol morphology. For instance, one frequently computes aerosol optical properties based on the assumption that all chemical aerosol components are contained in separate particles (externally mixed), and that each such particle can be approximated as a homogeneous sphere. As pointed out in Kahnert (2008); Benedetti et al. (2009), this approach is highly attractive from a practical point of view, because the aerosol optical observation operators, which map mixing ratios to radiometric properties, become linear functions of the mixing ratios of the different chemical species. A linearisation of the observation operator is a prerequisite for most of the commonly used data-assimilation methodologies, such as the variational method (e.g. Kahnert, 2008; Benedetti et al., 2009). However, such approximations can also introduce substantial errors. In the remote-sensing community awareness for this problem has been growing over the past 1–2 decades. As a result, one has developed retrieval methods for desert dust aerosol particles that are based on spheroidal model particles (e.g. Dubovik et al., 2006), which can mimic the optical properties of mineral dust particles better than homogeneous spheres (Kahnert, 2004; Nousiainen et al., 2006). In chemical data assimilation, the problem is still treated rather negligently. A few assimilation studies account for internal mixing (where several aerosol components can be contained within one particle) of different chemical components (e.g. Saide et al., 2013). But the particles are still assumed to be perfectly homogeneous spheres. To the best of our knowledge there are currently no aerosol optical observation operators in chemical transport models that take complex morphological properties of aerosol particles such as non-sphericity or inhomogeneous internal structure into account.

This study describes the coupling of two different aerosol-optics models to a regional CTM. One optics model is based on the simple external-mixture and homogeneous-sphere approximations. The second model takes both external and internal mixing of aerosol components into account. Also, it employs morphologically more realistic models for black carbon particles. Although black carbon contributes, on average, only some 5% to the mass mixing ratio of particulate matter over Europe, it can have a significant global radiative warming effect. Previous theoretical studies on the optical properties of black-carbon particles suggest that the use of homogeneous sphere models can introduce substantial errors in the absorption cross section and single scattering albedo of such particles (e.g. Kahnert, 2010a; Kahnert et al., 2013). Also, the largest mixing-state sensitivity in both regional and global radiative fluxes comes from black carbon according to Klingmüller et al. (2014).

The main goal of this study is to assess the impact of aerosol morphology and mixing state on radiometric quantities and radiative forcing simulated with a chemical transport model. To this end we compare the two optics models, and we gauge the significance of morphology by comparing the differences in the optics model output to other sources of error. As a gauge we use the impact of

including or omitting aerosol microphysical processes; this provides us with a reference which is
95 generally agreed to have a significant effect on aerosol transport models (Andersson et al., 2015;
Kokkola et al., 2008).

The CTM, its aerosol microphysic and mass transport set-ups, and the aerosol-optics models are
described in Sect. 2. There we also explain the methodology we employ for comparison of the optics
models. In Sect. 3 we present and discuss computational results for selected cases and for several
100 radiative and optical parameters. Concluding remarks are given in Sect. 4.

2 Model description and methods

2.1 General considerations and terminology

Aerosol particles typically originate from different emission sources, such as sea-salt particles com-
ing from marine sources, wind-blown dust from dry land surfaces, volcanic ash from magmatic
105 or phreatomagmatic eruptions, or black carbon produced during combustion of fossil fuel, biofuel,
or biomass. During atmospheric transport particles from different sources can be mixed, resulting
in heterogeneous aerosol populations consisting of particles of different morphologies, sizes, and
chemical composition. A mixture in which different chemical species are contained in separate par-
ticles is referred to as an *external mixture*. On the other hand, aerosol dynamic processes, such as
110 nucleation, condensation, and coagulation, give rise to the formation and growth of secondary par-
ticles from precursor gases, as well as to the condensation of precursor gases onto existing primary
particles. These processes result in particles in which several chemical species are mixed with each
other in one and the same particle. Such a population is referred to as an *internal mixture*. There are
two types of internal mixtures. If, e.g., hydrophilic liquid-phase components mix with each other, one
115 can obtain a *homogeneous internal mixture* of different chemical species. On the other hand, con-
densation of gas-phase species onto non-soluble primary particles, or cloud processing of aerosol
particles can result in liquid-phase material coating a solid core of, e.g., mineral dust or black car-
bon. We refer to the latter as an *inhomogeneous internal mixture*. Aerosol populations in nature are
often both externally and internally mixed, i.e., they contain particles that are composed of a single
120 chemical species as well as other particles that are composed of different chemical species, which
can be homogeneously or inhomogeneously internally mixed.

Aerosol optical properties are strongly dependent on not only the size and chemical composition,
but also on the mixing state, shape, and internal structure of particles. Therefore, before explaining
the aerosol-optics model, we first need to briefly describe the kind of information that can be pro-
125 vided by the aerosol transport model. In particular, we need to understand the level of detail with
which the size distribution, size-dependent chemical composition, and the mixing state of the aerosol
particles can be computed in a large-scale model.

2.2 Aerosol transport modelling with MATCH

As a regional model we employ the Multiple-scale Atmospheric Transport and CHemistry modelling system (MATCH), an offline Eulerian model developed by the Swedish Meteorological and Hydrological Institute (Andersson et al., 2007). For this study we have set up the MATCH model over the European domain with a $0.4 \times 0.4^\circ$ horizontal resolution and a rotated latitude-longitude grid, covering about 34° longitude and 42° latitude. The model has 40 vertical η layers with varying thickness depending on the topography, and it extends up to about 13 hPa. The meteorological input comes from the numerical weather-prediction model HIRLAM (High-Resolution Limited Area Model) (Unden et al., 2002).

The MATCH model allows us to choose between two aerosol model versions, a simpler mass-transport model, and a more sophisticated aerosol dynamic transport model.

2.2.1 Mass transport model

A simple version of the CTM MATCH, which we refer to as the “mass-transport model”, neglects all aerosol dynamic processes. It contains a photochemistry model that computes mass concentrations of secondary inorganic aerosols (SIA), which are formed from precursor gases. The SIA fraction of aerosol particles consists of ammonium sulphate ($(\text{NH}_4)_2\text{SO}_4$), ammonium nitrate (NH_4NO_3), other particulate sulphates (PSO_x), and other particulate nitrates (PNO_x). The mass transport model further contains a sea-salt module that computes NaCl emissions based on the parametrisations described in Mårtensson et al. (2003); Monahan et al. (1986). More details on the MATCH photochemistry model can be found in Robertson et al. (1999); Andersson et al. (2007); the MATCH sea-salt model is described in Foltescu et al. (2005). The mass transport model also contains a simple wind-blown dust model and a module for transport of primary particulate matter (PPM), i.e., aerosol particles other than sea-salt and wind-blown dust that are emitted as particles, rather than being formed from gas precursors. The size bins in the PPM model are flexible. In the current model set-up the sea-salt and PPM models were run for four size bins as shown in Table 1. We used gridded EMEP PPM emission data for the year 2007 in conjunction with black-carbon (BC) and organic-carbon (OC) emission data by Kupiainen and Klimont (2004, 2007). The latter provide BC and OC emissions per country and emission sector. We distributed these among the grid cells in the model domain according to the EMEP PPM gridded emissions. Thus, the BC and OC emissions vary among grid cells in accordance with the EMEP PPM emissions, while the sum of all BC and OC emissions over all grid cells per country and emission sector agrees with the corresponding BC and OC emissions, respectively, reported in Kupiainen and Klimont (2004, 2007). The remaining emissions (PPM-BC-OC) in each grid cell are interpreted as dust particles. The primary-particle emissions are distributed among the size bins; during atmospheric transport they remain chemically and dynamically inert in the model. Thus no chemical transformation, mixing processes with other compounds, or other size-

Table 1. Size bins (characterised by the radius r) and chemical species in the MATCH mass transport model (Andersson et al., 2007). The labels “p” and “s” refer to primary emitted particles and secondary particles generated from gas precursors.

size bin	r (nm)	OC	BC	Dust	other				other	other
					PPM	NaCl	(NH ₄) ₂ SO ₄	NH ₄ NO ₃	PSO _x	PNO _x
1	10–50	p	p		p	p	s	s	s	s
2	50–500	p	p		p	p	s	s	s	s
3	500–1250					p	s	s	s	s
4	1250–5000	p	p	p	p	p	s	s	s	s

transformation processes are included in the model. The SIA components are given as total mass concentrations without any information about their size distribution. In the optics model a fixed
165 size distribution is assumed to assign the total SIA mass to the four size bins. Water adsorption by particles is computed in the optics model as described in Sect. 2.3.1.

2.2.2 Aerosol microphysics module - SALSA

A more realistic description of particles can be achieved by accounting for aerosol microphysical processes. To this end the Sectional Aerosol module for Large Scale Applications (SALSA)
170 (Kokkola et al., 2008) has recently been coupled to the MATCH photochemistry model (Andersson et al., 2015). This model tracks mass concentrations of different species per size bin, and particle number concentrations. Thus, it provides size-dependent composition and mixing state of aerosol particles. The description of PNO_x, wind-blown dust, and secondary organic aerosols (SOA) is still
175 in an early development stage in MATCH-SALSA. In the current version, PNO_x is simply computed according to the same photochemistry-scheme as in the mass-transport model, and the PNO_x mass is assigned to size bin 15 (see Tab. 2). Wind-blown dust and SOA are absent in the present model version. The size distributions for the emitted particles can be found in table 4 and figure 6 of Andersson et al. (2013).

Table 2 shows the current model set-up with the number and range of the size bins. As is evident
180 from this table, MATCH-SALSA accounts for both internally and externally mixed aerosol particles. In total, there are 20 different size bins in MATCH-SALSA, each one of them representing a particle size range (volume-equivalent radius, r), mixing state, and composition. Some size bins have the same size range, but different mixing states and/or compositions. For instance, size bins 12, 15, and 18 describe the same size range (350–873 nm), but different internal mixtures of various species.
185 Similarly, bins 4 and 8 have the same size range (25–49 nm), but one describes an internal mixture, the other an external mixture of aerosol species.

As in the mass-transport model, "other PPM", i.e. primary particles other than BC and OC, are interpreted as dust particles. Note that water is not directly calculated as a prognostic variable in

Table 2. Size bins and chemical species in the MATCH-SALSA aerosol microphysical transport model. An “x” marks that the species is present in a particular size bin.

size bin	r (nm)	mixing state	other							
			OC	BC	PPM	NaCl	PSO _x	PNO _x	PNH _x	
1	1.5–3.8	internal	x					x		x
2	3.8–9.8	internal	x					x		x
3	9.8–25	internal	x					x		x
4	25–49	internal+H ₂ O	x	x	x	x	x	x		x
5	49–96	internal+H ₂ O	x	x	x	x	x	x		x
6	96–187	internal+H ₂ O	x	x	x	x	x	x		x
7	187–350	internal+H ₂ O	x	x	x	x	x	x		x
8	25–49	external	x	x				x		x
9	49–96	external	x	x				x		x
10	96–187	external	x	x				x		x
11	187–350	external	x	x	x			x		x
12	350–873	NaCl+H ₂ O					x			
13	873–2090	NaCl+H ₂ O					x			
14	2090–5000	NaCl+H ₂ O					x			
15	350–873	internal+H ₂ O	x	x	x			x	x	x
16	873–2090	internal+H ₂ O	x		x			x		x
17	2090–5000	internal+H ₂ O	x		x			x		x
18	350–873	internal+H ₂ O			x			x		x
19	873–2090	internal+H ₂ O			x			x		x
20	2090–5000	internal+H ₂ O			x			x		x

MATCH-SALSA. Rather, it is a diagnostic variable computed in the MATCH-optics model as explained in Sect. 2.3.2. The table merely indicates which size bins are assumed in the optics model to be internally mixed with adsorbed water. A more detailed description of the MATCH-SALSA model can be found in Andersson et al. (2015).

2.3 Aerosol-optics modelling

Aerosol-optics models coupled to a CTM have to make consistent use of the information provided by the CTM, while invoking assumptions on optically relevant parameters that are not provided by the CTM. The parameters that influence the particles’ optical properties are

- the aerosol size distribution;
- the refractive index of the materials of which the aerosol particles are composed;
- the morphology of the particles.

200 *Morphology* refers to both the overall shape of the particle, and, in case of inhomogeneously mixed particles, the variation of the refractive index inside the particle.

The information provided by the CTM depends on the level of detail in the process descriptions. In the MATCH mass transport model, we have size information for the primary particles, but only the total mass for secondary inorganic aerosols. Thus we have to invoke assumptions about the size
205 distribution of these particles. The MATCH optics models in conjunction with the MATCH mass transport model assume that 10 % of the SIA aerosol mass are in the smallest size bin (see Table 1), 60 % in the second, 20 % in the third, and 10 % in the fourth size bin. Also, the mass transport model lacks any information about the mixing state of the particles. We therefore have to invoke appropriate assumptions on whether the aerosol particles are externally or internally mixed. Both
210 the mass transport model and MATCH-SALSA lack information on whether the internally mixed particles are homogeneous or inhomogeneous. Also, neither model provides any information on the shape of the particles. The refractive indices of each chemical component in the aerosol phase and their spectral variation are listed in Appendix D. They can also be found in Fig. 4 in Kahnert (2010a). That reference also contains detailed information about the different literature sources from which
215 the refractive indices are taken.

2.3.1 Optics model for externally mixed aerosol particles

The simplest conceivable optics model assumes that all particles are homogeneous spheres, and that all chemical species are each in separate particles, i.e., externally mixed. As explained in (Kahnert, 2008), the external-mixture assumption results in a linear relation between the mass mixing ratios
220 and the optical properties. Owing to the linearity, this model is particularly attractive for data assimilation applications (e.g. Benedetti et al., 2009), which require linearised observation operators. However, this is also the crudest possible optics model, as it neglects both the effect of internal mixing and of particle morphology on optical properties.

The external-mixture model is implemented in the MATCH mass transport model, where it is
225 primarily being used in the MATCH 3DVAR data assimilation system (Kahnert, 2008). Optical properties are pre-computed for twelve wavelength bands ranging from the UV-C to the mid-IR. Dust and black carbon are assumed to be hydrophobic, while sea salt, OC, and SIA components can each mix internally with water. The water volume fraction depends on temperature and humidity; it is computed by use of the parametrisation given in Gerber (1985), which computes the particle's
230 wet-radius as a function of dry-radius, relative humidity, and temperature. The aerosol/water mixture is assumed to be homogeneous. The dielectric properties of a homogeneous mixture of two or more components are described by a complex effective refractive index m_{eff} , which is usually computed by effective medium theory (EMT) (although chemical transport modellers often use simple volume mixing rules, most likely because EMTs are not commonly known in that field). We
235 use Bruggemann's EMT (Bruggemann, 1935). More information of EMT is given in Appendix C.

Optical properties are pre-computed for eleven water volume fractions between 0 and 0.98; for intermediate volume fractions the optical properties are linearly interpolated. The optical properties contained in the database are the extinction cross section C_{ext} , the scattering cross section C_{sca} , the value of the phase function in the exact backscattering direction $p(180^\circ)$, and the asymmetry parameter g .

240

As explained in Kahnert (2008), size-averaged optical properties are pre-computed by averaging over a log-normal size distribution $n_i(r) = N_i^0 / (\sqrt{2\pi} r \ln \sigma_i) \exp[-\ln^2(r/R_i)/(2\ln^2 \sigma_i)]$ for each size bin i , where N_i^0 relates to the number density of particles in size bin i , r denotes the particle radius, $R_1 = 0.022 \mu\text{m}$, $R_2 = 0.158 \mu\text{m}$, $R_3 = 0.791 \mu\text{m}$, $R_4 = 2.5 \mu\text{m}$ are the geometric mean radii in each size mode, and the geometric standard deviation $\sigma_1 = \sigma_3 = \sigma_4 = 1.8$, $\sigma_2 = 1.5$ are based on measurements in Neusüß et al. (2002). Appendix A provides detailed explanations of how to convert mass mixing ratios computed in the model into particle number densities, how these are used in computing size-averaged optical properties, and how to obtain radiometric properties of the atmosphere, such as aerosol optical depth (AOD) or backscattering coefficient β_{bak} , from the particles' optical properties and from the MATCH aerosol fields.

245

250

2.3.2 Optics model for aerosol particles of different mixing states

The new MATCH-optics model accounts for both internally and externally mixed aerosol particles, and it contains both homogeneously and inhomogeneously mixed aerosol particles. Different shapes and morphologies are assumed for different types of particles.

255

1. Pure, externally mixed black carbon particles are assumed to have a fractal aggregate morphology as shown in Fig. 1. The fractal morphology can be described by the statistical scaling law $N_s = k_0 (R_g/a)^{D_f}$, where N_s denotes the number of spherical monomers in the aggregate, D_f and k_0 are the fractal dimension and fractal prefactor, a is the monomer radius, and $R_g = \sqrt{\sum_{n=1}^{N_s} r_n^2 / N_s}$ is the radius of gyration, where r_n describes the distance of the n th monomer from the aggregate's centre of mass. We use $D_f = 1.8$, $k_0 = 1.3$, which is based on the review in Bond and Bergstrom (2006). Although in the atmosphere black carbon aggregates may also have higher fractal dimensions (e.g. Adachi et al., 2007; Chakrabarty et al., 2014; China et al., 2014), assuming a fractal dimension around 1.8 yields mass-absorption cross sections at 550 nm wavelength that lie closer to experimental data, as was shown in Kahnert and Devasthale (2011). The monomer radius can vary within a range of 10-25 nm (Bond and Bergstrom, 2006). Here it is assumed to be $a = 25$ nm. This is consistent with field observations (Adachi and Buseck, 2008); also, it was shown (Kahnert, 2010b) that this choice of monomer radius in light scattering computations yields results for the single-scattering albedo of black carbon aggregates consistent with observations (Bond and Bergstrom, 2006).

260

265

270

The calculations in Kahnert and Devasthale (2011) were limited to aggregates up to $N_s = 600$. In order to cover the size range of externally mixed black carbon in SALSA we had to ex-




Figure 1. Examples of fractal aggregate model particles for computing optical properties of externally mixed black carbon. The aggregates consist of 1000, 1500, 2000, and 2744 monomers (in clockwise order, starting from upper left).

275 tend these calculations to aggregate sizes up to $N_s = 2744$, which corresponds to a volume-
 equivalent radius of $R_V = 350$ nm (compare with Table 2). We used the multiple-sphere T-
 matrix code (Mackowski and Mishchenko, 2011), which is based on the numerically exact
 superposition T-matrix method for solving Maxwell’s equations. Figure 2 shows some of
 the computed black carbon optical properties as a function of volume-equivalent particle ra-
 280 dius and wavelength. All optical properties are averaged over particle orientations, where the
 orientation-averaging is performed analytically (Khlebstov, 1992). The absorption cross sec-
 tion C_{abs} shows the characteristic decline at long wavelengths, where the refractive index of
 black carbon is changing only slowly (Chang and Charalampopoulos, 1990). Also, C_{abs} in-
 creases with particle size. For small particle sizes this increase goes as $\sim R_V^3$, which is typical
 for the Rayleigh scattering regime (Mishchenko et al., 2002).




Figure 2. Absorption cross section C_{abs} (upper left), single-scattering albedo SSA (upper right), asymmetry parameter g (bottom left), and backscattering cross section C_{bak} (bottom right) of black carbon aggregates as a function of volume-equivalent radius R_V and wavelength λ .

2. Black carbon particles that are internally mixed with other aerosol components are morphologically very complex. It is technically beyond the reach of our present capabilities to build an aerosol-optics database with the use of morphologically realistic model particles. However, it is possible to employ realistic model particles in reference computations for some selected cases. This has recently been done in different studies Kahnert et al. (2013); Scarnato et al. (2013, 2015). In the study by Kahnert et al. (2013), optical properties of encapsulated aggregate model particles, such as the one shown in Fig. 3 (left), were computed in the range of 100–500 nm (volume-equivalent radius), for different black carbon volume fractions, and for wavelengths from the UV-C to the mid-IR. The morphological parameters characterising these model particles were based on field observations (Adachi and Buseck, 2008); the coating material was sulphate. The computations were performed with the discrete dipole approximation (Yurkin and Hoekstra, 2007).

In Kahnert et al. (2013) the computational results were compared to those obtained with simple model particles, such as externally-mixed homogeneous spheres, internally-mixed homogeneous spheres, and concentric core-shell particles with a carbon core and a sulphate shell. The analysis revealed which morphological properties of the encapsulated aggregate particles had the dominant impact on the optical properties. There are two important properties: (1) the amount of carbon mass that interacts with the electromagnetic field has a major impact




Figure 3. Morphologically realistic encapsulated aggregate model for internally mixed black carbon (left), and core-grey-shell model (right).

on the absorption cross section C_{abs} . In a core-shell model as well as in a model based on externally mixed homogeneous spheres, all of the black carbon is concentrated in a single sphere. Owing to absorption of the electromagnetic field does not penetrate deeply into this sphere. Hence much of the carbon mass is shielded from interacting with the field, resulting in an underestimation of C_{abs} compared to the encapsulated aggregates, in which a much larger fraction of the carbon mass can contribute to the absorption of electromagnetic energy. By contrast, in a homogeneous internal-mixture model the black carbon is distributed evenly throughout the sulphate host, which allows too much of the carbon mass to interact with the field. This results in an overestimation of C_{abs} . (2) Compared to a bare-black carbon aggregate, a coated aggregate has a larger geometric cross section. Hence more light is intercepted by an internally-mixed particle and focused onto the black carbon inclusion, thus enhancing C_{abs} . This effect is neglected in the external-mixture model, resulting in an underestimation of C_{abs} . Note that in earlier studies (e.g. Jacobson (2000)) it was often tacitly assumed that a core-shell model would give the most accurate estimates of the aerosol optical properties, owing to its morphological similarity to encapsulated black-carbon particles. However, the results in Kahnert et al. (2013) have clearly shown the shortcomings of the conventional core-shell model. But once we understand which morphological properties are most essential, and which ones make a minor contribution to the optical properties, we can devise model particles that account for the most important morphological effects, yet are sufficiently simple for computing a look-up table for large-scale modelling. It was proposed in Kahnert et al. (2013) to use a core-shell model (hence accounting for the coating effect) in which only part of the carbon mass is contained in the core, and the remaining part is homogeneously mixed with the shell. The model particle is illustrated in Fig. 3 (right). The fraction f_c of the carbon mass located in the core is a free parameter, with which one can interpolate between the two extreme models

Table 3. Core fraction f_c in the core-grey-shell model as a function of wavelength λ .

λ [μm]	0.2000	0.2316	0.3040	0.3400	0.3550	0.3800	0.3932
f_c	0.7	0.7	0.7	0.7	0.7	0.6	0.6
λ [μm]	0.4400	0.5000	0.5320	0.5332	0.6750	0.7016	0.8700
f_c	0.6	0.5	0.5	0.5	0.5	0.5	0.2
λ [μm]	1.0101	1.0200	1.064	1.2705	1.4625	1.7840	2.0460
f_c	0.1	0.1	0.1	0.1	0.1	0.1	0.1
λ [μm]	2.3250	2.7885	3.4615	3.5000	8.0205	10.600	12.195
f_c	0.1	0.1	0.1	0.1	0.1	0.1	0.1

325 of the homogeneous mixture ($f_c = 0$, all carbon mass mixed with the shell) and the regular
core-shell model ($f_c = 1$, all carbon mass in the core). This model has been referred to as the
concentric core-grey-shell (CGS) model. The tuning of the free parameter f_c in the model
was done to fit the reference model of encapsulated aggregates as described in Kahnert et al.
(2013). It was found that f_c is independent of particle size, black-carbon volume fraction, and
330 of the optical property one wants to fit. However, f_c does depend on the wavelength of light.

The CGS model has been employed in generating the new MATCH-optics look-up table. The
shell material can be any mixture of water-soluble components. We use the same values of f_c
as those determined in Kahnert et al. (2013). Its dependence on wavelength is given in Table
3.

335 3. In the mass transport model, we assume that all SIA components and all sea salt is internally
mixed. We further assume that in size bins 1–4, 0, 70, 70, and 100 %, respectively, of the black
carbon, 0, 70, 70, and 70 % of the organic carbon, and 0, 1.3, 1.3, and 1.3 % of the dust are
internally mixed; the remaining BC, OC, and dust mass is externally mixed. In SALSA, the
mixing state depends on the size bin (see Table 2), and the mixing proportions are provided by
340 the model results. In both the mass transport model and in MATCH-SALSA, the contribution
to the effective refractive index of dust and black carbon is computed by the Maxwell–Garnett
EMT (Maxwell Garnett, 1904), while for all other components we use the Bruggemann EMT
(Bruggemann, 1935).

345 4. All other externally mixed particles not containing black carbon are assumed to be homoge-
neous spheres in the present version of the look-up table.

The look-up tables contain results for C_{ext} , C_{sca} , g , and C_{bak} in 28 wavelength bands from the
UV-C to the mid-IR. Computations with the CGS model were performed for 37 discrete BC volume
fractions, namely, $f_{\text{BC}} = 0.00, 0.01, \dots, 0.20, 0.25, \dots, 1.00$. For the shell material, as well as for
non-carbon containing particles, the table contains (depending on the wavelength band) up to 40
350 discrete values of the real part and up to 18 discrete values of the imaginary part of the refractive

index. The range of the refractive indices varies with wavelength; it is determined by those chemical components that, at each given wavelength, have the most extreme values of the refractive index. The optical properties are pre-averaged over particle sizes for each size bin. Thus we generated one look-up table each for the mass transport model with its four size bins, and for SALSA with its 20
355 size bins. In Salsa it is assumed that the number density is constant in each size bin.

The MATCH-optics model computes in each grid cell and for each size bin the effective refractive index of the internally mixed material by use of EMT. The corresponding optical properties are obtained by linearly interpolating the closest pre-computed results in the look-up table. Size-averaging is performed by weighing the optical cross sections as well as $g \cdot C_{\text{sca}}$ in each size bin with the number density per bin and adding over all bins. The integrated quantities are then divided by the total
360 particle number density; the integral over $g \cdot C_{\text{sca}}$ is also divided by the size-averaged scattering cross section.

2.4 Methodology for comparing the optics models

The new internal-mixture optics model with its BC fractal aggregate and core-grey-shell model particles accounts for significant morphological details in aerosol particles. The main question we want
365 to address is whether or not this high level of detail is really necessary, i.e., if it has any *significant* impact on optical properties modelled with a CTM. By *significant* we mean an impact that is comparable to other effects whose importance is well understood. Thus to make such an assessment we need to pick a well-understood effect that can serve as a gauge, i.e., to which we can compare the
370 impact of particle morphology on optical properties. We take the effect of aerosol microphysics as a gauge. As aerosol microphysics is well-known to have a substantial impact on aerosol concentrations and size distributions (Andersson et al., 2015; Matsui et al., 2013), this effect will provide us with a reference to which we can compare the impact of the morphological assumptions made in the aerosol-optics model. Thus we compute aerosol optical properties

- 375 1. with the MATCH mass-transport model (i.e., with aerosol microphysics switched off), in conjunction with the old optics model (abbreviated by MT-EXT, “mass-transport external mixture”);
2. with the MATCH mass-transport model in conjunction with the new optics model (MT-CGS, “mass-transport core-grey-shell”);
- 380 3. with the MATCH-SALSA model (i.e., with aerosol microphysics switched on), in conjunction with the old optics model (abbreviated SALSA-EXT, “MATCH-SALSA external mixture”);
4. with the MATCH-SALSA model, in conjunction with the new optics model (SALSA-CGS, “MATCH-SALSA core-grey-shell”).

We first perform a comparison of monthly and geographically averaged differences in aerosol optical
385 properties. More specifically, comparison of model set-ups MT-EXT with MT-CGS, or SALSA-EXT
with SALSA-CGS allows us to assess the impact of the morphological assumptions in the optics
model. Comparison of MT-EXT with SALSA-EXT, or MT-CGS with SALSA-CGS will give us
an estimate of how much the inclusion or omission of microphysical processes impacts modelling
results of aerosol radiometric properties.

390 While statistical analyses can uncover general trends, it is difficult to understand the underlying
physical reasons for model differences from an analysis of temporally and geographically averaged
model results. Therefore, we also consider a few case studies in some more detail. We take the
optical properties modelled with different MATCH-versions as input to a radiative transfer model
and analyse the total aerosol radiative forcing and the black-carbon radiative forcing. The main
395 goal is to understand how differences in single-scattering optical properties between the two optics
models impact the outcome of the radiative transfer simulations. To keep the case studies within
manageable bounds, we restrict ourselves to four geographic locations (two over land, two over the
ocean), two instances (one representing low-BC summer concentrations, one representing high-BC
winter conditions), and we limit ourselves to comparing the model set-ups MT-EXT, MT-CGS, and
400 SALSA-CGS. More specifically, we consider one site over Northern Italy (45.0° N, 8.5° E), one over
the Mediterranean Sea (37.5° N, 5.5° E), one over Poland (52.6° N, 21.0° E), and one over the North
Sea (52.0° N, 2.7° E). For the two instances, we pick 22 June 2007 12:00 UTC, and 22 December
2007 12:00 UTC. Radiative transfer calculations are performed for each of these four sites and for
both instances. Vertical profiles of the aerosol optical depth per layer, the single-scattering albedo,
405 and the asymmetry parameter are used as input to the libRadtran radiative transfer package (Kylling
et al., 1998), assuming a plane-parallel atmosphere. For the surface albedo of the ocean we assume
a spectrally constant value of 0.065, while for the spectrally varying surface albedo of the two land
locations we used MODIS observations for each of the two instances. The results were spectrally
integrated to obtain the broadband radiative fluxes. The radiative transfer simulations were repeated
410 for corresponding profiles of optical properties (with a 1 km resolution) in the absence of black car-
bon, as well as for clear-sky conditions. This allows us to compute differences in broadband radiative
fluxes, i.e., the radiative effect of black carbon, and the radiative effect of all aerosol components.
The results of this radiative transfer study are discussed in Sect. 3.2.

To further investigate the significance of the optics model on radiometric properties, we also look
415 at optical properties relevant for remote sensing, namely, backscattering coefficient and Ångström
exponent. These results are discussed in Sects. 3.3 and 3.4, respectively.




Figure 4. Aerosol optical depth over Europe on 22 December 2007, 12:00 UTC (noon). Results are shown for the mass transport model in conjunction with the old external-mixture optics model (1st to the left), and with the new internal-mixture/core-grey-shell/fractal BC aggregate model (2nd to the left), as well as for the MATCH-SALSA model in conjunction with the new optics model (3rd to the left) and old optics model (4th to the left). The circles indicate the four locations used for radiative transfer studies. Note the semi-logarithmic colour scale!

3 Results

3.1 Aerosol optical properties in MATCH

Figure 4 gives us a first impression of the differences among the four model configurations. The extinction AOD is shown for MT-EXT (1st from the left), MT-GCS (2nd), Salsa-CGS (3rd) and Salsa-EXT (4th). The overall spatial patterns are similar. This is expected, since all model configurations used the same EMEP emissions and HIRLAM meteorological data. However, the magnitude of the AOD results can differ significantly among the four model runs (note the semi-logarithmic scale!). It is also remarkable that the differences between the two optics models depend on whether we make this comparison within the mass-transport model (compare MT-EXT to MT-CGS), or within Salsa (compare SALSA-CGS and Salsa-EXT). It can also vary geographically. This merely confirms the complexity of aerosol-optics modelling. Replacing one optics model by another will not simply offset the resulting optical properties by some common factor; it will introduce a significant change in modelled optical properties, of which the magnitude and even the sign can be dependent on local conditions, such as the size-distribution and the chemical composition of the aerosol particles.

This is also evident from a comparative analysis of geographically and temporally averaged aerosol optical properties. Table 4 shows aerosol optical depth (AOD), backscattering coefficient (BSCA), single scattering albedo (SSA) and asymmetry parameter (g), each at three different wavelengths (355, 532, and 1064 nm), and each averaged over the whole model domain and over one month (December 2007). The columns show relative differences; for instance, $MT(EXT,CGS) = (MT-EXT - MT-CGS)/(MT-CGS)$ is the relative difference of MT-model results obtained with the EXT and CGS optics models.

Table 4. Averaged relative difference in aerosol optical depth (AOD), backscattering coefficient (BSCA), single scattering albedo (SSA) and asymmetry parameter (g), among the different model set-ups for December 2007. The average has been performed over a whole month and over all grid-cells (horizontally for AOD, horizontally and vertically for all other optical properties). Each number corresponds to a relative difference between two model set-ups. For instance, $MT(EXT,CGS) = (MT-EXT - MT-CGS)/(MT-CGS)$ compares results obtained with the mass transport model (MT) by using the two different optics models (EXT and CGS).

$X_{\lambda,[nm]}$	MT(EXT,CGS)	Salsa(EXT,CGS)	CGS(MT,Salsa)	EXT(MT,Salsa)
AOD ₃₅₅	0.16	-0.28	-0.50	-0.19
AOD ₅₃₂	0.08	-0.14	0.00	0.25
AOD ₁₀₆₄	0.18	-0.03	0.14	0.37
BSCA ₃₅₅	0.44	-0.01	-0.47	-0.23
BSCA ₅₃₂	0.26	-0.08	-0.19	0.11
BSCA ₁₀₆₄	0.99	-0.01	-0.36	0.28
SSA ₃₅₅	-0.02	0.04	0.03	-0.03
SSA ₅₃₂	-0.02	0.05	0.04	-0.02
SSA ₁₀₆₄	-0.07	0.08	0.08	-0.07
g ₃₅₅	0.06	-0.01	-0.10	-0.03
g ₅₃₂	0.10	-0.00	-0.06	0.04
g ₁₀₆₄	0.17	-0.02	-0.11	0.06

Comparison of the columns "MT(EXT,CGS)" and "Salsa(EXT,CGS)" illustrates the differences between the optics models in the absence and presence of aerosol-microphysical processes. As we
440 already suspected from inspection of Fig. 4, differences in the optics models defy simplistic explanations; both the magnitude and sign of these difference can be strongly dependent on the size-dependent chemical composition and mixing state of the aerosols, hence on the model version with which the optics models are being compared. In our case, we see that compared to the CGS model, the EXT-optics model predicts higher values of AOD, BSCA, and g in the MT model, and lower
445 values of SSA. In Salsa the roles of the CGS and EXT model are reversed.

The column "CGS(MT,Salsa)" shows differences between optical properties computed in the absence and presence of aerosol-microphysical processes, where the optics-computations have been performed with the CGS model. The column "EXT(MT,Salsa)" shows an analogous comparison, where the optics-computations have been performed with the EXT model. If we compare the magni-
450 tudes of the entries in the columns "MT(EXT,CGS)" and "Salsa(EXT,CGS)" with the corresponding magnitudes of the entries in the columns "CGS(MT,Salsa)" and "EXT(MT,Salsa)", then we see that the differences between the two optics models (EXT,CGS) are roughly on the same order as the differences between the two aerosol models (MT,Salsa). Thus, the main observation is that the choice of aerosol-optics model can have an effect on modelled optical properties that is of comparable
455 magnitude as the level of detail in the description of aerosol-microphysical processes.

While spatio-temporally averaged model results allows us to draw some general conclusions, it is difficult to understand the reasons for the observed differences from such an analysis. We will, therefore, complement this investigation in the following sections with a more detailed analysis of some selected case studies.

460 3.2 Optical properties and radiative forcing

In Sec. 2.3.2 we have discussed how morphological properties of aerosol particles can impact their optical properties. We now take this one step further and discuss how the optical properties of particles can impact the radiative properties of an aerosol-laden atmosphere. We will show results for a single wavelength near the maximum of the solar spectrum. The comparison will focus on three
465 model set-ups, MT-EXT, MT-CGS, and Salsa-CGS. We will use MT-CGS as a reference and compare it first to Salsa-CGS in order to investigate the impact of aerosol-microphysics (MT versus Salsa). Next, we compare MT-CGS to MT-EXT in order to investigate the impact of the optics model (CGS versus EXT).

The result for the optical properties obtained with the three model versions (AOD per layer, SSA
470 and g) at the wavelength 532(CGS)/500(EXT) nm, together with the radiative forcing for all aerosol component and black carbon, respectively, can be seen in Figs. 5–10 for Northern Italy and the Mediterranean on 22 June 2007. Each figure shows the differences in direct solar flux ΔF_s (top left), diffuse downwelling flux ΔF_d (top right), diffuse upwelling flux ΔF_u (centre left), and net radiative flux $\Delta F_{\text{net}} = \Delta F_s + \Delta F_d - \Delta F_u$ (centre right), where either the difference between aerosol-laden
475 and clear sky conditions are considered (Figs. 5 and 6), or the difference between fluxes in the presence and absence of black carbon (Figs. 9 and 10). Here, *downwelling* fluxes are obtained by integrating the radiance over all azimuthal angles, and over polar angles from 90° to 180° , where the positive z -axis is directed from the ground to the top-of-atmosphere (TOA). Similarly, the *upwelling* flux is obtained by integrating the radiance over all azimuthal angles, and over polar angles from 0 to
480 90° . TOA results for the other geographical locations are summarized in Table 5 in terms of aerosol forcing ($\Delta F_{\text{net}} = F_{\text{net}}(\text{with aerosol particles}) - F_{\text{net}}(\text{no aerosol particles})$), and in Table 6 in terms of black-carbon forcing ($\Delta F_{\text{net}} = F_{\text{net}}(\text{with black carbon}) - F_{\text{net}}(\text{no black carbon})$). The wavelengths 532(CGS)/500(EXT) nm are near the maximum of the solar spectrum. Each figure has a vertical span of 5 km, which comprises that part of the troposphere where most aerosol particles are concentrated
485 in the cases we picked.

3.2.1 Comparison of aerosol microphysics and mass-transport model

We start by comparing radiative fluxes in the presence and absence of all aerosol components, which we refer to as the “aerosol radiative effects”. Figures 5 and 6 show the aerosol radiative effects modelled over Northern Italy and over the Mediterranean north of Algiers, respectively. The general
490 patterns in both plots can be understood as follows. In the presence of all aerosol components,

optical extinction is stronger than in clean air. Thus the presence of all aerosol components gives a reduction ΔF_s of the direct solar flux (upper left). As the aerosol optical depth per atmospheric layer strongly increases near the ground, the magnitude of ΔF_s increases sharply with decreasing altitude. Further, extinction in the form of scattering results in the generation of diffuse flux; the downwelling diffuse flux accumulates downward, resulting in an increasing excess of downwelling flux ΔF_d in the presence of aerosol components as one approaches the surface. The upwelling flux F_u is generated by scattering in the atmosphere and reflection at the surface. Since aerosol extinction reduces the net radiative flux as one approaches the surface, less upwelling diffuse flux is generated at low altitudes; hence the difference in upwelling flux ΔF_u between an aerosol-laden and a clear sky atmosphere is negative near the surface. However, it increases with altitude, because at higher altitudes the magnitude of the difference (aerosol – clear sky) in the net radiative flux that can be converted into upwelling diffuse flux decreases at higher altitudes.

If we focus now on differences in the radiative net flux ΔF_{net} at high altitudes, i.e., the radiative forcing effect of aerosol particles, then we see significant differences between the mass transport model (MT, red) and SALSA (green) at both geographic locations. It is evident that the main cause for these are corresponding differences in the diffuse upwelling flux ΔF_u .

At both locations the diffuse upwelling flux is smaller for SALSA than for MT, but for different reasons. Over the Mediterranean (Fig. 6), the AOD is significantly smaller for SALSA than for MT, resulting in less extinction of the direct flux, hence less generation of diffuse flux, and a smaller radiative cooling effect for SALSA.

Over Northern Italy (Fig. 5), there is almost no difference in AOD between the two models. It can be seen from the AOD profile that the majority of aerosol particles reside in the lowest 1 km near the surface. However, above 1 km the results of ΔF_u obtained with SALSA and MT diverge with increasing altitude. This is a result of the reflection by the near-surface aerosol layer, which is different in the two models. In MT the SSA is higher than in SALSA, resulting in more scattering, thus in more diffuse radiation. The asymmetry parameter is larger in MT than in SALSA; correspondingly, the partitioning in MT between downwelling and upwelling radiation is somewhat shifted in favour of the former. However, this only partially counteracts the generation of a higher amount of diffuse upwelling radiation in MT due to the higher SSA. The net effect is a higher value of ΔF_u in MT, hence a larger radiative cooling effect at higher altitudes.

To further analyse the difference in optical properties between MT and SALSA, we look at the aerosol masses and the relative sizes of the particles. Figure 7 shows vertical profiles of the effective radius r_{eff} defined according to Eq. 1, in SALSA (green) and the MT model (black) over Northern Italy (left) and over the Mediterranean (right).

$$r_{\text{eff}} = \frac{\int_0^{\infty} n(r) \pi r^2 \cdot r dr}{\int_0^{\infty} n(r) \cdot \pi r^2 dr} \quad (1)$$

Figure 8 shows profiles for the total aerosol mass (1st row), BC (black carbon) (2nd row), sulphate (3rd row), and nitrate (4th row) for both Northern Italy (1st column) and Mediterranean (2nd column). We focus on the total aerosol mass, which is expected to impact the aerosol optical depth. The aerosol optical depth is dependent on the number density (which, in turn, increases with the mass mixing ratio), as well as on the extinction cross section (which generally increases with the effective radius of the particles). Over Northern Italy, the SALSA model predicts a larger mass mixing ratio than the MT model (Fig. 8, upper left) and a smaller particle size (Fig. 7, left). This results in a higher number density but a smaller extinction cross section. These two effects cancel almost exactly, resulting in nearly identical aerosol optical depths predicted with the two models (Fig. 5, bottom left).
530 By contrast, over the Mediterranean the two models predict similar mass mixing ratios (Fig. 8, upper right), while SALSA predicts a much lower effective radius than the MT model (Fig. 7, right). As a consequence, the optical depth is significantly lower in SALSA than in MT (Fig. 6, bottom left). The SSA is lower in SALSA than in MT. This is mainly caused by the fact that the effective radius is smaller in SALSA than in MT, since SSA is usually increasing with size.

540 For the other two geographical locations and the second time event, the TOA results are summarised in Table 5. Over the North Sea, Northern Italy, and the Mediterranean the TOA forcing is strongest in the MT-EXT model (mass transport with old optics model), it is smaller in the MT-CGS model (mass transport with new optics model), and weakest in the SALSA-CGS model (aerosol-microphysics with new optics model). However, over Poland the negative forcing in SALSA-CGS
545 is strongest among the three models in the summer, and second strongest in the winter. This can be explained by SALSA having a larger amount of aerosol particles throughout the column at that site, especially more sulphate, which, when externally mixed, contributes to a larger amount of scattering and therefore a higher SSA and a larger diffuse upwelling radiative flux.

We now compare radiative fluxes in the presence and absence of black carbon, which we refer
550 to as the “black carbon radiative effect”. Figures 9 and 10 show the radiative effect of black carbon together with the optical properties with and without black carbon. Again, differences in ΔF_{net} at TOA are mainly caused by corresponding differences in the upwelling diffuse radiative flux ΔF_{u} . In these figures, we have to focus on the difference in the optical properties when analysing the radiative fluxes. The general pattern can be seen in Fig. 9, which shows the differences in radiative
555 fluxes and in the optical properties over Northern Italy. The direct solar flux (upper left) decreases with decreasing altitude owing to extinction. The magnitude of the decrease mainly reflects the differences in optical depth in the presence and absence of black carbon (bottom left), which is larger in SALSA than in MT. The decrease in solar flux does not automatically result in an increase in the downwelling diffuse flux with decreasing altitude (upper right), as it was in the comparison
560 of fluxes in the presence and absence of *all* aerosol components. The situation is more complex now. Near the surface, where the optical depth is largest, the difference in SSA in the presence and absence of black carbon is large in the MT model (bottom centre, red lines), and smaller in SALSA

(green lines). As a result, absorption contributes more to the total extinction in the MT model than in SALSA (at least near the surface). Hence, the portion of the downwelling flux that is absorbed on its way downward is larger in the MT model than in SALSA, resulting in a decrease of the diffuse downwelling flux with decreasing altitude (upper right, red line). The differences between the two models in the diffuse upwelling flux are very small (centre left, red and green lines). This is the result of cancelling effects; for instance, there is less direct solar flux, but more diffuse downwelling flux in SALSA that can be converted into diffuse upwelling flux through scattering. As a result, the differences between both models in the net flux (centre right, red and green lines) are almost negligible.

Figure 10 shows the radiative effect of black carbon over the Mediterranean. Again, differences in ΔF_{net} at TOA are mainly caused by corresponding differences in the upwelling diffuse radiative flux ΔF_{u} . MT and SALSA only differ by a few mW/m^2 . The main reason is that the optical properties, especially the SSA, differ in the presence and absence of BC by almost the same amount in both models.

Table 6 shows the black carbon forcing for all the four geographical locations and both months. As a general trend, large differences in BC concentrations are visible as corresponding differences in BC forcing. For instance, near-surface BC winter- concentrations in Northern Italy are about a factor of 10 higher than in summer; summer-concentrations over the Mediterranean are more than a factor of 2 higher than in winter; in Northern Italy in winter, Salsa predicts more than 2 times higher BC-concentrations than the mass-transport model, while over the Mediterranean in winter the role of the two models is reversed (not shown). All of this corresponds with the respective BC-forcing rates in Table 6. However, when the differences in BC-concentrations are small, then there is no clear relation between the concentration-differences and the corresponding differences in BC-forcing rates. For instance, as we see in Fig. 8, there is almost no difference between the BC-concentrations computed with the two models over the Mediterranean in summer. But the table shows that the mass-transport model predicts a slightly higher forcing rate than SALSA. A possible cause is the difference in the size-distributions in SALSA and the mass-transport model.

3.2.2 Comparison of the two optics models

The comparison between SALSA and the MT model in the previous section served two purposes. First, it helped us to develop a basic understanding for the effects of aerosol particles and black carbon on radiative fluxes. Second, it provided us with a gauge for assessing the importance of the aerosol-optics model, which will be the subject of this section.

We compare the old EXT (blue line) and the new CGS (red line) optics models in Figs. 5 and 6, each used in conjunction with the MT-version of MATCH. The net radiative flux ΔF_{net} in the CGS model shows a weaker TOA cooling effect than the EXT model, both over Northern Italy and over the Mediterranean. Again, the upwelling flux has the dominant impact on the behaviour of the TOA

net radiative flux. Over Northern Italy (Fig. 5) the diffuse upwelling flux is larger for the EXT model
600 above 1 km, whereas it is smaller below 1 km and at the bottom of the atmosphere (BOA). The AOD
profile reveals that in the EXT model extinction is stronger than in the CGS model throughout the
tropospheric column. As a result, there is more diffuse downwelling flux being generated in EXT
than in CGS. At the bottom of the atmosphere (BOA) extinction of the direct flux is stronger than
generation of diffuse downwelling flux; hence less downwelling flux is reflected by the surface,
605 resulting in less BOA upwelling diffuse flux in EXT than in CGS. Higher up in the troposphere,
the upwelling diffuse flux is mostly generated by atmospheric scattering rather than reflection from
the surface. As the SSA in EXT is higher than in CGS, more diffuse flux is generated, resulting in
a stronger radiative cooling effect in EXT than in CGS.

Over the Mediterranean (Fig. 6), the EXT and CGS model have almost identical AOD profiles in
610 the green part of the spectrum. However, at longer wavelengths EXT predicts substantially higher
AOD values than CGS (see Appendix E, Fig. 16). For instance, at $\lambda = 1020$ nm the near-surface
AOD per layer computed with the EXT-model is about twice as high as that computed with the CGS-
model. This explains the larger amount of diffuse broadband radiation generated in the EXT model,
which results in a stronger negative TOA net flux in EXT as compared to the CGS model. Note
615 that the differences in SSA between EXT and CGS are less than 0.03, while the differences in g
are as much as 0.07. The higher values of g in EXT may contribute to the large amount of diffuse
downwelling radiation in that model; however, the dominant effect is likely to be the high optical
depth at red and IR wavelengths (see Appendix E).

Table 5 summarises the TOA net radiative flux at all four geographical locations for both June and
620 December. The largest differences among the models are seen in December at the two northernmost
locations, i.e., Poland and the North Sea. At these two places, the total aerosol amount (not shown)
is significantly higher than at the other two locations farther south, giving rise to larger absolute
changes in the aerosol forcing.

The black carbon forcing in Fig. 9 (Northern Italy) and 10 (Mediterranean) display different be-
625 haviours in radiative fluxes, comparing the EXT (blue) and CGS (red) model results. Over Northern
Italy, the net black carbon forcing is more significant than over the Mediterranean in Fig. 10 due to
higher levels of BC, see Fig. 8. As can be seen in Fig. 9, the differences in optical properties com-
puted with and without black carbon are larger in the CGS model than in the EXT model, particularly
for the SSA. This means that in the CGS model the presence of black carbon causes more absorption
630 than in the EXT model, thus generating less diffuse down- and upwelling flux by scattering. As a re-
sult, the CGS model predicts more radiative warming, i.e., a higher TOA radiative net flux than the
EXT model. The reason for this is that (i) the CGS model treats externally mixed soot as aggregates,
which have a lower SSA than the massive black carbon spheres in the EXT model; and (ii) the CGS
model treats internally mixed soot as a coated core-grey-shell model, which accounts for focusing of




Figure 5. Aerosol forcing and optical properties at 532(CGS)/500(EXT) nm over Northern Italy in June. Results are shown for the three model versions MT-EXT (blue), MT-CGS (red) and Salsa-CGS (green). The aerosol forcing is derived by taking the difference in radiative fluxes between an aerosol laden and a clear sky. The difference in direct (ΔF_s) and diffuse (ΔF_d) downwelling, as well as the diffuse upwelling (ΔF_u) and the net radiative flux (aerosol forcing) ($\Delta F_{\text{net}} = \Delta F_s + \Delta F_d - \Delta F_u$) are shown in the first four figures (1st and 2nd row of plots). The optical properties aerosol optical depth (AOD), single scattering albedo (SSA) and asymmetry parameter (g) are shown in the 3rd row of plots.

635 electromagnetic radiation onto the carbon core, thus enhancing absorption, i.e., lowering the SSA, while the EXT model treats *all* black carbon as externally mixed.

Let us now return to the main question of this study, namely, whether or not the level of detail in aerosol optics modelling has a significant impact on observable radiometric properties. We already saw in Table 4 that, on average, the effect of including aerosol microphysics on optical properties is
640 of comparable magnitude as the effect caused by the morphological assumptions in the aerosol optics model. However, we also saw that the magnitude and sign of these impacts can be quite variable and depending on several factors. We find this confirmed in the analysis of our radiative-transfer study. More specifically, we can compare in Figs. 5–10 the differences in radiative forcing between MT-CGS and Salsa-CGS (red and green lines) to the corresponding differences between MT-CGS and
645 MT-EXT (red and blue lines). We see that in some cases the choice of optics model has a stronger effect than the inclusion of aerosol microphysics (e.g. Fig. 9), while in other cases it is the other way round (e.g. Fig. 6). We can also inspect Tables 5 and 6 and arrive at the same result.




Figure 6. Same as figure 5 but over the Mediterranean.




Figure 7. Effective radius, r_{eff} , for the two chemical transport model versions MT and SALSA over Northern Italy and Mediterranean the 2007-06-22 at 12:00.




Figure 8. Vertical distribution of aerosol particles in Northern Italy and over the Mediterranean on 2007-06-22 at 12:00.

Table 5. The aerosol forcing at the top of the atmosphere (TOA), $\Delta F_{\text{net,TOA}}$ [W m^{-2}], for the four different geographical locations, one summer (2007-06-22, 12:00) and one winter (2007-12-22, 12:00) event, and three model versions, MT-EXT, MT-CGS and Salsa-CGS.

		MT-EXT	MT-CGS	SALSA-CGS
Summer	Poland	-0.21	-0.21	-0.77
	North Sea	-0.34	-0.29	-0.24
	Northern Italy	-0.06	-0.05	0.01
	Mediterranean	-1.20	-0.99	-0.30
Winter	Poland	-4.41	-2.00	-2.18
	North Sea	-2.85	-1.21	-0.86
	Northern Italy	-1.15	-0.53	-0.57
	Mediterranean	-0.09	-0.04	-0.03




Figure 9. Black carbon forcing and optical properties at 532(CGS)/500(EXT) nm over Northern Italy in June. Results are shown for the three model versions MT-EXT (blue), MT-CGS (red) and Salsa-CGS (green). The black carbon forcing is derived by taking the difference in radiative fluxes between an aerosol laden including black carbon and an aerosol laden sky omitting black carbon. The difference in direct (ΔF_s) and diffuse (ΔF_d) downwelling, as well as the diffuse upwelling (ΔF_u) and the net radiative flux (aerosol forcing) ($\Delta F_{\text{net}} = \Delta F_s + \Delta F_d - \Delta F_u$) are shown in the first four figures (1st and 2nd row of plots). The optical properties aerosol optical depth (AOD), single scattering albedo (SSA) and asymmetry parameter (g) are shown in the 3rd row of plots.

Table 6. Same as table 5, but for black carbon forcing.

		MT-EXT	MT-CGS	SALSA-CGS
Summer	Poland	1.02×10^{-2}	1.16×10^{-2}	1.20×10^{-2}
	North Sea	1.71×10^{-2}	1.54×10^{-2}	1.49×10^{-2}
	Northern Italy	4.61×10^{-2}	7.77×10^{-2}	7.86×10^{-2}
	Mediterranean	8.54×10^{-3}	6.45×10^{-3}	2.41×10^{-3}
Winter	Poland	4.03×10^{-2}	3.56×10^{-2}	6.83×10^{-2}
	North Sea	1.95×10^{-2}	2.28×10^{-2}	4.97×10^{-3}
	Northern Italy	6.73×10^{-2}	1.08×10^{-1}	1.46×10^{-1}
	Mediterranean	6.03×10^{-4}	1.34×10^{-3}	3.13×10^{-4}




Figure 10. Same as Figure 9, but over the Mediterranean.

In the following two subsections, we will focus on the selected case-studies and discuss the significance of the optics model for radiometric quantities that are relevant for remote sensing applications.

650 3.3 Backscattering coefficient

From ground-based and space-borne lidar measurements one can obtain the aerosol backscattering coefficient β , which is proportional to the backscattering cross section C_{bak} of the particles and the aerosol number density. Figure 11 shows vertical profiles of β computed at two locations and at two instances, as indicated in the figure headings. Each panel shows computational results obtained
 655 with the three different model versions. The figure shows results for the second Nd:YAG harmonic of 532 nm. Corresponding results computed for wavelengths of 355 and 1064 nm lead to similar conclusions.

We saw in Fig. 8 for June over Northern Italy (upper left) that SALSA predicts an aerosol mass mixing ratio, hence a particle number density, that is higher than that in the MT model. But we also
 660 saw in Fig. 7 (left) that SALSA predicts lower values of r_{eff} . This results in lower values of C_{bak} . We see in Fig. 11 (upper left) that the effect on β of the higher number density dominates over the effect of the lower r_{eff} , resulting in values of β that are about 30 % higher in SALSA (green line) than in the MT model (red line). Over the Mediterranean, both SALSA and the MT model predict similar mass mixing ratios (Fig. 8, upper right); but SALSA still predicts substantially lower values of r_{eff}




Figure 11. Backscattering coefficient at a wavelength of 532 nm at Northern Italy and Mediterranean and at two different time events (22/6 and 22/12-2007 at 12.00), computed with the three model versions, MT-EXT (blue), MT-CGS (red) and Salsa-CGS (green).

665 (Fig. 7, right). The result is that β computed with the MT model (red line) is almost twice as high as the corresponding results obtained with SALSA (green line) (Fig. 11, upper right).

A similar comparison of the two optics models (red and blue lines in Fig. 11) shows that the new CGS optics model consistently predicts substantially lower values of β than the old EXT optics model. This agrees with the comparison shown in Kahnert et al. (2013) between encapsulated black carbon aggregates and externally mixed homogeneous spheres. (In a retrieval algorithm, an optics model that overestimates the backscattering cross section would result in underestimated retrieval results for the particle number density.) The differences between the two optics models are on the same order of magnitude (and often even larger) than the corresponding differences between the SALSA and the MT versions of the aerosol transport model.

675 3.4 Ångström exponent

The Ångström exponent α in a wavelength interval $[\lambda_1, \lambda_2]$ is defined as

$$\alpha = -\frac{\log(\tau(\lambda_1)/\tau(\lambda_2))}{\log(\lambda_1/\lambda_2)}, \quad (2)$$

Table 7. Ångström exponent in the wavelength region 532–1064 nm for the four different geographical locations, one summer (2007-06-22, 12:00) and one winter (2007-12-22, 12:00) event, and three model versions, MT-EXT, MT-CGS and Salsa-CGS.

		MT-EXT	MT-CGS	SALSA-CGS
Summer	Poland	0.32×10^0	0.12×10^1	0.28×10^1
	North Sea	0.80×10^0	0.12×10^1	0.21×10^1
	Northern Italy	0.11×10^1	0.11×10^1	0.15×10^1
	Mediterranean	0.36×10^0	0.12×10^1	0.21×10^1
Winter	Poland	0.80×10^0	0.12×10^1	0.22×10^1
	North Sea	0.79×10^0	0.11×10^1	0.14×10^1
	Northern Italy	0.13×10^1	0.10×10^1	0.12×10^1
	Mediterranean	0.13×10^1	0.98×10^0	0.14×10^1

where τ denotes the extinction optical depth. This quantity is often used for obtaining particle size information (usually, the smaller the particle size, the larger α). Table 7 shows values of α for our different test cases computed with the three model versions in the wavelength interval 532–1064 nm. If we compare the columns labelled MT-CGS and SALSA-CGS, then we see that the mass-transport model consistently gives lower values of α . This is related to the high values of r_{eff} in that model, which we noted earlier. On the other hand, if we compare the columns labelled MT-EXT and MT-CGS, then we see that the new optics model (CGS) predicts higher values of α than the old model (EXT) in June (summer) for all four geographical locations and in December (winter) for the locations Poland and North Sea. This indicates that the errors introduced by the simple external-mixture model in computing α are unpredictable, even the sign of the error. When used in a size retrieval algorithm the retrieval errors caused by the EXT model would be equally hard to predict. The difference between the MT and SALSA model is larger, but not much larger, than the differences between the old and new optics models. Note that the performance of the MT model could be improved in comparison to SALSA by modifying the assumed size distribution in the MT model. By contrast, the differences between the two optics models are rather fundamental; it is caused by the simple treatment of aerosol morphology in the EXT model.

4 Conclusions

We have implemented a new aerosol-optics model in a regional chemical transport model. The new model differs from an earlier optics model described in Kahnert (2008) in three essential points. (i) While the old model treats all chemical components as externally mixed, the new model accommodates both external and internal mixtures of aerosol species. (ii) The old model treats black

carbon particles as homogeneous spheres; the new model assumes a fractal aggregate morphology
700 with fractal parameters based on observations. Mass absorption cross sections and single scattering
albedos computed with this model have previously been evaluated by comparison with measure-
ments (Kahnert, 2010b). (iii) The new model describes internally mixed black carbon particles by
a recently developed “core-grey-shell” model (Kahnert et al., 2013). This model accounts for the
inhomogeneous internal mixing state of black carbon aggregates encapsulated in a shell of liquid-
705 phase material. The model has been evaluated by comparison with reference computations based on
observation-derived realistic models for encapsulated fractal aggregates (Kahnert et al., 2013). Item
(i) has been incorporated in other CTMs earlier (e.g. Saide et al., 2013); however, to the best of our
knowledge, items (ii) and (iii) go significantly beyond the current state-of-the-art of aerosol-optics
models employed in CTMs. The main question of the present study is whether or not such a sub-
710 stantial level of detail in the description of aerosol morphology and optical properties is needed in
a CTM.

We first performed a comparison of optical properties averaged over the entire model domain and
over one month. To gauge the differences between the new and the old optics model, we further com-
pare two model versions of the CTM with different levels of detail in the aerosol process descriptions,
715 namely, one version that includes aerosol-dynamic processes, and a simpler mass-transport model,
in which aerosol microphysics is switched off. The importance of aerosol microphysics is well un-
derstood and can therefore serve as a reference. We found that the differences in optical properties
between the two optics models are on the same order as those between the versions that include
and exclude microphysical processes. For example, the aerosol optical depth computed with the two
720 optics models differs by $-25-18\%$; differences obtained by inclusion or omission of aerosol mi-
crophysics are between $-50-37\%$. Corresponding differences in the backscattering coefficient are
 $-8-99\%$ and $-47-28\%$, respectively. Analogous observations can be made for other radiometric
properties.

We further wanted to understand how the differences in optical properties impact radiative transfer
725 processes in an aerosol-laden atmosphere. To this end we compare radiative fluxes modelled with
the old and new optics models. The comparison showed that the differences in radiative net-fluxes
between the two optics models are of similar magnitude as corresponding differences between the
aerosol-microphysics and the mass-transport models.

These results strongly suggest that simplifications in the assumptions on aerosol morphology in
730 the optics model can introduce substantial errors in modelled radiative fluxes and observables rel-
evant to remote sensing. In chemistry-climate models such errors would enter into the simulation
of the direct aerosol radiative forcing effect and add to all other sources of error in the model. In
model evaluations that make use of remote sensing observations these errors would complicate the
comparison between model results and observations.

735 The modifications to the morphology-assumptions in the optics model were limited to black-carbon particles. There are many other aerosol particles with complex morphological properties, such as mineral dust, which our optics model still treats by a simple homogeneous-sphere model. The findings of our study should be an incentive for improving the description of dust and volcanic-ash optical modelling in CTMs. A recent review of our current state-of-knowledge on aerosol morphology and aerosol optics for a variety of different aerosol particles can be found in Kahnert et al. (2014).

The findings of this study are likely to have implications for chemical data assimilation. In data assimilation one employs an *observation operator* that maps the model results to observable quantities. In case of satellite-based observations of aerosol optical properties, the observation operator is just our aerosol-optics model, possibly coupled to a radiative transfer model. Many data-assimilation methodologies, such as the variational method, require a linear (or, at least, linearised) observation operator. In the old optics model, which assumes externally-mixed aerosol particles, the observation operator is, indeed, linear (Kahnert, 2008). This largely explains why external-mixture optics models are widely used in chemical data assimilation systems (e.g. Kahnert, 2008; Benedetti et al., 2009; Liu et al., 2011). However, the new optics model we introduced here does not provide us with a linear map from the aerosol concentrations to the optical parameters. To what extent one could linearise this model and make use of its Jacobian in a data assimilation system mainly depends on the degree of nonlinearity, which would need to be investigated thoroughly.

5 Code availability

755 The aerosol microphysics code SALSA is distributed under the Apache 2.0 license, while the chemistry transport model MATCH and the aerosol-optics data base are available upon request from SMHI.

Appendix A: Size-averaged optical properties in the external-mixture optics model

The external-mixture optics model is based on using four size bins that cover the dry-radius intervals $[r^{\min}, r^{\max}] = [0.01, 0.05] \mu\text{m}$, $[0.05, 0.5] \mu\text{m}$, $[0.5, 1.25] \mu\text{m}$, and $[1.25, 5] \mu\text{m}$. The geometric mean radius $R = \sqrt{r^{\min} r^{\max}} = 0.5(\log r^{\min} + \log r^{\max})$ is given in each of these intervals by $R_1 = 0.022 \mu\text{m}$, $R_2 = 0.158 \mu\text{m}$, $R_3 = 0.791 \mu\text{m}$, and $R_4 = 2.5 \mu\text{m}$. In each size bin it is assumed that the particle number density is given by a log-normal distribution

$$n_i(r) = N_i^0 / (\sqrt{2\pi} r \ln \sigma_i) \exp[-\ln^2(r/R_i)/(2\ln^2 \sigma_i)], \quad (\text{A1})$$

765 where $\sigma_1 = \sigma_3 = \sigma_4 = 1.8$, $\sigma_2 = 1.5$ are based on measurements in Neusüß et al. (2002). Here, N_i^0 would be the total number density per mode if each size-mode extended from $r = 0$ to $r = \infty$. However, since each mode is truncated at the bin-boundaries r^{\min} and r^{\max} , the number density N_i

of particles per size bin is obtained from integration over this finite interval, i.e.

$$N_i = \int_{r_i^{\min}}^{r_i^{\max}} n_i(r) dr = N_i^0 \frac{1}{2} [\operatorname{erf}(x_i^{\max}) - \operatorname{erf}(x_i^{\min})], \quad (\text{A2})$$

770 where $x_i^{\max} = \ln(r_i^{\max}/R_i)/(\sqrt{2}\ln\sigma_i)$, and similarly for x_i^{\min} . Analogously, one obtains the particle-mass density M_i in each size bin by integrating over the truncated log-normal mode, which yields

$$\begin{aligned} M_i &= \frac{4}{3}\pi\rho_i \int_{r_i^{\min}}^{r_i^{\max}} n_i(r)r^3 dr \\ &= \frac{4}{3}\pi R_i^3 \rho_i N_i^0 \exp\left(\frac{9}{2}\ln^2\sigma_i\right) \frac{1}{2} [\operatorname{erf}(y_i^{\max}) - \operatorname{erf}(y_i^{\min})], \end{aligned} \quad (\text{A3})$$

775 where $y_i^{\max} = x_i^{\max} - 3\ln\sigma_i/\sqrt{2}$, and similarly for y_i^{\min} , and where ρ_i is the density of the aerosol particles in the i th size bin. From this we obtain the desired relation for converting the mass-density M_i into the number-density N_i :

$$N_i = \frac{M_i}{\frac{4}{3}\pi R_i^3 \rho_i} \cdot \frac{\operatorname{erf}(x_i^{\max}) - \operatorname{erf}(x_i^{\min})}{\exp\left(\frac{9}{2}\ln^2\sigma_i\right) [\operatorname{erf}(y_i^{\max}) - \operatorname{erf}(y_i^{\min})]}. \quad (\text{A4})$$

Mass-mixing ratios X_i are simply converted into mass densities M_i according to $M_i = X_i\rho_{\text{air}}$, where ρ_{air} denotes the density of air.

780 In the external-mixture optics-database, optical properties are pre-computed by integrating optical properties at discrete sizes over the truncated log-normal size distribution. This integration is done numerically with a high size-resolution. The computation is performed for different refractive indices m , optical wavelengths λ , and for each size bin i . Thus, one obtains, e.g., extinction cross sections $C_{\text{ext}}(\lambda, m, i)$, which can be saved in a look-up table.

785 Secondary inorganic aerosols as well as organic aerosols and sea salt are assumed to be hydrophilic. We use the parameterisation by Gerber (1985) to compute the wet-radius r_w from the aerosol dry radius R , from which we obtain the volume-fraction of water $f_w = (r_w^3 - R^3)/R^3$. The effective refractive index m of the aerosol-water mixture is computed from that of the dry aerosol, m_a , and of water, m_w by use of effective-medium theory. In each grid cell, we obtain from the
790 MATCH model, for each size bin i and for each aerosol component k , the number density $N_i(k)$. From that we compute the ensemble-averaged extinction cross section

$$\bar{C}_{\text{ext}}(\lambda) = \frac{1}{N} \sum_k \sum_{i=1}^4 N_i(k) C_{\text{ext}}(\lambda, m(k), i), \quad (\text{A5})$$

where the total number density is given by

$$N = \sum_k \sum_{i=1}^4 N_i(k). \quad (\text{A6})$$

795 Note that the ensemble-average involves an average over both size and chemical composition. The ensemble-averaged scattering cross section $\bar{C}_{\text{sca}}(\lambda)$ is computed analogously. From this we obtain the averaged single-scattering albedo

$$\bar{\omega}(\lambda) = \frac{\bar{C}_{\text{sca}}(\lambda)}{\bar{C}_{\text{ext}}(\lambda)}. \quad (\text{A7})$$

The phase function $p(\Theta)$, hence its first Legendre-moment, known as the asymmetry parameter g are normalised quantities. Here Θ denotes the scattering angle. To average these quantities, one first needs to "de-normalise" by multiplying them with the scattering cross section. Thus

$$\bar{p}(\Theta; \lambda) = \frac{1}{N\bar{C}_{\text{sca}}(\lambda)} \sum_k \sum_{i=1}^4 N_i(k) C_{\text{sca}}(\lambda, m(k), i) p(\Theta, m(k), i; \lambda) \quad (\text{A8})$$

$$\bar{g}(\lambda) = \frac{1}{N\bar{C}_{\text{sca}}(\lambda)} \sum_k \sum_{i=1}^4 N_i(k) C_{\text{sca}}(\lambda, m(k), i) g(m(k), i; \lambda). \quad (\text{A9})$$

Once the ensemble-averaged optical properties in each grid cell of the model domain have been computed, one can compute radiometric observables, such as the extinction aerosol optical depth

$$\tau_{\text{ext}}(\lambda) = \sum_z N(z) \bar{C}_{\text{ext}}(\lambda, z) \Delta z, \quad (\text{A10})$$

or the backscattering coefficient

$$\beta_{\text{bak}}(\lambda, z) = \frac{1}{4\pi} N(z) \bar{C}_{\text{sca}}(\lambda, z) \bar{p}(180^\circ; \lambda, z), \quad (\text{A11})$$

where z labels grid cells in the vertical column, and Δz denotes the layer-thickness.

810 **Appendix B: Size-averaged optical properties in the internal-mixture model**

In SALSA the number-density as a function of particle radius, $n(r)$, is given by a step-function with $n_i(r) = \text{const}_i$ in each size bin i . This makes the pre-integration of optical properties over each size bin rather simple. On the other hand, we no longer assume that all aerosol components are externally mixed. Thus the ensemble-average over different chemical components k is no longer given by a simple summation $\sum_k \dots$, as it was in the external-mixture model. Rather, for each size bin in which several aerosol components are internally mixed one computes an effective refractive index, m_{eff} , by use of effective-medium theory. One then reads the optical properties for that refractive index from the look-up table. Finally, one computes ensemble-averaged optical properties by summing over all size bins, $\sum_i \dots$.

820 **Appendix C: Effective-medium theory**

In effective-medium theory (EMT) one considers a composite material consisting of two materials with refractive indices m_1 and m_2 and volume fractions f_1 and $f_2 = 1 - f_1$. One then invokes assumptions about the topology of the mixture and derives a formula for the effective refractive index,

m_{eff} , of the composite material. For instance, it is often the case that $f_1 \gg f_2$. In this case one can regard the first material as a host matrix that contains inclusions of the second material. This is the basis of the Maxwell-Garnett EMT (Maxwell Garnett, 1904). The resulting expression for m_{eff} is

$$m_{\text{eff}} = \sqrt{m_1^2 \frac{m_1^2(2 - 2f_2) + m_2^2(1 + 2f_2)}{m_1^2(2 + f_2) + m_2^2(1 - f_2)}}. \quad (\text{C1})$$

In the Bruggemann EMT (Bruggemann, 1935) one treats both materials more symmetrically; both components are assumed to be embedded in a host matrix with an effective refractive index given by

$$m_{\text{eff}} = \sqrt{\frac{1}{4}m_1^2(2 - 3f_2) + m_2^2(3f_2 - 1) + \sqrt{\frac{1}{16}[m_1^2(2 - 3f_2) + m_2^2(3f_2 - 1)]^2 + \frac{1}{2}m_1^2m_2^2}}. \quad (\text{C2})$$

Although not immediately manifest, this equation is symmetric under exchange of the two materials.

The volume-fraction is obtained from the mass concentrations M_1 and M_2 computed in the transport model, i.e. $f_2 = M_2/(M_1 + M_2)$. In SALSA, we apply the Maxwell-Garnett rule for an internal mixtures of mineral-dust inclusions in a host matrix of soluble compounds. Also, in the core-grey-shell model the effective refractive index of the grey shell, i.e., the homogeneous mixture of black carbon with soluble compounds, is computed with Maxwell-Garnett EMT. For mixtures of soluble compounds (sulphate, nitrate, ammonium, sea salt, organic compounds, and water) we use the Bruggemann EMT. If more than two components are mixed with each other, then the mixing rule is applied iteratively.

Appendix D: Refractive indices

The refractive indices that are used in the new optics model (and in the effective-medium calculations) are listed in Table 8.

Appendix E: Optical properties at different wavelengths in the considered case-studies

Figures 12–35 show vertical profiles of AOD, single-scattering albedo, and asymmetry parameter at the four geographic location, for the summer and winter incident, and for 12 different optical wavelengths.

Acknowledgements. E. Andersson acknowledge funding from the Swedish National Space Board within the OSCES project (No. 101/13). M. Kahnert has been funded by the Swedish Research Council (Vetenskapsrådet) within the AGES project (project 621-2011-3346).

Table 8. Refractive index m for each wavelength in the aerosol-optics look-up table and for various aerosol components.

λ [μm]	0.2000	0.2316	0.3040	0.3400
$m(\text{SO}_4)$	1.4840+0.1000E-07 i	1.4840+0.1000E-07 i	1.4676+0.1000E-07 i	1.4554+0.1000E-07 i
$m(\text{BC})$	0.9400+0.3500E+00 i	1.0717+0.5817E+00 i	1.3314+0.7523E+00 i	1.4471+0.7214E+00 i
$m(\text{OC})$	1.5300+0.5000E-02 i	1.5300+0.5000E-02 i	1.5300+0.5000E-02 i	1.5300+0.5000E-02 i
$m(\text{NaCl})$	1.5100+0.5000E-05 i	1.5100+0.5000E-05 i	1.5100+0.1866E-05 i	1.5100+0.6592E-06 i
$m(\text{Dust})$	1.5190+0.2070E-01 i	1.5190+0.2070E-01 i	1.5240+0.1947E-01 i	1.5272+0.1683E-01 i
$m(\text{H}_2\text{O})$	1.4517+0.1101E-06 i	1.4094+0.1092E-07 i	1.3701+0.3879E-08 i	1.3604+0.2758E-08 i
λ [μm]	0.3550	0.3800	0.3932	0.4400
$m(\text{SO}_4)$	1.4508+0.1000E-07 i	1.4448+0.1000E-07 i	1.4416+0.1000E-07 i	1.4336+0.1000E-07 i
$m(\text{BC})$	1.4954+0.7086E+00 i	1.5757+0.6871E+00 i	1.6181+0.6758E+00 i	1.6771+0.6586E+00 i
$m(\text{OC})$	1.5300+0.5000E-02 i	1.5300+0.5000E-02 i	1.5300+0.5000E-02 i	1.5300+0.5000E-02 i
$m(\text{NaCl})$	1.5090+0.2946E-06 i	1.5040+0.1476E-06 i	1.5014+0.6998E-07 i	1.5000+0.2544E-07 i
$m(\text{Dust})$	1.5239+0.1250E-01 i	1.5160+0.2500E-02 i	1.5147+0.2170E-02 i	1.5135+0.1400E-02 i
$m(\text{H}_2\text{O})$	1.3572+0.2416E-08 i	1.3528+0.1944E-08 i	1.3508+0.1702E-08 i	1.3449+0.9324E-09 i
λ [μm]	0.5000	0.5320	0.5332	0.6750
$m(\text{SO}_4)$	1.4310+0.1000E-07 i	1.4304+0.1000E-07 i	1.4303+0.1000E-07 i	1.4285+0.1860E-07 i
$m(\text{BC})$	1.7329+0.6414E+00 i	1.7626+0.6323E+00 i	1.7637+0.6319E+00 i	1.8097+0.5824E+00 i
$m(\text{OC})$	1.5300+0.5500E-02 i	1.5300+0.5500E-02 i	1.5300+0.5500E-02 i	1.5300+0.7091E-02 i
$m(\text{NaCl})$	1.5000+0.1550E-07 i	1.5000+0.1198E-07 i	1.5000+0.1185E-07 i	1.4900+0.1212E-06 i
$m(\text{Dust})$	1.5160+0.1200E-02 i	1.5167+0.1129E-02 i	1.5167+0.1126E-02 i	1.5170+0.9818E-03 i
$m(\text{H}_2\text{O})$	1.3394+0.9243E-09 i	1.3371+0.1818E-08 i	1.3370+0.1850E-08 i	1.3297+0.2187E-07 i
λ [μm]	0.7016	0.8700	1.0101	1.0200
$m(\text{SO}_4)$	1.4280+0.2214E-07 i	1.4253+0.2044E-06 i	1.4216+0.1749E-05 i	1.4213+0.1963E-05 i
$m(\text{BC})$	1.8175+0.5730E+00 i	1.8752+0.5645E+00 i	1.9210+0.5622E+00 i	1.9219+0.5643E+00 i
$m(\text{OC})$	1.5300+0.7333E-02 i	1.5300+0.9409E-02 i	1.5300+0.1327E-01 i	1.5300+0.1370E-01 i
$m(\text{NaCl})$	1.4900+0.2282E-06 i	1.4800+0.3027E-04 i	1.4700+0.1498E-03 i	1.4700+0.1584E-03 i
$m(\text{Dust})$	1.5170+0.9335E-03 i	1.5184+0.8000E-03 i	1.5190+0.7347E-03 i	1.5190+0.7261E-03 i
$m(\text{H}_2\text{O})$	1.3287+0.3624E-07 i	1.3243+0.3714E-06 i	1.3215+0.2657E-05 i	1.3213+0.2380E-05 i
λ [μm]	1.0640	1.2705	1.4625	1.7840
$m(\text{SO}_4)$	1.4197+0.2915E-05 i	1.4122+0.1621E-04 i	1.4045+0.1030E-03 i	1.3926+0.5308E-03 i
$m(\text{BC})$	1.9261+0.5738E+00 i	1.9457+0.6183E+00 i	1.9639+0.6597E+00 i	1.9943+0.7290E+00 i
$m(\text{OC})$	1.5285+0.1515E-01 i	1.5179+0.1721E-01 i	1.5068+0.1864E-01 i	1.4801+0.1337E-01 i
$m(\text{NaCl})$	1.4700+0.1966E-03 i	1.4692+0.3754E-03 i	1.4615+0.5382E-03 i	1.4500+0.7944E-03 i
$m(\text{Dust})$	1.5190+0.6853E-03 i	1.5188+0.6418E-03 i	1.5180+0.8000E-03 i	1.5180+0.9990E-03 i
$m(\text{H}_2\text{O})$	1.3205+0.1279E-05 i	1.3167+0.1090E-04 i	1.3128+0.3528E-03 i	1.3040+0.1270E-03 i
λ [μm]	2.0460	2.3250	2.7885	3.4615
$m(\text{SO}_4)$	1.3803+0.1490E-02 i	1.3580+0.2885E-02 i	1.3146+0.5669E-01 i	1.3669+0.1579E+00 i
$m(\text{BC})$	2.0192+0.7854E+00 i	2.0510+0.8453E+00 i	2.1099+0.9444E+00 i	2.1955+0.1088E+01 i
$m(\text{OC})$	1.4613+0.1000E-01 i	1.4554+0.9641E-02 i	1.4800+0.7724E-02 i	1.4800+0.7000E-02 i
$m(\text{NaCl})$	1.4482+0.1276E-02 i	1.4370+0.2950E-02 i	1.5339+0.7462E-02 i	1.4800+0.1757E-02 i
$m(\text{Dust})$	1.5180+0.1492E-02 i	1.5180+0.2610E-02 i	1.5180+0.8077E-02 i	1.5180+0.2805E-01 i
$m(\text{H}_2\text{O})$	1.2947+0.7103E-03 i	1.2756+0.5344E-03 i	1.1278+0.1055E+00 i	1.3913+0.1237E-01 i
λ [μm]	3.5000	8.0205	10.6000	12.1950
$m(\text{SO}_4)$	1.3760+0.1580E+00 i	1.1641+0.5511E+00 i	1.7200+0.3400E+00 i	1.7858+0.2517E+00 i
$m(\text{BC})$	2.2004+0.1097E+01 i	2.6572+0.1742E+01 i	2.9285+0.2063E+01 i	3.0719+0.2210E+01 i
$m(\text{OC})$	1.4800+0.7000E-02 i	1.1237+0.7906E-01 i	1.7600+0.7000E-01 i	1.6352+0.5117E-01 i
$m(\text{NaCl})$	1.4800+0.1600E-02 i	1.4080+0.1581E-01 i	1.5000+0.1400E-01 i	1.4383+0.1539E-01 i
$m(\text{Dust})$	1.5180+0.2973E-01 i	1.1798+0.1015E+00 i	1.9100+0.2500E+00 i	1.7614+0.4543E+00 i
$m(\text{H}_2\text{O})$	1.3840+0.9340E-02 i	1.2676+0.3436E-01 i	1.1531+0.7145E-01 i	1.0874+0.2243E+00 i




Figure 12. Aerosol optical depth over North Italy at 22-12-2007 for the 12 wavelengths in the CGS optics model and 5 of the 7 wavelengths in the EXT model. The wavelengths do not exactly overlap, but the EXT wavelengths that lies within 40 nm of the CGS wavelength are plotted in the same graph.




Figure 13. Same as Fig. 12, but over the Mediterranean.




Figure 14. Same as Fig. 12, but over Poland.




Figure 15. Same as Fig. 12, but over the North Sea




Figure 16. Same as Fig. 12, but 2007-06-22.




Figure 17. Same as Fig. 12, but 2007-06-22 and over the Mediterranean.




Figure 18. Same as Fig. 12, but 2007-06-22 and over Poland.




Figure 19. Same as Fig. 12, but 2007-06-22 and over the North sea.




Figure 20. Asymmetry parameter over North Italy at 22-12-2007 for the 12 wavelengths in the CGS optics model and 5 of the 7 wavelengths in the EXT model. The wavelengths do not exactly overlap, but the EXT wavelengths that lies within 40 nm of the CGS wavelength are plotted in the same graph.




Figure 21. Same as Fig. 20, but over the Mediterranean.




Figure 22. Same as Fig. 20, but over Poland.




Figure 23. Same as Fig. 20, but over the North sea.




Figure 24. Same as Fig. 20, but 2007-06-22.




Figure 25. Same as Fig. 20, but 2007-06-22 and over the Mediterranean.




Figure 26. Same as Fig. 20, but 2007-06-22 and over Poland.




Figure 27. Same as Fig. 20, but 2007-06-22 and over the North sea.




Figure 28. Single scattering albedo over North Italy at 22-12-2007 for the 12 wavelengths in the CGS optics model and 5 of the 7 wavelengths in the EXT model. The wavelengths do not exactly overlap, but the EXT wavelengths that lies within 40 nm of the CGS wavelength are plotted in the same graph.




Figure 29. Same as Fig. 28, but over the Mediterranean.




Figure 30. Same as Fig. 28, but over Poland.




Figure 31. Same as Fig. 28, but over the North sea.




Figure 32. Same as Fig. 28, but 2007-06-22.




Figure 33. Same as Fig. 28, but 2007-06-22 and over the Mediterranean.




Figure 34. Same as Fig. 28, but 2007-06-22 and over Poland.




Figure 35. Same as Fig. 28, but 2007-06-22 and over the North sea.

References

- Adachi, K. and Buseck, P. R.: Internally mixed soot, sulfates, and organic matter in aerosol particles from Mexico City, *Atmos. Chem. Phys.*, 8, 6469–6481, doi:10.5194/acp-8-6469-2008, 2008.
- Adachi, K., Chung, S. H., Friedrich, H., and Buseck, P. R.: Fractal parameters of individual soot particles
855 determined using electron tomography: Implications for optical properties, *J. Geophys. Res.*, 112, D14202, doi:10.1029/2006JD008296, 2007.
- Andersson, C., Langner, J., and Bergström, R.: Interannual variation and trends in air pollution over Europe due to climate variability during 1958–2001 simulated with a regional CTM coupled to the ERA40 reanalysis, *Tellus B*, 59, 77–98, 2007.
- 860 Andersson, C., Bergström, R., Bennet, C, Thomas, M., Robertson, L., Kokkola, H., Korhonen, H. and Lehtinen, K.: MATCH-SALSA - Multi-scale Atmospheric Transport and Chemistry model coupled to the SALSA aerosol microphysics model, SMHI Report RMK 115, available at: <http://www.smhi.se/publikationer/match-salsa-multi-scale-atmospheric-transport-and-chemistry-model-coupled-to-the-salsa-aerosol-microphysics-model-1.34623> (last accessed 10 February 2016), 2013.
- 865 Andersson, C., Bergström, R., Bennet, C., Robertson, L., Thomas, M., Korhonen, H., Lehtinen, K. E. J., and Kokkola, H.: MATCH-SALSA – Multi-scale Atmospheric Transport and CHEMISTRY model coupled to the SALSA aerosol microphysics model – Part 1: Model description and evaluation, *Geosci. Model Dev.*, 8, 171–189, doi:10.5194/gmd-8-171-2015, 2015.
- Benedetti, A., Morcrette, M. J.-J., Boucher, O., Dethof, A., Engelen, R. J., Huneeus, M. F. H. F. N., Jones, L.,
870 and S. Kinne, J. W. K., Mangold, A., Razinger, M., Simmons, A. J., and Suttie, M.: Aerosol analysis and forecast in the European Centre for Medium-Range Weather Forecasts Integrated Forecast System: 2. Data assimilation, *J. Geophys. Res.*, 114, D13205, doi:10.1029/2008JD011115, 2009.
- Bond, T. C. and Bergstrom, R. W.: Light absorption by carbonaceous particles: An investigative review, *Aerosol Sci. Tech.*, 40, 27–67, 2006.
- 875 Bruggemann, D. A. G.: Berechnung verschiedener physikalischer Konstanten von heterogenen Substanzen. 1. Dielektrizitätskonstanten und Leitfähigkeiten der Mischkörper aus isotropen Substanzen, *Ann. Phys.-Berlin*, 24, 636–664, 1935.
- China, S., Salvadori, N. and Mazzoleni, C.: Effect of Traffic and Driving Characteristics on Morphology of Atmospheric Soot Particles at Freeway On-Ramps, *Environ. Sci. Technol.*, 48 (6), 3128–3135,
880 doi:10.1021/es405178n, 2014.
- Chakrabarty, R.K., Beres, N.D., Moosmüller, H., China, S., Mazzoleni, C., Dubey, M.K., Liu, L., and Mishchenko, M.I.: Soot superaggregates from flaming wildfires and their direct radiative forcing, *Scientific Reports*, 4, 5508 EP-, doi:10.1038/srep05508, 2014.
- Chang, H. and Charalampopoulos, T. T.: Determination of the wavelength dependence of refractive indices of
885 flame soot, *P. R. Soc. A*, 430, 577–591, 1990.
- Dubovik, O., Sinyuk, A., Lapyonok, T., Holben, B. N., Mishchenko, M. I., Yang, P., Eck, T. F., Volten, H., Muñoz, O., Veihelmann, B., van der Zande, W. J., Leon, J.-F., Sorokin, M., and Slutsker, I.: Application of spheroid models to account for aerosol particle nonsphericity in remote sensing of desert dust, *J. Geophys. Res.*, 111, D11208, doi:10.1029/2005JD006619, 2006.

- 890 Foltescu, V., Pryor, S. C., and Bennet, C.: Sea salt generation, dispersion and removal on the regional scale, *Atmos. Environ.*, 39, 2123–2133, 2005.
- Gerber, H. E.: Relative-humidity parameterization of the Navy Aerosol Model (NAM), Tech. Rep. 8956, Naval Research Laboratory, Washington, DC, 1985.
- Jacobson, M. Z.: A physically-based treatment of elemental carbon optics: Implications for global direct forcing
895 of aerosols, *Geophys. Res. Lett.*, 27, 217–220, 2000.
- Jones, A. R.: Light scattering in combustion, in: *Light Scattering Reviews*, edited by: Kokhanovsky, A., Springer, Berlin, 393–444, 2006.
- Kahnert, F. M.: Reproducing the optical properties of fine desert dust aerosols using ensembles of simple model particles, *J. Quant. Spectrosc. Ra.*, 85, 231–249, 2004.
- 900 Kahnert, M.: Variational data analysis of aerosol species in a regional CTM: background error covariance constraint and aerosol optical observation operators, *Tellus B*, 60, 753–770, 2008.
- Kahnert, M.: Modelling the optical and radiative properties of freshly emitted light absorbing carbon within an atmospheric chemical transport model, *Atmos. Chem. Phys.*, 10, 1403–1416, doi:10.5194/acp-10-1403-2010, 2010a.
- 905 Kahnert, M.: On the discrepancy between modelled and measured mass absorption cross sections of light absorbing carbon aerosols, *Aerosol Sci. Tech.*, 44, 453–460, 2010b.
- Kahnert, M. and Devasthale, A.: Black carbon fractal morphology and short-wave radiative impact: a modelling study, *Atmos. Chem. Phys.*, 11, 11745–11759, doi:10.5194/acp-11-11745-2011, 2011.
- Kahnert, M., Nousiainen, T., and Lindqvist, H.: Models for integrated and differential scattering optical prop-
910 erties of encapsulated light absorbing carbon aggregates, *Opt. Express*, 21, 7974–7993, 2013.
- Kahnert, M., Nousiainen, T., and Lindqvist, H.: Review: Model particles in atmospheric optics, *J. Quant. Spectrosc. Ra.*, 146, 41–58, 2014.
- Khlebtsov, N. G.: Orientational averaging of light-scattering observables in the T-matrix approach, *Appl. Opt.*, 31, 5359–5365, 1992.
- 915 Klingmüller, K., Steil, B., Brühl, C., Tost, H., and Lelieveld, J.: Sensitivity of aerosol radiative effects to different mixing assumptions in the AEROPT 1.0 submodel of the EMAC atmospheric-chemistry-climate model, *Geosci. Model Dev.*, 7, 2503–2516, doi:10.5194/gmd-7-2503-2014, 2014.
- Kokkola, H., Korhonen, H., Lehtinen, K. E. J., Makkonen, R., Asmi, A., Järvenoja, S., Anttila, T., Partanen, A.-I., Kulmala, M., Järvinen, H., Laaksonen, A., and Kerminen, V.-M.: SALSA – a Sectional Aerosol module
920 for Large Scale Applications, *Atmos. Chem. Phys.*, 8, 2469–2483, doi:10.5194/acp-8-2469-2008, 2008.
- Kupiainen, K. and Klimont, Z.: Primary emissions of submicron and carbonaceous particles in Europe and the potential for their control, Tech. Rep. IR-04-079, IIASA, Laxenburg, Austria, 2004.
- Kupiainen, K. and Klimont, Z.: Primary emissions of fine carbonaceous particles in Europe, *Atmos. Environ.*, 41, 2156–2170, 2007.
- 925 Kylling, A., Bais, A. F., Blumthaler, M., Schreder, J., Zerefos, C. S., and Kosmidis, E.: The effect of aerosols on solar UV irradiances during the photochemical activity and solar ultraviolet radiation campaign, *J. Geophys. Res.*, 103, 26051–26060, 1998.

- Liu, Z., Liu, Q., Lin, H.-C., Schwartz, C. S., Lee, Y.-H., and Wang, T.: Three-dimensional variational assimilation of MODIS aerosol optical depth: Implementation and application to a dust storm over East Asia, *J. Geophys. Res.*, 116, D23206, doi:10.1029/2011JD016159, 2011.
- 930 Mackowski, D. W. and Mishchenko, M. I.: A multiple sphere T-matrix Fortran code for use on parallel computer clusters, *J. Quant. Spectrosc. Ra.*, 112, 2182–2192, 2011.
- Mårtensson, E. M., Nilsson, E. D., de Leeuw, G., Cohen, L. H., and Hansson, H.-C.: Laboratory simulations and parameterization of the primary marine aerosol production, *J. Geophys. Res.*, 108, 4297, doi:10.1029/2002JD002263, 2003.
- 935 Matsui, H., Koike, M., Kondo, Y., Moteki, N., Fast, J.D., and Zaveri, R.A.: Development and validation of a black carbon mixing state resolved three-dimensional model: Ageing processes and radiative impact, *J. Geophys. Res.*, 118, 2304–2326, doi:10.1029/2012JD018446, 2013.
- Maxwell Garnett, J. C.: Colours in metal glasses and in metallic films, *Philos. T. R. Soc. A*, 203, 385–420, 1904.
- 940 Mishchenko, M. I., Travis, L. D., and Lacis, A. A.: *Scattering, Absorption, and Emission of Light by Small Particles*, Cambridge University Press, Cambridge, 2002.
- Monahan, E. C., Spiel, D. E., and Davidson, K. L.: A model of marine aerosol generation via whitecaps and wave disruption, in: *Oceanic Whitecaps and their Role in Air-Sea Exchange*, edited by: Monahan, E. C. and Niocaill, G. M., D Reidel, Norwell, MA, 167–174, 1986.
- 945 Neusüß, C., Wex, H., Birmili, W., Wiedensohler, A., Koziar, C., Busch, B., Brüggemann, E., Gnauk, T., Ebert, M., and Covert, D. S.: Characterization and parameterization of atmospheric particle number-, mass-, and chemical-size distributions in central Europe during LACE 98 and MINT, *J. Geophys. Res.*, 107, LAC 9-1–LAC 9-13, doi:10.1029/2001JD000514, 2002.
- Nousiainen, T.: Optical modeling of mineral dust particles: a review, *J. Quant. Spectrosc. Ra.*, 110, 1261–1279, doi:10.1029/2009JD012611, 2009.
- 950 Nousiainen, T., Kahnert, M., and Veihelmann, B.: Light scattering modeling of small feldspar aerosol particles using polyhedral prisms and spheroids, *J. Quant. Spectrosc. Ra.*, 101, 471–487, 2006.
- Scarnato, B.V., Vahidinia1, S., Richard, D. T., and Kirchstetter, T. W.: Effects of internal mixing and aggregate morphology on optical properties of black carbon using a discrete dipole approximation model; *Atmos. Chem. Phys.*, 13, 5089–5101, doi:10.5194/acp-13-5089-2013, 2013.
- 955 Scarnato, B.V., China, S., Nielsen, K. and Mazzoleni, C.: Perturbations of the optical properties of mineral dust particles by mixing with black carbon: a numerical simulation study, *Atmos. Chem. Phys.*, 15, 6913–6928, doi:10.5194/acp-15-6913-2015, 2015.
- Robertson, L., Langner, J., and Engardt, M.: An Eulerian limited-area atmospheric transport model, *J. Appl. Meteorol.*, 38, 190–210, 1999.
- 960 Saide, P. E., Carmichael, G. R., Liu, Z., Schwartz, C. S., Lin, H. C., da Silva, A. M., and Hyer, E.: Aerosol optical depth assimilation for a size-resolved sectional model: impacts of observationally constrained, multi-wavelength and fine mode retrievals on regional scale analyses and forecasts, *Atmos. Chem. Phys.*, 13, 10425–10444, doi:10.5194/acp-13-10425-2013, 2013.
- 965 Undén, P., Rontu, L., Järvinen, H., Lynch, P., Calvo, J., Cats, G., Cuxart, J., Eerola, K., Fortelius, C., Garcia-Moya, J. A., Jones, C., Lenderink, G., McDonald, A., McGrath, R., Navascues, B., Nielsen, N. W., Ødegaard, V., Rodriguez, E., Rummukainen, M., Rööm, R., Sattler, K., Hansen Sass, B., Savijärvi, H., Schreur, B. W.,

Sigg, R., The, H., and Tijm, A.; HIRLAM-5 Scientific Documentation, available at <http://www.hirlam.org/index.php/documentation> (last accessed 2016-02-15), 2002.

970 Yurkin, M. A. and Hoekstra, A. G.: The discrete dipole approximation: an overview and recent developments, *J. Quant. Spectrosc. Ra.*, 106, 558–589, 2007.