

Response to Reviewer 1

This is not the venue for discussions about appropriate ... We agree with Reviewer 1 that the complexity of a model should always be questioned and so, justified. Adding processes and thus, complexity to a model is not a goal in itself. It should serve a specific scientific objective or should come from evidences that makes the representation of the different levels of complexity necessary. In that paper, we don't test all the different parameterizations that are included in PISCES. This is of course not feasible as it would require tens of experiments (probably even more since the model is not linear). Thus, we cannot assess the effect each individual parameterization has on the model results since it would require an enormous amount of work.

The complexity of PISCES can be partly explained by its history. This is a community model that has been developed and used over the last decade by different groups and for different purposes. Many parameterizations have been introduced during that period to serve these different needs and scientific objectives. Over time, the model has gained increasing levels of complexity which have been sometimes added in parallel. This means that at some state of the development of PISCES, some processes were tested independently of some other processes. One of the drawbacks of this approach is that the impact of these parameterizations on the model results can not be properly evaluated since biogeochemical models are highly non linear. Probably, something that will be required at some point of the model history is to reassess the usefulness of each class of processes in the light of the anticipated model utilization. We admit that this selection will be probably partly subjective since it will be based on the needs of a community which has very diverse scientific interests.

For the processes listed by Reviewer 1, we give here a justification for including them in the model and when known, the impact of those parameterizations on the model results is briefly explained.

acclimation of cells to local silicate (pg 1389) This parameterization was already present in PISCES-v1. The justification for using this parameterization was detailed in the submitted version of the manuscript on page 1389, lines 1-7. The impact of that parameterization on the model results was tested more than ten years ago and has never been re-tested since so that with the current version of the model, it is unknown how important the parameterization is.

stickiness of cells (pg 1389) The stickiness is not explicitly included in the parameterization. However, as explained in the submitted version of the manuscript, an increased stickiness would be equivalent to a higher aggregation coefficient. The scientific justification of that parameterization is given in the submitted version of the manuscript on page 1389, lines 12-15. The impact of this parameterization is rather strong when diatoms concentrations are high and are stressed. Higher aggregation coefficients lead to a more rapid decline of the blooms. The text has been modified in the revised version of the manuscript to further detail the impacts of that parameterization.

demands of iron for photosynthesis, respiration, ... The parameterization proposed by Flynn and Hipkin (1999) was adopted in PISCES as explained in the submitted version of the manuscript in page 1392, lines 5-9. This parameterization is fully detailed and tested in that paper.

variations in Si/C (pg 1393) The scientific justification for using that parameterization is quite extensively presented in pages 1392 to 1394 of the submitted version of the manuscript. It was already embedded in PISCES-v1 in a slightly simpler version. The silicate distribution is obviously highly sensitive to that parameterization since it directly controls the silicon utilization by diatoms.

oxygen effects on mortality (pg 1394) As explained in page 1394, lines 20-22, mortality of zooplankton is strongly increased in anoxic areas. This parameterization basically results in the exclusion of both microzooplankton and mesozooplankton from anoxic waters. This parameterization is also required to avoid respiration in waters with no oxygen. This justification has been added in the revised version of the manuscript at the bottom of page 1394 of the submitted version of the manuscript.

changing global grazing threshold (pg 1395) The choice of a grazing formulation is quite arbitrary as explained in the manuscript on page 1395, line 8. In PISCES, the value of the threshold is set in the namelist. For the reference simulation, we chose a value which gave realistic phytoplankton biomass in the oligotrophic areas. However, we did not extensively test the sensitivity of the results to that choice. We anticipate quite a strong sensitivity in the oligotrophic areas where phytoplankton concentrations are close to that threshold.

flux feeding (pg 1397) The impact of the flux feeding has been tested in (Gehlen et al., 2006). A reference to that study has been added in the revised version of the manuscript.

upper trophic levels (pg 1397) As explained in page 1397, lines 6-14, we adopted the formulation proposed by Anderson et al. (2013) to model the upper trophic levels. It is assumed that carnivores are ordered into an infinite chain. We have not tested the sensitivity of the model to that parameterization.

aggregation processes (pg 1400) As for flux feeding, these processes have been studied in (Gehlen et al., 2006). They were already included in PISCES-v1.

dissolution of silica in two phases (pg 1408) ... We agree with Reviewer 1 that since this process is quite uncertain, its inclusion in PISCES can be questioned. Nevertheless, an ongoing analysis comparing predicted biogenic silica profiles to observations show that this parameterization improves the agreement with observations (Marco van Hulst et al. (2015), in preparation).

variability of rain ratio (pg 1417) Reviewer 1 is right. The parameterization of the rain ratio is quite ad-hoc and we acknowledged this in the submitted version of the manuscript (see page 1418, lines 1-4). To answer the concern expressed by Reviewer 1, we have added in the revised version of the manuscript a statement which clearly explains that a good fit to the observation does not preclude that it is for wrong reasons.

dissolution of silica in guts (pg 1418) Reviewer 1 means probably calcite. The scientific justification of this parameterization is detailed in page 1418, lines 5-9. As for biogenic silica (see above), the analysis of the vertical distribution of calcite is currently performed in an on-going study.

There are some parameterizations not fully explained ... Several parameterizations are now better described, including the one quoted by Reviewer 1.

Some symbols are used for more than one thing ... We have changed several symbols that could be confusing. For instance, z is systematically used for depth whereas Z is restricted to zooplankton. t instead of T is used for time to avoid confusion with temperature. The symbol g now only defines grazing.

In the sensitivity experiments that are shown, it is not always clear ... The objective we had in mind when presenting these sensitivity tests was probably not clear enough in the submitted version of the manuscript. In PISCES-v2, we introduced some new formulations which represent new processes (SIZE, FOOD, PAR) or which propose an alternative description of an already existing process (LIGHT). To evaluate the sensitivity of the model results to these new parameterizations, we made a series of sensitivity experiments which are presented in section 7 of the manuscript. Our intention was not to unequivocally demonstrate that the new formulations produce better results but was rather to show the consequences of their utilization on the model behavior. In other words, these sensitivity tests were designed to illustrate the significance of the model solution to the proposed parameterizations. For instance, the assumption made for the light parameterization has profound impacts on the model behavior and therefore, this choice is not insignificant. On the other hand, the PAR experiment suggests that the model is rather insensitive to variations of the PAR fraction of shortwave radiation. Finally, the food quality is predicted to play an important role in the oligotrophic areas but not in the more productive areas. As our intention was not clear enough according to Reviewer 1, we have rewritten the introduction part of section 7 of the manuscript to better clarify the objectives of this section.

The parameterization of half saturation with cell-size is linear ... We agree with Reviewer 1 that the dependence of the half-saturation constant to cell size is not necessarily linear but may follow a power law function with an exponent different from 1. However, recent reviews give quite different values which range from about 0.84 to about 1 (Litchman et al., 2007; Edwards et al., 2012). Thus, as a conservative approach which can also be derived from simple volumetric considerations (surface-to-volume ratios), we assumed a linear relationship which remains within the range estimated from laboratory experiments. We have modified the text in section 4.1.1 around equations 7 in the revised version of the manuscript to better justify our choice.

Where numbers are used in equations, ... We now explain the numbers used in the equations when not obvious.

Relaxation of nutrients once a year (pg 1429, line 28) ... The sources and sinks of nutrients are not fully balanced in the ocean. As a consequence, the total inventory of these nutrients evolve with time. This may not be a problem when the model is run for only short experiments of the order of several decades but, in our case, the model has been run for 4000 years. Thus, a small drift can result in very strong changes in total inventories after this long integration. To overcome that problem, we chose to restore the total inventory of silicate, phosphate, alkalinity (we forgot to mention Alkalinity in the submitted version of the manuscript) and nitrate to the observed inventories computed from the World Ocean Atlas (for P, Si and N) and from GLODAP (for Alkalinity). In practice, this correction is done by scaling the 3-D concentrations with a constant uniform factor so that the total inventories match the observations. Thus, we do not restore the simulated 3-D distributions

to 3-D observed fields so that the predicted spatial and temporal patterns are not corrected to better match the observations. The correction is done once a year on the 1st of January (the frequency of this correction is specified in a namelist of the model). The correction is very small and corresponds to a relative change in the concentrations of the tracers of the order of 10^{-5} yr^{-1} . The last paragraph of section 6.1 has been substantially rewritten in the revised version of the manuscript to make this point less confusing. We now better detail the correction technique and indicate the magnitude of the correction.

Minimum threshold for Chl in the Pacific ... We think that Reviewer 1 refers to the minimum threshold below which Fe concentration can not decrease. Reviewer 2 had a similar concern and we address that concern in our response to Reviewer 2.

pg 1377, line 26 remove to be done.

pg 1378, line 1: do you mean internal? not external? Yes, we mean internal. We changed elemental to internal to make the text clearer.

pg 1380, line 10: explain IOM The meaning of IOM is now explained. See the responses to Reviewer 2.

pg 1381, line 3: I think you mean "nitrogen fixing" not "nutrient fixing" Reviewer 1 is right. We meant nitrogen fixing. This is corrected in the revised version of the manuscript.

pg 1382, lines 10-16: I don't follow the explanation here ... We mean that the N/P ratio in organic matter is constant and identical in all modeled organic pools. We make this point clearer in the text.

pg 1384: Eq 1 should it be P in the denominator Yes, Reviewer 1 is right. Equation has been modified accordingly.

pg 1384 (and onwards) - what is the currency in the main equations Most of the equations are in C units except when specified. This was detailed at the beginning of section 4 in page 1383, lines 20-25 of the submitted version of the manuscript. In the revised version of the manuscript, we added the explicit units used in the model.

pg 1384: which equation of 2 is used in the standard run Equation 2a is used in the standard experiment. This is now stated in the manuscript.

pg 1388: $\mu_{NO_3}^P$ needs clearer explanation of what this variable is We rewrote lines 1 to 3 of page 1388 to better explain what these parameters are. $\mu_{NO_3}^P$ is the net uptake rate of nitrate.

pg 1395: line 19 should be "with" done.

pg 1396, lines 7-9. I don't quite follow this ... Following Reviewer's 2 comment, this part has been partly rewritten. It should be now clearer. Reviewer 2 suggests to test that parameterization by a sensitivity study. This is exactly the objective of the FOOD experiment described in section 7.3.

pg 1397, line 9: grammar issues We are sorry but we are not able to detect these issues.

- pg 1398: Eq. 32 and line 6: use of N confusing here ...** N has been changed to I which makes this equation more consistent with previous equations too.
- pg 1400: line 8 - would be nice to have a reference for ...** This version of PISCES has never been published. We changed this line to precise that point.
- pg 1407: define sFe and bFe; also GOC turns up ...** GOC was defined in page 1400, line 22. We changed lines 4 to 6 in page 1406 to define SFe and bFe which are respectively the iron content of the small and big particles.
- pg 1425: What about iron deposition on top of ice?** We assumed in the model that dust deposition is not affected by sea ice. We recognize that this is quite a strong assumption which is not ideal. This will probably require some modifications of the model in the near future. We expect this assumption to have a small impact on the model results since high latitudes, especially in the Southern Ocean, are experiencing small deposition fluxes. This assumption is now mentioned in section 4.9.1 of the revised manuscript.
- pg 1426, line 3: would be nice to know how different results are with simple ...** We agree with Reviewer 1 that it would be interesting. However, testing the sediment model, which is not considered to be really part of the core of PISCES, was not the purpose of this study. Furthermore, the sediment model has been already described in Gehlen et al. (2006) and has not changed since.
- pg 1428, line 24: why GM only poleward of 10 degrees** In NEMO, the coefficient used in the GM parameterization is calculated from the local growth rate of baroclinic instability and is forced to decrease to vanish at the Equator following Tréguier et al. (1997).
- pg 1431, line 13-15: Doesn't this suggest grazing is too high?** Yes, Reviewer 1 is right. This might suggest that grazing by microzooplankton is too strong. However, the estimates are quite uncertain and thus, grazing remains poorly constrained. We modified the text in page 1431, lines 13-15.
- pg 1431, line 25: not sure why you use the word "Consequently" here** We remove this word from the text.
- Pg 1434, line 18: I believe the Dierssen and Smith (2000) algorithm is ...** Reviewer 1 is right. This algorithm has been constructed for the western Antarctic Peninsula waters and thus, its use over the whole Southern Ocean is questionable. Following his suggestion, we removed that part in the revised version of the manuscript.
- Pg 1436, line 25: Doesn't this suggest too few diatoms?** This does not necessarily suggest too few diatoms. this bias might also be due to a too low Si/C ratio in diatoms or a too strong dissolution of biogenic silica in the upper ocean.
- Pg 1437, line 18: DIC doesn't compare particularly well.** We agree with Reviewer 1 that mainly at the surface, the model does not compare very favorably to the GLODAP atlas. However, at the surface, observations include the contribution from anthropogenic carbon which is absent in the model since we set atmospheric CO₂ to its preindustrial value. To perform a more consistent comparison, we now compare the model output to DIC minus Anthropogenic DIC which are both available from GLODAP. This change improves the fit between the model and the observations.

Figure 12 has been changed as well as the Taylor plots on Figure 17 to account for the new dataset.

Pg 1443, line 14: As in point 4 above - are these results more realistic ... Please refer to our response to point 4.

Pg 1445, line 1: Do you mean "Chl" here ... Yes, we mean chlorophyll. This has been corrected in the revised version of the manuscript.

Pg 1445, line 9: grammar issues We are sorry but we are not able to correct this issue.

Pg 116, line 13: Not sure reliable is the right word here We changed reliable to reasonable.

Pg 1447, line 10: Grammar issues fixed.

Response to Reviewer 2

In many places, the authors describe an assumption using the word supposed ...

We have replaced supposed by assumed everywhere in the text.

Abstract: Replace silicium by silicon Done.

Page 3, suggest explicitly noting that balanced growth We are not sure to understand this comment. However, we clearly state now in the manuscript that Fe/C ratio varies greatly, in particular because of luxury uptake which is not the case for nitrogen.

Page 5 top, discussion of where PISCES is currently embedded PISCES is currently embedded and made available to the community in NEMO and ROMS. However, PISCES-v2 is only available in the NEMO system. This is now clearly stated in the text.

Page 5 bottom, note that is an overview of major changes ... This is now explained in the text.

Page 6 define IOM IOM is the new standard Input Output Manager of the NEMO modeling system. This is now defined in the text.

Page 13, middle, says nanophytoplankton, should say "diatom" in line 13 In fact, observations generally show that the mean cell size of the phytoplankton community tends to increase as the biomass increases. This is true for diatoms but also for other species.

Section 4.14 - What is the resulting range in Fe/C ratios or quotas? The Fe/C in PISCES varies between the minimum quota which is defined in equation 20 and the maximum quota which can be specified by the model user in the namelist. With the standard set of parameters, this quota varies between about 1 and 40 $\mu\text{mol Fe mol C}^{-1}$. This range is now mentioned in the text at the end of that section.

Section 4.15 - What is the resulting range in Si/C or Si quotas? The Si/C or Si quotas in diatoms vary between about 0 in the Si poor areas and more than 1 in the Southern Ocean. The global mean Si/C ratio in diatoms is predicted to be about 0.23 as mentioned in section 6.2.

For both of these are the general spatial patterns consistent with observations?

In the manuscript, we have not performed a validation of the Fe/C and Si/C ratios. Unfortunately, *in situ* data on the internal quotas of natural plankton communities are very sparse. Considering silicon, most of the observations report on the Si/C (or Si/N) ratio of the whole particulate materials or on the Si/N drawdown ratios which are only indirectly related to the quotas of diatoms. For instance, the Si/N drawdown ratio is mainly driven by the relative contribution of diatoms, the Si/N uptake ratios by diatoms, and by the recycling/dissolution efficiency of each nutrient. For iron, significant discrepancies exist between the different techniques used to measure the Fe/C or Fe/P quotas which lead to variations that span a 100-fold range, when applied to the same water samples (Twining et al., 2004; King et al., 2012). To overcome those difficulties, we made the basic assumption that the reproduction of the observed surface patterns of Si and Fe by PISCES should to some extent validate

the simulated temporal and spatial distribution of these quotas. This assumption may well be too strong since the surface distributions of Si and Fe depend on many other dynamical and biogeochemical processes, such as vertical advection, mixing, dissolution/reminerization, etc ...

Section 4.3 - Is DOM kept at the same quasi-Redfield ratio 122/16/1? Yes, the DOM pools are assigned the quasi-Redfield ratio 122/16/1 as explained in lines 7-9, page 24.

page 47 several typos in the references listed We are sorry but we have not found any typos in this page.

Page 48 seems somewhat problematic that a key limiting nutrient like Fe is not ...

The reviewer is right. Fe is not conserved with respect to sea ice. New ice is formed with an iron concentration equal to that of the underlying sea water but is melting with an iron concentration of 10 nM. Conserving iron would have required to model iron in the sea ice which was out of the scope of our work. Instead, we have based our very simple parameterization on the results of the work of Lancelot et al. (2009) as explained in section 4.9. In PISCES, iron is overall a net source of iron because the iron concentration in melting sea ice is much larger than the iron concentration in forming sea ice. Overall, the source is equal to 0.024 Gmol Fe yr⁻¹, several orders of magnitude less than from the sediments, that is 34 Gmol Fe yr⁻¹. We have added those two numbers in sections 4.9.5 and 4.9.3 respectively.

Page 52 - Could the authors also give the export flux at 100 m depth We decided not to chose the depth of 100 m because primary production and thus, the production of organic matter can still be substantial below that horizon, in particular in the oligotrophic subtropical gyres. Furthermore, in the literature, very diverse depths are chosen ranging generally from 100m to 200m, which means that there is no general consensus on a reference depth. However, to follow the reviewer's suggestion, we have added a footnote in Table 4 which gives the vertical flux of sinking particles at 100m depth, 7.3 GtC yr⁻¹.

Page 55 - To be clear, there is an arbitrary iron source that kicks in ... Yes, the reviewer is right. Iron concentration can not decrease below 0.01 nM, which is equivalent to adding a source of iron to prevent iron to decrease below that threshold. The impact of this assumption is discussed in Tagliabue et al. (2009) and Tagliabue and Voelker (2011) as mentioned in the manuscript page 60. We re-copy here the corresponding section in Tagliabue et al. (2009) which addresses this issue: *In the previous version of PISCES [Aumont and Bopp, 2006], a minimum Fe threshold (of 0.01nM) was necessary to accurately represent the biogeochemical dynamics of the equatorial Pacific region. If this parameterization is removed, all the Fe that is upwelled in the eastern equatorial Pacific is utilized close to the upwelling sites and the phytoplankton bloom cannot propagate westward (compare Figure 5a with Figures 5b and 5c). Therefore, such a parameterization (which represents an additional, yet unconstrained, Fe source) was deemed necessary to accurately represent the distribution of tropical Pacific chlorophyll a, especially in spring, as well as the transition between Fe and NO₃ limitation [see Aumont and Bopp, 2006].* In the revised version of the manuscript, we now also refer to the figure 5 shown in Tagliabue et al. (2009).

page 63 top, charlot ? This is a typo. We meant limiting.

page 67, I not that the N₂O cycling is not described in the primary literature ...

The description of the N₂O module was left for another study planned by a PhD student, J. Martinez-Rey. This study is now under review in Biogeosciences Discussion. It is now properly quoted in the manuscript.

Page 67, line 29, please revise this sentence We are sorry but we don't really see what should be changed in that sentence.

If there is one global metric I'd suggest adding to the paper it would be ... We

agree with Reviewer 2 that diagnostics on the OMZ characteristics are indeed very interesting because of the important role these regions play both for marine biogeochemistry and biology. On figure 1, we display the thickness of the OMZs as predicted by PISCES and as observed in the World Ocean Atlas 2009. The model does not a very good job at simulating both the location and the vertical extent of the OMZs. In fact, this result was anticipated since the oxygen distribution is extremely sensitive to both export production and the dynamics, especially the ventilation rates of the mode, intermediate and deep water masses. Previous modeling studies have shown that coarse resolution ocean biogeochemical models exhibit serious biases, especially in the tropical Pacific and Atlantic oceans (Moore et al., 2013; Bopp et al., 2013). In the latter ocean basin, models simulate a significant OMZ whereas concentrations below 20 μM are rarely observed. Experiments performed with high resolution models suggest that part of those deficiencies may be due to a poor representation of the equatorial zonal jets (Dietze and Loeptien, 2013; Getzlaff and Dietze, 2013). Thus, since OMZ distribution reflects the characteristics of marine biogeochemistry as much as ocean dynamics, we made the choice not to show and discuss that figure in the submitted version of the manuscript. If Reviewer 2 recommends to include that figure in the revised version of the manuscript, we would do so.

How is the routing of Fe handled when the prey Fe/C is lower ... This is described in equation 27b. When the Fe/C ratio is below the Fe/C ratio of zooplankton, the gross growth efficiency of zooplankton is decreased such that the extra carbon (and nutrients) ingested by zooplankton is routed to organic and inorganic loss processes. This is now clearly explained in section 4.2.1. There was although an error in equation 27b which has been corrected in the revised version of the manuscript.

In general, the text/labeling within figures is too small We have remade the figures and increased the size of the labels.

Figure 10, suggest adding at least one more low end contour done.

Figure 11. definitely needs more resolution/contours in the color scale on the low end, ...
The figure has been redone following reviewer's advice.

References

Bopp, L., Resplandy, L., Orr, J., Doney, S., Dunne, J., Gehlen, M., Halloran, P., Heinze, C., Ilyina, T., Séférian, R., et al.: Multiple stressors of ocean ecosystems in the 21st century: projections with CMIP5 models, *Biogeosciences*, 10, 6225–6245, 2013.




Figure 1: Annual mean thickness (in meters) of the Oxygen Minimum Zones as observed in the World Ocean Atlas 2009 (a) and as simulated by PISCES (b). Oxygen Minimum Zones are defined here as oxygen concentrations below $20 \mu\text{M}$.

- Dietze, H. and Loeptien, U.: Revisiting “nutrient trapping” in global coupled biogeochemical ocean circulation models, *Global Biogeochemical Cycles*, 27, 265–284, doi:10.1002/gbc.20029, 2013.
- Edwards, K. F., Thomas, M. K., Klausmeier, C. A., and Litchman, E.: Allometric scaling and taxonomic variation in nutrient utilization traits and maximum growth rate of phytoplankton, *Limnol. Oceanogr*, 57, 554–566, 2012.
- Gehlen, M., Bopp, L., Emprin, N., Aumont, O., Heinze, C., Kriest, I., and Ragueneau, O.: Reconciling surface ocean productivity, export fluxes and sediment composition in a global biogeochemical ocean model, *Biogeosciences*, 3, 521–537, 2006.
- Getzlaff, J. and Dietze, H.: Effects of increased isopycnal diffusivity mimicking the unresolved equatorial intermediate current system in an earth system climate model, *Geophysical Research Letters*, 40, 2166–2170, doi:10.1002/grl.50419, 2013.
- King, A. L., Sañudo Wilhelmy, S. A., Boyd, P. W., Twining, B. S., Wilhelm, S. W., Breene, C., Ellwood, M. J., and Hutchins, D. A.: A comparison of biogenic iron quotas during a diatom spring bloom using multiple approaches, *Biogeosciences*, 9, 667–687, doi:10.5194/bg-9-667-2012, 2012.
- Litchman, E., Klausmeier, C. A., Schofield, O. M., and Falkowski, P. G.: The role of functional traits and trade-offs in structuring phytoplankton communities: scaling from cellular to ecosystem level, *Ecology Letters*, 10, 1170–1181, doi:10.1111/j.1461-0248.2007.01117.x, 2007.
- Moore, J. K., Lindsay, K., Doney, S. C., Long, M. C., and Misumi, K.: Marine Ecosystem Dynamics and Biogeochemical Cycling in the Community Earth System Model [CESM1(BGC)]: Comparison of the 1990s with the 2090s under the RCP4.5 and RCP8.5 Scenarios, *Journal of Climate*, 26, 9291–9312, doi:10.1175/JCLI-D-12-00566.1, 2013.
- Twining, B. S., Baines, S. B., Fisher, N. S., and Landry, M. R.: Cellular iron contents of plankton during the Southern Ocean Iron Experiment (SOFeX), *Deep Sea Research Part I: Oceanographic Research Papers*, 51, 1827–1850, doi:10.1016/j.dsr.2004.08.007, 2004.

Aumont Ethé Tagliabue Bopp Gehlen
Manuscript prepared for Geosci. Model Dev. Discuss.
with version 4.1 of the L^AT_EX class copernicus_discussions.cls.
Date: 29 June 2015

PISCES-v2: an ocean biogeochemical model for carbon and ecosystem studies

O. Aumont¹, C. Ethé², A. Tagliabue³, L. Bopp⁴, and M. Gehlen⁴

¹Laboratoire d'Océanographie et de Climatologie: Expérimentation et Approches Numériques, IPSL, 4 Place Jussieu, 75005 Paris, France

²Institut Pierre et Simon Laplace, 4 Place Jussieu, 75005 Paris, France

³Dept. of Earth, Ocean and Ecological Sciences, School of Environmental Sciences, University of Liverpool, 4 Brownlow Street, Liverpool L69 3GP, UK

⁴Laboratoire des Sciences du Climat et de l'Environnement, IPSL, Orme des Merisiers, 91190 Gif-sur-Yvette, France

Correspondence to: O. Aumont (olivier.aumont@ird.fr)

Abstract

PISCES-v2 is a biogeochemical model which simulates the lower trophic levels of marine ecosystem (phytoplankton, microzooplankton and mesozooplankton) and the biogeochemical cycles of carbon and of the main nutrients (P, N, Fe, and Si). The model is intended to be used for both regional and global configurations at high or low spatial resolutions as well as for short-term (seasonal, interannual) and long-term (climate change, paleoceanography) analyses. There are twenty-four prognostic variables (tracers) including two phytoplankton compartments (diatoms and nanophytoplankton), two zooplankton size-classes (microzooplankton and mesozooplankton) and a description of the carbonate chemistry. Formulations in PISCES-v2 are based on a mixed Monod–Quota formalism: on one hand, stoichiometry of C/N/P is fixed and growth rate of phytoplankton is limited by the external availability in N, P and Si. On the other hand, the iron and ~~silicium~~ silicon quotas are variable and growth rate of phytoplankton is limited by the internal availability in Fe. Various parameterizations can be activated in PISCES-v2, setting for instance the complexity of iron chemistry or the description of particulate organic materials. So far, PISCES-v2 has been coupled to the NEMO and ROMS systems. A full description of PISCES-v2 and of its optional functionalities is provided here. The results of a quasi-steady state simulation are presented and evaluated against diverse observational and satellite-derived data. Finally, some of the new functionalities of PISCES-v2 are tested in a series of sensitivity experiments.

1 Introduction

Human activities have released large amounts of carbon into the atmosphere since the beginning of the industrial era leading to an increase in atmospheric CO₂ by more than 100 ppmv. The oceans play a major role in the carbon cycle and in its adjustment. [Sabine et al. \(2004\)](#) have estimated that the oceans have absorbed about one third of the anthropogenic emissions. This role is tightly controlled by the physical and biogeochemical states of the marine system, i.e. by the characteristics of the solubility and biological pumps. Yet, the role played by the ocean in

the carbon cycle is likely to be modified in response to climate and chemical changes induced by the anthropogenic carbon emissions (e.g., Orr et al., 2005; Steinacher et al., 2010; Bopp et al., 2013). Global ocean biogeochemical models represent powerful tools to study the carbon cycle and to predict its response to future and past climate and chemical changes. Since the pioneering work by Bacastow and Maier-Reimer (1990) based on a very simple description of the carbon cycle, the number and the complexity of models have rapidly increased (e.g., Six and Maier-Reimer, 1996; Moore et al., 2004; Quéré et al., 2005; Aumont and Bopp, 2006; Yool et al., 2011). However, a greater complexity of the models raises difficulties related to the lack of data for validation and to the theoretical justification of the parameterizations (e.g., Anderson, 2005, 2010).

PISCES is a biogeochemical model which simulates the marine biological productivity and that describes the biogeochemical cycles of carbon and of the main nutrients (P, N, Si, Fe). This model can be seen as one of the many Monod models (Monod, 1942) as opposed to the quota models (McCarthy, 1980; Droop, 1983) which are alternative kind of ocean biogeochemical models. Thus, it assumes a constant Redfield ratio and phytoplankton growth depends on the external concentration in nutrients. This choice was dictated by the computing cost as describing the internal pools of the different elements (necessary for a quota model) requires many more prognostic variables. Ultimately, PISCES was ~~supposed~~ assumed to be suited for a wide range of spatial and temporal scales, including typically several thousand years-long simulations on the global scale.

In contrast to the Monod approach, when modeling silicate, iron and/or chlorophyll, assuming constant ratios is not justified anymore as these ratios can vary substantially. For instance, the Fe/C ratio can vary by at least an order of magnitude (~~e.g., Sunda and Huntsman, 1995, 1997~~) to be, in particular as a result of luxury uptake, (e.g., Sunda and Huntsman, 1995, 1997) compared to the N/C ratio which varies by “only” two to three times. Equally, the Si/C ratio can vary significantly in response to the degree of iron stress (Hutchins and Bruland, 1998; Takeda, 1998). Thus, in PISCES, a compromise between the two classical types of ocean model was chosen. The Fe/C, Si/C and Chl/C ~~elemental~~ internal ratios are prognostically predicted based on the external concentrations of the limiting nutrients as in the quota approach. Phytoplankton

growth rates are predicted using simultaneously the Monod approach for N, P and Si and the quota approach for Fe. As a consequence, PISCES should be considered to be a mixed Monod-Quota model.

Historically, the development of PISCES started in 1997 with the release of the P3ZD model which was a simple NPZD model with semi-labile DOM (Aumont, 1998; Aumont et al., 2002). Phytoplankton growth rate was only limited by one nutrient, effectively phosphate and many shortcomings were apparent in this model, especially in the HNLC regions. This served to justify the development, beginning in 1999, of a more complex model that includes three limiting nutrients (Fe, Si, P), two phytoplankton and two zooplankton size-classes. This model was called HAMOCC5 (Aumont et al., 2003) as it was based on HAMOCC3.1 (Six and Maier-Reimer, 1996) and used in the LSG model (Maier-Reimer et al., 1993). When this code was embedded in the ocean model OPA (Madec et al., 1998), it required some major changes and improvements, partly because of the much finer vertical resolution. In addition to the numerical schemes, these changes were mostly an improved treatment of the optics and the separation of the particulate organic matter into two different size-classes. All these changes and the major recordings it required led us to adopt a new name for the model: PISCES. This name can be translated as fishes from Latin. It can also be considered as the following acronym: Pelagic Interactions Scheme for Carbon and Ecosystem Studies.

PISCES has been used so far to address a wide range of scientific questions. Unfortunately, a complete list of the studies which have been based on or made use of PISCES is not available but more than about hundred referenced studies explicitly rely directly or indirectly on this model. These range from process studies (Aumont and Bopp, 2006; Gehlen et al., 2006; Tagliabue et al., 2009a; Tagliabue and Völker, 2011) to operational oceanography (Brasseur et al., 2009). PISCES has been used to analyse intraseasonal (Gorgues et al., 2005; Resplandy et al., 2009) to interannual and decadal timescales (Raynaud et al., 2006; Rodgers et al., 2008). PISCES is part of the IPSL and CNRM Earth-System Models which contribute to the different IPCC-related activities including the CMIP5 modeling component (Séférian et al., 2013). Several studies have been conducted that consider the potential impact of climate change on ocean biogeochemistry (Dufresne et al., 2002; Bopp et al., 2005; Steinacher et al., 2010). Modeling studies focusing

on paleoceanography have been based on PISCES (Bopp et al., 2003; Tagliabue et al., 2009b). Finally, PISCES is also used in regional configurations to study specific regions such as the Peru upwelling (Echevin et al., 2008; Albert et al., 2010) or the Indian ocean (Resplandy et al., 2012).

PISCES is currently embedded into two modeling systems: NEMO (Madec, 2008) and ROMS_AGRIF (Penven et al., 2006; Debreu et al., 2011). It can be downloaded from their respective websites:

- <http://www.nemo-ocean.eu> for the NEMO ocean modeling framework
- <http://www.romsagrif.org> for the ROMS_AGRIF modeling framework.

However, PISCES-v2 is currently available only in the NEMO modeling system. The implementation of this updated version of PISCES in the ROMS_AGRIF modeling system is currently underway and should be finished and available by the end of 2015.

Since 2001, PISCES has undergone active developments. In 2004, a stable release of the model was made available to the community on the OPA website. Soon after, an earlier documentation of the model was published as Supplement to the study by Aumont and Bopp (2006). Since then, the model has significantly evolved without any update of the documentation and this has effectively rendered the earlier documentation obsolete. After six years of intense developments, it is more than appropriate at this point to provide the current or future users of the model with an updated and accurate description of the current state of PISCES, called PISCES-v2. The following document describes the main aspects of the model. At its end, a description of a climatological simulation is proposed using the standard set of parameters available when the model is downloaded. Finally, the impact of several new parameterizations is evaluated through the performance of a set of sensitivity experiments.

2 Changes from previous release

As already mentioned, PISCES as a research tool is in perpetual evolution. Numerous changes have been made relative to the previously documented version PISCES-v1. A brief list of the

main changes is made below, with these changes organized thematically. [These changes are detailed in the following sections.](#)

– Changes made to the code structure and design:

1. Transition to full native Fortran 90 coding. The model has also undergone a reorganization of its architecture and coding conventions following the evolution of NEMO.
2. I/O interface should now be set by default to IOM ([the new Input Output Manager of the NEMO modeling system](#)) to benefit from the major improvements this interface offers.
3. Memory and performance improvements have been made. This version should run slightly faster and take much less memory than v1.
4. The namelist now includes many more parameters that may thus be changed without recompiling the code.

– Changes made to the nutrients:

1. Iron chemistry can be described according to two different parameterizations: the simple old chemistry scheme based on one ligand and one inorganic species, and a new complex chemistry module based on five iron species and two ligands.
2. Scavenging of inorganic iron and coagulation of iron colloids have been redesigned.

– Changes made to the phytoplankton compartments:

1. Nutrients limitation terms now include a simple description of the impact of cell size.
2. Iron content and growth rate limitation by iron is modeled following the quota formalism. Luxury uptake of iron can be represented by this new formulation.
3. Redesign of silicification, calcification ~~and nutrient~~ [as well as nitrogen](#) fixation by diazotrophs.
4. The relationship between growth rate (primary production) and light can be chosen between two different formulations.

- Changes made to the zooplankton compartments:
 1. The microzooplankton grazing formulation is now identical to that of mesozooplankton.
 2. Thresholds can be selected for both total food or individual prey types.
 3. Food quality affects the gross growth efficiency of both zooplankton compartments.
- Changes made to dissolved organic matter and particulate materials:
 1. Two different schemes for the description of particulate organic matter can be chosen: the traditional two-compartment model or the Kriest model.
 2. Bacterial implicit description has been redesigned.
 3. Dissolution of biogenic silica assumes two different fractions.
 4. The dust distribution in the water column is modeled using a very crude parameterization.
 5. The numerics of vertical sedimentation has been improved (time splitting scheme).
- Changes made to the external sources of nutrients and to the treatment of the bottom of the water column:
 1. Spatially variable solubility of iron in dust can be specified from a file.
 2. River discharge of nutrients has been improved.
 3. Denitrification in sediments is now parameterized as well as variable preservation of calcite.

As a consequence of these changes, the user should be warned that results produced with PISCES-v1 cannot be reproduced by PISCES-v2. Furthermore, in the rest of this work, PISCES will designate PISCES-v2.

3 Model description

PISCES currently has twenty-four compartments (see Fig. 1). There are five modeled limiting nutrients for phytoplankton growth: Nitrate and Ammonium, Phosphate, Silicate and Iron. It should be mentioned that Phosphate and Nitrate+Ammonium are not really ~~independent~~ independent nutrients in PISCES. They are linked by a constant ~~Redfield-ratio-and identical~~ Redfield ratio in all the modeled organic compartments but the nitrogen pool undergoes nitrogen fixation and denitrification in the open ocean and the upper sediments. Furthermore, their external sources (rivers, dust deposition) are not linked by a constant ratio. This means that if the latter three processes (nitrogen fixation, denitrification, and external sources) are deactivated and if the initial distributions of Nitrate+Ammonium and Phosphate are identical, the simulated fields of both nutrients should remain identical.

Four living compartments are represented: two phytoplankton size classes/groups corresponding to nanophytoplankton and diatoms, and two zooplankton size classes which are microzooplankton and mesozooplankton. For phytoplankton, the prognostic variables are the carbon, iron, chlorophyll and silicon biomasses (the latter only for diatoms). This means that the Fe/C and Chl/C ratios of both phytoplankton groups as well as the Si/C ratio of diatoms are prognostically predicted by the model. For zooplankton, only the total biomass is modeled. For all species, the C/N/P/O₂ ratios are ~~supposed~~ assumed constant and are not allowed to vary. In PISCES, the Redfield ratios C/N/P are set to ~~122/16~~ 122/16/1 (Takahashi et al., 1985) and the -O/C ratio is set to 1.34 (Kortzinger et al., 2001). In addition, the Fe/C ratio of both zooplankton groups is kept constant. No silicified zooplankton is assumed. The bacterial pool is not yet explicitly modeled.

There are three non-living compartments: semi-labile dissolved organic matter, small sinking particles, and large sinking particles. As for the living compartments, the C, N and P pools are not distinctly modeled. Thus, constant Redfield ratios are imposed for C/N/P. On the other hand, the iron, silicon and calcite pools of the particles are explicitly modeled. As a consequence, their ratios are allowed to vary. The sinking speed of the particles is not altered by their content in calcite and biogenic silicate (“The ballast effect”, Honjo, 1996; Armstrong et al.,

2002). The latter particles are assumed to sink at the same speed as the large organic matter particles. An earlier version of PISCES had included a simple description of this “ballast effect” (Gehlen et al., 2006) but it has been abandoned since as observations don’t suggest a clear relationship between sinking speeds and mineral composition of particles (Lee et al., 2009). All the non-living compartments experience aggregation due to turbulence and differential settling as well as Brownian coagulation for DOM.

In addition to the ecosystem model, PISCES also simulates dissolved inorganic carbon, total alkalinity and dissolved oxygen. The latter tracer is also used to define the regions where oxic or anoxic degradation processes take place.

4 Model equations

The reader should be aware that in the following equations, the conversion ratios between the different elements (Redfield ratios) have been generally omitted except when particular parameterizations are defined. All phytoplankton and zooplankton biomasses are in carbon units (mol C L^{-1}) except for the silicon, chlorophyll and iron content of phytoplankton, which are respectively in Si, Chl and Fe units (mol Si L^{-1} , g Chl L^{-1} , and mol Fe L^{-1}). Finally, all parameters and their standard values in PISCES are listed in Tables 1a–1e at the end of this section.

4.1 Phytoplankton

4.1.1 Nanophytoplankton

$$\frac{\partial P}{\partial t} = (1 - \delta^P)\mu^P P - m^P \frac{P}{K_m + N} \frac{P}{K_m + P} P - \text{sh} \times w^P P^2 - g^Z(P)Z - g^M(P)M \quad (1)$$

In this equation, P is the nanophytoplankton biomass, and the 5 terms on the right-hand side represent growth, mortality, aggregation, and grazing by micro- and mesozooplankton, respectively. The mortality term is modulated by a hyperbolic function of P to avoid extinction of nanophytoplankton at very low growth rates.

In PISCES, the growth rate of nanophytoplankton (μ^P) can be computed according to two different parameterizations:

$$\mu^P = \mu_P f_1(L_{\text{day}}) \underline{g} \underline{f}_2(\underline{Z} z_{\text{mxl}}) \left(1 - \exp \left(\frac{-\alpha^P \theta^{\text{Chl}, P} \text{PAR}^P}{L_{\text{day}} (\mu_{\text{ref}} + b_{\text{resp}})} \right) \right) L_{\text{lim}}^P \quad (2a)$$

$$\mu^P = \mu_P f_1(L_{\text{day}}) \underline{g} \underline{f}_2(\underline{Z} z_{\text{mxl}}) \left(1 - \exp \left(\frac{-\alpha^P \theta^{\text{Chl}, P} \text{PAR}^P}{L_{\text{day}} \mu_P L_{\text{lim}}^P} \right) \right) L_{\text{lim}}^P \quad (2b)$$

where b_{resp} is a small respiration rate and μ_{ref} a reference growth rate, independent of temperature. All other terms in these equations are defined below. The choice between the two different formulations is made through a parameter in the namelist (`ln_newprod`). When `ln_newprod` is set to true, Eq. (2a) which is the default option of PISCES, Equation 2a is used. In the previous equations, L_{day} is day length ($\in [0, 1]$). $\underline{f}(L_{\text{day}}) \underline{f}_1(L_{\text{day}})$ expresses the dependency of growth rate to the length of the day (Gilstad and Sakshaug, 1990; Thompson, 1999). $\underline{Z} z_{\text{mxl}}$ is the depth of the mixed layer and $\underline{g}(\underline{Z} z_{\text{mxl}}) \underline{f}_2(z_{\text{mxl}})$ imposes an additional reduction of the growth rate when the mixed layer depth exceeds the euphotic depth:

$$\underline{f}_1(L_{\text{day}}) = 1.5 \frac{L_{\text{day}}}{0.5 + L_{\text{day}}} \quad (3a)$$

$$\underline{\Delta} \underline{Z} z = \max \left(0, \underline{Z} z_{\text{mxl}} - \underline{Z} z_{\text{eu}} \right) \quad (3b)$$

$$\underline{T} t_{\text{dark}} = (\underline{\Delta} \underline{Z} z)^2 / 86400 \quad (3c)$$

$$\underline{g} \underline{f}_2(\underline{Z} z_{\text{mxl}}) = 1 - \frac{\underline{T} t_{\text{dark}}}{\underline{T}_{\text{dark}}^P + \underline{T}_{\text{dark}}} \frac{t_{\text{dark}}}{\underline{t}_{\text{dark}}^P + t_{\text{dark}}} \quad (3d)$$

where $\underline{Z} z_{\text{eu}}$ is the depth of the euphotic zone defined as the depth at which there is 1% of surface PAR. $\underline{T}_{\text{dark}}^P \underline{t}_{\text{dark}}^P$ is set to 3 days for nanophytoplankton and 4 days for diatoms, as diatoms generally better cope with prolonged dark periods. $\underline{T}_{\text{dark}} \underline{t}_{\text{dark}}$ is an estimate of the mean

residence time of the phytoplankton cells within the unlit part of the mixed layer, assuming a vertical diffusion coefficient of $1 \text{ m}^2 \text{ s}^{-1}$. [86400 converts \$t_{\text{dark}}\$ from \$\text{s}^{-1}\$ to \$\text{day}^{-1}\$](#) . Figure 2 displays $g(z_{\text{mix}}) \cdot f_2(z_{\text{mix}})$ as a function of $\Delta Z \Delta z$.

μ_P is defined as follows (Eppley, 1972):

$$5 \quad f_P(T) = b_P^T \quad (4a)$$

$$\mu_P = \mu_{\text{max}}^0 f_P(T) \quad (4b)$$

In PISCES, vertical penetration of the Photosynthetic Available Radiation (PAR) is based on a simplified version of the model by Morel (1988), which is described in Lengaigne et al. (2007). Visible light is split into three wavebands: blue (400–500 nm), green (500–600 nm) and red (600–700 nm). For each waveband, the chlorophyll-dependent attenuation coefficients are fitted to the coefficients computed from the full spectral model of Morel (1988) (as modified in Morel and Maritorena, 2001) assuming the same power-law expression. At the sea surface, visible light is split equally between the three wavebands. PAR can be a constant or a variable fraction of the downwelling shortwave radiation, as specified in the namelist (`ln_varpar`).

$$\text{PAR}_1(0) = \text{PAR}_2(0) = \text{PAR}_3(0) = \frac{\rho_{\text{par}}}{3} \text{SW} \quad (5a)$$

$$\text{PAR}^P(z) = \beta_1^P \text{PAR}_1(z) + \beta_2^P \text{PAR}_2(z) + \beta_3^P \text{PAR}_3(z) \quad (5b)$$

Light absorption by phytoplankton depends on the waveband and on the species. The normalized coefficients β_i have been computed for each phytoplankton group by averaging and normalizing, for each waveband, the absorption coefficients published in Bricaud et al. (1995).

In PISCES, the nutrient limitation terms are defined as follows:

$$L_{\text{lim}}^P = \min(L_{\text{PO}_4}^P, L_{\text{N}}^P, L_{\text{Fe}}^P) \quad (6a)$$

$$L_{\text{PO}_4}^P = \frac{\text{PO}_4}{\text{PO}_4 + K_{\text{PO}_4}^P} \quad (6b)$$

$$25 \quad L_{\text{N}}^P = L_{\text{NO}_3}^P + L_{\text{NH}_4}^P \quad (6c)$$

$$L_{\text{NH}_4}^P = \frac{K_{\text{NO}_3}^P \text{NH}_4}{K_{\text{NO}_3}^P K_{\text{NH}_4}^P + K_{\text{NH}_4}^P \text{NO}_3 + K_{\text{NO}_3}^P \text{NH}_4} \quad (6d)$$

$$L_{\text{NO}_3}^P = \frac{K_{\text{NH}_4}^P \text{NO}_3}{K_{\text{NO}_3}^P K_{\text{NH}_4}^P + K_{\text{NH}_4}^P \text{NO}_3 + K_{\text{NO}_3}^P \text{NH}_4} \quad (6e)$$

$$L_{\text{Fe}}^P = \min \left(1, \max \left(0, \frac{\theta^{\text{Fe},P} - \theta_{\min}^{\text{Fe},P}}{\theta_{\text{opt}}^{\text{Fe},P}} \right) \right) \quad (6f)$$

5 As already stated in the introduction, PISCES is a mixed Monod–Quota model. Thus, N and P limitations are based on a Monod parameterization where growth depends on the external nutrient concentrations whereas Fe limitation is modeled according to a classical Quota approach. It should be noted here that for iron, an optimal quota ($\theta_{\text{opt}}^{\text{Fe},P}$) is used in the denominator which allows luxury uptake as in the model proposed by [Buitenhuis and Geider \(2010\)](#).

10 The choice of the half-saturation constants is rather difficult as observations show that they can vary by several orders of magnitude (e. g., [Perry, 1976](#); [Sommer, 1986](#); [Donald et al., 1997](#)). However, in general, these constants increase with the size of the phytoplankton cell as a consequence of a smaller surface-to-volume ratio (diffusive hypothesis) ([Eppley et al., 1969](#)). Thus, diatoms will tend to have larger half-saturation constants than nanophytoplankton.

15 However, in PISCES, phytoplankton are modeled by only two compartments, each of them encompassing a large range. Experiments performed with the model have shown that results are sensitive to the choice of these half-saturation constants.

Following these remarks, it appeared not appropriate to keep constant the nutrient half-saturation constants. It was then decided to make them vary with the phytoplankton biomass of each compartment because the observations show that the increase in biomass is generally due to the addition of larger size classes of phytoplankton (e.g., [Raimbault et al., 1988](#); [Armstrong, 1994](#); [Hurt and Armstrong, 1996](#)):

$$P_1 = \min(P, P_{\max}) \quad (7a)$$

$$P_2 = \max(0, P - P_{\max}) \quad (7b)$$

$$K_i^P = K_i^{P,\min} \frac{P_1 + S_{\text{rat}}^P P_2}{P_1 + P_2} \quad (7c)$$

where S_{rat}^P is the size ratio of the larger size class over the smaller size class. $K_i^{P,\min}$ is the half-saturation constant of the smaller size class. This parameterization assumes that half-saturation constants **increases-increase** linearly with size (Eppley et al., 1969). The size dependence of these constants with cell size is not necessarily linear but has been suggested to follow a power law function with an exponent lower than 1 (Litchman et al., 2007). However, in a recent review, Edwards et al. (2012) found an exponent close to 1 for nitrogen (linear relationship) and larger than 1 for phosphorus. Thus, considering these uncertainties, we decided to keep a linear relationship, which remains within the estimated range and which can also be derived from simple volumetric considerations (surface-to-volume ratio). The three parameters in this equation (P_{max} , $K_i^{P,\min}$, and S_{rat}^P) can be independently specified for each phytoplankton group. Finally, observations also suggest that these half-saturation constants should vary with the mean nutrient concentrations, probably as an acclimation to the local environment (Collos et al., 1980; Smith et al., 2009). This acclimation mechanism is not included in PISCES, except for the case of Silicate (see Sect. 4.1.2).

The distinction between new production based on nitrate and regenerated production based on ammonium is computed as follows (O'Neill et al., 1989):

$$\mu_{\text{NO}_3}^P = \mu^P \frac{L_{\text{NO}_3}^P}{L_{\text{NO}_3}^P + L_{\text{NH}_4}^P} \quad \mu_{\text{NH}_4}^P = \mu^P \frac{L_{\text{NH}_4}^P}{L_{\text{NO}_3}^P + L_{\text{NH}_4}^P} \quad (8)$$

where $\mu_{\text{NO}_3}^P$ and $\mu_{\text{NH}_4}^P$ are the uptake rates of nitrate and ammonium respectively.

The nanophytoplankton aggregation term w^P depends on the shear rate sh as the main driver of aggregation is the local turbulence. This shear rate is set to 1 s^{-1} in the mixed layer and to 0.01 s^{-1} below. This means that the aggregation is reduced by a factor of 100 below the mixed layer.

4.1.2 Diatoms

$$\frac{\partial D}{\partial t} = (1 - \delta^D)\mu^D D - m^D \frac{D}{K_m + D} D - \text{sh} \times w^D D^2 - g^Z(D)Z - g^M(D)M \quad (9)$$

In this equation, D is the nanophytoplankton biomass, and the 5 terms on the right-hand side represent growth, mortality, aggregation, and grazing by micro- and mesozooplankton, respectively.

As for nanophytoplankton, the absorption coefficients of diatoms depend on the considered waveband:

$$\text{PAR}^D = \beta_1^D \text{PAR}_1 + \beta_2^D \text{PAR}_2 + \beta_3^D \text{PAR}_3 \quad (10)$$

The production terms for diatoms are defined as for nanophytoplankton except that the limitation terms also include Si:

$$L_{\text{lim}}^D = \min(L_{\text{PO}_4}^D, L_{\text{N}}^D, L_{\text{Fe}}^D, L_{\text{Si}}^D) \quad (11a)$$

$$L_{\text{Si}}^D = \frac{\text{Si}}{\text{Si} + K_{\text{Si}}^D} \quad (11b)$$

As for the other nutrients, the half-saturation factor of Silicate can vary significantly over the ocean. In the tropical and temperate regions, this factor is around $1 \mu\text{M}$ whereas values as high as $88.7 \mu\text{M}$ have been measured for Antarctic species (Sommer, 1986; Martin-Jézéquel et al., 2000). In that case, rather than an effect of the cell size, these variations are a consequence of an acclimation of the cells to their local environment. When plotted against maximum local yearly concentration of silicate, a crude relationship can be inferred (Pondaven et al., 1998):

$$K_{\text{Si}}^D = K_{\text{Si}}^{D,\text{min}} + \frac{7\check{\text{Si}}^2}{(K_{\text{Si}})^2 + \check{\text{Si}}^2} \quad (12)$$

where \check{S}_i here is the maximum S_i concentration over a year (note that during the first year of a pluriannual simulation, \check{S}_i is set to a constant). For the other nutrients, we use the same parameterization as for nanophytoplankton (see Eq. 7).

The diatoms aggregation term w_p^D is increased in case of nutrient limitation because it has been shown that diatoms cells tend to excrete a mucus (exocellular polysaccharides, EPS) which increases their stickiness. As a consequence, collisions between cells yield to a more efficient aggregation process (Smetacek, 1985; Decho, 1990):

$$w^D = w^P + w_{\max}^D (1 - L_{\lim}^D) \quad (13)$$

Furthermore as for nanophytoplankton, the aggregation is multiplied by the shear rate. [Enhanced aggregation rates when diatoms are stressed result in a rapid decline of the diatoms blooms when nutrients become exhausted and produce strong export events.](#)

4.1.3 Chlorophyll in nanophytoplankton and diatoms

Chlorophyll biomass I^{Chl} (where I denotes P or D , typical units are $\mu\text{g Chl L}^{-1}$ or mg Chl m^{-3}) for both phytoplankton groups is parameterized using the photoadaptive model of Geider et al. (1997):

$$\begin{aligned} \frac{\partial I^{\text{Chl}}}{\partial t} = & (1 - \delta^I) (12\theta_{\min}^{\text{Chl}} + (\theta_{\max}^{\text{Chl},I} - \theta_{\min}^{\text{Chl}}) \rho^{I^{\text{Chl}}}) \mu^I I - m^I \frac{I}{K_m + I} I^{\text{Chl}} \\ & - \text{sh} \times w^I I I^{\text{Chl}} - \theta^{\text{Chl},I} g^Z(I) Z - \theta^{\text{Chl},I} g^M(I) M \end{aligned} \quad (14)$$

where I is the phytoplankton group and $\theta^{\text{Chl},I}$ is the chlorophyll-to-carbon ratio of the considered phytoplankton class. [12 represents the molar mass of carbon.](#) $\rho^{I^{\text{Chl}}}$ represents the ratio of energy assimilated to energy absorbed as defined by Geider et al. (1996):

$$\rho^{I^{\text{Chl}}} = \frac{144 \check{\mu}^I I}{\alpha^I I^{\text{Chl}} \frac{\text{PAR}^I}{L_{\text{day}}}} \quad (15a)$$

$$\check{\mu}^I = \mu_P \underline{g} \underline{f}_2(Z_{\text{mxl}}) \left(1 - \exp \left(\frac{-\alpha^I \theta^{\text{Chl},I} \text{PAR}^I}{L_{\text{day}} \mu_P L_{\text{lim}}^I} \right) \right) L_{\text{lim}}^I \quad (15b)$$

In this equation, 144 is the square of the molar mass of C and is used to convert from mol to mg as the standard unit for Chl is generally in mg Chl m⁻³. It should be noted that for chlorophyll synthesis, the second parameterization of phytoplankton growth is used to compute $\check{\mu}^I$ (see Eq. 2b). This is necessary because of the expression for ρ_{Chl}^I .

4.1.4 Iron in nanophytoplankton and diatoms

The temporal evolution of the iron biomass of phytoplankton I^{Fe} (model units are mol Fe L⁻¹), where I denotes P or D , is driven by the following equation:

$$\frac{\partial I^{\text{Fe}}}{\partial t} = (1 - \delta^I) \mu^{I^{\text{Fe}}} I - m^I \frac{I}{K_m + I} I^{\text{Fe}} - \text{sh} \times w^I I I^{\text{Fe}} - \theta^{\text{Fe},I} g^Z(I) Z - \theta^{\text{Fe},I} g^M(I) M \quad (16)$$

Iron in phytoplankton is modeled in PISCES according to a classical quota approach. However, to be consistent with chlorophyll and silica, we model the iron biomass of phytoplankton (I^{Fe}) rather than the iron quota ($\theta^{\text{Fe},I}$) directly. Growth rate of the iron biomass of phytoplankton is parameterized according to:

$$\mu^{I^{\text{Fe}}} = \theta_{\text{max}}^{\text{Fe},I} L_{\text{lim},1}^{I^{\text{Fe}}} L_{\text{lim},2}^{I^{\text{Fe}}} \frac{1 - \frac{\theta^{\text{Fe},I}}{\theta_{\text{max}}^{\text{Fe},I}}}{1.05 - \frac{\theta^{\text{Fe},I}}{\theta_{\text{max}}^{\text{Fe},I}}} \mu_P \quad (17)$$

As in Flynn and Hipkin (1999), Iron uptake is also downregulated via a feedback from $\theta^{\text{Fe},I}$ using a normalized inverse hyperbolic function with a small shape factor set to 0.05.

In the former equation, $L_{\text{lim},1}^{I^{\text{Fe}}}$ is the iron limitation term and is modeled as follows:

$$L_{\text{lim},1}^{I^{\text{Fe}}} = \frac{\text{bFe}}{\text{bFe} + K_{\text{Fe}}^{I^{\text{Fe}}}} \quad (18a)$$

$$K_{\text{Fe}}^{I^{\text{Fe}}} = K_{\text{Fe}}^{I^{\text{Fe},\text{min}}} \frac{I_1 + S_{\text{rat}}^I I_2}{I_1 + I_2} \quad (18\text{b})$$

$$I_2 = \max(0, I - I_{\text{max}}), I_1 = I - I_2 \quad (18\text{c})$$

where $b\text{Fe}$ is the concentration of bioavailable iron (see Sect. 4.5.3) The half-saturation constant for iron uptake is also increasing with phytoplankton biomass as for the other half-saturation constants (see Eq. 7).

At low iron concentrations, observations suggest that iron uptake might be enhanced, at least for some species (Harrison and Morel, 1986; Doucette and Harrison, 1991), giving surge uptake. Morel (1987) proposed a parameterization of both this surge uptake and the downregulation of iron uptake at high iron quota (see above) which has been included in the recent model of Buitenhuis and Geider (2010). In PISCES, a different parameterization has been chosen since downregulation is already included in Eq. (17):

$$L_{\text{lim},2}^{I^{\text{Fe}}} = \frac{4 - 4.5 L_{\text{Fe}}^I}{L_{\text{Fe}}^I + 0.5} \quad (19)$$

$L_{\text{lim},2}$ equals 4 at very low iron concentrations and 1 at high iron concentration. Overall, the downregulation in Eq. (17) together with the surge uptake induced by the previous equation results in a behavior of the system that is qualitatively equivalent to what results from the parameterization of Buitenhuis and Geider (2010).

The demands for iron in phytoplankton are for photosynthesis, respiration and nitrate/nitrite reduction. Following Flynn and Hipkin (1999), we assume that the rate of synthesis by the cell of new components requiring iron is given by the difference between the iron quota and the sum of the iron required by these three sources of demand, which we defined as the actual minimum iron quota:

$$\theta_{\text{min}}^{\text{Fe},I} = \frac{0.0016}{55.85} \theta^{\text{Chl},I} + \frac{1.21 \times 10^{-5} \times 14}{55.85 \times 7.625} L_{\text{N}}^P \times 1.5 + \frac{1.15 \times 10^{-4} \times 14}{55.85 \times 7.625} L_{\text{NO}_3}^P \quad (20)$$

In this equation, the first right term corresponds to photosynthesis, the second term corresponds to respiration and the third term estimates nitrate and nitrite reduction. The parameters used

in this equation are directly taken from Flynn and Hipkin (1999). The modeled iron quota in PISCES varies thus between this minimum quota $\theta_{\min}^{\text{Fe},I}$ and the maximum quota $\theta_{\max}^{\text{Fe},I}$, i.e. between about 1 and 40 $\mu\text{mol Fe (mol C)}^{-1}$ when using the standard set of parameters (see Table 1a).

5 4.1.5 Silicon in diatoms

$$\begin{aligned} \frac{\partial D^{\text{Si}}}{\partial t} = & \theta_{\text{opt}}^{\text{Si},D} (1 - \delta^D) \mu^D D - \theta^{\text{Si},D} g^M(D) M - \theta^{\text{Si},D} g^Z(D) Z - m^D \frac{D}{K_m + D} D^{\text{Si}} \\ & - \text{sh} \times w^D D D^{\text{Si}} \end{aligned} \quad (21)$$

The elemental ratio Si/C (or Si/N) has been observed to vary by a factor of about 4 to 5 over the global ocean with a mean value around $0.14 \pm 0.13 \text{ mol mol}^{-1}$ (Sarthou et al., 2005). Light, N, P, or Fe stress has been demonstrated to lead to heavier silicification (e.g., Takeda, 1998; Franck et al., 2000; Martin-Jézéquel et al., 2000). It has been suggested that these elevated elemental ratios result from the physiological adaptation of the silicon uptake by the cell depending on the growth rate and on the G2 cycle phase during which Si is incorporated (Martin-Jézéquel et al., 2000; Claquin et al., 2002). Lighter silicification can only result from silicate limitation.

We model the variations of the Si/C ratio following the parameterization proposed by Bucciarelli et al. (2002, unpublished manuscript):

$$\theta_{\text{opt}}^{\text{Si},D} = \theta_{\text{m}}^{\text{Si},D} L_{\text{lim},1}^{D^{\text{Si}}} \min \left(5.4, \left(4.4 \exp \left(-4.23 F_{\text{lim},1}^{D^{\text{Si}}} \right) F_{\text{lim},2}^{D^{\text{Si}}} + 1 \right) \left(1 + 2 L_{\text{lim},2}^{D^{\text{Si}}} \right) \right) \quad (22)$$

Relative to the original parameterization, an additional limitation term by Si has been added ($F_{\text{lim},2}^{D^{\text{Si}}}$) to produce a lighter silicification in case of Si exhaustion.

The different terms in Eq. (22) are defined as follows:

$$F_{\text{lim},1}^{D^{\text{Si}}} = \min \left(\frac{\mu^D}{\mu_P L_{\text{lim}}^D}, L_{\text{PO}_4}^D, L_{\text{N}}^D, L_{\text{Fe}}^D \right) \quad (23a)$$

$$F_{\text{lim},2}^{D^{\text{Si}}} = \min \left(1, 2.2 \max \left(0, L_{\text{lim},1}^{D^{\text{Si}}} - 0.5 \right) \right) \quad (23b)$$

$$L_{\text{lim},1}^{D\text{Si}} = \frac{\text{Si}}{\text{Si} + K_{\text{Si}}^{-1}} \quad (23\text{c})$$

$$L_{\text{lim},2}^{D\text{Si}} = \begin{cases} \frac{\text{Si}^3}{\text{Si}^3 + (K_{\text{Si}}^2)^3} & \text{if } \varphi < 0 \\ 0 & \text{if } \varphi > 0 \end{cases} \quad (23\text{d})$$

where φ is the latitude. In the Southern Ocean, observations show that diatoms are very heavily silicified. After correcting for the potential effects of iron limitation, silicification in the Southern ocean is at least three times stronger than in the tropical regions, which can only be explained by the diatoms morphological types (Baines et al., 2010). To reproduce those high Si/C ratios, we have introduced the term $L_{\text{lim},2}^{D\text{Si}}$ which increases the Si/C ratio by a factor of up to 3 when silicate concentrations are high, a specific characteristics of the Southern Ocean. This increase is restricted to the Southern Hemisphere and is controlled by the parameter K_{Si}^2 . This parameter is set in the namelist and thus, if it is set to a very high value, then no increase of Si/C at high silicate concentrations is predicted by the model.

4.2 Zooplankton

4.2.1 Microzooplankton

$$\frac{\partial Z}{\partial t} = e^Z (g^Z(P) + g^Z(D) + g^Z(\text{POC})) Z - g^M(Z)M - m^Z f_Z(T) Z^2 - r^Z f_Z(T) \left(\frac{Z}{K_m + Z} + 3\Delta(\text{O}_2) \right) Z \quad (24)$$

In this equation, Z is the microzooplankton biomass, and the 4 terms on the right-hand side represent growth, grazing by mesozooplankton, quadratic and linear mortalities, respectively.

The grazing rate depends on temperature according to a typical exponential relationship similar to what is used for phytoplankton:

$$g_m^Z = g_{\text{max}}^{0,Z} f_Z(T) \quad (25\text{a})$$

$$f_Z(T) = b_Z^T \quad (25b)$$

5 where $g_{\max}^{0,Z}$ is the maximum grazing rate at 0°C , b_Z is the temperature dependence and T is the temperature. In their review, [Buitenhuis et al. \(2010\)](#) have found a Q_{10} ($Q_{10} = b_Z^{10}$) between 1.7 and 2.2. Lower temperature dependences were found in laboratory experiments compared to what as been identified in the field. In PISCES, we have set Q_{10} to 2.14 which is close to the value found in the field but also close to the value chosen for mesozooplankton (see below). All terms driving the temporal evolution of microzooplankton have been assigned the same temperature dependence. Mortality is enhanced when oxygen is depleted. In other words, microzooplankton (but also mesozooplankton, see below) are treated as being unable to cope with anoxic waters. [This increased mortality also avoids respiration in waters devoid of oxygen.](#)

15 Grazing on each species I is defined as:

$$F = \sum_J p_J^Z \max(0, J - J_{\text{thresh}}^Z)$$

$$F_{\text{lim}} = \max(0, F - \min(0.5F, F_{\text{thresh}}^Z))$$

$$g^Z(I) = g_m^Z \frac{F_{\text{lim}} p_I^Z \max(0, I - I_{\text{thresh}}^Z)}{F K_G^Z + \sum_J p_J^Z J} \quad (26a)$$

20 where J denotes all the species microzooplankton can graze upon (P , D , and POC) and p_J^Z is the preference microzooplankton has for each J . In PISCES, we have chosen a Michaelis–Menten parameterization with no switching and a threshold (F_{thresh}^Z) ([Gentleman et al., 2003](#)). This choice is rather arbitrary. Another very popular formulation in models is the Michaelis–Menten parameterization with active switching introduced by [Fasham et al. \(1990\)](#). However, this parameterization exhibits anomalous dynamics such as sub-optimal feeding ([Gentleman et al., 2003](#)). In our parameterization, a threshold for each individual resource (J_{thresh}^Z) can be specified in addition to the global threshold (F_{thresh}^Z). For low food abundance, this global threshold is allowed to slowly decrease to 0 as a function of the total food level to maintain some grazing pressure, in particular in the ocean interior.

Responses of zooplankton to quality of their preys have been termed stoichiometric modulation of predation (SMP) by [Mitra and Flynn \(2005\)](#). A complete review of the different expected responses has been presented by [Mitra et al. \(2007\)](#). For instance, when confronted to [with](#) poor food quality, zooplankton can increase their ingestion rate ([Plath and Boersma, 2001](#); [Darchambeau and Thys, 2005](#)), or decrease it as the food can become deleterious ([Flynn and Davidson, 1993](#)). Accounting for the complexities of these different types of behavior has not been implemented within PISCES as this would require a model with flexible stoichiometry. Additionally, it would require a correct parameterization of the different potential responses and the apparently contradictory nature of observed responses implies that this task will be very complicated. In PISCES, food quality is [supposed-assumed](#) to only affect gross growth efficiency (e^Z): When food quality becomes poor (either the Fe/C ratio $\theta^{\text{Fe},I}$ or the N/C ratio $\theta^{\text{N},I}$ of the preys decreases), e^Z decreases:

$$e_N^Z = \min \left(1, \frac{\sum_I \theta^{\text{N},I} g^Z(I)}{\theta^{\text{N},C} \sum_I g^Z(I)}, \frac{\sum_I \theta^{\text{Fe},I} g^Z(I)}{\theta^{\text{Fe},Z} \sum_I g^Z(I)} \right) \quad (27a)$$

$$e^Z = e_N^Z \min \left(e_{\max}^Z, (1 - \sigma^Z) \frac{\min \left(\frac{\sum_I \theta^{\text{N},I} g^Z(I)}{\theta^{\text{N},C} \sum_I g^Z(I)}, \frac{\sum_I \theta^{\text{Fe},I} g^Z(I)}{\theta^{\text{Fe},Z} \sum_I g^Z(I)} \right)}{1} \right) \quad (27b)$$

When the Fe/C ratio of the ingested preys becomes lower than the zooplankton Fe/C ratio, the excess carbon (and nutrients) is lost as dissolved inorganic and organic carbon (and nutrients). This is described in PISCES by a decrease in the carbon gross growth efficiency (Equation 27b). By construction in PISCES, the N/C quota is constant, so this quota is estimated by solving the classical Droop equation assuming that it is at steady state (see above the definition of $\theta^{\text{N},I}$).

4.2.2 Mesozooplankton

$$\begin{aligned} \frac{\partial M}{\partial t} = & e^M (g^M(P) + g^M(D) + g^M(\text{POC}) + g_{\text{FF}}^M(\text{GOC}) + g_{\text{FF}}^M(\text{POC}) + g^M(Z)) M \\ & - m^M f_M(T) M^2 - r^M f_M(T) \left(\frac{M}{K_m + M} + 3\Delta(\text{O}_2) \right) M \end{aligned} \quad (28)$$

In this equation, M is the mesozooplankton biomass, and the 3 terms on the right-hand side represent growth, quadratic and linear mortalities, respectively. All terms in this equation have been assigned the same temperature dependence using a Q_{10} of 2.14 (Buitenhuis et al., 2005).

Parameterization of mesozooplankton grazing is similar to microzooplankton. In addition to the “conventional” concentration-dependent grazing described by Eq. (26), flux-feeding is also accounted for in PISCES. This type of grazing has been shown to be potentially very important for the fate of particles in the water column below the euphotic zone (Dilling and Alldredge, 2000; Stemmann et al., 2004). Flux feeding depends on the flux and thus, on the product of the concentration by the sinking speed. In PISCES, both the small and the large particles experience this type of grazing:

$$g_{\text{FF}}^M(\text{POC}) = g_{\text{FF}} f_M(T) w_{\text{POC}} \text{POC} \quad (29\text{a})$$

$$g_{\text{FF}}^M(\text{GOC}) = g_{\text{FF}} f_M(T) w_{\text{GOC}} \text{GOC} \quad (29\text{b})$$

This importance of flux feeding has been analyzed in PISCES by Gehlen et al. (2006). They have shown that flux feeding is the most important process that controls the flux of particulate organic carbon below the surface mixed layer.

In Eq. (28), the term with a quadratic dependency to mesozooplankton does not depict aggregation but grazing by the higher, non-resolved trophic levels. Following Anderson et al. (2013), the upper trophic levels are modeled assuming an infinite chain of carnivores. This assumption permits to easily compute the production of fecal pellets as well as the respiration and excretion by these non-resolved carnivores:

$$P_{\text{up}}^M = \sigma^M f_{\text{up}}(e_{\text{max}}^M) m^M f_M(T) M^2 \quad (30\text{a})$$

$$R_{\text{up}}^M = (1 - \sigma^M - e_{\text{max}}^M) f_{\text{up}}(e_{\text{max}}^M) m^M f_M(T) M^2 \quad (30\text{b})$$

where function $f_{\text{up}}(x)$ is:

$$f_{\text{up}}(x) = \sum_{i=0}^{\infty} x^i = \frac{1}{1-x} \quad \text{for } 0 < x < 1 \quad (31)$$

It should be noted here that a similar quadratic term is also included in the equation for microzooplankton (see Eq. 26) despite the fact that their predators are (at least partially) represented in PISCES. In that case, this term rather represents other density-dependent mortality factors such as viral diseases. As a consequence, the assumption of an infinite chain of carnivores is not used for microzooplankton and everything is routed to POC.

4.3 DOC

The temporal evolution of DOC is driven by the following equation:

$$\begin{aligned} \frac{\partial \text{DOC}}{\partial t} = & (1 - \gamma^Z)(1 - e^Z - \sigma^Z) \sum \underline{NI} g^Z(\underline{NI}) Z + (1 - \gamma^M)(1 - e^M - \sigma^M) \\ & \left(\sum \underline{NI} g^M Z(\underline{NI}) + g_{\text{FF}}^M(\text{GOC}) \right) M + \delta^D \mu^D D + \delta^P \mu^P P + \lambda_{\text{POC}}^* \text{POC} \\ & + (1 - \gamma^M) R_{\text{up}}^M - \text{Remin} - \text{Denit} - \Phi_1^{\text{DOC}} - \Phi_2^{\text{DOC}} - \Phi_3^{\text{DOC}} \end{aligned} \quad (32)$$

where \underline{NI} includes P , D and POC for microzooplankton and P , D , Z , and POC for mesozooplankton (see Eqs. 24 and 28, respectively). In the following, DOM and DOC will be used indifferently since the stoichiometric ratios in dissolved organic matter are supposed assumed constant in PISCES.

Marine DOM has traditionally been divided into several fractions characterized by their lability. DOM, which recycles over timescales of a few months to a few years, is called semi-labile DOM (Anderson and Williams, 1999). Transport of this pool of dissolved organic matter can make a significant part of the carbon pump (Carlson et al., 1994; Anderson and Williams, 1999). As a consequence, this important pool of DOM is modeled in PISCES. The labile and refractory pools of DOM are not explicitly modeled.

The degradation of semi-labile DOC is parameterized as follows:

$$\text{Remin} = \min \left(\frac{\text{O}_2}{\text{O}_2^{\text{ut}}}, \lambda_{\text{DOC}} f_P(T) (1 - \Delta(\text{O}_2)) L_{\text{lim}}^{\text{bact}} \frac{\text{Bact}}{\text{Bact}_{\text{ref}}} \text{DOC} \right) \quad (33a)$$

$$\text{Denit} = \min \left(\frac{\text{NO}_3}{r_{\text{NO}_3}^*}, \lambda_{\text{DOC}} f_P(T) \Delta(\text{O}_2) L_{\text{lim}}^{\text{bact}} \frac{\text{Bact}}{\text{Bact}_{\text{ref}}} \text{DOC} \right) \quad (33b)$$

Remineralization of DOC can be either oxic (Remin) or anoxic (Denit) depending on the local oxygen concentration. The distinction between the two types of organic matter degradation is performed using a factor $\Delta(\text{O}_2)$ that varies between 0 and 1 (see Sect. 4.5.1 for the formulation of this factor). It is assumed that the specific rates of degradation (λ_{DOC}) specified for respiration and denitrification are identical.

Depending on the quality of the organic matter, bacteria may take up nutrients from seawater (e.g., Goldman and Dennett, 1991; Thingstad and Lignell, 1997), and thus may be limited by their availability. Of course, bacterial production is also limited by the abundance of dissolved organic matter. Therefore, we parameterize the regulation of the degradation of DOM by bacterial activity (L^{bact}) according to:

$$L^{\text{bact}} = L_{\text{lim}}^{\text{bact}} L_{\text{DOC}}^{\text{bact}} \quad (34a)$$

$$L_{\text{DOC}}^{\text{bact}} = \frac{\text{DOC}}{\text{DOC} + K_{\text{DOC}}} \quad (34b)$$

$$L_{\text{lim}}^{\text{bact}} = \min(L_{\text{NH}_4}^{\text{bact}}, L_{\text{PO}_4}^{\text{bact}}, L_{\text{Fe}}^{\text{bact}}) \quad (34c)$$

$$L_{\text{Fe}}^{\text{bact}} = \frac{\text{bFe}}{\text{bFe} + K_{\text{Fe}}^{\text{bact}}} \quad (34d)$$

$$L_{\text{PO}_4}^{\text{bact}} = \frac{\text{PO}_4}{\text{PO}_4 + K_{\text{PO}_4}^{\text{bact}}} \quad (34e)$$

$$L_{\text{N}}^{\text{bact}} = L_{\text{NO}_3}^{\text{bact}} + L_{\text{NH}_4}^{\text{bact}} \quad (34f)$$

$$L_{\text{NH}_4}^{\text{bact}} = \frac{K_{\text{NO}_3}^{\text{bact}} \text{NH}_4}{K_{\text{NO}_3}^{\text{bact}} K_{\text{NH}_4}^{\text{bact}} + K_{\text{NH}_4}^{\text{bact}} \text{NO}_3 + K_{\text{NO}_3}^{\text{bact}} \text{NH}_4} \quad (34g)$$

$$L_{\text{NO}_3}^{\text{bact}} = \frac{K_{\text{NH}_4}^{\text{bact}} \text{NO}_3}{K_{\text{NO}_3}^{\text{bact}} K_{\text{NH}_4}^{\text{bact}} + K_{\text{NH}_4}^{\text{bact}} \text{NO}_3 + K_{\text{NO}_3}^{\text{bact}} \text{NH}_4} \quad (34h)$$

The half-saturation constants of the P and N limitation terms (K_i^{bact}) are set in the namelist.

In PISCES, bacterial biomass is not explicitly modeled; Instead, we use the following formulation:

$$z_{\max} = \max \left(Zz_{\text{mxl}}, Zz_{\text{eu}} \right) \quad (35a)$$

$$\text{Bact} = \begin{cases} \min(0.7(Z + 2M), 4 \mu\text{mol CL}^{-1}) & \text{if } z \leq z_{\max} \\ \text{Bact}(z_{\max}) \left(\frac{z_{\max}}{z} \right)^{0.683} & \text{Otherwise} \end{cases} \quad (35b)$$

In the previous equation, $0.7(Z + 2M)$ is a proxy for the bacterial concentration. This relationship has been constructed from [an unpublished version of PISCES \(already mentioned in Aumont and Bopp \(2006\)\)](#) that includes an explicit description of the bacterial biomass. Below a certain depth (z_{\max}), this biomass decreases with depth via a power-law function (Aristegui et al., 2009).

In Eq. (32), the terms Φ^{DOC} denote aggregation processes and are described hereafter (see Sect. 4.4.1). For DOM, we consider turbulence-induced as well as Brownian aggregation processes.

$$\Phi_1^{\text{DOC}} = \text{sh} \times (a_1 \text{DOC} + a_2 \text{POC}) \text{DOC} \quad (36a)$$

$$\Phi_2^{\text{DOC}} = \text{sh} \times a_3 \text{GOC} \times \text{DOC} \quad (36b)$$

$$\Phi_3^{\text{DOC}} = (a_4 \text{POC} + a_5 \text{DOC}) \text{DOC} \quad (36c)$$

4.4 Particulate organic matter

PISCES includes two different schemes for particulate organic matter:

- A simple model based on two different size-classes for particulate organic matter. In that case, particulate organic matter is modeled in PISCES using two tracers corresponding to the two size-classes: POC for the smaller class (1–100 μm) and GOC for the larger class (100–5000 μm).

- 5 – A more complex model proposed by [Kriest and Evans \(1999\)](#) in which the size-spectrum of the particulate organic matter can be represented by a power-law function. Here, particulate organic matter is represented by two variables: the first (POC) is the carbon concentration and the second (NUM) is the total number of aggregates by unit volume of water.

10 By default, the simplest parameterization is used. The Kriest model is activated by a cpp key `key_kriest`.

4.4.1 Two compartments model of POM

The temporal evolution of POC is written:

$$\begin{aligned}
 \frac{\partial \text{POC}}{\partial t} = & \sigma^Z \sum g^Z(X)Z + 0.5m^D \frac{D}{D + K_m} D + r^Z f_Z(T) \frac{Z}{Z + K_m} Z + m^Z f_Z(T) Z^2 \\
 & + (1 - 0.5R_{\text{CaCO}_3})(m^P \frac{P}{P + K_m} P + w^P P^2) \\
 & + \lambda_{\text{POC}}^* \text{GOC} + \Phi_1^{\text{DOC}} + \Phi_3^{\text{DOC}} - (g^M(\text{POC}) + g_{\text{FF}}^M(\text{POC})) M - g^Z(\text{POC}) Z \\
 & - \lambda_{\text{POC}}^* \text{POC} - \Phi - w_{\text{POC}} \frac{\partial \text{POC}}{\partial z}
 \end{aligned} \tag{37}$$

where w_{POC} is the vertical sinking speed. For POC, it is set to a constant value, in general to a small value of the order of a few meters per day. The fate of mortality and aggregation of nanophytoplankton depends on the proportion of the calcifying organisms (R_{CaCO_3}). We assume that 50 % of the organic matter of the calcifiers is associated with the shell. Since calcite is significantly denser than organic matter, 50 % of the biomass of the dying calcifiers is routed to the fast sinking particles. The same is assumed for the mortality of diatoms as a consequence of the denser density of biogenic silica.

5 The specific degradation rate λ_{POC}^* depends on temperature with a Q_{10} of about 1.9, the same as for phytoplankton. Furthermore, observations generally tend to show slower degradation rates when waters are anoxic ([Harvey et al., 1995](#); [Mooy et al., 2002](#)). In [Mooy et al. \(2002\)](#), the

attenuation coefficient (b) for the flux was found to be about 0.4 instead of the standard value 0.86 (Martin et al., 1987). This corresponds to a 45 % decrease of the degradation rate in anoxic waters relative to oxic waters, which is implemented as:

$$\lambda_{\text{POC}}^* = \lambda_{\text{POC}} f_P(T) (1 - 0.45 \Delta(\text{O}_2)) \quad (38)$$

POC experiences aggregation due to turbulence and differential settling:

$$\Phi = \text{sh} \times a_6 \text{POC}^2 + \text{sh} \times a_7 \text{POC} \times \text{GOC} + a_8 \text{POC} \times \text{GOC} + a_9 \text{POC}^2 \quad (39)$$

In this equation, the first two terms correspond to turbulent aggregation, and the two last terms to differential settling aggregation. The values of the parameters controlling these processes have been computed offline assuming a steady-state power-law size-spectrum for particles with an exponent of 3.6. Subsequently, the different coagulation kernels (e.g., Jackson, 1990; Krist and Evans, 1999) have been integrated over the size-ranges corresponding to the different compartments. A constant stickiness of 0.1 has been chosen.

The temporal evolution of GOC is written:

$$\begin{aligned} \frac{\partial \text{GOC}}{\partial t} = & \sigma^M \left(\sum_I g^M(I) + g_{\text{FF}}^M(\text{POC}) + g_{\text{FF}}^M(\text{GOC}) \right) M + r^M f_M(T) \frac{M}{M + K_m} M + P_{\text{up}}^M \\ & + 0.5 R_{\text{CaCO}_3} \left(m^P \frac{P}{P + K_m} P + w^P P^2 \right) + 0.5 m^D \frac{D}{D + K_m} D w^D D^2 \\ & + \Phi + \Phi_2^{\text{DOC}} - g_{\text{FF}}^M(\text{GOC}) M - \lambda_{\text{POC}}^* \text{GOC} - w_{\text{GOC}} \frac{\partial \text{GOC}}{\partial z} \end{aligned} \quad (40)$$

The equation controlling the temporal evolution of GOC is similar to that of POC. However, some observations have shown that the mean sinking speed of particulate organic matter increases with depth (e.g., Berelson, 2002). Such an increase is consistent with the power law formulation proposed by Martin et al. (1987). Such an increase in the settling speed is parameterized in PISCES for GOC as follows:

$$\underline{Zz}_{\text{max}} = \max(\underline{Zz}_{\text{eu}}, \underline{Zz}_{\text{mxl}}) \quad (41a)$$

$$w_{\text{GOC}} = w_{\text{GOC}}^{\min} + (200 - w_{\text{GOC}}^{\min}) \frac{\max(0, z - Z_{\max})}{5000} \frac{\max(0, z - z_{\max})}{5000} \quad (41b)$$

The parameters in this equation have been adjusted using a model of aggregation/disaggregation with multiple size classes (Gehlen et al., 2006). The maximum sinking speed is set to 200
 10 m d^{-1} and is reached at about 5000 m depth over most of the ocean since z_{\max} is generally less
than 100 m. We have not included any ballasting effect due to the higher density of biogenic
 silica or calcite (Klaas and Archer, 2002; Armstrong et al., 2002). In fact, observations are rather
 15 contradictory on this ballast effect (Lee et al., 2009). In particular, the greater efficiency of the
 vertical sedimentation of organic matter when associated with calcite and biogenic silica may
 be due rather to the protection of an organic matter fraction by the inorganic matrix (Moriceau
 et al., 2009; Engel et al., 2009).

4.4.2 Kriest model of particulate organic matter

Here we present a brief overview of the model of Kriest and Evans (1999). The reader is referred
 20 to the literature where the method has been presented (e.g., Kriest and Evans, 1999, 2000;
 Kriest, 2002) for more detail. The model postulates that the carbon content ($m(d_i)$), the sinking
 speed ($w(d_i)$) and the abundance of the aggregates ($n(d_i)$) can be described by power-law
 functions of their diameters (d_i):

$$m(d_i) = C d_i^{\zeta} \quad (42a)$$

$$w(d_i) = B d_i^{\nu} \quad (42b)$$

$$n(d_i) = A d_i^{\epsilon} \quad (42c)$$

It is also assumed, as in Kriest (2002), that aggregates above a certain size L have a constant
 sinking speed w_L .

The slope of the size spectrum can be computed from the total number of aggregates (NUM)
 5 and the total mass of particles (POC), which are the two state variables of the model:

$$\epsilon = \frac{(\zeta + 1)\text{POC} - m_L \text{NUM}}{\text{POC} - m_L \text{NUM}} \quad (43)$$

where m_l is the mass of the smallest aggregate (of size l).

Having ϵ , the average sinking speed of numbers (w_{NUM}) and mass (w_{POC}) can be computed following [Kriest \(2002\)](#):

$$10 \quad w_{\text{POC}} = w_l \frac{\zeta + 1 - \epsilon + \left(\frac{L}{l}\right)^{1+\nu+\zeta-\epsilon\nu}}{1 + \nu + \zeta - \epsilon} \quad (44a)$$

$$w_{\text{NUM}} = w_l \frac{1 - \epsilon + \left(\frac{L}{l}\right)^{1+\nu-\epsilon\nu}}{1 + \nu - \epsilon} \quad (44b)$$

15 The number of particles and the mass of particles change independently. For instance, sinking tends to remove larger particles. As a consequence, the relationship between the number of particles and their mass evolves with time and space and so does ϵ . As a result, the sinking speeds for both mass and number vary with space and time.

Aggregation (ξ) depends on the particle abundance, their size distribution, rate of turbulent shear and the difference in particle sinking speeds, as well as the stickiness (the probability that two particles stick together after contact). The approach implemented in PISCES follows that described in [Kriest \(2002](#), see there for term ξ and its computation). Currently it is assumed that turbulent shear rate is high in the mixed layer (1 m s^{-1}), and low below (0.01 m s^{-1}). Summing-up the number of collisions due to turbulent shear and differential settlement, C_{sh} and C_{ds} , respectively, the decrease of the number of particles due to aggregation then is:

$$25 \quad \xi = \text{Stick} \times (C_{\text{sh}} + C_{\text{ds}}) \quad (45)$$

In PISCES, the stickiness (the efficiency of the collisions) is set to a constant value in the namelist.

The temporal evolution of the mass of particles is given as:

$$5 \quad \frac{\partial \text{POC}}{\partial t} = \sigma^Z \sum g^Z(X)Z + \sigma^M \left(\sum_I g^M(I) + g_{\text{FF}}^M(\text{POC}) \right) M + m^P \frac{P}{P + K_m} P \\ + w^P P^2 + m^D \frac{D}{D + K_m} D + w^D D^2 + r^Z f_Z(T) \frac{Z}{Z + K_m} Z + m^Z f_Z(T) Z^2$$

$$\begin{aligned}
& + r^M f_M(T) \frac{M}{M + K_m} M + P_{\text{up}}^M - g^M(\text{POC})M - g^Z(\text{POC})Z - \lambda_{\text{POC}}^* \text{POC} \\
& + \Phi_1^{\text{DOC}} + \Phi_3^{\text{DOC}} - (g^M(\text{POC}) + g_{\text{FF}}^M(\text{POC})) M - g^Z(\text{POC})Z \\
& - \lambda_{\text{POC}}^* \text{POC} - w_{\text{POC}} \frac{\partial \text{POC}}{\partial z}
\end{aligned} \tag{46}$$

10 This is exactly equal to the the two equations used for the temporal evolution of POC and GOC in the two-compartments model of PISCES (see Eqs. 37 and 40).

$$\begin{aligned}
\frac{\partial \text{NUM}}{\partial t} = & \frac{\sigma^Z \sum g^Z(X)Z}{\bar{m}_Z} + \frac{\sigma^M (\sum_I g^M(I) + g_{\text{FF}}^M(\text{POC})) M}{\bar{m}_M} + \frac{m^P \frac{P}{P+K_m} P + w^P P^2}{\bar{m}_P} \\
& + \frac{m^D \frac{D}{D+K_m} D + w^D D^2}{\bar{m}_D} + \frac{r^Z f_Z(T) \frac{Z}{Z+K_m} Z + m^Z f_Z(T) Z^2}{\bar{m}_Z} \\
& + \frac{r^M f_M(T) \frac{M}{M+K_m} M + P_{\text{up}}^M}{\bar{m}_M} - \frac{(g^M(\text{POC}) + g_{\text{FF}}^M(\text{POC})) M}{\bar{m}_M} \\
& - \frac{g^Z(\text{POC})Z}{\bar{m}_Z} - \lambda_{\text{POC}}^* \text{POC} + \frac{\Phi_1^{\text{DOC}} + \Phi_3^{\text{DOC}}}{m_l} - \xi - w_{\text{NUM}} \frac{\partial \text{NUM}}{\partial z}
\end{aligned} \tag{47}$$

In this equation, each process affecting the mass of the particles is divided by the mean mass (\bar{m}) of the compartment exerting this process to convert to numbers.

4.4.3 Iron in particles

In this subsection, the description corresponds to the two-compartments version of the model. To obtain the Kriest version, the equations for both SFe and BFe, [the iron content of the small and big particles respectively](#), should be simply summed.

$$\begin{aligned}
\frac{\partial \text{SFe}}{\partial t} = & \sigma^Z \sum_I \theta^{\text{Fe},I} g^Z(I)Z + \theta^{\text{Fe},Z} (r^Z f_Z(T) \frac{Z}{Z + K_m} Z + m^Z f_Z(T) Z^2) \\
& + \lambda_{\text{GOC}}^* \text{BFe} + \theta^{\text{Fe},P} (1 - 0.5 R_{\text{CaCO}_3}) (m^P \frac{P}{P + K_m} P + \text{sh} \times w^P P^2)
\end{aligned}$$

5

$$\begin{aligned}
& + \theta^{\text{Fe,D}} 0.5m^D \frac{D}{D + K_m} D + \lambda_{\text{Fe}} \text{POC Fe}' + \text{Cgfe1} - \lambda_{\text{POC}}^* \text{SFe} - \theta^{\text{Fe,POC}} \Phi \\
& - \theta^{\text{Fe,POC}} (g^M(\text{POC}) + g_{\text{FF}}^M(\text{POC})) M + \kappa_{\text{Bact}}^{\text{SFe}} \text{Bactfe} - \theta^{\text{Fe,POC}} g^Z(\text{POC}) \\
& - w_{\text{POC}} \frac{\partial \text{SFe}}{\partial z}
\end{aligned} \tag{48}$$

$$\begin{aligned}
\frac{\partial \text{BFe}}{\partial t} = & \sigma^M \left(\sum_I \theta^{\text{Fe,I}} g^M(I) + \theta^{\text{Fe,POC}} g_{\text{FF}}^M(\text{POC}) + \theta^{\text{Fe,GOC}} g_{\text{FF}}^M(\text{GOC}) \right) M \\
& + \theta^{\text{Fe,M}} (r^M f_M(T) \frac{M}{M + K_m} M + P_{\text{up}}^M) + \theta^{\text{Fe,P}} 0.5 R_{\text{CaCO}_3} (m^P \frac{P}{P + K_m} P \\
& + \text{sh} \times w^P P^2) + \theta^{\text{Fe,D}} (0.5m^D \frac{D}{D + K_m} D + \text{sh} \times w^D D^2) + \kappa_{\text{Bact}}^{\text{BFe}} \text{Bactfe} \\
& + \lambda_{\text{Fe}} \text{GOC Fe}' + \theta^{\text{Fe,POC}} \Phi + \text{Cgfe2} - \theta^{\text{Fe,GOC}} g_{\text{FF}}^M(\text{GOC}) M - \lambda_{\text{POC}}^* \text{BFe} \\
& - w_{\text{GOC}} \frac{\partial \text{BFe}}{\partial z}
\end{aligned} \tag{49}$$

15 where Fe' is the free form of dissolved iron. Its determination is detailed in Sect. 4.5.3. Bactfe is the amount of iron taken up by bacteria which is lost as particulate organic iron. Its computation is detailed in Sect. 4.5.3.

The free form of dissolved iron Fe' is the only form of iron that is ~~supposed~~ assumed to be susceptible to scavenging. The scavenging rate of iron is made dependent upon the particulate load of the seawater as follows (e.g., Honeyman et al., 1988; Parekh et al., 2004):

$$\lambda_{\text{Fe}}^* = \lambda_{\text{Fe}}^{\text{min}} + \lambda_{\text{Fe}}(\text{POC} + \text{GOC} + \text{CaCO}_3 + \text{BSi}) + \lambda_{\text{Fe}}^{\text{dust}} \text{Dust} \tag{50a}$$

$$5 \text{ Scav} = \lambda_{\text{Fe}}^* \text{Fe}' \tag{50b}$$

Implicitly, in this equation, it is assumed that the affinity of iron for the different types of biogenic particles is the same. Iron is also scavenged by lithogenic particles originating from dust deposition as evidenced by mesocosm experiments (Wagner et al., 2010). The concentration of lithogenic particles is estimated as described in Eq. (84). Model estimates (Ye et al., 2011)

suggest a different affinity for these particles compared to biogenic particles, which justifies the split between biogenic and lithogenic materials in Eq. (50). The amount of iron that is scavenged by POC ($\lambda_{\text{Fe}}\text{POCFe}'$) and GOC ($\lambda_{\text{Fe}}\text{GOCFe}'$) is then allocated to SFe and BFe, respectively.

4.4.4 PSi

$$\begin{aligned}
 \frac{\partial \text{PSi}}{\partial t} = & \theta^{\text{Si},D} g^M(D) M + \theta^{\text{Si},D} g^Z(D) Z + \theta^{\text{Si},D} m^D \frac{D}{K_m + D} D^{\text{Si}} + \text{sh} \times w^D D D^{\text{Si}} \\
 & - \lambda_{\text{PSi}}^* \text{Diss}_{\text{Si}} \text{PSi} - w_{\text{GOC}} \frac{\partial \text{CaCO}_3}{\partial z}
 \end{aligned} \tag{51}$$

The dissolution rate of PSi depends on in situ temperature and on silicic acid saturation following the parameterization proposed by [Ridgwell et al. \(2002\)](#):

$$\begin{aligned}
 \text{Si}_{\text{eq}} &= 10^{6.44 - \frac{968}{T+273.15}} \\
 \text{Si}_{\text{sat}} &= \frac{\text{Si}_{\text{eq}} - \text{Si}}{\text{Si}_{\text{eq}}} \\
 \lambda_{\text{PSi}}^* &= \lambda_{\text{PSi}} \left[0.225 \left(1 + \frac{T}{15} \right) \text{Si}_{\text{sat}} + 0.775 \left(\left(1 + \frac{T}{400} \right)^4 \text{Si}_{\text{sat}} \right)^9 \right]
 \end{aligned} \tag{52}$$

The evolution of λ_{PSi}^* as a function of Si and of temperature is shown on Fig. 4.

Laboratory experiments show that the diatom frustule is made of two biogenic silica phases which dissolve simultaneously, but at different rates (e.g., [Kamatani et al., 1980](#); [Van Capellen et al., 2002](#); [Moriceau et al., 2009](#); [Loucaides et al., 2012](#)). The first phase dissolves significantly faster than the second phase. It is associated with membrane lipids and amino acids and represents about 1/3 of the frustule ([Moriceau et al., 2009](#)). However, the existence of these two phases is still a matter of debate as it has been hypothesized to be a result of the experimental design of the dissolution experiments ([Loucaides et al., 2012](#)). In PISCES, despite this uncertainty, we model silica dissolution using two phases. The proportion of the most “labile” phase

10 is set to a constant (χ_{lab}) in the upper ocean and is computed in the rest of the ocean assuming steady state:

$$\begin{aligned} \underline{Z}z_{\text{max}} &= \max(\underline{Z}z_{\text{cu}}, \underline{Z}z_{\text{mxi}}) \\ \chi_{\text{lab}} &= \begin{cases} \chi_{\text{lab}}^0 & \text{if } z \leq z_{\text{max}} \\ \chi_{\text{lab}}^0 \exp\left(-(\lambda_{\text{PSi}}^{\text{lab}} - \lambda_{\text{PSi}}^{\text{ref}}) \left(\frac{z - \underline{Z}z_{\text{max}}}{w_{\text{GOC}}} \frac{z - z_{\text{max}}}{w_{\text{GOC}}}\right)\right) & \text{Otherwise} \end{cases} \end{aligned} \quad (53a)$$

$$15 \lambda_{\text{PSi}} = \chi_{\text{lab}} \lambda_{\text{PSi}}^{\text{lab}} + (1 - \chi_{\text{lab}}) \lambda_{\text{PSi}}^{\text{ref}} \quad (53b)$$

4.5 Nutrients

4.5.1 Nitrate and ammonium

$$\frac{\partial \text{NO}_3}{\partial t} = \text{Nitrif} - \mu_{\text{NO}_3}^P P - \mu_{\text{NO}_3}^D D - R_{\text{NH}_4} \lambda_{\text{NH}_4} \Delta(\text{O}_2) \text{NH}_4 - R_{\text{NO}_3} \text{Denit} \quad (54)$$

$$\begin{aligned} \frac{\partial \text{NH}_4}{\partial t} &= \gamma^Z (1 - e^Z - \sigma^Z) \sum_I g^Z(I) Z + \gamma^M (1 - e^M - \sigma^M) \\ &\times \left(\sum_I g^M(I) + g_{\text{FF}}^M(\text{POC}) + g_{\text{FF}}^M(\text{GOC}) \right) M + \gamma^M R_{\text{up}}^M + \text{Remin} + \text{Denit} + N_{\text{fix}} \\ &- \text{Nitrif} - \lambda_{\text{NH}_4} \Delta(\text{O}_2) \text{NH}_4 - \mu_{\text{NH}_4}^P P - \mu_{\text{NH}_4}^D D \end{aligned} \quad (55)$$

Nitrification (Nitrif) corresponds to the conversion of ammonium to nitrate due to bacterial activity. It is assumed to be photoinhibited (e.g., [Horrihan et al., 1981](#); [Yoshioka and Saijo, 1984](#)) and reduced in suboxic waters:

$$5 \text{Nitrif} = \lambda_{\text{NH}_4} \frac{\text{NH}_4}{1 + \overline{\text{PAR}}} (1 - \Delta(\text{O}_2)) \quad (56a)$$

$$\overline{\text{PAR}} = \overline{\text{PAR}}_1 + \overline{\text{PAR}}_2 + \overline{\text{PAR}}_3 \quad (56b)$$

where $\overline{\text{PAR}}$ is the PAR averaged over the mixed layer and $\Delta(\text{O}_2)$ varies between 0 (oxic conditions, $\text{O}_2 > \text{O}_2^{\text{min},1}$) and 1 (anoxia) according to:

$$\Delta(\text{O}_2) = \min \left(1, \max \left(0, 0.4 \frac{\text{O}_2^{\text{min},1} - \text{O}_2}{\text{O}_2^{\text{min},2} + \text{O}_2} \right) \right) \quad (57)$$

When waters become suboxic, nitrate instead of oxygen is consumed during the remineralization of organic matter, i.e. denitrification (Denit). The N/C stoichiometric ratio of denitrification R_{NO_3} can be computed from $R_{-\text{O}_2/\text{NO}_3}$ and is found to be 0.86 (Paulmier et al., 2009). Equation (57), implies that denitrification stops at oxygen concentration above $6 \mu\text{M}$ (Lipschultz et al., 1990). We further assume complete oxidation by nitrate of the ammonia released from organic matter during denitrification. This oxidation rate has been arbitrarily set to the same value as nitrification rate (λ_{NH_4}).

Finally, nitrogen fixation is parameterized in PISCES as follows:

$$L_N^{\text{Dz}} = \begin{cases} 0.01 & \text{if } \underline{zL_N^P} \geq 0.8 \\ 1 - L_N^P & \text{Otherwise} \end{cases} \quad (58a)$$

$$N_{\text{fix}} = N_{\text{fix}}^m \max(0, \mu_P - 2.15) L_N^{\text{Dz}} \min \left(\frac{\text{bFe}}{K_{\text{Fe}}^{\text{Dz}} + \text{bFe}}, \frac{\text{PO}_4}{K_{\text{PO}_4}^{\text{P},\text{min}} + \text{PO}_4} \right) \left(1 - e^{-\frac{-\text{PAR}}{E_{\text{fix}}}} \right) \quad (58b)$$

This very crude parameterization is based on the following assumptions that have been inferred from studies of *Trichodesmium* (e.g., Mills et al., 2004; Masotti et al., 2007; Zehr, 2011):

- Nitrogen fixation is restricted to warm waters above 20°C ($\underline{\mu_P > \mu_P(20) = 2.15}$)
- Nitrogen fixation is restricted to areas with insufficient nitrogen ($\underline{L_N^P < 0.8}$)
- Nitrogen fixation requires iron and phosphorus
- Nitrogen fixation needs high light levels, i.e. E_{fix} is high.

The scaling factor N_{fix}^m is set from the namelist and thus, may be chosen by the user.

4.5.2 Phosphate

$$\begin{aligned} \frac{\partial \text{PO}_4}{\partial t} = & \gamma^Z (1 - e^Z - \sigma^Z) \sum_I g^Z(I) Z + \gamma^M (1 - e^M - \sigma^M) \left(\sum_I g^M(I) + \sum_I g_{\text{FF}}^M(I) \right) M \\ & + \gamma^M R_{\text{up}}^M + \text{Remin} + \text{Denit} - \mu^P P - \mu^D D \end{aligned} \quad (59)$$

All terms in this equation have been described previously.

4.5.3 Iron

$$\begin{aligned} \frac{\partial \text{Fe}}{\partial t} = & \max \left(0, (1 - \sigma^Z) \frac{\sum_I \theta^{\text{Fe},I} g^Z(I)}{\sum_I g^Z(I)} - e_{\text{N}}^Z \theta^{\text{Fe},Z} \right) \sum_I g^Z(I) Z \\ & + \max \left(0, (1 - \sigma^M) \frac{\sum_I \theta^{\text{Fe},I} g^M(I) + \sum_I \theta^{\text{Fe},I} g_{\text{FF}}^M(I)}{\sum_I g^M(I) + \sum_I g_{\text{FF}}^M(I)} - e_{\text{N}}^M \theta^{\text{Fe},Z} \right) \\ & \left(\sum_I g^M(I) + \sum_I g_{\text{FF}}^M(I) \right) M + \gamma^M \theta^{\text{Fe},Z} R_{\text{up}}^M + \lambda_{\text{POC}}^* \text{SFe} \\ & - (1 - \delta^P) \mu^{P^{\text{Fe}}} P - (1 - \delta^D) \mu^{D^{\text{Fe}}} D - \text{Scav} - \text{Cgfe1} - \text{Cgfe2} \\ & - \text{Aggfe} - \text{Bactfe} \end{aligned} \quad (60)$$

Iron scavenging (Scav) has been described previously in Sect. 4.4.3. Iron is present in seawater largely as colloids (e.g., Wu et al., 2001; Wu and Boyle, 2002; Boyd and Ellwood, 2010). These colloids may aggregate with dissolved organic matter as it forms gels. Thus, they may be transferred to the particulate pool, and settle to the ocean floor. Very few models have incorporated this potential important sink of dissolved iron (Ye et al., 2009, 2011). In PISCES, we model this process following the approach chosen for DOM (see Sect. 4.3):

$$\text{Cgfe1} = ((a_1 \text{DOC} + a_2 \text{POC}) \times \text{sh} + a_4 \text{POC} + a_5 \text{DOC}) \times \text{Fe}_{\text{coll}} \quad (61a)$$

$$10 \quad C_{\text{gfe}2} = a_3 \text{GOC} \times \text{sh} \times \text{Fe}_{\text{coll}} \quad (61\text{b})$$

Fe_{coll} is computed from the iron chemistry model (see below).

When dissolved iron concentration exceeds the total ligand concentration L_T , scavenging is enhanced as it is done in many other biogeochemical models (e.g., [Moore et al., 2004](#); [Dutkiewicz et al., 2005](#)):

$$15 \quad \text{Aggfe} = 1000 \lambda^{\text{Fe}} \max(0, \text{Fe} - L_T) \text{Fe}' \quad (62)$$

This scavenging loss term is ~~supposed~~ assumed to be definitive, i.e. iron is permanently removed from the ocean by this process.

Heterotrophic bacteria acquire iron from seawater using siderophore-based iron transport
 20 systems ([Haygood et al., 1993](#); [Martinez et al., 2000](#)). Observations show that they have quite elevated Fe/C ratios and account for a significant fraction of the total biological uptake of iron ([Tortell et al., 1996, 1999](#)). The bacterial uptake of iron is parameterized according to:

$$\text{Bactfe} = \mu_P L_{\text{lim}}^{\text{Bact}} \theta_{\text{max}}^{\text{Fe,Bact}} \frac{\text{Fe}}{K_{\text{Fe}}^{\text{B,I}} + \text{Fe}} \text{Bact} \quad (63)$$

25 where $\theta_{\text{max}}^{\text{Fe,Bact}}$ denotes the maximum Fe/C ratio of bacteria.

The different iron pools are computed using a chemistry model. Two different chemistry models are available in PISCES:

- A simple chemistry model based on one ligand L and two dissolved iron forms: dissolved inorganic iron Fe' and dissolved complexed iron FeL .
- The complex chemistry model of [Tagliabue and Arrigo \(2006\)](#) as modified by [Tagliabue and Völker \(2011\)](#). This model is based on two ligands (L_W and L_S) and five iron forms: free Fe(II) ($\text{Fe(II)}'$) and Fe(III) ($\text{Fe(III)}'$), Fe(III) bound to the weak ligand (FeL_W),
 5 Fe(III) bound to the strong ligand (FeL_S) and solid iron (Fe_P).

The complex iron model is activated in PISCES setting the boolean variable `ln_fechem` to true.

Our main purpose is not to provide a fully detailed description of both chemistry models as they have been described fairly extensively elsewhere. For the simple chemistry model, the reader should refer of [Aumont and Bopp \(2006\)](#) whereas the complex model is detailed in [Tagliabue and Völker \(2011\)](#). For the complex model, all chemical constants have identical values to what was chosen in [Tagliabue and Völker \(2011\)](#) and are thus not listed in Tables 1a–1e. Only a very brief description of both models will be given here, especially for the complex model. Both models are based on the assumption that chemical reactions are fast enough relative to the other biogeochemical processes affecting iron (for instance phytoplankton uptake) that they can be considered at equilibrium.

4.5.4 Simple chemistry model

Dissolved iron is assumed to be in the form of free inorganic iron Fe' and of “complexed” iron FeL . Both forms of iron are ~~supposed~~ assumed to be equally susceptible to consumption by phytoplankton despite recent observations suggest that this may be not the case ([Nishioka and Takeda, 2000](#); [Chen and Wang, 2001](#); [Chen et al., 2003](#)). In other words, the total bioavailable concentration of iron is equal to the total dissolved iron concentration (Fe). The chemical speciation of iron is deduced from the three following equations:

$$\begin{aligned} L_T &= \text{FeL} + L' \\ \text{Fe} &= \text{FeL} + \text{Fe}' \\ K_{\text{eq}}^{\text{Fe}} &= \frac{\text{FeL}}{L'\text{Fe}'} \end{aligned} \quad (64)$$

The chemical equilibrium constant $K_{\text{eq}}^{\text{Fe}}$ is computed from the formulation proposed by [Liu and Millero \(2002\)](#). Solving this set of equations is equivalent to solve a second-order polynomial equation in Fe' , whose solution is:

$$\Delta = 1 + K_{\text{eq}}^{\text{Fe}} L_T - K_{\text{eq}}^{\text{Fe}} \text{Fe}$$

$$\text{Fe}' = \frac{-\Delta + \sqrt{\Delta^2 + 4K_{\text{eq}}^{\text{Fe}}\text{Fe}}}{2K_{\text{eq}}^{\text{Fe}}} \quad (65)$$

Colloidal iron is assumed to [represent 50% of FeL](#):

$$10 \quad \text{Fe}_{\text{coll}} = 0.5\text{FeL} \quad (66)$$

The total ligand concentration L_T can be either constant over the ocean, using a value defined in the namelist or can be variable using the relationship proposed by [Tagliabue and Völker \(2011\)](#):

$$15 \quad L_T = \max(0.09(\text{DOC} + 40) - 3, 0.6) \quad (67)$$

where L_T is in nmol L^{-1} and DOC in $\mu\text{mol L}^{-1}$.

4.5.5 Complex chemistry model

The iron chemical system is governed by the following set of four equations:

$$20 \quad 0 = k_{\text{IW}}\text{Fe(III)'}L_W - k_{\text{bW}}\text{FeL}_W - k_{\text{phW}}\text{FeL}_W - k_{\text{th}}\text{FeL}_W \quad (68a)$$

$$0 = k_{\text{IS}}\text{Fe(III)'}L_S - k_{\text{bS}}\text{FeL}_S - k_{\text{phS}}\text{FeL}_S \quad (68b)$$

$$0 = k_{\text{phW}}\text{FeL}_W + k_{\text{phS}}\text{FeL}_S + k_{\text{th}}\text{Fe(III)'} - k_{\text{ox}}\text{Fe(II)'} \quad (68c)$$

$$0 = k_{\text{pcp}}\text{Fe(III)'} - k_r\text{Fe}_P \quad (68d)$$

A supplementary reaction has been added relative to the original set of equations. In the Pacific Ocean, thermal (dark) reduction of Fe(III) organic complexes has been shown to produce the accumulation of sizeable amount of Fe(II)' in the mesopelagic zone ([Hansard et al., 2009](#)).

5 Additional constraints are given by the conservation of total dissolved iron (Fe), L_{WT} and L_{ST} over the fast timescale:

$$\text{Fe} = \text{Fe(III)'} + \text{Fe(II)'} + \text{FeL}_W + \text{FeL}_S + \text{Fe}_P \quad (69a)$$

$$L_{WT} = FeL_W + L_W \quad (69b)$$

$$L_{ST} = FeL_S + L_S \quad (69c)$$

Solving this system of equations is equivalent to solve a third-order polynomial equation in $Fe(III)'$ (Eq. 16 in [Tagliabue and Völker, 2011](#)). Because thermal aphotic reduction of FeL_W has been added here, the definition of some coefficients in the original study has changed:

$$b = 1 + \frac{k_{phW} + k_{th}}{k_{ox}} \quad (70a)$$

$$K_W = \frac{k_{phW} + k_{th} + k_{bW}}{k_{lW}} \quad (70b)$$

where k_{th} has been set to 0.0048 h^{-1} . Then, knowing $Fe(III)'$, the other four iron species can be computed.

Observations suggest that the weak ligand (L_W) is ubiquitous in the water column and is probably produced by the degradation of organic matter sinking from the upper layers of the ocean. The strong ligand is present in the upper ocean and is most probably produced by autotrophic and heterotrophic bacteria (for instance siderophores) (e.g., [Boyd and Ellwood, 2010](#)). In PISCES, we assume that $2/3$ of the total ligand concentration above 0.6 nmol L^{-1} is going to L_S , the rest is attributed to L_W :

$$L_{WT} = 0.6 + \frac{1}{3}(L_T - 0.6) \quad (71a)$$

$$L_{ST} = \frac{2}{3}(L_T - 0.6) \quad (71b)$$

As in the simple chemistry model, the ligand concentration L_T can be either constant over the ocean, using a value defined in the namelist or can be variable using the relationship proposed by [Tagliabue and Völker \(2011\)](#) (see Eq. 67).

The rate constants required by the model are identical to those described by [Tagliabue et al. \(2009a\)](#) as modified by [Tagliabue and Völker \(2011\)](#). Furthermore, we have slightly changed

the formulation of the oxidation rate constant used in the original model:

$$k_{\text{ox}} = k'_{\text{ox}} \frac{\max(\text{O}_2, 1 \mu\text{molL}^{-1})}{\text{O}_{2\text{sat}}} \quad (72)$$

This avoids numerical problems in strongly anoxic areas where oxygen concentration is close to 0. Bioavailable iron can be defined either as $\text{Fe(II)}' + \text{Fe(III)}' + \text{FeL}_S$ or as $\text{Fe(II)}' + \text{Fe(III)}' + \text{FeL}_S + \text{FeL}_W$. k_{th} has assigned the value computed from the observations by [Hansard et al. \(2009\)](#), consistent with the data of [Pullin and Cabaniss \(2003\)](#). Colloidal iron and dissolved inorganic iron are defined as:

$$\text{Fe}_{\text{coll}} = 0.5(\text{Fe}_p + \text{FeL}_W + \text{FeL}_S) \quad (73a)$$

$$\text{Fe}' = \text{Fe(III)}' + \text{Fe(II)}' \quad (73b)$$

We assumed that 50% of the iron bound to ligands and of the particulate inorganic iron is colloidal iron.

4.5.6 Si

$$\frac{\partial \text{Si}}{\partial t} = \lambda_{\text{PSi}}^* \text{Diss}_{\text{Si}} \text{PSi} - \theta_{\text{opt}}^{D,\text{Si}} (1 - \delta^D) \mu^D D \quad (74)$$

All terms in this equation have been already defined previously.

4.6 Calcite

$$\frac{\partial \text{CaCO}_3}{\partial t} = P_{\text{CaCO}_3} - \lambda_{\text{CaCO}_3}^* \text{CaCO}_3 - w_{\text{GOC}} \frac{\partial \text{CaCO}_3}{\partial z} \quad (75)$$

In PISCES, calcium carbonate is ~~supposed~~assumed to exist only in the form of calcite. Thus, aragonite is not considered, for instance, for the computation of chemical dissolution in the water column.

The biological production of sinking calcite is defined as:

$$P_{\text{CaCO}_3} = R_{\text{CaCO}_3} \left(\eta^Z g^Z(P)Z + \eta^M g^M(P)M + 0.5(m^P \frac{P}{K_m + P} P + \text{sh} \times w^P P^2) \right) \quad (76)$$

The rain ratio R_{CaCO_3} is variable. We propose the following parameterization for this ratio:

$$\begin{aligned} R_{\text{CaCO}_3} = & r_{\text{CaCO}_3} L_{\text{lim}}^{\text{CaCO}_3} \frac{T}{0.1 + T} \max\left(1, \frac{P}{2}\right) \\ & \times \frac{\max(0, \text{PAR} - 1)}{4 + \text{PAR}} \frac{30}{30 + \text{PAR}} \\ & \times \left(1 + \exp\left(\frac{-(T - 10)^2}{25}\right)\right) \times \min\left(1, \frac{50}{Z_{\text{mixl}}} \frac{50}{z_{\text{mixl}}}\right) \end{aligned} \quad (77)$$

These parameterization is based on a set of very simple assumptions, mainly inferred from the review by [Zondervan \(2007\)](#):

- Coccolithophores are not very abundant in very oligotrophic waters.
- Calcification tends to be maximum at intermediate light levels and decrease at either high and low light levels, around 30 and 4 W m^{-2} respectively.
- Coccolithophores are not ~~very abundant at very low temperature~~ found when the temperature of sea water is below 0 °C.
- Coccolithophores are found in stratified waters. Their abundance decreases when the mixed layer depth (z_{mixl}) exceeds 50 m.
- Maximum levels of coccolithophores are found in the mid-latitudes, where temperature is around 10 °C.

We recognize that this parameterization is quite ad-hoc and may seem arbitrary. But as it will be shown, it simulates reasonable calcification patterns and alkalinity distribution ([yet we recognize that it could be for the wrong reasons](#)). Furthermore, it avoids an explicit modeling of the coccolithophores which is far from being trivial.

Only part (η^I) of the grazed shells are routed to sinking calcite. The rest is taken to dissolve in the acidic guts of zooplankton ([Jansen and Wolf-Gladrow, 2001](#)). This dissolution is still debated. However, observations tend to show that a significant proportion of the sinking shells is lost in the upper ocean, with this being associated with grazing as well as other mechanisms ([Milliman et al., 1999](#)).

The dissolution of calcite is modeled as in [Gehlen et al. \(2007\)](#):

$$\Delta\text{CO}_3^{2-} = \max\left(0, 1 - \frac{\text{CO}_3^{2-}}{\text{CO}_{3,\text{sat}}^{2-}}\right) \quad (78)$$

$$\lambda_{\text{CaCO}_3}^* = \lambda_{\text{CaCO}_3} (\Delta\text{CO}_3^{2-})^{\text{nca}} \quad (79)$$

4.7 The carbonate system

$$\begin{aligned} \frac{\partial\text{DIC}}{\partial t} &= \gamma^Z(1 - e^Z - \sigma^Z) \sum_I g^Z(I)Z + \gamma^M(1 - e^M - \sigma^M) \left(\sum_I g^M(I) + \sum_I g_{\text{FF}}^M(I) \right) M \\ &+ \gamma^M R_{\text{up}}^M + \text{Remin} + \text{Denit} + \lambda_{\text{CaCO}_3}^* \text{CaCO}_3 - P_{\text{CaCO}_3} - \mu^D D - \mu^P P \end{aligned} \quad (80)$$

$$\begin{aligned} \frac{\partial\text{Alk}}{\partial t} &= \theta^{\text{N,C}} \text{Remin} + \theta^{\text{N,C}} (r_{\text{NO}_3}^* + 1) \text{Denit} + \theta^{\text{N,C}} \gamma^Z (1 - e^Z - \sigma^Z) \sum_I g^Z(I)Z \\ &+ \theta^{\text{N,C}} \gamma^M (1 - e^M - \sigma^M) \left(\sum_I g^M(I) + \sum_I g_{\text{FF}}^M(I) + \theta^{\text{N,C}} \gamma^M R_{\text{up}}^M \right) M \\ &+ \theta^{\text{N,C}} \mu_{\text{NO}_3}^P P + \theta^{\text{N,C}} \mu_{\text{NO}_3}^D D + \theta^{\text{N,C}} N_{\text{fix}} + 2\lambda_{\text{CaCO}_3}^* \text{CaCO}_3 \\ &+ \theta^{\text{N,C}} \Delta(\text{O}_2) (r_{\text{NH}_4}^* - 1) \lambda_{\text{NH}_4} \text{NH}_4 - \theta^{\text{N,C}} \mu_{\text{NH}_4}^P P - \theta^{\text{N,C}} \mu_{\text{NH}_4}^D D \end{aligned}$$

$$-2\theta^{N,C}\text{Nitrif} - 2P_{\text{CaCO}_3} \quad (81)$$

All terms in the above equations have been described previously in this document. In addition to these biogeochemical fluxes, the ocean exchanges CO_2 with the atmosphere at the sea-surface. The gas exchange coefficient is computed from the relationship proposed by [Waninkhof \(1992\)](#). No exchange is allowed with the atmosphere across sea-ice:

$$k_{\text{gCO}_2} = k'_{\text{gCO}_2} \times (1 - \%_{\text{ice}}) \quad (82)$$

where $\%_{\text{ice}}$ is the concentration of sea-ice which varies between 0 and 1. The carbonate chemistry follows the OCMIP protocols (more information at <http://ocmip5.ipsl.jussieu.fr/OCMIP/>) except that it has been simplified to reduce the computing cost: alkalinity only includes carbonate, borate and water (H^+ , OH^-).

Atmospheric $p\text{CO}_2$ can be set as an external tunable parameter via a namelist parameter or read from a file. Its value is uniform over the global ocean (no spatial gradient) and is not allowed to vary in response to the air-sea fluxes. This means that PISCES does not include an interactive atmospheric (box or more complex) model (although this functionality can be added very easily). Finally, the impact of atmospheric pressure on $p\text{CO}_2$ can be accounted for by setting the Boolean `ln_presatm` to true in the namelist. In that case, the 2-D spatial distribution of atmospheric pressure should be read in a file.

4.8 Oxygen

$$\begin{aligned} \frac{\partial \text{O}_2}{\partial t} = & \text{O}_2^{\text{ut}}(\mu_{\text{NH}_4}^P P + \mu_{\text{NH}_4}^D D) + (\text{O}_2^{\text{ut}} + \text{O}_2^{\text{nit}})(\mu_{\text{NO}_3}^P P + \mu_{\text{NO}_3}^D D) + \text{O}_2^{\text{nit}} N_{\text{fix}} \\ & - \text{O}_2^{\text{ut}} \gamma^Z (1 - e^Z - \sigma^Z) \sum_I g^Z(I) Z - \text{O}_2^{\text{ut}} \gamma^M (1 - e^M - \sigma^M) \left(\sum_I g^M(I) \right. \\ & \left. + \sum_I g_{\text{FF}}^M(I) \right) M - \text{O}_2^{\text{ut}} \gamma^M R_{\text{up}}^M - \text{O}_2^{\text{ut}} \text{Remin} - \text{O}_2^{\text{nit}} \text{Nitrif} \end{aligned} \quad (83)$$

In this equation, the stoichiometric ratio O_2^{ut} represents the change in oxygen relative to carbon when ammonium is converted to organic matter, whereas O_2^{nit} denotes the consumption of oxygen during nitrification. Their values have been set respectively to 131/122 and 32/122 so that the typical Redfield ratio for oxygen is equal to 1.34 as proposed by [Kortzinger et al. \(2001\)](#).

Oxygen is exchanged with the atmosphere using the parameterization of [Wanninkhof \(1992\)](#) to compute the gas exchange coefficient. The atmospheric concentration of oxygen is constant over time and space and cannot be specified by the user. As for CO_2 , no air-sea fluxes are allowed when the ocean is covered by sea-ice (see Eq. 82).

4.9 External supply of nutrients

Nutrients are supplied to the ocean from five different sources: atmospheric dust deposition, rivers, sea ice, sediment mobilization, and hydrothermal vents.

4.9.1 Atmospheric deposition

The model can include the atmospheric supply of Fe, Si, P and N. The former three sources (Fe, Si and P) are dependent on each other as they are computed from the same dust input file. They are activated in PISCES by setting the Boolean `ln_dust` to true. Otherwise, no atmospheric source of Fe, P and Si is prescribed. Furthermore, in that case, the dust concentration in the ocean (used for instance in Eq. 50) is set to 0. The iron content of dust is set to a constant value specified in the namelist. Its default value is 3.5 % which is the average content of crustal material (e.g., [Taylor and McLennan, 1985](#); [Jickells and Spokes, 2001](#); [Jickells et al., 2005](#)). The solubility of dust iron in sea water can be either set to a constant value in the namelist or can be read from a file if `ln_solub` is set to true. Once it has left the surface layer, particulate inorganic iron from dust is still assumed to experience dissolution. The dissolution rate is computed assuming that mineral particles sink at a constant speed specified in the namelist and that about 0.01 % of the particulate iron dissolves in a day ([Bonnet and Guieu, 2004](#)). This is equivalent to a remineralization length scale of 20 000 m if the sinking speed is set to a typical value of 2 m day^{-1} , of the same order as the length scale prescribed for the same process by [Moore et al.](#)

(2004). Atmospheric deposition of Si is also considered following Moore et al. (2002b) and is restricted to the first layer of the model. Atmospheric deposition of P is computed from dust deposition assuming that the total phosphorus content of dust is 750 ppm (Mahowald et al., 2008) and that the solubility in surface sea water is 10 % (Ridame and Guieu, 2002; Mahowald et al., 2008). As for Si, deposition is restricted to the first level of the ocean model. Atmospheric deposition of N is treated separately from the deposition of the other nutrients and can be activated in the model by the boolean `ln_ndepo`. All nitrogen deposited at the ocean surface is ~~supposed~~ assumed to dissolve. We made the quite strong assumption for all nutrients that sea ice does not alter the deposition fluxes.

The dust (Dust) concentration in the ocean is modeled in a very simplistic way in PISCES. It is computed from dust deposition assuming a constant sinking speed (the same as the sinking speed used to compute iron dissolution from dust in the interior of the ocean). Furthermore, dust is not transported by the ocean currents. This assumption is made in PISCES to avoid adding another prognostic tracer in the model. As a consequence, the concentration of dust is computed as:

$$\text{Dust} = \frac{D_{\text{dust}}}{w_{\text{dust}}} \quad (84)$$

where D_{dust} is dust deposition at the surface and w_{dust} is the prescribed sinking speed of dust.

4.9.2 River discharge

River discharge is activated by setting the boolean variable `ln_river` to true in the namelist. The river discharge of the different elements is then read from a file that must be provided in that case by the user. The river supply of DIN, DIP, DON, DOP, Si, DIC, Alkalinity and DOC need to be provided. As DON, DOC, and DOP are not separately modeled in PISCES (fixed stoichiometry), dissolved organic matter is assumed to remineralize instantaneously at the river mouth and thus, DON, DOP, and DIC are added to DIN, DIP and DIC, respectively. As a default in PISCES, river supply of all elements but DIC and Alkalinity is taken from the GLOBAL-NEWS2 datasets (Mayorga et al., 2010). For DIC and Alkalinity, we use results from

the Global Erosion Model (GEM) of Ludwig et al. (1996), neglecting the POC delivery as most of it is lost in the estuaries and in the coastal zone (Smith and Hollibaugh, 1993). All fields are interpolated onto the ORCA grid and colocalized with the river runoff prescribed in the physical model. Iron is also delivered to the ocean by rivers. The amount of supplied iron is computed from the river supply of inorganic carbon, assuming a constant Fe/DIC ratio. This ratio is determined so that the total Fe supply equals $1.45 \text{ Tg Fe yr}^{-1}$ as estimated by Chester (1990).

4.9.3 Reductive mobilization of iron from marine sediments

Reductive mobilization of iron from marine sediments have been recognized as a significant source to the ocean (Johnson et al., 1999; de Baar and de Jong, 2001; Moore et al., 2004). Fe concentrations in the sediment pore waters are often several orders of magnitude larger than in the seawater. A large part of the iron released to the ocean either by diffusion or by resuspension is likely to be oxidized in insoluble forms and trapped back to the sediments, at least in oxygenated waters (de Baar and de Jong, 2001). Yet, some of this iron should escape as observations clearly show increasing concentration gradients of particulate and dissolved iron toward the coastal zones. Unfortunately, almost no quantitative information is available to parameterize this potentially important source. Observations from benthic chambers indicate that this source may be controlled by the oxygen concentrations overlying the sediments (Raiswell and Anderson, 2005; Severmann et al., 2010) and perhaps the magnitude of the organic carbon export to the sediments (Elrod et al., 2004). Such potential relationships are not yet embedded in PISCES.

In a way similar to Moore et al. (2004), we apply a maximum constant iron source from the sediments. Since anoxic sediments are more likely to release iron to the seawater, we have modulated this source by a factor (F_{sed}) computed from the metamodel of Middelburg et al. (1996):

$$z_{F_{\text{sed}}} = \min \left(8, \left(\frac{z}{500 \text{ m}} \right)^{-1.5} \right) \quad (85a)$$

$$\zeta_{\text{Fsed}} = -0.9543 + 0.7662 \ln(z_{\text{Fsed}}) - 0.235 (\ln(z_{\text{Fsed}}))^2 \quad (85b)$$

$$F_{\text{sed}} = \min(1, \exp(\zeta_{\text{Fsed}})/0.5) \quad (85c)$$

From this metamodel, it is possible to estimate the relative contribution of anaerobic processes to the total mineralization of organic matter in the sediments, and thus to have an indication on how well the sediment is oxygenated (Soetaert et al., 2000). Our modulation factor is simply set equal to this relative contribution. The maximum iron flux from the sediments has been set by default to $12 \mu\text{mol Fe m}^{-2} \text{d}^{-1}$ by adjusting the modeled iron distribution to the few iron observations available over the continental margins. This value is ~~close~~ identical to that used by Moore et al. (2004) in their model (~~2~~). The maximum iron flux constant can be specified in the namelist and thus, may be changed from the default value by the user.

Unfortunately, as a consequence of the relatively coarse resolution of ORCA2, the model bathymetry is not able to correctly represent the critical spatial scales of the ocean bathymetry. An example is the continental shelves, which typically have a width scale of 10–30 km, which can be approximately an order of magnitude less than the horizontal resolution of the model. In order to take sub-model gridscale bathymetric variations into account in the Fe source function, the model grid structure has been compared with the high-resolution ETOPO5 dataset. An algorithm was developed whereby for each and every horizontal grid cell, the corresponding region in the ETOPO5 dataset is considered. For each vertical level in the model corresponding to a particular horizontal gridpoint, the corresponding ocean bottom area from ETOPO5 (in fractional units) is saved, with the end result being a three dimensional array containing an equivalent area for the bottom bathymetry of the ocean for the ETOPO5 dataset. The iron flux computed as described above is then multiplied by this fractional area $\%_{\text{sed}}$ (which varies between 0 and 1):

$$F_{\text{Fe}}^{\text{sed}} = F_{\text{Fe,max}}^{\text{sed}} \times F_{\text{sed}} \times \%_{\text{sed}} \quad (86)$$

[This corresponds to a global flux of 34 Gmol Fe yr⁻¹.](#)

5 4.9.4 Iron from hydrothermalism

Recent studies have shown that hydrothermalism may deliver to the deep ocean a significant amount of dissolved iron (e.g., Mackey et al., 2002; Boyle and Jenkins, 2008; Bennett et al., 2008; Toner et al., 2009). Despite very large uncertainties, this source has been estimated, based on discrete data and a model, to 3 to 9×10^8 mol Fe yr⁻¹ globally (Bennett et al., 2008; Tagliabue et al., 2010). In PISCES, this source is included following the modeling study by Tagliabue et al. (2010) and may be activated by setting the boolean `ln_hydrofe` to true. The hydrothermal flux of iron has been computed based on observed correlations between ³He and dFe (Boyle et al., 2005; Boyle and Jenkins, 2008) and using a data compilation of dFe/³He (see the Supplement of Tagliabue et al., 2010). Then, the spatial distribution of this flux has been derived from previous modeling works on ³He, which relate the ³He flux to the ridge-spreading rates (Farley et al., 1995; Dutay et al., 2004). 0.2% of the delivered iron is ~~supposed~~assumed to be soluble.

4.9.5 Iron from sea ice

The last external source of nutrients which is taken into account in PISCES is the exchange of iron between the ocean and the sea ice associated with formation and melting. This source is activated by setting the boolean variable `ln_ironice` to true. The receding ice-edge is often characterized by intense phytoplankton abundance which can be explained by ocean stratification promoted by the melting of sea ice (Smith and Nelson, 1985) as well as the releases of iron accumulated in sea ice during winter (Sedwick and Di Tullio, 1997; Tagliabue and Arrigo, 2006). Measurements in sea ice have found iron concentrations of more than one order of magnitude higher than in adjacent sea water (Lannuzel et al., 2007, 2008). About 90% of this iron has been shown to be of oceanic origin (Lannuzel et al., 2007). Thus, iron is taken up from sea water when ice forms and is released back to the ocean when it melts. Lancelot et al. (2009) have studied the impact of this source in the Southern Ocean and shown that it is of primary importance in the seasonal ice zone. Their approach relies on the modeling of iron concentration within sea ice. In PISCES, we have simplified this model by assuming that iron concentration

in sea ice is constant. In that case, the iron fluxes between the ocean and the sea ice can be computed from the water fluxes between these two reservoirs:

$$F_{\text{Fe}}^{\text{ice},-} = \min(0, -\text{EP}_{\text{oi}}) \times \text{Fe} \quad (87\text{a})$$

$$F_{\text{Fe}}^{\text{ice},+} = \max(0, -\text{EP}_{\text{oi}}) \times \text{Fe}_{\text{ice}} \quad (87\text{b})$$

$$10 \quad F_{\text{Fe}}^{\text{ice}} = F_{\text{Fe}}^{\text{ice},-} + F_{\text{ice}}^{\text{ice},+} \quad (87\text{c})$$

where EP_{oi} is the water flux (in $\text{kg m}^{-2} \text{s}^{-1}$) from the ice to the ocean and Fe_{ice} is the iron concentration in sea ice which has been found to be of the order of 10 nmol L^{-1} . In this equation, $F_{\text{Fe}}^{\text{ice},-}$ is thus the loss of iron from the ocean when sea ice forms and $F_{\text{ice}}^{\text{ice},+}$ is the release of iron to ocean when sea ice melts. It should be noted here that since we don't model iron in sea ice, the exchange of iron between both reservoirs is not conservative. [In the model configuration presented here, ice represents a net source of iron of \$0.024 \text{ Gmol Fe yr}^{-1}\$.](#)

4.10 Bottom boundary conditions

At the bottom of the ocean, the exchange between the sediments and the ocean can be represented either with or without a sediment model. The sediment model is activated by using the cpp key `key_sed`. This model will not be described in this document. It is basically identical to the model of [Heinze et al. \(1999\)](#) with some modifications as described by [Gehlen et al. \(2006\)](#). The main modification is the addition of denitrification to the set of early diagenetic reactions. Parameter values are identical to those in [Heinze et al. \(1999\)](#).

25 When the sediment model is not activated, very basic but different treatments are applied at the bottom of the ocean depending on the tracer considered. For biogenic silica, the amount of particulate material that is permanently buried in the sediments is [supposed-assumed](#) to exactly balance the external input from dust deposition and river discharge, described in the previous section. Then, we assume that the part of biogenic silica that is not permanently buried redissolves back to the water column instantaneously.

5 For particulate organic carbon, we first determine the proportion of organic matter reaching the seafloor that is permanently buried. The burial efficiency is computed using the algorithm

proposed by [Dunne et al. \(2007\)](#):

$$E_{\text{burial}} = 0.013 + \frac{0.53F_{\text{OC}}^2}{(7.0 + F_{\text{OC}})^2} \quad (88)$$

where E_{burial} is the burial efficiency and F_{OC} is the flux of organic carbon at the bottom (in $\text{mmol C m}^{-2} \text{d}^{-1}$). We then use the metamodel by [Middelburg et al. \(1996\)](#) to determine the proportion of degradation of the remaining organic matter that is due to denitrification:

$$\begin{aligned} \log(P_{\text{denit}}) = & -2.2567 - 1.185 \log(F_{\text{OC}}) - 0.221 (\log(F_{\text{OC}}))^2 - 0.3995 \log(\text{NO}_3) \log(\text{O}_2) \\ & + 1.25 \log(\text{NO}_3) + 0.4721 \log(\text{O}_2) - 0.0996 \log(z) \\ & + 0.4256 \log(F_{\text{OC}}) \log(\text{O}_2) \end{aligned} \quad (89)$$

where the tracer concentrations are in $\mu\text{mol L}^{-1}$ and F_{OC} is the flux of organic carbon at the bottom (in $\mu\text{mol cm}^{-2} \text{d}^{-1}$). In this equation, oxygen and nitrate concentrations are not allowed to be below $10 \mu\text{mol L}^{-1}$ and $1 \mu\text{mol L}^{-1}$ respectively. Then, the fluxes of nitrate and oxygen to the sediment as a consequence of denitrification and oxic degradation, respectively, can be computed:

$$F_{\text{NO}_3}^{\text{denit}} = R_{\text{NO}_3} P_{\text{denit}} F_{\text{OC}} \quad (90a)$$

$$F_{\text{O}_2}^{\text{oxic}} = \text{O}_2^{\text{ut}} (1 - P_{\text{denit}}) F_{\text{OC}} \quad (90b)$$

Particulate organic carbon which has been degraded by denitrification and oxic processes is released in the bottom box as ammonium.

Calcite is treated specifically embedded in PISCES. The preservation of calcite in the sediments is represented as a function of the saturation level of the overlying waters:

$$\%_{\text{CaCO}_3} = \min \left(1, 1.3 \frac{0.2 - \Omega}{0.4 - \Omega} \right) \quad (91)$$

where Ω is the calcite saturation level. This relationship has been deduced from the study by [Archer \(1996\)](#). The permanent burial of calcite is modulated by $\%_{\text{CaCO}_3}$. The amount of calcite that is not buried, instantaneously dissolves back to the ocean.

5 **5 Model parameters and their default values**

Tables 1a–1e list model parameters, their respective units and default values as well as a brief description of each of them. Many of these parameters can be specified in the `namelist_pisces` file. As much as possible, the parameter values have been derived from the literature. However, many parameters, such as the mortality rates, are either not constrained at all, or only poorly constrained by the observations. Their values have been adjusted by successive simulations evaluated against the observational datasets presented below.

In addition to the parameters above, PISCES includes a number of control parameters defined as boolean variables that appear in the `namelist` file `namelist_pisces`. These variables either allow to switch between different functional forms or activate additional functionalities. These control parameters are listed in Table 2. Finally, some functionalities, such as the Kriest model of particulate organic matter, require a major reorganization of the code, for instance a change in the number of prognostic variables. In that case, these functionalities are activated through CPP keys which force the model to be recompiled. These CPP keys are listed in Table 3.

6 **6 Model results**

The objective of this section is not to present a full and exhaustive validation of the model results. This has already been presented in a wide range of publications using different configurations of the model (see the Introduction). Here we present instead a brief comparison of PISCES with available observations, in its standard global configuration. This configuration is the default setup available when downloading the code from the NEMO website (the standard `ORCA2_OFF_PISCES` configuration). All the necessary input files can be obtained from this website.

6.1 **6.1 Model setup**

The dynamical state of the ocean has been simulated using the ocean physical model ORCA2-LIM in its version 3.2 (Madec, 2008). This model is based on an ocean general circulation model

OPA9, coupled with the sea ice model Louvain-la-Neuve Ice Model LIM2 (Timmermann et al., 2005). The spatial resolution is about 2° by $2^\circ \cos \Phi$ (where Φ is the latitude) with a focusing of the meridional resolution to 0.5° in the equatorial domain. The model has 30 vertical layers, with an increased vertical thickness from 10 m at the surface to 500 m at 5000 m. Representation of the topography is based on the partial step thicknesses (Barnier et al., 2006; Penduff et al., 2007). Lateral mixing along isopycnal surfaces is performed both on tracers and momentum as in Lengaigne et al. (2003). The parameterization of Gent and McWilliams (1990) is applied poleward of 10° to represent the effects of non-resolved mesoscale eddies. Vertical mixing is parameterized using the turbulent kinetic energy (TKE) scheme of (~~Gaspar et al., 1990~~) Gaspar et al. (1990) as modified by Madec (2008).

The fields used to drive the ocean are identical to those used by Aumont and Bopp (2006). However, the resulting physical circulation state simulated by the ocean model is different as several new parameterizations and new algorithms have been included in ORCA2-LIM. Climatological atmospheric forcing fields have been constructed from various data sets consisting of daily NCEP/NCAR 2 m atmospheric temperature averaged over 1948–2003 (Kalnay et al., 1996), monthly relative humidity (Trenberth et al., 1989), monthly ISCCP total cloudiness averaged over 1983–2001 (Rossow and Schiffer, 1999), monthly precipitation averaged over 1979–2001 (Xin and Arkin, 1997), and weekly wind stress based on ERS satellite product and TAO observations (Menkes et al., 1998). Surface heat fluxes and evaporation are computed using empirical bulk formulas described by Goose (1997). To avoid any strong model drift, modeled sea surface salinity is restored to the monthly WOA01 data set (Conkright et al., 2002) with a nudging timescale of 40 days applied through local freshwater forcing (thereby conserving salt). The ocean dynamical model has been spun up for 200 years, starting from rest and from the climatology of (~~Conkright et al., 2002~~) Conkright et al. (2002) for temperature and salinity.

Phosphate, oxygen, nitrate and silicic acid distributions have been initialized at uniform concentrations inferred from observed climatologies (Garcia et al., 2010). Initial values for dissolved inorganic carbon and alkalinity are taken from the OCMIP guidelines (Orr, 1999). The ecological tracers are initialized uniformly to arbitrary low values. Iron concentrations are set everywhere to 0.6 nM. The model is then spun up offline for 4000 years using the circula-

Discussion Paper | Discussion Paper | Discussion Paper | Discussion Paper | Discussion Paper

tion state predicted by the dynamical model. Atmospheric $p\text{CO}_2$ is set to a preindustrial value of 278 ppm. After this integration, primary productivity as well as CO_2 fluxes drift by less than $0.001 \text{ Gt C yr}^{-1}$. As the external sources and sinks of nutrients are not fully balanced (see the model description), ~~global-mean-the global inventories of Phosphate, Nitrate and Silicate concentrations-~~, Alkalinity and Silicate are restored toward the observed ~~values inventories~~, once a year ~~-on the first of January. In practice, this correction is done by scaling the 3-D concentrations with a constant uniform factor so that the simulated total inventories don't drift away from the observed inventories. Thus, we do not restore the simulated 3-D distributions to 3-D observed fields so that the predicted spatial and temporal patterns are not corrected in any way to better match the observations. However, the predicted global inventories of P, N, Si and alkalinity can not be used to evaluate the model skill since they are not prognostically predicted. Anyhow, this correction is very small corresponding to a relative change in the concentration of the tracers of the order of $1-5 \cdot 10^{-5} \text{ yr}^{-1}$ so that no significant jump is introduced by this technique. The activation of this technique as well as the frequency at which it is applied are controlled by a boolean parameter and a parameter respectively, in the namelist file `namelist_pisces` (see Table 2).~~

6.2 Global budget

Table 4 presents the global carbon budget as simulated by PISCES, when embedded in ORCA2-LIM. The annual net predicted primary production is 44 Gt C yr^{-1} . This value falls on the lower bound of the broad estimates given by satellite observations which give values between 37 to 67 Gt C yr^{-1} (Longhurst et al., 1995; Antoine et al., 1996; Behrenfeld and Falkowski, 1997; Behrenfeld et al., 2005). Using PISCES in a higher resolution model would certainly produce a significantly larger number as mesoscale and submesoscale processes have been shown to stimulate biological productivity (McGillicuddy et al., 1998; Oschlies and Garçon, 1998; Lévy et al., 2001) and coastal regions, characterized by a intense primary productivity, are not properly resolved by the coarse grid.

About 17% of the primary production is due to diatoms. Global estimates of the contribution of diatoms to total production are rather uncertain and broad. Nelson et al. (1995) have

Discussion Paper | Discussion Paper | Discussion Paper

suggested that diatoms may be responsible for up to 40 % of the total primary production. However, as discussed by [Aumont and Bopp \(2006\)](#), this value is certainly overestimated. In recent years, algorithms, which attempt to retrieve the composition of phytoplankton from space, have been developed (e.g., [Alvain et al., 2005](#); [Uitz et al., 2006](#); [Hirata et al., 2008](#); [Brewin et al., 2010](#)). Only a few of these methods give quantitative estimates of the contribution of the different species or size-classes to total biomass or primary productivity ([Brewin et al., 2011](#)). The estimated global contribution of diatoms from these methods ranges from as low as 7 % to as high as 32 % of the total phytoplankton ([Uitz et al., 2010](#); [Hirata et al., 2011](#)) (if one assumes crudely that microphytoplankton are effectively equivalent to diatoms). Finally, ocean biogeochemical models predict the contribution of diatoms to be between 15 to 30 % (e.g., [Moore et al., 2002a](#); [Aumont et al., 2003](#); [Dutkiewicz et al., 2005](#); [Yool et al., 2011](#)).

Export production at 150 m is estimated to be 6.9 Gt C yr^{-1} . 86 % of this export is related to settling particles (one third by the small sinking particles and two third by the fast sinking particles). The remainder is due to vertical advection and diffusion of dissolved organic carbon, which occurs mainly in the mid-ocean gyres (vertical advection) and in the high latitude regions during winter (vertical diffusion). Constraining export production is rather difficult, if not impossible, considering the very broad range given by estimates either based on models or observations and the different definitions of export production, in particular the depth horizon at which it is estimated (e.g., [Eppley and Peterson, 1979](#); [Schlitzer, 2000](#); [Moore et al., 2002a](#); [Yool et al., 2011](#)). Mesozooplankton grazes about 9 % of total primary production. This value is close to other estimates either based on observations ([Calbet, 2001](#)) or models ([Moore et al., 2002a](#); [Buitenhuis and Geider, 2010](#)). Total grazing by mesozooplankton is predicted to be $11.2 \text{ Gt C yr}^{-1}$ by PISCES, quite similar to the value of $10.4 \pm 3.7 \text{ Gt C yr}^{-1}$ estimated by [Hernández-León and Ikeda \(2005\)](#) for the global respiration of mesozooplankton in the upper 200 m of the ocean. About 80 % of total primary production, i.e. $35.8 \text{ Gt C yr}^{-1}$, is consumed up by microzooplankton above the upper bound of the $25\text{--}33 \text{ Gt C yr}^{-1}$ given by [Buitenhuis et al. \(2010\)](#) when extrapolating observations. Despite estimates of grazing by microzooplankton are quite badly constrained, this might suggest that it is overestimated in the model.

Table 5 shows the calcite and silicon budgets for the upper 150 m of the ocean. Production of calcite and export at 150 m are simulated to be, respectively, about 1.6 and 0.8 Gt C yr^{-1} by PISCES. These numbers fall within the limits of the quite large range of $0.4\text{--}1.8 \text{ Gt C yr}^{-1}$ estimated either for global calcification or export of PIC (Murnane et al., 1999; Lee, 2001; Moore et al., 2002a; Balch et al., 2007; Berelson et al., 2007). For silicate, the model predicts a vertical export of biogenic silicate of $106 \text{ Tmol Si yr}^{-1}$. This value is within the $105 \pm 17 \text{ Tmol Si yr}^{-1}$ estimated for the global ocean (Tréguer and De La Rocha, 2012). Global production of biogenic silica by diatoms is $146 \text{ Tmol Si yr}^{-1}$ in our model. This value is quite low compared to the $239 \text{ Tmol Si yr}^{-1}$ given by Tréguer and De La Rocha (2012). Consequently, about About 27 % of biogenic silica dissolves in the top 150 m of the ocean, half the estimate of Nelson et al. (1995) and Tréguer and De La Rocha (2012). However, as already mentioned, because of its coarse resolution, the physical model configuration does not properly resolve the coastal zones. For the open ocean only (in a strict sense), Tréguer and De La Rocha (2012) estimated biogenic silica production to be about $103 \text{ Tmol Si yr}^{-1}$. Not surprisingly then, considering the limitations due to the spatial resolution, our modeled estimate is between the open ocean and global values. The mean Si/C for uptake of diatoms as predicted by PISCES is thus $0.270.23$, which is high relative to the optimal Si/C of 0.13 (Brzezinski, 1985). This suggests thus that over most of the ocean, diatom cells are stressed, not a very surprising result. Furthermore, a large part of the biogenic silica production occurs within the Southern Ocean, a region where diatom cells are very heavily silicified (Baines et al., 2010).

Table 6 presents the global nitrogen budget as simulated by PISCES. River discharge and Atmospheric deposition of nitrogen are given by the prescribed input fields to PISCES. By definition, burial in the sediments is set exactly equal to river discharge. Nitrogen fixation is predicted to be $111.8 \text{ Tg N yr}^{-1}$. This value is close to the mean value of about 140 Tg N yr^{-1} estimated from direct observations or nutrients analysis (Capone et al., 1997; Deutsch et al., 2007). Figure 6 shows a comparison between the spatial distribution of observed nitrogen fixation rates from the MAREDAT project and that as simulated by PISCES. This indicates that, despite a quite simplistic formulation, the model is able to capture the main observed patterns, at least on an annual mean basis. Modeled denitrification in the water column and in the sediments

are about 78 and 93 TgN yr^{-1} , respectively. Sediment denitrification estimates are significantly higher, in the range of 130–300 TgN yr^{-1} (Codispoti et al., 2001; Galloway et al., 2004; Gruber, 2004). However, considering the coarse spatial resolution of the model, this is expected as most of benthic denitrification occurs over the continental margins. The sources and sinks of nitrogen are slightly unbalanced, with the sources exceeding the sinks by about 21 TgN yr^{-1} .

6.3 Modeled tracer distributions

6.3.1 Chlorophyll

The modeled chlorophyll distribution is compared to GLOBCOLOUR satellite observations for two seasons on Fig. 7. The seasons have been defined to roughly correspond to bloom periods in the high latitudes. The observed patterns are qualitatively reproduced by the model. Slightly too low chlorophyll concentrations are simulated in the subtropical gyres. This discrepancy may be explained by the lack of acclimation dynamics to oligotrophic conditions in the model or by the assumption of constant stoichiometry either in phytoplankton or in organic matter (Ayata et al., 2013). Chlorophyll concentrations are quite strongly underestimated in the equatorial Atlantic and in the Arabian Sea. In the latter region, mesoscale and submesoscale processes have been shown to be of critical importance (Lee et al., 2000; Kawamiya, 2001; Hood et al., 2003). A model study, using PISCES coupled to a higher resolution version of NEMO, has been shown to simulate chlorophyll distribution in much better agreement with the observations (Koné et al., 2009). Chlorophyll concentrations are high in the eastern boundary upwelling systems. The sedimentary source of iron plays a critical role in these systems. When this iron source is not included in models, modeled chlorophyll concentrations are much lower (Aumont and Bopp, 2006; Moore and Braucher, 2008).

In two of the three main HNLC regions, i.e., the equatorial Pacific and the eastern subarctic Pacific, the model succeeds in reproducing the moderate chlorophyll concentrations. In spring, chlorophyll levels are strongly overestimated east of Japan. As in all coarse resolution models, the ocean circulation in this region is not correctly represented with an incorrect trajectory of the Kuroshio current (i.e., Gnanadesikan et al., 2002; Dutkiewicz et al., 2005; Aumont

and Bopp, 2006). Simulated mixed layer depths are too deep in winter and as a consequence, the spring bloom is very strong (similar features occur in the North Atlantic). In the equatorial Pacific ocean, a minimum threshold value has been imposed on iron (0.01 nmol L^{-1}) in the model. If not used, chlorophyll concentrations become much too low on both sides of the Equator, resulting in ~~a strong an~~ accumulation of macronutrients and a poleward migration of the southern (northern) boundary of the northern (southern) subtropical gyre (see Figure 5 in Tagliabue et al. (2009a)). The existence of such threshold suggests that either a minor but regionally important source of iron is missing in PISCES (for instance the dissolution of particulate inorganic iron) or that the standard iron chemistry is too simple (Tagliabue et al., 2009a; Tagliabue and Völker, 2011).

In the Southern Ocean, the third and largest of the principal HNLC regions, chlorophyll concentrations appear to be strongly overestimated by the model when evaluated against satellite-derived observational products, especially during summer. Furthermore, the increase in phytoplankton in late spring and early summer occurs too early. However, numerous studies comparing satellite chlorophyll to in situ data have shown that the standard algorithms used to deduce chlorophyll concentrations from reflectance tend to underestimate in situ observed values by a factor of about 2 to 2.5, especially for intermediate concentrations (e.g., Dierssen and Smith, 2000; Korb et al., 2004; Garcia et al., 2005; Kahru and Mitchell, 2010). ~~In Fig. ??, we have corrected chlorophyll concentrations from GLOBCOLOUR (based on algorithm OC2V4) using the polynomial relationship proposed by Dierssen and Smith (2000). From this perspective, modeled values are now underestimated, especially in the Atlantic sector of the Southern Ocean.~~ Clearly, evaluating the model in the Southern Ocean is quite challenging and requires a more thorough systematic analysis of both the model and the available datasets.

6.3.2 Iron

Figure 8 shows the distribution of iron at three different depth ranges for the model and for the observations. The observational distributions come from the recently published database of Tagliabue et al. (2012) augmented with about 1000 recent observations. The dataset can be

5 downloaded from <http://pcwww.liv.ac.uk/~atagliab>. A complete and exhaustive validation of the model is made difficult by the relative sparsity of the data.

As expected, the highest concentrations of iron in the open ocean are found in the subtropical North Atlantic ocean and in the Arabian Sea. Those high values are produced by the enhanced dust deposition, mainly emanating from the Sahara desert. The model tends to underestimate
10 the maximum values found in both basins. Interestingly, the local minimum, which is observed west off Mauritania just below the maximum Saharan dust plume, is well captured by the model. Such a minimum is explained by the combination of very low solubilities of the iron contained in the Saharan dust particles when they are close to their source region (Bonnet and Guieu, 2004; Luo et al., 2005) with enhanced scavenging by the dust particles deposited at the ocean surface
15 (Wagener et al., 2010). Very high iron concentrations, typically above 1 nmolL^{-1} are both observed and modeled along the coasts and over the continental margins as a result of sediment mobilization. As already mentioned in the previous section, this strong source of iron sustains the high productivity observed along the coasts (Johnson et al., 1999), in the eastern boundary upwelling systems (Bruland et al., 2005) but also downstream of the islands, especially in the
20 Southern Ocean (Blain et al., 2007; Pollard et al., 2007; Korb et al., 2008). In the rest of the open ocean, iron concentrations are typically low, generally below 0.2 nmolL^{-1} , especially in the HNLC regions. PISCES tends to exaggerate these low concentrations.

Iron concentrations increase with depth due to the remineralization of organic particles settling from the surface waters (Johnson et al., 1997; Moore and Braucher, 2008). However, ex-
25 cept near the coasts, concentrations rarely exceed 1 nmolL^{-1} . Again, PISCES captures the main observed patterns both at intermediate depths and in the deep ocean. In the Atlantic ocean and in the Arabian Sea, iron concentrations remain relatively elevated at intermediate depth in the observations and in the model. In the model, these high values are due to the slow but significant release of iron by the dust particles which sink out from the surface. In the Pacific ocean, the coastal signature extends far beyond the coastal domain. For instance, it has been proposed as a potential explanation for the episodic blooms observed at station P in the northeastern subarctic Pacific ocean (Lam et al., 2006; Misumi et al., 2011). In the deepest waters of the Pacific and
5 Indian oceans, iron concentrations tend to decrease to the bottom of the ocean and they often fall

below 0.6 nmol L^{-1} . Despite the fact that ligands concentrations in seawater are highly variable, they are typically larger than this value which is the uniform ligand concentration chosen in the model experiment shown here (e.g., [Wu and Luther, 1995](#); [Boyé et al., 2001, 2003](#); [Hunter and Boyd, 2007](#); [Ibsanmi et al., 2011](#)). The model explains this decrease by the aggregation of iron colloids which are transferred to the particulate pool and thus sink out of the ocean as hypothesized by several studies ([Wu et al., 2001](#); [Ye et al., 2009](#); [Gledhill and Kirsten, 2012](#)). The lowest iron concentrations in the intermediate and deep ocean are found in the Southern Ocean. Iron concentrations slowly increase with depth to reach about 0.4 nmol L^{-1} in the deep ocean. Higher values are found along Antarctica due to sediment mobilization.

6.3.3 Nutrients, oxygen, alkalinity and DIC

In this section, the simulated distributions of macro-nutrients, oxygen, alkalinity and DIC are evaluated against available observations. The observations comprise the World Ocean Atlas 2009 for nutrients and oxygen ([Garcia et al., 2010](#)), and the GLODAP database for DIC and alkalinity ([Key et al., 2004](#)).

Figures 9 and 10 show the surface distributions of Nitrate and Silicate and zonally averaged sections in the Atlantic and Pacific Oceans. At the surface, the model compares quite well with the observations, especially for Nitrate. Nitrate concentrations seem to be slightly overestimated along the Antarctic coast. However, as most of the data have been collected during the productive season in this region, the climatology is likely to be biased toward low values. The surface silicate distribution is less well represented by PISCES, in particular in the Southern Ocean. The silicate front (defined as the latitude at which silicate becomes exhausted) is located too far North in the model. At depth, both modeled nutrients exhibit the same deficiencies. In the Atlantic ocean, concentrations in the deep ocean are strongly overestimated. Too shallow North Atlantic deep waters (NADW), with strongly underestimated transport simulated for lower NADW, accounts for this problem ([Arsouze et al., 2008](#); [Griffies et al., 2009](#); [Smith et al., 2010](#)). As a result, Antarctic Bottom waters, characterized by high silicate and nitrate concentrations, tend to dominate over too large part of the deep Atlantic Ocean. In the Pacific Ocean,

5 both nitrate and silicate concentrations are underestimated in the deep waters of the Northern Hemisphere.

In Fig. 11, the modeled oxygen distribution is evaluated against observations. Not surprisingly, the surface distribution compares quite well to the observations as oxygen is close to its solubility value and is thus strongly constrained by sea surface temperature. At depth, the main deficiency is the overestimation of oxygen concentrations in the Pacific ocean. Ventilation along
10 Antarctica, mainly in the Ross and Weddell Seas, is too strong in the physical model. Inspection of the simulated mixed layer depths show that the mixed layer reaches the bottom of the ocean at several locations along Antarctica (not shown), which is not realistic (de Boyer-Montégut et al., 2004). The nearly homogeneous oxygen concentrations south of 60° S are a consequence
15 of this too intense winter mixing, which thus ventilates the deep ocean with too much oxygen.

Figures 12 and 13 display the modeled and observed distributions of DIC and alkalinity at the surface and along zonally averaged sections in both the Atlantic and the Pacific. ~~The modeled fields compare favorably to observations. Modeled DIC does not include the anthropogenic perturbation since atmospheric CO₂ was set to its pre-industrial value. We have estimated the observed pre-industrial distribution of DIC as the difference between total DIC and anthropogenic carbon, which are both available in GLODAP Key et al. (2004).~~ It should be also mentioned here that no observations were available north of 60° N. Values north of this latitude have been extrapolated for plotting purpose. At the surface, several modeled features are not visible in the observations. Very low alkalinity and DIC concentrations are predicted in the Bay of Bengal, in
20 the Gulf of Guinea, close to the Indonesian Islands and generally at the mouths of the tropical rivers. The lack of observations in these regions may explain this difference, as the GLODAP database is based on a rather coarse sampling coverage. In the deep ocean, the main deficiencies noticed for the macro-nutrients are apparent in the simulated distributions.

6.4 Skill assessment

In this section, we quantitatively estimate the model performance using Taylor diagrams (Taylor, 2001). Taylor diagrams evaluate both the correlation normalized by the observed SD (circumference axis) and the relative variability (radial axis) of model and observations. The dis-

5 tance between the model points and the (1,1) coordinate point (defined as the reference point) is equal to the standard root mean error, normalized by the observed SD. The closer the model is to the observations, the closer the points should be to the reference point. Although a number of means and diagnostics exist (Allen et al., 2007; Doney et al., 2009; Vichi and Masina, 2009), Taylor diagrams have become quite popular as they synthesize, in a quite convenient way, several statistical diagnostics.

10 Figures 14 and 15 show Taylor diagrams for surface chlorophyll and mesozooplankton averaged over the top 150 m of the ocean. The agreement is rather modest for both variables, especially for mesozooplankton. For chlorophyll, the model performs slightly better for annual-mean distributions, which suggests biases in the representation of the seasonal cycle. The Southern Ocean exhibits the poorest agreement. In particular, the model tends to strongly underestimate the spatial variability since the SD is smaller for the annual mean distribution than for seasonally varying fields. In the other basins, the variability is overestimated, especially in the Atlantic ocean where the spring blooms in the subarctic domain are too intense, at least relative to satellite observations (see Fig. 7). Mesozooplankton variability is strongly underestimated by PISCES in all basins. The use of a square closure scheme for mortality may partly explain this bias as this scheme tends to dampen extremes. Preliminary tests with PISCES coupled to the upper-trophic layer model APECOSM (Maury et al., 2007) produce a much greater spatial and temporal variability for mesozooplankton, especially in the high latitudes and along the continental margins.

20 Figure 16 shows Taylor diagrams for nutrients, oxygen, alkalinity and DIC. Overall, except for the carbonate system and iron, the model performs quite well, as expected from the comparison made in the previous section. The poorest agreement is found for both alkalinity and iron. For iron, the model tends to strongly underestimate the spatial variability, both at the surface and in the interior of the ocean. Through a re-inspection of Fig. 8, we can see that this weak bias is not surprising. In particular, the gradients from the coastal regions to the open ocean are generally too small. This suggests that the sediment source of iron is too small and should either be increased and/or made more variable. For the carbonate system, the predicted spatial variability is overestimated, in particular in the interior of the ocean. In fact, the data distri-

10
20
30
40
50
60
70
80
90
100
110
120
130
140
150
160
170
180
190
200
210
220
230
240
250
260
270
280
290
300
310
320
330
340
350
360
370
380
390
400
410
420
430
440
450
460
470
480
490
500
510
520
530
540
550
560
570
580
590
600
610
620
630
640
650
660
670
680
690
700
710
720
730
740
750
760
770
780
790
800
810
820
830
840
850
860
870
880
890
900
910
920
930
940
950
960
970
980
990
1000
1010
1020
1030
1040
1050
1060
1070
1080
1090
1100
1110
1120
1130
1140
1150
1160
1170
1180
1190
1200
1210
1220
1230
1240
1250
1260
1270
1280
1290
1300
1310
1320
1330
1340
1350
1360
1370
1380
1390
1400
1410
1420
1430
1440
1450
1460
1470
1480
1490
1500
1510
1520
1530
1540
1550
1560
1570
1580
1590
1600
1610
1620
1630
1640
1650
1660
1670
1680
1690
1700
1710
1720
1730
1740
1750
1760
1770
1780
1790
1800
1810
1820
1830
1840
1850
1860
1870
1880
1890
1900
1910
1920
1930
1940
1950
1960
1970
1980
1990
2000
2010
2020
2030
2040
2050
2060
2070
2080
2090
2100
2110
2120
2130
2140
2150
2160
2170
2180
2190
2200
2210
2220
2230
2240
2250
2260
2270
2280
2290
2300
2310
2320
2330
2340
2350
2360
2370
2380
2390
2400
2410
2420
2430
2440
2450
2460
2470
2480
2490
2500
2510
2520
2530
2540
2550
2560
2570
2580
2590
2600
2610
2620
2630
2640
2650
2660
2670
2680
2690
2700
2710
2720
2730
2740
2750
2760
2770
2780
2790
2800
2810
2820
2830
2840
2850
2860
2870
2880
2890
2900
2910
2920
2930
2940
2950
2960
2970
2980
2990
3000
3010
3020
3030
3040
3050
3060
3070
3080
3090
3100
3110
3120
3130
3140
3150
3160
3170
3180
3190
3200
3210
3220
3230
3240
3250
3260
3270
3280
3290
3300
3310
3320
3330
3340
3350
3360
3370
3380
3390
3400
3410
3420
3430
3440
3450
3460
3470
3480
3490
3500
3510
3520
3530
3540
3550
3560
3570
3580
3590
3600
3610
3620
3630
3640
3650
3660
3670
3680
3690
3700
3710
3720
3730
3740
3750
3760
3770
3780
3790
3800
3810
3820
3830
3840
3850
3860
3870
3880
3890
3900
3910
3920
3930
3940
3950
3960
3970
3980
3990
4000
4010
4020
4030
4040
4050
4060
4070
4080
4090
4100
4110
4120
4130
4140
4150
4160
4170
4180
4190
4200
4210
4220
4230
4240
4250
4260
4270
4280
4290
4300
4310
4320
4330
4340
4350
4360
4370
4380
4390
4400
4410
4420
4430
4440
4450
4460
4470
4480
4490
4500
4510
4520
4530
4540
4550
4560
4570
4580
4590
4600
4610
4620
4630
4640
4650
4660
4670
4680
4690
4700
4710
4720
4730
4740
4750
4760
4770
4780
4790
4800
4810
4820
4830
4840
4850
4860
4870
4880
4890
4900
4910
4920
4930
4940
4950
4960
4970
4980
4990
5000
5010
5020
5030
5040
5050
5060
5070
5080
5090
5100
5110
5120
5130
5140
5150
5160
5170
5180
5190
5200
5210
5220
5230
5240
5250
5260
5270
5280
5290
5300
5310
5320
5330
5340
5350
5360
5370
5380
5390
5400
5410
5420
5430
5440
5450
5460
5470
5480
5490
5500
5510
5520
5530
5540
5550
5560
5570
5580
5590
5600
5610
5620
5630
5640
5650
5660
5670
5680
5690
5700
5710
5720
5730
5740
5750
5760
5770
5780
5790
5800
5810
5820
5830
5840
5850
5860
5870
5880
5890
5900
5910
5920
5930
5940
5950
5960
5970
5980
5990
6000
6010
6020
6030
6040
6050
6060
6070
6080
6090
6100
6110
6120
6130
6140
6150
6160
6170
6180
6190
6200
6210
6220
6230
6240
6250
6260
6270
6280
6290
6300
6310
6320
6330
6340
6350
6360
6370
6380
6390
6400
6410
6420
6430
6440
6450
6460
6470
6480
6490
6500
6510
6520
6530
6540
6550
6560
6570
6580
6590
6600
6610
6620
6630
6640
6650
6660
6670
6680
6690
6700
6710
6720
6730
6740
6750
6760
6770
6780
6790
6800
6810
6820
6830
6840
6850
6860
6870
6880
6890
6900
6910
6920
6930
6940
6950
6960
6970
6980
6990
7000
7010
7020
7030
7040
7050
7060
7070
7080
7090
7100
7110
7120
7130
7140
7150
7160
7170
7180
7190
7200
7210
7220
7230
7240
7250
7260
7270
7280
7290
7300
7310
7320
7330
7340
7350
7360
7370
7380
7390
7400
7410
7420
7430
7440
7450
7460
7470
7480
7490
7500
7510
7520
7530
7540
7550
7560
7570
7580
7590
7600
7610
7620
7630
7640
7650
7660
7670
7680
7690
7700
7710
7720
7730
7740
7750
7760
7770
7780
7790
7800
7810
7820
7830
7840
7850
7860
7870
7880
7890
7900
7910
7920
7930
7940
7950
7960
7970
7980
7990
8000
8010
8020
8030
8040
8050
8060
8070
8080
8090
8100
8110
8120
8130
8140
8150
8160
8170
8180
8190
8200
8210
8220
8230
8240
8250
8260
8270
8280
8290
8300
8310
8320
8330
8340
8350
8360
8370
8380
8390
8400
8410
8420
8430
8440
8450
8460
8470
8480
8490
8500
8510
8520
8530
8540
8550
8560
8570
8580
8590
8600
8610
8620
8630
8640
8650
8660
8670
8680
8690
8700
8710
8720
8730
8740
8750
8760
8770
8780
8790
8800
8810
8820
8830
8840
8850
8860
8870
8880
8890
8900
8910
8920
8930
8940
8950
8960
8970
8980
8990
9000
9010
9020
9030
9040
9050
9060
9070
9080
9090
9100
9110
9120
9130
9140
9150
9160
9170
9180
9190
9200
9210
9220
9230
9240
9250
9260
9270
9280
9290
9300
9310
9320
9330
9340
9350
9360
9370
9380
9390
9400
9410
9420
9430
9440
9450
9460
9470
9480
9490
9500
9510
9520
9530
9540
9550
9560
9570
9580
9590
9600
9610
9620
9630
9640
9650
9660
9670
9680
9690
9700
9710
9720
9730
9740
9750
9760
9770
9780
9790
9800
9810
9820
9830
9840
9850
9860
9870
9880
9890
9900
9910
9920
9930
9940
9950
9960
9970
9980
9990
10000

bution which has been used to produce the observed climatology is rather coarse (Key et al., 2004). As a consequence, the interpolation procedure strongly smooths the DIC and Alkalinity distribution. Thus, the GLODAP database probably underestimates the real variability of these tracers. To avoid this problem, we should have used a non-interpolated data product as for iron or mesozooplankton. To estimate the potential uncertainty associated with the use of GLODAP, we have used another alkalinity database only available at the surface (Lee et al., 2006). The agreement between the model and this database is much better (see Fig. 16), confirming thus that interpolation in GLODAP potentially leads to a strong underestimate of the real spatial variability.

7 Sensitivity tests with some new parameterizations

A number of new parameterizations has been introduced in the current version of PISCES. The objective of this section is to briefly document the impact of some of these. To do so, we have run a series of sensitivity experiments for a duration of 10 years in which specific parameterizations have been either changed or removed. Table 7 summarizes the different performed experiments. The objective of these tests is not to unequivocally demonstrate that the new formulations improve the model skills but is rather to show the consequences of their utilization on the model behavior.

7.1 Dependence of growth rate on light

In the first two experiments, PAR and LIGHT, the sensitivity of the model results to the dependence of growth rate to light has been tested. In the PAR experiment, PAR is set as a constant fraction of incident shortwave radiation, here 43 %, as usually done in ocean biogeochemical models. Chlorophyll distribution is almost identical to the standard simulation (not shown). Furthermore, global primary production and export production remain almost unchanged (see Table 7). Model results are thus almost insensitive to the variability of the fraction of shortwave radiation that is PAR. In the second experiment, we use an alternative formulation of light lim-

5 itation which corresponds to the standard parameterization as proposed by Geider et al. (1997) (see Eq. 2b). In this formulation, the light saturation parameter E_k directly depends on temperature and nutrient limitation. Thus, since the Q_{10} of phytoplankton is close to 2, E_k is then predicted to be 6 to 8 times smaller in the very high latitudes than in the tropical domain. Furthermore, in the very oligotrophic regions such as the central subtropical gyres, E_k is close to
10 0 as a consequence of a very intense nutrient limitation. In the LIGHT experiment, the initial slopes of $P-I$ curves have been prescribed so that the resulting E_k are identical to those of the standard case at 15 °C for no nutrient limitation.

Figure 17a and b show the difference in chlorophyll between the LIGHT experiment and the standard case for two seasons. The alternative parameterization of light limitation produces
15 changes in surface chlorophyll at both seasons. In the very high latitudes of both hemispheres, surface chlorophyll is strongly increased during the corresponding growing season. The temperature dependence in the alternative parameterization produces lower light saturation parameters and thus, a weaker light limitation. On the contrary, in the mid to high latitudes of both hemispheres, surface chlorophyll is significantly lower, especially in the Southern Ocean and in the
20 Pacific Ocean. The temperature dependence of the light saturation parameter results in a weaker light limitation during Winter. As a consequence, chlorophyll concentrations and primary productivity are predicted to be higher during this season generating a significant consumption and export of nutrients. At the beginning of the growing season, the stock of nutrients in the upper ocean is then lower which leads to weaker and shorter spring blooms. In the very high latitudes,
25 the absence of light during Winter and the presence of sea ice explain the different modeled response. In the low latitudes, the differences are relatively small. Surface chlorophyll concentrations tend to be higher in HNLC and productive regions. The alternative formulation tends to produce a stronger light limitation in the subsurface and thus, reduces the nutrient uptake below the surface. More iron and macronutrients are advected into the surface layer (not shown) which results in higher chlorophyll concentrations and in some cases, in larger productive regions (for instance in the tropical Atlantic Ocean and in the Arabian Sea).

Figure 18 shows the day at which blooms reach their maximum intensity in the SEAWiFS
5 data, in the standard case and in LIGHT. Over the low and mid latitudes as well as in the North

Atlantic Ocean, the timing of the bloom maximum predicted by the standard model is in broad agreement with the satellite data. However, in the central part of the subarctic gyre of the North Pacific, the model simulates a bloom maximum which occurs much too early in the growing season, in January compared to August in the satellite observations. A similar bias is also predicted in part of the Southern Ocean, especially in the eastern part of the three sectors of this ocean. When the alternative parameterization of light limitation is used, the bloom timing remains unchanged over most of the ocean, except in the high latitudes in areas where the winter mixed layer remains relatively shallow. Such result is not surprising because the alternative formulation predicts a much lower light saturation parameter in cold waters which alleviates light limitation at the beginning of the growing season. As a consequence, the bloom occurs earlier in the growing season, which tends to worsen the model behavior in the high latitudes of both hemispheres. In the North Pacific, the strong bias is not modified by the alternative formulation which suggests that this bias is not related to an incorrect description of light limitation. In fact, the model predicts a very strong limitation of phytoplankton growth by iron during summer and thus, simulated chlorophyll concentrations are very low. In Winter, the mixed layer deepens supplying the surface with iron. However, it remains relatively shallow preventing thus phytoplankton from being severely light limited. Chlorophyll concentrations are then maximum during Winter and minimum during summer, which is identical to what is observed in the subtropical gyres, at BATS for instance (Lévy et al., 2005; Fernández I et al., 2005). Yet, it is completely out of phase relative to the observations, suggesting that in that region, the model either strongly overestimates iron limitation during summer or that iron-light co-limitations are incorrectly parameterized in PISCES.

The sensitivity experiment presented here shows that model results are very sensitive to how light limitation is parameterized. Primary production, export production as well as the magnitude of the bloom are strongly impacted by the choice of the formulation describing light limitation of phytoplankton growth. The parameterization proposed by Geider et al. (1997) shares some similarities with the Liebig's law of the minimum. When nutrients are very limiting, light limitation becomes negligible since E_k tends to 0. When light is strongly limiting, nutrients limitation becomes unimportant and growth rate becomes linearly related to light and

Chl/C. The parameterization used in the standard case is similar to the multiplicative description of the limiting factors. As a consequence, the standard parameterization predicts lower phytoplankton growth rates, smaller primary production and less intense blooms. On the other hand, the timing of the bloom maximum is much less sensitive to the formulation of light limitation, except in the strongly stratified areas of the high latitudes. At low latitudes, light limitation at the surface is of secondary importance, despite that light limitation in the subsurface appears to partly control the amount of nutrients supplied to the surface. In the mid and high latitudes, in areas characterized by deep winter mixed layers, the timing of the bloom maximum (but not its magnitude) appears to be virtually insensitive to the description of light limitation. This means that other factors, such as the timing of stratification, drive the timing of the bloom maximum.

7.2 Simple parameterization of cell size

In PISCES, a very basic parameterization of phytoplankton cell size has been developed to compute the values of the half-saturation coefficients for the different nutrients (see Eq. 7). This parameterization is based on the classical hypothesis, supported by observations, that the mean cell size of a phytoplankton community increases as the biomass increases (e.g., [Raimbault et al., 1988](#); [Armstrong, 1994](#); [Hurt and Armstrong, 1996](#)). In the SIZE experiment, this simple parameterization has been removed, i.e. the half-saturation constants are kept constant to their minimum values as specified in Table 1e.

Figure 17c and d display the differences in surface chlorophyll between the SIZE experiment and the standard configuration of the model. The largest differences are simulated in the high latitudes of both hemispheres, during the growing season. A closer inspection of the model results show that the largest changes occur at the end of the spring or Summer bloom, when the exhaustion in nutrients becomes a major limiting factor. In the standard experiment, the cell-size parameterization produces high half-saturation constants during the phytoplankton bloom since they directly depend on the biomass level. Thus, nutrient limitation occurs earlier and is more severe leading to a shorter and less intense bloom. In the eastern boundary upwelling systems, the biomass is also very high. However, unlike in the high latitudes, the phytoplankton biomass is mainly controlled by grazing so that nutrient concentrations are generally much higher than

the values of the high saturation constants. In the subtropical oligotrophic gyres, the impact is negligible since the mean cell size is predicted to be at its minimal value in the standard experiment, which is equivalent to what is imposed in the SIZE experiment.

10 The impact of the cell-size parameterization on nutrients is small, except for silicate in the equatorial Pacific Ocean (not shown). In this region, nanophytoplankton become strongly favored in the SIZE experiment because in the standard case, their cell size is not predicted to be minimum whereas for diatoms, such is the case. When the cell size parameterization is removed, nanophytoplankton biomass increases and completely outcompete diatoms. As a consequence, 15 silicate consumption in the Equatorial Pacific Ocean is strongly reduced which explains the simulated higher values in the SIZE experiment. However, the total chlorophyll concentration is nearly identical because the decrease in diatoms compensates for the increase in nanophytoplankton. Furthermore, the total chlorophyll biomass is regulated by the total supply in iron, whereas the contribution of the different phytoplankton species is driven by their competitive 20 abilities (here specified by the values of their half-saturation constants).

7.3 Food quality and grazing

Food quality may have profound impacts on the grazing activity by zooplankton as discussed by [Mitra et al. \(2007\)](#). When absorbing prey with poor nutritional value, zooplankton may have two different options: (1) increase the retention time of the prey to extract as many metabolites as they can ([Plath and Boersma, 2001](#)), or (2) decrease the retention time of the preys to maintain 25 the highest possible metabolite concentration in the digestive apparatus and thus to increase the probability to absorb valuable compounds ([Tirelli and Mayzaud, 2005](#); [Dutz et al., 2008](#)). In the first case, growth efficiency is increased whereas it is decreased in the second case. In PISCES, poor food quality is assumed to impair gross growth efficiency (e^Z) of both microzooplankton and mesozooplankton based on the stoichiometric ratios of their preys (Fe/C and N/C, see Eq. 27). In the FOOD sensitivity experiment, the effect of food quality on the gross growth efficiency has been removed, i.e. e_N^Z is set to 1.

5 Surface chlorophyll concentrations are almost unaltered when the impact of food quality is removed (see Fig. 17e and f). The only noticeable differences are simulated off the equatorial

Pacific Ocean where very strong iron limitation causes very low Fe/C ratios in phytoplankton. In the FOOD experiment, these low Fe/C ratios do not reduce zooplankton growth efficiency. Grazing pressure on phytoplankton is then higher. The nutrients distributions are also very close to those predicted in the standard experiment. Thus, food quality appears to have minimal consequences on ~~phytoplankton~~-chlorophyll and nutrients, at least in terms of their absolute values.

Figure 19 shows the relative changes in phytoplankton, microzooplankton and mesozooplankton biomasses (in carbon). A significant reduction in the carbon biomass of phytoplankton is predicted in the FOOD experiment. This reduction is maximum in the subtropical gyres where it may exceed 40 % because of more intense grazing by zooplankton. These changes are not perceptible in chlorophyll concentrations (at least with the color scale chosen on Fig. 17) because of the extremely low Chl/C in the gyres. Both on microzooplankton and mesozooplankton, the differences between the FOOD and the standard experiments are even more pronounced. Both zooplankton biomasses increase by more than 100 % in the subtropical gyres of all oceans and this increase even exceeds 200 % in the subtropical gyre of the South Pacific Ocean.

Food quality may thus have very important impacts on zooplankton, especially in the very oligotrophic regions. Furthermore, the importance of food quality is predicted to be more critical in regions depleted in nitrogen, characterized by very low N/C ratios in phytoplankton, than in iron limited areas. Several points may explain this greater sensitivity. First, even in the most severely iron limited areas, the Fe/C ratio in phytoplankton drops very rarely below half the value of the Fe/C ratio in zooplankton. In the central part of the subtropical gyres, where nitrogen limitation is the most intense, N/C ratios in phytoplankton can reach 0.04, that is about 3 times less than the N/C ratio of zooplankton. Second, the available food in the intense oligotrophic areas is much lower than in the iron limited regions. Chlorophyll concentrations in the typical HNLC regions are generally around 0.2 to 0.3 mg chl $a\ m^{-3}$ whereas it is below 0.1 mg chl $a\ m^{-3}$ in the subtropical gyres. As a consequence, zooplankton biomass is lower in the subtropical gyres which increases the magnitude of the relative changes.

8 Conclusions

5 In this paper, we have presented a full and thorough description of the current state of the ocean biogeochemical model PISCES, called PISCES-v2. Since the latest published version of the model (Aumont and Bopp, 2006), PISCES-v2 has undergone major changes both in terms of the modeled processes and of the model structure and performance. Relative to its previous version PISCES-v1, key changes are a major redesign of phytoplankton growth description, including
10 a quota-based representation of iron limitation, an improvement of the zooplankton compartment, a better description of the benthic processes and a simple description of nitrogen fixation by diazotrophs. A complete list of the changes made in PISCES-v2 relative to its previously published version is detailed in Sect. 2. The performance of the model has been then evaluated using a climatological simulation run to quasi-steady state. The model produces **reliable**
15 reasonable surface distributions of chlorophyll, mesozooplankton and nutrients (including iron) and simulates consistent vertical distributions of the main biogeochemical tracers. Some of the main deficiencies of the model are the spatial distribution of the Oxygen Minimum Zones, the silicic acid distribution in the Southern Ocean, too elevated nutrients concentrations in the deep Atlantic Ocean and an out-of-phase predicted seasonal cycle of chlorophyll in the subarctic
20 Pacific Ocean.

PISCES includes several optional parameterizations that may be activated from the namelist. In this study, we have presented the impacts of some of these optional formulations evaluated in a set of sensitivity experiments. The choice of the light limitation scheme has the largest effect on the model solution, especially on chlorophyll. The amplitude of the seasonal cycle in the high latitudes is profoundly impacted whereas the timing of the bloom maximum is in general
25 only very moderately altered. The effect of food quality on the growth efficiency of zooplankton has been shown to lead to important relative changes in the oligotrophic subtropical gyres. The model suggests that it is critical to maintain sufficiently high chlorophyll levels in these regions. It may also contribute to, at least partly, explain the too low primary productivity simulated by other biogeochemical models in the subtropical gyres (Yool et al., 2013).

The description of PISCES presented here has been restricted to the core scheme which can be obtained online from different SVN repositories depending on the dynamical framework in which it is embedded (see the Introduction for a list of these repositories). In addition to the description of the lower trophic levels of marine ecosystems, and the biogeochemical cycles of carbon and of the main nutrients (N, P, Si, Fe), as described in this manuscript, a few additional modules have been embedded into PISCES. These modules enable [the model](#) to compute the cycles of climate-relevant gases emitted by the ocean such as dimethylsulfide (DMS) ([Bopp et al., 2008](#)), and nitrous oxide (N₂O) ([Martinez-Rey et al., 2013](#)). An explicit representation of paleo-proxies, such as $\delta^{13}\text{C}$ ([Tagliabue et al., 2009b](#)), Pa/Th ([Dutay et al., 2009](#)), Nd ([Arsouze et al., 2009](#)), is also available.

PISCES is still in a phase of active developments despite that its development has started more than 10 years ago already. Avenues for future improvements are large and numerous and concern all aspects of the model. The challenges confronting marine biogeochemical modeling have been identified in many dedicated studies (e.g., [Doney, 1999](#); [Hood et al., 2006](#); [Merico et al., 2009](#); [Smith et al., 2011](#); [Mitra et al., 2014](#)). Setting priorities in a long list of potential necessary modifications is a rather difficult task which relies not only on the diagnostic of the major deficiencies of the current model but also on the future research scope envisioned for the model. In the coming years, PISCES will evolve along two main avenues. First, a more sophisticated treatment of phytoplankton physiology will replace the current relatively simple scheme. A main consequence is the representation of variable elemental ratios for all major elements (N, P, Fe, Si, C). Redfield-Monod models have been shown to exhibit serious deficiencies which advocate for their replacement by more detailed mechanistic schemes ([Flynn, 2010](#); [Smith et al., 2011](#)). Second, almost all marine biogeochemical models have been built on the classical distinction between phytoplanktonic autotrophic organisms and zooplanktonic heterotrophic organisms. However, this dichotomy has been increasingly challenged in the recent years as observations have shown that most protists, with the exception of diatoms probably, have to a lesser or greater degree a mixotrophic status (e.g., [Stoecker, 1998](#); [Flynn et al., 2013](#)). The conceptual schemes on which biogeochemical models, including PISCES, should then be revised, in particular the distinction between phytoplankton and microzooplankton.

Appendix A

5 Model structure

The model is being coded in FORTRAN 90. To activate PISCES, the cpp key `key_pisces` should be declared. Only the subroutines that compute the biological or chemical sources and sinks are considered to be part of PISCES. Thus, this excludes the computation of the advection-diffusion equation (the transport of the tracers), as it is not specific to PISCES. There are two types of subroutines: the initialization of the tracers and of the parameters and the computation of the various biogeochemical sources and sinks. The latter PISCES subroutines are called from within the ocean model timeloop.

The objective here is not to precisely detail the PISCES code but rather to list the different modules and to briefly describe their role. All the subroutines that compute the biogeochemical sources/sinks are called from `p4zsms` which is then the main PISCES subroutine.

- *p4zbio.F90*: Computation of the new tracer concentrations by summing up all the different sources and sinks.
- *p4zche.F90*: Computation of the various chemical constants.
- *p4zfechem.F90*: Computation of the iron chemistry. Scavenging of iron, aggregation of iron colloids.
- *p4zflx.F90*: Air-sea fluxes of CO₂ and O₂.
- *p4zint.F90*: Time interpolation of various terms (growth rate, ...).
- *p4zlim.F90*: Co-limitations of phytoplankton growth by the different nutrients.
- *p4zlys.F90*: Calcite chemistry and dissolution.
- *p4zmeso.F90*: Sources and sinks of mesozooplankton (mortality, grazing, ...).
- *p4zmicro.F90*: Sources and sinks of microzooplankton.

- *p4zmort.F90*: Computation of the various mortality terms of nanophytoplankton and diatoms.
- 5 – *p4zopt.F90*: Optical model and computation of the euphotic depth.
- *p4zprod.F90*: Growth rate of the two phytoplankton groups.
- *p4zrem.F90*: Remineralization of organic matter, dissolution of biogenic silica.
- *p4zsed.F90*: Top and bottom boundary conditions of the biogeochemical tracers (deposition, sedimentary losses, ...).
- 10 – *p4zsink.F90*: Aggregation of organic matter, computation of the particles sinking speeds. Vertical sedimentation of particles using a MUSCL advection scheme.
- *p4zsms.F90*: Main PISCES subroutine which calls the other subroutine.

Besides the subroutines listed above, several subroutines perform the model initialization. We will only discuss the initialization of the parameters necessary to PISCES. The tracers concentrations are excluded here as their initialization will of course vary with the ocean model.

- *trcini.pisces.F90*: Initialization of various biogeochemical parameters. Allocation of the arrays used in PISCES. This subroutine also calls all the initialization subroutines included in the PISCES subroutines listed above.
- *trcnam_pisces.F90*: This subroutine reads the informations necessary to write the netcdf files when IOM is not used.
- 20 – *par_pisces.F90*: It sets the PISCES parameters such as the number of tracers and the name of the indices, the number of additional diagnostics, ...
- *sms_pisces.F90*: This subroutine defines some general PISCES variables and arrays and allocates them.

Many parameter values of the model can be specified from the namelist `namelist_pisces`. When such is the case, the corresponding parameter name in the namelist file is indicated in Tables [A1a–A1d](#).

Acknowledgements. The authors are grateful to the whole community of PISCES users. The model would be useless without them. In particular, we thank Thomas Gorguès, Keith B. Rodgers, Coralie Peruche, Christophe Menkès, Vincent Echevin, Gildas Cambon and the whole NEMO system team for their precious help, expertise and support which made the release of this version possible. O. Aumont, L. Bopp, M. Gehlen were supported by ANR-CEP09 MACROES. Olivier Aumont will be eternally grateful to Ernst Maier-Reimer. Even if he did not directly participate to the design of PISCES, nothing would have been possible without him. PISCES was built upon HAMOCC3 and HAMOCC3.1 and still some parts of the current code of PISCES come from these models. This work is dedicated to his memory.

References

- Albert, A., Echevin, V., Lévy, M., and Aumont, O.: Impact of nearshore wind stress curl on coastal circulation and primary productivity in the Peru upwelling system, *J. Geophys. Res.*, 115, C12033, doi:[10.1029/2010JC006569](https://doi.org/10.1029/2010JC006569), 2010.
- Allen, J. I., Somerfield, P. J., and Gilbert, F. J.: Quantifying uncertainty in high-resolution coupled hydrodynamic-ecosystem models, *J. Marine Syst.*, 64, 3–14, 2007.
- Alvain, S., Moulin, C., Dandonneau, Y., and Bréon, F.-M.: Remote sensing of phytoplankton groups in case 1 waters from global SeaWiFS imagery, *Deep-Sea Res. Pt. I*, 52, 1989–2004, 2005.
- Anderson, T. R.: Plankton functional type modelling: running before we can walk?, *J. Plankton Res.*, 27, 1073–1081, 2005.
- Anderson, T. R.: Progress in marine ecosystem modelling and the “unreasonable effectiveness of mathematics”, *J. Marine Syst.*, 81, 4–11, 2010.
- Anderson, T. R. and Williams, P. J. B.: A one dimensional model of dissolved organic carbon cycling in the water column incorporating combined biological-photochemical decomposition, *Global Biogeochem. Cy.*, 13, 337–349, 1999.
- Anderson, T. R., Hessen, D. O., Mitra, A., Mayor, D. J., and Yool, A.: Sensitivity of secondary production and export flux to choice of trophic transfer formulation in marine ecosystem models, *J. Marine Syst.*, 125, 41–53, doi:[10.1016/j.jmarsys.2012.09.008](https://doi.org/10.1016/j.jmarsys.2012.09.008), 2013.

- Antoine, D., André, J. M., and Morel, A.: Oceanic primary production 2. Estimation at global scale from satellite (coastal zone color scanner) chlorophyll, *Global Biogeochem. Cy.*, 10, 57–69, 1996.
- Archer, D. E.: An atlas of the distribution of calcium carbonate in sediments of the deep sea, *Global Biogeochem. Cy.*, 10, 159–174, 1996.
- Aristegui, J., Gasol, J. M., Duarte, C. M., and Herndl, G. J.: Microbial oceanography of the dark oceans's pelagic realm, *Limnol. Oceanogr.*, 54, 1501–1529, 2009.
- Armstrong, R. A.: Grazing limitation and nutrient limitation in marine ecosystems: steady-state solutions of an ecosystem model with multiple food chains, *Limnol. Oceanogr.*, 39, 597–608, 1994.
- Armstrong, R. A., Lee, C., Hedges, J. I., Honjo, S., and Wakeham, S. G.: A new, mechanistic model for organic carbon fluxes in the ocean based on the quantitative association of POC with ballast minerals, *Deep-Sea Res. Pt. II*, 49, 219–236, 2002.
- Arsouze, T., Dutay, J.-C., Kageyama, M., Lacan, F., Alkama, R., Marti, O., and Jeandel, C.: A modeling sensitivity study of the influence of the Atlantic meridional overturning circulation on neodymium isotopic composition at the Last Glacial Maximum, *Clim. Past*, 4, 191–203, doi:[10.5194/cp-4-191-2008](https://doi.org/10.5194/cp-4-191-2008), 2008.
- Arsouze, T., Dutay, J.-C., Lacan, F., and Jeandel, C.: Reconstructing the Nd oceanic cycle using a coupled dynamical – biogeochemical model, *Biogeosciences*, 6, 2829–2846, doi:[10.5194/bg-6-2829-2009](https://doi.org/10.5194/bg-6-2829-2009), 2009.
- Aumont, O.: Etude du cycle naturel du carbone dans un modèle 3D de l'océan mondial, PhD thesis, Univ. Paris VI, Paris, 1998.
- Aumont, O. and Bopp, L.: Globalizing results from ocean in-situ iron fertilization studies, *Global Biogeochem. Cy.*, 20, GB2017, doi:[10.1029/2005GB002591](https://doi.org/10.1029/2005GB002591), 2006.
- Aumont, O., Belviso, S., and Monfray, P.: Dimethylsulfoniopropionate (DMSP) and dimethylsulfide (DMS) sea surface distributions simulated from a global 3-D ocean carbon cycle model, *J. Geophys. Res.*, 107, 4.1–4.19, doi:[10.1029/1999JC000111](https://doi.org/10.1029/1999JC000111), 2002.
- Aumont, O., Maier-Reimer, E., Blain, S., and Monfray, P.: An ecosystem model of the global ocean including Fe, Si, P co-limitation, *Global Biogeochem. Cy.*, 17, 1060, doi:[10.1029/2001GB001745](https://doi.org/10.1029/2001GB001745), 2003.
- Ayata, S. D., Lévy, M., Aumont, O., Sciandra, A., Sainte-Marie, J., Tagliabue, A., and Bernard, O.: Phytoplankton growth formulation in marine ecosystem models: should we take into account photo-acclimation and variable stoichiometry in oligotrophic areas?, *J. Marine Syst.*, 125, 29–40, doi:[10.1016/j.jmarsys.2012.12.010](https://doi.org/10.1016/j.jmarsys.2012.12.010), 2013.

- Bacastow, R. and Maier-Reimer, E.: Ocean-circulation model of the carbon cycle, *Clim. Dynam.*, 4, 95–125, 1990.
- 10 Baines, S. B., Twining, B. S., Brzezinski, M. A., Nelson, D. M., and Fisher, N. S.: Causes and biogeochemical implications of regional differences in silicification of marine diatoms, *Global Biogeochem. Cy.*, 24, GB4031, doi:[10.1029/2010GB003856](https://doi.org/10.1029/2010GB003856), 2010.
- Balch, W. M., Drapeau, D. T., Bowler, B. C., and Booth, E.: Prediction of pelagic calcification rates using satellite-measurements, *Deep-Sea Res. Pt. II*, 54, 478–495, 2007.
- 15 Barnier, B., Madec, G., Penduff, T., Molines, J.-M., Tréguier, A.-M., le Sommer, J., Beckmann, A., Biastoch, A., Böning, C., Dengg, J., Derval, C., Durand, E., Gulev, S., Remy, E., Talandier, C., Theetten, S., Maltrud, M., McClean, J., and Cuevas, B. D.: Impact of partial steps and momentum advection schemes in a global ocean circulation model at eddy-permitting resolution, *Ocean Dynam.*, 56, 543–567, doi:[10.1007/s10236-006-0082-1](https://doi.org/10.1007/s10236-006-0082-1), 2006.
- Behrenfeld, M. J. and Falkowski, P. G.: Photosynthetic rates derived from satellite-based chlorophyll concentration, *Limnol. Oceanogr.*, 42, 1–20, 1997.
- 20 Behrenfeld, M. J., Boss, E., Siegel, D. A., and Shea, D. M.: Carbon-based ocean productivity and phytoplankton physiology from space, *Global Biogeochem. Cy.*, 19, GB1006, doi:[10.1029/2004GB002299](https://doi.org/10.1029/2004GB002299), 2005.
- Bennett, S. A., Achterberg, E. P., Connelly, D. P., Statham, P. J., Fones, G. R., and German, C. R.: The distribution and stabilisation of dissolved Fe in deep-sea hydrothermal plumes, *Earth Planet. Sc. Lett.*, 270, 157–167, 2008.
- 25 Berelson, W. M.: Particle settling rates increase with depth in the ocean, *Deep-Sea Res. Pt. II*, 49, 237–251, 2002.
- Berelson, W. M., Balch, W. M., Najjar, R., Feely, R. A., Sabine, C., and Lee, K.: Relating estimates of CaCO_3 production, export, and dissolution in the water column to measurements of CaCO_3 rain into sediment traps and dissolution on the sea floor: a revised global carbonate budget, *Global Biogeochem. Cy.*, 21, GB1024, doi:[10.1029/2006GB002803](https://doi.org/10.1029/2006GB002803), 2007.
- 30 Blain, S., Quéguiner, B., Armand, L., Belviso, S., Bombled, B., Bopp, L., Bowie, A., Brunet, C., Brussaard, C., Carlotti, F., Christaki, U., Corbière, A., Durand, I., Ebersbach, F., Fuda, J.-L., Garcia, N., Gerringa, L., Griffiths, B., Guigue, C., Guillerm, C., Jacquet, S., Jeandel, C., Laan, P., Lefèvre, D., Monaco, C. L., Malits, A., Mosseri, J., Obernosterer, I., Park, Y.-H., Picheral, M., Pondaven, P., Remenyi, T., Sandroni, V., Sarthou, G., Savoye, N., Scouarnec, L., Souhaut, M., Thuiller, D., Timmermans, K., Trull, T., Uitz, J., van Beek, P., Veldhuis, M., Vincent, D., Viollier, E., Vong, L., and Wa-

- gener, T.: Effect of natural iron fertilization on carbon sequestration in the Southern Ocean, *Nature*, 446, 1070–1074, 2007.
- 10 Bonnet, S. and Guieu, C.: Dissolution of atmospheric iron in seawater, *Geophys. Res. Lett.*, 31, L03303, doi:[10.1029/2003GL018423](https://doi.org/10.1029/2003GL018423), 2004.
- Bopp, L., Kohfeld, K. E., Quéré, C. L., and Aumont, O.: Dust impact on marine biota and atmospheric $p\text{CO}_2$ during glacial periods, *Paleoceanography*, 18, 1046, doi:[10.1029/2002PA000810](https://doi.org/10.1029/2002PA000810), 2003.
- 15 Bopp, L., Aumont, O., Cadule, P., Alvain, S., and Gehlen, M.: Response of diatoms distribution to global warming and potential implications: a global model study, *Geophys. Res. Lett.*, 32, L19606, doi:[10.1029/2005GL023653](https://doi.org/10.1029/2005GL023653), 2005.
- Bopp, L., Aumont, O., Belviso, S., and Blain, S.: Modelling the effect of iron fertilization on dimethylsulphide emissions in the Southern Ocean, *Deep-Sea Res. Pt. II*, 55, 901–912, doi:[10.1016/j.dsr2.2007.12.002](https://doi.org/10.1016/j.dsr2.2007.12.002), 2008.
- 20 Bopp, L., Resplandy, L., Orr, J. C., Doney, S. C., Dunne, J. P., Gehlen, M., Halloran, P., Heinze, C., Ilyina, T., Séférian, R., Tjiputra, J., and Vichi, M.: Multiple stressors of ocean ecosystems in the 21st century: projections with CMIP5 models, *Biogeosciences*, 10, 6225–6245, doi:[10.5194/bg-10-6225-2013](https://doi.org/10.5194/bg-10-6225-2013), 2013.
- Boyd, P. W. and Ellwood, M. J.: The biogeochemical cycle of iron in the ocean, *Nat. Geosci.*, 3, 675–682, 2010.
- 25 Boyé, M., van den Berg, C. M. G., de Jong, J. T. M., Leach, H., Croot, P., and de Baar, H. J. W.: Organic complexation of iron in the Southern Ocean, *Deep-Sea Res. Pt. I*, 48, 1477–1497, 2001.
- Boyé, M., Aldrich, A. P., and J. T. M. M. de Jong, C. M. G. v., Veldhuis, M., and de Baar, H. J. W.: Horizontal gradient of the chemical speciation of iron in surface waters of the northeast Atlantic Ocean, *Mar. Chem.*, 80, 129–143, 2003.
- 30 Boyle, E. A. and Jenkins, W. J.: Hydrothermal iron in the deep western South Pacific, *Geochim. Cosmochim. Ac.*, 72, A107, 2008.
- Boyle, E. A., Bergquist, B. A., Kayser, R. A., and Mahowald, N.: Iron, manganese, and lead at Hwawaii Ocean Time-series station ALOHA: temporal variability and an intermediate water hydrothermal plume, *Geochim. Cosmochim. Ac.*, 69, 933–952, 2005.
- Brasseur, P., Gruber, N., Barciela, R., Brander, K., Doron, M., El Moussaoui, A., Hobday, A. J., Huret, M., Kremer, A.-S., Lehodey, P., Matear, R., Moulin, C., Murtugudde, R., Senina, I., and Svendsen, E.: Integrating biogeochemistry and ecology into ocean data assimilation systems, *Oceanography*, 22, 206–215, 2009.
- 5

- Brewin, R. J. W., Sathyendranath, S., Hirata, T., Lavender, S., Baraciela, R. M., and Hardman-Mountford, N.: A three-component model of phytoplankton size class for the Atlantic ocean, *Ecol. Model.*, 221, 1472–1483, 2010.
- 10 Brewin, R. J. W., Hardman-Mountford, N., Lavender, S. J., Raitsoo, D., Hirata, T., Uitz, J., Devred, E., Bricaud, A., Ciotti, A., and Gentili, B.: An intercomparison of bio-optical techniques for detecting dominant phytoplankton size class from satellite remote sensing, *Remote Sens. Environ.*, 115, 325–339, 2011.
- 15 Bricaud, A., Babin, M., Morel, A., and Claustre, H.: Variability in the chlorophyll-specific absorption coefficients of natural phytoplankton: analysis and parameterization, *J. Geophys. Res.*, 100, 13321–13332, 1995.
- Bruland, K. W., Rue, E. L., Smith, G. J., and DiTullio, G. R.: Iron, macronutrients and diatom blooms in the Peru upwelling regime: brown and blue waters of Peru, *Mar. Chem.*, 93, 81–103, 2005.
- Brzezinski, M. A.: The Si : C : N ratio of marine diatoms: interspecific variability and the effect of some environmental variables, *J. Phycol.*, 21, 347–357, 1985.
- 20 Buitenhuis, E. T. and Geider, R. J.: A model of phytoplankton acclimation to iron-light colimitation, *Limnol. Oceanogr.*, 55, 714–724, 2010.
- Buitenhuis, E., Le Quéré, C., Aumont, O., Bunker, A., Hirst, A., Ikeda, T., O'Brien, T., Pontkiovski, S., and Straile, D.: Biogeochemical fluxes through mesozooplankton, *Global Biogeochem. Cy.*, 20, GB2003, doi:[10.1029/2009GB003601](https://doi.org/10.1029/2009GB003601), 2005.
- 25 Buitenhuis, E. T., Rivkin, R. B., Saille, S., and Le Quéré, C.: Biogeochemical fluxes through microzooplankton, *Global Biogeochem. Cy.*, 24, GB4015, doi:[10.1029/2005GB002511](https://doi.org/10.1029/2005GB002511), 2010.
- Calbet, A.: Mesozooplankton grazing effect on primary production: a global comparative analysis in marine ecosystems, *Limnol. Oceanogr.*, 46, 1824–1830, 2001.
- Capone, D. G., Zehr, J. P., Paerl, H. W., Berman, B., and Carpenter, E. J.: *Trichodesmium*, a globally significant marine cyanobacterium, *Science*, 276, 1221–1229, 1997.
- 30 Carlson, C. A., Ducklow, H. W., and Michaels, A. F.: Annual flux of dissolved organic carbon from the euphotic zone in the northwestern Sargasso Sea, *Nature*, 371, 405–408, 1994.
- Chen, M. and Wang, W.-X.: Bioavailability of natural colloidal-bound iron to marine phytoplankton: influences of colloidal size and aging, *Limnol. Oceanogr.*, 46, 1956–1967, 2001.
- Chen, M., Dei, R. C. H., Wang, W.-X., and Guo, L.: Marine diatom uptake of iron bound with natural colloids of different origins, *Mar. Chem.*, 81, 177–189, 2003.
- 5 Chester, R.: *Marine Geochemistry*, Unwin Hyman, London, 698 pp., 1990.

- Claquin, P., Martin-Jézéquel, V., Kromkamp, J. C., Veldhuis, M., and Kraay, G.: Uncoupling of silicon compared to carbon and nitrogen metabolism, and the role of the cell cycle, in continuous cultures of *Thalassiosira Pseudonana* (Bacillariophyceae), under light, nitrogen, and phosphorus control, *J. Phycol.*, 38, 922–930, 2002.
- 10 Codispoti, L. A., Brandes, J. A., Christensen, J. P., Devol, A. H., Naqvi, S. W. A., Pearl, H. W., and Yoshinari, T.: The oceanic fixed nitrogen and nitrous oxide budgets: moving targets as we enter the anthropocene?, *Sci. Mar.*, 65, 85–105, 2001.
- Collos, Y., Vaquer, A., and Souchu, P.: Acclimation of nitrate uptake by phytoplankton to high substrate levels, *J. Phycol.*, 41, 466–478, 1980.
- 15 Conkright, M. E., Locarnini, R. A., Garcia, H. E., O'Brien, T. D., Boyer, T. P., Stephens, C., and Antonov, J.: World Ocean Atlas 2001: Objective Analyses, Data Statistics and Figures, CD-ROM Documentation, Tech. rep., National Oceanographic Data Centre, Silver Spring, MD, USA, 2002.
- Darchambeau, F. and Thys, I.: In situ filtration responses of *Daphnia galeata* to changes in food quality, *J. Plankton Res.*, 27, 227–236, 2005.
- 20 de Baar, H. J. W. and de Jong, J. T. M.: Distributions, sources and sinks of iron in seawater, in: The Biogeochemistry of Iron in Seawater, edited by: Turner, D. and Hunter, K., John Wiley, Hoboken, NJ, 85–121, 2001.
- de Boyer-Montégut, C., Madec, G., Fischer, A. S., Lazar, A., and Iudicone, D.: Mixed layer depth over the global ocean: an examination of profile data and a profile-based climatology, *J. Geophys. Res.*, 25 109, C12003, doi:[10.1029/2004JC002378](https://doi.org/10.1029/2004JC002378), 2004.
- Debreu, L., Marchesiello, P., Penven, P., and Cambon, G.: Two-way nesting in split-explicit ocean models: algorithms, implementation and validation, *Ocean Model.*, 49–50, 1–21, 2011.
- Decho, A. W.: Microbial exopolymer secretions in ocean environments: their role(s) in food web and marine processes, *Oceanogr. Mar. Biol.*, 28, 73–153, 1990.
- 30 Deutsch, C., Sarmiento, J. L., Sigman, D. M., Gruber, N., and Dunne, J. P.: Spatial coupling of nitrogen inputs and losses in the ocean, *Nature*, 445, 163–167, doi:[10.1038/nature05392](https://doi.org/10.1038/nature05392), 2007.
- Dierssen, H. M. and Smith, R. C.: Bio-optical properties and remote sensing ocean color algorithms for Antarctic Peninsula waters, *J. Geophys. Res.*, 105, 26301–26312, doi:[10.1029/1999JC000296](https://doi.org/10.1029/1999JC000296), 2000.
- Dilling, L. and Alldredge, A. L.: Fragmentation of marine snow by swimming macrozooplankton: a new process impacting carbon cycling in the sea, *Deep-Sea Res. Pt. I*, 47, 1227–1245, 2000.
- Donald, K. M., Scanlan, D. J., Carr, N. G., Mann, N. H., and Joint, I.: Comparative phosphorus nutrition of the marine cyanobacterium *Synechococcus* WH7803 and the marine diatom *Thalassiosira* 5 *Weissflogii*, *J. Plankton Res.*, 19, 1793–1814, 1997.

- Doney, S. C.: Major challenges confronting marine biogeochemical modeling, *Global Biogeochem. Cy.*, 13, 705–714, 1999.
- Doney, S. C., Lima, I., Moore, J. K., Lindsay, K., Behrenfeld, M. J., Westberry, T. K., Mahowald, N., Glover, D. M., and Takahashi, T.: Skill metrics for confronting global upper ocean ecosystem-biogeochemistry models against field and remote sensing data, *J. Marine Syst.*, 76, 95–112, 2009.
- 10 Doucette, G. J. and Harrison, P. J.: Aspects of iron and nitrogen nutrition in the red tide dinoflagellate *Gymnodinium sanguineum*, *Mar. Biol.*, 110, 175–182, 1991.
- Droop, M. R.: 25 years of algal growth kinetics, *Bot. Mar.*, 26, 99–112, 1983.
- Dufresne, J. L., Friedlingstein, P., Berthelot, M., Bopp, L., Ciais, P., Fairhead, L., LeTreut, H., and Monfray, P.: Effects of climate change due to CO₂ increase on land and ocean carbon uptake, *Geophys. Res. Lett.*, 29, 1405, doi:[10.1029/2001GL013777](https://doi.org/10.1029/2001GL013777), 2002.
- 15 Dunne, J. P., Sarmiento, J. L., and Gnanadesikan, A.: A synthesis of global particle export from the surface ocean and cycling through the ocean interior and on the seafloor, *Global Biogeochem. Cy.*, 21, GB4006, doi:[10.1029/2006GB002907](https://doi.org/10.1029/2006GB002907), 2007.
- 20 Dutay, J.-C., Jean-Baptiste, P., Campin, J.-M., Ishida, A., Maier-Reimer, E., Matear, R. J., Mouchet, A., Totterdell, I. J., Yamanaka, Y., Rodgers, K., Madec, G., and Orr, J. C.: Evaluation of OCMIP-2 ocean models' deep circulation with mantle helium-3, *J. Marine Syst.*, 48, 15–36, 2004.
- Dutay, J.-C., Lacan, F., Roy-Barman, M., and Bopp, L.: Influence of particle size and type on 231Pa and 230Th simulation with a global coupled biogeochemical-ocean general circulation model: a first approach, *Geochem. Geophys. Geosy.*, 10, Q01011, doi:[10.1029/2008GC002291](https://doi.org/10.1029/2008GC002291), 2009.
- 25 Dutkiewicz, S., Follows, M. J., and Parekh, P.: Interactions of the iron and phosphorus cycles: a three-dimensional model study, *Global Biogeochem. Cy.*, 19, GB1021, doi:[10.1029/2004GB002342](https://doi.org/10.1029/2004GB002342), 2005.
- Dutz, J., Koski, M., and Jonasdóttir, S.: Copepod reproduction is unaffected by diatom aldehydes or lipid composition, *Limnol. Oceanogr.*, 53, 225–235, doi:[10.4319/lo.2008.53.1.0225](https://doi.org/10.4319/lo.2008.53.1.0225), 2008.
- 30 Echevin, V., Aumont, O., Ledesma, J., and Flores, G.: The seasonal cycle of surface chlorophyll in the Peruvian upwelling system: a modelling study, *Prog. Oceanogr.*, 79, 167–176, 2008.
- [Edwards, K. F., Thomas, M. K., Klausmeier, C. A., and Litchman, E.: Allometric scaling and taxonomic variation in nutrient utilization traits and maximum growth rate of phytoplankton, *Limnol. Oceanogr.*, 57, 554–566, 2012.](https://doi.org/10.1016/j.limnol.2012.05.001)
- Elrod, V. A., Berelson, W. M., Coale, K. H., and Johnson, K. S.: The flux of iron from continental shelf sediments: a missing source for global budgets, *Geophys. Res. Lett.*, 31, L12307, doi:[10.1029/2004GL020216](https://doi.org/10.1029/2004GL020216), 2004.

- 5 Engel, A., Szlosek, J., Abramson, L., Liu, Z., and Lee, C.: Investigating the effect of ballasting by CaCO_3 in *Emiliania huxleyi*: I. Formation, settling velocities and physical properties of aggregates, *Deep-Sea Res. Pt. II*, 56, 1396–1407, doi:[10.1016/j.dsr2.2008.11.027](https://doi.org/10.1016/j.dsr2.2008.11.027), 2009.
- Eppley, R. W.: Temperature and phytoplankton growth in the sea, *Fish. Bull.*, 70, 1063–1085, 1972.
- Eppley, R. W. and Peterson, B. J.: Particulate organic matter flux and planktonic new production in the
10 deep ocean, *Nature*, 282, 677–680, 1979.
- Eppley, R. W., Rogers, J. N., and McCarthy, J. J.: Half-saturation constants for uptake of nitrate and ammonium by marine phytoplankton, *Limnol. Oceanogr.*, 14, 912–920, 1969.
- Farley, K. A., Maier-Reimer, E., Schlosser, P., and Broecker, W. S.: Constraints on mantle ^3He fluxes and deep-sea circulation from an oceanic general circulation model, *J. Geophys. Res.-Sol. Ea.*, 100,
15 3829–3839, 1995.
- Fasham, M. J. R., Ducklow, H. W., and McKelvie, S. M.: A nitrogen-based model of plankton dynamics in the oceanic mixed layer, *J. Mar. Res.*, 48, 591–639, 1990.
- Fernández I, C., Raimbault, P., Garcia, N., Rimmelin, P., and Caniaux, G.: An estimation of annual new production and carbon fluxes in the northeast Atlantic Ocean during 2001, *J. Geophys. Res.-Oceans*,
20 110, C07S13, doi:[10.1029/2004JC002616](https://doi.org/10.1029/2004JC002616), 2005.
- Flynn, K. J.: Ecological modelling in a sea of variable stoichiometry: dysfunctionality and the legacy of Redfield and Monod, *Prog. Oceanogr.*, 84, 52–65, doi:[10.1016/j.pocan.2009.09.006](https://doi.org/10.1016/j.pocan.2009.09.006), 2010.
- Flynn, K. J. and Davidson, K.: Predator–prey interactions between *Isochrysis galbana* and *Oxyrrhis marina*. I I. Release of non-protein amines and faeces during predation of *Isochrysis*, *J. Plankton Res.*,
25 15, 893–905, 1993.
- Flynn, K. J. and Hipkin, C. R.: Interactions between iron, light, ammonium and nitrate: insights from the construction of a dynamic model of algal physiology, *J. Phycol.*, 35, 1171–1190, 1999.
- Flynn, K. J., Stoecker, D. K., Mitra, A., Raven, J. A., Glibert, P. M., Hansen, P. J., Granéli, E., and Burkholder, J. M.: Misuse of the phytoplankton–zooplankton dichotomy: the need to assign organisms as mixotrophs within plankton functional types, *J. Plankton Res.*, 35, 3–11,
30 doi:[10.1093/plankt/fbs062](https://doi.org/10.1093/plankt/fbs062), 2013.
- Franck, V. M., Brzezinski, M. A., Coale, K. H., and Nelson, D. M.: Iron and silicic acid concentrations regulate Si uptake north and south of the Polar Frontal Zone in the Pacific Sector of the Southern Ocean, *Deep-Sea Res. Pt. II*, 47, 3315–3338, 2000.
- Galloway, J. N., Dentener, F. J., Capone, D. G., Boyer, E. W., Howarth, R. W., Setzinger, S. P., Asner, G. P., Cleveland, C. C., Green, P. A., Holland, E. A., Karl, D. M., Michaels, A. F., and Porter, J. H.: Nitrogen cycles: past, present, future, *Biogeochemistry*, 70, 153–226, 2004.

- 5 Garcia, C. A. E., Garcia, V. M. T., and McClain, C. R.: Evaluation of SeaWiFS chlorophyll algorithms in the Southwestern Atlantic and Southern Oceans, *Remote Sens. Environ.*, 95, 125–137, 2005.
- Garcia, H. E., Locarnini, R. A., Boyer, T. P., Antonov, J. I., Baranova, O. K., Zweng, M. M., and Johnson, D. R.: *World Ocean Atlas 2009, Volume 4: Nutrients (Phosphate, Nitrate, Silicate)*, U.S. Government Printing Office, Washington, D.C., noaa atlas nesdis 71st Edn., 398 pp., 2010.
- 10 Gaspar, P., Gregoris, Y., and Lefevre, J. M.: A simple eddy kinetic energy model for simulations of the ocean vertical mixing: tests at station Papa and Long-Term Upper Ocean Study Site, *J. Geophys. Res.*, 95, 16179–16193, 1990.
- Gehlen, M., Bopp, L., Emprin, N., Aumont, O., Heinze, C., and Ragueneau, O.: Reconciling surface ocean productivity, export fluxes and sediment composition in a global biogeochemical ocean model, *Biogeosciences*, 3, 521–537, doi:[10.5194/bg-3-521-2006](https://doi.org/10.5194/bg-3-521-2006), 2006.
- 15 Gehlen, M., Gangstø, R., Schneider, B., Bopp, L., Aumont, O., and Ethe, C.: The fate of pelagic CaCO₃ production in a high CO₂ ocean: a model study, *Biogeosciences*, 4, 505–519, doi:[10.5194/bg-4-505-2007](https://doi.org/10.5194/bg-4-505-2007), 2007.
- Geider, R. J., MacIntyre, H. L., and Kana, T. M.: A dynamic model of photoadaptation in phytoplankton, *Limnol. Oceanogr.*, 41, 1–15, 1996.
- 20 Geider, R. J., MacIntyre, H. L., and Kana, T. M.: A dynamic model of phytoplankton growth and acclimation: responses of the balanced growth and Chlorophyll *a* : carbon ratio to light, nutrient-limitation and temperature, *Mar. Ecol.-Prog. Ser.*, 148, 187–200, 1997.
- Gent, P. R. and McWilliams, J. C.: Isopycnal mixing in ocean circulation models, *J. Phys. Oceanogr.*, 20, 150–155, 1990.
- 25 Gentleman, W., Leising, A., Frost, B., Strom, S., and Murray, J.: Functional responses for zooplankton feeding on multiple resources: a review of assumptions and biological dynamics, *Deep-Sea Res. Pt. II*, 50, 2847–2875, 2003.
- Gilstad, M. and Sakshaug, E.: Growth rates of ten diatoms species from the Barents Seas at different irradiance and day lengths, *Mar. Ecol.-Prog. Ser.*, 64, 169–173, 1990.
- 30 Gledhill, M. and Kirsten, K. N.: The organic complexation of iron in the marine environment: a review, *Frontiers in Microbiology*, 3, doi:[10.3389/fmicb.2012.00069](https://doi.org/10.3389/fmicb.2012.00069), 2012.
- Gnanadesikan, A., Slater, R. J., Gruber, N., and Sarmiento, J. L.: Oceanic vertical exchange and new production: a comparison between models and observations, *Deep-Sea Res. Pt. II*, 49, 363–401, 2002.
- Goldman, J. C. and Dennett, M. R.: Ammonium regeneration and carbon utilization by marine bacteria grown on mixed substrates, *Mar. Biol.*, 109, 369–378, 1991.

- 5 Goose, H.: Modelling the large-scale behaviour of the coupled ocean-sea-ice system, PhD thesis, Université catholique de Louvain, Louvain-La-Neuve, Belgium, 231 pp., 1997.
- Gorgues, T., Menkes, C., Aumont, O., Vialard, J., Dandonneau, Y., and Bopp, L.: Biogeochemical impact of tropical instability waves in the Equatorial Pacific, *Geophys. Res. Lett.*, 32, L24615, doi:[10.1029/2005GL024110](https://doi.org/10.1029/2005GL024110), 2005.
- 10 Griffies, S. M., Biastoch, A., Böning, C., Bryan, F., Danabasoglu, G., Chassignet, E. P., England, M. H., Gerdes, R., Haak, H., Hallberg, R. W., Hazeleger, W., Jungclaus, J., Large, W. G., Madec, G., Pirani, A., Samuels, B. L., Scheinert, M., Gupta, A. S., Severijns, C. A., Tréguier, H. L. S. A.-M., Winton, M., Yeager, S., and Yin, J.: Coordinated Ocean-ice Reference Experiments (COREs), *Ocean Model.*, 26, 1–46, 2009.
- 15 Gruber, N.: The dynamics of the marine nitrogen cycle and atmospheric CO₂, in: *Carbon Climate Interactions*, edited by: Oguz, T. and Follows, M., Kluwer, Dordrecht, 97–148, 2004.
- Hansard, S. P., Landing, W. M., Measures, C. I., and Voelker, B. M.: Dissolved iron(II) in the Pacific Ocean: measurements from the PO₂ and P16N CLIVAR/CO₂ repeat hydrography expeditions, *Deep-Sea Res. Pt. I*, 56, 1117–1129, 2009.
- 20 Harrison, P. J. and Morel, F. M. M.: Response of the marine diatom *Thalassiosira weissflogii* to iron stress, *Limnol. Oceanogr.*, 31, 989–997, 1986.
- Harvey, H. R., Tuttle, J. H., and Bell, J. T.: Kinetics of phytoplankton decay during simulated sedimentation: changes in biochemical composition and microbial activity under oxic and anoxic conditions, *Geochim. Cosmochim. Ac.*, 59, 3367–3377, 1995.
- 25 Haygood, M. G., Holt, P. D., and Butler, A.: Aerobactin production by a planktonic marine *Vibrio* sp., *Limnol. Oceanogr.*, 38, 1091–1097, 1993.
- Heinze, C., Maier-Reimer, E., Winguth, A. M. E., and Archer, D.: A global oceanic sediment model for long-term climate studies, *Global Biogeochem. Cy.*, 13, 221–250, 1999.
- Hernández-León, S. and Ikeda, T.: A global assessment of mesozooplankton respiration in the ocean, *J. Plankton Res.*, 27, 153–158, 2005.
- 30 Hirata, T., Aiken, J., Hardman-Mountford, N., Smyth, T. J., and Barlow, R.: An absorption model to determine phytoplankton size classes from satellite ocean colour, *Remote Sens. Environ.*, 112, 3153–3159, 2008.
- Hirata, T., Hardman-Mountford, N. J., Brewin, R. J. W., Aiken, J., Barlow, R., Suzuki, K., Isada, T., Howell, E., Hashioka, T., Noguchi-Aita, M., and Yamanaka, Y.: Synoptic relationships between surface Chlorophyll-*a* and diagnostic pigments specific to phytoplankton functional types, *Biogeosciences*, 8, 311–327, doi:[10.5194/bg-8-311-2011](https://doi.org/10.5194/bg-8-311-2011), 2011.
- 5

- Honeyman, B., Balistrieri, L., and Murray, J.: Oceanic trace metal scavenging and the importance of particulate concentration, *Deep-Sea Res. Pt. I*, 35, 227–246, 1988.
- Honjo, S.: Fluxes of particles to the interior of the open oceans, in: *Particle Flux to the Ocean*, edited by: Ittekkot, V., Schäfer, P., Honjo, S., and Depetris, P. J., Vol. 57 of SCOPE, Wiley, New York, 91–254, 1996.
- Hood, R. R., Kohler, K. E., McCreary, J. P., and Smith, S. L.: A four-dimensional validation of a coupled physical-biological model of the Arabian Sea, *Deep-Sea Res. Pt. II*, 50, 2917–2945, 2003.
- Hood, R. R., Laws, E. A., Armstrong, R. A., Bates, N. R., Brown, C. W., Carlson, C. A., Chai, F., Doney, S. C., Falkowski, P. G., Feely, R. A., Friedrichs, M. A. M., Landry, M. R., Keith Moore, J., Nelson, D. M., Richardson, T. L., Salihoglu, B., Schartau, M., Toole, D. A., and Wiggert, J. D.: Pelagic functional group modeling: progress, challenges and prospects, *Deep-Sea Res. Pt. II*, 53, 459–512, doi:[10.1016/j.dsr2.2006.01.025](https://doi.org/10.1016/j.dsr2.2006.01.025), 2006.
- Horrigan, S. G., Carlucci, A. F., and Williams, P. M.: Light inhibition of nitrification in sea-surface waters, *J. Mar. Res.*, 39, 557–565, 1981.
- Hunter, K. A. and Boyd, P. W.: Iron-binding ligands and their role in the ocean biogeochemistry of iron, *Environ. Chem.*, 4, 221–232, 2007.
- Hurtt, G. C. and Armstrong, R. A.: A pelagic ecosystem model calibrated with BATS data, *Deep-Sea Res.*, 43, 653–683, 1996.
- Hutchins, D. A. and Bruland, K. W.: Iron-limited diatom growth and Si:N uptake ratios in a coastal upwelling regime, *Nature*, 393, 561–564, 1998.
- Ibisanmi, E., Sander, S. G., Boyd, P. W., Bowie, A. R., and Hunter, K. A.: Vertical distributions of iron-(III) complexing ligands in the Southern Ocean, *Deep-Sea Res. Pt. II*, 58, 2113–2125, 2011.
- Jackson, G. A.: A model of the formation of marine algal flocs by physical coagulation processes, *Deep-Sea Res.*, 37, 1197–1211, 1990.
- Jansen, H. and Wolf-Gladrow, D. A.: Carbonate dissolution in copepod guts: a numerical model, *Mar. Ecol.-Prog. Ser.*, 221, 199–207, 2001.
- Jickells, T. D. and Spokes, L. J.: Atmospheric iron inputs to the oceans, in: *The Biogeochemistry of Iron in Seawater*, edited by: Turner, D. and Hunter, K., John Wiley, Hoboken, NJ, 85–121, 2001.
- Jickells, T. D., An, Z. S., Andersen, K. K., Baker, A. R., Bergametti, G., Brooks, N., Cao, J. J., Boyd, P. W., Duce, R. A., Hunter, K. A., Kawahata, H., Kubilay, N., laRoche, J., Liss, P. S., Mahowald, N., Prospero, J. M., Ridgwell, A. J., Tegen, I., and Torres, R.: Global iron connections between desert dust, ocean biogeochemistry, and climate, *Nature*, 308, 67–71, doi:[10.1126/science.1105959](https://doi.org/10.1126/science.1105959), 2005.

- Johnson, K. S., Gordon, R. M., and Coale, K. H.: What controls dissolved iron concentrations in the world ocean?, *Mar. Chem.*, 57, 137–161, 1997.
- Johnson, K. S., Chavez, F. P., and Friederich, G. E.: Continental-shelf sediment as a primary source of iron for coastal phytoplankton, *Nature*, 398, 697–700, 1999.
- 10 Kahru, M. and Mitchell, B. G.: Blending of ocean colour algorithms applied to the Southern Ocean, *Remote Sensing Letters*, 1, 119–124, 2010.
- Kalnay, E. C., Kanamitsu, M., Kistler, R., Collins, W., Deaven, D., Gandin, L., Iredell, M., Saha, S., White, G., Woollen, J., Zhu, Y., Chelliah, M., Ebisuzaki, W., Higgins, W., Janowiak, J., Mo, K. C., Ropelewski, C., Wang, J., Letmaa, A., Reynolds, R., Jenne, R., and Joseph, D.: The NCEP/NCAR Reanalysis project, *B. Am. Meteorol. Soc.*, 77, 437–471, 1996.
- 15 Kamatani, A., Riley, J. P., and Shirrow, G.: The dissolution of opaline silica of diatom tests in sea water, *Journal of the Oceanographical Society of Japan*, 36, 201–208, 1980.
- Kawamiya, M.: Mechanism of offshore nutrient supply in the Western Arabian Sea, *J. Mar. Res.*, 59, 675–696, 2001.
- 20 Key, R. M., Kozyr, A., Sabine, C. L., Lee, K., Wanninkhof, R., Bullister, J., Feely, R. A., Millero, F., Mordy, C., and Peng, T.-H.: A global ocean carbon climatology: results from GLODAP, *Global Biogeochem. Cy.*, 18, GB4031, doi:[10.1029/2004GB002247](https://doi.org/10.1029/2004GB002247), 2004.
- Klaas, C. and Archer, D. E.: Association of sinking organic matter with various types of mineral ballast in the deep sea: implications for the rain ratio, *Global Biogeochem. Cy.*, 16, 1116, doi:[10.1029/2001GB001765](https://doi.org/10.1029/2001GB001765), 2002.
- 25 Koné, V., Aumont, O., Lévy, M., and Resplandy, L.: Physical and biogeochemical controls of the phytoplankton seasonal cycle in the Indian Ocean: a modeling study, *Geophysical Monograph*, 185, 147–166, 2009.
- Korb, R. E., Whitehouse, M. J., and Ward, P.: SeaWiFS in the southern ocean: spatial and temporal variability in phytoplankton biomass around South Georgia, *Deep-Sea Res. Pt. II*, 51, 99–116, 2004.
- 30 Korb, R. E., Whitehouse, M. J., Atkinson, A., and Thorpe, S. E.: Magnitude and maintenance of the phytoplankton bloom at South Georgia: a naturally iron-replete environment, *Mar. Ecol.-Prog. Ser.*, 368, 75–91, 2008.
- Kortzinger, A., Hedges, J. I., and Quay, P. D.: Redfield ratios revisited: removing the biasing effect of anthropogenic CO₂, *Limnol. Oceanogr.*, 46, 964–970, 2001.
- Kriest, I.: Different parameterizations of marine snow in a 1-D model and their influence on representation of marine snow, nitrogen budget and sedimentation, *Deep-Sea Res. Pt. I*, 49, 2133–2162, 2002.
- 5

- Kriest, I. and Evans, G. T.: Representing phytoplankton aggregates in biogeochemical models, *Deep-Sea Res. Pt. I*, 46, 1841–1859, 1999.
- Kriest, I. and Evans, G.: A vertically resolved model for phytoplankton aggregation, *Proceedings of the Indian Academy of Science, Earth and Planetary Science*, 109, 453–469, 2000.
- 10 Lam, P. J., Bishop, J. K. B., Henning, C. C., Marcus, M. A., Waychunas, G. A., and Fung, I. Y.: Wintertime phytoplankton bloom in the subarctic Pacific supported by continental margin iron, *Global Biogeochem. Cy.*, 20, GB1006, doi:[10.1029/2005GB002557](https://doi.org/10.1029/2005GB002557), 2006.
- Lancelot, C., de Montety, A., Goosse, H., Becquevort, S., Schoemann, V., Pasquer, B., and Vancoppenolle, M.: Spatial distribution of the iron supply to phytoplankton in the Southern Ocean: a model study, *Biogeosciences*, 6, 2861–2878, doi:[10.5194/bg-6-2861-2009](https://doi.org/10.5194/bg-6-2861-2009), 2009.
- 15 Lannuzel, D., Schoemann, V., de Jong, J. T. M., Tison, J., and Chou, L.: Distribution and biogeochemical behaviour of iron in the East Antarctic sea ice, *Mar. Chem.*, 106, 18–32, 2007.
- Lannuzel, D., Schoemann, V., de Jong, J. T. M., Chou, L., Delille, B., Becquevort, S., and Tison, J.-L.: Iron study during a time series in the western Weddell pack ice, *Mar. Chem.*, 108, 85–95, 2008.
- 20 Lee, C. M., Jones, B. H., Brink, K. H., and Fischer, A. S.: The upper-ocean response to monsoonal forcing in the Arabian Sea: seasonal and spatial variability, *Deep-Sea Res. Pt. II*, 47, 1177–1226, 2000.
- Lee, C., Peterson, M. L., Wakeman, S. G., Armstrong, R. A., Cochran, J. K., Miquel, J. M., Fowler, S. W., Hirschberg, D., Beck, A., and Xue, J.: Particulate organic matter and ballast fluxes measured using time-series and settling velocity sediment traps in the northwestern Mediterranean Sea, *Deep-Sea Res. Pt. II*, 56, 1420–1436, 2009.
- 25 Lee, K.: Global net community production estimated from the annual cycle of surface water total dissolved inorganic carbon, *Limnol. Oceanogr.*, 46, 1287–1297, 2001.
- Lee, K., Tong, L. T., Millero, F. J., Sabine, C. L., Dickson, A. G., Goyet, C., Park, G.-H., Wanninkhof, R., Feely, R. A., and Key, R. M.: Global relationships of total alkalinity with salinity and temperature in surface waters of the world's oceans, *Geophys. Res. Lett.*, 33, L19605, doi:[10.1029/2006GL027207](https://doi.org/10.1029/2006GL027207), 2006.
- Lengaigne, M., Madec, G., Menkes, C., and Alory, G.: Effect of isopycnal diffusion in the tropical Pacific Ocean, *J. Geophys. Res.*, 108, 3345, doi:[10.1029/2002JC001704](https://doi.org/10.1029/2002JC001704), 2003.
- Lengaigne, M., Menkes, C., Aumont, O., Gorgues, T., Bopp, L., André, J.-M., and Madec, G.: Influence of the oceanic biology on the tropical Pacific climate in a coupled general circulation model, *Clim. Dynam.*, 28, 503–516, doi:[10.1007/s00382-006-0200-2](https://doi.org/10.1007/s00382-006-0200-2), 2007.
- 5

- Lévy, M., Klein, P., and Tréguier, A.-M.: Impact of sub-mesoscale physics on production and subduction of phytoplankton in an oligotrophic regime, *J. Mar. Res.*, 59, 535–565, 2001.
- Lévy, M., Lehahn, Y., André, J.-M., Mémerly, L., Loisel, H., and Heifetz, E.: Production regimes in the northeast Atlantic: a study based on Sea-viewing Wide Field-of-view Sensor (SeaWiFS) chlorophyll and ocean general circulation model mixed layer depth, *J. Geophys. Res.-Oceans*, 110, C07S10, doi:[10.1029/2004JC002771](https://doi.org/10.1029/2004JC002771), 2005.
- Lipschultz, F., Wofsy, S., Ward, B., Codispoti, L., Friederich, G., and Elkins, J.: Bacterial transformations of inorganic nitrogen in the oxygen-deficient waters of the eastern tropical South Pacific Ocean, *Deep-Sea Res. Pt. I*, 37, 1513–1541, 1990.
- [Litchman, E., Klausmeier, C. A., Schofield, O. M., and Falkowski, P. G.: The role of functional traits and trade-offs in structuring phytoplankton communities: scaling from cellular to ecosystem level, *Ecology Letters*, 10, 1170–1181, doi:\[10.1111/j.1461-0248.2007.01117.x\]\(https://doi.org/10.1111/j.1461-0248.2007.01117.x\), 2007.](#)
- Liu, X. and Millero, F. J.: The solubility of iron in seawater, *Mar. Chem.*, 77, 43–54, 2002.
- Longhurst, A., Sathyendranath, S., Platt, T., and Caverhill, C.: An estimate of global primary production in the ocean from satellite radiometer data, *J. Plankton Res.*, 17, 1245–1271, 1995.
- Loucaides, S., Van Cappellen, P., Roubeix, V., Moriceau, B., and Ragueneau, O.: Controls on the recycling and preservation of biogenic silica from biomineralization to burial, *Silicon*, 4, 7–22, 2012.
- Ludwig, W., Probst, J. L., and Kempe, S.: Predicting the oceanic input of organic carbon by continental erosion, *Global Biogeochem. Cy.*, 10, 23–41, 1996.
- Luo, C., Mahowald, N., Meskhidze, N., Chen, Y., Siefert, R., Baker, A., and Johansen, A.: Estimation of iron solubility from observations and a global aerosol model, *J. Geophys. Res.*, 110, D23307, doi:[10.1029/2005JD006059](https://doi.org/10.1029/2005JD006059), 2005.
- Mackey, D. J., O’Sullivan, J. E., and Watson, R. J.: Iron in the western Pacific: a riverine or hydrothermal source for iron in the equatorial undercurrent?, *Deep-Sea Res. Pt. I*, 49, 877–893, 2002.
- Madec, G.: “NEMO Ocean Engine”, Note du Pôle de Modélisation 27, Institut Pierre-Simon Laplace (IPSL), France, 2008.
- Madec, G., Delecluse, P., Imbard, M., and Lévy, C.: OPA8.1 Ocean General Circulation Model Reference Manual, Notes du pôle de modélisation, IPSL, 1998.
- Mahowald, N., Jickells, T. D., Baker, A. R., Artaxo, P., Benitez-Nelson, C. R., Bergametti, G., Bond, T. C., Chen, Y., Cohen, D. D., Herut, B., Kubilay, N., Losno, R., Luo, C., Maenhaut, W., McGee, K. A., Okin, G. S., Siefert, R. L., and Tsukuda, S.: Global distribution of atmospheric phosphorus sources, concentrations and deposition rates, and anthropogenic impacts, *Global Biogeochem. Cy.*, 22, GB4026, doi:[10.1029/2008GB003240](https://doi.org/10.1029/2008GB003240), 2008.

- Maier-Reimer, E., Mikolajewicz, U., and Hasselmann, K.: Mean circulation of the Hamburg LSG OGCM and its sensitivity to the thermohaline surface forcing, *J. Phys. Oceanogr.*, 23, 731–757, 1993.
- Martin, J. H., Knauer, G. A., Karl, D. M., and Broenkow, W. W.: VERTEX: carbon cycling in the north-east Pacific, *Deep-Sea Res.*, 34, 267–285, 1987.
- 10 Martin-Jézéquel, V., Hildebrand, M., and Brzezinski, M.: Silicon metabolism in diatoms: implications for growth, *J. Phycol.*, 36, 1–20, 2000.
- Martinez, J. S., Zhang, G. P., Holt, P. D., Jung, H.-T., Carrano, C. J., Haygood, M. G., and Butler, A.: Self-assembling amphiphilic siderophores from marine bacteria, *Science*, 287, 1245–1247, doi:[10.1126/science.287.5456.1245](https://doi.org/10.1126/science.287.5456.1245), 2000.
- 15 Martinez-Rey, J., Bopp, L., Gehlen, M., ~~and Tagliabue, A., and Gruber, N.~~: Oceanic N₂O emissions in the 21st century, ~~*Geophys. Res. Abstr.*, EGU2013-10424, EGU General Assembly 2013, Vienna, Austria, 2013.~~ [Biogeosciences Discuss.](https://doi.org/10.5194/bgd-11-16703-2014), 11, 16703–16742, doi:[10.5194/bgd-11-16703-2014](https://doi.org/10.5194/bgd-11-16703-2014), 2014.
- Masotti, I., Ruiz-Pino, D., and Le Bouteiller, A.: Photosynthetic characteristics of *Trichodesmium* in the southwest Pacific Ocean: importance and significance, *Mar. Ecol.-Prog. Ser.*, 338, 47–59, 2007.
- 20 Maury, O., Faugeras, B., Shin, Y.-J., Poggiale, J.-C., Ben Ari, T., and Marsac, F.: Modeling environmental effects on the size-structured energy flow through marine ecosystems. Part 1: The model, *Prog. Oceanogr.*, 74, 479–499, 2007.
- Mayorga, E., Seitzinger, S. P., Harrison, J. A., Dumont, E., Beusen, A. H. W., Bowman, A. F., Fekete, B. M., Kroeze, C., and Van Drecht, G.: Global Nutrient Export from WaterSheds 2 (NEWS 2): model development and implementation, *Journal of Environ. Modell. Softw.*, 25, 837–853, 2010.
- McCarthy, J. J.: The kinetics of nutrient utilization, *Can. B. Fish. Aquat. Sci.*, 210, 211–233, 1980.
- McGillicuddy, O. J., Robinson, A. R., Siegel, D. A., Jannasch, H. W., Johnson, R., Dickey, T. D., McNell, H., Michaels, A. F., and Knap, A. H.: Influence of mesoscale eddies on new production in the Sargasso Sea, *Nature*, 394, 263–266, 1998.
- 30 Menkes, C., Boulanger, J.-P., Busalacchi, A. J., J. Vialard, J., Delecluse, P., McPhaden, M. J., Hackert, E., and Grima, N.: Impact of TAO vs. ERS wind stresses onto simulations of the tropical Pacific Ocean during the 1993–1998 period by the OPA OGCM, in: Climatic Impact of Scale Interactions for the Tropical Ocean-Atmosphere System, EuroClivar Workshop Report, 46–48, 1998.
- Merico, A., Bruggeman, J., and Wirtz, K.: A trait-based approach for downscaling complexity in plankton ecosystem models, *Ecol. Model.*, 220, 3001–3010, doi:[10.1016/j.ecolmodel.2009.05.005](https://doi.org/10.1016/j.ecolmodel.2009.05.005), 2009.
- Middelburg, J. J., Soetaert, K., Herman, P. M. J., and Heip, C.: Denitrification in marine sediments: a model study, *Global Biogeochem. Cy.*, 10, 661–673, 1996.

- 5 Milliman, J. D., Troy, P. J., Balsch, W. M., Adams, A. K., Li, Y.-H., and Mackenzie, F. T.: Biologically mediated dissolution of calcium carbonate above the chemical lysocline?, *Deep-Sea Res. Pt. I*, 46, 1653–1669, 1999.
- Mills, M. M., Ridame, C., Davey, M., La Roche, J., and Geider, R. J.: Iron and phosphorus co-limit nitrogen fixation in the eastern tropical North Atlantic, *Nature*, 429, 292–294, 2004.
- 10 Misumi, K., Tsumune, D., Yoshida, Y., Uchimoto, K., Nakamura, T., Nishioka, J., Mitsudera, H., Bryan, F. O., Lindsay, K., Moore, J. K., and Doney, S. C.: Mechanisms controlling dissolved iron distribution in the North Pacific: a model study, *J. Geophys. Res.*, 116, G03005, doi:[10.1029/2010JG001541](https://doi.org/10.1029/2010JG001541), 2011.
- Mitra, A. and Flynn, K. J.: Predator–prey interactions: is “ecological stoichiometry” sufficient when good food goes bad?, *J. Plankton Res.*, 27, 393–399, 2005.
- 15 Mitra, A., Flynn, K. J., and Fasham, M. J. R.: Accounting for grazing dynamics in nitrogen-phytoplankton-zooplankton models, *Limnol. Oceanogr.*, 52, 649–661, 2007.
- Mitra, A., Castellani, C., Gentleman, W. C., Jónasdóttir, S. H., Flynn, K. J., Bode, A., Halsband, C., Kuhn, P., Licandro, P., Agersted, M. D., Cakbet, A., Lindeque, P. K., Koppelman, R., Møller, E. F., 20 Gislason, A., Nielsen, T. G., and St. John, M.: Bridging the gap between marine biogeochemical and fisheries sciences; configuring the zooplankton link, *Prog. Oceanogr.*, 129, 176–199, 2014.
- Monod, J.: *Recherches sur la Croissance des Cultures Bactériennes*, Hermann, Paris, 1942.
- Moore, J. K. and Braucher, O.: Sedimentary and mineral dust sources of dissolved iron to the world ocean, *Biogeosciences*, 5, 631–656, doi:[10.5194/bg-5-631-2008](https://doi.org/10.5194/bg-5-631-2008), 2008.
- 25 Moore, J. K., Doney, S. C., Glover, D. M., and Fung, I. Y.: Iron cycling and nutrient limitation patterns in surface waters of the world ocean, *Deep-Sea Res. Pt. II*, 49, 463–507, 2002a.
- Moore, J. K., Doney, S. C., Kleypas, J. A., Glover, D. M., and Fung, I. Y.: An intermediate complexity marine ecosystem model for the global domain, *Deep-Sea Res. Pt. II*, 49, 403–462, 2002b.
- Moore, J. K., Doney, S. C., and Lindsay, K.: Upper ocean ecosystem dynamics and iron cycling in a 30 global three-dimensional model, *Global Biogeochem. Cy.*, 18, GB4028, doi:[10.1029/2004GB002220](https://doi.org/10.1029/2004GB002220), 2004.
- Mooy, B. A. S. V., Keil, R. G., and Devol, A. H.: Impact of suboxia on sinking particulate organic carbon: enhanced carbon flux and preferential degradation of amino acids via denitrification, *Geochim. Cosmochim. Ac.*, 3, 457–465, 2002.
- Morel, A.: Optical modeling of the upper ocean in relation to its biogenous matter content, *J. Geophys. Res.*, 93, 709–722, 1988.

- 5 Morel, A. and Maritorena, S.: Bio-optical properties of oceanic waters: a reappraisal, *J. Geophys. Res.-Oceans*, 106, 7163–7180, doi:[10.1029/2000JC000319](https://doi.org/10.1029/2000JC000319), 2001.
- Morel, F. M. M.: Kinetics of nutrient uptake and growth in phytoplankton, *J. Phycol.*, 23, 137–150, 1987.
- Moriceau, B., Goutx, M., Guigue, C., Lee, C., Armstrong, R. A., Duflos, M., Tamburini, C., Charrière, B., and Ragueneau, O.: Si-C interactions during degradation of the diatom *Skeletonema marinoi*, *Deep-Sea Res. Pt. II*, 56, 1381–1395, doi:[10.1016/j.dsr2.2008.11.026](https://doi.org/10.1016/j.dsr2.2008.11.026), 2009.
- 10 Murnane, R. J., Sarmiento, J. L., and Le Quééré, C.: Spatial distribution of air-sea CO₂ fluxes and the interhemispheric transport of carbon by the oceans, *Global Biogeochem. Cy.*, 13, 287–305, 1999.
- Nelson, D. M., Tréguer, P., Brzezinski, M. A., Leynaert, A., and Quéguigner, B.: Production and dissolution of biogenic silica in the ocean: revised global estimates, comparison with regional data and relationship to biogenic sedimentation, *Global Biogeochem. Cy.*, 9, 359–372, 1995.
- 15 Nishioka, J. and Takeda, S.: Change in the concentrations of iron in different size-fractions during growth of the oceanic diatom *Chaetoceros* sp.: importance of small colloidal iron, *Mar. Biol.*, 137, 231–238, 2000.
- O’Neill, R. V., Angelis, D. L., Pastor, J. J., Jackson, B. J., and Post, W. M.: Multiple nutrient limitation in ecological models, *Ecol. Model.*, 46, 147–163, 1989.
- 20 Orr, J. C.: On ocean carbon-cycle model comparison, *Tellus B*, 51, 509–510, 1999.
- Orr, J. C., Fabry, V. J., Aumont, O., Bopp, L., Doney, S. C., Feely, R. A., Gnanadesikan, A., Gruber, N., Ishida, A., Joos, F., Key, R. M., Lindsay, K., Maier-Reimer, E., Matear, R., Monfray, P., Mouchet, A., Najjar, R. G., Plattner, G.-K., Rodgers, K. B., Sabine, C. L., Sarmiento, J. L., Schlitzer, R., Slater, R. D., Totterdell, I. J., Weirig, M.-F., Yamanaka, T., and Yool, A.: Anthropogenic ocean acidification over the twenty-first century and its impact on calcifying organisms, *Nature*, 437, 681–686, 2005.
- Oschlies, A. and Garçon, V.: Eddy-induced enhancement of primary production in a model of the North Atlantic Ocean, *Nature*, 394, 266–269, 1998.
- 30 Parekh, P., Follows, M. J., and Boyle, E. A.: Decoupling of iron and phosphate in the global ocean, *Global Biogeochem. Cy.*, 19, GB002280, doi:[10.1029/2004GB002280](https://doi.org/10.1029/2004GB002280), 2004.
- Paulmier, A., Kriest, I., and Oschlies, A.: Stoichiometries of remineralisation and denitrification in global biogeochemical ocean models, *Biogeosciences*, 6, 923–935, doi:[10.5194/bg-6-923-2009](https://doi.org/10.5194/bg-6-923-2009), 2009.
- Penduff, T., Le Sommer, J., Barnier, B., Treguier, A.-M., Molines, J.-M., and Madec, G.: Influence of numerical schemes on current-topography interactions in 1/4° global ocean simulations, *Ocean Sci.*, 3, 509–524, doi:[10.5194/os-3-509-2007](https://doi.org/10.5194/os-3-509-2007), 2007.

- Penven, P., Debreu, L., Marchesiello, P., and McWilliams, J. C.: Evaluation and application of the ROMS
1-way embedding procedure to the central California upwelling system, *Ocean Model.*, 12, 157–187,
2006.
- Perry, M. J.: Phosphate utilization by an oceanic diatom in phosphorus-limited chemostat culture and in
the oligotrophic waters of the central north Pacific, *Limnol. Oceanogr.*, 21, 88–107, 1976.
- Plath, K. and Boersma, M.: Mineral limitation of zooplankton: stoichiometric constraints and optimal
foraging, *Ecology*, 82, 1260–1269, 2001.
- Pollard, R., Sanders, R., Lucas, M., and Statham, P.: The Crozet Natural Iron Bloom and Export Experiment
(CROZEX), *Deep-Sea Res. Pt. II*, 54, 1905–1914, 2007.
- Pondaven, P., Fravallo, C., Ruiz-Pino, D., Tréguer, P., Quéguiner, B., and Jeandel, C.: Modelling the silica
pump in the permanently open ocean zone of the Southern Ocean, *J. Marine Syst.*, 17, 587–619, 1998.
- Pullin, M. J. and Cabaniss, S. E.: The effects of pH, ionic strength, and iron-fulvic acid interactions on
the kinetics of non-photochemical iron transformations II: the kinetics of thermal reduction, *Geochim.
Cosmochim. Ac.*, 67, 4079–4089, 2003.
- Quééré, C. L., Harrison, S. P., Prentice, I. C., Buitenhuis, E. T., Aumont, O., Bopp, L., Cotrim Da
Cunha, L., Geider, R., Giraud, X., Klaas, C., Kohfeld, K. E., Legendre, L., Manizza, M., Platt, T.,
Rivkin, R. B., Sathyendranath, S., Watson, J. U. A. J., and Wolf-Gladrow, D.: Ecosystem dynamics
based on plankton functional types for global ocean biogeochemistry models, *Glob. Change Biol.*, 11,
2016–2040, 2005.
- Raimbault, P., Rodier, M., and Taupier-Letage, I.: Size-fraction of phytoplankton in the Ligurian Sea and
the Algerian Basin (Mediterranean Sea): size distribution versus total concentration, *Marine Microbial
Food Web*, 3, 1–7, 1988.
- Raiswell, R. and Anderson, T. F.: Reactive iron enrichment in sediments deposited beneath euxinic bot-
tom waters: constraints on supply by shelf recycling, *Geol. Soc. Spec. Publ.*, 248, 179–194, 2005.
- Raynaud, S., Orr, J. C., Aumont, O., Rodgers, K. B., and Yiou, P.: Interannual-to-decadal variability of
North Atlantic air-sea CO₂ fluxes, *Ocean Sci.*, 2, 43–60, doi:[10.5194/os-2-43-2006](https://doi.org/10.5194/os-2-43-2006), 2006.
- Resplandy, L., Vialard, J., Lévy, M., Aumont, O., and Dandonneau, Y.: Seasonal and intraseasonal bio-
geochemical variability in the thermocline ridge of the southern tropical Indian Ocean, *J. Geophys.
Res.*, 114, C07024, doi:[10.1029/2008JC005246](https://doi.org/10.1029/2008JC005246), 2009.
- Resplandy, L., Lévy, M., Bopp, L., Echevin, V., Pous, S., Sarma, V. V. S. S., and Kumar, D.: Con-
trolling factors of the oxygen balance in the Arabian Sea's OMZ, *Biogeosciences*, 9, 5095–5109,
doi:[10.5194/bg-9-5095-2012](https://doi.org/10.5194/bg-9-5095-2012), 2012.

- Ridame, C. and Guieu, C.: Saharan input of phosphate to the oligotrophic water of the open western Mediterranean Sea, *Limnol. Oceanogr.*, 47, 856–869, 2002.
- 5 Ridgwell, A. J., Watson, A. J., and Archer, D. E.: Modelling the response of the oceanic Si inventory to perturbations and consequences for atmospheric CO₂, *Global Biogeochem. Cy.*, 16, 1071, doi:[10.1029/2002GB001877](https://doi.org/10.1029/2002GB001877), 2002.
- Rodgers, K. B., Aumont, O., Menkes, C., and Gorgues, T.: Decadal variations in equatorial Pacific ecosystems and ferrocline/pycnocline decoupling, *Global Biogeochem. Cy.*, 22, GB2019, doi:[10.1029/2006GB002919](https://doi.org/10.1029/2006GB002919), 2008.
- 10 Rossow, W. B. and Schiffer, R. A.: Advances in understanding clouds from ISCCP, *B. Am. Meteorol. Soc.*, 80, 2261–2288, 1999.
- Sabine, C. L., Feely, R. A., Gruber, N., Key, R. M., Lee, K., Bullister, J. L., Wanninkhof, R., Wong, C., Wallace, D. W., Tilbrook, B., Millero, F. J., Peng, T.-H., Kozyr, A., Ono, T., and Aida, F. R.: The oceanic sink for anthropogenic CO₂, *Science*, 305, 367–371, 2004.
- 15 Sarthou, G., Timmermans, K., Blain, S., and Treguer, P.: Growth physiology and fate of diatoms in the ocean: a review, *J. Sea Res.*, 53, 25–42, 2005.
- Schlitzer, R.: Applying the adjoint method for biogeochemical modeling: export of particulate organic matter in the World Ocean, in: *Inverse Methods in Biogeochemical Cycles*, edited by: Kasibhata, P., Vol. 114 of AGU Monograph, American Geophysical Union, 107–124, 2000.
- [Sedwick, P. N. and Di Tullio, G. R.: Regulation of algal blooms in Antarctic shelf waters by the release of iron from melting sea ice, *Geophysical Research Letters*, 24, 2515–2518, 1997.](https://doi.org/10.1029/2000GL013628)
- 20 Séférian, R., Bopp, L., Gehlen, M., Orr, J. C., Ethé, C., Cadule, P., Aumont, O., Méliá, D. S. y., Voldoire, A., and Madec, G.: Skill assessment of three earth system models with common marine biogeochemistry, *Clim. Dynam.*, 40, 2549–2573, doi:[10.1007/s00382-012-1362-8](https://doi.org/10.1007/s00382-012-1362-8), 2013.
- Severmann, S., MacManus, J., Berelson, W. M., and Hammond, D. E.: The continental shelf benthic iron flux and its isotope composition, *Geochim. Cosmochim. Ac.*, 74, 3984–4004, 2010.
- Six, K. D. and Maier-Reimer, E.: Effects of plankton dynamics on seasonal carbon fluxes in an ocean general circulation model, *Global Biogeochem. Cy.*, 10, 559–583, 1996.
- 30 Smetacek, V.: Role of sinking in diatom life-history cycles: ecological, evolutionary and geological significance, *Mar. Biol.*, 84, 239–251, 1985.
- Smith, G. C., Haines, K., Kanzow, T., and Cunningham, S.: Impact of hydrographic data assimilation on the modelled Atlantic meridional overturning circulation, *Ocean Sci.*, 6, 761–774, doi:[10.5194/os-6-761-2010](https://doi.org/10.5194/os-6-761-2010), 2010.

- Smith, S. L., Yamanaka, Y., Pahlow, M., and Oschlies, A.: Optimal uptake kinetics: physiological acclimation explains the pattern of nitrate uptake by phytoplankton in the ocean, *Mar. Ecol.-Prog. Ser.*, 384, 1–12, doi:[10.3354/meps08022](https://doi.org/10.3354/meps08022), 2009.
- 5 Smith, S. L., Pahlow, M., Merico, A., and Wirtz, K. W.: Optimality-based modeling of planktonic organisms, *Limnol. Oceanogr.*, 56, 2080–2094, doi:[10.4319/lo.2011.56.6.2080](https://doi.org/10.4319/lo.2011.56.6.2080), 2011.
- Smith, S. V. and Hollibaugh, J. T.: Coastal Metabolism and the oceanic organic carbon balance, *Rev. Geophys.*, 31, 75–89, 1993.
- Smith, W. O. S. and Nelson, D. M.: Phytoplankton bloom produced by a receding ice-edge in the Ross Sea: spatial coherence with the density field, *Science*, 227, 163–166, 1985.
- 10 Soetaert, K., Middelburg, J. J., Herman, P. M. J., and Buis, K.: On the coupling of benthic and pelagic biogeochemical models, *Earth-Sci. Rev.*, 51, 173–201, 2000.
- Sommer, U.: Nitrate and silicate competition among antarctic phytoplankton, *Mar. Biol.*, 91, 345–351, 1986.
- 15 Steinacher, M., Joos, F., Frölicher, T. L., Bopp, L., Cadule, P., Cocco, V., Doney, S. C., Gehlen, M., Lindsay, K., Moore, J. K., Schneider, B., and Segschneider, J.: Projected 21st century decrease in marine productivity: a multi-model analysis, *Biogeosciences*, 7, 979–1005, doi:[10.5194/bg-7-979-2010](https://doi.org/10.5194/bg-7-979-2010), 2010.
- Stemmann, L., Jackson, G. A., and Gorsky, G.: A vertical model of particle size distributions and fluxes in the mid-water column that includes biological and physical processes—Part II Application to a three year survey in the NW Mediterranean Sea, *Deep-Sea Res. Pt. I*, 51, 885–908, 2004.
- 20 Stoecker, D. K.: Conceptual models of mixotrophy in planktonic protists and some ecological and evolutionary implications, *Eur. J. Protistol.*, 34, 281–290, doi:[10.1016/S0932-4739\(98\)80055-2](https://doi.org/10.1016/S0932-4739(98)80055-2), 1998.
- Sunda, W. G. and Huntsman, S. A.: Iron uptake and growth limitation in oceanic and coastal phytoplankton, *Mar. Chem.*, 50, 189–206, 1995.
- 25 Sunda, W. G. and Huntsman, S. A.: Interrelated influence of iron, light and cell size on marine phytoplankton growth, *Nature*, 390, 389–392, 1997.
- Tagliabue, A. and Arrigo, K. R.: Processes governing the supply of iron to phytoplankton in stratified seas, *J. Geophys. Res.*, 111, C06019, doi:[10.1029/2005JC003363](https://doi.org/10.1029/2005JC003363), 2006.
- 30 Tagliabue, A. and Völker, C.: Towards accounting for dissolved iron speciation in global ocean models, *Biogeosciences*, 8, 3025–3039, doi:[10.5194/bg-8-3025-2011](https://doi.org/10.5194/bg-8-3025-2011), 2011.
- Tagliabue, A., Bopp, L., Aumont, O., and Arrigo, K.: Influence of light and temperature on the marine iron cycle: from theoretical to global modeling, *Global Biogeochem. Cy.*, 23, C06019, doi:[10.1029/2005JC003363](https://doi.org/10.1029/2005JC003363), 2009a.

- Tagliabue, A., Bopp, L., Roche, D. M., Bouttes, N., Dutay, J.-C., Alkama, R., Kageyama, M., Michel, E., and Paillard, D.: Quantifying the roles of ocean circulation and biogeochemistry in governing ocean carbon-13 and atmospheric carbon dioxide at the last glacial maximum, *Clim. Past*, 5, 695–706, doi:[10.5194/cp-5-695-2009](https://doi.org/10.5194/cp-5-695-2009), 2009b.
- Tagliabue, A., Bopp, L., Dutay, J.-C., Bowie, A. R., Chever, F., Jean-Baptiste, P., Bucciarelli, E., Lanzuzel, D., Sarthou, G., Aumont, O., Gehlen, M., and Jeandel, C.: Hydrothermal contribution to the oceanic dissolved iron inventory, *Nat. Geosci.*, 3, 252–256, doi:[10.1038/ngeo818](https://doi.org/10.1038/ngeo818), 2010.
- Tagliabue, A., Mtshali, T., Aumont, O., Bowie, A. R., Klunder, M. B., Roychoudhury, A. N., and Swart, S.: A global compilation of dissolved iron measurements: focus on distributions and processes in the Southern Ocean, *Biogeosciences*, 9, 2333–2349, doi:[10.5194/bg-9-2333-2012](https://doi.org/10.5194/bg-9-2333-2012), 2012.
- Takahashi, T., Broecker, W. S., and Langer, S.: Redfield ratio based on chemical data from isopycnal surfaces, *J. Geophys. Res.*, 90, 6907–6924, 1985.
- Takeda, S.: Influence of iron availability on nutrient consumption ratio of diatoms in oceanic waters, *Nature*, 393, 774–777, 1998.
- Taylor, K. E.: Summarizing multiple aspects of model performance in single diagram, *J. Geophys. Res.*, 106, 7183–7193, 2001.
- Taylor, S. R. and McLennan, S. M.: *The Continental Crust: Its Composition and Evolution*, Blackwell, Malden, Mass, 1985.
- Thingstad, T. F. and Lignell, R.: Theoretical models for the control of bacterial growth rate, abundance diversity and carbon demand, *Aquat. Microb. Ecol.*, 13, 19–27, 1997.
- Thompson, P.: The response of growth and biogeochemical composition to variations in daylength, temperature and irradiance in the marine diatom *Thalassiosira Pseudonana* (Bacillariophyceae), *J. Phycol.*, 35, 1215–1223, 1999.
- Timmermann, R., Goose, H., Madec, G., Fichet, T., Ethé, C., and Dulière, V.: On representation of high latitude processes in the ORCALIM global coupled sea ice-ocean model, *Ocean Model.*, 8, 175–201, 2005.
- Tirelli, V. and Mayzaud, P.: Relationship between functional response and gut transit time in the calanoid copepod *Acartia clausi*: role of food quantity and quality, *J. Plankton Res.*, 27, 557–568, doi:[10.1093/plankt/fbi031](https://doi.org/10.1093/plankt/fbi031), 2005.
- Toner, B. M., Fakra, S. C., Manganini, S. J., Santelli, C. M., Marcus, M. A., Moffett, J. W., Rouxel, O., German, C. R., and Edwards, K. J.: Preservation of iron(II) by organic-rich matrices in a hydrothermal plume, *Nat. Geosci.*, 2, 197–201, doi:[10.1038/ngeo433](https://doi.org/10.1038/ngeo433), 2009.

- Tortell, P. D., Maldonado, M. T., and Price, N. M.: The role of heterotrophic bacteria in iron-limited ocean ecosystems, *Nature*, 383, 330–332, doi:[10.1038/383330a0](https://doi.org/10.1038/383330a0), 1996.
- 5 Tortell, P. D., Maldonado, M. T., Granger, J., and Price, N. M.: Marine bacteria and biogeochemical cycling of iron in the oceans, *FEMS Microbiol. Ecol.*, 29, 1–11, 1999.
- Tréguer, P., and De La Rocha, C. L.: The World Ocean Silica Cycle, *Annual Review of Marine Science*, 5, 477–501, doi:[10.1146/annurev-marine-121211-172346](https://doi.org/10.1146/annurev-marine-121211-172346), 2012.
- 10 Trenberth, K. E., Olson, J. G., and Large, W. G.: A Global Ocean Wind Stress Climatology Based on the ECMWF Analyses, NCAR/TN-338+STR NCAR/TN-338+STR, National Center for Atmospheric Research, Boulder, USA, 1989.
- Uitz, J., Claustre, H., Morel, A., and Hooker, S. B.: Vertical distribution of phytoplankton communities in open ocean, An assessment based on surface chlorophyll, *J. Geophys. Res.*, 111, GB3016, doi:[10.1029/2005JC003207](https://doi.org/10.1029/2005JC003207), 2006.
- 15 Uitz, J., CLaustre, H., Gentili, B., and Stramski, D.: Phytoplankton class-specific primary production in the world's oceans: seasonal and interannual variability from satellite observations, *Global Biogeochem. Cy.*, 24, GB3016, doi:[10.1029/2009GB003680](https://doi.org/10.1029/2009GB003680), 2010.
- Van Capellen, P., Dixit, S., and Van Beusekom, J. E. E.: Biogenic silica dissolution in the oceans: reconciling experimental and field-based dissolution rates, *Global Biogeochem. Cy.*, 16, 1075, doi:[10.1029/2001GB001431](https://doi.org/10.1029/2001GB001431), 2002.
- 20 Vichi, M. and Masina, S.: Skill assessment of the PELAGOS global ocean biogeochemistry model over the period 1980–2000, *Biogeosciences*, 6, 2333–2353, doi:[10.5194/bg-6-2333-2009](https://doi.org/10.5194/bg-6-2333-2009), 2009.
- Wagener, T., Guieu, C., and Leblond, N.: Effects of dust deposition on iron cycle in the surface Mediterranean Sea: results from a mesocosm seeding experiment, *Biogeosciences*, 7, 3769–3781, doi:[10.5194/bg-7-3769-2010](https://doi.org/10.5194/bg-7-3769-2010), 2010.
- 25 Wanninkhof, R.: Relationship between wind speed and gas exchange over the ocean, *J. Geophys. Res.*, 97, 7373–7382, 1992.
- Wu, J. and Luther, G. W.: Complexation of Fe(III) by natural organic ligands in the Northwest Atlantic Ocean by a competitive ligand equilibration method and a kinetic approach, *Mar. Chem.*, 50, 159–177, 1995.
- 30 Wu, J., Boyle, E., Sunda, W., and Wen, L. S.: Soluble and colloidal iron in the oligotrophic North Atlantic and North Pacific, *Science*, 293, 847–849, 2001.
- Wu, K. and Boyle, E.: Iron in the Sargasso Sea: implications for the processes controlling dissolved Fe distribution in the ocean, *Global Biogeochem. Cy.*, 16, GB1086, doi:[10.1029/2001GB001453](https://doi.org/10.1029/2001GB001453), 2002.

- 5 Xin, P. and Arkin, P. A.: Global precipitation: a 17-year monthly analysis based on gauge observations, satellite estimations, and numerical model inputs, *B. Am. Meteorol. Soc.*, 78, 2539–2558, 1997.
- Ye, Y., Völker, C., and Wolf-Gladrow, D. A.: A model of Fe speciation and biogeochemistry at the Tropical Eastern North Atlantic Time-Series Observatory site, *Biogeosciences*, 6, 2041–2061, doi:[10.5194/bg-6-2041-2009](https://doi.org/10.5194/bg-6-2041-2009), 2009.
- 10 Ye, Y., Wagener, T., Völker, C., Guieu, C., and Wolf-Gladrow, D. A.: Dust deposition: iron source or sink? a case study, *Biogeosciences*, 8, 2107–2124, doi:[10.5194/bg-8-2107-2011](https://doi.org/10.5194/bg-8-2107-2011), 2011.
- Yool, A., Popova, E. E., and Anderson, T. R.: Medusa-1.0: a new intermediate complexity plankton ecosystem model for the global domain, *Geosci. Model Dev.*, 4, 381–417, doi:[10.5194/gmd-4-381-2011](https://doi.org/10.5194/gmd-4-381-2011), 2011.
- 15 Yool, A., Popova, E. E., and Anderson, T. R.: MEDUSA-2.0: an intermediate complexity biogeochemical model of the marine carbon cycle for climate change and ocean acidification studies, *Geosci. Model Dev.*, 6, 1767–1811, doi:[10.5194/gmd-6-1767-2013](https://doi.org/10.5194/gmd-6-1767-2013), 2013.
- Yoshioka, Y. and Saijo, Y.: Photoinhibition and recovery of NH_4^+ -oxidizing bacteria and NO_2^- -oxidizing bacteria, *J. Gen. Appl. Microbiol.*, 30, 151–166, 1984.
- Zehr, J. P.: Nitrogen fixation by marine cyanobacteria, *Trends Microbiol.*, 19, 162–173, 2011.
- 20 Zondervan, I.: The effects of light, macronutrients, trace metals and CO_2 on the production of calcium carbonate and organic carbon in coccolithophores? A review, *Deep-Sea Res. Pt. II*, 54, 521–537, 2007.

Table 1a. Model parameters for phytoplankton with their default values in PISCES.

Parameter	Units	Value	Description
μ_{\max}^0	d^{-1}	0.6	Growth rate at 0°C
μ_{ref}	d^{-1}	1.0	Growth rate reference for light limitation
b_{resp}	d^{-1}	0.033	Basal respiration rate
b_P	–	1.066	Temperature sensitivity of growth
α^I	$(\text{W m}^{-2})^{-1} \text{d}^{-1}$	2; 2	Initial slope of P - I curve
δ^I	–	0.05; 0.05	Exudation of DOC
β_1^I	–	2.1; 1.6	Absorption in the blue part of light
β_2^I	–	0.42; 0.69	Absorption in the green part of light
β_3^I	–	0.4; 0.7	Absorption in the red part of light
$K_{\text{PO}_4}^{I,\text{min}}$	nmol P L^{-1}	0.8; 2.4	Minimum half-saturation constant for phosphate
$K_{\text{NH}_4}^{I,\text{min}}$	$\mu\text{mol N L}^{-1}$	0.013; 0.039	Minimum half-saturation constant for ammonium
$K_{\text{NO}_3}^{I,\text{min}}$	$\mu\text{mol N L}^{-1}$	0.13; 0.39	Minimum half-saturation constant for nitrate
$K_{\text{Si}}^{D,\text{min}}$	$\mu\text{mol Si L}^{-1}$	1	Minimum half-saturation constant for silicate
K_{Si}^I	$\mu\text{mol Si L}^{-1}$	16.6	Parameter for the half-saturation constant
$K_{\text{Si}}^{I,\text{min}}$	$\mu\text{mol Si L}^{-1}$	2; 20	Parameters for Si/C
$K_{\text{Fe}}^{I,\text{min}}$	nmol Fe L^{-1}	1; 3	Minimum half-saturation constant for iron uptake
S_{rat}^I	–	3; 3	Size ratio of Phytoplankton
$\theta_{\text{Si},D}^{I,\text{opt}}$	$\text{mol Si (mol C)}^{-1}$	0.159	Optimal Si/C uptake ratio of diatoms
$\theta_{\text{opt}}^{\text{Fe},I}$	$\mu\text{mol Fe (mol C)}^{-1}$	7; 7	Optimal iron quota
$\theta_{\text{max}}^{\text{Fe},I}$	$\mu\text{mol Fe (mol C)}^{-1}$	40; 40	Maximum iron quota
m^I	d^{-1}	0.01; 0.01	phytoplankton mortality rate
w^P	$\text{d}^{-1} \text{mol C}^{-1}$	0.01	Minimum quadratic mortality of phytoplankton
w_{max}^D	$\text{d}^{-1} \text{mol C}^{-1}$	0.03	Maximum quadratic mortality of diatoms
$\theta_{\text{max}}^{\text{Chl},I}$	$\text{mg Chl (mg C)}^{-1}$	0.033; 0.05	Maximum Chl/C ratios of phytoplankton
$\theta_{\text{min}}^{\text{Chl}}$	$\text{mg Chl (mg C)}^{-1}$	0.0033	Minimum Chl/C ratios of phytoplankton
I_{max}	$\mu\text{mol C L}^{-1}$	1; 1	Threshold concentration for size dependency

Table 1b. Model parameters for zooplankton with their default values in PISCES.

Parameter	Units	Value	Description
b_Z	–	1.079; 1.079	Temperature sensitivity term
e_{\max}^I	–	0.3; 0.35	Maximum growth efficiency of zooplankton
σ^I	–	0.3; 0.3	Non-assimilated fraction
γ^I	–	0.6; 0.6	Excretion as DOM
g_m^I	d^{-1}	3; 0.75	Maximum grazing rate
g_{FF}^M	$(\text{mmolL}^{-1})^{-1}$	2×10^3	Flux feeding rate
K_G^I	μmolCL^{-1}	20; 20	Half-saturation constant for grazing
p_P^I	–	1; 0.3	Preference for nanophytoplankton
p_D^I	–	0.5; 1	Preference for diatoms
p_{POC}^I	–	0.1, 0.3	Preference for POC
p_Z^M	–	1.0	Preference for microzooplankton
F^I_{thresh}	μmolCL^{-1}	0.3; 0.3	Food threshold for zooplankton
J^Z_{thres}	μmolCL^{-1}	0.001	Specific food thresholds for microzooplankton
J^M_{thres}	μmolCL^{-1}	0.001	Specific food thresholds for mesozooplankton
m^I	$(\mu\text{molCL}^{-1})^{-1}\text{d}^{-1}$	0.004; 0.03	Zooplankton quadratic mortality
r^I	d^{-1}	0.03, 0.005	Zooplankton linear mortality
K_m	μmolCL^{-1}	0.2	Half-saturation constant for mortality
ν^I	–	0.5; 0.75	Fraction of calcite that does not dissolve in guts
$\theta_{\text{Fe,Zoo}}^I$	$\mu\text{molFemolC}^{-1}$	10	Fe/C ratio of zooplankton

Table 1c. Model parameters for DOM with their default values in PISCES.

Parameter	Units	Value	Description
λ_{DOC}	d^{-1}	0.3	Remineralization rate of DOC
K_{DOC}	$\mu\text{mol CL}^{-1}$	417	Half-saturation constant for DOC remin.
$K_{\text{NO}_3}^{\text{Bact}}$	$\mu\text{mol NL}^{-1}$	0.03	NO_3 half-saturation constant for DOC remin.
$K_{\text{NH}_4}^{\text{Bact}}$	$\mu\text{mol NL}^{-1}$	0.003	NH_4 half-saturation constant for DOC remin.
$K_{\text{PO}_4}^{\text{Bact}}$	$\mu\text{mol PL}^{-1}$	0.003	PO_4 half-saturation constant for DOC remin.
$K_{\text{Fe}}^{\text{Bact}}$	nmol Fe L^{-1}	0.01	Fe half-saturation constant for DOC remin.
a_1	$(\mu\text{mol CL}^{-1})^{-1}\text{d}^{-1}$	0.37	Aggregation rate (turbulence) of DOC \rightarrow POC
a_2	$(\mu\text{mol CL}^{-1})^{-1}\text{d}^{-1}$	102	Aggregation rate (turbulence) of DOC \rightarrow POC
a_3	$(\mu\text{mol CL}^{-1})^{-1}\text{d}^{-1}$	3530	Aggregation rate (turbulence) of DOC \rightarrow GOC
a_4	$(\mu\text{mol CL}^{-1})^{-1}\text{d}^{-1}$	5095	Aggregation rate (Brownian) of DOC \rightarrow POC
a_5	$(\mu\text{mol CL}^{-1})^{-1}\text{d}^{-1}$	114	Aggregation rate (Brownian) of DOC \rightarrow POC

Table 1d. Model parameters for particulate organic and inorganic matter with their default values in PISCES.

Parameter	Units	Value	Description
λ_{POC}	d^{-1}	0.025	Degradation rate of POC
w_{POC}	m d^{-1}	2	Sinking speed of POC
$w_{\text{GOC}}^{\text{min}}$	m d^{-1}	30	Minimum sinking speed of GOC _b
w_{dust}	m s^{-1}	2	Sinking speed of dust
a_6	$(\mu\text{mol CL}^{-1})^{-1}\text{d}^{-1}$	25.9	Aggregation rate (turbulence) of POC→GOC
a_7	$(\mu\text{mol CL}^{-1})^{-1}\text{d}^{-1}$	4452	Aggregation rate (turbulence) of POC→GOC
a_8	$(\mu\text{mol CL}^{-1})^{-1}\text{d}^{-1}$	3.3	Aggregation rate (settling) of POC→GOC
a_9	$(\mu\text{mol CL}^{-1})^{-1}\text{d}^{-1}$	47.1	Aggregation rate (settling) of POC→GOC
$\lambda_{\text{Fe}}^{\text{min}}$	d^{-1}	3×10^{-5}	Minimum scavenging rate of iron
λ_{Fe}	$\text{d}^{-1} \mu\text{mol}^{-1} \text{L}$	0.005	Slope of the scavenging rate of iron $\times \lambda_{\text{am1}}$
$\lambda_{\text{Fe}}^{\text{dust}}$	$\text{d}^{-1} \text{mg}^{-1} \text{L}$	150	Scavenging rate of iron by dust
λ_{CaCO_3}	d^{-1}	0.197	Dissolution rate of calcite
n_{ca}	–	1	Exponent in the dissolution rate of calcite
χ_{lab}^0	–	0.5	Proportion of the most labile phase in PSi
$\lambda_{\text{PSi}}^{\text{slow}}$	d^{-1}	0.003	Slow dissolution rate of BSi
$\lambda_{\text{PSi}}^{\text{fast}}$	d^{-1}	0.025	Fast dissolution rate of BSi

Table 1e. Model parameters for various processes with their default values in PISCES.

Parameter	Units	Value	Description
λ_{NH_4}	d^{-1}	0.05	Maximum nitrification rate
$\text{O}_2^{\text{min},1}$	$\mu\text{mol O}_2 \text{L}^{-1}$	1	Half-saturation constant for denitrification
$\text{O}_2^{\text{min},2}$	$\mu\text{mol O}_2 \text{L}^{-1}$	6	Half-saturation constant for denitrification
L_{T}	nmol L^{-1}	0.6	Total concentration of iron ligands
$N_{\text{fix}}^{\text{m}}$	$\mu\text{mol N L}^{-1} \text{d}^{-1}$	0.013	Maximum rate of nitrogen fixation
$K_{\text{Fe}}^{\text{Dz}}$	nmol Fe L^{-1}	0.1	Fe half-saturation constant of nitrogen fixation
E_{fix}	W m^{-2}	50	Photosynthetic parameter of nitrogen fixation
Fe_{ice}	nmol Fe L^{-1}	15	Iron concentration in sea ice
$F_{\text{Fe},\text{min}}^{\text{sed}}$	$\mu\text{mol Fe m}^{-2} \text{d}^{-1}$	2	Maximum sediment flux of Fe
Sol_{Fe}	–	0.02	Solubility of iron in dust
O_2^{it}	$\text{mol O}_2 (\text{mol C})^{-1}$	133/122	O/C for ammonium-based processes
O_2^{nit}	$\text{mol O}_2 (\text{mol C})^{-1}$	32/122	O/C ratio of nitrification
$r_{\text{NH}_4}^*$	$\text{mol N} (\text{mol C})^{-1}$	3/5	C/N ratio of ammonification
$r_{\text{NO}_3}^*$	$\text{mol N} (\text{mol C})^{-1}$	105/16	C/N ratio of denitrification
$\theta_{\text{N},\text{C}}^{\text{N}}$	$\text{mol N} (\text{mol C})^{-1}$	16/122	N/C Redfield ratio
r_{CaCO_3}	–	0.3	Rain-ratio parameter

Table 2. Boolean variables in the namelist. These variables activate functionalities of PISCES.

Boolean name	Description
ln_co2int	Read atmospheric pco2 from a file (T) or constant (F)
ln_presatm	Constant atmospheric pressure (F) or from a file (T)
ln_varpar	PAR made a variable fraction of shortwave (T) or not (F)
ln_newprod	Use Eq. (2a) (T) or Eq. (2b) for phytoplankton growth
ln_dust	Dust input from the atmosphere (T)
ln_solub	Variable solubility of iron in dust (T)
ln_river	River discharge of nutrient (T)
ln_ironsed	Sedimentary source of iron (T)
ln_ironice	Iron input from sea ice (T)
ln_hydrofe	Iron input from hydrothermalism (T)
ln_pisdmp	Relaxation of some tracers to a mean value (T) ^a
ln_check_mass	Check mass conservation (T)

^a The frequency at which the restoring technique is applied is specified by the parameter nn_pisdmp.

Table 3. Available CPP keys in PISCES.

CPP Key	Description
<code>key_pisces</code>	Activate the PISCES model
<code>key_kriest</code>	Activate the Kriest model (see Sect. 4.4)
<code>key_sed</code>	Activate the sediment model (see Sect. 4.10)

Table 4. Global annual budget of C in the top 150 m of the ocean.

Carbon budget ^a	
Primary production in the top 150 m of the ocean	
7.5	Primary production by diatoms
36.8	Primary production by nanophytoplankton
44.3	Global total primary production
Export from the top 150 m of the ocean	
3.9	Vertical flux due to sinking big POC
2	Vertical flux due to sinking small POC
1	Advective/diffusive vertical flux of organic matter
6.9 ^b	Total vertical flux of organic matter
Various fluxes in the top 150 m of the ocean	
35.8	Grazing by microzooplankton on phytoplankton
40.2	Total grazing by microzooplankton
4	Grazing by mesozooplankton on phytoplankton
11.2	Total grazing by mesozooplankton
51.2	Total grazing by zooplankton
22.3	Remineralization of DOC

^a Carbon fluxes are all in Gt C yr⁻¹.

^b The total vertical flux due to sinking POC is 7.3 Gt C yr⁻¹ at 100m depth.

Table 5. Global annual budget of Calcite and Si in the top 150 m of the ocean.

Calcite budget ^a		
	1.6	Production of calcite
	0.8	Dissolution of calcite
0.8		Vertical flux of sinking calcite particles
Biogenic silica budget ^b		
	145.8	Production of BSi
	39.6	Dissolution of BSi
106.2		Vertical flux of dissolved BSi

^aCalcite fluxes are all in Gt C_{yr}⁻¹.

^bBiogenic silica fluxes are all in Tmol Si yr⁻¹.

Table 6. Annual budget* of N over the global ocean.

Sources of Nitrogen to the ocean	
36	River discharge
67	Atmospheric deposition
111.8	Nitrogen fixation
214.8	Total input of Nitrogen
Sinks of Nitrogen from the ocean	
77.6	Denitrification in the water column
92.8	Denitrification in the sediments
23.2	Permanent burial in the sediments
193.6	Total loss of Nitrogen
21.2	Net budget of Nitrogen (Sources minus Sinks)

* All nitrogen fluxes are in Tg N yr^{-1} .

Table 7. Sensitivity experiments performed with PISCES to evaluate the impact of specific parameterizations. Primary Production (PP) and Export Production at 150 m (EP) are in Gt C yr^{-1} .

Experiment	Description	Parameterization choices	PP	EP
PAR	Impact of variable PAR fraction	<code>ln_varpar = .false.</code>	44.4	5.8
LIGHT	Impact of light limitation	Eq. (2b)	42.6	7.3
SIZE	Impact of variable cell sizes	<code>xsizern, xsizerd = 1</code>	44.8	6.2
FOOD	Impact of food quality	$\theta^{N,I} = 0.136$	43.4	6.1

Table A1a. Model parameters for phytoplankton with their default values in PISCES.

Parameter	Coding name
b_{resp}	bresp
α^I	pislope; pislope2
δ^I	excret; excret2
$K_{\text{NH}_4}^{I,\text{min}}$	concnh4; concdnh4
$K_{\text{NO}_3}^{I,\text{min}}$	concnno3; concdno3
K_{Si}^1	xksil
K_{Si}^2	xksi2
$K_{\text{Fe}}^{I,\text{min}}$	concnfer; concdfer
S_{rat}^I	xsizern; xsizerd
$\theta_{\text{m}}^{\text{Si},D}$	grosip
$\theta_{\text{opt}}^{\text{Fe},I}$	qnfelim; qdfelim
$\theta_{\text{max}}^{\text{Fe},I}$	fecnm; fecdm
m^I	mprat; mprat2
w^P	wchl
w_{max}^D	wchld
K_{m}	xkmort
$\theta_{\text{max}}^{\text{Chl},I}$	chlcnm; chlcdm
$\theta_{\text{min}}^{\text{Chl}}$	chlcmin
I_{max}	xsizephy; xsizedia

Table A1b. Model parameters for zooplankton with their default values in PISCES.

Parameter	Coding name
e_{\max}^I	epsher; epsher2
σ^I	unass; unass2
γ^I	sigma; sigma2
g_m^I	graze; graze2
g_{FF}^M	grazflux
K_G^I	xkgraz; xkgraz2
p_P^I	xpref2p; xprefp
p_D^I	xpref2d; xprefc
p_{POC}^I	xpref2c; xprefpoc
p_Z^M	xprefz
F_{thresh}^I	xthresh; xthresh2
m^I	mzrat; mzrat2
r^I	resrat; resrat2
ν^I	part; part2
$\theta^{\text{Fe,Zoo}}$	ferat3

Table A1c. Model parameters for organic and inorganic matter with their default values in PISCES.

Parameter	Coding name
λ_{DOC}	xremik
K_{DOC}	xkdoc
$K_{\text{NO}_3}^{\text{Bact}}$	concbno3
$K_{\text{NH}_4}^{\text{Bact}}$	concbnh4
$K_{\text{Fe}}^{\text{Bact}}$	concbfe
λ_{POC}	xremip
w_{POC}	wsbio
$w_{\text{GOC}}^{\text{min}}$	wsbio2
w_{dust}	wdust
λ_{Fe}	xlam1
$\lambda_{\text{Fe}}^{\text{dust}}$	xlamdust
λ_{CaCO_3}	kdca
nca	nca
χ_{lab}^0	xsilab
$\lambda_{\text{PSi}}^{\text{slow}}$	xsirem
$\lambda_{\text{PSi}}^{\text{fast}}$	xsiremlab

Table A1d. Model parameters for various processes with their default values in PISCES.

Parameter	Coding name
λ_{NH_4}	nitrif
$\text{O}_2^{\text{min},1}$	oxymin
L_T	ligand
$N_{\text{fix}}^{\text{m}}$	nitrfix
$K_{\text{Fe}}^{\text{Dz}}$	concfediaz
E_{fix}	diazolight
Fe_{ice}	icefeinput
$F_{\text{Fe},\text{min}}^{\text{sed}}$	sedfeinput
Sol_{Fe}	dustsolub
r_{CaCO_3}	caco3r




Fig. 1. Architecture of PISCES. This figure only shows the ecosystem model omitting thus oxygen and the carbonate system. The elements which are explicitly modeled are indicated in the left corner of each box.




Fig. 2. Reduction of growth rate when the mixed layer depth exceeds the euphotic depth for nanophytoplankton (continuous line) and diatoms (dashed line). Depth corresponds to ΔZ .




Fig. 3. $\theta_{\text{opt}}^{\text{Si},D}$ as a function of Si concentration and $F_{\text{lim},1}^{D^{\text{Si}}}$. The vertical axis corresponds to $\log(\text{Si})$.




Fig. 4. Dissolution rate of PSI (λ_{PSi}^*) normalized to its value at 0 °C with no silicate. Temperature is in °C.




Fig. 5. Sediment source of iron as a function of depth. This plot displays the vertical variation of F_{sed} (see Eq. 85 for the definition of this factor).




Fig. 6. Annual-mean depth averaged N_2 fixation rates in $\mu\text{molN m}^{-2} \text{d}^{-1}$. (a) Database from the MAREMIP project (Luo et al., 2013); (b) model predictions.




Fig. 7. Surface seasonal mean Chlorophyll concentrations ($\text{mg chl } a \text{ m}^{-3}$) in April–May–June (panels **a** and **c**) and November–December–January (panels **b** and **d**). Panels (**a**) and (**b**) display satellite observations from GLOBCOLOUR. Panels (**c**) and (**d**) are model results.

Surface seasonal-mean Chlorophyll concentrations (mgchl *a*) in November–December–January for the Southern Ocean (south of 35S). Panel (a) corresponds to satellite observations from GLOBCOLOUR corrected using the algorithm developed by Dierssen and Smith (2000). Panel (b) shows model results.




Fig. 8. Spatial distribution of annual mean iron concentrations (in nmol L^{-1}) as observed (left column) and as simulated by PISCES (right column). On panels **(a)** and **(b)**, iron has been averaged over the top 50 m of the ocean. On panels **(b)** and **(c)**, iron has been averaged over 200–1000 m. The bottom two panels display the iron distributions average over the depth range 1000–5000 m. Model values have been sampled at the same location and month as the data.




Fig. 9. Annual mean NO₃ concentrations in $\mu\text{mol N L}^{-1}$. Observations are from the World Ocean Atlas 2009 (Garcia et al., 2010). (a) Observed surface. (b) Model run surface. (c) Observed transect zonally averaged over the Atlantic. (d) Same as (c) but for the model. (e) Observed transect zonally averaged over the Pacific. (f) Same as (e) but for the model.




Fig. 10. Annual mean SiO_3 concentrations in $\mu\text{mol Si L}^{-1}$. Observations are from the World Ocean Atlas 2009 (Garcia et al., 2010). Panels are the same as on Fig. 9.




Fig. 11. Annual mean O₂ concentrations in $\mu\text{mol L}^{-1}$. Observations are from the World Ocean Atlas 2009 (Garcia et al., 2010). Panels are the same as on Fig. 9.




Fig. 12. Annual mean natural DIC concentrations in $\mu\text{mol L}^{-1}$. Observations are from GLODAP. The pre-industrial distribution of DIC has been estimated in GLODAP as the difference between total DIC and anthropogenic carbon. Panels are the same as on Fig. 9.




Fig. 13. Annual mean Alkalinity concentrations in $\mu\text{mol eq L}^{-1}$. Observations are from GLODAP. Panels are the same as on Fig. 9.




Fig. 14. Taylor diagrams of model-observation comparisons for surface chlorophyll (log10-transformed) using monthly mean fields **(a)** and annual mean fields **(b)**. Black dot corresponds to global comparison; red dot to the Atlantic ocean, green dot to the Pacific ocean, brown dot to the Indian ocean and gray dot to the Southern Ocean (south of 45° S).




Fig. 15. Taylor diagram of model-observation comparisons for mesozooplankton using monthly mean fields. Data come from the Green Ocean Project website. Black dot corresponds to the global ocean; red dot to the Atlantic ocean, green dot to the Pacific ocean, brown dot to the Indian ocean and gray dot to the Southern Ocean (south of 45° S).




Fig. 16. Taylor diagrams of model-observation comparisons for nutrients using monthly mean fields. The data are identical to those used in previous plots. Panel (a) corresponds to the global ocean. Panel (b) shows the comparison restricted to the top 100 m of the ocean. Black dot corresponds to NO_3 , brown dot to O_2 , red dot to PO_4 , green dot to SiO_3 , light blue dot to DIC, purple dot to Alkalinity and gray dot to iron. The additional purple dot labeled as Alk-Lee uses the database constructed by Lee et al. (2006) to compare with the model.




Fig. 17. Surface seasonal mean Chlorophyll anomaly ($\text{mg chl } a \text{ m}^{-3}$) relative to the standard simulation in April–May–June (left column) and November–December–January (right column). Panels (a) and (b) correspond to the LIGHT test; panels (c) and (d) show to the SIZE test; panels (e) and (f) display the FOOD test.




Fig. 18. Day of the year at which sea surface chlorophyll is maximum. Panel (a) corresponds to the observations; panel (b) displays the standard simulation. Panel (c) shows the difference between the LIGHT and the standard experiments. Only the regions where the amplitude of the seasonal cycle exceeds $0.1 \text{ mg chl } a \text{ m}^{-3}$ are shown.




Fig. 19. Annual-mean relative change in the surface carbon biomass of total phytoplankton (panel a), microzooplankton (panel b), and mesozooplankton (panel c) in the FOOD experiment compared to the standard case.