

Validating a 1D SVAT Model in a Range of Ecosystems in USA and Australia: Evidence Towards its use as a Tool to Study Earth's System Interactions

George P. Petropoulos^{1*}, Matthew R. North¹, Gareth Ireland¹, Prashant K. Srivastava^{2,3}, Daisy V. Rendall¹

¹Department of Geography and Earth Sciences, Aberystwyth University, Aberystwyth, SY23 3DB, UK

²Hydrological Sciences Branch, NASA Goddard Space Flight Center, Greenbelt, Maryland, USA

³Earth System Science Interdisciplinary Center, University of Maryland, Maryland, USA

*Correspondence to: george.petropoulos@aber.ac.uk; Tel: +44-0-1970-621861

ABSTRACT

This paper describes the validation of the SimSphere SVAT model conducted at a range of sites in the USA and Australia representative of different ecosystem types. Specific focus was given to examining the models' ability in predicting Shortwave Incoming Solar Radiation (R_g), Net Radiation (R_{net}), Latent Heat (LE), Sensible Heat (H), Air Temperature at 1.3m ($T_{air\ 1.3m}$) and Air Temperature at 50m ($T_{air\ 50m}$). Model predictions were compared against corresponding *in-situ* measurements acquired for a total of 72 selected days of the year 2011 obtained from 8 sites belonging to the AmeriFlux (USA) and OzFlux (Australia) monitoring networks. Selected sites were representative of a variety of environmental, biome and climatic conditions, to allow for the inclusion of contrasting conditions in the model evaluation.

Results generally showed a good agreement between the model predictions and the *in-situ* measurements, particularly so for the R_g , R_{net} , $T_{air\ 1.3m}$ and $T_{air\ 50m}$ parameters. The simulated R_g parameter exhibited a Root Mean Square Deviation (RMDS) within 25% of the observed fluxes for 58 of the 72 selected days respectively, whereas an RMSD within ~24% of the observed fluxes was reported for the R_{net} parameter for all days of study (RMSD = 58.69 Wm^{-2}). A systematic underestimation of R_g and R_{net} (Mean Bias Error (MBE) = -19.48 Wm^{-2} and -16.46 Wm^{-2}) was also found. Simulations for the $T_{air\ 1.3m}$ and $T_{air\ 50m}$ parameters showed good agreement with the *in-situ* observations, exhibiting RMSD's of 3.23°C and 3.77°C (within ~15% and ~18% of the observed) for all days of analysis respectively. Comparable, yet slightly less satisfactory simulation accuracies were exhibited for the H and LE parameters (RMSDs = 38.47 Wm^{-2} and 55.06 Wm^{-2} , ~34% and ~28% of the observed). Highest simulation accuracies were obtained for the open woodland savannah and mulga woodland sites for most of the compared parameters. Nash-Sutcliffe efficiency index for all parameters ranging from 0.720 to 0.998, suggesting a very good model representation of the observations.

To our knowledge, this study presents the most detailed evaluation of SimSphere validation done so far, and the first validation of it conducted in Australian ecosystem types. Findings are important and timely, given the expanding use of the model both as an educational and research tool today. This includes ongoing research by different Space Agencies examining its synergistic use with Earth Observation data towards the development of global operational products.

41 **Keywords:** *SimSphere, SVAT, Net Radiation, Latent Heat, Sensible Heat, Air Temperature,*
42 *Shortwave Incoming Solar Radiation, Land Surface Interactions, FLUXNET*

43

44 1. INTRODUCTION

45 The importance of studying land surface-atmosphere interactions to develop a better
46 understanding of Earth's physical processes and feedbacks is evident from several
47 investigations. Today, particularly so in the face of climate change, it has been recognised by the
48 global scientific community as a topic requiring further attention and investigation (Battrick et
49 al. 2006; Petropoulos et al., 2014). This is documented by the fact that it is of crucial importance
50 to help address directives such as the European Parliament "Directive 2000/60/EC", aimed at
51 establishing a framework for community action in the field of water policy", namely the EU
52 Water Framework Directive. On this basis, the need to develop a holistic understanding of how
53 land surface parameters characterising the planet's energy and water budget in different
54 ecosystems has never been more important (WMO, 2002; ESA, 2014).

55 Land surface parameterization schemes (LSPs, also known as land surface models (LSMs)) are
56 one of the preferred scientific tools to quantify at fine spatial and temporal resolutions Earth
57 system interactions. Those simulate a number of parameters characterising land surface
58 interactions within the lower atmospheric boundary from a predefined set of surface
59 characteristics (i.e. properties of soil, vegetation and water). Often LSP's are utilised, amongst
60 others, to assess water resources, to evaluate the hydrological impacts of changes in climate and
61 land use, to model land-atmosphere exchanges and emissions of aerosols (Prentice et al., 2014).
62 Recent developments in mathematical modelling have been driven primarily by the progress in
63 computer technology, the expansion of modelling into new fields and disciplines and the need
64 for increased accuracy in model predictions (Bellocchi et al., 2010). As a result, LSPs have
65 advanced considerably to include detailed parameterisations of momentum, energy, mass and
66 biogeochemistry (Rosolem et al., 2013).

67 One group of LSPs include the Soil-vegetation-atmosphere transfer (SVAT) models. Those are
68 mathematical representations of vertical 'views' of the physical mechanisms controlling energy
69 and mass transfers in the soil/vegetation/atmosphere continuum. These deterministic models
70 are able to provide estimates of the time course of soil and vegetation state variables at time-
71 steps compatible with the dynamics of atmospheric processes. During the last number of
72 decades SVAT models have evolved from simple energy balance parameterisations e.g. the
73 bucket schemes adopted by Manabe (1969), through the schemes of Deardorff (1978), to the
74 biosphere-atmosphere transfer scheme (BATS) of Dickinson et al. (1986) and the simple
75 biosphere (SiB) model of Sellers et al. (1986). At present, SVATs are able to describe the
76 multifarious transfer processes through varying degrees of complexity, including the energy,
77 water and carbon dioxide (CO₂) fluxes between the ground surface covered by different
78 vegetation types and the atmosphere over different temporal and spatial scales (Olchev et al.,
79 2008). These require an application context constrained by input variables (atmospheric
80 forcing and vegetation) and input parameters (soil and vegetation properties, initialisation) to
81 simulate the water and energy budget at the surface (Coudert et al., 2008; Ridler et al., 2012).

82 However, before applying a computer simulation model to perform any kind of analysis or
83 operation, a variety of validity tests need to be executed. The process of validating a
84 mathematical model's performance, coherence and representation of the natural environment
85 is regarded as an essential step in its development. This allows an evaluation of its ability to
86 systematically reproduce the system being simulated (model reliability) and the level of
87 accuracy in which the model reproduces the natural environment (model usefulness) (Huth

88 and Holzworth, 2005; Wallach, 2006). Numerous model validation techniques exist; for a
89 comprehensive overview on the topic see for example Bellocchi et al. (2010). The procedures to
90 perform the task of validation appear in several forms, depending on data availability, system
91 characteristics and researchers' opinion (Hsu et al., 1999). A common strategy is to examine the
92 model's simulation versus actual observations acquired from the real world using common
93 statistical metrics, and several validation studies of this type of have been undertaken globally
94 (Henderson-Sellers et al., 1995; Viterbo and Beljaars, 1995; Liang et al., 1998; Wang et al., 2007;
95 Abramowitz et al., 2008; Slevin et al., 2015). In addition, Kramer *et al.* (2002) in an attempt to
96 holistically assess the capability of a model of portraying a real world system, has proposed a
97 set of model assessment criteria, namely: accuracy, generality and realism. Accuracy is
98 described by Kramer et al. (2002) as the 'goodness of fit' to *in-situ* measurements. Generality is
99 described as the applicability of the model in numerous ecosystems. Realism is described as the
100 ability of the model to address relationships between modelled phenomena.

101 The SimSphere land biosphere model is one example of a SVAT model. Formerly known as the
102 Penn-State University Biosphere-Atmosphere Modelling Scheme (PSUBAMS) (Carlson and
103 Boland, 1978; Carlson et al. 1981; Lynn and Carlson, 1990), this 1-d model was considerably
104 modified to its current state by Gillies et al. (1997) and Petropoulos et al. (2013a). Since its
105 early development, the model has become highly variable in its applicational use (for a recent
106 overview of the model use and its applications see Petropoulos et al., 2009a). Amongst others,
107 it has been involved in studies concerning the study of land surface interactions (Todhunter
108 and Terjung, 1987; Ross and Oke, 1988) and the examination of hypothetical scenarios
109 examining feedback processes (Wilson et al., 1999; Grantz et al., 1999). Furthermore, its use
110 synergistically with Earth Observation (EO) data is being considered at present for the
111 development of operational products of energy fluxes and/or soil moisture on a global scale
112 (Chauhan et al., 2003; ESA STSE, 2012). These investigations have been based around the
113 implementation of a technique commonly termed in the literature as the 'triangle' (Carlson,
114 2007; Petropoulos & Carlson, 2011). A variant of this method, which though is not using
115 SimSphere, it is already deployed over Spain to operationally deliver surface soil moisture at 1
116 km spatial resolution from ESAs own SMOS satellite (Piles et al., 2011).

117 As SimSphere's use is rapidly expanding worldwide as both a research and educational tool, its
118 validation and establishment of its coherence and correspondence to what it has been built to
119 simulate is of paramount importance. In this respect, a series of SA experiments have already
120 been conducted on the model (Olioso et al., 1996; Petropoulos et al., 2009b; Petropoulos et al.,
121 2013 a-c). Such studies have allowed quantifying the relative influence of each model input to
122 the simulation of key parameters by the model, rank them in order of importance and
123 understand how different parts of the model interplay. Yet, to our knowledge, validation studies
124 involving direct comparisons of SimSphere predictions against *in-situ* observations have as yet
125 been scarce and incomprehensive. Such validation exercises have so far only been performed
126 over a very small range of land use/cover types and on earlier versions of the model when it
127 was still under development (e.g. Todhunter and Terjung, 1987; Ross and Oke, 1988).
128 Furthermore, to our knowledge, very few studies, if any, have acted to specifically validate
129 SimSphere to numerous global ecosystems, for example, over Australian ecosystems. In this
130 context, and given SimSphere's currently expanding global use, a fully inclusive and
131 comprehensive validation of the model is now of fundamental importance.

132 In preview of the above, the main objective of this study was to evaluate SimSphere's ability to
133 model key parameters characterising land surface interactions. In this context, the main focus of
134 this study has been to understand specifically the models' ability in predicting Shortwave
135 Incoming Radiation (R_g), Net Radiation (R_{net}), Latent Heat (LE), Sensible Heat (H), and Air
136 temperature (T_{air}) at a height of 1.3m and 50m. Model validation is assessed through a
137 comparison of the model results with corresponding observations from actual *in-situ*
138 measurements acquired at local scale from 8 experimental sites (72 days in total) belonging to
139 the OzFlux (Australia) and AmeriFlux (USA) global monitoring networks. This allowed including
140 contrasting conditions in the model evaluation.

141

142 2. SIMSPHERE MODEL DESCRIPTION

143 This work deals with the SimSphere 1D boundary layer model devoted to the study of energy
144 and mass interactions of the Earth system. Formerly known as the Penn-State University
145 Biosphere-Atmosphere Modelling Scheme (PSUBAMS) (Carlson and Boland, 1978; Carlson et al.
146 1981), this model was considerably modified to its current state by Gillies et al. (1997) and
147 Petropoulos et al. (2013a). It is currently maintained and freely distributed from Aberystwyth
148 University, United Kingdom (<http://www.aber.ac.uk/simsphere>). Further details about the
149 model architecture can be found in Gillies (1993). In brief, the *physical* components ultimately
150 determine the microclimate conditions in the model and are grouped into three categories,
151 radiative, atmospheric and hydrological. The primary forcing of this component is the available
152 clear sky radiant energy reaching the surface or the plant canopy, calculated as a function of sun
153 and earth geometry, atmospheric transmission factors for scattering and absorption, the
154 atmospheric and surface emissivities and surface (including soil and plant) albedoes.

155 The *vertical structure* effectively correspond to the components of the Planetary Boundary
156 Layer (PBL) that are divided into four layers - a *surface mixing layer*, a *surface of constant flux*
157 *layer*, a *surface of vegetation or bare soil* layer. The depths of all three layers are somewhat
158 variable with time. The top of the mixing layer is identified by the presence of a temperature
159 inversion that caps the air in convective contact with the surface layer. At night, the situation is
160 reversed as the Earth cools down more rapidly than the atmosphere. The surface "constant
161 flux" layer evolves in the model as a series of equilibrium states between the transition layer
162 below and the mixing layer above. Heat and moisture are assumed to be instantaneously
163 conveyed between the surface and the top of the surface layer, which is chosen to be at a height
164 of 50 meters. In reality this height varies between about 20 and 50 meters. The transition layer
165 applies to a layer in which the vertical exchanges are dominated by molecular and radiative
166 effects as well as by vertical wind changes. In the case of vegetation, the transition layer is
167 represented by the microclimate within and at the top of the vegetation canopy. The substrate
168 layer refers to the depth of the soil over which heat and water is conducted. It consists of two
169 layers, a surface layer and a root zone. Water flows from the surface and the root zone to the
170 atmosphere respectively by direct evaporation or through the plants as well as between the
171 two layers. Soil water content is specified by assigning a fractional volume of field capacity,
172 which essentially is the "soil moisture availability". Five layers are used to compute the flow of
173 heat in the substrate. An initial soil temperature profile is assigned on the basis of the initial
174 surface temperature (furnished from a meteorological sounding) and a climatological substrate
175 temperature, which one obtains from mean data. A governing parameter for heat conduction is

176 the "thermal inertia" that contains both soil conductivity and soil diffusivity (or alternately, the
177 volumetric heat content). This parameter is the one that also governs the rate of H flux to or
178 from the atmosphere through the soil surface.

179 The *horizontal* component of the model is composed of 4 parts: (i) *Planetary Boundary Layer*
180 (*PBL*), (ii) *Surface Layer*, (iii) *Transition Layer* and (iv) *Substrate Layer*. Due to SimSphere
181 simulating parameters in a 1-dimensional vertical column, the model is restricted horizontally
182 only to areas representative of its initialised conditions, therefore the model has an undefined
183 spatial coverage. The vegetation component is dormant at night, that is, after radiation sunset.
184 The night time dynamics for the surface fluxes differ from those during the day time. Heat and
185 moisture fluxes are exchanged between both the ground and foliage, between plant and inter-
186 plant airspaces through stomatal and cuticular resistances in the leaf (for water vapour) and
187 the air, between soil and the interplant air spaces and between the entire vegetation canopy
188 and the air. A separate component exists for the bare soil fluxes between the surface and the
189 air. Vegetation and soil fluxes meld at the top of the vegetation canopy, their relative weights
190 depending on the fractional vegetation cover, which is specified as an input to the model. As
191 such, SimSphere is thus referred to as a form of two-stream or two-source model. The soil
192 hydraulic parameters are prescribed from the Clapp and Hornberger (1978) classification. The
193 soil surface turbulent fluxes are determined following the Monin and Obukov (1954) similarity
194 theory which takes into account atmospheric stability.

195 SimSphere represents various physical processes taking place in a column that extends from
196 the root zone below the soil surface up to a level well above the surface canopy, the top of the
197 surface mixing layer. The processes and interactions simulated by the model are allowed to
198 develop over a 24-h cycle at a chosen time step (typically 30'), starting from a set of initial
199 conditions given in the early morning. For its parameterisation, input parameters are
200 categorised into 7 defined groups; time and location, vegetation, surface, hydrological,
201 meteorological, soil and atmospheric (Table 1). From initialisation, over a 24-hour cycle
202 SimSphere assesses the evolution of more than 30 variables associated with the radiative,
203 hydrological and atmospheric physical domains.

204

205 **3. EXPERIMENTAL SET UP**

206 A total of 5 AmeriFlux and 3 OzNet experimental sites were used, providing a comprehensive
207 dataset of measured micrometeorological parameters together with general meteorological
208 observations. **The potential use of several FLUXNET sites was evaluated before deciding on the**
209 **final 8 experimental sites used in the study. Sites were excluded from analysis based on the**
210 **requirement to fulfil specific criteria, namely a) sites needed to incorporate different land cover**
211 **types for the evaluation of the model's ability to simulate fluxes over different land cover/land**
212 **use types, b) sites were required to show homogeneous land cover, invariable topography and**
213 **limited anthropogenic intervention, and c) site data needed to include measurements of the 6**
214 **parameters validated in the study simultaneously for the same day, any sites which did not**
215 **successfully meet this criteria were excluded. Experimental days were further excluded**
216 **following the pre-processing steps outlined in section 4.1.** Table 2 provides an overview of the
217 characteristics of the experimental sites used in this study. At each site, micrometeorological
218 measurements of various parameters are acquired including the turbulent fluxes of heat and
219 moisture, shortwave incoming radiation (R_g), net radiation (R_{net}) (at the surface) and air

220 temperature (T_{air}) (often at different heights). Flux measurements methods and calculations
221 performed within the FLUXNET sites are designed with the same specifications at all sites. All
222 collected data are quality-controlled and standard procedures for error corrections are
223 prescribed. Details on the FLUXNET measurements and the raw data processing can be found in
224 Aubinet et al. (2000).

225 The sites were representative of a range of ecosystem types with markedly different site
226 characteristics to include contrasting conditions in the model evaluation. All *in-situ* data
227 acquired from each site was collected covering the year 2011, allowing for a sufficient database
228 for model parameterisation and validation to be developed. All data was obtained from the
229 FLUXNET database (<http://fluxnet.ornl.gov/obtain-data>) at Level 2 processing, to allow
230 consistency and interoperability. This processing level includes the originally acquired *in-situ*
231 data from which any erroneous data caused by obvious instrumentation error have been
232 removed. Additionally, atmospheric *in-situ* data was collected from the freely distributed
233 University of Wyoming's weather balloon data archive
234 (<http://weather.uwyo.edu/upperair/sounding.html>). Local profiles of temperature, dew point
235 temperature, wind direction, wind speed and atmospheric pressure were taken from nearest
236 possible experimental sites which were also used in model parameterisation.

237

238 **4 SIMSPHERE PARAMETERISATION AND VALIDATION**

239 This section provides a synopsis of the methodology followed in parameterising and
240 subsequently evaluating SimSphere's ability to simulate key parameters characterising land
241 surface interactions. An overview of the main steps included is furnished in Figure 1.

242

243 **Figure 1:** Flowchart of the overall methodology followed

244 **4.1 Datasets Pre-processing**

245 Following data acquisition, further analysis was implemented aimed at identifying the specific
246 days for which SimSphere would be parameterised and validated for each experimental site.
247 Initially, for each site, cloudy days were identified and eliminated from any further analysis.
248 Judgment on which days (or time-periods) were cloud-free was based on the observation of R_g
249 diurnal observation, where cloud-free days were flagged as those having smoothly symmetrical
250 R_g curves, a property signifying clear-sky conditions (e.g. Carlson et al. 1991).

251 Subsequently, for the subset of days which included only the cloud-free days, the energy
252 balance closure (EBC) was computed. EBC evaluation has been accepted as a valid method for
253 accuracy assessment of turbulent fluxes derived from eddy covariance measurements (Wilson
254 et al., 2002; Barr et al., 2006). Energy imbalance provides important information on how they
255 should be compared with model simulations (e.g. Twine et al., 2000; Culf et al., 2002). In this
256 study, EBC was principally evaluated by performing a regression analysis (e.g. see Wilson and
257 Baldocchi, 2000; Wilson et al. 2002; Castellvi et al., 2006). The linear regression coefficients
258 (slope and intercept) as well as the coefficient of determination (R^2) were calculated from the
259 ordinary least squares (OLS) relationship between the 30-min estimates of the dependent flux
260 variables (LE+H) and the independently derived available energy ($R_{\text{net-G-S}}$). In addition to this,

261 the Energy Balance Ratio (EBR) parameter was computed by cumulatively summing $R_{net}-G-S$
262 and $LE+H$ from the 30-min mean average surface energy flux components, and then rationing
263 each of the cumulative sums as follows (e.g. Wilson et al. 2002 ; Liu et al., 2006):

264

$$265 \quad EBR = \frac{\sum (LE + H)}{\sum (Rn - G - S)} \quad (1)$$

266 In the above equation, G refers to the soil surface heat flux and S refers to the above ground
267 heat storage in the vegetation. This index ranges generally from zero to one, with values closer
268 to one highlighting a satisfactory diurnal energy closure, indicating a good quality of *in-situ*
269 measurements. All days with poor EBC ($EBR < 0.750$, slope < 0.85 , $R^2 < 0.930$) were excluded
270 from further analysis.

271 Further conditions were subsequently employed to ensure that selected days were of the
272 highest possible class in terms of *in-situ* data quality. Firstly, all days selected were within the
273 same year to eliminate effects ascribed from inter-annual variability in vegetation phenology or
274 climatic conditions. Secondly, selected simulation days were assessed for atmospheric stable
275 conditions, namely low wind speeds and low available energy (Maayar *et al.*, 2001). Such
276 conditions were identified by the evaluation of the *in-situ*, where direct measurements of wind
277 speed and energy flux amplitude and diurnal trend were used as indicators of atmospherically
278 stable conditions. As a result, a final set of a total of 72 non-consecutive days from the selected
279 experimental sites were identified as being suitable to be included in this study.

280 4.2 Model Parameterisation

281 SimSphere was parameterised to the daily conditions existent at the flux tower for each of the
282 selected days. *In-situ* data sets provided measurements of soil water content, temperature, wind
283 speed, wind direction and atmospheric pressure at the corresponding time of initialisation,
284 6.00am (local time). Ancillary parameters, critical for the models' initialisation, were largely
285 acquired through either the sites respective PI (for the case of OzFlux), or the FLUXNET
286 database (for the case of AmeriFlux). Such measurements included detailed information on the
287 vegetation (LAI, FVC, vegetation height, cuticle resistance), pedological (soil morphology and
288 soil classification) and topographical (slope, aspect, surface roughness) characteristics of each
289 site. If no further ancillary information was available, specific parameters were acquired
290 through the analysis of standard literature sources (e.g. Mascart et al., 1991; Carlson et al.,
291 1991). The soil type parameters were obtained using the soil texture data provided at each
292 FLUXNET test site and information supplied in some instances by the experimental site
293 managers themselves. This was also the case for the topographical information required in
294 model initialisation. Wind and water vapour sounding profiles which were attained at 06:00
295 GMT from the University of Wyoming database to correspond to the models' initialisation were
296 also used in model parameterisation. Upon completion of its initialisation, the model was
297 executed for each site/day forced by observations acquired from each site on which it had been
298 parameterised. The 30' average value of each of the targeted model outputs per site for the
299 period 0530-2330 hours was subsequently exported in SPSS to validate the model predictions.

300 4.4 Model performance assessment

301 A series of statistical terms included to evaluate the agreement between the in-situ and the
 302 SimSphere predictions, including the mean bias error (MBE, or bias- eq. 2) and mean standard
 303 deviation (MSD, or scatter- eq. 3) of the observed and modelled values, the root mean square
 304 difference (RMSD) (eq. 4), the mean absolute difference (MAD) (eq. 5) the linear regression fit
 305 model coefficient of determination (R^2) (eq. 6) and the Nash-Sutcliffe (1970) (denoted as Nash)
 306 index (eq. 7):

307

$$Bias = MBE = \frac{1}{N} \sum_{i=1}^N (P_i - O_i) \quad (2)$$

$$Scatter = MSD = \frac{1}{(N-1)} \sum_{i=1}^N (P_i - O_i - \overline{(P_i - O_i)})^2 \quad (3)$$

$$RMSD = \sqrt{bias^2 + scatter^2} \quad (4)$$

$$MAD = N^{-1} \sum_{i=1}^N |P_i - O_i| \quad (5)$$

$$R^2 = \left[\frac{\sum_{i=1}^N (P_i - \bar{P})(O_i - \bar{O})}{\left[\sum_{i=1}^N (O_i - \bar{O})^2 \sum_{i=1}^N (P_i - \bar{P})^2 \right]^{0.5}} \right]^2 \quad (6)$$

$$NASH = 1 - \frac{\sum_{i=1}^N (O_i - P_i)^2}{\sum_{i=1}^N (O_i - \bar{O})^2} \quad (7)$$

308 P denotes the “predicted” values obtained from SimSphere and O denotes the “observed” values
 309 from the selected OzFlux and AmeriFlux site-days.

310 The utilisation of these statistics has been widely demonstrated in a number of previous studies
 311 comparing model outputs to observational networks (e.g. Alexandris & Kerkides, 2003;
 312 Marshall et al., 2013). All statistical metrics were computed from comparisons performed at
 313 identical 0.5 hourly intervals between the two datasets for each day of comparison. In addition,
 314 these statistical parameters, where appropriate, were also computed for each site, providing a
 315 summary of the model predictions per experimental site on which the model was validated.

316 5. RESULTS

317 The main results from the comparisons between the SimSphere predictions and the
 318 corresponding in-situ data for the different parameters evaluated in this study are summarised
 319 in Tables 3 to 8. In addition, Figure 2 provides a graphical illustration of the agreement between
 320 the simulated values and *in-situ* measurements per parameter for all sites together and **Figure 3**
 321 **illustrates the diurnal agreement between the modelled outputs and *in-situ* observed fluxes for**
 322 **a selected site and days.** The detailed findings from the comparisons performed are made
 323 available next.

324 *Figure 2: Scatterplot comparison of SimSphere predicted and in-situ for all parameters*

325 5.1 Incoming Shortwave Radiation (R_g) at the surface

326 Simulation accuracy of R_g was largely accurate, exhibited by low RMSD (within ~19% of the
327 observed fluxes) and MAE values (RMSD = 67.83 Wm^{-2} , MAE = 46.43 Wm^{-2}) (Table 3 and Figure
328 2). A moderate underestimation of the observed fluxes was also evident (MBE= -19.48 Wm^{-2}).
329 Notably, R_g yielded the highest correlated results of all parameters assessed ($R^2 = 0.971$, NASH =
330 0.963), further illustrated in Figure 2, where the distribution of points within the feature space
331 were predominantly centred on the 1:1 line, showing a strong relationship between both
332 variables.

333 On a per site basis, the highest simulation accuracies were attained within the US_Moz
334 deciduous broadleaf site in comparison to all other sites (RMSD= 50.36 Wm^{-2} , within ~15% of
335 the observed fluxes, MAE= 36.57 Wm^{-2}). The Howard Springs woody savannah site also attained
336 comparably high simulation accuracies (RMSD= 52.53 Wm^{-2} , within ~16% of the observed
337 fluxes, MAE= 33.79 Wm^{-2}). Contrarily, model predictions of R_g for the Australian Calperum
338 grazing pasture site were significantly lower, indicating a weaker model performance (RMSD=
339 100.65 Wm^{-2} , within ~25% of the observed fluxes, MAE = 61.91 Wm^{-2}), closely followed by the
340 US_Whs shrubland site (RMSD= 90.45 Wm^{-2} , within ~25% of the observed fluxes, MAE = 46.09
341 Wm^{-2}). Within the majority of sites, model simulation consistently underestimated the *in-situ*
342 measurements (MBE = -4.85 Wm^{-2} to -56.40 Wm^{-2}), with the US_Moz deciduous forest site being
343 the only exception (MBE = 16.47 Wm^{-2}). That is, the true change (*in-situ* observations), for 6 of
344 the 7 sites tends to be larger than the model-based estimates. Inter-site variability was minimal
345 for the simulation of this parameter, with only a difference of ~9% between the minimum and
346 maximum RMSD as a percentage of the observed fluxes on a per site basis.

347 Evidently, agreement over the Australian sites generally increased for the period between
348 February to June, with a significant decrease in accuracy from August to early February. For
349 example, over the Calperum grazing pasture site, RMSD ranged from 24.14 to 53.78 Wm^{-2} (or
350 within ~6% to ~21% of the observed fluxes) for all the test days located within the period from
351 24/02/2011 to 24/04/2011. In contrast, for the same site, RMSD varied from 84.41 Wm^{-2} to
352 149.29 Wm^{-2} (or within ~41% to ~ 53% of the observed fluxes) for all the test days for the
353 period between 22/07/2011 to 29/12/2011. Similar trends were observed for all other
354 Australian sites, although some anomalies were present. In relation to the US sites the adverse
355 was found; highest simulation accuracies were predominantly derived for the test days located
356 during the period between October and late April. Clearly, periods of highest simulation
357 accuracy for both the Australian and US sites correspond to their respective summer season,
358 and are thus consistent between the two continents. Generally the results for the US sites
359 suggested that the conditions prevalent within the wet season (October to May) may have had
360 an influence on model accuracy.

361 5.2 Net Radiation (R_{net}) at the surface

362 Table 4 and Figure 2 indicate a high overall performance in the models' ability to accurately
363 predict R_{net} , confirmed by the high simulation accuracy (RMSD = 58.69 Wm^{-2} , within ~24% of
364 the observed fluxes, MAE = 46.42 Wm^{-2}) reported for all sites. Furthermore, comparisons of R_{net}
365 for all days of simulation showed a low average MSD of 54.44 Wm^{-2} , indicating the model's
366 capability to precisely represent the amplitude of the R_{net} flux, with low dispersion of variance
367 from the *in-situ* trends, as evidenced in Figure 2. MBE results indicated a moderate

368 underestimation of the *in-situ* measurements by the model (-16.49 Wm⁻²), with 7 of the 8 site
369 averages showing an underestimation of the in-situ trends (negative MBE values in a range of -
370 0.09 to -46.10 Wm⁻²). A much larger inter-site variability was reported for the model simulation
371 accuracies of the R_{net} parameter, where RMSD ranged between 33.90 Wm⁻² to 78.03 Wm⁻² (also
372 reflected in the RMSD as a percentage of observed fluxes ranging between ~16% and ~30% on
373 a per site average basis) showing to some extent a deficiency in the capability of the model to
374 capture the land surface process over varying land cover types. The R_{net} results exhibited largely
375 similar statistical agreement to those observed for those of the R_g parameter.

376 Most noticeably, in correspondence with the R_g parameter results, SimSphere showed superior
377 simulation accuracy within the Alice Springs mulga woodland site in comparison to the other
378 land cover types, with the reported accuracies significantly above the overall average (RMSD =
379 33.90 Wm⁻², within ~16% of the observed fluxes, MAE = 26.25 Wm⁻²). Moreover, the woody
380 savannah site of Howard Springs again exhibited high simulation accuracies (RMSD = 47.05
381 Wm⁻², within ~21% of the observed fluxes, MAE = 35.74 Wm⁻²), with comparable accuracies to
382 the simulation of the R_g parameter. Conversely, the model showed an inferior performance
383 when simulating R_{net} within the US_Ton wooded savannah site where a systematic and more
384 pronounced underestimations of R_{net} was evident (MBE = -46.10 Wm⁻²). This constant
385 underestimation by the model led to a poorer agreement between the model predictions and *in-*
386 *situ* observations for the US_Ton site as reflected in the statistical analysis (RMSD= 78.03 Wm⁻²,
387 within ~30% of the observed fluxes, MAE = 65.22 Wm⁻²). It should be noted that the accuracy of
388 the model estimations on a per site basis did not correlate between both the R_g and R_{net}
389 parameter estimations, with only the US_Whs shrubland site exhibiting weaker simulation
390 accuracies for both parameters, and as indicated above, a relatively high simulation accuracy for
391 the Howard Springs woody savannah site.

392 Evidently, as indicated by Table 4, trends in simulation accuracy dependent on test day were
393 apparent. Although comparable; the trends were not as prominent as those exhibited for the R_{net}
394 parameter. Within the Australian sites, low RMSD was exhibited predominantly for the test days
395 within the period of March to July, although some discrepancies were present during specific
396 days. For example, the 27th of May simulation date for the Howard Springs site reported an
397 RMSD of 70.60 Wm⁻² (within ~38% of the observed fluxes) indicating a day of unusually high
398 error for this period. However, such anomalies were limited. Generally, for the US sites, highest
399 RMSD was exhibited for the period concurrent to the wet season (October to April), with the
400 highest error rates exhibited during the dry period, for example the 27th of February simulation
401 date for the US_Ton site (RMSD = 113.80 Wm⁻², within ~73% of the observed fluxes), although
402 again, anomalies in such trends were notable yet uncommon.

403 5.3 Latent Heat (LE)

404 As presented in Table 5, highest RMSD in relation to the observed fluxes was reported for the LE
405 parameter in comparison to all other parameters evaluated (RMSD = 39.47 Wm⁻²), where the
406 model showed some deficiencies when reproducing LE fluxes in varying land cover, both in
407 terms of their seasonal and diurnal evolution. An average R² value of 0.700 is also indicative of a
408 poorer correlation between the predictions and observations of LE (Figure 2). When averaged
409 over all days and sites, the model-based estimates tended towards a conservative
410 overestimation of the observed fluxes, indicated by an average MBE of 2.84 Wm⁻².

411 On a site by site basis the US_ib1 cropland site consistently yielded the highest statistical
412 agreement between model predicted and observed values, with low error and high correlation
413 results (RMSD = 52.54 Wm⁻², within 20% of the observed fluxes, MAE = 15.16 Wm⁻², R² = 0.827,
414 NASH = 0.945). Notably, all other sites exhibited poorer agreement, with RMSD values in
415 relation to the observed fluxes above 30% for 6 of the 8 sites (RMSD within percentage of the
416 observed fluxes varying between ~34% and 83%). Generally, each site exhibited a significant
417 range of MBE, from -11.49 Wm⁻² (US_Whs) to 25.65 Wm⁻² (US_Moz), suggesting high variability
418 between the partitioning of LE in each ecosystem. Peak LE flux values exhibited high inter-site
419 variability, with both the US_ib1 (Cropland) and US_Moz (deciduous broadleaf forest) sites
420 containing the highest LE flux peaks of 458.5 Wm⁻² and 376 Wm⁻² respectively. In comparison, a
421 maximum LE flux peak of just 143.7 Wm⁻² was reported for the US_Whs (Shrubland) site,
422 suggesting a substantial range of 314.8 Wm⁻² between lowest daily peak LE and maximum daily
423 peak LE. Noticeably, trends in simulation accuracy dependent on test day were comparable to
424 both the R_g and R_{net} parameter results, however with significantly higher inter-site variability in
425 RMSD ranges.

426 5.4 Sensible Heat (H)

427 SimSphere showed a satisfactory ability to accurately simulate H fluxes in numerous
428 ecosystems for the 72 days included in this study, with an average RMSD and R² values of 55.06
429 Wm⁻², within ~28% of the observed fluxes, and 0.829 respectively. Results were largely similar
430 to that of the LE flux simulation accuracies, although model performance for the LE parameter
431 underperformed that of the H flux for the majority of statistical metrics computed herein.

432 Average RMSD values ranged from 38.07 Wm² to 69.94 Wm² (US_Var and US_Whs) and within
433 ~17% to ~68% of the observed fluxes (US_Var and US_ib1) when analysed on a site by site
434 basis, underlining the greatest inter-site variability was reported for this parameter. In addition,
435 R² values ranged from 0.73 (US_ib1) to 0.94 (US_VAR). The latter was suggestive that model
436 predictions were in generally in good agreement to the *in-situ* measurements, showing a strong
437 relationship between both variables. The grassland site (US_Var) consistently showed superior
438 model performance in comparison to all other sites, with values indicating an excellent
439 agreement to the observed diurnal evolution (RMSD = 38.07 Wm⁻², within ~17% of the
440 observed fluxes, MAE = 28.35 Wm⁻²). MSD values reported for US_Var were 19.41 Wm⁻² lower
441 than the all site average, suggesting a systematically accurate representation of H fluxes at this
442 site. MSD for H flux were directly comparable to the overall average MSD values reported for R_g
443 and R_{net}, yet significantly higher than the LE fluxes. Simulation accuracy were comparably high
444 for the simulated H fluxes for 5 of the 8 sites, with RMSD values in relation to the observed
445 fluxes above 30% (RMSD within percentage of the observed fluxes varying between ~17% and
446 30%). Notably, results for the US_ib1 site exhibited significant error, with RMSD and MSD values
447 (69.94 Wm⁻², within ~68% of the observed fluxes, and 67.73 Wm⁻² respectively).

448 For the Australian sites, no significant trends were evident dependent on simulation day, with
449 generally comparable accuracy ranges for the specific test days including anomalistic days
450 which exhibited significantly higher error ranges. For example, the Howard Springs woody
451 savannah site indicated RMSD for the majority of simulation days ranging between 28.29 Wm⁻²
452 and 50.31 Wm⁻² (within ~15% to ~21% of the observed fluxes) on a per test day basis, with the
453 18th of April and 13th of May experimental days exhibiting an RMSD of 75.86 Wm⁻² and 96.93

454 Wm^{-2} (within ~52% and ~65% respectively. Similar intra-site variability was notable for the US
455 sites.

456 **5.5 Air Temperature 1.3m ($T_{air\ 1.3m}$)**

457 SimSphere showed a high capability in simulating $T_{air\ 1.3m}$ with an average RMSD as low as
458 3.23°C (within ~15% of the observed) and relatively high R^2 value of 0.843, see Table 7.
459 Furthermore, $T_{air\ 1.3m}$ exhibited neither a consistent over or underestimation, with an overall
460 average MBE of 0.28°C. Simulation accuracy for $T_{air\ 1.3m}$ was relatively stable, with a low range of
461 RMSD values reported over all sites. RMSD values ranged from 2.17°C (within ~9% of the
462 observed) in the woodland savannah site of Howard Springs, and 4.74°C (within ~25% of the
463 observed) in the grazing pasture site of Calperum. Overall, agreement between the predictions
464 and observations was greatest for the Howard springs site, with results confirming a high
465 overall correlation to the observed diurnal evolution of $T_{air\ 1.3m}$. The deciduous broadleaf site of
466 US_Moz also exhibited comparably high simulation accuracy (RMSD = 2.38°C, within ~11% of
467 the observed, MAE = 1.84°C, NASH = 0.853). The Calperum site exhibited the weakest
468 agreement of $T_{air\ 1.3m}$ with an average RMSD 1.51°C higher than the all site average. The R^2
469 analysis further appraised the models ability to accurately simulate air temperature, with a
470 range of values indicating high correlation between model predicted and observed $T_{air\ 1.3m}$ (0.74
471 to 0.93). MSD displayed a high range of values (2.1°C to 3.76°C), showing to some extent the
472 inability of the model to consistently predict $T_{air\ 1.3m}$ with a high level of precision. The trends in
473 simulation accuracy dependent on test day were again insignificant for the $T_{air\ 1.3m}$ parameter,
474 exhibiting similar patterns to those indicated for the H flux parameter.

475 **5.6 Air Temperature 50m ($T_{air\ 50m}$)**

476 The model showed a slightly inferior performance in predicting $T_{air\ 50m}$ (RMSD = 3.77°C, within
477 ~18% of the observed) when compared to $T_{air\ 1.3m}$, with an average RMSD difference of 0.54°C
478 (~3% percentage difference in relation to the observed) (Table 8 and Figure 2). A lower average
479 R^2 value of 0.775 is reported compared to that of $T_{air\ 1.3m}$ ($R^2 = 0.843$), indicating a weaker, yet
480 close, agreement between both variables. However, the values reported still showed a highly
481 acceptable correlation between the modelled estimates and the *in-situ* measurements, as
482 indicated by an average NASH value of 0.825. Once averaged, $T_{air\ 50m}$ exhibited a minor
483 underestimation of -0.38°C; however the range of MBE reported between sites was significantly
484 less (2.1°C), suggesting a more consistent simulation of T_{air} at 50m compared to at 1.3m by
485 SimSphere. In contrast, agreement between the simulated $T_{air\ 50m}$ and *in-situ* measurements
486 resulted in a higher MSD than that reported for the $T_{air\ 1.3m}$ parameter, with the exception of the
487 Howard Springs site. When analysed on a per site basis, notably, in correspondence with the T_{air}
488 $_{1.3m}$ parameter, agreement between the estimated and measured values over both the Howard
489 Springs and US_Moz sites exhibited highest simulation accuracy (RMSD = 2.04°C and 2.85°C,
490 within ~8% and ~13% of the observed, respectively). Moreover, weakest agreement was
491 reported over the Calperum site, again in correspondence with the results of the $T_{air\ 1.3m}$
492 parameter. No systematic trends were apparent in the inter-site variability of simulation
493 accuracy dependent on test day.

494 **6. DISCUSSION**

495 The present study evaluated the ability of the SimSphere SVAT model to accurately represent
496 key parameters characterising land surface interactions within eight ecosystems in two

497 continents. A total of 72 days (10 days per site of the 8 sites selected) from year 2011 were
498 selected from Australia and USA to validate the model's ability to predict Shortwave Incoming
499 Radiation (R_g), Net Radiation (R_{net}), Latent Heat (LE), Sensible Heat (H), and Air temperature
500 (T_{air}) at a height of 1.3m and 50m.

501 Variable model performance was clearly evident when simulating both the LE and H fluxes
502 within contrasting land cover types. For example, as discussed, highest simulation accuracy was
503 attained within the grassland study sites. In contrast, simulation accuracy within forested
504 ecosystems was less satisfactory. The deciduous forest stand (US_Moz), with an average canopy
505 height of 24.2m attained significantly low simulation accuracy, and was also outperformed by
506 the Mulgia forested ecosystem (Alice Springs), characterised by a sparse canopy at a height of
507 6.5m. Such results suggest that the increased complexity and heterogeneity of forested
508 environments, particularly those with understory vegetation, can have profound effects on the
509 overall exchange of mass and energy which cannot be represented within the models
510 parameterisation and hence can impact influence LE and H outputs. The partitioning of LE and
511 H fluxes are also highly susceptible to a number of other factors. Small changes in the moisture
512 availability, most particularly from the deep layer soil water content (SWC) can have a strong
513 influence (Carlson and Lynn, 1991; Olioso et al., 2000), but also to the representativeness of the
514 radiosonde data to the existent local conditions (Taconet et al. 1986). As reported by Taconet et
515 al. (1986), an error of just $\sim 2^\circ\text{C}$ in the sounding profile temperature can cause a variation of
516 $\sim 45 \text{ Wm}^{-2}$ in the corresponding fluxes, most particularly so for H flux. SimSphere was forced
517 with surface moisture and root zone moisture availability data taken directly from the *in-situ*
518 data, as well as only nearby representative sounding profiles used an accurate representation
519 of the local conditions were attained. These highly influential parameters were consistently
520 misrepresented within the models' parameterisation, providing a possible reason in part for the
521 lower simulation accuracies attained.

522 R_g was estimated by the model to a high level of accuracy (**error within $\sim 19\%$ of the observed**
523 **fluxes**), where an R^2 value of 0.971 and a NASH value of 0.960 reported for all days of analysis
524 suggests that model predictions had excellent correlation to the observed dataset. This indicates
525 that SimSphere was able to simulate the trend of R_g well. A possible reason for the
526 underestimation of R_g by the model is perhaps linked to the solar transmission model and/or
527 the surface albedo calculation in the model, as has also been pointed out previously by
528 Todhunter and Terjung (1978). Furthermore, previous sensitivity analysis studies undertaken
529 upon the model confirm that R_g is significantly influenced by the sites aspect (Petropoulos et al.,
530 2014). Therefore simulation accuracy may partly be related to the models representation of
531 sites topographical characteristics.

532 In the majority of the experimental sites a general underestimation of R_{net} was attained by the
533 model, which led to a mean RMSD and R^2 value of 58.69 Wm^{-2} and 0.960 respectively. These
534 results are also comparable to those reported in other analogous validation studies (Carlson
535 and Boland, 1978; Todhunter and Terjung, 1987; Ross & Oke, 1988). Todhunter and Terjung
536 (1987) compared predicted R_{net} from the model versus corresponding R_{net} values obtained from
537 the literature from Los Angeles, USA, and showed both daytime and night time simulations to be
538 in agreement within the range reported in the literature. Ross and Oke (1988) also confirmed
539 the capability of the model in simulating the day-to-day variation of R_{net} for comparisons using
540 eighteen cloud-free days over an urban area of Vancouver, B.C. in Canada. Ross and Oke (1988)
541 reported an overall average RMSD error of 43 Wm^{-2} for comparisons for all cloud-free days, a

542 minor improvement on the RMSD of 58.69 Wm^{-2} presented herein. Disparity in the results
543 between this study and those studies could be the results of utilising model simulations over
544 dissimilar land cover types, where it is largely accepted that R_{net} partitioning into LE and H
545 fluxes is highly dependable on the vegetation and surface characteristics of the site (Oliosio et al.,
546 2000). Previous sensitivity analysis studies undertaken on the SimSphere further confirm this
547 observation (Petropoulos et al., 2014). Similarly to R_g , simulation accuracy of R_{net} was described
548 by Ross and Oke (1988) to be a factor of long wave radiation, mainly the values of atmospheric
549 and surface emissivities (which effect the surface temperature estimation). Increased
550 representation of the surface optical properties and long wave radiation estimation of the
551 model could greatly enhance simulation accuracy.

552 Overall simulation accuracies were lower for estimates of $T_{\text{air } 50\text{m}}$ compared to estimates of $T_{\text{air } 1.3\text{m}}$
553 in all but one site, Howard Springs. One possible explanation for this may be the
554 fundamental problem that model estimates of $T_{\text{air } 50\text{m}}$ could only be validated against ancillary
555 air temperature data obtained directly from the sites flux tower, thus direct comparison
556 specifically at 50m could not be achieved. Similarly to the LE and H fluxes, variable simulation
557 accuracies dependent on land cover types were also evident. Three sites: Calperum, US_Var and
558 US_IB1, all exhibit noticeably weaker simulation accuracies in comparison to the remaining
559 sites. On further investigation, all 3 sites show an ecosystem which is characterised by high
560 inter-annual variability of vegetation phenology, such as vegetation height, leaf width, FVC etc.
561 Modelled T_{air} peaked between 10.30 and 14.30 local time. For instances where time-lag between
562 the predicted and observed T_{air} comparisons is observed, such effects may be linked with the
563 energy storage in the vegetation and the air, as it is not taken into account in the SimSphere
564 simulations. This may partly explain some of the inaccuracies reported for T_{air} estimation in
565 Alice Springs and US_MOZ as this effect is most important for forested sites. Carlson and Boland
566 (1978) and Carlson et al. (1991) also described a similar hysteresis effect in comparisons which
567 they performed for different vegetation canopies and environmental conditions (urban and
568 rural environments). Carlson and Boland (1978) suggested thermal inertia to be related
569 proportionally to an increase in the time lag between solar noon and the time of maximum H
570 flux and T_s , whereas Carlson et al. (1991) admitted that they were unable to practically explain
571 this “hysteresis” trend. Through comprehensive sensitivity analysis studies undertaken by
572 Petropoulos et al. (2009b; 2013a-c; 2014), parameters closely associated to vegetation
573 phenology have been previously outlined to have a highly influential control on air temperature
574 magnitude and extent. Conversely, sites which show relatively stable vegetation phenology such
575 as US_Ton (wooded savannah) exhibited more accurate temperature estimates. Furthermore,
576 the air temperature of the site covered by the dead forest had greater daily fluctuation
577 compared to the stands covered by mature forest which generally had the smallest daily
578 fluctuations. However, more studies is required in this direction in categorising the dead forest
579 from mature forest, currently which is not possible in the given land cover database. A more
580 improved land cover information can provide more in turn behind the performances during the
581 validation. As SimSphere model assume a homogenous canopy layer, some discrepancies may
582 occur in the air temperature simulation, which also the case over here. Furthermore a very
583 important point to also consider in the overall interpretation of the results is that the model
584 does not account for advective conditions which might be important for instance when strong
585 winds exist. Yet, generally, air temperature at 1.3m and 50m were well represented by the
586 model with results obtained showing a significant improvement on values reported in previous
587 validation attempts (Carlson and Boland, 1978; Carlson et al. 1991).

588 All in all, SimSphere demonstrated a high capability of simulating parameters associated with
589 the Earth's energy balance. It is also apparent that the model fulfils 3 of Kramer et al.'s (2002)
590 model assessment criteria, *namely accuracy, generality and realism* (see also section 1) In
591 regards to accuracy, no significant systematic prediction errors occurred within all of the fluxes
592 analysed, with the exception of a consistent underestimation of R_g and R_{net} . Additionally,
593 simulated peak heat and water flux values were in high accordance with the *in-situ* data,
594 typically at 12:30 – 13:30 LST, with a slight lag for LE and H fluxes (13:00-14:00 LST). In terms
595 of *generality*, the model has shown high levels of generality, with acceptable simulation
596 accuracies attained in the majority of sites validated. In order to improve the models generality,
597 the inclusion of more forested environments would comprehensively assess the models
598 applicability to different land cover types, particularly heterogeneous forest stands where
599 simulation accuracy tends to be lower. Finally, *realism* in the model has been most notable in
600 the simulation of LE, H and T_{air} fluxes, where slight change in the vegetation phenology or SWC
601 was accountable for characterising the diurnal evolution of fluxes in all sites validated.

602 This study can advance our understanding on SimSphere's capability to simulate the
603 interactions between different components of our Earth system and related land surface
604 processes. As no model is perfect some discrepancies between predictions and measurements
605 will always appear. Identification of these discrepancies are most interesting, because they can
606 teach us more about causes of model uncertainties in the prediction of hydro-meteorological
607 variables, and help us to improve the model structure and performance. Some large
608 discrepancies between the simulated and observed datasets could be due to model
609 parameterisation. Apart from environmental factors, some instrumentation error in tower flux
610 indicated by the presence of many spikes (too large or too small values) measurements can also
611 affect the accuracy, even if model simulated results are in agreement with actual conditions. The
612 other possible reasons is the presence of spikes in the fluxes, observed particularly on the days
613 of low agreement, which could occurred from horizontal advection, footprint changes as well as
614 a non-stationarity of turbulent regimes (Papale et al., 2006). Unfortunately, such conditions
615 cannot be captured and replicated by SimSphere.

616 In overall, it is important to recognise that uncertainty is inevitable in any model, will never be
617 as complex as the reality it portrays. In this way the model fulfills its objective as a tool as it
618 identifies the patterns of change, expected, if not always the magnitudes, indicating its
619 usefulness in practical applications either as a stand-alone tool or in combination with remote
620 sensing as done for instance through the implementation of the "triangle" technique. On this
621 basis, validation efforts presented herein are particularly important, where ensuring that all
622 model outputs are in close coherence to the physical processes being modelled are imperative
623 to the successful development of such applications.

624 **7. CONCLUSIONS**

625 This study evaluated the ability of the SimSphere land biosphere model in predicting a number
626 of parameters characterising land surface interactions for eight sites from the global terrestrial
627 monitoring network, FLUXNET. A rigorous comparison was performed for 72 selected days in
628 year 2011. The main findings of this study are concluded as follows:

629 Overall, SimSphere estimates of instantaneous energy fluxes and air temperature showed good
630 agreement in all ecosystems evaluated, apart from a minor underestimation of R_g and R_{net} (MBE

631 = -19.48 Wm⁻² and -16.49 Wm⁻² respectively). Some ecosystems exhibited poorer simulation
632 accuracies than others, most noticeably cropland (US_lb1) and grazing pasture (Calperum);
633 whilst the woodland savannah (Howard Springs) and mulga woodland (Alice Springs)
634 ecosystems both attained the highest overall simulation accuracies. Comparisons showed a
635 good agreement between modelled and measured fluxes, especially for the days with smoothed
636 daily flux trends. Very high values of the Nash-Sutcliffe efficiency index were also reported for
637 all parameters ranging from 0.720 to 0.998, suggesting, in overall, a very good model
638 representation of the observations. Highest simulation accuracies were obtained for the open
639 woodland savannah and mulga woodland sites for most of the compared parameters.

640 The process of validating any physical model is imperative to understand its representation of
641 real world scenarios. It helps identifying any deficiencies in the models' predictive ability and
642 helps identify any possible sources of error and uncertainty associated with a model. To our
643 knowledge, very few studies, if any, have acted to specifically validate SimSphere to numerous
644 ecosystems in the USA and Australia. On this basis, with the use of the model as either a
645 standalone research or educational tool, or for its synergy with EO data, its validation is not only
646 timely, but essential. SimSphere, despite its inherent architectural limitations can be applied in
647 the future for solving various theoretical and applied tasks. There is certainly room for further
648 improvements on the model in developing it further in terms of its representation of the various
649 physical processes characterising land surface interactions. This is a promising research
650 direction on which model development efforts should be focused in the future.

651 **Acknowledgements**

652 This research was conducted with the support of the European Commission under the Marie
653 Curie Re-Integration Grant "TRANSFORM-EO" project. Authors gratefully acknowledge the
654 funding body for their financial support. Authors would also like to extend thank you to the PI's
655 from the AmeriFlux and OzFlux networks for the data provision which made this study possible.

656

657 **References**

- 658 **Abramowitz, G., Leuning, R., Clark, M., & Pitman, A.: Evaluating the performance of land surface**
659 **models. *J. Climate*, 21(21), 5468-5481, 2008.**
- 660 Akkermans, T., Thiery, W., and Van Lipzig, N. P.: The regional climate impact of a realistic future
661 deforestation scenario in the Congo Basin, *J. Climate*, 27, 2714-2734, 2014.
- 662 Alexandris, S. and P. Kerkides.: New empirical formula for hourly estimations of reference
663 evapotranspiration, *Agric. Water Manag.*, 60, 157-180, 2003.
- 664 Aubinet, M., Grelle A., Ibrom, A., Rannik, Ü., Moncrieff, J., Foken, T., Kowalski, A. S., Martin, P. H.,
665 Berbigier, P., Bernhofer, Ch., Clement, R., Elbers, J., Granier, A., Grünwald, T.,
666 Morgenstern, K., Pilegaard, K., Rebmann, C., Snijders, W., Valentini, R., and Vesala, T.:
667 Estimates of the Annual Net Carbon and Water Exchange of Forests: The EUROFLUX
668 Methodology', *Adv. Ecol. Res.* 30, 113-175, 2000.
- 669 Barr, A. G., Morgenstern, K., Black, T. A., McCaughey, J. H., and Nesic, Z.: Surface energy balance
670 closure by the eddy covariance method above three boreal forest stands and
671 implications for the measurement of the CO₂ flux, *Agric. Forest Meteor.*, 322-337, 2006.
- 672 Battrick, B., and Herland, E. A.: The changing Earth. New scientific challenges for ESA's Living
673 Planet Programme, ESA SP-1304, ESA, Publications Division, ESTC, The Netherlands,
674 2006.

675 Bellocchi, G., Rivington, M., Donatelli, M., and Matthews, K.: Validation of biophysical models:
676 issues and methodologies, A review. *Agron. Sustain. Dev.*, 30, 109-130, 2010.

677 Calperum Mallee SuperSite, Terrestrial Ecosystem Research Network: available at:
678 <http://www.tern-supersites.net.au/index.php/calperum>, 2014. last access: 26
679 November 2014.

680 Carlson, T. N.: An overview of the "triangle method" for estimating surface evapotranspiration
681 and soil moisture from satellite imagery, *Sensors*, 7, 1612–1629, 2007.

682 Carlson, T. N., and Boland, F. E.: Analysis of urban-rural canopy using a surface heat
683 flux/temperature model, *J. Appl. Meteorol.*, 17, 998-1013, 1978.

684 Carlson, T. N., Dodd, J. K., Benjamin, S. G., and Cooper, J. N.: Satellite estimation of the surface
685 energy balance, moisture availability and thermal inertia, *J. Appl. Meteorol.*, 20, 6-87,
686 1981.

687 Carlson, T. N., and Lynn, B.: The effects of plant water storage on transpiration and radiometric
688 surface temperature, *Agr. Forest Meteorol.*, 57, 171-186, 1991.

689 Castellvi, F., Martinez-Cob, A., and Perez-Coveta, O.: Estimating sensible and latent heat fluxes
690 over rice using surface renewal, *Agric. Forest Meteorol.*, 139, 164-169, 2006.

691 Chauhan, N.S., Miller, S. and Ardanuy, P.: Spaceborne soil moisture estimation at high resolution:
692 a microwave-optical/IR synergistic approach, *Int. J. Remote Sens.*, 22, 4599–4646, 2003.

693 Clapp, R. B. and Hornberger, G. M.: Empirical equations for some soil hydraulic-properties,
694 *Water Resour. Res.*, 14, 601-604, 1978.

695 Cui, X., & Graf, H. F.: Recent land cover changes on the Tibetan Plateau: a review, *Climatic
696 Change*, 94, 47-61, 2009.

697 Culf, A.D., Folken, T. and J.H.C. Gash.: The energy balance closure problem, In: *Vegetation, Water,
698 Humans and the Climate*, Berlin:Springer-Verlag, 2002.

699 Coudert, B., Ottlé, C., and Briottet, X.: Monitoring land surface processes with thermal infrared
700 data: Calibration of SVAT parameters based on the optimisation of diurnal surface
701 temperature cycling features, *Remote Sens. Environ.*, 112, 872-887, 2008.

702 Deardorff, J. W.: Efficient prediction of ground surface temperature and moisture, with inclusion
703 of a layer of vegetation, *J. Geophys. Res-Oceans*, 83, 1889-1903, 1978.

704 Dickinson, R. E., and Henderson-Sellers, A.: Modelling tropical deforestation: A study of GCM
705 land-surface parametrizations, *Q. J. Roy. Meteor. Soc.*, 114, 439-462, 1988.

706 Fermi Agricultural Full Site Info, AmeriFlux: available at:
707 <http://ameriflux.ornl.gov/fullsiteinfo.php?sid=46>, 2014. last access: 26 November 2014.

708 Fermi National Accelerator Laboratory - (Agricultural site), FLUXNET: available at:
709 <http://fluxnet.ornl.gov/site/899>, 2014. last access: 26 November 2014.

710 Gillies, R. R.: A physically-based land use classification scheme using remote solar and thermal
711 infrared measurements suitable for describing urbanisation. PhD Thesis, University of
712 Newcastle, UK, 121pp, 1993.

713 Gillies, R. R., Kustas, W. P., and Humes, K. S.: A verification of the 'triangle' method for obtaining
714 surface soil water content and energy fluxes from remote measurements of the
715 Normalized Difference Vegetation Index (NDVI) and surface e, *Int. J. Remote Sens.*, 18,
716 3145-3166, 1997.

717 Granz, D., Zhang, X. and T. N. Carlson.: Observations and model simulations link stomatal
718 inhabitation to impaired hydraulic conductance following ozone exposure in cotton,
719 *Plant, Cell and Environ.*, 22, , 1999.

720 **Henderson-Sellers, A., Pitman, A. J., Love, P. K., Irannejad, P., & Chen, T. H.: The project for
721 intercomparison of land surface parameterization schemes (PILPS): Phases 2 and 3.
722 *Bulletin of the American Meteorological Society*, 76(4), 489-503, 1995.**

723 Hsu, M. H., Kuo, A. Y., Kuo, J. T., and Liu, W. C.: Procedure to calibrate and verify numerical
724 models of estuarine hydrodynamics, *J. Hydraul. Eng.*, 125, 166-182, 1999.

725 Huth, N., and Holzworth, D.: Common sense in model testing, in: Zenger A., Argent R.M. (Eds.),
726 *Proc. MODSIM 2005 International Congress on Modelling and Simulation: Advances and
727 applications for management and decision making*, 12–15 December, Melbourne,
728 Australia, 2804–2809, 2005.

729 Farouki, O.T.: The thermal properties of soils in cold regions. *Cold Regions Sci. and Tech.*, 5:67-
730 75, 1981.

731 Kramer, K., Iinonen, I., Bartelink, H., Berbigier, P., Borgnetti, M., Bernhofer, C., Cienciala, E.,
732 Dolman, A. J., Froer, O., Gracia, A., Granier, A., Grunwald, T., Hari, P., Jans, W., Kellomaki, S.,
733 Loustau, D., Magnani, F., Markkanen, T., Matteucci, G., Mohren, G. M., Moors, E., Nissenen,
734 A., Peltola, H., Sabate, S., Sanchez, A., Sontag, M. Valentini, R. and T., Vesala.: Evaluation of
735 six-process-based forest growth models using eddy-covariance measurements of CO₂
736 and H₂O fluxes at six forest sites in Europe, *Global Change Biol.*, 8, 213-230, 2002.

737 Koirala, S., Yeh, P. J. F., Hirabayashi, Y., Kanae, S., and Oki, T.: Global-scale land surface hydrologic
738 modelling with the representation of water table dynamics, *J. Geophys. Res-Atmos.*, 119,
739 75-89, 2014.

740 Liang, X., Wood, E. F., Lettenmaier, D. P., Lohmann, D., Boone, A., Chang, S., ... & Zeng, Q. C.: The
741 Project for Intercomparison of Land-surface Parameterization Schemes (PILPS) phase 2
742 (c) Red-Arkansas River basin experiment:: 2. Spatial and temporal analysis of energy
743 fluxes. *Global and planetary change*, 19(1), 137-159, 1998.

744 Liu, Y., Hiyama, T., and Yamaguchi, Y.: Scaling of land surface temperature using satellite data: A
745 case examination on ASTER and MODIS products over a heterogeneous terrain area,
746 *Rem. Sens. Envir.*, 105, 115-128, 2006.

747 Lucky Hills Shrubland Full Site Info, AmeriFlux: available at:
748 <http://ameriflux.ornl.gov/fullsiteinfo.php?sid=216>, 2014. last access: 26 November
749 2014.

750 Lynn, B. H., and Carlson, T. N.: A stomatal resistance model illustrating plant vs. external control
751 of transpiration, *Agr. Forest Meteorol.*, 52, 5-43, 1990.

752 Maayar, M., Price, D. T., Delire, C., Foley, J. A., Black, T. A., and Bessemoulin, P.: Validation of the
753 Integrated Biosphere Simulator over Canadian deciduous and coniferous boreal forest
754 stands, *J. Geophys. Res-Atmos.*, 106, 14339-14355, 2001.

755 Marshall, M., Tu, K., Funk, C., Michaelsen, J., Williams, P., Williams, C., and Kutsch, W.: Improving
756 operational land surface model canopy evapotranspiration in Africa using a direct
757 remote sensing approach, *Hydrol Earth Syst. Sc.*, 17, 1079-1091, 2013.

758 Mascart P, Taconet O, Pinty J.P. and Mehrez M., B.: Canopy resistance formulation and its effect
759 in mesoscale models: a HAPEX perspective, *Agric. For. Meteorol.*, 54, 319-351, 1991.

760 Missouri Ozark Full Site Info, AmeriFlux. available at:
761 <http://ameriflux.ornl.gov/fullsiteinfo.php?sid=64>, 2014. last access: 26 November 2014.

762 Monin, A. S., & Obukhov, A.: Basic laws of turbulent mixing in the surface layer of the
763 atmosphere, *Contrib. Geophys. Inst. Acad. Sci. USSR*, 151, 163-187, 1954.

764 Monitoring Sites - Alice Springs, OzFlux: available at:
765 <http://www.ozflux.org.au/monitoringsites/alicesprings/index.html#intro>, 2014. last
766 access: 26 November 2014.

767 Monitoring Sites - Calperum, OzFlux: available at:
768 http://www.ozflux.org.au/monitoringsites/calperum/calperum_description.html, 2014.
769 last access: 26 November 2014.

770 Monitoring Sites - Howard Springs, OzFlux: available at:
771 [http://www.ozflux.org.au/monitoringsites/howardsprings/howardsprings_description.](http://www.ozflux.org.au/monitoringsites/howardsprings/howardsprings_description.html)
772 [html](http://www.ozflux.org.au/monitoringsites/howardsprings/howardsprings_description.html), 2014. last access: 26 November 2014.

773 Nash, J., and Sutcliffe, J. V.: River flow forecasting through conceptual models part I - A
774 discussion of principles, *J. hydrol.*, 10, 282-290, 1970.

775 Olchev, A., Ibrom, A., Ross, T., Falk, U., Rakkibu, G., Radler, K., Grotea, S., Kreileina, H., and
776 Gravenhorst, G.: A modelling approach for simulation of water and carbon dioxide
777 exchange between multi-species tropical rain forest and the atmosphere, *Ecol. Model.*,
778 212, 122-130, 2008.

779 Olioso, A., T. N. Carlson and N., Brisson.: Simulation of diurnal transpiration and photosynthesis
780 of a water stressed soybean crop, *Agric. Forest Meteorol.*, 81, 41-59, 1996.

781 Olioso, A., Braud, I., Chanzy, A., Courault, D., Demarty, J., Kergoat, L., and Wigneron, J. P.: SVAT
782 modeling over the Alpilles-ReSeDA experiment: comparing SVAT models over wheat

783 fields, *Agronomie-Sciences des Productions Vegetales et de l'Environnement*, 22, 651-
784 668, 2002.

785 Papale D., Reichman, M., Aubinet, M., Canfora, E., Bernhofer, C., Kutsch, W., and Longdoz. B.:
786 Towards a standardized processing of Net Ecosystem Exchange measured with eddy
787 covariance technique: Algorithms and uncertainty estimation, *Biogeosciences*, 3, 571-
788 583, 2006.

789 Pedinotti, V.: The SWOT satellite mission: Contribution of the large swath altimetry for
790 improving the hydrological and hydrodynamic processes of a large scale model
791 (Doctoral dissertation), 2013.

792 Petropoulos, G., Carlson, T. and Wooster, M. J.: An Overview of the Use of the SimSphere Soil
793 vegetation Atmospheric Transfer (SVAT) Model for the Study of Land Atmosphere
794 Interactions, *Sensors*, 9, 4286-4308, 2009a.

795 Petropoulos, G., Wooster, M. J., Kennedy, K., Carlson, T.N. and Scholze, M.: A global sensitivity
796 analysis study of the 1d SimSphere SVAT model using the GEM SA software, *Ecol. Model.*,
797 220, 2427-2440, 2009b.

798 Petropoulos G. and Carlson, T. N.: Retrievals of turbulent heat fluxes and soil moisture content
799 by remote sensing, in: *Advances in Environmental Remote Sensing: Sensors, Algorithms,*
800 *and Applications*, Ed. Taylor and Francis, 556, 667-502, 2011.

801 Petropoulos, G. P., Griffiths, H. M., and Tarantola, S.: Towards Operational Products Development
802 from Earth Observation: Exploration of SimSphere Land Surface Process Model
803 Sensitivity using a GSA approach, 7th International Conference on Sensitivity Analysis of
804 25 Model Output, 1-4 July 2013, Nice, France, 2013a.

805 Petropoulos G., Griffiths, H. M., and Ioannou-Katidis, P.: Sensitivity exploration of SimSphere
806 land surface model towards its use for operational products development from Earth
807 observation data, chapter 14, in: *Advancement in Remote Sensing for Environmental*
808 *Applications*, edited by: Mukherjee, S., Gupta, M., Srivastava, P. K., and Islam, T., Springer,
809 2013b.

810 Petropoulos, G. P., Griffiths, H., and Tarantola, S.: Sensitivity analysis of the SimSphere SVAT
811 model in the context of EO-based operational products development, *Environ. Modell.*
812 *Softw.*, 49, 166-179, 2013c.

813 Petropoulos, G. P., Konstas, I., and Carlson, T. N: Automation of SimSphere Land Surface Model
814 Use as a Standalone Application and Integration with EO Data for Deriving Key Land
815 Surface Parameters, European Geosciences Union, 7-12 April 2013, Vienna, Austria,
816 2013d.

817 Petropoulos, G.P., Griffiths, H.M., Carlson, T.N., Ioannou-Katidis., P. and Holt, T.: SimSphere
818 Model Sensitivity Analysis Towards Establishing its Use for Deriving Key Parameters
819 Characterising Land Surface Interactions, *Geoscientific Model Development Discussions*,
820 7, 1873-1887, 2014.

821 Petropoulos G.P., H. Griffiths, W. Dorigo, A. Xaver & A. Gruber: Surface Soil Moisture Estimation:
822 Significance, Controls and Conventional Measurement Techniques Chapter 2, pages 29-
823 48, in "Remote Sensing of Energy Fluxes and Soil Moisture Content", by G.P. Petropoulos,
824 Taylor and Francis, ISBN: 978-1-4665-0578-0, 2013.

825 Piles, M., Camps, A., Vall-Llossera, M., Corbella, I., Panciera, R., Rudiger, C., Kerr, Y. H., and
826 Walker, J.: Downscaling SMOS-derived soil moisture using MODIS visible/infrared data,
827 *IEEE T. Geosci. Remote.*, 49, 3156-3166, 2011.

828 Prentice, I. C., Liang, X., Medlyn, B. E., and Wang, Y. P.: Reliable, robust and realistic: the three R's
829 of next-generation land surface modelling, *Atmos. Chem. Phys. Discussions*, 14, 24811-
830 24861, 2014.

831 Ridler, M. E., Sandholt, I., Butts, M., Lerer, S., Mougin, E., Timouk, F., Kergoat, L., and Madsen, H.:
832 Calibrating a soil-vegetation-atmosphere transfer model with remote sensing estimates
833 of surface temperature and soil surface moisture in a semi-arid environment, *J. Hydrol.*,
834 436, 1-12, 2012.

835 Rosolem, R., Gupta, H. V., Shuttleworth, W. J., Gonçalves, L. G. G., and Zeng, X.: Towards a
836 comprehensive approach to parameter estimation in land surface parameterization
837 schemes, *Hydrol. Process.*, 27, 2075-2097, 2013.

838 Ross, S. L., and Oke, T. R.: Tests of three urban energy balance models, *Bound-Lay. Meteorol.*, 44,
839 73-96, 1988.

840 Second Space for Hydrology Workshop, European Space Agency (ESA): available at:
841 <http://earth.esa.int/hydrospace07/>, 2014. last accessed: 16 December 2014.

842 Sellers, P. J., Mintz, Y. C. S. Y., Sud, Y. E. A., and Dalcher, A.: A simple biosphere model (SiB) for use
843 within general circulation models, *J. Atmos. Sci.*, 43, 505-531, 1986.

844 Sheikh, V., Visser, S. and Stroosnijder, L.: A simple model to predict soil moisture: Bridging Event
845 and Continuous Hydrological (BEACH) modelling, *Environ. Modell. Softw.*, 24, 542-556,
846 2009.

847 Slevin, D., Tett, S. F. B., & Williams, M.; Multi-site evaluation of the JULES land surface model
848 using global and local data. *Geoscientific Model Development*, 8(2), 295-316, 2015.

849 Stoyanova, J. S., and Georgiev, C. G.: SVAT modelling in support to flood risk assessment in
850 Bulgaria, *Atmos. Res.*, 123, 384-399, 2013.

851 Taconet, O., Carlson, T., Bernard, R. and Vidal-Madjar, D.: Evaluation of a surface/vegetation
852 parameterisation using satellite measurements of surface temperature, *J. Clim. Appl.*
853 *Meteorol.*, 25, 1752-1767, 1986.

854 Todhunter, P. E., and Terjung, W. H.: Intercomparison of three urban climate models, *Bound-*
855 *Lay. Meteorol.*, 42, 181-205, 1987.

856 Tonzi Ranch Full Site Info, AmeriFlux: available at:
857 <http://ameriflux.ornl.gov/fullsiteinfo.php?sid=29>, 2014. last access: 26 November 2014.

858 Twine, T. E., Kustas, W. P., Norman, J. M., Cook, D. R., Houser, P., Meyers, T. P., Prueger, J. H.,
859 Tarks, P. J., and Wesley, M. L.: Correcting eddy-covariance flux underestimates over a
860 grassland, *Agr. Forest. Meteorol.*, 103, 279-300, 2000.

861 Vaira Ranch Full Site Info, AmeriFlux: available at:
862 <http://ameriflux.ornl.gov/fullsiteinfo.php?sid=30>, 2014. last access 26 November 2014.

863 Verbeeck, H., Samson, R., Granier, A., Montpied, P. and Lemeur, R.: Multi-year model analysis of
864 GPP in a temperate beech forest in France, *Ecol. Model.*, 210, 85-109, 2008.

865 Viterbo, P., & Beljaars, A. C.: An improved land surface parameterization scheme in the ECMWF
866 model and its validation. *Journal of Climate*, 8(11), 2716-2748, 1995.

867 Wallach, D.: Evaluating crop models, in: Wallach D., Makowski D., Jones J. W. (Eds.), Working
868 with dynamic crop models, Elsevier, Amsterdam, The Netherlands, 11-53, 2006.

869 Walnut Gulch Experimental Watershed, USDA: available at:
870 <http://ars.usda.gov/PandP/docs.htm?docid=10978&page=2>, 2014. last access: 26
871 November 2014.

872 Wang, Y. P., Baldocchi, D., Leuning, R. A. Y., Falge, E. V. A., & Vesala, T.: Estimating parameters in
873 a land-surface model by applying nonlinear inversion to eddy covariance flux
874 measurements from eight Fluxnet sites. *Global Change Bio.*, 13(3), 652-670, 2007.

875 Wilson, L. G., Everett, L. G., & Cullen, S. J.: *Handbook of vadose zone characterization &*
876 *monitoring.* CRC Press, 1994.

877 Wilson, K., Carlson, T., and Bunce, J. A.: Feedback significantly influences the simulated effect of
878 CO2 on seasonal evapotranspiration from two agricultural species, *Global Change Biol.*,
879 5, 903-917, 1999.

880 Wilson, K.B. and Baldocchi, D.D.: Seasonal and inter-annual variability of energy fluxes over a
881 broadleaved temperate deciduous forest in North America, *Agric. For. Meteorol.*, 100, 1-
882 18, 2000.

883 Wilson, K., Goldstein, A., Falge, E., Aubinet, M., Baldocchi, D., Berbigier, P. and Verma, S.: Energy
884 balance closure at FLUXNET sites, *Agric. For. Meteorol.*, 113, 223-243, 2002.

885

886

887

888 **Table 1:** Summary of the main SimSphere inputs. In Parentheses are also provided the units of
 889 each of the model inputs where applicable.

890

NAME OF THE MODEL INPUT	PROCESS IN WHICH PARAMETER IS INVOLVED	MIN VALUE	MAX VALUE
Slope (<i>degrees</i>)	TIME & LOCATION	0	45
Aspect (<i>degrees</i>)	TIME & LOCATION	0	360
Station Height (<i>meters</i>)	TIME & LOCATION	0	4.92
Fractional Vegetation Cover (%)	VEGETATION	0	100
LAI (m^2m^{-2})	VEGETATION	0	10
Foliage emissivity (<i>unitless</i>)	VEGETATION	0.951	0.990
[Ca] (external [CO ₂] in the leaf) (<i>ppmv</i>)	VEGETATION	250	710
[Ci] (internal [CO ₂] in the leaf) (<i>ppmv</i>)	VEGETATION	110	400
[O ₃] (ozone concentration in the air) (<i>ppmv</i>)	VEGETATION	0.0	0.25
Vegetation height (<i>meters</i>)	VEGETATION	0.021	20.0
Leaf width (<i>meters</i>)	VEGETATION	0.012	1.0
Minimum Stomatal Resistance (sm^{-1})	PLANT	10	500
Cuticle Resistance (sm^{-1})	PLANT	200	2000
Critical leaf water potential (<i>bar</i>)	PLANT	-30	-5
Critical solar parameter (Wm^{-2})	PLANT	25	300
Stem resistance (sm^{-1})	PLANT	0.011	0.150
Surface Moisture Availability (<i>vol/vol</i>)	HYDROLOGICAL	0	1
Root Zone Moisture Availability (<i>vol/vol</i>)	HYDROLOGICAL	0	1
Substrate Max. Volum. Water Content (<i>vol/vol</i>)	HYDROLOGICAL	0.01	1
Substrate climatol. mean temperature ($^{\circ}C$)	SURFACE	20	30
Thermal inertia ($Wm^{-2}K^{-1}$)	SURFACE	3.5	30
Ground emissivity (<i>unitless</i>)	SURFACE	0.951	0.980
Atmospheric Precipitable water (<i>cm</i>)	METEOROLOGICAL	0.05	5
Surface roughness (<i>meters</i>)	METEOROLOGICAL	0.02	2.0
Obstacle height (<i>meters</i>)	METEOROLOGICAL	0.02	2.0
Fractional Cloud Cover (%)	METEOROLOGICAL	1	10
RKS (satur. thermal conduct. ($Wm^{-1} K^{-1}$) (<i>Farouki 1981</i>)	SOIL	0	10
Cosby B (unitless parameter)(<i>see Cosby et al., 1984</i>)	SOIL	2.0	12.0
THM (satur.vol. water cont.) (<i>vol/vol</i>) (<i>Cosby et al., 1984</i>)	SOIL	0.3	0.5
PSI (satur. water/matric potential) (kPa) (<i>Wilson et al., 1994</i>)	SOIL	1	7
Wind direction (<i>degrees</i>)	WIND SOUNDING PROFILE	0	360
Wind speed (<i>knots</i>)	WIND SOUNDING PROFILE	---	---
Altitude (<i>1000's feet</i>)	WIND SOUNDING PROFILE	---	---
Pressure (<i>mBar</i>)	MOISTURE SOUNDING PROFILE	---	---
Temperature (<i>Celsius</i>)	MOISTURE SOUNDING PROFILE	---	---
Temperature-Dewpoint Temperature (<i>Celsius</i>)	MOISTURE SOUNDING PROFILE	---	---

891

Table 2: Site Descriptions of chosen sites

Site Name	Site Abbreviation	Country	Geographic Location	PFT	Ecosystem Type	Dominant Species	Elevation	Climate
Alice Springs	-	Australia	-22.283/133.249	MWO	Mulga Woodland	<i>Acacia aneura</i>	606m	Desert : hot and dry summers and cold winters
Calperum	-	Australia	-34.003/140.588	PAS	Grazing Pasture	<i>Eucalyptus stricta</i>	200m	Subtropical Dry Summer
Howard Springs	-	Australia	-12.495/131.15	WSV	Woody Savannah	<i>Eucalyptus miniata</i> and <i>Eucalyptus tentrodonata</i>	64m	Tropical wet and dry: hot and humid summers
Vaira Ranch	US_VAR	USA	38.406/-120.950	GRA	Grassland	<i>Brachypodium distachyon</i> , <i>Hypochaeris glabr</i> , <i>Trifolium dubium</i>	129m	Mediterranean: hot and dry summers, wet and cold winters
Missouri Ozark	US_MOZ	USA	38.7441/-92.200	DBL	Deciduous Broadleaf	<i>Quercus alba</i> , <i>Quercus velutina</i> , <i>Carya ovata</i>	219m	Temperate continental
Fermi Agricultural	US_IB1	USA	41.8593/-88.2227	CRO	Cropland	Soybean (C3)	225m	Wet and hot summers and mild winters
Tonzi Ranch	US_TON	USA	38.4316/-120.9660	WSV	Woody Savannah	<i>Quercus douglasii</i> , <i>Pinus sabiniana</i> , <i>Brachypodium distachyon</i>	169m	Mediterranean: hot and dry summers, wet and cold winters
Lucky Hills Shrubland	US_WHS	USA	31.7438/-110.0522	SHR	Shrubland	<i>Larrea tridentate</i> , <i>Acacia constricta</i> , <i>Flourensia cernua</i>	1372m	Semi-Arid

892

893

894

Table 3: Daily simulation accuracy and average site simulation accuracy for R_g fluxes. Bias, catter, RMSD and MAE are expressed in Wm^{-2} . NASH index is unitless.

Location	Date	Bias	Scatter	RMSD	RMSD as % of Observed	MAE	NASH
Alice Springs	23/03/2011	-5.53	33.38	33.83	8.25	24.74	0.998
	15/04/2011	13.56	28.84	31.87	8.90	19.10	0.956
	23/04/2011	3.96	29.62	29.88	8.40	19.37	0.974
	10/05/2011	1.82	20.40	20.48	6.37	13.41	0.979
	24/05/2011	-16.47	25.45	30.32	10.29	20.29	0.924
	31/05/2011	-13.52	21.89	25.73	8.73	17.08	0.996
	18/06/2011	-26.93	32.75	42.40	15.37	28.03	0.949
	25/06/2011	-35.78	39.47	53.27	19.01	35.84	0.993
	18/07/2011	-34.00	33.93	48.04	16.73	34.00	1.000
	20/08/2011	-48.38	40.44	63.06	17.87	48.38	0.975
	Average		-19.48	62.36	67.825	21.20	46.29
Calperum	24/02/2011	9.68	23.06	25.01	5.85	19.077	0.994
	02/03/2011	8.41	22.63	24.14	5.71	18.314	0.979
	31/03/2011	30.48	28.25	41.56	12.30	30.482	0.986
	24/04/2011	41.93	33.67	53.78	20.58	41.932	0.975
	22/07/2011	-58.28	61.06	84.41	40.79	60.624	0.978
	28/07/2011	-67.87	71.01	98.22	46.28	70.950	0.974
	28/08/2011	-108.13	102.92	149.29	52.81	110.484	0.889
	01/12/2011	-110.33	75.49	133.69	26.40	112.586	0.899
	23/12/2011	-76.00	62.66	98.50	19.34	78.332	0.978
	29/12/2011	-74.10	62.08	96.67	18.56	76.348	0.991
	Average		-40.42	80.91	90.45	24.52	61.91
Howard Springs	18/04/2011	18.24	20.76	27.64	7.64	18.78	0.975
	23/04/2011	7.81	15.15	17.04	4.67	11.64	0.978
	13/05/2011	-0.93	20.24	20.26	5.91	15.11	0.989
	27/05/2011	24.47	29.62	38.42	12.84	25.10	0.978
	03/06/2011	-8.37	34.64	35.64	10.94	27.60	0.935
	14/06/2011	-20.95	43.62	48.39	14.86	35.50	0.974
	22/06/2011	-15.48	42.38	45.12	14.31	33.86	0.976
	22/07/2011	-37.30	56.85	67.99	21.94	48.96	0.982
	28/07/2011	-63.83	69.49	94.36	28.24	67.30	0.989
	27/09/2011	-52.80	51.87	74.01	19.51	54.04	0.979
	Average		-14.913	50.367	52.528	15.64	33.789
US_MOZ	28/06/2011	-48.13	51.40	70.42	15.12	59.86	0.976
	01/08/2011	-5.55	34.91	35.35	8.87	24.81	0.976
	18/08/2011	-2.57	35.53	35.63	8.84	27.93	0.991
	31/08/2011	42.46	42.04	59.76	17.57	42.46	0.974
	01/09/2011	34.48	30.62	46.11	13.23	34.48	0.978
	07/09/2011	4.83	41.09	41.38	10.62	30.60	0.987
	12/09/2011	16.18	33.51	37.21	10.52	24.67	0.969
	30/09/2011	29.14	34.46	45.10	14.38	29.22	0.988
	29/09/2011	42.10	34.04	54.14	23.88	42.10	0.978
	11/11/2011	48.52	44.14	65.59	33.89	48.52	0.972
	Average		16.50	47.58	50.36	14.67	36.57
US_IB1	30/05/2011	-70.94	67.44	97.88	22.57	70.94	0.939
	07/06/2011	-64.46	68.10	93.76	21.27	65.04	0.898
	28/06/2011	-69.64	69.19	98.17	20.03	72.25	0.899
	08/07/2011	-55.80	74.50	93.08	19.71	67.98	0.937
	24/08/2011	7.96	56.42	56.98	15.31	38.42	0.986
	13/09/2011	12.64	43.93	45.71	12.96	31.17	0.978
	15/09/2011	-2.54	43.42	43.50	12.71	29.90	0.940
	01/10/2011	13.80	42.18	44.38	12.00	27.31	0.977
	15/10/2011	12.39	47.00	48.61	17.53	29.42	0.949

	24/10/2011	15.15	45.93	48.365	19.38	28.51	0.997
	Average	-20.15	68.20	71.114	18.71	46.09	0.950
US_TON	27/02/2011	39.37	24.89	46.58	37.72	39.68	0.961
	17/03/2011	-88.37	74.91	115.85	37.22	88.37	0.899
	24/05/2011	-77.28	51.05	92.61	20.19	77.28	0.961
	24/06/2011	-62.15	40.59	74.23	15.30	62.15	0.965
	30/07/2011	-10.44	17.10	20.04	4.62	15.34	0.973
	07/08/2011	-19.86	27.43	33.87	7.76	24.87	0.984
	28/08/2011	-1.79	19.71	19.79	4.83	14.83	0.991
	15/09/2011	46.82	36.15	59.15	17.80	46.82	0.974
	01/11/2011	66.77	55.13	86.59	40.25	66.77	0.925
	16/11/2011	58.47	50.65	77.36	43.03	58.47	0.941
		Average	-4.85	69.54	69.71	20.59	49.46
US_WHS	08/02/2011	-119.41	122.29	170.92	35.60	119.474	0.899
	16/02/2011	-124.62	114.72	169.39	55.35	124.624	0.845
	25/03/2011	-141.67	114.86	182.38	44.63	141.666	0.880
	22/06/2011	-73.15	48.54	87.79	17.72	73.152	0.937
	13/07/2011	-77.12	63.05	99.61	20.11	78.604	0.913
	02/08/2011	-42.92	63.54	76.68	17.01	59.743	0.986
	28/08/2011	-21.54	47.97	52.59	12.80	41.999	0.983
	03/08/2011	-11.92	36.71	38.59	9.59	29.599	0.997
	05/10/2011	-1.32	35.02	35.04	10.04	24.874	0.985
	20/10/2011	11.97	27.15	29.67	9.50	18.541	0.991
		Average	-56.40	83.36	100.65	24.48	67.45
All Sites Average		-19.48	62.36	67.83	19.19	46.42	0.963

897

898

899

900

901

902

903

904

905

906

907

908

909

910

911

912

913

914

915

916

Table 4: Daily simulation accuracy and average site simulation accuracy for R_{net} fluxes. Bias, scatter, RMSD and MAE are expressed in Wm^{-2} . NASH index is unitless.

Location	Date	Bias	Scatter	RMSD	RMSD as % of Observed	MAE	NASH Index
Alice Springs	23/03/2011	-47.84	39.66	62.14	18.48	49.88	0.989
	15/04/2011	5.37	20.58	21.27	8.37	15.35	0.978
	23/04/2011	5.82	20.03	20.86	8.93	15.03	0.982
	10/05/2011	0.24	19.92	19.92	10.08	16.86	0.981
	24/05/2011	15.02	14.52	20.89	11.86	17.07	0.968
	31/05/2011	-16.37	18.30	24.55	14.32	20.45	0.991
	18/06/2011	-32.89	21.07	39.06	22.95	34.37	0.974
	25/06/2011	-40.45	18.12	44.32	27.28	40.62	0.979
	18/07/2011	-17.88	11.17	21.08	11.86	18.28	0.998
	20/08/2011	-34.57	13.29	37.04	16.38	34.57	0.964
Average	-16.35	29.69	33.90	16.23	26.25	0.980	
Calperum	24/02/2011	28.31	33.37	43.76	14.33	38.93	0.979
	02/03/2011	2.23	22.55	22.66	7.88	17.92	0.998
	31/03/2011	10.28	26.72	28.63	13.03	24.49	0.982
	24/04/2011	36.99	44.56	57.91	36.50	49.76	0.981
	22/07/2011	-62.63	39.68	74.14	69.82	62.63	0.968
	28/07/2011	-42.48	38.93	57.62	53.47	42.56	0.964
	28/08/2011	-76.72	58.52	96.49	55.19	76.72	0.945
	01/12/2011	-70.84	52.79	88.34	23.33	74.16	0.911
	23/12/2011	-18.27	33.56	38.21	10.26	26.07	0.965
	29/12/2011	-40.99	41.01	57.98	15.64	42.62	0.971
Average	-23.41	56.46	61.12	24.63	45.59	0.966	
Howard Springs	18/04/2011	22.80	32.62	39.79	13.93	32.82	0.963
	23/04/2011	17.03	30.42	34.86	11.58	28.66	0.944
	13/05/2011	40.73	28.01	49.44	21.98	40.77	0.956
	27/05/2011	54.63	44.72	70.60	38.42	56.14	0.939
	03/06/2011	20.03	27.17	33.75	17.42	25.21	0.985
	14/06/2011	16.26	33.68	37.39	19.99	29.82	0.985
	22/06/2011	10.77	39.44	40.89	22.60	29.58	0.989
	22/07/2011	-0.61	34.49	34.50	17.89	26.80	0.967
	28/07/2011	-51.75	47.36	70.15	32.05	57.36	0.995
	27/09/2011	-26.45	29.78	39.82	14.85	30.20	0.997
Average	10.35	45.89	47.05	21.03	35.74	0.972	
US_VAR	10/05/2011	-32.46	19.86	38.05	12.24	32.46	0.974
	23/06/2011	-36.76	33.67	49.85	14.69	44.40	0.987
	19/07/2011	-10.81	34.63	36.28	11.48	31.93	0.989
	30/07/2011	-2.93	49.87	49.95	17.07	43.81	0.974
	07/08/2011	4.39	40.18	40.42	14.71	32.47	0.911
	27/08/2011	40.92	61.81	74.13	32.86	68.51	0.978
	22/09/2011	43.98	65.16	78.61	49.50	72.56	0.946
	07/10/2011	-2.19	85.26	85.29	52.10	78.18	0.998
	26/11/2011	3.42	61.11	61.21	74.33	54.67	0.996
	19/12/2011	-8.42	47.35	48.09	102.45	43.57	0.996
Average	-0.09	58.64	58.64	26.52	50.26	0.975	
US_MOZ	28/06/2011	-88.46	58.74	106.19	26.25	91.19	0.957
	01/08/2011	-8.96	31.83	33.07	9.28	23.32	0.984
	18/08/2011	-29.16	31.88	43.20	13.01	38.60	0.989

	31/08/2011	-7.51	36.16	36.93	12.47	31.74	0.969
	01/09/2011	5.45	26.09	26.65	9.00	20.74	0.968
	07/09/2011	-26.40	51.75	58.09	20.09	43.98	0.964
	12/09/2011	-2.30	29.74	29.83	10.55	23.89	0.981
	30/09/2011	-17.85	46.09	49.42	22.39	37.06	0.991
	29/09/2011	33.28	35.39	48.58	34.83	33.77	0.905
	11/11/2011	54.81	64.20	84.28	87.69	56.09	0.886
	Average	-13.25	49.83	51.56	19.00	38.46	0.959
US_IB1	30/05/2011	-86.39	70.85	111.73	26.35	86.39	0.842
	07/06/2011	-35.43	40.05	53.47	14.38	37.86	0.986
	28/06/2011	-38.58	33.74	51.25	13.39	40.59	0.972
	08/07/2011	-52.02	19.96	55.72	15.01	52.02	0.976
	24/08/2011	19.23	54.20	57.51	18.55	41.64	0.946
	13/09/2011	15.26	54.05	56.16	18.53	48.64	0.977
	15/09/2011	-1.69	70.25	70.27	27.59	59.80	0.899
	01/10/2011	15.91	58.94	61.05	23.83	45.12	0.985
	15/10/2011	24.75	73.02	77.10	43.41	68.48	0.978
	24/10/2011	-28.90	73.82	79.27	51.27	71.18	0.996
Average	-16.79	67.54	69.59	23.15	55.17	0.956	
US_TON	27/02/2011	-101.40	51.67	113.80	73.72	101.40	0.911
	17/03/2011	-88.31	35.39	95.13	46.41	88.31	0.913
	24/05/2011	-70.18	38.19	79.89	21.08	70.18	0.952
	24/06/2011	-83.36	42.99	93.79	24.11	83.36	0.962
	30/07/2011	-65.26	42.12	77.67	21.57	66.65	0.986
	07/08/2011	-53.89	54.31	76.51	22.52	58.28	0.965
	28/08/2011	-39.97	57.08	69.69	22.73	58.79	0.971
	15/09/2011	2.42	38.27	38.35	16.01	30.94	0.966
	01/11/2011	26.56	47.53	54.45	51.09	46.09	0.984
	16/11/2011	12.42	48.78	50.34	52.70	48.18	0.963
	Average	-46.10	62.96	78.03	30.30	65.22	0.957
US_WHS	08/02/2011	-56.66	73.69	92.95	36.12	66.57	0.912
	16/02/2011	-71.45	65.15	96.69	61.91	75.32	0.872
	25/03/2011	-70.67	57.33	91.00	39.10	75.11	0.874
	22/06/2011	-55.39	72.62	91.33	34.65	59.76	0.929
	13/07/2011	-10.84	27.38	29.45	10.03	23.78	0.985
	02/08/2011	-15.37	36.24	39.37	11.95	30.58	0.964
	28/08/2011	5.33	26.54	27.07	10.11	18.49	0.996
	03/08/2011	-24.34	51.80	57.24	22.11	41.30	0.996
	05/10/2011	48.88	27.23	55.95	29.01	48.88	0.968
	20/10/2011	8.07	52.60	53.22	34.52	50.05	0.978
Average	-26.24	64.52	69.653	28.94	50.271	0.947	
All Sites Average		-16.49	54.44	58.69	23.81	45.90	0.964

919

920

921

922

923

924

Table 5: Daily simulation accuracy and average site simulation accuracy for LE fluxes. Bias, scatter, RMSD and MAE are expressed in Wm^{-2} . NASH index is unitless.

Location	Date	Bias	Scatter	RMSD	RMSD as % of Observed	MAE	NASH Index
Alice Springs	23/03/2011	-23.75	45.45	51.28	18.34	36.85	0.997
	15/04/2011	-17.30	23.04	28.81	23.21	19.96	0.992
	23/04/2011	2.76	23.88	24.04	30.85	14.13	0.989
	10/05/2011	20.87	19.88	28.82	87.51	21.32	0.935
	24/05/2011	4.59	4.68	6.56	36.44	5.44	0.969
	31/05/2011	5.12	8.63	10.04	51.10	6.65	0.968
	18/06/2011	-0.34	8.61	8.61	26.74	6.70	0.979
	25/06/2011	3.25	9.22	9.77	44.77	7.45	0.950
	18/07/2011	12.90	13.33	18.55	124.66	13.42	0.914
	20/08/2011	19.44	14.83	24.45	145.53	19.44	0.758
	Average	2.75	24.59	24.75	36.03	15.16	0.945
Calperum	24/02/2011	-9.77	31.40	32.89	20.08	23.06	0.995
	02/03/2011	-13.83	25.93	29.39	25.35	21.17	0.992
	31/03/2011	-8.48	18.35	20.21	22.22	13.19	0.994
	24/04/2011	-8.26	17.96	19.76	32.63	13.20	0.990
	22/07/2011	-7.97	15.53	17.45	54.76	10.97	0.979
	28/07/2011	-9.24	13.33	16.22	35.06	11.54	0.983
	28/08/2011	-17.69	24.64	30.33	63.45	19.45	0.979
	01/12/2011	-5.22	20.11	20.78	21.99	15.76	0.988
	23/12/2011	24.57	39.14	46.21	31.35	31.75	0.993
	29/12/2011	-11.57	30.29	32.43	21.10	24.78	0.993
Average	-6.75	27.20	28.02	29.41	18.49	0.989	
Howard Springs	18/04/2011	-31.86	46.21	56.13	22.11	40.76	0.997
	23/04/2011	-17.90	77.00	79.06	24.84	46.29	0.998
	13/05/2011	-5.36	23.19	23.80	14.63	17.17	0.997
	27/05/2011	35.70	44.91	57.37	71.24	39.41	0.970
	03/06/2011	26.12	37.60	45.78	74.56	29.79	0.976
	14/06/2011	7.11	16.14	17.64	30.44	12.01	0.984
	22/06/2011	31.51	35.67	47.60	52.70	36.33	0.982
	22/07/2011	13.30	29.13	32.02	30.11	20.23	0.993
	28/07/2011	-10.94	20.67	23.39	15.82	17.39	0.996
	27/09/2011	-25.35	70.48	74.90	32.73	39.03	0.965
Average	2.23	50.06	50.11	22.23	29.84	0.986	
US_VAR	10/05/2011	-9.01	13.06	15.87	12.82	12.66	0.968
	23/06/2011	29.67	38.13	48.31	76.27	31.90	0.978
	19/07/2011	23.91	29.52	37.99	193.52	25.48	0.928
	30/07/2011	27.99	31.61	42.22	357.06	29.02	0.292
	07/08/2011	22.12	25.56	33.80	354.25	22.98	0.654
	27/08/2011	24.33	29.46	38.21	532.37	24.56	0.665
	22/09/2011	17.85	21.54	27.97	403.04	17.85	0.414
	07/10/2011	6.59	27.20	27.98	43.26	19.53	0.979
	26/11/2011	-2.67	13.20	13.47	27.84	8.58	0.992
	19/12/2011	-2.61	10.60	10.91	34.99	7.21	0.985
Average	13.817	28.93	32.06	92.96	19.98	0.786	
US_MOZ	28/06/2011	-11.80	56.09	57.32	16.41	43.455	0.912
	01/08/2011	66.84	84.61	107.83	42.92	73.193	0.912

	18/08/2011	25.06	59.74	64.79	22.93	45.616	0.937
	31/08/2011	37.95	49.68	62.51	39.44	41.24	0.912
	01/09/2011	46.76	62.26	77.87	49.50	53.78	0.927
	07/09/2011	21.02	48.81	53.14	38.55	38.27	0.869
	12/09/2011	40.56	50.34	64.65	49.34	45.22	0.945
	30/09/2011	15.96	38.19	41.39	37.46	28.55	0.974
	29/09/2011	16.38	35.63	39.22	119.95	35.57	0.945
	11/11/2011	28.35	32.97	43.48	115.51	32.72	0.841
	Average	25.65	55.92	61.52	37.32	42.02	0.917
US_IB1	30/05/2011	-28.88	61.84	68.26	16.15	54.17	0.899
	07/06/2011	40.29	71.27	81.87	28.77	65.32	0.927
	28/06/2011	32.16	51.86	61.02	31.59	49.59	0.982
	08/07/2011	-35.32	28.67	45.49	17.58	35.36	0.947
	24/08/2011	1.74	37.11	37.15	9.69	31.07	0.972
	13/09/2011	-1.04	50.50	50.51	15.28	43.88	0.821
	15/09/2011	-6.30	15.45	16.68	6.14	13.25	0.998
	01/10/2011	0.80	37.23	37.24	16.76	28.78	0.964
	15/10/2011	38.31	53.74	66.00	43.70	52.64	0.979
	24/10/2011	-14.13	17.31	22.35	14.22	18.56	0.978
	Average	2.76	52.47	52.54	19.64	39.26	0.947
US_TON	27/02/2011	-5.85	22.86	23.60	31.85	17.43	0.981
	17/03/2011	-16.50	43.06	46.11	33.43	32.99	0.969
	24/05/2011	-56.28	73.75	92.78	39.70	62.52	0.899
	24/06/2011	-3.14	35.44	35.58	21.81	27.23	0.948
	30/07/2011	6.05	29.06	29.68	41.56	20.93	0.969
	07/08/2011	2.09	20.96	21.06	24.63	16.99	0.990
	28/08/2011	0.90	16.51	16.54	31.22	11.71	0.985
	15/09/2011	7.75	22.49	23.79	63.47	14.02	0.983
	01/11/2011	-2.22	14.10	14.28	20.75	11.12	0.991
	16/11/2011	4.30	10.10	10.98	30.59	7.15	0.987
	Average	-6.29	38.27	38.79	40.36	22.21	0.970
US_WHS	08/02/2011	9.61	12.40	15.69	217.20	10.35	0.886
	16/02/2011	1.03	7.80	7.87	102.72	4.61	0.946
	25/03/2011	-0.038	5.98	5.98	103.62	4.22	0.925
	22/06/2011	-2.64	6.02	6.57	63.29	4.47	0.913
	13/07/2011	-5.69	21.22	21.97	42.26	16.75	0.956
	02/08/2011	-43.53	36.74	56.96	27.02	44.83	0.975
	28/08/2011	-39.80	37.57	54.73	46.11	41.24	0.979
	03/08/2011	-12.72	15.97	20.41	25.42	15.11	0.986
	05/10/2011	-13.01	17.25	21.61	51.87	13.88	0.973
	20/10/2011	0.18	7.57	7.57	40.99	4.81	0.966
	Average	-11.49	25.52	27.99	50.61	15.36	0.951
All Sites Average		2.836	37.870	39.472	33.70	25.591	0.936

927

928

929

930

931

Table 6: Daily simulation accuracy and average site simulation accuracy for H fluxes. Bias, scatter, RMSD and MAE are expressed in Wm^{-2} . NASH index is unitless.

Location	Date	Bias	Scatter	RMSD	RMSD as % of Observed	MAE	NASH Index
Alice Springs	23/03/2011	-24.28	61.35	65.98	36.42	56.40	0.996
	15/04/2011	25.00	28.48	37.90	16.51	29.89	0.963
	23/04/2011	2.38	42.43	42.50	17.16	32.46	0.965
	10/05/2011	-24.02	64.04	68.40	28.09	53.23	0.975
	24/05/2011	9.20	27.77	29.25	12.40	24.61	0.921
	31/05/2011	-17.74	44.73	48.12	20.25	34.45	0.932
	18/06/2011	-16.03	37.98	41.22	19.41	28.27	0.983
	25/06/2011	-11.18	39.11	40.68	21.86	26.44	0.998
	18/07/2011	-7.95	28.68	29.76	12.63	22.79	0.999
	20/08/2011	-37.00	65.84	75.52	26.10	54.33	0.973
	Average	-10.16	49.35	50.39	22.57	36.29	0.970
Calperum	24/02/2011	58.73	62.79	85.97	41.06	69.62	0.981
	02/03/2011	4.58	46.74	46.96	16.77	35.21	0.963
	31/03/2011	8.70	42.43	43.31	20.97	30.60	0.899
	24/04/2011	67.41	72.42	98.93	70.00	74.96	0.991
	22/07/2011	-19.03	34.44	39.34	29.72	25.54	0.997
	28/07/2011	-1.21	32.85	32.88	30.46	25.32	0.998
	28/08/2011	-14.37	31.47	34.60	19.36	22.87	0.998
	01/12/2011	-20.74	38.84	44.02	11.19	36.18	0.986
	23/12/2011	-15.69	33.46	36.96	11.17	30.30	0.951
	29/12/2011	-12.29	38.80	40.70	12.26	32.77	0.932
	Average	5.61	54.53	54.81	23.70	38.34	0.970
Howard Springs	18/04/2011	56.78	50.31	75.86	51.67	58.88	0.995
	23/04/2011	24.08	34.73	42.26	32.73	29.46	0.996
	13/05/2011	69.81	67.25	96.93	65.29	70.17	0.995
	27/05/2011	12.17	32.14	34.36	16.66	24.12	0.973
	03/06/2011	12.11	42.25	43.95	21.14	30.03	0.963
	14/06/2011	19.13	46.53	50.31	21.14	34.01	0.932
	22/06/2011	-18.82	44.08	47.93	26.97	34.39	0.998
	22/07/2011	-9.05	26.81	28.29	15.32	19.52	0.937
	28/07/2011	-14.96	43.91	46.39	25.46	31.70	0.974
	27/09/2011	3.94	39.00	39.20	20.99	29.47	0.912
	Average	15.52	51.92	54.19	29.97	36.18	0.967
US_VAR	10/05/2011	37.64	40.41	55.22	23.14	41.20	0.889
	23/06/2011	-5.64	26.33	26.93	8.81	19.04	0.987
	19/07/2011	10.05	25.86	27.74	8.07	22.16	0.931
	30/07/2011	-7.48	31.14	32.03	9.83	23.88	0.847
	07/08/2011	11.30	24.19	26.70	8.75	21.24	0.869
	27/08/2011	29.36	37.65	47.74	19.25	37.53	0.899
	22/09/2011	34.80	28.53	45.00	24.92	38.05	0.899
	07/10/2011	29.17	25.74	38.90	25.23	30.29	0.997
	26/11/2011	28.17	32.33	42.88	67.81	30.92	0.984
	19/12/2011	13.81	18.96	23.46	40.82	19.18	0.994
	Average	13.82	33.48	38.07	17.13	28.35	0.930
US_MOZ	28/06/2011	-9.39	35.77	36.98	34.11	26.10	0.943
	01/08/2011	-34.10	58.25	67.49	50.95	44.07	0.926
	18/08/2011	19.00	35.01	39.83	23.82	29.07	0.911

	31/08/2011	-5.01	61.27	61.48	36.50	45.51	0.954
	01/09/2011	-14.39	60.86	62.54	36.40	47.65	0.938
	07/09/2011	-20.00	83.89	86.24	38.73	70.20	0.847
	12/09/2011	-1.37	45.67	45.69	25.26	36.45	0.970
	30/09/2011	-16.75	79.20	80.95	44.61	62.64	0.899
	29/09/2011	31.91	47.11	56.91	41.40	40.83	0.964
	11/11/2011	12.38	39.64	41.52	45.15	35.47	0.745
	Average	1.24	57.63	57.64	42.44	42.44	0.910
US_IB1	30/05/2011	43.82	42.74	61.21	96.12	55.53	0.912
	07/06/2011	-26.18	35.35	43.99	32.53	35.86	0.938
	28/06/2011	-21.76	24.51	32.77	13.97	26.23	0.981
	08/07/2011	27.47	13.96	30.82	26.27	27.47	0.987
	24/08/2011	66.89	39.50	77.69	74.73	67.52	0.949
	13/09/2011	40.24	33.83	52.57	86.18	43.64	0.945
	15/09/2011	44.11	35.65	56.71	99.42	44.87	0.974
	01/10/2011	70.61	49.18	86.05	60.18	70.61	0.960
	15/10/2011	20.11	36.1	41.37	37.97	31.27	0.958
	24/10/2011	36.48	24.821	44.12	120.21	36.85	0.987
	Average	30.18	46.56	55.48	68.45	43.99	0.959
US_TON	27/02/2011	-31.49	54.12	62.62	47.89	48.24	0.974
	17/03/2011	-32.30	53.99	62.91	42.14	41.69	0.949
	24/05/2011	20.70	66.34	69.49	25.01	50.30	0.891
	24/06/2011	-29.63	48.44	56.79	18.84	38.08	0.963
	30/07/2011	-26.67	65.91	71.10	21.16	49.32	0.964
	07/08/2011	-33.82	59.47	68.42	20.66	51.35	0.985
	28/08/2011	1.24	58.79	58.80	19.55	44.20	0.961
	15/09/2011	18.72	47.12	50.70	21.14	36.56	0.979
	01/11/2011	43.03	29.34	52.08	68.88	45.21	0.894
	16/11/2011	26.49	28.39	38.82	43.20	28.90	0.979
	Average	-4.37	59.77	59.93	26.84	43.39	0.954
US_WHS	08/02/2011	-18.24	59.82	62.54	21.88	47.84	0.896
	16/02/2011	-32.83	49.03	59.01	30.47	46.02	0.921
	25/03/2011	-27.28	38.85	47.47	16.42	38.03	0.973
	22/06/2011	-43.74	88.41	98.64	34.20	62.97	0.954
	13/07/2011	11.17	38.21	39.81	13.40	26.23	0.970
	02/08/2011	66.41	49.29	82.71	53.07	66.83	0.931
	28/08/2011	68.22	63.93	93.49	50.47	70.74	0.929
	03/08/2011	18.90	36.66	41.24	17.56	30.47	0.974
	05/10/2011	77.51	66.79	102.31	48.15	77.81	0.969
	20/10/2011	36.28	40.16	54.12	29.51	41.09	0.997
	Average	17.47	67.73	69.94	30.07	48.97	0.951
All Sites Average		8.66	52.62	55.06	28.40	40.14	0.951

934

935

936

937

938

939

940
941
942

Table 7: Daily simulation accuracy and average site simulation accuracy for T_{air} 1.3m. Bias, scatter, RMSD and MAE are expressed in Celsius. NASH index is unitless.

Location	Date	Bias	Scatter	RMSD	RMSD as % of Observed	MAE	NASH Index
Alice Springs	23/03/2011	-1.19	1.81	2.16	9.42	1.87	0.822
	15/04/2011	0.56	2.60	2.66	11.95	1.99	0.842
	23/04/2011	3.70	1.87	4.14	21.71	3.72	0.839
	10/05/2011	-0.09	2.75	2.75	17.22	2.52	0.871
	24/05/2011	2.97	3.48	4.58	30.80	3.06	0.850
	31/05/2011	-1.66	2.20	2.76	21.86	2.37	0.927
	18/06/2011	-0.07	2.41	2.41	17.78	2.15	0.911
	25/06/2011	-2.97	2.68	3.99	26.59	3.34	0.915
	18/07/2011	-1.25	1.92	2.29	14.21	2.08	0.911
	20/08/2011	-0.33	2.10	2.13	12.55	1.93	0.917
	Average	-0.03	3.11	3.11	18.34	2.50	0.881
Calperum	24/02/2011	-3.28	2.68	4.24	15.08	3.69	0.874
	02/03/2011	0.82	2.26	2.40	12.84	1.68	0.914
	31/03/2011	1.01	3.31	3.46	21.74	2.65	0.886
	24/04/2011	-0.45	3.47	3.50	21.99	3.21	0.903
	22/07/2011	-2.56	1.58	3.01	38.32	2.61	0.904
	28/07/2011	-3.21	2.76	4.24	30.76	3.51	0.867
	28/08/2011	-7.92	3.43	8.63	61.07	7.98	0.791
	01/12/2011	-3.30	1.50	3.63	18.09	3.30	0.785
	23/12/2011	-5.55	2.91	6.26	22.00	5.64	0.833
	29/12/2011	-4.45	1.77	4.79	18.18	4.45	0.835
	Average	-2.89	3.76	4.74	25.05	3.87	0.859
Howard Springs	18/04/2011	1.80	0.88	2.01	7.62	1.86	0.743
	23/04/2011	-0.03	0.78	0.78	2.71	0.68	0.915
	13/05/2011	0.39	1.59	1.64	7.20	1.26	0.923
	27/05/2011	2.14	2.01	2.93	12.70	2.60	0.813
	03/06/2011	2.11	1.98	2.89	12.40	2.70	0.826
	14/06/2011	1.27	2.41	2.72	14.25	2.47	0.794
	22/06/2011	-0.98	1.90	2.13	9.04	2.01	0.871
	22/07/2011	0.17	2.14	2.15	8.85	1.82	0.888
	28/07/2011	-1.38	1.74	2.22	8.61	2.08	0.851
	27/09/2011	0.07	1.10	1.10	3.88	0.95	0.910
Average	0.56	2.10	2.17	8.83	1.84	0.853	
US_VAR	10/05/2011	-3.70	2.79	4.64	25.05	3.91	0.862
	23/06/2011	1.37	2.61	2.94	11.44	1.94	0.939
	19/07/2011	-0.69	2.34	2.44	9.69	2.16	0.927
	30/07/2011	2.53	3.34	4.19	17.02	3.21	0.915
	07/08/2011	0.55	2.85	2.90	12.09	2.27	0.933
	27/08/2011	-0.79	2.80	2.90	10.38	2.63	0.926
	22/09/2011	-3.78	2.99	4.82	16.48	4.14	0.884
	07/10/2011	0.08	2.95	2.95	19.95	2.73	0.846
	26/11/2011	1.93	1.49	2.44	23.92	1.99	0.863
	19/12/2011	1.42	1.28	1.92	27.01	1.56	0.890
Average	-0.11	3.34	3.35	16.14	2.66	0.898	
US_MOZ	28/06/2011	-0.70	0.75	1.03	4.28	0.97	0.821
	01/08/2011	1.67	1.04	1.97	7.04	1.68	0.909
	18/08/2011	-0.49	1.09	1.19	4.73	1.03	0.898
	31/08/2011	-0.97	1.21	1.55	5.05	1.23	0.903
	01/09/2011	3.87	2.58	4.65	14.55	3.87	0.631
	07/09/2011	1.14	1.67	2.02	10.73	1.45	0.890

	12/09/2011	1.73	0.91	1.96	7.72	1.73	0.883
	30/09/2011	0.70	2.03	2.14	12.43	1.79	0.830
	29/09/2011	-2.59	1.31	2.90	23.41	2.65	0.844
	11/11/2011	-1.70	2.12	2.72	21.14	2.45	0.924
	Average	0.23	2.37	2.38	10.52	1.84	0.853
US_IB1	30/05/2011	1.81	1.82	2.57	9.36	1.81	0.753
	07/06/2011	0.49	1.19	1.29	4.18	1.01	0.923
	28/06/2011	3.82	2.17	4.39	19.44	3.82	0.585
	08/07/2011	0.88	3.72	3.82	14.94	3.04	0.782
	24/08/2011	4.18	1.67	4.50	17.50	4.18	0.752
	13/09/2011	8.40	4.44	9.50	32.96	8.40	0.625
	15/09/2011	2.83	2.96	4.09	23.65	2.96	0.768
	01/10/2011	2.18	0.93	2.37	24.16	2.19	0.710
	15/10/2011	4.08	1.41	4.31	34.43	4.08	0.272
	24/10/2011	0.98	2.67	2.84	25.42	2.49	0.850
	Average	3.01	3.44	4.57	21.57	3.44	0.702
US_TON	27/02/2011	-1.68	0.94	1.93	25.67	1.71	0.833
	17/03/2011	-1.68	2.13	2.71	26.02	2.33	0.837
	24/05/2011	-0.69	1.34	1.51	9.06	1.18	0.922
	24/06/2011	1.51	1.36	2.03	8.59	1.79	0.906
	30/07/2011	1.47	2.03	2.51	10.34	1.86	0.923
	07/08/2011	3.11	2.78	4.18	17.63	3.11	0.875
	28/08/2011	2.08	2.42	3.20	14.78	2.12	0.919
	15/09/2011	4.26	3.15	5.30	24.52	4.29	0.788
	01/11/2011	1.27	2.14	2.49	14.90	2.27	0.873
	16/11/2011	0.39	0.96	1.03	7.08	0.82	0.919
	Average	1.00	2.77	2.94	16.30	2.15	0.880
US_WHS	08/02/2011	-1.32	1.92	2.33	7.79	2.05	0.901
	16/02/2011	0.79	1.89	2.05	11.97	1.79	0.869
	25/03/2011	-1.21	1.45	1.89	13.17	1.50	0.924
	22/06/2011	-0.56	2.59	2.66	8.27	2.07	0.880
	13/07/2011	2.26	2.24	3.18	11.24	2.98	0.745
	02/08/2011	0.55	1.37	1.48	4.98	1.17	0.907
	28/08/2011	0.65	1.35	1.50	5.11	1.20	0.940
	03/08/2011	2.76	4.31	5.12	17.83	4.27	0.739
	05/10/2011	0.56	1.23	1.35	6.61	1.10	0.934
	20/10/2011	-0.91	2.34	2.51	11.18	2.02	0.909
	Average	0.49	2.56	2.61	10.34	1.99	0.875
All Sites Average		0.28	2.93	3.23	15.37	2.54	0.850

943
944
945
946
947
948
949
950
951
952
953

954
955
956

Table 8: Daily simulation accuracy and average site simulation accuracy for T_{air} 50m. Bias, scatter, RMSD and MAE are expressed in Celsius. NASH index is unitless.

Location	Date	Bias	Scatter	RMSD	RMSD as % of Observed	MAE	NASH Index
Alice Springs	23/03/2011	-2.14	2.23	3.09	13.45	2.55	0.758
	15/04/2011	-0.05	3.10	3.10	13.91	2.71	0.785
	23/04/2011	3.49	2.91	4.54	23.82	3.49	0.849
	10/05/2011	-1.02	3.49	3.64	22.77	3.34	0.829
	24/05/2011	1.89	4.15	4.56	30.73	3.37	0.835
	31/05/2011	-2.59	3.05	4.00	31.72	3.32	0.898
	18/06/2011	-0.87	3.14	3.26	24.08	2.92	0.880
	25/06/2011	-3.61	3.41	4.97	33.05	3.96	0.899
	18/07/2011	-2.28	2.49	3.38	20.98	2.87	0.877
	20/08/2011	-1.28	3.01	3.27	19.24	2.95	0.872
	Average	-0.84	3.74	3.84	22.65	3.15	0.848
Calperum	24/02/2011	-4.35	3.88	5.83	20.74	4.91	0.833
	02/03/2011	0.15	3.03	3.03	16.23	2.58	0.868
	31/03/2011	0.78	4.36	4.43	27.79	3.77	0.837
	24/04/2011	-1.19	4.67	4.82	30.33	4.56	0.862
	22/07/2011	-2.09	2.81	3.50	44.57	2.73	0.900
	28/07/2011	-3.91	3.27	5.10	37.00	4.14	0.843
	28/08/2011	-8.46	4.52	9.59	67.82	8.76	0.771
	01/12/2011	-4.36	2.73	5.14	25.63	4.36	0.717
	23/12/2011	-6.68	3.54	7.56	26.56	6.78	0.800
	29/12/2011	-5.29	2.57	5.88	22.32	5.31	0.803
	Average	-3.54	4.57	5.78	30.56	4.79	0.823
Howard Springs	18/04/2011	0.85	1.20	1.47	5.58	1.07	0.852
	23/04/2011	-0.70	1.46	1.62	5.61	1.37	0.828
	13/05/2011	-0.52	1.57	1.66	7.29	1.47	0.910
	27/05/2011	2.14	1.19	2.44	10.57	2.15	0.845
	03/06/2011	1.92	1.07	2.19	9.40	1.92	0.876
	14/06/2011	0.82	1.07	1.35	7.05	1.20	0.900
	22/06/2011	-1.38	1.97	2.40	10.18	2.18	0.860
	22/07/2011	-0.39	2.24	2.27	9.38	1.93	0.881
	28/07/2011	-1.90	2.01	2.76	10.69	2.33	0.833
	27/09/2011	-0.30	1.65	1.68	5.93	1.44	0.863
Average	0.05	2.04	2.04	8.30	1.71	0.865	
US_VAR	10/05/2011	-4.69	3.78	6.02	32.55	5.17	0.818
	23/06/2011	0.64	3.98	4.03	15.67	3.19	0.899
	19/07/2011	-1.89	3.44	3.93	15.60	3.46	0.884
	30/07/2011	1.58	4.43	4.70	19.12	3.55	0.906
	07/08/2011	-0.43	4.00	4.03	16.78	3.42	0.898
	27/08/2011	-1.79	4.01	4.39	15.70	4.00	0.888
	22/09/2011	-4.33	4.06	5.94	20.32	4.89	0.863
	07/10/2011	-0.80	3.62	3.71	25.06	3.45	0.805
	26/11/2011	1.66	2.41	2.92	28.69	2.45	0.831
	19/12/2011	1.16	1.89	2.22	31.27	1.88	0.867
Average	-0.89	4.24	4.34	20.92	3.55	0.866	
US_MOZ	28/06/2011	-1.44	1.26	1.91	7.97	1.772	0.674

	01/08/2011	1.38	1.69	2.18	7.79	1.677	0.910
	18/08/2011	-1.44	1.70	2.22	8.81	1.83	0.819
	31/08/2011	-1.78	1.86	2.58	8.40	2.02	0.842
	01/09/2011	3.49	3.43	4.89	15.29	3.62	0.655
	07/09/2011	0.23	2.35	2.37	12.54	2.07	0.843
	12/09/2011	1.09	1.81	2.11	8.33	1.59	0.893
	30/09/2011	0.12	2.82	2.82	16.35	2.50	0.762
	29/09/2011	-3.44	1.58	3.79	30.60	3.44	0.798
	11/11/2011	-1.96	1.75	2.63	20.50	2.14	0.934
	Average	-0.46	2.81	2.85	12.56	2.22	0.813
US_IB1	30/05/2011	1.23	2.41	2.71	9.87	1.83	0.750
	07/06/2011	0.43	2.35	2.39	7.75	2.09	0.840
	28/06/2011	3.08	3.14	4.40	19.47	3.12	0.661
	08/07/2011	-0.19	4.09	4.10	16.03	3.61	0.741
	24/08/2011	4.36	3.29	5.46	21.23	4.36	0.741
	13/09/2011	8.20	5.50	9.88	34.27	8.20	0.491
	15/09/2011	1.86	3.84	4.26	23.98	3.32	0.740
	01/10/2011	1.76	1.50	2.31	23.63	1.76	0.767
	15/10/2011	4.10	2.34	4.73	37.73	4.10	0.267
	24/10/2011	0.33	3.17	3.19	27.71	2.84	0.829
	Average	2.52	4.11	4.82	22.69	3.52	0.683
US_TON	27/02/2011	-2.08	1.44	2.53	33.73	2.08	0.797
	17/03/2011	-1.98	2.84	3.46	33.20	2.93	0.795
	24/05/2011	-1.41	2.13	2.55	15.30	2.37	0.844
	24/06/2011	0.81	2.51	2.64	11.17	1.96	0.897
	30/07/2011	0.60	3.14	3.19	13.17	2.52	0.895
	07/08/2011	2.45	4.01	4.70	19.85	3.04	0.878
	28/08/2011	1.17	3.62	3.80	17.59	2.92	0.889
	15/09/2011	3.41	4.21	5.42	25.07	3.63	0.821
	01/11/2011	0.53	2.69	2.74	16.40	2.51	0.859
	16/11/2011	-0.13	1.57	1.58	10.84	1.49	0.853
	Average	0.34	3.42	3.43	19.02	2.55	0.853
US_WHS	08/02/2011	-1.43	2.64	3.00	10.03	2.65	0.872
	16/02/2011	1.15	2.02	2.32	13.55	1.79	0.870
	25/03/2011	-1.61	2.54	3.01	21.01	2.52	0.873
	22/06/2011	-1.00	3.04	3.20	9.97	2.81	0.838
	13/07/2011	1.25	2.59	2.88	10.18	2.21	0.811
	02/08/2011	-0.37	2.15	2.18	7.34	2.01	0.841
	28/08/2011	-0.32	2.10	2.13	7.26	1.94	0.903
	03/08/2011	1.84	4.70	5.05	17.59	4.16	0.746
	05/10/2011	-0.67	2.04	2.15	10.54	1.93	0.884
	20/10/2011	-1.43	3.13	3.44	15.34	3.02	0.864
	Average	-0.19	3.03	3.04	12.03	2.51	0.850
All Sites Average		-0.376	3.496	3.766	17.90	3.00	0.825

957

958

959

960

List of Figures




Figure 1: Flowchart of the overall methodology followed




Figure 2: Scatterplot comparison of SimSphere predicted and in situ a) R_g Flux, b) R_{net} Flux, c) LE Flux, d) H Flux, e) T_{air} 1.3m, f) T_{air} 50m.




Figure 3: Simulated and observed fluxes for the Alice Spring site (Shrubland), a) illustrates the diurnal trend of the simulated fluxes from SimSphere against the observed in-situ fluxes for the 15th of April 2011 (Spring), b) illustrates the diurnal trend of the simulated fluxes from SimSphere against the observed in-situ fluxes for the 20th of August 2011 (Summer).