

Reply to the Editor’s Decision for Manuscript gmdd-8-3197-2015
“Parameterization of the snow-covered surface albedo in the Noah-MP Version
1.6 by implementing vegetation effects”

Topical Editor Decision: Reconsider after major revisions (18 Sep 2015)

Comments to the Author: Based on continued concerns of the Reviewer, I recommend further major revisions. Additionally, based on this Reviewer’s request and in my capacity as Editor I have forwarded the manuscript to Noah-MP developers Guo-Yue Niu and Zon-Liang Yang for an informal review of the results. These will not be considered formal reviews but will be considered as data points in any final decision regarding this manuscript.

⇒ The authors appreciate the Topical Editor’s effort to contact the developer of Noah-MP, Dr. Gue-Yue Niu. We have made an extra experiment with the Noah-MP, as Dr. Niu suggested, using the observed LAI and the prescribed SAI. For the observed LAI, we have used the AVHRR GIMSS LAI3g data and took an average for winter during 2001-2010 for each vegetation type within the domain of 40 – 60°N and 105 – 145°E. Since SAI is not observed, we have taken the prescribed monthly SAI value in the Noah-MP for each vegetation type. The model results are shown as the square symbols in Fig. 1. The RMS errors in albedo are reduced with more realistic LAI and SAI, but still quite significant compared to those from the new parameterization. Therefore, we suggest to use the new parameterization of the leaf and stem area which considers the structure of vegetation with a forest shading effect to improve the snow-covered surface albedo in the Noah-MP.

Based on this extra experiment, suggested by the developer of Noah-MP, we believe that our new parameterization produces the snow-covered surface albedo with much less errors than the current parameterization in the Noah-MP.




Figure 1: Comparison of RMSE values of albedo with the original minimum value of LAI and SAI (dashed lines) vs. new LI and SI (solid lines) vs. observed LAI and prescribed SAI (square symbols) for three radiation options for (a) BATS and (b) CLASS radiation schemes. OPT_RAD1 is MTSA, OPT_RAD2 is two-stream radiation option with no canopy gap, and OPT_RAD3 is the scheme that calculates the gap from vegetation fraction.

Reply to the Comments by Referee #2 for Manuscript gmdd-8-3197-2015
“Parameterization of the snow-covered surface albedo in the Noah-MP Version
1.6 by implementing vegetation effects”

General comments: (1) The presented modification gives a strong improvement in the simulation of snow-covered surface albedo by the Noah-MP model, but the results are rather specific to Noah-MP and are achieved by a rather simple tuning of vegetation parameters; I don't see anything that could be described as “an advanced conceptual frame through a clear understanding of vegetation effects on snow albedo”. (2) The authors have not fully answered my request for a “brief discussion of how Noah-MP predicts leaf and stem indices”; the new figures 3 and 4 are useful in showing that the Noah-MP dynamic vegetation option produces unrealistic results (large seasonal variations in leaf area for evergreen forests and stem area for all forest types) which will certainly lead to poor albedo simulations, but what I was getting at was why does the model behave this way? (3) Equations 1 and 2 are not informative without information on how `m_leaf` and `m_stem` are calculated. (4) Have the model developers been contacted to check that this really is the expected and intended behaviour?

⇒ The authors appreciate the comprehensive review by the referee. Below please find the item-by-item response by the authors.

- (1) We agree with the reviewer that the results here are specific to Noah-MP, but it is clearly specified in the title. Please note that the title specifically mention that this study focuses on the Noah-MP Version 1.6, which implies that the conclusions in this study are only related to the Noah-MP. Although it may look like a simple tuning, we have done lots of experiments to identify the problem in calculating the snow albedo in the Noah-MP. It is not just a simple tuning but a new parameterization that considers the vegetation structure with a forest shading effect, which results in significant improvement of the model-calculated albedo.
- (2) As the reviewer has admitted, the new Figs. 3 and 4 are useful in showing that the Noah-MP dynamic vegetation option produces unrealistic results (large seasonal variations in leaf area for evergreen forests and stem area for all forest types). Please note that the Topical Editor has contacted the developer of Noah-MP, Dr. Gue-Yue Niu, and the developer also admitted this problem:

“I basically agree with the reviewer that the model has problems in predicting LAI and SAI, especially for evergreen needleleaf trees (the modeled is not very realistic with so large seasonal variations, although in reality it has seasonal variations)”.

As he mentioned. the seasonal variation is realistic but the *large* seasonal variation is unrealistic. Although he had not provided explanation why the model behave in that way, we believe that it is due to the use of LAI and SAI which are defined as the photosynthetically active indices. Probably our community may need to study deeply on this issue, but it is not within the scope of this study.

- (3) In the following, we provide explanation on how `m_leaf` and `m_stem` are calculated:

The simulated LAI and SAI have large seasonal variations, even in the case of evergreen forests. This is because LAI and SAI are defined as the photosynthetic active indices. When

the growing index is off, the foliage photosynthesis parameter becomes zero and the leaf assimilation after removing the respiration losses also becomes zero. Thus, in winter, the leaf mass is reduced. The stem mass is calculated in the same way. The equations for calculating the leaf and stem mass are shown:

$$\begin{aligned}
ADDNPPLF &= \text{MAX}(0., LEAFPT * CARBFX - GRLEAF - RSLEAF) \\
ADDNPPST &= \text{MAX}(0., STEMPT * CARBFX - GRSTEM - RSSTEM) \\
NPPL &= \text{MAX}(ADDNPPLF, -LFDEL) \\
NPPS &= \text{MAX}(ADDNPPST, -STDEL) \\
LFMASS &= LFMASS + (NPPL - LFTOVR - DIELF) * DT \\
STMASS &= STMASS + (NPPS - STTOVR - DIEST) * DT
\end{aligned}$$

where $ADDNPPLF$ ($ADDNPPST$) is the leaf (stem) assimilation after removing the respiration losses (g/m^2), $NPPL$ ($NPPS$) is the leaf (stem) net primary productivity ($\text{g/m}^2/\text{s}$), and $LFMASS$ ($STMASS$) is the leaf (stem) mass (g/m^2). The $LEAFPT$ is fraction of carbon allocated to leaves, $STEMPT$ is fraction of carbon flux to stem, and $CARBFX$ is carbon assimilated per model step (g/m^2). The $GRLEAF$ ($GRSTEM$) is the growth respiration rate for leaf (stem) ($\text{g/m}^2/\text{s}$), $RSLEAF$ is the leaf maintenance respiration per time step (g/m^2), and $RSSTEM$ is the stem respiration (g/m^2). The $LFDEL$ ($STDEL$) is the maximum leaf (stem) mass available to change ($\text{g/m}^2/\text{s}$), $LFTOVR$ ($STTOVR$) is the leaf (stem) turnover per time step (g/m^2), $DIELF$ ($DIEST$) is the death of leaf (stem) mass per time step (g/m^2) and DT is a time step (s).

For calculating albedo, the vegetation structure is more important than the photosynthetic capacity, especially in winter. Therefore, we suggest new parameters that include non-photosynthetic leaves and stems.

- (4) The Topical Editor has contacted the model developer, Dr. Gue-Yue Niu, and received the comments as follows:

“I would like to suggest them to do an extra experiment: using observed LAI & SAI to rerun the model to see the modeled albedo. If the results are still not comparable to observations, then propose new schemes to improve albedo simulations.”

Following Dr. Niu’s suggestion, we have done an extra experiment with the Noah-MP, using the observed LAI and the prescribed SAI. For the observed LAI, we have used the AVHRR GIMSS LAI3g data and took an average for winter during 2001-2010 for each vegetation type within the domain of $40 - 60^\circ\text{N}$ and $105 - 145^\circ\text{E}$. Since SAI is not observed, we have taken the prescribed monthly SAI value in the Noah-MP for each vegetation type. The model results are shown as the square symbols in Fig. 1 (we have replaced Fig. 7 in the revised manuscript with this one). The RMS errors in albedo are reduced with more realistic LAI and SAI, but still quite significant compared to those from the new parameterization. Therefore, we suggest to use the new parameterization of the leaf and stem area which considers the structure of vegetation with a forest shading effect to improve the snow-covered surface albedo in the Noah-MP.




Figure 1: Comparison of RMSE values of albedo with the original minimum value of LAI and SAI (dashed lines) vs. new LI and SI (solid lines) vs. observed LAI and prescribed SAI (square symbols) for three radiation options for (a) BATS and (b) CLASS radiation schemes. OPT_RAD1 is MTSA, OPT_RAD2 is two-stream radiation option with no canopy gap, and OPT_RAD3 is the scheme that calculates the gap from vegetation fraction.

Parameterization of the snow-covered surface albedo in the Noah-MP Version 1.6 by implementing vegetation effects

S. Park^{1,3,4} and S. K. Park^{1,2,3,4}

¹Department of Atmospheric Science and Engineering, Ewha Womans University, Seoul, Republic of Korea

²Department of Environmental Science and Engineering, Ewha Womans University, Seoul, Republic of Korea

³Center for Climate/Environment Change Prediction Research, Ewha Womans University, Seoul, Republic of Korea

⁴Severe Storm Research Center, Ewha Womans University, Seoul, Republic of Korea

Correspondence to: S. K. Park (spark@ewha.ac.kr)

Abstract. The snow-covered surface albedo varies with many factors, including snow grain size, snow cover thickness, snow age, forest shading factor, etc., and its parameterization is still under great uncertainty. For the snow-covered surface condition, albedo of forest is typically lower than that of short vegetation; thus snow albedo is dependent on the spatial distributions of characteristic land cover and on the canopy density and structure. In the Noah land surface model with multiple physics options (Noah-MP), almost all vegetation types in East Asia during winter have the minimum values of leaf area index (LAI) and stem area index (SAI), which are too low and do not consider the vegetation types. Because LAI and SAI are represented in terms of photosynthetic activeness, stem and trunk in winter are not well represented with only these parameters. We found that such inadequate representation of the vegetation effect is mainly responsible for the large positive bias in calculating the winter surface albedo in the Noah-MP. In this study, we investigated the vegetation effect on the snow-covered surface albedo from observations and improved the model performance by implementing a new parameterization scheme. We developed new parameters, called leaf index (LI) and stem index (SI), which properly manage the effect of vegetation structure on the snow-covered surface albedo. As a result, the Noah-MP's performance in the winter surface albedo has significantly improved – the root mean square error is reduced by approximately 69 %.

1 Introduction

Snow albedo is very important in calculating the energy budget at the land surface, but the vegetation effects on adequate parameterization of the snow surface albedo are still under great uncertainty.

Vegetation influences both snow cover and albedo that can be summarized in three points. First, the canopy changes snow depth because leaves and branches can intercept part of the snow. Second, vegetation generally has a larger roughness than bare soil. Normally just a small amount of snow is sufficient to cover a bare soil, resulting in high albedo. However, the same snow amount above a grass field cannot cover all the grass because some individual elements of grass can be higher than snow depth. Thus, in case of snow, total albedo over a grass is lower than that over a bare soil. With the same amount of snow, total albedo becomes much lower over a forest. This can be the case due to litter in snow beneath trees (Hardy et al., 2000); however, it is mostly related to snow depth to cover vegetation. For example, in order to fully cover a tree 10 m high, snow should accumulate to more than 10 m. Although the tree top can be intercepted by snow with an accumulated amount lower than 10 m, the shading effect of tree through its structure still remains at non-zero solar zenith angle. Lastly, vegetation can change heat flux with different temperature from a bare soil. Moreover, vegetation changes the longwave radiation as a tree re-emits radiation downwards. For instance, when the air temperature is just below 0 °C, there is no frost and snow melts earlier under trees.

Previous studies have addressed the apparent relationship between snow cover over different vegetation types and the snow surface albedo through field measurements and satellite observations as well (Henderson-Sellers and Wilson, 1983; Jin et al., 2002; Gao et al., 2005). Gao et al. (2005) found that the maximum snow-covered albedos of non-forest types are typically higher than those of forest types, showing the shading effect of the density and vertical structure of canopy on snow cover. Forest shading is caused by leaves, stems, branches and trunks, and has a direct and quantifiable effect on albedo. Despite the spatial distribution of albedo generally follows the patterns of land cover type (Jin et al., 2002), many land surface models (LSMs) do not consider the vegetation effect on snow albedo or use unrealistic vegetation parameters; thus resulting in no significant differences in snow-covered albedo over different land surface (Essery, 2013). In numerical models, albedo under snow condition is usually parameterized through separate treatments for different surfaces (i.e., snow-covered vs. snow-free), which are weighted by the snow cover fraction. Thus, the snow cover fraction is also important for accurate calculation of albedo.

In this study, we examine how vegetation effects can be adequately considered for computation of albedo during winter in the Noah land surface model with multiple physics options (Noah-MP) (Niu et al., 2011; Yang et al., 2011). In the Noah-MP, the formula for albedo includes a sum of leaf area index (LAI) and stem area index (SAI). The details on how LAI and SAI depend on albedo are explained in Sect. 3.2.1. In most cases, the grid box values of LAI and SAI in winter are set to the same minimum values over all vegetation types (i.e., $LAI_{min} = 0.05$ and $SAI_{min} = 0.01$) and are too low compared to Asner et al. (2003). The crucial point to note is that both LAI and SAI are represented as photosynthetically active structures in the Noah-MP. Being photosynthetically active leaves and stems absent in winter, the model cannot simulate albedo accurately because nonphotosynthetic vegetation structures are not parameterized at all. In winter, these nonphotosynthetic parts

are very important for surface albedo through shadowing. Such deficient parameterizations of non-photosynthetic vegetation structures can be a major cause of the large positive bias errors of albedo in the Noah-MP. Therefore, we improved the model-calculated albedo with a new parameterization of vegetation parameters, based on an advanced conceptual frame through a clear understanding of vegetation effects on snow albedo.

2 Model and data description

2.1 The Noah-MP

The Noah-MP has evolved from the Noah land surface model and has a variety of potentials to accelerate physically-based ensemble climate prediction model runs and identification of both the optimal scheme combinations and the critical processes controlling the coupling strength (Niu et al., 2011). In this study, the model has been used in offline mode, simulating the land surface processes with atmospheric forcing. The Noah-MP has 12 different scheme sets representing various physical processes. We have selected the default options that were verified for the global river basins by Yang et al. (2011). The dynamic vegetation option is employed to assess the minimum leaf and stem area indices. For this study, we have conducted experiments using the Biosphere-Atmosphere Transfer Scheme (BATS; Dickinson et al., 1993; Yang et al., 1997) as the option of snow surface albedo. We have also used the Canadian Land Surface Scheme (CLASS; Verseghy, 1991; Verseghy et al., 1993) with new vegetation parameters for testing model performance of albedo. In terms of the albedo options in the Noah-MP, the CLASS scheme simply computes the overall snow albedo depending on fresh snow albedo and snow age while the BATS one calculates snow albedo for direct and diffuse radiation in visible and near-infrared broadband accounting for several additional parameters such as grain size growth, impurity, and especially solar zenith angle (SZA) (Niu et al., 2011).

The computational domain on which we have run the model covers $4000 \text{ km} \times 4000 \text{ km}$, with a grid size of approximately 30 km, in the East Asia region ($105^{\circ}\text{--}145^{\circ} \text{ E}$, $20^{\circ}\text{--}60^{\circ} \text{ N}$). However, most analyses have been performed over north of 40° N , where snow falls moderately. The model has run during the years 2001–2010, with a spin-up time of 6 months. Soil temperature and moisture, snow cover has been initialized by having a spin-up period of 6 months. Simulations started on 00:00 UTC 01 June.

2.2 Data sets

2.2.1 Atmospheric forcing

The atmospheric data are required to force the land surface processes in LSMs. For the Noah-MP, the Global Land Data Assimilation System (GLDAS) (Rodell et al., 2004) data have been used to drive the model during the period 2001–2010. The atmospheric forcing data set is a combination

of NOAA/GDAS atmospheric analysis field, spatially and temporally disaggregated NOAA Climate Prediction Center Merged Analysis of Precipitation (CAMP) fields, and observation-based downward shortwave and longwave radiation fields derived using the method of the Air Force Weather Agency's AGRicultural METeorological modeling system (AGRMET). The data consist of 8 forcing fields: precipitation, downward shortwave and longwave radiation, near-surface air temperature, near-surface specific humidity, near-surface zonal and meridional wind, and surface pressure. The temporal resolution is 3 h and the spatial resolution is 0.25° .

2.2.2 MODIS albedo

The MODerate-resolution Imaging Spectroradiometer (MODIS) albedo product (MCD43C3), installed on both TERRA and AQUA polar satellites, is evaluated every 16 days in a Level 3 data set, projected to a 0.05° latitude/longitude Climate Modelling Grid (CMG) (Schaaf et al., 2002). We use total shortwave broadband for white-sky albedo (bihemispherical reflectance under conditions of isotropic illumination), and quality flags that include the percentage of snow and the percentage contribution of fine resolution data. Cescatti et al. (2012) validated MODIS albedo retrievals against in situ measurements across 53 FLUXNET sites, and found a good agreement in mean yearly values between retrievals and measurements with a high correlation ($r^2 = 0.82$).

2.2.3 USGS land use and land cover

The yearly MODIS land cover and land use data within the International Geosphere–Biosphere Programme (IGBP) global vegetation classification scheme has been slightly modified to fit into the land cover classification of the U.S. Geological Survey (USGS) (Anderson et al., 1976). In this study, land cover types are grouped into 27 types according to the USGS classification, as shown in Table 1.

3 Results

3.1 Physical properties of snow-covered vegetation

For figuring out the difference of albedo among various vegetation types, we select 10 areas within $40^\circ\text{--}60^\circ\text{ N}$ and $105^\circ\text{--}145^\circ\text{ E}$ (Fig. 1) where a single type or similar two types of vegetation occupies more than 70 % in each area (Table 2). In order to minimize the effects of snow cover change, we have averaged the MODIS white-sky albedo over dominating vegetation type in each area for winter time (i.e., 273–129 as Julian day) in total shortwave broadband during 10 years (2001–2010) with a 100 % snow cover fraction (SCF) (see Fig. 2). The number of days in each area with a 100 % SCF is sufficient for comparison, as shown in Table 3. It is evident that the snow-covered albedo values are distributed over a wide range and relatively low for various forest types. The snow-covered surface albedo is different when the snow is over the ground surface vs. over the canopy, mainly due to

an uneven structure of the canopy and the forest shading effect. When the growing season is over, leaves disappear for some vegetation (e.g., deciduous trees) and remain unchanged for others (e.g., evergreen trees), while most of stems remain unchanged – reaching the minimum values. Thus, this change at the end of the growing season and the amount of stems depend on vegetation types and make distinctions of albedos over the forest types in winter.

Compared to the MODIS observation, the Noah-MP snow-covered albedos are overestimated over all vegetation types in winter and have little difference between forest and short vegetation (see Table 4). This is mainly due to the use of LAI and SAI, which are not able to quantify leaves and stems representing the forest masking in winter. In the Noah-MP, LAI and SAI are computed as follows:

$$\text{LAI} = \max(m_{\text{leaf}} \cdot \text{LAPM}, \text{LAI}_{\min}) \quad (1)$$

$$\text{SAI} = \max(m_{\text{stem}} \cdot \text{SAPM}, \text{SAI}_{\min}) \quad (2)$$

where m_{leaf} (m_{stem}) is the leaf (stem) mass per unit area (in g m^{-2}) and LAPM (SAPM) is the leaf (stem) area per unit mass (in $\text{m}^2 \text{g}^{-1}$). The subscript min implies the minimum value. The default values of LAI_{\min} and SAI_{\min} are 0.05 and $0.01 \text{ m}^2 \text{m}^{-2}$, respectively, for all vegetation types. During most of the winter period, both LAI and SAI remain equal to their minimum values (i.e., LAI_{\min} and SAI_{\min}). Tian et al. (2004) indicated that discrepancies in the winter albedos between the MODIS observation and LSMs were related to the uncertainty in quantifying LAI and SAI in the model. They also mentioned that stems would have different single-scattering albedo than green leaves, and hence it might be inadequate for the model to treat LAI and SAI the same way in the albedo parameterization.

Figure 3 shows a monthly averaged LAI, calculated from Noah-MP, during the period of 2001-2010 for deciduous broadleaf forest (Fig. 3a), deciduous needleleaf forest (Fig. 3b), and evergreen needleleaf forest and mixed forest (Fig. 3c). The red lines represent the minimum, mean and maximum LAI of the reference values from Asner et al. (2003) for each vegetation type. It is notable that the model-evaluated LAI values are much smaller than the reference values for all vegetation types in winter. In Fig. 4, the forest SAI has a seasonal cycle in the Noah-MP, but the magnitude is negligibly small.

The LAI and SAI in Figs. 3 and 4 have large seasonal variations, even in the case of evergreen needleleaf forest. Such large seasonal variations are directly related to the leaf and stem mass, represented by Eqs. (1) and (2) – see Appendix A for the detailed calculation steps. The leaf and stem mass are reduced in winter because they consider the photosynthetic capacity. However, for calculating albedo, the vegetation structure is more important than the photosynthetic capacity, especially in winter.

As previously stated, both LAI and SAI are linked to photosynthetic activeness in the Noah-MP. Compared to the reference value of LAI_{\min} (e.g., Asner et al., 2003), the model-calculated LAI_{\min} is highly underestimated for all forest types during a winter season, and SAI_{\min} is even lower than LAI_{\min} . These uncertainties are due to disregard of non-photosynthetic process in defining LAI in

160 the model. Note that the observed LAI represents all leaves regardless of photosynthetic capacity. Hence, the observed LAI is higher than the model-evaluated LAI. Actually, the structure and density of all leaves and stems have effects on albedo. Observed LAI includes all leaves, regardless of their ability to express photosynthesis; hence it is higher than the one evaluated by the model, which is related to photosynthesis. Therefore, it is necessary to properly parameterize the vegetation effects
165 on the snow-covered albedo.

3.2 Parameterization of the vegetation effects on the snow surface albedo

We have introduced new parameters in the model – leaf index (LI) and stem index (SI). LI represents a sum of LAI defined in the Noah-MP (i.e., photosynthetic leaves) and LAI of nonphotosynthetic leaves. We have substituted LI with the reference minimum values (see Asner et al., 2003) for four
170 forest types, as shown in Table 5 in Sect. 3.2.2, in order to draw realistic SI effect. SI represents a sum of SAI defined in the model (i.e., photosynthetic stems) and SAI of nonphotosynthetic stems. In defining SI, we assumed that trees were mature and focused on winter when the growing season is over; thus SI having no seasonal cycle. To figure out how albedo responses to stems, we examine the sensitivity of winter albedo to SI over forest types in the Noah-MP, and then validate albedo with
175 the optimal SI value.

3.2.1 Sensitivity of the snow-covered albedo to SI

We have focused on testing the albedo scheme in idealized cases. To avoid the effects of other parameters such as snow cover fraction and snow age, we have assumed a fresh snow with the snow cover fraction set to 1, the wetted fraction of both LI and SI to 1, and snow depth to 1 m. We have
180 imposed SI to vary from 0.1 to 10.0 with an interval of 0.1, as well as SZA 0, 30, 45, 60 and 75°. In the Noah-MP, options for two-stream radiation transfer decide the values of canopy gap probability for direct and diffuse beam. Here the canopy gap probability is defined as the chance that a photon penetrates through the vegetation without being intercepted by any crowns (Niu and Yang, 2004). The modified two-stream approximation (MTSA), which is the first option of the two-stream radiation transfer scheme, explicitly includes the three-dimensional structure of the vegetation canopy by
185 calculating the total canopy gap probability for direct beam, P_c . It equals to the sum of the between-crown gap probability, P_{bc} , which is a function of crown geometric properties and the SZA, and the within-crown gap probability, P_{wc} , which is parameterized on the basis of a modified version of Beer’s law:

$$190 \quad P_{bc} = e^{-\rho_c \pi R^2 / \cos(\theta')}$$

$$P_{wc} = (1 - P_{bc})e^{-0.5 F_a H_d / \cos \theta} \quad (4)$$

$$P_c = \min(1 - f_{veg}, P_{bc} + P_{wc}) \quad (5)$$

where ρ_t is the crown density (stems m^{-2}), R is the horizontal crown radius, θ is the solar zenith angle, $\theta' = \tan^{-1}[(b/R)\tan\theta]$, and b is the vertical crown radius. F_a is the foliage area volume density (m^{-1}) and is equal to $\text{LSAI}/(\frac{4}{3}\pi R^2 b \rho_t)$, where LSAI is the effective leaf and stem area index, through which the effect of clumping of needles into shoots is included (Chen et al., 1991; Niu and Yang, 2004). H_d is the crown depth. f_{veg} is the green vegetation fraction, and ranges from zero to 1. Therefore, if we apply new LI and SI, LSAI is changed and then the canopy gap probability is changed.

Figure 5 depicts the sensitivity of the snow-covered surface albedo and each term in albedo equation in the Noah-MP to SI, averaged over four forest types for different SZA. Total albedo by vegetation and ground, f_{re} , is evaluated as:

$$f_{\text{re}} = \begin{cases} \alpha_{\text{dc}}(1 - P_c) + \alpha_{\text{d}}P_c & \text{(for direct beam)} \\ \alpha_{\text{ic}}(1 - K_{\text{open}}) + \alpha_{\text{i}}K_{\text{open}} & \text{(for diffuse beam)} \end{cases} \quad (6)$$

where α_{d} and α_{dc} are the direct albedo of the underlying surface and canopy, respectively, and α_{i} and α_{ic} are the diffuse albedo of the underlying surface and canopy, respectively. The parameterization of canopy albedo is explained in detail in Sellers (1985). K_{open} is the between-crown gap probability for diffuse radiation, and here it has been set to 0.05. Calculation of the total albedo based on the four radiation components, shown in Fig. 5 (i.e., direct vs. diffuse albedo for visible and direct vs. diffuse albedo for near-infrared broadband), is explained in detail in Appendix AB.

As expected, total albedo over four forest types generally decreases with increasing SI because snow albedo over the vegetated surface is lower than that over the bare soil surface (Fig. 5a). At a fixed SI, albedo represents different patterns for different SZA – with increasing SZA, albedo decreases at relatively low SI while it increases at relatively high SI. Note that there is sufficient ground surface at relatively low SI that can be shaded by the vegetative canopy as SZA increases (Fig. 5b). Thus, at low SI, albedo is highest when the shadow area of underlying snow-covered surface is the smallest, that is, at local noon. Wang and Zeng (2010) also pointed out this feature using the Community Land Model 3.0. Canopy albedo also decreases with SI due to the increasing optical depth of direct and diffuse beam through leaf and stem area (Fig. 5c–f).

3.2.2 Validation of surface albedo with the optimal SI value

For quantifying the forest shading effect through SI in winter, we have compared the Noah-MP albedo with observations. We have performed model runs by repeatedly changing SI from 0.0 to 3.0 in order to find the optimal SI for each forest type that is able to reduce the bias of albedo between observation and model output. LAI and SAI were used to calculate fluxes of carbon, radiation, turbulent heat, etc., during the growing season. Hence, we have applied LI and SI when the growing season index was off. Albedo has then been averaged for 10 years in specific winter days (i.e., 337, 353, and 1, 17, and 33 as Julian day) for vegetation types in the East Asia region, in order to

compare with the MODIS 16-day observations. We have to priorly evaluate LI for drawing realistic SI effect because LI and SI are considered together for calculating albedo. Here we have assigned the observation values (Asner et al., 2003) to the LI values. Asner et al. (2003) collected more than 1,000 estimated LAI from literature and then made a global LAI dataset through a statistical analysis. The LAI dataset of forest types have been compiled from plenty of robust samples, thus they are sufficiently reliable to be used as the reference values of LI.

Figure 6 shows the bias errors of albedo between observations and model output (i.e., absolute values of model minus observation) with SI values for four forest types (i.e., deciduous broadleaf, deciduous needleleaf, evergreen needleleaf and mixed forest) and three short vegetation types (i.e., grassland, shrubland and mixed shrubland/grassland). In case of forest types, the bias errors of albedo decrease with increasing SI. On the contrary, the bias errors of albedo for short vegetation types decrease slightly or even increase with increasing SI. Since short vegetation does not have high SI in winter, modification of albedo over short vegetation by increasing SI is meaningless. The optimized SI values, effective for reduction of bias errors in albedo, are 1.3 for deciduous broadleaf forest, 1.5 for deciduous needleleaf forest, 2.3 for evergreen needleleaf forest, and 2.0 for mixed forest (see Table 5). The bias errors of albedo decrease very slowly after a certain value of SI; hence SI is considered to be optimized when the difference of bias between two consecutive SI is less than 0.005. The other land cover types are not optimized and the default values of SAI_{min} are used. The reason why the errors do not decrease below a certain level is possibly due to other parameters such as snow age, fresh snow albedo, and snow cover fraction that are not validated in the model.

The model performance with the new LI and SI has been evaluated by calculating the root mean square errors (RMSEs) of albedo ~~with observation~~ against observation (MODIS), as shown in Fig. 7. ~~The Here, the RMSEs of albedo, simulated with the observed LAI and the prescribed monthly SAI, are also depicted (square symbols). For the observed LAI, we used the AVHRR GIMSS LAI3g data and took an average for winter during 2001 – 2010 for each vegetation type within the domain of 40 – 60°N and 105 – 145°E. Since SAI is not observed, we have taken the prescribed monthly SAI value in the Noah-MP for each vegetation type. Even with the observed vegetation parameter, the RMSEs in simulated albedo are still significant. It is noteworthy that the~~ performance of the Noah-MP in calculating albedo has greatly improved with the new parameterization – RMSE has been reduced by approximately 69 %. Although we optimized SI with the MTSA, Fig. 7 shows how the parameterization affects albedo with other options as well. The simulations of albedo are improved for all two-stream radiation transfer and snow surface albedo schemes – BATS (Fig. 7a) and CLASS (Fig. 7b) with RMSEs reduced by approximately 70 % on the average. Albedo with other radiation options is also overestimated, due to underestimated LAI and SAI.

There are three radiation options in the Noah-MP for calculating energy fluxes (see Fig. 7): The first is MTSA, employed for parameterization in this study, that calculates canopy gaps from three-dimensional structure and solar zenith angle (OPT_RAD1). The second option is two-stream ra-

diation option with no canopy gap that means leaves are evenly distributed within the grid-cell with a 100 % vegetation fraction (OPT_RAD2). The last is the tile approach that computes energy fluxes in vegetated fraction and bare fraction separately and then sum them up weighted by fraction (OPT_RAD3). The optimal LI and SI obtained through the MTSA had the similar improving effect on albedo when applied to the other options. The RMSEs with the original minimum values of LAI and SAI increase until mid-winter (e.g., the 17th Julian day), and decrease after that. During the winter, albedo is dominantly influenced by the snow cover and forest masking (Bonan, 2008; Essery et al., 2009; Brovkin et al., 2013). The Noah-MP overestimates snow cover fraction and underestimates vegetation parameters (i.e., LAI and SAI) related to albedo, therefore it makes albedo be greatly overestimated. This error is significantly reduced by applying new parameters that consider all the forest structure effect with realistic values.

4 Conclusions

In winter, albedo has a large variation due to snow cover; however, in the forest region, the snow-covered albedo remains low because of two reasons. First, when the snow covers a forest canopy, the incident radiation is diffused rather than reflected due to irregular surfaces. Second, vegetation shields the snow-covered surfaces. In addition, under the forest, temperature is relatively high and snow tends to melt earlier. This effect reduces albedo further, causing more radiation to be absorbed by the ground; thus resulting in a strong positive feedback (Qu and Hall, 2007). Therefore, accurate calculation of albedo is very influential in the land surface processes. We have addressed the noticeable relationship between the vegetation types and the snow surface albedo through satellite observations. Nevertheless, in the Noah land surface model with multiple physics options (Noah-MP) as well as many land surface models, albedo was calculated without considering the vegetation effects properly. In order to apply the vegetation effect on the snow-covered albedo, we have introduced new parameters, called leaf index (LI) and stem index (SI). We focused on the SI effect because stems are more critical than leaves in the winter albedo. The performance of the Noah-MP in calculating albedo has remarkably improved with simple parameterization for all radiation options and snow surface albedo schemes. However, there is a limitation to enhancing the accuracy of albedo by changing only vegetation indices. Thus, it is required to assess the other parameters, too, such as snow cover fraction and fresh snow albedo, which are not validated against observations.

Appendix A: Calculation of the leaf and stem mass

The simulated LAI and SAI in the Noah-MP have large seasonal variations, even in the case of evergreen forests. This is because LAI and SAI are defined as the photosynthetic active indices. When the growing index is off, both the foliage photosynthesis parameter and the value of the carbon assimilated (CARBFX; g/m^2) becomes zero, then the leaf assimilation after removing the respiration

losses (ADDNPPLF; g/m^2) also becomes zero. Thus, in winter, the leaf mass is reduced. The stem mass is calculated in the same way. The equations for calculating leaf and stem mass are shown in the following:

```

ADDNPPLF = MAX(0., LEAFPT*CARBFX - GRLEAF-RSLEAF)
ADDNPPST = MAX(0., STEMPT*CARBFX - GRSTEM-RSSTEM)
NPPL = MAX(ADDNPPLF, -LFDEL)
NPPS = MAX(ADDNPPST, -STDEL)
LFMASS = LFMASS + (NPPL-LFTOVR-DIELF)*DT
STMASS = STMASS + (NPPS-STTOVR-DIEST)*DT

```

where ADDNPPST is the stem assimilation after removing the respiration losses (g/m^2), NPPL(NPPS) is the leaf (stem) net primary productivity ($\text{g/m}^2/\text{s}$), and LFMASS (STMASS) is the leaf (stem) mass (g/m^2). The LEAFPT is fraction of carbon allocated to leaves and STEMPT is fraction of carbon flux to stem. The GRLEAF (GRSTEM) is the growth respiration rate for leaf (stem) ($\text{g/m}^2/\text{s}$), RSLEAF is the leaf maintenance respiration per time step (g/m^2), and RSSTEM is the stem respiration (g/m^2). The LFDEL (STDEL) is the maximum leaf (stem) mass available to change ($\text{g/m}^2/\text{s}$), LFTOVR (STTOVR) is the leaf (stem) turnover per time step (g/m^2), DIELF (DIEST) is the death of leaf (stem) mass per time step (g/m^2) and DT is a time step (s).

Appendix B: Calculation of total albedo and weights by the radiation components

We here discuss how the radiation components in Fig. 5 are weighted to calculate the total albedo. First, the downward solar radiation (SWDOWN; W/m^2) is divided into four parts – direct visible (SOLAD (1)) and diffuse visible (SOLAI (1)) radiation, and direct near-infrared (SOLAD (2)) and diffuse near-infrared (SOLAI (2)) radiation, as shown in the following equations:

```

SOLAD (1) = SWDOWN*0.7*0.5      ! direct vis
SOLAD (2) = SWDOWN*0.7*0.5      ! direct nir
SOLAI (1) = SWDOWN*0.3*0.5      ! diffuse vis
SOLAI (2) = SWDOWN*0.3*0.5      ! diffuse nir

```

Second, four albedo components are weighted to calculate the total radiation as follows:

```

RVIS = ALBD (1)*SOLAD (1) + ALBI (1)*SOLAI (1)
RNIR = ALBD (2)*SOLAD (2) + ALBI (2)*SOLAI (2)
FSR = RVIS + RNIR

```

where ALBD (1) and ALBD (2) are albedos from direct visible bands and direct near-infrared bands, respectively, and ALBI (1) and ALBI (2) are albedos from diffuse visible bands and diffuse near-infrared bands, respectively. RVIS and RNIR are reflected radiative fluxes from visible bands and

near-infrared bands, respectively, and FSR is total reflected radiative flux. Finally, the total albedo is obtained through the formula:

$$\text{ALBEDO} = \text{FSR} / \text{SWDOWN},$$

as the ratio of total reflected radiative flux to the downward solar radiation.

335 *Acknowledgements.* This work is supported by the National Research Foundation of Korea grant (No. 2009-0083527) funded by the Korean government (MSIP). The authors thank Seungbum Hong at the National Institute of Ecology, Korea, for assistance with the Noah-MP.

References

- Anderson, J. R.: A Land Use and Land Cover Classification System for Use With Remote Sensor Data, Vol. 964,
 340 US Government Printing Office, Washington, 1976.
- Asner, G. P., Scurlock, J. M., and Hicke, J. A.: Global synthesis of leaf area index observations: implications
 for ecological and remote sensing studies, *Global Ecol. Biogeogr.*, 12, 191–205, 2003.
- Bonan, G. B.: Forests and climate change: forcings, feedbacks, and the climate benefits of forests, *Science*, 320,
 1444–1449, 2008.
- 345 Brovkin, V., Boysen, L., Raddatz, T., Gayler, V., Loew, A., and Claussen, M.: Evaluation of vegetation
 cover and land-surface albedo in MPI-ESM CMIP5 simulations, *J. Adv. Model. Earth Syst.*, 5, 48–57,
 doi:10.1029/2012MS000169, 2013.
- Cescatti, A., Marcolla, B., Vannan, S. K. S., Pan, J. Y., Roman, M. O., Yang, X., Ciais, P., Cook, R. B.,
 Law, B. E., Matteucci, G., Migliavacca, M., Moors, E., Richardson, A. D., Seufert, G., and Schaaf, C. B.: In-
 350 tercomparison of MODIS albedo retrievals and in situ measurements across the global FLUXNET network,
Remote Sens. Environ., 121, 323–334, 2012.
- Dickinson, R. E., Henderson-Sellers, A., and Kennedy, P. J.: Biosphere–Atmosphere Transfer Scheme (BATS)
 version 1e as coupled to the NCAR Community Climate Model, NCAR Tech. Note NCAR/TN-387+STR,
 Natl. Cent. for Atmos. Res., Boulder, CO, 80 pp., 1993.
- 355 Essery, R.: Large-scale simulations of snow albedo masking by forests. *Geophys. Res. Lett.*, 40, 5521–5525,
 doi:10.1002/grl.51008, 2013.
- Essery, R., Rutter, N., Pomeroy, J., Baxter, R., Stahli, M., Gustafsson, D., Barr, A., Bartlett, P., and Elder, K.:
 An evaluation of forest snow process simulations, *B. Am. Meteorol. Soc.*, 90, 1120–1135, 2009.
- Gao, F., Schaaf, C. B., Strahler, A. H., Roesch, A., Lucht, W., and Dickinson, R.: MODIS bidirectional re-
 360 flectance distribution function and albedo Climate Modeling Grid products and the variability of albedo for
 major global vegetation types, *J. Geophys. Res.*, 110, D01104, doi:10.1029/2004JD005190, 2005.
- Hardy, J. P., Melloh, R., Robinson, P., and Jordan, R.: Incorporating effects of forest litter in a snow process
 model. *Hydrol. Proc.*, 14, 3227–3237, 2000.
- Henderson-Sellers, A. and Wilson, M. F.: Surface albedo data for climatic modeling, *Rev. Geophys.*, 21, 1743–
 365 1778, doi:10.1029/RG021i008p01743, 1983.
- Jin, Y., Schaaf, C. B., Gao, F., Li, X., Strahler, A. H., Zeng, X., and Dickinson, R. E.: How does snow impact
 the albedo of vegetated land surfaces as analyzed with MODIS data?, *J. Geophys. Res.*, 29, 12.1–12.4,
 doi:10.1029/2001GL014132, 2002.
- Leonard, R. E. and Eschner, A. R.: Albedo of intercepted snow, *Water Resour. Res.*, 4, 931–935,
 370 doi:10.1029/WR004i005p00931, 1968.
- Niu, G.-Y. and Yang, W.-L.: Effects of vegetation canopy processes on snow surface energy and mass bal-
 ances, *J. Geophys. Res.*, 109, D23111, doi:10.1029/2004JD004884, 2004.
- Niu, G.-Y. and Yang, Z.-L.: An observation-based formulation of snow cover fraction and its evaluation over
 large North American river basins, *J. Geophys. Res.*, 112, D21101, doi:10.1029/2007JD008674, 2007.
- 375 Niu, G.-Y., Yang, Z.-L., Mitchell, K. E., Chen, F., Ek, M. B., Barlage, M., Kumar, A., Manning, K., Niyogi, D.,
 Rosero, E., Tewari, M., and Xia, Y.: The community Noah land surface model with multiparameterization

options (Noah-MP): 1. model description and evaluation with local-scale measurements, *J. Geophys. Res.*, 116, D12109, doi:10.1029/2010JD015139, 2011.

Norman, J. R.: *Microclimate: The Biological Environment*, John Wiley, Canada, 1983.

380 Qu, X. and Hall, A.: What controls the strength of snow-albedo feedback?, *J. Climate*, 20, 3971–3981, doi:10.1175/JCLI4186.1, 2007.

Rodell, M., Houser, P. R., Jambor, U., Gottschalck, J., Mitchell, K., Meng, C- J., Arsenault, K., Cosgrove, B., Radakovich, J., Bosilovich, M., Entin, J. K., Walker, J. P., Lohmann, D., and Toll, D.: The Global Land Data Assimilation System, *B. Am. Meteorol. Soc.*, 85, 381–394, doi:10.1175/BAMS-85-3-381, 2004.

385 Schaaf, C. B., Gao, F., Strahler, A. H., Lucht, W., Li, X., Tsang, T., Strugnell, N. C., Zhang, X., Jina, Y., Muller, J.-P., Lewis, P., Barnsley, M., Hobson, P., Disney, M., Roberts, G., Dunderdale, M., Doll, C., d’Entremont, R. P., Hu, B., Liang, S., Privette, J. L., and Roy, D.: First operational BRDF, albedo nadir reflectance products from MODIS, *Remote Sens. Environ.*, 83, 135–148, 2002.

Sellers, P. J.: Canopy reflectance, photosynthesis and transpiration, *Int. J. Remote Sens.*, 6, 1335–1372, 1985.

390 Tian, Y., Dickinson, R. E., Zhou, L., Zeng, X., Dai, Y., Myneni, R. B., Knyazikhin, Y., Zhang, X., Friedl, M., Yu, H., Wu, W., and Shaikh, M.: Comparison of seasonal and spatial variations of leaf area index and fraction of absorbed photosynthetically active radiation from Moderate Resolution Imaging Spectroradiometer (MODIS) and Common Land Model, *J. Geophys. Res.*, 109, D01103, doi:10.1029/2003JD003777, 2004.

Verseghy, D. L.: CLASS-A Canadian land surface scheme for GCMS: I. Soil model, *Int. J. Climatol.*, 11, 111–133, doi:10.1002/joc.3370110202, 1991.

395 Verseghy, D. L., McFarlane, N. A., and Lazare, M.: CLASS: A Canadian Land Surface Scheme for GCMs. II. Vegetation model and coupled runs, *Int. J. Climatol.*, 13, 347–370, 1993.

Wang, Z. and Zeng, X.: Evaluation of snow albedo in land models for weather and climate studies, *J. Clim. Appl. Meteorol.*, 49, 363–380, 2010.

400 Yang, Z.-L. and Dickinson, R. E.: Validation of the snow submodel of the Biosphere–atmosphere transfer scheme with Russian snow cover and meteorological observation data, *J. Climate*, 10, 353–373, 1997.

Yang, Z.-L., Niu, G.-Y., Mitchell, K. E., Chen, F., Ek, M. B., Barlage, M., Longuevergne, L., Manning, K., Niyogi, D., Tewari, M., and Xia, Y.: The community Noah land surface model with multiparameterization options (Noah-MP): 2. Evaluation over global river basins, *J. Geophys. Res.*, 116, D12110, doi:10.1029/2010JD015140, 2011.

405

Table 1. The USGS land cover classification.

1-Urban and Built-up Land	15-Mixed Forest
2-Dryland Cropland and Pasture	16-Water bodies
3-Irrigated Cropland and Pasture	17-Herbaceous Wetland
4-Mixed Dryland/Irrigated Cropland and Pasture	18-Wooded Wetland
5-Cropland/Grassland Mosaic	19-Barren or Sparsely Vegetated
6-Cropland/Woodland Mosaic	20-Herbaceous Tundra
7-Grassland	21-Wooded Tundra
8-Shrubland	22-Mixed Tundra
9-Mixed Shrubland/Grassland	23-Bare Ground Tundra
10-Savanna	24-Snow or Ice
11-Deciduous Broadleaf Forest	25-Playa
12-Deciduous Needleleaf Forest	26-Lava
13-Evergreen Broadleaf Forest	27-White Sand
14-Evergreen Needleleaf Forest	

Table 2. Geographic location, vegetation type and percentage of dominant vegetation type for selected areas.

Area	Longitude	Latitude	Vegetation type	Percentage
1	105.00–107.25° E	56.50–58.25° N	Mixed Forest	71.4
2	116.25–120.00° E	55.50–57.75° N	Shrubland	80.0
			Mixed Shrubland/Grassland	
3	122.50–127.75° E	57.50–60.00° N	Deciduous Needleleaf Forest	85.7
4	133.75–136.50° E	55.75–60.00° N	Deciduous Needleleaf Forest	90.4
5	138.50–140.75° E	56.25–60.00° N	Shrubland	82.6
			Mixed Shrubland/Grassland	
6	121.25–126.50° E	53.75–55.75° N	Deciduous Needleleaf Forest	93.5
7	107.75–111.50° E	49.00–51.00° N	Mixed Forest	81.7
8	113.75–117.75° E	45.00–49.00° N	Grassland	97.2
9	123.75–127.75° E	46.75–50.25° N	Dryland Cropland and Pasture	77.2
10	135.00–137.75° E	45.00–47.74° N	Mixed Forest	93.8

Table 3. Number of days with a 100 % snow cover fraction (SCF) in each area in the period of 2001–2010.

Area	1	2	3	4	5	6	7	8	9	10
No. of days	1483	6159	13235	12391	4033	8201	328	9318	7973	248

Table 4. The values of mean and standard deviation (SD) of snow albedo from the MODIS observation and the Noah-MP results for the shortwave broadband for different categories of the MODIS land cover. The data are averaged for winter time in 2001-2010 over each vegetation type within 40–60° N and 105–145° E with corresponding SD when SCF equals 100 % and other snow conditions are the same.

MODIS land cover	MODIS		Noah-MP	
	Mean	SD	Mean	SD
Dryland Cropland and Pasture	0.515	0.088	0.795	0.012
Grassland	0.595	0.080	0.787	0.020
Shrubland	0.432	0.079	0.805	0.029
Mixed Shrubland/Grassland	0.474	0.085	0.806	0.026
Deciduous Broadleaf Forest	0.428	0.076	0.748	0.075
Deciduous Needleleaf Forest	0.346	0.080	0.678	0.134
Evergreen Needleleaf Forest	0.296	0.097	0.761	0.069
Mixed Forest	0.334	0.093	0.721	0.115

Table 5. Minimum value of LAI, reference values (LI), the default minimum value of SAI, and optimized SI values for selected USGS land cover type (forest). The optimized values are based on the sensitivity test.

USGS Land Cover type	Minimum of LAI (default)	LI (reference)	Minimum of SAI (default)	SI (optimized)
11-Deciduous Broadleaf Forest	0.05	0.6	0.01	1.3
12-Deciduous Needleleaf Forest	0.05	0.5	0.01	1.5
14-Evergreen Needleleaf Forest	0.05	0.5	0.01	2.3
15-Mixed Forest	0.05	0.5	0.01	2.0




Figure 1. Geographical locations of the study domain. Each blue area has a dominant vegetation type as explained in Table 2.




Figure 2. The MODIS white-sky albedo for total shortwave broadband averaged for winter time in 2001–2010 (dots) and corresponding standard deviation (bars) when snow cover fraction equals 100 %. Descriptions on the areas are provided in Fig. 1 and Table 2.




Figure 3. Monthly averaged LAI in the Noah-MP during the period of 2001–2010 for (a) deciduous broadleaf forest (DecB), (b) deciduous needleleaf forest (DecN), (c) evergreen needleleaf forest (EverN) and mixed forest (Mix). The red lines represent the minimum, mean and maximum LAI of the reference values. In (c), the reference values are shown only for EverN.




Figure 4. Monthly averaged SAI in the Noah-MP during the period of 2001–2010 for deciduous broadleaf forest (DecB), deciduous needleleaf forest (DecN), evergreen needleleaf forest (EverN) and mixed forest (Mix).




Figure 5. Sensitivity of the snow-covered surface albedo and each term in the albedo equation in the Noah-MP to SI averaged over four forest types: **(a)** total albedo, **(b)** canopy gap probability, **(c)** direct albedo and **(d)** diffuse albedo for visible broadband, and **(e)** direct albedo and **(f)** diffuse albedo for near-infrared broadband.




Figure 6. Sensitivity of the winter-averaged albedo to SI over four forest types and three short vegetation types in the Noah-MP for 2001–2010. The optimized value for each forest type is indicated with a larger symbol.




Figure 7. Comparison of RMSE values of albedo with the original minimum value of LAI and SAI (dashed lines) vs. new LI and SI (solid lines) vs. observed LAI and prescribed SAI (square symbols) for three radiation options for (a) BATS and (b) CLASS radiation schemes. OPT_RAD1 is MTSA, OPT_RAD2 is two-stream radiation option with no canopy gap, and OPT_RAD3 is the scheme that calculate the gap from vegetation fraction.