

Plant functional type classification for Earth System Models: Results from the European Space Agency's Land Cover Climate Change Initiative

Benjamin Poulter^{1,2*}, Natasha MacBean¹, Andrew Hartley³, Iryna Khlystova⁴, Olivier Arino⁵,
Richard Betts³, Sophie Bontemps⁶, Martin Boettcher⁷, Carsten Brockmann⁷, Pierre Defourny⁶,
Stefan Hagemann⁴, Martin Herold⁸, Grit Kirches⁷, Celine Lamarche⁶, Dimitri Lederer⁶, Catherine
Ottlé¹, Marco Peters⁷ and Philippe Peylin¹

¹*Laboratoire des Sciences du Climat et de l'Environnement, LSCE-IPSL (CEA-CNRS-UVSQ), 91191
Gif-sur-Yvette, France*

²*Department of Ecology, Montana State University, Bozeman 59717, Montana, USA*

³*Met Office Hadley Centre, FitzRoy Road, Exeter, EX1 3PB, United Kingdom*

⁴*Max Planck Institute for Meteorology, Bundesstrasse 53, 20146 Hamburg, Germany*

⁵*ESA-ESRIN, 00044, Frascati, Italy*

⁶*Université catholique de Louvain, Earth and Life Institute, 1348 Louvain-la-Neuve, Belgium*

⁷*Brockmann-Consult GmbH, Max-Planck Str. 2, 21502 Geesthacht, Germany*

⁸*Laboratory of Geo-Information Science and Remote Sensing, Wageningen University,*

Droevendaalsesteeg 3, Wageningen 6708 PB, The Netherlands

*Email: benjamin.poulter@montana.edu, phone: +1 (406) 551-3969

Abstract

Global land cover is a key variable in the earth system with feedbacks on climate, biodiversity and natural resources. However, global land-cover datasets presently fall short of user needs in providing detailed spatial and thematic information that is consistently mapped over time and easily transferable to the requirements of earth system models. In 2009, the European Space Agency launched the Climate Change Initiative (CCI), with land cover (LC_CCI) as one of thirteen Essential Climate Variables targeted for research development. The LC_CCI was implemented in three phases, first responding to a survey of user needs, then developing a global, moderate resolution, land-cover dataset for three time periods, or epochs, 2000, 2005, and 2010, and the last phase resulting in a user-tool for converting land cover to plant functional type equivalents. Here we present the results of the LC_CCI project with a focus on the mapping approach used to convert the United Nations Land Cover Classification System to plant functional types (PFT). The translation was performed as part of consultative process among map producers and users and resulted in an open-source conversion tool. A comparison with existing PFT maps used by three-earth system modeling teams shows significant differences between the LC_CCI PFT dataset and those currently used in earth system models with likely consequences for modeling terrestrial biogeochemistry and land-atmosphere interactions. The main difference between the new LC_CCI product and PFT datasets used currently by three different dynamic global vegetation modeling teams is a reduction in high latitude grassland cover, a reduction in tropical tree cover, and an expansion in temperate forest cover in Europe. The LC_CCI tool is flexible for users to

Plant functional type classification

- 45 modify land cover to PFT conversions and will evolve as Phase 2 of the European Space Agency
- 46 CCI program continues.

47 **Introduction**

48 Terrestrial ecosystems are characterized by a wide variety of biomes covering arctic to tropical
49 vegetation and extending over almost 150 million square kilometers, about 30% of the earth's
50 surface (Olson et al., 2001). Land surface features associated with terrestrial ecosystems vary
51 greatly across the earth due to climate, soil and disturbance conditions. Some of these features,
52 like Leaf Area Index (LAI), surface roughness and albedo exert a strong control on the exchange
53 of biogeochemical fluxes, including carbon, water and nutrients, as well as energy fluxes between
54 vegetation and the atmosphere (Bonan, 2008). These fluxes have an influence on multiple
55 atmospheric processes that function over various temporal and spatial scales (Sellers et al.,
56 1996). Because of the importance of land-cover feedbacks on climate, a detailed and accurate
57 description of global vegetation types and their patterns is thus a key component in dynamic
58 global vegetation models (DGVM) and earth system models (ESM), with relevance for both
59 weather and climate prediction. Presently, there are several global datasets of land cover
60 available for modeling purposes, including MODIS-based land cover (Friedl et al., 2010),
61 GLC2000 (Bartholome and Belward, 2005), and GLOBCOVER (Arino et al., 2008). However, the
62 current generation of global land-cover datasets provides little consistency in terms of time
63 period of observations, spatial resolution, thematic resolution and accuracy standards. This
64 presents various challenges for earth system modeling applications that require recent and
65 consistent time series of land-cover and particular thematic information regarding land-cover
66 categories (Giri et al., 2005; Herold et al., 2008; Neumann et al., 2007; Poulter et al., 2011;
67 Wullschleger et al., 2014).

68

Plant functional type classification

69 To address these challenges, the European Space Agency established the Land Cover component
70 of the Climate Change Initiative (LC_CCI) and surveyed the land-surface modeling community to
71 define user requirements for developing a new global land-cover dataset (Bontemps et al., 2012;
72 Herold et al., 2011; Hollmann et al., 2013). The LC_CCI addressed these data needs by
73 implementing an improved approach for mapping moderate-resolution global land cover
74 consistently through time using surface-reflectance from the MERIS and VEGETATION 1 and 2
75 sensors aboard ENVISAT and SPOT 4 and 5, respectively. The final LC_CCI product resulted in the
76 development of three global land-cover datasets, one for each of three epochs (1998-2002, 2003-
77 2007 and 2008-2012) using a spectral classification approach derived from that of GLOBCOVER
78 (Arino et al., 2008), yet with improved algorithms (Radoux et al., 2014). More importantly, its
79 implementation to multi-year and multi-sensor time series ensured temporal consistency across
80 epochs (Bontemps et al., 2012). The LC_CCI land-cover maps depict the permanent features of
81 the land surface by providing information on land-cover classes defined by the United Nations
82 Land Cover Classification System (UNLCCS). It also delivers land surface seasonality products in
83 response to the needs of the ESM and DGVM communities for dynamic information about land-
84 surface processes (Bontemps et al., 2012). Land surface seasonality products provide for each
85 pixel the climatology describing, on a weekly basis, seasonal dynamics of snow cover, vegetation
86 “greenness” based on the normalized difference vegetation index and burned area. Of particular
87 relevance to the needs of the ESM modeling community, the LC_CCI developed a framework to
88 convert the categorical land-cover classes to the fractional area of plant functional types,
89 available at various spatial scales relevant to the respective ESMs.

90

Plant functional type classification

91 Plant functional types, or PFTs, are a key feature of current generation ESMS and represent
92 groupings of plant species that share similar structural, phenological, and physiological traits,
93 and can be further distinguished by climate zone (Bonan et al., 2002). Typically, 5-15 PFTs are
94 included in an earth system model simulation (Table 1), including natural and managed grasses
95 with either C3 or C4 photosynthetic pathways, broadleaf or needleleaf trees with deciduous,
96 evergreen or 'raingreen' phenology, and shrubs (Alton, 2011; Krinner et al., 2005; Sitch et al.,
97 2003). The PFT concept was originally proposed as a non-phylogenetic classification system
98 partly to reduce computational complexity of ESMS but also to maintain a feasible framework for
99 hypothesis testing. For example, interpreting the outcome of interactions for 5-15 PFTs
100 following a model simulation is much more tractable than interpreting interactions among the
101 thousands of plant species found throughout the world. The PFT concept also provides a
102 practical solution to the problem that many of the plant traits required to parameterize a model
103 at a species level are difficult to obtain (Ustin and Gamon, 2010). Second generation DGVMs are
104 currently addressing some of the limitations posed by the PFT concept as plant trait data become
105 more widely available (Kattge et al., 2011), as model structure becomes more computationally
106 efficient (Fisher et al., 2010), or as modeling concepts move toward adaptive trait rather than
107 'fixed' values (Pavlick et al., 2013; Scheiter and Higgins, 2009).

108
109 This paper describes the LC_CCI land-cover classification and presents a conversion scheme that
110 'cross-walks' the categorical UNLCCS land-cover classes to their PFT fractional equivalent. This
111 work is one of several LC_CCI publications that have previously described the need for consistent
112 land-cover mapping (Bontemps et al., 2012), the user-requirements (Tsendbazar et al., 2014),
113 and the processing of remote sensing data (Radoux et al., 2014). Land-cover to PFT conversion is

Plant functional type classification

114 a complex task and until the mapping of plant functional traits at global scale becomes possible
115 (i.e., via 'optical types', Ustin and Gamon, 2010), the cross-walking approach remains a viable
116 alternative for generating vegetation requirements for ESM and DGVM modeling approaches
117 (Bonan et al., 2002; Faroux et al., 2013; Gotangco Castillo et al., 2013; Jung et al., 2006; Lawrence
118 et al., 2011; Lawrence and Chase, 2007; Poulter et al., 2011; Verant et al., 2004; Wullschleger et
119 al., 2014). The LC_CCI conversion scheme described here provides users with a transparent
120 methodology as well as the flexibility to modify the cross-walking approach to fit the needs of
121 their study region. The conversion scheme has been derived as part of a consultative process
122 among experts involved in deriving the land cover map data and three ESM modeling groups as
123 part of Phase 1 of the project. With consensus for the thematic translation scheme, a conversion
124 tool has been designed to spatially resample PFT fractions to various model grid formats
125 common to the climate modeling community. The cross-walking table is expected to be
126 periodically updated by the LC_CCI team, i.e., Phase 2 of LC_CCI began in 2014, and will be
127 revised to include modifications and improvements related to the classification scheme and
128 mapping procedure.

129

130 **Methods**

131 *LC_CCI Land Cover Mapping Scheme*

132 The LC_CCI combined spectral data from 300-m full and 1000-m reduced resolution MERIS
133 surface reflectance (and SPOT-VEGETATION for the pre-MERIS era) to classify land cover into 22
134 Level 1 classes and 14 Level 2 sub-classes following the UN LCCS legend (Di Gregorio and Jansen,
135 2000). The whole archive of full and reduced resolution MERIS data, 2003-2012, was first pre-
136 processed in a series of steps that include radiometric and geometric corrections, cloud

Plant functional type classification

137 screening and atmospheric correction with aerosol retrieval before being merged to 7-day
138 composites. An automated classification process, combining supervised and unsupervised
139 algorithms, was then applied to the full time series to serve as a baseline to derive land-cover
140 maps that were representative of three 5-year periods, referred to as epochs, for 2000 (1998-
141 2002), 2005 (2003-2007) and 2010 (2008-2012). The classification process was achieved
142 through back- and up-dating methods using the full resolution SPOT-VEGETATION and MERIS
143 time series. The three global land-cover maps described all the terrestrial areas by 22 land cover
144 classes explicitly defined by a set of classifiers according to the UNLCCS, each classifier referring
145 to vegetation life form, leaf type and leaf longevity, flooding regime, non-vegetated cover types
146 and artificiality. Inland open water bodies and coastlines were mapped using Wide Swath Mode,
147 Image Mode at Medium-resolution (150 m) and Global Monitoring Image Mode (1 km) acquired
148 by the Advanced Synthetic Aperture Radar (ASAR) sensor aboard ENVISAT satellite for a single
149 period (2005-2010).

150

151 In addition to the land cover classification, the land surface seasonality products describe, for 1
152 km² rather than 300 meter resolution, the average behavior and the inter-annual variability of
153 the seasonal normalized difference vegetation index (NDVI), the burned area, and the snow
154 occurrence, computed over the 1998-2012 period. These seasonality products were spatially
155 coherent with the land cover classification and were provided at weekly intervals averaged over
156 this 15-year period and were based on existing independent products: SPOT-VEGETATION NDVI
157 daily time series, MODIS burned area (MCD64A1), and MODIS snow cover (MOD10A2). All
158 products are provided to users in NetCDF and geotiff file format referenced to Plate Carrée
159 projection using the World Geodetic System (WGS 84) and are available from

Plant functional type classification

160 <http://maps.elie.ucl.ac.be/CCI/viewer/>. Detailed descriptions of each component in the
161 processing chain can be found on the European Space Agency Land Cover Climate Change
162 Initiative web site <http://www.esa-landcover-cci.org>.

163

164 *Cross-walking land cover to PFTs*

165 The conversion of land cover classes to PFTs is a non-trivial task that is made more complicated
166 by the fact that the number and description of PFTs are not standardized across DGVMs. In the
167 past, land cover (and other) information has been used to derive PFT maps based on individual
168 model PFT descriptions. The method used to convert the land cover to PFTs has not always been
169 documented in detail for each model. The aim of the approach taken here was to develop a
170 general framework that could easily be adapted to the specific PFT description of any individual
171 model. In consultation with the three climate modeling teams engaged in the LC_CCI project,
172 Laboratoire des Sciences du Climat et de l'Environnement (LSCE), Met Office Hadley Centre
173 (MOHC) and Max Planck Institute for Meteorology (MPI), 10 PFT groups were defined based on
174 their phenology (needleleaf or broadleaf, evergreen or deciduous), physiognomy (tree, shrub, or
175 grass), and grassland management status (natural or managed). Three additional non-PFT
176 classes were added for bare soil, water and snow/ice. The cross-walking methodology is based
177 on the approach of Poulter et al. (2011) and assumes that each UNLCCS category could be split
178 into one or more PFT classes according to the LC class description at the per pixel level (Table 2).
179 For example, the 'cropland' UNLCCS land cover class was assigned as 100% managed grass,
180 whereas the UNLCCS 'tree cover, needleleaved evergreen, open (15-30%)' class was assigned to
181 30% needleleaved evergreen, 5% broadleaved deciduous shrub, 5% needleleaved evergreen
182 shrub, and 15% natural grass. Of note, wet tropical forest vegetation, mainly the UNLCCS class

Plant functional type classification

183 'tree cover, broadleaved evergreen, closed to open (>15%)', was assigned to the PFT categories
184 of 'broadleaf evergreen' tree (90%) and deciduous (5%), evergreen shrub (5%) following
185 observations that moist tropical forests tend to have indeterminate phenology rather than
186 distinct periods of onset and offset (Borchert et al., 2002; Fontes et al., 1995; Reich and Borchert,
187 1984). The derivation of Table 2 was the result of consultative process among the producers of
188 the land cover map and the three modeling groups that reached a consensus on the PFT fractions
189 for each LCCS-defined land cover class. The aim of this process was to gain a fuller
190 understanding of the methods behind, and implications of, the respective vegetation
191 classifications (LC and PFT). For example, previous LC class descriptions have included "semi-
192 deciduous" in the description of broadleaved evergreen trees, as in tropical rainforests in
193 particular, phenological strategies of certain species result in more pronounced seasonal leaf
194 dynamics. However, such subtle differences in functionality are not currently incorporated into
195 DGVMs, and tropical rainforests are considered to be 100% evergreen. Thus, in the cross-
196 walking table derived in this study, the relevant LC class was mapped only evergreen trees and
197 shrubs (see LC class 50 in Table 2). Other issues that were discussed included how different
198 vegetation types are treated within a grid cell for DGVMs and the lack of representation of over-
199 and understory canopies, which both had implications for how to deal with mosaic and open-
200 cover classes.

201 For the most part, the cross-walking approach followed the definitions of the UNLCCS classes,
202 where fixed proportions of land cover were split using a one to one rule for the respective PFT
203 categories, as described above. In cases where the UNLCCS class was defined by a large range of
204 tree cover and with no upper bound, i.e., ">15%" (Table 2) the uncertainties in this conversion
205 can be considered larger than compared with other categories. In these cases, the land cover

Plant functional type classification

206 remote sensing team of experts provided the criteria for the conversion approach, taking into
207 account their improved understanding of the constraints of DGVMs. The impact of these
208 uncertainties on the final PFT fractions, and on the simulated variables, is beyond the scope of
209 this study. Here we purely aim to properly document a new, generic method for mapping
210 between LC classes and PFT fractions that can be used for all DGVMs. However, the issue of
211 uncertainty in the cross-walking procedure is currently being investigated in Phase 2 of the
212 LC_CCI project.

213

214 *The LC_CCI conversion tool*

215 The LC_CCI land cover and seasonality products are initially downloaded in full spatial
216 resolution, i.e., 300-meter grid cells for land cover, and 1km grid cells for the seasonality
217 products, at global extent in Plate Carrée projection. In order to fulfil a range of ESM
218 requirements, the LC_CCI project team developed the LC_CCI user tool to allow users to adjust
219 parameters of the LC products in a way that is suitable to their model set-up, including modifying
220 the spatial resolution and converting the LC_CCI classes to fractional PFT area. The BEAM Earth
221 Observation Toolbox and Development Platform, designed for visualization and analysis of
222 ENVISAT products, was selected to provide the basis of the conversion software. A list of
223 resampling resolution and coordinate system options are provided in Table 3. The coordinate re-
224 projection and aggregation of the LC_CCI data uses slightly different resampling algorithms
225 depending on whether the tool is used on the land-cover or seasonality products. The tool
226 converts the original LC_CCI geotiff file to target files produced in NetCDF-4 format and following
227 CF (Climate and Forecast) conventions, more commonly used in numerical modelling. The open-
228 source BEAM tool (source code at <https://github.com/bcdev>) can be run independently using

Plant functional type classification

229 either Windows or Unix-based operating systems and the compiled operational tool can be
230 downloaded from <http://maps.elie.ucl.ac.be/CCI/viewer/download.php>.

231

232 *Re-sampling algorithm for LC_CCI land cover*

233 For the land cover classes, the resampling algorithm produces an aggregated LC_CCI dataset that
234 in addition to the fractional area of each PFT, also includes the fractional area of each LC_CCI
235 UNLCCS class, the majority (dominant) LC_CCI UNLCCS class, and the overall accuracy of the
236 aggregated classification. The majority class n is defined as the LC_CCI class which has the rank n
237 of sorted list of LC_CCI classes by fractional area in the target cell (see Figure 1). The number of
238 majority classes computed is a parameter, which can be defined by user, so that the full number
239 of LCCS classes can be reduced to a user-defined subset, i.e., the top 3. Each original, valid land,
240 water, snow or ice pixel contributes to the final target cell according to its area percentage
241 contribution. The accuracy is calculated by the median of the land cover classification probability
242 values weighted by the fractional area.

243

244 *Re-sampling algorithm for LC_CCI seasonality products*

245 The aggregation of LC_CCI seasonality products is specific for NDVI (i.e., greenness), burned
246 areas, and snow cover. In the case of the LC_CCI NDVI condition, the mean NDVI over all valid
247 NDVI observations are included in the aggregated product. The burned area and snow cover
248 LC_CCI products also contain 3 different layers: the proportion of area (in %) covered by burned
249 or snow area, the average frequency of the burned area or snow area detected over the
250 aggregated zone and the sum of all valid observations of burned or snow area. Similar to
251 aggregation rules for land-cover, each original pixel contributes to the target cell according to its

Plant functional type classification

252 area percentage but the value of a pixel will only be considered if its value falls within its valid
253 range, i.e., zero to one for NDVI.

254

255 *Extension to specific model needs*

256 The LC_CCI tool provides users with a zero-order classification, that is, the PFT classes are
257 defined as broadly as possible so that users have the advantage to continue to aggregate to the
258 requirements of their model (Figure 2). For example, models that do not include shrub PFTs can
259 merge shrub and tree categories together to create a single woody PFT category. Modeling
260 groups that require climatic distinctions for PFTs, for example, temperate versus tropical versus
261 boreal types can use their own climate or biome datasets such as Koeppen-Geiger or Trewartha
262 ecological zones (Baker et al., 2010; Kottek, 2006; Peel et al., 2007) and define classification rules
263 based on temperature thresholds, for example (Poulter et al., 2011). Most models also require a
264 distinction between the C3 and C4 photosynthetic pathways for different grass species, where C4
265 is more common in warm and dry climates (Edwards et al., 2010; Still et al., 2003). The
266 photosynthetic biochemistry of C4 grasses is very different to C3 grasses and their distribution
267 can be mapped either according to climate (Poulter et al., 2011) or to some combination of
268 remote sensing, ground-based observations and ecosystem modeling (Still et al., 2003). The
269 LC_CCI managed grassland PFT category represents all non-irrigated, irrigated and pasture lands
270 and so drawing finer thematic distinctions between these must come from country or sub-
271 country statistics similar to downscaling work made by Hurtt et al. (2006), Klein Goldewijck
272 (1997) and others (Monfreda et al., 2008; Ramankutty and Foley, 1998).

273

274 *Analysis and comparison to PFT maps*

Plant functional type classification

275 For analysis and demonstration of the tool, we compare the LC_CCI PFTs with the original PFTs
276 used by the Land Surface Model (LSM) components of the ESMs from the three modeling centers
277 developing ORCHIDEE at LSCE (Krinner et al., 2005), JULES at MOHC (Clark et al., 2011; Cox et
278 al., 2000; Pacifico et al., 2011), and JSBACH at MPI (Knorr, 2000; Pongratz et al., 2009; Reick et
279 al., 2013). The original ORCHIDEE PFT map, based on 12 PFTs plus bare soil, has its origins in the
280 Olson land cover dataset from the 1980's (Olson et al., 1983) and the International Geosphere
281 Biosphere Program (IGBP) DISCover dataset for the period 1992-93 (Loveland and Belward,
282 1997). This was implemented within ORCHIDEE using a look-up table approach to estimate PFT
283 fractions (Verant et al., 2004). The JULES model also uses PFT distributions derived from the
284 IGBP DISCover dataset to estimate fractional coverage of 5 PFTs and 4 non-vegetated surfaces
285 (water, urban, snow/ice and bare soil). JSBACH uses original data from Wilson and Henderson-
286 Sellers (1985) and continuous tree fractions from Defries (1999) to represent the distribution
287 and abundance of 12 PFTs. The LC_CCI Epoch 2010 was converted to 0.5 degree resolution using
288 the LC_CCI user tool and compared with the individual default model PFT maps to illustrate
289 regional differences and biases between products and to provide a baseline of how the LC_CCI
290 products may improve land surface model performance.

291

292 **Results**

293 *Global summary of LC_CCI*

294 The global land areas covered by the aggregated 0.5 degree LC_CCI PFT equivalents (Figure 3)
295 are dominated by barren and bare soil (39 Mkm²), followed by forests (30 Mkm²), managed
296 grasslands, croplands and pasture (25 Mkm²), natural grasslands (18 Mkm²), and shrublands (14
297 Mkm²). For comparison, the MODIS Collection 5 land cover product developed by Friedl et al.

Plant functional type classification

298 (2010), report for barren area 18 Mkm², forest and savanna at 49 Mkm², a shrubland area of 22
299 Mkm², and 12 Mkm² for croplands. With reference to the Food and Agriculture Organization
300 (FAO) statistics, forest area is reported as 38 Mkm² (FAO and JRC, 2012), cropland area as
301 approximately 15 Mkm² (Monfreda et al., 2008) and pasture lands of 28 Mkm² (Ramankutty et
302 al., 2008). While part of the areal differences are explained by the spatial resolution between the
303 moderate-resolution MODIS data (500m) in comparison to the 0.5-degree LC_CCI data, thematic
304 differences introducing uncertainty in aggregating to forest, grassland, etc. classes, and factors
305 stemming from different definitions of forest cover thresholds used to categorize forest land
306 between the UNLCCS approach (10% cover) and the IGBP (60%) approach used for MODIS. In
307 addition, the UNLCCS to PFT conversion approach considers assumptions related to plant
308 community level variability, and so a bare soil fraction is introduced during the conversion (see
309 Table 3) increasing its global area and partially explaining the difference with MODIS land cover.

310

311 *Comparison with original PFT maps*

312 Differences between the LC_CCI PFT datasets and the original PFT datasets were specific for each
313 ESM (Figure 4) largely because the original reference data were different per modeling group.
314 Another challenge was that different PFT classification schemes were used for each model (Table
315 1), introducing further aggregation uncertainties in the comparison between LC_CCI and the
316 original PFT data.

317

318 For all modeling teams, grasslands PFT distributions showed the largest changes, with
319 significant reductions in northern latitudes for ORCHIDEE and JULES (Figure 6). For ORCHIDEE,
320 the grassland PFT reductions were associated with an increase in bare soil, together with a shift

Plant functional type classification

321 from C3 grasses to (boreal) forest in the mid-to-high latitudes (Figure 5). Agricultural PFTs, not
322 included in JULES, were similar for the original ORCHIDEE and LC_CCI inputs at regional scales,
323 but showed increases in tropical regions where deforestation activities were high, e.g., the
324 Brazilian arc of deforestation region. JSBACH generally had a reduction in cropland area,
325 especially over North America and the North African arid regions.

326

327 Over arid regions, in comparison to the original PFT map, JULES decreased in C4 grasses over
328 Australia, with an associated increase in the fractional cover of shrubs and bare soil. In the Sahel,
329 apparent differences in the definition of natural and managed C4 grass account for differences
330 found between ORCHIDEE and JSBACH. The inclusion of the LC_CCI product resulted in a large
331 increase in the C4 grass fraction over the Sahel in ORCHIDEE, whereas no significant change in
332 the C4 grass fraction has been found over these areas for JSBACH. Instead, an increase in C4
333 crops was found over the Sahel for JSBACH. Since the JSBACH conversion also accounts for
334 pasture, this difference may be well the result of the pasture definition, which is a weighted part
335 of all herbaceous PFTs. This also partly explains why the JSBACH C4 pasture PFT decreases
336 exactly in the same areas where the C4 crops increase due to the use of the LC_CCI data. In JULES,
337 the C4 types over Sahel shift to bare soil.

338

339 In the tropics, reductions in broadleaved tropical tree cover were largely consistent across all 3
340 ESMs, although increases in broadleaf forest area were found for some parts of African Congo
341 Basin for JULES (Figure 6). Needleleaved forest area increased compared to the reference
342 dataset for both JULES and JSBACH for boreal Europe and Australia (shrubland PFTs). The
343 increase in needleleaved PFTs in boreal Europe was partially associated with a decrease in

Plant functional type classification

344 broadleaves (Figure 6a and 6b) for all three models, but also a decrease in natural grassland
345 cover.

346

347 **Discussion**

348 *Advantages of the LC_CCI for ESM modeling*

349 The LC_CCI approach provides the ESM modeling community with a flexible tool for using up-to-
350 date land-cover information consistently provided over time. Following the requests of the user
351 survey, the land-cover dataset is available across multiple spatial domains, conforms to standard
352 file formats used in numerical models, and includes information on classification confidence
353 levels for the land cover classes and resulting PFT fractions. The standardized conversion tool
354 provides users with a consistent documented approach for aggregating land cover classes and
355 thus overcomes limitations associated with consensus approaches, for example (Tuanmu and
356 Jetz, 2014). Of particular importance is that the multi-temporal LC_CCI mapping approach
357 facilitated more accurate mapping leading to improved remote sensing observations of
358 deforested areas in the tropics, the treeline-tundra boundary in the high latitudes, and better
359 distinctions between managed and non-managed grasslands in Africa. Additionally, the SAR-
360 based water bodies and coastline delineation helped to standardize the physical boundaries
361 between terrestrial and water systems for all models. Using this standardized PFT mapping
362 approach for ESMs can be expected to reduce model ensemble uncertainty as attempted by
363 recent inter-model comparison efforts (Huntzinger et al., 2013).

364

365 *Opportunities for Phase 2*

Plant functional type classification

366 During Phase 1 of the LC_CCI project (2011-2014) several limitations of the conversion scheme
367 and tool were recognized and have been targeted for improvement in Phase 2, where
368 improvements to the land cover thematic classes and to the conversion scheme will be made. For
369 example, in the high latitudes, a reduction in grassland fractional cover was observed with the
370 LC_CCI product for all models, and on further investigation, it was recognized that a better
371 representation of lichens and moss vegetation (Class 140, Table 3) would be an improvement for
372 the Sparse Vegetation category (Class 150), especially in the high latitudes. Conversion of high-
373 latitude land cover classes to PFT equivalents has been a challenge in several recent regional
374 studies (Ottlé et al., 2013; Wullschleger et al., 2014) where discriminating spectrally between
375 shrubs and trees, or grass and non-vascular plant species, remains difficult. Accurate mapping of
376 high-latitude vegetation can be particularly important for modeling wildfire (Yue et al., 2014)
377 where the spread of tundra fire is sensitive to fuel loading. In the tropics, the seasonal cycle of
378 forest canopies continues to be a contentious issue (Morton et al., 2014; Myneni et al., 2007;
379 Poulter and Cramer, 2009; Ryan et al., 2014) with the binary distinction between evergreen and
380 deciduous phenology proving to be overly-simplistic where semi-deciduous traits are perhaps
381 more appropriate (Borchert et al., 2002) and thus the development of tropical phenology traits
382 that correspond to recent observations is a high priority (Bi et al., 2015). More specifically, Phase
383 2 will target i) improved thematic accuracy with a specific focus on transition areas (e.g.
384 grassland-sparse vegetation-bare soil, tree-shrub-grassland) and the distinction between C3 and
385 C4 grasses, ii) create a historical land cover time series to cover the 1990s using 1km AVHRR
386 NDVI surface reflectances, iii) include more detailed change detection, with more classes, i.e.,
387 IPCC land categories (forests, agriculture, grassland, settlement, wetland, other land) as targets,
388 and iv) deliver an albedo and/or LAI seasonality product.

Plant functional type classification

389

390 Physiological traits such as nitrogen fixation and different photosynthetic pathways, C3, C4 or
391 Crassulacean Acid Metabolism (CAM), are presently not detectable from surface reflectance
392 values, and so broad climate-based assumptions must be made to split into these groups. These
393 assumptions can lead to large uncertainties that can impact a chain of ecosystem processes and
394 land surface properties. While the LC_CCI dataset provides updated information on inland water
395 bodies, the seasonality of water bodies and wetlands is yet to be represented and only
396 considered in radar based surveys (Schroeder et al., In preparation). Finally, the existing 22
397 UNLCCS land-cover classes currently do not include pastures whereas the importance of grazing
398 on biogeochemical cycling is becoming increasingly recognized (Foley et al., 2005). Instead,
399 pastures are currently mapped as croplands or grasslands according to their degree of
400 management. Better thematic discrimination between these 3 classes would clearly improve the
401 carbon cycle modeling as agriculture, in the broadest sense, is a significant contributor to land
402 degradation and anthropogenic global greenhouse gas emissions (Haberl et al., 2007). Earth
403 observation products are generally limited in to mapping land surface structural properties
404 rather than functional one, and model-data fusion approaches can help reconcile problems that
405 might arise from this limitation, especially in the case of grassland systems which may be
406 managed or unmanaged, or may have different photosynthetic pathways. Nevertheless, remote
407 sensing of land *management* categories remains a challenging task since existing classification
408 approaches have yet to demonstrate an ability to capture the whole range of rangelands and
409 crops diversity at global scale.

410

411 *Earth System Modeling challenges*

Plant functional type classification

412 Updating PFT datasets used in ESMs will clearly lead to improvements in the realism of the
413 patterns of biogeography and have important feedbacks on simulating ecosystem processes and
414 interactions with the atmosphere. Available PFT datasets used in ESMs remain outdated, using
415 land cover information from the 1980s mainly because of a lack of tools available for cross-
416 walking land cover to PFTs. The LC_CCI scheme and tool fills a critical data need for improving
417 the representation of carbon, water and energy cycles being developed by the modeling
418 community, however, extensive model benchmarking and calibration activities may now be
419 necessary before the new PFT datasets result in model improvement. For example, model
420 processes may be calibrated to some extent to produce performance metrics under outdated
421 land cover information, and thus a range of benchmarks should be considered when
422 transitioning to new PFT information.

423

424 *Summary*

425 The LC_CCI has made significant progress in responding to the ESM community data needs
426 (Tsendbazar et al., 2014). These include:

- 427 - New land-cover classifications for 3 Epochs using consistent algorithms and based on the
428 UNLCCS system.
- 429 - A user-friendly tool that can map the UNLCCS classes into user-defined PFT classes and at
430 most grid resolutions used by the ESM community.
- 431 - Seasonality products describing average weekly conditions for burned area, NDVI and
432 snow cover.
- 433 - Confidence information for each of the UNLCCS classes and a median estimate for the
434 converted PFT legend.

Plant functional type classification

435 The UNLCCS-PFT conversion tool and the land cover products will continue to be improved
436 during Phase 2 of the LC_CCI with updates made periodically and described at [http://www.esa-](http://www.esa-landcover-cci.org)
437 [landcover-cci.org](http://www.esa-landcover-cci.org).

438

439 **Acknowledgements**

440 The LC_CCI project was funded by the European Space Agency Climate Change Initiative Phase 1.
441 The authors appreciate the support and comments from Frank Martin Seifert, Vasileos Kalogirou
442 and Fabrizio Ramoino.

443

444 **References**

- 445 Alton, P. B.: How useful are plant functional types in global simulations of the carbon, water, and
446 energy cycles?, *Journal of Geophysical Research*, 116, G01030, 2011.
- 447 Arino, O., Bicheron, P., Achard, F., Latham, J., Witt, R., and Weber, J. L.: GLOBCOVER The most
448 detailed portrait of Earth, *ESA Bulletin-European Space Agency*, 136, 24-31, 2008.
- 449 Baker, B., Diaz, H., Hargrove, W., and Hoffman, F. M.: Use of the Köppen–Trewartha climate
450 classification to evaluate climatic refugia in statistically derived ecoregions for the
451 People’s Republic of China, *Clim. Change*, 98, 113-131, 2010.
- 452 Bartholome, E. and Belward, A. S.: GLC2000: a new approach to global land cover mapping from
453 Earth observation data, *Int. J. Remote Sens.*, 26, 1959-1977, 2005.
- 454 Bi, J., Knyazikhin, Y., Choi, S., Park, T., Barichivich, J., Ciais, P., Fu, R., Ganguly, S., Hall, F., Hilker, T.,
455 Huete, A., Jones, M., Kimball, J., Lyapustin, A. I., Möttus, M., Nemani, R. R., Piao, S., Poulter,
456 B., Saleska, S. R., Saatchi, S. S., Xu, L., Zhou, L., and Myneni, R. B.: Sunlight mediated
457 seasonality in canopy structure and photosynthetic activity of Amazonian rainforests,
458 *Environmental Research Letters*, 10, 064014, 2015.
- 459 Bonan, G. B.: Forests and climate change: Forcings, feedbacks, and the climate benefits of forests,
460 *Science*, 320, 1444-1449, 2008.
- 461 Bonan, G. B., Levis, S., Kergoat, L., and Oleson, K. W.: Landscapes as patches of plant functional
462 types: An integrating concept for climate and ecosystem models, *Global Biogeochemical*
463 *Cycles*, 16, 5-21-25-23, 2002.
- 464 Bontemps, S., Herold, M., Kooistra, L., van Groenestijn, A., Hartley, A., Arino, O., Moreau, I., and
465 Defourny, P.: Revisiting land cover observation to address the needs of the climate
466 modeling community, *Biogeosciences*, 9, 2145-2157, 2012.
- 467 Borchert, R., Rivera, G., and Hagnauer, W.: Modification of vegetative phenology in a tropical
468 semideciduous forest by abnormal drought and rain, *Biotropica*, 34, 381-393, 2002.
- 469 Clark, J. S., Bell, D. M., and Hersh, M.: Climate change vulnerability of forest biodiversity: climate
470 and competition tracking of demographic rates, *Global Change Biol.*, 17, 1834-1849, 2011.
- 471 Cox, P. M., Betts, R. A., Jones, C. D., Spall, S. A., and Totterdell, I. J.: Acceleration of global warming
472 due to carbon-cycle feedbacks in a coupled climate model, *Nature*, 408, 184-187, 2000.

Plant functional type classification

- 473 DeFries, R., Townshend, J. R. G., and Hansen, M. C.: Continuous fields of vegetation characteristics
474 at the global scale at 1-km resolution, *Journal of Geophysical Research*, 104, 911-916,
475 1999.
- 476 Di Gregorio, A. and Jansen, L.: *Land Cover Classification System (LCCS): Classification Concepts
477 And User Manual*, Rome, Italy, 2000.
- 478 Edwards, E. J., Osborne, C. P., Stromberg, C. A. E., Smith, S. A., and Consortium, C. G.: The Origins of
479 C4 Grasslands: Integrating Evolutionary and Ecosystem Science, *Science*, 328, 587-591,
480 2010.
- 481 FAO and JRC: *Global forest land-use change 1990–2005*, Food and Agriculture Organization of
482 the United Nations and European Commission Joint Research Centre, Rome, FAO, 2012.
- 483 Faroux, S., Kaptue Tchente, A. T., Roujean, J. L., Masson, V., Martin, E., and Le Moigne, P.:
484 ECOCLIMAP-II/Europe: a twofold database of ecosystems and surface parameters at 1 km
485 resolution based on satellite information for use in land surface, meteorological and
486 climate models, *Geoscientific Model Development*, 6, 563-582, 2013.
- 487 Fisher, R. A., McDowell, N., Purves, D., Moorcroft, P., Sitch, S., Cox, P. M., Huntingford, C., Meir, P.,
488 and Woodward, F. I.: Assessing uncertainties in a second-generation dynamic vegetation
489 model caused by ecological scale limitations, *New Phytol.*, 187, 666-681, 2010.
- 490 Foley, J. A., Defries, R., Asner, G. P., Barford, C., Bonon, G., Carpenter, S. R., Chapin, F. S., Coe, M. T.,
491 Daily, G. C., Gibbs, H. K., Helkowski, J. H., Holloway, T., Howard, E. A., Kucharik, C. J.,
492 Monfreda, C., Patz, J. A., Prentice, I. C., Ramankutty, N., and Snyder, P. K.: Global
493 consequences of land use, *Science*, 309, 570-574, 2005.
- 494 Fontes, J., Gastellu-Etchegorry, J. P., Amram, O., and Fluzat, G.: A Global Phenological Model of the
495 African Continent, *Ambio*, 24, 297-303, 1995.
- 496 Friedl, M. A., Sulla-Menashe, D., Tan, B., Schneider, A., Ramankutty, N., Sibley, A., and Huang, X.:
497 MODIS Collection 5 Global Land Cover: Algorithm refinements and characterization of
498 new datasets, *Remote Sens. Environ.*, 114, 168-182, 2010.
- 499 Giri, C., Zhu, Z., and Reed, B.: A comparative analysis of the Global Land Cover 2000 and MODIS
500 land cover data sets, *Remote Sens. Environ.*, 94, 123-132, 2005.
- 501 Gotangco Castillo, C. K., Levis, S., and Thornton, P.: Evaluation of the New CNDV Option of the
502 Community Land Model: Effects of Dynamic Vegetation and Interactive Nitrogen on CLM4
503 Means and Variability*, *J. Clim.*, 25, 3702-3714, 2013.
- 504 Gurney, K. R., Law, R. M., Denning, A. S., Rayner, P., Baker, D., Bousquet, P., Bruhwiler, L., Chen, Y.
505 H., Ciais, P., Fan, S. M., Fung, I. Y., Gloor, M., Heimann, M., Higuchi, N., John, J., Maki, T.,
506 Maksyutov, S., Masarie, K., Peylin, P., Prather, M., Pak, B. C., Randerson, J. T., Sarmiento, J.,
507 Taguchi, S., Takahashi, T., and Yuen, C. W.: Towards robust regional estimates of CO₂
508 sources and sinks using atmospheric transport models, *Nature*, 415, 626-630, 2002.
- 509 Haberl, H., Erb, K. H., Krausmann, F., Gaube, V., Bondeau, A., Plutzar, C., Gingrich, S., Lucht, W., and
510 Fischer-Kowalski, M.: Quantifying and mapping the human appropriation of net primary
511 production in earth's terrestrial ecosystems, *Proceedings of the National Academy of
512 Science*, 104, 12942-12947, 2007.
- 513 Herold, M., Mayaux, P., Woodcock, C. E., Baccini, A., and Schmullius, C.: Some challenges in global
514 land cover mapping: An assessment of agreement and accuracy in existing 1 km datasets,
515 *Remote Sens. Environ.*, 112, 2538-2556, 2008.
- 516 Herold, M., van Groenestijn, A., Kooistra, L., Kalogirou, V., and Arino, O.: User Requirements
517 documents: Land Cover CCI, Université catholique de Louvain (UCL)-Geomatics, Louvain-
518 la-Neuve, Belgium., 2011.

Plant functional type classification

- 519 Hollmann, R., Merchant, C., Saunders, R., Downy, C., Buchwitz, M., Cazenave, A., Chuvieco, E.,
520 Defourny, P., de Leeuw, G., Forsberg, R., Holzer-Popp, T., Paul, F., Sandven, S.,
521 Sathyendranath, S., van Roozendaal, M., and Wagner, W.: The ESA climate change
522 initiative: Satellite data records for essential climate variables, *Bulletin of the American*
523 *Meteorology Society*, 94, 1541–1552, 2013.
- 524 Huntzinger, D. N., Schwalm, C., Michalak, A. M., Schaefer, K., King, A. W., Wei, Y., Jacobson, A., Liu,
525 S., Cook, R. B., Post, W. M., Berthier, G., Hayes, D., Huang, M., Ito, A., Lei, H., Lu, C., Mao, J.,
526 Peng, C. H., Peng, S., Poulter, B., Ricciuto, D., Shi, X., Tian, H., Wang, W., Zeng, N., Zhao, F.,
527 and Zhu, Q.: The North American Carbon Program Multi-Scale Synthesis and Terrestrial
528 Model Intercomparison Project – Part 1: Overview and experimental design, *Geosci.*
529 *Model Dev.*, 6, 2121-2133, 2013.
- 530 Hurtt, G. C., Frothingham, S., Fearon, M. G., Moore, B., Shevliakova, E., Malyshev, S., Pacala, S., and
531 Houghton, R. A.: The underpinnings of land-use history: three centuries of global gridded
532 land-use transitions, wood-harvest activity, and resulting secondary lands, *Global Change*
533 *Biol.*, 12, 1208-1229, 2006.
- 534 Jung, M., Henkel, K., Herold, M., and Churkina, G.: Exploiting synergies of global land cover
535 products for carbon cycle modeling, *Remote Sens. Environ.*, 101, 534-553, 2006.
- 536 Kattge, J. and Diaz, S. and Lavorel, S. and Prentice, I. C. and Leadley, P. and BÖNisch, G. and
537 Garnier, E. and Westoby, M. and Reich, P. B. and Wright, I. J. and Cornelissen, J. H. C. and
538 Violle, C. and Harrison, S. P. and Van Bodegom, P. M. and Reichstein, M. and Enquist, B. J.
539 and Soudzilovskaia, N. A. and Ackerly, D. D. and Anand, M. and Atkin, O. and Bahn, M. and
540 Baker, T. R. and Baldocchi, D. and Bekker, R. and Blanco, C. C. and Blonder, B. and Bond, W.
541 J. and Bradstock, R. and Bunker, D. E. and Casanoves, F. and Cavender-Bares, J. and
542 Chambers, J. Q. and Chapin Iii, F. S. and Chave, J. and Coomes, D. and Cornwell, W. K. and
543 Craine, J. M. and Dobrin, B. H. and Duarte, L. and Durka, W. and Elser, J. and Esser, G. and
544 Estiarte, M. and Fagan, W. F. and Fang, J. and FernÁNdez-MÉNdez, F. and Fidelis, A. and
545 Finegan, B. and Flores, O. and Ford, H. and Frank, D. and Freschet, G. T. and Fyllas, N. M.
546 and Gallagher, R. V. and Green, W. A. and Gutierrez, A. G. and Hickler, T. and Higgins, S. I.
547 and Hodgson, J. G. and Jalili, A. and Jansen, S. and Joly, C. A. and Kerkhoff, A. J. and Kirkup,
548 D. and Kitajima, K. and Kleyer, M. and Klotz, S. and Knops, J. M. H. and Kramer, K. and
549 KÜHN, I. and Kurokawa, H. and Laughlin, D. and Lee, T. D. and Leishman, M. and Lens, F.
550 and Lenz, T. and Lewis, S. L. and Lloyd, J. and LlusiÀ, J. and Louault, F. and Ma, S. and
551 Mahecha, M. D. and Manning, P. and Massad, T. and Medlyn, B. E. and Messier, J. and
552 Moles, A. T. and MÜLLer, S. C. and Nadrowski, K. and Naeem, S. and Niinemets, Ü. and
553 NÖLlert, S. and NÜSke, A. and Ogaya, R. and Oleksyn, J. and Onipchenko, V. G. and Onoda,
554 Y. and OrdoÑEz, J. and Overbeck, G. and Ozinga, W. A. and PatiÑO, S. and Paula, S. and
555 Pausas, J. G. and PeÑUelas, J. and Phillips, O. L. and Pillar, V. and Poorter, H. and Poorter, L.
556 and Poschlod, P. and Prinzing, A. and Proulx, R. and Rammig, A. and Reinsch, S. and Reu, B.
557 and Sack, L. and Salgado-Negret, B. and Sardans, J. and Shiodera, S. and Shipley, B. and
558 Siefert, A. and Sosinski, E. and Soussana, J. F. and Swaine, E. and Swenson, N. and
559 Thompson, K. and Thornton, P. and Waldram, M. and Weiher, E. and White, M. and White,
560 S. and Wright, S. J. and Yguel, B. and Zaehle, S. and Zanne, A. E. and Wirth, C.: TRY – a
561 global database of plant traits, *Global Change Biol.*, 17, 2905-2935, 2011.
- 562 Klein Goldewijk, K. and Batjes, J. J.: A hundred year (1890-1990) database for integrated
563 environmental assessments (HYDE, version 1.1), Bilthoven, the Netherlands, 1997.

Plant functional type classification

- 564 Knorr, W.: Annual and interannual CO₂ exchanges of the terrestrial biosphere: process-based
565 simulations and uncertainties, *Global Ecol. Biogeogr.*, 9, 225-252, 2000.
- 566 Kottek, M., J. Grieser, C. Beck, B. Rudolf, and F. Rubel, 2006: . *Meteorol. Z.*, 15, 259-263.: World
567 Map of the Köppen-Geiger climate classification updated, *Meteorologische Zeitschrift*, 15,
568 259-263, 2006.
- 569 Krinner, G., Viovy, N., de Noblet-Ducoudré, N., Ogeé, J., Polcher, J., Friedlingstein, P., Ciais, P., Sitch,
570 S., and Prentice, I. C.: A dynamic global vegetation model for studies of the coupled
571 atmosphere-biosphere system, *Global Biogeochemical Cycles*, 19, GB1015,
572 doi:10.1029/2003GB002199, 2005.
- 573 Lawrence, D. M., Oleson, K. W., Flanner, M. G., Thornton, P. E., Swenson, S. C., Lawrence, P. J., Zeng,
574 X., Yang, Z. L., Levis, S., Sakaguchi, K., Bonan, G. B., and Slater, A. G.: Parameterization
575 Improvements and Functional and Structural Advances in Version 4 of the Community
576 Land Model, *Journal of Advances in Modeling Earth Systems*, 3, M03001, 2011.
- 577 Lawrence, P. J. and Chase, T. N.: Representing a MODIS Consistent Land Surface in the
578 Community Land Model (CLM 3.0): Part 1 Generating MODIS Consistent Land Surface
579 Parameters, *Journal of Geophysical Research*, 112, G01023, 2007.
- 580 Loveland, T. R. and Belward, A. S.: The IGBP-DIS global 1km land cover data set, DISCover: First
581 results, *Int. J. Remote Sens.*, 18, 3289-3295, 1997.
- 582 Monfreda, C., Ramankutty, N., and Foley, J. A.: Farming the planet: 2. Geographic distribution of
583 crop areas, yields, physiological types, and net primary production in the year 2000,
584 *Global Biogeochemical Cycles*, 22, GB1022, 2008.
- 585 Morton, D. C., Nagol, J., Carabahal, C. C., Rosette, J., Palace, M., Cook, B. D., Vermote, E. F., Harding,
586 D. J., and North, P. R. J.: Amazon forests maintain consistent canopy structure and
587 greenness during the dry season, *Nature*, 506, 221-224, 2014.
- 588 Myneni, R. B., Yang, W., Nemani, R. R., Huete, A. R., Dickinson, R. E., Knyazikhin, Y., Didan, K., Fu,
589 R., Negron Juarez, R. I., Saatchi, S. S., Hashimoto, H., Shabanov, N. V., Tan, B., Ratana, P.,
590 Privette, J. L., Morisette, J. T., Vermote, E. F., Roy, D. P., Wolfe, R. E., Fiedl, M. A., Running, S.
591 W., Votava, P., El-Saleous, N., Devadiga, S., Su, Y., and Salomonson, V. V.: Large seasonal
592 swings in leaf area of Amazon rainforests, *Proceedings of the National Academy of*
593 *Science*, 104, 4820-4823, 2007.
- 594 Neumann, K., Herold, M., Hartley, A., and Schmillius, C.: Comparative assessment of CORINE2000
595 and GLC2000: Spatial analysis of land cover data for Europe, *Journal of Applied Earth*
596 *Observation and Geoinformation*, 9, 425-437, 2007.
- 597 Olson, D. M., Dinerstein, E., Wikramanaye, E. D., Burgess, N. D., Powell, G. V. N., Underwood, E. C.,
598 D'Amico, J. A., Itoua, I., Strand, H. E., Morrison, J. C., Loucks, C. J., Allnutt, T. F., Ricketts, T.
599 H., Kura, Y., Lamoreux, J. F., Wettengel, W. W., Hedao, P., and Kassem, K. R.: Terrestrial
600 ecoregions of the world: A new map of life on Earth, *Bioscience*, 51, 933-938, 2001.
- 601 Olson, J., Watts, J. A., and Allison, L. J.: Carbon in Live Vegetation of Major World Ecosystems,
602 ORNL-5862. Oak Ridge National Laboratory, Oak Ridge, Tennessee, pp. 164. , 1983.
- 603 Ottlé, C., Lescure, J., Maignan, F., Poulter, B., Wang, T., and Delbart, N.: Use of various remote
604 sensing land cover products for plant functional type mapping over Siberia, *Earth System*
605 *Science Data*, 5, 331-348, 2013.
- 606 Pacifico, F., Harrison, S. P., Jones, C. D., Arneth, A., Sitch, S., Weedon, G. P., Barkley, M. P., Palmer, P.
607 I., Serça, D., Potosnak, M., Fu, T. M., Goldstein, A., Bai, J., and Schurgers, G.: Evaluation of a
608 photosynthesis-based biogenic isoprene emission scheme in JULES and simulation of

Plant functional type classification

- 609 isoprene emissions under present-day climate conditions, *Atmos. Chem. Phys.*, 11, 4371-
610 4389, 2011.
- 611 Pavlick, R., Drewry, D. T., Bohn, K., Reu, B., and Kleidon, A.: The Jena Diversity-Dynamic Global
612 Vegetation Model (JeDi-DGVM): a diverse approach to representing terrestrial
613 biogeography and biogeochemistry based on plant functional trade-offs, *Biogeosciences*,
614 10, 4137-4177, 2013.
- 615 Peel, M. C., Finlayson, B. L., and McMahon, T. A.: Updated world map of the Köppen-Geiger
616 climate classification, *Hydrology and Earth System Sciences*, 11, 1633-1644, 2007.
- 617 Pongratz, J., Reick, C. H., Raddatz, T., and Claussen, M.: Effects of anthropogenic land cover
618 change on the carbon cycle of the last millennium, *Global Biogeochemical Cycles*, 23,
619 GB4001, 2009.
- 620 Poulter, B., Ciais, P., Hodson, E. L., Lischke, H., Maignan, F., Plummer, S., and Zimmermann, N. E.:
621 Plant functional type mapping for earth system models, *Geoscientific Model Development*,
622 4, 993-1010, 2011.
- 623 Poulter, B. and Cramer, W.: Satellite remote sensing of tropical forest canopies and their seasonal
624 dynamics, *Int. J. Remote Sens.*, 30, 6575-6590, 2009.
- 625 Radoux, J., Lemarche, C., Van Bogaert, E., Bontemps, S., Brockmann, C., and Defourny, P.:
626 Automated Training Sample Extraction for Global Land Cover Mapping, *Remote Sensing*,
627 6, 3965-3987, 2014.
- 628 Ramankutty, N., Evan, A. T., Monfreda, C., and Foley, J. A.: Farming the planet: 1. Geographic
629 distribution of global agricultural lands in the year 2000, *Global Biogeochemical Cycles*,
630 22, GB1003, 2008.
- 631 Ramankutty, N. and Foley, J. A.: Characterizing patterns of global land use: An analysis of global
632 croplands data, *Global Biogeochemical Cycles*, 12, 667-685, 1998.
- 633 Reich, P. B. and Borchert, R.: Water stress and tree phenology in a tropical dry forest in the
634 lowlands of Costa Rica, *J. Ecol.*, 72, 61-74, 1984.
- 635 Reick, C. H., Raddatz, T., Brovkin, V., and Gayler, V.: Representation of natural and anthropogenic
636 land cover change in MPI-ESM, *Journal of Advances in Modeling Earth Systems*, 5, 1-24,
637 2013.
- 638 Ryan, C. M., Williams, M., Hill, T. C., Grace, J., and Woodhouse, I. H.: Assessing the phenology of
639 southern tropical Africa: A comparison of hemispherical photography, scatterometry, and
640 optical/NIR remote sensing, *IEEE Transactions on Geoscience and Remote Sensing*, 52,
641 519-528, 2014.
- 642 Scheiter, S. and Higgins, S. I.: Impacts of climate change on the vegetation of Africa: an adaptive
643 dynamic vegetation modelling approach, *Global Change Biol.*, 15, 2224-2246, 2009.
- 644 Schroeder, R., McDonald, K., Chan, S., Chapman, B., Podest, E., Bohn, T., Jones, L., Kimball, J.,
645 Zimmermann, R., and Küppers, M.: Development and evaluation of a multi-year global
646 inundated area dataset derived from combined active/passive microwave remote
647 sensing, In preparation. In preparation.
- 648 Sellers, P., Randall, D. A., Collatz, G. J., Berry, J. A., Field, C. B., Dazlich, D., Zhang, C., Collelo, G. D.,
649 and Bounoua, L.: A revised land surface parameterization (SiB2) for atmospheric GCMs.
650 Part I: Model formulation, *J. Clim.*, 9, 676-705, 1996.
- 651 Sitch, S., Smith, B., Prentice, I. C., Arneth, A., Bondeau, A., Cramer, W., Kaplan, J. O., Levis, S., Lucht,
652 W., Sykes, M. T., Thonicke, K., and Venevsky, S.: Evaluation of ecosystem dynamics, plant
653 geography and terrestrial carbon cycling in the LPJ dynamic global vegetation model,
654 *Global Change Biol.*, 9, 161-185, 2003.

Plant functional type classification

- 655 Still, C. J., Berry, J. A., Collatz, G. J., and DeFries, R.: Global distribution of C3 and C4 vegetation:
656 Carbon cycle implications, *Global Biogeochemical Cycles*, 17, 6-1-6-14, 2003.
- 657 Tsendbazar, N. E., de Bruin, S., and Herold, M.: Assessing global land cover reference datasets for
658 different user communities, *ISPRS Journal of Photogrammetry and Remote Sensing*, 2014.
659 2014.
- 660 Tuanmu, M.-N. and Jetz, W.: A global 1-km consensus land-cover product for biodiversity and
661 ecosystem modelling, *Global Ecol. Biogeogr.*, 9, 1031-1045, 2014.
- 662 Ustin, S. L. and Gamon, J. A.: Remote sensing of plant functional types, *New Phytol.*, 186, 795-816,
663 2010.
- 664 Verant, S., Laval, K., Polcher, J., and De Castro, M.: Sensitivity of the continental hydrological cycle
665 to the spatial resolution over the Iberian Peninsula, *Journal of Hydrometeorology*, 5, 267-
666 285, 2004.
- 667 Wilson, M. F. and Henderson-Sellers, A.: A global archive of land cover and soils data for use in
668 general circulation climate models, *Journal of Climatology*, 5, 119-143, 1985.
- 669 Wullschleger, S. D., Epstein, H. E., Box, E. O., Euskirchen, E. S., Goswami, S., Iverson, C. M., Kattge,
670 J., Norby, R. J., van Bodegom, P. M., and Xu, X.: Plant functional types in Earth system
671 models: past experiences and future directions for application of dynamic vegetation
672 models in high-latitude ecosystems, *Ann. Bot.*, 2014. 2014.
- 673 Yue, C., Ciais, P., Cadule, P., Thonicke, K., Archibald, S., Poulter, B., Hao, W. M., Hantson, S.,
674 Mouillot, F., Friedlingstein, P., Maignan, F., and Viovy, N.: Modelling fires in the terrestrial
675 carbon balance by incorporating SPITFIRE into the global vegetation model ORCHIDEE –
676 Part 1: Simulating historical global burned area and fire regime, *Geoscientific Model
677 Development Discussions*, 7, 1-51, 2014.
678
679

Plant functional type classification

680 Table 1: Plant functional types used by three earth system models and mapped by the LC_CCI
 681 Initiative.

ORCHIDEE	JSBACH	JULES	ESA LC_CCI
Tropical broadleaf evergreen	Tropical broadleaf evergreen	Broadleaf trees	Broadleaf evergreen tree
Tropical broadleaf deciduous	Tropical broadleaf deciduous	Needleleaf trees	Broadleaf deciduous tree
Temperate needleleaf evergreen	Extra-tropical evergreen	C3 grass	Needleleaf evergreen tree
Temperate broadleaf deciduous	Extra-tropical deciduous	C4 grass	Needleleaf deciduous tree
Temperate broadleaf summergreen	Rain-green shrubs	Shrubs	Broadleaf evergreen shrub
Boreal needleleaf evergreen	Deciduous shrubs		Broadleaf deciduous shrub
Boreal broadleaf summergreen	Tundra		Needleleaf evergreen shrub
Boreal needleleaf summergreen	Swamp		Needleleaf deciduous shrub
C3 grass	C3 grass		Natural grass
C4 grass	C4 grass		Managed grass
C3 crops	C3 crops		
C4 crops	C4 crops		

682

Plant functional type classification


686 Table 3: Minimum set of projections and spatial resolutions included in the re-projection,
 687 aggregation, subset and conversion tool developed by the LC_CCI project - LC_CCI user tool

Regional subset ID	Predefined regional subset
	Free specification of regional subset (4 corner coordinates)
Spatial resolution	Original resolution 0.25 degree 0.5 degree 1 degree 1.875 degree 1.875 x 1.25 degree 3.75 x 2.5 degree
Projection	Original projection (Plate-Carrée) Gaussian grid, Rotated lat/lon grid
Conversion of LC_CCI classes to PFT	LC_CCI standard cross table User defined cross table

688

Plant functional type classification

689 Figure 1: - Visualization of the pixel aggregation from the spatial resolution of original LC_CCI
690 map product into the user-defined spatial resolution of the aggregated LC_CCI map product.
691




692

	Area	Majority class
class a	~ 8/16	1
class b	~ 5/16	2
class c	~ 2/16	3
class d	~ 1/16	4

693
694
695


Plant functional type classification

696 Figure 2: The LC_CCI land cover conversion tool processing chain requires converting the
697 thematic legend and resampling the grid resolution to user defined PFT and coordinate system.
698 Independent of the LC_CCI tool, users can append climate classes to the PFT aggregation.
699
700
701



Plant functional type classification


702 Figure 3: Fractional coverage of plant functional types, at 0.5-degree spatial resolution,
703 calculated from original 300-meter LC_CCI dataset, epoch 2008-2012, using the LC_CCI
704 conversion tool



705
706

Plant functional type classification


707 Figure 4: Global PFT coverage comparing the LC_CCI and original datasets for a) ORCHIDEE, b)
 708 JULES, and c) JSBACH. Where 'Br' is broadleaf, 'Ne' is needleleaf, 'Ev' is evergreen, 'De' is
 709 deciduous, 'ManGr' is managed grassland, 'NatGr' is natural grassland, and 'barren' includes bare
 710 soil or ice. Note JSBACH has no bare soil category.



711

Plant functional type classification


712 Figure 5: Difference in fractional coverage between the LC_CCI (epoch 2008-2012) and original
713 ORCHIDEE PFT dataset, based on Olson et al. (1983).



714


Plant functional type classification

715 Figure 6: Regional correlations between the original ESM PFT coverage and the LC_CCI, epoch
 716 2008-2012, coverage for a) broadleaved trees, b) needleleaved trees, c) natural grasslands, and
 717 d) managed grasslands. The regions follow the TRANSCOM Experiment biome boundary
 718 definitions, which partition terrestrial ecosystems into 13 regions of similar vegetation (see
 719 Appendix 1).
 720



Plant functional type classification

721 Appendix 1: TRANSCOM experiment biome boundaries from Gurney et al. (2002). The codes
722 from Figure 6 are Boreal North America (NAMBO), Temperate North America (NAMTE), Tropical
723 South America (SAMTR), Temperate South America (SAMTE), North Africa (NAf), South Africa
724 (SAf), Boreal Eurasia (EuBO), Temperate Eurasia (EuTE), Tropical Asia (AsTR), Australi (AUST),
725 Europe (EURO), Arid North Africa (NAfarid), Arid South Africa (SAfarid).



726