

Reaching the lower stratosphere: Validating an extended vertical grid for COSMO

Johannes Eckstein, Sarah Schmitz, and Roland Ruhnke

Karlsruhe Institute of Technology, Institute of Meteorology and Climate Research,
Herrmann-von-Helmholtz-Platz 1, 76344 Eggenstein-Leopoldshafen, Germany

Correspondence to: Johannes Eckstein (johannes.eckstein@kit.edu)

Abstract. This study presents an extended vertical grid for the regional atmospheric model COSMO reaching up to 33km. The extended setup has been used to stably simulate eleven months in a domain covering central and northern Europe. Temperature and relative humidity have been validated using radio sonde data in polar and temperate latitudes, focussing on the polar and mid-latitude stratosphere over Europe. Temperature values are reproduced very well by the model. Relative humidity could only be met in the mean over the whole time period after excluding data from Russian stations, which showed significantly higher values. A sensitivity study shows the stability of the model against different forcing intervals and damping layer heights.

1 Introduction

10 The upper troposphere and lowermost stratosphere is a place of sharp gradients in many constituents of air and of the physical parameters used to describe its state. Temperature and ozone are textbook examples, but methane, water and many more species also show a strong gradient. At the same time, being the boundary to the lower atmosphere, this is an area where small scale fluctuations can have a strong influence on the stratosphere and its composition (Zahn et al., 2014).

15 In order to simulate this highly vulnerable and influential layer directly, a model with high vertical and horizontal resolution is needed. Global models usually are too coarsely resolved and cannot model the small scale processes. In extending the vertical layering of the regional model COSMO to 33km, we present here a model that can fill the gap. As we planned to apply the extended setup to simulations covering polar spring and the associated ozone loss with the coupled chemistry model
20 COSMO-ART (Vogel et al., 2009), we focus here on polar latitudes, but always refer to temperate regions also.

After an introduction to the model and an exact definition of the extended vertical grid in Sect. 2, the measurement data is introduced in Sect. 3. COSMO is shown to be able to run stably with the extended layering. Using radio sonde data and regrided data from meteorological reanalyses, it is shown that the model is able to reproduce temperatures very well (Sect. 4.2) while relative humidity is more difficult (Sect. 4.3) and only its mean value could be reproduced. Two runs with different boundary conditions were performed to test the influence on the model result.

Additionally, three more runs were done in order to test the stability of the model against an increased boundary forcing interval set to 12h and 24h instead of 6h and against increasing the thickness of the damping layer by setting its lower end down to 22km instead of 28km. Sect. 5 presents the results of this sensitivity study, showing that the model will still run stably.

2 The model: vertical grid, boundary data and domain

This section gives a short introduction to COSMO and explains the changes made to the standard vertical grid as well as the boundary data used and the specified domain.

2.1 Introduction to the model

COSMO (COntortium for Small-scale MOdelling) is a regional atmospheric model that has been developed by a consortium lead by the German weather service DWD (Deutscher Wetterdienst). DWD uses the model for its regional numerical weather forecast of Europe and Germany with a resolution of 7km and 2.8km respectively (Baldauf et al., 2011b). Many extensions have been developed for the model, for example COSMO-ART including chemistry and aerosols (Vogel et al., 2009). For this study, the model was set up to run in forecast mode to simulate several months in form of a hindcast using reanalysis data as boundary forcing.

The standard setup of COSMO used for the forecast of central Europe (DWD domain COSMO-DE) reaches to a height of 22.0km (Baldauf et al., 2011a). This is the vertical grid referred to as the standard vertical setup or grid in this study, well aware of the fact the vertical grid used to simulate a larger European domain (COSMO-EU) that reaches up to 23.6km (Schulz and Schättler, 2009) is just as frequently used by DWD. The model has also been used to study greater heights in tropical latitudes in the AMMA (African Monsoon Multidisciplinary Analyses) project (Gantner and Kalthoff, 2010), reaching 28.0km, and a tropical setup reaching up to 30.0km has also been developed (Krähenmann et al., 2013). With the extended vertical grid presented in this study, it becomes possible to simulate the lowermost stratosphere in polar latitudes. This validation study opens the door to new applications of COSMO.




Figure 1. The vertical grids of the COSMO model considered in this study. The damping layers are also given as shaded areas.

2.2 The extended vertical grid

The standard vertical grid of the COSMO model reaches up to 22.0 km in 50 layers. The vertical structure is visible from Fig. 1, exact values are given in Tab. 1. The damping layer in the top layers begins at 11.357 m in standard setup.

The vertical layering of the new grid introduced in this study is also given in Fig. 1 and Tab. 1. It is focused on the lower stratosphere, with the highest of the 60 layers at 33 km, the damping layer beginning at 28 km ($rdheight = 28000.0$ in the namelist). The top layer of the extended grid about 10 km above that of the standard grid and the distance between the layers is slightly smaller in all heights above the lowest kilometer, as is also visible in Fig. 1.

In order to test the sensitivity of the model to the size of the damping layer, an additional model run was done, for which the lower boundary of the damping layer was set to 22 km ($rdheight = 22000.0$ in the namelist), which is just the top of the standard grid. The damping layer then spans one third of the model layers.

2.3 The analyses used as boundary data

In order to examine the influence of different boundary data on the model results, the model was run twice, using ERA-Interim and NCEP reanalysis data for starting and boundary values. The vertical layering of the two reanalyses is displayed in Fig. 2. In order to better evaluate the model, the reanalysis data was also interpolated to the vertical grid used for the output of the model.




Figure 2. The vertical structure of the NCEP and ERA-Interim reanalysis used as boundary conditions.

The reanalysis project of the National Center for Environmental Prediction (NCEP) provides data starting on the first of January, 1948, giving global fields every six hours (0, 6, 12 and 18 UTC) at a resolution of T62, which corresponds to 1.875° (192 points on a latitude) (Kalnay and Coauthors, 1996). The upper boundary is at 2.7 hPa, approximately 42 km in the U.S. standard atmosphere (Sissenwine et al., 1962). So the new vertical grid reaching up to 33 km is still within the vertical limits of the NCEP reanalysis data.

ERA-Interim is the reanalysis project of the European Center for Medium Range Weather Forecast (ECMWF) (Dee et al., 2011). The data was used in this study at a resolution of T255 (corresponding to 0.7° , 512 points on a latitude) and up to 0.1 hPa. So both the vertical and horizontal resolution are higher than those of the NCEP reanalysis. ERA-Interim is available for the same timestamps as the NCEP reanalysis.

In standard setup, the reanalysis data was used in a six hourly interval ($hincbound = 6.0$ in the namelist) to force the model. The sensitivity of the model to this interval of boundary forcing was tested by performing two additional model runs using the ERA-Interim reanalysis data and using it as forcing every 12 and 24 hours ($hincbound = 12.0$ or $hincbound = 24.0$ respectively).

2.4 The model domain

The model domain used in this study is shown in Fig. 3. It covers most of Europe with a focus on the polar latitudes, stretching from northern Africa in the south and covering Svalbard, east of Greenland at 74°N , in the north. The resolution was set to 0.2° . The COSMO model is operationally used by DWD to produce regional weather forecasts for central Europe, but not in northern hemisphere polar latitudes (Baldauf et al., 2011a).




Figure 3. The model domain and the radio sonde stations used in this study. The domain is displayed as gray shading, the radiosonde stations are numbered from south to north, numbers also referring to Tab. 3. Russian stations are marked in red.

So the domain chosen here can be used to assess the performance of the model in polar latitudes, since a direct comparison to an area of regular use is possible. The required namelist parameters needed to reproduce the model domain are given in Tab. 2.

95 The first timestep simulated by the model runs used in this study is October 1, 2010, 0:00 UTC and the last output is for September 1, 2011, 0:00 UTC. The cold temperatures that can be expected in the polar stratosphere especially in winter and the warming in spring both lay well within the simulated time. Output was produced on an hourly basis, the model timestep was set to 60s, using the namelist parameter $dt = 60.0$. It could be shown that the model runs stably in this setup by validating the
 100 whole timeperiod with radiosonde data.

The timespan of eleven months is due to the time limit applied to the calculation. The model was run with a time limit of two days, reaching a total number of 8076 output hours. The last output then turns out to be on September 2, 2011, at 11:00 UTC, but the authors decided to perform this study for the exact eleven months, as given above.

105 3 Measurements

This study validates the output of the COSMO model using the temperature (T) and relative humidity (rH) recorded by radiosondes of stations within the model domain. T and rH are regularly observed values and are here considered basic physical parameters whose distribution well represents the physical state of the model. The measurement data used in this study was taken from the ESRL (Earth System Research Laboratory) radiosonde database provided by NOAA (National Oceanic and Atmospheric Administration) (Schwartz and Govett, 1992).

The location of the 24 stations is given in Fig. 3, exact values and the names being given in Tab. 3. This choice includes all polar stations in the domain and the same number of temperate stations with good data coverage.

115 All stations typically release one radiosonde every twelve hours, at 0 UTC and 12 UTC, so 671 ascents can be expected from each station during the period of 335 simulated days. The actual number of ascents for each station is also given in Tab. 3. All stations except Ny Alesund, which has a little more than one ascent per day, come close to or exceed this number, the average being at 673 ascents. Model and regrided reanalysis data was only considered at times when there was an ascent
120 at the specific station, so approximately every twelve hours.

In order to compare sonde and model data, the grid point closest to each station was used to compare the simulation with measurements. Since the resolution is only 0.2° , the error made by this simple identification is small. The latitude and longitude of the closest grid point can also be found in Tab. 3. An interpolation to the exact location was not considered necessary as the radio sondes drift
125 with the wind, an effect not accountable, since the exact geographic location of each measurement taken by the sonde is not available. This is also the reason why no interpolation in the vertical was done.

In each ascent, the value closest to each model output layer at even kilometers was identified with the height of that layer, the maximum difference allowed having been set to 500 m. Since there are
130 typically more than 20 measurements taken in an ascent, the error was much smaller than this value, reaching only 156.0 m on average, with a standard deviation of 126.3 m.

The data was used as downloaded from the server, only excluding values in $rH > 100\%$. It was found that all stations in Russia give much higher humidity values than the other stations, which is the reason why the humidity data of all Russian stations were excluded from the investigation. This
135 will be further discussed in Sect. 4.3.1.

4 Results

This sections presents the results of the model validation study. Two questions are to be answered: Is the model able to simulate the polar latitudes and the stratospheric heights? And what is the

influence of the boundary data on these results? Following the questions, the answers will also have
140 to be twofold.

After presenting the output grid, the results in temperature are presented. Those of relative humidity are described in the following section. The latter is preceded by the explanation why it seemed reasonable to exclude the data of Russian stations when examining relative humidity.

4.1 The output grid

145 In order to compare the model results to the measurements, model output on a vertical grid of whole kilometers from 8 km to 33 km was used. The values given out above 27 km are already within the damping layer and the results can no longer be considered to come genuinely from the model, so measurements were only compared up to 27 km.

As noted above, the boundary data was also interpolated onto the output grid, using the same
150 program that is used to prepare the boundary data for running the model, called INT2LM (Schättler, 2013). COSMO uses terrain following coordinates. Above a certain value specified in the namelist, the layers become smooth and are no longer terrain following. This height has to be higher than the highest mountain tops in the domain and in this case was set to $vflat = 7000.0$, given in the namelist in m. This is the reason why all analyses done in this study only start at 8 km.

155 4.2 Temperature

To begin the discussion, a look at Fig. 4 exemplifies the basis of this study. It shows all the soundings of the station Jan Mayen during the time considered here. The warming at the end of the polar winter is well visible. Most striking are the many white areas in the image, showing the lack of measurement data. The bottom figure shows the corresponding result of the model run with boundary data by ERA-
160 Interim. The image is filled, but the data was only used for the following analysis if measurements were also available at the timestamp.

Fig. 5 gives exemplary timeseries of Jan Mayen and Madrid in 26 km height, approximately 2.5 km above the model top of the standard vertical COSMO grid for both model runs. When comparing the two figures, temperature values reflect the different latitude: winter temperatures above
165 Jan Mayen are much colder than above Madrid, the warming in spring much more pronounced. The good correspondance of model and measurement not only shows that the two model runs and also the boundary data are very similar, but also that the model performance does not change during the whole simulated period. There is no greater offset in the end than in the beginning.

To compare the data in a more quantitative manner, Fig. 6 shows the mean ascent at Jan Mayen
170 for both model runs. The boundary data is also included in the image. All three soundings lay on top of each other. The minimum temperature in the lowermost stratosphere is well reproduced. In order to compare to a temperate station, Fig. 6 also gives the mean ascent of the station in Madrid.




Figure 4. Temperature values of all soundings of the station Jan Mayen, station no. 20. Measurements are displayed on the top, the image below shows the corresponding model values. Note that this is not a timeseries plot. The dates along the abscissa hold true only for the location they indicate and do not define exact time in between. Dates only increase from left to right, but they are not evenly spaced in time.

The minimum is more pronounced, but also reproduced by the model. There is no difference visible between the model run forced by ERA-Interim and that forced by NCEP reanalysis data.

175 In order to further compare the performance of COSMO, Fig. 7 shows the scatterplots of all measured against modelled temperature values with colorcoded height intervals for all polar stations. The variability in higher altitudes is lower, which is why the scatter is reduced with height. Both model runs with different boundary data simulate temperature very well, reaching about $r^2 = 0.98$. The results of the model in temperate latitudes was just as good and the correlation does not reach
 180 higher values when using the regridded boundary data (not shown).




Figure 5. Timeseries of measured and modelled temperature, 26 km above Jan Mayen (top) and Madrid (bottom). Interpolated reanalysis data is also shown.

When reducing the data to values of descriptive statistics, all stations can be easily compared. Fig. 8 shows the mean of $T_{\text{model}} - T_{\text{meas}}$ and $T_{\text{bound}} - T_{\text{meas}}$ for all levels and for stratospheric levels with $z \geq 11$ km. The stratospheric layers are also those layers added when using the extended instead of the standard vertical grid. In both cases, the values are well reproduced by the model. When considering all layers, the mean values of the boundary data are lower than those of measurement, the model output actually being closer to the measurement. When considering the new stratospheric layers, the model performance is just as good as it is when considering all layers. The boundary data is now closer to measurements than for all levels. Overall, COSMO is able to reproduce measurements in temperate as well as polar latitudes in all heights, the mean difference never exceeding

185

190 0.5K.




Figure 6. Mean temperature values at each height for the station on Jan Mayen, station no. 20, on the top, and for Madrid, station no. 1, on the bottom, showing results of the run forced by ERA-Interim (left) and NCEP (right). The horizontal lines give the 1σ standard deviation.

The spatial distribution for the run forced by ERA-Interim is shown in Fig. 9, the figure being very similar when looking at the results of the run using the NCEP reanalysis as boundary data. It now becomes clear that the slight outliers of stations 7, 16 and 21 also visible in Fig. 8 are all close to the eastern border of the model domain. By looking at the stations used to examine the problem of Russian humidity data however, it could be shown that this effect is not visible when considering more eastern stations. It is not due to the relative location of the three stations within the model domain but more likely to the measurement data.

Another aspect when comparing the model output to measurements and regridded reanalysis data is the variability of the model in between those times when measurements or reanalysis data is available. Model output was saved every hour, while measurement or reanalysis data is available at most every six hours, as explained in Sect. 2.3 and 3. In order to assess this variability, Fig. 10 shows a shorter time series of only ten days of the three datasets, including all existing model and reanalysis data. It becomes obvious that the model shows an internal variability that is not present in the less frequent measurement or reanalysis data. The greater variability is linked to physical processes




Figure 7. Scatter plot of modelled against measured temperature for polar stations when forcing the model with ERA-Interim (top) and NCEP reanalysis data (bottom). The data was color coded by height to visually inspect the variability in each height section. The statistics in the upper left hand corner refer to the whole dataset.

205 that happen on short timescales of only hours or less. These cannot be captured by regridding the reanalysis data to a finer grid.

4.3 Relative humidity

4.3.1 Excluding the Russian humidity data

When examining the relative humidity of the 24 stations chosen for the validation of the model, it
 210 became apparent that the model could not reproduce the relative humidity data of any station within




Figure 8. Mean difference in temperature over all heights (top) and heights with $z \geq 11$ km (bottom) for each station. The dashed line corresponds in color to the full line is always half the standard deviation of the difference above and below the mean value. See Tab. 3 for a list of the stations corresponding to the numbers.

Russia (or of Gomel, the only station in Belarus with data during the modelled period, as became clear when examining more stations).

As there was no apparent reason for this offset and only 7 stations lay within Russia in the original set (5 polar and 2 temperate), this issue needed further investigation. The data of all available 23 Russian stations well within the model domain and Gomel in Belarus (see Tab. 4) was compared with 24 other stations in the eastern part of the domain but not in Russia or Belarus (see Tab. 5). The result is best illustrated by the mean over all rH values of all ascents in each group. Fig. 11 shows the result for the Russian stations and the 24 stations outside of Russia that had been chosen. While

Meas-COSMO by ERA, 01.10.10 to 31.08.11, all levels




Figure 9. Mean difference of model values and measurements of temperature for each station over all levels when using ERA-Interim as forcing data. The picture is similar when using NCEP reanalysis data.

the model reproduces the values of the stations outside of Russia, the measurement values of those
 220 stations within Russia are very different from the model values but also from the regrided analysis
 or the measurements of those stations outside of Russia.

In addition to the mean, the station Kaliningrad (no. 8), surrounded by the non-Russian stations
 Leba (no. 11), Kaunas (no. 12), Visby (no. 13) and Tallin (no. 16), also allows a spatial investigation.
 While the results of Kaliningrad are similar to the mean of Russian stations, the mean ascents of the
 225 surrounding stations are all similar to the mean of the non-Russian stations.

These two findings are in line with Balagurov et al. (2006) and Moradi et al. (2013). The authors
 of these studies come to the conclusion that the measurement technique used in radio sondes of
 Russia give values for relative humidity that are significantly too high for low pressure. Altogether,
 this lead to the decision to exclude Russian stations from the further investigation of the performance
 230 of COSMO with respect to relative humidity.

4.3.2 Results when excluding Russian data

When excluding the Russian stations (no. 7, 10, 13, 16-18 and 21), 10 temperate and 7 polar stations
 remain to examine relative humidity.




Figure 10. Timeseries of measured and modelled temperature as well as the regridded boundary data, 10km (top) and 23km (bottom) above Scoresbysund, station no. 19, on ten days at the end of January, 2011. All datapoints available in each dataset are included.

The mean values of the ascents of temperate and polar stations for both model runs is given in
 235 Fig. 12. The low stratospheric values are well reproduced by the model for polar and temperate
 stations and both runs, while the tropospheric offset is larger. In heights lower than 13km, the model
 is too humid on average, the values being approximately 10% too high. The mean of tropospheric
 values seems to be better reproduced for polar stations when using the NCEP reanalysis. The bias
 is of measurements and model data is also present in the forcing reanalysis data, these being dryer
 240 than measurements on average. The model reduces this bias and produces a wetter atmosphere than
 that of the reanalyses. So the bias is combination of model physics, boundary data and maybe also
 measurement problems. Overall, model results fit measurements better than the reanalysis data.




Figure 11. Mean relative humidity values of the 23 Russian stations and Gomel (BY) on the top, 24 stations outside of Russia but in the eastern part of the domain on the bottom. The horizontal lines give the 1σ standard deviation.

However, when looking at the scatter plot of the polar stations, given in Fig. 13, it becomes clear that the model is only able to reproduce a mean value that is similar to the measurements. There is no notable correlation in any height. The variability in the measurements is simply too high to be reproduced by the model. This is also visible in the figures showing the mean ascents. The standard deviation of the model and the regrided analysis is much smaller than that of the measurements in stratospheric layers. Fig. 14 shows the timeseries of relative humidity in 10km and 21km height. In 21km height, the values are very low most of the time. While the small scale variations in the




Figure 12. Mean values of relative humidity for polar (top) and temperate (bottom) stations for the model run forced by ERA-Interim (left) an NCEP (right) reanalysis data. Russian stations were excluded from this analysis, as described in the text. The horizontal lines give the 1σ standard deviation.

250 troposphere are not reproduced by the model, the stratospheric variability is well captured by the model.

Fig. 15 shows the spatial distribution of mean $rH_{\text{meas}} - rH_{\text{model}}$ over all layers. The Russian stations have been excluded, but two other stations also show an offset compared to the other stations: Thorshaven (no. 11) and Scoresbysund (no. 23). The modelled values are higher than measurements, 255 with $\Delta rH = 4\%$. This again is probably not an effect of the model, but more likely of the measurements since surrounding stations do not show similar effects. The value fits the range of 2-6% of dry bias reported by Wang et al. (2013) for radio sondes of type Vaisala RS92, but the type of sonde is not known for any of the stations in this study.

260 Relative humidity is on the one side very variable, so that it becomes hard to model exactly, on the other side seems not an easy parameter to measure, as shows the problems first found in Russian data, but apparently also present in the data of other stations.

Similar to examining temperature, a closer look at shorter time period in form of a time series can give information on the internal variability of relative humidity in the model. Fig. 16 shows the




Figure 13. Scatterplots of modelled against measured relative humidity for the run forced by ERA-Interim (top) and NCEP (bottom). The data was color coded by height to visually inspect the variability in each height section. The statistics in the upper left hand corner refer to the whole dataset.

time series of relative humidity at Scoresbysund for ten days at the end of January, 2011. The model
 265 shows a great variability on short time scales that is not present in the other data sets. The coarsely
 time-resolved measurements cannot be used to judge the fluctuations happening in the model on
 short time scales. It becomes understandable that especially relative humidity is difficult to compare
 to radio sonde data, as the variability in the field is just so large that the model cannot be expected to
 reproduce the exact values that were measured at a specific site.




Figure 14. Timeseries of relative humidity in 10 km (top) and 21 km (bottom) height above Jan Mayen for the model forced by ERA-Interim data.

270 5 Sensitivity study

5.1 Boundary forcing interval

This section describes the results of the two model runs that were performed with less frequent boundary forcing of 12 (called int12 in plots) and 24 hours (int24) relative to the other runs with six-hourly forcing (called int6). Both of these runs ran stably and the setups were used to simulate
 275 the same time period as the run with six-hourly forcing.

In order to compare the three runs, Tab. 6 gives the correlation coefficients of model and measured temperature and relative humidity (excluding Russian stations) for all three runs, listed separately

Meas-COSMO by ERA, 01.10.10 to 31.08.11, all levels




Figure 15. Mean difference of measurements and model values of relative humidity for each station when using ERA-Interim as forcing data. The picture is similar when using NCEP reanalysis data.

for polar and temperate stations. The correlation is slightly weaker for both variables with the increased boundary forcing interval, the coefficient becoming smaller as the interval increases. This is expected, as the forcing interval determines how strongly the model is influenced by the boundary values that represent a realistic meteorology. But the decrease is not very strong and measured temperature can still be seen as very well reproduced even by the run that uses only one boundary input field per day.

In addition to comparing each run with measurement data, the runs can be directly compared with one another. For this, the six-hourly time series data that was prepared at each station presents a good database. The difference between the model runs does not increase with simulation time (not shown). The mean difference between the separate stations and a mean of all stations in each height is presented in Fig. 17. In all heights and for both variables, the run with 24-hourly forcing shows a larger difference to the original run than the run with 12-hourly forcing.

5.2 Extending the damping layer

In a second test, the sensitivity of the model to the extent of the damping layer was investigated with an additional model run. For this run, the lower end of the damping layer was set to 22 km (called




Figure 16. Timeseries of relative humidity in 10 km (top) and 23 km (bottom) height above Scoresbysund for the model, the forcing ERA-Interim reanalysis and the measurement data at the end of January, 2011.

rdh22 in plots) 6 km lower than in the original run (rdh28). It then extends one third of total model height of 33 km.

295 Another test run had been planned for which the model height was increased to 42 km, leaving the damping layer as is. This setup ran only for a few days before numerical instabilities lead to the breakdown of the model. The reasons for these instabilities were not investigated further, but this also showcases that it is not a trivial task to find a vertical grid with which the model runs stably.

The setup with *rdheight* = 22.0 on the other hand ran stably for the time period considered in this study. Tab. 7 lists the correlation coefficient of model against measurement data for temperate and polar stations, including all layers up to 21 km. The differences are only marginally small and the runs can be considered to reproduce measurements equally well.

300




Figure 17. Difference between the model runs with 12 and 24-hourly forcing to the original run with 6-hourly forcing for T (top) and rH (bottom). Shown is one profile for each station and the mean of all stations.

In order to assess the difference between the model runs, the six-hourly data generated for each station is again used to calculate a profile of the difference of the two model runs for each station and for the whole dataset. The result of the analysis is shown in Fig. 18. The shapes of the curves are similar to those of Fig. 17, where the boundary input interval was varied. The overall difference is small and similar in magnitude to the difference when doubling the boundary forcing interval to 12 hours. Just where the damping layer starts to be active, a kink is visible in the profile of T , showing the necessity to stop evaluation of the model below the damping layer height when wanting to compare measurements and the model.




Figure 18. Difference between the model run with the lowest extent of the damping layer at 28 km to the standard with $rdheight = 22$ for T (top) and rH (bottom). Shown is one profile for each station and the mean of all stations.

6 Summary and conclusions

This study presents a new, extended vertical grid for the regional model COSMO. The extended grid reaches up to 33 km, almost 10 km above the model top of the standard vertical setup used for the forecast of central Europe by DWD in the domain COSMO-DE. By reducing the magnitude of the damping layer to 5 km, the added layer that can be considered to be free running reaches 28 km, compared to 11 km in the standard setup. This is already well in the lowermost stratosphere.

The extended vertical grid is planned to be used for simulations covering polar spring and the associated ozone loss, which is why it was tested using a domain spreading over central and northern

Europe. To assess the influence of different boundary conditions, two model runs were compared
320 with measurements, using ERA-Interim or NCEP reanalysis as boundary conditions for the model.
Both model runs covered the same period, from October 1, 2010 to September 1, 2011. The model
simulated this period stably. Additionally, three more runs using ERA-Interim as boundary forcing
were done, two with an increased boundary forcing interval of 12 and 24 hours and one with an
increased damping reaching down to 22 km.

325 The output was compared with measurements of temperature and relative humidity from all 12
polar radio sonde stations in the domain and as many in temperate latitudes.

The measurements of temperatures are well reproduced by the model for all stations and heights.
This is not only true for the mean, but also for the comparison of single ascents. The error in heights
above 11 km is even smaller than that when considering all layers, probably because the variability is
330 not as high as when including the tropospheric values. The mean error made by the model is smaller
than 0.5 K for all stations. The boundary data, which was regridded to the output grid, reaches similar
values.

When comparing relative humidity values, it was found that Russian stations (and Gomel in Be-
larus) had systematically submitted higher values. This finding was strengthened by comparing all
335 23 Russian stations in the domain and Gomel to 24 stations not in Russia, but in the eastern part
of the domain and considering model and boundary data. After excluding Russian stations from the
analysis of relative humidity, it became apparent that the model is not capable of reproducing the
exact values of each measurement, and neither is the regridded boundary data. But it does reproduce
the low stratospheric values and fits measurements well when taking a mean over the whole time
340 period. In the tropospheric layers, the model values are more humid than measurements.

The sensitivity study using longer boundary forcing intervals shows how the model reacts to this
factor. The difference to measurements increases with increasing the interval, just like the difference
to the original model run. The stability of the model when using the extended vertical layering does
not depend on short boundary forcing intervals. The results of the run with an increased damping
345 layer height reaching down to 22 km do not differ much from the original setup. The height of the
damping layer does influence the results of the model, but differences reach only about 1 K to the
case of T , for example.

The vertical grid for COSMO presented in this study seems a good alternative to the standard
vertical layering of the COSMO-DE domain when focusing on the upper troposphere and lower
350 stratosphere in polar latitudes. It has been shown to run stably, simulating almost a year. By com-
paring with data from synoptic radio sondes and regridded reanalysis data, it could be shown that
the model is able to reproduce measurements of temperature well and produce reasonable values of
relative humidity. The enlarged time series show a small scale variability in the model that is not
present in the measurements and cannot be expected from regridding the boundary data. The sta-
355 bility against varying the boundary forcing interval and the extent of the damping layer was shown

with three additional model runs. Using this extended vertical grid expands the possible applications of COSMO into the stratosphere. With its high resolution it could be used to study cross-tropopause transport or simulate the chemistry of the lower stratosphere in polar latitudes when also including COSMO-ART.

360 **Appendix A: Model specifications**

This part of the appendix specifies the model setup. It gives the namelist settings for the preprocessor int2lm needed to reproduce the geographic model domain in Tab. 2 and the exact values of the vertical grids - the new, extended grid as well as the standard grid used for COSMO-DE - in Tab. 1.

Appendix B: Specifications of the stations

365 This part of the appendix specifies the stations of which data was used in this study. Tab. 3 lists the information for those stations used for the original study, while Tab. 4 and Tab. 5 list the information of those 48 stations that were used to investigate the bias in relative humidity of the stations in Russia.

Acknowledgements. We acknowledge support by Deutsche Forschungsgemeinschaft and Open Access Publishing Fund of Karlsruhe Institute of Technology.

Table 1. Heights of the layers of the standard and the extended COSMO grid, specified in m.

no.	extended	standard	no.	extended	standard
0	0.00	0.00	31	8711.53	7539.64
1	70.00	20.00	32	9255.31	8080.00
2	151.86	51.43	33	9818.03	8642.86
3	245.82	94.64	34	10399.91	9228.57
4	352.10	150.00	35	11001.17	9837.50
5	470.92	217.86	36	11622.05	10470.00
6	602.52	298.57	37	12262.76	11126.43
7	747.13	392.50	38	12923.55	11807.14
8	904.97	500.00	39	13604.64	12512.50
9	1076.27	621.43	40	14306.25	13242.86
10	1261.25	757.14	41	15028.62	13998.57
11	1460.15	907.50	42	15771.97	14780.00
12	1673.20	1072.28	43	16536.53	15587.50
13	1900.61	1253.57	44	17322.52	16421.43
14	2142.63	1450.00	45	18130.19	17282.14
15	2399.47	1662.50	46	18959.74	18170.00
16	2671.37	1891.43	47	19811.42	19085.36
17	2958.56	2137.14	48	20685.45	20028.57
18	3261.25	2400.00	49	21582.05	21000.00
19	3579.68	2680.36	50	22501.46	22000.00
20	3914.09	2978.57	51	23443.90	
21	4264.68	3295.00	52	24409.61	
22	4631.70	3630.00	53	25398.80	
23	5015.37	3983.93	54	26411.71	
24	5415.92	4357.14	55	27448.57	
25	5833.58	4750.00	56	28509.60	
26	6268.57	5162.86	57	29595.03	
27	6721.12	5596.07	58	30705.08	
28	7191.47	6050.00	59	31840.00	
29	7679.83	6525.00	60	33000.00	
30	8186.44	7021.43			

370 References

Balagurov, A., Kats, A., Krestyannikova, N., and Schmidlin, F.: WMO Radiosonde humidity sensor intercomparison, Instruments and observing methods report No. 85 WMO/TD-No. 1305, WMO, 2006.

Table 2. Namelist parameters of the preprocessor int2lm needed to reproduce the model domain.

namelist block	parameter	value
LMGRID	ivctype	2
	irefatm	2
	lnewVGrid	.TRUE.
	ielm_tot	190
	jelm_tot	255
	kelm_tot	60
	pollat	30.0
	pollon	-170.0
	polgam	0.0
	dlon	0.2
	dlat	0.2
	startlat_tot	-29.0
	startlon_tot	-19.0
	vflat	18000.0
DATA	ie_ext	200
	je_ext	265

Baldauf, M., Förstner, J. F., Klink, S., Reinhardt, T., Schraff, C., Seifert, A., and Stephan, K.: Kurze Beschreibung des Lokal-Modells Kurzestfrist COSMO-DE (LMK) und seiner Datenbanken auf dem Datenserver des
 375 DWD, Tech. rep., DWD, 2011a.

Baldauf, M., Seifert, A., Förstner, J., Majewski, D., Raschendorfer, M., and Reinhardt, T.: Operational convective-scale numerical weather prediction with the COSMO model: description and sensitivities, Monthly Weather Review, 139, 3887–3905, 2011b.

Dee, D. P., Uppala, S. M., Simmons, A. J., Berrisford, P., Poli, P., Kobayashi, S., Andrae, U., Balmaseda,
 380 M. A., Balsamo, G., Bauer, P., Bechtold, P., Beljaars, A. C. M., van de Berg, L., Bidlot, J., Bormann, N., Delsol, C., Dragani, R., Fuentes, M., Geer, A. J., Haimberger, L., Healy, S. B., Hersbach, H., Hólm, E. V., Isaksen, L., Kállberg, P., Köhler, M., Matricardi, M., McNally, A. P., Monge-Sanz, B. M., Morcrette, J.-J., Park, B.-K., Peubey, C., de Rosnay, P., Tavolato, C., Thépaut, J.-N., and Vitart, F.: The ERA-Interim reanalysis: configuration and performance of the data assimilation system, Quarterly Journal of the Royal
 385 Meteorological Society, 137, 553–597, doi:10.1002/qj.828, 2011.

Gantner, L. and Kalthoff, N.: Sensitivity of a modelled life cycle of a mesoscale convective system to soil conditions over West Africa, Quarterly Journal of the Royal Meteorological Society, 136, 471–482, doi:10.1002/qj.425, 2010.

Kalnay, E. and Coauthors: The NCEP/NCAR 40-Year Reanalysis Project, Bull. Amer. Meteor. Soc., 77,
 390 437–471, 1996.

Table 3. Specifications of the stations of which data was used in this study. Stations 1-12 are in temperate, 13-24 in polar latitudes. The international countrycode is also given. Real coordinates are those of the true location, model coordinates those of the closest grid point used to compare measurements and model data.

no.	name	country	WMO no.	lat real	lat model	lon real	lon model	ascents
1	Madrid	ES	8221	40.470	40.494	-3.580	-3.521	654
2	Pratica di Mare	IT	16245	41.650	41.562	12.430	12.537	995
3	Bucharest	RO	15420	44.500	44.554	26.130	26.168	670
4	Stuttgart	DE	10739	48.830	48.796	9.200	9.107	674
5	Legionowo	PL	12374	52.400	52.428	20.970	21.112	671
6	Castor Bay	IE	3918	54.300	54.247	-6.190	-6.178	495
7	Moscow	RU	27612	55.750	55.859	37.570	37.458	633
8	Stavanger	SE	1415	58.870	58.929	5.670	5.735	623
9	Jokioinen	FI	2963	60.820	60.721	23.500	23.588	652
10	Kargopol	RU	22845	61.500	61.441	38.930	38.903	593
11	Thorshavn	DK	6011	62.020	62.007	-6.770	-6.783	651
12	Keflavik	IS	4018	63.970	63.951	-22.600	-22.593	649
13	Kandalaksa	RU	22217	67.150	67.136	32.350	32.366	670
14	Bodo Vi	NO	1152	67.250	67.137	14.400	14.601	651
15	Sodankyla	FI	2836	67.370	67.390	26.650	26.677	663
16	Nar'Jan Mar	RU	23205	67.650	67.662	53.020	52.948	636
17	Sojna	RU	22271	67.880	67.946	44.130	44.126	650
18	Murmansk	RU	22113	68.970	68.963	33.050	33.004	672
19	Scoresbysund	GL	4339	70.480	70.642	-21.970	-22.020	657
20	Jan Mayen	NO	1001	70.930	70.911	-8.670	-8.860	1040
21	Malye Karmakuly	RU	20744	72.380	72.285	52.730	52.609	591
22	Bjornoya	NO	1028	74.520	74.640	19.020	18.792	986
23	Danmarkshavn	GL	4320	76.770	76.759	-18.670	-18.470	644
24	Ny Alesund	NO	1004	78.920	78.994	11.930	11.981	352

Krähenmann, S., Kothe, S., Panitz, H.-J., and Ahrens, B.: Evaluation of daily maximum and minimum 2-m temperatures as simulated with the Regional Climate Model COSMO-CLM over Africa, *Meteorologische Zeitschrift*, 22, 297–316, <http://dx.doi.org/10.1127/0941-2948/2013/0468>, 2013.

395 Moradi, I., Soden, B., Ferraro, R., Arkin, P., and Vömel, H.: Assessing the quality of humidity measurements from global operational radiosonde sensors, *Journal of Geophysical Research: Atmospheres*, 118, 8040–8053, doi:10.1002/jgrd.50589, 2013.

Schättler, U.: A Description of the Nonhydrostatic Regional COSMO-Model Part V: Preprocessing: Initial and Boundary Data for the COSMO-Model, Tech. rep., DWD, 2013.

Table 4. Specifications of the Russian stations of which data was used in this study, liste from south to north. Real coordinates are those of the true location, model coordinates those of the closest grid point used to compare measurements and model data.

no.	name	country	WMO no.	lat real	lat model	lon real	lon model	ascents
1	Voronez	RU	34122	51.670	51.608	39.270	39.392	640
2	Kursk	RU	34009	51.770	51.865	36.170	36.056	603
3	Gomel	BY	33041	52.450	52.595	31.000	30.948	468
4	Suhinici	RU	27707	54.120	53.983	35.330	35.341	587
5	Rjazan	RU	27730	54.630	54.651	39.700	39.578	668
6	Kaliningrad	RU	26702	54.700	54.696	20.620	20.733	442
7	Smolensk	RU	26781	54.750	54.680	32.070	32.131	671
8	Moscow	RU	27612	55.750	55.859	37.570	37.458	633
9	Niznij Novgorod	RU	27459	56.270	56.330	44.000	43.869	654
10	Velikie Luki	RU	26477	56.380	56.450	30.600	30.566	649
11	Bologoe	RU	26298	57.900	57.877	34.050	34.220	639
12	Vologda	RU	27037	59.230	59.217	39.870	39.908	300
13	St. Petersburg	RU	26063	59.970	60.054	30.300	30.348	656
14	Kargopol	RU	22845	61.500	61.441	38.930	38.903	593
15	Sykytyvkar	RU	23804	61.720	61.672	50.830	50.748	668
16	Petrozavodsk	RU	22820	61.820	61.926	34.270	34.313	666
17	Arhangelsk	RU	22550	64.530	64.405	40.580	40.568	296
18	Kem	RU	22522	64.980	65.083	34.800	34.658	645
19	Pecora	RU	23418	65.120	65.044	57.100	57.081	670
20	Kandalaksa	RU	22217	67.150	67.136	32.350	32.366	670
21	Nar'Jan Mar	RU	23205	67.650	67.662	53.020	52.948	636
22	Sojna	RU	22271	67.880	67.946	44.130	44.126	650
23	Murmansk	RU	22113	68.970	68.963	33.050	33.004	672
24	Malye Karmakuly	RU	20744	72.380	72.285	52.730	52.609	589

Schulz, J.-P. and Schättler, U.: Kurze Beschreibung des Lokal-Modells Europa COSMO-EU (LME) und seiner
400 Datenbanken auf dem Datenserver des DWD, Tech. rep., DWD, 2009.

Schwartz, B. and Govett, M.: A Hydrostatically Consistent North American Radiosonde Data Base At The
Forecast Systems Laboratory, 1946-Present, Tech. rep., NOAA, Forecast Systems Laboratory, 1992.

Sissenwine, N., Dubin, M., and Wexler, H.: The U.S. Standard Atmosphere, 1962, Journal of Geophysical Re-
search, 67, 3627–3630, doi:10.1029/JZ067i009p03627, http://dx.doi.org/10.1029/JZ067i009p03627, 1962.

405 Vogel, B., Vogel, H., Bäumer, D., Bangert, M., Lundgren, K., Rinke, R., and Stanelle, T.: The comprehensive
model system COSMO-ART – Radiative impact of aerosol on the state of the atmosphere on the regional
scale, Atmospheric Chemistry and Physics, 9, 8661–8680, doi:10.5194/acp-9-8661-2009, 2009.

Table 5. Same as Tab. 4 but for those stations outside of Russia used to compare to those in Russia.

no.	name	country	WMO no.	lat real	lat model	lon real	lon model	ascents
1	Bucharest	RO	15420	44.500	44.554	26.130	26.168	670
2	Cluj Napoca	RO	15120	46.780	46.839	23.570	23.496	336
3	Poprad	PL	11952	49.030	49.073	20.320	20.240	672
4	Prostejov	PL	11747	49.450	49.337	17.130	17.256	656
5	Prague	CZ	11520	50.000	49.896	14.450	14.589	1341
6	Wroclaw	PL	12425	51.130	51.169	16.980	16.949	668
7	Lin	DE	10393	52.220	52.118	14.120	14.197	1348
8	Legionowo	PL	12374	52.400	52.428	20.970	21.112	671
9	Greifswald	DE	10184	54.100	54.149	13.400	13.399	668
10	Schleswig	DE	10035	54.530	54.599	9.550	9.656	671
11	Leba	PL	12120	54.750	54.747	17.530	17.609	667
12	Kaunas	LT	26629	54.880	54.757	23.880	23.914	336
13	Visby	SE	2591	57.650	57.725	18.350	18.255	594
14	Goteborg	SE	2527	57.670	57.580	12.300	12.237	331
15	Stavanger	NO	1415	58.870	58.929	5.670	5.735	623
16	Tallin	EE	26038	59.450	59.574	24.800	24.733	333
17	Jokioinen	FI	2963	60.820	60.721	23.500	23.588	652
18	Jyvaskayla	FI	2935	62.400	62.346	25.670	25.642	670
19	Sundsvall	SE	2365	62.530	62.610	17.470	17.398	598
20	Orland	NO	1241	63.700	63.599	9.600	9.551	667
21	Lulea	SE	2185	65.550	65.542	22.130	22.085	331
22	Bodo Vi	NO	1152	67.250	67.137	14.400	14.601	638
23	Sodankyla	FI	2836	67.370	67.390	26.650	26.677	663
24	Bjornoya	NO	1028	74.520	74.640	19.020	18.792	986

Table 6. Correlation coefficients for the three model runs forced by ERA-Interim against measurements, using 6, 12 or 24 hourly boundary forcing for polar and temperate stations and both variables, T and rH .

forcing interval	temperate			polar		
	6	12	24	6	12	24
T	0.961	0.957	0.946	0.982	0.979	0.973
rH	0.754	0.740	0.707	0.747	0.735	0.706

Wang, J., Zhang, L., Dai, A., Immler, F., Sommer, M., and Vömel, H.: Radiation dry bias correction of Vaisala RS92 humidity data and its impacts on historical radiosonde data, *Journal of Atmospheric and Oceanic Technology*, 30, 197–214, 2013.

Table 7. Correlation coefficients for the two model runs forced by ERA-Interim against measurements, using 28 km or 22 km as lowest extent of the damping layer for polar and temperate stations and both variables, T and rH .

	temperate		polar	
damp. height	28	22	28	22
T	0.961	0.962	0.982	0.982
rH	0.754	0.758	0.747	0.747

Zahn, A., Christner, E., van Velthoven, P. F. J., Rauthe-Schöch, A., and Brenninkmeijer, C. A. M.: Processes controlling water vapor in the upper troposphere/lowermost stratosphere: An analysis of 8 years of monthly measurements by the IAGOS-CARIBIC observatory, *Journal of Geophysical Research: Atmospheres*, 119, 11,505–11,525, doi:10.1002/2014JD021687, <http://dx.doi.org/10.1002/2014JD021687>, 2014.