

Dear Dr. Tomomichi Kato,

We just uploaded the revised manuscript by M. Fader, W. von Bloh, S. Shi, A. Bondeau, and W. Cramer: “Modelling Mediterranean agro-ecosystems by including agricultural trees in the LPJmL model”.

We have carefully taken the reviewers’ comments into consideration in preparing our revision, and we believe to have responded satisfactorily to all suggestions. Please see below for a point by point response to the reviewers' suggestions and the track marked version of the manuscript.

Please do not hesitate to contact me if you have any questions.

Thank you and kind regards,

Marianela Fader

Institut Méditerranéen de Biodiversité et d'Ecologie marine et continentale (IMBE)
Europole de l'Arbois
Bâtiment Villemin, BP 80
F-13545 Aix-en-Provence cedex 04, France.
marianela.fader@imbe.fr
0033 (0)4 42 90 84 73
<http://www.imbe.fr/marianela-fader.html>

Anonymous Referee #1

Fader and co-authors improved the representation of Mediterranean crops or agricultural trees in a land surface model LPJmL. This makes the model better capable of simulating the crop activity, water cycle and other important environmental variables (such as soil organic carbon) for the Mediterranean region which has important environmental and economic linkages with the rest of the world. Generally I found that the paper is fairly easy to follow and the authors have done a good job in putting the current work in the context of existing crop/agronomic/biogeochemical models. I am convinced that this development is necessary as there is a lack of large-scale vegetation model that is able to simulate both natural vegetation and perennial crops in good detail for the Mediterranean region.

Thank you very much for your comments and suggestions. We will address them point by point below and describe the changed made in the manuscript.

The authors went further to construct a new data set that's needed for their simulation and validated the model simulation in three respects: agricultural yield, irrigation requirement and soil organic carbon. The results section often includes not only the results by the authors themselves but also the comparison with other studies thus seems a little like being mixed with some discussion, but I feel this is a necessary way to organize the material and it facilitates the understanding by readers so quite reasonable.

Thanks for this. We are glad that you see this in the same way we do. We thought that having the comparison next to the results would facilitate the understanding because the reader has in mind the results he just read and can better evaluate the model performance.

The paper represents a fair amount of work. Considering all these points, despite the fact the so-called development in this paper is essentially re-parameterization of existing agricultural trees in the LPJmL model, which differed from the traditional “development” that involves a large amount of coding work and inclusion of new processes, I suggest acceptance of this paper for publishing after the authors address the following (minor) comments.

One general comment on the language: The paper is generally easy to follow, however the language could still be improved. The authors could consider further careful polishing the language by themselves, or inviting a native speaker to improve the overall language quality (I believe the latter is better).

We asked one English-proficient person to go through the paper. He made several small changes that improved the readability and expressions of the manuscript.

For example, on page 5001, Lines 7-8: “Some models are very advanced in other processes, like it is the case of STICS for biochemical cycles”. I doubt the “other processes” should be “some processes”, because there are no processes being discussed before this sentence (When we say other, there must be something to be compared with). “like it is the case of ...” could be abbreviated to “like the case of ...” or “such as the case of ...”. There are other places that the texts could be understood and they don't read fluently.

Some other technical comments:

p4998-L19: pave → paves

p4999-L2: commonly → often

p4999-L3: strengthen → strengthened

p4999-L5: as well → as well as

p4999-L15: like → such as

1 p5000-L6: apples → apple
2 p5001-L4: revision → review?
3 p5005-L4: proportional → proportionally?
4 P5015-L17: Summarizing → To summarize
5 p5016-L21: besides France → except France
6 p5018-L6: establishing → by establishing
7 p5019-L10: in this moment → at this moment
8 p5019-L28: will not be → are not?
9
10 ***All changed, thanks.***
11
12 p4999-L9: “is having already at present ...”, complex expression hard to understand, please
13 rephrase.
14
15 ***We rephrased to: "Environmental degradation involving soil erosion, biodiversity loss and***
16 ***pollution is negatively affecting natural and societal systems and is expected to intensify***
17 ***even more in future due to urbanization, industrialization and population growth (Doblas-***
18 ***Miranda et al., 2015; Lavorel et al., 1998; Scarascia-Mugnozza et al., 2000; Schröter et al.,***
19 ***2005; Zdruli, 2014)."***
20
21 p5006-L11: Please define this “mean absolute percent error”.
22
23 ***We did: "Our results are in good agreement with their numbers with a mean absolute***
24 ***percent error of 26% (MAPE, calculated as the sum of percentage differences between their***
25 ***values and our values, divided by their values, and finally multiplied by 1 over the sample***
26 ***size)."***
27
28 p5011-L25: Again, please define “Willmott coefficient” and justify why the authors chose it.
29
30 ***In order to keep the flow of the sentence this was added as a foot note: " The Willmott***
31 ***coefficient was developed by Willmott (1982) as a tool for testing model performance***
32 ***against independent data and is calculated by $1 - \frac{\sum(p-o)^2}{\sum(p-\bar{o} + o - \bar{o})^2}$, where o is the***
33 ***independent data, p the LPJmL simulated data, and \bar{o} denotes the mean of independent***
34 ***data."***
35
36 p5015-L3: What's the source of this “independent data” and why we should trust it?
37
38 ***Sentence was completed: "Despite the difference in the period of time analysed and the***
39 ***methodology, Fig. 4 shows that our results agree well for Israel and Spain for the***
40 ***independent data (from the water commissioner of Israel and the executive secretary of the***
41 ***International Commission of Irrigation and Drainage, respectively), while they found their***
42 ***values to be overestimated for both countries"***
43
44 p5015-L24: “in a unit of area with natural and agricultural vegetation”, this is difficult to
45 understand, do you mean “in an area mixed with natural and agricultural vegetations”?
46
47 ***Rephrased to: "This is the case for carbon sequestration by soils in a unit of area that has***
48 ***both natural and agricultural vegetation –a very important variable for the climate debate***
49 ***and the ecosystem service research domain."***
50

P5003: Methods, the 2nd and 3rd paragraph. Maybe add one more paragraph to give more details on how the simulation is done. Did you do a spin-up run then followed by a full transient run? And also, after reconstructing the land use data by merging MMR with HYDE in 2.1, did you apply the land use change when doing the simulation?

Following sentence was added: "The model was spun-up for 5000 years with dynamic natural vegetation in order to bring the carbon pools in equilibrium and additionally 390 years with natural and agricultural vegetation. The spin-up simulations were followed by a transient run from 1901 to 2010 using the land use patterns described in the next section."

P5019, section 4.2, first paragraph: many of the applications listed here could be achieved by other land surface models, though they cannot be done by agronomic models as pointed out by the authors. Also perhaps only the these applications are limited to Mediterranean regions, LPJmL has the advantage thanks to the developments presented in this paper. Please point out these two points, otherwise the claim "The inclusion of perennial crops in LPJmL presented in the current study opens up the possibility for a number of large-scale applications and research studies ..." is not completely correct.

A sentence was added: "Some of these applications could also be performed by land surface models, but LPJmL has now the advantage of considering perennial crops in detail, which allows more precise quantifications not only in the Mediterranean region, but also in other macro-regions where agriculture is partially dominated by tree crops, such as Australia, South Africa, Chile, Western Argentina and California"

Figure 1: I guess the unit "FM" for the yield means "fresh matter", but please explain this explicitly.

We added this to the caption and in the following sentence: "Figure 1 shows LPJmL simulated yields in metric tonnes fresh matter per hectare for the calibrated run for all new crops where FAOSTAT had data in the Mediterranean region, averaged for the period 2000-2009"

Figure 3, Figure S3: Although the colorbars currently used have similar colors for small values around zero that facilitate the interpretation of the figure. However one cannot easily distinguish positive versus negative values. Please think to change to some contrasting colorbars that have different colors for positive and negative values (now they're both bluish or greenish).

The scales were changed to contrasting colours (light blue and light yellow).

Anonymous Referee #2

Received and published: 25 August 2015

Review of paper: "Modelling Mediterranean agro-ecosystems by including agricultural trees in the LPJmL model" by Fader et al.

This is an especially well written paper, presented in a very thoughtful way, and should be published with little change. It carefully combines together model projections, tests them against comprehensive data, and illustrates the implications geographically. The reference list is long and well-considered, and in that context, serves to bring papers to the debate of environmental change that might otherwise be missed.

Thank you very much for your comments and suggestions. We will address them point by point below and describe the changes made in the manuscript.

Much of the Methods is devoted to determining accurate known patterns of land use for the contemporary period. In a changing climate, these areas can be expected to change geographically. Maybe the authors could add a couple more sentences in Discussion, as to how an interactive assessment of future land-use can be calculated (rather than relying on its prescription).

A sentence was added: "Another input-related issue is the need of scenarios of future crop patterns as a consequence of climatic and socio-economic change. With climate change very likely affecting the potential growing areas of agricultural trees and their profitability, studies aiming at a future quantification of agricultural, biochemical and hydrological variables would profit from coupling between models like LPJmL and land use models."

An especially interesting feature of this paper is that it introduces so many other aspects of environmental consideration, besides the climate change angle. This includes matters of biodiversity, pest-control and irrigation. Would the authors be willing to write a little more in the Discussion section as to how knowledge of some of these factors could aid stabilisation of food provision in a changing climate? In other words, better knowledge of crop response to other drivers – as this paper presents – might aid adaptation planning.

The first paragraph of section 4.2. was completed and rephrased for accounting for this: "The inclusion of perennial crops in LPJmL presented in the current study opens up the possibility for a number of large-scale applications and research studies that cannot be performed with empirical and/or input-intensive agronomic models. These include assessments of climate change impacts on hydrological variables, agricultural production and carbon sequestration as well as evaluation of consequences of land use change (including expansion of irrigated areas). Some of these applications could also be performed by land surface models, but LPJmL has now the advantage of considering perennial crops in detail, which allows more precise quantifications not only in the Mediterranean region, but also in other macro-regions where agriculture is partially dominated by tree crops, such as Australia, South Africa, Chile, Western Argentina and California. Moreover, having a more accurate representation of perennial crops also allows studies on shifts in suitable growing areas for agricultural trees, diversity of diets, resilience of agricultural systems, needs for climate change adaptation and implications for food security as well as assessments on ecosystem services provided by perennial cultures, for example habitat provision for avifauna. "

Where there remains large uncertainties in parameter settings, it might be appropriate to re-iterate these in the Discussion. So for example, lines 24, 25, p5007, where the time-before-harvest is difficult to predict due to issues of fertilisation and available water use. And for other similar examples through the paper, as this will enable the Discussion to highlight what is next required to improve models further.

In the paragraph about parameters uncertainties and prospects for further refinements we included this sentence: "Another benefit of more precise management information would

be the possibility of differentiating parameters that are assumed to be static and equal for all plants in the present study, such as the number of years that a perennial plantation stays in production before being renewed and the period of time from the planting until the first harvest (see section 2.2)."

This paper may have relevance regarding refugee movements, which clearly is an issue at the moment, with many attempting to leave the Southern shores of the Mediterranean. Although the current movement of people is mainly due to other reasons, it is not inconceivable that future levels of climate change could trigger similar events, should it aggravate food scarcity issues. In that context, this paper has particular importance by creating a model capable of projecting different crop yields. So a sentence, in general terms, about better predictive modelling of food security aids global planning for potential matters of migration (i.e. its possible prevention, if due to well considered avoidance of poor food levels) may make this paper also highly relevant in socio-economic discussions?

We modify a sentence to mention the issue of migration and added another one to point out possible applications in the direction of food security:

"Timely and appropriate coping with the combination of these challenges will need collaboration between the Mediterranean countries and local communities in a number of issues, including advanced development of adaptation options, common plans on energy transition, environmental policy and best-practice rules in nature conservation and agricultural management. Collaboration will have to be designed in a framework that allows taking into account the larger European and global picture in terms of environmental change, dynamics of ecological systems, foreign investments and migration movements. [...]"

"Moreover, having a more accurate representation of perennial crops also allows studies on shifts in suitable growing areas for agricultural trees, diversity of diets, resilience of agricultural systems, needs for climate change adaptation and implications for food security as well as assessments on ecosystem services provided by perennial cultures, for example habitat provision for avifauna. "

A line in the Conclusions about feasibility of routinely seeing LPJmL, and with these new agricultural crops, either directly at the bottom of a GCM or forced with CMIP5 diagnostics, would be helpful. Is interactive prescription of land use possible, and maybe the next step in LPJmL development? (i.e. where climate is projected to change, new/lost areas for the crops of this paper can be determined). I realise the paper is predominantly about the contemporary period, but much of it has multiple direct implications for impacts of different scenarios (e.g. RCP85 vs RCP60: : :.)

Section 4.2. names now a lot of applications about these issues. Moreover, one additional sentence refers now specifically to the issue of land use patterns. The way we imagine this is rather a coupling with land use models that have socio-economic variables:

"Another input-related issue is the need of scenarios of future crop patterns as a consequence of climatic and socio-economic change. With climate change very likely affecting the potential growing areas of agricultural trees and their profitability, studies aiming at a future quantification of agricultural, biochemical and hydrological variables would profit from coupling between models like LPJmL and land use models."

Another sentence refers to an example of an application with GCMs: "A first application of this model development was presented by Fader et al. (2015), pointing out that irrigation water needs of perennial crops in the Mediterranean region might increase significantly under climate change and some countries may face constraints to meet the higher water demands."

Minor points: There are a few very small typos in places, so another read of the manuscript before publication (or by the editors) would be beneficial. Just very small things e.g. "paves" not "pave" line 19, p4998.

Some of the paragraphs are very long, and might just be easier for the reader if they are split up a bit e.g. paragraph starting "In this context the need to perform Mediterranean-wider assessments..."

We gave the manuscript to one English-proficient person that went through it and improved the phrasing in several parts of the manuscript.

Moreover we cut up 2 long paragraphs into 2 pieces. The ones starting with "The first step for the compilation of the land use dataset [...]" and "This procedure yields a gridded, global dataset at 30 arc minutes spatial resolution [...]" However, we decided to let the one starting with "To support adaptation and mitigation efforts for climate change and environmental degradation [...]" as one long paragraph in order to cluster there the complete literature review.

The use references is impressive, but is there a single reference that lists the majority of the Equations? Hence as a temporary measure until the documentation (as mentioned in Section 5) is available?

Unfortunately, there is no single publication that lists all the features, modules and function of the model at this moment. This is why we decided to include a bit of the development history in the introduction to allow the reader to know the publications related to every part of the model. At present we are working not only on the model documentation but also on a new LPJmL-paper that should include all developments until present.

Line 28, p5002. Maybe to help the reader, state what "a new input dataset" contains, i.e. what is need to drive the model. Obviously meteorological conditions, but what else. This is important should anyone be considering potential application of this model in an Earth System Model. [OK, can see this is given beginning of Section "Methods"]

That sentence was completed as: "The following section outlines the methodology applied, including the compilation of a new input dataset with land use patterns which was needed for the validation and application of the model, the description of the modelling approach and parametrisation of the new crops, as well as the computation of irrigation requirements and soil carbon densities"

In the methods at the end of the first paragraph the required inputs are named: "Required inputs are: a) gridded, monthly climate variables (temperature, cloudiness interpolated to daily values and precipitation and rainy days converted through a weather generator in daily values); b) atmospheric CO₂ concentrations; c) gridded soil texture as described in Schaphoff et al. (2013); d) a gridded dataset of land use patterns prescribing which crops are grown where and whether they are irrigated or rain-fed. "

Section 2.3. Give units for all quantities (or if dimensionless, then say that or show units as [.]). I assume it is standard for this area of science to give variable names as acronyms e.g. WHC. As opposed to a Greek character in equations, with “WHC” as a subscript for instance?

Yes, it is standard for this area to use acronyms.

We added the units as you suggested:

"[...]

fRil [fraction] is the proportion of roots in the irrigated layer.

wil [fraction] is the water content in the irrigated layer.

wr [fraction] is the water content weighted with the root density for the soil column.

[...]

"

Some of the outliers in Fig 1 are explained in the main text (p5012). Is there anything from this that might also be appropriate in Discussion i.e. what is needed to get predictions for contemporary periods even more accurate, especially where the model currently fails?

The section 4.2 details the potential improvements related to input, parameters and inclusion of processes. Following paragraph, which was extended, refers specially to the outliers of Fig. 1:

"For national-scale studies re-parametrisation with different representative plants for the groups (e.g. using hazelnuts instead of almonds for nuts trees Turkey) is possible without any difficulty. For this group of improvements data on management is essential, including harvesting times, post-harvest uses, planting densities, planted varieties, etc. Another benefit of more precise management information would be the possibility of differentiating parameters that are assumed to be static and equal for all plants in the present study, such as the number of years that a perennial plantation stays in production before being renewed and the period of time from the planting until the first harvest (see section 2.2)"

Fig 1, is there some way of showing which countries the circles refer to.

We do show the country names for the outliers. We tried to put them in every bubble but the overlap avoids understanding almost all names and decreases the overview of the positions of the bubbles. So we prefer to keep it like it is, i.e. country names only for outliers.

Also, is the size of yields relative to the radius, or the area of the “bubbles”?

To the radius. We added this to the caption.

Only if the authors have time, but is it possible to tidy up the maps so they are on a similar projection? For instance, Figure 6, looks “squashed” in the latitude direction. Fig 3 looks on a slightly different projection to Fig 5.

Actually the projection in the plotting program is the same, the inclusion of broader and thinner legends had squeezed the figures. So we separated legends and figures to solve this.

Modelling Mediterranean agro-ecosystems by including agricultural trees in the LPJmL model

Fader, M.¹; von Bloh, W.²; Shi, S.³; Bondeau, A.¹; Cramer, W.¹

[1] Institut Méditerranéen de Biodiversité et d'Ecologie marine et continentale. Aix-Marseille Université, [CNRS, IRD, Avignon Université, Technopôle Arbois-Méditerranée UMR-CNRS 7263 / IRD-237 / UAPV](#), Bâtiment Villemin, ~~Europole de l'Arbois~~ BP 80, F-13545 Aix-en-Provence cedex 04, France.

~~[2] Laboratory of Excellence OT Med. Europôle Méditerranéen de l'Arbois, Bâtiment Gérard MEGIE, Avenue Louis PHILIBERT, 13857 Aix-en-Provence Cedex 3, France.~~

[3] Potsdam Institute for Climate Impact Research, Telegraphenberg, [D-14473](#) Potsdam-~~D-14473~~, Germany.

[4] Research Software Development Group, Research IT Services, University College London. Podium Building (1st Floor), Gower Street, London WC1E 6BT, United Kingdom

Abstract

~~In the Mediterranean region, Climate-climate~~ and land use change ~~in the Mediterranean region~~ ~~is~~are expected to ~~affect impact on~~ natural and agricultural ecosystems by ~~warming, reduced rainfall, direct degradation of ecosystems and decreases in precipitation, increases in temperature as well as biodiversity loss and anthropogenic degradation of natural resources.~~ ~~Human population Demographic~~ growth ~~and socioeconomic changes, notably in on~~ the Eastern and Southern shores, will require increases in food production and put additional pressure on agro-ecosystems and water resources. Coping with these challenges requires informed decisions that, in turn, require assessments by means of a comprehensive agro-ecosystem and hydrological model. This study presents the inclusion of 10 Mediterranean agricultural plants, mainly perennial crops, in an agro-ecosystem model (LPJmL): nut trees, date palms, citrus trees, orchards, olive trees, grapes, cotton, potatoes, vegetables and fodder grasses.

The model was successfully tested in three model outputs: agricultural yields, irrigation requirements and soil carbon density. With the development presented in this study, LPJmL is now able to simulate in good detail and mechanistically the functioning of Mediterranean agriculture with a comprehensive representation of ecophysiological processes for all vegetation types (natural and agricultural) and in a consistent framework that produces

estimates of carbon, agricultural and hydrological variables for the entire Mediterranean basin.

This development paves the way for further model extensions aiming at the representation of alternative agro-ecosystems (e.g. agroforestry), and opens the door for a large number of applications in the Mediterranean region, for example assessments on the consequences of land use transitions, the influence of management practices and climate change impacts.

1 Introduction

The Mediterranean region is a transitional zone between the subtropical and temperate zones with high intra- and interannual variability (Lionello et al., 2006). This region has been identified as one of the regional climate change hotspots, with a high likelihood of experiencing more frequent and more intensive heat waves, ~~often commonly~~ combined with and strengthened by more intensive and longer droughts (IPCC, 2012; Kovats et al., 2014, Diffenbaugh and Giorgi, 2012). This will likely have adverse implications for the food and energy producing sectors as well as for human health, tourism, labour productivity and ecosystem services (Kovats et al., 2014; Skurras and Psaltopoulos, 2012). However, climate change is only a part of the future challenges that the Mediterranean region will face in the near future. Environmental degradation involving soil erosion, biodiversity loss and pollution is negatively affecting having already at present detrimental effects for natural and societal systems and is expected to intensify even more in future due to urbanization, industrialization and population growth (Doblas-Miranda et al., 2015; Lavorel et al., 1998; Scarascia-Mugnozza et al., 2000; Schröter et al., 2005; Zdruli, 2014).

Most aspects of Climate-change and environmental degradation will strongly affect the Mediterranean agricultural sector directly. This Agriculture sector plays a very important role, not only for food security within the region itself, but also through its economic integration in other regions, such as through the significant export of products to like the rest of Europe (Hervieu, 2006). Agriculture plays therefore a key role for This importance is based on the influence of the sector in the national economies, making a part of the dependence of rural population rely on it upon agriculture for their livelihood and creating the linkages with other issues and sectors, such as food security, culture and tourism (Verner, 2012, Hervieu, 2006). Human population Demographic growth and socioeconomic developments in on the Eastern and Southern shores, as well as the already high dependence of the region on the international food markets will increase the need for local food production. Additionally, and potential resource allocation trade-offs, (especially for water and land), will put Mediterranean agriculture under increased pressure, calling for more and more efficient production practices (Verner, 2012; World Bank, 2013).

To support adaptation and mitigation efforts for climate change and environmental degradation, In this context the need to perform Mediterranean-wide assessments on of the state of agriculture and the likely consequences of global change becomes evident are required. These would have to be complemented by analyses on the potential developments and future

difficulties of the agricultural sector and its interactions with the environment. The large scale character of such assessments and the necessity of looking into possible future scenarios require the utilization of modelling tools that cover the essential characteristics of the dominant agro-ecosystem models in the region. However, ~~a~~At present, ~~there is no~~ suitable modelling framework for this task exists. Given the range of conditions in the region, such a tool should be large-scale, process-based and integrated modelling framework for the major crop types agriculture, grasslands and natural vegetation, taking into account the carbon cycle and hydrology that considers the special structure of Mediterranean agriculture, which is largely dominated by perennial crops of them. Notably, the presence of perennial, woody species is a characteristic of Mediterranean agro-ecosystems and they deliver 45% of agricultural outputs (Lobianco & Esposti, 2006). Most large ~~Existing agro-ecosystem or~~ crop model ~~families~~ have implemented some tree crops and in some cases applications in Mediterranean environments (mostly small scale) were published. For example the crop model STICS has been used is able to simulate the growth of vineyards and apple trees (García de Cortázar-Atauri, 2006, Nesme et al., 2006, Valdés-Gómez et al., 2009); and in the CropSyst model pears, apples, vineyards and peaches are included (Marsal et al., 2013, 2014, Marsal and Stöckle, 2012). Other ~~large~~ modelling frameworks offer general and specific formulation for horticultural systems that have been applied in other regions, mainly in Anglo-Saxon countries. This is the case of the EPIC/SWAT/SWIM families (Neitsch et al., 2004, Gerik et al., 2014) for cotton and apple, and the APSIM model, for cotton and vineyards (Holzworth et al., 2014). ~~In California,~~ another region with Mediterranean climate, there is a dynamic modelling community assessing climate change impacts on horticulture by process-based (Gutiérrez et al., 2006) and empirical models (Lobell et al., 2007). At the global scale. The the GAEZ model approach offers potential growing areas for citrus, olives and cotton (IIASA/FAO, 2012) at global scale. The GCWM model is probably the most complete model in terms of perennial crops, comprising citrus, cotton, date palm and grapes (Siebert & Döll, 2008, 2010). Other ~~smaller communities and single scholars~~ have presented developments of models for fruit trees, such as the inclusion of like it is the case of kiwi, vineyards and apples in the SPASMO model (Clothier et al., 2012), walnuts in CAN-WALNUT (Baldocchi & Wong 2006) and date palm by Sperling (2013), and the A model by Villalobos et al. (2013) focused focuses on transpiration of apricot, apple, citrus, olive, peach, pistachio and walnut trees.

The ~~for~~ goals of these applications are diverse, going from simple reproduction of experiments, over epidemiological analysis (causes and patterns of diseases), to the simulation of potential phenological changes, and the influence of management practices on agricultural production. Concerning the and impacts of climate change. ~~Regarding the later point,~~ Moriondo et al. (2015) presented a detailed review on empirical and process-based models for olives and vineyards. They concluded that process-based models are better suited for climate impact studies but they have to be completed and improved to account for the perennial nature of these crops, the effect of higher CO₂ in the atmosphere, dynamic growing periods and the effect of management practices.

~~The revision r~~ Reviewing the existing studies reveals of literature shows two important points. First, there is no single model or model family comprising all major agricultural plants of the

Mediterranean region. Second, there is no model combining dynamic simulation of natural vegetation and agro-ecosystems for the Mediterranean region. Some models are very advanced in ~~some other~~ processes, like ~~it is~~ the case of STICS for biochemical cycles. ~~And someOther~~ models have unique features, such as the detailed consideration of hydrology in the unsaturated zone, salt and leaching transport in the WOFOST model coupled with the SWAP and PEARL models (Kroes & Van Dam, 2003). ~~But, as written above, there is a lack of models combining dynamic simulation of natural vegetation and agro-ecosystems for the Mediterranean region.~~ The CENTURY model, which focuses on soil organic carbon computation, offers a forest module and general formulations that can be adapted to horticulture but only one small-scale application was presented for the Mediterranean region (Álvaro-Fuentes et al., 2011). Without the integrated modelling of natural vegetation and agro-ecosystems in a comprehensive framework, there are many questions that cannot be answered. Notably, some of them are of extreme relevance for the Mediterranean region and concern water requirements and availability for the agricultural sector, sustainable food production potentials under climate change, environmental consequences of land use (including biodiversity loss), and soil carbon sequestration patterns, including responses to land use change.

To better assess the potential responses of Mediterranean agro-ecosystems to these forcings, we have extended ~~In this context we present here~~ the ~~improved~~ representation of Mediterranean agriculture in the Lund-Potsdam-Jena managed land model (LPJmL). LPJmL ~~was the result of including agriculture in a dynamic global vegetation model—the LPJ model—which had been designed and developed based on the BIOME model family. The LPJ objectives were to compute the dynamics distribution of natural vegetation, annual crops and natural grasslands, by considering as well as carbon pools and fluxes, hydrological variables, and coupled photosynthesis and &transpiration (Sitch et al., 2003, Gerten et al., 2004, Bondeau et al., 2007). The model has undergone~~ This model was further developed by Bondeau et al. (2007) into the LPJmL model by including agro-ecosystems (annual crops and managed grassland). Major developments and validation efforts were also undertaken with respect to in the hydrology of the model, including a river routing and irrigation scheme (Rost et al., 2008), the management of dams and reservoirs (Biemans et al., 2011) and a five soil layer hydrology (Schaphoff et al. 2013). The representation of agricultural systems has also been improved by including bioenergy systems (tree and grass bioenergy plantations, jatropha, sugar cane, Behringer et al., 2008, Lapola et al., 2009). ~~—a~~ An agricultural management module has been added, representing the combined influence of management practices on plant and stand development (e.g. fertilizer inputs, mechanization, use of high-yielding varieties, weed and pest control, etc., Fader et al., 2010), and better representation of sowing dates and multiple cropping systems (Waha et al., 2012, 2013). ~~to name some examples.~~ LPJmL is widely recognized as a state-of-the-art agro-ecosystem and hydrology model and ~~It has undergone the broadest possible range of validation efforts against experimental and observational data.~~

Consequently, LPJmL is a state-of-the-art agriculture and hydrology model. In a recent intercomparison study of global hydrological models and global gridded crop models ~~on for the~~ the assessment of future irrigation water availability, Elliot et al. (2014) indicated that LPJmL

is unique in that it ~~falls-performs well into~~ both categories. LPJmL is intensively used at the global and macro-regional scale in various research fields, particularly for questions related to future food security, land use change, and adaptation to climate change ([Gerten et al., 2008](#), [Rost et al., 2009](#), [Lapola et al., 2010](#), [Fader et al., 2010, 2013](#), [Waha et al., 2013](#), [Müller et al., 2014](#)).

~~Since However, most of the~~ Mediterranean-specific crops ~~were-have been~~ lacking in LPJmL. Thus, we ~~here present in this study~~an extension that ~~the inclusion includes of~~ 10 new crop ~~classes-functional types~~ that are especially important in the ~~Mediterranean~~ region. ~~Most of these may be called, being most of them~~ “agricultural trees”: nut trees, date palms, citrus trees, orchards, olive trees, grapes, cotton, potatoes, vegetables and fodder grasses. ~~Their inclusion made possible to account for ~88% of irrigated areas of the Mediterranean instead of ~50%. The relevance of the development is then demonstrated by presenting and comparing the resulting simulations on agricultural yields, irrigation requirements and soil carbon densities in the Mediterranean region.~~

The ~~next-following~~ section ~~comprises-outlines~~ the methodology applied, including the compilation of a new input dataset ~~with land use patterns~~ which was needed for the validation and application of the model, the description of the modelling approach and parametrisation of the new crops, as well as the computation of irrigation requirements and soil carbon densities. The results section details exemplarily the performance of the model in simulating yields, soil carbon and irrigation water requirements. Finally, the paper is closed by a discussion on perspectives for future developments, potential applications and further refinements.

2 Methods

~~As a function of climatic conditions and agricultural management,~~ LPJmL simulates, spatially ~~-explicitly~~ and at a daily to yearly temporal resolution, growing periods (sowing and harvest dates), net and gross primary productivity, carbon sequestration in plants' compartments and soil, ~~agricultural production,~~ heterotrophic and autotrophic respiration, ~~agricultural production,~~ as well as a number of hydrological variables, such as runoff, soil evaporation, plant transpiration, plants' interception, percolation, infiltration, river discharge, irrigation water requirements, water stress and soil water content.

~~Required LPJmL-~~inputs ~~consist in:~~ a) gridded, monthly climate variables (temperature, cloudiness interpolated to daily values and precipitation and rainy days converted through a weather generator in daily values); b) ~~global-atmospheric~~ CO₂ concentrations; c) gridded soil texture as described in Schaphoff et al. (2013); d) a gridded dataset of land use patterns prescribing which crops are grown where and whether they are irrigated or rain-fed.

~~For the present study, we used Climate-climate~~ inputs at 30 arc minutes spatial resolution and global CO₂ concentrations ~~were~~ derived from the CRU 3.10 datasets ([Harris et al., 2014](#)). ~~The~~

land use patterns needed for the implementation of the new Mediterranean crops in LPJmL had to be compiled from different sources, as explained in the following section. The model was spun-up for 5000 years with dynamic natural vegetation in order to bring the carbon pools in equilibrium and additionally 390 years with natural and agricultural vegetation. The spin-up simulations were followed by a transient run from 1901 to 2010 using the land use patterns described in the next section. ~~The land use patterns needed for the implementation of the new crops in LPJmL had to be compiled from different sources, as explained in the following section.~~

2.1 New land use patterns for used in LPJmL

LPJmL needs irrigated and rain-fed physical (as opposed to harvested) areas for each simulated crop. Physical and harvested areas differ through multiple cropping practices, i.e. when one area is used twice in a year the harvested area is double as large as the physical area. For the present model development a new land use dataset had to be compiled from different sources: Portmann et al. (2011), hereafter *MIRCA*, Monfreda et al. (2008), hereafter *MON*, Klein Goldewijk et al. (2011), hereafter *HYDE*, and Ramankutty et al. (2008), hereafter *RAM*.

The first step for the compilation of the land use dataset was determining the harvested areas of all LPJmL classes, including the new Mediterranean crops, for the present time. For crops present in *MIRCA*, which differentiate between irrigated and rain-fed areas, we took the all values in that study were used directly. For the missing crops, *MON* corresponding classes (see Table S1) were compared at grid-cell level with the *MIRCA* classes "other perennial" and "other annual", thereby to splitting the harvested areas into the rain-fed and irrigated part. This procedure was done for olives, non-citrus orchards, nuts trees and vegetables. For the first three groups, the *MIRCA* class "other perennial" was used for the grid-cell specific splitting between rain-fed and irrigated areas. In the case of For vegetables, rain-fed and irrigated areas were derived through comparison with the class "other annual".

Large inconsistencies were found at the grid-cell level between *MIRCA* and *MON*, for example cases where a single crop area in *MON* is was larger than the sum of the rain-fed and irrigated corresponding "other" classes in *MIRCA*. And Also the absence or small extent of the class irrigated "other perennial" in areas that intuitively may be assumed as irrigated, like it is in the case of orchards and olives in Egypt. The author of *MIRCA* (Felix F. Portmann, personal communication, 2013) assumes that most of these inconsistencies are due to issues of scale, being the grid-cell differences potentially large but presenting a good agreement at the administrative level. Grasslands, representing meadows, were taken directly from *RAM* and assumed to be always rain-fed.

After deriving harvested areas for around the year 2000, the calculation followed the flow chart shown in Fig. S1. Harvested areas were compared with cropland and grassland areas from *RAM* in order to exclude multiple cropping and derive physical cultivation areas.

Mis en forme : Allemand (Allemagne)

Decadal cropland data from *HYDE* were interpolated to derive annual values and then used for extrapolating the land use patterns of ~2000 to the past, until 1700 (see below, Equations 1 to 5). Historical irrigation fractions were determined as explained in Fader et al. (2010).

Due to inconsistencies between *HYDE* and the dataset combining *MIRCA*, *MON* and *RAM* (hereafter *MMR*), proportionally changing cell fractions using *HYDE* historical trend would have given an unrealistic global-overall trend of cropland areas. For this reason crop-specific bias corrections had to be performed in between, as follows:

First, the global (*g*) area difference (*D*) was calculated:

$$D_{2000} = HYDE_{g,2000} - MMR_{g,2000} \quad (1)$$

Bias correction of *HYDE* global values was performed by:

$$HYDE_{g,y,bias_corrected} = HYDE_{g,y} - D_{2000} \quad (2)$$

Where *y* represents years from 1700 to 2000.

Cell (*c*) bias correction was done by:

$$HYDE_{corrected,c,y} = \frac{HYDE_{c,y}}{HYDE_{g,y}} * HYDE_{g,y,bias_corrected} \quad (3)$$

Proportional temporal change of *MMR* cell values was done from 2000 backwards by:

$$MMR_{y,c,proportional} = \frac{HYDE_{corrected,c,y}}{HYDE_{corrected,c,y+1}} * MMR_{y,c} \quad (4)$$

And final cell values (LU_{LPJmL}) were calculated by:

$$LU_{LPJmL} = \frac{HYDE_{corrected,g,y}}{MIRCA_{proportional,g,y}} * MMR_{y,c,proportional} \quad (5)$$

Cell fractions from 2001 to 2010 follow the trend between 1950 and 2000.

This procedure yields a gridded, global dataset at 30 arc minutes spatial resolution of the cultivation areas of 24 crops from 1700 to 2010. Table 1 shows the resulting areas for each crop class of LPJmL. Expanding LPJmL for modelling Mediterranean crops made possible to account for ~88% of irrigated areas of the Mediterranean instead of ~50%. For rain-fed areas the improvement was from ~21% to 73%. The remaining areas are mostly fallow land and were included in the model class "others" which is parametrised as grasslands.

There is a general lack of information about the planting areas for agricultural trees at national and subnational level. Nevertheless, it was possible to make two comparisons. First, EUROSTAT offers information about olive trees areas at country level for 8 countries of the Northern Mediterranean. Our results are in good agreement with their numbers with a mean absolute percent error of 26% (MAPE, calculated as the sum of percentage differences

1 | ~~between their values and our values, divided by their values, and finally multiplied by 1 over~~
2 | ~~the sample size~~) of 26%. Second, national harvested areas as reported by FAOSTAT for
3 | dates, olives, cotton seed, grapes and potatoes as an average of 2000 to 2009 could be
4 | compared with our dataset. The agreement is high (MAPE <30%) for all classes except ~~foring~~
5 | olives and dates (MAPE 47% and 46% respectively) where our dataset has mainly smaller
6 | areas. This is due to the fact that *MIRCA*, *RAM* and *MON* have been compiled for the year
7 | 2000 and the FAO shows strong an accelerated expansion of areas from 2000 to 2010, for
8 | example 54% for olives in Morocco, 45% for dates in Egypt and Turkey. Overall the input
9 | dataset compiled here has no better alternatives and ~~is appears to be broadly~~ suitable for
10 | applications until newer versions of the land use data used as sources are released.

11 | 2.2 Implementation, calibration and parametrisation of Mediterranean agricultural 12 | trees and crops

13 | 12 crops were already present in LPJmL (temperate cereals, rice, tropical cereals, maize,
14 | temperate roots, tropical roots, pulses, rapeseed, soybeans, sunflower, sugar cane, others). In
15 | this study we included nut trees, date palms, citrus trees, orchards, olive trees, grapes,
16 | potatoes, cotton, vegetables and fodder grasses.

17 | For each new crop a representative ~~plant-species~~ was selected ~~after-for~~ which the
18 | parameterisation was performed (see Table 2 for details on the parameters described in the
19 | following sentences). Potatoes were introduced as an annual crop following the ~~same~~
20 | approaches ~~as~~ described in Bondeau et al. (2007) for other annual crops. Potatoes are planted
21 | in early spring in cooler climates and late winter in warmer regions (FAO, 2008). In LPJmL
22 | they are sown each year in the areas indicated by the land use input taking into account the
23 | seasonality of rainfall and temperature and the experience of farmers (see Waha et al., 2012
24 | ~~for more information~~). In case of no water stress, leaf area index (LAI) development follows a
25 | prescribed curve (as in SWAT) with inflexions ~~s~~ points according to the parameters shown in
26 | Table 2 (~~Pphu~~ ~~1a/2b~~, ~~Lmax~~ ~~1a/2b~~, ~~Phusen~~), but LAI ~~will be is~~ reduced in ~~the~~ case of water
27 | stress by scaling it ~~with-to~~ the difference between atmospheric demand and water supply.
28 | Phenology and maturity ~~is-are~~ modelled after the heat unit theory: when the accumulated
29 | difference between daily temperatures and base temperature reaches a prescribed total
30 | growing degree amount (called hereafter PHU for potential heat units) ~~then~~ the potatoes are
31 | ripe and ~~they~~ are harvested. Absorbed photosynthetically active radiation drives assimilation,
32 | ~~and carbon~~ ~~Carbon~~ allocation to different parts of the plant is a function of PHU development.
33 | The PHU parameter used depends on the mean temperature for spring varieties and on the
34 | sowing date for the winter/fall varieties and ranges from 1500 to 2400°Cd (with lower PHU in
35 | cooler climates).

36 | Agricultural trees –including grapes and cotton which are modelled in the present study as
37 | small trees– are planted as samplings with 2.3 grams carbon in sapwood and ~~4-6a~~ LAI ~~of 1.6~~
38 | in the growing areas indicated by the land use input. Each agricultural tree has a country and
39 | tree-specific planting density, and a tree-specific parameter determines the number of years
40 | that are needed for trees to grow before the first harvest. The ~~lat~~ter parameter depends ~~in~~
41 | ~~reality~~ not only on the varieties used and on the biophysical situation, but also on ~~the chosen~~

management, especially on the usage of fertilizers and irrigation. ~~Due to this complexity~~
~~There is insufficient quantitative data a general lack of information~~ on this issue, and ~~we~~
~~therefore, thus, this parameter is~~ assumed ~~this parameter~~ to be 4 years for all agricultural
trees. After the ~~escat number of~~ years, a plant-specific portion (HI, "harvest ratio" or "harvest
index") of the net primary productivity (NPP) of the tree is harvested every year. Thus, fruit
growth is represented by a carbon accumulation that equals the multiplication of HI and NPP.
An additional tree-specific parameter determines the replanting cycles of trees. Since there is
no data available on this, ~~it was we~~ assumed that plantations are renewed (replanted) after 40
years, ~~and the cycle begins one more time~~. Most agricultural trees have chilling requirements,
i.e. ~~they~~ need a period of low temperatures before flowering. This is modelled using the
parameter T_{lim} shown in Table 2. 20 years running average of the coldest month maximum
and minimum temperatures are compared with these values and define the bioclimatic limits
of each species. ~~–~~ Hence, temperature warming above these limits would inhibit the
establishment and survival of the perennial crops.

For deciduous trees the active phase starts when the daily temperature is higher than the base
temperature and it is assumed that fruit growth occurs in the second half of the active phase of
the year, i.e. when the phenological scalar (fraction of the maximum leave coverage) is > 0.5
and before leaf senescence starts. Leaf senescence occurs when daily temperatures fall below
the base temperature.

Following Sitch et al. (2003), evergreen trees are assumed to have constant ~~leave-leaf~~
coverage and ~~leave-leaf~~ longevities > 1 year. In this case, the accumulation ~~of carbohydrates~~
in the fruits occurs ~~in-on~~ days where temperature is above a tree-specific base temperature
until a tree-specific threshold is reached (GDD).

Grass grows in the same areas of agricultural trees, excepting for cotton ~~and~~ grape ~~plantations~~
and orchards ~~plantations~~. For these three classes we assumed that grasses and weeds ~~are not~~
~~supposed to do not~~ grow, ~~thereby in order to avoiding~~ competition with the crops, ~~and~~
~~thus implying~~ that ~~the any~~ ground cover is eradicated through some sort of weed control. This
is the dominant practice in reality, although exceptions to this rule are gaining in importance.

The categories "vegetables" and "fodder grass" are modelled following the modelling
approach of C3 grass described in Sitch et al. (2003). This is very appropriate for fodder
grasses in the Mediterranean region since ~~they these~~ are mainly alfalfa and clover. ~~In case of~~
~~the For~~ vegetables, this parametrisation ~~allows accountings~~ for the very large physiological
and allometric heterogeneity of vegetables, and also for multiple harvests per year, a fact that
is well represented by a constant cover of the areas. Following the implementation of
temperate herbaceous PFTs in Sitch et al. (2003), the photosynthesis in vegetables and fodder
grasses is assumed to be optimal between 10 and 30°C. Vegetables and fodder grasses are
harvested ~~when once their~~ phenology is complete (i.e. the growing degree day accumulation
determined by a parameter was reached) and the biomass increment is equal or greater than
200 g C m⁻² since the last harvest event. At that time, 50% of the aboveground biomass is
transferred to the harvest compartment. This ~~assumption is an average and might may~~ be
rather low for some vegetables ~~like such as e.g.~~ lettuce, and rather high for others, ~~such as like~~
~~e.g.~~ beans. For the conversion from dry to fresh matter it was assumed that vegetables have an

average water content of 40%, which, again, is rather low for some cases, e.g. cucumbers, but rather high for e.g. garlic. The moisture content of fodder grass varies approximately between 10% and 75% depending on whether it is reported for hay, silage or fresh fodder. Here we represent fodder grass for hay production and assume, thus, a moisture content of 10%.

The standard calibration process for agricultural management in LPJmL crops was extended in order to include agricultural trees. For annual crops this procedure consists in performing a set of runs with systematically modified management parameters representing the heterogeneity of fields, high-yielding varieties and the maximal achievable LAI (see more details in Fader et al., 2010). Similarly, 10 runs systematically modifying the tree and country-specific plantation density parameter were performed for calibrating the management of agricultural trees plantations. Planting densities range from 25% to 230% of the standard values, which were derived from literature research (see Table 2). For grapes the range was prescribed between 2000 and 15000 vines per hectare. The tree density for each country was then chosen based on the best matching with reported FAO yields.

2.3 Irrigation water requirements and soil carbon

The computation of net and gross irrigation requirements in LPJmL (NIR and GIR, respectively, see below) is explained in detail in Rost et al. (2008) and Rohwer et al. (2006). The functioning of soil decomposition, soil biochemistry and soil hydrology, including soil organic carbon (SOC), is explained in Schaphoff et al. (2013) and Sitch et al. (2003). In the following paragraphs a short and simplified summary of these procedures will be given.

Irrigation is triggered in irrigated areas when soil water content is lower than 90% of field capacity in the upper 50 cm of the soil (here “irrigated layer”). The Plants-plants' NIR is modelled in LPJmL as the amount of water that plants need, taking into account the water holding capacity of the irrigated layer and the relative soil moisture (Rost et al., 2008):

$$NIR[mm\ d^{-1}] = \min\left(\frac{1}{f_{Ril}}\left(\frac{D}{S_y} - w_r\right), 1 - w_{il}\right) WHC \quad (6)$$

Where:

D [mm d⁻¹] is the atmospheric demand, which depends on potential evapotranspiration and canopy conductance.

S_y [mm d⁻¹] is the soil water supply, which equals to a crop's specific maximum transpirational rate if the soil is saturated or declines linearly with soil moisture.

f_{Ril} [fraction] is the proportion of roots in the irrigated layer.

w_{il} [fraction] is the water content in the irrigated layer.

w_r [fraction] is the water content weighted with the root density for the soil column.

1 *WHC* [mm] is the field capacity of the irrigated layer (water holding capacity).

2 |
3 *GIR*, also called water withdrawal or extraction, is obtained by dividing *NIR* by the project
4 efficiencies (*EP*):

5
6
$$GIR [mm d^{-1}] = \frac{NIR}{EP} \quad (7)$$

7 |
8 *EP* is a country-specific parameter calculated for LPJmL by Rohwer et al. (2006) after the
9 approach described in the FAO irrigation manual (Savva and Frenken, 2002). It takes into
10 account reported data on conveyance efficiency (*EC*), field application efficiency (*EA*) and a
11 management factor (*MF*):

12
13
$$EP [0 to < 1] = EC * EA * MF \quad (8)$$

14 |
15 *EA* represents the water use efficiency on the fields and [its](#) increase from surface irrigation
16 systems, over sprinkler systems, to drip irrigation systems. *EC* represents the water use
17 efficiency in the distribution systems and is assumed to be linked to irrigation systems (lower
18 for open channels than for pressurized pipelines). *MF* varies between 0.9 and 1 and is higher
19 in pressurized and small scale systems under the assumption that large-scale systems are more
20 difficult to manage (see more details in Rohwer et al. 2006).

21 | [The soil column in](#) LPJmL has 5 hydrologically and thermally active layers (20 cm, 30 cm, 50
22 cm, 1 m and 1 m thickness) where roots have access to water. Infiltration depends on the soil
23 water content of the first layer (water that does not infiltrate runs off) and percolation
24 between the layers was simulated following the storage routine technique (see Schaphoff et
25 al., 2013 for more details). Excess water over the saturation level is assumed to feed
26 subsurface runoff. LPJmL has two soil carbon pools, with intermediate and fast turnover
27 (0.001 and 0.03 rate of turnover per year at 10°C). The maximum decomposition rate is
28 reached around field capacity and decreases afterwards due to decreased soil oxygen content.
29 A simple energy balance model is used for the thermal soil module. It includes a one-
30 dimensional heat conduction equation, convection of latent heat, thawing and sensible heat
31 (see Schaphoff et al., 2013 for more details).

32 **3 Results**

33 The performance of the improved LPJmL version was tested by simulating agricultural yields,
34 irrigation water requirements and soil carbon density [and comparing the results to published](#)
35 [observations](#).

3.1 Agricultural yields

Figure 1 shows LPJmL simulated yields in metric tonnes fresh matter per hectare for the calibrated run for all new crops where FAOSTAT had data in the Mediterranean region, averaged for the period 2000-2009. LPJmL simulates all nuts, olives, fruits and potatoes in a very good agreement with FAO reported values, showing Willmott coefficients¹ of ≥ 0.6 in all cases. Only two cases with large planting areas and significant differences are visible, both in Turkey, for grapes and nut trees. The latter is due to the chosen representative tree for the parametrisation of this group (almonds) which does not represent the majority of nut plantations in Turkey. In 2010 almost 70% of nut trees in Turkey were hazelnuts and only 3% almonds (FAO, 2015a). The underestimation of grape yield in Turkey might be related to more than one factor, including the fact that the wine sector is very dynamic in recent years there, with increases in production but decreases in area harvested (FAO, 2015a). FAO calculates yields by dividing national production by harvested area and calculates, thus, an increase in yields and a higher average over the years analysed. Our input dataset shows a slight increase in grape areas with relative constant production, hence, we calculated a lower yield average over the years. Also the general parametrisation for European grapes probably cannot represent the special character of local Turkish varieties that are well adapted to sandy soils and high altitudes.

Validating subnational patterns of yields is very difficult due to a general lack of data on this and important differences with other estimations in terms of scale, methods and time frames. However, we included in Fig. S2 a comparison with the yields from Monfreda et al. (2008) for the new crops where this study offers subnational data (note that their estimates are for the period of time around the year 2000 and at the administration level). LPJmL reproduces correctly a number of spatial patterns, such as some high yielding regions: olives in Greece, vineyards in Israel, Lebanon, Southern Spain, the Po valley and the Italian provinces of Emilia Romagna and Latium, potatoes in Turkey, Greece, Egypt, Morocco, Israel, Lebanon and Algeria, as well as cotton yields in Southern Spain, Greece, Turkey, Egypt, Israel and Lebanon. Also some low-yielding zones are in good agreement, as it is the case for potatoes in the Balkans, Portugal and Tunisia, olives in Morocco, Algeria and Tunisia, and cotton in Tunisia. However, some few patterns shown by Monfreda et al. (2008) are not shown in LPJmL simulations, including the North-South pattern of olives in Spain, the high yielding zone of olives in Southern Italy and the grapes and olives yields in Egypt. The first case is due to the extremely higher management intensity in Southern Spain that-which is not captured by the national calibration of planting densities (Scheidel and Krausmann, 2011). The latter case is originated by differences between the *MON* and *MIRCA* land use datasets that produce, in turn, a lack of irrigated areas of grapes and olives in Egypt in our dataset (see section 2.2 for more details). The same is the case of the gaps in potatoes, e.g. in France. Overall there is a large agreement with no systematic differences between the spatial patterns shown by Monfreda et al. (2008) and the ones computed in the present study.

¹ The Willmott coefficient was developed by Willmott (1982) as a tool for testing model performance against independent data and is calculated by $1 - \frac{\sum(p-o)^2}{\sum((p-\bar{o})^2 + (o-\bar{o})^2)}$, where o is the independent data, p the LPJmL simulated data, and \bar{o} denotes the mean of independent data.

3.2 Irrigation water requirements

Figure 2 shows LPJmL-simulated NIR per hectare, which presents a clear North-South pattern that follow the climate-driven patterns of potential evapotranspiration. Konzmann et al. (2013) presented simulated irrigation requirements globally for around 10 crop functional types with a former version of the LPJmL model where tree plantations were represented as mowed grasslands. Their Fig. 1 shows a ~~similar~~ grid-cell pattern ~~broadly than similar to~~ ours but ~~the our~~ more detailed representation of Mediterranean crops ~~in the present study leads~~ to higher values in various regions, including Algeria, Tunisia, Israel, Lebanon, Greece and the Iberian Peninsula. This is in good agreement with a general tendency of trees to absorb and transpire more water than grasslands (Belluscio, 2009).

Siebert et al. (2010) computed present irrigation consumptive water use for subnational administrative units by means of the GCWM model that compares well with our NIR estimates (Fig. 3). However, they estimated higher values in Egypt, Libya, Greece and Portugal and lower values in the Po Valley and Southern France. These differences are mainly linked to disparities in the land use dataset used as inputs: Siebert et al. (2010) areas equipped for irrigation are larger than the ones used in the present study in the first group of countries, and smaller in the second group. The comparison in absolute terms (Fig. S3) shows a similar pattern but with additional high differences in the Spanish province of Andalucía. These ~~differences~~ may be linked to ~~the~~ model approach, for example ~~the difference in the~~ crops considered (olives, orchards, nuts ~~are~~ only considered in the present study by LPJmL), different methods ~~for to model~~ evapotranspiration (Penman-Monteith versus Priestley-Taylor) and differences in growing periods (e.g. dynamic versus static sowing dates for annual crops). Since Andalucía is a region strongly ~~linked to characterized by~~ horticulture and ~~taking into account that we due to our~~ parametrised ~~ation of~~ vegetables ~~as through~~ C3 grasses, it is worthwhile to look in more detail into this class, also because neither Siebert et al. (2010) nor the present study accounts for cultivation of vegetables in greenhouses. We computed independent irrigation water requirements of $2357.2 \text{ m}^3 \text{ ha}^{-1}$ based on the values presented in Table 2 from Gallardo & Thompson (2013) that concerns various vegetables and water melon grown in greenhouses. Based on the crop cycles described in their publication, we assumed the possibility of planting 3 vegetables types per year using the same area (multiple cropping). Vegetables are planted on around 1.6 Mha in the Mediterranean region. This yields total water consumption of 11.3 km^3 . The present study computed 9.7 km^3 , thus, a very similar value.

In total, the agricultural sector in the Mediterranean was simulated to withdraw approx. 223 km^3 of water per year for irrigation (average 2000-2009), with GIR being especially high in the Nile Delta, the Eastern Mediterranean and in some Spanish regions (not shown). Our national GIR values are in good agreement with AQUASTAT data (FAO, 2015b) (Fig. 4, squares), with some differences. It is ~~however~~ difficult to evaluate the quality of the AQUASTAT data. For example the values of three countries with large differences to our estimates (Algeria, Lebanon and Jordan) are in fact not reported data but modelled data. Assuming that the modelling was performed by the FAO's model CROPWAT, our estimates might be more accurate since we perform a ~~validated~~ process-based calculation of

transpiration instead of prescribing crop water coefficients. Another example of uncertainty is shown in the case of Egypt. In Fig. 4 (red symbols) it is evident that the estimates for Egypt vary largely, e.g. Rayan & Djebedjian (2004) presented much lower estimates than AQUASTAT.

Döll and Siebert (2002) were probably the first authors ~~who quantifying-quantified~~ irrigation water requirements at the global level while distinguishing two crop classes (rice and non-rice). Their Table 5 shows GIR for Egypt, Israel and Spain according to independent data and their calculations (using irrigation areas from 1995 and climate from 1961 to 1990). Despite the difference in the period of time analysed and the methodology, Fig. 4 shows that our results agree well for Israel and Spain for the independent data ~~(from the water commissioner of Israel and the executive secretary of the International eCommission of Irrigation and dDrainage, respectively)~~, while they found their values to be overestimated for both countries. For Egypt, independent data delivers 30% higher GIR than our calculations and 27% lower than the values of Döll and Siebert (2002). However, ~~the one more time, it must be asked how reliability of these are~~ reported water use data ~~cannot be established, especially in the case of~~ ~~for~~ Egypt, ~~where these numbers are~~ ~~relevant in-for~~ negotiations on water allocation treaties with upstream countries of the Nile river.

A report by Cánovas Cuenca (2013), quoting an unpublished paper by Cánovas & del Campo (2006), shows in its Table 17 irrigation water requirements for Mediterranean countries ~~when~~ assuming that every hectare agriculture needs 6176 m³ of water per year. Our analysis shows that while this number delivers a fair estimate of irrigation requirements for some Northern Mediterranean countries, it strongly underestimates irrigation requirements of dry Mediterranean countries (Fig. 4, dots). This confirms that the environmental and climate diversity of the Mediterranean region requires spatial-explicit modelling approaches.

~~SummarizingTo summarize~~, LPJmL computes Mediterranean irrigation water requirements in the range of former studies, even if comparisons are a challenge due to inconsistent model inputs, differences in modelling approaches and due to the fact that to our best knowledge, this is the first study with a complete representation of Mediterranean crops.

3.3 Soil ~~Carbon-carbon~~ density

As mentioned in the introduction, some assessments can only be performed with a model that includes natural and agricultural ecosystems with a fair detail in the hydrological cycle. This is the case ~~of assessingfor~~ carbon sequestration by soils in a unit of area ~~that has both with~~ natural and agricultural vegetation –a very important variable for the climate debate and the ecosystem service research domain.

Soils under forest, grasslands and cropland show different ~~soil~~-carbon densities depending on ~~climatic variablese~~, vegetation's ~~characteristics~~, soil structure, and ~~the management-of-the system~~. Generally speaking, forests have higher proportions of soil organic carbon (SOC) compared to mowed grasslands and they, in turn, have higher values compared to cropland planted with annual crops (Jobbágy and Jackson, 2000, Ecclesia et al., 2012, Wert et al., 2005). Also evergreen broadleaved forests and plantations have usually higher SOC than deciduous

1 | forests and plantations in semi-arid climates (Doblas-Miranda et al., 2013). Agricultural trees
2 | plantations have a lower tree density than forest, generally a regular distribution of trees that
3 | increase soil evaporation and they are subject to removal of biomass (harvest). These factors
4 | lead ~~generally~~ to lower SOC values in tree plantations compared to forest. However,
5 | management of tree plantations, including irrigation input, planting density, presence or
6 | eradication of grass strips and mulching can strongly increase or decrease SOC. Putting all
7 | this information together and ~~being assuming~~ that management and environmental factors ~~are~~
8 | comparable, the SOC of agricultural tree plantations is expected to be generally higher than
9 | the SOC of mowed grasslands and generally lower than the SOC of natural forests and native
10 | grasslands. This is especially true for evergreen tree plantations and managed grasslands with
11 | high-frequency mowing. Hence, the implementation of agricultural trees in LPJmL should
12 | ~~have produced~~ higher SOC over the entire soil profile in many Mediterranean areas. This is
13 | because before the implementation of agricultural trees, the areas corresponding to these agro-
14 | ecosystems were simulated as mowed grasslands.

15 | As expected, Fig. 5 shows that implementing agricultural trees in LPJmL increased the carbon
16 | stock in soils in the whole Mediterranean ~~besides except~~ France. This exception is due to the
17 | fact that there are only two new implemented crops in France with significant areas: non-
18 | citrus orchards and grapes. Both are deciduous trees, with high soil evaporation at the
19 | beginning and end of the active period affecting carbon decomposition. Also orchards have a
20 | relatively low planting density in France (1300 trees per hectare) which reduces shadow
21 | effects and ~~vegetation carbon litter~~ input that, in turn, results in a lower soil carbon compared
22 | to high density mowed grass.

23 | Validation of the new SOC patterns is very challenging since SOC measurements are spatially
24 | discontinuous as well as dependent on local conditions, sampling method and small scale
25 | drainage conditions. Comparison with empirically or process-based modelled SOC is also
26 | difficult due to differences in approaches, parameters, processes considered and issues of
27 | scale. Nevertheless, we compared our SOC estimates before (LPJmL_{Old}) and after
28 | (LPJmL_{New}) the implementation of agricultural trees with the organic carbon density from the
29 | HWSD database (Hiederer & Köchy, 2012). These data are produced ~~by~~ establishing
30 | functions between SOC and soil type, topography, climate variables and land use situation.
31 | For this comparison we ~~built-calculated~~ the difference of absolute differences ($LPJmL_{Old} -$
32 | $HWSD - LPJmL_{New} - HWSD$). ~~The result is shown in~~ Fig. 6). Considering significant
33 | differences (results $\leq 1 \text{ t ha}^{-1}$), the number of grid-cells with decreased differences to the
34 | HWSD estimates almost doubles the number of grid-cells with increased differences (767
35 | versus 460 grid-cells). This means that the development presented in this study moved
36 | LPJmL's results for SOC closer to HWSD values.

37 | As mentioned before, ~~options for~~ comparison with other SOC estimates ~~is complex~~
38 | ~~are limited~~. The documentation of HWSD estimates (Hiederer & Köchy, 2012) offers an
39 | impressive effort in comparing their estimates with other assessments. They found large,
40 | spatially diverse differences to other estimates; some of them could be associated with
41 | differences in approaches. ~~While-When~~ comparing LPJmL results with HWSD estimates, it is
42 | necessary to bear in mind that HWSD offers a more detailed spatial scale and representation

of processes linked to soil types, while LPJmL has a more detailed influence of land use history, seasonality of temperature and types of crops in SOC formation.

4 Discussion

4.1 Advances through consistent carbon-water-agriculture modelling

Environmental degradation, ~~future~~-climate change, and population growth will put Mediterranean agriculture and natural ecosystems under ~~enormous~~ pressure (IPCC, 2012, Kovats et al., 2014, Diffenbaugh and Giorgi 2012, Skurras and Psaltopoulos, 2012, Doblas-Miranda et al., 2015). Timely and appropriate coping with the combination of these ~~issues~~ challenges will need collaboration between the Mediterranean countries and local communities in a number of issues, including advanced development of adaptation options, common plans on energy transition, environmental policy and best-practice rules in nature conservation and agricultural management. Collaboration will have to be designed in a framework that allows taking into account the larger European and global picture in terms of environmental change, ~~and~~ dynamics of ecological systems, foreign investments and migration movements. This calls for new tools that are able to be applied at large scale and can account for the interlinkages between agricultural systems, carbon cycle and water resources.

~~LPJmL is a state-of-the-art, global ecosystem model that needs a minimal set of inputs, uses a rather limited amount of parameters and requires limited computational resources.~~ With the model development presented in the current study, LPJmL is now a tool suitable to support Mediterranean decisions makers ~~and further advance our understanding of the Earth system as a managed space~~. The inclusion of Mediterranean crops in LPJmL not only increased substantially the proportion of agricultural areas for which quantitative assessments are possible, considered but it also improved the potential for computation of irrigation requirements and soil carbon. ~~The outcome is in this way the development delivered~~ a model with a comprehensive representation of ecophysiological processes for all vegetation types (natural and agricultural) in a consistent and validated framework that produces estimates of carbon, agricultural and hydrological variables for the entire Mediterranean basin. As such, LPJmL is especially suitable for analyses on water issues. Taking into account the projected water scarcity due to climate change in the Mediterranean area, the continuously dropping groundwater tables due to overexploitation, and the projected increases in irrigation water demand (Fischer et al., 2007, Konzmann et al., 2013, Wada et al., 2010, Arnell 2004), this constitutes a promising area for future model applications and further development. A first application of this model development was presented by Fader et al. (2015), pointing out that irrigation water needs of perennial crops in the Mediterranean region might increase significantly under climate change and some countries may face constraints to meet the higher water demands.

4.2 Potential applications and perspective for further research

The inclusion of perennial crops in LPJmL presented in the current study opens up the possibility for a number of large-scale applications and research studies that cannot be performed with empirical and/or input-intensive agronomic models. These include assessments of ~~the~~ climate change impacts on hydrological variables, agricultural production and carbon sequestration; ~~investigation of shifts in suitable growing areas for agricultural trees, as well as~~ evaluation of consequences of land use change (including expansion of irrigated areas); ~~and assessments on ecosystem services provided by perennial cultures, for example habitat provision for avifauna.~~ Some of these applications could also be performed by land surface models, but LPJmL has now the advantage of considering perennial crops in detail, which allows more precise quantifications not only in the Mediterranean region, but also in other macro-regions where agriculture is partially dominated by tree crops, such as Australia, South Africa, Chile, Western Argentina and California. Moreover, having a more accurate representation of perennial crops also allows studies on shifts in suitable growing areas for agricultural trees, diversity of diets, resilience of agricultural systems, needs for climate change adaptation and implications for food security as well as assessments on ecosystem services provided by perennial cultures, for example habitat provision for avifauna.

~~However, besides the improvements that are ongoing at in this moment in the various LPJmL working groups, there are many possible~~ Further improvements and refinements of LPJmL that can ~~and should be performed envisaged~~ for some applications in the Mediterranean area. We can divide these potential improvements in three groups, enumerated from least to most complex and work-intensive: a) input-related, b) parameter-related, and c) inclusion of new processes.

The most important input-related improvement concerns national and subnational studies and is related to the need of increasing the spatial resolution in all inputs used by LPJmL. ~~However, the~~ The limited availability of climate data and scenarios in a higher spatial resolution for the whole basin as well as missing detailed flow direction maps, especially for North Africa, has constrained this refinement until now. Nevertheless, work on data interpolation and downscaling is ongoing to bring the model at the 15 arc minute resolution. Another input-related issue is the need of scenarios of future crop patterns as a consequence of climatic and socio-economic change. With climate change very likely affecting the potential growing areas of agricultural trees and their profitability, studies aiming at a future quantification of agricultural, biochemical and hydrological variables would profit from coupling between models like LPJmL and land use models.

Small-scale application aiming at comparing and analysing single crops may require parameter-related changes such as re-parametrisation allowing differentiation of harvesting times after uses and varieties (e.g. varieties of grapes, difference between table olives and olives for oil), grid-cell specific planting densities and its differentiation between irrigated and rain-fed conditions, as well as crop-specific setting up of fruits, which at present depends on the phenological development but ~~will not be are not~~ differentiated for different crops. For national-scale studies re-parametrisation with different representative plants for the groups (e.g. using hazelnuts instead of almonds for nuts trees Turkey) is possible without any

difficulty. For this group of improvements data on management is essential, including harvesting times, post-harvest uses, planting densities, planted varieties, etc. Another benefit of more precise management information would be the possibility of differentiating parameters that are assumed to be static and equal for all plants in the present study, such as the number of years that a perennial plantation stays in production before being renewed and the period of time from the planting until the first harvest (see section 2.2).

Some refinements in modelled-processes should be undertaken for studies aiming at detecting year-to-year phenological changes or sub-yearly patterns of carbon allocation in agricultural trees. These may include improved representation of chilling requirements, for example implementing the chilling units approach (Byrne & Bacon, 2015), variable harvest index depending on the special conditions of the year, implementation of dwarf trees, and daily update of carbon partitioning. Also, including a more differentiated approach of agricultural management, such as discretizing practices, typology and processes affected, may be necessary for assessments on climate change adaptation and soil carbon. Also connected to soil carbon, the inclusion of erosion and salinization would be essential since this process plays an important role in the semi-arid, hilly and terraced landscapes of the Mediterranean area (García-Orenes et al., 2012, Poessen and Hooke, 1997).

Finally, the inclusion of horticulture in LPJmL opens the door for further large developments aiming at the assessment of alternative agro-ecosystem managements and their environmental performance. One clear example for this would be the link between agroforestry systems and biodiversity conservation.

Code availability and technicalities

LPJmL is written in the C programming language and is run mainly under UNIX-like systems. Inputs and outputs are in binary format. Depending on the size of the region analysed and on the desired spatial resolution, it may require a high-performing computational structure. The version used as base for the present development was the 3.5.003 and the revision number 213340 from 17/5/2014.

The main site for downloads of different model versions can be found under:

<https://www.pik-potsdam.de/research/projects/activities/biosphere-water-modelling/lpjml/versions>

There, downloads are free of charge and possible after registration. However, the latest model version that includes the agricultural module and the Mediterranean development is not available yet since the different working groups are still compiling a complete technical documentation and merging the last developments into one unique model version. Please contact the first author of this publication if you ~~plan have in mind an urgent application of the model and envisage that can be done in the framework of a longer-term scientific, close~~ collaboration.

Acknowledgments

This work is a contribution to the Labex OT-Med (n° ANR-11-LABX-0061) funded by the French Government «Investissements d’Avenir» program of the French National Research Agency (ANR) through the A*MIDEX project (n° ANR-11-IDEX-0001-02). [AB and WC receive support from the European Union’s Seventh Framework Programme for research, technological development and demonstration under the projects OPERAs \(grant agreement number 308393\) and LUC4C \(grant agreement number 603542\).](#)

We thank the Joint Research Centre of the European Commission for giving us access to the dataset “Global Soil Organic Carbon Estimates” of the European Soil Data Centre (ESDAC).

We thank the LPJmL group in the Potsdam Institute for Climate Impact Research for discussions and technical support. We also thank Stefan Siebert for the provision of data for the validation effort in Fig. 3.

References

- Aguilera, F., Ruiz, L., Fornaciari, M., Romano, B., Galán, C., Oteros, J., Ben Dhiab, A., Msallem, M., and Orlandi, F.: Heat accumulation period in the Mediterranean region: phenological response of the olive in different climate areas (Spain, Italy and Tunisia). *International Journal of Biometeorology*, 58, 867-876, doi: 10.1007/s00484-013-0666-7, 2014.
- Alasalvar, C. and Shahidi, F. (eds.): *Tree Nuts: Composition, Phytochemicals, and Health Effects* (Nutraceutical Science and Technology). Taylor & Francis. 340 pp., 2008.
- Al-Khayri, J. M. and Niblett, C. L.: Envision of an international consortium for palm research. *Emirates Journal of Food and Agriculture*, 24(5), 470-479, 2012.
- Álvaro-Fuentes, J., Easter, M., Cantero-Martínez, C., and Paustian, K.: Modelling soil organic carbon stocks and their changes in the northeast of Spain. *European Journal of Soil Science*, 62(5), 685–695, doi: 10.1111/j.1365-2389.2011.01390.x, 2011.
- Arnell, N. W.: Climate change and global water resources: SRES emissions and socio-economic scenarios. *Global Environmental Change* 14, 31–52, doi: 10.1016/j.gloenvcha.2003.10.006, 2004.
- Baldocchi, D. and Wong, S.: *An Assessment of Impacts of Future CO₂ and Climate on Agriculture*. California Climate Change Center, 40 pp., 2006.
- Bastin, S. and Henken, K.: *Water Content of Fruits and Vegetables*. University of Kentucky. <http://www2.ca.uky.edu/enri/pubs/enri129.pdf>, accessed 13.3.2015, 1997.
- Belluscio, A.: Planting trees can shift water flows. *Nature News* 7th November 2009. <http://www.nature.com/news/2009/091107/full/news.2009.1057.html>, accessed 22.4.2015, 2009.
- Beringer, T. and Lucht, W.: Nachhaltiges globales Bioenergiepotenzial. Commissioned expert study for the German Advisory Council on Global Change (WBGU) as a contribution to the flagship report *World in Transition - Future Bioenergy and Sustainable Land Use*, 2008.
- Biemans, H., Haddeland, I., Kabat, P., Ludwig, F., Hutjes, R. W. A., Heinke, J., von Bloh, W., and Gerten, D.: Impact of reservoirs on river discharge and irrigation water supply during the 20th century. *Water Resources Research*, 47, W03509, doi: 10.1029/2009WR008929, 2011.
- Bondeau, A., Smith, P., Zaehle, S., Schaphoff, S., Lucht, W., Cramer, W., Gerten, D., Lotze-Campen, H., Müller, C., Reichstein, M., and Smith, B.: Modelling the role of agriculture for the 20th century global terrestrial carbon balance. *Global Change Biology*, 13, 1-28, doi: 10.1111/j.1365-2486.2006.01305.x, 2007.
- Byrne, D. H., and Bacon, T.: Chilling accumulation: its importance and estimation. Dept. Of Horticultural Sciences, Texas A&M University. <http://aggie-horticulture.tamu.edu/stonefruit/chillacc.html>, accessed 21.09.2015, 2015.
- California Rare Fruit Growers: Olive. *Olea Europaea* L. <http://www.crfg.org/pubs/ff/olive.html>. accessed 13.3.2015, 1997.
- Cannell, M. G. R.: Dry matter partitioning in tree crops. In: Cannell, M. G. R., and Jackson, J. E., (eds.) *Attributes of trees as crop plants*. Abbotts Ripton, Institute of Terrestrial Ecology, 160-193. <http://nora.nerc.ac.uk/7081/1/N007081CP.pdf>, accessed 13.3.2015, 1985.

1 Cánovas Cuenca, J.: Report on water desalination status in the Mediterranean countries.
2 Instituto Murciano de Investigación y Desarrollo Agrario y Alimentario. Consejería de
3 Agricultura y Agua de la Región de Murcia, 308 pp., 2012.

4 Clothier, B., Hall, A. and Green, S.: Chapter 6. Horticulture. Adapting the horticultural and
5 vegetable industries to climate change. [http://www.climatecloud.co.nz/CloudLibrary/2012-](http://www.climatecloud.co.nz/CloudLibrary/2012-33-CC-Impacts-Adaptation_SLMACC-Chapter6.pdf)
6 [33-CC-Impacts-Adaptation_SLMACC-Chapter6.pdf](http://www.climatecloud.co.nz/CloudLibrary/2012-33-CC-Impacts-Adaptation_SLMACC-Chapter6.pdf), accessed 20.3.2015, 2012.

7 Diffenbaugh, N. S. and Giorgi, F.: Climate change hotspots in the CMIP5 global climate
8 model ensemble. *Climatic Change*, 114, 813-822, doi: 10.1007/s10584-012-0570-x , 2012.

9 Doblas-Miranda, E., Martínez-Vilalta, J., Lloret, F., Álvarez, A., Ávila, A., Bonet, F.,
10 Brotons, L., Castro, J., Curiel, J., Díaz, M., Ferrandis, P., García-Hurtado, E., Iriondo, J.,
11 Keenan, T., Latron, J., Llusà, J., Loepfe, L., Mayol, M., Moré, G., Moya, D., Peñuelas, J.,
12 Pons, X., Poyatos, R., Sardans, J., Sus, O., Vallejo, V., Vayreda, J., and Retana, J.:
13 Reassessing global change research priorities in Mediterranean terrestrial ecosystems: how
14 far have we come and where do we go from here? *Global Ecology and Biogeography* 24,
15 25–43, doi: 10.1111/geb.12224, 2015.

16 Doblas-Miranda, E., Rovira, P., Brotons, L., Martínez-Vilalta, J., Retana, J., Pla, M., and
17 Vayreda, J.: Soil carbon stocks and their variability across the forests, shrublands and
18 grasslands of peninsular Spain. *Biogeosciences*, 10, 8353-8361, doi: 10.5194/bg-10-8353-
19 2013, 2013.

20 Döll, P. and Siebert, S.: Global modeling of irrigation water requirements. *Water Resources*
21 *Research*, 38(4), 1037, doi: 10.1029/2001WR000355, 2002.

22 Duke, J. A.: Handbook of energy crops. Purdue University. Center for New Crops and Plant
23 Products. http://www.hort.purdue.edu/newcrop/duke_energy/refa-f.html, accessed
24 13.3.2015, 1983.

25 Ecclesia, R. P., Jobbagy, E. G., Jackson, R. B., Biganzoli, F., and Piñeiro, G.: Shifts in soil
26 organic carbon for plantation and pasture establishment in native forests and grasslands of
27 South America. *Global Change Biology*, 18, 3237–3251, doi: 10.1111/j.1365-
28 2486.2012.02761.x, 2012.

29 Elshibli, S.: Genetic Diversity and Adaptation of Date Palm (*Phoenix dactylifera* L.). PhD
30 Thesis University of Helsinki, 2009.

31 Fader, M., Gerten, D., Krause, M., Lucht, W., and Cramer, W.: Spatial decoupling of
32 agricultural production and consumption: quantifying dependence of countries on food
33 imports due to domestic land and water constraints. *Environmental Research Letters* 8,
34 014046, doi: 10.1088/1748-9326/8/1/014046, 2013.

35 Fader, M., Rost, S., Müller, C., Bondeau, A., and Gerten, D.: Virtual water content of
36 temperate cereals and maize: Present and potential future patterns. *Journal of Hydrology*,
37 384(3–4), 218–231, doi: 10.1016/j.jhydrol.2009.12.011, Green-Blue Water Initiative
38 (GBI), 2010.

39 Fader, M., von Bloh, W., Shi, S., Bondeau, A., and Cramer, W.: Mediterranean irrigation
40 under climate change: More efficient irrigation needed to compensate increases in
41 irrigation water requirements. *Hydrology and Earth System Sciences Discussions*, 12,
42 8459–8504, doi: 10.5194/hessd-12-8459-2015, 2015.

43 FAO: AQUASTAT. <http://www.fao.org/nr/water/aquastat/main/index.stm>, accessed
44 01.07.2014, 2015b.

1 FAO: Crop Water Information: Citrus. http://www.fao.org/nr/water/cropinfo_citrus.html,
2 accessed 13.3.2015, 2013a.

3 FAO: Crop Water Information: Grape. http://www.fao.org/nr/water/cropinfo_grape.html,
4 accessed 13.3.2015, 2013c.

5 FAO: Crop Water Information: Olive. http://www.fao.org/nr/water/cropinfo_olive.html,
6 accessed 13.3.2015, 2013b.

7 FAO: Cultivation of Potatoes. <http://www.fao.org/potato-2008/en/potato/cultivation.html>,
8 accessed 13.3.2015, 2008.

9 FAO: Cultivation of Potatoes. <http://www.fao.org/potato-2008/en/potato/cultivation.html>,
10 accessed 13.3.2015, 2008.

11 FAO: Date palm cultivation. FAO Plant production and protection paper 156, Rev. 1. Edited
12 and compiled by A. Zaid. <http://www.fao.org/docrep/006/y4360e/y4360e00.HTM>,
13 accessed 13.3.2015, 2002.

14 FAO: FAOSTAT. <http://faostat.fao.org/site/567/default.aspx#ancor>, accessed 01.07.2014,
15 2015a.

16 FAO: Tree crops. Guidelines for estimating area data. Statistics division.
17 [http://www.fao.org/fileadmin/templates/ess/ess_test_folder/documents/Production_trade/d](http://www.fao.org/fileadmin/templates/ess/ess_test_folder/documents/Production_trade/definitions/Tree_crops_guidelines_for_estaimating_area.doc)
18 efinitions/Tree_crops_guidelines_for_estaimating_area.doc, accessed 13.3.2015, 2011.

19 Fischer, G., Tubiello, F. N., van Velthuizen, H., and Wiberg, D. A.: Climate change impacts
20 on irrigation water requirements: Effects of mitigation, 1990–2080. *Technological*
21 *Forecasting and Social Change*, 74, 1083–1107, doi: 10.1016/j.techfore.2006.05.021, 2007.

22 Gallardo, M. and Thompson, R. B.: Water requirements and irrigation management in
23 Mediterranean greenhouses: the case of the southeast coast of Spain. In: FAO-AGP and
24 ISHS-CMPC (eds) *Good Agricultural Practices for Greenhouse Vegetable Crops:*
25 *Principles for Mediterranean Climate Areas*. FAO, Rome, Italy. pp. 109-136, 2013.

26 Garcia de Cortázar-Atauri, I.: Adaptation du modèle STICS à la vigne (*Vitis vinifera* L.):
27 utilisation dans le cadre d'une étude d'impact du changement climatique à l'échelle de la
28 France. PhD thesis ENSAM, 292 pp., 2006.

29 García-Orenes, F., Roldán, A., Mataix-Solera, J., Cerda, A., Campoy, M., Arcenegui, V., and
30 Caravaca, F.: Soil structural stability and erosion rates influenced by agricultural
31 management practices in a semi-arid Mediterranean agro-ecosystem. *Soil Use and*
32 *Management* 28, 571-579, doi: 10.1111/j.1475-2743.2012.00451.x, 2012.

33 Gerik, T., Williams, J., Francis, L., Greiner, J., Magre, M., Meinardus, A., Steglich, E., and
34 Taylor, R.: EPIC-Environmental Policy Integrated Climate Model. User's Manual Version
35 0810, 2014.

36 Gerten, D., Luo, Y., Le Maire, G., Parton, W. J., Keough, C., Weng, E., Beier, C., Ciais, P.,
37 Cramer, W., Dukes, J. S., Sowerby, A., Hanson, P. J., Knapp, A., Linder, S., Nepstad, D.,
38 and Rustad, L.: Modelled effects of precipitation, on ecosystem carbon and water
39 dynamics in different climatic zones. *Global Change Biology* 14:1-15, doi :
40 10.1111/j.1365-2486.2008.01651.x, 2008

41 Gerten, D., Schaphoff, S., Haberlandt, U., Lucht, W., and Sitch, S.: Terrestrial vegetation and
42 water balance - hydrological evaluation of a dynamic global vegetation model. *Journal of*
43 *Hydrology*, 286, 249–270, doi: 10.1016/j.jhydrol.2003.09.029, 2004.

44 Gordon, R., Brown, D. M., and Dixon, M. A.: Estimating potato leaf area index for specific
45 cultivars. *Potato Research*, 40(3), 251-266, doi: 10.1007/BF02358007, 1997.

Gutierrez, A. P., Ponti, L., Ellis, C. K., and d'Oultremont, T.: Analysis of climate effects on agricultural systems: A report to the Governor of California. 43 pp. http://www.climatechange.ca.gov/climate_action_team/reports/index.html, accessed 20.3.2015, 2006.

Harris, I., Jones, P. D., Osborn, T. J., and Lister, D. H.: Updated high-resolution grids of monthly climatic observations - the CRU TS3.10 Dataset. *International Journal of Climatology* 34, 623-642, doi:10.1002/joc.3711, 2014.

Haverkort, A. J. and MacKerron, D. K. L. (eds.): *Potato ecology and modelling of crops under conditions limiting growth*. Kluwer Academic Publishers, The Netherlands, 1995.

Hervieu, B.: *Agriculture: a strategic sector in the Mediterranean area*. CIHEAM analytic note N°18, 2006.

Hiederer, R. and Köchy, M.: Global Soil Organic Carbon Estimates and the Harmonized World Soil Database. EUR Scientific and Technical Research series – ISSN 1831-9424 (online), doi: 10.2788/13267, 2012.

Holzworth, D. P., Huth, N. I., deVoil, P. G., Zurcher, E. J., Herrmann, N. I., McLean, G., Chenu, K., van Oosterom, E. J., Snow, V., Murphy, C., Moore, A. D., Brown, H., Whish, J. P. M., Verrall, S., Fainges, J., Bell, L. W., Peake, A. S., Poulton, P. L., Hochman, Z., Thorburn, P. J., Gaydon, D. S., Dalgliesh, N. P., Rodriguez, D., Cox, H., Chapman, S., Doherty, A., Teixeira, E., Sharp, J., Cichota, R., Vogeler, I., Li, F. Y., Wang, E., Hammer, G. L., Robertson, M. J., Dimes, J. P., Whitbread, A. M., Hunt, J. van Rees, H., McClelland, T., Carberry, P. S., Hargreaves, J. N.G., MacLeod, N., McDonald, C., Harsdorf, J., Wedgwood, S., and Keating, B. A.: APSIM – Evolution towards a new generation of agricultural systems simulation. *Environmental Modelling & Software*, 62, 327-350, doi: 10.1016/j.envsoft.2014.07.009, 2014.

IIASA/FAO: *Global Agro-ecological Zones (GAEZ v3.0)*. IIASA, Laxenburg, Austria and FAO, 2012.

IPCC: Summary for Policymakers. In: *Managing the Risks of Extreme Events and Disasters to Advance Climate Change Adaptation* [Field, C.B., Barros, V., Stocker, T.F., Qin, D., Dokken, D.J., Ebi, K.L., Mastrandrea, M.D., Mach, K.J., Plattner, G.-K., Allen, S.K., Tignor, M., and Midgley, P.M. (eds.)]. A Special Report of Working Groups I and II of the Intergovernmental Panel on Climate Change. Cambridge University Press, Cambridge, UK, and New York, NY, USA, pp. 3-21, 2012.

Janick, J. and Paull, R. E. (eds.): *Encyclopedia of Fruit and Nuts*. Purdue University, USA. CAB International. 972 pp, 2008.

Kailis, S. and Harris, D.: *Producing Table Olives*. CSIRO Publishing, Collingwood, 2007.

Kiranga, N. A.: Morpho-argro-physio-karyotypic characterization of wild cotton (*Gossypium* spp.). Germplasm from selected counties in Kenya. <http://ir-library.ku.ac.ke/bitstream/handle/123456789/9050/Njagi%20Anthony%20Kiranga.pdf?sequence=1>, accessed 13.3.2015, 2013.

Klein Goldewijk, K., Beusen, A., de Vos, M., and van Drecht, G.: The HYDE 3.1 spatially explicit database of human induced land use change over the past 12,000 years. *Global Ecology and Biogeography*, 20(1), 73-86, doi: 10.1111/j.1466-8238.2010.00587.x, 2011.

Konzmann, M., Gerten, G., and Heinke, J.: Climate impacts on global irrigation requirements under 19 GCMs, simulated with a vegetation and hydrology model, *Hydrological Sciences Journal*, 58(1), 88-105, doi: 10.1080/02626667.2013.746495, 2013.

- Kovats, R. S., Valentini, R., Bouwer, L. M., Georgopoulou, E., Jacob, D., Martin, E., Rounsevell, M. and Soussana, J.-F.: Europe. In: Barros, V.R., Field, C.B., Dokken, D.J., Mastrandrea, M.D., Mach, K.J., Bilir, T.E., Chatterjee, M., Ebi, K.L., Estrada, Y.O., Genova, R.C., Girma, B., Kissel, E.S., Levy, A.N., MacCracken, S., Mastrandrea, P.R., and White, L.L. (eds.): Climate Change 2014: Impacts, Adaptation, and Vulnerability. Part B: Regional Aspects. Contribution of Working Group II to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change. Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, pp. 1267-1326, 2014.
- Kroes, J. G. and van Dam, J. C.: Reference Manual SWAP version 3.0.3. Alterra-report 773 3. Alterra, Green World Research, Wageningen, 211 pp, 2003.
- Ku, S.-B., Edwards, G. E., and Tanner, C. B.: Effects of light, carbon dioxide, and temperature on photosynthesis, oxygen inhibition of photosynthesis, and transpiration in *Solanum tuberosum*. Plant Physiology, 59, 868-872, 1977.
- Lapola, D.M., Priess, J.A., and Bondeau, A.: Modeling the land requirements and potential productivity of sugarcane and jatropha in Brazil and India using the LPJmL dynamic global vegetation model. Biomass and Bioenergy, 33(8), 1087–95, doi: 10.1016/j.biombioe.2009.04.005, 2009.
- Lavorel, S., Canadell, J., Rambal, S. and Terradas, J.: Mediterranean terrestrial ecosystems: Research priorities on global change effects. Global Ecology & Biogeography Letters, 7(3), 157-166, doi: 10.1046/j.1466-822X.1998.00277.x, 1998.
- Lionello, P., Malanotte-Rizzoli, P., Boscolo, R., Alpert, P., Artale, V., Li, L., Luterbacher, J., May, W., Trigo, R., Tsimplis, M., Ulbrich, U. and Xoplaki, E.: The Mediterranean Climate: An Overview of the Main Characteristics and Issues. In: Lionello, P., Malanotte-Rizzoli, P., Boscolo, R. (eds.): Mediterranean Climate Variability. Developments in Earth and Environmental Sciences 4, 2006.
- Lobell, D. B., Nicholas Cahill, K., and Field, C. B.: Historical effects of temperature and precipitation on California crop yields. Climatic Change, 81, 187–203, doi: 10.1007/s10584-006-9141-3, 2007.
- Lobianco, A., and Esposti, R.: The regional model for Mediterranean agriculture. IDEMA research project paper. https://lobianco.org/antonello/_media/academic:pubs:idema_deliverable17.pdf. Accessed 21.09.2015.
- Marsal, J. and Stöckle, C. O.: Use of CropSyst as a decision support system for scheduling regulated deficit irrigation in a pear orchard. Irrigation Science, 30, 139–147, doi: 10.1007/s00271-011-0273-5, 2012.
- Marsal, J., Girona, J., Casadesus, J., López, G., and Stöckle, C. O.: Crop coefficient (K_c) for apple: comparison between measurements by a weighing lysimeter and prediction by CropSyst. Irrigation Science, 31, 455–463, doi: 10.1007/s00271-012-0323-7, 2013.
- Marsal, J., Johnson, S., Casadesus, J., López, G., Girona, J., and Stöckle, C.: Fraction of canopy intercepted radiation relates differently with crop coefficient depending on the season and the fruit tree species. Agricultural and Forest Meteorology, 184, 1-11, doi: 10.1016/j.agrformet.2013.08.008, 2014.
- Meier, U.: Growth stages of mono- and dicotyledonous plants. BBCH Monograph. Federal Biological Research Centre for Agriculture and Forestry. Blackwell Wissenschafts-Verlag. 622 pp., 2001.

1 Ministry of Agriculture, Food and Rural Affairs: Wine grape production outside traditional
2 areas in Ontario. http://www.omafra.gov.on.ca/english/crops/facts/info_grapeprod.htm,
3 accessed 13.3.2015, 2013.

4 Monfreda, C., Ramankutty, N., and Foley, J. A.: Farming the planet: 2. Geographic
5 distribution of crop areas, yields, physiological types, and net primary production in the
6 year 2000. *Global Biogeochemical Cycles*, 22, GB1022, doi: 10.1029/2007GB002947,
7 2008.

8 Morales Sierra, A.: A model of productivity for olive orchards. MSc Thesis Plant Production
9 Systems, Wageningen University, 2012.

10 Moriondo, M., Ferrise, R., Trombi, G., Brilli, L., Dibari, C., and Bindi, M.: Modelling olive
11 trees and grapevines in a changing climate. *Environmental Modelling & Software*, doi:
12 10.1016/j.envsoft.2014.12.016, in press

13 Neitsch, S. L., Arnold, J. G., Kiniry, J. R., Srinivasan, R., and Williams J. R.: Assessment
14 Tool. Input/output file documentation. Version 2005. Grassland, Soil and Water Research
15 Laboratory, Temple, Texas. <http://swat.tamu.edu/media/1291/swat2005io.pdf>, accessed
16 13.3.2015, 2004.

17 Neitsch, S. L., Arnold, J. G., Kiniry, J. R., Srinivasan, R., and Williams J.R.: Assessment
18 Tool. Input/output file documentation. Version 2005. Grassland, Soil and Water Research
19 Laboratory, Temple, Texas. <http://swat.tamu.edu/media/1291/swat2005io.pdf>, accessed
20 13.3.2015, 2004.

21 Nesme, T., Brisson, N., Lescourret, F., Bellon, S., Crété, X., Plénet, D., and Habib. R.:
22 Epistics: A dynamic model to generate nitrogen fertilisation and irrigation schedules in
23 apple orchards, with special attention to qualitative evaluation of the model. *Agricultural*
24 *Systems*, 90(1-3), 202-225, doi: 10.1016/j.agsy.2005.12.006, 2006.

25 Netafim: Irrigation. Almond best practices. [https://www.netafim.com/crop/almond/best-](https://www.netafim.com/crop/almond/best-practice)
26 [practice](https://www.netafim.com/crop/almond/best-practice), accessed 13.3.2015, 2013.

27 Orlandi, F., Garcia-Mozo, H., Ben Dhiab, A., Galán, C., Msallem, M., and Fornaciari, M.:
28 Olive tree phenology and climate variations in the Mediterranean area over the last two
29 decades. *Theoretical and Applied Climatology*, 115(1-2), 207-218, doi: 10.1007/s00704-
30 013-0892-2, 2014.

31 Orwa, C., Mutua, A., Kindt, R., Jamnadass, R., and Anthony, S.: Agroforestry Database: a
32 tree reference and selection guide version 4.0. World Agroforestry Centre, Kenya, 2009.

33 Ostberg, S., Lucht, W., Schaphoff, S., and Gerten, D.: Critical impacts of global warming on
34 land ecosystems. *Earth System Dynamics*, 4, 347-357, doi: 10.5194/esd-4-347-2013, 2013.

35 Perry, L.: Cold Climate Fruit Trees. University of Vermont Extension Department of Plant
36 and Soil Science. <http://perrysperennials.info/articles/coldfruit.html>, accessed 13.3.2015,
37 2011.

38 Poesen, J. W. A. and Hooke, J. M.: Erosion, flooding and channel management in
39 Mediterranean Environments of southern Europe. *Progress in Physical Geography*, 21(2),
40 157-199, doi: 10.1177/030913339702100201, 1997.

41 Pontificia Universidad Católica de Chile: El Almendro. [https://climafrutal.wordpress.com/el-](https://climafrutal.wordpress.com/el-almendro/)
42 [almendro/](https://climafrutal.wordpress.com/el-almendro/), accessed 13.3.2015, 2008.

43 Portmann, F., Siebert, S., and Döll, P.: MIRCA 2000 – Global monthly irrigated and rainfed
44 crop areas around the year 2000: A new high-resolution data set for agricultural and

hydrological modelling. *Global Biogeochemical Cycles*, 24, GB1011, doi: 10.1029/2008GB003435, 2011.

Ramankutty, N., Evan, A. T., Monfreda, C., and Foley, J. A.: Farming the planet: 1. Geographic distribution of global agricultural lands in the year 2000. *Global Biogeochemical Cycles*, 22, GB1003, doi: 10.1029/2007GB002952, 2008.

Rayan, M. A. and Djebdjan, B.: Egypt's Water Demand, Supply and Management Policies. International Water Demand Management Conference, May 30 – June 3, 2004, Dead Sea – Jordan.
http://www.academia.edu/2085529/Egypt_s_Water_Demand_Supply_and_Management_Policies, accessed 13.3.2015, 2004.

Rayan, M. A. and Djebdjan, B.: Egypt's Water Demand, Supply and Management Policies. International Water Demand Management Conference, May 30 – June 3, 2004, Dead Sea – Jordan.
http://www.academia.edu/2085529/Egypt_s_Water_Demand_Supply_and_Management_Policies, accessed 13.3.2015, 2004.

Rohwer, J., Gerten, D., and Lucht, W.: Development of functional irrigation types for improved global crop modelling. PIK Report 104. Potsdam, 98 pp., 2006.

Rost, S., Gerten, D., Bondeau, A., Lucht, W., Rohwer, J., and Schaphoff, S.: Agricultural green and blue water consumption and its influence on the global water system. *Water Resources Research*, 44, W09405, doi: 10.1029/2007WR006331, 2008.

Roussos, P.A.: Training and Pruning Olives. Proceedings of the MGS Symposium: Dry Gardening – Philosophy and Practice. Agricultural University of Athens, Greece.
<http://www.mediterraneangardensociety.org/olives.html>, accessed 13.3.2015, 2007.

Sakin, E.: Net primary productivity of southeast Anatolia Region (SAR) in Turkey. *International Journal of Agriculture & Biology*, 14, 617–620, 2012.

Savva, A. P. and Frenken, K.: Monitoring the Technical and Financial Performance of an Irrigation Scheme. *Irrigation Manual Module 14*. 58 pp, 2002.

Scarascia-Mugnozza, G., Oswald, H., Piussi, P., and Radoglou, K.: Forests of the Mediterranean region: gaps in knowledge and research needs. *Forest Ecology and Management*, 132, 97-109, doi: 10.1016/S0378-1127(00)00383-2, 2000.

Schaphoff, S., Heyder, U., Ostberg, S., Gerten, D., Heinke, J., and Lucht, W.: Contribution of permafrost soils to the global carbon budget. *Environmental Research Letters*, 8, 014026, doi: 10.1088/1748-9326/8/1/014026, 2013.

Scheidel, A. and Krausmann, F.: Diet, trade and land use: a socio-ecological analysis of the transformation of the olive oil system. *Land Use Policy*, 28, 47–56, doi: 10.1016/j.landusepol.2010.04.008, 2011.

Schröter, D., Cramer, W., Leemans, R., Prentice, I.C., Araújo, M. B., Arnell, N. W., Bondeau, A., Bugmann, H., Carter, T. R., Garcia, C. A., de la Vega-Leinert, A. C., Erhard, M., Ewert, F., Glendining, M., House, J.I., Kankaanpää, S., Klein, R. J. T., Lavorel, S., Lindner, M., Metzger, M. J., Meyer, J., Mitchell, T. D., Reginster, I., Rounsevell, M., Sabaté, S., Sitch, S., Smith, B., Smith, J., Smith, P., Sykes M. T., Thonicke, K., Thuiller, W., Tuck, G., Zaehle, S., and Zierl, B.: Ecosystem service supply and vulnerability to global change in Europe. *Science*, 310, 1333-1337, doi: 10.1126/science.1115233, 2005.

- 1 Siebert, S. and Döll, P.: Quantifying blue and green virtual water contents in global crop
2 production as well as potential production losses without irrigation. *Journal of Hydrology*
3 384, 198-217, doi: 10.1016/j.jhydrol.2009.07.031, 2010.
- 4 Siebert, S. and Döll, P.: The Global Crop Water Model (GCWM): Documentation and first
5 results for irrigated crops. Frankfurt Hydrology Paper 07, Institute of Physical Geography,
6 University of Frankfurt, Frankfurt am Main, Germany, 2008.
- 7 Siebert, S., Burke, J., Faures, J. M., Frenken, K., Hoogeveen, J., Döll, P. D., and Portmann, F.
8 T.: Groundwater use for irrigation – a global inventory. *Hydrology and Earth System*
9 *Sciences*, 14, 1863–1880, doi: 10.5194/hess-14-1863-2010, 2010.
- 10 Sitch, S., Smith, B., Prentice, C., Arneth, A., Bondeau, A., Cramer, W., Kaplan, J.O., Levis,
11 S., Lucht, W., Sykes, M.T., Thonicke, K. and Venevsky, S.: Evaluation of ecosystem
12 dynamics, plant geography and terrestrial carbon cycling in the LPJ dynamic global
13 vegetation model. *Global Change Biology*, 9, 161-185, doi: 10.1046/j.1365-
14 2486.2003.00569.x, 2003.
- 15 Skuras, D. and Psaltopoulos, D.: A broad overview of the main problems derived from
16 climate change that will affect agricultural production in the Mediterranean area. In:
17 Meybeck, A., Lankoski, J., Redfern, S., Azzu, N. and Gitz, V.: *Building Resilience for*
18 *Adaptation to Climate Change in the Agriculture Sector. Proceedings of a Joint*
19 *FAO/OECD Workshop 23–24 April, 2012.*
- 20 Slovak wine academy Pezinok: Viticulture and viniculture. Study material. Elementary
21 Seminar. Module 1. European Commission, Leonardo Davinci Transfer of Innovation
22 Programm. 50 pp.
23 http://svapezinok.sk/files/gallery/20130103172711_WEB_SVA%20Zakladny%20seminar
24 [_EN_01.pdf](#), accessed 13.3.2015, 2009.
- 25 Sperling, O.: Water Relations in Date Palm Trees – a Combined Approach using Water, Plant,
26 and Atmospheric Data. Ph.D. Thesis, Ben-Gurion University, 2013.
- 27 Strik, B. C: Growing Table Grapes. EC 1639. Oregon State University, Extension Service.
28 http://smallfarms.oregonstate.edu/sites/default/files/publications/growing_table_grapes_ec
29 [1639_may_2011.pdf](#), accessed 13.3.2015, 2011.
- 30 Toky, O. M., Kumar, P., and Kumar, P.: Structure and function of traditional agroforestry
31 systems in the western Himalaya. I. Biomass and productivity. *Agroforestry Systems*, 9,
32 47-70, doi: 10.1007/BF00120155, 1989.
- 33 Tsiros, E., Domenikiotis, C., and Dalezios, N. R.: Assessment of cotton phenological stages
34 using agroclimatic indices: An innovative approach. *Italian Journal of Agrometeorology*,
35 2009(1), 50-55, 2009.
- 36 Valdés-Gómez, H., Celette, F., García De Cortázar-Atauri, I., Jara-Rojas, F., and Gary C.:
37 Modelling soil water content and grapevine growth and development with the STICS crop-
38 soil model under two different water management strategies. *Journal International des*
39 *Sciences de la Vigne et du Vin* 43(1), 13-28, 2009.
- 40 Verner, D.: *Adaptation to a Changing Climate in the Arab Countries: A Case for Adaptation*
41 *Governance and Leadership in Building Climate Resilience. The World Bank:*
42 *Washington, DC., 2012.*
- 43 Villalobos, F. J., Testi, L., Orgaz, F., García-Tejera, O., Lopez-Bernal, A., González-Dugo,
44 M. V., Ballester-Lurbe, C., Castel, J. R., Alarcón-Cabañero, J. J., Nicolás-Nicolás, E.,
45 Girona, J., Marsal, J., and Fereres, E.: Modelling canopy conductance and transpiration of

fruit trees in Mediterranean areas: a simplified approach. *Agricultural and Forest Meteorology*, 171-172, 93-103, doi: 10.1016/j.agrformet.2012.11.010, 2013.

Wada, Y., Beek, L. P. H. van, van Kempen, C. M., Reckman, J. W. T. M., Vasak, S., and Bierkens, M. F. P.: Global depletion of groundwater resources. *Geophysical Research Letters*, 37, L20402, doi: 10.1029/2010GL044571, 2010.

Waha, K., Müller, C., Bondeau, A., Dietrich, J. P., Kurukulasuriya P, Heinke, J., and Lotze-Campen, H.: Adaptation to climate change through the choice of cropping system and sowing date in sub-Saharan Africa. *Global Environmental Change*, 23(1), 130–43, doi: 10.1016/j.gloenvcha.2012.11.001, 2013.

Waha, K., van Bussel, L. G. J., Müller, C., and Bondeau, A.: Climate-driven simulation of global crop sowing dates. *Global Ecology and Biogeography*, 12, 2, 247-259, doi: 10.1111/j.1466-8238.2011.00678.x, 2012.

Werth, M., Brauckmann, H.-J., Broll, G., and Schreiber, K.-F.: Analysis and simulation of soil organic-carbon stocks in grassland ecosystems in SW Germany. *Journal of Plant Nutrition and Soil Science*, 168(4), 472–482, doi: 10.1002/jpln.200421704, 2005.

Willmott, C. J., 1982. Some comments on the evaluation of model performance. *Bulletin American Meteorological Society*. Pp. 1309-1313

World Bank: Population Growth (Annual %). *World Development Indicators*, Washington, DC. <http://data.worldbank.org/indicator/SP.POP.GROW>, accessed 1.12. 2013, 2013.

Wright, S., Hutmacher, B., Shrestha, A., Banuelos, G., Keeley, M., Delgado, R., and Elam, S.: Double Row and Conventional Cotton in Tulare County, California. In: Fischer, R. A.: *New directions for a diverse planet. Proceedings of the 4th International Crop Science Congress*. Brisbane, Australia, 26 September - 1 October 2004. http://www.regional.org.au/au/asa/2004/poster/2/7/4/1774_wrightsd.htm, accessed 13.3.2015, 2014.

Wünsche, J. N. and Lakso, A. N.: *Apple Tree Physiology–Implications for Orchard and Tree Management*. Presented at the 43rd Annual IDFTA Conference, February 6-9, 2000, Napier, New Zealand. <http://virtualorchard.net/IDFTA/cft/2000/july/cftjuly2000p82.pdf>, accessed 13.3.2015, 2000.

Yan, H., Cao, M., Liu, J., and Tao, B.: Potential and sustainability for carbon sequestration with improved soil management in agricultural soils of China. *Agriculture, Ecosystems & Environment*, 121(4), 325–335, doi: 10.1016/j.agee.2006.11.008, 2007.

Zamski, E. and Schaffer, A. A.: *Photoassimilate distribution in plants and crops: source-sink relationships*. Marcel Dekker, Inc., New York, 33 pp., 1996.

Zanotelli, D., Montagnani, L., Manca, G., and Tagliavini, M.: Net primary productivity, allocation pattern and carbon use efficiency in an apple orchard assessed by integrating Earth eddy covariance, biometric and continuous soil chamber measurements. *Biogeosciences*, 10, 3089–3108, doi: 10.5194/bg-10-3089-2013, 2013.

Zdruli, P.: Land resources of the Mediterranean: Status, pressures, trends and impacts of future regional development. *Land Degradation & Development*, 25, 373-384, doi: 10.1002/ldr.2150, 2014.

1 **Table 1:** Areas of LPJmL crops in the Mediterranean region. The stars indicate crops that are
2 implemented in this study.

	Rainfed		Irrigated		Total	
	10 ⁶ ha	%	10 ⁶ ha	%	10 ⁶ ha	%
Temp.Cer.	50.11	12.0	6.38	24.8	56.49	12.7
Maize	13.84	3.3	3.37	13.1	17.21	3.9
Fodder Grass*	9.90	2.4	0.48	1.8	10.38	2.3
Trop.Cer.	7.64	1.8	0.25	1.0	7.89	1.8
Pulses	6.03	1.4	0.68	2.6	6.71	1.5
Olives*	4.86	1.2	1.61	6.3	6.47	1.5
Vegetables*	4.24	1.0	1.60	6.2	5.84	1.3
Orchards*	4.27	1.0	1.26	4.9	5.53	1.2
Sunflower	4.40	1.1	0.43	1.7	4.82	1.1
Grapes*	4.09	1.0	0.59	2.3	4.68	1.1
Potatoes*	1.40	0.3	0.66	2.6	2.06	0.5
Cotton*	0.34	0.1	1.67	6.5	2.01	0.5
Nuts*	1.45	0.3	0.52	2.0	1.97	0.4
Temp.Roots	1.17	0.3	0.74	2.9	1.91	0.4
Rapeseed	1.72	0.4	0.05	0.2	1.77	0.4
Groundnuts	1.56	0.4	0.11	0.4	1.67	0.4
Rice	0.14	0.0	0.88	3.4	1.02	0.2
Citrus*	0.12	0.0	0.86	3.3	0.98	0.2
Soybeans	0.63	0.2	0.17	0.7	0.80	0.2
Trop.Roots	0.59	0.1	0.00	0.0	0.59	0.1
Date Palm*	0.03	0.0	0.24	0.9	0.27	0.1
Sugar Cane	0.16	0.0	0.09	0.3	0.24	0.1
Others	43.83	10.5	3.11	12.1	46.94	10.6
Grasslands	254.86	61.1	0.00	0.0	254.86	57.5
Total	417.36	100.0	25.74	100.0	443.10	100.0
Total without grasslands	162.50		25.74		188.25	
Crop area considered before/after development (%)	21/73		51/88		23/75	

3
4

Table 2: Key parameters of agricultural trees and potatoes. **R**: representative tree/plant for parametrisation; **K_{est}**: tree density range; **HI**: harvest ratio/index; **T_b**: base temperature; **GDD**: growing degree day requirements to grow full leaf coverage (in deciduous trees) or to reach ripeness of fruit (in evergreen trees); **T_{lim}**: lower and upper coldest monthly mean temperature; **Ph_{opt}**: lower and upper temperature optimum for photosynthesis; **HE**: maximal height of tree; **Phu_{1/2}**: fraction of potential heat units accumulated in the first (1) or second (2) inflexion point of the optimal leaf area curve; **Lmax_{1/2}**: fraction of maximal leaf area index accumulated in the first (1) or second (2) inflexion point of the optimal leaf area curve; **Phusen**: fraction of growing season at which senescence becomes the dominant process; **Lai_{ha}**: fraction of leaf area index still present at harvest; **PHU**: potential heat units for ripeness; **WCF**: water content factor for conversion from dry to fresh matter.

Crop	R	Seasonality	K _{est} (trees ha ⁻¹)	HI (frac)	T _b (°C)	GDD (acc.°C)	Ph _{opt} (°C)	T _{lim} (°C)	HE (m)	WCF (% of DM)	
Citrus	Orange tree	Evergreen broadleaved	500-4600 ^g	0.5 ^h	18 ^a	1000 ⁱ	23 to 30 ^g	-10 to 40 ^g	10 ^j	13 ^k	
Olive trees	Olive Trees	Evergreen broadleaved	75-690 ^{n,o}	0.45 ^f	15 ^l	1000 ^p	15 to 30 ^{n,m}	-10 to 15 ^{n,m}	15 ^m	30 ^q	
Date palm	Date palm	Evergreen broadleaved	100-920 ^r	0.5 ^u	14 ^s	1700 ^u	25 to 35 ^s	-10 to 40 ^s	20 ^s	70 ^{s,t}	
Orchards	Apple tree	Deciduous broadleaved	325-2990 ^{aa}	0.49 ^v	7 ^a	400 ^x	15 to 25 ^a	-15 to 15 ^z	13 ^y	16 ^k	
Nut trees	Almond tree	Deciduous broadleaved	100-920 ^{ac}	0.6 ^w	7 ^{ab}	300 ^{ad}	20 to 25 ^{ab}	-10 to 15 ^{ab}	10 ^{ae}	90 ^{af}	
Grapes	Vine plants	Deciduous broadleaved	2000-15000 ^{ak}	0.6 ^{al}	10 ^{ai}	300 ^{ah}	17 to 20 ^{ag}	10 to 15 ^{ag}	2 ^{aj}	20 ^k	
Cotton	Cotton plants	Deciduous broadleaved	20000-184000 ^{aq}	0.19 ^{am}	15 ^{ao}	300 ^{ao}	16 to 22 ^{ao}	-10 to 40 ^{ao}	3 ^{ap}	91 ^{an}	
Crop	R	LAI curve parameters (frac)				T _b	PHU	Ph _{opt}	WCF		
		Phu_1	Lmax_1	Phu_2	Lmax_2	Phusen	Lai_ha	(°C)	(acc.°C)	(°C)	(% of DM)
Potatoes	Potatoes	0.15 ^a	0.01 ^a	0.5 ^a	0.95 ^a	0.9 ^a	0.0 ^b	0.0 ^c	1500-2400 ^a	16 to 25 ^{a,d,e}	20 ¹¹
a Neitsch et al., 2004 (SWAT model). For Tb of citrus the Tb of poplar and oak was taken as a proxy.			p Orlandi et al., 2014 ⁴³ .			aa FAO, 2011.			Mis en forme : Espagnol (Argentine)		
b Gordon et al., 1997.			q Kailis & Harris, 2007.			ab Pontificia Universidad Católica de Chile, 2008.			Mis en forme : Espagnol (Argentine)		
c Haverkort & MacKerron, 1995 compiled different base temperatures in different studies and exposed that a base temperature of 0.0°C describes tuberisation rate well in temperate zones.			r Al-Khayri & Niblet, 2012.			ac Netafim, 2013.			Mis en forme : Allemand (Allemagne)		
d FAO, 2008.			s FAO, 2002.			ad Janick & Paull, 2008.					
e Ku et al., 1977.			t Elshibli, 2009.			ae Duke, 1983.					
f Morales Sierra, 2012.			u Large variations and lack of data lead to estimation of these parameters assuming relatively small, high yielding varieties of palms, flowering around one month, developing fruits in around 4 months, with 1000 kg weight and yields of 500 kg per palm.			af Alasalvar & Shahidi, 2008.					
g FAO, 2013a. For T _{lim} of citrus it was assumed that the rest can be induced by water deficit, not only by low temperatures.			v Zanotelli et al., 2013.			ag FAO, 2013c.					
h Cannell, 1985.			w Toki et al., 1989.			ah Meier, 2001.					
i Based on FAO, 2013a information that indicates that citrus need 7 to 14 months from flowering to maturity.			x Wünsche & Lakso, 2000.			ai Strick, 2011.					
j Orwa et al., 2009.			y Parametrised as standard apple tree and not dwarf.			aj Ministry of agriculture, food and rural affairs, 2013.					
k Bastin & Henken, 1997.						ak Slovak Wine Academy Pezinok, 2009.					
l Aguilera et al., 2014 ⁴³ .						al Zamski & Schaffer, 1996.					
m California Rare Fruit Growers, 1997.						am Sakin, 2012.					
n FAO, 2013b.						an Yan et al., 2007.					
						ao Tsiros et al., 2009.					
						ap Kiranga, 2013. In wild conditions cotton can be up to 5 meters high. In crops, up to 1.5 meters.					
						aq Wright et al., 2004; Paytas, 2011. Planting density varies largely.					
o Roussos, 2007.			z Perry, 2011. Apple, pear and cherry trees can survive lower winter temperatures, but other fruit trees (i.e. fig, peach and apricot trees) cannot.								




Figure 1: Scatterplot of LPJmL-simulated yields (averages over 2000–2009) versus reported yields from FAOSTAT for Mediterranean countries. The line represents the 1:1 line. **FM** stands for fresh matter. The **radius of the bubbles** indicates the relative size of the harvested area in the respective country.




Figure 2: LPJmL-simulated net irrigation water requirements (NIR), as average over the period 2000-2009.




Figure 3: Comparison of irrigation consumptive water use from Siebert et al. (2010) and net irrigation water requirements computed in this study as percentage of the Siebert et al. (2010) values. Negative (positive) values indicate higher (lower) values in their study.




Figure 4: Scatterplot of LPJmL-simulated gross irrigation water requirements (hm^3 , averages over 2000–2009) and the estimates presented by other studies. The line represents the 1:1 line. Egypt and Spain have been coloured in red and green, respectively, to visualize the disparities between different studies. (Note that Cánovas Cuenca, 2013, assumes a fixed requirement of $6176 \text{ m}^3 \text{ ha}^{-1}$).




Figure 5: Difference in LPJmL-simulated soil organic carbon (average 2000 to 2009) before and after the implementation of agricultural trees described in this study. Negative (positive) values indicate that the development decreased (increased) soil organic content.




Figure 6: Difference between the HWSD soil carbon density and LPJmL-simulated soil carbon density before and after the model development presented here ($(LPJmL_{Old} - HWSD) / (LPJmL_{New} - HWSD)$). Positive values indicate an improvement in simulation of soil carbon stock due to the implementation of Mediterranean crops described in the present study.