

Simulating the thermal regime and thaw processes of ice-rich permafrost ground with the land-surface model CryoGrid 3

Westermann, Sebastian¹, Langer, Moritz^{2,3,4}, Boike, Julia², Heikenfeld, Max², Peter, Maria^{1,2}, Eitzelmüller, Bernd¹, and Krinner, Gerhard^{3,4}

¹Department of Geosciences, University of Oslo, P.O. Box 1047, Blindern, 0316 Oslo, Norway

²Alfred Wegener Institute Helmholtz Center for Polar and Marine Research, Telegrafenberg A43, 14473 Potsdam, Germany

³CNRS, LGGE (UMR5183), 38041 Grenoble, France

⁴Université Grenoble Alpes, LGGE (UMR5183), 38041 Grenoble, France

Correspondence to: Sebastian Westermann
(sebastian.westermann@geo.uio.no)

Abstract. Thawing of permafrost in a warming climate is governed by a complex interplay of different processes of which only conductive heat transfer is taken into account in most model studies. However, observations in many permafrost landscapes demonstrate that lateral and vertical movement of water can have a pronounced influence on the thaw trajectories, creating distinct landforms like thermokarst ponds and lakes even in areas where permafrost is otherwise thermally stable. Novel process parameterizations are required to include such phenomena in future projections of permafrost thaw and hereby triggered climatic feedbacks. In this study, we present a new land-surface scheme designed for permafrost applications, CryoGrid 3, which constitutes a flexible platform to explore new parameterizations for a range of permafrost processes. We document the model physics and employed parameterizations for the basis module CryoGrid 3, and compare model results with in-situ observations of surface energy balance, surface temperatures, and ground thermal regime from the Samoylov permafrost observatory in NE Siberia. The comparison suggests that CryoGrid 3 can not only model the evolution of the ground thermal regime in the last decade, but also consistently reproduce the chain of energy transfer processes from the atmosphere to the ground. In addition, we demonstrate a simple 1D parameterization for thaw process in permafrost areas rich in ground ice, which can phenomenologically reproduce both formation of thermokarst ponds and subsidence of the ground following thawing of ice-rich subsurface layers. Long-term simulation from 1901-2100 driven by reanalysis data and climate model output demonstrate that the hydrological regime can both accelerate and delay permafrost thawing. If meltwater from thawed ice-rich layers can drain, the ground subsides while at the same time the formation of a talik is delayed. If the melt-

water pools at the surface, a pond is formed which enhances heat transfer in the ground and leads to the formation of a talik. The model results suggest that the trajectories of future permafrost thaw are strongly influenced by the cryostratigraphy, as determined by the late quaternary history of a site.

1 Introduction

25 In the past decade, the fate of permafrost areas in a changing climate has rapidly moved into the
focus of climate researchers. This has led to significant improvements in the representation of per-
mafrost and frozen ground processes in the land-surface schemes employed in Earth System mod-
eling schemes (e.g. Lawrence et al., 2008). These efforts have to a large part been motivated by the
potential release of the greenhouse gases carbon dioxide and methane from thawing carbon-rich per-
mafrost soils (Walter et al., 2006; Zimov et al., 2006). However, including this possibly significant
30 climatic feedback (Schaefer et al., 2011) in future climate predictions requires a correct description
of both the climate forcing of the global permafrost area and the physical processes governing the
transformation of the local permafrost systems. However, there is a striking gap between the tempo-
ral and spatial scales on which the forcing and the impacts occur (e.g. Lee et al., 2014). This implies
35 that the actual mobilization of carbon is not directly accessible through process modeling on large
grid cells, as it is generally done in atmospheric modeling. As an example, thermokarst and active
layer detachments can trigger strong mass wasting of ice- and organic-rich soils in time periods of
days to years at spatial scales of meters to hundreds of meters. Carbon fluxes from hotspots can sig-
nificantly exceed the background fluxes (e.g. Walter et al., 2006; Abnizova et al., 2012; Langer et al.,
40 2015). Integrating such highly localized and non-linear processes in large-scale grid-based models
requires the development of up- and downscaling strategies, which, despite of the emerging stronger
collaboration between field scientists and modelers, is still in its infancy.

Many permafrost areas are dominated by ice-rich ground (Brown et al., 1997). Due to ice wedges,
segregation ice or ice lenses, the volumetric ice content can exceed the natural pore volume of the
45 soil (i.e. excess ground ice) which in case of thawing will lead to a settling and consolidation of the
ground material. The released meltwater is either lost as surface or subsurface runoff leading to an
effective subsidence of the surface, or it pools up, thus forming a thermokarst pond or lake. In either
case, the ground and surface properties are modified by thawing of excess ground ice which can in
turn influence the trajectories of subsequent permafrost thaw. The vast lowland permafrost areas in
50 Siberia, Alaska and Canada dominated by thermokarst lakes and depressions are ample evidence
that thawing of excess ground ice has played a major role in the landscape development throughout
the Holocene. To predict future landscape development in a warming climate which is intimately
linked to the carbon cycle of these areas new and robust model approaches are needed which can
potentially be implemented in the land-surface schemes of Earth System Models (ESMs) used for
55 climate projections.

At the moment, there are two main directions in permafrost modeling on larger spatial scales: ded-
icated ground thermal models focus on the physical variables characterizing the state of the per-
mafrost, and have been successfully applied to map permafrost distribution (e.g. Gislås et al., 2013;
Zhang, 2013; Zhang et al., 2013; Westermann et al., 2013; Fiddes et al., 2015) and derive the ground
60 temperature evolution for past and future climate conditions (e.g. Zhang et al., 2008; Jafarov et al.,

2012). They are characterized by vertical modeling domains of one hundred to several hundred meters with a vertical resolution of centimeters within the active layer. As a consequence, they provide a sufficient accuracy to resolve the annual dynamics of active layer thawing and refreezing, as well as the evolution of ground temperatures in deeper layers. For validation purposes, they can thus
65 be by directly compared to field measurements, which can usually be reproduced to within 0.2 m (active layer thickness) or 1.5 K (ground temperature), depending on the type of input data (e.g. Langer et al., 2013; Westermann et al., 2013; Jafarov et al., 2012).

On the other hand, ESM land-surface schemes are directed towards adequately providing the lower boundary for the atmospheric transfer scheme, i.e. the surface fluxes of radiation, momentum and
70 sensible and latent heat. There is a huge spread in the modeled present permafrost area in the CMIP5 results, with values between 1.4 (factor 10 underestimation) and 27.3 (factor 2 overestimation) million km² (Koven et al., 2013). While latent heat effects due to soil freezing and thawing have been included in a number of ESMs (e.g. Gouttevin et al., 2012), the conductive heat transfer in the ground remains poorly represented due to the limited number of soil grid cells and insufficient depth of the
75 soil column. Therefore, projections on timescales of many decades over which permafrost thaw will occur can be strongly biased, which may at least partly explain the large inter-model spread in the sensitivity of permafrost extent, ground temperatures and active layer depth towards climate change (Koven et al., 2013).

In this study, we describe a new permafrost modeling scheme, CryoGrid 3, which aims to narrow the
80 gap between traditional transient permafrost models and land-surface schemes employed in Earth System modeling. CryoGrid 3 adds land-atmosphere coupling to the subsurface heat transfer model CryoGrid 2 (Westermann et al., 2013), which has been employed to simulate ground temperatures in Norway (Westermann et al., 2013) and NE Greenland (Westermann et al., 2015). In this way, it is possible to run CryoGrid 3 with the same forcing data as ESM land-surface schemes and compare
85 results of present and future simulations of the ground thermal state. Moreover, the numerical structure is designed to be as simple as possible which eases the requirements for adding and investigating parameterizations for permafrost processes. Following a similar philosophy as Essery (2015), CryoGrid 3 is dedicated to “active users” with the programming skills to modify and extend the code, rather than being a “black box tool” with a predefined and thus limited set of options. To illustrate
90 the potential of the model scheme, we implement simple parameterizations for ground subsidence and initial thermokarst formation and demonstrate the effect on future simulations of the ground thermal regime for a permafrost site in NE Siberia.

2 CryoGrid 3 - model description

CryoGrid 3 builds upon the subsurface thermal model CryoGrid 2 (Westermann et al., 2013), supplemented by the calculation of the surface energy balance as upper boundary condition, as well

as modifications of the snow scheme. As in CryoGrid 2, the energy transfer in the ground is governed by heat conduction, while soil hydrology is not explicitly accounted for. An overview of the processes implemented in the different models is given in Table 1. In the following, we provide a detailed description of the CryoGrid 3 model physics and the numerical scheme.

100 2.1 Driving variables

CryoGrid 3 is driven by time series of (i) air temperature T_{air} [$^{\circ}\text{C}$], relative or absolute humidity (RH [-] or q [-]) and wind speed u [ms^{-1}] at a known height h [m] above ground, (ii) incoming short-wave and long-wave radiation (S_{in} and L_{in}) [Wm^{-2}], (iii) air pressure p [Pa], and (iv) rates of snowfall and rainfall (P_s and P_r) [ms^{-1}].

105 2.2 The surface energy balance

The energy input to the uppermost grid cell is derived from the surface energy balance, i.e. the fluxes of short-wave radiation ($S_{\text{in}}, S_{\text{out}}$) and long-wave radiation ($L_{\text{in}}, L_{\text{out}}$), as well as the sensible, latent and ground heat fluxes, Q_h, Q_e and Q_g (see Sect. 2.4 for the latter). In the following, superscript numbers denote the number of a grid cell (increasing with depth, with grid cell 1 located at the surface). The equations follow Foken (2008), with variables and constants explained in Appendix A. The energy input to the uppermost grid is computed as:

$$\frac{\partial E^1}{\partial t} = S_{\text{in}} + S_{\text{out}} + L_{\text{in}} + L_{\text{out}} + Q_h + Q_e + Q_g. \quad (1)$$

While incoming short-wave and long-wave radiation are directly provided (Sect. 2.1), the other fluxes are parameterized based on the driving data sets. With the albedo α , the outgoing short-wave is calculated as

$$S_{\text{out}} = -\alpha S_{\text{in}}, \quad (2)$$

while the outgoing long-wave radiation L_{out} is derived from Kirchhoff's and Stefan-Boltzmann Law as

$$L_{\text{out}} = (\varepsilon - 1)L_{\text{in}} - \varepsilon\sigma (T^1 + 273.15\text{K})^4. \quad (3)$$

120 The turbulent fluxes of sensible and latent heat, Q_h and Q_e , are parameterized in terms of gradients of air temperature and absolute humidity (see Appendix B) between the air at height h above ground and at the surface following Monin-Obukhov similarity theory (Monin and Obukhov, 1954):

$$Q_h = -\frac{\rho_a c_p}{r_a^{\text{H}}} (T_{\text{air}}(h) - T^1) \quad (4)$$

$$Q_e = -\frac{\rho_a L_{\text{lg}}}{r_a^{\text{W}} + r_s} (q(h) - q(z_0)) \quad (5)$$

125 The resistance terms r constitute the aerodynamic resistance, r_a , characterizing the strength of the turbulent exchange, and the surface resistance against evapotranspiration, r_s . The latter is an empirical parameter that can be adjusted to account for the fact that the water vapor pressure above soil surfaces is lower than the saturation vapor pressure above a water surface (Eq. B3) for non-saturated surface soils.

130 The aerodynamic resistances r_a for sensible heat H and latent heat W,

$$r_a^{\text{H,W}} = (\kappa u_*)^{-1} \left[\ln \frac{h}{z_0} - \Psi_{\text{H,W}} \left(\frac{h}{L_*}, \frac{z_0}{L_*} \right) \right], \quad (6)$$

account for the atmospheric stability by including the integrated atmospheric stability functions $\Psi_{\text{H,W}}$ which describe deviations from the logarithmic profile of the neutral atmospheric stratification (Appendix C). They are inversely proportional to the friction velocity

$$135 \quad u_* = u(h) \kappa \left[\ln \frac{h}{z_0} - \Psi_M \left(\frac{h}{L_*}, \frac{z_0}{L_*} \right) \right]^{-1}, \quad (7)$$

which depends on the wind speed u and the integrated stability function for momentum, Ψ_M (Appendix C). The variable L_* is the Obukhov length, which in itself is a function of the turbulent fluxes Q_h and Q_e as well as the friction velocity u_* :

$$L_* = \frac{\rho_a c_p (T_{\text{air}}(h) + 273.15 \text{ K})}{\kappa g} \frac{u_*^3}{Q_h + 0.61 \frac{c_p}{L_{\text{lg}}} (T_{\text{air}}(h) + 273.15 \text{ K}) Q_e}. \quad (8)$$

140 Eqs. 1 to 8 thus constitute a coupled non-linear equation system, for which unique solutions Q_h , Q_e , u_* , and L_* exist.

The surface emissivity ε , the roughness length z_0 and the surface resistance against evapotranspiration r_s are assigned different values, depending on whether the surface is snow-free or covered by snow. The same applies to the albedo α for which a constant value is used for snow-free surfaces,

145 while for snow-covered surfaces the albedo changes with the age of the surface snow (Sect. 2.3).

2.3 Surface energy balance of snow

For snow-covered surfaces, the governing equations for the surface energy balance remain widely unchanged, with a few notable exceptions: first, penetration of short-wave radiation in the snow is accounted for by assuming exponential damping with depth (e.g. O'Neill and Gray, 1972). The short-wave radiation flux at depth d below the snow surface is given by

$$S_{\text{in}}(d) = (1 - \alpha_{\text{snow}}) S_{\text{in}}(0) e^{-\beta_{\text{SW}} d}, \quad (9)$$

with a bulk value for the entire spectral range for the SW-radiation extinction coefficient β_{SW} [m^{-1}] assumed. At the bottom of the snow layer, the remaining short-wave radiation flux is assigned as

155 energy input to the uppermost soil grid cell.

The albedo of the snow is calculated as a function of the the difference between time t and the time t_{SF} when the last significant snow fall occurred, following the formulations used in the ERA–interim reanalysis (Dee et al., 2011; ECMWF, 2006). The time of the last significant snow fall t_{SF} is set to t if the snowfall in the period $[t - 1 \text{ d}, t]$ exceeds a snow water equivalent of SWE_{SF} . Furthermore, 160 the snow albedo is set to the albedo of fresh snow, α_{max} , from which it decreases with time towards the albedo of old snow, α_{min} . For non-melting conditions, a linear decrease is assumed,

$$\alpha_{\text{snow}} = \begin{cases} \alpha_{\text{min}} - \tau_{\alpha, f}(t - t_{\text{SF}})(\alpha_{\text{max}} - \alpha_{\text{min}}) & \text{for } \tau_{\alpha, f}(t - t_{\text{SF}}) \leq 1 \\ \alpha_{\text{min}} & \text{for } \tau_{\alpha, f}(t - t_{\text{SF}}) > 1 \end{cases}, \quad (10)$$

while an exponential decrease is assumed for melting snow,

$$\alpha_{\text{snow}} = \alpha_{\text{min}} + e^{-\tau_{\alpha, m}(t - t_{\text{SF}})}(\alpha_{\text{max}} - \alpha_{\text{min}}). \quad (11)$$

165 The values for the time constants $\tau_{\alpha, f}$ and $\tau_{\alpha, m}$ [s^{-1}] (Table 4) are selected according to ECMWF (2006) and correspond to an albedo change of 0.1 in 12.5 days for non-melting conditions and an e-folding time of about 4 days for melting conditions.

According to Eq. 1, positive temperatures can occur for snow grid cells, which will trigger melting of the snow. In this case, the temperature of the concerned grid cell is set to $T = 0^\circ\text{C}$, and the energy 170 input $\partial E_i / \partial t$ is partitioned into energy leading to a temperature change and energy leading to snow melt, as detailed in Sect. 2.5. For the uppermost grid cell, all components of the surface energy balance are calculated according to Eqs. 3-8 with a fixed temperature $T_1 = 0^\circ\text{C}$.

2.4 Subsurface heat transfer

The subsurface thermal scheme of CryoGrid 3 is based on conductive heat transfer as given by 175 Fourier’s Law,

$$c_{\text{eff}}(z, T) \frac{\partial T}{\partial t} - \frac{\partial}{\partial z} \left(k(z, T) \frac{\partial T}{\partial z} \right) = 0, \quad (12)$$

where the generation and consumption of latent heat due to the phase change of soil water is taken into account through an effective volumetric heat capacity $c_{\text{eff}}(z, T)$ [$\text{Jm}^{-3}\text{K}^{-1}$] featuring a sharp peak in the freezing range of the soil water (Westermann et al., 2013). Both the effective heat capacity and the thermal conductivity $k(z, T)$ [$\text{Wm}^{-1}\text{K}^{-1}$] of the soil are functions of the volumetric fractions of the soil constituents water, ice, air, mineral and organic (Cosenza et al., 2003; 180 Westermann et al., 2013). At the lower boundary of the soil domain, a “geothermal heat flux” F_{lb} is assigned, which is usually assumed constant over time.

Vertical or lateral movement of water is not considered in the here described basis version of Cryo- 185 Grid 3, so that the sum of the volumetric contents of ice and water remains constant (for simplicity, the densities of ice and water are assumed equal). The phase change of the soil water is determined by the freezing characteristic $\theta_w(z, T)$ (see Dall’Amico et al., 2011) in dependency of the soil type

(three classes, sand, silt and clay), which can change with depth. The employed parameterizations for $c_{\text{eff}}(z, T)$, $k(z, T)$ and $\theta_w(z, T)$ are identical to the ones employed in CryoGrid 2, documented in detail in Westermann et al. (2013). The subsurface thermal properties are specified as a stratigraphy of the volumetric contents of soil mineral and organic material, air and the sum of ice and water (as in Tables 2, 3).

2.5 Energy transfer and mass balance of the snow pack

In the current version of CryoGrid 3, a constant density ρ_{snow} is assumed for the entire snow pack, with the exception of density modifications through refreezing rain and melt-water (see below). A change in the snow water equivalent ΔSWE due to snowfall, sublimation or melt (Sect. 2.3) leads to a change in snow depth Δd_{snow} as

$$\Delta d_{\text{snow}} = \rho_{\text{snow}} / \rho_{\text{ice}} \Delta\text{SWE}. \quad (13)$$

To account for the build-up and disappearance of the snow cover, the position of the upper boundary is allowed to change dynamically by adding or removing grid cells. The initial temperature of a newly added grid cell is set to the air temperature T_{air} .

As in soil, a thermal model based on heat conduction is employed for the snow pack, in conjunction with a 1D “cold-hydrology” scheme for percolation of melt and rain water in snow, following Westermann et al. (2011). However, only a limited number of natural processes occurring in the snow pack are accounted for, and CryoGrid 3 is therefore not comparable to sophisticated snow schemes, such as SNOWPACK (Bartelt and Lehning, 2002; Lehning et al., 2002a, b) or CROCUS (Vionnet et al., 2012). In contrast to the soil domain, where latent heat effects are accounted for by the effective heat capacity, two state variables are required to describe the energy content of a snow grid cell, its temperature T and the volumetric water content θ_w . Similar to the soil domain, the principal means of energy transfer within the snow pack is conductive heat transfer. We define two auxiliary variables T^* and θ_w^* , which are equal to T and θ_w at the beginning of each time step (see Sect. 2.6). We subsequently update T^* according to the heat conduction equation

$$c_{\text{snow}}(z, T) \frac{\partial T^*}{\partial t} - \frac{\partial}{\partial z} \left(k_{\text{snow}}(z, T) \frac{\partial T^*}{\partial z} \right) = 0, \quad (14)$$

with the snow heat capacity c_{snow} and thermal conductivity k_{snow} being functions of the snow density ρ_{snow} as

$$c_{\text{snow}} = c_{\text{ice}} \frac{\rho_{\text{snow}}}{\rho_{\text{ice}}} \quad (15)$$

and (Yen, 1981)

$$k_{\text{snow}} = c_{\text{ice}} \left(\frac{\rho_{\text{snow}}}{\rho_{\text{water}}} \right)^{1.88}. \quad (16)$$

In this first step, θ_w^* is unchanged. Eq. 14 allows both for temperatures T^* above the melting point of ice, $T_m = 0^\circ\text{C}$, and for $T^* < T_m$ for non-zero water content $\theta_w^* \neq 0$. The temperature T and water

content θ_w of a cell are thus adjusted according to the energy content $E = c_{\text{snow}}(T^* - T_m) + L_{\text{sl}}\theta_w^*$,

$$(T|\theta_w) = \begin{cases} \left(T_m \left| \theta_w^* + (T^* - T_m) \frac{c_{\text{snow}}}{L_{\text{sl}}} \right. \right) & \text{for } E > 0 \\ \left(T^* + \theta_w^* \frac{L_{\text{sl}}}{c_{\text{snow}}} \left| 0 \right. \right) & \text{for } E \leq 0 \end{cases} \quad (17)$$

Hereby, $E = 0$ corresponds to ice (i.e. $\theta_w = 0$) at temperature T_m . Since a decrease in the water content, $\theta_w < \theta_w^*$, corresponds to refreezing of water, the ice content θ_i is increased by $\theta_w^* - \theta_w$.

225 The water flux at the upper boundary is given by the rainfall rate, which is added to the water content of the first grid cell. If the water content in a grid cell exceeds the field capacity of the snow, θ_w^{fc} (i.e. the maximum volumetric water content the snow can hold) following melt or rainfall, the water is infiltrated in the snow cover in a routing scheme similar to the one employed in Westermann et al. (2011). Hence, a snow grid cell with temperature T can hold the field capacity, plus the amount of
 230 water required to increase its temperature to T_m , i.e. $\theta_w^{\text{fc}} - (T - T_m) c_{\text{snow}} / L_{\text{sl}}$. Excess water is routed to the next cell, until the cell at the bottom of the snow pack is reached, where infiltration is allowed until saturation. The water then starts to pool up from bottom to top up, until it reaches the top of the snow pack, where excess water is removed from the system as runoff. The infiltration routine updates the water contents θ_w for each grid cell. In a final step, the temperature T is recalculated by
 235 applying Eq. 17 to account for the refreezing of the infiltrated water.

2.6 Time integration

In CryoGrid 3, only the physical process of heat conduction and the associated surface energy balance are integrated in time, while all other processes, such as the infiltration and refreezing of melt-
 240 water (Sect.2.5), occur instantly at the end of each time step in the form of a “physical consistency check”. To make the code intuitive to use and modify, the simplest possible time integration scheme is employed, first-order forward Euler. Since it is neither computationally efficient nor numerically stable, an adaptive integration time step is used which prevents too large changes of the internal energy of a grid cell and thus allows a certain degree of control over the accuracy of the solution.

245 As a first step, the method of lines (Schuesser, 1991) is used to discretize the spatial derivatives in Eqs. 12 and 14. The change of the internal energy E^i of grid cell i ($i \neq 1, N$) over time is given by

$$\frac{\partial E^i}{\partial t} = \frac{1}{\Delta z^i} \times \left(k^{i-1,i}(t) \frac{T^{i-1}(t) - T^i(t)}{\Delta z^{i-1,i}} - k^{i,i+1}(t) \frac{T^i(t) - T^{i+1}(t)}{\Delta z^{i,i+1}} \right), \quad (18)$$

where Δz^i is the thickness of the i -th grid cell, while $\Delta z^{i-1,i}$ is the spacing between and $k^{i-1,i}$ the thermal conductivity assigned to the midpoint of the $i-1$ -th and the i -th grid cells. For the lowermost
 250 grid cell, $i = N$, the geothermal heat flux is used so that

$$\frac{\partial E^N}{\partial t} = \frac{1}{\Delta z^N} \times \left(k^{N-1,N}(t) \frac{T^{N-1}(t) - T^N(t)}{\Delta z^{N-1,N}} + F_{\text{lb}} \right). \quad (19)$$

For the uppermost grid cell, $i = 1$, the discretized form of Eq. 1 is used, i.e.

$$\frac{\partial E^1}{\partial t} = \frac{1}{\Delta z^1} \times \left(S_{\text{in}} + S_{\text{out}} + L_{\text{in}} + L_{\text{out}} + Q_h + Q_e - k^{1,2}(t) \frac{T^1(t) - T^2(t)}{\Delta z^{1,2}} \right). \quad (20)$$

Using the first-order forward Euler scheme to integrate Eqs. 18-20 in time (as given by the first term
255 in Eqs. 12 and 14), the temperature of the grid cell i at time $t + \Delta t$ is obtained as

$$T^i(t + \Delta t) = T^i(t) + \frac{\Delta t}{c_{\text{eff}}^i(T^i(t))} \frac{\partial E^i(t)}{\partial t}. \quad (21)$$

The integration time step Δt is adapted so that change of the internal energy of a grid cell between
time steps does not exceed a threshold ΔE_{max} as

$$\Delta t = \begin{cases} \Delta t_{\text{max}} & \text{for } \max \left(\Delta t_{\text{max}} \frac{\partial E^i}{\partial t} \right) \leq \Delta E_{\text{max}} \\ \Delta E_{\text{max}} / \max \left(\frac{\partial E^i}{\partial t} \right) & \text{for } \max \left(\Delta t_{\text{max}} \frac{\partial E^i}{\partial t} \right) > \Delta E_{\text{max}}. \end{cases} \quad (22)$$

260 To avoid the computational costs of solving the coupled equation system for the surface energy
balance (Eqs. 1-8), the Obukhov length L_* of the time step t is employed so that the friction velocity
 u_* and the fluxes Q_h and Q_e can be computed for the time step $t + \Delta t$ following Eqs. 1-7. The
Obukhov length is subsequently updated with the new values of u_* , Q_h and Q_e of time step $t + \Delta t$.
While this procedure introduces numerical errors and potential instabilities, it is adequate for short
265 time steps Δt during which the components of the energy balance and the associated Obukhov
lengths do not change strongly.

2.7 CryoGrid 3 Xice - a simple representation of permafrost thaw processes in ice-rich permafrost

Excess ground ice melt: In many permafrost landscapes, ground layers super-saturated with ice occur,
270 i.e. the volumetric ice content exceeds the porosity of the matrix material when thawed. Such
layers are stable when frozen, but become unstable upon thawing, with the water from the melted
ground ice becoming mobile. If the meltwater can drain laterally, e.g in sloping terrain, this will result
in subsidence of the surface, while the water can pool up at the surface and create a thermokarst
pond in areas with poor drainage. Either process has the potential to drastically modify the surface
275 energy balance and ground heat transfer and thus change the trajectories of permafrost thaw considerably.
CryoGrid 3 Xice facilitates modeling of such phenomena by introducing vertical fluxes of
water when a grid cell supersaturated with ice thaws for the first time (Fig. 1). Hereby, we follow
the physically-based subsidence scheme proposed by Lee et al. (2014), but extend it further with a
simple representation of energy transfer within shallow water bodies. We emphasize that CryoGrid
280 3 Xice can only account for the melting of excess ground ice, but not to its aggregation. e.g. due to
long-term sedimentation processes, growth of peat, or formation of ice wedges, ice lens and segregation
ice.

In the following, we demonstrate the scheme for the case of grid cells with equal spacing (Fig. 1). For

each grid cell, a so-called natural porosity θ_{nat} (e.g. Hopkins, 1949; Nicou et al., 1993) is assigned
 285 which is the porosity of the consolidated soil after excess ground ice has melted and the meltwater
 has been removed. For saturated soils without excess ground ice, the natural porosity is equal to the
 ice/water content $\theta_{i,w}$. If a previously frozen grid cell with index i and $\theta_{\text{nat}} < \theta_i + \theta_w$ thaws for
 the first time (i.e. it reaches a temperature of more than 0°C and all ice has melted, Fig. 1 b₁/c₁), a
 column [m] of water of

$$290 \quad h_w^{i \rightarrow i-1} = \Delta z^i (\theta_w^i - \theta_{\text{nat}}^i) \quad (23)$$

is moved upwards to grid cell $i - 1$. To conserve volume, a column of mineral and organic ground
 material of

$$h_{m,o}^{i-1 \rightarrow i} = \Delta z^i \frac{\theta_{m,o}^{i-1}}{\theta_m^{i-1} + \theta_o^{i-1}} (\theta_w^i - \theta_{\text{nat}}^i) \quad (24)$$

is moved downwards from grid cell $i - 1$ to i which ensures that the porosity of grid cell i is now
 295 equal to the natural porosity (Fig. 1 b₂/c₂). The process is recursively repeated from the initial grid
 cell upwards, i.e. the increase of water and decrease of mineral/organic contents in grid cell $i - 1$
 will again trigger the condition $\theta_{\text{nat}} < \theta_i + \theta_w$ (Fig. 1 b₂/c₂). Hence, water is moved further upwards,
 while mineral and organic contents move down. When an unsaturated layer is reached, the air is dis-
 placed and moved upwards by the water added to the grid cell, similar to Eq. 23. If sufficient air is
 300 present in the soil column, repeated application of the excess ice routine can lead to an uppermost
 grid cell which consists entirely of air. In this case, the grid cell is removed from the soil domain
 (similar to the procedure during snow melt, Sect. 2.5) and the ground subsides by one grid cell. After
 all soil air is removed by this mechanism, ongoing melting of excess ice layers triggers the water to
 pool up at the surface (i.e. the uppermost grid cell consists of 100% water, Fig. 1 c₇). If this water
 305 is removed through lateral surface or subsurface runoff, the ground subsides (Fig. 1 c₇^S), while a
 thermokarst pond will form in flat landscapes where lateral fluxes of water are small (Fig. 1 c₇). A
 combination of both processes is possible in landscapes with pronounced microtopography, in that
 thawing ice-rich ground initially subsides until it reaches the level of a surrounding wetland, after
 which a thermokarst pond is formed. To simulate such effects, a water table depth can be defined in
 310 CryoGrid 3 Xice above which all surface water is removed, followed by the formation of a pond at
 the given water table depth.

When lateral water fluxes lead to ground subsidence, it is evident that this will also change the soil
 moisture regime in the active layer. However, in the absence of an explicit treatment of the water
 balance, we cannot simulate such effects in CryoGrid 3 Xice. For simplicity, we assume that the
 315 ground remains fully saturated after the excess ice meltwater reaches the surface, i.e. only surface
 water is removed. We assume the surface cover to be unaffected by excess ice meltwater, so that the
 parameters related to the surface energy balance, such as the albedo α and the roughness length z_0
 remain unchanged.

Energy transfer in water bodies: If water pools up and forms a thermokarst pond, the surface energy balance and the heat transfer in the ground are modified systematically (Boike et al., 2015). If the uppermost grid cell consists entirely of water, we therefore reduce the surface albedo and set the surface resistance to evapotranspiration to zero. Shallow water bodies such as thermokarst ponds and lakes are generally well mixed during summer (Boike et al., 2015) which strongly increases the energy transfer from the surface to the bottom of the water body. We phenomenologically simulate this process by assigning a uniform temperature to all unfrozen water layers when the uppermost grid cell is unfrozen. In a first step, the temperatures of all grid cells are advanced to the next time step by the normal heat transfer scheme of CryoGrid 3. Secondly, the weighted average of their temperatures is assigned to all unfrozen water grid cells, thus achieving a uniform profile while conserving the internal energy within the water column. When the surface freezes, a stable stratification forms in the water column which significantly reduces the heat transfer (Boike et al., 2015). If the uppermost grid cell is frozen, the normal heat transfer scheme of CryoGrid 3 is employed without modifications (i.e. without applying any mixing) which assigns the relatively low thermal conductivity of immobile water ($0.57 \text{ Wm}^{-1}\text{K}^{-1}$) to all water layers. In the model, melting of the winter ice cover in spring occurs from top to bottom, i.e. the ice does not float to the surface as observed in nature. However, since the mixing is applied to all unfrozen grid cells above and below the remaining ice layer, heat is transferred to the bottom of the ice layer and leads to a rapid melting from both sides. In essence, the normal heat conduction scheme of CryoGrid 3 is left unchanged for water bodies, with the only modification being the “mixing” of water temperatures for summer conditions. While this simple scheme can account for the strong asymmetry of the heat transfer into the water body between winter and summer, it excludes potentially important processes such as radiative heat transfer and buoyancy-driven thermal stratifications in the water column. In Sect. 5.1, the scheme is validated against in-situ measurements of lake bottom temperatures from Boike et al. (2015).

3 Study area

3.1 The Lena River Delta

At its mouth in the Laptev Sea, the Lena River forms one of the largest deltas in the Arctic, covering around $25\,000 \text{ km}^2$. Continuous cold permafrost with a mean annual near-surface ground temperature on the order of -10°C underlies the Lena Delta to about 400 and 600 m below the surface (Yershov et al., 1991). The landscape has been shaped by the river through erosion and sedimentation (Fedorova et al., 2015), but also permafrost-related thermokarst processes which have created a large number of water bodies. Within the modern delta, three main geomorphological units can be distinguished according to their late quaternary genesis which has led to distinctly different subsurface (cryo-)stratigraphies.

The first river terrace, which covers large parts of the eastern and central delta, is shaped by Holocene

river erosion and sedimentation processes. The sediment mainly consists of silty sands, while organic
355 matter has accumulated in thick peat layers (Schwamborn et al., 2002) which feature volumetric or-
ganic contents of 5-10% and mineral contents of 20-40% (Kutzbach et al., 2004; Zubrzycki et al.,
2012). Such sediments are in general super-saturated with ground ice, with average volumetric
ice contents reaching 60-80% and massive ice wedges extending up to 9 m deep in the ground
(Grigoriev et al., 1996; Schwamborn et al., 2002; Langer et al., 2013).

360 *The second river terrace* in the NW part of the delta was created in a continental setting before and
during the transition to the Holocene when the sea level was lower and the gradient of the Lena
River steeper than today. It therefore features coarse-grained sandy deposits which are characterized
by low ground ice and organic contents (Schwamborn et al., 2002). Near the surface, an organic up-
per horizon is often lacking and about 50% of the area is free of vegetation (Rachold and Grigoriev,
365 1999; Ulrich et al., 2009; Schneider et al., 2009). The second terrace is often referred to as “Arga
complex”, after Arga Island in the NW part of the delta.

The third river terrace is the oldest geomorphological unit of the Lena Delta, shaped by the ex-
tremely cold climate of the last glacial period when the area was not glaciated. Often referred to
as “Ice Complex” or “Yedoma”, the ground in the third terrace is extremely rich in ground ice and
370 organic matter, which has not been eroded by the Lena River in the Holocene (Grigoriev, 1993;
Zubrzycki et al., 2012). Massive ice wedges reach depths of 20-25 m (Schirrmeister et al., 2003;
Grigoriev, 1993; Schwamborn et al., 2002) and in many parts, the ground ice can reach volumetric
contents of 90% (Schirrmeister et al., 2011). The upper soil horizon features a thick organic layer.

The three geomorphological units offer the opportunity to conduct a sensitivity study along a strong
375 gradient of ground ice contents, with little to no excess ground ice in the second terrace, while the
first and third terrace feature medium and high excess ground ice contents, respectively.

3.2 The Samoylov permafrost observatory

Samoylov Island (72°22'N, 126°28'E) with its Russian-German research station has been in the fo-
cus of intense scientific investigations for more than two decades, with focus on energy and carbon
380 cycling (Abnizova et al., 2012; Kutzbach et al., 2004, 2007; Sachs et al., 2008, 2010; Wille et al.,
2008; Zhang et al., 2012), the characteristics of microbial communities (Wagner et al., 2007, 2009),
validation of remote sensing products (Langer et al., 2010) and ESM development (e.g Ekici et al.,
2014; Yi et al., 2014; Chadburn et al., 2015). The island is located in the south-central part of the
Lena River Delta in the first terrace within one of the main river channels (Sect. 3.1).

385 Samoylov Island provides the unique opportunity to force CryoGrid 3 with a long-term in-situ
record, while at the same time providing comprehensive high-quality observations for validation.
An automatic weather station (AWS) has been operational since 1998 (Boike et al., 2008), mea-
suring air temperature, net radiation, humidity and wind speed at hourly intervals. Since 2009, all
components of the radiation balance are measured, i.e. incoming and outgoing short and longwave

390 radiation. Eddy covariance measurements of sensible and latent heat fluxes are ongoing since 2007
in close proximity of the AWS (Langer et al., 2011a, b), and gaps in the time series of the climate
station have, when possible, been filled with data from the eddy covariance stations. Furthermore,
records of temporary climate stations operated in different time periods on Samoylov Island have
been used to fill gaps in the AWS record (for details, see Boike et al., 2013).

395 The spatial distribution and the onset and termination of the snow cover were recorded with an au-
tomated camera. Systematic measurements of thaw depth have been conducted since 2002 on a 150
point grid, using a steel rod pushed vertically into the soil to the depth at which ice-bonded soil pro-
vides firm resistance. The grid points on the polygonal tundra include land surfaces classified as dry
and wet tundra and overgrown water (Fig. 12 in Boike et al., 2013). The maximum thaw depth at
400 the end of the summer shows a large interannual variability, with an overall increase of thaw depths
since the start of the measurements in 2002. Active layer temperatures were measured since 2002
in three profiles installed across the low centered ice wedge polygon in the center, rim and slope
(the area with a very small topographic gradient between center and rim). The temperatures were
recorded using thermistors (107, Campbell Scientific Ltd.) which are calibrated at 0°C so that the
405 absolute error was less than 0.1°C over the temperature range $\pm 30^\circ\text{C}$. The temperature data used
for validation of CryoGrid 3 are from the deepest sensor at 0.40 m depth below surface in the center
of the polygon which is always water-saturated and mostly comprised of peat. Additional detailed
information concerning the climate, permafrost, land cover, vegetation, and soil characteristics on
Samoylov Island can be found in Boike et al. (2013).

410 **4 CryoGrid 3 runs for the Lena River Delta**

To evaluate the performance of CryoGrid 3 and demonstrate the excess ice scheme, two model runs
were performed: *validation runs* for the period 1979-2014 were forced by measured data from the
Samoylov AWS and downscaled ERA–interim reanalysis data to benchmark the model performance.
To assess the potential impact of excess ground ice on the trajectories of permafrost thaw, *long-term*
415 *thaw susceptibility runs* from 1901 to 2100 were conducted based on reanalysis data and climate
model output.

A first validation run is conducted for a subsurface stratigraphy typical for Samoylov Island on the
first river terrace so that it can be compared to in-situ observations. Since the assumed snow den-
sity has a pronounced influence on ground temperatures (see also Langer et al., 2013), we conduct
420 two runs each for confining values of 200 kg m^{-3} and 250 kg m^{-3} , the range obtained from in-situ
measurements (Boike et al., 2013). An additional run is conducted to benchmark the performance
of the simple water body scheme (Sect. 2.7) against measurements of lake bottom temperatures on
Samoylov Island (Boike et al., 2015). The model set-up is unchanged for this “lake validation run”,
but the subsurface stratigraphy of a well-developed thermokarst lake is applied.

425 The long-term thaw susceptibility runs investigate the influence of cryostratigraphy on thaw trajectories, using the three geomorphological units in the Lena Delta to perform a sensitivity analysis guided by real-world conditions. Hereby, two scenarios for future warming (RCP 4.5 and 8.5, see below) and two scenarios for ground drainage are considered: in the “subsidence” scenario, excess water pooling at the surface is completely removed (corresponding to good drainage), while no water
430 is removed for the “thermokarst” scenario so that a pond forms following thawing of excess ground ice (corresponding to poor drainage). For these runs, a snow density of 225kg m^{-3} is employed.

4.1 Surface, snow and ground properties

Subsurface properties: To investigate the effect of different (excess) ground ice contents in the long-term thaw susceptibility runs, a typical stratigraphy is compiled for each of the three river terraces,
435 oriented at available observations (Sect. 3.2). In the following, these are referred to as “Samoylov” (first terrace), “Arga” (second terrace), and “Ice Complex” (third terrace) stratigraphy .

The three stratigraphies are documented in Table 2. We assume that the terraces differ in their near-surface composition whereas in greater depth ($>20\text{m}$) the ground is assumed uniform with sandy sediments lacking excess ground ice (Schirrmeyer et al., 2011; Schwamborn et al., 2002).
440 For “Arga”, sandy mineral soil saturated with ice/water (but without excess ground ice) is assumed close to the surface, corresponding to the lack of an upper organic horizon and the general sparsity of vegetation (Sect. 3.1). For “Samoylov”, we assign a typical stratigraphy for a medium wet polygon center, in which soil temperature measurements employed for validation are conducted (Sect. 3.2). On top, a medium dry upper organic layer is assumed, followed by a saturated layer with a high wa-
445 ter/ice content which extends to a depth of 0.9m , well below the modeled and measured active layer thicknesses in 2010-2015 (Sect. 5). In this layer, we assume that the soil matrix structure of mineral and organic material can support the high soil porosity, so that thawing of ground ice does not lead to vertical or lateral water movement (natural porosity is equal to the true porosity, Table 2). This is changed in the layer below, extending to 9m , a typical depth of ice wedges (Sect. 3.1), in which
450 we assume the soil matrix can only support a porosity of 40%, so that the excess water becomes mobile upon thawing. The “Ice Complex” stratigraphy is similar to “Samoylov”, but features a drier upper organic layer followed by a saturated layer with high porosity which can yet be supported by the soil matrix. The super-saturated layer below features very high ice contents and extends to a typical depth of ice wedges in the third terrace (Sect. 3.1). For the “Samoylov” and “Ice Complex”
455 stratigraphies, the depth below which excess ground ice is assumed has been adjusted, so that it is below the modeled maximum thaw depth for the years 1901-present (Sect. 5) and thermokarst or subsidence do not occur before present-day.

For the validation runs (Table 3), we employ the “Samoylov” stratigraphy as described above, as well an estimated stratigraphy for the thermokarst lake “Sa_Lake_1” (Boike et al., 2015) on Samoylov
460 Island. For the latter, we prescribe the true water depth of 6m (Table 3), slightly deeper than the

thermokarst lake that would form after the melting of all the excess ground ice in the “Samoylov” stratigraphy (approx. 3.75 m depth). Furthermore, we assume that layers of excess ground ice are no longer present under the lake so that the lake depth remains constant over time.

As far as possible, surface properties derived from measurements on Samoylov Island are employed (Table 4), as published in Langer et al. (2011a, b) and Boike et al. (2013). In addition, the maximum snow depth in CryoGrid 3 is restricted to 0.45 m, the approximate height of the polygon rims above the centers. Snow depths observations of Boike et al. (2013) show that the maximum snow height in polygon centers rarely exceeds this values, since further snowfall is largely blown away by consistently strong winds.

4.2 Model forcing data

Validation run: The forcing data for the validation run are based on the available in-situ record of incoming short- and long-wave radiation S_{in} and L_{in} , air temperature T_{air} , absolute humidity q (computed from relative humidity RH , Appendix B) and wind speed u from the AWS on Samoylov Island. To fill remaining gaps in the measurement time series and to generate a record for model spin-up prior to the measurement period, surface fields of the ERA-interim reanalysis (Simmons et al., 2007; Dee et al., 2011) were statistically downscaled for Samoylov Island. For each day of the year, a linear regression was fitted between the ERA-interim variable and the available measurements for a period of one month around the respective date. Using the regression results for each day of year (if a measurement was not available), a time series from June 1, 1979, to December 31, 2014, was compiled for the the above mentioned variables. The rates of snowfall were used directly from the reanalysis, since reliable measurements are lacking. However, the snow depth is limited to the approximate height of the polygons (Sect. 4.1), which ensures realistic maximum annual snow depths independent of the snowfall rates. The largest uncertainties can hence be expected during the build-up phase of the snow cover, before the threshold height is reached.

CryoGrid 3 was initialized to steady-state conditions of the forcing for the first ten model years by re-running the model with the 1979-1989 forcing until the change of temperatures in the uppermost 100 m was less than 0.05°C between subsequent 10-year runs. The first initial condition was determined as described in Westermann et al. (2013).

Long-term thaw susceptibility run: In order to force the model from 1901 to 2100 with a consistent data set, model-derived forcing data building on an anomaly approach were employed which were subsequently downscaled for Samoylov Island similar to the validation run. Past and present-day forcing data were taken from the CRU-NCEP dataset (<http://dods.extra.cea.fr/data/p529viov/cruncep/>) which combines high-frequency variability from the National Centers for Environmental Prediction/National Center for Atmospheric Research reanalysis (Kalnay et al., 1996) with the monthly mean climatologies from the Climatic Research Unit temperature and precipitation datasets (Harris et al., 2014). The meteorological forcing for future climate change experiments was constructed following

the Representative Concentration Pathway (RCP) 4.5 and 8.5 climate change projections obtained with the CCSM4 coupled climate model (Meehl et al., 2012). Monthly climate anomalies (for the meteorological state variables: air temperature, specific humidity, surface pressure and wind speed) and scale factors (for the meteorological flux variables: precipitation, incoming shortwave and long-wave radiation) were extracted from this model output (http://www.earthsystemgrid.org/dataset/ucar.cgd.cesm4.permafrostRCN_pro)

500 The high-frequency variability in the future meteorological forcing was obtained by looping over the 1996-2005 CRU-NCEP forcing data. The base climatological period which is used to calculate the monthly anomalies/scale factors is also 1996-2015. These forcing data were also used in Koven et al. (2015) and in an ongoing international intercomparison exercise within the framework of the Permafrost Carbon Network (<http://www.permafrostcarbon.org/>, work in progress).

As for the validation run, the data from the closest model grid cell were downscaled for Samoylov Island using the in-situ record from the AWS. Since the match with the measured data was significantly worse compared to the ERA-interim reanalysis (in particular for the CCSM4 data, but also for CRU-NCEP), the entire data set for the overlap periods 2010-2012 for CRU-NCEP and 2012-2014 for CCSM4 was used to determine a linear regression for L_{in} , T_{air} and q . For the incoming short-wave radiation S_{in} which follows a defined diurnal pattern, the procedure was conducted separately for each of the time steps 0000, 0600, 1200 and 1800. Values with $S_{in}=0$ were excluded from the regression.

510 Downscaling of wind speeds was challenging, since there was no statistically significant correlation to measured data. Therefore, the distribution of wind speeds from CRU-NCEP/CCSM4 was scaled to fit the distribution obtained from the in-situ record. For this purpose, the wind speeds were sorted in ascending order in bins containing 20% of the available data each. The CRU-NCEP/CCSM4 data were then scaled linearly so that the average wind speed of each bin matched the average wind speed of the in-situ data in the respective bin. As for the validation run, the snowfall data from CRU-NCEP/CCSM4 were used without modification, but as in the validation run, the maximum height of the snow cover is limited to the approximate height of the polygon rims. The model initialization procedure was identical to the validation run.

520

5 Model results

5.1 Comparison to in-situ data for Samoylov Island

525 *Polygon center stratigraphy*: The model results of the validation run for the polygon center stratigraphy (Table 3) are compared to published in-situ measurements of the surface energy balance and the ground thermal regime from Samoylov Island (Langer et al., 2011a, b; Boike et al., 2013). CryoGrid 3 is very well capable of reproducing measured surface temperatures, except for a slight cold-bias for temperatures colder than -30°C (Fig. 2). The spread of the data points around the 1:1 line could to some extent be explained by small temporal offsets between the “real” measured data and the ERA-interim reanalysis upon which part of forcing data are based (Sect. 4.2). Furthermore, the Bowen

530

ratio (Q_h/Q_e) during the summer months agrees well with published values from Langer et al. (2011a), who found average values of 0.35 for 2007 and 0.50 for 2008. The modeled values for the corresponding periods are 0.3 and 0.55, respectively, independent of the assumed snow density.

535 Fig. 3 displays a comparison of the absolute flux values for a summer and winter period. CryoGrid 3 is very well capable of reproducing the observed energy partitioning at the surface, although the modeled values in winter are slightly smaller in magnitude. The satisfactory agreement of surface energy balance and surface temperatures indicates that CryoGrid 3 can consistently reproduce the land-atmosphere interactions on Samoylov Island, despite the simple treatment of summer evapo-
540 transpiration with a constant value for the surface resistance against evapotranspiration r_s .

Boike et al. (2013) provide snow start and end dates, as determined visually from images taken by an automatic camera system. A comparison to the corresponding dates modeled in CryoGrid 3 is not entirely straight-forward in particular for the snow melt, since the dates obtained from the camera images represent spatial averages, while the model calculates the snowmelt for only a single
545 realization of snow depth. On the average of the years 2001-2010, CryoGrid 3 captures the start and the end of the snow season quite well: the measured and modeled onset date of the snow cover deviate by $+3.8 \pm 19.2$ days, while the snow end dates differ by $+1.7 \pm 12.2$ days. However, the large standard deviations are evidence of significant deviations in some years, which can amount to up to two weeks for the snowmelt and more than 1 month for the onset of the snowcover. The latter
550 is completely determined by the forcing data of snowfall (which are obtained from the reanalysis), so that the bias is explained by shortcomings in distinguishing snow- and rainfall in the reanalysis in fall. Significant biases in the snowmelt date can be explained by two reasons: first, the modeled maximum snow depth may be biased due to an erroneous amount of snowfall (which is again determined from the reanalysis). Secondly, the melting of the snow cover usually occurs in episodes
555 which are interrupted by cold periods, during which the melting ceases or even new snow falls. Even a small bias in the melt rates can therefore cause a considerable deviation in the termination of the snowmelt, for instance when the snow in the model erroneously persists until the beginning of a cold spell and only melts in the next warm period.

CryoGrid 3 results of the validation run are further compared to measurements of ground temperatures and thaw depths, with a particular focus on the annual dynamics and interannual differences.
560 Fig. 4 displays the comparison of modeled ground temperatures at a depth of 0.4 m in a wet polygon center. In most years, the simulations can successfully reproduce the annual temperature range of about 20 to 25°C, while the length of the zero curtain, when ground temperatures are confined to 0°C, is in some years underestimated by up to one week. While the reason for this is not entirely
565 clear, we note that the uncertainty of the snow forcing data is largest during the onset and build-up of the annual snow cover (Sect. 4.2) which could at least partly explain biases of modeled ground temperatures during fall freezing. An example is the considerably too early end of of the zero curtain in fall 2010 in the modeled active layer temperatures (Fig. 4), which coincides with a modeled onset

of the snow cover 27 days later than observed by Boike et al. (2013).

570 In the majority of the considered years, CryoGrid 3 results successfully capture both the absolute values of thaw depths and the timing of the thaw progression (Fig. 5). With the exception of the year 2002, the agreement between measurements and model results is excellent, and even small interannual variations on the order of a few centimeters are reproduced (e.g. 2012 vs. 2014). There is no significant difference in modeled thaw depths between model runs for different snow densities, 575 which is in agreement with a study of model sensitivity for Samoylov Island (Langer et al., 2013). We emphasize that the average of 150 point measurements of thaw depth is displayed in Fig. 5, while a considerable spatial variability occurs (average standard deviation - 1σ - of 0.06 m) due to variable ground and surface properties.

We conclude that the good agreement between the modeled and measured ground thermal regime 580 can be linked to the surface energy balance, surface temperatures and the snow dynamics all of which are adequately reproduced in the model.

Thermokarst lake stratigraphy: For the thermal regime and the melting of excess ground ice layers below water bodies, the temperature at the lake bottom is a crucial variable. For validation of the simple water body scheme (Sect. 2.7), we compare the three-year in-situ record of bottom temperatures from the 6 m deep “Sa_Lake_1” on Samoylov Island presented in Boike et al. (2015) to lake 585 bottom temperatures modeled with CryoGrid Xice (see Sect. 4.1 for details). Fig. 6 is clear evidence that the model can reproduce the main features of the annual temperature regime, in particular the strong asymmetry between winter and summer temperatures. The model can furthermore reproduce 590 the characteristic spike-like dip in temperatures visible in the measurements each fall: at this time, the surface is not frozen yet and persistent well-mixed conditions lead to a cooling of the entire water column towards 0°C. When the surface freezes, the heat transfer in the water column switches to comparatively inefficient conductive heat transfer, so that the heat loss towards the surface is strongly reduced. The lake bottom is then warmed by the heat stored in the sediments below, before it slowly 595 cools again in the course of winter. During this time, CryoGrid 3 Xice slightly underestimates the measured temperatures, most likely because buoyancy-driven thermal stratifications of the water column are not accounted for by the simplified model physics. During ice break-up in spring, CryoGrid 3 Xice shows a bottom cooling to 0°C, which is not observed in the measurements, but the timing of the subsequent strong warming at the beginning of summer is once again captured by the model. 600 While the model slightly underestimates the true annual average temperatures, the comparison to measured lake bottom temperatures suggests that the simple water body scheme of CryoGrid 3 Xice can in general reproduce the thermal regime at the bottom of Sa_Lake_1. It is remarkable that rather modest modifications of the model physics (namely the mixing of temperatures in the water column during summer) and changes to model parameters (e.g. the albedo) facilitate reproducing the ther-

605 mal regime both in cold permafrost ground (Fig. 4, mean annual temperature about -9°C) and at the
bottom of a thermokarst lake (Fig. 6, mean annual temperature about $+3.5^{\circ}\text{C}$).

5.2 Long-term thaw susceptibility runs with CryoGrid 3 Xice

The CRU-NCEP/CCSM4 data set facilitates simulations of the ground thermal state over a period
of 200 years from 1901 to 2100 for different future warming scenarios. Since these forcing data are
610 derived from a coupled climate model, they feature considerable biases for some forcing variables,
although partly corrected as described in Sect. 4.2. Therefore, the ground thermal regime modeled
for Samoylov Island deviates to a certain extent from the observations. Using the stratigraphy for
Samoylov Island (Table 2), the modeled maximum thaw depths for the period 2002-2014 is on av-
erage 0.66 m, approximately 0.1 m more than for the more realistic forcing of the validation run
615 (Fig. 5).

The stratigraphy of the ground has a pronounced influence in particular on the thaw depth: for the
Ice Complex stratigraphy, a significantly lower average maximum thaw depth of 0.43 m was ob-
tained, which is a result of the higher ice/water contents underneath the insulating unsaturated top
layer (Table 2). For the mineral soils of the “Arga” stratigraphy, a much deeper active layer with
620 maximum thaw depth of on average 1.29 m was modeled. While systematic measurements of thaw
depths are lacking for these two geomorphological units, the modeled values are in general agree-
ment with the few existing observations: on the Ice Complex on Kurunghak Island approx. 10 km
SW of Samoylov Island, thaw depths have been determined to 0.2-0.3 m in August 2013 which is
significantly shallower than at the same time in the first terrace on Samoylov Island (Fig. 5). For the
625 second terrace, a point observation on Arga Island ($73^{\circ}29'39.2''\text{N}$, $124^{\circ}22'33.1''\text{E}$) about 150 km
NW of Samoylov Island from 11 Aug, 2010, yielded a thaw depth of just below 1 m, almost double
compared to Samoylov Island on the first terrace. We emphasize that all CryoGrid 3 model runs are
based on the forcing data set downscaled for Samoylov Island, so that the results cannot be strictly
compared to observations on the second and third terrace. However, the differences in both modeled
630 and measured thaw depths between the three geomorphological units are systematic and significantly
larger than the interannual variability on Samoylov Island, which suggests that CryoGrid 3 can re-
produce existing patterns of thaw depths for the three geomorphological units in the Lena Delta.

Figs. 7 and 8 display the long-term evolution of the annual freeze-thaw state for the three different
stratigraphies and two future GHG emission scenarios. An important feature common to all is that
635 stable permafrost is modeled for a period of more than a century, with the active layer undergoing an-
nual freeze-thaw cycles on top of frozen ground. For the “Arga” stratigraphy lacking excess ground
ice, neither ground subsidence nor thermokarst can occur and the simulations yield identical results
for the two cases. For RCP 4.5, permafrost is thermally stable until 2100, with a gradual increase of
maximum thaw depth and a ground warming in 10 m depth of around 5°C (Fig. 9). For RCP 8.5,
640 the warming of the ground accelerates after 2050, which results in the formation of a talik towards

the end of the model period, with temperatures at 10 m depth approaching 0°C. This remarkable ground warming can be linked to a strong increase of incoming long-wave radiation and air temperatures in the CCSM4 data, in particular in fall and early winter. Air temperatures increase by about 1°C/decade for RCP 8.5, which can at least partly explain that the permafrost which features modeled 10 m-temperatures between -8 and -5°C in 2010, reaches the thaw threshold in 2100. The ground warming is amplified by a particularly pronounced temperature increase in fall and early winter, before the annual snow cover has reached its full thickness and thus insulating effect. In the month of November, the warming is almost twice compared to the annual average, with average temperatures increasing by approximately 15°C from 2010 and 2100, which leads to delayed freezing of the active layer and eventually the formation of a talik.

If layers with excess ground ice are present, the modifications of the model physics implemented in CryoGrid 3 Xice become relevant, as soon as the maximum thaw extends into these layers. For the “Samoylov” stratigraphy, this is the case around 2060, while thawing of ground layers supersaturated with ice occurs considerably earlier for “Ice Complex”. For the subsidence scenario and RCP 4.5 (Fig. 7), the ground subsides on the order of 1 m in less than 10 years, but subsequently stabilizes since the substantial fractions of mineral and organic material in the ground layers supersaturated with ground ice (Table 2) consolidate and thus form a thicker buffer over the ice-rich layers, so that the annual ground thawing does not reach into the zone with excess ground ice any more. For the “Ice Complex” stratigraphy, on the other hand, ground subsidence continues at a reduced rate after the strong initial thaw, since the formation of the buffer layer is slower than the increase in annual thaw depths due to the low contents of mineral and organic material in the excess ice layers. For the future warming scenario RCP 8.5 (Fig. 8), the ground warming is strong enough that a talik can form within the buffer layer at the end of the simulation period for the “Samoylov” stratigraphy. For “Ice Complex”, however, the simulations do not display talik formation until 2100, in despite of a significant increase in thaw depths. The energy input to the ground which for the “Arga” and “Samoylov” stratigraphies leads to talik formation, is to a large extent absorbed by thawing of ice-rich ground in relative proximity to the surface. This process, on the other hand, leads to a drastic subsidence of more than 5 m, accompanied by a warming of mean annual ground temperatures in the permanently frozen ground layers to close to 0°C. For the subsidence scenario, ground temperatures at 10 m depth below the subsided surface remain below 0°C in all four cases. We conclude that hydrological conditions leading to drainage of meltwater from excess ground ice delay or even prevent the onset of talik formation compared to a ground stratigraphy without excess ground ice.

For the “thermokarst” scenario, CryoGrid 3 Xice delivers entirely different trajectories of permafrost thaw. As a consequence of thawing of ground layers with excess ice, a thermokarst pond forms which rapidly reaches a depth of 1 m and more. As observed in nature (Boike et al., 2012), the pond initially freezes to its bottom each winter for a certain period in the RCP 4.5 scenario (Fig. 7). Upon further warming, the pond does not freeze to its bottom anymore, and a talik forms at the bottom of the

pond and the underlying sediment layer. The maximum modeled thickness of the winter ice layer on the pond is on the order of 1-2 m, which is in qualitative agreement with maximum ice thicknesses
680 of around 2 m observed for current climate conditions on thermokarst lakes in Samoylov Island (Boike et al., 2012, 2015). For the “Samoylov” stratigraphy, a significantly shallower pond forms compared to the “Ice Complex” stratigraphy, but the thickness of the unfrozen sediments underneath the pond is larger due to higher mineral and organic contents in the excess ground ice layers. For RCP 8.5 and “Samoylov”, the entire excess ice layer extending to a depth of 9 m is thawed by around
685 2090, so that the lake depth does not increase further. For “Ice Complex” the excess ice layers extend deeper and the thermokarst lake continues to deepen until the end of the simulation period, with a final depth of around 10 m. Following the initial development of a pond, ground temperatures at 10 m depth rapidly increase (Fig. 9) and eventually become positive, except for the “Samoylov” stratigraphy and RCP4.5.

690 The model results suggest that hydrological conditions preventing the drainage of meltwater from excess ground ice lead to the formation of thermokarst ponds and finally talik development already for a the moderate warming scenarios RCP 4.5. Compared to the “Arga” stratigraphy without layers with excess ground ice, permafrost thaw is considerably accelerated and taliks occur earlier, while expanding more rapidly in deeper ground layers.

695 **6 Discussion and Outlook**

6.1 CryoGrid 3 - an experimental platform for permafrost process parameterizations

The structure of CryoGrid 3 is designed to be similar to ESM land-surface schemes, so that algorithms and process parameterizations can easily be transferred. While an adaptive time step is employed in the time integration procedure (Sect. 2.6), constant time steps as typical in large-scale
700 modeling approaches are feasible, but the numerical stability cannot be guaranteed with the simple Forward Euler scheme in this case. Nevertheless, this choice guarantees maximum simplicity and readability of the CryoGrid 3 code, so that the model can be easily modified to test the effect of adding different process parameterizations to the basis module. As such, the goal of CryoGrid 3 is different from existing land-surface schemes, such as GEOtop (Endrizzi et al., 2013), COUP
705 (Jansson and Karlberg, 2004) or SURFEX (Masson et al., 2013), which are aimed at users running the models with a predefined set of model options, largely without modifying the program code itself. In contrast, CryoGrid 3 is optimized to encourage “users” with the necessary programming skills to modify and extend the program code, thus providing an open platform to explore the effect of different process parameterizations on model results.

710 In this study, CryoGrid 3 has been applied to cold, continuous permafrost in Siberia. While the satisfactory agreement of model results and in-situ observations is encouraging, it is likely that the model performance would be less satisfactory for some environmental conditions encountered in

permafrost areas. We see three major shortcomings of the current version of CryoGrid 3 which should be addressed prior to application in a wider range of permafrost environments:

- 715 1. *Snow physics*: CryoGrid 3 employs a constant density for freshly fallen snow and does not account for snow microstructural processes leading to changes of density and thermal properties. While this simple scheme seems adequate for the conditions on Samoylov Island with relatively low snow depths and spread of snow densities (Boike et al., 2013), a more sophisticated treatment of snow processes is warranted if CryoGrid 3 shall be applied e.g. to mountain environments characterized by a much higher snow cover. Hereby, in particular advanced snow pack models developed for avalanche forecasting, such as CROCUS (Vionnet et al., 2012) or SNOWPACK (Bartelt and Lehning, 2002), have the potential to significantly improve the performance of CryoGrid 3, but also more flexible, modular approaches like the Factorial Snowpack Model (Essery, 2015) could be incorporated.
- 720
- 725 2. *Vegetation cover*: While CryoGrid 3 can successfully reproduce energy transfer and ground thermal regime in the low-vegetated study area in NE Siberia, it is unlikely that it can account for the near-surface energy transfer in areas with higher vegetation, in particular forests. Many existing schemes simply add a vegetation/canopy layer with defined exchange coefficients for the fluxes of mass and energy on top of the existing soil grid, which is feasible in various levels of complexity (e.g. Zhang et al., 2003; ECMWF, 2006).
- 730
3. *Water balance*: In the study area, seasonal changes of the water level are generally low, in agreement with estimates of the summer water balance showing that evapotranspiration on average more or less balances precipitation (Boike et al., 2013). In many permafrost systems, however, seasonal changes in soil moisture play a pronounced role for energy transfer in the ground (e.g. Hollesen et al., 2011). While other land-surface schemes dedicated to modeling permafrost temperatures, such as GEOtop (Endrizzi et al., 2013), explicitly account for the dynamics of soil water transport by solving Richard's Equation, a more simple bucket approach (e.g. Budyko, 1961, 1974; Zaitarian, 1992) may be highly feasible in permafrost regions.
- 735

6.2 Comparison to field and model studies on ground subsidence

740 The CryoGrid 3 Xice scheme constitutes a simple way of incorporating ground subsidence and thermokarst in ESM land-surface schemes. With respect to ground subsidence, it is to a large degree similar to the ground subsidence scheme CLM EXICE (Lee et al., 2014) which was employed to model ground subsidence in Arctic permafrost areas until 2100. While Lee et al. (2014) find a maximum ground subsidence on the order of 0.5 m in NE Siberia, significantly larger values are obtained

745 in this study, which can be explained by the fact that much lower total excess ice contents were assigned in the large-scale study (maximum 0.5 m water equivalent opposed to 2 m for Samoylov, and almost 10 m for Ice Complex).

In the long-term thaw susceptibility runs for Samoylov Island, ground subsidence does not occur at or before present-day which is in agreement with in-situ observations (Boike et al., 2013). A validation of the modeled future subsidence rates is therefore not possible in a strict sense. However, the study of Günther et al. (2015) conducted for Muostakh Island off the coast of the Lena River Delta (150 km SE of Samoylov Island, dominated by Pleistocene sediments similar to the “Ice Complex” stratigraphy), where strong ground subsidence has occurred in the last 50 years, facilitates comparing measured and modeled properties of ground subsidence in a qualitatively way. The site has experienced significant warming most likely related to changes in the sea ice conditions in the past 50 years, and Günther et al. (2015) document ground subsidence rates of up to 5 m/50 y which is comparable in magnitude to the simulations for the Ice Complex stratigraphy for the next 50 years. Furthermore, the study can clearly document an inverse relationship between active layer thickness and subsidence rates (Fig. 10, Günther et al., 2015) which they relate to the differences in the ground ice content below the active layer. They document a ground subsidence of on average 2 m/50 y at a thaw depth of 0.4 m, about half this value for a thaw depth of 0.65 m, and close to no subsidence for active layer depths exceeding 0.8 m. In our simulations for the three stratigraphies, we can qualitatively reproduce this relationship. We model a current active layer thickness of about 0.4 m with future ground subsidence rates 6-8 m/50 y (Ice Complex stratigraphy) and 0.65 m with 4 m/50 y subsidence for the Samoylov stratigraphy. For the Arga stratigraphy lacking excess ground ice, the modeled active layer thickness exceeds 1 m, while no subsidence occurs. The exact subsidence rates necessarily depend on the applied forcing (which is certainly different for the past 50 years on Muostakh Island than for the next 50 years on Samoylov Island), but the qualitative agreement with the long-term observations suggests that CryoGrid 3 Xice can capture crucial dependencies on subsurface parameters that have been observed in nature.

6.3 Towards modeling permafrost landscape dynamics in regions with ice-rich ground

CryoGrid 3 Xice is based on a 1D-approximation for processes which in nature are strongly influenced by lateral fluxes of energy and water. A thermokarst pond can only extend with depth in the model, while it will also expand laterally due to erosion at the shores. Conversely, cooling from surrounding thermally stable permafrost could delay or stop further thawing of the ground underneath an established water body. Such processes have been described in dedicated thermokarst lake models (Plug and West, 2009; Kessler et al., 2012), but the coupled 3D-framework of heat conduction and water and mass movement requiring high spatial resolutions of around 10 m precludes application in ESM land-surface schemes. The algorithm describing excess ice melt in CryoGrid 3 Xice is on the other hand simple and highly practical for implementation in large-scale models: if a layer with excess ground ice thaws for the first time, the properties of the above-lying grid cells are rearranged while at the same time conserving mass. As demonstrated by Lee et al. (2014) for CLM, such a step could easily implemented in an ESM land-surface scheme. When a thermokarst pond or lake has

formed, heat conduction (as already implemented in the land-surface scheme) continues to be the
785 only means of energy transfer, while mixing of the water column during summer is simulated by
assigning an average temperature for the water body after each time step. In this way, well-mixed
conditions as typical during summer (Boike et al., 2015) are simulated, while the relatively low ther-
mal conductivity of liquid water limits the heat transfer for the stratified water column under an ice
cover (Boike et al., 2015). During winter, the heat transfer in the ground below the lake is most likely
790 biased since the stratification with the densest water at the bottom is not reproduced in CryoGrid 3
Xice, which is agreement with the cold-bias of modeled lake bottom temperatures (Fig. 6). How-
ever, it is not clear how this will affect the seasonal dynamics and the long-term evolution of the
water body, which should be clarified in a comparison with dedicated lake schemes, such as FLake
(Mironov et al., 2010).

795 Furthermore, it must be emphasized that the satisfactory performance of the CryoGrid 3 Xice wa-
ter body scheme was only confirmed for a well-developed thermokarst lake which represents the
endpoint of the thermokarst simulations for the Ice Complex and Samoylov stratigraphies (Figs. 7,
8). It is not ultimately clear how the scheme would perform for the wide range of smaller and
medium-sized ponds on Samoylov Island (Muster et al., 2012; Langer et al., 2015) which necessar-
800 ily represent initial stages for the development of larger thermokarst lakes. Therefore, we consider
the modeled dynamics of the transition from terrestrial permafrost to a deeper lake to be highly un-
certain and recommend more research on the factors governing the thermal regime in particular of
smaller water bodies.

Another critical missing step to describing landscape evolution is to understand the 3D-variability
805 of near-surface ground ice content and active layer thickness, as well as the resulting lateral fluxes
of heat and water. As demonstrated in Sect. 5.2, thermokarst/subsidence processes will preferably
occur in areas, where layers with high excess content exist close to the base of the active layer. In
such areas, already a small increase of active layer thickness could trigger melting of excess ice and
thus initiate transformative landscape changes.

810 **6.4 Threshold behavior and metastable states in permafrost systems?**

The simulations suggest that ground thawing could display a pronounced threshold behavior in ice-
rich permafrost landscapes. Once the active layer extends into layers with excess ground ice, irre-
versible ice melting with pronounced impact on the ground surface is initiated: when the meltwater
is removed, the ground subsides, bringing initially deeper layers with excess ground ice in proximity
815 to the thaw front. The CryoGrid 3 Xice simulations suggest that this process can remove several me-
ters of ground ice within less than a decade (“Ice Complex” stratigraphy, Figs. 7, 8). Such depressed
areas would become preferential flow paths for surface waters and eventually develop into incised
drainage networks (e.g. Fortier et al., 2007). On the other hand, accumulation of sediments from the
melted out ice layers will eventually create a protective layer without excess ground ice on top of the

820 remaining layers, so that the permafrost system can return to a thermally stable state with increased active layer thickness, as for the RCP 4.5 simulation for the “Samoylov” stratigraphy (Fig. 7). A similar “metastable state” could be reached by a thermokarst pond in which the entire water and ground column refreeze during winter, as typical for small ponds on Samoylov Island (Langer et al., 2015; Boike et al., 2015). In the CryoGrid 3 Xice simulations, the ponds generally form taliks within
825 10-15 years of their creation (Figs. 7, 8), but that occurs in conjunction with ongoing warming. It should therefore be investigated whether CryoGrid 3 Xice could also yield such metastable states for present-day climate forcing on Samoylov Island.

The present-day landscape in NE Siberia is evidence that melting of ground ice has shaped many permafrost landscapes since the last glacial period (e.g. Schirrmeister et al., 2008). First results are
830 encouraging that CryoGrid 3 Xice may be developed further into a modeling tool with which such observed landscape changes can be linked to the development of the climate forcing. The presented simulations with CryoGrid 3 Xice indicate that the trajectories of permafrost thaw are crucially influenced by the vertical and lateral distribution of ground ice and processes following excess ground ice thaw. The stark contrast between the different thaw scenarios for the “subsidence” and “thermokarst”
835 cases suggests that simulations of permafrost thaw based exclusively on conductive heat transfer, as employed in the vast majority of studies, are highly questionable in regions with ice-rich ground.

7 Conclusions

In this study, we present a novel land-surface scheme designed to reproduce the ground thermal regime in permafrost regions. The numerical structure of CryoGrid 3 is simple, using a first-order
840 Forward Euler time integration scheme. This relative simplicity makes the new model scheme a flexible platform to explore new parameterizations for a range of permafrost processes. At the upper boundary, CryoGrid 3 is forced by the surface energy balance, while conductive heat transfer is the principal means of energy transfer in the ground. Model results forced with measured time series of meteorological variables from the Samoylov permafrost observatory agree well with in-situ records
845 of ground temperatures and active layer thickness, while at the same time consistently reproducing measured surface temperatures and components of the surface energy balance.

In a second step, we introduce a simple parameterization for thawing of excess ground ice, which redistributes the generated meltwater within the 1D soil column. In a simplified way, the scheme is capable of simulating both ground subsidence and formation of thermokarst ponds. In a long-term
850 simulation from 1901-2100, we retrace the rapid transition of a permafrost ground column that is thermally stable throughout the 20th century to a thermokarst pond which eventually develops an underlying talik in the course of the 21st century warming. These initial model results suggest that the cryostratigraphy of the ground and the hydrological regime of a site have a pronounced impact on the permafrost thaw trajectory. In general, ground subsidence due to drainage of meltwater appears

855 to delay the onset of talik formation, while the formation of a thermokarst pond has the opposite
effect. In the simulations, the differences between the subsidence and the thermokarst scenario are
on the order of the differences between the two investigated future anthropogenic emission scenarios,
RCP 4.5 and 8.5. The results support earlier findings that melting of excess ground ice and resulting
hydrological processes deserve critical attention in modeling efforts on the future evolution of the
860 world-wide permafrost state.

Code availability

The CryoGrid 3 source code is available on request from the two first authors of the study: Sebastian
Westermann, sebastian.westermann@geo.uio.no; Moritz Langer, mlanger@awi.de.

Appendix A: Definitions and constants

865 α : surface albedo [-]
 ε : Kirchhoff emissivity [-]
 σ : Stefan-Boltzmann constant [$\text{W m}^{-2}\text{K}^{-4}$]
 ρ : density [kg m^{-3}]
 c_p : specific heat capacity of air at constant pressure [$\text{J kg}^{-1}\text{K}^{-1}$]
870 ρ_a : density of air [kg m^{-3}]
 $\kappa = 0.4$: von Kármán-constant (e.g. Foken, 2008)
 u_* : friction velocity [m s^{-1}]
 z_0 : aerodynamic roughness length [m]
 L_* : Obukhov length [m]
875 $\Psi_{M,H,W}(\zeta_1, \zeta_2)$: integrated stability functions for momentum, heat and water vapor [-]
 g : gravitational constant [m s^{-2}]
 r_s : surface resistance against evapotranspiration [s m^{-1}]
 $L_{sl} = 0.33 \text{ MJ kg}^{-1}$: specific latent heat of fusion of water
 $L_{lg} = 2.5 \text{ MJ kg}^{-1}$: specific latent heat of vaporization of water
880

Appendix B: Calculation of absolute humidity

The absolute humidity q can be derived from the relative humidity RH and the saturation vapor pressure e^* accessible through the empirical August-Roche-Magnus formula:

$$q(z_0) = 0.622 \frac{e^*(T^1)}{p} \quad (\text{B1})$$

$$885 \quad q(h) = 0.622 \frac{RH(h)e^*(T_{\text{air}}(h))}{p} \quad (\text{B2})$$

$$e^*(T) = a_1 \exp\left(\frac{a_2 T}{T + a_3}\right). \quad (\text{B3})$$

We assign $a_1 = 611$ Pa, $a_2 = 17.62$ and $a_3 = 243.12$ K for the saturation vapor pressure over water surfaces and $a_1 = 611$ Pa, $a_2 = 22.46$ and $a_3 = 272.62$ K over ice surfaces ($T_1 < 0^\circ\text{C}$) (Sonntag, 1990).

890 Appendix C: Atmospheric stability functions

The integrated stability functions $\Psi_{M,H,W}$ are determined by integrating the universal functions $\varphi_{M,H,W}(\zeta)$ as

$$\Psi_{M,H,W}(\zeta_1, \zeta_2) = \int_{\zeta_2}^{\zeta_1} d\zeta \varphi_{M,H,W}(\zeta). \quad (\text{C1})$$

For neutral and unstable atmospheric stratifications ($h/L_* \leq 0$), we assume the commonly employed
895 functions by Högström (1988)

$$\varphi_M(\zeta) = (1 - 19\zeta)^{-1/4} \quad (\text{C2})$$

$$\varphi_{H,W}(\zeta) = 0.95(1 - 11.6\zeta)^{-1/2}, \quad (\text{C3})$$

while functions compiled from data of the SHEBA campaign (Uttal et al., 2002) over Arctic sea ice are employed for stable stratification conditions (Grachev et al., 2007), $h/L_* > 0$

$$900 \quad \varphi_M(\zeta) = 1 + \frac{6.5\zeta(1 + \zeta)^{1/3}}{1.3 + \zeta} \quad (\text{C4})$$

$$\varphi_{H,W}(\zeta) = 1 + \frac{5\zeta(1 + \zeta)}{1 + 3\zeta + \zeta^2}. \quad (\text{C5})$$

Acknowledgements. This work was funded by CryoMet (project no. 214465; Research Council of Norway), Page21 (contract no. GA282700; European Union FP7-ENV), COUP (project no. 244903/E10; JPI Climate; Research Council of Norway), SatPerm (project no. 239918; Research Council of Norway), a Feodor-Lynen
905 Grant awarded to M. Langer by the Alexander-von-Humboldt Foundation, Germany, and the Department of Geosciences, University of Oslo, Norway. We gratefully acknowledge the support of the Russian-German research station on Samoylov Island. We thank Günther Stoof, Sofia Antonova and Sascha Niemann for their

contributions to fieldwork, and Isabelle Gouttevin for valuable discussions in the planning stages of CryoGrid
3.

910 References

- Abnizova, A., Siemens, J., Langer, M., and Boike, J.: Small ponds with major impact: The relevance of ponds and lakes in permafrost landscapes to carbon dioxide emissions, *Global Biogeochemical Cycles*, 26, GB2041, doi:10.1029/2011GB004237, 2012.
- Bartelt, P. and Lehning, M.: A physical SNOWPACK model for the Swiss avalanche warning. Part I: numerical
915 model, *Cold Regions Science and Technology*, 35, 123–145, 2002.
- Boike, J., Wille, C., and Abnizova, A.: Climatology and summer energy and water balance of polygonal tundra in the Lena River Delta, Siberia, *Journal of Geophysical Research: Biogeosciences* (2005–2012), 113, doi:10.1029/2007JG000540, 2008.
- Boike, J., Langer, M., Lantuit, H., Muster, S., Roth, K., Sachs, T., Overduin, P., Westermann, S., and McGuire,
920 A. D.: Permafrost–Physical Aspects, Carbon Cycling, Databases and Uncertainties, in: *Recarbonization of the Biosphere*, pp. 159–185, Springer, 2012.
- Boike, J., Kattenstroth, B., Abramova, K., Bornemann, N., Chetverova, A., Fedorova, I., Fröb, K., Grigoriev, M., Grüber, M., Kutzbach, L., Langer, M., Minke, M., Muster, S., Piel, K., Pfeiffer, E.-M., Stoof, G., Westermann, S., Wischnewski, K., Wille, C., and Hubberten, H.-W.: Baseline characteristics of climate, permafrost and
925 land cover from a new permafrost observatory in the Lena River Delta, Siberia (1998–2011), *Biogeosciences*, 10, 2105–2128, doi:10.5194/bg-10-2105-2013, <http://www.biogeosciences.net/10/2105/2013/>, 2013.
- Boike, J., Georgi, C., Kirilin, G., Muster, S., Abramova, K., Fedorova, I., Chetverova, A., Grigoriev, M., Bornemann, N., and Langer, M.: Thermal processes of thermokarst lakes in the continuous permafrost zone of northern Siberia – observations and modeling (Lena River Delta, Siberia), *Biogeosciences*, 12, 5941–5965,
930 doi:10.5194/bg-12-5941-2015, <http://www.biogeosciences.net/12/5941/2015/>, 2015.
- Brown, J., Ferrians Jr, O., Heginbottom, J., and Melnikov, E.: *Circum-Arctic map of permafrost and ground-ice conditions*, US Geological Survey Circum-Pacific Map, 1997.
- Budyko, M. I.: The heat balance of the earth’s surface, *Soviet Geography*, 2, 3–13, 1961.
- Budyko, M. I.: *Climate and life*, Academic Press, 1974.
- 935 Burt, W. V.: Albedo over wind-roughened water, *Journal of Meteorology*, 11, 283–290, 1954.
- Chadburn, S., Burke, E., Essery, R., Boike, J., Langer, M., Heikenfeld, M., Cox, P., and Friedlingstein, P.: An improved representation of physical permafrost dynamics in the JULES land-surface model, *Geoscientific Model Development*, 8, 1493–1508, doi:10.5194/gmd-8-1493-2015, <http://www.geosci-model-dev.net/8/1493/2015/>, 2015.
- 940 Cosenza, P., Guerin, R., and Tabbagh, A.: Relationship between thermal conductivity and water content of soils using numerical modelling, *European Journal of Soil Science*, 54, 581–588, 2003.
- Dall’Amico, M., Endrizzi, S., Gruber, S., and Rigon, R.: A robust and energy-conserving model of freezing variably-saturated soil, *The Cryosphere*, 5, 469–484, 2011.
- Dee, D. P., Uppala, S. M., Simmons, A. J., Berrisford, P., Poli, P., Kobayashi, S., Andrae, U., Balmaseda, M. A., Balsamo, G., Bauer, P., Bechtold, P., Beljaars, A. C. M., van de Berg, L., Bidlot, J., Bormann, N., Delsol, C., Dragani, R., Fuentes, M., Geer, A. J., Haimberger, L., Healy, S. B., Hersbach, H., Hólm, E. V., Isaksen, I., Kållberg, P., Köhler, M., Matricardi, M., McNally, A. P., Monge-Sanz, B. M., Morcrette, J.-J., Park, B.-K., Peubey, C., de Rosnay, P., Tavolato, C., Thépaut, J.-N., and Vitart, F.: The ERA-Interim

- reanalysis: configuration and performance of the data assimilation system, *Quarterly Journal of the Royal Meteorological Society*, 137, 553–597, doi:10.1002/qj.828, 2011.
- 950 ECMWF: IFS DOCUMENTATION - Cy31r1, PART IV: PHYSICAL PROCESSES, Reading, UK, <https://software.ecmwf.int/wiki/display/IFS/CY31R1+Official+IFS+Documentation>, 2006.
- Ekici, A., Beer, C., Hagemann, S., and Hauck, C.: Simulating high-latitude permafrost regions by the JSBACH terrestrial ecosystem model, *Geoscientific Model Development*, 7, 631–647, 2014.
- 955 Endrizzi, S., Gruber, S., Dall’Amico, M., and Rigon, R.: GEOTop 2.0: simulating the combined energy and water balance at and below the land surface accounting for soil freezing, snow cover and terrain effects, *Geosci. Model Dev. Discuss*, 6, 6279–6341, 2013.
- Essery, R.: A Factorial Snowpack Model (FSM 1.0), *Geoscientific Model Development Discussions*, 8, 6583–6609, doi:10.5194/gmdd-8-6583-2015, <http://www.geosci-model-dev-discuss.net/8/6583/2015/>, 2015.
- 960 Fedorova, I., Chetverova, A., Bolshiyarov, D., Makarov, A., Boike, J., Heim, B., Morgenstern, A., Overduin, P., Wegner, C., Kashina, V., et al.: Lena Delta hydrology and geochemistry: long-term hydrological data and recent field observations, *Biogeosciences*, 12, 345–363, 2015.
- Fiddes, J., Endrizzi, S., and Gruber, S.: Large-area land surface simulations in heterogeneous terrain driven by global data sets: application to mountain permafrost, *The Cryosphere*, 9, 411–426, doi:10.5194/tc-9-411-2015, <http://www.the-cryosphere.net/9/411/2015/>, 2015.
- 965 Foken, T.: *Micrometeorology*, Springer, Berlin Heidelberg, Germany, doi:10.1007/978-3-540-74666-9, 2008.
- Fortier, D., Allard, M., and Shur, Y.: Observation of rapid drainage system development by thermal erosion of ice wedges on Bylot Island, Canadian Arctic Archipelago, *Permafrost and Periglacial Processes*, 18, 229–243, 2007.
- 970 Gislén, K., Eitzelmüller, B., Farbrøt, H., Schuler, T., and Westermann, S.: CryoGRID 1.0: Permafrost distribution in Norway estimated by a spatial numerical model, *Permafrost and Periglacial Processes*, 24, 2–19, 2013.
- Gouttevin, I., Krinner, G., Ciais, P., Polcher, J., and Legout, C.: Multi-scale validation of a new soil freezing scheme for a land-surface model with physically-based hydrology, *The Cryosphere*, 6, 407–430, doi:10.5194/tc-6-407-2012, <http://www.the-cryosphere.net/6/407/2012/>, 2012.
- 975 Grachev, A., Andreas, E., Fairall, C., Guest, P., and Persson, P.: SHEBA flux–profile relationships in the stable atmospheric boundary layer, *Boundary-Layer Meteorology*, 124, 315–333, 2007.
- Grigoriev, M.: *Cryomorphogenesis of the Lena River mouth area*, Russian Academy of Science, Siberian Branch, Yakutsk, Russia (in Russian), 1993.
- 980 Grigoriev, M., Imaev, V., Imaeva, L., Kozmin, B., Kunitzkiy, V., Lationov, A., Mikulenko, K., Skryabin, R., and Timirsin, K.: *Geology, seismicity and cryogenic processes in the arctic areas of Western Yakutia*, Yakut Scientific Centre, Russian Academy of Science, Siberian Branch, Yakutsk, Russia (in Russian), 1996.
- Günther, F., Overduin, P. P., Yakshina, I. A., Opel, T., Baranskaya, A. V., and Grigoriev, M. N.: Observing Muostakh disappear: permafrost thaw subsidence and erosion of a ground-ice-rich island in response to arctic summer warming and sea ice reduction, *The Cryosphere*, 9, 151–178, 2015.
- 985 Harris, I., Jones, P., Osborn, T., and Lister, D.: Updated high-resolution grids of monthly climatic observations—the CRU TS3. 10 Dataset, *International Journal of Climatology*, 34, 623–642, 2014.

- Høgstrøm, U.: Non-dimensional wind and temperature profiles in the atmospheric surface layer: A re-evaluation, *Boundary-Layer Meteorology*, 42, 55–78, 1988.
- 990 Hollesen, J., Elberling, B., and Jansson, P.-E.: Future active layer dynamics and carbon dioxide production from thawing permafrost layers in Northeast Greenland, *Global Change Biology*, 17, 911–926, 2011.
- Hopkins, D. M.: Thaw lakes and thaw sinks in the Imuruk Lake area, Seward Peninsula, Alaska, *The Journal of Geology*, pp. 119–131, 1949.
- Jafarov, E. E., Marchenko, S. S., and Romanovsky, V. E.: Numerical modeling of permafrost dynamics in Alaska
995 using a high spatial resolution dataset, *The Cryosphere*, 6, 613–624, 2012.
- Jansson, P. and Karlberg, L.: COUP manual: Coupled heat and mass transfer model for soil-plant-atmosphere systems, Technical manual for the CoupModel, pp. 1–453, 2004.
- Järvinen, O. and Leppäranta, M.: Transmission of solar radiation through the snow cover on floating ice, *Journal of Glaciology*, 57, 861–870, 2011.
- 1000 Kalnay, E., Kanamitsu, M., Kistler, R., Collins, W., Deaven, D., Gandin, L., Iredell, M., Saha, S., White, G., Woollen, J., Zhu, Y., Leetmaa, A., Reynolds, R., Chelliah, M., Ebisuzaki, W., Higgins, W., Janowiak, J., Mo, K., Ropelewski, C., Wang, J., Jenne, R., and Joseph, D.: The NCEP/NCAR 40-year reanalysis project, *Bulletin of the American Meteorological Society*, 77, 437–472, 1996.
- Kessler, M., Plug, L. J., and Walter Anthony, K.: Simulating the decadal-to millennial-scale dynamics of morphology and sequestered carbon mobilization of two thermokarst lakes in NW Alaska, *Journal of Geophysical Research: Biogeosciences* (2005–2012), 117, G00M06, doi:10.1029/2011JG001796, 2012.
1005
- Koven, C. D., Riley, W. J., and Stern, A.: Analysis of permafrost thermal dynamics and response to climate change in the CMIP5 Earth System Models, *Journal of Climate*, 26, 1877–1900, 2013.
- Koven, C. D., Lawrence, D. M., and Riley, W. J.: Permafrost carbon- climate feedback is sensitive to deep
1010 soil carbon decomposability but not deep soil nitrogen dynamics, *Proceedings of the National Academy of Sciences*, 112, 3752–3757, 2015.
- Kutzbach, L., Wagner, D., and Pfeiffer, E.-M.: Effect of microrelief and vegetation on methane emission from wet polygonal tundra, Lena Delta, Northern Siberia, *Biogeochemistry*, 69, 341–362, 2004.
- Kutzbach, L., Wille, C., and Pfeiffer, E.-M.: The exchange of carbon dioxide between wet arctic tundra and the atmosphere at the Lena River Delta, Northern Siberia, *Biogeosciences*, 4, 869–890,
1015 doi:10.5194/bg-4-869-2007, <http://www.biogeosciences.net/4/869/2007/>, 2007.
- Langer, M., Westermann, S., and Boike, J.: Spatial and temporal variations of summer surface temperatures of wet polygonal tundra in Siberia-implications for MODIS LST based permafrost monitoring, *Remote Sensing of Environment*, 114, 2059–2069, 2010.
- 1020 Langer, M., Westermann, S., Muster, S., Piel, K., and Boike, J.: The surface energy balance of a polygonal tundra site in northern Siberia Part 1: Spring to fall, *The Cryosphere*, 5, 151–171, 2011a.
- Langer, M., Westermann, S., Muster, S., Piel, K., and Boike, J.: The surface energy balance of a polygonal tundra site in northern Siberia Part 2: Winter, *Cryosphere*, 5, 509–524, 2011b.
- Langer, M., Westermann, S., Heikenfeld, M., Dorn, W., and Boike, J.: Satellite-based modeling of permafrost
1025 temperatures in a tundra lowland landscape, *Remote Sensing of Environment*, 135, 12–24, 2013.

- Langer, M., Westermann, S., Walter Anthony, K., Wischniewski, K., and Boike, J.: Frozen ponds: production and storage of methane during the Arctic winter in a lowland tundra landscape in northern Siberia, Lena River delta, *Biogeosciences*, 12, 977–990, 2015.
- Lawrence, D., Slater, A., Romanovsky, V., and Nicolsky, D.: Sensitivity of a model projection of near-surface permafrost degradation to soil column depth and representation of soil organic matter, *Journal of Geophysical Research*, 113, F02011, doi:10.1029/2007JF000883, 2008.
- 1030
- Lee, H., Swenson, S. C., Slater, A. G., and Lawrence, D. M.: Effects of excess ground ice on projections of permafrost in a warming climate, *Environmental Research Letters*, 9, 124006, doi:10.1088/1748-9326/9/12/124006, 2014.
- 1035
- Lehning, M., Bartelt, P., Brown, B., and Fierz, C.: A physical SNOWPACK model for the Swiss avalanche warning. Part III: meteorological forcing, thin layer formation and evaluation, *Cold Regions Science and Technology*, 35, 169–184, 2002a.
- Lehning, M., Bartelt, P., Brown, B., Fierz, C., and Satyawali, P.: A physical SNOWPACK model for the Swiss avalanche warning:: Part II. Snow microstructure, *Cold Regions Science and Technology*, 35, 147–167, 2002b.
- 1040
- Lei, R., Leppäranta, M., Erm, A., Jaatinen, E., Pärn, O., et al.: Field investigations of apparent optical properties of ice cover in Finnish and Estonian lakes in winter 2009, *Estonian Journal of Earth Sciences*, 60, 50, 2011.
- Masson, V., Le Moigne, P., Martin, E., Faroux, S., Alias, A., Alkama, R., Belamari, S., Barbu, A., Boone, A., Bouyssel, F., Brousseau, P., Brun, E., Calvet, J.-C., Carrer, D., Decharme, B., Delire, C., Donier, S.,
- 1045
- Essaouini, K., Gibelin, A.-L., Giordani, H., Habets, F., Jidane, M., Kerdraon, G., Kourzeneva, E., Lafaysse, M., Lafont, S., Lebeaupin Brossier, C., Lemonsu, A., Mahfouf, J.-F., Marguinaud, P., Mokhtari, M., Morin, S., Pigeon, G., Salgado, R., Seity, Y., Taillefer, F., Tanguy, G., Tulet, P., Vincendon, B., Vionnet, V., and Voldoire, A.: The SURFEXv7.2 land and ocean surface platform for coupled or offline simulation of earth surface variables and fluxes, *Geoscientific Model Development*, 6, 929–960, doi:10.5194/gmd-6-929-2013, <http://www.geosci-model-dev.net/6/929/2013/>, 2013.
- 1050
- Meehl, G. A., Washington, W. M., Arblaster, J. M., Hu, A., Teng, H., Tebaldi, C., Sanderson, B. N., Lamarque, J.-F., Conley, A., Strand, W. G., , and White III, J.: Climate system response to external forcings and climate change projections in CCSM4, *Journal of Climate*, 25, 3661–3683, 2012.
- Mironov, D., HEIST, E., Kourzeneva, E., Ritter, B., Schneider, N., and Terzhevik, A.: Implementation of the lake parameterisation scheme FLake into the numerical weather prediction model COSMO, *Boreal environment research*, 15, 218–230, 2010.
- 1055
- Monin, A. and Obukhov, A.: Basic laws of turbulent mixing in the ground layer of the atmosphere, *Tr. Geofiz. Inst. Akad. Nauk SSSR*, 151, 163–187, 1954.
- Muster, S., Langer, M., Heim, B., Westermann, S., and Boike, J.: Subpixel heterogeneity of ice-wedge polygonal tundra: a multi-scale analysis of land cover and evapotranspiration in the Lena River Delta, Siberia, *Tellus B*, 64, 17301, doi:10.3402/tellusb.v64i0.17301, 2012.
- 1060
- Nicou, R., Charreau, C., and Chopart, J.-L.: Tillage and soil physical properties in semi-arid West Africa, *Soil and tillage research*, 27, 125–147, 1993.

- O'Neill, A. and Gray, D. M.: Solar radiation penetration through snow, in: *The Role of Snow and Ice in Hydrology*, Proceedings of the Banff Symposium, International Association of Hydrological Sciences, 107, pp. 227–240, 1972.
- 1065 Plug, L. J. and West, J.: Thaw lake expansion in a two-dimensional coupled model of heat transfer, thaw subsidence, and mass movement, *Journal of Geophysical Research: Earth Surface* (2003–2012), 114, F01 002, doi:10.1029/2006JF000740, 2009.
- 1070 Rachold, V. and Grigoriev, M. N.: Russian-German Cooperation SYSTEM LAPTEV SEA 2000: The Lena Delta 1998 Expedition, *Berichte zur Polarforschung* (Reports on Polar Research), 315, 1–259, 1999.
- Sachs, T., Wille, C., Boike, J., and Kutzbach, L.: Environmental controls on ecosystem-scale CH₄ emission from polygonal tundra in the Lena River Delta, Siberia, *Journal of Geophysical Research: Biogeosciences* (2005–2012), 113, G00A03, doi:10.1029/2007JG000505, 2008.
- 1075 Sachs, T., Giebels, M., Boike, J., and Kutzbach, L.: Environmental controls on CH₄ emission from polygonal tundra on the microsite scale in the Lena river delta, Siberia, *Global Change Biology*, 16, 3096–3110, 2010.
- Schaefer, K., Zhang, T., Bruhwiler, L., and Barrett, A.: Amount and timing of permafrost carbon release in response to climate warming, *Tellus B*, 63, 165–180, doi:10.1111/j.1600-0889.2011.00527.x, 2011.
- Schuesser, W.: *The numerical method of lines: integration of partial differential equations*, vol. 212, Academic Press, San Diego, USA, 1991.
- 1080 Schirrmeister, L., Grosse, G., Schwamborn, G., Andreev, A., Meyer, H., Kunitsky, V. V., Kuznetsova, T. V., Dorozhkina, M., Pavlova, E., Bobrov, A., et al.: Late Quaternary history of the accumulation plain north of the Chekanovsky Ridge (Lena Delta, Russia): a multidisciplinary approach, *Polar Geography*, 27, 277–319, 2003.
- 1085 Schirrmeister, L., Grosse, G., Kunitsky, V., Magens, D., Meyer, H., Dereviagin, A., Kuznetsova, T., Andreev, A., Babiy, O., Kienast, F., et al.: Periglacial landscape evolution and environmental changes of Arctic lowland areas for the last 60 000 years (western Laptev Sea coast, Cape Mamontov Klyk), *Polar Research*, 27, 249–272, 2008.
- Schirrmeister, L., Grosse, G., Schnelle, M., Fuchs, M., Krbetschek, M., Ulrich, M., Kunitsky, V., Grigoriev, M., Andreev, A., Kienast, F., et al.: Late Quaternary paleoenvironmental records from the western Lena Delta, arctic Siberia, *Palaeogeography, Palaeoclimatology, Palaeoecology*, 299, 175–196, 2011.
- 1090 Schneider, J., Grosse, G., and Wagner, D.: Land cover classification of tundra environments in the Arctic Lena Delta based on Landsat 7 ETM+ data and its application for upscaling of methane emissions, *Remote Sensing of Environment*, 113, 380–391, 2009.
- 1095 Schwamborn, G., Rachold, V., and Grigoriev, M. N.: Late Quaternary sedimentation history of the Lena Delta, *Quaternary International*, 89, 119–134, 2002.
- Simmons, A., Uppala, S., Dee, D., and Kobayashi, S.: ERA-Interim: New ECMWF reanalysis products from 1989 onwards, *ECMWF Newsletter*, 110, 25–35, 2007.
- Sonntag, D.: Important new values of the physical constants of 1986, vapour pressure formulations based on the ITS-90, and psychrometer formulae, *Zeitschrift für Meteorologie*, 40, 340–344, 1990.
- 1100 Ulrich, M., Grosse, G., Chabrillat, S., and Schirrmeister, L.: Spectral characterization of periglacial surfaces and geomorphological units in the Arctic Lena Delta using field spectrometry and remote sensing, *Remote Sensing of Environment*, 113, 1220–1235, 2009.

- Uttal, T., Curry, J., McPhee, M., Perovich, D., Moritz, R., Maslanik, J., Guest, P., Stern, H., Moore, J., Turenne,
1105 R., et al.: Surface heat budget of the Arctic Ocean, *Bulletin of the American Meteorological Society*, 83,
255–275, 2002.
- Vionnet, V., Brun, E., Morin, S., Boone, A., Faroux, S., Le Moigne, P., Martin, E., and Willemet, J.-M.: The
detailed snowpack scheme Crocus and its implementation in SURFEX v7.2, *Geoscientific Model Develop-*
ment, 5, 773–791, doi:10.5194/gmd-5-773-2012, <http://www.geosci-model-dev.net/5/773/2012/>, 2012.
- 1110 Wagner, D., Gattinger, A., Embacher, A., Pfeiffer, E., Schloter, M., and Lipski, A.: Methanogenic activity and
biomass in Holocene permafrost deposits of the Lena Delta, Siberian Arctic and its implication for the global
methane budget, *Global Change Biology*, 13, 1089–1099, 2007.
- Wagner, D., Kobabe, S., and Liebner, S.: Bacterial community structure and carbon turnover in permafrost-
affected soils of the Lena Delta, northeastern Siberia, *Canadian journal of microbiology*, 55, 73–83, 2009.
- 1115 Walter, K., Zimov, S., Chanton, J., Verbyla, D., and Chapin III, F.: Methane bubbling from Siberian thaw lakes
as a positive feedback to climate warming, *Nature*, 443, 71–75, 2006.
- Westermann, S., Boike, J., Langer, M., Schuler, T., and Etzelmüller, B.: Modeling the impact of wintertime rain
events on the thermal regime of permafrost, *The Cryosphere*, 5, 1697–1736, 2011.
- Westermann, S., Schuler, T., Gislås, K., and Etzelmüller, B.: Transient thermal modeling of permafrost condi-
1120 tions in Southern Norway, *The Cryosphere*, 7, 719–739, 2013.
- Westermann, S., Elberling, B., Højlund Pedersen, S., Stendel, M., Hansen, B., and Liston, G.: Future permafrost
conditions along environmental gradients in Zackenberg, Greenland, *The Cryosphere*, 9, 719–735, 2015.
- Wille, C., Kutzbach, L., Sachs, T., Wagner, D., and Pfeiffer, E.: Methane emission from Siberian arctic polygo-
nial tundra: eddy covariance measurements and modeling, *Global Change Biology*, 14, 1395–1408, 2008.
- 1125 Yen, Y.-C.: Review of thermal properties of snow, ice and sea ice, Tech. rep., DTIC Document, 1981.
- Yershov, E., Kondrat'yeva, K., Loginov, V., and Sychev, I.: Geocryological map of Russia and neighbouring
republics, Faculty of Geology, Chair of Geocryology, Lomonosov Moscow State University, 1991.
- Yi, S., Wischnewski, K., Langer, M., Muster, S., and Boike, J.: Freeze/thaw processes in complex permafrost
landscapes of northern Siberia simulated using the TEM ecosystem model: impact of thermokarst ponds and
1130 lakes, *Geoscientific Model Development*, 7, 1671–1689, 2014.
- Zaitarian, V.: A land-surface hydrology parameterization with sub-grid variability for general circulation mod-
els, *Journal of geophysical research*, 97, 2717–2728, 1992.
- Zhang, Y.: Spatio-temporal features of permafrost thaw projected from long-term high-resolution modeling for
a region in the Hudson Bay Lowlands in Canada, *Journal of Geophysical Research: Earth Surface*, 118,
1135 542–552, 2013.
- Zhang, Y., Chen, W., and Cihlar, J.: A process-based model for quantifying the impact of climate change
on permafrost thermal regimes, *Journal of Geophysical Research: Atmospheres* (1984–2012), 108, 4695,
doi:10.1029/2002JD00335, 2003.
- Zhang, Y., Chen, W., and Riseborough, D. W.: Transient projections of permafrost distribution in Canada during
1140 the 21st century under scenarios of climate change, *Global and Planetary Change*, 60, 443–456, 2008.
- Zhang, Y., Sachs, T., Li, C., and Boike, J.: Upscaling methane fluxes from closed chambers to eddy covariance
based on a permafrost biogeochemistry integrated model, *Global Change Biology*, 18, 1428–1440, 2012.

- Zhang, Y., Wang, X., Fraser, R., Olthof, I., Chen, W., McLennan, D., Ponomarenko, S., and Wu, W.: Modelling and mapping climate change impacts on permafrost at high spatial resolution for an Arctic region with complex terrain, *The Cryosphere*, 7, 1121–1137, doi:10.5194/tc-7-1121-2013, <http://www.the-cryosphere.net/7/1121/2013/>, 2013.
- 1145
- Zimov, S., Schuur, E., and Chapin III, F.: Permafrost and the global carbon budget, *Science*, 312, 1612–1613, 2006.
- Zubrzycki, S., Kutzbach, L., Pfeiffer, E., and Vakhrameeva, P.: Variability of Soil Organic Carbon Stocks of Different Permafrost-Affected Soils: Initial Results from a North-South Transect in Siberia, in: *Proceedings of the Tenth International Conference on Permafrost*, Salekhard, Russia, pp. 485–490, 2012.
- 1150

Table 1: Overview of the different processes that are accounted for in the models CryoGrid (abbreviated CG) 2 (see Westermann et al., 2013) and 3, as well as in the excess ice scheme Xice. Simplified parameterizations in brackets.

	CG2	CG3	CG3 Xice
<hr/>			
upper boundary condition			
surface temperature	×	-	-
surface energy balance	-	×	×
<hr/>			
snow			
heat conduction	×	×	×
dynamic build-up	×	×	×
ablation due to melt	-	×	×
ablation due to sublimation	-	×	×
compaction	-	-	-
water infiltration	-	×	×
water refreezing	-	×	×
<hr/>			
ground/subsurface			
heat conduction	×	×	×
latent heat effects	×	×	×
water balance	-	-	-
excess ice melt	-	-	×
subsidence	-	-	(×)
heat transfer in water bodies	-	-	(×)

Table 2: Sediment stratigraphies of the long-term susceptibility runs, with volumetric fractions of the soil constituents and soil type for each layer given. Layers super-saturated with ice in bold.

depth [m]	water/ice	mineral	organic	air	natural porosity	type
Arga, no excess ground ice						
0-10	0.4	0.6	0.0	0.0	0.4	sand
>10	0.3	0.7	0.0	0.0	0.3	sand
Samoylov Island, medium excess ground ice content						
0-0.15	0.4	0.1	0.15	0.35	0.75	sand
0.15-0.9	0.65	0.3	0.05	0.0	0.65	silt
0.9-9	0.65	0.3	0.05	0.0	0.4	sand
>9	0.3	0.7	0.0	0.0	0.3	sand
Ice Complex, high excess ground ice content						
0-0.15	0.3	0.1	0.1	0.5	0.8	sand
0.15-0.55	0.7	0.25	0.05	0.0	0.7	sand
0.55-20	0.9	0.05	0.05	0.0	0.4	sand
>20	0.3	0.7	0.0	0.0	0.3	sand

Table 3: Sediment stratigraphies of the validation runs, with volumetric fractions of the soil constituents and soil type for each layer given. Layers super-saturated with ice in bold.

depth [m]	water/ice	mineral	organic	air	natural porosity	type
Samoylov Island, moist polygon center						
0-0.15	0.4	0.1	0.15	0.35	0.75	sand
0.15-0.9	0.65	0.3	0.05	0.0	0.65	silt
0.9-9	0.65	0.3	0.05	0.0	0.4	sand
>9	0.3	0.7	0.0	0.0	0.3	sand
Samoylov Island, thermokarst lake Sa_Lake_1 (Boike et al., 2015)						
0-6	1.0	0.0	0.0	0.0	1.0	sand
6-9	0.4	0.55	0.05	0.0	0.4	sand
>9	0.3	0.7	0.0	0.0	0.3	sand

Table 4: Model parameters assumed in all CryoGrid 3 and CryoGrid 3 Xice simulations.

snow density	ρ_{snow}	200-250	kg m^{-3}	Boike et al. (2013)
albedo snow-free surface	α_{summer}	0.2		Langer et al. (2011a)
albedo pond/lake	α_{lake}	0.07		Burt (1954)
albedo fresh snow	α_{max}	0.85		ECMWF (2006)
albedo old snow	α_{min}	0.50		ECMWF (2006)
time constant of snow albedo change - non-melting	$\tau_{\alpha,f}$	0.008	day^{-1}	ECMWF (2006)
time constant of snow albedo change - melting	$\tau_{\alpha,m}$	0.24	day^{-1}	ECMWF (2006)
SW-radiation extinction coefficient in snow	β_{SW}	25	m^{-1}	Lei et al. (2011), Järvinen and Leppäranta (2011)
new snowfall threshold	SWE_{SF}	0.0025	m	ECMWF (2006)
surface resistance snow-free surface	r_s	50	sm^{-1}	Langer et al. (2011a)
roughness length snow-free surface	$z_{0,\text{summer}}$	10^{-3}	m	Langer et al. (2011a)
roughness length snow	$z_{0,\text{winter}}$	5×10^{-4}	m	Langer et al. (2011b)
emissivity snow-free surface	$\varepsilon_{\text{snow}}$	0.97		Langer et al. (2010)
emissivity snow	$\varepsilon_{\text{snow}}$	0.99		Langer et al. (2011b)
geothermal heat flux	F_{lb}	0.05	Wm^{-2}	Langer et al. (2013)
thermal conductivity mineral soil fraction	k_{mineral}	3.0	$\text{Wm}^{-1}\text{K}^{-1}$	Langer et al. (2013)




Figure 1: The scheme for thawing excess ground ice for an example with saturated ground conditions and equally spaced grid cells (with indicated fractions of ice, water and mineral): when a grid cell with excess ground ice thaws from time step t_1 to t_2 (a \rightarrow b₁), excess water is successively moved upwards from grid cell, while the respective amounts of mineral and organics are moved downwards (b₁ \rightarrow b₆). Following repeated thawing of excess ice grid cells, the uppermost grid cell consists entirely of water (c₇) and an initial pond has formed. In case of the subsidence scenario, such grid cells are removed (c₈^S) and the ground surface is lowered.




Figure 2: Modeled vs. measured daily average surface temperatures for in total 2066 days between 2002 and 2009. The measured surface temperatures are obtained from measurements of outgoing long-wave radiation at the Samoylov climate station (Langer et al., 2011a, b; Boike et al., 2013). The black line represents the 1:1 line.




Figure 3: Modeled and measured average surface energy balance for a) the summer period 7 Jun to 30 Aug 2008 and b) the winter period 1 Dec 2007 to 30 Jan 2008, as documented in Langer et al. (2011a, b). Shown are the net radiation (Q_{net}), sensible, latent, and ground heat flux (Q_h , Q_e , Q_g), and the residual of the surface energy balance C (0 in the model). The modeled fluxes are taken from the validation run with low snow density (200 kg m^{-3}), the values for the high snow density agree within 2 Wm^{-2} . The measured fluxes correspond to Fig. 9 in Langer et al. (2011b).




Figure 4: Modeled and measured ground temperatures at a depth of 0.4 m at a wet polygon center on Samoylov Island (Boike et al., 2013). The blue area depicts the spread between model runs with snow densities of 200 and 250 kg m⁻³.




Figure 5: Modeled and measured thaw depths on Samoylov Island. The measurements correspond to the average of 150 locations on Samoylov Island (Boike et al., 2013). The average standard deviation of the measurements (i.e. the spatial variability of thaw depths) is 0.06 m. The blue area depicts the spread between model runs with snow densities of 200 and 250 kg m⁻³.




Figure 6: Modeled and measured temperature at the bottom of a 6 m deep thermokarst lake on Samoylov Island for the years 2009 to 2012. The measurements are described in detail in Boike et al. (2015), the model results are derived from validation runs for a lake stratigraphy with snow densities of 200 kg m^{-3} and of 250 kg m^{-3} (see Sect. 5.1).




Figure 7: Annual freeze-thaw state for three different cryostratigraphies (top to bottom: Arqa, Samoylov, Ice Complex) and the drainage scenarios “subsidence” (left) and “thermokarst” (right), as simulated by CryoGrid 3 Xice forced by CRU-NCEP/CCSM4 and the moderate future warming scenario RCP 4.5.




Figure 8: Annual freeze-thaw state for three different cryostratigraphies (top to bottom: Arqa, Samoylov, Ice Complex) and the drainage scenarios “subsidence” (left) and “thermokarst” (right), as simulated by CryoGrid 3 Xice forced by CRU-NCEP/CCSM4 and the future warming scenario RCP 8.5.




Figure 9: Annual average temperatures at 10 m below the ground surface (i.e. the actual ground surface in case of subsidence) for the years 1901-2100; two future warming scenarios and three different (cryo-)stratigraphies of the ground.