

# Upscaling methane emission hotspots in boreal peatlands

Fabio Cresto Aleina<sup>1</sup>, Benjamin R. K. Runkle<sup>2,3</sup>, Tim Brücher<sup>1,4</sup>, Thomas Kleinen<sup>1</sup>, and Victor Brovkin<sup>1</sup>

<sup>1</sup>Max Planck Institute for Meteorology, Hamburg, Germany.

<sup>2</sup>Institute of Soil Science, Center for Earth System Research and Sustainability. Universität Hamburg, Hamburg, Germany.

<sup>3</sup>Department of Biological and Agricultural Engineering, University of Arkansas, Fayetteville, AR, USA.

<sup>4</sup>GEOMAR Helmholtz Centre for Ocean Research, Kiel, Germany.

Correspondence to: Fabio Cresto Aleina (fabio.cresto-aleina@mpimet.mpg.de)

**Abstract.** Upscaling properties and effects of small-scale surface heterogeneities to larger scales is a challenging issue in land surface modeling. We developed a novel approach to upscale local methane emissions in a boreal peatland from the micro-topographic scale to the landscape-scale. We based this new parameterization on the analysis of the water table pattern generated by the Hummock-Hollow model, a micro-topography resolving model for peatland hydrology. We introduce this parameterization of methane hotspots in a global model-like version of the Hummock-Hollow model that underestimates methane emissions. We tested the robustness of the parameterization by simulating methane emissions for the next century, forcing the model with three different RCP scenarios. The *Hotspot* parameterization, despite being calibrated for the 1976-2005 climatology, mimics the output of the micro-topography resolving model for all the simulated scenarios. The new approach bridges the scale gap of methane emissions between this version of the model and the configuration explicitly resolving micro-topography.

pel, 2008). In turn, small-scale heterogeneity could influence the land-atmosphere fluxes on larger scale. Several studies have addressed the hydrological cycle in drylands, where water recycled by vegetation may play an important role in the local water budget (Dekker and Rietkerk, 2007; Janssen et al., 2008). In particular, Baudena et al. (2013) showed that the amount of water transferred through transpiration may change up to 10 % if one considers different vegetation patterns, even with the same biomass density and the same spatial scale. Recent efforts have also been focused on downscaling remote sensing information to simulate subgrid surface heterogeneities (e. g., Peng et al., 2015; Stoy and Quaife, 2015), and to scale up information across scales using network techniques (Baudena et al., 2015).

Effects of small-scale heterogeneities on land-atmosphere fluxes are of especial interest in northern peatlands because of the great amount of carbon stored in the soil (Hugelius et al., 2013; Tarnocai et al., 2009). Recent studies have shown that greenhouse gas fluxes, in particular of methane, strongly depend on the micro-topographic features of such environments (Gong et al., 2013; Couwenberg and Fritz, 2012), and that local hydrology is regulated by micro-reliefs (Shi et al., 2015; Gong et al., 2012; Bohn et al., 2013; Van der Ploeg et al., 2012). In particular, a typical feature of methane emitting landscapes is the non linear relationship between fluxes and emitting surface area. A small fraction of the total landscape can therefore function as a "hotspot" for methane fluxes. Recent eddy covariance measurements in northern peatlands showed how the saturated surface, with water table near to the surface level, despite covering only 10 % of the total landscape, is responsible for up to 45 % of the total methane emissions (Sachs et al., 2010).

This "hotspot" feature of methane emissions potentially constitutes a large local and even regional feedback to the climate system, which is neglected in the current Global Circu-

## 1 Introduction

Land surface is a heterogeneous mixture of vegetation types, lakes, wetlands, and bare soil. Correct representation of such small-scale heterogeneities in climate system models is a challenge. How can models better account for the small-scale features in the large-scale climate system? Proposing a new parameterization to fill a scaling gap between local and larger scales is the main focus of this paper. Many recent studies have focused on different approaches to simulate local small-scale characteristics of the land surface, with climate enforcing evolution of different soil surface heterogeneities and small-scale vegetation patterns (Shur and Jorgenson, 2007; Couwenberg and Joosten, 2005; Rietkerk and van de Kop-

lation Models (GCMs), as shown by e. g. Baird et al. (2009). Because of the complexity of the small scale biogeochemical and hydrological interactions that regulate this "hotspot" effect, it is computationally feasible to represent such non-linear phenomena only in local mechanistic models (i. e., Nungesser, 2003; Acharya et al., 2015; Cresto Aleina et al., 2013), with a fine grained resolution ( $10^{-2} - 10^0$  m). The "hotspot" effect is due to the nonlinear relationships between decomposition and its drivers (e. g., soil temperature and water level), and therefore a spatially explicit model able to identify such "hotspots" is likely to perform better in representing methane emissions (Schmidt et al., 2011).

Cresto Aleina et al. (2015) developed the Hummock-Hollow (HH) model, a model for resolving micro-relief features in a typical boreal peatland (hummocks and hollows) and coupled this hydrological model to a process-based model for methane emissions developed by Walter and Heimann (2000). They found that a micro-topography representation is necessary to correctly capture hydrology dynamics and methane fluxes, as the water table position regulates the depth of the oxic zone, where part of the methane coming from the anoxic zone is oxidized and emitted to atmosphere as  $\text{CO}_2$ .

Global land surface models such as JSBACH (Raddatz et al., 2007; Reick et al., 2013), the land component of the Max Planck Institute Earth System Model MPI-ESM (Giorgetta et al., 2013), operate at a spatial resolution analogous to the atmospheric one, which is of about  $50 \text{ km} \times 50 \text{ km}$  at the finest feasible scale. To include a representation of the "hotspot effect" on this scale, new sub-grid scale parameterizations are needed.


In the present paper we propose a novel method to fill the scaling gap from local mechanistic models to large-scale mean field approximations, using the output of the local fine grained model to tune and modify the coarse grained bucket-like model, in order to upscale the local information ( $10^0 - 10^1$  m) to the landscape-scale (e. g.,  $10^3$  m).

We present an application of this upscaling method to the HH model, where we analyze the dynamics of the area which we assume being a hotspot for methane emissions. We then use this information to modify a version of the HH model without micro-topography representation, which originally failed to represent the magnitude of methane fluxes. In this paper we present (i) results for the average climatology of the past 30 years, for which we calibrated the parameterization, and (ii) for the next century, testing the robustness of the parameterization under a different forcing.

## 2 Methods

### 2.1 The HH Model

The Hummock-Hollow (HH) model (Cresto Aleina et al., 2015) simulates peatland micro-topographic controls on



**Figure 1.** Schematics of the HH model showing two grid cells, a hummock and a hollow. The model represents a  $1 \text{ km} \times 1 \text{ km}$  peatland, and works at a  $1 \text{ m} \times 1 \text{ m}$  grid cell. It is therefore able to resolve the micro-topographical features such as hummocks and hollows. The figure shows two typical grid cells, a hummock and a hollow, and the variables needed for the water table dynamics (Equation 1 in the text). Each grid cell has an elevation which is randomly assigned from the distribution of elevation data collected in situ. For each grid cell we simulate a dynamical water table, which changes with snowmelt (Sn), precipitation (P), evapotranspiration (ET), and lateral runoff among the different grid cells ( $R_{\text{hummock/hollow}}$ ). These quantities regulate the change in water table depth (W).

land-atmosphere fluxes. It is suited to work at a  $1 \text{ m} \times 1 \text{ m}$  resolution, which is the typical spatial scale of peatland micro-topography. Each grid cell of the HH model represents just one micro-topographic feature, namely a hummock or a hollow. The model simulates a  $1 \text{ km} \times 1 \text{ km}$  peatland and its parameters are tuned with values for a typical peatland in Northwest Russia. In particular, we use the model to simulate the Ust-Pojeg mire in the Komi Republic ( $61^\circ 56' \text{N}$ ,  $50^\circ 13' \text{E}$ ,  $119 \text{ m a.s.l.}$ ). The site has been extensively studied, and recent efforts described peat characteristics (Pluchon et al., 2014), fluxes of water vapor (Runkle et al., 2012), carbon dioxide (Schneider et al., 2012), and methane (Gažovič et al., 2010), as well as energy and water balance (Runkle et al., 2014) and spatial distribution of dissolved organic carbon (DOC) (Avagyan et al., 2014, 2015). The micro-topography is initialized with micro-topographic data collected through surveying with a theodolite. An elevation distribution is derived from the data, and it is possible to randomly assign an elevation at each grid cell (for more information, Cresto Aleina et al., 2015). Depending on the elevation, the grid cell is therefore either a hollow or a hummock (Fig. 1).

For each grid cell (i. e., for each micro-topographic unit) we compute the water balance as:

$$\frac{dW_{i,j}}{dt} = \frac{Sn + P - ET_{i,j} - R_{i,j}}{s_{i,j}} \quad (1)$$

Configuration	Properties	Resolution
<i>Microtopography</i>	Explicitly resolves micro-topography. Computationally expensive and requires fine scale data for initialization.	1 m × 1 m
<i>Single Bucket</i>	Averages quantities over the domain. Does not consider micro-topography. Computationally fast and requires minimal information for initialization.	1 km × 1 km
<i>Hotspot</i>	Averages quantities over the domain. Considers micro-topographic information. Computationally fast and requires minimal information for initialization.	1 km × 1 km

**Table 1.** Description of the different configurations of the Hummock-Hollow (HH) model used in the present paper.

where  $W_{i,j}$  is the water table level in the grid cell at the position  $(i,j)$  relative to the surface level,  $S_n$  is the snowmelt,  $P$  is the precipitation input,  $ET_{i,j}$  is the evapotranspiration,  $R_{i,j}$  is the lateral runoff,  $s_{i,j}$  is the drainable porosity, and  $t$  is time. The time step is  $\delta t = 1$  day. Terms without the indices  $(i,j)$  are applied uniformly over the model domain. Water table is computed in respect to the micro-topographic surface, and it is positive above the surface, and negative below it. For a description of the parameterization of  $S_n$  and  $ET_{i,j}$  see Appendix A. This version of the model with the explicit representation of hummocks and hollows is called the *Microtopography* configuration.

The HH model can also run in a *Single Bucket* configuration, where all quantities are averaged over the model domain. Equation 1 becomes therefore:

$$\frac{dW}{dt} = \frac{S_n + P - ET - R}{s} \quad (2)$$

The lateral flux is implemented in the same way in the two versions, but in the *Microtopography* version the water can flow from cell to cell, while in the *Single Bucket* version the water simply flows out of the system. Cresto Aleina et al. (2015) showed that the *Single Bucket* configuration, despite being computationally much faster, fails to represent the peatland hydrology, constantly underestimating the water table position in comparison to measurements. This is due to the strong runoff that washes away the water at the beginning of the simulation. Because of the more rugged, hummocky surface represented in the *Microtopography* version, the runoff is delayed. This behavior better agrees with in situ measurements for water table position (Schneider et al., 2012), whereas the water table position simulated by the HH model in the *Single Bucket* configuration is too low. Table 1 describes the main differences between the two configurations of the HH model, and the *Hotspot* parameterization we present in this paper.

## 2.2 Coupling to a process-based methane emission model

The HH model is coupled to a process-based model for methane emissions, in order to quantify the effect of surface heterogeneities on GHG fluxes. The model developed by Walter and Heimann (2000) is a quite general model for methane emissions, and can be applied to peatlands in different environments. It is the same model that is used and coupled with some Dynamical Global Vegetation Models (DGVMs) (e.g., Kleinen et al., 2012; Schuldt et al., 2013; Petrescu et al., 2008; Zhang et al., 2002). We tuned the model to perform in a typical peatland at the latitude of the Ust-Pojeg mire complex. In the *Microtopography* configuration, we computed methane fluxes locally and we averaged over the model domain in order to upscale the local fluxes at the landscape-scale. The process-based model for methane emissions provides an output of methane fluxes  $F_{CH_4}^{i,j}$  as a function of the water table (computed by the HH model), net primary productivity (NPP), and soil temperature (T):

$$F_{CH_4}^{i,j}(t) = f(W_{i,j}(t), NPP(t), T(t)) \quad (3)$$

Where  $W_{i,j}$  is the water table depth with respect to the surface computed at each position  $(i,j)$ . All variables are represented at the daily time step. We force the model with time series of T and NPP taken from CMIP5 experiments performed by the MPI-ESM model. We then considered the model output for the grid cell which corresponds to the Ust-Pojeg mire (see Sect. 2.4). The amount of methane which is emitted by each kind of surface class changes according to the relative position of water table and surface. In the process-based methane emission model developed by Walter and Heimann (2000), the water table is a key variable in methane fluxes, because of the oxidation processes simulated as the water table drops below the surface and as the oxic zone deepens. The HH model in the *Microtopography*

configuration reasonably represents the hydrological interactions among hummocks and hollows and the variability of emissions within the peatland. In the *Single Bucket* configuration the water table drops quickly below the surface after the snow melt due to a strong runoff, and thus most of the methane transported from below ground is oxidized. Parameters for the methane emission model are described in Appendix B.

### 2.3 The *Hotspot* parameterization

The HH Model has a critical scale of about 0.01 km<sup>2</sup> at which seasonal results do not change for finer resolutions (Cresto Aleina et al., 2015). Even at this resolution it is unfeasible to include a micro-topography parameterization in the current GCMs.

The general purpose of our *Hotspot* parameterization is to upscale information from the local to the atmospheric scale. The HH model identifies different surface types depending on the relative position of the water table  $W$  and the surface:

$$\begin{aligned} W > \epsilon_a &\Rightarrow \text{wet surface} \\ -\epsilon_b \leq W \leq \epsilon_a &\Rightarrow \text{saturated surface} \\ W < -\epsilon_b &\Rightarrow \text{dry surface} \end{aligned}$$

Here we assume, after Couwenberg and Fritz (2012):

$$\begin{aligned} \epsilon_a &= 15 \text{ cm} \\ \epsilon_b &= 10 \text{ cm} \end{aligned}$$

because of the importance of such thresholds for methane emissions. We assume the *saturated* surface to be the surface class which dominates the methane emission dynamics, as a water table near to the surface prevents oxidation.

After obtaining the seasonal behavior of the desired surface class, we aim to parameterize of the area covered by the *saturated* surface class with a fractional number  $q$ , which represents the fraction of the total surface which is saturated at each time step. This information results in a different water table behavior which in turns controls methane emissions. By knowing the fraction  $q$  of saturated surface at each time step  $t$  we implicitly subdivide the domain of the HH model in the *Single Bucket* version  $A$  in unsaturated surface  $A_{unsat}$  and saturated surface  $A_{sat}$ :

$$A = (1 - q)A_{unsat} + qA_{sat} \quad (4)$$

The position of the water table in  $A_{sat}$  stays between  $-\epsilon_b \leq W_t^s \leq \epsilon_a$ , which is given by the definition of the saturated surface, and therefore we assume:

$$W_t^s = -\epsilon_b + (\epsilon_a + \epsilon_b)r \quad (5)$$

where  $r$  is a random number between 0 and 1. The position of the water table in  $A_{unsat}$ , instead, is the one computed by the HH model in the *Single Bucket* configuration, i. e.,

$W$  in Eq. 2, which responds to precipitation and evapotranspiration. Methane fluxes are calculated as a function of the water table assuming a linear relationship between emitting area and methane fluxes:

$$F_{CH_4} = (1 - q)F_{CH_4}^{SB}(W) + qF_{CH_4}^{sat}(W_t^s) \quad (6)$$

where  $F_{CH_4}$  is the methane flux from the whole domain,  $F_{CH_4}^{SB}$  the flux from the HH model in the *Single Bucket* version, and  $F_{CH_4}^{sat}$  the flux from the saturated area  $A_{sat}$ . The saturated area fraction  $q$  is defined in Eq. 4. The other forcing variables for  $F_{CH_4}$  stay unchanged, as in Eq. 3.

The specific form of  $q$  as a function of time will be inferred by the analysis of the saturated area dynamics, an output of the HH model in the *Microtopography* configuration.

### 2.4 Forcing data

The HH model is forced with prescribed snowmelt, precipitation, and evapotranspiration (Equation 1). The simulated  $Sn$  is a stochastic input that functions as initialization parameter for the water table. It is parameterized to gain the same magnitude of the observational data (Schneider et al., 2012; Runkle et al., 2014). Evapotranspiration is simulated according to observations of Runkle et al. (2014) using an empirical parameterization. All parameterizations are described in more detail in the Appendices. In Equation 1 we assumed  $Sn$  and  $P$  to be uniform over the whole simulated domain and we did not apply any downscaling further.

We forced the process-based model for methane emissions developed by Walter and Heimann (2000) (Equation 3) and the water balance (Equation 1) with prescribed time series of NPP and T, and of precipitation  $P$  respectively. The time series are computed from simulations performed for the CMIP5 experiments with the MPI-ESM model at T63 resolution for the grid cell which corresponds to the Ust-Pojeg mire. The potential bias introduced by using NPP of C3 grasses and not the one for mosses (not included in the MPI-ESM model) is negligible as discussed by Cresto Aleina et al. (2015).

We used the  $P$ , T, and NPP from the last 30 years of the IPCC historical simulations and forced the model to infer a parameterization of the saturated area (Equations 4 and 6) for the past 30-year climatology. To assess the robustness of our parameterization for future simulations we chose three Representative Concentration Pathways (RCP) scenarios (Taylor et al., 2012), and we therefore considered the identical set of variables from the RCP2.6, RCP4.5, and RCP8.5 experiments from year 2006 to 2099 on daily resolution (Giorgetta et al., 2013).


## 3 Results and Discussion

### 3.1 Hotspot area dynamics

By averaging the output of the model over 30 years of simulations, from 1976 to 2005 we calculated the average dy-

Symbol	Meaning	Value
$t_0$	Initial day of simulation	79
$t_1$	Initial day of maximum saturation	110
$t_2$	Final day of maximum saturation	170
$t_3$	Initial day of minimum saturation	260
$q_{in}$	Initial saturation area density	0.52
$q_{max}$	Maximum saturation area density	0.8
$q_{min}$	Minimum saturation area density	0.5

**Table 2.** Parameter values for Equation 7. We infer the values from the dynamics of the grid cells belonging to the *saturated* surface class as in Figure 2. Days are computed according to the Julian calendar.



**Figure 2.** Area densities for *dry* (red line), *wet* (blue line), and *saturated* (green line) grid cells. The solid lines represent the different surface class dynamics averaged over 30 years, from 1976 to 2005. Shaded areas represent standard deviations over the same period of time. The dynamics of the *saturated* grid cells are mimicked by the empirical *Hotspot* parameterization (black dotted line), Equation 7 in the text.

namics of the three surface classes, wet, saturated, and dry. In particular, we are interested in the 30-year average of the *saturated* area  $A_{sat}$  dynamics (Eq. 4). After snowmelt, most of the simulated peatland surface is either *saturated*, or *wet* (Fig. 2). As the simulations continue, surface and subsurface runoff wash water out of the peatland, changing the relative composition of the area densities. More and more cells become *dry* by having a water table lower than 10 cm below the surface. Grid cells belonging to the *wet* surface class, with a high water table, become saturated and towards the beginning of August virtually no grid cell displays a water table higher than 15 cm above the surface level. At the end of the simulations, almost in all grid cells the water table lays more than 10 cm below the surface level, and the peatland is relatively dry by the end of October.

We used the output of the spatially explicit HH model to describe the dynamics of methane emission hotspots, assuming that the *saturated* grid cells are the ones where methane emissions are higher. We therefore infer the dynamics of the *saturated* grid cells from Figure 2, and obtain the following parameterization for methane emission hotspots:


$$q(t) = \begin{cases} q_{in} + \frac{q_{max} - q_{in}}{t_1 - t_0} (t - t_0) & \text{if } t \leq t_1 \\ q_{max} & \text{if } t_1 < t \leq t_2 \\ q_{max} + \frac{q_{min} - q_{max}}{t_3 - t_2} (t - t_2) & \text{if } t_2 < t \leq t_3 \\ q_{min} & \text{otherwise} \end{cases} \quad (7)$$

where  $t$  is the daily time step of the simulation, and the parameters  $t_i$  and  $q_j$  are tuned quantities obtained according to the dynamics of *saturated* grid cells in Figure 2. Values for the parameterization are described in Table 2. We slightly overestimate the amount of *saturated* grid cells in order to take into account the potential methane emission hotspots belonging to the *wet* surface class.

We illustrate the empirical parameterization of the area density computed by Equation 7 in Figure 2 (black dotted line). This parameterization represents the average dynamics of methane emission hotspots for the 30-year-period 1976-2005.

### 3.2 Methane emissions for 1976-2005

We compared methane emissions from the Ust-Pojeg mire simulated over a 30 year period (1976-2005) in the three versions of the HH model (Table 1). We then averaged the 30 simulations and studied the differences in dynamics among the different HH model versions. The *Microtopography* configuration (black line in Figure 3) produces seasonal fluxes that more than double the cumulative methane fluxes produced by the HH model in the *Single Bucket* configuration (red line in Figure 3). In particular towards July and August, when temperatures are higher and methane fluxes larger, the



**Figure 3.** Methane emissions from the HH model coupled with the Walter and Heimann (2000) model. Solid lines are averages over 30 years (1976-2005) and shaded areas represent standard deviations. Emissions are computed using the HH model in the *Microtopography* configuration (black line), in the *Single Bucket* configuration (red line), and in the *Single Bucket* configuration with the *Hotspot* parameterization (blue line).

Symbol	Meaning	Value	Units
$\text{CH}_4^{SB}$	Cumulative emissions from the <i>Single Bucket</i> configuration	$1.70 \pm 0.11 \times 10^4$	$\text{mg m}^{-2}$
$\text{CH}_4^{Mic}$	Cumulative emissions from the <i>Microtopography</i> configuration	$3.82 \pm 0.30 \times 10^4$	$\text{mg m}^{-2}$
$\text{CH}_4^{HS}$	Cumulative emissions from the <i>Single Bucket</i> configuration with the <i>Hotspot</i> parameterization	$3.47 \pm 0.25 \times 10^4$	$\text{mg m}^{-2}$

**Table 3.** Cumulative emissions from different model configurations. The *Single Bucket* configuration produces less than the half of the cumulative methane emissions with respect to the model with micro-topography representation. By inserting a simple parameterization of the *saturated* surface dynamics, we improve significantly the seasonal methane emissions.

two versions of the HH model diverge in flux estimation 360 and the *Single Bucket* configuration largely underestimates methane fluxes (Cresto Aleina et al., 2015).

Combining Equations 4 and 6, and the empirical parameterization of the hotspot area density  $q(t)$  (Equation 7), we obtain a new flux dynamics (blue line in figure 3). The new 365 parameterized fluxes display similar magnitude and dynam-

ics as the fluxes simulated by the *Microtopography* configuration, but at a much lower computational cost. The main difference between the emissions from the *Single Bucket* and the *Microtopography* configuration is the large underestimation in the central part of the summer season, i. e. in July and August. The *Hotspot* parameterization, by changing the saturated area, improved this feature. The visual improvement

is confirmed by the large differences in the seasonally cumulated methane emissions. The differences in cumulative emissions from the three model configurations are summarized in Table 3.

The *Hotspot* parameterization mimics the general magnitude and dynamics of the emissions from the *Microtopography* configuration, but fails to capture the whole amplitude of methane emissions at the beginning and at the end of the simulations. Such discrepancies might be caused by other variables which, differently from the water table, remain averaged over the domain. In particular, peat depth is uniform and the model does not have a heterogeneous peat profile as in the *Microtopography* configuration. This difference may influence the carbon available for methane emissions.

The *Hotspot* parameterization doubles the cumulative fluxes over the season with respect to the *Single Bucket* configuration, despite its low computational costs. From an ecological perspective, modeling CH<sub>4</sub> fluxes more accurately will improve our estimates of carbon stocks, which may help constrain dynamic vegetation models, bacterial C consumption models, and potential feedbacks with the atmosphere. Also, modeling hydroecological effects of "slower" runoff from a peatland can potentially influence vegetation dynamics of mosses in models including moss dynamics, e. g., Porada et al. (2013). The HH model is novel in the physical representation of lateral fluxes of water among hummocks and hollows, but other models representing surface heterogeneity controls on water table (e. g., Shi et al., 2015) and methane fluxes (e. g., Bohn et al., 2013), display similar effects. Therefore, and because of the process-based nature of the HH model, we are confident in hypothesizing similar results if a *Hotspot*-like parameterization was to be applied to other models.

### 3.3 Future projections with the *Hotspot* parameterization


The *Hotspot* parameterization mimics the simulated methane emissions of the *Microtopography* configuration for the 1976-2005 period for which it has been tuned. We now force the model for the 2006-2099 period with data from the CMIP5 experiments. The HH model does not simulate an increasing trend for methane emissions for the next 100 years, despite the generally higher temperatures (panels (a), (c), and (e) in Figure 4). Even in the RCP8.5 scenario, despite an increase of 4 K in average temperature in year 2099 in respect to the RCP4.5 and the RCP2.6 simulations, we can not find any significant trend. This result is in agreement with the findings from the Wetland and Wetland CH<sub>4</sub> Inter-comparison of Models Project (WETCHIMP) experiments (Melton et al., 2013), which did not find a large significant trend in methane emissions simulated by the models participating in the inter-comparison project because of increased temperature or of precipitation trends. We use these two variables to force the HH model coupled with the methane emis-

sion model. Such an increase is suggested to reduce stomatal conductance, with the same amount of evapotranspiration, thus increasing waterlogged surface area. In particular, Melton et al. (2013) did not find a large significant trend in methane emissions simulated by the models participating in the inter-comparison project because of increased temperature or precipitation trends, which are the two variables we use to force the HH model coupled with the methane emission model.

Moreover, future changes in precipitation could potentially affect the water table position and therefore the saturated area fraction which could not correspond to the one described in the *Hotspot* parameterization. In the RCP simulations, even if precipitation changes in respect to present day and among the scenarios, the differences are not so large to cause significant effects on methane emissions. In panels (a), (c), and (e) of Figure 4, the outputs of the HH model in the *Microtopography* and in the *Single Bucket* configurations, i.e., the black and red lines, respectively, have water table explicitly depending on precipitation simulated in the RCP scenarios. The the *Hotspot* parameterization (i. e., blue lines), despite using the saturated area dynamics for the years 1976-2005, are quite close to the methane emissions from the *Microtopography* configuration. We then conclude that the potential bias introduced by using a fixed saturated area dynamics (the one for the period 1976-2005) and not a dynamic one is negligible.

The *Single Bucket* configuration estimates 42.8 - 50.8 % of the methane emissions cumulated over the season simulated by the *Microtopography* configuration with the RCP8.5 scenario forcing. These estimates are very similar with forcing from the RCP4.5 scenario (44.3 - 50.4 %) and from the RCP2.6 scenario (43.0 - 50.6 %). If we include the *Hotspot* parameterization, the simulated annual methane emissions range from  $[2.831 - 4.321] \times 10^4$  mg m<sup>-2</sup> with the RCP8.5 forcing. This is 83.9 - 101.5 % of the emissions simulated by the *Microtopography* configuration. As for the *Single Bucket* configuration, numbers are similar for the other forcing scenarios. The simulated emissions range from  $[2.771 - 4.056] \times 10^4$  mg m<sup>-2</sup> (88.4 - 100.1 % of the emissions in the *Microtopography* configuration) for the RCP4.5 scenario, and  $[2.648 - 4.102] \times 10^4$  mg m<sup>-2</sup> (87.7 - 104.3 % of the emissions in the *Microtopography* configuration) for the RCP2.6 scenario. Amplitude and timing of year-to-year variability of cumulative methane emissions with the *Hotspot* parameterization are also comparable to the ones simulated by the *Microtopography* configuration in all simulated scenarios.

These results increase the applicability of the *Hotspot* parameterization. Despite being tuned for the 1976-2005 climatology, it works for the next century of simulations under very different forcing scenarios. This is due to the large differences in hydrological representations between the *Microtopography* and *Single Bucket* configurations. Such differences are almost totally overcome with the use of the *Hotspot* parameterization. These improvements make the parameter-



**Figure 4.** Performances of the three configurations of the HH model for future projections in different scenarios. Panels (a), (c), and (e) represent seasonally cumulated methane emissions computed by the HH model forced with CMIP5 data for the time period 2006-2099 from the RCP8.5, 4.5, and 2.6 experiments respectively. Panels (b), (d), and (f) represent the seasonal effectiveness of the *Hotspot* parameterization for future projections, forced with CMIP5 data for the time period 2006-2099 from the RCP8.5, 4.5, and 2.6 experiments respectively. We illustrate the ratio of methane emissions with respect to the *Microtopography* configuration for the *Hotspot* parameterization (in blue) and the *Single Bucket* configuration (in red). We averaged each day of simulation over the 2006-2099 period.



475 ization applicable also for future time slices, despite the dif-  
ferences in temperature, precipitation, and NPP forcing be-  
tween the time period used for the parameterization tuning  
and the scenario projections. 530

We also tested the effectiveness of the *Hotspot* parameter-  
480 ization over the seasonal cycle. We averaged for each simu-  
lated day the methane emissions over the 2005-2099 period  
for all model configurations, and for all scenarios. We then  
divided the daily emissions from the *Single bucket* configura- 535  
tion and from the *Hotspot* parameterization by the emissions  
from the *Microtopography* configuration to investigate the  
485 impact of the new parameterization on the seasonal cycle. In  
all simulated scenarios, the *Hotspot* parameterization works  
very well during the mid season. From mid-May till the be- 540  
ginning of October, when methane emissions are higher, the  
ratio between the *Hotspot* parameterization and the *Microtopo-*  
*graph* parameterization is near one (panels (b), (d), and  
490 (f) in Figure 4). The ratio between emissions from the *Single*  
*Bucket* configuration and the *Microtopography* configuration 545  
reaches its maximum only towards the end of the simula-  
495 tions, therefore missing the larger methane emissions peaks  
in June, July, and August.

#### 4 Summary and Conclusions 550

We developed a new parameterization to bridge the scal-  
ing gap between a process-based, small-scale hydrologi-  
500 cal model for peatlands, and a mean field approximation, 555  
analogous to a large-scale parameterization in a DGVM.  
The *Hotspot* parameterization uses the output of the HH  
(Hummock-Hollow) model (Cresto Aleina et al., 2015)  
which simulates a 1 km  $\times$  1 km peatland. The HH model can  
505 work in both configurations, a spatially explicit one work- 560  
ing at 1 m  $\times$  1 m scale, simulating explicitly hummocks  
and hollows (the *Microtopography* configuration) and a mean  
field approximation of it, where all quantities are averaged  
over the domain (the *Single Bucket* configuration). If cou-  
510 pled to a process-based methane emission model (Walter and 565  
Heimann, 2000) the *Microtopography* configuration simu-  
lates more realistic methane fluxes because of the better rep-  
resentation of hydrology due to the explicit description of  
processes at 1 m scale, but at a much higher computational  
515 cost. We assumed that the lack of representation of saturated 570  
areas in the *Single Bucket* configuration, which are methane  
emission hotspots, diminish the cumulative emissions over  
the season by half.

We inferred a parameterization of this hotspot area for  
520 emissions for the period 1976-2005, which are the last 30 575  
years of the historical simulations from the CMIP5 experi-  
ments. We analyzed the spatial pattern of the HH model out-  
put in the *Microtopography* configuration averaged over the  
30 simulated years. We introduced this information in the  
525 *Single Bucket* configuration, modifying the hydrology of the 580  
mean field approximation, obtaining the *Hotspot* parameter-

ization. This novel approach that takes into account the in-  
formation from the spatially explicit simulations bridges the  
gaps between the simulated methane emissions. The *Hotspot*  
parameterization, due to its higher modified water table, is  
able to mimic the general magnitude and dynamics of the  
emissions from the model with micro-topography represen-  
tation.

By forcing the model with time series of temperature, NPP,  
and precipitation for the next century from CMIP5 experi-  
ments in the RCP8.5, RCP4.5, and RCP2.6 scenarios, we  
assessed the robustness of the *Hotspot* parameterization un-  
der forcing for which it was not originally calibrated. The  
parameterization holds for years 2006-2099 for all three sce-  
narios. Overall, the ratio between the seasonally cumulated  
emissions from the HH model in the *Microtopography* con-  
figuration and the ones simulated by the *Hotspot* parameter-  
ization ranges between 0.84 and 1.04. This is a substantial  
improvement in comparison to the methane emissions simu-  
lated by the *Single Bucket* configuration, which only pro-  
duces between 43 and 51 % of the seasonally cumulated  
methane emissions. The *Hotspot* parameterization at almost  
no computational costs therefore qualitatively changes and  
improves the simulated system response for methane emis-  
sions. 550

We only applied this method to the HH model simulating  
a single peatland in west Russia. This method, though, uses  
the information of a mechanistic, spatially explicit model and  
it is a significant first step towards a full parameterization  
of the micro-topographic impacts on complex ecosystems at  
the DGVM-scale. In order to develop such a parameteriza-  
tion we would need a comprehensive and statistical analy-  
sis on the response of the mechanistic local-scale model to  
different climatic forcing, i. e. we would need HH-like mod-  
els working at the micro-topographic scale applied at differ-  
ent peatlands in other climatic zones. Another limitation of  
the applicability of this study is its dependency on the avail-  
ability of data to calibrate the original HH model in its *Mi-*  
*crotopography* configuration, as accurate measurements of  
peatland micro-relief are needed to initialize surface height.  
While it is not realistic to have theodolite micro-topographic  
measurements globally, other methods and products could  
help provide similar information. Aerial photographs pro-  
vide some information on micro-topography, but generally  
at an overly coarse scale. Statistical downscaling methods as  
the ones used, e. g., by Muster et al. (2012) and Stoy and  
Quaife (2015) are therefore needed to infer information on  
surface heterogeneities, but they are not necessarily useful  
in identifying micro-topography distribution. Airborne mea-  
surements could aid in giving qualitative and stochastic infor-  
mation also on structural peatland patterns, such as the ones  
described by Couwenberg and Joosten (2005). This informa-  
tion could be used by the HH model to generate non ran-  
dom configurations, potentially investigating the influence of  
structured patterns on hydrology and methane emissions.

Introducing the analysis of spatial patterns produced by different mechanistic models in multiple ecosystems is a powerful method to infer landscape-scale dynamics and characteristics of patterns.

625

## 585 Appendix A: Climatology parameterization

Cresto Aleina et al. (2015) did not consider the snow melt  $S_n$ , since they started the simulations later on, and therefore initialized the water table to match the observations at the end of April.  $S_n$  represents the water input at the beginning 630 of the warm season. The cold season is not represented in the model, because we assume that snow covers the area (almost) uniformly. We compute the snowmelt as a random number varying between 200 and 300 mm. We used this range for  $S_n$  in order to obtain an initial water table level on the same order of magnitude of the one observed by Schneider et al. 595 (2012) and Runkle et al. (2014).

Evapotranspiration is dependent on the soil dryness and patchiness. We refer to former studies (Nichols and Brown, 1980), which extensively analyzed the evaporation rate from Sphagnum moss surfaces. The evapotranspiration rate depends on the day of the season, the surface wetness, and on the micro-topographic features. 600 640

$$ET_{i,j} = \begin{cases} \frac{ET_{i,j}^{max}}{fr(W_{i,j})} \sin\left(\frac{(t-4t_0)\pi}{6t_0}\right) & \text{if } 180 < t < 300 \\ \frac{ET_{i,j}^{max}}{fr(W_{i,j})} & \text{otherwise} \end{cases} \quad (A1) \quad 645$$

where  $t$  is the daily time step in days of Julian calendar,  $t_0 = 30$  days is a time constant and  $ET_{i,j}^{max}$  is a function of the micro-topographic features for the cell at the position  $i, j$ : 605 650

$$ET_{i,j}^{max} = \begin{cases} 6 \text{ mm d}^{-1} & \text{if Hummock} \\ 3 \text{ mm d}^{-1} & \text{if Hollow} \end{cases} \quad (A2) \quad 655$$

$fr(W_{i,j})$  takes into account the fact that evaporation takes place at a higher rate if the water table is above the surface: 655

$$fr(W_{i,j}) = \begin{cases} 1 & \text{if } W_{i,j} \text{ above the surface level} \\ 2 & \text{if } W_{i,j} \text{ below the surface level} \end{cases} \quad (A3) \quad 660$$

We use this very simple parameterization of the evapotranspiration rate in order to study the general response of the model to random climatic conditions and to produce quantities in the order of the ones measured by Runkle et al. (2014). 665

## 615 Appendix B: Parameters for the methane emission model

We tuned the parameters used in the process-based model for methane emissions (Walter and Heimann, 2000) in order to apply the model at the latitude of the Ust-Pojeg Mire complex, in the Komi Republic, Russia ( $61^\circ 56'N$ ,  $50^\circ 13'E$ ,  $119$  670

m a.s.l.). Walter and Heimann (2000) used a tuning parameter for the model,  $R_0$ , which we fix at 0.30. Another parameter needed for the coupling with the methane emission model is the soil depth, which we fixed at 150 cm following in situ observation.

*Acknowledgements.* F. Cresto Aleina, B. R. K. Runkle, T. Brücher and V. Brovkin have been supported by the Cluster of Excellence "CliSAP" (EXC177) of the University of Hamburg, as funded by the German Research Foundation (DFG). The authors would like to acknowledge Lars Kutzbach for the help and the fruitful discussions which led to significant model improvements. The authors also thank the three anonymous reviewers that with their comments greatly helped in increasing the clarity and the quality of the paper.

## References

- Acharya, S., Kaplan, D. A., Casey, S., Cohen, M. J., and Jawitz, J. W.: Coupled local facilitation and global hydrologic inhibition drive landscape geometry in a patterned peatland, *Hydrology and Earth System Sciences*, 19, 2133–2144, doi:10.5194/hess-19-2133-2015, <http://www.hydrol-earth-syst-sci.net/19/2133/2015/>, 2015.
- Avagyan, A., Runkle, B., Hartmann, J., and Kutzbach, L.: Spatial Variations in Pore-Water Biogeochemistry Greatly Exceed Temporal Changes During Baseflow Conditions in a Boreal River Valley Mire Complex, Northwest Russia, *Wetlands*, 34, 1171–1182, doi:10.1007/s13157-014-0576-4, <http://dx.doi.org/10.1007/s13157-014-0576-4>, 2014.
- Avagyan, A., Runkle, B. R. K., Hennings, N., Haupt, H., Virtanen, T., and Kutzbach, L.: Dissolved organic matter dynamics during the spring snowmelt at a boreal river valley mire complex in Northwest Russia, *Hydrological Processes*, 2015.
- Baird, A. J., Belyea, L. R., and Morris, P. J.: *Carbon Cycling in Northern Peatlands*, vol. 184 of *Geophysical Monograph Series*, American Geophysical Union, Washington, D. C., doi:10.1029/GM184, <http://www.agu.org/books/gm/v184/2008GM000826/2008GM000826.shtml>, 2009.
- Baudena, M., von Hardenberg, J., and Provenzale, A.: Vegetation patterns and soil-atmosphere water fluxes in drylands, *Advances in Water Resources*, 53, 131–138, doi:10.1016/j.advwatres.2012.10.013, <http://linkinghub.elsevier.com/retrieve/pii/S0309170812002758>, 2013.
- Baudena, M., Sánchez, A., Georg, C.-P., Ruiz-Benito, P., Rodriguez, M. A., Zavala, M. A., and Rietkerk, M.: Revealing patterns of local species richness along environmental gradients with a novel network tool, *Scientific Reports*, 5, doi:doi:10.1038/srep11561, 2015.
- Bohn, T. J., Podest, E., Schroeder, R., Pinto, N., McDonald, K. C., Glagolev, M., Filippov, I., Maksyutov, S., Heimann, M., Chen, X., and Lettenmaier, D. P.: Modeling the large-scale effects of surface moisture heterogeneity on wetland carbon fluxes in the West Siberian Lowland, *Biogeosciences*, 10, 6559–6576, doi:10.5194/bg-10-6559-2013, <http://www.biogeosciences.net/10/6559/2013/>, 2013.
- Couwenberg, J. and Fritz, C.: Towards developing IPCC methane 'emission factors' for peatlands (organic soils), *Mires and*

- 675 Peat, 10, 1–17, [http://mires-and-peat.net/map10/map\\_10\\_03.pdf](http://mires-and-peat.net/map10/map_10_03.pdf), 2012. 735
- Couwenberg, J. and Joosten, H.: Self-organization in raised bog patterning: the origin of microtope zonation and mesotope diversity, *Journal of Ecology*, 93, 1238–1248, doi:10.1111/j.1365-2745.2005.01035.x, <http://doi.wiley.com/10.1111/j.1365-2745.2005.01035.x>, 2005. 740
- Cresto Aleina, F., Brovkin, V., Muster, S., Boike, J., Kutzbach, L., Sachs, T., and Zuyev, S.: A stochastic model for the polygonal tundra based on Poisson-Voronoi diagrams, *Earth System Dynamics*, 4, 187–198, doi:10.5194/esd-4-187-2013, 2013.
- 685 Cresto Aleina, F., Runkle, B. R. K., Kleinen, T., Kutzbach, L., Schneider, J., and Brovkin, V.: Modeling micro-topographic controls on boreal peatland hydrology and methane fluxes, *Biogeosciences*, 12, 5689–5704, doi:10.5194/bg-12-5689-2015, <http://www.biogeosciences.net/12/5689/2015/>, 2015. 745
- 690 Dekker, S. and Rietkerk, M.: Coupling microscale vegetation – soil water and macroscale vegetation – precipitation feedbacks in semiarid ecosystems, *Global change biology*, pp. 1–8, doi:10.1111/j.1365-2486.2006.01327.x, <http://onlinelibrary.wiley.com/doi/10.1111/j.1365-2486.2007.01327.x/full>, 2007. 750
- 695 Gažovič, M., Kutzbach, L., Schreiber, P., Wille, C., and Wilmking, M.: Diurnal dynamics of CH<sub>4</sub> from a boreal peatland during snowmelt, *Tellus B*, 62, 133–139, doi:10.1111/j.1600-0889.2010.00455.x, <http://dx.doi.org/10.1111/j.1600-0889.2010.00455.x>, 2010. 755
- 700 Giorgetta, M. A., Jungclaus, J., Reick, C. H., Legutke, S., Bader, J., Böttinger, M., Brovkin, V., Crueger, T., Esch, M., Fieg, K., Glushak, K., Gayler, V., Haak, H., Hollweg, H.-D., Ilyina, T., Kinne, S., Kornbluh, L., Matei, D., Mauritsen, T., Mikolajewicz, U., Mueller, W., Notz, D., Pithan, F., Raddatz, T., Rast, S., Redler, R., Roeckner, E., Schmidt, H., Schnur, R., Segschneider, J., Six, K. D., Stockhause, M., Timmreck, C., Wegner, J., Widmann, H., Wieners, K.-H., Claussen, M., Marotzke, J., and Stevens, B.: Climate and carbon cycle changes from 1850 to 2100 in MPI-ESM simulations for the Coupled Model Inter-comparison Project phase 5, *Journal of Advances in Modeling Earth Systems*, 5, 572–597, doi:10.1002/jame.20038, <http://dx.doi.org/10.1002/jame.20038>, 2013. 760
- 705 Gong, J., Wang, K., Kellomäki, S., Zhang, C., Martikainen, P. J., and Shurpali, N.: Modeling water table changes in boreal peatlands of Finland under changing climate conditions, *Ecological Modelling*, 244, 65 – 78, doi:<http://dx.doi.org/10.1016/j.ecolmodel.2012.06.031>, <http://www.sciencedirect.com/science/article/pii/S0304380012003213>, 2012. 765
- 710 Gong, J., Kellomäki, S., Wang, K., Zhang, C., Shurpali, N., and Martikainen, P. J.: Modeling {CO<sub>2</sub>} and {CH<sub>4</sub>} flux changes in pristine peatlands of Finland under changing climate conditions, *Ecological Modelling*, 263, 64 – 80, doi:<http://dx.doi.org/10.1016/j.ecolmodel.2013.04.018>, <http://www.sciencedirect.com/science/article/pii/S0304380013002330>, 2013. 770
- 715 Hugelius, G., Tarnocai, C., Broll, G., Canadell, J. G., Kuhry, P., and Swanson, D. K.: The Northern Circumpolar Soil Carbon Database: spatially distributed datasets of soil coverage and soil carbon storage in the northern permafrost regions, *Earth System Science Data*, 5, 3–13, doi:10.5194/essd-5-3-2013, <http://www.earth-syst-sci-data.net/5/3/2013/>, 2013. 775
- Janssen, R. H. H., Meinders, M. B. J., van NES, E. H., and Schef-fer, M.: Microscale vegetation-soil feedback boosts hysteresis in a regional vegetation–climate system, *Global Change Biology*, 14, 1104–1112, doi:10.1111/j.1365-2486.2008.01540.x, <http://doi.wiley.com/10.1111/j.1365-2486.2008.01540.x>, 2008.
- Kleinen, T., Brovkin, V., and Schuldt, R. J.: A dynamic model of wetland extent and peat accumulation: results for the Holocene, *Biogeosciences*, 9, 235–248, doi:10.5194/bg-9-235-2012, <http://www.biogeosciences.net/9/235/2012/>, 2012.
- Melton, J. R., Wania, R., Hodson, E. L., Poulter, B., Ringeval, B., Spahni, R., Bohn, T., Avis, C. A., Beerling, D. J., Chen, G., Eliseev, A. V., Denisov, S. N., Hopcroft, P. O., Lettenmaier, D. P., Riley, W. J., Singarayer, J. S., Subin, Z. M., Tian, H., Zürcher, S., Brovkin, V., van Bodegom, P. M., Kleinen, T., Yu, Z. C., and Kaplan, J. O.: Present state of global wetland extent and wetland methane modelling: conclusions from a model inter-comparison project (WETCHIMP), *Biogeosciences*, 10, 753–788, doi:10.5194/bg-10-753-2013, <http://www.biogeosciences.net/10/753/2013/>, 2013.
- Muster, S., Langer, M., Heim, B., Westermann, S., and Boike, J.: Subpixel heterogeneity of ice-wedge polygonal tundra : a multi-scale analysis of land cover and evapotranspiration in the Lena River Delta , Siberia, *Tellus Series B Chemical And Physical Meteorology*, 1, 1–19, doi:10.3402/tellusb.v64i0.17301, 2012.
- Nichols, D. S. and Brown, J. M.: Evaporation from a sphagnum moss surface, *Journal of Hydrology*, 48, 289 – 302, doi:[http://dx.doi.org/10.1016/0022-1694\(80\)90121-3](http://dx.doi.org/10.1016/0022-1694(80)90121-3), <http://www.sciencedirect.com/science/article/pii/0022169480901213>, 1980.
- Nungesser, M. K.: Modelling microtopography in boreal peatlands: hummocks and hollows, *Ecological Modelling*, 165, 175–207, doi:10.1016/S0304-3800(03)00067-X, <http://linkinghub.elsevier.com/retrieve/pii/S030438000300067X>, 2003.
- Peng, J., Loew, A., Zhang, S., and Wang, J.: Spatial downscaling of satellite soil moisture data using a temperature vegetation dryness index, *IEEE Transactions on Geoscience and Remote Sensing*, Accepted for publication, 2015.
- Petrescu, A. M. R., van Huissteden, J., Jackowicz-Korczynski, M., Yurova, A., Christensen, T. R., Crill, P. M., Bäckstrand, K., and Maximov, T. C.: Modelling CH<sub>4</sub> emissions from arctic wetlands: effects of hydrological parameterization, *Biogeosciences*, 5, 111–121, doi:10.5194/bg-5-111-2008, <http://www.biogeosciences.net/5/111/2008/>, 2008.
- Pluchon, N., Hugelius, G., Kuusinen, N., Kuhry, P., Pluchon, N., Hugelius, G., and Kuusinen, N.: The Holocene storage in two boreal peatlands of Northeast European Russia, *The Holocene*, doi:10.1177/0959683614523803, 2014.
- Porada, P., Weber, B., Elbert, W., Pöschl, U., and Kleidon, A.: Estimating global carbon uptake by lichens and bryophytes with a process-based model, *Biogeosciences*, 10, 6989–7033, doi:10.5194/bg-10-6989-2013, <http://www.biogeosciences.net/10/6989/2013/>, 2013.
- Raddatz, T., Reick, C., Knorr, W., Kattge, J., Roeckner, E., Schnur, R., Schnitzler, K.-G., Wetzell, P., and Jungclaus, J.: Will the tropical land biosphere dominate the climate–carbon cycle feedback during the twenty-first century?, *Climate Dynamics*, 29, 565–574, doi:10.1007/s00382-007-0247-8, <http://dx.doi.org/10.1007/s00382-007-0247-8>, 2007.

- Reick, C., Raddatz, T., Brovkin, V., and Gayler, V.: Representation of natural and anthropogenic land cover change in MPI-ESM, *Journal of Advances in Modeling Earth Systems*, 5, 459–482, 2013.
- Rietkerk, M. and van de Koppel, J.: Regular pattern formation in real ecosystems., *Trends in ecology & evolution* (Personal edition), 23, 169–75, doi:10.1016/j.tree.2007.10.013, <http://www.ncbi.nlm.nih.gov/pubmed/18255188>, 2008.
- Runkle, B., Wille, C., Gažovič, M., Wilmking, M., and Kutzbach, L.: The surface energy balance and its drivers in a boreal peatland fen of northwestern Russia, *Journal of Hydrology*, 511, 359 – 373, doi:<http://dx.doi.org/10.1016/j.jhydrol.2014.01.056>, <http://www.sciencedirect.com/science/article/pii/S002216941400078X>, 2014.
- Runkle, B. R. K., Wille, C., Gažovič, M., and Kutzbach, L.: Attenuation Correction Procedures for Water Vapour Fluxes from Closed-Path Eddy-Covariance Systems, *Boundary-Layer Meteorology*, 142, 401–423, doi:10.1007/s10546-011-9689-y, <http://link.springer.com/10.1007/s10546-011-9689-y>, 2012.
- Sachs, T., Giebels, M., Boike, J., and Kutzbach, L.: Environmental controls on CH<sub>4</sub> emission from polygonal tundra on the microsite scale in the Lena river delta, Siberia, *Global Change Biology*, 16, 3096–3110, doi:10.1111/j.1365-2486.2010.02232.x, <http://doi.wiley.com/10.1111/j.1365-2486.2010.02232.x>, 2010.
- Schmidt, M. W. I., Torn, M. S., Abiven, S., Dittmar, T., Guggenberger, G., Janssens, I. A., Kleber, M., Kogel-Knabner, I., Lehmann, J., Manning, D. A. C., Nannipieri, P., Rasse, D. P., Weiner, S., and Trumbore, S. E.: Persistence of soil organic matter as an ecosystem property., *Nature*, 478, 49–56, doi:10.1038/nature10386, 2011.
- Schneider, J., Kutzbach, L., and Wilmking, M.: Carbon dioxide exchange fluxes of a boreal peatland over a complete growing season, Komi Republic, NW Russia, *Biogeochemistry*, 111, 485–513, doi:10.1007/s10533-011-9684-x, <http://dx.doi.org/10.1007/s10533-011-9684-x>, 2012.
- Schuldt, R. J., Brovkin, V., Kleinen, T., and Winderlich, J.: Modelling Holocene carbon accumulation and methane emissions of boreal wetlands – an Earth system model approach, *Biogeosciences*, 10, 1659–1674, doi:10.5194/bg-10-1659-2013, <http://www.biogeosciences.net/10/1659/2013/>, 2013.
- Shi, X., Thornton, P. E., Ricciuto, D. M., Hanson, P. J., Mao, J., Sebestyen, S. D., Griffiths, N. A., and Bisht, G.: Representing northern peatland microtopography and hydrology within the Community Land Model, *Biogeosciences*, 12, 6463–6477, doi:10.5194/bg-12-6463-2015, <http://www.biogeosciences.net/12/6463/2015/>, 2015.
- Shur, Y. L. and Jorgenson, M. T.: Patterns of permafrost formation and degradation in relation to climate and ecosystems, *Permafrost and Periglacial Processes*, 18, 7–19, doi:10.1002/ppp.582, <http://dx.doi.org/10.1002/ppp.582>, 2007.
- Stoy, P. C. and Quaife, T.: Probabilistic Downscaling of Remote Sensing Data with Applications for Multi-Scale Biogeochemical Flux Modeling, *PLoS ONE*, 10, e0128935, doi:10.1371/journal.pone.0128935, 2015.
- Tarnocai, C., Canadell, J. G., Schuur, E. A. G., Kuhry, P., Mazhitova, G., and Zimov, S.: Soil organic carbon pools in the northern circumpolar permafrost region, *Global Biogeochemical Cycles*, 23, doi:10.1029/2008GB003327, <http://dx.doi.org/10.1029/2008GB003327>, 2009.
- Taylor, K. E., Stouffer, R., and Meehl, G. A.: An overview of cmip5 and the experiment design, *Bull. Amer. Meteor. Soc.*, 93, 485–498, doi:<http://dx.doi.org/10.1175/BAMS-D-11-00094.1>, 2012.
- Van der Ploeg, M. J., Appels, W. M., Cirkel, D. G., Oosterwoud, M. R., Witte, J.-P., and van der Zee, S.: Microtopography as a Driving Mechanism for Ecohydrological Processes in Shallow Groundwater Systems, *Vadose Zone Journal*, 11, <https://www.soils.org/publications/vzj/abstracts/11/3/>, 2012.
- Walter, B. P. and Heimann, M.: A process-based, climate-sensitive model to derive methane emissions from natural wetlands: Application to five wetland sites, sensitivity to model parameters, and climate, *Global Biogeochemical Cycles*, 14, 745–765, 2000.
- Zhang, Y., Li, C., Trettin, C. C., Li, H., and Sun, G.: An integrated model of soil, hydrology, and vegetation for carbon dynamics in wetland ecosystems, *Global Biogeochemical Cycles*, 16, 9–1–9–17, doi:10.1029/2001GB001838, <http://dx.doi.org/10.1029/2001GB001838>, 1061, 2002.