



# UNIVERSITY OF LEEDS

## SCHOOL OF EARTH AND ENVIRONMENT

University of Leeds  
Woodhouse Lane  
Leeds, LS2 9JT, UK  
Tel: +44 (0)113 34 39086  
E-mail: r.ivanovic@leeds.ac.uk  
<http://homepages.see.leeds.ac.uk/~earri>  
[www.see.leeds.ac.uk/people/r.ivanovic](http://www.see.leeds.ac.uk/people/r.ivanovic)

2 June 2016

Dear Jeremy,

Please find attached our revised paper with changes from the previous version tracked. We have made the minor revisions requested by the second reviewer, addressed your request to include a version number, and have also added the required 'data availability' section (and modified the associated manuscript text to point to it). Therefore, we hope the manuscript is now ready to be published in *GMD*.

In addition to these revisions, we have updated the solar constant to match the recently established protocol for CMIP6 simulations (<http://solarisheppa.geomar.de/cmip6>). This had not been fixed at the time of submission.

A detailed response to the reviewer and editorial comments follows. Below that is a version of the manuscript with changes tracked.

Yours sincerely,

A handwritten signature in black ink, appearing to read "Ruža F. Ivanović".

Ruža F. Ivanović

(Line and page numbers in the response below refer to the new version of the manuscript with changes accepted.)

## Reviewer 1

### **Reviewer's summary:**

'The paper describes the design of the coordinated Core simulation over 21-9 ka with time varying orbital forcing, greenhouse gases, ice sheets and other geographical changes. The choice of two ice sheet reconstructions is given and now the change of meltwater flux is taken into account after the paper revision. I

recommend the paper for publication when the following information is provided for completion.'

**1. Reviewer's comment:** The information that PMIP modellers would need is the difference in the two ice sheet boundary condition since the runs demand enormous amount of computer resource and not all can run their model using each ice sheet. For other information such as Greenhouse Gas contents, the paper provides good information on the difference from the previous LGM PMIP and the reason of the discrepancy among data. Since this time two options are given for the ice sheets, PMIP modellers like to know in section 2.4 the advantage and disadvantage of each ice sheet reconstruction (ICE-6G and GLAC) and the main reason of difference to decide which ice sheet one chooses (ex., Why the GLAC is smaller/thinner than ICE-6G?). Also the difference map of Figure 2 would help (difference between the right and left).'

**Authors' response:** We have extended the text in section 2.4 and updated Figures 2 and 4 to add the requested information. Specifically, we have added a description of how ICE-6G\_C was constructed (lines 1-17, page 18); GLAC-1D was already described in the manuscript (lines 18-31 of page 18 and lines 1-5 of page 19). We have also included a section that explains the differences in the reconstructions (lines 15-26, page 19) including the advantages and disadvantages of each. We have updated Figures 2 and 4 with a third column (c) showing the difference between the two reconstructions.

**Reviewer's comment continued:** 'The influence of the different ice sheets could be investigated later in sensitivity studies as is proposed in the paper and PMIP4.

'After adding this information, the paper and the PMIP4 deglaciation experimental design are very useful for many modellers which are eager to contribute to CMIP6. Good Luck for the project!'

## Topical Editor

**Additional request:** 'Could you please add the version number for all relevant boundary conditions to the title/manuscript, according to the standard GMD recommendation for experiment description papers? This will clarify both the present manuscript, and place any future updates to your protocol in good context. Thanks.'

**Authors' response:** We have added the version number (1) back into the title of the manuscript and have clarified the version number in the text (the heading and opening paragraph of section 2) and table captions, as requested. The data repository is also labelled with the version number.

1   **Transient climate simulations of the deglaciation 21-9**  
2   **thousand years before present, [version 1](#); PMIP4 Core**  
3   **experiment design and boundary conditions.**

4   **Ruza F. Ivanovic<sup>1</sup>, Lauren J. Gregoire<sup>1</sup>, Masa Kageyama<sup>2</sup>, Didier M. Roche<sup>2,3</sup>,**  
5   **Paul J. Valdes<sup>4</sup>, Andrea Burke<sup>5</sup>, Rosemarie Drummond<sup>6</sup>, W. Richard Peltier<sup>6</sup>,**  
6   **Lev Tarasov<sup>7</sup>**

7   [1]{School of Earth & Environment, University of Leeds, Woodhouse Lane, Leeds, LS2 9JT,  
8   United Kingdom}

9   [2]{Laboratoire des Sciences du Climat et de l'Environnement, LSCE/IPSL, CEA-CNRS-  
10   UVSQ, Université Paris-Saclay, F-91191 Gif-sur-Yvette, France}

11   [3]{Earth and Climate Cluster, Faculty of Earth and Life Sciences, Vrije Universiteit  
12   Amsterdam, Amsterdam, the Netherlands}

13   [4]{School of Geographical Sciences, University of Bristol, University Road, Bristol, BS8  
14   1SS, United Kingdom}

15   [5]{Department of Earth and Environmental Sciences, Irvine Building, University of St.  
16   Andrews, St. Andrews, KY16 9AL, United Kingdom}

17   [6]{Department of Physics, University of Toronto, 60 St George Street, Toronto, Ontario,  
18   Canada M5S 1A7}

19   [7]{Department of Physics and Physical Oceanography, Memorial University of  
20   Newfoundland and Labrador, St. John's, NL, Canada A1B 3X7}

21   Correspondence to: Ruza F. Ivanovic (r.ivanovic@leeds.ac.uk)

22   **Abstract**

23   The last deglaciation, which marked the transition between the last glacial and present  
24   interglacial periods, was punctuated by a series of rapid (centennial and decadal) climate  
25   changes. Numerical climate models are useful for investigating mechanisms that underpin the  
26   climate change events, especially now that some of the complex models can be run for

multiple millennia. We have set up a Paleoclimate Modelling Intercomparison Project (PMIP) working group to coordinate efforts to run transient simulations of the last deglaciation, and to facilitate the dissemination of expertise between modellers and those engaged with reconstructing the climate of the last 21 thousand years. Here, we present the design of a coordinated Core experiment over the period 21-9 thousand years before present (ka) with time varying orbital forcing, greenhouse gases, ice sheets, and other geographical changes. A choice of two ice sheet reconstructions is given, and we make recommendations for prescribing ice meltwater (or not) in the Core experiment. Additional *focussed* simulations will also be coordinated on an ad-hoc basis by the working group, for example to investigate more thoroughly the effect of ice meltwater on climate system evolution, and to examine the uncertainty in other forcings. Some of these *focussed* simulations will target shorter durations around specific events in order to understand them in more detail and allow the more computationally expensive models to take part.

## 14 1 Introduction

### 15 1.1 Climate evolution over the last deglaciation

16 The last deglaciation is a period of major climate change, when Earth transitioned from its last  
17 full glacial state to the current interglacial climate. The *Last Glacial Maximum* (LGM)  
18 marked the culmination of the last glacial cycle when vast ice sheets covered large regions of  
19 the Northern Hemisphere, stretching over North America and Eurasia (e.g. Boulton et al.,  
20 2001; Dyke et al., 2002; Peltier et al., 2015; Svendsen et al., 2004; Tarasov et al., 2012), and  
21 the Antarctic Ice Sheet expanded to the edge of the continental shelf (Argus et al., 2014;  
22 Briggs et al., 2014; Lambeck et al., 2014 and references therein). Changes in the ice sheets  
23 resulted in a total sea level rise of ~115-130 m between LGM and the late Holocene  
24 (Lambeck et al., 2014; Peltier and Fairbanks, 2006) depending upon the time assumed to  
25 correspond to the LGM, and ~100 m from 21 ka to 9 ka (the period of focus for this  
26 manuscript).

27 Historically, the EPILOG group defined the LGM as having occurred 23-19 ka (21 ka centre  
28 point), when climate was generally cool and ice sheets were more or less at their largest,  
29 based on ice core and sea level records (Mix et al., 2001). It represents the time of maximum

1 terrestrial ice volume. More recently, the last sea level lowstand has been found to have  
2 occurred either around 26 ka (Peltier and Fairbanks, 2006) or 21 ka (Lambeck et al., 2014)  
3 with relatively stable (low) sea level between those dates. Nearly all ice sheets were at or  
4 close to their maximum extent between 26 ka and 19 ka (Clark et al., 2009).

5 During the LGM, global annual mean surface temperatures are estimated to have been around  
6  $4.0 \pm 0.8$  °C colder than today (Annan and Hargreaves, 2013). The Earth began warming  
7 towards its present state from around 19 ka (Fig. 1h; Buizert et al., 2014; Jouzel et al., 2007),  
8 as summer insolation at northern high latitudes and global atmospheric greenhouse gas  
9 concentrations gradually increased (Fig. 1c-f; Bereiter et al., 2015; Berger, 1978; Loulergue et  
10 al., 2008; Marcott et al., 2014). By 9 ka, although the northern ice sheets had not quite  
11 retreated (or disappeared) to their present day configuration, most of the Northern Hemisphere  
12 deglaciation had taken place (Clark et al., 2012; Lambeck et al., 2014; Peltier et al., 2015;  
13 Tarasov et al., 2012; Figures 1g and 2), with both surface air temperatures (Fig. 1h-i) and  
14 atmospheric greenhouse gases (Fig. 1d-f) approaching present day values. However, much of  
15 Antarctica remained heavily glaciated well into the Holocene, with the majority of its  
16 deglacial ice loss taking place between 12 and 6 ka (Argus et al., 2014; Briggs et al., 2014;  
17 Mackintosh et al., 2014). Antarctica's total contribution to post-glacial eustatic sea level is  
18 poorly constrained, but recent studies have not supported LGM contributions greater than  
19 about 15 m eustatic sea level equivalent (Bentley et al., 2014; Briggs et al., 2014; Golledge et  
20 al., 2013; Mackintosh et al., 2011; Philippon et al., 2006; Whitehouse et al., 2012),  
21 emphasising the dominance of North American and Eurasian Ice Sheet dynamics in the global  
22 sea level record during the last deglaciation (Argus et al., 2014; Lambeck et al., 2014; Peltier  
23 et al., 2015). It should be noted that there is some controversy over whether deglacial ice  
24 sheet reconstructions close the global sea level budget (Clark and Tarasov, 2014), with a  
25 potential LGM shortfall of 'missing ice'.

26 The last deglaciation is not only an interesting case study for understanding multi-millennial  
27 scale processes of deglaciation, but also provides the opportunity to study shorter and more  
28 dramatic climate changes. Superimposed over the gradual warming trend (EPICA Community  
29 Members, 2004; Jouzel et al., 2007; Petit et al., 1999; Stenni et al., 2011) are several abrupt  
30 climate transitions lasting from a few years to a few centuries (examples of which are given

1 below) and it remains a challenge to reconstruct or understand the chain of events surrounding  
2 these instances of rapid cooling and warming.

3 *Heinrich Event 1* (approx. 16.8 ka; Hemming, 2004) occurred during the relatively cool  
4 Northern Hemisphere Heinrich Stadial 1 (~18-14.7 ka). It was characterised by the release of  
5 a vast number of icebergs from the North American and Eurasian ice sheets into the open  
6 North Atlantic, where they melted. The existence of these iceberg ‘armadas’ is evidenced by a  
7 high proportion of ice rafted debris in North Atlantic sediments between 40° N and 55° N,  
8 predominantly of Laurentide (Hudson Strait) provenance (Hemming, 2004 and references  
9 therein). There are several competing theories for the cause of Heinrich Event 1. There is a  
10 substantial body of evidence suggesting that it occurred during or was precursory to a period  
11 of Atlantic Meridional Overturning Circulation (AMOC) slow down (e.g. Hall et al., 2006;  
12 Hemming, 2004; McManus et al., 2004) and weak North Atlantic Deep Water (NADW)  
13 formation (e.g. Keigwin and Boyle, 2008; Roberts et al., 2010) under a relatively cold,  
14 Northern Hemisphere surface climate (Shakun et al., 2012). Even though the interpretation of  
15 a cause and effect link between Heinrich Event 1 and the diminished strength of the AMOC  
16 remains rather compelling (e.g. Kageyama et al., 2013), it is increasingly being suggested that  
17 the melting icebergs might not have caused the recorded AMOC slow down, but may have  
18 provided a positive feedback to amplify or prolong AMOC weakening and widespread North  
19 Atlantic cooling (e.g. Álvarez-Solas et al., 2011; Barker et al., 2015).

20 During the subsequent 14.2-14.7 ka interval, Northern Hemisphere temperatures are seen to  
21 have risen by as much as  $14.4 \pm 1.9$  °C in just a few decades (Buijzer et al., 2014; Goujon et  
22 al., 2003; Kindler et al., 2014; Lea et al., 2003; Severinghaus and Brook, 1999), with a  
23 dramatic shift in some components of Greenland climate taking place in as little as one to  
24 three years (Steffensen et al., 2008). This abrupt event is termed the *Bølling Warming* or  
25 *Bølling Transition* (Severinghaus and Brook, 1999). At roughly the same time (~14.6 ka),  
26 there was a rapid jump in global sea level of 12-22 metres in around 350 years or less, known  
27 as *Meltwater Pulse 1a* (MWP1a; Deschamps et al., 2012). It is not known exactly which ice  
28 mass(es) contributed this  $40 \text{ mm yr}^{-1}$  (or greater) flux of water to the oceans (e.g. Lambeck et  
29 al., 2014; Peltier, 2005). Some studies have mainly attributed it to a southern source (Bassett  
30 et al., 2005, 2007; Carlson, 2009; Clark et al., 1996, 2002; Weaver et al., 2003), whereas more  
31 recent work has suggested that at most, less than 4.3 metres eustatic sea level equivalent of

1 meltwater could have come from Antarctica (Argus et al., 2014; Bentley et al., 2010, 2014;  
2 Briggs et al., 2014; Golledge et al., 2012, 2013, 2014; Licht, 2004; Mackintosh et al., 2011,  
3 2014; Whitehouse et al., 2012) and that Northern Hemisphere ice was the primary contributor  
4 (Aharon, 2006; Gregoire et al., 2012; Keigwin et al., 1991; Marshall and Clarke, 1999;  
5 Peltier, 2005; Tarasov et al., 2012; Tarasov and Peltier, 2005). Exactly how the Bølling  
6 Warming and MWP1a are linked, or what triggered either, remains uncertain.

7 Ice core records of  $\delta D$  indicate that from around 14.5 ka to 12.8 ka, the general trend of  
8 increasing Southern Hemisphere warming, temporarily stalled (Jouzel et al., 2007; ice core  
9 chronology from Veres et al., 2013) for a period known as the *Antarctic Cold Reversal*  
10 (Jouzel et al., 1995). Southern Hemisphere cooling is thought to have been relatively  
11 widespread, extending from the South Pole to the southern mid-latitudes, with glacial  
12 readvance (or stall in glacial retreat) recorded to have peaked 13.0-14.2 ka in Patagonia  
13 (García et al., 2012; Kaplan et al., 2011; Strelin et al., 2011) and ~13.0 ka in New Zealand  
14 (Putnam et al., 2010; Rother et al., 2014). There are several hypotheses for the cause of the  
15 Antarctic Cold Reversal. For example, some have linked it to a change in ocean circulation  
16 induced by the delivery of Antarctic ice melt to the Southern Ocean (Menviel et al., 2010,  
17 2011), or possibly as a bipolar response to AMOC recovery and Northern Hemisphere  
18 warming during the Bølling Warming (Menviel et al., 2011; Stocker, 1998). Using a CMIP5  
19 level coupled atmosphere-ocean model, Peltier and Vettoretti (2014) and Vettoretti and Peltier  
20 (2015) have recently shown that ice core inferred southern hemisphere cooling and northern  
21 hemisphere warming could have been caused by a nonlinear salt oscillator mechanism. Others  
22 have argued that a change in Southern Hemisphere winds and ocean circulation is the  
23 explanation; for example, a simultaneous northward migration of the southern Subtropical  
24 Front and northward expansion of cold water originating in the Southern Ocean (Putnam et  
25 al., 2010). The ongoing disagreement over the timing, duration and extent of the Antarctic  
26 Cold Reversal means that its cause is difficult to pin down.

27 The next event of particular interest is the *Younger Dryas cooling*, when Northern  
28 Hemisphere temperatures are thought to have dropped by several degrees at 12.8-11.7 ka and  
29 most prominently in high latitudes (Büizert et al., 2014; Heiri et al., 2007; Lea et al., 2003;  
30 Liu et al., 2012; Simonsen et al., 2011; Steffensen et al., 2008). The event presents a  
31 conceptual paradox; the magnitude of the cooling is difficult to reconcile with rising

1 atmospheric CO<sub>2</sub> (approximately +10 ppm compared to the earlier Bølling period ~ 14.5 ka;  
2 Bereiter et al., 2015) and increasing boreal summer insolation (Berger and Loutre, 1991). It is  
3 possible that changes in the atmospheric hydrological cycle, such as a shift in source moisture  
4 region, could be partly responsible for the δ<sup>18</sup>O signal, requiring a smaller temperature  
5 anomaly to match the records (Liu et al., 2012). For the climate cooling itself, a rerouting of  
6 North American freshwater discharge to the Arctic and/or Atlantic Oceans might have caused  
7 a reduction in NADW formation (Broecker et al., 1989; Condron and Winsor, 2012; Tarasov  
8 and Peltier, 2005). Simulating this period within the context of the preceding climate  
9 evolution could be key to understanding exactly what the surface climate and deep ocean  
10 changes were during the Younger Dryas, and how these relate to contemporaneous proxy  
11 records (e.g. Buizert et al., 2014).

12 In this description, we have sought to capture some of the last deglaciation's main climatic  
13 events, but there are others that could shape the focus of further study in the working group.  
14 For example, early on in the period there is evidence of around 10 m sea level rise taking  
15 place in 500-800 years around 20-19 ka (Clark et al., 2004; Clark and Mix, 2002; De Deckker  
16 and Yokoyama, 2009; Yokoyama et al., 2001a, 2001b). Whilst the event itself remains  
17 somewhat controversial (Cabioc'h et al., 2003; Hanebuth et al., 2000, 2009; Peltier and  
18 Fairbanks, 2006; Shennan and Milne, 2003), it could be the expression of accelerating  
19 deglacial ice melt following the Last Glacial Maximum. More recently, the Barbados record  
20 of relative sea level history indicates that following the Younger Dryas cooling episode, there  
21 may have been another meltwater pulse (Fairbanks, 1989; Peltier and Fairbanks, 2006),  
22 referred to as Meltwater Pulse 1b. Significant debate surrounds the magnitude and timing of  
23 Meltwater Pulse 1b (Bard et al., 1996; Cabioc'h et al., 2003; Cutler et al., 2003; Edwards et al.,  
24 1993; Shennan, 1999; Stanford et al., 2011) and even its existence, because similar to the 19  
25 ka event, it is not seen in all sea level records spanning the interval (e.g. Bard et al., 1996,  
26 2010; Hanebuth et al., 2000). However, evidence of rapid Antarctic retreat around the time of  
27 the event could provide a possible cause for this late deglacial rapid sea level rise (Argus et  
28 al., 2014).

## 1    1.2 Transient modelling of the last deglaciation

2    Transient modelling of the last deglaciation is valuable for examining dynamic and threshold  
3    behaviours (Braconnot et al., 2012) endemic to the Earth's non-stationary climate system,  
4    especially ice-ocean-atmosphere interactions. It is the best tool for reaching a comprehensive  
5    understanding of complex and interrelating climate processes with specific regard to chains of  
6    events.

7    Such simulations are useful for examining the effect of temporally varying climate forcings  
8    across the globe and in different environmental systems: what geographical patterns arise and  
9    how are they connected, how do these vary through time from seasonal to millennial time  
10   scales, and how long does it take before a change in forcing is manifested in a climate  
11   response? The spatial coherency of specific events can be investigated to identify processes  
12   for simultaneous change as well as lead/lag mechanisms. For example, Roche et al. (2011)  
13   investigated patterns of spatial variability in the deglaciation as caused by long-term changes  
14   in orbital parameters, atmospheric greenhouse gas concentrations, and ice sheet  
15   extent/topography. The results indicated a simultaneous onset of hemispheric warming in the  
16   North and South, showing that obliquity forcing was the main driver of the early deglacial  
17   warming. In the same investigation, it was found that sea-ice covered regions were the first  
18   parts of the world to exhibit significant rises in temperature, implying that a better knowledge  
19   of sea-ice evolution could be key to fully understanding the trigger for widespread  
20   deglaciation and warming feedbacks. A further example of the insights available into lead-lag  
21   relationships provided by long, transient climate simulations under glacial boundary  
22   conditions is provided by the previously referenced Dansgaard-Oeschger oscillation-related  
23   analyses of Peltier and Vettoretti (2014) and Vettoretti and Peltier (2015), which appear to  
24   mimic the Heinrich Stadial 1 to Bølling transition.

25   Through comparison to geological timeseries data, transient simulations enable the  
26   'fingerprinting' of specific climate processes to find out what mechanisms [in the model] can  
27   cause recorded climate signals. Comparing complex, global-scale models to combined  
28   geological records can provide multiple 'fingerprints' in different variables from different  
29   archives and in different locations to help narrow down plausible scenarios. For example,  
30   Men viel et al. (2011) ran a suite of simulations, varying oceanic meltwater fluxes through the  
31   last deglaciation in order to identify which freshwater-forcing scenarios reproduce the

1 Atlantic Ocean circulation state implied by sedimentary records of AMOC strength/depth and  
2 ventilation age (Gherardi et al., 2005; McManus et al., 2004 with ages shifted as per Alley,  
3 2000; Thornalley et al., 2011) as well as the Northern Hemisphere surface climate (Alley,  
4 2000; Bard, 2002; Bard et al., 2000; Heiri et al., 2007; Lea et al., 2003; Martrat et al., 2004,  
5 2007). It was argued that such climate simulations could be used to improve constraints on the  
6 timing, duration, magnitude, and location of meltwater inputs to the global ocean.

7 Liu et al. (e.g. 2009) used climate ‘fingerprinting’ to identify possible mechanisms for the  
8 abrupt Bølling Warm Event, finding that in their model, a forced cessation of freshwater  
9 inputs to the North Atlantic (representing ice sheet melt) superimposed on a steady increase in  
10 atmospheric CO<sub>2</sub> caused an abrupt resumption in the strength of the AMOC (almost matching  
11 a record produced by McManus et al., 2004). This in turn induced a rapid warming in  
12 Northern Hemisphere surface climate (close to records from Bard et al., 2000; Cuffey and  
13 Clow, 1997; and Waelbroeck et al., 1998) and an increase in tropical rainfall over the Cariaco  
14 Basin (comparable to Lea et al., 2003), whilst Antarctic surface temperatures remained  
15 relatively stable (similar to Jouzel et al., 2007). Using a suite of simulations from the same  
16 model, Otto-Bliesner et al. (2014) went on to suggest that a combination of rapid  
17 strengthening of NADW seen by Liu et al. (e.g. 2009) and rising greenhouse gas  
18 concentrations was responsible for increased African humidity around 14.7 ka, matching the  
19 model output to a range of regional climate proxies (including deMenocal et al., 2000;  
20 Tierney et al., 2008; Tjallingii et al., 2008; Verschuren et al., 2009; Weijers et al., 2007).

21 Thus, climate proxy fingerprinting can be useful for understanding the spatial coherency of  
22 climatic changes and their underlying mechanisms. However, correlation between model and  
23 geological data does not guarantee that the correct processes have been simulated; there is  
24 always the problem of *equifinality*, whereby the same end state can be reached by multiple  
25 means. In a process sense, this may be particularly uncertain when a model does not  
26 reproduce the full chain of events that led to a distinguishable climatic signal. For example,  
27 mechanisms for many of the major changes in oceanic freshwater inputs proposed by Liu et  
28 al. (2009) and Meniel et al. (2011) have not yet been directly simulated (e.g. by dynamic ice  
29 sheet models). In both studies, they are imposed as model boundary conditions. Further  
30 simulations with different forcing scenarios and from a range of models would help to address  
31 such uncertainties.

1 Transient simulations of the last deglaciation also provide necessary boundary conditions for  
2 modelling a variety of Earth System components that may not be interactively coupled to the  
3 climate model being used. For example, Gregoire et al. (2015) drove a dynamic ice sheet  
4 model with climate data produced by a similar set of simulations to Roche et al. (2011). Using  
5 a low resolution GCM, individual climate forcings – including orbit, greenhouse gases, and  
6 meltwater fluxes – were isolated so that their relative contribution to melting the modelled  
7 North American ice sheets could be examined. The work concluded that the last deglaciation  
8 was primarily driven by changes in Northern Hemisphere insolation, causing around 60% of  
9 the North American Ice Sheet melt, whilst increasing CO<sub>2</sub> levels were responsible for most of  
10 the remaining changes (Gregoire et al., 2015). The sufficiency of these two forcings for North  
11 American glaciation/deglaciation had previously also been identified with fully coupled  
12 glaciological and energy balance climate models (Tarasov and Peltier, 1997). Gregoire et al  
13 (2012) were also able to highlight a ‘saddle-collapse’ mechanism, whereby gradual warming  
14 trends could result in abrupt ice sheet melting events, when a threshold in ice mass balance  
15 was crossed, which could have occurred during MWP1a and the 8.2 kyr event..

16 A further example is given by Liu et al. (2012), who carried out an asynchronous (or ‘offline’)  
17 coupling between simulated sea surface temperatures and an isotope-enabled atmospheric  
18 model to investigate the Younger Dryas cooling event (~12 ka). The results revised the  
19 presupposed Greenland temperatures at this time by 5 °C, demonstrating that changes in  
20 moisture source must be an important consideration for the robust interpretation of Greenland  
21 ice core δ<sup>18</sup>O records and our understanding of high-latitude climate sensitivity. More  
22 recently, the same methodology was applied to understanding Chinese cave records of the  
23 East Asian Summer Monsoon 21-0 ka (Liu et al., 2014), not only to better interpret what the  
24 speleothem δ<sup>18</sup>O tells us about regional hydroclimate variability, but also to understand the  
25 wider teleconnections controlling those patterns.

26 In addition, there are now transient simulations of the last deglaciation from climate models  
27 that have been interactively coupled with dynamic ice sheet models (Bonelli et al., 2009;  
28 Heinemann et al., 2014) and isotope systems (Caley et al., 2014). Furthermore, a fast Earth  
29 System Model of Intermediate Complexity (EMIC) that includes an interactive ice sheet  
30 model has been used to look at Earth System dynamics (the role of orbital cycles, aeolian  
31 dust, subglacial regolith properties, the carbon cycle, and atmospheric trace gases) on much

1 longer, glacial-interglacial timescales >120 ka and encompassing the last deglaciation (Bauer  
2 and Ganopolski, 2014; Brovkin et al., 2012; Ganopolski et al., 2010; Ganopolski and Calov,  
3 2011). However, the older, uncoupled climate-ice sheet model approach discussed above  
4 remains useful because it enables a wider suite of models to be employed than would  
5 otherwise be feasible due to limited computational efficiency (e.g. of state-of-the-art, high  
6 resolution/complexity models) or software engineering capability. It may also allow for the  
7 same Earth System component model (e.g. of ice sheets or  $\delta^{18}\text{O}$ ) to be driven by multiple  
8 climate models, in order to examine the range of responses and assess [climate] model  
9 performance.

10 With sufficient computational power to make long simulations of the last deglaciation a  
11 feasible undertaking, it is timely to coordinate new efforts to ensure that a framework exists to  
12 (i) utilise the cutting edge science in climate modelling and palaeoclimate reconstruction, and  
13 (ii) robustly intercompare simulations run with different models by different groups and  
14 palaeoclimatic data.

### 15 **1.3 Establishing a new PMIP working group**

16 For more than twenty years, the Paleoclimate Modeling Intercomparison Project (PMIP) has  
17 been internationally coordinating multi-model simulations with complex climate models in  
18 order to evaluate model performance and better understand [past] climate changes (Braconnot  
19 et al., 2007, 2012; PMIP website, 2007). Currently entering its fourth phase, PMIP is a  
20 growing organisation that continues to contribute towards other coordinated efforts to  
21 understand present day climate change; including the Coupled Model Intercomparison Project  
22 (Taylor et al., 2011a, CMIP; e.g. 2011b) and the Intergovernmental Panel on Climate  
23 Change's (IPCC) Assessment Reports (e.g. the Fifth Assessment Report; Flato et al., 2013;  
24 Masson-Delmotte et al., 2013). It encompasses a broad range of models, from very fast, lower  
25 resolution EMICS, through a range of coupled GCMs to the latest generation of higher  
26 resolution and complexity Earth System Models. Thus, the main challenges for the fourth  
27 Phase of PMIP include: designing experiments that are suitable for all of its participants;  
28 addressing sufficiently fundamental questions to be of interest to the EMIC community;  
29 defining adequately focused scope for the feasible participation of the latest generation of  
30 ESMs; and prescribing flexible model setups that can be implemented in this range of models,

1 whilst maintaining the ability to robustly compare results. In addition, a continuing challenge  
2 for PMIP is to assemble suitable palaeoclimatic datasets for comparison to model results.

3 One of the most recent working groups to be established in PMIP is the Last Deglaciation  
4 Working Group. With the aim of coordinating transient simulations of the last deglaciation,  
5 the challenge of including the full range of PMIP models is at the forefront of our experiment  
6 design. The experiment will be partitioned into three phases (Fig. 1b and Sect. 4), which will  
7 form milestones for managing its long duration (12 thousand years) as well as for scheduling  
8 any shorter, alternative simulations to the Core.

9 The aim of this paper is to outline the model setup for the transient Core experiment for the  
10 last deglaciation, specifically for the sub-period of 21-9 ka. Prescribed boundary conditions  
11 include orbital parameters, atmospheric trace gases and ice sheets. In association with the ice  
12 sheet reconstructions, we also provide bathymetric, orographic and land-sea mask evolution,  
13 as well as make recommendations for freshwater forcing (or global ocean salinity changes)  
14 through the period.

#### 15 **1.4 Approach**

16 One of the roles of PMIP has been to systematically study the ability of climate models to  
17 retrodict different past climates for which there are ‘observational’ data from geological  
18 archives (e.g. Braconnot et al., 2000, 2007, 2012; Haywood et al., 2010; Joussaume et al.,  
19 1999; Kageyama et al., 2006; Kohfeld and Harrison, 2000; Masson-Delmotte et al., 2006;  
20 Otto-Bliesner et al., 2009; Weber et al., 2007). In this vein, many palaeoclimate model  
21 intercomparison projects have been designed to facilitate the robust comparison of results  
22 from the same ‘experiment’ (i.e. simulation set) across a range of different models, usually  
23 taking a prescriptive approach to model setup to ensure that any differences observed in the  
24 results are attributable to differences in model structure and not to differences in chosen  
25 ‘boundary conditions’ and climate forcings. However, as Schmidt et al. (2011) point out, the  
26 choice of one particular configuration from a range of plausible boundary conditions and  
27 forcings is often arbitrary and does not account for uncertainties in the data used for  
28 developing the forcings/boundary conditions. Moreover, in designing the PMIP last  
29 deglaciation experiment, we have attempted to strike a balance between establishing a  
30 framework within which to assess model differences and performance, and taking the

1 opportunity to utilise the full range of PMIP climate models (Earth System, General  
2 Circulation and Intermediate Complexity) to examine uncertainties in deglacial forcings,  
3 trigger-mechanisms and dynamic feedbacks. In short, when we do not precisely know the  
4 climate forcing for an event, or the temporal evolution of model boundary conditions, it is  
5 more efficient to compare the results from models that use different forcings with geological  
6 and palaeoclimatic data than to run one scenario with all models and all scenarios with all  
7 models. The aim is to use the results of the comparison to narrow down the range of  
8 uncertainty in the forcings/boundary conditions and reach a better understanding of  
9 underlying climate mechanisms.

10 Consequently, forcings/boundary conditions that are relatively well established (atmospheric  
11 trace gases and orbital parameters) are tightly constrained in the Core experiment design.  
12 Others are given with multiple precisely described possibilities to choose from (ice sheet  
13 reconstructions) and the remainder (e.g. freshwater/salinity, aerosols and vegetation) are left  
14 to the discretion of individual participants. Recommendations will be made for the latter  
15 grouping of forcings/boundary conditions; for example, freshwater/global salinity fluxes that  
16 are consistent with the provided ice sheet evolutions, and the use of preindustrial aerosol  
17 and/or vegetation values when they are not model prognostics; but a flexible approach is  
18 advantageous not only scientifically (i.e. for examining the climatic response to uncertain  
19 forcings, see above), but also practically (for accommodating the wide range of participating  
20 models). Further to this, it will be left to the expert user to decide how often to make manual  
21 updates to those boundary conditions that cannot evolve automatically in the model, such as  
22 bathymetry, orography and land sea mask. This is also necessary because of the specific  
23 technical and resource requirements associated with setting up and running each participant  
24 model.

25 In addition to the Core, we will coordinate a series of experiments that are designed to:  
26 (i) explore uncertainties in the boundary conditions and climate forcings,  
27 (ii) test specific hypotheses for mechanisms of climate change and to explain individual  
28 events,  
29 (iii) focus on shorter time periods (for example, abrupt events) and thus include  
30 computationally expensive models for which a twelve thousand year simulation is  
31 unfeasible.

1 These optional simulations will be referred to as *focussed* experiments, and participants are  
2 encouraged to contribute towards the design and coordination of these simulations within the  
3 working group ([there is a dedicated wiki page to coordinate these: PMIP Last Deglaciation](#)  
4 [Working Group, 2016; https://pmip4.lsce.ipsl.fr/doku.php/exp\\_design:degla](#)) ([dedicated Wiki](#)  
5 [page](#) to coordinate these [here](#):  
6 [https://wiki.lsce.ipsl.fr/pmip3/doku.php/pmip3:wg:degla:index](#)).

7 The start date for the experiment has been chosen to be in line with PMIP's historical  
8 definition of the LGM; 21 ka (Abe-Ouchi et al., 2015; e.g. Braconnot et al., 2000; Kohfeld  
9 and Harrison, 2000). However, we are aware that some groups may prefer to begin their  
10 simulations from the earlier date of 26 ka (around the last sea level lowstand; Clark et al.,  
11 2009; Lambeck et al., 2014; Peltier and Fairbanks, 2006) and both orbital and atmospheric  
12 trace gas parameters will be provided from this earlier date. Although the working group's  
13 focus will at least initially be 21-9 ka, boundary conditions for the Core simulations will be  
14 provided from 21 ka to the preindustrial (26 ka to the preindustrial for orbital insolation and  
15 trace gases).

16 The following is not meant to be an exhaustive review of climate forcing reconstructions  
17 through the last deglaciation. Instead, our intention is to consolidate the current knowledge in  
18 a practical experiment design for a range of climate models. Within this coordinated context,  
19 the aim is to explore the forcings and underlying feedback mechanisms for the rapid climate  
20 events that punctuated the gradual warming and deglaciation of the Earth.

21 The paper is structured so that Sect. 2 outlines the model boundary conditions and climate  
22 forcings for the Core experiment. Section 3 presents how we will ensure the feasible  
23 participation of a range of climate models with different complexity and computational  
24 efficiency, as well as the plan to run additional, targeted, hypothesis- and sensitivity-led  
25 simulations. Section 4 discusses the three phases of the long Core experiment.

## 26 **2 Core experiment (21 ka to 9 ka), [version 1](#)**

27 The Core simulations of the last deglaciation ([version 1](#)) will focus on the period from 21 ka  
28 to 9 ka, although there will also be the option to spin up the simulation with time-evolving  
29 orbital and trace gas parameters from 26 ka and all boundary conditions will be available  
30 from 21 ka to the preindustrial. Recommendations for the initialisation state at 21 ka are

1 summarised in Table 1 and described below (Sect. 2.1). Prescribed boundary conditions  
2 include insolation via the Earth's astronomical parameters (Sect. 2.2), atmospheric trace gases  
3 (Sect. 2.3), ice sheets (Sect. 2.4), meltwater fluxes (Sect. 2.5), and orography/bathymetry  
4 (Sect. 2.6), as summarised in Table 2. Boundary condition data for [version 1 of](#) the Core  
5 experiment are provided on the PMIP4 [wiki](#) ([PMIP Last Deglaciation Working Group, 2016](#);  
6 [https://pmip4.lsce.ipsl.fr/doku.php/exp\\_design:deglac](https://pmip4.lsce.ipsl.fr/doku.php/exp_design:deglac));  
7 <https://wiki.lsce.ipsl.fr/pmip3/doku.php/pmip3:wg:deglac:bc:core> ([PMIP Last Deglaciation](#)  
8 [Working Group, 2015](#)).

## 9 **2.1 Last Glacial Maximum spinup**

10 There is a choice of two possibilities for starting the last deglaciation Core simulations. Either  
11 the simulation should be initialised from the end of a spun-up, PMIP-compliant LGM (21 ka)  
12 simulation, or a simulation with transient orbital and trace gas forcing should be run from an  
13 earlier time period (orbital and trace gas parameters will be provided from 26 ka onwards).  
14 Whichever method is applied, we require that it is comprehensively documented along with  
15 information on the model's state of spinup at 21 ka (e.g. timeseries of surface climates,  
16 maximum strength of the North Atlantic Meridional Overturning Circulation stream function,  
17 net radiation at the top of the atmosphere).

### 18 **2.1.1 Equilibrium-type spinup (21 ka)**

19 For setting up an equilibrium-type spinup, please make sure to use the following constraints,  
20 which may differ from other PMIP 21 ka simulation protocols:

- 21 - Insolation should be set so that eccentricity is 0.018994, obliquity is  $22.949^\circ$ ,  
22 perihelion- $180^\circ$  is  $114.42^\circ$ , the date of the vernal equinox is 21st March at noon.  
23 [These data are consistent with previous PMIP LGM boundary conditions \(PMIP](#)  
24 [LGM Working Group, 2010\)](#). [T](#)he solar constant is the same as for the  
25 preindustrial (e.g.  $1361.0 \pm 0.5 \text{ W m}^{-2}$ ; Mamajek et al., 2015; as per CMIP6  
26 [version 3.1](#)(e.g.  $1365 \text{ W m}^{-2}$ , as in the PMIP3-CMIP5 preindustrial experiment).  
27 [These are consistent with previous PMIP LGM boundary conditions \(PMIP LGM](#)  
28 [Working Group, 2010\)](#).

- 1            - Prescribed atmospheric trace gases should be as follows: CO<sub>2</sub> at 190 ppm, CH<sub>4</sub> at  
2            375 ppb, N<sub>2</sub>O at 200 ppb (Fig. 3), with CFCs at 0 and O<sub>3</sub> at the PMIP3-CMIP5  
3            preindustrial value (e.g. 10 DU). This is to be compatible with the time-evolving  
4            boundary conditions for the Core simulations (Sect. 2.3). Note that the LGM  
5            atmospheric CO<sub>2</sub> and CH<sub>4</sub> concentrations have changed slightly from earlier LGM  
6            experiments (e.g. PMIP3, which used 185 ppm and 350 ppb, respectively; PMIP  
7            LGM Working Group, 2010). However, N<sub>2</sub>O remains at 200ppb, which is more  
8            representative of the longer glacial period than the 187 ppb concentration recorded  
9            at 21 ka (Fig. 3c). These updates are in line with the latest ice core age model,  
10            (AICC2012; Veres et al., 2013) and records (Bereiter et al., 2015; Schilt et al.,  
11            2010), which are also used for the transient forcings described below (Sect. 2.3).
- 12            - Prescribed ice sheets should use either the GLAC-1D or ICE-6G\_C reconstruction  
13            at 21 ka (see Sect. 2.4). The associated topography and coastlines should be used  
14            as per the chosen ice sheet reconstruction. Beyond maintaining consistency with  
15            the coastlines, it is optional whether or not to implement the associated bathymetry  
16            and participants should adapt the bathymetry according to their model's  
17            capabilities (for example, depending on whether the spatial resolution allows for it  
18            or makes this a useful adaptation). These data will be provided with the ice sheet  
19            reconstructions. Whichever ice sheet reconstruction is chosen for the LGM spinup  
20            should be carried through to the Core transient simulation.
- 21            - Global ocean salinity should be +1 psu, compared to preindustrial, to account for  
22            the increased terrestrial ice mass at the LGM (PMIP LGM Working Group, 2015).
- 23            - Any other boundary conditions should be set to be consistent with the Core  
24            transient simulation to follow (Sect. 2.2-2.7).

25 On the freshwater budget, PMIP advises groups to ‘carefully check the fresh water budget in  
26 their LGM experiments in order to avoid unnecessary drifts of the ocean salinity. It can be  
27 necessary to route the snow which has fallen in excess on the ice sheets to the ocean. Given  
28 the change in coastlines, it is also sometimes necessary to relocate the large river estuaries on  
29 the coast’ (PMIP LGM Working Group, 2015). Tarasov and Peltier (2006) provide a  
30 glaciological example of the possible re-routings for North America. As they become  
31 available, routing maps for the Last Glacial Maximum continents will be provided on the

1 PMIP4 last deglaciation PMIP\_Wiki—(PMIP Last Deglaciation Working Group,  
2 2016)([address above](#)).

3 The integration time required for spinning up the LGM climate state should be decided on a  
4 case-by-case basis by the user ([see comments by Kageyama et al., 2016, on spin-up and](#)  
5 [duration of experiments](#)). Groups may choose to initialise their equilibrium-type simulation  
6 from other PMIP LGM runs. However, caution is advised. Some of the boundary conditions  
7 for previous PMIP LGM simulations are different to the setup outlined here, specifically in  
8 [terms of relation to](#) ice sheets and trace gases concentrations, and therefore need to be adapted  
9 to match these requirements. The protocol for the PMIP4-CMIP6 (being finalised at the time  
10 of writing) is currently compatible with the LGM spin-up described here. Therefore, provided  
11 that either the ICE-6G\_C or GLAC-1D ice sheet reconstruction is used for both the LGM  
12 spin-up and transient run, the PMIP4-CMIP6 LGM simulation can be used to initialise  
13 transient simulations of the last deglaciation without alteration. Please provide timeseries data  
14 for the diagnosis of model [dis]equilibrium at 21 ka (introduction to Sect. 2.1).

### 15 2.1.2 Transient orbital and trace gas parameters (26-21 ka)

16 If this is the preferred option to initialise the Core, it is recommended that the simulation is  
17 setup as per Sect. 2.1.1, but with time-evolving orbital and trace gas parameters instead of  
18 fixed ones. Specifically for orbit, the eccentricity, obliquity, perihelion–180° and date of the  
19 vernal equinox values listed above should be replaced with their transient equivalents, as per  
20 Berger (1978). For the atmospheric trace gases, carbon dioxide, methane and nitrous oxide  
21 values should be replaced with the transient equivalents provided on the PMIP4 last  
22 deglaciation wiki (PMIP Last Deglaciation Working Group,  
23 2016)(<https://wiki.lsce.ipsl.fr/pmip3/doku.php/pmip3:wg:deglabecore>), which are set and  
24 according to Bereiter et al. (2015), Loulergue et al. (2008) and Schilt et al. (2010),  
25 respectively, on the AICC2012 chronology (Veres et al., 2013); Fig. 3.

26 In this case, all other boundary conditions should remain fixed in line with the LGM  
27 equilibrium-type experiment design until 21 ka, when the fully transient Core simulations  
28 begin. This transient spin-up can be initialised from a spun-up previous LGM, cold ocean,  
29 preindustrial, or observed present day ocean simulation.

1    **2.2 Insolation (21-9 ka)**

2    As per Sect. 2.1, the solar constant should be fixed to the established preindustrial conditions  
3    ([e.g.  \$1361.0 \pm 0.5 \text{ W m}^{-2}\$ ; Mamajek et al., 2015](#)) ([e.g.  \$1365 \text{ W m}^{-2}\$](#) ) throughout the run, which  
4    is [in line with](#) the PMIP preindustrial experiment setup (PMIP LGM Working Group, 2015).  
5    However, the orbital parameters should be time-evolving through the deglaciation to follow  
6    Berger (1978); e.g. Fig. 1c.

7    **2.3 Atmospheric trace gases (21-9 ka)**

8    For the deglaciation, CFCs should be fixed at 0, and O<sub>3</sub> should be set to PMIP3-CMIP5  
9    preindustrial values (e.g. 10 DU), as used for the LGM. When a model is not running with  
10   dynamic atmospheric chemistry, the remaining trace gases should be time-evolving, with CO<sub>2</sub>  
11   following Bereiter et al. (2015), CH<sub>4</sub> following Loulergue et al. (2008) and N<sub>2</sub>O following  
12   Schilt et al. (2010), all adjusted to the AICC2012 chronology (Veres et al., 2013); Fig. 1d-f.

13   The atmospheric CO<sub>2</sub> concentrations provided by Bereiter et al. (2015) is a composite dataset,  
14   combining previous Antarctic ice core records and composites (for the period 26-0 ka: Ahn  
15   and Brook, 2014; Lüthi et al., 2008; MacFarling Meure et al., 2006; Marcott et al., 2014;  
16   Rubino et al., 2013; Siegenthaler et al., 2005) on the AICC2012 timescale of Veres et al.  
17   (2013) to produce a high resolution record that is consistent with the other, lower resolution  
18   trace gas records used in this experiment (CH<sub>4</sub> and N<sub>2</sub>O as discussed above). Groups are free  
19   to decide on the temporal resolution of trace gas model inputs based on these records and if  
20   lower resolution is employed, the method used to smooth or create a spline through the data  
21   should be fully documented. Exploring the influence of CO<sub>2</sub> resolution on the climate system  
22   may form the basis of a coordinated additional simulation, which will be optional for  
23   participant groups. The details of the setup for such *focussed* simulations (also discussed in  
24   Sect. 3) will be discussed and determined at a later date.

25   It is noted that the N<sub>2</sub>O value from Schilt et al. (2010) and Veres et al. (2013) does not match  
26   the previously defined LGM N<sub>2</sub>O concentration (Sect. 2.1.1); 187 ppb compared to 200 ppb  
27   (Fig. 3c). This is because the N<sub>2</sub>O record is highly variable during the last glacial lowstand  
28   (26-21 ka), with a range of ~33 ppb (183-216 ppb) and a mean of 201 ppb. Thus 200 ppb  
29   seems a reasonably representative N<sub>2</sub>O concentration for the spinup phase of the simulation,

1 although the Core simulations will start with the more chronologically accurate value of 187  
2 ppb.

3 **2.4 Ice sheet reconstructions (21-9 ka)**

4 For the Core experiment, ice sheet extent and topography should be prescribed from one of  
5 two possible reconstructions: ICE-6G\_C (Fig. 2a and Fig. 4a) and GLAC-1D (Fig. 2b and  
6 Fig. 4b).

7 The ICE-6G\_C reconstruction is fully published (Argus et al., 2014; Peltier et al., 2015), and  
8 the reader is directed to this literature for further information. The ICE-6G C (VM5a) model  
9 has been designed such that its total mass and local thickness variations provide excellent fits  
10 to all of the data that may be invoked to constrain it. In particular the total mass is constrained  
11 by the globally averaged (eustatic) rise of sea level that is well approximated by the coral  
12 based record of relative sea level history from the island of Barbados in the Caribbean Sea  
13 (Peltier and Fairbanks, 2006). On the other hand, the local variations of ice thickness are  
14 constrained to fit not only the very large number of radiocarbon dated histories of relative sea  
15 level change from the regions that were once ice covered, but also by the voluminous records  
16 of present day vertical motion of the crust that are now available from North America, Eurasia  
17 and Antarctica, based upon space-based Global Positioning System measurements (Argus et  
18 al., 2014; Peltier et al., 2015). Furthermore, the reconstruction includes a history of Antarctic  
19 glaciation that correctly includes the expansion of ice cover in the Ross Sea and Weddell Sea  
20 embayments and out to the shelf break at the LGM. Stuhne and Peltier (2015) assess the  
21 compatibility of ICE-6G C with current understanding of ice dynamical processes using data  
22 assimilation methods. The model is unique internationally insofar as the range of  
23 observational constraints that it has been shown to reconcile. Since the ICE-6G C  
24 reconstruction is fully published (Argus et al., 2014; Peltier et al., 2015), the reader is directed  
25 to this literature for further detailed information.

Formatted: Font color: Auto

26 The GLAC-1D reconstruction is combined from different sources (Briggs et al., 2014;  
27 Tarasov et al., 2012; Tarasov and Peltier, 2002) and whilst it is mostly published, there are  
28 some new components; therefore, a short description follows. The Eurasian and North  
29 American components are from Bayesian calibrations of a glaciological model (Tarasov et al.,  
30 2012; this study), the Antarctic component is from a scored ensemble of 3344 glaciological

1 model runs (Briggs et al., 2014) and the Greenland component is the hand-tuned  
2 glaciological model of Tarasov and Peltier (2002) updated to the GICC05 age chronology  
3 (Rasmussen et al., 2006). All four of the GLAC-1D ice sheet components employ dynamical  
4 ice sheet models that have been constrained with relative sea level data. Where available, they  
5 have also been constrained by geologically-inferred deglacial ice margin chronologies, pro-  
6 glacial lake levels, ice core temperature profiles, present-day vertical velocities, past ice  
7 thickness, and present day ice configuration. Details of exactly how these constraints were  
8 derived and applied are given in the relevant references above. The four components (North  
9 American, Eurasia, Antarctica and Greenland) were combined under Glacial Isostatic  
10 Adjustment (GIA) post-processing for a near-gravitationally self-consistent solution (Tarasov  
11 and Peltier, 2004), which was tested against complete Glacial Isostatic Adjustment solutions.  
12 The topography in the global combined solution was adjusted in Patagonia and Iceland  
13 following ICE-5G (Peltier, 2004), but the changes in these ice caps are not reflected in the ice  
14 mask.

15

16 Both datasets include ice extent and topography at intervals of 1,000 years or less through the  
17 deglaciation. Specifically, the ICE 6G\_C reconstruction is provided at 1,000 year intervals  
18 for the period spanning 26–21 ka and 500 year intervals for 21–0 ka. For GLAC-1D, the data  
19 are at 100 year intervals 21–0 ka. In both reconstructions, ice extent is provided as a fractional  
20 ice mask.

21 The two reconstructions incorporate similar constraints for North American ice sheet extent•  
22 (i.e. Dyke, 2004). For Eurasia, ICE-6G\_C follows the ice extent provided by Gyllencreutz et  
23 al. (2007), whereas GLAC-1D uses data from Hughes et al. (2015). The reconstructions only  
24 differ slightly in their ice extent evolution (Figures 2 and 4), for example the Barents Sea  
25 deglaciates earlier in GLAC-1D than in ICE-6G\_C (Fig. 2). The main differences between the  
26 reconstructions are in the shape and volume of individual ice sheets. In particular, the North  
27 American Ice Sheet reaches an elevation of 4000 m in ICE-6G\_C, but is only 3500 m high in  
28 GLAC-1D. Similarly, the shape and thickness of the Barents Sea Ice Sheet are not the same in  
29 the two reconstructions. The ICE 6G\_C dataset is provided at both 1 degree horizontal  
30 resolution and 10 minute horizontal resolution, GLAC-1D is provided at 1 degree (longitude)  
31 \* 0.5 degree (latitude) horizontal resolution.

**Formatted:** Line spacing: 1.5 lines, Widow/Orphan control,  
Adjust space between Latin and Asian text, Adjust space  
between Asian text and numbers

1 ICE-6G C has overall better fit to relative sea level constraints given the degrees of freedom  
2 from local hand-tuning. GLAC-1D was derived with fewer degrees of freedom given the  
3 internal constraints of glacial physics and assumptions in the climate forcing (which in part  
4 depends on climatologies derived from PMIP2 and PMIP3 results). GLAC-1D incorporates  
5 additional constraints that are inapplicable to purely geophysical constrained models, such as  
6 ICE-6G C. These include inferred pro-glacial lake levels for North America as well as  
7 proximity to the present-day observed Antarctic ice sheet after a transient, multi-glacial cycle  
8 simulation with the underlying ice-earth model. However, this comes at the cost of a 10 to 15  
9 m shortfall in far-field relative sea level for GLAC-1D compared to ICE-6G C (part of the so  
10 called 'missing ice' problem; Clark and Tarasov, 2014). All reconstructions are subject to as  
11 yet unquantified uncertainties, such as the impact of lateral inhomogeneity in the viscous  
12 structure of the Earth.

13 The ICE-6G C dataset is provided at both 1 degree horizontal resolution and 10 minute  
14 horizontal resolution, GLAC-1D is provided at 1 degree (longitude)  $\times$  0.5 degree (latitude)  
15 horizontal resolution. Both datasets include ice extent and topography at intervals of 1,000  
16 years or less through the deglaciation. Specifically, the ICE-6G C reconstruction is provided  
17 at 1,000-year intervals for the period spanning 26-21 ka and 500-year intervals for 21-0 ka.  
18 For GLAC-1D, the data are at 100-year intervals 21-0 ka. In both reconstructions, ice extent is  
19 provided as a fractional ice mask.

20  
21 Ice surface elevation (topography) should be implemented as an anomaly from present day  
22 topography and added to the model's present day topography after regridding onto the model  
23 resolution, following the previous LGM experimental protocol (PMIP LGM Working Group,  
24 2010, 2015). Land surface properties will need to be adjusted for changes in ice extent. Where  
25 ice retreats, land surface should be initialised as bare soil if a dynamic vegetation model is  
26 used, otherwise use prescribed vegetation (see Sect. 2.7) with appropriate consideration of  
27 soil characteristics. Where ice is replaced by ocean, it is advised to follow the procedure for  
28 changing coastlines described in Sect. 2.7. Inland lakes can be prescribed based on the ice  
29 sheet and topography reconstructions, but this is not compulsory. It is also optional whether to  
30 include changes in river routing basins (i.e. catchments) and outlets, which can either be  
31 calculated from the provided topography and land-sea mask data (see Sect. 2.6), or can be

1 manually set to follow routing maps, which will be provided on the [PMIP4](#) last deglaciation  
2 [PMIP wiki \(PMIP Last Deglaciation Working Group, 2016\)](#).

3 Groups are free to choose how often to update ice extent and elevation. This could be done at  
4 regular intervals (e.g. the sub-1000 year time slices provided) or at specific times during the  
5 deglaciation, as was done in the TraCE-21 ka experiment (Liu et al., 2009). Changes in ice  
6 extent can have a large impact on climate through ice albedo changes and feedbacks. We thus  
7 recommend that when possible, ice sheets are not updated at times of abrupt regional or  
8 global climate change, particularly the events that the working group will focus on, as this  
9 could artificially introduce stepped shifts in climate. Groups are also advised to consider that  
10 ice sheet associated boundary conditions (ice extent and elevation, land-sea mask,  
11 bathymetry) may need to be updated more often at times of rapid ice retreat. The timing and  
12 way in which land ice changes are implemented must be documented.

13 Alternative ice sheet reconstructions or simulations can be used to test the sensitivity of  
14 climate to this boundary condition. Simulations with coupled ice sheet-climate models are  
15 also welcomed. Although these will not form part of the Core, for which ICE-6G\_C or  
16 GLAC-1D should be used, they will be coordinated as important supplementary *focussed*  
17 simulations.

18 [For technical notes advising on the implementation of the ice sheet reconstructions in](#)  
19 [palaeoclimate models, see Kageyama et al. \(2016\)](#).

20 **2.5 Ice meltwater**

21 The Core experiment protocol is flexible on whether or not to include prescribed ice melt (i.e.  
22 freshwater fluxes) delivered from the ice sheets to the ocean and how to do it. It is  
23 recommended to run at least one version of the Core experiment with ice melt included, since  
24 around 110 m of ice-volume equivalent sea-level is thought to have melted 26-9 ka (e.g.  
25 Lambeck et al., 2014) and considering the historical importance attached to the influence of  
26 [de]glacial freshwater fluxes on climate (e.g. Broecker et al., 1989; Condron and Winsor,  
27 2012; Ganopolski and Rahmstorf, 2001; Liu et al., 2009; Rahmstorf, 1995, 1996; Teller et al.,  
28 2002; Thornalley et al., 2010; Weaver et al., 2003). However, it is also important to note the  
29 ongoing debate over the extent to which catastrophic freshwater fluxes brought about abrupt  
30 deglacial climate change; several alternative or complementary mechanisms have been

1 proposed (e.g. Adkins et al., 2005; Álvarez-Solas et al., 2011; Barker et al., 2010, 2015;  
2 Broecker, 2003; Hall et al., 2006; Knorr and Lohmann, 2003, 2007; Roche et al., 2007;  
3 Rogerson et al., 2010; Thiagarajan et al., 2014). Moreover, a thorough investigation of the  
4 extent to which non-freshwater-forced climate evolution matches the geological records has  
5 merit in its own right; can abrupt deglacial changes be simulated without ice-meltwater? To  
6 what extent can ‘observed’ patterns be attributed to better constrained forcings, such as  
7 atmospheric CO<sub>2</sub> and Earth’s orbit? It is for all of these reasons that a flexible protocol is  
8 required.

9 Freshwater forcing scenarios consistent with the ice sheet reconstructions and which hence  
10 conserve salinity throughout the deglacial experiment are provided in two formats (the ‘*melt*-’  
11 scenarios described below). In addition, there is the option to run without any ice meltwater  
12 (‘*no-melt*’) to provide a robust reference for simulations that include uncertain meltwater  
13 fluxes. Thus, at least one Core simulation should be run using one of the following ice sheet  
14 meltwater scenarios:

15 *melt-uniform*: a globally uniform freshwater flux (or salinity target) through time, designed to  
16 conserve ocean salinity based on changing terrestrial ice mass. Fluxes  
17 consistent with the ice sheet reconstructions are provided.

18 *melt-routed*: a distributed routing that is consistent with the geographic evolution of the ice  
19 sheet reconstructions (GLAC-1D and ICE-6G\_C; Sect. 2.4) and gives the flux  
20 through time at individual meltwater river outlets along the coast. Again,  
21 versions of this scenario are provided.

22 *no-melt*: no ice meltwater is included in the core; neither a globally integrated ocean  
23 salinity target (*melt-uniform*) nor a distributed routing at the coastlines (*melt-*  
24 *routed*) is implemented. This is best implemented as a sensitivity-type  
25 experiment to account for model-specificness and meltwater flux uncertainty  
26 when also implementing *melt*- scenarios in accompanying versions of the Core  
27 simulation.

28 Multiple Core simulations exploring more than one of these scenarios are welcomed.

29 Data for the *melt*- scenarios will be available from the PMIP4 last deglaciation [wiki \(PMIP](#)  
30 [Last Deglaciation Working Group, 2016\)](#). The data for *melt-uniform* are available at the time

1 of writing (following the respective ice volume changes from ICE-6G\_C and GLAC1-D; Fig.  
2 1g), data for *melt-routed* will be made available as they are produced (anticipated by  
3 [April/May/June/July](#) 2016). These *melt-* scenarios represent a ‘best-estimate’ approach to  
4 resolving the yet unknown geographically- and temporally-precise freshwater fluxes of the  
5 last deglaciation, and they are also consistent with the ice sheet reconstructions employed in  
6 the core. As such, they provide robust and justifiable boundary conditions for simulations that  
7 will be assessed against palaeoclimate reconstructions.

8 However, participants do not have to use the [recommended] versions of *melt-uniform* or  
9 *melt-routed* that are consistent with ICE-6G\_C and GLAC-1D, and can instead use their own  
10 scenarios to explore uncertainty in the ice sheet meltwater flux forcing. This is because the  
11 working group aims to use the full suite of PMIP climate models to examine forcing/boundary  
12 condition uncertainty (see discussion of model intercomparison project approaches in Sect.  
13 1.4). Please note that in some ice melt (including *no-melt*) scenarios, global water budget may  
14 not be balanced through time (as is also true for *no-melt*). Therefore, it is advised to also use  
15 at least one scenario that falls within geological constraints (such as the ICE-6G\_C or GLAC-  
16 1D consistent scenarios for *melt-uniform* and *melt-routed*).

17 Regardless of which scenario is employed, it is important that meltwater fluxes are prescribed  
18 as time-evolving model boundary conditions; rather than as step-wise adjustments at the same  
19 time as the ice sheets are updated, for example. Unless they are intentional conditions of the  
20 scenario, there should be no sudden jumps in the freshwater being applied. Furthermore, we  
21 invite participants to upload the boundary condition data for other freshwater flux scenarios  
22 along with appropriate documentation as/when they become available, and to contribute  
23 towards the coordination of *focussed* experiments (see Sect. 3) that will test specific  
24 hypotheses associated with model and climate sensitivity to the location, duration and  
25 magnitude of freshwater fluxes.

## 26 2.6 Topography, bathymetry, coastlines and rivers

27 [Participants are recommended to note the advice set out by Kageyama et al. \(2016\) for](#)  
28 [implementing these boundary conditions in PMIP4-CMIP6 Last Glacial Maximum and](#)  
29 [Pliocene equilibrium-type experiments.](#)

Formatted: No bullets or numbering

1 Changes in the ice sheets and their glacial eustatic and isostatic influence affected continental  
2 topography and ocean bathymetry, which in turn shifted the coordinates of river mouths and  
3 the coastal outline throughout the deglaciation. Hence time-varying topographic, bathymetric  
4 and land-sea mask fields that match the chosen ice sheet from Sect. 2.4 (i.e. ICE-6G\_C or  
5 GLAC-1D) should be used; these are provided within the ice sheet reconstruction datasets.

6 Topography should be updated at the same time as the model's ice sheet is updated; this is  
7 mainly implicit to implementing the ice sheet reconstruction because the major orographic  
8 changes through the deglaciation relate directly to ice sheet evolution. This said, due to glacial  
9 isostatic adjustment components in the ice sheet reconstructions, there is evolution in  
10 continental topography that is not directly the lowering/heightening of the ice surface, and it  
11 is up to individuals whether they incorporate this or mask only the changes in ice sheet  
12 orography.

13 Ocean bathymetry will be provided. When deemed possible, this boundary condition should  
14 be varied through time. Where differences in the land-sea mask require extra land to fill up  
15 coastal regions, or land to be cut away into ocean as sea level rises (see next paragraph on  
16 coastlines), the model must be changed accordingly, because it is important to adequately  
17 represent the changing land-sea mask; for example, in order to include overlying grounded  
18 ice.

19 Following on from this, coastlines will need to be varied according to changes in global sea  
20 level (and each model's horizontal grid resolution). It will be left to the discretion of  
21 participants to decide how often to update either boundary condition, and when deciding on  
22 their frequency it is recommended that groups consider the implications for opening/closing  
23 seaways and their effect on ocean circulation and climate. Furthermore, the frequency need  
24 not be regular and may instead focus on key 'events' in the marine [gateway] realm.  
25 However, whenever possible and foreseeable, groups are encouraged to avoid making  
26 stepwise changes to model boundary conditions that would interfere with signals of abrupt  
27 climate change; particularly those events that the working group aims to focus on (e.g.  
28 Heinrich Event 1, the Bølling Warming, MWP1a, the Younger Dryas etc.) unless the forcing  
29 (e.g. opening of a gateway) is assumed to be linked with the event.

30 If groups wish, model river networks can be remapped to be consistent with this and updated  
31 on the same timestep as the ice sheet reconstruction, either manually or by the model.

1 However, it is appreciated that the technical challenges associated with such a methodology  
2 would be impractical for many. Therefore, following the recommendation of the PMIP3 LGM  
3 Working Group (2010) [and Kageyama et al. \(2016\)](#)[and Kageyama et al. \(in prep.\)](#), ‘river  
4 pathways and basins should be at least adjusted so that fresh water is conserved at the Earth’s  
5 surface and care should be taken that rivers reach the ocean’ at every timestep that the  
6 bathymetry is adjusted; for example, when sea levels were lower, some river mouths may  
7 need to be displaced towards the [new] coastline to make sure they reach the ocean.

8

## 9 **2.7 Vegetation, land surface and other forcings**

10 In this section, recommendations are made for last deglaciation vegetation, land surface and  
11 aerosol (dust) parameters in the model.

12 There are three recommended options for setting up the Core experiment’s vegetation and  
13 land surface parameters, they can either be: (i) computed using a dynamical vegetation model  
14 (e.g. coupled to the atmospheric component of the model); (ii) prescribed to match the CMIP5  
15 preindustrial setup (Taylor et al., 2011a, 2011b) with fixed vegetation types and fixed plant  
16 physiology (including leaf area index); or (iii) prescribed to match the CMIP5 preindustrial  
17 setup (Taylor et al., 2011a, 2011b) with fixed vegetation types and interactive plant  
18 physiology if running with an enabled carbon cycle. If prescribing vegetation and land  
19 surface, i.e. using option (ii) and (iii), groups should be aware that coastal land will be  
20 emerged compared to preindustrial because of the increased terrestrial ice volume and  
21 associated lower eustatic sea level (with the maximum during the early stages of the Core).  
22 Therefore, vegetation/land surface will need to be interpolated onto the emerged land from  
23 preindustrial grid cells, for example using nearest neighbour methods.

24 For models with prognostic aerosols, the parameters for dust [forcing] can be computed  
25 dynamically. Alternatively, it is recommended that Core simulations fix the associated  
26 parameters according to the CMIP5 preindustrial simulation (Taylor et al., 2011a, 2011b),  
27 with no temporal variation. Examining the influence of different transient aerosol scenarios  
28 (for those models that do not include prognostic dust, for example) could constitute a further  
29 suite of sensitivity simulations for comparison with the Core

1 There is no last deglaciation protocol for setting up other forcings, transient or fixed in time.  
2 For all simulations, groups are required to fully document their methods, including  
3 experiment design and especially when different or with additional components to the setup  
4 described here.

### 5 **3 Coordinating further simulations**

6 As already discussed, we are faced with the challenge of designing an experiment that is  
7 suitable to be run with a wide range of models, from the more computationally efficient class  
8 of intermediate complexity models, to state-of-the-art Earth System Models. One particular  
9 difficulty is enabling the most complex and highest resolution climate models to participate in  
10 this 12 thousand year long experiment when for some, even the integration to reach the LGM  
11 spinup state demands a huge amount of computational resource. There is no easy solution and  
12 our approach will be to augment the Core simulations with shorter *focussed* simulations that  
13 target specific questions, mechanisms and time periods. Whilst the most computationally  
14 expensive models (e.g. the latest generation of Earth System Models) may not be able to  
15 participate in the Core, they will be included in the shorter subset of *focussed* simulations.  
16 Similarly, alternative full-deglaciation simulations can be coordinated for the less  
17 computationally expensive models in the working group (e.g. low resolution General  
18 Circulation Models, and Earth System Models of Intermediate Complexity).

19 One line of investigation relating to meltwater inputs from ice sheets and icebergs is to carry  
20 out a suite of sensitivity simulations examining different injection sites. These simulations  
21 would help to address some of the uncertainty in freshwater flux scenarios. For example,  
22 geochemical evidence suggests that smaller and more localised discharges of freshwater than  
23 have traditionally been considered in climate models may have an important influence on  
24 ocean circulation (e.g. Hall et al., 2006), implying that precise freshwater fluxes are needed in  
25 the models to examine their effect. Certainly, others have shown that the location of injection  
26 is a controlling factor on the impact of freshwater delivery to the ocean, not just laterally (e.g. Roche  
27 et al., 2007). A set of coordinated simulations exploring a range of uncertainty in the  
28 freshwater forcing (location, depth, duration, magnitude, and physical characteristics such as  
29 temperature and density) would be well suited for the *focussed* experiments, thus building on

1 the Core simulations, which may themselves indicate interesting avenues for investigation;  
2 partly the purpose of a flexible meltwater approach.

3 However, freshwater is not the only issue and other *focussed* experiments could include the  
4 influence of greenhouse gas record, differences in ice sheet reconstructions (e.g. the PMIP3  
5 merged ice sheet from Abe-Ouchi et al., 2015; ICE-6G\_C; GLAC-1D) or simulations with  
6 [coupled] ice-sheet models, the relative importance of different forcings (e.g. insolation vs.  
7 trace gases vs. ice sheet evolution), sensitivity to dust-forcing scenarios, the influence of  
8 changes in tidal energy dissipation (Schmittner et al., 2015), event-specific hypothesis testing,  
9 and shorter-term variability within the climate system.

10 Based on on-going discussions, it is likely that the first sets of *focussed* simulations will be:

- 11 • Sensitivity and hypothesis-driven simulations that compare results from uniformly  
12 distributed meltwater fluxes to results from river-routed meltwater fluxes to examine  
13 the impact of the regional specificity of freshwater forcing upon climate system  
14 evolution.
- 15 • Sensitivity simulations that are free from ice meltwater fluxes to provide information  
16 on what climate evolution was caused by processes other than freshwater fluxes to the  
17 ocean.
- 18 • A hypothesis-driven investigation of the possible mechanisms for preconditioning the  
19 glacial ocean for the relatively cool Heinrich Stadial 1 and ensuing catastrophic  
20 iceberg discharge (Barker et al., 2015).
- 21 • Sensitivity experiments examining the role of trace gas forcing resolution on climate  
22 evolution; for example, smoothing the record provided by Bereiter et al. (2015).

23 We have described the plans for *focussed* simulations to highlight the depth of the working  
24 group's aims and to properly contextualise the Core simulations, but the purpose of this  
25 manuscript is to outline the model setup for the Core experiment. The design for subsequent  
26 *focussed* simulations will be described at a later date on the PMIP4 [Last Deglaciation Working Group](#),  
27 [Working Group Wiki \(PMIP Last Deglaciation Working Group, 2016\)](#) (<https://wiki.legi.enpc.fr/pmip3/doku.php/pmip3:wg:deglac:index>) and we welcome  
28 contributions to the discussion of what further simulations to coordinate there.  
29

1   **4 Working group phases**

2   The experiment will be split into three phases that are designed to run seamlessly into each  
3   other (Fig. 1a). Phase one begins at the LGM (21 ka) and will finish at the abrupt Bølling  
4   Warming event, which is where Phase 2 picks up, encompassing the Bølling Warming. Phase  
5   3 begins at the start of the Younger Dryas cooling and is currently planned to continue  
6   through to the end of the Core experiment at 9 ka.

7   Perhaps most importantly, this affords near-future milestones for managing the ultimate  
8   completion of the long full deglacial simulation across all participant groups. It will provide a  
9   timetabled framework for beginning and continuing the longer simulations; for scheduling  
10   shorter, event- or challenge-specific transient simulations by more computationally expensive  
11   models (see discussion in Sect. 3); and for the analysis and publication of results as the  
12   milestones are reached. Another motivation is to ensure that the experiment design for later  
13   periods of the last deglaciation is updated according to knowledge gained from simulations of  
14   the preceding time period; for example, changes in ocean and climate states, which have  
15   previously been shown to have a strong influence on climate trajectories (e.g. Kageyama et  
16   al., 2010; Timm and Timmermann, 2007). This is particularly important for setting up shorter,  
17   event-specific *focussed* simulations, but it is not planned to be explicitly used to influence the  
18   Core. Splitting the period into phases also provides the opportunity to update model boundary  
19   conditions and climate forcing data with cutting edge palaeoclimate reconstructions, as they  
20   emerge during the lifespan of the multi-model experiment. However, care will be taken to  
21   ensure that these are physically consistent between phases, and these updates will not  
22   compromise the Core simulations described in this manuscript. This is so as not to  
23   disadvantage more computationally efficient models that may have already completed  
24   simulating the full 21–9 ka (or beyond) period. Instead, the information will be incorporated  
25   into *focussed* versions of the last deglaciation simulations; possibly spun-off sub-periods that  
26   do not have to start again at the LGM.

27   Each phase will encompass at least one distinguishable climate event; Heinrich Stadial 1 and  
28   Heinrich Event 1 in Phase 1 following on from the LGM; MWP1a, the Bølling Warming and  
29   the Antarctic Cold Reversal in Phase 2; and the Younger Dryas cooling in Phase 3 (Fig. 1b).  
30   As outlined in Sect. 3, simulations of these shorter events can be coordinated in the *focussed*  
31   simulations. This is to engage the higher complexity/resolution models, which are unable to

- 1 run longer simulations, but can use the wider framework of the working group to provide
- 2 valuable knowledge on rapid climate changes known to have taken place in the last 21 ka.

### 3 **5 Summary**

4 The last deglaciation presents a host of exciting opportunities to study the Earth System and  
5 in particular, to try to understand a range of abrupt climate changes that occurred over just a  
6 few years to centuries within the context of more gradual trends. Numerical climate models  
7 provide useful tools to investigate the mechanisms that underpin the events of this well-  
8 studied time period, especially now that technological and scientific advances make it  
9 possible to run multi-millennium simulations with some of the most complex models. Several  
10 recent modelling studies have begun this task, but many questions and untested hypotheses  
11 remain. Therefore, under the auspices of the Paleoclimate Modelling Intercomparison Project  
12 (PMIP), we have set up an initiative to coordinate efforts to run transient simulations of the  
13 last deglaciation, and to facilitate the dissemination of expertise between modellers and those  
14 engaged with reconstructing the climate of the last 21 thousand years.

15 The first step has been to design a Core experiment suitable for a range of PMIP models; from  
16 relatively fast and coarse resolution Earth System Models of Intermediate Complexity, to new  
17 generations of the more complex and higher resolution General Circulation and Earth System  
18 Models. The setup for this Core experiment, is based on an approach that tries to combine a  
19 traditional Model Intercomparison Project method of strictly prescribing boundary conditions  
20 across all models, and the philosophy of utilising the breadth of participants to address  
21 outstanding uncertainty in the climate forcings, model structure and palaeoclimate  
22 reconstructions. Accordingly, we have made recommendations for the initialisation conditions  
23 for the simulation and have stated our minimum requirements for the transient experiment  
24 design, as summarised in Table 1 and 2, respectively.

25 However, there are some uncertainties that the Core is not designed to deal with directly or  
26 exhaustively; two examples discussed in this manuscript being the effect of trace gas record  
27 resolution and the influence of ice melt on the oceans and climate, respectively. We know that  
28 the Core simulations will not tackle all of our questions, and are likely to give rise to others.  
29 Therefore, additional *focussed* simulations will also be coordinated on an ad-hoc basis by the  
30 working group. Many of these will build on and be centred around the Core; often taking

1 shorter snapshots in time, thus including the most computationally expensive models in the  
2 experiment, or presenting twelve-thousand year alternatives to the Core for faster models to  
3 contribute. Not all simulations will be suitable for all models, but the aim is that taken as a  
4 whole, the experiment can utilise the wide range of PMIP model strengths and hence  
5 minimise individual weaknesses.

6 Essentially, the Core experiment has been designed to be inclusive, taking into account the  
7 best compromise between uncertainties in the geological data and model limitations. The  
8 hypothesis-driven *focussed* experiments will go further than the Core to target the questions  
9 that remain. It is hoped that this exciting initiative will improve our individual efforts,  
10 providing new opportunities to drive the science forwards towards understanding this  
11 fascinating time period, specific mechanisms of rapid climate warming, cooling and sea level  
12 change, and Earth's climate system more broadly.

### 13 Author Contributions

14 RFI and LJG lead the PMIP Last Deglaciation Working Group, for which AB, MK, DMR and  
15 PJV act as the advisory group. RFI, LJG, MK, DMR, PJV and AB collaboratively designed  
16 the working group's aims, structure, Core experiment and additional experiments in  
17 consultation with the wider community. RD, WRP and LT provided the ice sheet  
18 reconstructions, plus associated boundary conditions. RFI and LJG collated these and all other  
19 boundary condition data for the simulations. RFI and LJG wrote the manuscript and produced  
20 the figures with contributions from all authors.

Formatted: Heading 5

### 21 Data availability

22 [All boundary condition data required for running the last deglaciation Core experiment](#)  
23 [version 1 \(summarised by Table 1 and Table 2\) can be downloaded from the PMIP4 last](#)  
24 [deglaciation wiki \(PMIP Last Deglaciation Working Group, 2016;](#)  
25 [https://pmip4.lsce.ipsl.fr/doku.php/exp\\_design:degla](https://pmip4.lsce.ipsl.fr/doku.php/exp_design:degla).

Formatted: Heading 5

### 26 Acknowledgments

27 RFI is funded by a NERC Independent Research Fellowship [#NE/K008536/1]. Data  
28 processing for boundary condition preparation was carried out using the computational

1 facilities of the Palaeo@Leeds modelling group, University of Leeds, UK. All authors would  
2 like to thank everyone who has taken the time to discuss the Working Group's aims and  
3 experiments with us. We are especially grateful to Jean-Yves Peterschmitt ([Laboratoire des](#)  
4 [Sciences du Climat et de l'Environnement LSCE](#), France) for [managing and](#) archiving the  
5 boundary conditions, [as well as setting up and maintaining the PMIP wiki pages;](#) [to](#) Emilie  
6 Capron ([British Antarctic Survey](#), UK) for help with the ice core data and [to](#) Bette Otto-  
7 Bliesner ([National Center for Atmospheric Research](#), USA) for useful comments on an earlier  
8 version of this manuscript. Specific thanks also go to Anders Carlson, Eric Wolff, Andreas  
9 Schmittner, Shawn Marshall and an anonymous reviewer for valuable comments on the  
10 manuscript, [and as well as](#) to Jeremy Fyke for editorial handling.

## 11 References

- 12 [Abe-Ouchi, A., Saito, F., Kageyama, M., Braconnot, P., Harrison, S. P., Lambeck, K., Otto-](#) ·  
13 [Bliesner, B. L., Peltier, W. R., Tarasov, L., Peterschmitt, J.-Y. and Takahashi, K.: Ice-sheet](#)  
14 [configuration in the CMIP5/PMIP3 Last Glacial Maximum experiments, Geosci Model Dev](#)  
15 [Discuss, 8\(6\), 4293–4336, doi:10.5194/gmdd-8-4293-2015, 2015.](#)
- 16 [Adkins, J. F., Ingersoll, A. P. and Pasquero, C.: Rapid climate change and conditional](#)  
17 [instability of the glacial deep ocean from the thermobaric effect and geothermal heating, Quat.](#)  
18 [Sci. Rev., 24\(5–6\), 581–594, doi:10.1016/j.quascirev.2004.11.005, 2005.](#)
- 19 [Aharon, P.: Entrainment of meltwaters in hyperpycnal flows during deglaciation super floods](#)  
20 [in the Gulf of Mexico, Earth Planet. Sci. Lett., 241\(1–2\), 260–270,](#)  
21 [doi:10.1016/j.epsl.2005.10.034, 2006.](#)
- 22 [Ahn, J. and Brook, E. J.: Siple Dome ice reveals two modes of millennial CO<sub>2</sub> change during](#)  
23 [the last ice age, Nat. Commun., 5, 3723, doi:10.1038/ncomms4723, 2014.](#)
- 24 [Alley, R. B.: The Younger Dryas cold interval as viewed from central Greenland, Quat. Sci.](#)  
25 [Rev., 19\(1–5\), 213–226, doi:10.1016/S0277-3791\(99\)00062-1, 2000.](#)
- 26 [Álvarez-Solas, J., Montoya, M., Ritz, C., Ramstein, G., Charbit, S., Dumas, C., Nisancioglu,](#)  
27 [K., Dokken, T. and Ganopolski, A.: Heinrich event 1: an example of dynamical ice-sheet](#)  
28 [reaction to oceanic changes, Clim Past, 7\(4\), 1297–1306, doi:10.5194/cp-7-1297-2011, 2011.](#)

**Formatted:** Bibliography, Widow/Orphan control, Adjust space between Latin and Asian text, Adjust space between Asian text and numbers

**Field Code Changed**

- 1 [Annan, J. D. and Hargreaves, J. C.: A new global reconstruction of temperature changes at](#)  
2 [the Last Glacial Maximum. Clim Past, 9\(1\), 367–376, doi:10.5194/cp-9-367-2013, 2013.](#)
- 3 [Argus, D. F., Peltier, W. R., Drummond, R. and Moore, A. W.: The Antarctica component of](#)  
4 [postglacial rebound model ICE-6G\\_C \(VM5a\) based on GPS positioning, exposure age dating](#)  
5 [of ice thicknesses, and relative sea level histories. Geophys. J. Int., ggu140,](#)  
6 [doi:10.1093/gji/ggu140, 2014.](#)
- 7 [Bard, E.: Climate shock- Abrupt changes over millenial time scales. Phys. Today, 55, 32–38,](#)  
8 [2002.](#)
- 9 [Bard, E., Hamelin, B., Arnold, M., Montaggioni, L., Cabioch, G., Faure, G. and Rougerie, F.:](#)  
10 [Deglacial sea-level record from Tahiti corals and the timing of global meltwater discharge.](#)  
11 [Nature, 382\(6588\), 241–244, doi:10.1038/382241a0, 1996.](#)
- 12 [Bard, E., Rostek, F., Turon, J.-L. and Gendreau, S.: Hydrological Impact of Heinrich Events](#)  
13 [in the Subtropical Northeast Atlantic. Science, 289\(5483\), 1321–1324,](#)  
14 [doi:10.1126/science.289.5483.1321, 2000.](#)
- 15 [Bard, E., Hamelin, B. and Delanghe-Sabatier, D.: Deglacial Meltwater Pulse 1B and Younger](#)  
16 [Dryas Sea Levels Revisited with Boreholes at Tahiti. Science, 327\(5970\), 1235–1237,](#)  
17 [doi:10.1126/science.1180557, 2010.](#)
- 18 [Barker, S., Knorr, G., Vautravers, M. J., Diz, P. and Skinner, L. C.: Extreme deepening of the](#)  
19 [Atlantic overturning circulation during deglaciation. Nat. Geosci., 3\(8\), 567–571,](#)  
20 [doi:10.1038/ngeo921, 2010.](#)
- 21 [Barker, S., Chen, J., Gong, X., Jonkers, L., Knorr, G. and Thornalley, D.: Icebergs not the](#)  
22 [trigger for North Atlantic cold events. Nature, 520\(7547\), 333–336, doi:10.1038/nature14330,](#)  
23 [2015.](#)
- 24 [Bassett, S. E., Milne, G. A., Mitrovica, J. X. and Clark, P. U.: Ice Sheet and Solid Earth](#)  
25 [Influences on Far-Field Sea-Level Histories. Science, 309\(5736\), 925–928,](#)  
26 [doi:10.1126/science.1111575, 2005.](#)
- 27 [Bassett, S. E., Milne, G. A., Bentley, M. J. and Huybrechts, P.: Modelling Antarctic sea-level](#)  
28 [data to explore the possibility of a dominant Antarctic contribution to meltwater pulse IA.](#)  
29 [Quat. Sci. Rev., 26\(17–18\), 2113–2127, doi:10.1016/j.quascirev.2007.06.011, 2007.](#)

- 1 [Bauer, E. and Ganopolski, A.: Sensitivity simulations with direct shortwave radiative forcing](#)  
2 [by aeolian dust during glacial cycles, Clim Past, 10\(4\), 1333–1348, doi:10.5194/cp-10-1333-](#)  
3 [2014, 2014.](#)
- 4 [Bentley, M. J., Fogwill, C. J., Brocq, A. M. L., Hubbard, A. L., Sugden, D. E., Dunai, T. J.](#)  
5 [and Freeman, S. P. H. T.: Deglacial history of the West Antarctic Ice Sheet in the Weddell](#)  
6 [Sea embayment: Constraints on past ice volume change, Geology, 38\(5\), 411–414,](#)  
7 [doi:10.1130/G30754.1, 2010.](#)
- 8 [Bentley, M. J., Ó Cofaigh, C., Anderson, J. B., Conway, H., Davies, B., Graham, A. G. C.,](#)  
9 [Hillenbrand, C.-D., Hodgson, D. A., Jamieson, S. S. R., Larter, R. D., Mackintosh, A., Smith,](#)  
10 [J. A., Verleyen, E., Ackert, R. P., Bart, P. J., Berg, S., Brunstein, D., Canals, M., Colhoun, E.](#)  
11 [A., Crosta, X., Dickens, W. A., Domack, E., Dowdeswell, J. A., Dunbar, R., Ehrmann, W.,](#)  
12 [Evans, J., Favier, V., Fink, D., Fogwill, C. J., Glasser, N. F., Gohl, K., Golledge, N. R.,](#)  
13 [Goodwin, I., Gore, D. B., Greenwood, S. L., Hall, B. L., Hall, K., Hedding, D. W., Hein, A.](#)  
14 [S., Hocking, E. P., Jakobsson, M., Johnson, J. S., Jomelli, V., Jones, R. S., Klages, J. P.,](#)  
15 [Kristoffersen, Y., Kuhn, G., Leventer, A., Licht, K., Lilly, K., Lindow, J., Livingstone, S. J.,](#)  
16 [Massé, G., McGlone, M. S., McKay, R. M., Melles, M., Miura, H., Mulvaney, R., Nel, W.,](#)  
17 [Nitsche, F. O., O'Brien, P. E., Post, A. L., Roberts, S. J., Saunders, K. M., Selkirk, P. M.,](#)  
18 [Simms, A. R., Spiegel, C., Stolldorf, T. D., Sugden, D. E., van der Putten, N., van Ommen,](#)  
19 [T., Verfaillie, D., Vyverman, W., Wagner, B., White, D. A., Witte, A. E. and Zwartz, D.: A](#)  
20 [community-based geological reconstruction of Antarctic Ice Sheet deglaciation since the Last](#)  
21 [Glacial Maximum, Quat. Sci. Rev., 100, 1–9, doi:10.1016/j.quascirev.2014.06.025, 2014.](#)
- 22 [Bereiter, B., Eggleston, S., Schmitt, J., Nehrbass-Ahles, C., Stocker, T. F., Fischer, H.,](#)  
23 [Kipfstuhl, S. and Chappellaz, J.: Revision of the EPICA Dome C CO<sub>2</sub> record from 800 to](#)  
24 [600 kyr before present, Geophys. Res. Lett., 42\(2\), 2014GL061957,](#)  
25 [doi:10.1002/2014GL061957, 2015.](#)
- 26 [Berger, A.: Long-Term Variations of Daily Insolation and Quaternary Climatic Changes, J.](#)  
27 [Atmospheric Sci., 35\(12\), 2362–2367, doi:10.1175/1520-](#)  
28 [0469\(1978\)035<2362:LTVODI>2.0.CO;2, 1978.](#)
- 29 [Berger, A. and Loutre, M. F.: Insolation values for the climate of the last 10 million years,](#)  
30 [Quat. Sci. Rev., 10\(4\), 297–317, doi:10.1016/0277-3791\(91\)90033-Q, 1991.](#)

1 Bonelli, S., Charbit, S., Kageyama, M., Woillez, M.-N., Ramstein, G., Dumas, C. and  
2 Quiquet, A.: Investigating the evolution of major Northern Hemisphere ice sheets during the  
3 last glacial-interglacial cycle, *Clim Past*, 5(3), 329–345, doi:10.5194/cp-5-329-2009, 2009.  
4 Boulton, G. S., Dongelmans, P., Punkari, M. and Broadgate, M.: Palaeoglaciology of an ice  
5 sheet through a glacial cycle:: the European ice sheet through the Weichselian, *Quat. Sci.*  
6 *Rev.*, 20(4), 591–625, doi:10.1016/S0277-3791(00)00160-8, 2001.  
7 Braconnot, P., Joussaume, S., de Noblet, N. and Ramstein, G.: Mid-Holocene and Last  
8 Glacial Maximum African monsoon changes as simulated within the Paleoclimate Modelling  
9 Intercomparison Project, *Glob. Planet. Change*, 26(1–3), 51–66, doi:10.1016/S0921-  
10 8181(00)00033-3, 2000.  
11 Braconnot, P., Otto-Bliesner, B., Harrison, S., Joussaume, S., Peterchmitt, J.-Y., Abe-Ouchi,  
12 A., Crucifix, M., Driesschaert, E., Fichefet, T., Hewitt, C. D., Kageyama, M., Kitoh, A.,  
13 Laîné, A., Loutre, M.-F., Marti, O., Merkel, U., Ramstein, G., Valdes, P., Weber, S. L., Yu,  
14 Y. and Zhao, Y.: Results of PMIP2 coupled simulations of the Mid-Holocene and Last  
15 Glacial Maximum – Part 1: experiments and large-scale features, *Clim Past*, 3(2), 261–277,  
16 doi:10.5194/cp-3-261-2007, 2007.  
17 Braconnot, P., Harrison, S. P., Kageyama, M., Bartlein, P. J., Masson-Delmotte, V., Abe-  
18 Ouchi, A., Otto-Bliesner, B. and Zhao, Y.: Evaluation of climate models using palaeoclimatic  
19 data, *Nat. Clim. Change*, 2(6), 417–424, doi:10.1038/nclimate1456, 2012.  
20 Briggs, R. D., Pollard, D. and Tarasov, L.: A data-constrained large ensemble analysis of  
21 Antarctic evolution since the Eemian, *Quat. Sci. Rev.*, 103, 91–115,  
22 doi:10.1016/j.quascirev.2014.09.003, 2014.  
23 Broecker, W. S.: Does the Trigger for Abrupt Climate Change Reside in the Ocean or in the  
24 Atmosphere?, *Science*, 300(5625), 1519–1522, doi:10.1126/science.1083797, 2003.  
25 Broecker, W. S., Kennett, J. P., Flower, B. P., Teller, J. T., Trumbore, S., Bonani, G. and  
26 Wolfli, W.: Routing of meltwater from the Laurentide Ice Sheet during the Younger Dryas  
27 cold episode, *Nature*, 341(6240), 318–321, doi:10.1038/341318a0, 1989.

- 1 Brovkin, V., Ganopolski, A., Archer, D. and Munhoven, G.: Glacial CO<sub>2</sub> cycle as a  
2 succession of key physical and biogeochemical processes, *Clim. Past.*, 8(1), 251–264,  
3 doi:10.5194/cp-8-251-2012, 2012.
- 4 Buizert, C., Gkinis, V., Severinghaus, J. P., He, F., Lecavalier, B. S., Kindler, P.,  
5 Leuenberger, M., Carlson, A. E., Vinther, B., Masson-Delmotte, V., White, J. W. C., Liu, Z.,  
6 Otto-Bliesner, B. and Brook, E. J.: Greenland temperature response to climate forcing during  
7 the last deglaciation, *Science*, 345(6201), 1177–1180, doi:10.1126/science.1254961, 2014.
- 8 Cabioch, G., Banks-Cutler, K. A., Beck, W. J., Burr, G. S., Corrège, T., Lawrence Edwards,  
9 R. and Taylor, F. W.: Continuous reef growth during the last 23 cal kyr BP in a tectonically  
10 active zone (Vanuatu, SouthWest Pacific), *Quat. Sci. Rev.*, 22(15–17), 1771–1786,  
11 doi:10.1016/S0277-3791(03)00170-7, 2003.
- 12 Caley, T., Roche, D. M. and Renssen, H.: Orbital Asian summer monsoon dynamics revealed  
13 using an isotope-enabled global climate model, *Nat. Commun.*, 5, doi:10.1038/ncomms6371,  
14 2014.
- 15 Carlson, A. E.: Geochemical constraints on the Laurentide Ice Sheet contribution to  
16 Meltwater Pulse 1A, *Quat. Sci. Rev.*, 28(17–18), 1625–1630,  
17 doi:10.1016/j.quascirev.2009.02.011, 2009.
- 18 Clark, P. U. and Mix, A. C.: Ice sheets and sea level of the Last Glacial Maximum, *Quat. Sci.  
Rev.*, 21(1–3), 1–7, doi:10.1016/S0277-3791(01)00118-4, 2002.
- 20 Clark, P. U. and Tarasov, L.: Closing the sea level budget at the Last Glacial Maximum, *Proc.  
21 Natl. Acad. Sci.*, 111(45), 15861–15862, doi:10.1073/pnas.1418970111, 2014.
- 22 Clark, P. U., Alley, R. B., Keigwin, L. D., Licciardi, J. M., Johnsen, S. J. and Wang, H.:  
23 Origin of the first global meltwater pulse following the Last Glacial Maximum,  
24 *Paleoceanography*, 11(5), 563–577, doi:10.1029/96PA01419, 1996.
- 25 Clark, P. U., Mitrovica, J. X., Milne, G. A. and Tamisiea, M. E.: Sea-Level Fingerprinting as  
26 a Direct Test for the Source of Global Meltwater Pulse IA, *Science*, 295(5564), 2438–2441,  
27 doi:10.1126/science.1068797, 2002.

1 [Clark, P. U., McCabe, A. M., Mix, A. C. and Weaver, A. J.: Rapid Rise of Sea Level 19,000](#)  
2 [Years Ago and Its Global Implications, Science, 304\(5674\), 1141–1144,](#)  
3 [doi:10.1126/science.1094449, 2004.](#)

4 [Clark, P. U., Dyke, A. S., Shakun, J. D., Carlson, A. E., Clark, J., Wohlfarth, B., Mitrovica, J.](#)  
5 [X., Hostetler, S. W. and McCabe, A. M.: The Last Glacial Maximum, Science, 325\(5941\),](#)  
6 [710–714, doi:10.1126/science.1172873, 2009.](#)

7 [Clark, P. U., Shakun, J. D., Baker, P. A., Bartlein, P. J., Brewer, S., Brook, E., Carlson, A. E.,](#)  
8 [Cheng, H., Kaufman, D. S., Liu, Z., Marchitto, T. M., Mix, A. C., Morrill, C., Otto-Btiesner,](#)  
9 [B. L., Pahnke, K., Russell, J. M., Whitlock, C., Adkins, J. F., Blois, J. L., Clark, J., Colman,](#)  
10 [S. M., Curry, W. B., Flower, B. P., He, F., Johnson, T. C., Lynch-Stieglitz, J., Markgraf, V.,](#)  
11 [McManus, J., Mitrovica, J. X., Moreno, P. I. and Williams, J. W.: Global climate evolution](#)  
12 [during the last deglaciation, Proc. Natl. Acad. Sci., 109\(19\), E1134–E1142,](#)  
13 [doi:10.1073/pnas.1116619109, 2012.](#)

14 [Condron, A. and Winsor, P.: Meltwater routing and the Younger Dryas, Proc. Natl. Acad.](#)  
15 [Sci., 109\(49\), 19928–19933, doi:10.1073/pnas.1207381109, 2012.](#)

16 [Cuffey, K. M. and Clow, G. D.: Temperature, accumulation, and ice sheet elevation in central](#)  
17 [Greenland through the last deglacial transition, J. Geophys. Res., 102\(C12\), 26383–26,396,](#)  
18 [doi:10.1029/96JC03981, 1997.](#)

19 [Cutler, K. B., Edwards, R. L., Taylor, F. W., Cheng, H., Adkins, J., Gallup, C. D., Cutler, P.](#)  
20 [M., Burr, G. S. and Bloom, A. L.: Rapid sea-level fall and deep-ocean temperature change](#)  
21 [since the last interglacial period, Earth Planet. Sci. Lett., 206\(3–4\), 253–271,](#)  
22 [doi:10.1016/S0012-821X\(02\)01107-X, 2003.](#)

23 [De Deckker, P. and Yokoyama, Y.: Micropalaeontological evidence for Late Quaternary sea-](#)  
24 [level changes in Bonaparte Gulf, Australia, Glob. Planet. Change, 66\(1–2\), 85–92,](#)  
25 [doi:10.1016/j.gloplacha.2008.03.012, 2009.](#)

26 [deMenocal, P., Ortiz, J., Guilderson, T., Adkins, J., Sarnthein, M., Baker, L. and Yarusinsky,](#)  
27 [M.: Abrupt onset and termination of the African Humid Period:: rapid climate responses to](#)  
28 [gradual insolation forcing, Quat. Sci. Rev., 19\(1–5\), 347–361, doi:10.1016/S0277-](#)  
29 [3791\(99\)00081-5, 2000.](#)

1 Deschamps, P., Durand, N., Bard, E., Hamelin, B., Camoin, G., Thomas, A. L., Henderson,  
2 G. M., Okuno, J. 'ichi and Yokoyama, Y.: Ice-sheet collapse and sea-level rise at the Bolling  
3 warming 14,600 years ago, *Nature*, 483(7391), 559–564, doi:10.1038/nature10902, 2012.

4 Dyke, A. S.: An outline of North American deglaciation with emphasis on central and  
5 northern Canada, in *Quaternary Glaciations-Extent and Chronology - Part II: North America*,  
6 vol. Volume 2, Part 2, pp. 373–424, Elsevier. [online] Available from:  
7 [https://www.lakeheadu.ca/sites/default/files/uploads/53/outlines/2014-15/NECU5311/Dyke\\_2004\\_DeglaciationOutline.pdf](https://www.lakeheadu.ca/sites/default/files/uploads/53/outlines/2014-15/NECU5311/Dyke_2004_DeglaciationOutline.pdf) (Accessed 20 October 2015), 2004.

8 Dyke, A. S., Andrews, J. T., Clark, P. U., England, J. H., Miller, G. H., Shaw, J. and Veillette,  
9 J. J.: The Laurentide and Innuitian ice sheets during the Last Glacial Maximum, *Quat. Sci.  
Rev.*, 21(1–3), 9–31, doi:10.1016/S0277-3791(01)00095-6, 2002.

10 Edwards, R. L., Beck, J. W., Burr, G. S., Donahue, D. J., Chappell, J. M. A., Bloom, A. L.,  
11 Druffel, E. R. M. and Taylor, F. W.: A Large Drop in Atmospheric  $^{14}\text{C}/^{12}\text{C}$  and Reduced  
12 Melting in the Younger Dryas, Documented with  $^{230}\text{Th}$  Ages of Corals, *Science*, 260(5110),  
13 962–968, doi:10.1126/science.260.5110.962, 1993.

14 EPICA Community Members: Eight glacial cycles from an Antarctic ice core, *Nature*,  
15 429(6992), 623–628, doi:10.1038/nature02599, 2004.

16 Fairbanks, R. G.: A 17,000-year glacio-eustatic sea level record: influence of glacial melting  
17 rates on the Younger Dryas event and deep-ocean circulation, *Nature*, 342(6250), 637–642,  
18 doi:10.1038/342637a0, 1989.

19 Flato, G., Marotzke, J., Abiodun, B., Braconnot, P., Chou, S. C., Collins, W., Cox, P.,  
20 Driouech, F., Emori, S., Eyring, V., Forest, C., Glecker, P., Guilyardi, E., Jackob, C., Kattsov,  
21 V., Reason, C. and Rummukainen, M.: Evaluation of Climate Models, in *Climate Change  
2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment  
Report of the Intergovernmental Panel on Climate Change*, edited by T. F. Stocker, D. Qin, G.  
22 K. Plattner, M. Tignor, S. K. Allen, J. Boschung, A. Nauels, Y. Xia, V. Bex, and P. M.  
23 Midgley, pp. 741–866, Cambridge University Press, Cambridge, United Kingdom and New  
24 York, NY, USA. [online] Available from: [http://www.ipcc.ch/pdf/assessment-report/ar5/wg1/WG1AR5\\_Chapter09\\_FINAL.pdf](http://www.ipcc.ch/pdf/assessment-report/ar5/wg1/WG1AR5_Chapter09_FINAL.pdf) (Accessed 20 October 2015), 2013.

- 1 [Ganopolski, A. and Calov, R.: The role of orbital forcing, carbon dioxide and regolith in 100](#)  
2 [kyr glacial cycles, Clim Past, 7\(4\), 1415–1425, doi:10.5194/cp-7-1415-2011, 2011.](#)
- 3 [Ganopolski, A. and Rahmstorf, S.: Rapid changes of glacial climate simulated in a coupled](#)  
4 [climate model, Nature, 409\(6817\), 153–158, doi:10.1038/35051500, 2001.](#)
- 5 [Ganopolski, A., Calov, R. and Claussen, M.: Simulation of the last glacial cycle with a](#)  
6 [coupled climate ice-sheet model of intermediate complexity, Clim Past, 6\(2\), 229–244,](#)  
7 [doi:10.5194/cp-6-229-2010, 2010.](#)
- 8 [García, J. L., Kaplan, M. R., Hall, B. L., Schaefer, J. M., Vega, R. M., Schwartz, R. and](#)  
9 [Finkel, R.: Glacier expansion in southern Patagonia throughout the Antarctic cold reversal,](#)  
10 [Geology, 40\(9\), 859–862, doi:10.1130/G33164.1, 2012.](#)
- 11 [Gherardi, J.-M., Labeyrie, L., McManus, J. F., Francois, R., Skinner, L. C. and Cortijo, E.: Evidence from the Northeastern Atlantic basin for variability in the rate of the meridional overturning circulation through the last deglaciation, Earth Planet. Sci. Lett., 240\(3–4\), 710–723, doi:10.1016/j.epsl.2005.09.061, 2005.](#)
- 15 [Golledge, N. R., Fogwill, C. J., Mackintosh, A. N. and Buckley, K. M.: Dynamics of the last glacial maximum Antarctic ice-sheet and its response to ocean forcing, Proc. Natl. Acad. Sci., 109\(40\), 16052–16056, doi:10.1073/pnas.1205385109, 2012.](#)
- 18 [Golledge, N. R., Levy, R. H., McKay, R. M., Fogwill, C. J., White, D. A., Graham, A. G. C., Smith, J. A., Hillenbrand, C.-D., Licht, K. J., Denton, G. H., Ackert Jr., R. P., Maas, S. M. and Hall, B. L.: Glaciology and geological signature of the Last Glacial Maximum Antarctic ice sheet, Quat. Sci. Rev., 78, 225–247, doi:10.1016/j.quascirev.2013.08.011, 2013.](#)
- 22 [Golledge, N. R., Menziel, L., Carter, L., Fogwill, C. J., England, M. H., Cortese, G. and Levy, R. H.: Antarctic contribution to meltwater pulse 1A from reduced Southern Ocean overturning, Nat. Commun., 5, doi:10.1038/ncomms6107, 2014.](#)
- 25 [Goujon, C., Barnola, J.-M. and Ritz, C.: Modeling the densification of polar firn including heat diffusion: Application to close-off characteristics and gas isotopic fractionation for Antarctica and Greenland sites, J. Geophys. Res. Atmospheres, 108\(D24\), 4792, doi:10.1029/2002JD003319, 2003.](#)

- 1 Gregoire, L. J., Payne, A. J. and Valdes, P. J.: Deglacial rapid sea level rises caused by ice-  
2 sheet saddle collapses, *Nature*, 487(7406), 219–222, doi:10.1038/nature11257, 2012.
- 3 Gregoire, L. J., Valdes, P. J. and Payne, A. J.: The relative contribution of orbital forcing and  
4 greenhouse gases to the North American deglaciation, *Geophys. Res. Lett.*, 42(22),  
5 2015GL066005, doi:10.1002/2015GL066005, 2015.
- 6 Gyllencreutz, R., Mangerud, J., Svendsen, J.-I. and Lohne, Ø.: DATED—a GIS-based  
7 reconstruction and dating database of the Eurasian deglaciation, *Appl. Quat. Res. Cent. Part*  
8 *Glaciat. Terrain Geol. Surv. Finl. Spec. Pap.*, 46, 113–120, 2007.
- 9 Hall, I. R., Moran, S. B., Zahn, R., Knutz, P. C., Shen, C.-C. and Edwards, R. L.: Accelerated  
10 drawdown of meridional overturning in the late-glacial Atlantic triggered by transient pre-H  
11 event freshwater perturbation, *Geophys. Res. Lett.*, 33(16), L16616,  
12 doi:10.1029/2006GL026239, 2006.
- 13 Hanebuth, T., Stattegger, K. and Grootes, P. M.: Rapid Flooding of the Sunda Shelf: A Late-  
14 Glacial Sea-Level Record, *Science*, 288(5468), 1033–1035,  
15 doi:10.1126/science.288.5468.1033, 2000.
- 16 Hanebuth, T. J. J., Stattegger, K. and Bojanowski, A.: Termination of the Last Glacial  
17 Maximum sea-level lowstand: The Sunda-Shelf data revisited, *Glob. Planet. Change*, 66(1–2),  
18 76–84, doi:10.1016/j.gloplacha.2008.03.011, 2009.
- 19 Haywood, A. M., Dowsett, H. J., Otto-Bliesner, B., Chandler, M. A., Dolan, A. M., Hill, D.  
20 J., Lunt, D. J., Robinson, M. M., Rosenbloom, N., Salzmann, U. and Sohl, L. E.: Pliocene  
21 Model Intercomparison Project (PlioMIP): experimental design and boundary conditions  
22 (Experiment 1), *Geosci. Model Dev.*, 3(1), 227–242, doi:10.5194/gmd-3-227-2010, 2010.
- 23 Heinemann, M., Timmermann, A., Elison Timm, O., Saito, F. and Abe-Ouchi, A.: Deglacial  
24 ice sheet meltdown: orbital pacemaking and CO<sub>2</sub> effects, *Clim Past*, 10(4), 1567–1579,  
25 doi:10.5194/cp-10-1567-2014, 2014.
- 26 Heiri, O., Cremer, H., Engels, S., Hoek, W. Z., Peeters, W. and Lotter, A. F.: Lateglacial  
27 summer temperatures in the Northwest European lowlands: a chironomid record from  
28 Hijkemeer, the Netherlands, *Quat. Sci. Rev.*, 26(19–21), 2420–2437,  
29 doi:10.1016/j.quascirev.2007.06.017, 2007.

1 [Hemming, S. R.: Heinrich events: Massive late Pleistocene detritus layers of the North](#)  
2 [Atlantic and their global climate imprint, Rev. Geophys., 42\(1\), RG1005,](#)  
3 [doi:10.1029/2003RG000128, 2004.](#)

4 [Hughes, A. L. C., Gyllencreutz, R., Lohne, Ø. S., Mangerud, J. and Svendsen, J. I.: The last](#)  
5 [Eurasian ice sheets – a chronological database and time-slice reconstruction, DATED-1,](#)  
6 [Boreas, doi:10.1111/bor.12142, 2015.](#)

7 [Joussaume, S., Taylor, K. E., Braconnot, P., Mitchell, J. F. B., Kutzbach, J. E., Harrison, S.](#)  
8 [P., Prentice, I. C., Broccoli, A. J., Abe-Ouchi, A., Bartlein, P. J., Bonfils, C., Dong, B., Guiot,](#)  
9 [J., Herterich, K., Hewitt, C. D., Jolly, D., Kim, J. W., Kislov, A., Kitoh, A., Loutre, M. F.,](#)  
10 [Masson, V., McAvaney, B., McFarlane, N., de Noblet, N., Peltier, W. R., Peterschmitt, J. Y.,](#)  
11 [Pollard, D., Rind, D., Royer, J. F., Schlesinger, M. E., Syktus, J., Thompson, S., Valdes, P.,](#)  
12 [Vettoretti, G., Webb, R. S. and Wyputta, U.: Monsoon changes for 6000 years ago: Results of](#)  
13 [18 simulations from the Paleoclimate Modeling Intercomparison Project \(PMIP\), Geophys.](#)  
14 [Res. Lett., 26\(7\), 859–862, doi:10.1029/1999GL900126, 1999.](#)

15 [Jouzel, J., Vaikmae, R., Petit, J. R., Martin, M., Duclos, Y., Stievenard, M., Lorius, C., Toots,](#)  
16 [Mélières, M. A., Burckle, L. H., Barkov, N. I. and Kotlyakov, V. M.: The two-step shape](#)  
17 [and timing of the last deglaciation in Antarctica, Clim. Dyn., 11\(3\), 151–161,](#)  
18 [doi:10.1007/BF00223498, 1995.](#)

19 [Jouzel, J., Masson-Delmotte, V., Cattani, O., Dreyfus, G., Falourd, S., Hoffmann, G., Minster,](#)  
20 [B., Nouet, J., Barnola, J. M., Chappellaz, J., Fischer, H., Gallet, J. C., Johnsen, S.,](#)  
21 [Leuenberger, M., Louergue, L., Luethi, D., Oerter, H., Parrenin, F., Raisbeck, G., Raynaud,](#)  
22 [D., Schilt, A., Schwander, J., Selmo, E., Souchez, R., Spahni, R., Stauffer, B., Steffensen, J.](#)  
23 [P., Stenni, B., Stocker, T. F., Tison, J. L., Werner, M. and Wolff, E. W.: Orbital and](#)  
24 [Millennial Antarctic Climate Variability over the Past 800,000 Years, Science, 317\(5839\),](#)  
25 [793–796, doi:10.1126/science.1141038, 2007.](#)

26 [Kageyama, M., Laîné, A., Abe-Ouchi, A., Braconnot, P., Cortijo, E., Crucifix, M., de Vernal,](#)  
27 [A., Guiot, J., Hewitt, C. D., Kitoh, A., Kucera, M., Martí, O., Ohgaito, R., Otto-Bliesner, B.,](#)  
28 [Peltier, W. R., Rosell-Melé, A., Vettoretti, G., Weber, S. L. and Yu, Y.: Last Glacial](#)  
29 [Maximum temperatures over the North Atlantic, Europe and western Siberia: a comparison](#)

- 1 between PMIP models, MARGO sea–surface temperatures and pollen-based reconstructions,  
2 *Quat. Sci. Rev.*, 25(17–18), 2082–2102, doi:10.1016/j.quascirev.2006.02.010, 2006.
- 3 Kageyama, M., Paul, A., Roche, D. M. and Van Meerbeeck, C. J.: Modelling glacial climatic  
4 millennial-scale variability related to changes in the Atlantic meridional overturning  
5 circulation: a review, *Quat. Sci. Rev.*, 29(21–22), 2931–2956,  
6 doi:10.1016/j.quascirev.2010.05.029, 2010.
- 7 Kageyama, M., Merkel, U., Otto-Bliesner, B., Prange, M., Abe-Ouchi, A., Lohmann, G.,  
8 Ohgaito, R., Roche, D. M., Singarayer, J., Swingedouw, D. and X Zhang: Climatic impacts of  
9 fresh water hosing under Last Glacial Maximum conditions: a multi-model study, *Clim Past*,  
10 9(2), 935–953, doi:10.5194/cp-9-935-2013, 2013.
- 11 Kageyama, M., Braconnot, P., Harrison, S. P., Haywood, A. M., Jungclaus, J., Otto-Bliesner,  
12 B. L., Peterschmitt, J.-Y., Abe-Ouchi, A., Albani, S., Bartlein, P. J., Brierley, C., Crucifix, M.,  
13 Dolan, A., Fernandez-Donado, L., Fischer, H., Hopcroft, P. O., Ivanovic, R. F., Lambert, F.,  
14 Lunt, D. J., Mahowald, N. M., Peltier, W. R., Phipps, S. J., Roche, D. M., Schmidt, G. A.,  
15 Tarasov, L., Valdes, P. J., Zhang, Q. and Zhou, T.: PMIP4-CMIP6: the contribution of the  
16 Paleoclimate Modelling Intercomparison Project to CMIP6, *Geosci. Model Dev. Discuss.*, 1–  
17 46, doi:10.5194/gmd-2016-106, 2016.
- 18 Kaplan, M. R., Strelin, J. A., Schaefer, J. M., Denton, G. H., Finkel, R. C., Schwartz, R.,  
19 Putnam, A. E., Vandergoes, M. J., Goehring, B. M. and Travis, S. G.: In-situ cosmogenic  
20  $^{10}\text{Be}$  production rate at Lago Argentino, Patagonia: Implications for late-glacial climate  
21 chronology, *Earth Planet. Sci. Lett.*, 309(1–2), 21–32, doi:10.1016/j.epsl.2011.06.018, 2011.
- 22 Keigwin, L. D. and Boyle, E. A.: Did North Atlantic overturning halt 17,000 years ago?,  
23 *Paleoceanography*, 23(1), PA1101, doi:10.1029/2007PA001500, 2008.
- 24 Keigwin, L. D., Jones, G. A., Lehman, S. J. and Boyle, E. A.: Deglacial meltwater discharge,  
25 North Atlantic Deep Circulation, and abrupt climate change, *J. Geophys. Res. Oceans*,  
26 96(C9), 16811–16826, doi:10.1029/91JC01624, 1991.
- 27 Kindler, P., Guillevic, M., Baumgartner, M., Schwander, J., Landais, A. and Leuenberger, M.:  
28 Temperature reconstruction from 10 to 120 kyr b2k from the NGRIP ice core, *Clim Past*,  
29 10(2), 887–902, doi:10.5194/cp-10-887-2014, 2014.

- 1 Knorr, G. and Lohmann, G.: Southern Ocean origin for the resumption of Atlantic  
2 thermohaline circulation during deglaciation, *Nature*, 424(6948), 532–536,  
3 doi:10.1038/nature01855, 2003.
- 4 Knorr, G. and Lohmann, G.: Rapid transitions in the Atlantic thermohaline circulation  
5 triggered by global warming and meltwater during the last deglaciation, *Geochem. Geophys.*  
6 *Geosystems*, 8(12), Q12006, doi:10.1029/2007GC001604, 2007.
- 7 Kohfeld, K. . and Harrison, S. : How well can we simulate past climates? Evaluating the  
8 models using global palaeoenvironmental datasets, *Quat. Sci. Rev.*, 19(1-5), 321–346,  
9 doi:10.1016/S0277-3791(99)00068-2, 2000.
- 10 Lambeck, K., Rouby, H., Purcell, A., Sun, Y. and Sambridge, M.: Sea level and global ice  
11 volumes from the Last Glacial Maximum to the Holocene, *Proc. Natl. Acad. Sci.*,  
12 doi:10.1073/pnas.1411762111, 2014.
- 13 Lea, D. W., Pak, D. K., Peterson, L. C. and Hughen, K. A.: Synchronicity of Tropical and  
14 High-Latitude Atlantic Temperatures over the Last Glacial Termination, *Science*, 301(5638),  
15 1361–1364, doi:10.1126/science.1088470, 2003.
- 16 Licht, K. J.: The Ross Sea's contribution to eustatic sea level during meltwater pulse 1A,  
17 *Sediment. Geol.*, 165(3–4), 343–353, doi:10.1016/j.sedgeo.2003.11.020, 2004.
- 18 Liu, Z., Otto-Bliesner, B. L., He, F., Brady, E. C., Tomas, R., Clark, P. U., Carlson, A. E.,  
19 Lynch-Stieglitz, J., Curry, W., Brook, E., Erickson, D., Jacob, R., Kutzbach, J. and Cheng, J.:  
20 Transient Simulation of Last Deglaciation with a New Mechanism for Bølling-Allerød  
21 Warming, *Science*, 325(5938), 310–314, doi:10.1126/science.1171041, 2009.
- 22 Liu, Z., Carlson, A. E., He, F., Brady, E. C., Otto-Bliesner, B. L., Briegleb, B. P.,  
23 Wehrenberg, M., Clark, P. U., Wu, S., Cheng, J., Zhang, J., Noone, D. and Zhu, J.: Younger  
24 Dryas cooling and the Greenland climate response to CO<sub>2</sub>, *Proc. Natl. Acad. Sci.*,  
25 doi:10.1073/pnas.1202183109, 2012.
- 26 Liu, Z., Wen, X., Brady, E. C., Otto-Bliesner, B., Yu, G., Lu, H., Cheng, H., Wang, Y.,  
27 Zheng, W., Ding, Y., Edwards, R. L., Cheng, J., Liu, W. and Yang, H.: Chinese cave records  
28 and the East Asia Summer Monsoon, *Quat. Sci. Rev.*, 83, 115–128,  
29 doi:10.1016/j.quascirev.2013.10.021, 2014.

1 Louergue, L., Schilt, A., Spahni, R., Masson-Delmotte, V., Blunier, T., Lemieux, B.,  
2 Barnola, J.-M., Raynaud, D., Stocker, T. F. and Chappellaz, J.: Orbital and millennial-scale  
3 features of atmospheric CH<sub>4</sub> over the past 800,000 years, *Nature*, 453(7193), 383–386,  
4 doi:10.1038/nature06950, 2008.

5 Lüthi, D., Le Floch, M., Bereiter, B., Blunier, T., Barnola, J.-M., Siegenthaler, U., Raynaud,  
6 D., Jouzel, J., Fischer, H., Kawamura, K. and Stocker, T. F.: High-resolution carbon dioxide  
7 concentration record 650,000–800,000 years before present, *Nature*, 453(7193), 379–382,  
8 doi:10.1038/nature06949, 2008.

9 MacFarling Meure, C., Etheridge, D., Trudinger, C., Steele, P., Langenfelds, R., van Ommen,  
10 T., Smith, A. and Elkins, J.: Law Dome CO<sub>2</sub>, CH<sub>4</sub> and N<sub>2</sub>O ice core records extended to  
11 2000 years BP, *Geophys. Res. Lett.*, 33(14), L14810, doi:10.1029/2006GL026152, 2006.

12 Mackintosh, A., Golledge, N., Domack, E., Dunbar, R., Leventer, A., White, D., Pollard, D.,  
13 DeConto, R., Fink, D., Zwart, D., Gore, D. and Lavoie, C.: Retreat of the East Antarctic ice  
14 sheet during the last glacial termination, *Nat. Geosci.*, 4(3), 195–202, doi:10.1038/ngeo1061,  
15 2011.

16 Mackintosh, A. N., Verleyen, E., O'Brien, P. E., White, D. A., Jones, R. S., McKay, R.,  
17 Dunbar, R., Gore, D. B., Fink, D., Post, A. L., Miura, H., Leventer, A., Goodwin, I., Hodgson,  
18 D. A., Lilly, K., Crosta, X., Golledge, N. R., Wagner, B., Berg, S., van Ommen, T., Zwart,  
19 D., Roberts, S. J., Vyverman, W. and Masse, G.: Retreat history of the East Antarctic Ice  
20 Sheet since the Last Glacial Maximum, *Quat. Sci. Rev.*, 100, 10–30,  
21 doi:10.1016/j.quascirev.2013.07.024, 2014.

22 Mamajek, E. E., Prsa, A., Torres, G., Harmanec, P., Asplund, M., Bennett, P. D., Capitaine,  
23 N., Christensen-Dalsgaard, J., Depagne, E., Folkner, W. M., Haberreiter, M., Hekker, S.,  
24 Hilton, J. L., Kostov, V., Kurtz, D. W., Laskar, J., Mason, B. D., Milone, E. F., Montgomery,  
25 M. M., Richards, M. T., Schou, J. and Stewart, S. G.: IAU 2015 Resolution B3 on  
26 Recommended Nominal Conversion Constants for Selected Solar and Planetary Properties,  
27 ArXiv151007674 Astro-Ph [online] Available from: <http://arxiv.org/abs/1510.07674>  
28 (Accessed 19 May 2016), 2015.

29 Marcott, S. A., Bauska, T. K., Buzert, C., Steig, E. J., Rosen, J. L., Cuffey, K. M., Fudge, T.  
30 J., Severinghaus, J. P., Ahn, J., Kalk, M. L., McConnell, J. R., Sowers, T., Taylor, K. C.,

- 1 [White, J. W. C. and Brook, E. J.: Centennial-scale changes in the global carbon cycle during](#)  
2 [the last deglaciation, Nature, 514\(7524\), 616–619, doi:10.1038/nature13799, 2014.](#)
- 3 [Marshall, S. J. and Clarke, G. K. C.: Modeling North American Freshwater Runoff through](#)  
4 [the Last Glacial Cycle, Quat. Res., 52\(3\), 300–315, doi:10.1006/qres.1999.2079, 1999.](#)
- 5 [Martrat, B., Grimalt, J. O., Lopez-Martinez, C., Cacho, I., Sierro, F. J., Flores, J. A., Zahn, R.,](#)  
6 [Canals, M., Curtis, J. H. and Hodell, D. A.: Abrupt Temperature Changes in the Western](#)  
7 [Mediterranean over the Past 250,000 Years, Science, 306\(5702\), 1762–1765,](#)  
8 [doi:10.1126/science.1101706, 2004.](#)
- 9 [Martrat, B., Grimalt, J. O., Shackleton, N. J., Abreu, L. de, Hutterli, M. A. and Stocker, T. F.:](#)  
10 [Four Climate Cycles of Recurring Deep and Surface Water Destabilizations on the Iberian](#)  
11 [Margin, Science, 317\(5837\), 502–507, doi:10.1126/science.113994, 2007.](#)
- 12 [Masson-Delmotte, V., Kageyama, M., Braconnot, P., Charbit, S., Krinner, G., Ritz, C.,](#)  
13 [Guilyardi, E., Jouzel, J., Abe-Ouchi, A., Crucifix, M., Gladstone, R. M., Hewitt, C. D., Kitoh,](#)  
14 [A., LeGrande, A. N., Marti, O., Merkel, U., Motoi, T., Ohgaito, R., Otto-Bliesner, B., Peltier,](#)  
15 [W. R., Ross, I., Valdes, P. J., Vettoretti, G., Weber, S. L., Wolk, F. and Yu, Y.: Past and](#)  
16 [future polar amplification of climate change: climate model intercomparisons and ice-core](#)  
17 [constraints, Clim. Dyn., 26\(5\), 513–529, doi:10.1007/s00382-005-0081-9, 2006.](#)
- 18 [Masson-Delmotte, V., Schulz, M., Abe-Ouchi, A., Beer, J., Ganopolsk, A., González Rouco,](#)  
19 [J. F., Jansen, E., Lambeck, K., Luterbacher, J., Naish, T., Osborn, T., Otto-Bliesner, B.,](#)  
20 [Quinn, T., Ramesh, R., Rojas, M., Shao, X. and Timmermann, A.: Information from](#)  
21 [Paleoclimate Archives, in Climate Change 2013: The Physical Science Basis. Contribution of](#)  
22 [Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate](#)  
23 [Change, edited by T. F. Stocker, D. Qin, G. K. Plattner, M. Tignor, S. K. Allen, J. Boschung,](#)  
24 [A. Nauels, Y. Xia, V. Bex, and P. M. Midgley, pp. 383–464, Cambridge University Press,](#)  
25 [Cambridge, United Kingdom and New York, NY, USA. \[online\] Available from:](#)  
26 [http://www.ipcc.ch/pdf/assessment-report/ar5/wg1/WG1AR5\\_Chapter05\\_FINAL.pdf](http://www.ipcc.ch/pdf/assessment-report/ar5/wg1/WG1AR5_Chapter05_FINAL.pdf)  
27 [\(Accessed 20 October 2015\), 2013.](#)
- 28 [McManus, J. F., Francois, R., Gherardi, J.-M., Keigwin, L. D. and Brown-Leger, S.: Collapse](#)  
29 [and rapid resumption of Atlantic meridional circulation linked to deglacial climate changes,](#)  
30 [Nature, 428\(6985\), 834–837, doi:10.1038/nature02494, 2004.](#)

- 1 [Menviel, L., Timmermann, A., Timm, O. E. and Mouchet, A.: Climate and biogeochemical](#)  
2 [response to a rapid melting of the West Antarctic Ice Sheet during interglacials and](#)  
3 [implications for future climate, Paleoceanography, 25\(4\), PA4231,](#)  
4 [doi:10.1029/2009PA001892, 2010.](#)
- 5 [Menviel, L., Timmermann, A., Timm, O. E. and Mouchet, A.: Deconstructing the Last](#)  
6 [Glacial termination: the role of millennial and orbital-scale forcings, Quat. Sci. Rev., 30\(9–](#)  
7 [10\), 1155–1172, doi:10.1016/j.quascirev.2011.02.005, 2011.](#)
- 8 [Mix, A. C., Bard, E. and Schneider, R.: Environmental processes of the ice age: land, oceans,](#)  
9 [glaciers \(EPILOG\), Quat. Sci. Rev., 20\(4\), 627–657, doi:10.1016/S0277-3791\(00\)00145-1,](#)  
10 [2001.](#)
- 11 [Monnin, E., Steig, E. J., Siegenthaler, U., Kawamura, K., Schwander, J., Stauffer, B., Stocker,](#)  
12 [T. F., Morse, D. L., Barnola, J.-M., Bellier, B., Raynaud, D. and Fischer, H.: Evidence for](#)  
13 [substantial accumulation rate variability in Antarctica during the Holocene, through](#)  
14 [synchronization of CO<sub>2</sub> in the Taylor Dome, Dome C and DML ice cores, Earth Planet. Sci.](#)  
15 [Lett., 224\(1–2\), 45–54, doi:10.1016/j.epsl.2004.05.007, 2004.](#)
- 16 [Otto-Bliesner, B. L., Schneider, R., Brady, E. C., Kucera, M., Abe-Ouchi, A., Bard, E.,](#)  
17 [Braconnot, P., Crucifix, M., Hewitt, C. D., Kageyama, M., Marti, O., Paul, A., Rosell-Melé,](#)  
18 [A., Waelbroeck, C., Weber, S. L., Weinelt, M. and Yu, Y.: A comparison of PMIP2 model](#)  
19 [simulations and the MARGO proxy reconstruction for tropical sea surface temperatures at last](#)  
20 [glacial maximum, Clim. Dyn., 32\(6\), 799–815, doi:10.1007/s00382-008-0509-0, 2009.](#)
- 21 [Otto-Bliesner, B. L., Russell, J. M., Clark, P. U., Liu, Z., Overpeck, J. T., Konecky, B.,](#)  
22 [deMenocal, P., Nicholson, S. E., He, F. and Lu, Z.: Coherent changes of southeastern](#)  
23 [equatorial and northern African rainfall during the last deglaciation, Science, 346\(6214\),](#)  
24 [1223–1227, doi:10.1126/science.1259531, 2014.](#)
- 25 [Peltier, W. R.: Global glacial isostasy and the surface of the Ice-Age Earth: The ICE-5G](#)  
26 [\(VM2\) model and GRACE, Annu. Rev. Earth Planet. Sci., 32\(1\), 111–149,](#)  
27 [doi:10.1146/annurev.earth.32.082503.144359, 2004.](#)
- 28 [Peltier, W. R.: On the hemispheric origins of meltwater pulse 1a, Quat. Sci. Rev., 24\(14–15\),](#)  
29 [1655–1671, doi:10.1016/j.quascirev.2004.06.023, 2005.](#)

- 1 [Peltier, W. R. and Fairbanks, R. G.: Global glacial ice volume and Last Glacial Maximum](#)  
2 [duration from an extended Barbados sea level record, Quat. Sci. Rev., 25\(23–24\), 3322–3337,](#)  
3 [doi:10.1016/j.quascirev.2006.04.010, 2006.](#)
- 4 [Peltier, W. R. and Vettoretti, G.: Dansgaard-Oeschger oscillations predicted in a](#)  
5 [comprehensive model of glacial climate: A “kicked” salt oscillator in the Atlantic, Geophys.](#)  
6 [Res. Lett., 41\(20\), 2014GL061413, doi:10.1002/2014GL061413, 2014.](#)
- 7 [Peltier, W. R., Argus, D. F. and Drummond, R.: Space geodesy constrains ice age terminal](#)  
8 [deglaciation: The global ICE-6G C \(VM5a\) model, J. Geophys. Res. Solid Earth,](#)  
9 [2014JB011176, doi:10.1002/2014JB011176, 2015.](#)
- 10 [Petit, J. R., Jouzel, J., Raynaud, D., Barkov, N. I., Barnola, J.-M., Basile, I., Bender, M.,](#)  
11 [Chappellaz, J., Davis, M., Delaygue, G., Delmotte, M., Kotlyakov, V. M., Legrand, M.,](#)  
12 [Lipenkov, V. Y., Lorius, C., PÉpin, L., Ritz, C., Saltzman, E. and Stievenard, M.: Climate and](#)  
13 [atmospheric history of the past 420,000 years from the Vostok ice core, Antarctica, Nature,](#)  
14 [399\(6735\), 429–436, doi:10.1038/20859, 1999.](#)
- 15 [Philippon, G., Ramstein, G., Charbit, S., Kageyama, M., Ritz, C. and Dumas, C.: Evolution of](#)  
16 [the Antarctic ice sheet throughout the last deglaciation: A study with a new coupled climate—](#)  
17 [north and south hemisphere ice sheet model, Earth Planet. Sci. Lett., 248\(3–4\), 750–758,](#)  
18 [doi:10.1016/j.epsl.2006.06.017, 2006.](#)
- 19 [PMIP Last Deglaciation Working Group: PMIP4 Last Deglaciation experiment Design Wiki,](#)  
20 [\[online\] Available from: \[https://pmip4.lsce.ipsl.fr/doku.php/exp\\\_design:degla\]\(https://pmip4.lsce.ipsl.fr/doku.php/exp\_design:degla\) \(Accessed 2](#)  
21 [June 2016\), 2016.](#)
- 22 [PMIP LGM Working Group: PMIP3-CMIP5 Last Glacial Maximum experiment design,](#)  
23 [\[online\] Available from: <https://wiki.lsce.ipsl.fr/pmip3/doku.php/pmip3:design:21k:final> \(Accessed 9 April 2015\), 2010.](#)
- 25 [PMIP LGM Working Group: PMIP-CMIP6 Last Glacial Maximum experiment design,](#)  
26 [\[online\] Available from:](#)  
27 <https://wiki.lsce.ipsl.fr/pmip3/doku.php/pmip3:cmip6:design:21k:index> (Accessed 9 April  
28 [2015\), 2015.](#)

1 PMIP website: Paleoclimate Model Intercomparison Project, [online] Available from:  
2 <http://pmip.lsce.ipsl.fr/> (Accessed 13 November 2014), 2007.

3 Putnam, A. E., Denton, G. H., Schaefer, J. M., Barrell, D. J. A., Andersen, B. G., Finkel, R.  
4 C., Schwartz, R., Doughty, A. M., Kaplan, M. R. and Schlüchter, C.: Glacier advance in  
5 southern middle-latitudes during the Antarctic Cold Reversal, *Nat. Geosci.*, 3(10), 700–704,  
6 doi:10.1038/ngeo962, 2010.

7 Rahmstorf, S.: Bifurcations of the Atlantic thermohaline circulation in response to changes in  
8 the hydrological cycle, *Nature*, 378(6553), 145–149, doi:10.1038/378145a0, 1995.

9 Rahmstorf, S.: On the freshwater forcing and transport of the Atlantic thermohaline  
10 circulation, *Clim. Dyn.*, 12(12), 799–811, doi:10.1007/s003820050144, 1996.

11 Rasmussen, S. O., Andersen, K. K., Svensson, A. M., Steffensen, J. P., Vinther, B. M.,  
12 Clausen, H. B., Siggaard-Andersen, M.-L., Johnsen, S. J., Larsen, L. B., Dahl-Jensen, D.,  
13 Bigler, M., Röhlisberger, R., Fischer, H., Goto-Azuma, K., Hansson, M. E. and Ruth, U.: A  
14 new Greenland ice core chronology for the last glacial termination, *J. Geophys. Res.*  
15 *Atmospheres*, 111(D6), D06102, doi:10.1029/2005JD006079, 2006.

16 Roberts, N. L., Piotrowski, A. M., McManus, J. F. and Keigwin, L. D.: Synchronous  
17 Deglacial Overturning and Water Mass Source Changes, *Science*, 327(5961), 75–78,  
18 doi:10.1126/science.1178068, 2010.

19 Roche, D. M., Renssen, H., Weber, S. L. and Goosse, H.: Could meltwater pulses have been  
20 sneaked unnoticed into the deep ocean during the last glacial?, *Geophys. Res. Lett.*, 34(24),  
21 n/a–n/a, doi:10.1029/2007GL032064, 2007.

22 Roche, D. M., Renssen, H., Paillard, D. and Levavasseur, G.: Deciphering the spatio-temporal  
23 complexity of climate change of the last deglaciation: a model analysis, *Clim Past*, 7(2), 591–  
24 602, doi:10.5194/cp-7-591-2011, 2011.

25 Rogerson, M., Colmenero-Hidalgo, E., Levine, R. C., Rohling, E. J., Voelker, A. H. L., Bigg,  
26 G. R., Schönfeld, J., Cacho, I., Sierro, F. J., Löwemark, L., Reguera, M. I., Abreu, L. de and  
27 Garrick, K.: Enhanced Mediterranean-Atlantic exchange during Atlantic freshening phases,  
28 *Geochem. Geophys. Geosystems*, 11, 22 PP., doi:201010.1029/2009GC002931, 2010.

- 1 [Rother, H., Fink, D., Shulmeister, J., Mifsud, C., Evans, M. and Pugh, J.: The early rise and](#)  
2 [late demise of New Zealand's last glacial maximum. Proc. Natl. Acad. Sci., 111\(32\), 11630–](#)  
3 [11635, doi:10.1073/pnas.1401547111, 2014.](#)
- 4 [Rubino, M., Etheridge, D. M., Trudinger, C. M., Allison, C. E., Battle, M. O., Langenfelds, R.](#)  
5 [L., Steele, L. P., Curran, M., Bender, M., White, J. W. C., Jenk, T. M., Blunier, T. and](#)  
6 [Francey, R. J.: A revised 1000 year atmospheric  \$\delta^{13}\text{C}\$ -CO<sub>2</sub> record from Law Dome and](#)  
7 [South Pole, Antarctica. J. Geophys. Res. Atmospheres, 118\(15\), 8482–8499,](#)  
8 [doi:10.1002/jgrd.50668, 2013.](#)
- 9 [Schilt, A., Baumgartner, M., Schwander, J., Buiron, D., Capron, E., Chappellaz, J.,](#)  
10 [Loulergue, L., Schüpbach, S., Spahni, R., Fischer, H. and Stocker, T. F.: Atmospheric nitrous](#)  
11 [oxide during the last 140,000 years. Earth Planet. Sci. Lett., 300\(1–2\), 33–43,](#)  
12 [doi:10.1016/j.epsl.2010.09.027, 2010.](#)
- 13 [Schmidt, G. A., Jungclaus, J. H., Ammann, C. M., Bard, E., Braconnot, P., Crowley, T. J.,](#)  
14 [Delaygue, G., Joos, F., Krivova, N. A., Muscheler, R., Otto-Bliesner, B. L., Pongratz, J.,](#)  
15 [Shindell, D. T., Solanki, S. K., Steinhilber, F. and Vieira, L. E. A.: Climate forcing](#)  
16 [reconstructions for use in PMIP simulations of the last millennium \(v1.0\). Geosci Model Dev.](#)  
17 [4\(1\), 33–45, doi:10.5194/gmd-4-33-2011, 2011.](#)
- 18 [Schmittner, A., Green, J. a. M. and Wilmes, S.-B.: Glacial ocean overturning intensified by](#)  
19 [tidal mixing in a global circulation model. Geophys. Res. Lett., 42\(10\), 2015GL063561,](#)  
20 [doi:10.1002/2015GL063561, 2015.](#)
- 21 [Severinghaus, J. P. and Brook, E. J.: Abrupt Climate Change at the End of the Last Glacial](#)  
22 [Period Inferred from Trapped Air in Polar Ice. Science, 286\(5441\), 930–934,](#)  
23 [doi:10.1126/science.286.5441.930, 1999.](#)
- 24 [Shakun, J. D., Clark, P. U., He, F., Marcott, S. A., Mix, A. C., Liu, Z., Otto-Bliesner, B.,](#)  
25 [Schmittner, A. and Bard, E.: Global warming preceded by increasing carbon dioxide](#)  
26 [concentrations during the last deglaciation. Nature, 484\(7392\), 49–54,](#)  
27 [doi:10.1038/nature10915, 2012.](#)
- 28 [Shennan, I.: Global meltwater discharge and the deglacial sea-level record from northwest](#)  
29 [Scotland. J. Quat. Sci., 14\(7\), 715–719, doi:10.1002/\(SICI\)1099-](#)  
30 [1417\(199912\)14:7<715::AID-JQS511>3.0.CO;2-G, 1999.](#)

- 1 [Shennan, I. and Milne, G.: Sea-level observations around the Last Glacial Maximum from the](#)  
2 [Bonaparte Gulf, NW Australia, Quat. Sci. Rev., 22\(14\), 1543–1547, doi:10.1016/S0277-](#)  
3 [3791\(03\)00088-X, 2003.](#)
- 4 [Siegenthaler, U., Stocker, T. F., Monnin, E., Lüthi, D., Schwander, J., Stauffer, B., Raynaud,](#)  
5 [D., Barnola, J.-M., Fischer, H., Masson-Delmotte, V. and Jouzel, J.: Stable Carbon](#)  
6 [Cycle&#150;Climate Relationship During the Late Pleistocene, Science, 310\(5752\), 1313–](#)  
7 [1317, doi:10.1126/science.1120130, 2005.](#)
- 8 [Simonsen, S. B., Johnsen, S. J., Popp, T. J., Vinther, B. M., Gkinis, V. and Steen-Larsen, H.](#)  
9 [C.: Past surface temperatures at the NorthGRIP drill site from the difference in firn diffusion](#)  
10 [of water isotopes, Clim Past, 7\(4\), 1327–1335, doi:10.5194/cp-7-1327-2011, 2011.](#)
- 11 [Smith, R. S. and Gregory, J. M.: A study of the sensitivity of ocean overturning circulation](#)  
12 [and climate to freshwater input in different regions of the North Atlantic, Geophys. Res. Lett.,](#)  
13 [36\(15\), L15701, doi:10.1029/2009GL038607, 2009.](#)
- 14 [Spahni, R., Chappellaz, J., Stocker, T. F., Loulergue, L., Hausmann, G., Kawamura, K.,](#)  
15 [Flückiger, J., Schwander, J., Raynaud, D., Masson-Delmotte, V. and Jouzel, J.: Atmospheric](#)  
16 [Methane and Nitrous Oxide of the Late Pleistocene from Antarctic Ice Cores, Science,](#)  
17 [310\(5752\), 1317–1321, doi:10.1126/science.1120132, 2005.](#)
- 18 [Stanford, J. D., Hemingway, R., Rohling, E. J., Challenor, P. G., Medina-Elizalde, M. and](#)  
19 [Lester, A. J.: Sea-level probability for the last deglaciation: A statistical analysis of far-field](#)  
20 [records, Glob. Planet. Change, 79\(3–4\), 193–203, doi:10.1016/j.gloplacha.2010.11.002, 2011.](#)
- 21 [Steffensen, J. P., Andersen, K. K., Bigler, M., Clausen, H. B., Dahl-Jensen, D., Fischer, H.,](#)  
22 [Goto-Azuma, K., Hansson, M., Johnsen, S. J., Jouzel, J., Masson-Delmotte, V., Popp, T.,](#)  
23 [Rasmussen, S. O., Röhlisberger, R., Ruth, U., Stauffer, B., Siggaard-Andersen, M.-L.,](#)  
24 [Sveinbjörnsdóttir, Á. E., Svensson, A. and White, J. W. C.: High-Resolution Greenland Ice](#)  
25 [Core Data Show Abrupt Climate Change Happens in Few Years, Science, 321\(5889\), 680–](#)  
26 [684, doi:10.1126/science.1157707, 2008.](#)
- 27 [Stenni, B., Buiron, D., Frezzotti, M., Albani, S., Barbante, C., Bard, E., Barnola, J. M.,](#)  
28 [Baroni, M., Baumgartner, M., Bonazza, M., Capron, E., Castellano, E., Chappellaz, J.,](#)  
29 [Delmonte, B., Falourd, S., Genoni, L., Iacumin, P., Jouzel, J., Kipfstuhl, S., Landais, A.,](#)  
30 [Lemieux-Dudon, B., Maggi, V., Masson-Delmotte, V., Mazzola, C., Minster, B., Montagnat,](#)

- 1 [M., Mulvaney, R., Narcisi, B., Oerter, H., Parrenin, F., Petit, J. R., Ritz, C., Scarchilli, C.,](#)  
2 [Schilt, A., Schüpbach, S., Schwander, J., Selmo, E., Severi, M., Stocker, T. F. and Udisti, R.:](#)  
3 [Expression of the bipolar see-saw in Antarctic climate records during the last deglaciation,](#)  
4 [Nat. Geosci., 4\(1\), 46–49, doi:10.1038/ngeo1026, 2011.](#)
- 5 [Stocker, T. F.: The Seesaw Effect, Science, 282\(5386\), 61–62,](#)  
6 [doi:10.1126/science.282.5386.61, 1998.](#)
- 7 [Strelin, J. A., Denton, G. H., Vandergoes, M. J., Ninnemann, U. S. and Putnam, A. E.:](#)  
8 [Radiocarbon chronology of the late-glacial Puerto Bandera moraines, Southern Patagonian](#)  
9 [Icefield, Argentina, Quat. Sci. Rev., 30\(19–20\), 2551–2569,](#)  
10 [doi:10.1016/j.quascirev.2011.05.004, 2011.](#)
- 11 [Stuhne, G. R. and Peltier, W. R.: Reconciling the ICE-6G C reconstruction of glacial](#)  
12 [chronology with ice sheet dynamics: The cases of Greenland and Antarctica, J. Geophys. Res.](#)  
13 [Earth Surf., 120\(9\), 2015JF003580, doi:10.1002/2015JF003580, 2015.](#)
- 14 [Svendsen, J. I., Alexanderson, H., Astakhov, V. I., Demidov, I., Dowdeswell, J. A., Funder,](#)  
15 [S., Gataullin, V., Henriksen, M., Hjort, C., Houmark-Nielsen, M., Hubberten, H. W.,](#)  
16 [Ingólfsson, Ó., Jakobsson, M., Kjær, K. H., Larsen, E., Lokrantz, H., Lunkka, J. P., Lyså, A.,](#)  
17 [Mangerud, J., Matiouchkov, A., Murray, A., Möller, P., Niessen, F., Nikolskaya, O., Polyak,](#)  
18 [L., Saarnisto, M., Siegert, C., Siegert, M. J., Spielhagen, R. F. and Stein, R.: Late Quaternary](#)  
19 [ice sheet history of northern Eurasia, Quat. Sci. Rev., 23\(11–13\), 1229–1271,](#)  
20 [doi:10.1016/j.quascirev.2003.12.008, 2004.](#)
- 21 [Tarasov, L. and Peltier, W. R.: Terminating the 100 kyr ice age cycle, J. Geophys. Res.](#)  
22 [Atmospheres, 102\(D18\), 21665–21693, doi:10.1029/97JD01766, 1997.](#)
- 23 [Tarasov, L. and Peltier, W. R.: Greenland glacial history and local geodynamic consequences,](#)  
24 [Geophys. J. Int., 150\(1\), 198–229, doi:10.1046/j.1365-246X.2002.01702.x, 2002.](#)
- 25 [Tarasov, L. and Peltier, W. R.: A geophysically constrained large ensemble analysis of the](#)  
26 [deglacial history of the North American ice-sheet complex, Quat. Sci. Rev., 23\(3–4\), 359–](#)  
27 [388, doi:10.1016/j.quascirev.2003.08.004, 2004.](#)
- 28 [Tarasov, L. and Peltier, W. R.: Arctic freshwater forcing of the Younger Dryas cold reversal,](#)  
29 [Nature, 435\(7042\), 662–665, doi:10.1038/nature03617, 2005.](#)

- 1 Tarasov, L. and Peltier, W. R.: A calibrated deglacial drainage chronology for the North  
2 American continent: evidence of an Arctic trigger for the Younger Dryas, *Quat. Sci. Rev.*,  
3 25(7–8), 659–688, doi:10.1016/j.quascirev.2005.12.006, 2006.
- 4 Tarasov, L., Dyke, A. S., Neal, R. M. and Peltier, W. R.: A data-calibrated distribution of  
5 deglacial chronologies for the North American ice complex from glaciological modeling,  
6 *Earth Planet. Sci. Lett.*, 315–316, 30–40, doi:10.1016/j.epsl.2011.09.010, 2012.
- 7 Taylor, K. E., Stouffer, R. J. and Meehl, G. A.: An Overview of CMIP5 and the Experiment  
8 Design, *Bull. Am. Meteorol. Soc.*, 93(4), 485–498, doi:10.1175/BAMS-D-11-00094.1, 2011a.
- 9 Taylor, K. E., Stouffer, R. J. and Meehl, G. A.: A Summary of the CMIP5 Experiment  
10 Design, [online] Available from: [http://cmip-](http://cmip-pcmdi.llnl.gov/cmip5/docs/Taylor_CMIP5_design.pdf)  
11 [pcmdi.llnl.gov/cmip5/docs/Taylor\\_CMIP5\\_design.pdf](http://cmip-pcmdi.llnl.gov/cmip5/docs/Taylor_CMIP5_design.pdf) (Accessed 13 November 2014b), 2011.
- 12 Teller, J. T., Leverington, D. W. and Mann, J. D.: Freshwater outbursts to the oceans from  
13 glacial Lake Agassiz and their role in climate change during the last deglaciation, *Quat. Sci.*  
14 *Rev.*, 21(8–9), 879–887, doi:10.1016/S0277-3791(01)00145-7, 2002.
- 15 Thiagarajan, N., Subhas, A. V., Southon, J. R., Eiler, J. M. and Adkins, J. F.: Abrupt pre-  
16 Bolling-Allerod warming and circulation changes in the deep ocean, *Nature*, 511(7507), 75–  
17 78, doi:10.1038/nature13472, 2014.
- 18 Thornalley, D. J. R., McCave, I. N. and Elderfield, H.: Freshwater input and abrupt deglacial  
19 climate change in the North Atlantic, *Paleoceanography*, 25(1), PA1201,  
20 doi:10.1029/2009PA001772, 2010.
- 21 Thornalley, D. J. R., Barker, S., Broecker, W. S., Elderfield, H. and McCave, I. N.: The  
22 Deglacial Evolution of North Atlantic Deep Convection, *Science*, 331(6014), 202–205,  
23 doi:10.1126/science.1196812, 2011.
- 24 Tierney, J. E., Russell, J. M., Huang, Y., Damsté, J. S. S., Hopmans, E. C. and Cohen, A. S.:  
25 Northern Hemisphere Controls on Tropical Southeast African Climate During the Past 60,000  
26 Years, *Science*, 322(5899), 252–255, doi:10.1126/science.1160485, 2008.
- 27 Timm, O. and Timmermann, A.: Simulation of the Last 21 000 Years Using Accelerated  
28 Transient Boundary Conditions\*, *J. Clim.*, 20(17), 4377–4401, doi:10.1175/JCLI4237.1,  
29 2007.

- 1 [Tjallingii, R., Claussen, M., Stuut, J.-B. W., Fohlmeister, J., Jahn, A., Bickert, T., Lamy, F.](#)  
2 [and Röhl, U.: Coherent high- and low-latitude control of the northwest African hydrological](#)  
3 [balance, Nat. Geosci., 1\(10\), 670–675, doi:10.1038/ngeo289, 2008.](#)
- 4 [Veres, D., Bazin, L., Landais, A., Toyé Mahamadou Kele, H., Lemieux-Dudon, B., Parrenin,](#)  
5 [F., Martinerie, P., Blayo, E., Blunier, T., Capron, E., Chappellaz, J., Rasmussen, S. O., Severi,](#)  
6 [M., Svensson, A., Vinther, B. and Wolff, E. W.: The Antarctic ice core chronology](#)  
7 [\(AICC2012\): an optimized multi-parameter and multi-site dating approach for the last 120](#)  
8 [thousand years, Clim Past, 9\(4\), 1733–1748, doi:10.5194/cp-9-1733-2013, 2013.](#)
- 9 [Verschuren, D., Sinninghe Damsté, J. S., Moernaut, J., Kristen, I., Blaauw, M., Fagot, M.,](#)  
10 [Haug, G. H., Geel, B. van, Batist, M. D., Barker, P., Vuille, M., Conley, D. J., Olago, D. O.,](#)  
11 [Milne, I., Plessen, B., Eggemont, H., Wolff, C., Hurrell, E., Ossebaar, J., Lyaruu, A., Plicht,](#)  
12 [J. van der, Cumming, B. F., Brauer, A., Rucina, S. M., Russell, J. M., Keppens, E., Hus, J.,](#)  
13 [Bradley, R. S., Leng, M., Mingram, J. and Nowaczyk, N. R.: Half-precessional dynamics of](#)  
14 [monsoon rainfall near the East African Equator, Nature, 462\(7273\), 637–641,](#)  
15 [doi:10.1038/nature08520, 2009.](#)
- 16 [Vettoretti, G. and Peltier, W. R.: Interhemispheric air temperature phase relationships in the](#)  
17 [nonlinear Dansgaard-Oeschger oscillation, Geophys. Res. Lett., 42\(4\), 2014GL062898,](#)  
18 [doi:10.1002/2014GL062898, 2015.](#)
- 19 [Waelbroeck, C., Labeyrie, L., Duplessy, J. C., Guiot, J., Labracherie, M., Leclaire, H. and](#)  
20 [Duprat, J.: Improving past sea surface temperature estimates based on planktonic fossil](#)  
21 [faunas, Paleoceanography, 13\(3\), 272–283, doi:10.1029/98PA00071, 1998.](#)
- 22 [Weaver, A. J., Saenko, O. A., Clark, P. U. and Mitrovica, J. X.: Meltwater Pulse 1A from](#)  
23 [Antarctica as a Trigger of the Bølling-Allerød Warm Interval, Science, 299\(5613\), 1709–](#)  
24 [1713, doi:10.1126/science.1081002, 2003.](#)
- 25 [Weber, S. L., Drijfhout, S. S., Abe-Ouchi, A., Crucifix, M., Eby, M., Ganopolski, A.,](#)  
26 [Murakami, S., Otto-Bliesner, B. and Peltier, W. R.: The modern and glacial overturning](#)  
27 [circulation in the Atlantic ocean in PMIP coupled model simulations, Clim Past, 3\(1\), 51–64,](#)  
28 [doi:10.5194/cp-3-51-2007, 2007.](#)

- 1 [Weijers, J. W. H., Schefuß, E., Schouten, S. and Damsté, J. S. S.: Coupled Thermal and](#)  
2 [Hydrological Evolution of Tropical Africa over the Last Deglaciation, Science, 315\(5819\),](#)  
3 [1701–1704, doi:10.1126/science.1138131, 2007.](#)
- 4 [Whitehouse, P. L., Bentley, M. J. and Le Brocq, A. M.: A deglacial model for Antarctica:](#)  
5 [geological constraints and glaciological modelling as a basis for a new model of Antarctic](#)  
6 [glacial isostatic adjustment, Quat. Sci. Rev., 32, 1–24, doi:10.1016/j.quascirev.2011.11.016,](#)  
7 [2012.](#)
- 8 [Yokoyama, Y., Esat, T. M. and Lambeck, K.: Last glacial sea-level change deduced from](#)  
9 [uplifted coral terraces of Huon Peninsula, Papua New Guinea, Quat. Int., 83–85, 275–283,](#)  
10 [doi:10.1016/S1040-6182\(01\)00045-3, 2001a.](#)
- 11 [Yokoyama, Y., De Deckker, P., Lambeck, K., Johnston, P. and Fifield, L. K.: Sea-level at the](#)  
12 [Last Glacial Maximum: evidence from northwestern Australia to constrain ice volumes for](#)  
13 [oxygen isotope stage 2, Palaeogeogr. Palaeoclimatol. Palaeoecol., 165\(3–4\), 281–297,](#)  
14 [doi:10.1016/S0031-0182\(00\)00164-4, 2001b.](#)
- 15 [Abe-Ouchi, A., Saito, F., Kageyama, M., Braeckman, P., Harrison, S. P., Lambeck, K., Otto-](#)  
16 [Bliesner, B. L., Peltier, W. R., Tarasov, L., Peterschmitt, J. Y. and Takahashi, K.: Ice sheet](#)  
17 [configuration in the CMIP5/PMIP3 Last Glacial Maximum experiments, Geosci Model Dev](#)  
18 [Discuss, 8\(6\), 4293–4336, doi:10.5194/gmdd-8-4293-2015, 2015.](#)
- 19 [Adkins, J. F., Ingersoll, A. P. and Pasquero, C.: Rapid climate change and conditional](#)  
20 [instability of the glacial deep ocean from the thermobaric effect and geothermal heating, Quat.](#)  
21 [Sci. Rev., 24\(5–6\), 581–594, doi:10.1016/j.quascirev.2004.11.005, 2005.](#)
- 22 [Aharon, P.: Entrainment of meltwaters in hyperpycnal flows during deglaciation superfloods](#)  
23 [in the Gulf of Mexico, Earth Planet. Sci. Lett., 241\(1–2\), 260–270,](#)  
24 [doi:10.1016/j.epsl.2005.10.034, 2006.](#)
- 25 [Ahn, J. and Brook, E. J.: Siple Dome ice reveals two modes of millennial CO<sub>2</sub> change during](#)  
26 [the last ice age, Nat. Commun., 5, 3723, doi:10.1038/ncomms4723, 2014.](#)
- 27 [Alley, R. B.: The Younger Dryas cold interval as viewed from central Greenland, Quat. Sci.](#)  
28 [Rev., 19\(1–5\), 213–226, doi:10.1016/S0277-3791\(99\)00062-1, 2000.](#)

- 1 Álvarez-Solas, J., Montoya, M., Ritz, C., Ramstein, G., Charbit, S., Dumas, C., Nisancioğlu,  
2 K., Dokken, T. and Ganopolski, A.: Heinrich event 1: an example of dynamical ice sheet  
3 reaction to oceanic changes, *Clim Past*, 7(4), 1297–1306, doi:10.5194/cp-7-1297-2011, 2011.
- 4 Annan, J. D. and Hargreaves, J. C.: A new global reconstruction of temperature changes at  
5 the Last Glacial Maximum, *Clim Past*, 9(1), 367–376, doi:10.5194/cp-9-367-2013, 2013.
- 6 Argus, D. F., Peltier, W. R., Drummond, R. and Moore, A. W.: The Antarctica component of  
7 postglacial rebound model ICE-6G-C (VM5a) based on GPS positioning, exposure age dating  
8 of ice thicknesses, and relative sea level histories, *Geophys. J. Int.*, ggu140,  
9 doi:10.1093/gji/ggu140, 2014.
- 10 Bard, E.: Climate shock—Abrupt changes over millennial time scales, *Phys. Today*, 55, 32–38,  
11 2002.
- 12 Bard, E., Hamelin, B., Arnold, M., Montaggioni, L., Cabioch, G., Faure, G. and Rougerie, F.:  
13 Deglacial sea-level record from Tahiti corals and the timing of global meltwater discharge,  
14 *Nature*, 382(6588), 241–244, doi:10.1038/382241a0, 1996.
- 15 Bard, E., Rostek, F., Turon, J. L. and Gendreau, S.: Hydrological Impact of Heinrich Events  
16 in the Subtropical Northeast Atlantic, *Science*, 289(5483), 1321–1324,  
17 doi:10.1126/science.289.5483.1321, 2000.
- 18 Bard, E., Hamelin, B. and Delanghe Sabatier, D.: Deglacial Meltwater Pulse 1B and Younger  
19 Dryas Sea Levels Revisited with Boreholes at Tahiti, *Science*, 327(5970), 1235–1237,  
20 doi:10.1126/science.1180557, 2010.
- 21 Barker, S., Knorr, G., Vautravers, M. J., Diz, P. and Skinner, L. C.: Extreme deepening of the  
22 Atlantic overturning circulation during deglaciation, *Nat. Geosci.*, 3(8), 567–571,  
23 doi:10.1038/ngeo921, 2010.
- 24 Barker, S., Chen, J., Gong, X., Jonkers, L., Knorr, G. and Thornalley, D.: Icebergs not the  
25 trigger for North Atlantic cold events, *Nature*, 520(7547), 333–336, doi:10.1038/nature14330,  
26 2015.
- 27 Bassett, S. E., Milne, G. A., Mitrovica, J. X. and Clark, P. U.: Ice Sheet and Solid Earth  
28 Influences on Far Field Sea Level Histories, *Science*, 309(5736), 925–928,  
29 doi:10.1126/science.1111575, 2005.

- 1 Bassett, S. E., Milne, G. A., Bentley, M. J. and Huybrechts, P.: Modelling Antarctic sea level  
2 data to explore the possibility of a dominant Antarctic contribution to meltwater pulse IA,  
3 Quat. Sci. Rev., 26(17–18), 2113–2127, doi:10.1016/j.quascirev.2007.06.011, 2007.
- 4 Bauer, E. and Ganopolski, A.: Sensitivity simulations with direct shortwave radiative forcing  
5 by aeolian dust during glacial cycles, Clim Past, 10(4), 1333–1348, doi:10.5194/cp-10-1333-  
6 2014, 2014.
- 7 Bentley, M. J., Fogwill, C. J., Brocq, A. M. L., Hubbard, A. L., Sugden, D. E., Dunai, T. J.  
8 and Freeman, S. P. H. T.: Deglacial history of the West Antarctic Ice Sheet in the Weddell  
9 Sea embayment: Constraints on past ice volume change, Geology, 38(5), 411–414,  
10 doi:10.1130/G30754.1, 2010.
- 11 Bentley, M. J., Ó Cofaigh, C., Anderson, J. B., Conway, H., Davies, B., Graham, A. G. C.,  
12 Hillenbrand, C. D., Hodgson, D. A., Jamieson, S. S. R., Larter, R. D., Mackintosh, A., Smith,  
13 J. A., Verleyen, E., Ackert, R. P., Bart, P. J., Berg, S., Brunstein, D., Canals, M., Colhoun, E.  
14 A., Crosta, X., Dickens, W. A., Domack, E., Dowdeswell, J. A., Dunbar, R., Ehrmann, W.,  
15 Evans, J., Favier, V., Fink, D., Fogwill, C. J., Glasser, N. F., Gohl, K., Golledge, N. R.,  
16 Goodwin, I., Gore, D. B., Greenwood, S. L., Hall, B. L., Hall, K., Hedding, D. W., Hein, A.  
17 S., Hocking, E. P., Jakobsson, M., Johnson, J. S., Jomelli, V., Jones, R. S., Klages, J. P.,  
18 Kristoffersen, Y., Kuhn, G., Leventer, A., Licht, K., Lilly, K., Lindow, J., Livingstone, S. J.,  
19 Massé, G., McGlone, M. S., McKay, R. M., Melles, M., Miura, H., Mulvaney, R., Nel, W.,  
20 Nitsche, F. O., O'Brien, P. E., Post, A. L., Roberts, S. J., Saunders, K. M., Selkirk, P. M.,  
21 Simms, A. R., Spiegel, C., Stolldorf, T. D., Sugden, D. E., van der Putten, N., van Ommen,  
22 T., Verfaillie, D., Vyverman, W., Wagner, B., White, D. A., Witus, A. E. and Zwartz, D.: A  
23 community-based geological reconstruction of Antarctic Ice Sheet deglaciation since the Last  
24 Glacial Maximum, Quat. Sci. Rev., 100, 1–9, doi:10.1016/j.quascirev.2014.06.025, 2014.
- 25 Bereiter, B., Eggleston, S., Schmitt, J., Nehrbass Ahles, C., Stoeker, T. F., Fischer, H.,  
26 Kipfahl, S. and Chappellaz, J.: Revision of the EPICA Dome C CO<sub>2</sub> record from 800 to  
27 600 kyr before present, Geophys. Res. Lett., 42(2), 2014GL061957,  
28 doi:10.1002/2014GL061957, 2015.

- 1 Berger, A.: Long Term Variations of Daily Insolation and Quaternary Climatic Changes, *J.  
2 Atmospheric Sci.*, 35(12), 2362–2367, doi:10.1175/1520-  
3 0469(1978)035<2362:LTVO>2.0.CO;2, 1978.
- 4 Berger, A. and Loutre, M. F.: Insolation values for the climate of the last 10 million years,  
5 *Quat. Sci. Rev.*, 10(4), 297–317, doi:10.1016/0277-3791(91)90033-Q, 1991.
- 6 Benelli, S., Charbit, S., Kageyama, M., Woillez, M. N., Ramstein, G., Dumas, C. and  
7 Quinet, A.: Investigating the evolution of major Northern Hemisphere ice sheets during the  
8 last glacial interglacial cycle, *Clim Past*, 5(3), 329–345, doi:10.5194/cp-5-329-2009, 2009.
- 9 Boulton, G. S., Dongelmans, P., Punkari, M. and Broadgate, M.: Palaeoglaciology of an ice  
10 sheet through a glacial cycle:: the European ice sheet through the Weichselian, *Quat. Sci.  
11 Rev.*, 20(4), 591–625, doi:10.1016/S0277-3791(00)00160-8, 2001.
- 12 Braconnot, P., Joussaume, S., de Noblet, N. and Ramstein, G.: Mid Holocene and Last  
13 Glacial Maximum African monsoon changes as simulated within the Paleoclimate Modelling  
14 Intercarparison Project, *Glob. Planet. Change*, 26(1–3), 51–66, doi:10.1016/S0921-  
15 8181(00)00033-3, 2000.
- 16 Braconnot, P., Otto-Bliesner, B., Harrison, S., Joussaume, S., Peterchmitt, J. Y., Abe-Ouchi,  
17 A., Crucifix, M., Driesschaert, E., Fichefet, T., Hewitt, C. D., Kageyama, M., Kitoh, A.,  
18 Lainé, A., Loutre, M. F., Marti, O., Merkel, U., Ramstein, G., Valdes, P., Weber, S. L., Yu,  
19 Y. and Zhao, Y.: Results of PMIP2 coupled simulations of the Mid Holocene and Last  
20 Glacial Maximum—Part 1: experiments and large scale features, *Clim Past*, 3(2), 261–277,  
21 doi:10.5194/cp-3-261-2007, 2007.
- 22 Braconnot, P., Harrison, S. P., Kageyama, M., Bartlein, P. J., Masson-Delmotte, V., Abe-  
23 Ouchi, A., Otto-Bliesner, B. and Zhao, Y.: Evaluation of climate models using palaeoclimatic  
24 data, *Nat. Clim. Change*, 2(6), 417–424, doi:10.1038/nclimate1456, 2012.
- 25 Briggs, R. D., Pollard, D. and Tarasov, L.: A data constrained large ensemble analysis of  
26 Antarctic evolution since the Eemian, *Quat. Sci. Rev.*, 103, 91–115,  
27 doi:10.1016/j.quascirev.2014.09.003, 2014.
- 28 Broecker, W. S.: Does the Trigger for Abrupt Climate Change Reside in the Ocean or in the  
29 Atmosphere?, *Science*, 300(5625), 1519–1522, doi:10.1126/science.1083797, 2003.

- 1 Broecker, W. S., Kennett, J. P., Flower, B. P., Teller, J. T., Trumbore, S., Bonani, G. and  
2 Wolfli, W.: Routing of meltwater from the Laurentide Ice Sheet during the Younger Dryas  
3 cold episode, *Nature*, 341(6240), 318–321, doi:10.1038/341318a0, 1989.
- 4 Broekin, V., Ganopolski, A., Archer, D. and Munhoven, G.: Glacial CO<sub>2</sub> cycle as a  
5 succession of key physical and biogeochemical processes, *Clim. Past*, 8(1), 251–264,  
6 doi:10.5194/cp-8-251-2012, 2012.
- 7 Buizert, C., Gkinis, V., Severinghaus, J. P., He, F., Leesavalier, B. S., Kindler, P.,  
8 Leuenberger, M., Carlson, A. E., Vinther, B., Masson-Delmotte, V., White, J. W. C., Liu, Z.,  
9 Otto-Bliesner, B. and Brook, E. J.: Greenland temperature response to climate forcing during  
10 the last deglaciation, *Science*, 345(6201), 1177–1180, doi:10.1126/science.1254961, 2014.
- 11 Cabioch, G., Banks-Cutler, K. A., Beck, W. J., Burr, G. S., Corrège, T., Lawrence-Edwards,  
12 R. and Taylor, F. W.: Continuous reef growth during the last 23 cal kyr BP in a tectonically  
13 active zone (Vanuatu, SouthWest Pacific), *Quat. Sci. Rev.*, 22(15–17), 1771–1786,  
14 doi:10.1016/S0277-3791(03)00170-7, 2003.
- 15 Caley, T., Roehe, D. M. and Renssen, H.: Orbital Asian summer monsoon dynamics revealed  
16 using an isotope-enabled global climate model, *Nat. Commun.*, 5, doi:10.1038/ncomms6371,  
17 2014.
- 18 Carlson, A. E.: Geochemical constraints on the Laurentide Ice Sheet contribution to  
19 Meltwater Pulse 1A, *Quat. Sci. Rev.*, 28(17–18), 1625–1630,  
20 doi:10.1016/j.quascirev.2009.02.011, 2009.
- 21 Clark, P. U. and Mix, A. C.: Ice sheets and sea level of the Last Glacial Maximum, *Quat. Sci. Rev.*,  
22 21(1–3), 1–7, doi:10.1016/S0277-3791(01)00118-4, 2002.
- 23 Clark, P. U. and Tarasov, L.: Closing the sea level budget at the Last Glacial Maximum, *Proc. Natl. Acad. Sci.*,  
24 111(45), 15861–15862, doi:10.1073/pnas.1418970111, 2014.
- 25 Clark, P. U., Alley, R. B., Keigwin, L. D., Licciardi, J. M., Johnsen, S. J. and Wang, H.:  
26 Origin of the first global meltwater pulse following the Last Glacial Maximum,  
27 *Paleoceanography*, 11(5), 563–577, doi:10.1029/96PA01419, 1996.

- 1 Clark, P. U., Mitrovica, J. X., Milne, G. A. and Tamisiea, M. E.: Sea Level Fingerprinting as  
2 a Direct Test for the Source of Global Meltwater Pulse 1A, *Science*, 295(5564), 2438–2441,  
3 doi:10.1126/science.1068797, 2002.
- 4 Clark, P. U., McCabe, A. M., Mix, A. C. and Weaver, A. J.: Rapid Rise of Sea Level 19,000  
5 Years Ago and Its Global Implications, *Science*, 304(5674), 1141–1144,  
6 doi:10.1126/science.1094449, 2004.
- 7 Clark, P. U., Dyke, A. S., Shakun, J. D., Carlson, A. E., Clark, J., Wohlfarth, B., Mitrovica, J.  
8 X., Hostetler, S. W. and McCabe, A. M.: The Last Glacial Maximum, *Science*, 325(5941),  
9 710–714, doi:10.1126/science.1172873, 2009.
- 10 Clark, P. U., Shakun, J. D., Baker, P. A., Bartlein, P. J., Brewer, S., Brook, E., Carlson, A. E.,  
11 Cheng, H., Kaufman, D. S., Liu, Z., Marchitto, T. M., Mix, A. C., Morrill, C., Otto Bliesner,  
12 B. L., Pahnke, K., Russell, J. M., Whitlock, C., Adkins, J. F., Blois, J. L., Clark, J., Colman,  
13 S. M., Curry, W. B., Flower, B. P., He, F., Johnson, T. C., Lynch Stieglitz, J., Markgraf, V.,  
14 McManus, J., Mitrovica, J. X., Moreno, P. I. and Williams, J. W.: Global climate evolution  
15 during the last deglaciation, *Proc. Natl. Acad. Sci.*, 109(19), E1134–E1142,  
16 doi:10.1073/pnas.1116619109, 2012.
- 17 Condron, A. and Winsor, P.: Meltwater routing and the Younger Dryas, *Proc. Natl. Acad.  
Sci.*, 109(49), 19928–19933, doi:10.1073/pnas.1207381109, 2012.
- 19 Cuffey, K. M. and Clow, G. D.: Temperature, accumulation, and ice sheet elevation in central  
20 Greenland through the last deglacial transition, *J. Geophys. Res.*, 102(C12), 26383–26,396,  
21 doi:10.1029/96JC03981, 1997.
- 22 Cutler, K. B., Edwards, R. L., Taylor, F. W., Cheng, H., Adkins, J., Gallup, C. D., Cutler, P.  
23 M., Burr, G. S. and Bloom, A. L.: Rapid sea-level fall and deep-ocean temperature change  
24 since the last interglacial period, *Earth Planet. Sci. Lett.*, 206(3–4), 253–271,  
25 doi:10.1016/S0012-821X(02)01107-X, 2003.
- 26 De Deckker, P. and Yokoyama, Y.: Micropalaeontological evidence for Late Quaternary sea-  
27 level changes in Bonaparte Gulf, Australia, *Glob. Planet. Change*, 66(1–2), 85–92,  
28 doi:10.1016/j.gloplacha.2008.03.012, 2009.

1 deMenocal, P., Ortiz, J., Guilderson, T., Adkins, J., Sarnthein, M., Baker, L. and Yarusinsky,  
2 M.: Abrupt onset and termination of the African Humid Period: rapid climate responses to  
3 gradual insolation forcing, *Quat. Sci. Rev.*, 19(1–5), 347–361, doi:10.1016/S0277-  
4 3791(99)00081-5, 2000.

5 Desehamps, P., Durand, N., Bard, E., Hamelin, B., Camoin, G., Thomas, A. L., Henderson,  
6 G. M., Okuno, J. 'ichi and Yokoyama, Y.: Ice sheet collapse and sea level rise at the Bolling  
7 warming 14,600 years ago, *Nature*, 483(7391), 559–564, doi:10.1038/nature10902, 2012.

8 Dyke, A. S.: An outline of North American deglaciation with emphasis on central and  
9 northern Canada, in *Quaternary Glaciations Extent and Chronology – Part II: North America*,  
10 vol. Volume 2, Part 2, pp. 373–424, Elsevier. [online] Available from:  
11 [https://www.lakeheadu.ca/sites/default/files/uploads/53/outlines/2014-15/NECU5311/Dyke\\_2004\\_DeglaciationOutline.pdf](https://www.lakeheadu.ca/sites/default/files/uploads/53/outlines/2014-15/NECU5311/Dyke_2004_DeglaciationOutline.pdf) (Accessed 20 October 2015), 2004.

13 Dyke, A. S., Andrews, J. T., Clark, P. U., England, J. H., Miller, G. H., Shaw, J. and Veillette,  
14 J. J.: The Laurentide and Innuitian ice sheets during the Last Glacial Maximum, *Quat. Sci. Rev.*,  
15 21(1–3), 9–31, doi:10.1016/S0277-3791(01)00095-6, 2002.

16 Edwards, R. L., Beck, J. W., Burr, G. S., Donahue, D. J., Chappell, J. M. A., Bloom, A. L.,  
17 Druffel, E. R. M. and Taylor, F. W.: A Large Drop in Atmospheric  $^{14}\text{C}/^{12}\text{C}$  and Reduced  
18 Melting in the Younger Dryas, Documented with  $^{230}\text{Th}$  Ages of Corals, *Science*, 260(5110),  
19 962–968, doi:10.1126/science.260.5110.962, 1993.

20 EPICA Community Members: Eight glacial cycles from an Antarctic ice core, *Nature*,  
21 429(6992), 623–628, doi:10.1038/nature02599, 2004.

22 Fairbanks, R. G.: A 17,000-year glacio-eustatic sea-level record: influence of glacial melting  
23 rates on the Younger Dryas event and deep-ocean circulation, *Nature*, 342(6250), 637–642,  
24 doi:10.1038/342637a0, 1989.

25 Flato, G., Marotzke, J., Abiodun, B., Braconnot, P., Chou, S. C., Collins, W., Cox, P.,  
26 Driouech, F., Emori, S., Eyring, V., Forest, C., Glecker, P., Guilyardi, E., Jackob, C., Kattsov,  
27 V., Reason, C. and Rummukainen, M.: Evaluation of Climate Models, in *Climate Change  
2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment  
Report of the Intergovernmental Panel on Climate Change*, edited by T. F. Stocker, D. Qin, G.  
30 Plattner, M. Tignor, S. K. Allen, J. Boschung, A. Nauels, Y. Xia, V. Bex, and P. M.

- 1 Midgley, pp. 741–866, Cambridge University Press, Cambridge, United Kingdom and New  
2 York, NY, USA. [online] Available from: [http://www.ipcc.ch/pdf/assessment-report/ar5/wg1/WG1AR5\\_Chapter09\\_FINAL.pdf](http://www.ipcc.ch/pdf/assessment-report/ar5/wg1/WG1AR5_Chapter09_FINAL.pdf) (Accessed 20 October 2015), 2013.
- 4 Ganopolski, A. and Calov, R.: The role of orbital forcing, carbon dioxide and regolith in 100  
5 kyr glacial cycles, *Clim Past*, 7(4), 1415–1425, doi:10.5194/cp-7-1415-2011, 2011.
- 6 Ganopolski, A. and Rahmstorf, S.: Rapid changes of glacial climate simulated in a coupled  
7 climate model, *Nature*, 409(6817), 153–158, doi:10.1038/35051500, 2001.
- 8 Ganopolski, A., Calov, R. and Claussen, M.: Simulation of the last glacial cycle with a  
9 coupled climate ice sheet model of intermediate complexity, *Clim Past*, 6(2), 229–244,  
10 doi:10.5194/cp-6-229-2010, 2010.
- 11 García, J. L., Kaplan, M. R., Hall, B. L., Schaefer, J. M., Vega, R. M., Schwartz, R. and  
12 Finkel, R.: Glacier expansion in southern Patagonia throughout the Antarctic cold reversal,  
13 *Geology*, 40(9), 859–862, doi:10.1130/G33164.1, 2012.
- 14 Gherardi, J. M., Labeyrie, L., McManus, J. F., Francois, R., Skinner, L. C. and Cortijo, E.:  
15 Evidence from the Northeastern Atlantic basin for variability in the rate of the meridional  
16 overturning circulation through the last deglaciation, *Earth Planet. Sci. Lett.*, 240(3–4), 710–  
17 723, doi:10.1016/j.epsl.2005.09.061, 2005.
- 18 Golledge, N. R., Fogwill, C. J., Mackintosh, A. N. and Buckley, K. M.: Dynamics of the last  
19 glacial maximum Antarctic ice sheet and its response to ocean forcing, *Proc. Natl. Acad. Sci.*,  
20 109(40), 16052–16056, doi:10.1073/pnas.1205385109, 2012.
- 21 Golledge, N. R., Levy, R. H., McKay, R. M., Fogwill, C. J., White, D. A., Graham, A. G. C.,  
22 Smith, J. A., Hillenbrand, C. D., Lieft, K. J., Denton, G. H., Aekert Jr., R. P., Maas, S. M.  
23 and Hall, B. L.: Glaciology and geological signature of the Last Glacial Maximum Antarctic  
24 ice sheet, *Quat. Sci. Rev.*, 78, 225–247, doi:10.1016/j.quascirev.2013.08.011, 2013.
- 25 Golledge, N. R., Menzel, L., Carter, L., Fogwill, C. J., England, M. H., Cortese, G. and Levy,  
26 R. H.: Antarctic contribution to meltwater pulse 1A from reduced Southern Ocean  
27 overturning, *Nat. Commun.*, 5, doi:10.1038/ncomms6107, 2014.
- 28 Goujon, C., Barnola, J. M. and Ritz, C.: Modeling the densification of polar firn including  
29 heat diffusion: Application to close off characteristics and gas isotopic fractionation for

- 1 Antarctica and Greenland sites, *J. Geophys. Res. Atmospheres*, 108(D24), 4792,  
2 doi:10.1029/2002JD003319, 2003.
- 3 Gregoire, L. J., Payne, A. J. and Valdes, P. J.: Deglacial rapid sea level rises caused by ice  
4 sheet saddle collapses, *Nature*, 487(7406), 219–222, doi:10.1038/nature11257, 2012.
- 5 Gregoire, L. J., Valdes, P. J. and Payne, A. J.: The relative contribution of orbital forcing and  
6 greenhouse gases to the North American deglaciation, *Geophys. Res. Lett.*, 42(22),  
7 2015GL066005, doi:10.1002/2015GL066005, 2015.
- 8 Gyllencreutz, R., Mangerud, J., Svendsen, J. I. and Lohne, Ø.: DATED a GIS based  
9 reconstruction and dating database of the Eurasian deglaciation, *Appl. Quat. Res. Cent. Part*  
10 *Glaciat. Terrain Geol. Surv. Finl. Spec. Pap.*, 46, 113–120, 2007.
- 11 Hall, I. R., Moran, S. B., Zahn, R., Knutz, P. C., Shen, C. C. and Edwards, R. L.: Accelerated  
12 drawdown of meridional overturning in the late glacial Atlantic triggered by transient pre-H  
13 event freshwater perturbation, *Geophys. Res. Lett.*, 33(16), L16616,  
14 doi:10.1029/2006GL026239, 2006.
- 15 Hanebuth, T., Stattegger, K. and Grootes, P. M.: Rapid Flooding of the Sunda Shelf: A Late-  
16 Glacial Sea Level Record, *Science*, 288(5468), 1033–1035,  
17 doi:10.1126/science.288.5468.1033, 2000.
- 18 Hanebuth, T. J. J., Stattegger, K. and Bojanowski, A.: Termination of the Last Glacial  
19 Maximum sea level lowstand: The Sunda Shelf data revisited, *Glob. Planet. Change*, 66(1–2),  
20 76–84, doi:10.1016/j.gloplacha.2008.03.011, 2009.
- 21 Haywood, A. M., Dowsett, H. J., Otto-Bliesner, B., Chandler, M. A., Dolan, A. M., Hill, D.  
22 J., Lunt, D. J., Robinson, M. M., Rosenbloom, N., Salzmann, U. and Sohl, L. E.: Pliocene  
23 Model Intercomparison Project (PlioMIP): experimental design and boundary conditions  
24 (Experiment 1), *Geosci. Model Dev.*, 3(1), 227–242, doi:10.5194/gmd-3-227-2010, 2010.
- 25 Heinemann, M., Timmermann, A., Elison Timm, O., Saito, F. and Abe-Ouchi, A.: Deglacial  
26 ice sheet meltdown: orbital pacemaking and CO<sub>2</sub> effects, *Clim. Past*, 10(4), 1567–1579,  
27 doi:10.5194/cp-10-1567-2014, 2014.
- 28 Heiri, O., Cremer, H., Engels, S., Hoek, W. Z., Peeters, W. and Lotter, A. F.: Lateglacial  
29 summer temperatures in the Northwest European lowlands: a chironomid record from

- 1 Hijkermeer, the Netherlands, *Quat. Sci. Rev.*, 26(19–21), 2420–2437,  
2 doi:10.1016/j.quascirev.2007.06.017, 2007.
- 3 Hemming, S. R.: Heinrich events: Massive late Pleistocene detritus layers of the North  
4 Atlantic and their global climate imprint, *Rev. Geophys.*, 42(1), RG1005,  
5 doi:10.1029/2003RG000128, 2004.
- 6 Hughes, A. L. C., Gyllencreutz, R., Lohne, Ø. S., Mangerud, J. and Svendsen, J. I.: The last  
7 Eurasian ice sheets – a chronological database and time slice reconstruction, DATED 1,  
8 *Boreas*, doi:10.1111/bor.12142, 2015.
- 9 Joussaume, S., Taylor, K. E., Braconnot, P., Mitchell, J. F. B., Kutzbach, J. E., Harrison, S.  
10 P., Prentice, I. C., Broeckli, A. J., Abe-Ouchi, A., Bartlein, P. J., Bonfils, C., Dong, B., Guiot,  
11 J., Herterich, K., Hewitt, C. D., Jolly, D., Kim, J. W., Kislov, A., Kitoh, A., Loutre, M. F.,  
12 Masson, V., McAvaney, B., McFarlane, N., de Noblet, N., Peltier, W. R., Peterchmitt, J. Y.,  
13 Pollard, D., Rind, D., Royer, J. F., Schlesinger, M. E., Syktus, J., Thompson, S., Valdes, P.,  
14 Vettoretti, G., Webb, R. S. and Wyputta, U.: Monsoon changes for 6000 years ago: Results of  
15 18 simulations from the Paleoclimate Modeling Intercomparison Project (PMIP), *Geophys.  
16 Res. Lett.*, 26(7), 859–862, doi:10.1029/1999GL900126, 1999.
- 17 Jouzel, J., Vaikmae, R., Petit, J. R., Martin, M., Duclos, Y., Stievenard, M., Lorius, C., Toots,  
18 M., Molières, M. A., Burek, L. H., Barkov, N. I. and Kotlyakov, V. M.: The two-step shape  
19 and timing of the last deglaciation in Antarctica, *Clim. Dyn.*, 11(3), 151–161,  
20 doi:10.1007/BF00223498, 1995.
- 21 Jouzel, J., Masson-Delmotte, V., Cattani, O., Dreyfus, G., Falourd, S., Hoffmann, G., Minster,  
22 P., Nouet, J., Barnola, J. M., Chappellaz, J., Fischer, H., Gallet, J. C., Johnsen, S.,  
23 Leuenberger, M., Louergue, L., Luethi, D., Oerter, H., Parrenin, F., Raisbeck, G., Raynaud,  
24 D., Schilt, A., Schwander, J., Selmo, E., Souchez, R., Spahni, R., Stauffer, B., Steffensen, J.  
25 P., Stenni, B., Stocker, T. F., Tison, J. L., Werner, M. and Wolff, E. W.: Orbital and  
26 Millennial Antarctic Climate Variability over the Past 800,000 Years, *Science*, 317(5839),  
27 793–796, doi:10.1126/science.1141038, 2007.
- 28 Kageyama, M., Braconnot, P., Peterchmitt, J. Y., Harrison, S. P., Jungclaus, J. H., Otto-  
29 Bliesner, B., Brierley, C., Crucifix, M., Haywood, A. M., Ivanovic, R. F., Lunt, D. J., Phipps,

- 1 S. J., Roche, D. M., Schmidt, G. A., Zhang, Q. and Zhou, Y.: Overview of the PMIP4 CMIP6  
2 experiments, *Geosci. Model Dev.*, in prep.
- 3 Kageyama, M., Lainé, A., Abe Ouchi, A., Braconnot, P., Cortijo, E., Crucifix, M., de Vernal,  
4 A., Guiot, J., Hewitt, C. D., Kitoh, A., Kucera, M., Martí, O., Ohgaito, R., Otto Bliesner, B.,  
5 Peltier, W. R., Rosell Melé, A., Vettoretti, G., Weber, S. L. and Yu, Y.: Last Glacial  
6 Maximum temperatures over the North Atlantic, Europe and western Siberia: a comparison  
7 between PMIP models, MARGO sea surface temperatures and pollen-based reconstructions,  
8 *Quat. Sci. Rev.*, 25(17–18), 2082–2102, doi:10.1016/j.quascirev.2006.02.010, 2006.
- 9 Kageyama, M., Paul, A., Roche, D. M. and Van Meerbeeck, C. J.: Modelling glacial climatic  
10 millennial scale variability related to changes in the Atlantic meridional overturning  
11 circulation: a review, *Quat. Sci. Rev.*, 29(21–22), 2931–2956,  
12 doi:10.1016/j.quascirev.2010.05.029, 2010.
- 13 Kageyama, M., Merkel, U., Otto Bliesner, B., Prange, M., Abe Ouchi, A., Lohmann, G.,  
14 Ohgaito, R., Roche, D. M., Singarayer, J., Swingedouw, D. and X Zhang: Climatic impacts of  
15 fresh water hosing under Last Glacial Maximum conditions: a multi-model study, *Clim Past*,  
16 9(2), 935–953, doi:10.5194/cp-9-935-2013, 2013.
- 17 Kaplan, M. R., Strelin, J. A., Schaefer, J. M., Denton, G. H., Finkel, R. C., Schwartz, R.,  
18 Putnam, A. E., Vandergoes, M. J., Goehring, B. M. and Travis, S. G.: In situ cosmogenic  
19  $^{10}\text{Be}$  production rate at Lago Argentino, Patagonia: Implications for late glacial climate  
20 chronology, *Earth Planet. Sci. Lett.*, 309(1–2), 21–32, doi:10.1016/j.epsl.2011.06.018, 2011.
- 21 Keigwin, L. D. and Boyle, E. A.: Did North Atlantic overturning halt 17,000 years ago?,  
22 *Paleoceanography*, 23(1), PA1101, doi:10.1029/2007PA001500, 2008.
- 23 Keigwin, L. D., Jones, G. A., Lehman, S. J. and Boyle, E. A.: Deglacial meltwater discharge,  
24 North Atlantic Deep Circulation, and abrupt climate change, *J. Geophys. Res. Oceans*,  
25 96(C9), 16811–16826, doi:10.1029/91JC01624, 1991.
- 26 Kindler, P., Guillemin, M., Baumgartner, M., Schwander, J., Landais, A. and Leuenberger, M.:  
27 Temperature reconstruction from 10 to 120 kyr b2k from the NGRIP ice core, *Clim Past*,  
28 10(2), 887–902, doi:10.5194/cp-10-887-2014, 2014.

- 1 Knorr, G. and Lohmann, G.: Southern Ocean origin for the resumption of Atlantic  
2 thermohaline circulation during deglaciation, *Nature*, 424(6948), 532–536,  
3 doi:10.1038/nature01855, 2003.
- 4 Knorr, G. and Lohmann, G.: Rapid transitions in the Atlantic thermohaline circulation  
5 triggered by global warming and meltwater during the last deglaciation, *Geochem. Geophys.*  
6 *Geosystems*, 8(12), Q12006, doi:10.1029/2007GC001604, 2007.
- 7 Kehfeld, K. . and Harrison, S. .: How well can we simulate past climates? Evaluating the  
8 models using global palaeoenvironmental datasets, *Quat. Sci. Rev.*, 19(1–5), 321–346,  
9 doi:10.1016/S0277-3791(99)00068-2, 2000.
- 10 Lambeck, K., Rouby, H., Purcell, A., Sun, Y. and Sambridge, M.: Sea level and global ice  
11 volumes from the Last Glacial Maximum to the Holocene, *Proc. Natl. Acad. Sci.*,  
12 doi:10.1073/pnas.1411762111, 2014.
- 13 Lea, D. W., Pak, D. K., Peterson, L. C. and Hughen, K. A.: Synchronicity of Tropical and  
14 High-Latitude Atlantic Temperatures over the Last Glacial Termination, *Science*, 301(5638),  
15 1361–1364, doi:10.1126/science.1088470, 2003.
- 16 Licht, K. J.: The Ross Sea's contribution to eustatic sea level during meltwater pulse 1A,  
17 *Sediment. Geol.*, 165(3–4), 343–353, doi:10.1016/j.sedgeo.2003.11.020, 2004.
- 18 Liu, Z., Otto Bliesner, B. L., He, F., Brady, E. C., Tomas, R., Clark, P. U., Carlson, A. E.,  
19 Lynch Stieglitz, J., Curry, W., Brook, E., Erickson, D., Jacob, R., Kutzbach, J. and Cheng, J.:  
20 Transient Simulation of Last Deglaciation with a New Mechanism for Bølling Allerød  
21 Warming, *Science*, 325(5938), 310–314, doi:10.1126/science.1171041, 2009.
- 22 Liu, Z., Carlson, A. E., He, F., Brady, E. C., Otto Bliesner, B. L., Briegleb, B. P.,  
23 Wehrenberg, M., Clark, P. U., Wu, S., Cheng, J., Zhang, J., Noone, D. and Zhu, J.: Younger  
24 Dryas cooling and the Greenland climate response to CO<sub>2</sub>, *Proc. Natl. Acad. Sci.*,  
25 doi:10.1073/pnas.1202183109, 2012.
- 26 Liu, Z., Wen, X., Brady, E. C., Otto Bliesner, B., Yu, G., Lu, H., Cheng, H., Wang, Y.,  
27 Zheng, W., Ding, Y., Edwards, R. L., Cheng, J., Liu, W. and Yang, H.: Chinese cave records  
28 and the East Asia Summer Monsoon, *Quat. Sci. Rev.*, 83, 115–128,  
29 doi:10.1016/j.quascirev.2013.10.021, 2014.

- 1 Louergue, L., Schilt, A., Spahni, R., Masson Delmotte, V., Blunier, T., Lemieux, B.,  
2 Barnola, J. M., Raynaud, D., Stocker, T. F. and Chappellaz, J.: Orbital and millennial-scale  
3 features of atmospheric CH<sub>4</sub> over the past 800,000 years, *Nature*, 453(7193), 383–386,  
4 doi:10.1038/nature06950, 2008.
- 5 Lüthi, D., Le Floch, M., Bereiter, B., Blunier, T., Barnola, J. M., Siegenthaler, U., Raynaud,  
6 D., Jouzel, J., Fischer, H., Kawamura, K. and Stocker, T. F.: High resolution carbon dioxide  
7 concentration record 650,000–800,000 years before present, *Nature*, 453(7193), 379–382,  
8 doi:10.1038/nature06949, 2008.
- 9 MacFarling Meure, C., Etheridge, D., Trudinger, C., Steele, P., Langenfelds, R., van Ommen,  
10 T., Smith, A. and Elkins, J.: Law Dome CO<sub>2</sub>, CH<sub>4</sub> and N<sub>2</sub>O ice core records extended to  
11 2000 years BP, *Geophys. Res. Lett.*, 33(14), L14810, doi:10.1029/2006GL026152, 2006.
- 12 Mackintosh, A., Golledge, N., Domack, E., Dunbar, R., Leventer, A., White, D., Pollard, D.,  
13 DeConto, R., Fink, D., Zwart, D., Gore, D. and Lavoie, C.: Retreat of the East Antarctic ice  
14 sheet during the last glacial termination, *Nat. Geosci.*, 4(3), 195–202, doi:10.1038/ngeo1061,  
15 2011.
- 16 Mackintosh, A. N., Verleyen, E., O'Brien, P. E., White, D. A., Jones, R. S., McKay, R.,  
17 Dunbar, R., Gore, D. B., Fink, D., Post, A. L., Miura, H., Leventer, A., Goodwin, I., Hodgson,  
18 D. A., Lilly, K., Crosta, X., Golledge, N. R., Wagner, B., Berg, S., van Ommen, T., Zwart,  
19 D., Roberts, S. J., Vyverman, W. and Masse, G.: Retreat history of the East Antarctic Ice  
20 Sheet since the Last Glacial Maximum, *Quat. Sci. Rev.*, 100, 10–30,  
21 doi:10.1016/j.quascirev.2013.07.024, 2014.
- 22 Marcott, S. A., Bauska, T. K., Buizert, C., Steig, E. J., Rosen, J. L., Cuffey, K. M., Fudge, T.,  
23 J., Severinghaus, J. P., Ahn, J., Kalk, M. L., McConnell, J. R., Sowers, T., Taylor, K. C.,  
24 White, J. W. C. and Brook, E. J.: Centennial-scale changes in the global carbon cycle during  
25 the last deglaciation, *Nature*, 514(7524), 616–619, doi:10.1038/nature13799, 2014.
- 26 Marshall, S. J. and Clarke, G. K. C.: Modeling North American Freshwater Runoff through  
27 the Last Glacial Cycle, *Quat. Res.*, 52(3), 300–315, doi:10.1006/qres.1999.2079, 1999.
- 28 Martrat, B., Grimalt, J. O., Lopez Martinez, C., Cacho, I., Sierro, F. J., Flores, J. A., Zahn, R.,  
29 Canals, M., Curtis, J. H. and Hodell, D. A.: Abrupt Temperature Changes in the Western

- 1 Mediterranean over the Past 250,000 Years, *Science*, 306(5702), 1762–1765,  
2 doi:10.1126/science.1101706, 2004.
- 3 Martrat, B., Grimalt, J. O., Shackleton, N. J., Abreu, L. de, Hutterli, M. A. and Stocker, T. F.:  
4 Four Climate Cycles of Recurring Deep and Surface Water Destabilizations on the Iberian  
5 Margin, *Science*, 317(5837), 502–507, doi:10.1126/science.1139994, 2007.
- 6 Masson-Delmotte, V., Kageyama, M., Braeck, P., Charbit, S., Krinner, G., Ritz, C.,  
7 Guilyardi, E., Jouzel, J., Abe-Ouchi, A., Crucifix, M., Gladstone, R. M., Hewitt, C. D., Kitoh,  
8 A., LeGrande, A. N., Martí, O., Merkel, U., Motoi, T., Ohgaito, R., Otto-Bliesner, B., Peltier,  
9 W. R., Ross, I., Valdes, P. J., Vettoretti, G., Weber, S. L., Wolk, F. and Yu, Y.: Past and  
10 future polar amplification of climate change: climate model intercomparisons and ice core  
11 constraints, *Clim. Dyn.*, 26(5), 513–529, doi:10.1007/s00382-005-0081-9, 2006.
- 12 Masson-Delmotte, V., Schulz, M., Abe-Ouchi, A., Beer, J., Ganopolski, A., González-Rouco,  
13 J. F., Jansen, E., Lambeck, K., Luterbacher, J., Naish, T., Osborn, T., Otto-Bliesner, B.,  
14 Quinn, T., Ramesh, R., Rojas, M., Shao, X. and Timmermann, A.: Information from  
15 Paleoclimate Archives, in *Climate Change 2013: The Physical Science Basis. Contribution of  
Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate  
Change*, edited by T. F. Stocker, D. Qin, G. K. Plattner, M. Tignor, S. K. Allen, J. Boschung,  
A. Nauels, Y. Xia, V. Bex, and P. M. Midgley, pp. 383–464, Cambridge University Press,  
Cambridge, United Kingdom and New York, NY, USA. [online] Available from:  
[http://www.ipcc.ch/pdf/assessment-report/ar5/wg1/WG1AR5\\_Chapter05\\_FINAL.pdf](http://www.ipcc.ch/pdf/assessment-report/ar5/wg1/WG1AR5_Chapter05_FINAL.pdf)  
(Accessed 20 October 2015), 2013.
- 22 McManus, J. F., Francois, R., Gherardi, J. M., Keigwin, L. D. and Brown-Leger, S.: Collapse  
23 and rapid resumption of Atlantic meridional circulation linked to deglacial climate changes,  
24 *Nature*, 428(6985), 834–837, doi:10.1038/nature02494, 2004.
- 25 Menzel, L., Timmermann, A., Timm, O. E. and Mouchet, A.: Climate and biogeochemical  
26 response to a rapid melting of the West Antarctic Ice Sheet during interglacials and  
27 implications for future climate, *Paleoceanography*, 25(4), PA4231,  
28 doi:10.1029/2009PA001892, 2010.

- 1 Menziel, L., Timmermann, A., Timm, O. E. and Mouchet, A.: Deconstructing the Last  
2 Glacial termination: the role of millennial and orbital scale forcings, *Quat. Sci. Rev.*, 30(9–  
3 10), 1155–1172, doi:10.1016/j.quascirev.2011.02.005, 2011.
- 4 Mix, A. C., Bard, E. and Schneider, R.: Environmental processes of the ice age: land, oceans,  
5 glaciers (EPILOG), *Quat. Sci. Rev.*, 20(4), 627–657, doi:10.1016/S0277-3791(00)00145-1,  
6 2001.
- 7 Monnin, E., Steig, E. J., Siegenthaler, U., Kawamura, K., Schwander, J., Stauffer, B., Stocker,  
8 T. F., Morse, D. L., Barnola, J. M., Bellier, B., Raynaud, D. and Fischer, H.: Evidence for  
9 substantial accumulation rate variability in Antarctica during the Holocene, through  
10 synchronization of CO<sub>2</sub> in the Taylor Dome, Dome C and DML ice cores, *Earth Planet. Sci.  
11 Lett.*, 224(1–2), 45–54, doi:10.1016/j.epsl.2004.05.007, 2004.
- 12 Otto Bliesner, B. L., Schneider, R., Brady, E. C., Kucera, M., Abe-Ouchi, A., Bard, E.,  
13 Braeckman, P., Crucifix, M., Hewitt, C. D., Kageyama, M., Marti, O., Paul, A., Rosell Melé,  
14 A., Waelbroeck, C., Weber, S. L., Weinelt, M. and Yu, Y.: A comparison of PMIP2 model  
15 simulations and the MARGO proxy reconstruction for tropical sea surface temperatures at last  
16 glacial maximum, *Clim. Dyn.*, 32(6), 799–815, doi:10.1007/s00382-008-0509-0, 2009.
- 17 Otto Bliesner, B. L., Russell, J. M., Clark, P. U., Liu, Z., Overpeck, J. T., Konecky, B.,  
18 deMenocal, P., Nicholson, S. E., He, F. and Lu, Z.: Coherent changes of southeastern  
19 equatorial and northern African rainfall during the last deglaciation, *Science*, 346(6214),  
20 1223–1227, doi:10.1126/science.1259531, 2014.
- 21 Peltier, W. R.: Global glacial isostasy and the surface of the Ice Age Earth: The ICE-5G  
22 (VM2) model and GRACE, *Annu. Rev. Earth Planet. Sci.*, 32(1), 111–149,  
23 doi:10.1146/annurev.earth.32.082503.144359, 2004.
- 24 Peltier, W. R.: On the hemispheric origins of meltwater pulse 1a, *Quat. Sci. Rev.*, 24(14–15),  
25 1655–1671, doi:10.1016/j.quascirev.2004.06.023, 2005.
- 26 Peltier, W. R. and Fairbanks, R. G.: Global glacial ice volume and Last Glacial Maximum  
27 duration from an extended Barbados sea level record, *Quat. Sci. Rev.*, 25(23–24), 3322–3337,  
28 doi:10.1016/j.quascirev.2006.04.010, 2006.

- 1 Peltier, W. R. and Vettoretti, G.: Dansgaard-Oeschger oscillations predicted in a  
2 comprehensive model of glacial climate: A “kicked” salt oscillator in the Atlantic, *Geophys.*  
3 *Res. Lett.*, 41(20), 2014GL061413, doi:10.1002/2014GL061413, 2014.
- 4 Peltier, W. R., Argus, D. F. and Drummond, R.: Space geodesy constrains ice age terminal  
5 deglaciation: The global ICE 6G\_C (VM5a) model, *J. Geophys. Res. Solid Earth*,  
6 2014JB011176, doi:10.1002/2014JB011176, 2015.
- 7 Petit, J.-R., Jouzel, J., Raynaud, D., Barkov, N. I., Barnola, J. M., Basile, I., Bender, M.,  
8 Chappellaz, J., Davis, M., Delaygue, G., Delmotte, M., Kotlyakov, V. M., Legrand, M.,  
9 Lipenkov, V. Y., Lorius, C., Pépin, L., Ritz, C., Saltzman, E. and Stievenard, M.: Climate and  
10 atmospheric history of the past 420,000 years from the Vostok ice core, *Antarctica, Nature*,  
11 399(6735), 429–436, doi:10.1038/20859, 1999.
- 12 Philippon, G., Ramstein, G., Charbit, S., Kageyama, M., Ritz, C. and Dumas, C.: Evolution of  
13 the Antarctic ice sheet throughout the last deglaciation: A study with a new coupled climate—  
14 north and south hemisphere ice sheet model, *Earth Planet. Sci. Lett.*, 248(3–4), 750–758,  
15 doi:10.1016/j.epsl.2006.06.017, 2006.
- 16 PMIP Last Deglaciation Working Group: PMIP3 Last Deglaciation experiment design,  
17 [online] Available from: <https://wiki.lsce.ipsl.fr/pmip3/doku.php/pmip3:wg:deglac:index> (Accessed 9 April 2015), 2015.
- 19 PMIP LGM Working Group: PMIP3-CMIP5 Last Glacial Maximum experiment design,  
20 [online] Available from: <https://wiki.lsce.ipsl.fr/pmip3/doku.php/pmip3:design:21k:final> (Accessed 9 April  
21 2015), 2010.
- 22 PMIP LGM Working Group: PMIP CMIP6 Last Glacial Maximum experiment design,  
23 [online] Available from:  
24 <https://wiki.lsce.ipsl.fr/pmip3/doku.php/pmip3:cmip6:design:21k:index> (Accessed 9 April  
25 2015), 2015.
- 26 PMIP website: Paleocean Model Intercomparison Project, [online] Available from:  
27 <http://pmip.lsce.ipsl.fr/> (Accessed 13 November 2014), 2007.
- 28 Putnam, A. E., Denton, G. H., Schaefer, J. M., Barrell, D. J. A., Andersen, B. G., Finkel, R.  
29 C., Schwartz, R., Doughty, A. M., Kaplan, M. R. and Schlüchter, C.: Glacier advance in

- 1 southern middle latitudes during the Antarctic Cold Reversal, *Nat. Geosci.*, 3(10), 700–704,  
2 doi:10.1038/ngeo962, 2010.
- 3 Rahmstorf, S.: Bifurcations of the Atlantic thermohaline circulation in response to changes in  
4 the hydrological cycle, *Nature*, 378(6553), 145–149, doi:10.1038/378145a0, 1995.
- 5 Rahmstorf, S.: On the freshwater forcing and transport of the Atlantic thermohaline  
6 circulation, *Clim. Dyn.*, 12(12), 799–811, doi:10.1007/s003820050144, 1996.
- 7 Rasmussen, S. O., Andersen, K. K., Svensson, A. M., Steffensen, J. P., Vinther, B. M.,  
8 Clausen, H. B., Siggaard Andersen, M. L., Johnsen, S. J., Larsen, L. B., Dahl Jensen, D.,  
9 Bigler, M., Röhlisberger, R., Fischer, H., Goto Azuma, K., Hansson, M. E. and Ruth, U.: A  
10 new Greenland ice core chronology for the last glacial termination, *J. Geophys. Res.  
Atmospheres*, 111(D6), D06102, doi:10.1029/2005JD006079, 2006.
- 12 Roberts, N. L., Piotrowski, A. M., McManus, J. F. and Keigwin, L. D.: Synchronous  
13 Deglacial Overturning and Water Mass Source Changes, *Science*, 327(5961), 75–78,  
14 doi:10.1126/science.1178068, 2010.
- 15 Roche, D. M., Renssen, H., Weber, S. L. and Goosse, H.: Could meltwater pulses have been  
16 sneaked unnoticed into the deep ocean during the last glacial?, *Geophys. Res. Lett.*, 34(24),  
17 n/a–n/a, doi:10.1029/2007GL032064, 2007.
- 18 Roche, D. M., Renssen, H., Paillard, D. and Levavasseur, G.: Deciphering the spatio-temporal  
19 complexity of climate change of the last deglaciation: a model analysis, *Clim Past*, 7(2), 591–  
20 602, doi:10.5194/cp-7-591-2011, 2011.
- 21 Rogerson, M., Colmenero Hidalgo, E., Levine, R. C., Rohling, E. J., Voelker, A. H. L., Bigg,  
22 Schönfeld, J., Cacho, I., Sierro, F. J., Löwemark, L., Reguera, M. I., Abreu, L. de and  
23 Garrick, K.: Enhanced Mediterranean-Atlantic exchange during Atlantic freshening phases,  
24 *Geochem. Geophys. Geosystems*, 11, 22 PP., doi:201010.1029/2009GC002931, 2010.
- 25 Rother, H., Fink, D., Shulmeister, J., Mifsud, C., Evans, M. and Pugh, J.: The early rise and  
26 late demise of New Zealand's last glacial maximum, *Proc. Natl. Acad. Sci.*, 111(32), 11630–  
27 11635, doi:10.1073/pnas.1401547111, 2014.
- 28 Rubino, M., Etheridge, D. M., Trudinger, C. M., Allison, C. E., Battle, M. O., Langenfelds, R.  
29 L., Steele, L. P., Curran, M., Bender, M., White, J. W. C., Jenk, T. M., Blunier, T. and

- 1 Francey, R. J.: A revised 1000 year atmospheric  $\delta^{13}\text{C}$  CO<sub>2</sub> record from Law Dome and  
2 South Pole, Antarctica, *J. Geophys. Res. Atmospheres*, 118(15), 8482–8499,  
3 doi:10.1002/jgrd.50668, 2013.
- 4 Schilt, A., Baumgartner, M., Schwander, J., Buiron, D., Capron, E., Chappellaz, J.,  
5 Louergue, L., Schüpbach, S., Spahni, R., Fischer, H. and Stocker, T. F.: Atmospheric nitrous  
6 oxide during the last 140,000 years, *Earth Planet. Sci. Lett.*, 300(1–2), 33–43,  
7 doi:10.1016/j.epsl.2010.09.027, 2010.
- 8 Schmidt, G. A., Jungclaus, J. H., Ammann, C. M., Bard, E., Braconnot, P., Crowley, T. J.,  
9 Delaygue, G., Joos, F., Krivova, N. A., Muscheler, R., Otto Bliesner, B. L., Pongratz, J.,  
10 Shindell, D. T., Solanki, S. K., Steinhilber, F. and Vieira, L. E. A.: Climate forcing  
11 reconstructions for use in PMIP simulations of the last millennium (v1.0), *Geosci Model Dev.*,  
12 4(1), 33–45, doi:10.5194/gmd.4.33–2011, 2011.
- 13 Schmittner, A., Green, J. a. M. and Wilmes, S. B.: Glacial ocean overturning intensified by  
14 tidal mixing in a global circulation model, *Geophys. Res. Lett.*, 42(10), 2015GL063561,  
15 doi:10.1002/2015GL063561, 2015.
- 16 Severinghaus, J. P. and Brook, E. J.: Abrupt Climate Change at the End of the Last Glacial  
17 Period Inferred from Trapped Air in Polar Ice, *Science*, 286(5441), 930–934,  
18 doi:10.1126/science.286.5441.930, 1999.
- 19 Shakun, J. D., Clark, P. U., He, F., Marcott, S. A., Mix, A. C., Liu, Z., Otto Bliesner, B.,  
20 Schmittner, A. and Bard, E.: Global warming preceded by increasing carbon dioxide  
21 concentrations during the last deglaciation, *Nature*, 484(7392), 49–54,  
22 doi:10.1038/nature10915, 2012.
- 23 Shennan, I.: Global meltwater discharge and the deglacial sea level record from northwest  
24 Scotland, *J. Quat. Sci.*, 14(7), 715–719, doi:10.1002/(SICI)1099-  
25 1417(199912)14:7<715::AID-JQS511>3.0.CO;2-G, 1999.
- 26 Shennan, I. and Milne, G.: Sea-level observations around the Last Glacial Maximum from the  
27 Bonaparte Gulf, NW Australia, *Quat. Sci. Rev.*, 22(14), 1543–1547, doi:10.1016/S0277-  
28 3791(03)00088-X, 2003.

- 1 Siegenthaler, U., Stocker, T. F., Monnin, E., Lüthi, D., Schwander, J., Stauffer, B., Raynaud,  
2 D., Barnola, J. M., Fischer, H., Masson Delmotte, V. and Jouzel, J.: Stable Carbon  
3 Cycle&#150;Climate Relationship During the Late Pleistocene, *Science*, 310(5752), 1313–  
4 1317, doi:10.1126/science.1120130, 2005.
- 5 Simonsen, S. B., Johnsen, S. J., Popp, T. J., Vinther, B. M., Gkinis, V. and Steen Larsen, H.  
6 C.: Past surface temperatures at the NorthGRIP drill site from the difference in firn diffusion  
7 of water isotopes, *Clim Past*, 7(4), 1327–1335, doi:10.5194/cp-7-1327-2011, 2011.
- 8 Smith, R. S. and Gregory, J. M.: A study of the sensitivity of ocean overturning circulation  
9 and climate to freshwater input in different regions of the North Atlantic, *Geophys. Res. Lett.*,  
10 36(15), L15701, doi:10.1029/2009GL038607, 2009.
- 11 Spahni, R., Chappellaz, J., Stocker, T. F., Louergue, L., Hausmann, G., Kawamura, K.,  
12 Flückiger, J., Schwander, J., Raynaud, D., Masson Delmotte, V. and Jouzel, J.: Atmospheric  
13 Methane and Nitrous Oxide of the Late Pleistocene from Antarctic Ice Cores, *Science*,  
14 310(5752), 1317–1321, doi:10.1126/science.1120132, 2005.
- 15 Stanford, J. D., Hemingway, R., Rohling, E. J., Challener, P. G., Medina Elizalde, M. and  
16 Lester, A. J.: Sea level probability for the last deglaciation: A statistical analysis of far-field  
17 records, *Glob. Planet. Change*, 79(3–4), 193–203, doi:10.1016/j.gloplacha.2010.11.002, 2011.
- 18 Steffensen, J. P., Andersen, K. K., Bigler, M., Clausen, H. B., Dahl Jensen, D., Fischer, H.,  
19 Goto Azuma, K., Hansson, M., Johnsen, S. J., Jouzel, J., Masson Delmotte, V., Popp, T.,  
20 Rasmussen, S. O., Röhlisberger, R., Ruth, U., Stauffer, B., Siggaard Andersen, M. L.,  
21 Sveinbjörnsdóttir, Á. E., Svensson, A. and White, J. W. C.: High Resolution Greenland Ice  
22 Core Data Show Abrupt Climate Change Happens in Few Years, *Science*, 321(5889), 680–  
23 684, doi:10.1126/science.1157707, 2008.
- 24 Stenni, B., Buiron, D., Frezzotti, M., Albani, S., Barbante, C., Bard, E., Barnola, J. M.,  
25 Baroni, M., Baumgartner, M., Bonazza, M., Capron, E., Castellano, E., Chappellaz, J.,  
26 Delmonte, B., Falourd, S., Genoni, L., Iacumin, P., Jouzel, J., Kipfstuhl, S., Landais, A.,  
27 Lemieux-Dufour, B., Maggi, V., Masson Delmotte, V., Mazzola, C., Minster, B., Montagnat,  
28 M., Mulvaney, R., Narcisi, B., Oerter, H., Parrenin, F., Petit, J. R., Ritz, C., Scarchilli, C.,  
29 Schilt, A., Schüpbach, S., Schwander, J., Selmo, E., Severi, M., Stocker, T. F. and Udisti, R.:

- 1 Expression of the bipolar see saw in Antarctic climate records during the last deglaciation,  
2 *Nat. Geosci.*, 4(1), 46–49, doi:10.1038/ngeo1026, 2011.
- 3 Stoecker, T. F.: The Seesaw Effect, *Science*, 282(5386), 61–62,  
4 doi:10.1126/science.282.5386.61, 1998.
- 5 Strelin, J. A., Denton, G. H., Vandergoes, M. J., Ninnemann, U. S. and Putnam, A. E.:  
6 Radioisotope chronology of the late glacial Puerto Bandera moraines, Southern Patagonian  
7 Icefield, Argentina, *Quat. Sci. Rev.*, 30(19–20), 2551–2569,  
8 doi:10.1016/j.quascirev.2011.05.004, 2011.
- 9 Svendsen, J. I., Alexanderson, H., Astakhov, V. I., Demidov, I., Dowdeswell, J. A., Funder,  
10 S., Gataullin, V., Henriksen, M., Hjort, C., Houmark-Nielsen, M., Hubberten, H. W.,  
11 Ingólfsson, Ó., Jakobsson, M., Kjær, K. H., Larsen, E., Lokrantz, H., Lunkka, J. P., Lyså, A.,  
12 Mangerud, J., Matiouchkov, A., Murray, A., Möller, P., Niessen, F., Nikolskaya, O., Polyak,  
13 L., Saarnisto, M., Siegert, C., Siegert, M. J., Spielhagen, R. F. and Stein, R.: Late Quaternary  
14 ice sheet history of northern Eurasia, *Quat. Sci. Rev.*, 23(11–13), 1229–1271,  
15 doi:10.1016/j.quascirev.2003.12.008, 2004.
- 16 Tarasov, L. and Peltier, W. R.: Terminating the 100-kyr ice age cycle, *J. Geophys. Res.  
Atmospheres*, 102(D18), 21665–21693, doi:10.1029/97JD01766, 1997.
- 18 Tarasov, L. and Peltier, W. R.: Greenland glacial history and local geodynamic consequences,  
19 *Geophys. J. Int.*, 150(1), 198–229, doi:10.1046/j.1365-246X.2002.01702.x, 2002.
- 20 Tarasov, L. and Peltier, W. R.: A geophysically constrained large ensemble analysis of the  
21 deglacial history of the North American ice sheet complex, *Quat. Sci. Rev.*, 23(3–4), 359–  
22 388, doi:10.1016/j.quascirev.2003.08.004, 2004.
- 23 Tarasov, L. and Peltier, W. R.: Arctic freshwater forcing of the Younger Dryas cold reversal,  
24 *Nature*, 435(7042), 662–665, doi:10.1038/nature03617, 2005.
- 25 Tarasov, L. and Peltier, W. R.: A calibrated deglacial drainage chronology for the North  
26 American continent: evidence of an Arctic trigger for the Younger Dryas, *Quat. Sci. Rev.*,  
27 25(7–8), 659–688, doi:10.1016/j.quascirev.2005.12.006, 2006.

- 1 Tarasov, L., Dyke, A. S., Neal, R. M. and Peltier, W. R.: A data calibrated distribution of  
2 deglacial chronologies for the North American ice complex from glaciological modeling,  
3 *Earth Planet. Sci. Lett.*, 315–316, 30–40, doi:10.1016/j.epsl.2011.09.010, 2012.
- 4 Taylor, K. E., Stouffer, R. J. and Meehl, G. A.: An Overview of CMIP5 and the Experiment  
5 Design, *Bull. Am. Meteorol. Soc.*, 93(4), 485–498, doi:10.1175/BAMS-D-11-00094.1, 2011a.
- 6 Taylor, K. E., Stouffer, R. J. and Meehl, G. A.: A Summary of the CMIP5 Experiment  
7 Design, [online] Available from: [http://cmip-](http://cmip-pcmdi.llnl.gov/cmip5/docs/Taylor_CMIP5_design.pdf)  
8 [pcmdi.llnl.gov/cmip5/docs/Taylor\\_CMIP5\\_design.pdf](http://cmip-pcmdi.llnl.gov/cmip5/docs/Taylor_CMIP5_design.pdf) (Accessed 13 November 2014b), 2011.
- 9 Teller, J. T., Leverington, D. W. and Mann, J. D.: Freshwater outbursts to the oceans from  
10 glacial Lake Agassiz and their role in climate change during the last deglaciation, *Quat. Sci.*  
11 *Rev.*, 21(8–9), 879–887, doi:10.1016/S0277-3791(01)00145-7, 2002.
- 12 Thiagarajan, N., Subhas, A. V., Southon, J. R., Eiler, J. M. and Adkins, J. F.: Abrupt pre-  
13 Bolling-Allerod warming and circulation changes in the deep ocean, *Nature*, 511(7507), 75–  
14 78, doi:10.1038/nature13472, 2014.
- 15 Thornalley, D. J. R., McCave, I. N. and Elderfield, H.: Freshwater input and abrupt deglacial  
16 climate change in the North Atlantic, *Paleoceanography*, 25(1), PA1201,  
17 doi:10.1029/2009PA001772, 2010.
- 18 Thornalley, D. J. R., Barker, S., Broecker, W. S., Elderfield, H. and McCave, I. N.: The  
19 Deglacial Evolution of North Atlantic Deep Convection, *Science*, 331(6014), 202–205,  
20 doi:10.1126/science.1196812, 2011.
- 21 Tierney, J. E., Russell, J. M., Huang, Y., Damsté, J. S. S., Hopmans, E. C. and Cohen, A. S.:  
22 Northern Hemisphere Controls on Tropical Southeast African Climate During the Past 60,000  
23 Years, *Science*, 322(5899), 252–255, doi:10.1126/science.1160485, 2008.
- 24 Timm, O. and Timmermann, A.: Simulation of the Last 21 000 Years Using Accelerated  
25 Transient Boundary Conditions\*, *J. Clim.*, 20(17), 4377–4401, doi:10.1175/JCLI4237.1,  
26 2007.
- 27 Tjallingii, R., Claussen, M., Staat, J. B. W., Fohlmeister, J., Jahn, A., Bickert, T., Lamy, F.  
28 and Röhl, U.: Coherent high- and low-latitude control of the northwest African hydrological  
29 balance, *Nat. Geosci.*, 1(10), 670–675, doi:10.1038/ngeo289, 2008.

- 1 Veres, D., Bazin, L., Landais, A., Toyé Mahamadou Kele, H., Lemieux-Dudon, B., Parrenin,  
2 F., Martinerie, P., Blayo, E., Blunier, T., Capron, E., Chappellaz, J., Rasmussen, S. O., Severi,  
3 M., Svensson, A., Vinther, B. and Wolff, E. W.: The Antarctic ice core chronology  
4 (AICC2012): an optimized multi-parameter and multi-site dating approach for the last 120  
5 thousand years, *Clim Past*, 9(4), 1733–1748, doi:10.5194/cp-9-1733-2013, 2013.
- 6 Verschuren, D., Sinninghe Damsté, J. S., Moernaut, J., Kristen, I., Blaauw, M., Fagot, M.,  
7 Haug, G. H., Geel, B. van, Barist, M. D., Barker, P., Vuille, M., Conley, D. J., Olago, D. O.,  
8 Milne, I., Plessen, B., Eggmont, H., Wolff, C., Hurrell, E., Ossebaar, J., Lyaruu, A., Plicht,  
9 J. van der, Cumming, B. F., Brauer, A., Rucina, S. M., Russell, J. M., Keppens, E., Hus, J.,  
10 Bradley, R. S., Leng, M., Mingram, J. and Nowaczyk, N. R.: Half precessional dynamics of  
11 monsoon rainfall near the East African Equator, *Nature*, 462(7273), 637–641,  
12 doi:10.1038/nature08520, 2009.
- 13 Vettoretti, G. and Peltier, W. R.: Interhemispheric air temperature phase relationships in the  
14 nonlinear Dansgaard-Oeschger oscillation, *Geophys. Res. Lett.*, 42(4), 2014GL062898,  
15 doi:10.1002/2014GL062898, 2015.
- 16 Waelbroeck, C., Labeyrie, L., Duplessy, J. C., Guiot, J., Labracherie, M., Leclaire, H. and  
17 Duprat, J.: Improving past sea surface temperature estimates based on planktonic fossil  
18 faunas, *Paleoceanography*, 13(3), 272–283, doi:10.1029/98PA00071, 1998.
- 19 Weaver, A. J., Saenko, O. A., Clark, P. U. and Mitrovica, J. X.: Meltwater Pulse 1A from  
20 Antarctica as a Trigger of the Bølling Allerød Warm Interval, *Science*, 299(5613), 1709–  
21 1713, doi:10.1126/science.1081002, 2003.
- 22 Weber, S. L., Drijfhout, S. S., Abe-Ouchi, A., Crucifix, M., Eby, M., Ganopolski, A.,  
23 Murakami, S., Otto-Bliesner, B. and Peltier, W. R.: The modern and glacial overturning  
24 circulation in the Atlantic ocean in PMIP coupled model simulations, *Clim Past*, 3(1), 51–64,  
25 doi:10.5194/cp-3-51-2007, 2007.
- 26 Weijers, J. W. H., Schefuß, E., Schouten, S. and Damsté, J. S. S.: Coupled Thermal and  
27 Hydrological Evolution of Tropical Africa over the Last Deglaciation, *Science*, 315(5819),  
28 1701–1704, doi:10.1126/science.1138131, 2007.
- 29 Whitehouse, P. L., Bentley, M. J. and Le Brocq, A. M.: A deglacial model for Antarctica:  
30 geological constraints and glaciological modelling as a basis for a new model of Antarctic

1 glacial isostatic adjustment, *Quat. Sci. Rev.*, 32, 1–24, doi:10.1016/j.quascirev.2011.11.016,  
2 2012.

3 Yokoyama, Y., Esat, T. M. and Lambeck, K.: Last glacial sea level change deduced from  
4 uplifted coral terraces of Huon Peninsula, Papua New Guinea, *Quat. Int.*, 83–85, 275–283,  
5 doi:10.1016/S1040-6182(01)00045-3, 2001a.

6 Yokoyama, Y., De Deckker, P., Lambeck, K., Johnston, P. and Fifield, L. K.: Sea level at the  
7 Last Glacial Maximum: evidence from northwestern Australia to constrain ice volumes for  
8 oxygen isotope stage 2, *Palaeogeogr. Palaeoclimatol. Palaeoecol.*, 165(3–4), 281–297,  
9 doi:10.1016/S0031-0182(00)00164-4, 2001b.

10

## 1 Tables

Formatted: Heading 6, No page break before

2 Table 1. Summary of recommended model boundary conditions to spin up the last  
3 deglaciation Core experiment [version 1](#) (pre 21 ka); see text for details. Participants are not  
4 required to follow the recommendation for these boundary conditions, but must document the  
5 method used, including information on the simulation's state of spinup at the point when the  
6 Core is started. Data are available from PMIP4 [Last dDeglaciation Working Group Wiki](#)  
7 ([PMIP Last Deglaciation Working Group, 2016](#)):  
8 <https://wiki.lsce.ipsl.fr/pmip3/doku.php/pmip3:wg-deglai:index>. Boundary condition group  
9 headings are in bold.

Spinup type	Boundary condition	Description
<b>Last Glacial Maximum (LGM; 21 ka)</b>	<b>Insolation</b>	Preindustrial (e.g. <a href="#">1361.0 ± 0.54365 W m⁻²</a> )
	Solar constant	0.018994
	Eccentricity	22.949°
	Obliquity	114.42°
	Perihelion–180°	Noon, 21 <sup>st</sup> March
	<b>Trace gases</b>	
	Carbon dioxide (CO <sub>2</sub> )	190 ppm
	Methane (CH <sub>4</sub> )	375 ppb
	Nitrous oxide (N <sub>2</sub> O)	200 ppb
	Chlorofluorocarbon (CFC)	0
	Ozone (O <sub>3</sub> )	Preindustrial (e.g. 10 DU)
	<b>Ice sheets, orography and coastlines</b>	21 ka data from either: - ICE-6G_C (references in text) - GLAC-1D (references in text)
	<b>Bathymetry</b>	Keep consistent with the coastlines, using either: - Data associated with the ice sheet - Preindustrial bathymetry
	<b>Global ocean salinity</b>	+ 1 psu, relative to preindustrial
<b>Transient orbit and trace gases (26-21 ka)</b>	<b>Orbital parameters</b>	All orbital parameters should be transient, as per Berger (1978) 26-21 ka
	<b>Trace gases</b>	Adjusted to the AICC2012 (Veres et al., 2013)
	Carbon dioxide (CO <sub>2</sub> )	Transient, as per Bereiter et al. (2015)
	Methane (CH <sub>4</sub> )	Transient, as per Loulergue et al. (2008)
	Nitrous oxide (N <sub>2</sub> O)	Transient, as per Schilt et al. (2010)
	<b>All others</b>	As per LGM (21 ka) spinup type.


Formatted: Font: 10 pt

1 **Table 2.** Summary of required model boundary conditions for the last deglaciation Core  
 2 experiment 21-9 ka [version 1](#); optional boundary conditions are labelled as such. Data are  
 3 available from PMIP4 [Last deglaciation Working Group Wiki](#):  
 4 [https://wiki.lsce.ipsl.fr/pmip3/doku.php/pmip3:wg:degl:index\(PMIP Last Deglaciation Working Group, 2016\)](https://wiki.lsce.ipsl.fr/pmip3/doku.php/pmip3:wg:degl:index(PMIP Last Deglaciation Working Group, 2016)). See text for details. Boundary condition group headings are in bold.

Boundary condition	Description
<b>Initial conditions (pre 21 ka)</b>	Recommended (optional) to use either: <ul style="list-style-type: none"> <li>- Last Glacial Maximum (LGM; 21 ka) equilibrium simulation, including +1 psu global ocean salinity</li> <li>- Transient orbit and trace gases (26-21 ka) and all other boundary conditions fixed as per equilibrium LGM</li> </ul> See Table 1 for details. The method must be documented, including information on the state of spinup
<b>Insolation</b>	Solar constant Orbital parameters
	Preindustrial (e.g. $136.1.0 \pm 0.55$ W m <sup>-2</sup> ) Transient, as per Berger (1978)
<b>Trace gases</b>	Adjusted to the AICC2012 age model (Veres et al., 2013): Carbon dioxide (CO <sub>2</sub> ) Methane (CH <sub>4</sub> ) Nitrous oxide (N <sub>2</sub> O) Chlorofluorocarbon (CFC) Ozone (O <sub>3</sub> )
	Transient, as per Bereiter et al. (2015) Transient, as per Loulergue et al. (2008) Transient, as per Schilt et al. (2010) 0 Preindustrial (e.g. 10 DU)
<b>Ice sheet</b>	Transient, with a choice of either : <ul style="list-style-type: none"> <li>- ICE-6G_C reconstruction (references in text)</li> <li>- GLAC-1D reconstruction (references in text)</li> </ul> How often to update the ice sheet is optional
<b>Orography and coastlines</b>	Transient. To be consistent with the choice of ice sheet. Orography is updated on the same timestep as the ice sheet. It is optional how often the land-sea mask is updated, but ensure consistency with the ice sheet reconstruction is maintained
<b>Bathymetry</b>	Keep consistent with the coastlines and otherwise use either: <ul style="list-style-type: none"> <li>- Transient data associated with the chosen ice sheet; it is optional how often the bathymetry is updated</li> <li>- Preindustrial bathymetry</li> </ul>
<b>River routing</b>	Ensure that rivers reach the coastline It is recommended (optional) to use one of the following: <ul style="list-style-type: none"> <li>- Preindustrial configuration for the model</li> <li>- Transient routing provided with the ice sheet reconstruction (<a href="#">if available</a>)</li> <li>- Manual/model calculation of river network to match topography</li> </ul>
<b>Freshwater fluxes</b>	At participant discretion. Three options are: <i>melt-uniform</i> , <i>melt-routed</i> and <i>no-melt</i> (see text). It is recommended (optional) to run at least one Core simulation with a scenario consistent with the chosen ice sheet reconstruction to conserve salinity (e.g. as provided). See text for full details (Sect. 2.5)
<b>Other (optional)</b>	Vegetation and land cover Aerosols (dust)
	Prescribed preindustrial cover or dynamic vegetation model Prescribed preindustrial distribution or prognostic aerosols

## 1 Figures


2 **Figure 1.** The last deglaciation; 3 forcings and events. (a) The three 4 phases of the Core experiment 5 [version 1](#) (Sect. 4). (b) Climate 6 events/periods discussed in the 7 text; Last Glacial Maximum 8 (LGM; 23-19 ka as according to 9 the EPILOG definition; Mix et al., 10 2001), Heinrich Stadial 1 (HS1), 11 Heinrich Event 1 (H1), Bølling 12 Warming (BW) and Meltwater 13 Pulse 1a (MWP1a), Antarctic 14 Cold Reversal (ACR) and the 15 Younger Dryas cooling (YD). (c) 16 June insolation at 60° N and 17 December insolation at 60° S 18 (Berger, 1978). (d) Atmospheric 19 carbon dioxide concentration 20 (recent composite of EPICA 21 Dome C, Vostok, Taylor Dome, 22 Siple Dome and West Antarctic 23 Ice Sheet Divide records, 24 Antarctica; Bereiter et al., 2015); 25 black dashed line shows 26 preindustrial concentration. (e) 27 Atmospheric methane concentration 28 (EPICA Dome C, 29 Antarctica; Loulergue et al., 30 2008); green dashed line shows



Formatted: Heading 6, No page break before

1 preindustrial concentration. (f) Atmospheric nitrous oxide concentration (Talos Dome,  
2 Antarctica; Schilt et al., 2010); brown dashed line shows preindustrial concentration. (g)  
3 Volume of the ice sheets according to the ICE-6G\_C reconstruction (solid lines; Argus et al.,  
4 2014; Peltier et al., 2015) and the GLAC-1D reconstruction (dashed lines; Briggs et al., 2014;  
5 Tarasov et al., 2012; Tarasov and Peltier, 2002). Associated meltwater scenarios *melt-uniform*  
6 and *melt-routed* (see Sect. 2.5) are consistent with these; all ice mass loss shown is supplied  
7 as freshwater to the ocean. (h) Greenland temperature reconstruction with  $\pm 1 \sigma$  shaded  
8 (averaged GISP2, NEEM and NGRIP records; Buizert et al., 2014). (i) Antarctic  $\delta D$  (EPICA  
9 Dome C; Jouzel et al., 2007). (d)-(f) and (h)-(i) are given on the AICC2012 timescale (Veres  
10 et al., 2013).

11




1   **Figure 2.** Northern Hemisphere ice sheet elevation at 21, 18, 15, 12 and 9 ka for: (a) the ICE  
2   6G\_C reconstruction at 10 arcminute horizontal resolution, elevation is plotted where the  
3   fractional ice mask is more than 0.5 (Peltier et al., 2015); (b) the GLAC-1D reconstruction at

4   1° (longitude) × 0.5° (latitude) horizontal resolution, elevation is plotted where the fractional  
5   ice mask is more than 0.5 (Briggs et al., 2014; Tarasov et al., 2012; Tarasov and Peltier,  
6   2002; this study); (c) the difference in elevation above sea level between the two

7   reconstructions where there is ice present in both (ICE-6G\_C minus GLAC-1D).


8



1 **Figure 3.** Atmospheric trace gases through the last deglaciation from Antarctic ice cores. (a)  
 2 Core experiment carbon dioxide according to a recent composite record from EPICA Dome C  
 3 (EDC), West Antarctic Ice Sheet Divide (WDC), Vostok, Taylor Dome and Siple Dome  
 4 (thick black line; Bereiter et al., 2015), which was produced on the AICC2012 chronology  
 5 (Veres et al., 2013). Also shown for comparison is an older composite record from EDC,  
 6 Vostok and Taylor Dome (thin blue line; Lüthi et al., 2008, adjusted to the AICC2012  
 7 chronology), as well as the original EDC CO<sub>2</sub> record (green line; Monnin et al., 2004) and the  
 8 recent, higher resolution WDC CO<sub>2</sub> record (dark red line; Marcott et al., 2014); which were  
 9

1 incorporated into the newer composite by Bereiter et al. (2015). **(b)** Methane according to the  
2 EPICA Dome C (EDC) record (Loulergue et al., 2008), shown both on the original EDC1  
3 chronology (green line; Spahni et al., 2005) and adjusted to the more recent AICC2012  
4 chronology for the Core experiment (thick black line; Veres et al., 2013). **(c)** Nitrous oxide  
5 according to the Talos Dome (TALDICE) record (Schilt et al., 2010), adjusted to the  
6 AICC2012 chronology for the Core experiment (thick black line; Veres et al., 2013). For  
7 comparison, the earlier EPICA Dome C (EDC) record on the EDC1 chronology is also shown  
8 (green line; Spahni et al., 2005). The nearest measured N<sub>2</sub>O concentration to 21 ka is from  
9 21.089 ka; hence the small offset between the slightly earlier concentration (187 ppb) used for  
10 the Core and the interpolated value plotted at 21 ka. For **(a)-(c)** 21 ka concentrations  
11 according to the AICC2012 age model (red dots) are shown in contrast to previous PMIP3  
12 LGM concentrations (blue dots; PMIP LGM Working Group, 2010). If using an equilibrium-  
13 type spinup for the start of a transient Core simulation at 21 ka (Sect. 2.1.1), use 190 ppm  
14 CO<sub>2</sub>, 375 ppb CH<sub>4</sub> and 200 ppb N<sub>2</sub>O.

15



1  
2 **Figure 4.** Southern Hemisphere ice sheet elevation at 21, 12 and 9 ka for: (a) the ICE-6G\_C  
3 reconstruction at 10 arcminute horizontal resolution, ice elevation is plotted where the  
4 fractional ice mask is more than 0.5 (Argus et al., 2014; Peltier et al., 2015); (b) the GLAC-  
5 1D reconstruction at  $1^{\circ}$  (longitude)  $\times$   $0.5^{\circ}$  (latitude) horizontal resolution, ice elevation is  
6 plotted where fractional ice mask is more than 0.5 (Briggs et al., 2014; Tarasov et al., 2012;  
7 Tarasov and Peltier, 2002); (c) the difference in elevation above sea level between the two  
8 reconstructions where there is ice present in both (ICE-6G\_C minus GLAC-1D).